Ice nucleating ability of particulate emissions from solid biomassfired cookstoves: an experimental study

Kimmo Korhonen¹, Thomas Bjerring Kristensen², John Falk², Robert Lindgren³, Christina Andersen⁴,

5 Ricardo Luis Carvalho^{3,a}, Vilhelm Malmborg⁴, Axel Eriksson⁴, Christoffer Boman³, Joakim Pagels⁴, Birgitta Svenningsson², Mika Komppula⁵, Kari E.J. Lehtinen¹ and Annele Virtanen¹

¹University of Eastern Finland, Dept. Applied Physics. P.O. box 1627, FI-70211 Kuopio, Finland ²Lund University, Department of Physics, SE-22100, Lund, Sweden

³Umeå University, Thermochemical Energy Conversion Laboratory, SE-90187, Umeå, Sweden
 ⁴Lund University, Ergonomics and Aerosol Technology, Box 118, Lund SE-22100, Sweden
 ⁵Finnish Meteorological Institute, Atmospheric Research Centre of Eastern Finland, P.O. box 1627, FI-70211 Kuopio, Finland

^a Now at: Centre for Environmental and Marine Studies, University of Aveiro, Department of Environment and Planning,

15 PT-3810-193, Aveiro, Portugal

Correspondence to: Kimmo Korhonen (Kimmo.Korhonen@uef.fi)

Abstract. This research was part of the Salutary Umeå Study of Aerosols in Biomass Cookstove Emissions (SUSTAINE) laboratory experiment campaign. We studied ice-nucleating abilities of particulate emissions from solid fuel burning cookstoves, using a portable ice nuclei counter Spectrometer for Ice Nuclei (SPIN). These emissions were generated from

- 20 two traditional cookstove types commonly used for household cooking in sub-Saharan Africa and two advanced gasifier stoves under research to promote sustainable development alternatives. The solid fuels studied included biomass from two different African tree species, Swedish softwood, and agricultural residue products relevant to the region. Measurements were performed with a modified version of the standard water boiling test on polydisperse samples from flue gas during burning and size-selected accumulation mode soot particles from a 15-m³ aerosol-storage chamber. The studied soot particle
- sizes in nm were 250, 260, 300, 350, 400, 450 and 500. From this chamber, the particles were introduced to watersupersaturated freezing conditions (-32 °C to -43 °C) in the SPIN.

Accumulation mode soot particles generally produced an ice-activated fraction of 10^{-3} in temperatures 1-1.5 °C higher than that required for homogeneous freezing at fixed $RH_w = 115\%$. In five special experiments, the combustion performance of one cookstove was intentionally modified. Two of these exhibited a significant increase in the ice-nucleating ability of the

particles, resulting in a 10⁻³ ice activation at up to 5.9 °C higher temperatures than homogeneous freezing and the observed increased ice nucleating ability. We investigated six different physico-chemical properties of the emission particles but found no clear correlation between them and increasing ice-nucleating ability. We conclude that the freshly emitted combustion aerosols form ice via immersion/condensation freezing at temperatures only moderately above homogeneous freezing conditions.

1 Introduction

35

Mixed-phase clouds (MPCs) play an essential role for climate and the hydrological cycle (Korolev et al., 2017; Mülmenstädt et al., 2015). Cloud droplets freeze homogeneously at temperatures near -38 °C (Pruppacher and Klett, 1997), but ice nucleating particles (INPs) may catalyze freezing of supercooled cloud droplets at higher temperatures. A wide range of

40 MPC properties including radiative properties and lifetime are sensitive to the formation of solid ice (Matus and L'Ecuyer, 2017), but there are significant gaps in our knowledge regarding INP concentrations within the MPCs and these important ice formation processes.

INPs active in immersion freezing at a temperature of -30 °C are relatively rare in the lower troposphere with concentrations

- 45 at the order of 0.01 cm⁻³ in many regions (DeMott et al., 2010;). Ambient INPs include dust and biological particles and potentially soot particles (Hoose and Möhler, 2012; Kanji et al., 2017). Soot particles from an acetylene burner, a kerosene burner and a soot generator have been reported to be active in immersion and condensation freezing at temperatures up to 24 °C (DeMott, 1990), -20 °C (Diehl and Mitra, 1998) and -10 °C (Gorbunov et al., 2001), respectively. However, a wide range of soot particles have been reported to be inefficient as INPs in immersion and condensation mode (e.g. Hoose and
- 50 Möhler, 2012), and the available parameterisations used to estimate the soot ice nucleating ability span several orders of magnitude (Vergara-Temprado et al., 2018). The radiative forcing associated with the impact of fossil fuel soot particles on MPCs has been reported to range from about 0.1 to about 1 Wm⁻² depending on their ice nucleating ability, which is uncertain (Yun et al., 2013). The Intergovernmental Panel on Climate Change (IPCC) expressed in their latest assessment report (Boucher et al., 2013) a great need for additional research with respect to the role of soot particles in heterogeneous
- 55 ice nucleation (Bond et al., 2013).

Soot particles are produced from incomplete combustion (combustion with insufficient oxygen supply) and ambient soot particle properties, such as morphology and chemical composition, are highly variable and influenced by the combustion conditions and atmospheric ageing (Ferry et al, 2002; Popovicheva et al., 2008; Koehler et al., 2009; Corbin et al., 2015;

60 Mahrt et al., 2018; Bhandari et al., 2019). It is not entirely clear which soot particle properties influence the ice nucleating ability of the particles. Chemical groups on the soot particle surface with the ability to form hydrogen bonds with water molecules are likely to be of importance (Gorbunov et al., 2001). In addition, the soot particle nanostructure may also be of

importance, with highly ordered graphene structures being more efficient in supporting ice nucleation relative to lowly ordered graphene structures (Häusler et al., 2018).

65

70

75

Biomass combustion for cooking is an important global source of energy, being also the major environmental health risk worldwide (Lim et al., 2012). It has been estimated that approximately 2.8 billion people depend on solid fuel combustion for daily cooking worldwide, mostly in the developing part of the World (Bonjour et al., 2013). Therefore, solid biomass combustion is a significant source of particulate matter (PM) and soot particles on regional to global scales. However, studies of the associated ice nucleating ability are scarce. Ambient measurements indicate that biomass combustion is a source of INPs for MPC conditions (Twohy et al., 2010; McCluskey et al., 2014), Detectable condensation/immersion freezing INP concentrations for a temperature of -30 °C have been reported for simulated wildfires in 9 of 21 and 13 of 22 experiments, respectively (Petters et al., 2009; Levin et al., 2016). Refractory black carbon has been observed to be associated with a significant fraction of the emitted INPs (Levin et al., 2016). Combustion with a modern log wood burner produced particles that acted as INPs at T = -35 °C, but not at T = -30 °C for condensation/immersion freezing (Chou et al., 2013). Huang et al. (2018) modeled the indirect climate impact of solid fueled cookstove aerosol emissions and reported a

- potentially significant climate impact in increase of high clouds in their model runs where black carbon (BC) particles acted as INPs. Therefore, soot particle emissions can potentially affect MPC conditions and thus climate.
- 80 The most common way of using biomass is still in traditional cookstoves/open fires, so called 3-stone fires, used in poorly ventilated spaces, resulting in severe emissions of air pollutants. However, an efficient utilization of modern biomass fuels in efficient biomass cookstoves can constitute an alternative to mitigate household air pollution (HAP) worldwide, and aerosol emission effect on radiative forcing (Carvalho et al. 2016). In Sub-Saharan Africa, the extensive use of firewood and charcoal also leads to considerable forest and soil degradation. Thus, densification of various locally produced biomass feedstocks and residues in the form of fuel pellets can contribute to a more sustainable utilization of biomass in households when comparing with the traditional harvesting and combustion of wood from natural forests (Carvalho et al. 2019). Additionally, the combustion of biomass pellets in micro-gasifier cookstoves appears as a relatively clean solution, and their
- 90 In this study, we present the first measurements of the ice nucleating ability of freshly emitted aerosol particles from biomass-fired cookstoves, we investigated the aerosol emissions from four different cookstoves representing different advances in technology and using a wide range of biomass fuel types. The ice nucleating ability of aerosol particles generated during standard water boiling tests under well-controlled laboratory conditions was measured online with a continuous flow diffusion chamber (CFDC) during the Salutary Umeå Study of Aerosols in Biomass Cookstove Emissions

emission performance can be similar to that achieved by gas stoves (Champion and Grieshop, 2019).

95 (SUSTAINE) campaign.

2 Experiment methodology

2.1 Experiment set-up

Cooking simulations were carried out at the Thermochemical Energy Conversion Laboratory (TEC-lab) of Umeå University, and the sampling set-up and relevant supporting instrumentation for the ice nucleation (IN) experiments are presented in Fig.

- 100 1. The emissions were generated using a modified version of the standardized water boiling test (WBT, version 4.2.4), where the cookstoves and fuels were used to heat up 5 liters of water from room temperature in a metal cooking pot. A typical simulation time was 60-85 min in total, including two phases of the WBT process: cold start (phase duration 15-40 min) and after that simmering (duration 45 min). The cooking pot was set to a designated height for each cook stove used in all but three special experiments. Solid biomass fuels were lit using 12 g of ethanol in each experiment, and water vapor from the
- 105 boiling was diverted away from the sample collector dome to prevent excess humidity entering the sampling lines. The sample aerosol was diluted by 1:100, cooled down to room temperature and dried to $< 10\% RH_w$ using ejector dilution (marked "ED" in Fig. 1) on dry and filtered compressed air to avoid saturation of the analyzers. The supportive data were collected using multiple instruments in parallel with the CFDC instrument. Particle number size distributions were monitored during the WBT using a Fast Particle Analyzer (FPA, Cambustion DMS 500) that was connected to the transient line "A" in
- 110 Fig. 1. The FPA allowed real-time monitoring of the particle size number distribution with its data output resolution of 10 Hz. Other instruments in the line "A" were a high-resolution time-of flight soot particle aerosol mass spectrometer (SP-AMS, Aerodyne Inc.) for measuring the physico-chemical composition of the aerosol particles, and a seven-wavelength aethalometer (Magee Scientific AE33, sampling rate 1 Hz) measuring the absorption by particulate matter. The two latter instruments were connected interchangeably to the chamber sampling line "B" (Fig. 1) for chamber experiments. An SMPS
- 115 system (classifier TSI 3082 equipped with aerosol neutralizer TSI 3088, and condensation particle counter (CPC), TSI 3775) measured particle number size distributions in chamber experiments at scan range in mobility diameters from 15.7 nm to 615.3 nm, with continuously repeated 180-second scans during aerosol injection and sampling at an aerosol to sheath flow ratio of 0.3 vs. 3 litres per min, respectively. Other instruments used in chamber experiments were a cloud condensation nuclei counter (CCNC, Droplet Measurement Technologies) that sampled in parallel with an aerosol particle mass analyzer
- 120 (APM, Kanomax) downstream of a differential mobility analyzer (TSI model 3071, marked DMA_{CCN} in Fig. 1) whose voltage was systematically varied for obtaining APM and CCNC scans at mobility diameters of 65 nm, 100 nm, 200 nm and at times also 350 nm. The time resolution of this measurement was approximately seven minutes between changes in DMA_{CCN} voltage, which enabled obtainment of one APM spectrum and 1-2 full scan cycles on the CCNC.
- 125 The SP-AMS data were measured using dual vaporizer measurements employing flash vaporization at 600 °C and infrared (IR) laser vaporization, followed by 70 eV electron ionization (Onasch et al. 2012). The IR laser vaporization allows detection of constituents from refractory black carbon (rBC) cores. Organic aerosol (OA) measured in this way includes nonrefractory OA and refractory oxygen containing species associated with the rBC core (Corbin et al. 2015). The OA was

derived from high resolution data using the software PIKA 1.22F and included the range 13-313 amu. A relative ionisation

- 130 efficiency of 1.4 and a collection efficiency of 1 were assumed. Information related to the soot maturity (Malmborg et al. 2019) was obtained from analysis of the refractory carbon fragments. The ratio of mid-carbon fragments ($C_6^+-C_{29}^+$) to the sum of all carbon fragments ($C_1^+-C_{59}^+$) was used as a marker of soot maturity. Information related to refractory oxygen species from the rBC cores (e.g. surface oxides) were derived from the ratio of the $C_3O_2^+$ ion to C_3^+ . The $C_3O_2^+$ is exclusively detected during laser vaporization, and thus not found when the laser is off. The strongest signals from surface
- 135 oxides are CO^+ and CO_2^+ . However, these are prone to interferences from gas phase species and non-refractory OA (Corbin et al. 2015; Nielsen et al. 2017).

The set-up allowed two different types of IN experimentation: transient and chamber experiments. The transient ones were conducted during the cooking simulation where for this purpose sample line "A" of Fig. 1 was used. The purpose of transient

- 140 experiments was to study the possible IN potential from fresh, and polydisperse cookstove emissions during the simmering phase of the WBT. The duration of each cooking simulation was at least 30 minutes (simmering phase), which enabled one scan on the CFDC, over an *RH* range for one sampling temperature. A more detailed description of the CFDC experiments is presented in Sect. 2.4. The chamber experiments were carried out to extend sampling time up to slightly more than two hours, and for this polydisperse aerosol was ejected into the chamber after preliminary drying and dilution so the sampling
- 145 could be continued after the end of the WBTs. The chamber used in the experiments was a 15-m³ stainless steel aerosol storage chamber. The chamber was purged with filtered air prior to each experiment. It was filled following the sampling set-up of Fig. 1 to a mass concentration of $10-100 \ \mu g \ m^{-3}$ before the IN experiments with the CFDC was started. A typical aerosol injection time that was required to reach a sufficient mass concentration in the chamber was 10-40 minutes, where we aimed to that the emissions represent one full combustion cycle. After filling, the typical sampling time was
- 150 approximately 2 hours before the background signal of the CFDC increased too high or deposition effects, such as coagulation and wall losses, reduced the aerosol concentration too much for optimal sampling. Deposition through wall losses was strongest for ultrafine particle sizes due to diffusion, and longer sampling times were possible for accumulation mode particles only.
- 155 A CPC (Airmodus model A20) was used in parallel to the CFDC and it recorded typical concentrations of 30-200 cm⁻³ at the 250-500 nm size range after size-selection of the sample particles. The focus of the chamber experiments was to investigate the role of accumulation mode soot particles as INPs, and the samples were size-selected using a Vienna type DMA (DMA_{IN} in Fig. 1) prior to the CFDC and the CPC, aiming for the largest particle size that was present with a reasonable number concentration (>30 cm⁻³). It is worth emphasizing that the chamber was used only for storing and mixing aerosol potentially representing different combustion phases and no atmospheric aging via e.g. oxidation or interaction with other chemicals
- was applied to any emission samples that were studied using the CFDC.

2.2 Biomass feedstocks and cookstoves

The fuels used in this study consisted of firewood species available in large areas of sub-Saharan Africa and which are commonly used in residential cooking, or alternatively they represent regionally relevant agricultural residues. For instance,

- 165 the *Casuarina equisetifolia* (CAS), commonly known as Horsetail tree, is a species which is commonly used in agroforestry systems in the southern part of the African continent (Potgieter et al., 2014). In this work, the CAS was selected because it is a tree species that can be used in the implementation of sustainable agroforestry systems in East Africa. According to the Food and Agriculture Organization (FAO), agroforestry is a powerful tool in enhancing various ecosystem services, including the enhancement of food productivity, landscape regeneration and woody biomass production (Food and Agriculture Organization of the United Nations, 2017). Furthermore, the *Sesbania sesban* (SES) is also a tree applied to
- support the implementation of agroforestry systems, being native in most parts of sub-Saharan Africa and widespread in Kenya where the African fuels were collected for this study.

In addition to firewood, four types of agricultural residues were studied. Africa produces annually approximately one million 175 tons of coffee (International Coffee Organization, 2018), the largest coffee exporting countries being located in the sub-Saharan regions of the continent. During processing, agricultural residues such as tree branches and leaves form up to 50% of the total weight of coffee products (Oliveira and Franca, 2015) and investigations of using these residues to support the sustainable development of sub-Saharan Africa are ongoing. The coffee husk (CH) used in this study were collected from Kenya where *Coffea arabica* is the dominant coffee species. Moreover, bagasse (BG) is a by-product of sugar production 180 which has also demonstrated to be suitable for bioenergy production. This residue is usually disposed when the juice is

- extracted from sugarcanes. In this study, we focus on the utilization of sugarcane bagasse for biomass fuel production and use, as sugarcane (*Saccharum officinarum*) is an important agricultural product in most parts of Sub-Saharan African countries and the biomass potential of this residue is also studied to promote sustainability. Additionally, the Water Hyacinth (WH), *Eichhornia crassipes*, is an indigenous species from South America, but it was introduced to Africa during the
- 185 colonial period. It is a very invasive species, which requires constant population control, hence its utilization is proposed for energy purposes. Consequently, the removal of some of these plants from African lakes contributes to the sustainability of these water systems, producing significant amounts of combustible waste whose potential as a biofuel is also under research.
- Furthermore, rice husk (RC) is an agricultural residue from processing African rice, *Oryza glaberrima*, when forming cereal
 products. Rice production has been growing steadily in Africa for decades and processing the plants from paddy rice to
 milled rice produces approximately 25 weight percent of husks (Muthayya *et al.*, 2014), whose potential as a biofuel like BG is being studied for sustainable energy use. Finally, typical Swedish softwood pellets (SW) that consist of a 50-50 mix
 between pine and spruce (*Pinus sylvestris* and *Picea abies*, respectively) were used as a reference fuel for a pelletized pure stemwood based wooden fuel. This enabled better comparability between the emission properties. The ash rich fuels, CH,

BG, WH and RC, were mixed with SW on 50-50 mass ratio to improve the suitability and combustibility of the biomass fuels for cookstove combustion applications, including a better control of the ash content (i.e. certain inorganic elements). This was done to enhance the stability of the combustion process and burning time.

The simplest cookstove used was the three-stone fire (3S), which is a ring of three bricks around an open fire to hold the 200 cooking pot above fire. The rocket stove (RS) design is an improved stove model. However, the design is also fairly simple: the fuel sticks are inserted into the fuel shelf through the horizontal metal pipe and natural air draft causes a hot burning flame in the vertical section, just below where the cooking pot is located during operation. The more advanced concepts included two pellet gasifier stoves where fuel pellets are loaded in a vertical pipe beneath the cooking pot and lit from above. During the operation, the pyrolysis front moves down on the fuel bed and releases volatile components. When the volatile 205 gases reach the top of the cookstove, they are mixed with secondary air where they create a hot burning flame beneath the cooking pot. Although the operational principle was similar in both gasifier stoves, their air supply method differs in the following way: the natural-draft gasifier stove (NDGS) uses natural air draft, while the forced-draft gasifier stove (FDGS) is

equipped with an adjustable fan to force primary and secondary air flows to allow a more efficient burn-out of soot and volatile organic matter.

210

Prior to experimentation, the fuels were pre-processed for the different types of cookstoves as follows: the fuel sticks used were chopped to pieces approximately 2 cm in diameter, 17 cm and 13 cm in length for the 3S and the RS, respectively, and dried at room temperature. The pellet fuels were pelletized at the TEC-lab into dimensions of 8 mm x 15 mm (diameter x length, respectively). Fuel usage was thus standardized for the repeated WBT experiments according to cookstove type. The

3S and the RS were loaded with 100 g of stick fuel in the beginning of each WBT, and re-fueled slowly and continuously throughout the cooking simulation to maintain the flaming combustion. A typical number of re-fueling events was between five and ten. The gasifier stoves were loaded with 1 kg of pellets before lighting the fire, since they are batch-fired appliances where fuel is not added during operation.

2.3 The CFDC instrument

220 Ice nucleation experiments were conducted with the spectrometer for ice nuclei (SPIN, DMT Inc., Colorado, USA), described by Garimella et al. (2016). The main difference relative to the older version used by Ignatius et al. (2016) is improved temperature control. The main difference relative to the older version used by Ignatius et al. (2016) is improved temperature control via e.g. by an increase in the total number of thermocouples from 8 to 32 in the main chamber (growth section).

225

Briefly, the SPIN is a CFDC instrument using parallel plate geometry similar to the Portable Ice Nuclei Chamber (PINC) introduced by Chou et al. (2011) and the Zürich Ice Nuclei Chamber (ZINC) described by Stetzer et al. (2008). The sample

flow at 1 standard liter per minute (SLPM), is sandwiched between two sheath flows of 4.5 SLPM each, and the residence time in the region where ice nucleation can take place is approximately 10 seconds. The diffusional flux of water vapor

- 230 across the chamber is created via setting the ice-covered plates to different sub-zero temperatures, and the relative humidity with respect to water and ice (RH_w and RH_i , respectively) can be adjusted through the temperature difference between the plates. The aerosol is exposed to an isothermal evaporation section, upstream the detection with an optical particle counter (OPC). The temperature of the evaporation section was set to follow the temperature of the aerosol sample in the growth section of the chamber. The temperature, relative humidities and the ideal path of the sample flow, from here on referred as
- 235 lamina flow, are modelled according to a 1D flow model by Rogers et al. (1988). The ice layer was created to the IN chamber through cooling the walls to -32 °C and filling the chamber with de-ionized water, which resulted in a thin layer of ice on the walls of the chamber. After filling the IN chamber was purged of excess water and vacuumed down to 70 mbar for 3 minutes, which removed loose ice and reduced roughness of the ice layer and thus the background signal. Here, the background signal refers to unwanted OPC counts in the ice crystal size range, which may be due to break up of frost on the
- 240 ice-covered plates or alternatively due to tiny leaks in the instrument chamber. Typically, the background signal was below 1 particle per liter in the beginning of the experiment and the IN chamber was re-iced when it exceeded 10-15 particles per liter. This protocol, together with averaging the instrument data over 10-second periods as was done in data analysis, allowed detection of the ice-activated fraction from the order of 10⁻⁶ upwards in experiments where the sample concentration exceeded 2000 cm⁻³. Data averaging enabled investigation of large droplet/ice crystal populations against their corresponding
- 245 background signals, thus decreasing the lowest detection limit.

The CPC that was operated parallel to the SPIN monitored the sample concentration and thus provided required information for calculation of the ice-activated fraction. The ice-activated fraction α is defined as:

$$\alpha = \frac{N_{ice}}{N_{CPC}} \qquad (1)$$

where N_{ice} is the background-corrected concentration of ice crystals detected by the OPC, considering the dilution resulting from aerosol to sheath flow within SPIN, and N_{CPC} is the concentration of sample particles detected by the CPC. Background correction means that the frequency of background counts is linearly interpolated between background checks and the corresponding temporal values are subtracted from the measured signal. The uncertainty calculations in activated fraction assume that the OPC and CPC particle counts are randomly distributed, with the standard deviation then given by the inverse of the square root of counts, respectively. The total random uncertainty of α is based on the propagation of errors of the variables involved in α calculations (Eq. 1), namely IN counts, IN background before and after each run, and the total particle count. The depicted error bars represent ±1 standard deviation of the inferred random uncertainties.

We define the ice onset as an ice-activated fraction of 10^{-3} . Freezing conditions with $RH_w < 100$ % at the lamina flow are named as deposition mode and freezing above water saturation is referred to as immersion freezing (Vali et al., 2015), because liquid droplet formation is expected prior to freezing. Condensation and immersion freezing modes are indistinguishable in this instrument. The experiments were carried out using automated sequences that are available in this version of the SPIN, to ensure comparability between experiments on different fuels and cookstoves.

2.4 Ice nucleation experiments

265

The transient experiments were performed during the modified WBT using the transient sampling line, denoted A in Fig. 1. Fresh, polydisperse emission aerosol was introduced to the SPIN after desiccation to $RH_w < 5$ %. The concentrations were diluted to the order of 10³ cm⁻³ to avoid vapor depletion inside the SPIN. *RH*-scans were used for the transient experiments at constant T = -32 °C, scanning the *RH_i* from 100% up to 160%, which is close to the procedure used by Petters et al. (2009) and Levin et al., (2016). This scan included studying both deposition and immersion modes because *RH_w* ranged from 73 % to 115 %. Two experiment scans where T was fixed at *T* = -28 °C had similar *RH_i* range. The experimental approach for the transient experiments is a standard operation procedure often applied in previous studies (e.g. Petters et al., 2009; Welti et al., 2009).

We introduced a different experimental approach with a focus on immersion freezing for the experiments involving sampling of combustion aerosol from the chamber. This procedure is illustrated in Fig. 2, where a homogeneous freezing experiment on highly diluted ammonium sulfate (AS) droplets was performed. The dry seed particles with a mobility diameter of 350 nm were introduced to the SPIN in number concentration of approximately 150 cm⁻³, and droplets formed in the growth chamber of the SPIN due to operation with super-saturated conditions on water. All experiments were carried out on similar automated *T*-scan ramps containing the following steps: First, the internal background was checked via sampling filtered dry air for at least 5 minutes before the *T*-scan began. As *T* is descending, homogeneous freezing causes detection of ice crystals at about -37.9 °C. When *T* reaches the lowest set point of -43 °C the background is checked again via filtering sample flow at

- 285 the inlet for 3 minutes, before opening it again and scanning the *T* back to -32 °C (ascending ramp). The phase change back to droplets is detected at -38.9 °C, when the ice-activated fraction decreases to below 10^{-3} . The difference in ice onset and offset temperatures is most likely caused by cold pockets that may occur in the IN chamber during cooling in descending ramps. When the temperature control uncertainty is defined as one standard deviation from the averaged lamina temperature, it was observed that the temperatures typically deviate 0.9-1.0 °C and 0.2-0.3 °C from the set value of the lamina
- 290 temperature (see the shaded area around lamina temperature graph in Fig. 2) in descending and ascending ramps, respectively. Therefore, all data obtained from the descending ramps were omitted from the analysis and only ascending ramps were studied. The total sampling time of about 2 hours, which was used in chamber experiments, enabled up to three

repetitions for each WBT emission sample. This sampling time allowed 2-3 ascending T scans for each chamber experiment, and the reproducibility of the experiments will be discussed in the Sect. 3.4.

295

Separation between liquid droplets and ice crystals was carried out using a basic particle size-threshold method in the immersion mode experiments. Based on the homogeneous freezing experiments and all the other experiments we carried out with the same operation conditions, the very largest droplets detected with the OPC approached a diameter of 6 μ m. This can be seen in Fig. 2 between 20.38-20.49 hrs and 20.75-20.87 hrs when the inlet is open, and the lamina *T* is above homogeneous freezing temperature. Therefore, particles larger than 6 μ m are considered ice crystals in all these experiments

300 homogeneous freezing temperature. Therefore, particles larger than 6 µm are considered ice crystals in all these experiments and the choice of this size threshold will be discussed in Sect. 3.4.

The approach with operating the SPIN with potential co-existence of droplets and ice crystals is unusual, and it has to our knowledge previously only been applied for investigating the homogeneous freezing temperature (Ignatius et al., 2016). In this context, it should be noted that the evaporation section of the SPIN is less efficient than the ones in some other CFDCs. The SPIN OPC was designed to detect for particle depolarization to be used for discrimination of droplets versus ice crystals on a particle-by-particle basis. However, we found that the depolarization data obtained with the SPIN5 OPC during this campaign were not of a quality sufficient for such an approach. We never observed any droplets with optical diameters larger than 6 µm in any experiment with the approach described above, so we found it meaningful to infer an ice-active fraction by use of that size threshold, since the ice crystal mode typically was centered around larger sizes (Fig. 2). The potential biases in the ice-active fraction associated with this approach are discussed in more detail in Sect. 3.4. One advantage associated with this approach was that we observed formation of a significant droplet mode for all experiments, which clearly indicated that the supersaturation sufficed to produce cloud droplets in order to investigate immersion freezing – even for particles with a low CCN activity.

315

The chamber experiments served two major purposes, (i) it allowed for studying aerosol emissions representing an average over several combustion phases, and (ii) it allowed for focused ice nucleation studies of the soot particles. Regarding that typical aerosol injection time was up to 40 minutes, the aerosol emissions from the 3S and the RS stoves are highly variable depending on the combustion phase e.g. fuel addition, flaming and smoldering. In contrast, the aerosol emissions from the 320 gasifier stoves were constant during most of a given transient experiment except for the ignition and the extinction phases, which allowed for SPIN scans over almost constant aerosol over time. Most aerosol injections into the aerosol storage chamber were carried out over roughly a full combustion cycle representing the different combustion phases. Typical particle mass concentrations in the storage chamber was at the order of 10-100 µg m⁻³, and the *RH*_w was typically at the order of 20-25%.

The immersion mode was studied on ramps with fixed $RH_w = 115$ %, and the chamber experiments are listed in Table 2. In a few selected experiments, we investigated how modified combustion conditions possibly would influence the ice nucleating activity of the emitted soot particles. Five different modified experiments with the FDGS and SW pellets were used to inject aerosol particles into the chamber from which SPIN sampled size selected soot particles. Two of them were carried out with

- the secondary air supply being intentionally blocked, and the pot height was adjusted to increase the particle size in three experiments. In the experiments with blocked secondary air supply, the FDGS was modified to simulate usage of a poorly functioning stove, and the experiment was repeated twice on SW fuel. This simulation was done by reducing the secondary air supply, which resulted in more inefficient and incomplete combustion and increased the production of large soot particles significantly. The "*pot height*" experiments were done using different, less favorable heights of the cooking pot. The three such experiments were done on designated operation conditions of the FDGS and SW pellets, but the cooking pot was
- intentionally lifted to achieve production of larger particles: the real-time size distribution was monitored on the FPA throughout the experiment and the pot height was adjusted in a way that an increased production of large soot particles was observed. In typical cases, an offset of 8-10 cm above designated pot height affected production of large soot particles significantly.

340

It has previously been shown that the CFDC ice concentrations often are biased low by a factor of 3 and potentially up to a factor of 10 depending on the operation conditions [DeMott et al., 2015; DeMott et al., 2017; Garimella et al., 2017]. This bias is due to non-ideal behaviour of instruments, when the sample and sheath flows cannot follow the theoretical streamlines ideally. Hence, we will also estimate this effect and its possible effect on the results of this study.

345 2.5 Emission particle characterization

It is still largely an open question, which soot particle parameters influence the ice nucleating ability. However, the extensive online characterization of the aerosol applied in this work allowed us to investigate such potential links. The CCN activity, reported as the apparent hygroscopicity parameter κ_a, was inferred as described by Petters and Kreidenweis (2007) and the effective density ρ_e in a similar fashion as done by Rissler et al. (2013). The aethalometer was used for studying optical properties of the emission particles, measuring the Absorption Angstrom Exponent (AAE) in the wavelength interval from 370 nm to 950 nm. Furthermore, three physico-chemical properties were inferred from the SP-AMS measurements: organic aerosol to black carbon (OA/BC) ratio, relative abundance of refractory oxygen species and indirect information about the nanostructure of soot particles (Malmborg et al., 2019). The aim of this characterization was to study if the aforementioned six physico-chemical properties can explain variability in IN abilities from repeated experiments on different cookstoves,

355 fuels and combustion conditions.

3 Results and discussion

We present the IN results from transient and chamber experiments, compared to homogeneous freezing experiments whenever the experimentation included sample temperatures near or below homogeneous freezing temperature from the droplet freezing test. A result is considered positive if heterogeneous ice nucleation is observed under freezing conditions

that are distinguishable from homogeneous freezing, and negative if ice crystals have not been observed or if there is no

360

3.1 Transient experiments

distinguishable difference to homogeneous freezing conditions.

Transient experiments are summarized in Table 1 and they all were conducted on fresh, polydisperse samples whose size distributions were measured simultaneously during the WBTs. All samples were dominated by ultrafine (< 100 nm) particles by number, and the fraction of particles larger than 250 nm was typically below 5% of the total particle number concentration in all experiments. It can be expected that the size distributions and emission properties varied more on the 3S and the RS than on gasifier stoves due to re-fueling, i.e. addition of fuel sticks during the simmering phase. Regarding that the particle number concentration introduced to the SPIN was typically 1000-2500 cm⁻³ in transient experiments, ice nucleation was detectable with ice active fractions from the order of 10⁻⁶ upwards. Large accumulation mode particles accounted for concentrations up to about 100 cm⁻³ within the used sample concentrations. Ice crystal formation was not observed within the detection limit in any experiment at T = -32 °C or above. These 11 observations indicate that the fresh aerosol emission from these four stoves are active ice nucleators in neither deposition nor immersion/condensation mode at the given experiment temperatures. The emissions from all experiments were dominated by ultrafine particles by number, which clearly had no contribution to heterogeneous ice nucleation for the given temperatures.

375 **3.2 Chamber experiments**

The particle number size distributions were measured using the SMPS from the storage chamber, and therefore the sample sizes, as Fig. 3 shows, are based on dry particle mobility. The number size distributions were typically highly dominated by ultrafine and largely inorganic hygroscopic particles as indicated by CCN measurements (although not presented in this study). All ice nucleation experiments involving polydisperse aerosol indicated that the ultrafine particles did not play a role as DIPs for the investigated and the study.

- as INPs for the investigated conditions. The chamber experiments focused on quasi-monodisperse size selected soot particles. We aimed for sampling the largest possible particle mobility diameter (250-500 nm) with reasonable number concentrations (30-200 cm⁻³) resulting in minor contributions of multiply charged particles for any given experiment (see Table 2). The presented size distributions represent the situation when an activated fraction of 10⁻³ was observed in the SPIN in each shown cookstove and fuel combination. Particles larger than 250 nm accounted for up to 15 % of the total number
- 385 concentration in all chamber experiments involving the SPIN, which indicates that large soot particles were present in all studied cases.

The IN efficiencies of the emissions are presented in the following paragraphs. All experiment results are presented when 10^{-3} activated fraction was observed during the first ascending *T* ramp, after the particle storage period in the chamber has been

390

close to one hour. The bias regarding activated fraction due to restrictions in real-life CFDC instruments is being discussed separately in Sect. 3.4, along with reproducibility of the observations.

Starting from the simplest cookstove concept, the 3S produced 10⁻³ ice activation between -37.8 °C and -38.6 °C (see Table 3). Although these values are close to the observed homogeneous freezing temperature, comparison to a similar experiment on diluted AS droplets reveals that ice onset due to homogeneous freezing occurs at -38.9 °C with this operation of the SPIN. The observed homogeneous freezing with 10⁻³ ice activation at an average lamina temperature of -38.9 °C may occur at slightly lower temperature than expected from previous results reported in the literature. It is not clear at which temperature homogeneous freezing in SPIN can be expected based alone on theoretical considerations. This is due to the sample aerosol being exposed to a span in temperature and RH, and uncertainties related to droplet sizes and residence times above the evaporation section as discussed by Ignatius et al. (2016). We ascribe this relatively lower and reproducible homogeneous freezing temperature for the current version of SPIN to highly improved temperature control relative to earlier versions of SPIN (Ignatius et al., 2016). Potential biases and errors are discussed in more detail later.

Fig. 4 shows the results on two fuels, CAS and SES combusted with the 3S, and three different sample sizes. The errors
related to the average lamina temperature are presented above as the maximum span from the ideal lamina temperature, based on the uppermost 13 pairs of thermocouples on the warm and cold plates, respectively (Garimella et al., 2016). The statistical uncertainty of the activated fraction (vertical error bars) was found to be more prominent with low *α* values close to 10⁻⁴; in general, this uncertainty was at the order of 10-20% when the activated fraction was above 10⁻³, which agree with values reported by Garimella et al. (2017). The uncertainty analyses confirm that there is a significant difference in 10⁻³ onset
temperatures between the 3S experiments and the homogeneous freezing experiment. Good experiment reproducibility (see Table 3) that will be discussed in Sect. 3.4 indicates very good agreement between repeated ramps and thus strengthens the validity of these observations. Size dependency that could be distinguished from the instrument uncertainty between 250, 300 and 500 nm particles was not observed, which may be due to differences in physico-chemical properties between the individual experiments.

415

The RS emissions were slightly more active INPs than the ones from the 3S, the 10⁻³ activated fraction was achieved between -37.9 °C and -38.1 °C which is 0.8-1.0 °C higher than observed homogeneous freezing in the AS experiment, as can be seen from Table 3. Indications of size-dependency was also not observed in these experiments. The results from regular gasifier stove experiments are presented in Fig. 5, and conditions for ice onset were reached at -37.8 °C on the FDGS

420 with CH+SW pellets, and at -38.1 °C on the NDGS with SW pellets during standard operation conditions. Once again, a

clear size-dependency was not observed, the 260 nm particles from the experiment on the FDGS appear to be slightly more IN active than 400 nm particles from the NDGS. When compared to standard tests on the 3S and the RS, these observations consistently show that the soot particles emitted from burning the various fuels can act as heterogeneous INP at temperatures only slightly higher than those needed for homogeneous freezing of solution droplets.

425

430

AMS data.

The results from the five special experiments with modified combustion conditions are presented in Fig. 6. Two of the samples induced heterogeneous ice nucleation at temperatures about 4 °C (blocked secondary air supply) and 6 °C (elevated pot height) higher than observed for homogeneous freezing and an ice active fraction of 10^{-3} , respectively. These experiments were repeated, and the ice activity was comparable to homogeneous freezing for other intendedly identical experiments. This indicates that very minor changes in combustion conditions significantly influence the ice nucleating ability of freshly produced soot particles.

3.3 Properties of the emission particles

Fig. 7 presents six different physico-chemical particle properties for the aerosol samples included in Fig. 6. The CCN activity presented as the apparent κ (κ_a) and the effective density (ρ_{eff}) that were inferred for particles with a mobility diameter of 350 nm, and the results are presented in the two upper panels of Fig. 7. The κ_a values are very low for all the 435 samples with only minor variations, and the ρ_{eff} values are also low for all the samples with minor variations. These values of $\rho_{\rm eff}$ (0.2-0.3) are typical for fractal-like rBC dominated agglomerates from several different sources (Rissler et al. 2013). We observed a tendency towards decreasing κ_a and ρ_{eff} with increasing mobility diameter in the soot mode throughout the campaign, so the presented values can be considered upper estimates of what may be relevant for the larger quasi-440 monodisperse soot particles studied with SPIN. The differences in κ_a and ρ_{eff} reported for a mobility size of 350 nm are expected to be representative for relative differences for the entire soot mode, which is supported by strong correlations between these properties inferred for 200 and 350 nm soot particles and further supported by the powerlaw models for ρ_{eff} with respect to soot particles as presented by Rissler et al. (2013). Information about optical properties (AAE in the wavelength interval 370 to 950 nm), relative abundance of refractory oxygen species ($C_3O_2^+/C_3^+$), the ratio of organic aerosol 445 mass (OA) (in the m/z range 13-330 Th) to rBC mass, and indirect information about the nanostructure (C_{mid}/C_{all}) of the poly-disperse aerosol is presented in the lower panels in Fig. 7, with the three latter properties being inferred from the SP-

It is noteworthy that both cases with blocked secondary air supply (sample sizes in nm 450 and 500) showed elevated 450 absorption in the UV region (elevated AAE), despite the low OA to BC ratios. These two cases were also associated with an increased fraction of carbon fragments in the mid-carbon range C_6^+ to C_{29}^+ in the refractory carbon mass spectrum detected upon laser vaporization with the SP-AMS. Malmborg et al. (2019) and Török et al. (2018) linked increased UV absorption to

the occurrence of large carbon fragments in the SP-AMS mass spectrum, decreased fringe length detected by HR-TEM and

refractory (partly pyrolyzing) organic carbon in thermal-optical analysis for emissions from the Mini-CAST flame soot 455 generator. These findings suggest that such relationships are also present for refractory carbon dominated particles from cookstoves. Particularly, the two experiments with blocked secondary air inlet in the FDGS facilitated emissions of refractory carbon species with elevated UV absorption. The ratio of $C_3O_2^+/C_3^+$ showed only small variations between experiments (0.0059-0.0069) suggesting that the occurrence of refractory oxygen species (for example surface oxides) did not vary much between experiments.

460

465

Our detailed aerosol characterization presented in Fig. 7 indicates that we did produce particles with slightly different properties for intendedly identical experiments, as also was indicated by the observed difference in the ice nucleating ability. The low hygroscopicity, the low effective density and the optical properties indicate that the studied soot particles were not covered or coated by significant amounts of soluble material despite the presence of some organic compounds. It is noteworthy that the two more ice active soot samples (450 nm on blocked secondary air supply and the first repetition of the *"pot height"* experiment with 400 nm) are significantly different with regards to several properties. The only minor visible trend with ice-activity is a slightly lower ratio of refractory oxygen species ($C_3O_2^+/C_3^+$) for the two more ice active soot samples. However, that trend is not significant in a direct comparison to the freezing temperatures, and the differences in refractory oxygen species between the samples are not significant considering the errors.

470 In general, these supportive data indicate that either (i) the inferred properties are not determining the ice nucleating ability, or (ii) a potential for complex combinations of different soot particle properties being of relevance for the ice nucleating ability. Additional studies are needed to address these questions in more detail.

3.4 Experiment reproducibility and bias in ice crystal detection

In this section, we discuss the reproducibility of the observations to validate the results and to evaluate the measurement precision of the SPIN. The results from repeated ramps are summarized in Table 3 where we present the ice onset temperatures and their uncertainties at the lamina *T*. Unlike in Figs. 4-6, we present the uncertainty as one standard deviation from the average lamina *T* in this context, to show the overall precision of the temperature control. Observation numbers #1, #2 and #3 represent the IN onset after chamber residence times of approximately 20, 40 and 60 minutes, respectively, after filling the chamber was completed. These comparisons show that the results from repeated ramps are generally in good agreement with each other, and the differences in onset temperature remain within the typical deviation of the SPIN (see Sect. 2.3). It can be expected that the most prominent chamber effect has been coagulation of the ultrafine particles to large soot particles, but this has not affected the ice nucleating ability.

It should also be mentioned that across the width of the lamina, there is an additional span in temperature typically at the order of ± 0.4 °C, so a fraction of the aerosol will be exposed to further lower or higher temperatures, respectively, compared to just the variability in average lamina temperature as presented in Fig. 2. However, we find that the variability in average lamina temperature is a reasonable estimate of the error in sample temperature for the experiments presented in Figs. 4-6 considering the high reproducibility of the results presented in Table 3.

- 490 Furthermore, the ice-active fraction is in the current study biased low for two reasons: (i) only a fraction of the sample aerosol is focused in the lamina (Garimella et al., 2017), and (ii) only the larger size fraction of the ice crystals (optical diameter >6 µm) is included in the ice-active fraction (Fig. 2). Garimella et al. (2017) reported that the fraction of sample particles being focused in the lamina for comparable operation conditions of SPIN to be from 10% to 30% in most cases. In our homogeneous freezing experiment, the 'raw' ice-activated fraction as presented e.g. in Fig. 4 is about 0.12 for an average lamina temperature of -41°C. At that temperature, we would ideally expect an ice-activated fraction close to 1, which would be reached applying a correction factor of 8 in that case. For slightly higher lamina temperatures, the correction factor could potentially be lower, so we consider a correction factor in the range from 3 to 8 to be possible for this experiment for the
 - lamina temperature in the range from -41 to -38.0 °C. Application of such correction factors for the homogeneous freezing experiment will result in 10-3 ice activation between -38.5 °C and -38.0 °C in average lamina temperature.

500

The presented ice active fractions are biased low for two main reasons as mentioned above. As it can be seen from Figs. 4-6, the ice active fractions for a temperature of -41 °C is typically at the order of 0.1 to 0.2, while we would expect an ice active fraction close to 1 for homogeneous freezing at that temperature below -40 °C. These observations confirm that the presented ice active fractions are biased low. Previous studies show that only a fraction of the sample aerosol is focused in 505 the lamina. DeMott et al. (2015) found that the ice active fraction for mineral dust particles should be upscaled on average by a correction factor of about 3 for a CFDC over a range of operation conditions including approximately similar operation conditions (e.g. T \approx -35 °C and *RHw* about 106-108% at the lamina) to what we apply in our study. However, for those specific operation conditions (lamina T = -35 °C and just before droplet break through with an expected RHw about 106-108%) there is quite some scatter in the appropriate correction factor and a larger correction factor is often observed (DeMott 510 et al., 2015). The offset can be ascribed to only a fraction of the sample particles being focused on the lamina. Garimella et al. (2017) reported upscaling factors for SPIN in the range from 1.5 to 10 depending on the operation conditions due to only a fraction of the sample aerosol being focused in the lamina. This is also evident in this study at laminar temperatures below -40 °C when all droplets are expected to freeze homogeneously: the maxima of the inferred activated fractions (see Figs. 4-6) are between approximately 0.07 and 0.2, which indicates that correction factors in the range from 5 and up to 13 may be of

515 relevance for our approach and these specific conditions at low temperatures.

In the current study, we have droplets and ice crystals coexisting after the evaporation section for a significant range of conditions, and with our established ice crystal threshold (optical diameter >6 μ m), only a (varying) fraction of the ice crystals were included in our reported ice active fractions. The ice crystal size modes had maxima close to an optical

520 diameter of 8 µm in the T-scan experiments, and a fraction of the ice crystal size mode would be present below 6 µm

potentially overlapping with the droplet mode. Lognormal distributions were fitted to the ice crystal size mode for different lamina temperatures higher than -36°C for the most ice-active sample studied in order to estimate the fraction of of ice crystals with optical sizes below 6 μ m. We find that the ice-active fraction defined from the 6 μ m threshold would have to be multiplied by a factor of 1.4 \pm 0.2 in order to obtain a reasonable estimate of the ice-activated fraction considering the entire

- 525 ice crystal size mode when the ice crystal mode was observed to be well-defined and ascribed to heterogeneous immersion freezing. The uncertainty range is inferred from a sensitivity study including a range of reasonable fits, and it is due to noise in the data. It is not clear to which extent this correction factor is independent from the correction factor related to the sample aerosol not being entirely focused in the lamina. In any case, the bias associated with a relatively high ice size threshold is likely to be significantly lower than the expected bias due to only a fraction of the sample being focused in the lamina.
- 530

The underestimation based on the size threshold of 6 μ m turned more significant when the sample temperature was low enough to induce freezing of droplets smaller than 3-4 μ m and/or possibly hydrated particles appearing at sizes well below the ice crystal threshold. Investigation of the depolarization ratio of particles with an optical diameter in the sub-micrometer range with the SPIN OPC confirms that ice dominates in this size range at the lowest temperatures < -40 °C. Hence, the ice

535 active fraction inferred from our procedure is underestimated, which is likely to be significantly more pronounced in the lowest temperature as can be seen from Fig. 2. Furthermore, the ice active fraction may also be underestimated due to (i) losses of ice crystals immediately above the OPC, which to our knowledge has not been studied in detail so far for SPIN, and (ii) a relatively small fraction of particles activating into droplets inside SPIN – leading to a reduced fraction for detection in the immersion mode. The potential losses have to our knowledge not yet been characterized in detail – while the ice-active fractions observed for homogeneous freezing of dilute ammonium sulfate droplets and droplets formed on hydrophobic

particles typically resulted in very similar ice-active fractions (e.g. Fig. 6).

The considerations discussed above lead us to estimate a likely upscaling factor at the order of ~3-5 of the ice active fraction for lamina temperatures around -33^oC, while the upscaling factor is likely to approach 5-10 for temperatures significantly below homogenous freezing, around -40 °C. Further studies are needed to assess these biases in more detail – and their relative importance for different operation conditions of the CFDC instrument.

4 Summary and conclusions

The SUSTAINE experiment campaign provided an excellent opportunity to study the IN efficiencies of emissions from different biomass-fired cookstove designs under well-controlled laboratory conditions, which enabled comparability between

550 individual experiments. Two cookstoves, the 3S and RS, represented designs that are commonly used in daily household cooking in sub-Saharan Africa. The two more sophisticated designs, the NDGS and the FDGS, are currently under research for sustainable development and their popularity can be expected to increase along with economic development in this and similar regions. Based on observations from five special experiments on FDGS, this study shows that even small changes in

combustion conditions can significantly affect the IN abilities of emission particles. Extensive regional usage of cookstoves

555 suggests that the solid biomass burning emissions from cookstoves are a significant source of atmospheric particulate matter in sub-Saharan Africa. The CFDC instrument in this study, the SPIN, showed relatively good performance in temperature control, which enabled high reproducibility of experiments.

We conclude that the studied polydisperse aerosol emissions from transient experiments with different cookstoves showed 560 no indication of heterogeneous freezing above the detection limit (ice-activated fraction of about 10⁻⁵) for the investigated conditions (T = -28 °C and -32 °C and ~75-105% RH_w). However, it should be kept in mind that those aerosol emissions typically were heavily dominated by ultrafine particles by number. Hence, the measurement sensitivity with respect to the accumulation mode particles alone was relatively low in those experiments. Therefore, we also carried out a number of immersion freezing experiments focused on size selected accumulation mode soot particles sampled from an aerosol storage 565 chamber. The results from chamber experiments on accumulation mode soot particles show that they can induce heterogeneous ice nucleation in higher temperatures than what is required for homogeneous freezing, as happened in most of the experiments. Therefore, we conclude that usage of the cookstoves can emit potential INPs in the atmosphere, and thus affect cloud properties such as the frozen fraction in low temperature MPCs. The chamber experiments also support the outcome of transient experiments that included the same cookstove-fuel combinations, because the experimental procedures 570 included sampling at corresponding RH_w of 115% but in lower temperatures. With these observations combined, we conclude that the fresh cookstove emissions that were tested both transiently and from the chamber do not contain components that are active INPs above a temperature of -32 °C. All but two chamber experiments show an obvious difference to homogeneous freezing experiment, and therefore these emissions may well be of relevance for INP number concentrations in regions where other sources of ambient INPs are absent. Despite the chamber experiments consistently 575 showing IN activity above homogeneous freezing temperature, it still needs to be noted that the studied emissions were relatively fresh: atmospheric ageing processes can affect the IN properties (Brooks, et al., 2014) before the emissions reach the upper atmosphere, as is being supported by Häusler at al. (2018). The effect of atmospheric ageing can be studied via using e.g. oxidation reactors in future studies.

580 Deterioration of combustion efficiency was observed to increase the INP potential of the emission particles, which is likely due to elevated large soot particle production in incomplete combustion of the tested biomass fuels. This, when combined with results from transient experiments and ones with standard combustion conditions, indicates that even modest changes in combustion efficiency can drastically alter the ice-nucleating capabilities of the emissions. Our analysis on physico-chemical properties of the emissions revealed that the soot particle non-refractory organic coating, carbon nanostructure, and light absorption characteristics were different between the least and the most ice-active results, yet none of these properties could define the IN efficiency alone. We recommend further studies that aim to investigate the relationships between the soot potential as INPs and soot nanostructure, refractory surface-bound oxygen, and non-refractory organic composition and coating thickness. It is clearly a possibility that the IN efficiency of soot is determined by a complex combination of multiple particle properties. It is therefore possible that co-correlated microphysical changes in the soot structure (on orders of 10-

590 1000 nm) blurred the effects from changes in the non-refractory organic composition, nanostructure or oxygen bound to the soot surfaces.

It is worth emphasizing that all experiments of this study were carried out in well-controlled laboratory conditions and using standardized test procedures. Contrary to that, real-life use can differ significantly from these experiments in many factors, such as in fuel pre-processing, fuel properties, technical stove conditions and practical cooking procedures. These all affect the combustion and emission performance that can have a prominent effect on ice-nucleating abilities of emitted aerosol particles, which this study shows. Therefore, the relevance between these observations and real-life use should be explored before the contribution of biomass-burning emissions can be further evaluated in global perspective.

Data availability

600 The data set is available upon request from Kimmo Korhonen (kimmo.korhonen@uef.fi).

Competing interests

The authors declare that they have no conflict of interest.

605 Author contribution

TBK planned the ice-nucleation experiments on the SPIN instrument that was operated by KK during the campaign. AV, JF, KK, MK and TBK participated in data analysis of ice-nucleation experiments and/or interpretation of results. RL and RLC prepared the experimental set-up for water boiling tests and operated the combustion facility during the campaign. AE, CA,

610 JF, JP, TBK and VM participated in collection of supportive data and/or interpretation of results. BS, CB and JP participated as the organizers and supervisors of the SUSTAINE experiment campaign. AV and KEJL achieved funding for the SPIN instrument. All authors participated in scientific discussions on this study and reviewed/edited the manuscript during its preparation process. KK prepared the manuscript with contributions from all co-authors.

615 Acknowledgements

This work was financially supported by the Swedish Research Council FORMAS through the project Sustainable Biomass Utilization in Sub-Saharan Africa for an Improved Environment and Health (Dnr. 942-2015-1385) and Atmospheric cloud droplet formation and ice formation of wood combustion aerosols (2015-992). T.B. Kristensen gratefully acknowledges

620 funding from the Swedish Research Council (VR) grant no. 2017-05016. LU-EAT researchers acknowledge financial support from the Swedish Research Councils VR (projects 2018-04200 and 2013-05021) and FORMAS (project 2013-

01023). University of Eastern Finland and Finnish Meteorological Institute acknowledge Academy of Finland, Centre of Excellence (grant no. 272041) and North-Savo Council - European Regional Development Fund's (project no. A32350). Ricardo Carvalho acknowledges the Postdoctoral grant JCK-1516 funded by the Kempe Foundation.

625 References

630

640

Bhandari, J., China, S., Chandrakar, K.K., Kinney, K., Cantrell, W., Shaw, R.A., Mazzoleni, L.R., Girotto, G., Sharma, N., Gorkowski, K., Gilardoni, S., Decesari, S., Facchini, M.C., Zanca, N., Pavese, G., Esposito, F., Dubey, M.K., Aiken, A.C., Chakrabarty, R.K., Moosmüller, H., Onasch, T.B., Zaveri, R.A., Scarnato, B.V., Fialho, P. and Mazzoleni, C., Extensive Soot Compaction by Cloud Processing from Laboratory and Field Observations. *Sci. Rep.* 9, 11824, doi:10.1038/s41598-019-48143-y, 2019.

Bond, T.C., Doherty, S.J., Fahey, D.W., Forster, P.M., Berntsen, T., DeAngelo, B.J., Flanner, M.G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P.K., Sarofirm, M.C., Schultz, M.G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S.K., Hopke, P.K., Jacobson, M.Z., Kaiser, J.W., Klimont, Z., Lohmann, U., Schwarz, L.B., Gli, J.H. D., Sterner, T., Warre, G.G., L.Z., Le, G.G., Dan, K., Kaiser, J.W., Klimont, Z., Lohmann, U., Schwarz, L.B., Gli, J.H. D., Sterner, T., Warre, G.G., L.Z., Le, G.G., Dan, K., Kaiser, J.W., Klimont, Z., Lohmann, V., Schwarz, K., Kaiser, J.W., Klimont, Z., Lohmann, V., Schwarz, K., Kaiser, J.W., Klimont, Z., Lohmann, V., Schwarz, K., Kaiser, J.W., Klimont, Z., Kaiser, J.W., Klimont, Z., Lohmann, V., Schwarz, K., Kaiser, J.W., Klimont, Z., Kaiser, J.K., Kai

635 J.P., Shindell, D., Storelvmo, T., Warren, S.G. and Zender, C.S., Bounding the role of black carbon in the climate system: A scientific assessment, J. Geophys. Res. Atmos., 118, 5380–5552, doi:10.1002/jgrd.50171, 2013.

Bonjour, S., Adair-Rohani, H., Wolf, J., Bruce, N.G., Mehta, S., Prüss-Ustün, A., Lahiff, M., Rehfuess, E.A., Mishra, V. and Smith, K.R.: Solid fuel use for household cooking: Country and regional estimates for 1980-2010, *Environmental Health Perspectives*, 121(7), 784–790, doi: 10.1289/ehp.1205987, 2013.

Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S.K., Sherwood, S., Stevens, B. and Zhang, X.Y.: Clouds and Aerosols. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the
Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.

Brooks, S.D., Suter, K. and Olivarez, L., Effects of Chemical Aging on the Ice Nucleation Activity of Soot and Polycyclic 650 Aromatic Hydrocarbon Aerosols, *J. Phys. Chem.* A, 118, 43, 10036-10047, doi: 10.1021/jp508809y, 2014. Carvalho, R.L., Lindgren, R., García-Lópeza, N., Nyambane, A., Nyberg, G., Diaz-Chavez, R. and Boman, C.: Household air pollution mitigation with integrated biomass/cookstove strategies in Western Kenya, *Energy Policy*, 131, 168–186, doi: 10.1016/j.enpol.2019.04.026, 2019.

655

Carvalho, R.L., Jensen, O.M. and Tarelho, L.A.C.: Mapping the performance of wood-burning stoves by installations worldwide, *Energy & Buildings*, 127, 658–679, doi: 10.1016/j.enbuild.2016.06.010, 2016.

Champion, W.M. and Grieshop, A.P.: Pellet-Fed Gasifier Stoves Approach Gas-Stove Like Performance during in-Home Use in Rwanda, *Env. Sci. Tech.*, 53, 6570–6579, doi: 10.1021/acs.est.9b00009, 2019.

Chou, C., Stetzer, O., Weingartner, E., Jurányi, Z., Kanji, Z. A., and Lohmann, U.: Ice nuclei properties within a Saharan dust event at the Jungfraujoch in the Swiss Alps, *Atmos. Chem. Phys.*, 11, 4725–4738, doi: 10.5194/acp-11-4725-2011, 2011.

665 Coffee Market Report, September 2018, International Coffee Organization, 2018.

Corbin, J. C., Lohmann, U., Sierau, B., Keller, A., Burtscher, H. and Mensah, A. A.: Black carbon surface oxidation and organic composition of beech-wood soot aerosols, *Atmos Chem. Phys*, 15(20), 11885-11907, doi:10.5194/acp-15-11885-2015, 2015.

670

DeMott, P., An exploratory study of ice nucleation by soot aerosols, J. Appl. Met., 29, 1072-1079, 1990.

DeMott, P.J. Petters, M.D., Prenni, A.J., Carrico, C.M., Kreidenweis, S.M., Collett Jr., J.L. and Hoosmüller, H., Ice nucleation behavior of biomass combustion particles at cirrus temperatures, *J. Geophys. Res.*, 114, D16205, doi: 10.1029/2009JD012036, 2009.

DeMott, P. J., Prenni, A.J., Liu, X., Kreidenweis, S.M., Petters M.D., Twohy, C.H., Richardson, M.S., Eidhammer, T., and Rogers, D.C., Predicting global atmospheric ice nuclei distributions and their impacts on climate, *PNAS*, 107, 11217-11222, https://doi.org/10.1073/pnas.0910818107, 2010.

DeMott, P.J., Hill, T.C.J., Petters, M.D., Bertram, A.K., Tobo, Y., Mason, R.H., Suski, K.J., McCluskey, C.S., Levin, E.J.T., Schill, G.P., Boose, Y., Rauker, A.M., Miller, A.J., Zaragoza, J., Rocci, K., Rothfuss, N.E., Taylor, H.P., Hader, J.D., Chou, C., Huffman, J.A., Pöschl, U., Prenni, A.J. and Kreidenweis S.M., Comparative measurements of ambient atmospheric concentrations of ice nucleating particles using multiple immersion freezing methods and a continuous flow diffusion

685 chamber, Atmos. Chem. Phys., 17, 11227–11245, doi: 10.5194/acp-17-11227-2017, 2017.

⁶⁸⁰

DeMott, P.J., Prenni, A.J., McMeeking, G.R., Sullivan, R.C., Petters, M.D., Tobo, Y., Niemand, M., Möhler, O., Snider, J.R., Wang, Z., Kreidenweis, S.M., Integrating laboratory and field data to quantify the immersion freezing ice nucleation activity of mineral dust particles, *Atmos. Chem. Phys.*, 15, 393–409, 10.5194/acp-15-393-2015, 2015.

690

Diehl, K and Mitra, S.K.: A laboratory study of the effects of a kerosene-burner exhaust on ice nucleation and the evaporation rate of ice crystals, *Atmos. Env.*, 32, 18, 3145-3151,1998.

Food and Agriculture Organization of the United Nations, 2017. Agroforestry for landscape restoration, Rome.

695

Garimella, S., Kristensen, T. B., Ignatius, K., Welti, A., Voigtländer, J., Kulkarni, G. R., Sagan, F., Kok, G. L., Dorsey, J., Nichman, L., Rothenberg, D., Rösch, M., Kirchgäßner, A., Ladkin, R., Wex, H., Wilson, T. W., Ladino, L. A., Abbatt, J. P. D., Stetzer, O., Lohmann, U., Stratmann, F., and Cziczo, D. J.: The Spectrometer for Ice Nuclei (SPIN): An instrument to investigate ice nucleation, *Atmos. Meas. Tech.*, 9, 2781–2795, doi: 10.5194/amt-9-2781-2016, 2016.

700

Garimella, S., Rothenberg, D.A., Wolf, M.J., David, R.O., Kanji, Z.A., Wang, C., Rösch, M. and Cziczo, D.J., Uncertainty in counting ice nucleating particles with continuous flow diffusion chambers, *Atmos. Chem. Phys.*, 17, 10855–10864, doi: 10.5194/acp-17-10855-2017, 2017.

705 Gorbunov, B., Baklanov, A., Kakutkina, N., Windsor, H.L. and Toumi, R., Ice nucleation on soot particles, *Aer. Sci.*, 32, 199-215, doi: 10.1016/S0021-8502(00)00077-X, 2001.

Ferry, D., Suzanne, J., Nitsche, S., Popovicheva, O.B. and Shonija, N.K., Water adsorption and dynamics on kerosene soot under atmospheric conditions. *J. Geophys. Res.*, 107, D23, 4734, doi:10.1029/2002JD002459, 2002.

710 Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 9817–9854, doi:10.5194/acp-12-9817-2012, 2012.

Huang, Y., Unger, N., Storelvmo, T., Harper, K., Zheng, Y. and Heyes, C., Global radiative effects of solid fuel cookstove aerosol emissions, *Atmos. Chem. Phys.*, 18, 5219-5233, doi: 10.5194/acp-189-5219-2018, 2018.

715

Häusler, T., Gebhardt, P., Iglesias, D., Rameshan, C., Marchesan, S., Eder, D. and Grothe, H.: Ice Nucleation Activity of Graphene and Graphene Oxides, *J. Phys. Chem.*, 122, 8182-8190, 10.1021/acs.jpcc.7b10675, 2018.

Ignatius, K., Kristensen, T. B., Järvinen, E., Nichman, L., Fuchs, C., Gordon, H., Herenz, P., Hoyle, C. R., Duplissy, J.,

- 720 Garimella, S.,Dias, A., Frege, C., Höppel, N., Tröstl, J., Wagner, R., Yan, C.,Amorim, A., Baltensperger, U., Curtius, J., Donahue, N. M., Gallagher, M. W., Kirkby, J., Kulmala, M., Möhler, O., Saathoff, H., Schnaiter, M., Tomé, A., Virtanen, A., Worsnop, D., and Stratmann, F.: Heterogeneous ice nucleation of viscous secondary organic aerosol produced from ozonolysis of a-pinene, *Atmos. Chem. Phys.*, 16, 6495–6509, doi: 10.5194/acp-16-6495-2016, 2016.
- 725 Kanji, Z., Ladino, L.A., Wex, H., Boose, Y., Burkert-Kohn, M., Czizco, D. and Krämer, M., Overview of Ice Nucleating Particles, *Meteorol. Monogr.*, 58, 1.1-1.33., doi: 10.1175/AMSMONOGRAPHS-D-16-0006.1, 2017.

Koehler, K.A., DeMott, P.J., Kreidenweiss, S.M., Popovicheva, O.B., Petters, M.D., Carrico, C.M., Kireeva, E.D., Khokhlova, T.D. and Shonija, N.D., Cloud condensation nuclei and ice nucleation activity of hydrophobic and hydrophilic soot particles. *Phys. Chem. Chem. Phys.*, 2009.11, 7906-7920, doi: 10.1039/B905334B, 2009.

Korolev, A., Farquhar, G., Field, P.R., Franklin, C., Lawson, P., Wand, Z., Williams, E., Abel, S.J., Axisa, D., Borrmann, S., Crosier, J., Fugal, J., Krämer, M., Lohmann, U., Schlenzek, O., Schnaiter, M. and Wendisch, M., Mixed-Phase Clouds: Progress and Challenges, *Meteorol.Monogr.*, 58, 5.1-5.50., doi: 10.1175/AMSMONOGRAPHS-D-17-0001.1, 2017.

735

730

Levin, E. J. T., McMeeking, G.R., DeMott, P.J., McCluskey C.S., Carrico C.M., Nakao S., Jayrathne T., Stone E.A., Stockwell C.E., Yokelson R.J. and Kreidenweis S.M., Ice-nucleating particle emissions from biomass combustion and the potential importance of soot aerosol, *J. Geophys. Res. Atmos.*, 121, 5888–5903, doi:10.1002/2016JD024879, 2016.

- Lim, S.S, Vos, T., Flaxman, A.D., Danaei, G., Shibuya, K., Adair-Rohani, H., AlMazroa, M.A., Amann, M., Anderson, H.R., Andrews, K.G., Aryee, M., Atkinson, C., Bacchus, L.J., Bahalim, A.N., Balakrishnan, K., Balmes, J., Barker-Collo, S., Baxter, A., Bell, M.L., Blore, J.D., Blyth, F., Bonner, C., Borges, G., Bourne, R., Boussinesq, M., Brauer, M., Brooks, P., Bruce, N.G., Brunekreef, B., Bryan-Hancock, C., Bucello, C., Buchbinder, R., Bull, F., Burnett, R.T., Byers, T.E., Calabria, B., Carapetis, J., Carnahan, E., Chafe, Z., Charlson, F., Chen, H., Chen, J.S., Chen, A.T-A., Child, J.C., Cohen, A., Colson, K.E., Cowie, B.C., Darby, S., Darling, S., Davis, A., Degenhardt, L., Dentener, F., Des Jarlais, D.C., Devries, K., Dherani, M., Ding, E.L., Dorsey, E.R., Driscoll, T., Edmond, K., Ali, S.E., Engell, R.E., Erwin, P.J., Fahimi, S., Falder, G., Farzadfar, F., Ferrari, A., Finucane, M.M., Flaxman, S., Fowkes, F.G., Freedman, G., Freeman, M.K., Gakidou, E., Ghosh, S., Giovannucci, E., Gmel, G., Graham, K., Grainger, R., Grant, B., Gunnell, D., Gutierrez, H.R., Hall, W., Hoek, H.W., Hogan, A., Hosgood, H.D. 3rd, Hoy, D., Hu, H., Hubbell, B.J., Hutchings, S.J., Ibeanusi, S.E., Jacklyn, G.L., Jasrasaria, R., Jonas,
- 750 J.B., Kan, H., Kanis, J.A., Kassebaum, N., Kawakami, N., Khang, Y.H., Khatibzadeh, S., Khoo, J.P., Kok, C., Laden, F., Lalloo, R., Lan, Q., Lathlean, T., Leasher, J.L., Leigh, J., Li, Y., Lin, J.K., Lipshultz, S.E., London, S., Lozano, R., Lu, Y., Mak, J., Malekzadeh, R., Mallinger, L., Marcenes, W., March, L., Marks, R., Martin, R., McGale, P., McGrath, J., Mehta, S.,

Mensah, G.A., Merriman, T.R., Micha, R., Michaud, C., Mishra, V., Mohd Hanafiah K., Mokdad, A.A., Morawska, L., Mozaffarian, D., Murphy, T., Naghavi, M., Neal, B., Nelson, P.K., Nolla, J.M., Norman, R., Olives, C., Omer, S.B.,

- 755 Orchard, J., Osborne, R., Ostro, B., Page, A., Pandey, K.D., Parry, C.D., Passmore, E., Patra, J., Pearce, N., Pelizzari, P.M., Petzold, M., Phillips, M.R., Pope, D., Pope, C.A. 3rd, Powles, J., Rao, M., Razavi, H., Rehfuess, E.A., Rehm, J.T., Ritz, B., Rivara, F.P., Roberts, T., Robinson, C., Rodriguez-Portales, J.A., Romieu, I., Room, R., Rosenfeld, L.C., Roy, A., Rushton, L., Salomon, J.A., Sampson, U., Sanchez-Riera, L., Sanman, E., Sapkota, A., Seedat, S., Shi, P., Shield, K., Shivakoti, R., Singh, G.M., Sleet, D.A., Smith, E., Smith, K.R., Stapelberg, N.J., Steenland, K., Stöckl, H., Stovner, L.J., Straif, K.,
- Straney, L., Thurston, G.D., Tran, J.H., Van Dingenen, R., van Donkelaar, A., Veerman, J.L., Vijayakumar, L., Weintraub, R., Weissman, M.M., White, R.A., Whiteford, H., Wiersma, S.T., Wilkinson, J.D., Williams, H.C., Williams, W., Wilson, N., Woolf, A.D., Yip, P., Zielinski, J.M., Lopez, A.D., Murray, C.J., Ezzati, M., AlMazroa, M.A. and Memish, Z.A.: A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*, 380(9859), 2224–60, doi: 10.1016/S0140-6736(12)61766-8, 2012.

Mahrt, F., Marcolli, C., David, R. O., Grönquist, P., Barthazy Meier, E. J., Lohmann, U., and Kanji, Z. A.: Ice nucleation abilities of soot particles determined with the Horizontal Ice Nucleation Chamber, *Atmos. Chem. Phys.*, 18, 13363–13392, https://doi.org/10.5194/acp-18-13363-2018, 2018.

770

Malmborg, V. B., Eriksson, A. C., Török, S., Zhang, Y., Kling, K., Martinsson, J., Fortner, E.C., Gren, L., Kook, S., Onasch, T.B., Bengtsson, P-E. and Pagels J.: Relating aerosol mass spectra to composition and nanostructure of soot particles. *Carbon*, 142, 535-546, doi: 10.1016/j.carbon.2018.10.072, 2019.

775 Matus, A. V. and L'Ecuyer, T.S., The role of cloud phase inEarth's radiation budget, J. Geophys. Res. Atmos., 122, 2559– 25, doi:10.1002/2016JD025951, 2017.

McCluskey, C.S., DeMott, P.J., Prenni, A.J., Levin, E.J.T., McMeeking, G.R., Sullivan, A. P., Hill, T. C. J., Nakao, S., Carrico, C. M. and Kreidenweis, S.M.: Characteristics of atmospheric ice nucleating particles associated with biomass burning in the US: Prescribed burns and wildfires, *J. Geophys. Res.Atmos.*, 119, 10458–10470, doi:10.1002/2014JD021980,

2014.

Muthayya S., Sugimoto, J.D., Montgomery, S and Maberlys, G.F.: An overview of global rice production, supply, trade, and consumption, *Ann. N.Y. Acad. Sci.* 1324, 7–14, doi: 10.1111/nyas.12540, 2014.

785

Mülmenstädt, J., Sourdeval, O., Delanoë, J. and Quaas, J., Frequency of occurrence of rain from liquid-, mixed-, and icephase clouds derived from A-Train satellite retrievals, *Geophys. Res. Lett.*, 42, 6502-6509, doi:10.1002/2015GL064604, 2015.

790 Möhler, O., Büttner S., Linke C., Schnaiter M., Saathoff H., Stetzer O., Wagner R., Krämer M., Mangold A., Ebert V. and Shurath U., Effect of sulfuric acid coating on heterogeneous ice nucleation by soot aerosol particles, *J. Geophys. Res.*, 110, D11210, doi:10.1029/2004JD005169, 2005.

Nielsen, I. E., Eriksson, A. C., Lindgren, R., Martinsson, J., Nyström, R., Nordin, E. Z., Sadiktsis, I., Boman, C., Nøjgaard,
J.K. and Pagels, J.: Time-resolved analysis of particle emissions from residential biomass combustion–Emissions of refractory black carbon, PAHs and organic tracers, *Atmos. Env.*, 165, 179-190, doi:10.1016/j.atmosenv.2017.06.033, 2017.

Oliveira, L.S. and Franca, A.S.; Coffee in Health and Disease Prevention - 1st Edition, Chapter 31 - An Overview of the Potential Uses for Coffee Husks, *Coffee in Health and Disease Prevention*, 283-291, doi: 10.1016/B978-0-12-4095175.00031-0, 2015.

Onasch, T. B., Trimborn, A., Fortner, E. C., Jayne, J. T., Kok, G. L., Williams, L. R., Davidovits, P. and Worsnop, D. R.: Soot particle aerosol mass spectrometer: development, validation, and initial application. *Aeros. Sci. Tech.*, 46(7), 804-817, 2012.

805

815

Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, doi:10.5194/acp-7-1961-2007, 2007.

Petters, M. D., Parsons, M.T., Prenni, A.J., DeMott P.J., Kreidenweis, S.M., Carrico C.M., Sullivan A.P., McMeeking G.R.,
Levin, E., Wold, C.E., Collett Jr., J.L. and Moosmüller, H., Ice nuclei emissions from biomass burning, *J. Geophys. Res.*, 114, D07209, doi:10.1029/2008JD011532, 2009.

Popovicheva O., Kireeva E., Persiantseva, N., Khokhlova, T., Shonija, N., Tishkova, V. and Demirdijan, B., Effect of soot on immersion freezing of water and possible atmospheric implications, *Atmos. Res.*, 90, 326–337, doi: 10.1016/j.atmosres.2008.08.004, 2008.

Potgieter, L.J., Richardson, D.M. and Wilson, J.R.U., *Casuarina*: biogeography and ecology of an important tree genus in a changing world, *Biol.*, 16, 609-633, doi: 10.1007/s10530-013-0613-x, 2014.

820 Pruppacher, H. R. and Klett, J. D.: Microphysics of Clouds and Precipitation, Atmospheric and Oceanographic Sciences Library, Kluwer Academic Publishers, Dordrecht, the Netherlands, 1997.

Rigg, Y.J., Alpert, P.A. and Knopf, D.A., Immersion freezing of water and aqueous ammonium sulfate droplets initiated by humic-like substances as a function of water activity, *Atmos. Chem. Phys.*, 13, 6603-6622, doi:10.5194/acp-13-6603-2013, 2013.

Rissler, J., Messing, M. E., Malik, A. I., Nilsson, P. T., Nordin, E. Z., Bohgard, M., Sanati, M and Pagels, J.: Effective density characterization of soot agglomerates from various sources and comparison to aggregation theory. *Aer. Sci. Tech.*, 47(7), 792-805,doi: 10.1080/02786826.2013.791381, 2013.

830

825

Rogers, D. C.: Development of a continuous flow thermal gradient diffusion chamber for ice nucleation studies, *Atmos. Res.*, 22, 149–181, 1988.

Stetzer, O., Baschek, B., Lüönd, F., and Lohmann, U.: The Zurich Ice Nucleation Chamber (ZINC) – a new instrument to
investigate atmospheric ice formation, *Aerosol Sci. Tech.*, 42, 64–72,
2008.

Twohy, C.H., DeMott, P.J., Pratt, K.A., Subramanian, R, Kok, G.L., Murphy, S.M., Lersch, T., Heymsfield, A.J., Wang, Z.,
Prather, K.A. and Seinfeld, J.H.: Relationships of Biomass-Burning Aerosols to Ice in Orographic Wave Clouds, *J. Atm. Sci.*,
67, 2437-2450, doi: 10.1175/2010JAS3310.1, 2010.

Török, S., Malmborg, V. B., Simonsson, J., Eriksson, A., Martinsson, J., Mannazhi, M., Pagels, J. and Bengtsson, P. E.: Investigation of the absorption Ångström exponent and its relation to physicochemical properties for mini-CAST soot. *Aer. Sci. Tech.*, 52(7), 757-767, doi: 10.1080/02786826.2018.1457767, 2018.

845 Vali, G., DeMott, P.J., Möhler, O. and Whale, T.F., Technical Note: A proposal for ice nucleation terminology, *Atmos. Chem. Phys.*, 15, 10263–10270, doi:10.5194/acp-15-10263-2015, 2015.

Vergara-Temprado, J., Holden, M.A., Orton, T.R., O'Sullivan, D., Umo, N.S., Browse, J., Reddington, C., Baeza-Romero, M.T., Jones, J.M., Lea-Langton, A., Williams, A., Carslaw, K.S. and Murray, B.J., Is Black Carbon an Unimportant Ice-Nucleating Particle in Mixed-Phase Clouds?, *J. Geophys. Res.*, 123, 4273-4283, doi: 10.1002/2017JD027831, 2018.

Welti A., Lüönd, F., Stetzer, O. and Lohmann, U., Influence of particle size on the ice nucleating ability of mineral dusts, *Atmos. Chem. Phys.*, 9, 6705–6715, doi: 10.5194/acp-9-6705-2009, 2009.

Wiedensohler, A.: An approximation of the bipolar charge distribution for particles in the submicron size range, *J. Aer. Sci.*, 19, 387-389, doi:10.1016/0021-8502(88)90278-9, 1988.

Yun, Y., Penner, J.E. and Popovicheva, O.: The effects of hygroscopicity on ice nucleation of fossil fuel combustion aerosols in mixed-phase clouds, *Atmos. Chem. Phys.*, 13, 4339–4348, doi: 10.5194/acp-13-4339-2013, 2013.



Figure 1: Schematic of the experimental set-up used to conduct the WBT modified version relevant to the IN experiments. Sample
lines A and B were used for transient and chamber experiments, respectively. Blue squares with letter "V" indicate valves which were used for switching between the sampling lines. The red and blue arrows show directions of sample flow related to transient and chamber experiments, respectively. The purple arrows show the flow direction used in both types of experiments, depending on position of the relevant switch valve. The acronyms are defined as follows: APM = aerosol particle mass analyzer, CCNC = cloud condensation nuclei counter, CFDC = continuous flow diffusion chamber ice nucleus counter, CPC = condensation particle counter, DMA = differential mobility analyzer, ED = ejector dilution, FPA = fast particle analyzer, SMPS = scanning mobility particle sizer, SP-AMS = soot particle aerosol mass spectrometer. The dimensions and sample line lengths are not in scale to each

other.



Figure 2: The immersion mode experiment procedure example from homogeneous freezing experiment with 350 nm ammonium sulfate seeds. Upper panel: Symbols Tw, Tc, and Ta represent average temperatures for warm wall, cold wall and sample aerosol (lamina), respectively. The shaded black area in the top graph represents one standard deviation from the averaged lamina temperature. Lower panel: Events marked "BGD" depict background signal checks. The color bar indicates the logarithmic intensity of particle counts and the dashed black line the ice threshold size of 6 µm. The gap in detection near 3 µm is a specific artefact in the OPC in this SPIN unit.



Figure 3: Examples of particle mobility size distributions from chamber experiments on different cookstoves, during the time 880 when an activated fraction of 10⁻³ was observed in the SPIN.



Figure 4: Ice-activation spectra of emissions from the 3-stone fire at $RH_w = 115\%$, chamber experiments. Each shaded area of respective color represents the T-span across the lamina (maximum uncertainty) during each observation. The error bars present combined relative standard deviations in particle detection by the CPC and OPC of the SPIN. The solid black line presents the 10⁻³ activation threshold. The ice-activation spectrum for homogeneous freezing is included for comparison.



Figure 5: Ice-activation spectra of emissions from gasifier stoves at $RH_w = 115\%$ at regular operation conditions. Each shaded area of respective color represents the *T*-span across the lamina (maximum uncertainty) during each observation. The error bars present combined relative standard deviations in particle detection by the CPC and OPC of the SPIN. The solid black line presents the 10^{-3} activation threshold. The ice-activation spectrum for homogeneous freezing is included for comparison.



Figure 6: Ice-activation spectra of emissions from combustion of SW pellets in forced-draft gasifier stove, with modified combustion conditions at $RH_w = 115\%$. Each shaded area of respective color represents the *T*-span across the lamina (maximum uncertainty) during each observation. The error bars present combined relative standard deviations in particle detection by the CPC and OPC of the SPIN. The solid black line presents the 10^{-3} activation threshold. The ice-activation spectrum for homogeneous freezing is included for comparison.



900 Figure 7: Physico-chemical properties of soot particles in special experiments where the combustion conditions of SW in the FDGS were modified. The five aerosol samples presented have been re-ordered in a way that the samples with higher ice nucleating ability are located further to the right. The properties from the top to the bottom represent the CCN activity (κ_a), the effective density (ρ_{eff}), Absorption Angstrom Exponent in the wavelength interval 370 to 950 nm (AAE₃₇₀₋₉₅₀), the ratio of organic aerosol mass to refractive BC mass (OA/BC), indirect qualitative information about the nanostructure (C_{mid}/C_{all}) and the relative abundance of refractory oxygen species ($C_3O_2^+/C_3^+$). The two uppermost panels are for quasi-monodisperse soot particles with a mobility diameter of 350 nm, while the other properties are inferred for polydisperse aerosol.

Table 1: Summary of transient experiments on polydisperse sample particles on RH_i scan 100-160 %, and the ice crystal detection910limit. The detection limit depends on sample concentration and is defined as the lowest activated fraction when detection of ice
crystals is distinguishable from the background signal in data averaged over 10-second periods.

Cookstove	Fuel	Temperature	Detection limit
3-stone fire	BIR	-32 °C	6.6x10 ⁻⁶
Rocket stove	CAS	-32 °C	7.3x10 ⁻⁶
Rocket stove	CAS	-32 °C	4.0x10 ⁻⁶
ND gasifier	SW	-32 °C	5.0x10 ⁻⁶
ND gasifier	CH+SW	-32 °C	1.3x10 ⁻⁶
ND gasifier	BG+SW	-28 °C	5.0x10 ⁻⁶
ND gasifier	BG+SW	-32 °C	3.4x10 ⁻⁶
FD gasifier	SW	-32 °C	1.2x10 ⁻⁵
FD gasifier	BG+SW	-32 °C	6.2x10 ⁻⁶
FD gasifier	WH+SW	-32 °C	3.9x10 ⁻⁶
FD gasifier	RC+SW	-32 °C	4.4x10 ⁻⁶

915 Table 2: Summary of chamber experiments. The "*pot height*" experiments represent modified combustion conditions when the cooking pot was intentionally placed above the designated height of the respective cookstove. The doubly charged particle fractions and sizes are determined following Wiedensohler (1988). In cases where the measurements were not performed in the size range relevant for multiply charged particles, the estimates are obtained using lognormal fits to the detected parts of the soot modes.

Cookstove	Fuel	Sample size (nm)	Doubly charged fraction	
3-stone fire	SES	250	0.08	
3-stone fire	CAS	300	0.08	
3-stone fire	CAS	500	0.03	
Rocket stove	CAS	250	0.20	
Rocket stove	CAS	350	0.06	
Rocket stove	CAS	450	0.04	
ND gasifier	SW	400	-	
FD gasifier	CH+SW	260	0.13	
Modified FD gasifier	SW	450	0.08	
Modified FD gasifier	SW	500	0.06	
FD gasifier, pot height #1	SW	400	0.1	
FD gasifier, pot height #2	SW	400	0.18	
FD gasifier, pot height #3	SW	500	-	

Table 3: Ice onset (10^{-3} activation) temperatures when ascending temperature ramps were repeated during each chamber experiment, calculated from background-corrected signal. The last entry shows results from repeated ramps in the homogeneous freezing test. The uncertainties are equal to \pm one standard deviation in the lamina temperature at the moment of detection. The "*pot height*" experiments represent modified combustion conditions when the cooking pot was intentionally placed above the designated height of the respective cookstove. Acronyms ND and FD refer to natural-draft and forced-draft, respectively.

Cookstove	Fuel	Sample	Ice onset T [°C],	Ice onset T [°C],	Ice onset T [°C],
		size (nm)	ramp #1	ramp #2	ramp #3
3-stone fire	SES	250	-38.3±0.2	-38.2±0.2	-38.2±0.4
3-stone fire	CAS	300	-38.3±0.5	-38.2±0.3	-38.6±0.4
3-stone fire	CAS	500	-37.8±0.3	-38.0±0.2	-38.1±0.3
Rocket stove	CAS	250	-38.2±0.3	-38.1±0.3	-
Rocket stove	CAS	350	-37.8±0.3	-38.1±0.3	-
Rocket stove	CAS	450	-37.9±0.4	-37.9±0.3	-
ND gasifier	SW	400	-38.2±0.3	-38.2±0.3	-
FD gasifier	CH+SW	260	-37.8±0.3	-38.1±0.3	-
Modified FD gasifier	SW	450	-34.7±0.3	-35.1±0.4	-35.1±0.4
Modified FD gasifier	SW	500	-38.1±0.3	-	-
FD gasifier, pot height #1	SW	400	-33.4±0.3	-33.9±0.3	-34.0±0.3
FD gasifier, pot height #2	SW	400	-38.9±0.4	-38.9±0.4	-
FD gasifier, pot height #3	SW	500	-39.3±0.4	-	-
-	AS	350	-38.9±0.3	-39.1±0.3	-38.9±0.3