

Interactive comment on “Simultaneous Measurement of Urban and Rural Single Particles in Beijing, Part I: Chemical Composition and Mixing State” by Yang Chen et al.

Yang Chen et al.

chenyang@cigit.ac.cn

Received and published: 23 May 2020

Dear reviewer,

The authors are thankful for the reviewer's comments, suggestions, and time. We have prepared a point-by-point response highlighted in blue. We hope our efforts in revising the manuscript can improve it for the selection of ACP.

Please also see the Supplement for a pdf file.

Anonymous Referee #3

This work did a nice job in measuring the chemical composition and mixing states of

C1

[Printer-friendly version](#)

[Discussion paper](#)



Interactive comment

aerosols at an urban and a rural site in Beijing. More than 4 million particles were detected at each site, of which the chemically analyzed particles were grouped and analyzed to investigate the potential sources and atmospheric processing. The authors found that the urban particles were influenced significantly by rural processing and transport. The paper is generally well-written, and I recommend it for publication after some addressing the following comments.

Comments: 1. The authors listed several studies in paragraph 2 in Introduction, but didn't contain any conclusion. What does it mean by "discrepancies remain among these studies"? Authors need to be more specific about what discrepancies exist among the studies. Please revise/rephrase and be more specific.

Ans: We have added the following passage to illustrate the discrepancies and gaps in knowledge (lines 53-62): "For example, the mass loading of PM2.5 can rapidly increase to hundreds $\mu\text{g m}^{-3}$. Both Wang et al. (2016) and Cheng et al. (2016) suggested the secondary formation of sulfate from the oxidation of NO₂, while Guo et al. (2014) proposed a mechanism of particle formation and growth. Different from local secondary formation and accumulation, Li et al. (2015) proposed that particles via long-range transport cause the elevation of PM2.5. According to Sun et al. (2014) and Zhai et al. (2016), regional transport played an important role during heavy haze episodes. However, most studies have focused on the urban areas of Beijing, with limited attention paid to rural areas. To illustrate the sources, evolution, and transport of particles, the investigation of rural areas around Beijing is necessary."

2. It should be noted that chemical bias of SPAMS might introduce uncertainties in representativeness of ambient particles and even in classification of chemically analyzed particles. I think there is a need to mention this caveat in your paper, and caution the readers that uncertainties may be expected for the results. This can be provided either in Introduction or Discussions.

Ans: We completely agree with the reviewer that the limitations of the instrumentation

[Printer-friendly version](#)

[Discussion paper](#)



should be addressed (lines70-72). “Due to the nature of laser desorption/ionization (LDI), the instrument is very sensitive to dust and other types of particles containing sodium and potassium, and this may cause bias in the particle matrix (Pratt and Prather, 2012).”

3. Please clarify the differences between this study and previous studies. The significance of this study is not well written in the paper.

Ans: Associated with what we mentioned in Lines 76-83, we have added the following paragraph: “Organics, sulfate, nitrate, ammonium and other species have been found internally mixed in the atmospheric particles, and these particle types are mostly from the combustion of fuel or biomass. The abundance of secondary species can indicate the degree of aging during atmospheric processing. Particles are more secondary species with deeper processing. However, these studies lack the use of this data to provide a view of the dynamic particulate processing. Therefore, we used the relative abundance of secondary species to adequately illustrate the process of single particles at both sites, providing a tracing system on a regional scale.”

4. L150: The EC category has four types including EC-Nitrate (EC-Nit), EC-Sulfate (EC-Sul), and EC-Nit-Sul. Is it “four”, not “three”? Or there is another type?

Ans: Sorry, that was is a typo. We have changed it to “three.”

5. According to line 243, K-rich is one of branches of k-rich category. According to Figure 6a, K-rich means BB (BB in figure 6 caption, and K-rich in the figure) According to line 366, BB-related particle means K-rich category. The use of K-rich category, K-rich type, BB and BB-related particle is confusing. Please make the description more clear, specific and concrete.

Ans: Thank you very much for this suggestion. The K-rich category is from BB-related particles, and we will stick to this term. We have changed our description in the text, as well as the caption of Figure 6a.

[Printer-friendly version](#)

[Discussion paper](#)



6. As you mentioned in Line 253, the household BB is prohibited in urban Beijing, which is inconsistent with figure 6c indicating that the highest number counts of K-rich were observed when wind speed was less than 2 m s⁻¹, is that possible K-rich_PKU can also from other sources?

Ans: This is a very interesting question.

The urban sampling site is around the 4th ring expressway of Beijing, 10 km from the 5th ring expressway. In these areas, there are still villages in which biofuels are used. Therefore, these K-rich_PKU were also from biomass burning.

7. L295: "mass spectra of NaK category contained f Na+, K+ . . ." Please correct. Ans: We have made the correction (line 319).

References

Cheng, Y., Zheng, G., Wei, C., Mu, Q., Zheng, B., Wang, Z., Gao, M., Zhang, Q., He, K., and Carmichael, G.: Reactive nitrogen chemistry in aerosol water as a source of sulfate during haze events in China, *Science Advances*, 2, e1601530, 2016.

Guo, S., Hu, M., Zamora, M. L., Peng, J., Shang, D., Zheng, J., Du, Z., Wu, Z., Shao, M., Zeng, L., Molina, M. J., and Zhang, R.: Elucidating severe urban haze formation in China, *Proc Natl Acad Sci U S A*, 111, 17373-17378, 10.1073/pnas.1419604111, 2014.

Li, P., Yan, R., Yu, S., Wang, S., Liu, W., and Bao, H.: Reinstate regional transport of PM2.5 as a major cause of severe haze in Beijing, *Proc Natl Acad Sci U S A*, 112, 2015.

Pratt, K. A., and Prather, K. A.: Mass spectrometry of atmospheric aerosols—recent developments and applications. Part II: On-line mass spectrometry techniques, *Mass Spectrom. Rev.*, 31, 17-48, 10.1002/mas.20330, 2012.

Sun, Y., Jiang, Q., Wang, Z., Fu, P., Li, J., Yang, T., and Yin, Y.: Investigation of the

[Printer-friendly version](#)

[Discussion paper](#)



sources and evolution processes of severe haze pollution in Beijing in January 2013, J. Geophys. Res. Atmos., 119, 4380-4398, 10.1002/2014jd021641, 2014.

Wang, G., Zhang, R., Gomez, M. E., Yang, L., Zamora, M. L., Hu, M., Lin, Y., Peng, J., Guo, S., and Meng, J.: Persistent sulfate formation from London Fog to Chinese haze, Proceedings of the National Academy of Sciences, 113, 13630-13635, 2016.

Zhai, S., An, X., Zhao, T., Sun, Z., Hou, Q., and Wang, C.: Detecting critical PM2.5 emission sources and their contributions to a heavy haze episode in Beijing, China by using an adjoint model, Atmospheric Chemistry and Physics Discussions, 2016, 1-20, 10.5194/acp-2016-911, 2016.

Please also note the supplement to this comment:

<https://www.atmos-chem-phys-discuss.net/acp-2019-933/acp-2019-933-AC1-supplement.pdf>

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2019-933>, 2020.

Printer-friendly version

Discussion paper

