

1 **Supporting Information:**

2 **Contribution of HONO to the atmospheric oxidation capacity at an industrial zone**
3 **in the Yangtze River Delta region of China**

4 Jun Zheng^{1*}, Xiaowen Shi¹, Yan Ma^{1,2}, Xinrong Ren^{3,4,5}, Halim Jabbour¹, Yiwei Diao^{1,4}, Weiwei Wang⁴, Yifeng
5 Ge¹, Yuchan Zhang¹, and Wenhui Zhu¹

6 ¹Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, Nanjing University
7 of Information Science & Technology, Nanjing 210044, China

8 ²NUIST Reading Academy, Nanjing University of Information Science & Technology, Nanjing 210044, China

9 ³Air Resources Laboratory, National Oceanic and Atmospheric Administration, College Park, Maryland, USA

10 ⁴Department of Atmospheric and Oceanic Science, University of Maryland, College Park, MD 20742, USA

11 ⁵Cooperative Institute for Climate and Satellites, University of Maryland, College Park, Maryland, USA

12 ⁶Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Department of
13 Atmospheric Physics, Nanjing University of Information Science and Technology, Nanjing 210044, China

14

15 *Correspondence to: Dr. Jun Zheng (zheng.jun@nuist.edu.cn)*


16 *Address: School of Environmental Science and Engineering, Nanjing University of Information*

17 *Science & Technology, Nanjing 210044, China*

18 *Tel.: +86-18251919852*


19 *Fax: +86-25-58731090*

20 Reaction scheme of conversion of HONO into Azo dye before being detected:



(SA)

(Diazonium)



(NED)

(Azo dye)