

## Final author response

→ Thank you for your review and comments concerning our manuscript entitled “Significant seasonal changes in optical properties of brown carbon in the mid-latitude atmosphere.” We sincerely appreciate all valuable comments and suggestion. We thank all reviewers for taking the time and effort necessary to review our manuscript. We have concerned all your comments carefully, and all comments were addressed in the revised manuscript.

### Reviewer # 1

#### General comments:

This paper presents the results of stationary measurements of biomass burning aerosol in South Korea over the course of multiple seasons. The authors identify biomass burning as a major contributor to light-absorbing components of the aerosol, and assert that UV-radiation can explain the lower absorptivity of summer vs winter samples, with irradiation of winter samples resulting in properties similar to observed summer samples. The paper is generally well-written and clear, and the figures are appropriate for the claims being made. In some cases, the language used is too conclusive, and some places need clarifying, as requested below.

→ Thank you for your valuable comment. All your comments were carefully taken into account in the revised version.

#### Specific comments:

Abstract line 10: seasonal changes in sources and sinks of HULIS and WSOC (what was directly measured rather than approximated). The next sentence then relates the HULIS to total BrC.

→ We measured both HULIS (water-soluble BrC) and MeOH-soluble BrC for all samples (Fig. 2b and Fig. S4b). For the UV radiation experiments, we only measured HULIS as an index of BrC. Thus, we kept the wording unchanged.

Abstract line 17: this was supported by laboratory UV radiation experiments (confirm is too strong of language)

→ We changed the word “confirmed” to “supported.”

Lines 23–24: Black carbon aerosol should not be included in the light absorbing organic aerosol category, as black carbon is primarily soot (inorganic carbon)

→ We clarified this sentence in the revised version (lines 24–31).

Lines 92–93: Since the K fraction and WSON are determined using parameterizations instead of direct measurements, “estimated” is a more accurate term than “calculated.” For the K/Al\_crust parameterization, please provide the value used in addition to the reference from where it came.

→ We changed the word “calculated” to “estimated.” We also provide the values used for the parameterization, in addition to the reference in the revised version (lines 102–106).

Line 95: The sea spray fraction is calculated assuming a certain ratio of Na to Cl, (1.4486:1) – how does this ratio compare to the measured Cl/Na ratio? Since the Cl and Na is assumed to come entirely from seawater, this assumption can be verified.

→ Here, 1.4486 is the ratio of the concentration of all elements except Cl to the Na concentration in seawater (Maenhaut et al., 2007). The mass ratio of Cl/Na is 1.79 for seawater. The measured Cl/Na mass ratio ranged from 0.01 to 1.78. Since the maximum sea-salt concentration in our samples was about 0.1% of seawater, the highest value (1.78) verifies that our assumption is valid. Since this is commonly referred to and sea spray fraction is negligible in this study, we do not explain this in the text.

Line 98: Similarly, provide the crustal V/A; parameterization used, and replace “calculated” with “estimated.”

→ We provided the value and changed the word “calculated” to “estimated.”

Line 201: While Pearson’s r-squared value of 0.5 does represent a decent correlation in atmospheric data, by definition this means that 50% of the variation in HULIS is attributable to levoglucosan, not a majority, so the phrase “was the major source” might be slightly misleading.

→ We changed the word “major” to “significant” (lines 214–216).

Line 228: Similarly, although the negative correlation with r-squared of 0.5 indicates a strong relationship between UV radiation and HULIS, it only explains half of the variance in HULIS, and the system is more complex than it sounds from the current language.

→ Yes, we agree. Thus, we just state similar trends for UV and HULIS. Then, we state that “these results are in agreement with the following laboratory results” instead of linking UV and HULIS at this stage.

Line 242: the phrase “in the winter samples with high HULIS concentrations” leads one to conclude that there were winter samples with low HULIS as well as high HULIS. If, as I assume, the authors simply want to reinforce the idea that winter had higher HULIS, then perhaps this would be clearer, “in the winter samples (characterized by higher average HULIS concentrations), while no significant changes occurred in the summer samples (characterized by low average HULIS concentrations)”

→ We clarified this paragraph in the revised version (lines 252–255).

Line 253–254: Please rephrase this sentence so that it is easier to understand. I don’t quite understand what the authors are saying here.

→ We argue that the trend of “light-absorbing property” is similar to that of “fluorescence property.” We clarified this sentence in the revised version (lines 261–262).

Line 264: Given this interesting result (that irradiated winter samples look like summer samples” I wonder if there is more to be done with predicting summer properties from winter observations. It seems harder to back-calculate the properties of the aerosol emitted in summer (when still fresh), but I assume it would be similar to the winter samples. This interesting point deserves more attention, and I wonder if any previous studies have come to this conclusion, or a similar conclusion.

→ Thank you for your insightful comment. Although we cannot back-calculate the summer properties based on unknown UV radiation intensity, exposure time, and other atmospheric conditions, we assume a similar initial property based on other tracers including non-crystal K, air mass back trajectories (Fig. S1), and light absorption property in this study.

Line 271: The use of word “confirmed” here to describe how the laboratory studies can explain the observed differences in fluorescence is too strong. The studies certainly support the hypothesis, and they go a long way toward suggesting the mechanism, but given that other processes in the atmosphere (oxidative, or possibly aqueous aging) could also result in the same bleaching, the language should be softened until there are many further studies, or until signatures of bleaching can be tied specifically to the mechanism studied here.

→ We changed the word “confirmed” to “supported.”

#### Technical corrections:

Line 21: “and THE global climate system”

Line 28: remove plural on “light absorbing aerosols”

Line 40–41: remove space between number and percent symbol in all instances

Line 153: “originates”

Line 204: omit “the” before “BrC”

Line 244: there must be an extra word in this sentence “exhibited followed corresponding”

Line 248: remove plural on “carbon contents”

Line 251: omit “a” before “photo-labile”

Line 267: omit “the” before “different seasons”

→ We made all above technical corrections as suggested.

#### **Reviewer # 2**

Han and coauthors describe off-line analysis of filter samples collected over four seasons in Seoul, Korea in order to investigate relationships between brown carbon (BrC) and water-soluble organic carbon (WSOC) and how they change throughout the year. Potential sources are also discussed using complementary off-line techniques including high performance liquid chromatography (HPLC),

excitation-emission matrix (EEM) analysis and isotope ratio mass spectrometry (IRMS). The seasonal behaviour of the relationship between humic-like substances (HULIS) and WSOC is interesting, with a pronounced increase in the HULIS/WSOC ratio in the winter months. The authors propose that this pattern is due to more effective reduction of HULIS mas through photodegradation during atmospheric transport in the summertime when photolysis rates and photochemical activity are higher. The absence of a seasonal pattern for potassium, in particular adds weight to the argument that the HULIS/WSOC ratio seasonality cannot be explained only by changes in biomass burning source contributions throughout the year. Although levoglucosan exhibits higher wintertime values, this tracer can also be degraded through photochemical processing. Overall I find the manuscript to be well written and structured and only have some minor comments:

→ Thank you for your review and valuable comments. All your comments were addressed in the revised manuscript.

In the introduction it is mentioned that “the origin of BrC is often attributed predominantly to HULIS”. It would be helpful to discuss known sources of HULIS here.

→ We added more about the sources of HULIS in the introduction section in the revised version (lines 35–42).

Through EEM analysis, the extracted HULIS component agreed temporally with the extracted HULIS concentration as shown in S4. How do the other extracted EEM factors agree temporally with HULIS concentration? If the agreement is strong between all three factors, then it is likely that they are associated with the same source. If C2 is characterized by tryptophan then is vegetative debris a likely source for this factor? Are C1 and C3 interpreted to have the same source but differ only in their EEM characteristics? This section (3.1) could be clearer.

→ Since the other EEM spectra are highly influenced by one dominant fluorophore HULIS (C1) as shown in Figure S7 (right), we cannot reliably characterize C2 component using PARAFAC model. Thus, we only use C1 component in this study. This is mentioned in the text (lines 165–170).

Either in the Introduction or Discussion it would be helpful to discuss the findings here in the context of previous on-line measurement studies that have observed similar photodegradation of brown carbon from biomass burning sources during atmospheric summer time transport, for example (Forrister et al., 2015; Selimovic et al., 2019; Healy et al., 2019).

→ Yes, this is mentioned in the discussion section in the revised version (lines 244–247).

Define “RU”

→ The detailed definition about “RU” is added in the revised manuscript (lines 133–136).

Line 186: What is the likely source of the non-crystal V- oil combustion, shipping? It would be useful to discuss

→ Since vanadium is mostly removed during the refining processes, the use of raw materials such as crude oil and coal can be the source of non-crystal vanadium. This is mentioned in the revised manuscript (lines 231–234).

Line 195: Define C3 plant-origin materials

→ We mentioned about C3 plant-origin materials in the revised manuscript (lines 205–207).


Line 209-210: The last line of this paragraph is unclear and should be rephrased.

→ Yes, we rephrased this more clearly in the revised version (lines 221–222).

Line 244: Remove “followed”

→ removed in the revised version.

[END]



# Significant seasonal changes in optical properties of brown carbon in the mid-latitude atmosphere

Heejun Han<sup>1</sup>, Guebuem Kim<sup>1</sup>, [Hojong Seo<sup>1</sup>](#), Kyung-Hoon Shin<sup>2</sup>, Dong-Hun Lee<sup>2</sup>

<sup>1</sup>School of Earth and Environmental Sciences/RIO, Seoul National University, Seoul, 08826, Korea

5 <sup>2</sup>Department of Marine Sciences and Convergent Technology, Hanyang University, Ansan, 15588, Korea

Correspondence to: Guebuem Kim (gkim@snu.ac.kr)

**Abstract.** Atmospheric brown carbon (BrC) plays significant roles in the light absorption and photochemistry of the atmosphere. Although the [sources and occurrences](#) of BrC have been studied extensively, its removal processes and optical characteristics in the atmosphere have been poorly understood. In this study, we examined the seasonal changes in sources and sinks of BrC and water-soluble organic carbon (WSOC) in the atmosphere of Seoul, Korea. Our results showed that the concentrations of BrC and WSOC decreased by approximately 80% and 30%, respectively, from the cold season (Oct–Jan) to the warm season (Jun–Sep). Excitation–emission matrix (EEM) spectra showed that the humic-like substance (HULIS) was the dominant fraction of BrC as the other components were not measurable. The air mass back trajectories of fire burning practices and the variations in [non-crustal potassium \(K\)](#) and [vanadium \(V\)](#) contents in the water-soluble aerosols during all seasons showed no measurable decrease in input of biomass-burning sources in summer. However, there was a significant shift in photo-resistivity of light-absorbing organic aerosols in the summer, indicating larger removals of ultraviolet (UV) degradable BrC. This [trend is supported by laboratory UV radiation experiments on the optical property changes of BrC and WSOC in aerosol samples](#). Thus, our results suggest that the photo-degradation has dominant roles in controlling the quantity and quality of light-absorbing organic aerosols in the different seasons in the mid-latitude atmosphere.

## 1 Introduction

Organic aerosols play a significant role in atmospheric chemistry and [the global climate system](#) (Ghan and Schwartz, 2007; Duarte and Duarte, 2013; Kirillova et al., 2014a). Most organic aerosols consist of a significant fraction of carbonaceous organic aerosols absorbing radiation (Laskin et al., 2015). These light-absorbing organic aerosols, [referred to as brown carbon \(BrC\)](#), have attracted increasing attention owing to their significant roles in the radiative forcing of the global climate system by directly absorbing solar radiation (Andreae and Gelencsér, 2006; Graber and Rudich, 2006; Ramanathan et al., 2007; Kirillova et al., 2014a; Saleh et al., 2014) and indirectly acting as cloud condensation nuclei (CCN) for cloud formation (Kanakidou et al., 2005; Andreae and Gelencsér, 2006; Graber and Rudich, 2006; Kirillova et al., 2014a). [The light-absorbing property of BrC, which absorbs solar radiation in the range of ultraviolet \(UV\) to visible wavelengths,](#) is

Author  
Deleted:

Author  
Deleted: occurrence and

Author  
Deleted:

Author  
Deleted:

Author  
Deleted: wa

Author  
Deleted: confirmed

Author  
Deleted: (or light-absorbing carbonaceous aerosols) including black carbon (BC) and

Author  
Deleted: BC is the most commonly known light-absorbing aerosols, which absorbs solar radiation over a wide spectral range, from ultraviolet (UV) to near-infrared radiation, while BrC highly absorbs radiation in the range of UV to visible wavelengths (Feng et al., 2013; Laskin et al., 2015).

Author  
Moved down [1]: (Feng et al., 2013; Laskin et

Author  
Deleted: Although t

50 expected to be considerably weaker than that of the [black carbon \(BC\)](#), which absorbs over a wide spectral range from UV to near-infrared radiation ([Feng et al., 2013](#); [Laskin et al., 2015](#)). However, its contribution would be significant due to its higher abundances over source regions ([Hoffer et al., 2006](#); [Gustafsson et al., 2009](#); [Feng et al., 2013](#)). Previous studies have estimated that the BrC contributes to approximately 19% of the total atmospheric radiative forcing ([Feng et al., 2013](#)).

55 The origin of BrC is attributed predominantly to atmospheric humic-like substance (HULIS), which has properties similar to [terrestrial and aquatic humic and fulvic acids](#) ([Andreae and Gelencsér, 2006](#); [Graber and Rudich, 2006](#); [Lukács et al., 2007](#); [Laskin et al., 2015](#); [Yan and Kim, 2017](#)). Previous studies have identified that HULIS has both direct emission sources from biomass burning, marine aerosols, and soil re-suspension, and indirect contribution from secondary formation ([Graber and Rudich, 2006](#); [Hoffer et al., 2006](#); [Lin et al., 2010](#); [Yan and Kim, 2017](#)). Recently, residential coal combustion was suggested as an important source of HULIS during gold seasons ([Tan et al., 2016](#); [Voliotis et al., 2017](#); [Li et al., 2019](#)). Besides HULIS, biological aerosols (e.g., bacteria, algae, pollen, spores, fungi, plant debris, and animal tissue) are another important sources of BrC ([Després et al., 2012](#); [Pöhlker et al., 2012](#)). In general, the atmospheric HULIS contributes to approximately 70% of the water-soluble organic carbon (WSOC) ([Laskin et al., 2015](#); [Park and Son, 2017](#)), while the WSOC contributes to 10–80% of the total organic carbon contents in aerosols ([Kirillova et al., 2014a](#); [Kirillova et al., 2014b](#); [Fu et al., 2015](#)). However, current understanding of the optical properties, chemical compositions, and degradation processes of light-absorbing organic aerosols still remain uncertain.

65 In this study, we evaluated the seasonal changes in optical and chemical characteristics, including photo-resistivity, of light-absorbing organic aerosols in an urban region. Recently, the method of excitation–emission matrix (EEM) characterization combined with a parallel factor analysis (PARAFAC) model has been employed for BrC studies ([Kieber et al., 2006](#); [Mladenov et al., 2011](#); [Matos et al., 2015](#); [Chen et al., 2016](#); [Yan and Kim, 2017](#)). This unique multi-analysis method can be used to identify individual chromophores of aerosol samples (i.e., HULIS versus protein-like substance) and light-absorbing properties of each sample. In addition, the stable carbon isotope ratios of WSOC ( $\delta^{13}\text{C}_{\text{WSOC}}$ ) and various chemical constituents were measured to identify the potential sources of organic aerosols in different seasons. Furthermore, we conducted laboratory experiments on the direct photochemical degradations of aerosol samples to verify the impacts of the UV radiation on the optical properties and carbon compositions of BrC and WSOC in the atmosphere.

## 2 Experimental methods

### 75 2.1 Study site and sample collection

Seoul is the capital city of South Korea and one of the largest metropolitan cities in the world. Korea has been highly affected by severe dust storms known as Asian dust or the Yellow dust originated from the Chinese and Mongolian deserts during the spring (Mar–May) and often during the winter (Dec–Jan) ([Lin et al., 2012](#)). The increase in contents of fine

Author

Deleted: solar radiation

Author

Moved (insertion) [1]

Author

Deleted:

Author

Deleted: In general, BrC is known to originate from various sources such as biomass burning, incomplete combustion process, secondary formation, volatile organic compounds, and soil humics ([Andreae and Gelencser, 2006](#); [Graber and Rudich, 2006](#); [Hoffer et al., 2006](#); [Ramanathan et al., 2007](#); [Kivácsy et al., 2008](#); [Saleh et al., 2014](#); [Laskin et al., 2015](#); [Jiang et al., 2019](#)).

Author

Deleted: winter periods

Author

Deleted:

Author

Deleted:

Author

Deleted: of

Author

Deleted: and changes in photo-resistivity of light-absorbing organic aerosols in the atmosphere

95 aerosols associated with anthropogenic emissions is of great concern and a focus of major environmental studies in this region (Seinfeld et al., 2004; Park et al., 2007).

100 Aerosol samples ( $N=78$ ) were collected using a high volume air sampler (HV-1000, SHIBATA) from March 2015 to January 2016 in Seoul, Korea ( $37.5^\circ$  N,  $127.0^\circ$  E; 20 m above ground level) (Fig. 1). The samples were collected for 24 h at a constant flow rate of  $1000 \text{ L min}^{-1}$  through a pre-combusted ( $450^\circ\text{C}$  for 5 h) glass microfiber filter (GF/F,  $8 \times 10$  inch,  $2 \mu\text{m}$  pore size, Whatman). A blank sample was collected by shortly exposing a blank filter at the study site and analyzed in the same manner as that for the other samples. The total suspended particulates (TSP) of the aerosol samples were measured by using the mass differences in desiccated filters between the pre- and post-sampling. The collected samples were covered with an aluminium foil, placed in a polyethylene bag, and stored in the dark at  $-20^\circ\text{C}$ .

105 Meteorological parameters of the study site including the temperature and UV radiation rate were obtained from the ambient air quality monitoring network named AirKorea in the Korea Environmental Corporation (KECO) and Korea Meteorological Administration (KMA). Fire maps around the study site were obtained by using the moderate resolution imaging spectroradiometer (MODIS) fire location data provided by NASA's fire information for resource management system (FIRMS) (Fig. S1).

115 A ten-day air mass back trajectory was drawn by using the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPPLIT) model to determine the source regions and transport pathways of air masses to the study site (Stein et al., 2015) (Fig. 1). Due to a regional meteorology of the study site, which is dominated by the East Asian Monsoonal effect, most air masses are transported from the arid and semi-arid regions in the Asian continent for most of the year (Yan and Kim, 2012; Yan and Kim, 2017)

## 2.2 Aerosol extraction and chemical analyses

120 For the analyses of water-soluble organic aerosols, a small portion of filter paper was cut into small pieces and placed in a pre-HCl-rinsed bottle. The organic components in the sample were extracted using Milli-Q water ( $18.2 \text{ M}\Omega \text{ cm}$ ) shaken at 125 rpm for 4 h (Wozniak et al., 2012). Then, the extracts were filtered through a syringe filter ( $0.45 \mu\text{m}$  pore size Nucleopore, Whatman) and analyzed to evaluate the WSOC and total dissolved nitrogen (TDN) by a high-temperature oxidation method using a total organic carbon (TOC) analyzer (TOC-V<sub>CPH</sub>, Shimadzu). The major ion species ( $\text{NO}_x$ ,  $\text{NH}_4^+$ ,  $\text{SO}_4^{2-}$ ,  $\text{Ca}^{2+}$ ,  $\text{Cl}^-$ ,  $\text{Na}^+$ , and  $\text{K}^+$ ) were analyzed by using a high performance liquid chromatography (HPLC, Waters 2695) system equipped with a conductivity detector (Waters 432) (Yan and Kim, 2015). The subsamples for trace element analyses were acidified to 2% in ultra-pure  $\text{HNO}_3$  and analyzed by using a high-resolution inductively coupled plasma mass spectrometer (HR-ICP-MS, Thermo Element 2). In order to correct the instrumental instability, Rh was included as internal

Author  
Deleted:

Author  
Deleted: ) (

Author  
Deleted: system

Author  
Moved (insertion) [2]

Author  
Deleted: dissolved

standard. The molecular marker compound, levoglucosan, was measured by using a gas chromatography-mass spectrometry (GC-MS, 7890N, Agilent) system coupled with a fused silica capillary column (HP-5MS, 25 m length, 0.25 mm i.d. and film thickness of 0.10 μm). Monthly representative filter samples were used for the analysis (N=25). The water-soluble organic nitrogen (WSON) concentration was calculated by using the concentration difference between the TDN and sum of inorganic nitrogen species (NO<sub>2</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>, and NH<sub>4</sub><sup>+</sup>).

The non-crustal potassium (K) and vanadium (V) fractions were estimated by using the following equation (Yan and Kim, 2012):

$$[non - crustal X] = [X] - ([X]/[Al])_{crust} \times [Al]_{aerosol} \quad (1)$$

For these calculations, the element compositions of the upper continental crust for K, Al, and V are assumed to be 2.80, 8.04, and 0.006 wt. %, respectively (Taylor and McLennan, 1995). The sea-spray fraction was estimated by using Cl<sup>-</sup> and Na<sup>+</sup> concentrations assuming that all Cl<sup>-</sup> and Na<sup>+</sup> are originated from seawater: sea-spray = Cl<sup>-</sup> + 1.4486 Na<sup>+</sup> (Maenhaut et al., 2007).

The value of δ<sup>13</sup>C<sub>WSOC</sub> was measured by using an isotope ratio mass spectrometer (IRMS) (Isoprime, Elementar) combined with a TOC analyzer (Vario TOC cube, Elementar) (Panetta et al., 2008; Troyer et al., 2010; Kim et al., 2015; Yan and Kim, 2017). The isotopic composition δ<sup>13</sup>C<sub>WSOC</sub> was determined using the following equation:

$$\delta^{13}C = \left( \frac{(^{13}C/^{12}C)_{Sample}}{(^{13}C/^{12}C)_{Standard}} - 1 \right) \times 1000 \text{ ‰} \quad (2)$$

where Vienna Pee Dee Belemnite (VPDB) was used as the isotope standard (Kelly et al., 2005; Troyer et al., 2010; Fu et al., 2015). Analytical tests were made with IAEA-CH6 sucrose (δ<sup>13</sup>C = -10.45±0.03 ‰) and Suwannee River Fulvic Acid (SRFA) (δ<sup>13</sup>C = -27.6±0.12 ‰; International Humic Substances Society) to evaluate the recovery and the accuracy of the measurements (Panetta et al., 2008; Troyer et al., 2010).

The HULIS fraction was separated by solid-phase extraction (SPE) by using the DEAE column (GE Healthcare®, HiTrap™ DEAE FF, 0.7 cm ID × 2.5 cm) to validate the humic fraction obtained by using the EEM-PARAFAC results (Baduel et al., 2009). The sample solutions were passed through the column at a constant flow rate of 1.0 mL min<sup>-1</sup>. The neutral components and hydrophobic bases, and mono-, di-carboxylic acids, and inorganic anions were removed with Milli-Q water and 0.04 M NaOH, respectively. The polycharged compounds, HULIS, were eluted with 1M NaCl. The final fractions were analyzed for the HULIS quantification using a TOC analyzer.

### 2.3 Fluorescence and absorbance measurements

Author

**Moved (insertion) [3]**

Author

**Deleted:** ) (

Author

**Moved (insertion) [4]**

Author

**Deleted:** as

Author

**Deleted:** calculated

Author

**Deleted:** [nc - K] = [K] - ([K]/[Al])<sub>crust</sub> × [Al]<sub>aerosol</sub>

Author

**Moved up [4]:** The water-soluble organic nitrogen (WSON) concentration was calculated by using the concentration difference between the TDN and sum of inorganic nitrogen species (NO<sub>2</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>, and NH<sub>4</sub><sup>+</sup>).

Author

**Deleted:** calculated estimated

Author

**Deleted:** ted

Author

**Moved up [2]:** The subsamples for trace element analyses were dissolved in ultra-pure HNO<sub>3</sub> and analyzed by using a high-resolution inductively coupled plasma mass spectrometer (HR-ICP-MS, Thermo Element 2).

Author

**Deleted:** ). The non-crustal vanadium (V) fraction was calculated by using the following equation: [nc - V] = [V] - ([V]/[Al])<sub>crust</sub> × [Al]<sub>aerosol</sub> (Taylor and McLennan, 1995; Yan and Kim, 2012). The molecular marker compound, levoglucosan, was measured by using a gas chromatography-mass spectrometry (GC-MS) (7890N, Agilent) system coupled with a fused silica capillary column (HP-5MS, 25 m length, 0.25 mm i.d. and film thick... [1]

Author

**Moved up [3]:** The molecular marker compound, levoglucosan, was measured by using a gas ... [2]

Author

**Deleted:** The HULIS fraction was separated by solid-phase extraction (SPE) by using DEAE ... [3]

Author

**Deleted:** l

Author

**Deleted:** analyse

Author

**Deleted:** i



230 Fluorescence EEM and absorbance spectra of the aerosol samples were measured by using a spectrophotometer (Aqualog, Horiba). The emission and excitation wavelength ranges were 240 to 700 nm and 250 to 500 nm, respectively, with scanning intervals of 1 nm. The PARAFAC model was performed using the Solo software in order to determine the [different](#) fluorescent components in aerosol [samples](#). The results were validated by split-half analysis and random initialization (Bro, 1997; Zepp et al., 2004; Stedmon and Bro, 2008). [The Raman and Rayleigh scattering signals, inner-filter effect, and blank subtraction were corrected using the Solo software](#). The [fluorescence intensity was normalized by Raman peak area of water](#) and stated as Raman Unit (RU) (Lawaetz and Stedmon, 2009). [This is a simple calibration method for quantitative fluorescence spectroscopy using the integrated area of water Raman peak at an excitation wavelength of 350 nm \(Lawaetz and Stedmon, 2009\)](#).

240 The absorption coefficient was calculated using the following equation:

$$a = 2.303 A/L, \quad (3)$$

where  $a$  is the absorption coefficient ( $\text{m}^{-1}$ );  $A$  is the absorbance;  $L$  is the optical path length of the quartz cuvette (m). The spectral slope ( $S$ ) was calculated using following equation:

$$a_\lambda = a_{\lambda_{ref}} e^{-S(\lambda - \lambda_{ref})}, \quad (4)$$

245 where  $a$  is the [Napierian](#) absorption coefficient ( $\text{m}^{-1}$ );  $\lambda$  is the wavelength;  $\lambda_{ref}$  is the reference wavelength (Twardowski et al., 2004; Helms et al., 2008). In this study, two distinct spectral ranges were selected in the shorter wavelength (275–295 nm) and the longer wavelength (350–400 nm).

#### 2.4 Photochemical degradation experiments for light-absorbing organic aerosols

250 Laboratory experiments were conducted for two different durations ([12 h and 42 d](#)) using small portion of the aerosol filter samples inside of the incubator equipped with internal UV lamp (UV-B; 280–315 nm; 15 W). The filter samples for the experiment were duplicated or triplicated to rule out any experimental bias. The incubator was maintained under positive air pressure with a constant temperature at 20°C. Aerosol samples were directly exposed to the UV radiation as its original particulate form, and each of the non-irradiated aerosol samples was evaluated for the initial conditions. For a long-term test, two representative aerosol filter samples were randomly selected from each of the higher and lower fluorescent groups, which had a similar range of WSOC concentration (winter = 19  $\mu\text{M}$ ; summer = 16  $\mu\text{M}$ ). After [the UV irradiation](#), filter samples were completely covered to prevent exposure to any light. [The extraction of organic components was performed](#) using the same procedure as described above. Each sample was analysed for optical and chemical properties after the irradiation.

### 260 3 Results

#### 3.1 BrC compositions

Author  
Deleted: The number of unique components was identified from the combined EEM data of aerosol samples.

Author  
Deleted: EEM results

Author  
Deleted: ere

Author  
Deleted: with

Author  
Deleted: Raman and Rayleigh scattering signals, inner-filter effect, and blank subtraction were corrected using the Solo software.

Author  
Deleted:  $\alpha$

Author  
Deleted: 2

Author  
Deleted: and

Author  
Deleted: 3

Author  
Deleted:

Author  
Deleted: and

Author  
Deleted:

Author  
Deleted: stimulated

Author  
Deleted: and directly

Author  
Deleted: extract

Author  
Deleted: ed



The PARAFAC model identified two humic-like fluorescent components and one protein-like fluorescent component (Fig. S2). The spectral characteristics of component 1 (C1; Ex/Em = 305/416 nm) and component 3 (C3; Ex/Em = 365/484 nm) are known to be highly associated with the atmospheric HULIS (Chen et al., 2016; Yan and Kim, 2017). Component 2 (C2; Ex/Em = 290/340 nm) is associated with a protein-like (tryptophan-like) component, which originates mostly from biological production (Coble, 2007; Birdwell and Engel, 2010; Yan and Kim, 2017) (Table. S1). However, C2 seems to be largely influenced by the dominant fluorophore HULIS, as shown by the EEM spectra (Fig. S7). A good correlation between C1 and C2 ( $r^2 = 0.9$ ;  $p < 0.05$ ) was obtained, although, in general, these two components have different sources and sinks in the atmosphere (Yan and Kim, 2017) (Fig. S3). Thus, we assumed that the HULIS (C1 and C3) was the dominant component of BrC in these samples, and C1 was used as a representative component of BrC in this study, as C1 and C3 exhibited a good correlation ( $r^2 = 0.8$ ;  $p < 0.05$ ) (Fig. S3).

The HULIS component obtained by the EEM-PARAFAC model agreed very well with the extracted HULIS concentration obtained by using the DEAE column ( $r^2 = 0.81$ ;  $p < 0.05$ ), indicating that the C1 component represents the actual HULIS (Fig. S4a). In addition, we compared the extraction efficiencies of HULIS (water-soluble BrC) to that of the MeOH-soluble BrC (Fig. S4b). Although the efficiency of the water-soluble BrC was approximately 20% lower than that of the MeOH-soluble BrC, they exhibited a good correlation ( $r^2 = 0.9$ ;  $p < 0.05$ ) (Fig. S4b). Therefore, we conclude that the HULIS is a good representative component of BrC in this study.

### 3.2 Seasonal variations in BrC and WSOC

The WSOC concentrations exhibited seasonal variations in the range of 3 to 40  $\mu\text{g m}^{-3}$  (average =  $16 \pm 7 \mu\text{g m}^{-3}$ ) with higher values during the cold seasons (Oct–Jan) (average =  $18 \pm 7 \mu\text{g m}^{-3}$ ) and lower values during the warm seasons (Jun–Sep) (average =  $13 \pm 3 \mu\text{g m}^{-3}$ ) (Fig. 2a). Similarly, the HULIS concentrations exhibited seasonal variations in the range of 13 to 294 RU (average =  $108 \pm 77 \text{RU}$ ) with higher values during the cold seasons (average =  $152 \pm 76 \text{RU}$ ) and lower values during the warm seasons (average =  $46 \pm 20 \text{RU}$ ) (Fig. 2b). After the HULIS content was normalized to the WSOC contents, the seasonal trend of the ratio of the HULIS content to the WSOC content was similar to that of the HULIS concentration, indicating that there was pronounced decrease in the average fraction of HULIS relative to WSOC from the cold to the warm seasons (Fig. 2c).

The  $\delta^{13}\text{C}_{\text{WSOC}}$  values ranged from -21.0 to -27.5 ‰ (average =  $-24.0 \pm 1.5 \text{‰}$ ), with no such seasonal variation trend (Fig. 2d). The levoglucosan concentration was in the range of 0.5 to 2.2  $\mu\text{g m}^{-3}$  (average =  $1.1 \pm 0.5 \mu\text{g m}^{-3}$ ) with relatively higher values during the cold seasons and the lower values during Aug–Sep (Fig. 2e). The temporal variations in major ion species concentrations did not show such seasonal behaviours throughout the year (Fig. 2f–2k). The temporal variation in  $\text{Ca}^{2+}$  concentration (average =  $0.8 \pm 0.2 \text{mg L}^{-1}$ ) exhibited a relatively constant level throughout the year (Fig. 2f). The  $\text{SO}_4^{2-}$

Author  
Deleted: thus

Author  
Deleted: ,

Author  
Deleted: which

Author  
Deleted: es

Author  
Deleted:

Author  
Deleted: ,

Author  
Deleted: which

Author  
Deleted: es

Author  
Deleted: in

Author  
Deleted: ,

Author  
Deleted: in the

Author  
Deleted: of

Author  
Deleted: ,

Author  
Deleted: exhibited

Author  
Deleted: exhibit

330 concentrations (average =  $12.0 \pm 10.3 \text{ mg L}^{-1}$ ) exhibited the highest value during the period of spring to summer and were slightly decreased during the cold seasons (Fig. 2g). The concentrations of  $\text{NO}_x$  (average =  $11.7 \pm 8.7 \text{ mg L}^{-1}$ ) were high during the warm periods (Jun–Oct) (Fig. 2h). The concentrations of non-crystal K (average =  $0.4 \pm 0.2 \text{ mg L}^{-1}$ ) also showed a relatively constant level throughout the year (Fig. 2i). The non-crystal V concentrations (average =  $0.01 \pm 0.01 \text{ mg L}^{-1}$ ) showed no such seasonal variations (Fig. 2j). The sea-spray concentrations ranged from 4.0 to  $33.6 \text{ mg L}^{-1}$  (average =  $17.3 \pm 6.4 \text{ mg L}^{-1}$ ), showing the highest concentration in July (Fig. 2k). Both temperature and UV radiation rate showed typical patterns of middle latitudes (Fig. 2l).

#### 4 Discussion

340 In order to characterize the sources of HULIS and WSOC in Seoul, which may cause the seasonal variabilities, we analyzed various tracers including  $\delta^{13}\text{C}_{\text{WSOC}}$ , major ions, and molecular marker as source indicators (Gabriel et al., 2002; Kelly et al., 2005; Kirillova et al., 2014b; Fu et al., 2015; Yan and Kim, 2015). The average  $\delta^{13}\text{C}_{\text{WSOC}}$  ( $-24.0 \pm 1.5 \text{ ‰}$ ) suggests that the burning activity of terrestrial C3 plant-origin materials (e.g., grassland, crop-residue, fuel-wood) could be a major source of WSOC in this region (Kelly et al., 2005; Das et al., 2010; Kawashima and Haneishi, 2012; Fu et al., 2015). The average  $\delta^{13}\text{C}_{\text{WSOC}}$  was in good agreement with that of HULIS ( $-25.4 \pm 1.6 \text{ ‰}$ ) extracted from precipitation in Seoul, Korea (Yan and Kim, 2017).

The good correlation ( $r^2 = 0.6$ ;  $p > 0.05$ ) between the WSOC and WSON concentrations suggests that these two variables are highly associated with a common organic source (Yan and Kim, 2015) (Fig. 3a). The WSOC and HULIS concentrations also exhibited a good correlation ( $r^2 = 0.5$ ;  $p < 0.05$ ) (Fig. 3b). A good linear correlation ( $r^2 = 0.5$ ;  $p < 0.05$ ) was also observed 350 between the HULIS and levoglucosan concentrations (Fig. 3c). This relationship indicates that biomass burning is a significant source of HULIS in Seoul, as levoglucosan is commonly used for the tracing of biomass burning (Fu et al., 2015; Kuang et al., 2015). This is consistent with the previous study, which has demonstrated that the BrC in the precipitation was primarily derived from biomass burning and terrestrial biogenic emissions ( $>70\%$ ), with minor contributions from fossil-fuel combustion, based on the  $^{14}\text{C}$  values in the HULIS (Yan and Kim, 2017). However, we observed significant decreases 355 ( $35 \pm 2\%$ ) in levoglucosan concentrations by the UV irradiation for 12 to 24 h in the laboratory experiments (Fig. S5). This is consistent with the previous studies showing that levoglucosan could be oxidized by the hydroxyl radical in the atmosphere (Hennigan et al., 2010; Hoffmann et al., 2010). Thus, we suggest that the major source of HULIS in our samples was from biomass burning, but low concentrations in WSOC and HULIS in summer could not be evaluated by using the levoglucosan.

360 In order to indirectly evaluate the biomass burning during the study period, we compiled the fire maps in combination with air mass back trajectories (Fig. S1). An evidence for the biomass burning effect is provided by the fire maps, where higher occurrences of fire spots correspond to agricultural burning practices in the East Asia continents (Fig. S1). The burning

Author  
Deleted: exhibited

Author  
Deleted: and

Author  
Deleted: ed

Author  
Deleted: gradually changed during the different seasons

Author  
Deleted: wa

Author  
Deleted: the major

Author  
Deleted:

Author  
Deleted:

Author  
Deleted: conclud

Author  
Deleted: e

Author  
Deleted: the summer decreases

Author  
Deleted: concentrations

Author  
Deleted: se tracers

practices mainly occurred in the spring and summer in the East Asia continents. No significant influence of open burning activity was observed in the winter (Fig. S1). These results suggest that the distinctive biomass burning was not linked to the seasonal variations in WSOC and HULIS concentrations in this study region. On the other hand, non-crystal K, which is an indicator of biomass or fossil fuel burning (Gabriel et al., 2002; Baduel et al., 2010), did not exhibit a seasonal trend (Fig. 2i). The correlation between the HULIS and non-crystal K was also insignificant ( $r^2 = 0.3$ ;  $p < 0.05$ ) (Fig. 3d). The K/V ratio is often used to trace V-purified fossil fuels, since V is mostly removed during the refining process (Yan et al., 2012). The used of raw materials such as crude oil and coal could be the source of non-crystal V (Tsukuda et al., 2005; Yan and Kim, 2012). However, the correlation between HULIS and non-crystal V was also insignificant ( $r^2 = 0.04$ ;  $p < 0.05$ ) (Fig. 3e), with K/V ratio in the range of 17–673 (average =  $124 \pm 154$ ), in our samples. This result suggests that the V-purified fossil fuel contents in our samples were insignificant (Yan et al., 2012). No significant correlations were observed between the HULIS and  $Ca^{2+}$  ( $r^2 = 0.1$ ;  $p < 0.05$ ),  $NO_x$  ( $r^2 = 0.1$ ;  $p > 0.05$ ),  $SO_4^{2-}$  ( $r^2 = 0.02$ ;  $p < 0.05$ ), and sea-spray fraction ( $r^2 = 0.02$ ;  $p > 0.05$ ) (Fig. 3f–3h). This also indicates that the seasonal changes in HULIS concentrations were insignificantly influenced by crustal minerals, sea salts, and fossil fuels. All these tracers suggest that the summer decreases in WSOC and HULIS concentrations were not associated with the changes in source inputs.

A significant negative correlation ( $r^2 = 0.5$ ;  $p < 0.05$ ) was observed between HULIS and the UV radiation (Fig. 3i). The HULIS was greatly reduced during the warm seasons (Jun–Sep) when the solar UV radiation and the temperature reached the annual maxima (Fig. 2b, 2l). It has been widely accepted that photochemical degradation is an important process for the efficient removal or alteration of photo-resistivity and compositions of chromophoric dissolved organic matter (CDOM) in rainwater and aerosols (Kieber et al., 2006; Mladenov et al., 2009; Forrister et al., 2015; Dasari et al., 2019; Healy et al., 2019). Significant seasonal changes in ratios of the HULIS concentration to the WSOC concentration, although the WSOC and HULIS concentrations exhibited similar seasonal changes, also suggest that HULIS is more effectively removed by the UV radiation in summer (Fig. 2c). These results are in agreement with the following laboratory experimental results.

In order to quantify the UV-degradable HULIS, we performed the UV radiation experiments for short-term (12 h; 48 samples) and long-term (42 d; 2 samples). After the 12 h of UV irradiation, the HULIS concentrations in the winter samples (characterized by higher average HULIS concentrations,  $N=35$ ) were greatly reduced by 13% (Fig. 4a). Similar decreases were observed for the WSOC concentrations for the winter samples (Fig. 4b). However, negligible changes (<2%) were observed in the summer samples (characterized by low average HULIS concentrations,  $N=13$ ) (Fig. 4a). After the 42 d of UV irradiation, the HULIS and WSOC concentrations in the winter sample decreased by approximately 52% and 25%, respectively (Fig. S6 and S7). However, there was no significant change in HULIS and WSOC concentrations in the summer sample (Fig. S6 and S7). This result suggests that the summer sample might consist largely of a photo-refractory WSOC pool owing to the high UV radiation, while the winter sample consisted largely of a photo-labile WSOC.

Author  
Moved (insertion) [5]

Author  
Deleted: T...e correlation between HULIS ... [4]

Author  
Moved up [5]: The K/V ratio is often used to trace V-purified fossil fuels (Yan et al., 2012).

Author  
Deleted: The average .../V ratio was ...n t... [5]

Author  
Deleted: that absorbs and fluoresces at certain wavelengths of light under solar radiation in aquatic ecosystems, which alters absorption in the UV region, spectral shape, and CDOM composition (Helms et al., 2013; Mladenov et al., 2009). In addition, previous studies have demonstrated the significant photochemical degradation of CD... [6]

Author  
Deleted: UV radiation experiments... Aftc... [7]

Author  
Deleted: with high HULIS concentrations, while no significant changes occurred in the summer samples with low HULIS concentrations (Fig. 4). The HULIS concentration ...reas...greatly r... [8]

Author  
Formatted: Font:Italic

485 The change in light-absorbing property of HULIS for the UV radiation experiment also exhibited a similar change with that of fluorescence property (Fig. 5). The absorption coefficient of the summer sample was not changed after the irradiation, while that of the winter sample exhibited an approximately twofold change (Fig. 5). The degradation of HULIS was considerably more effective at shorter wavelengths ( $S_{275-295}$ ), while no significant change was observed at longer wavelengths ( $S_{350-400}$ ), as expected from the fact that the absorption losses are higher in shorter wavelengths than the longer wavelengths (Del Vecchio and Blough, 2002). In order to look at the changes in light absorption property, the spectral slope ratios ( $S_R$ ) of shorter wavelength ( $S_{275-295}$ ) relative to longer wavelength ( $S_{350-400}$ ) were compared. In general, the  $S_R$  value increases with increasing irradiation (Del Vecchio and Blough, 2002; Helms et al., 2008). The  $S_R$  values of non-irradiated samples were 1.1 in the winter and 1.4 in the summer, and the irradiated samples were 1.4 in both winter and summer (Fig. 5). The  $S_R$  values of non-irradiated and irradiated samples in summer showed no measureable change during the UV experiment (Fig. 5). In addition, if the input of fresh HULIS is constant throughout the seasons,  $S_R$  should be constant.

490

495 However,  $S_R$  values of non-irradiated samples were 1.1 in the winter and 1.4 in the summer (Fig. 5). The  $S_R$  value of the non-irradiated summer sample was similar to that of the irradiated winter sample (1.4) (Fig. 5). Thus, this suggests that the lower summer HULIS concentration was associated to the UV degradation rather than to the reduced source inputs.

## 5 Conclusions

500 The seasonal variations in optical and chemical properties of the HULIS and WSOC were monitored in different seasons in the urban region. The air mass back trajectories combined with the fire maps and various chemical analyses demonstrated that biomass burning is a significant source contributing to HULIS in all seasons in Seoul, Korea. Our results suggest that photo-induced degradation plays a significant role in BrC quantity and quality in the atmosphere. In addition, the important role of photochemical degradation as a removal mechanism was supported by the laboratory experiments. This may provide

505 important pieces regarding the hidden life cycle of light-absorbing organic aerosols in the atmosphere.

The light-absorbing organic aerosols are considerably associated with climate sensitivity owing to their significant roles in the radiative forcing and global climate balance. Thus, this photochemical process needs to be considered in the modeling of climate forcing and biogeochemical cycles of the Earth's surface. Furthermore, the deposition fluxes of organic aerosols with different optical and chemical characteristics can have significant impacts as potential sources of organic carbon on surface waters and soils, which can lead to major effects on the global carbon cycle.

510

*Author contribution.* H.Han and G.Kim involved in planning the research and designing the experiment. H.Han collected the data and performed the analyses and experiment. H.Han and G.Kim involved in analyzing the results and writing the manuscript. H.Seo involve in sample analysis of trace element. K.-H.Shin and D.-H.Lee contributed to the sample analysis of molecular compounds. All authors contributed to the final version of the manuscript.

515

Author  
Deleted: trend of

Author  
Deleted: absorption

Author  
Deleted: consistent

Author  
Deleted: ies during the UV radiation experiments

Author  
Deleted: measurable

Author  
Formatted: Font:Italic

Author  
Deleted: e  $S_R$  value

Author  
Deleted: wa

Author  
Deleted: major

Author  
Deleted: confirmed

Author  
Deleted: e

Author  
Deleted: and chemical

Author  
Deleted: and soils

530 *Competing interests.* The authors declare no competing financial interests.

*Acknowledgements.* This work was supported by the National Research Foundation (NRF) of Korea (NRF-2018 R1A2B3001147) funded by the Korean government. We are grateful to Jeonghyun Kim for his support and assistance. [We acknowledge the use of data and imagery from LANCE FIRMS operated by NASA's EARTH Science Data and Information System \(ESDIS\) with funding provided by NASA Headquarters.](#) We also gratefully acknowledge the NOAA Air Resources Laboratory for use of the HYSPLIT model.

## References

- Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131–3148, <https://doi.org/10.5194/acp-6-3131-2006>, 2006.
- 540 Baduel, C., Voisin, D., and Jaffrezo, J. L.: Comparison of analytical methods for Humic Like Substances (HULIS) measurements in atmospheric particles, *Atmos. Chem. Phys.*, 9, 5949–5962, <https://doi.org/10.5194/acp-9-5949-2009>, 2009.
- Baduel, C., Voisin, D., and Jaffrezo, J. L.: Seasonal variations of concentrations and optical properties of water soluble HULIS collected in urban environment, *Atmos. Chem. Phys.*, 10, 4085–4095, [https://doi.org/10.5194/acp-10-4085-](https://doi.org/10.5194/acp-10-4085-2010)
- 545 2010, 2010.
- Birdwell, J. E. and Engel, A. S.: Characterization of dissolved organic matter in cave and spring waters using UV-Vis absorbance and fluorescence spectroscopy, *Org. Geochem.*, 41, 270–280, <https://doi.org/10.1016/j.orggeochem.2009.11.002>, 2010.
- Bro, R.: PARAFAC. Tutorial and applications, *Chemometr. Intell. Lab.*, 38, 149–171, [https://doi.org/10.1016/S0169-](https://doi.org/10.1016/S0169-7439(97)00032-4)
- 550 7439(97)00032-4, 1997.
- Chen, Q., Ikemori, F., and Mochida, M.: Light absorption and excitation-emission fluorescence of urban organic aerosol components and their relationship to chemical structure, *Environ. Sci. Technol.*, 50, 10859–10868, <https://doi.org/10.1021/acs.est.6b02541>, 2016.
- Coble, P. G.: Marine optical biogeochemistry: the chemistry of ocean color, *Chem. Rev.*, 107, 402–418, <https://doi.org/10.1021/cr050350+>, 2007.
- 555 [Das, O., Wang, Y., and Hsieh, Y.-P.: Chemical and carbon isotopic characteristics of ash and smoke derived from burning of C<sub>3</sub> and C<sub>4</sub> grasses, \*Org. Geochem.\*, 41, 263–269, <https://doi.org/10.1016/j.orggeochem.2009.11.001>, 2010.](#)
- Dasari, S., Andersson, A., Bikkina, S., Holmstrand, H., Budhavant, K., Satheesh, S., Asmi, E., Kesti, J., Backman, J., Salam, A., Bisht, D. S., Tiwari, S., Hameed, Z., and Gustafsson, Ö.: Photochemical degradation affects the light absorption of

- 560 water-soluble brown carbon in the South Asian outflow, *Sci. Adv.*, 5, eaau8066,  
https://doi.org/10.1126/sciadv.aau8066, 2019.
- Del Vecchio, R. and Blough, N. V.: Photobleaching of chromophoric dissolved organic matter in natural waters: Kinetics  
and modelling, *Mar. Chem.*, 78, 231–253, https://doi.org/10.1016/S0304-4203(02)00036-1, 2002.
- 565 [Després, V. R., Alex Huffman, J., Burrows, S. M., Hoose, C., Safatov, A. S., Buryak, G., Fröhlich-Nowoisky, J., Elbert, W.,  
Andreae, M. O., Pöschl, U., and Jaenicke, R.: Primary biological aerosol particles in the atmosphere: a review, \*Tellus\*  
\*B\*, 64, 15598, https://doi.org/10.3402/tellusb.v64i0.15598, 2012.](#)
- Duarte, R. M. B. O. and Duarte, A. C.: Atmospheric organic matter, *eMagRes*, 2, 415–426,  
https://doi.org/10.1002/9780470034590.emrstm1331, 2013.
- Feng, Y., Ramanathan, V., and Kotamarthi, V. R.: Brown carbon: A significant atmospheric absorber of solar radiation,  
570 *Atmos. Chem. Phys.*, 13, 8607–8621, https://doi.org/10.5194/acp-13-8607-2013, 2013.
- [Forrister, H., Liu, J., Scheuer, E., Dibb, J., Ziemba, L., Thornhill, K. L., Anderson, B., Diskin, B., Perring, A. E., Schwarz, J.  
P., Campuzano-Jost, P., Day, D. A., Palm, B. B., Jimenez, J. L., Nenes, A., and Weber, R. J.: Evolution of brown  
carbon in wildfire plumes, \*Geophys. Res. Lett.\*, 42, 4623–4630, https://doi.org/10.1002/2015GL063897, 2015.](#)
- 575 Fu, P., Kawamura, K., Chen, J., Qin, M., Ren, L., Sun, Y., Wang, Z., Barrie, L. A., Tachibana, E., Ding, A., and Yamashita,  
Y.: Fluorescent water-soluble organic aerosols in the High Arctic atmosphere, *Sci. Rep.*, 5, 9845,  
https://doi.org/10.1038/srep09845, 2015.
- Gabriel, R., Mayol-Bracero, O. L., and Andreae, M. O.: Chemical characterization of submicron aerosol particles collected  
over the Indian Ocean, *J. Geophys. Res.*, 107, 8005, https://doi.org/10.1029/2000JD000034, 2002.
- Ghan, S. J. and Schwartz, S. E.: Aerosol properties and processes: A path from field and laboratory measurements to global  
580 climate models, *Amer. Meteor. Soc.*, 88, 1059–1083, 2007.
- Graber, E. R. and Rudich, Y.: Atmospheric HULIS: How humic-like are they? A comprehensive and critical review, *Atmos.  
Chem. Phys.*, 6, 729–735, https://doi.org/10.5194/acp-6-729-2006, 2006.
- Gustafsson, Ö., Kruså, M., Zencak, Z., Sheesley, R. J., Granat, L., Engström, E., Praveen, P. S., Rao, P. S. P., Leck, C., and  
585 Rodhe, H.: Brown clouds over South Asia: biomass or fossil fuel combustion?, *Science*, 323, 495–498,  
https://doi.org/10.1126/science.1164857, 2009.
- [Healy, R. M., Wang, J. M., Sofowote, U., Su, Y., Debosz, J., Noble, M., Munoz, A., Jeong, C.-H., Hilker, N., Evans, G. J.,  
and Doerkson, G.: Black carbon in the Lower Fraser Valley, British Columbia: Impact of 2017 wildfires on local air  
quality and aerosol optical properties, \*Atmos. Environ.\*, 217, 116976, https://doi.org/10.1016/j.atmosenv.2019.116976,  
2019.](#)
- 590 Helms, J. R., Stubbins, A., Ritchie, J. D., and Minor, E. C.: Absorption spectral slopes and slope ratios as indicators of  
molecular weight, source, and photobleaching of chromophoric dissolved organic matter, *Limnol. Oceanogr.*, 33, 955–  
969, https://doi.org/10.4319/lo.2008.53.3.0955, 2008.

Author

Formatted: Font color: Text 1

Author

**Deleted:** Helms, J. R., Stubbins, A., Perdue, E. M., Green, N. W., Chen, H., and Mopper, K.: Photochemical bleaching of oceanic dissolved organic matter and its effect on absorption spectral slope and fluorescence, *Mar. Chem.*, 155, 81–91, 2013. -

- 600 Hennigan, C. J., Sullivan, A. P., Collett, J. L., and Robinson, A. L.: Levoglucosan stability in biomass burning particles exposed to hydroxyl radicals, *Geophys. Res. Lett.*, 37, L09806, <https://doi.org/10.1029/2010GL043088>, 2010.
- Hoffer, A., Gelencsér, A., Guyon, P., Kiss, G., Schmid, O., Frank, G. P., Artaxo, P., and Andreae, M. O.: Optical properties of humic-like substances (HULIS) in biomass-burning aerosols, *Atmos. Chem. Phys.*, 6, 3563–3570, <https://doi.org/10.5194/acp-6-3563-2006>, 2006.
- Hoffmann, D., Tilgner, A., Iinuma, Y., and Herrmann, H.: Atmospheric stability of levoglucosan: a detailed laboratory and modeling study, *Environ. Sci. Technol.*, 44, 694–699, <https://doi.org/10.1021/es902476f>, 2010.
- 605 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, <https://doi.org/10.5194/acp-5-1053-2005>, 2005.
- 610 Kawashima, H. and Haneishi, Y.: Effects of combustion emissions from the Eurasian continent in winter on seasonal  $\delta^{13}\text{C}$  of elemental carbon in aerosols in Japan, *Atmos. Environ.*, 46, 568–579, <https://doi.org/10.1016/j.atmosenv.2011.05.015>, 2012.
- Kelly, S. D., Stein, C., and Jickelle, T. D.: Carbon and nitrogen isotopic analysis of atmospheric organic matter, *Atmos. Environ.*, 39, 6007–6011, <https://doi.org/10.1016/j.atmosenv.2005.05.030>, 2005.
- 615 Kieber, R. J., Whitehead, R. F., Reid, S. N., Willey, J. D., and Seaton, P. J.: Chromophoric dissolved organic matter (CDOM) in rainwater, Southeastern North Carolina, USA, *J. Atmos. Chem.*, 54, 21–41, 2006.
- Kim, T.-H., Kim, G., Lee, S.-A., and Dittmar, T.: Extraordinary slow degradation of dissolved organic carbon (DOC) in a cold marginal sea, *Sci. Rep.*, 5, 13808, <https://doi.org/10.1038/srep13808>, 2015.
- Kirillova, E. N., Andersson, A., Han, J., Lee, M., and Gustafsson, Ö.: Sources and light absorption of water-soluble organic carbon aerosols in the outflow from northern China, *Atmos. Chem. Phys.*, 14, 1413–1422, <https://doi.org/10.5194/acp-14-1413-2014>, 2014a.
- Kirillova, E. N., Andersson, A., Tiwari, S., Sribastava, A. K., Bisht, D. S., and Gustafsson, Ö.: Water-soluble organic carbon aerosols during a fall New Delhi winter: Isotope-based source apportionment and optical properties, *J. Geophys. Res. Atmos.*, 119, 3476–3485, <https://doi.org/10.1002/2013JD020041>, 2014b.
- 625 Kuang, B. Y., Lin, P., Huang, X. H. H., and Yu, J. Z.: Sources of humic-like substances in the Pearl River Delta, China: positive matrix factorization analysis of  $\text{PM}_{2.5}$  major components and source markers, *Atmos. Chem. Phys.*, 15, 1995–2008, <https://doi.org/10.5194/acp-15-1995-2015>, 2015.
- Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, *Chem. Rev.*, 115, 4335–4382, <https://doi.org/10.1021/cr5006167>, 2015.
- 630 Lawaetz, A. J. and Stedmon, C. A.: Fluorescence intensity calibration using the raman scatter peak of water, *Appl. Spectrosc.*, 63, 936–940, <https://doi.org/10.1366/000370209788964548>, 2009.

**Author**

**Deleted:** Jiang, H., Frie, A. L., Lavi, A., Chen, J. Y., Zhang, H., Bahreini, R., and Lin, Y.-H.: Brown carbon formation from nighttime chemistry of unsaturated heterocyclic volatile organic compounds, *Environ. Sci. Technol. Lett.*, 6, 184–190, 2019. -

**Author**

**Deleted:** Kivácsy, Z., Kiss, G., Ceburnis, D., Jennings, G., Maenhaut, W., Salma, I., and Shooter, D.: Study of water-soluble atmospheric humic matter in urban and marine environments, *Atmos. Rea.*, 87, 1–12, 2008. -



- 645 Lin, C. Y., Sheng, Y. F., Chen, W. N., Wang, Z., Kuo, C. H., Chen, W. C., and Yang, T.: The impact of channel effect on Asian dust transport dynamics: a case in southeastern Asia, *Atmos. Chem. Phys.*, 12, 271–285, <https://doi.org/10.5194/acp-12-271-2012>, 2012.
- Lin, P., Huang, X.-F., He, L.-Y., and Yu, J. Z.: Abundance and size distribution of HULIS in ambient aerosols at a rural site in South China, *J. Aerosol Sci.*, 41, 74–87, <https://doi.org/10.1016/j.jaerosci.2009.09.001>, 2010.
- 650 Lukács, H., Gelencsér, A., Hammer, S., Puxbaum, H., Pio, C., Legrand, M., Kasper-Giebl, A., Handler, M., Limbeck, A., Simpson, D., and Preunkert, S.: Seasonal trends and possible sources of brown carbon based on 2-year aerosol measurements at six sites in Europe, *J. Geophys. Res.*, 122, D23S18, <https://doi.org/10.1029/2006JD008151>, 2007.
- Maenhaut, W., Raes, N., Chi, X., Cafmeyer, J., and Wang, W.: Chemical composition and mass closure for PM<sub>2.5</sub> and PM<sub>10</sub> aerosols at K-pusztá, Hungary, in summer 2006, *X-RAY Spectrom.*, 37, 193–197, <https://doi.org/10.1002/xrs.1062>, 2007.
- 655 Matos, J. T. V., Freire, S. M. S. C., Duarte, R. M. B. O., and Duarte, A. C.: Natural organic matter in urban aerosols: comparison between water and alkaline soluble components using excitation-emission matrix fluorescence spectroscopy and multiway data analysis, *Atmos. Environ.*, 102, 1–10, <https://doi.org/10.1016/j.atmosenv.2014.11.042>, 2015.
- 660 Mladenov, N., Lopez-Ramos, J., McKnight, D. M., and Reche, I.: Alpine lake optical properties as sentinels of dust deposition and global change, *Limnol. Oceanogr.*, 54, 6–2, 2386–2400, [https://doi.org/10.4319/lo.2009.54.6\\_part\\_2.2386](https://doi.org/10.4319/lo.2009.54.6_part_2.2386), 2009.
- Mladenov, N., Sommaruga, R., Morales-Baquero, R., Laurion, I., Camarero, L., Diéguez, M. C., Camacho, A., Delgado, A., Torres, O., Chen, Z., Felip, M., and Reche, I.: Dust inputs and bacteria influence dissolved organic matter in clear alpine lakes, *Nat. Commun.*, 2, 405, <https://doi.org/10.1038/ncomms1411>, 2011.
- 665 Panetta, R. J., Ibrahim, M., and Gélinas, Y.: Coupling a high-temperature catalytic oxidation total organic carbon analyzer to an isotope ratio mass spectrometer to measure natural-abundance  $\delta^{13}\text{C}$ -dissolved organic carbon in marine and freshwater samples, *Anal. Chem.*, 80, 5232–5239, <https://doi.org/10.1021/ac702641z>, 2008.
- Park, S. S., Kim, Y. J., Cho, S. Y., and Kim, S. J.: Characterization of PM<sub>2.5</sub> aerosols dominated by local pollution and Asian dust observed at an urban site in Korea during aerosol characterization experiments (ACE)-Asia project, *J. Air Waste Manage.*, 57, 2007.
- 670 Park, S. S. and Son, S.-C.: Relationship between carbonaceous components and aerosol light absorption during winter at an urban site of Gwangju, Korea, *Atmos. Res.*, 185, 73–83, <https://doi.org/10.1016/j.atmosres.2016.11.005>, 2017.
- Pöhlker, C., Huffman, J. A., and Pöschl, U.: Autofluorescence of atmospheric bioaerosols – fluorescent biomolecules and potential interferences, *Atmos. Meas. Tech.*, 5, 37–71, <https://doi.org/10.5194/amt-5-37-2012>, 2012.
- 675 Ramanathan, V., Ramana, M. V., Roberts, G., Kim, D., Corrigan, C., Chung, C., and Winker, D.: Warming trends in Asia amplified by brown cloud solar absorption, *Nature*, 448, 575–578, <https://doi.org/10.1038/nature06019>, 2007.

- Saleh, R., Robinson, E. S., Tkacik, D. S., Ahern, A. T., Liu, S., Aiker, A. C., Sullivan, R. C., Presto, A. A., Dubey, M. K., Yokelson, R. J., Donahue, N. M., and Robinson, A. L.: Brownness of organics in aerosols from biomass burning linked to their black carbon content, *Nat. Geosci.*, 7, 36940, <https://doi.org/10.1038/ngeo2220>, 2014.
- 680 Seinfeld, J. H., Carmichael, G. R., Arimoto, R., Conant, W. C., Brechtel, F. J., Bates, T. S., Cahill, T. A., Clarke, A. D., Doherty, S. J., Flatau, P. J., Huebert, B. J., Kim, J., Markowicz, K. M., Quinn, P. K., Russell, L. M., Russell, P. B., Shimizu, A., Shinzuka, Y., Song, C. H., Tang, Y., Uno, I., Vogelmann, A. M., Weber, R. J., Woo, J.-H., and Zhang, X. Y.: ACE-ASIA: Regional climatic and atmospheric chemical effects of Asian dust and pollution, *B. Am. Meteorol. Soc.*, 85, 367–380, 2004.
- 685 Stedmon, C. A. and Bro, R.: Characterizing dissolved organic matter fluorescence with parallel factor analysis: a tutorial, *Limnol. Oceanogr. Meth.*, 6, 11, 572–579, <https://doi.org/10.4319/lom.2008.6.572>, 2008.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Amer. Meteor. Soc.*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- 690 Taylor, S. R. and McLennan, S. M.: The geochemical evolution of the continental crust, *Rev. Geophys.*, 33, 241–265, <https://doi.org/10.1007/s10201-006-0194-9>, 1995.
- Troyer, I. D., Bouillon, S., Barker, S., Perry, C., Coorevits, K., and Merckx, R.: Stable isotope analysis of dissolved organic carbon in soil solutions using a catalytic combustion total organic carbon analyzer-isotope ratio mass spectrometer with a cryofocusing interface, *Rapid Commun. Mass Spectrom.*, 24, 365–374, <https://doi.org/10.1002/rcm.4403>, 2010.
- 695 [Tsukuda, S., Sugiyama, M., Harita, Y., and Nishimura, K.: Atmospheric bulk deposition of soluble phosphorus in Ashiu Experimental Forest, Central Japan: source apportionment and sample contamination problem, \*Atmos. Environ.\*, 39, 823–836, <https://doi.org/10.1016/j.atmosenv.2004.10.028>, 2005.](https://doi.org/10.1016/j.atmosenv.2004.10.028)
- Twardowski, M. S., Boss, E., Sullivan, J. M., and Donaghay, P. L.: Modeling the spectral shape of absorbing chromophoric dissolved organic matter, *Mar. Chem.*, 89, 69–88, <https://doi.org/10.1016/j.marchem.2004.02.008>, 2004.
- 700 Wozniak, A. S., Bauer, J. E., and Dickhut, R. B.: Characteristics of water-soluble organic carbon associated with aerosol particles in the eastern United States, *Atmos. Environ.*, 46, 181–188, <https://doi.org/10.1016/j.atmosenv.2011.10.001>, 2012.
- Yan, G., Cho, H.-M., Lee, I., and Kim, G.: Significant emissions of <sup>210</sup>Po by coal burning into the urban atmosphere of Seoul, Korea, *Atmos. Environ.*, 54, 80–85, <https://doi.org/10.1016/j.atmosenv.2012.02.090>, 2012.
- 705 Yan, G. and Kim, G.: Dissolved organic carbon in the precipitation of Seoul, Korea: Implications for global wet depositional flux of fossil-fuel derived organic carbon, *Atmos. Environ.*, 59, 117–124, <https://doi.org/10.1016/j.atmosenv.2012.05.044>, 2012.
- Yan, G. and Kim, G.: Sources and fluxes of organic nitrogen in precipitation over the southern East Sea/Sea of Japan, *Atmos. Chem. Phys.*, 15, 2761–2774, <https://doi.org/10.5194/acp-15-2761-2015>, 2015.

710 Yan, G. and Kim, G.: Speciation and sources of brown carbon in precipitation at Seoul, Korea: Insights from excitation-  
emission matrix spectroscopy and carbon isotopic analysis, *Environ. Sci. Technol.*,  
<https://doi.org/10.1021/acs.est.7b02892>, 2017.

715 Zepp, R. G., Sheldon, W. M., and Moran, M. A.: Dissolved organic fluorophores in southeastern US coastal waters:  
correction method for eliminating Rayleigh and Raman scattering peaks in excitation-emission matrices, *Mar. Chem.*,  
89, 15–36, <https://doi.org/10.1016/j.marchem.2004.02.006>, 2004.


720

725


730

735

740



750 **Figure 1:** Map of the geographical region around the study site and air mass transport pathways in the different seasons, spring (green), summer (red), fall (blue), and winter (brown). The ten-day air mass back trajectory was drawn by using the HYSPLIT model from March 2015 to January 2016 in Seoul.




760

Figure 2: Temporal variations in (a) WSOC concentration, (b) fluorescence intensity of HULIS, (c) ratio of HULIS to WSOC, (d)  $\delta^{13}\text{C}_{\text{WSOC}}$  values, (e) levoglucosan, (f)  $\text{Ca}^{2+}$ , (g)  $\text{SO}_4^{2-}$ , (h)  $\text{NO}_x$ , (i) non-crystal K, (j) non-crystal V, (k) sea-spray concentrations, (l) UV radiation rate, and temperature from March 2015 to January 2016 in Seoul, Korea.

765



795



800

805

Figure 4: Plots of the (a) HULIS versus UV-degradable HULIS and the (b) ratio of HULIS to WSOC versus UV-degradable HULIS in the different seasons, spring (green), summer (red), fall (blue), and winter (brown). The dashed lines represent the regression lines.

810

815

820

Author

Deleted: Correlations between

Author

Deleted: and

Author

Deleted: and

Author


Deleted: over



830

835

840



845 | Figure 5: The absorption coefficients of the aerosol samples collected in the winter (30.Dec) (blue) and summer (14.Aug) (red) in 2015. The solid lines represent the initial values for the non-irradiated samples, and the dashed lines represent the final values for the irradiated samples.

Author  
Deleted: -

850