



Sensitivity of Age of Air Trends on the derivation method for non-linear increasing tracers

Frauke Fritsch^{1,2}, Hella Garny^{1,2}, Andreas Engel³, Harald Bönisch⁴, and Roland Eichinger^{1,2}

¹Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

²Ludwig Maximilians University of Munich, Meteorological Institute Munich, Munich, Germany

³Institute for Atmospheric and Environmental Sciences, Goethe University Frankfurt am Main, Frankfurt am Main, Germany

⁴Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

Correspondence: Frauke Fritsch (frauke.fritsch@dlr.de)

Abstract. Mean age of air (AoA) is a diagnostic of transport along the stratospheric Brewer-Dobson circulation. While models consistently show negative trends, long-term time series (1975-2016) of AoA derived from observations show non-significant positive trends in mean AoA in the northern hemisphere. This discrepancy between observed and modeled mean AoA trends is still not resolved. There are uncertainties and assumptions required when deriving AoA from trace gas observations. At the same time, AoA from climate models is subject to uncertainties, too.

In this paper, we focus on the uncertainties due to the parameter selection in the method that is used to derive mean AoA from SF₆ measurements in Engel et al. (2009) and Engel et al. (2017). To correct for the non-linear increase in SF₆ concentrations, a quadratic fit to the time-series at the reference location, i.e. the tropical surface, is used. For this derivation, the width of the AoA distribution (age spectrum) has to be assumed. In addition, to choose the number of years the quadratic fit is performed for, the fraction of the age spectrum to be considered has to be assumed. Even though the uncertainty range due to all different aspects has already been taken into account for the total errors on the AoA values, the systematic influence of the parameter selection on AoA trends is described for the first time in the present study.

For this, we use the climate model EMAC (ECHAM MESSy Atmospheric Chemistry) as a test bed, where AoA derived from a linear tracer is available as a reference and modeled age spectra exist to diagnose the actual spatial age spectra widths. The comparison of mean AoA from the linear tracer with mean AoA from a SF₆ tracer shows systematic deviations specifically in the trends due to the selection of the parameters. However for an appropriate parameter selection, good agreement for both mean AoA and its trend can be found with deviations of about 1 % in mean AoA and 12 % in AoA trend.

In addition, a method to derive mean AoA is evaluated that applies a convolution to the reference time series. The resulting mean AoA and its trend only depend on an assumption about the ratio of moments. Also in that case, it is found that the larger the ratio of moments, the more the AoA trend gravitates towards the negative. The linear tracer and SF₆ AoA is found to agree within 0.3 % in the mean and 6 % in the trend.

The different methods and parameter selections were then applied to the balloon borne SF₆ and CO₂ observations. We found the same systematic changes in mean AoA trend dependent on the specific selection. When applying a parameter choice that is suggested by the model results, the AoA trend is reduced from 0.15 years/decade to 0.07 years/decade. It illustrates that



correctly constraining those parameters is crucial for correct mean AoA and trend estimates and still remains a challenge in the real atmosphere.

1 Introduction

The Brewer-Dobson circulation (BDC) is the slow, overturning equator-to-pole mass circulation in the stratosphere (Butchart, 2014). As such it influences how chemical species are distributed in the atmosphere. For example, in the case of ozone depleting substances it is crucial to understand their transport to high latitudes, where they most severely impact ozone. As the BDC cannot be measured directly, age of air (AoA) has been established as a common measure to quantify it (Hall and Plumb, 1994). Trace gas measurements can be used to derive AoA (Volk et al., 1997). Both the residual circulation transport and mixing contribute to AoA (Garny et al., 2014). From a conceptual point of view, AoA can be illustrated as a number of parcels that constitute an air mass and each of those parcels takes a different pathway from one reference point to an observation point. This idea was first introduced by Kida (1983). Those different pathways lead to a distribution of transport times, which is called the age spectrum of the specific air mass. The first moment of this age spectrum is the mean AoA.

Analyzing global chemistry-climate models very consistently shows a speeding up of the BDC in the cause of climate change (Oberländer et al., 2013; Shepherd and McLandress, 2011; Garcia and Randel, 2008). In contrast, both satellite and balloon borne observations show positive trends in AoA, which indicate a slowing of the BDC (Engel et al., 2017; Stiller et al., 2012; Haenel et al., 2015; Bönisch et al., 2011). Both observations and models have uncertainties in describing AoA.

Here, we want to focus on the uncertainties that arise from assumptions that are required to derive AoA from measurements. Those assumptions are necessary, as all tracers that are approximated as inert and measured in the atmosphere increase non-linearly. We will consider two methods here. First, the quadratic fit method was applied in Engel et al. (2009, 2017) which is highly relevant work regarding AoA trends. It considers the slope of the reference time series, which will be closer explained in Sec. 2.1.2. The different contributions of uncertainty were considered very thoroughly for the total uncertainty of mean AoA in Engel et al. (2009, 2017). Another option to treat the non-linearity of the reference time series is the convolution with a specific Green's function (i.e. age spectrum) to deduce mean age, as for example used in Ray et al. (2017). This also allows to employ more detailed approaches to the Green's function as shown in Andrews et al. (2001) and Bönisch et al. (2009). Leedham Elvidge et al. (2018) provides great insight to the total errors of mean AoA, applying both methods on measurements of various trace gases.

The latter also discusses the stratospheric lifetime of SF₆ and its effect on mean AoA. Both Leedham Elvidge et al. (2018) and Ray et al. (2017) find the lifetime of SF₆ to be lower than described so far. Up in the mesosphere SF₆ is depleted by photolysis and electron attachment. The depleted air moves into the polar vortex and into the whole stratosphere as the vortex breaks down, leading to AoA biases varying across the atmosphere. The effect on mean AoA in stratospheric mid-latitudes is estimated to be around 0.75 years.

In addition, Garcia et al. (2011) and Ray et al. (2014) worked on confining the uncertainty of mean AoA derived from SF₆ and CO₂ observations. Garcia et al. (2011) investigated AoA and its trends from a linear tracer and a SF₆ tracer with sinks in



WACCM (Whole Atmosphere Community Climate Model). They corrected the non-linearity of the SF₆ tracer (and the CO₂ tracer) with an exponential growth correction. They find negative trends for AoA from the linear and non-linear tracers, with the non-linear being smaller. Ray et al. (2014) narrowed down the error bar of the observed trends by including additional measurements and correcting for differences in measurement locations. Even with those corrections, the trend remains of similar magnitude as calculated by Engel et al. (2009), albeit with smaller error bars.

Still, the systematic uncertainty of the trend of mean AoA due to every single assumptions of the specific methods are not yet described. Investigating the method in a chemistry-climate model allows to use diagnostics that are not available for the real atmosphere and thereby to better understand the uncertainties due to the particular assumptions.

In Sec. 2 we will explain the methods under discussion, the trend calculation and provide details on the analyzed model simulations. We will present modeled age spectra and an analysis of the sensitivity of the trends on the parameter choices for a non-linear increasing tracer (SF₆) in Sec. 3. Sec. 4 discusses the results and relates them to the balloon borne observations. In Sec. 5 we will conclude our results.

2 Methodology

2.1 Age derivation

A well established way to derive mean AoA (Γ), in models is from a linearly increasing inert tracer. In this case, the mean age at a point \mathbf{x} is equal to the time lag τ of the tracer concentration χ relative to the concentration χ_0 at a selected reference area, which can be written as

$$\chi(\mathbf{x}, t) = \chi_0(t - \Gamma(\mathbf{x})) \quad (1)$$

2.1.1 Age spectrum

Additional information on the distribution of transport times is given by the actual age spectrum. The age spectrum is the distribution of transport times of trace gas concentrations to the location of interest. Thus, it is equated to the Green's function (G) that describes the propagation of the tracer concentration from the reference location to the observed location. This can be expressed with the following equation.

$$\chi(\mathbf{x}, t) = \int_0^{\infty} \chi_0(t - t') G(\mathbf{x}, t') dt' \quad (2)$$

Considering this equation, the Green's function can be obtained from a delta peak in tracer concentration. In the case of a non-stationary atmosphere, i.e. one with seasons, a set of inert tracer pulses at a reference surface is required. This method was already described and implemented for different models in e.g. Li et al. (2012) and Ploeger and Birner (2016).



2.1.2 Quadratic fit to the reference time series

In the real world, no perfectly linear increasing inert tracer is available. Often, SF₆ is used as a proxy for such a tracer. As the SF₆ emissions increase non-linearly, a common approach to derive mean AoA is a second order approximation that considers the slope of the reference time series. The general concept of the derivation is described in Volk et al. (1997). Engel et al. (2009) and Engel et al. (2017) apply it as an iterative procedure to optimize the length of the fit interval to the reference time series. First, the second order fit is performed on the reference time series for a set fit interval, e.g. 15 years. This yields the polynomial coefficients a , b and c . The fit is performed backwards from the reference time as follows:

$$\chi_0(t) = a + b(t - t_0) + c(t - t_0)^2 \quad (3)$$

Secondly, a first guess for the mean age is calculated using the coefficients applying the equation

$$\Gamma = \frac{b}{2c} - 2 \cdot C_1 \pm \sqrt{\left(\frac{b}{2c} - 2 \cdot C_1\right)^2 + \tau^2 - \frac{b}{2c} \cdot \tau} \quad (4)$$

Δ is the width of the age spectrum and C_1 is the ratio of first and second moments of the age spectrum, i.e. $C_1 = \Delta^2/\Gamma$. C_1 parameterizes the width of the spectrum relative to its mean. Again, τ is the lag time between the concentrations χ and χ_0 . To be consistent with Engel et al. (2017), the lag time is determined by approximating the reference time series by a second order function. In the third step, a fit interval t_{fit} is determined, which considers a given fraction of the age spectrum, e.g. 98%. This can be written as:

$$\int_0^{t_{\text{fit}}} G(t') dt' \stackrel{!}{=} \text{fraction of input} \quad (5)$$

The age spectrum assumed in this step is the inverse Gaussian that was proposed by Hall and Plumb (1994) based on a one dimensional diffusion analog equation.

$$G(t) = \sqrt{\frac{\Gamma^3}{4\pi\Delta^2 t^3}} \cdot \exp\left(-\frac{\Gamma(t - \Gamma)^2}{4\Delta^2 t}\right) \quad (6)$$

This equation uses the previous mean age value and the parameterization for the width. In the last step the value for the fit interval is used to determine once more the polynomial coefficients and recalculate the mean age value according to Eq. 4.

To perform this procedure one needs to choose the reference point, the fraction of the considered input and the ratio of the moments. Our aim is to systematically test the sensitivity of the derived mean AoA and AoA trends on those parameters.

2.1.3 Convolution Method

Another method to infer mean AoA from a non-linear increasing tracer is using a convolution. Considering Eq. 2, the reference time series χ_0 is convoluted with an assumed age spectrum G , i.e. Eq. 6 which gives an observed mixing ratio χ . Inserting



different mean age values Γ into Eq. 6, creates a look-up table of concentrations $\chi(\Gamma)$ that can be compared to the actual observed mixing ratio to pick the appropriate mean age value. This approach requires only an assumption about the ratio of moments. An assumption about the considered amount of reference input is not needed as the age spectrum weights the reference input according to the relative influence it has at the observed location. Therefore, the reference input can simply be considered for a long time interval, e.g. 25 years. Thus, this approach is expected to be quite physical and robust. Ray et al. (2017) and Leedham Elvidge et al. (2018) apply this approach to derive mean AoA from SF₆ and other non-linear increasing tracers, respectively.

2.2 EMAC Simulations

In the present study, we use the results of two simulations of the chemistry-climate model EMAC (ECHAM/MESSy Atmospheric Chemistry, Jöckel et al., 2010), which are both in a T42 horizontal (2.8 x 2.8 degrees) resolution with 90 layers in the vertical and explicitly resolved middle atmosphere dynamics (T42L90MA). In this setup, the uppermost model layer is centered at around 0.01 hPa and the vertical resolution in the upper troposphere lower stratosphere region (UTLS) is 500-600 m. MESSy (Modular Earth Submodel System) provides an infrastructure with generalized interfaces for standardized coupling of Earth System Model (ESM) components (dynamical cores, physical parameterizations, chemistry packages, diagnostics etc.). The dynamical core of EMAC is the European Center Hamburg general circulation model (ECHAM5, Roeckner et al., 2006).

The first simulation is a free running transient hindcast simulation using EMACv2.51 with full interactive chemistry stretching from 1960 to 2011. This simulation was performed within the ESCiMo project (Earth System Chemistry integrated Modeling, Jöckel et al., 2016) and is referred to as RC1-base-07. The setup of this simulation follows the REF-C1 scenario as defined by the Chemistry-Climate Model Initiative (CCMI, Eyring et al., 2013). Details on the setup and an evaluation of this chemistry-climate simulation can be found in Jöckel et al. (2016) and in Morgenstern et al. (2017).

The second simulation, termed tranPul, is an EMACv2.53.0 run which is designed to resemble the RC1-base-07 simulation as close as possible, however for cost efficiency reasons, without using interactive chemistry. This means that only the EMAC modules for dynamics, physics and diagnostics were used, namely AEROPT, CLOUD, CLOUDOPT, CONVECT, CVTRANS, E5VDIFF, ORBIT, OROGW, PTRAC, RAD, SURFACE, TNUDGE, TROPOP, VAXTRA. For details on these submodels refer to Jöckel et al. (2005, 2010, 2016). In that simulation, we transiently prescribed the monthly mean fields of the main greenhouse gases CO₂, CH₄, N₂O and O₃ from the RC1-base-07 simulation as zonal mean, monthly mean fields. As both simulations do not include a coupled ocean, SSTs and SICs have been prescribed following the global data set HadISST provided by the UK Met Office Hadley Centre (available via <http://www.metoffice.gov.uk/hadobs/hadisst/>, Rayner et al., 2003). Although in the L90 setup of EMAC, the quasi-biennial-oscillation (QBO) is internally generated (Giorgetta et al., 2002), the zonal winds near the equator were slightly nudged in the simulations to get the correct phasing of the observed QBO. Moreover, atmospheric aerosol (including volcanic aerosol) has been prescribed to take into account the interactions with radiation (in both simulations) and with heterogeneous chemistry (in the RC1-base-07 simulation).

The RC1-base-07 simulation is used to study the methods that are used to derive AoA from non-linear tracers, such as SF₆. The SF₆ time series prescribed in the simulation follows the RCP 6.0 scenario (Eyring et al., 2013). Its global mean is



shown in Fig. 1 as it is prescribed in the simulations to emulate emissions. The SF_6 time series as applied with the calculations from measurements is also displayed (see Levin et al., 2010; Maiss and Levin, 1994). From 1978 on, the time series is well constrained by NOAA measurements at Cape Grim. Before that, SF_6 is estimated from emissions. Prior to 1961, the data is completely interpolated. Both SF_6 time series differ slightly, as can be seen from Fig. 1. However, as explained in detail in Sec. 3.2, this does not impact our method evaluation.

In the tranPul simulation, periodical pulses of an inert tracer were released at the surface. This means that every three months, the lower boundary conditions of that tracer are set to 1 for one month at the tropical surface. After 9.5 years, the tracers are set to zero and started again to avoid large computational expenses. This operation is performed with the specifically designed MESSy submodule TPULSE. A similar EMAC setup was already applied in Hauck et al. (2019). The pulses allow to derive time evolving age spectra in EMAC as described for CLaMS (Chemical Lagrangian Model of the Stratosphere) in Ploeger and Birner (2016).




Figure 1. Globally averaged SF_6 time series on the ground as used in the RC1-base-07 simulation (blue) and as used with the observations calculations (red).



3 Results

3.1 Age spectra

It is fairly easy to derive mean AoA from linearly increasing tracers in models, as described in Sec. 2.1. To give an example of mean AoA in EMAC, Fig. 2 shows mean AoA derived from the idealized linear increasing tracer for July 2001-2010 from the RC1-base-07 simulation.




Figure 2. Mean AoA derived from the idealized tracer from the RC1-base-07 ESCiMo simulation. July mean for the years 2001-2010.

5

Since the derivation of mean AoA from measurements requires an assumption about the AoA spectrum, we derived spectra for EMAC in a transient simulation. Fig. 3 shows the seasonal evolution of the modeled spectra at 59° N and 106 hPa. This is close to 400 K and 60° N potential temperature, as in Figure 5 in the study by Ploeger and Birner (2016). The MAM, JJA and SON spectra show a similar propagation to Ploeger and Birner (2016). DJF rather shows one major peak instead of two as in Ploeger and Birner (2016). Our coarse spectra resolution of 3 months could be the reason for this, since the actual maxima of the age spectrum may fall in between a three month period such that they are not visible. For comparison, the spectrum calculated from the Hall and Plumb parameterization (see Eq. 6) is shown, using the values from the annual mean spectrum, that is a mean age value of 1.6 years (vertical line) and a ratio of moments of 1.8 years. For the chosen parameters, the modal age of the theoretical Hall and Plumb spectrum is lower than for the actual spectrum and the tail is broader. However, how well the modal age compares depends strongly on the applied ratio of moments.

10

15




Figure 3. Age spectra at 59° N and 106 hPa from the tranPul simulation. The dashed lines are the seasonal mean spectra. The pink solid line is the annual mean age spectrum. The solid yellow line is the spectrum calculated from the Hall and Plumb parameterization using the values from the annual mean, that is a mean age value of 1.6 years (vertical line) and a ratio of moments of 1.8 years

The ratio of moments calculated directly from the spectra as shown in Fig. 4 (left) are small (considering the length of the spectra of 9.5 years), considering values of 0.7 to 1.25 years are proposed for age calculations in Hall and Plumb (1994). Following Ploeger and Birner (2016), the spectra were extended to 50 years using an exponential fit on the second half of the spectrum. The ratio of moments in that case (Fig. 4 right) shift to larger values of up to 1.7 years in the annual mean covering the full simulation. This is consistent with e.g. Hauck et al. (2019), who also find ratio of moments larger than the upper limit of 1.25 years derived from the one dimensional diffusion analog. More details on the interannual variability of the ratio of moments will be discussed in Sec. 3.2. Also in that section, we will use the ratio of moments from the modeled spectra in the age calculations.

3.2 Sensitivity of mean AoA and AoA trends on parameter choice in the fit method

10 As described in Sec. 2.1, mean AoA and its trend were calculated from SF_6 and an ideal linear tracer for the RC1-base-07 simulation for a range of parameters and references for the fit method (explained in Sec. 2.1.2). Since we are considering a self-consistent model world and idealized SF_6 without sinks implemented, we can expect perfect agreement between the trends derived from the two tracers.




Figure 4. Mean ratio Δ^2/Γ derived from age spectra in the tranPul simulation. On the left for the regular 9.5 year long spectra, on the right with exponentially extended tail as in Ploeger and Birner (2016).

The fraction of input was varied between 90 %, 95 % and 98 % and the ratio of moments between 0.7, 0.8, 1.0 and 1.25 years. This covers the range for the ratio of moments proposed in Hall and Plumb (1994). Also, the pair of values 98 % and 0.7 years, as applied to the measurements in Engel et al. (2009) and Engel et al. (2017), is considered in this selection.

The resulting trends for 40° N and a selection of pressure levels are shown in Fig. 5. Other latitudes show similar behavior and are therefore not shown. The trend calculation was performed on deseasonalized values and considered autocorrelation (Thompson et al., 2015). The trend of the linear tracer is included in the plots for each parameter choice to make comparison easier, though it does not have any sensitivity to them. Overall, the trends derived from SF₆ are more negative the larger the ratio of moments and closer to zero the larger the fraction of input that is considered. Furthermore, the sensitivity of the trends to the ratio of moments increases when using a lower fraction of input. Intuitively, one might think that the results would be better the more of the reference input is considered (larger fraction of input). However, the fit performed on the reference time series is not weighted by, e.g. the spectrum. Therefore taking very long fit intervals takes into account more old air, which only has a small contribution to the actual mean AoA.

Fig. 5 also shows that depending on the pressure level, different selections of ratio of moment and reference input lead to agreement between the trends of the idealized and the SF₆ tracer. The influence of transport and mixing on the age distribution can explain that the best choice for the ratio of moments varies by location.

In the next step, we also consider a spatial distribution of ratio of moments. Even though we do not have age spectra available for the RC1-base-07 simulation, those from the tranPul simulation are a suitable first estimate for EMAC as a model. We considered the ratio of moments for the 9.5 year long spectra (PulRash) and the extended spectra (PulRa) (see Fig. 4). Fig. 6 shows the mean absolute deviation of mean AoA derived from SF₆ and the linear tracer as colors for different selections of parameters in the derivation. The white contours show the difference in trends in 10⁻¹ years/decade.




Figure 5. Overview of the trend in mean AoA derived from SF₆ (dotted lines) and a linear tracer (solid line) for different ratio of moments (x-axis) and fractions of input (colors) at 40° N. The panels show different pressure levels. The trend calculation considered the time series from 1975 to 2011. The error bars show 1σ which is the 68 % confidence levels.

It is clearly evident from Fig. 6 that using fixed ratio of moments for the whole latitude-height domain always leads to deviations in mean AoA and its trends in some region (Panels a-f). Using the ratio of moments derived from the 9.5 year spectra reduces the spatial pattern of deviations (panel g). However, an offset to the actual mean AoA values and trends remains as overall the ratios of moments are underestimated when neglecting the tail. In the range of the balloon borne observations (5-30 hPa and 32-51° N) this means an average deviation of 4 % in AoA values and 56 % in AoA trend. The ratio of moments derived from the extended age spectra leads to the best agreement across the widest spatial range for both mean AoA and its trend (Panel h). Both annual mean ratio of moments and transient ratio of moments, were used, though the transient ratio of moments do not show strong further improvement over the annual means (Panel i). Using PulRa and 95 % fraction of input provides 1 % average deviation in age values and 12 % deviation in trends in the region of the observations. The spatial dependency of an appropriate ratio of moments might play a role for the measurement application considering that the measurements in Engel et al. (2009) were analyzed between 5-30 hPa and 32-51° N.

Fig. 7 shows an example of the transient ratio of moments at 20 hPa and 40° N. Its trend is drawn, even though it is not significant as the trend is small compared to the variability. The trend of the ratio of moments is negative at all locations,




Figure 6. Mean absolute deviation of mean AoA calculated from SF₆ from AoA derived from an ideal tracer for different ratio of moments and fraction of input (colors). The white contours are the difference in trends in 10^{-1} years/decade. PulRa denotes the ratio of moments derived from the tranPul simulation annual mean extended age spectra, PulRa sh is from the 9.5 year spectra. PulRa tran uses the transient extended spectra from that simulation. White areas indicate where the derivation is not applicable. Black stippling show insignificance of the SF₆ trend from zero on the 95% confidence level. The trend calculation considered the time series from 1975 to 2011. The tropopause and the range 30 K above are excluded, as the AoA calculation is not defined in this region due to the different entry regions into the stratosphere that have to be considered.

but mostly not significant. The cycles visible in the ratio of moments can origin from the prescribed QBO and the solar cycle inherent in the prescribed ozone. It is beyond the scope of the current study to examine the large variability in the ratio of moments, but in particular the strong variation with periodicity of about 10 years calls for future investigations. Considering for example Labitzke (2007) it is anticipated that the solar cycle influences the BDC and thereby the shape of the age spectrum. The

5 AoA calculation typically considers about 10 years of reference input. That equates roughly to one cycle in strong variation in ratio of moments which are integrated. That might explain that most improvement in the AoA derivation comes from applying an appropriate spatial distribution of ratio of moments and not from using a transient one.

As shown in Fig. 6, Panel f, the combination of 98% and 0.7 years applied to derive AoA from observations leads to systematic offsets in both mean AoA and the AoA trends between the linear and SF₆ tracer in EMAC. In Fig. 8 the mean AoA

10 trends are shown, which further illustrates the offset in trends. The AoA trends derived from SF₆ with this parameter choice




Figure 7. Ratio of moments derived from extended age spectra from the tranPul simulation at 20 hPa and 40° N

are slightly negative or even positive and insignificant across a wide domain. Calculating the average AoA trend for 98 % and 0.7 years across the spatial range of the observations, i.e. 5-30 hPa and 32-51° N, yields

- $-0.13 \text{ years/dec} \pm 0.025 \text{ years/dec}$ for the linear tracer
- $-0.057 \text{ years/dec} \pm 0.033 \text{ years/dec}$ for the SF₆ tracer.

5 Hence, averaged across that range, the mean AoA trend derived using those parameters stretches to positive on the 2σ range (95 % confidence). Expressed relatively, these represent average deviations of 4 % in AoA values and 58 % in AoA trend.




Figure 8. Trends of mean AoA from 1975 to 2011 derived from a linear tracer (left panel) and from SF₆ for parameter selections as described in Fig. 6. White areas are the areas that can not be analyzed for a trend and black stippling shows insignificance of the trend from zero on the 95 % confidence level.

Since the reference location for the AoA calculation can also be varied in model calculations, we checked the effect on the resulting AoA and its trends. However, the different reference locations that have been tested do not show a strong influence on the resulting mean AoA or its trends. For example, an average of the concentration in the area between 20° S and 20° N



on the ground and the same range at 100 hPa were used. In that case, the differences in mean AoA are less than half a year and represents the offset time that the air traveled between the different reference locations (not shown). The differences in the trends are below 0.01 year/decade. Both statements refer to the results we obtained for different selections of fraction of input and ratio of moments, i.e. also considering selections that do not perform well overall. However, the reference location is important for quite specific tracer initialization on the ground, as for example if the linear AoA tracer is initialized from the poles. In such cases a reference location closer to the troposphere should be used. Since this is not the case for our simulations, we used the ground as reference in all calculations shown here as it is more applicable to measurements.

We find that the sensitivity of the AoA trends derived from SF₆ to the parameter choice arises predominantly from deviations in mean AoA in earlier years (1970s and 80s). As an example for this, the left of Fig. 9 shows the mean AoA time series and the derived trend at 20 hPa and 40° N for the combinations 98% and 0.7 years (red) and 95% and 1.25 years (blue). The linear tracer is given in black. 95% and 1.25 years represents a combination with relatively good agreement with the linear tracer in AoA whereas 98% and 0.7 years shows a strong deviation. For the mean AoA values at this location the length of the fit to the reference time series is around 11 years. The right panel of Fig. 9 shows the differences of the SF₆ derived mean AoA and the linear tracer derived mean AoA.




Figure 9. On the left mean AoA and its trend derived from SF₆ for two sets of parameters (red and blue) and from the linear tracer (black) at 20 hPa and 40° N from 1975 to 2011 from the RC1-base-07 Simulation. On the right accordingly mean AoA from the SF₆ tracer minus mean AoA from the linear tracer.

Considering Eq. 4, one explanation for the deviations might be that at the beginning of the reference time series the specific fit coefficients c and b of the second order polynomial are very sensitive to the fit interval length. Therefore Fig. 10 shows the coefficients dependent on the start date of the fit and the number of years fitted back. The parameters vary slightly stronger for fits early in the time series. As can be seen from Fig. 10 and already from the SF₆ time series in Fig. 1, the first part of the SF₆ time series is rather linear. In the eighties the second order contribution is strongest. Afterwards, the slope is rather linear again. Still, the times which show more variation in the coefficients with changing fit interval length do not clearly correspond to the times that the mean AoA derived from SF₆ deviates more from AoA derived from the linear tracer.




Figure 10. Coefficients b and c of the second order polynomial fit to the SF_6 reference time series dependent on the chosen length of fit interval.

Another explanation for the variation in mean AoA might be the sensitivity of the lag time τ to the fit interval length. To be consistent with Engel et al. (2009) and Engel et al. (2017), the lag time is determined by approximating the reference time series as a second order polynomial function of the concentration. Thus depending on the number of years considered, the obtained lag time might vary. This is counterintuitive to what one would expect if the lag time was directly looked up from the concentrations. Fig. 11 shows the lag times found for the SF_6 time series for fit interval lengths from 9 to 15 years at the same location as the example in Fig. 9. From roughly 1983 to 1990, τ depends stronger on the length of the fit interval and varies up to 12 % for the shown range of fit intervals.

As described when discussing the polynomial coefficients, the SF_6 reference time series slightly bends in the 1980s which makes the lag time calculation around the late 1980s more sensitive to the fit interval length. The specific years of stronger variation in lag time shift slightly for different locations, dependent on the transport times to that location. As seen from Fig. 9 (right), 1983 to 1990 is also the time of strongest deviation for the non-suitable parameter selection in Fig. 9. The variation in lag time contributes to the variation in mean age in the beginning of the time series dependent on the selection of parameters in the derivation.

The latter analyses were done for both SF_6 time series, the one prescribed in the simulation and the one used for calculations from observations. Only the results obtained from the SF_6 time series prescribed in the model are shown here as the results were similar.

3.3 AoA calculations via the Convolution Method

As described in Sec. 2.1.3, applying the convolution method weights the reference input according to the strength of its influence at a certain observation location. This avoids the influence of an assumption of the fraction of input, so one would expect to see more reliable agreement between AoA and its trends than when applying the fit method. Thus, the convolution method primarily requires an assumption about the ratio of moments of the assumed spectrum. The amount of years of reference




Figure 11. Lag time τ for fit interval length 9 to 15 years calculated for the SF₆ time series at 20 hPa and 40° N

can just be considered relatively high, e.g. 25 years. As a first example, Fig. 12 shows mean AoA and its trend calculated for ratio of moments 0.7 years and 1.25 years from the SF₆ tracer and from the linear tracer. One can see that for 1.25 years of ratio of moment, the agreement between AoA calculated from the linear tracer and from SF₆ becomes very good. However, for a ratio of moments of 0.7 years the disagreement prevails. Similar to the fit method, deviations in mean AoA are found in the beginning of the AoA time series maximizing in the 1980s, resulting in an AoA trend that is less negative.




Figure 12. Left, mean AoA and its trend derived from SF₆ using the convolution method for 0.7 and 1.25 years ratio of moments (red and blue) and from the linear tracer (black) at 20 hPa and 40° N from 1975 to 2011 from the RC1-base-07 simulation. Right, the difference in mean AoA derived from SF₆ and the linear tracer.



Similar to Fig. 6, Fig. 13 shows, height-latitude plots of the mean absolute deviation between mean AoA calculated using the linear increasing tracer and the SF₆ tracer in color and the difference in their trends in 10⁻¹ years/decade in white contours. The location of the balloon borne observations by Engel et al. (2017) is indicated by the red boxes. Here one can see that overall, a ratio of moments of 0.7 years, which was applied to the balloon borne observations so far, leads to too small negative or even positive trends, which are insignificant. Specifically, in the range of the observation this is a deviation in mean AoA of 2.8 % and in trend of 54 %. This remains similar to applying this ratio of moments in the fit method. On the other hand, using the ratio of 1.25 years yields very good agreement in the range of the balloon borne observations (0.3 % in mean AoA, 6 % in trend). Again, using the spatial distribution of ratio of moments derived from the extended model spectra, finds good agreement, too (0.4 % in mean AoA, 14 % in AoA trend in the range of the observations).




Figure 13. As Fig. 6, in color are the mean absolute deviation of mean AoA calculated from SF₆ from AoA derived from an ideal tracer for different ratio of moments using the convolution method. The white contours are the difference in trends in 10⁻¹ years/decade. PulRa denotes the ratio of moments derived from the tranPul simulation annual mean extended age spectra. White areas indicate where the derivation is not applicable. Black stippling shows insignificance of the SF₆ trend from zero on the 95% confidence level. The trend calculation considered the time series from 1975 to 2011. The red box shows the range of the balloon observations.

10 4 Discussion

4.1 Model Results

It has been shown here, that the variation of the mean AoA trend due to the selection of parameters is larger than the internal variability (e.g. Fig. 5). One might suggest that models still have spin-up effects as the deviations in mean AoA are often in the beginning of the time series. Both simulations, tranPul and RC1-base-07, were started in 1950, but the first ten years were considered as spin-up. So the data is used starting in 1960, such that the first mean AoA values are calculated for 1975, as the mean AoA calculation fits the reference time series backwards. Previous tests on the effects of spin-up time on AoA in EMAC support that our mean AoA values should not be considerably influenced by spin-up effects for this amount of spin-up. Hence,



the differences between models and the observations are not due to spin-up effects in models. We find that the major source of variation in SF₆-derived mean AoA especially using the fit method is the selection of parameters.

Another question we consider is whether a second order fit is appropriate to describe the SF₆ reference time series. Testing higher order fits shows, that the 3rd order coefficients would be two orders of magnitude smaller than the second order coefficients and the higher ones even smaller. Therefore, we remain with the second order assumption. Especially since higher order calculations would require even more assumptions about the transport time distribution which would be even more difficult to constrain in measurement applications.

The convolution method allows to avoid uncertainties about the amount of time considered and the function that is fitted to the time series. Still, there is a high sensitivity to the ratio of moments and it remains open which ratio of moments is describing the atmosphere best. Still, it is rather certain, that larger ratios of moments are more realistic.

4.2 Recalculation of AoA from balloon borne observations

It is most valuable to put the results derived from the model data concerning parameters and methods of AoA calculations into context by testing whether we see the same tendencies in calculations of observed AoA. The mean age values are recalculated from the observations by Engel et al. (2017) with the fit method with different parameters and the convolution method with different ratio of moments. The results for the trends are given in Fig. 14 and Table 1.




Figure 14. Trend in mean AoA derived from the observations for the fit and the convolution method for the respective fraction of input and ratio of moments.



Table 1. AoA trends recalculated from observations using fit method and convolution method for different parameters. Within the presented decimals, the errors are equal.

Method	Trend [years decade ⁻¹] calculated from ratio		
	0.7 years	1.25 years	2.0 years
Fit, 98 %	0.15 (±0.16)	0.15 (±0.16)	0.14 (±0.16)
Fit, 90 %	0.12 (±0.16)	0.07 (±0.16)	-0.007 (±0.16)
Convolution	0.11 (±0.16)	0.07 (±0.16)	0.008 (±0.16)

In agreement to what we presented for the model data, there is little influence of the ratio of moments on the trend using the fit method for 98% fraction of input, as was used in the calculations so far. For 90% fraction of input, the trend becomes less positive and eventually even slightly negative for larger ratio of moments. Thus, we see the same tendencies towards smaller, respectively more negative trends for larger ratio of moments and smaller fraction of input.

- 5 Applying the convolution method for larger ratio of moments gives smaller positive trends, too. This confirms what we have seen from the model data. For both methods for the larger ratio of moments negative trends as seen in the models lay within the margins of error.




Figure 15. Left, example AoA time series derived from the observations using the stated parameters. Right, the difference between maximum and minimum mean AoA found from the observations for the considered parameters (0.7, 0.8, 1.0 and 1.25 years for the convolution method and 90% 95% and 98% fit method).

Fig. 15 presents the AoA time series recalculated from the observations. On the left, the former parameter selection of 98% and 0.7 years for the fit method is displayed together with the recalculated version for 90% and 1.25 years. The right shows the spread in mean AoA for the different parameter and methods, that is 0.7, 0.8, 1.0 and 1.25 years for convolution method and 90% 95% and 98% for the fit method. The difference between the largest and smallest AoA value is shown. It becomes clear from the two plots, that the beginning of the SF₆ time series is more sensitive in the calculation, as seen for the model data.



Overall our results show that the selection of parameters helps resolve the difference between model and observation AoA trends. Furthermore, it should be mentioned that Volk et al. (1997) found good agreement with Waugh et al. (1997) using a ratio of moments of 1.25 years for airborne in-situ measurements of SF₆. Applying this ratio of moments to the balloon borne observations results in mean AoA trends that are the same as those of the model, within the margins of error.

5 However, one should consider that the trend was derived from both SF₆ and CO₂ measurements. For calculations using CO₂ measurements methane oxidation is considered as a minor source. In our discussion, we focused on the specific derivation methods themselves, such that we did not consider CO₂. Consequently, it was favorable to use a SF₆ tracer without mesospheric sinks. In particular, since the balloon borne observations we want to understand are located between 5-30 hPa and 32-51° N
10 and SF₆ balloon borne observations would make an influence of SF₆ sinks on the mean AoA trend less clear. However, the SF₆ sinks will be important to consider when discussing the specific differences between mean AoA derived from the global MIPAS satellite SF₆ observations and models, especially in the light of recent studies that report shorter lifetimes for SF₆ than assumed so far (Leedham Elvidge et al., 2018; Ray et al., 2017).

5 Conclusions

15 We investigated the sensitivity of the derivation of mean AoA to parameter choices that are necessary in the fit and convolution methods using a model as test-bed. References available in a model such as a linearly growing AoA tracer and age spectra allow specific testing of the methods.

In our analysis we were able to find a systematic variation of the mean AoA trend derived from SF₆ dependent on the ratio of moments and the fraction of input used in the AoA calculation when applying the fit method. The larger the ratio of moments
20 is, the more negative is the AoA trend. In addition, the larger the fraction of input is, the more towards positive the trend. While the sensitivity to the ratio of moments was included in the error range by Engel et al. (2009), the usage of a fixed fraction of input of 98 % reduced the sensitivity to the ratio of moments in their study.

Investigating another method to derive mean AoA, namely the convolution method, showed the same relation of smaller negative AoA trends for larger ratio of moments. The convolution method only assumes the ratio of moments. It provides very
25 good agreement between AoA from SF₆ and the linear tracer for the correct ratio of moment.

Applied to the balloon borne observations was the fit method with a small ratio of moments (0.7 years) and a big fraction of input (98 %), which both tend to shift the trend towards more positive values. Based on model results, we find better agreement with a linear AoA tracer for ratio of moments of about 1.25 and fraction of input of 95 %. Therefore, the parameter selection helps to resolve why the mean AoA trends found in Engel et al. (2009) and Engel et al. (2017) are positive, while models show
30 negative trends.

Our model results suggest using ratio of moments larger than 0.7 years. Using ratio of moments within the range 1.25-2 years is in line with what other 3d-model studies suggest (Waugh et al., 1997; Hauck et al., 2019). Specifically, using a spatial distribution is favorable as it considers aspects such as the broadening of the spectra due to mixing lower in the stratosphere.



Testing the impact of the parameter choice on AoA derived from the balloon borne observations, shows the same sensitivities for fit and convolution method as seen in the model. Within the margins of error the trend in AoA from observations and EMAC agree when assuming large ratio of moments (2 years). However, as the ratio of moments in the real atmosphere is currently unknown, we cannot conclude whether such a large ratio of moments is realistic. Overall, understanding the systematic influence of the parameter selection on the trend of mean AoA provides further insight in addition to the existing work on the uncertainties of mean AoA and its trends.

Data availability. The ESCiMo data used in this study can be obtained through the British Atmospheric Data Centre (BADC) archive (<ftp://ftp.ceda.ac.uk>). For instructions for access to both archives see <http://blogs.reading.ac.uk/ccmi/badc-data-access>. The tranPul data and AoA recalculation data from observations are available upon request.

10 *Author contributions.* The initial concepts to derive mean AoA were initialized by HB and AE. HG and FF initiated the method evaluation. The simulation execution was done by RE with development contributions by FF. HB further contributed discussion and insight to the method evaluations. AE did the recalculation of the observational data. FF performed the data analysis and wrote the main parts of the paper. All authors contributed in finalizing the paper.

Competing interests. The authors declare that they have no conflict of interest.

5 *Acknowledgements.* This study was funded by the Helmholtz Association under grant VH-NG-1014 (Helmholtz-Hochschul-Nachwuchsforschergruppe MACClim). We also acknowledge the project ESCiMo (Earth System Chemistry integrated Modelling), within which the EMAC simulations were conducted at the German Climate Computing Centre DKRZ through support from the Bundesministerium für Bildung und Forschung (BMBF).



References

- Andrews, A. E., Boering, K. A., Wofsy, S. C., Daube, B. C., Jones, D. B., Alex, S., Loewenstein, M., Podolske, J. R., and Strahan, S. E.: Empirical age spectra for the midlatitude lower stratosphere from in situ observations of CO₂: Quantitative evidence for a subtropical “barrier” to horizontal transport, *Journal of Geophysical Research: Atmospheres*, 106, 10 257–10 274, <https://doi.org/10.1029/2000JD900703>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2000JD900703>, 2001.
- 5 Bönisch, H., Engel, A., Curtius, J., Birner, T., and Hoor, P.: Quantifying transport into the lowermost stratosphere using simultaneous in-situ measurements of SF₆ and CO₂, *Atmospheric Chemistry and Physics*, 9, 5905–5919, <https://doi.org/10.5194/acp-9-5905-2009>, <https://www.atmos-chem-phys.net/9/5905/2009/>, 2009.
- Bönisch, H., Engel, A., Birner, T., Hoor, P., Tarasick, D. W., and Ray, E. A.: On the structural changes in the Brewer–Dobson circulation after 2000, *Atmospheric Chemistry and Physics*, 11, 3937–3948, <https://doi.org/10.5194/acp-11-3937-2011>, <https://www.atmos-chem-phys.net/11/3937/2011/>, 2011.
- 10 Butchart, N.: The Brewer–Dobson circulation, *Reviews of Geophysics*, 52, 157–184, <https://doi.org/10.1002/2013RG000448>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013RG000448>, 2014.
- Engel, A., Möbius, T., Bönisch, H., Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S., Nakazawa, T., Sugawara, S., et al.: Age of stratospheric air unchanged within uncertainties over the past 30 years, *Nature Geoscience*, 2, 28, 2009.
- 15 Engel, A., Bönisch, H., Ullrich, M., Sitals, R., Membrive, O., Danis, F., and Crevoisier, C.: Mean age of stratospheric air derived from AirCore observations, *Atmospheric Chemistry and Physics*, 17, 6825–6838, <https://doi.org/10.5194/acp-17-6825-2017>, <https://www.atmos-chem-phys.net/17/6825/2017/>, 2017.
- Eyring, V., Lamarque, J.-F., Hess, P., Arfeuille, F., Bowman, K., Chipperfield, M. P., Duncan, B., Fiore, A., Gettelman, A., Giorgetta, M., Granier, C., Hegglin, M., Kinnison, D., Kunze, M., Langematz, U., Luo, B., Martin, R., Matthes, K., Newman, P., Peter, T., Peter, T., Robock, A., Ryerson, T., Saiz-Lopez, A., Salawitch, R., Schultz, M., Shepherd, T. G., Shindell, D., Staehelin, J., Tegtmeier, S., Thomason, L., Tilmes, S., Vernier, J.-P., Waugh, D., and Young, P.: Overview of IGAC/SPARC Chemistry-Climate Model Initiative (CCMI) Community Simulations in Support of Upcoming Ozone and Climate Assessments, *eyring et al. (2013), Overview of IGAC/SPARC Chemistry-Climate Model Initiative (CCMI) Community Simulations in Support of Upcoming Ozone and Climate Assessments, SPARC Newsletter no. 40, WMO-WRCP, Geneva, Switzerland, 48-66., 2013.*
- 20 Eyring, V., Lamarque, J.-F., Hess, P., Arfeuille, F., Bowman, K., Chipperfield, M. P., Duncan, B., Fiore, A., Gettelman, A., Giorgetta, M., Granier, C., Hegglin, M., Kinnison, D., Kunze, M., Langematz, U., Luo, B., Martin, R., Matthes, K., Newman, P., Peter, T., Peter, T., Robock, A., Ryerson, T., Saiz-Lopez, A., Salawitch, R., Schultz, M., Shepherd, T. G., Shindell, D., Staehelin, J., Tegtmeier, S., Thomason, L., Tilmes, S., Vernier, J.-P., Waugh, D., and Young, P.: Overview of IGAC/SPARC Chemistry-Climate Model Initiative (CCMI) Community Simulations in Support of Upcoming Ozone and Climate Assessments, *SPARC Newsletter no. 40, WMO-WRCP, Geneva, Switzerland, 48-66., 2013.*
- 25 Garcia, R. R. and Randel, W. J.: Acceleration of the Brewer–Dobson Circulation due to Increases in Greenhouse Gases, *Journal of the Atmospheric Sciences*, 65, 2731–2739, <https://doi.org/10.1175/2008JAS2712.1>, <https://doi.org/10.1175/2008JAS2712.1>, 2008.
- Garcia, R. R., Randel, W. J., and Kinnison, D. E.: On the Determination of Age of Air Trends from Atmospheric Trace Species, *Journal of the Atmospheric Sciences*, 68, 139–154, <https://doi.org/10.1175/2010JAS3527.1>, <https://doi.org/10.1175/2010JAS3527.1>, 2011.
- Garny, H., Birner, T., Bönisch, H., and Bunzel, F.: The effects of mixing on age of air, *Journal of Geophysical Research: Atmospheres*, 119, 7015–7034, <https://doi.org/10.1002/2013JD021417>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JD021417>, 2014.
- 30 Giorgetta, M. A., Manzini, E., and Roeckner, E.: Forcing of the quasi-biennial oscillation from a broad spectrum of atmospheric waves, *Geophysical Research Letters*, 29, 86–1–86–4, <https://doi.org/10.1029/2002GL014756>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2002GL014756>, 2002.
- Haenel, F. J., Stiller, G. P., von Clarmann, T., Funke, B., Eckert, E., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., and Reddmann, T.: Reassessment of MIPAS age of air trends and variability, *Atmospheric Chemistry and Physics*, 15, 13 161–13 176, <https://doi.org/10.5194/acp-15-13161-2015>, <https://www.atmos-chem-phys.net/15/13161/2015/>, 2015.



- Hall, T. M. and Plumb, R. A.: Age as a diagnostic of stratospheric transport, *Journal of Geophysical Research*, 99, 1059, <https://doi.org/10.1029/93jd03192>, 1994.
- Hauck, M., Fritsch, F., Garny, H., and Engel, A.: Deriving stratospheric age of air spectra using an idealized set of chemically active trace gases, *Atmospheric Chemistry and Physics*, 19, 5269–5291, <https://doi.org/10.5194/acp-19-5269-2019>, <https://www.atmos-chem-phys.net/19/5269/2019/>, 2019.
- 5 Jöckel, P., Sander, R., Kerkweg, A., Tost, H., and Lelieveld, J.: Technical Note: The Modular Earth Submodel System (MESSy) - a new approach towards Earth System Modeling, *Atmospheric Chemistry and Physics*, 5, 433–444, <https://doi.org/10.5194/acp-5-433-2005>, <https://www.atmos-chem-phys.net/5/433/2005/>, 2005.
- Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geoscientific Model Development*, 3, 717–752, <https://doi.org/10.5194/gmd-3-717-2010>,
10 <https://www.geosci-model-dev.net/3/717/2010/>, 2010.
- Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C. A. M., Brinkop, S., Cai, D. S., Dyroff, C., Eckstein, J., Frank, F., Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, M., Meul, S., Neumaier, M., Nützel, M., Oberländer-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth System Chemistry integrated Modelling (ESCiMo) with the Modular Earth Submodel System (MESSy) version 2.51, *Geoscientific Model Development*, 9, 1153–1200,
15 <https://doi.org/10.5194/gmd-9-1153-2016>, <https://www.geosci-model-dev.net/9/1153/2016/>, 2016.
- Kida, H.: General Circulation of Air Parcels and Transport Characteristics Derived from a Hemispheric GCM, *Journal of the Meteorological Society of Japan. Ser. II*, 61, 510–523, https://doi.org/10.2151/jmsj1965.61.4_510, 1983.
- Labitzke, K.: Solar Variation and Stratospheric Response, *Space Science Reviews*, 125, 247–260, <https://doi.org/10.1007/s11214-006-9061-6>, 2007.
- 20 Leedham Elvidge, E., Bönisch, H., Brenninkmeijer, C. A. M., Engel, A., Fraser, P. J., Gallacher, E., Langenfelds, R., Mühle, J., Oram, D. E., Ray, E. A., Ridley, A. R., Röckmann, T., Sturges, W. T., Weiss, R. F., and Laube, J. C.: Evaluation of stratospheric age of air from CF₄, C₂F₆, C₃F₈, CHF₃, HFC-125, HFC-227ea and SF₆; implications for the calculations of halocarbon lifetimes, fractional release factors and ozone depletion potentials, *Atmospheric Chemistry and Physics*, 18, 3369–3385, <https://doi.org/10.5194/acp-18-3369-2018>, <https://www.atmos-chem-phys.net/18/3369/2018/>, 2018.
- 25 Levin, I., Naegler, T., Heinz, R., Osusko, D., Cuevas, E., Engel, A., Ilmberger, J., Langenfelds, R. L., Neining, B., Rohden, C. v., Steele, L. P., Weller, R., Worthly, D. E., and Zimov, S. A.: The global SF₆ source inferred from long-term high precision atmospheric measurements and its comparison with emission inventories, *Atmospheric Chemistry and Physics*, 10, 2655–2662, <https://doi.org/10.5194/acp-10-2655-2010>, <https://www.atmos-chem-phys.net/10/2655/2010/>, 2010.
- Li, F., Waugh, D. W., Douglass, A. R., Newman, P. A., Pawson, S., Stolarski, R. S., Strahan, S. E., and Nielsen, J. E.: Seasonal variations of stratospheric age spectra in the Goddard Earth Observing System Chemistry Climate Model (GEOSCCM), *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016877>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011JD016877>, 2012.
- 30 Maiss, M. and Levin, I.: Global increase of SF₆ observed in the atmosphere, *Geophysical Research Letters*, 21, 569–572, <https://doi.org/10.1029/94gl00179>, 1994.
- 35 Morgenstern, O., Heglin, M. I., Rozanov, E., O'Connor, F. M., Abraham, N. L., Akiyoshi, H., Archibald, A. T., Bekki, S., Butchart, N., Chipperfield, M. P., Deushi, M., Dhomse, S. S., Garcia, R. R., Hardiman, S. C., Horowitz, L. W., Jöckel, P., Josse, B., Kinnison, D., Lin, M., Mancini, E., Manyin, M. E., Marchand, M., Maréchal, V., Michou, M., Oman, L. D., Pitari, G., Plummer, D. A., Revell, L. E.,



- Saint-Martin, D., Schofield, R., Stenke, A., Stone, K., Sudo, K., Tanaka, T. Y., Tilmes, S., Yamashita, Y., Yoshida, K., and Zeng, G.: Review of the global models used within phase 1 of the Chemistry–Climate Model Initiative (CCMI), *Geoscientific Model Development*, 10, 639–671, <https://doi.org/10.5194/gmd-10-639-2017>, <https://www.geosci-model-dev.net/10/639/2017/>, 2017.
- Oberländer, S., Langematz, U., and Meul, S.: Unraveling impact factors for future changes in the Brewer–Dobson circulation, *Journal of Geophysical Research: Atmospheres*, 118, 10,296–10,312, <https://doi.org/10.1002/jgrd.50775>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/jgrd.50775>, 2013.
- Ploeger, F. and Birner, T.: Seasonal and inter-annual variability of lower stratospheric age of air spectra, *Atmospheric Chemistry and Physics*, 16, 10 195–10 213, <https://doi.org/10.5194/acp-16-10195-2016>, <http://www.atmos-chem-phys.net/16/10195/2016/>, 2016.
- Ray, E. A., Moore, F. L., Rosenlof, K. H., Davis, S. M., Sweeney, C., Tans, P., Wang, T., Elkins, J. W., Bönnisch, H., Engel, A., Sugawara, S., Nakazawa, T., and Aoki, S.: Improving stratospheric transport trend analysis based on SF₆ and CO₂ measurements, *Journal of Geophysical Research: Atmospheres*, 119, 14,110–14,128, <https://doi.org/10.1002/2014JD021802>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014JD021802>, 2014.
- Ray, E. A., Moore, F. L., Elkins, J. W., Rosenlof, K. H., Laube, J. C., Röckmann, T., Marsh, D. R., and Andrews, A. E.: Quantification of the SF₆ lifetime based on mesospheric loss measured in the stratospheric polar vortex, *Journal of Geophysical Research: Atmospheres*, 122, 4626–4638, <https://doi.org/10.1002/2016jd026198>, 2017.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002670>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2002JD002670>, 2003.
- Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornblüeh, L., Manzini, E., Schlese, U., and Schulzweida, U.: Sensitivity of Simulated Climate to Horizontal and Vertical Resolution in the ECHAM5 Atmosphere Model, *Journal of Climate*, 19, 3771–3791, <https://doi.org/10.1175/JCLI3824.1>, <https://doi.org/10.1175/JCLI3824.1>, 2006.
- Shepherd, T. G. and McLandress, C.: A Robust Mechanism for Strengthening of the Brewer–Dobson Circulation in Response to Climate Change: Critical-Layer Control of Subtropical Wave Breaking, *Journal of the Atmospheric Sciences*, 68, 784–797, <https://doi.org/10.1175/2010JAS3608.1>, <https://doi.org/10.1175/2010JAS3608.1>, 2011.
- Stiller, G. P., von Clarmann, T., Haenel, F., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., and López-Puertas, M.: Observed temporal evolution of global mean age of stratospheric air for the 2002 to 2010 period, *Atmospheric Chemistry and Physics*, 12, 3311–3331, <https://doi.org/10.5194/acp-12-3311-2012>, <https://www.atmos-chem-phys.net/12/3311/2012/>, 2012.
- Thompson, D. W. J., Barnes, E. A., Deser, C., Foust, W. E., and Phillips, A. S.: Quantifying the Role of Internal Climate Variability in Future Climate Trends, *Journal of Climate*, 28, 6443–6456, <https://doi.org/10.1175/JCLI-D-14-00830.1>, <https://doi.org/10.1175/JCLI-D-14-00830.1>, 2015.
- Volk, C. M., Elkins, J. W., Fahey, D. W., Dutton, G. S., Gilligan, J. M., Loewenstein, M., Podolske, J. R., Chan, K. R., and Gunson, M. R.: Evaluation of source gas lifetimes from stratospheric observations, *Journal of Geophysical Research: Atmospheres*, 102, 25 543–25 564, <https://doi.org/10.1029/97JD02215>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/97JD02215>, 1997.
- Waugh, D. W., Hall, T. M., Randel, W. J., Rasch, P. J., Boville, B. A., Boering, K. A., Wofsy, S. C., Daube, B. C., Elkins, J. W., Fahey, D. W., Dutton, G. S., Volk, C. M., and Vohralik, P. F.: Three-dimensional simulations of long-lived tracers using winds from MACCM2, *Journal of Geophysical Research: Atmospheres*, 102, 21 493–21 513, <https://doi.org/10.1029/97JD00793>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/97JD00793>, 1997.