

The manuscript entitled “Marine organic matter in the remote environment of the Cape Verde Islands – An introduction and overview to the MarParCloud campaign” by van Pinxteren et al. describes the conditions and early results for a campaign designed to study the chemical interactions between the ocean and the marine boundary layer, including the formation of boundary layer clouds. It represents an interdisciplinary study of the sort that are increasingly seen in the field as the tight connections between ocean ecology, marine chemistry, atmospheric chemistry, and cloud formation are accepted by the community. The manuscript is generally well-written, but could benefit from more specific language in some places, as described in the comments below. In general, I am excited to see the results that come from this study in greater detail in the future, as some of the results provide new perspectives. The manuscript is recommended for publication in Atmospheric Chemistry and Physics upon consideration of the comments provided.

We thank the Reviewer for the evaluation and the constructive comments. Replies to the specific Referee’s comments are provided below in red and new parts included in the manuscript are marked in *italic*:

Introduction and Motivation: Overall this section could be more specific about the gaps in knowledge that exist, rather than vaguely saying that broad swaths of information are ‘elusive’, for example. We do know quite a lot about ocean surface chemistry as a result of the prior work of many investigators, including many who are a part of this project. Some additional specific comments on this particular subject are listed below

We agree and adopted the suggestions from the reviewer to sharpen the introduction and be more precise as shown in the following.

Lines 96-101 – the word ‘changing’ is used often, but it is not clear what is inducing the change, and from what state a ‘change’ is referenced. Hence the sentence is rather vague. Consider using more active or specific verbs to be more precise about how marine OM will influence each of the important areas listed.

We agree that “changing” was used too often and replaced this word accordingly. In addition, we explained the effects in some more detail. However, we want to underline that these are general important characteristics of the OM in the aerosols but not specifically investigated within this work. Therefore, we did not want to draw too much attention on this by adding detailed studies but rather mention these effects in general and give appropriate citations.

It now reads: “In particular, the role of marine organic matter (OM) with its sources and contribution to marine aerosol particles, is still *elusive*. For example, where this particle fraction might lead to a variety of effects such as *impacting health through the generation of reactive oxygen species, OM composition increasing or decreasing the absorption of solar radiation and therefore radiative properties, and impacting marine ecosystems via atmospheric deposition* (e.g. Abbatt, et al. 2019; Brooks and Thornton 2018; Burrows, et al. 2013; Gantt and Meskhidze 2013; Pagnone, et al. 2019; Patel and Rastogi, 2020).”

Lines 101-103 – the impact of primary aerosol production at the ocean surface has been explored more deeply in recent years (within the last 10 yrs or so) and, while we don't know everything, I'd say that the ability for particles to act as CCN or INP is not 'elusive' – indicating that we don't know much of anything. I would suggest that we are missing certain important pieces of information about CCN activity and also about what makes marine particles good INP (or not).

We agree and specified this part.

It now reads: “Furthermore, knowledge on the properties of marine organic aerosol particles and their ability to act as cloud condensation nuclei (CCN) or ice nucleating particle (INP) is *not fully understood. The fraction of marine CCN made up of sea spray aerosol is still debated and suggested to comprise about 30% on a global scale (excluding the high southern latitudes) (Quinn, et al. 2017) and important pieces of information about marine CCN are still missing (e.g. Bertram, et al. 2018).* Ocean-derived INPs were suggested to play a dominating role in determining INP concentrations in near-surface-air over the remote areas such as the Southern Ocean, however their source strength in other oceanic regions *as well as knowledge about which physicochemical properties determine the INP efficiency* are still largely unknown (Burrows, et al. 2013; McCluskey, et al. 2018a; McCluskey, et al. 2018b).”

Line 122-123 – “or through a more direct transfer of OM from ocean compartments to the marine particles”. This is vague – other than bubble bursting aerosol generation, what mechanisms are being discussed here?

We agree that this expression was not straightforward and changed it to:

“The SML is involved in the generation of sea-spray (or primary) particles including their organic fraction by transfer of OM to rising bubbles before they burst out *to jet droplets and film droplets (de Leeuw, et al. 2011).*”

Line 133 – “occurs as particulate and chromophoric dissolved organic matter” Imposing such a view limits one's conception of a vast array of chemicals existing in the SML, as is the case. There are fractions that are non-chromophoric and also not defined as particulate.

We agree that the phrasing was not comprehensive and rephrased it to:

Within the SML, OM is a mixture of different compounds *including polysaccharides, amino acids, proteins, lipids, and chromophoric dissolved organic matter (CDOM) that are either dissolved or particulate.*

It is great to see the bullet list of questions in this section. I have a few small comments/questions about them, hopefully in the search for greater clarity.

Line 174 – Perhaps this is vague on purpose, but “To what extent” by which measure? By absolute mass? By absolute number of particles containing organic matter? What about the particle mixing state? As a fraction of total ambient aerosol mass? The way this question is asked has a profound effect on the environmental effects that it will speak to.

We thank for these suggestions and addressing also the suggestions of referee #2, we included the specific metrics that we investigated here, and it now reads:

- *To what extent is seawater a source of OM to aerosol particles (regarding number, mass, chemical composition, CCN and INP concentration) and in cloud water?*

Line 176 – What are “OM groups”? Can a more specific wording be used here? Are you referring particular chemical functionality (fatty acids, carbohydrates, etc), operationally defined classes of material (DOM, POM, etc), or some other definition?

In agreement with the suggestions of referee #2 we changed the question to:

- *What are the important chemically-defined OM groups (proteins, lipids, carbohydrates - as sum parameters and on molecular level) in oceanic surface films, aerosol particles and cloud water and how are they linked?*

Line 189 – Are the authors asking whether the presence of marine OM in the surface ocean drives the concentration of CCN in the MBL? As written, the question does not presume the ‘direction’ of the relationship...presumably the implied direction is (in general terms): ‘Does the ocean influence the atmosphere?’ Being specific about this directionality allows this question to commute better with the final question about emission parameterizations.

We made the question more specific and took the reviewers’ suggestion and re-phrased this it to

- *Does the presence of marine OM in the surface ocean drive the concentration of CCN in the MBL?*

Experimental Line 304 – Were cloud droplets actively removed from aerosol particle measurements? If so, which measurements? The reader may be able to deduce this from the size cuts (e.g., PM_{2.5}, lower stages of cascade impactors) but it’s less clear for some of the more indiscriminate samplers.

For the aerosol samplers with defined size cut, the size of sampled particles/droplet is defined via the size cut. The size-resolved aerosol particles, sampled with the Berner impactor were conditioned via a 3 m heated tube. The PM₁₀ sampler without defined size cut it is possible that during cloud events, cloud droplets were sampled to some extent. We included this information in the SI part “Aerosol particle sampling and chemical analysis of inorganic ions, OC/EC and WSOC” and it reads now:

“To avoid condensation of atmospheric water on the surface of the aluminium foils, a conditioning unit was mounted between the impactor inlet and the sampling unit consisting of a 3 m tube. By heating the sampled air, high relative humidity of the ambient air was reduced to 75-80% before the collection of the aerosol particles. The temperature difference between the ambient air at the impactor inlet and the sampled air after the conditioning unit was below 9 K.”

In addition, in the experimental part we added “It needs to be pointed out that the aerosol particle samplers run continuously and aerosol particles were also sampled during cloud events. *The cloud droplets were efficiently removed due to the pre-conditioning of the aerosol*

particles sampled with the Berner impactor (more information in the SI) and due to the size cut the PM₁ sampler. However, for aerosol particles sampled with the PM₁₀ sampler, small cloud droplets can be collected as well. In addition, the particles sampled with the low volume TSP sampler can be influenced by cloud droplets to some extent.”

Conditions Section 4.2: In addition to primary producers, the state and dynamics of other portions of the microbial ecosystem have been shown to be key aspects that control how marine OM is incorporated into aerosol. Were any other biological metrics (heterotrophic bacteria, zooplankton, etc) assessed? If not, for clarity, please make a statement so that other readers and interested parties will understand the extent of data availability. [***Note: the reviewer later saw the section on marine bacteria in the ‘Measurements’ section. Perhaps this means that section 4.2 should be presented as ‘Measurements’ as well. The comment above is clearly addressed by the authors in the manuscript, but I’ll leave it here as a signal that a reader may be similarly confused about the order of presentation.]

We thank the referee for pointing this out. We understood that the order of presentation may lead to some confusion and have re-structured this part (following also the suggestions of reviewer #2). In the revised version, we mention the chl-a concentrations in section 4.2. and shifted the discussion of the pigment results to section 5.4.1. To underline, that pigments as well as bacteria were analysed we:

i) renamed section 4.2.1 to “Pigment and bacteria concentration in seawater”

ii) included this information in section 4.2.1 and it now reads “*Chl-a concentrations varied between 0.11 µg L⁻¹ and 0.6 µg L⁻¹, and are more thoroughly discussed together with the pigment composition in section 5.4.1. Moreover, as other but phytoplankton organisms can contribute to the OM pool, bacterial abundance was analysed in the SML and bulk water samples and these data are reported in section 5.7.3.*”

iii) concluded section 5.4.1. with: “First analyses show that the DOC concentrations were not directly linked to the increasing chl-a concentrations, however their relation to single pigments, to the microbial abundance, to the background dust concentrations and finally to wind speed and solar radiation will be further resolved to elucidate potential biological and meteorological controls on the concentration and enrichment of DOC.”

Due to the inclusion of the pigment discussion in chapter 5.4.1. (that therefore became very long) we have added another subsection for the comparison of DOC data from the two sampling techniques: 5.4.2 DOC concentrations: manual glass plate vs. MarParCat sampling

Measurements and Selected Results Section 5.1.1: Does the small fraction of sea spray aerosol determined by size distribution measurements comport with the chemical measurements in Section 5.1.2? While this may be the subject of a future paper, a evidence that different aerosol measurements agree in a general sense is an important point to make in a paper of this type.

Thank you for this advice. We included more information on CCN in general in Chapter 5.1.1., and to address the comparison between physical and chemical aerosol measurements, we added information concerning the particle hygroscopicity in the following text:

“The hygroscopicity parameter kappa (κ) averaged 0.28, suggesting the presence of OM in the particles (see Gong et al., 2020a). Particle sizes for which κ was determined (i.e., the critical diameters determined during CCN analysis) were roughly 50 to 130 nm. The low value determined for κ is in line with the fact that sodium chloride from sea salt was below detection limit in the size segregated chemical analysis for particles in this size range (Figure 10), while insoluble EC and WSOM made up 30% of the main constituents at CVAO on average. A thorough statistical analysis of N_{CCN} and particle hygroscopicity concerning different aerosol types is reported in Gong, et al. (2020a).”

In chapter 5.1.2. we respectively added the following:

“...the chemical composition aligned well with the κ value from the hygroscopicity measurements (Gong, et al. 2020a).”

As well as:

“From the chemical composition no indications for anthropogenic influences was found as concentrations of elemental carbon and submicron potassium were low (see Tab. S5). However, according to the dust concentrations (Table 2) and the air mass origins (Fig. S1), as well as the PNSD (Gong et al. 2020a), the air masses during this period experienced low dust influences, that was however not visible from the main chemical constituents studied here and warrants more detailed chemical investigations (like size-resolved dust measurements), a distinction between mass-based and number-based analysis as well as detailed source investigations that are currently ongoing.”

We added in the conclusion:

“These findings underline that further in depth studies differentiating between submicron and supermicron particles as well as between aerosol number and aerosol mass are strongly required.”

Section 5.1.2: It is stated, somewhat simply, that the mass fraction of sodium and chloride in cloud water illustrates that supermicron particles were a dominant factor in cloud formation. However, mass-based chemical analysis, especially when not size resolved, is very easy to misinterpret. The key metric in cloud formation is the number concentration of droplets. The mass of supermicron particles increases as roughly the cube of the diameter, so particles that are 1 micron vs 100 nm contribute 1000x more mass to the cloud water! At the same time, particles larger than $d = 200$ nm activate at just about any relevant supersaturation relevant to this environment. The ‘control’ on the cloud drop number concentration, therefore, comes from the size range under 200 nm (down to, say 50 nm or so). Mass measurements will, therefore not tell us much about which particles actually control differences in cloud optical thickness, CDNC or droplet size – the climate relevant microphysical properties. Perhaps a clearer message concerning what we would like to learn from these measurements would help the description of the results, otherwise the results may be misinterpreted by the less well-initiated reader.

We thank the reviewer for these comments and fully agree that the mass vs. number discussion was not clearly addressed. We accordingly included the suggestions in Chapter 5.1.2 and it now reads:

“These compounds were also observed in the coarse fraction of the aerosol particles, *suggesting that the coarse mode particles served as efficient CCN and were efficiently transferred to the cloud water. To emphasize, these chemical analyses are based on mass, but the control of the cloud droplet number concentration comes from CCN number concentrations, including all particles with sizes of roughly above 100 nm. As larger particles contribute more to the total mass, chemical bulk measurements give no information about a direct influence of sea spray particles on cloud droplet concentrations, but it can show that the chemical composition is consistent with an (expected) oceanic influence on cloud water.*”

And at the end of the paragraph:

“In summary, cloud water chemical composition seemed to be *dominated* by coarse mode aerosol particle composition, and the presence of inorganic marine tracers (sodium, methane sulfonic acid) *shows that material from the ocean is transported to the atmosphere where it can become immersed in cloud droplets. More detailed investigations on the chemical composition, including comparison of constituents from submicron aerosol particles and the SML with the cloud water composition are planned.*”

Another aspect that suggests an influence of marine-derived particles on cloud processes is the finding of TEP (particles that are clearly of marine origin), in submicron aerosol particles. While these data are preliminary, the occurrence of these ocean-derived compounds in very small particles might be related to cloud processes that will be investigated in future studies.

We have mentioned this in chapter 5.7.2 and underlined it more clearly:

“Interestingly, a major part of TEP seems to be located in the sub-micron aerosol particles (Fig. 19). Sub-micron aerosol particles represent the longest living aerosol particle fraction and have a high probability to reach cloud level and *to contribute to cloud formation* and the occurrence of TEP in cloud water, which strongly underlines a possible vertical transport of these ocean-derived compounds.”

Finally, we included the particle mass/number issue in the conclusion as follows:

We clearly see a link between the ocean and the atmosphere as (i) the particles measured at the surface are well mixed within the marine boundary layer up to cloud level and (ii) ocean-derived compounds can be found in the (*submicron*) aerosol particles at mountain height and in the cloud water. The organic measurements will be implemented in a new source function for the oceanic emission of OM. From a perspective of particle number concentrations, the marine contributions to both CCN and INP are, however, rather limited. *These findings underline that further in depth studies differentiating between submicron and supermicron particles as well as between aerosol number and aerosol mass are strongly required.*

Line 618-619: The dominance of C16:0 and C18:0 fatty acids aligns well with the findings of Cochran et al. (Environ Sci Technol, 2016) from sea spray tank studies. Making this type of connection for the reader may help their interpretation of the rather quick overview of this topic in this paper.

Thank you for this recommendation. We have included this reference, and it now reads: “This result aligns well with the findings of Cochran, et al. (2016) from sea spray tank studies that connected the transfer of lipid-like compounds to their physicochemical properties such as solubility and surface activity.”

Figure 20: There is an error in the axis label for panel (a). The exponent is missing.

This is actually not an error. We explain in the figure caption that “*In panel a, please note the different power values between SML/ ULW (10^6 cells mL^{-1}) and cloud water samples (10^4 cells mL^{-1})*”.

Section 5.7.3: Based on using just a few measurements of total bacterial abundance without process-level context, it is challenging to see that the question about the involvement of bacteria in this ocean surface/marine boundary layer system will be adequately addressed. While general reporting of the numbers is provided and is another data point to compare with across the globe, the identity of the bacteria involved is going to be of utmost importance. Some bacteria that are in high abundance are not very productive (reduced carbon utilization), so bulk number concentrations may mask the active processes that control the overall behavior driven by bacteria as a whole.

Thank you for this comment. We absolutely agree about the limitations of bacterial abundance data to describe the direct influence of bacteria on air-sea interaction processes. During our campaign, we performed several incubation experiments with aerosol samples to determine the oxygen turnover of aerolised bacterial communities. Although we used samples with varying biomass collected (i.e. aerosol sampling times) and performed different incubation times (up to > 1 day) during these experiments, most rates obtained were close to/at the detection limit, hinting for a low (due to low abundance and/or high stress?), but yet to be determined activity pattern of aerosol communities. We do not present these data, as we want to avoid mis- or overinterpretation of these activity measurements. In line with the reviewers' suggestion, however, we are currently analysing the bacterial community structure by 16S rDNA amplicon sequencing from a subsequent study. These results will be used to study the selective transport of bacteria from the ocean, through the SML into the aerosol fraction. We envision similar approaches with our water and aerosol samples taken at the Cape Verde. This approach will certainly allow a better understanding of bacterial involvement in air-interaction processes, e.g. OM turnover. In order to actually assess the role of microorganism in air-sea interaction processes, a direct link between activity and identity would be needed in future studies, such as applying stable isotope probing in combination with molecular biology techniques (e.g. RNA-SIP).

We address this point by including the following sentence in the revised manuscript in chapter 5.7.3 and added: “*Further ongoing investigations aim to determine the bacterial community composition by 16S sequencing approaches. The resulting comparison of water and aerosol*

particle samples will help to better understand the specificity of the respective communities and to gain insights into the metabolic potential of abundant bacterial taxa in aerosol particles.”

Additional information: Data availability: We uploaded our data on World Data Centre PANGAEA (<https://www.pangaea.de/>) and included the respective DOIs in the revised manuscript.

Replies to Referee 2:

The manuscript presents an overview of the MarParCloud campaign at Cape Verde Islands in Sept – Oct 2017. Several interesting new scientific findings are reported in brief or just mentioned, but the main scope here is to present a synopsis of all oceanographic and atmospheric observations that have been carried out during the field campaign. If the Authors' intention was to give a flavour of a very multidisciplinary experiment, I think this emerges quite clearly from the paper. I have only two major remarks about the science (see my major comments below). Besides, since most of the results are object of specific papers in preparation, my remarks are mainly on the quality of the presentation.

We thank the Reviewer for the evaluation and the constructive comments. Replies to the specific Referee's comments are provided below in red and new parts included in the manuscript are marked in *italics*:

Major comments:

1. The Abstract is rather descriptive but it ends up with a strong statement: “from a perspective of particle number concentrations, marine contributions to both CCN and INP are rather limited”. I think that the Authors here should make an effort to a) be more quantitative (or refer to other papers either published or in preparation) and b) make sure that there is enough information in the Abstract as well as in the main body of the manuscript to support such conclusion. I have not found any sections in the paper dealing with CCN, just a brief mention at line 560 focusing on the proportion of CCN at low supersaturations accounted for by coarse particles, but what about the estimated contribution by primary marine aerosols?

We agree that the CCN results and the respective marine contributions were not discussed in sufficient detail and we extended this part. Additional information on CCN was added to Chapter 5.1.1., including also a new Figure (Fig. 9) with CCN number concentrations for a

range of super-saturations and for dust and marine air masses. In addition, the fraction of sea spray particles in all CCN and in all particles is now given. The respective text in the second paragraph of Chapter 5.1.1. now reads as follows:

“ N_{CCN} at different supersaturations were compared during dust and marine periods, as shown in Figure 9. During dust periods, the aerosol particles show a great enhancement in Aitken, accumulation and coarse mode number concentrations, such that overall N_{CCN} increases distinctly. N_{CCN} at a supersaturation of 0.30% (proxy for the supersaturation encountered in clouds present during the campaign) during the strongest observed dust periods is about 2.5 times higher than that during marine periods. As suggested by Modini, et al. (2015), Wex, et al. (2016) and Quinn, et al (2017), the coarse mode aerosol particles can be attributed to sea spray aerosol (SSA) in a marine environment. In these studies, the fraction of sea spray aerosol was determined based on three-modal fits from which the particle number concentrations in the different modes were determined. A similar analysis was done for this study. During marine periods, SSA accounted for about 3.7% of CCN number concentrations at 0.30% supersaturation and for 1.1% to 4.4% of N_{total} (total particle number concentration). The hygroscopicity parameter kappa (κ) averaged 0.28, suggesting the presence of OM in the particles.”

In addition, in order to be more quantitative, we included results from a study of free amino acids in all marine compartments (seawater, aerosol particles, cloud water) that was very preliminary at the time of writing the first version of the current manuscript but are currently more advanced and published at ACPD (Triesch, et al. 2020) within this SI. We referred to the results that are discussed in the separate paper but included the main findings in the overview paper in section 5.7.1 and it reads: *“A more comprehensive set of samples was analysed for FAA on molecular level as important organic nitrogen- containing compounds (Triesch, et al. 2020). The FAA, likely resulting from the ocean, were strongly enriched in the submicron aerosol particles ($EF_{aer (FAA)} 10^2$ - 10^4) and to a lower extent enriched in the supermicron aerosol particles ($EF_{aer (FAA)} 10^1$). The cloud water contained the FAA in significantly higher concentrations compared to their respective seawater concentrations and they were enriched by a factor of $4 \cdot 10^3$ compared to the SML. These high concentrations cannot be currently explained and possible sources such as biogenic formation or enzymatic degradation of proteins, selective enrichment processes or pH dependent chemical reactions are subject to future work. The presence of high concentrations of FAA in submicron aerosol particles and in cloud water together with the presence of inorganic marine tracers (sodium, methane sulfonic acid) point to an influence of oceanic sources on the local clouds (Triesch, et al. 2020).”*

This further shows a (qualitative) link between ocean-derived compounds being transferred to the atmosphere up to cloud level. In the abstract, such information had been added before but are specified now and it reads *“Organic nitrogen compounds (free amino acids) were enriched by several orders of magnitude in submicron aerosol particles and in cloud water compared to seawater.”*

With these modifications (and the upcoming ones about INP, see 2.) we included more solid facts in the manuscript and we think there is now enough information in the abstract as well as in the main body of the manuscript to support our conclusion. Finally, we want to point out that this manuscript is intended to provide an overview about the MarParCloud campaign and give a basis and orientation to the single papers on the specific topics of MarParCloud that are partly under revision and partly being currently finalized.

2. There is instead a section about INP (5.7.4) providing a short summary of the study of Gong et al. (2019b) and concluding that primary marine INP should be four orders of magnitude more abundant to account for the ambient INP concentrations measured in cloud and aerosol samples. If this is the basis for the final statement included in the Abstract, I suggest to report it along with the main hypothesis made by the Authors who assume “INP not enriched or altered during the production of sea spray” from the SML (lines 966 – 967), which is a strong assumption, in my opinion. Otherwise, more supporting information can be extracted from Gong et al..

We agree with the referee that a more comprehensive discussion about the INP findings is needed and thus more details and explanations on the INP analysis were added as follows:

In the main text (section 5.7.4.):

“ N_{INP} in PM_1 were generally lower than those in PM_{10} and, furthermore, N_{INP} in PM_1 at CVAO did not show elevated N_{INP} at warm temperatures, in contrast to N_{INP} in PM_{10} . These elevated concentrations in PM_{10} decreased upon heating the samples, clearly pointing to a biogenic origin of these INP. Therefore, ice active particles in general and biologically active INP in particular were mainly present in the supermicron particles, and particles in this size range are not suggested to undergo strong enrichment of OM during oceanic transfer via bubble bursting (Quinn et al., 2015 and refs. therein). “

Furthermore, we pointed out that INP are assumingly not enriched or altered in the supermicron mode during the production of sea spray as follows:

*“Assuming sea salt and the INP to be similarly distributed in both sea and cloud water (i.e., assuming that INP would not be enriched or altered during the production of *supermicron* sea spray particles), N_{INP} is at least four orders of magnitude higher than what would be expected if all airborne INP would originate from sea spray.”*

We performed changes and clarifications concerning the INP measurements in the abstract:

“However, INP measurements indicated also a significant contribution of other non-marine sources to the local INP concentration, as (biologically active) INP were mainly present in supermicron aerosol particles that are not suggested to undergo strong enrichment during ocean-atmosphere transfer.”

And in the conclusion:

“However, based on the findings that (biologically active) INP were mainly present in supermicron aerosol particles that are not suggested to undergo strong enrichment during ocean-atmosphere transfer as well as the INP abundance in seawater and in cloud water, other non-marine sources most likely significantly contributed to the local INP concentration.”

Specific comments:

1. Three of the seven research questions (page 5) should be retuned:

a. Question #1: specify what are the metrics of interest (number, mass, CCN, etc.).

We included the specific metrics of interest and it now reads:

- To what extent is seawater a source of OM on aerosol particles (regarding number, mass, chemical composition, CCN and INP concentration) and in cloud water?

b. Question #2: do “OM groups” mean source contributions or chemically-defined classes?

We mean chemically defined classes and changed it to:

- What are the important chemically-defined OM groups (proteins, lipids, carbohydrates - as sum parameters and on molecular level) in oceanic surface films, aerosol particles and cloud water and how are they linked?

c. Question #3: this reads like a rhetorical question; it should be restructured into something like “What are the main biological and physical factors responsible for the occurrence and accumulation of OM etc.”

We changed question #3 to:

- What are the main biological and physical factors responsible for the occurrence and accumulation of OM in the surface film and in other marine compartments (aerosol particles, cloud water)?

2. Line 225: Hg cannot be considered a good “example for trace metals”, as it exhibits unique chemical properties.

We agree that this sentence was not well phased and replaced it with: “Ocean surface mercury (Hg) associated with OM was investigated.”

3. Line 226: “pigments [...] were captured..”, I do not think the verb is appropriate.

We agree and replaced “captured” by “analysed”.

4. Line 227: “...(DMS), VOCs..”; as DMS is a VOC, this should be better put as “..(DMS), other VOCs..”.

We changed the sentence as suggested.

5. Line 439: “These issues will be further analysed”, does it mean elsewhere in the paper or in a future publication? Please specify.

We specified it and it now reads: “These issues will be analysed in further studies.”

6. Section 4.1.4: What is the relevance/representativeness of the five cloud scenes?

The different cloud times have shown to affect the in-cloud time of an air parcel that in turn affects in-cloud chemical processes (e.g. Lelieveld and Crutzen, 1991). For example, stratocumulus clouds is the time an air parcel spends in a cloud higher (at low mixing) compared to the in-cloud time of an air parcel in cumuli clouds. Moreover, it has been shown that the formation of MSA is enhanced when strong in-cloud processing occurs. We have added these considerations and therefore the relevance together with the respective references in a concluding sentence in chapter 4.1.4 and it now reads:

“The different cloud scenes reflect typical situations observed in conditions with either weaker or stronger winds. The average in-cloud time of an air parcel might depend on cloud type and cloud cover that in turn impacts in-cloud chemical processes (e.g. Lelieveld and Crutzen, 1991), such as the formation of methane-sulfonic acid and other organic acids (Hoffmann et al., 2016 und Chen et al., 2018). Future studies will relate the chemical composition of the aerosol particles and cloud water to the cloud scenes and their respective oxidation capacity. However, the rather coarse horizontal resolution of the satellite sensor and the missing information about time-resolved vertical profiles of thermodynamics and cloud condensate limits a further detailed characterization of these low-level cloud fields and their formation processes. A synergistic combination with ground-based in-situ and remote sensing measurements would be highly beneficial for future investigations to elucidate how cloud chemistry might be different for the varying cloud scenes depending on horizontal cloud patterns and vertical cloud structures.”

7. Lines 478 – 480: “A synergic combination with ground-based in-situ and remote sensing measurements would be highly beneficial for future investigations”. The sense of this sentence is rather obscure (“beneficial” for what? For which of the seven re-search questions listed in the introduction??), please clarify.

As mentioned above, we have changed the statement and added: “A synergistic combination with ground-based in-situ and remote sensing measurements would be highly beneficial for future investigations to elucidate how cloud chemistry might be different for the varying cloud scenes depending on horizontal cloud patterns and vertical cloud structures.”

The cloud types implicitly belong to these two (revised) research questions:

- To what extent is seawater a source of OM to aerosol particles (regarding number, mass, chemical composition, CCN and INP concentration) and in cloud water?
- What are the important chemically-defined OM groups (proteins, lipids, carbohydrates - as sum parameters and on molecular level) in oceanic surface films, aerosol particles and cloud water and how are they linked?

Although studies of in-cloud chemistry in relation to the cloud scenes were not the aim of the campaign, the differentiation between the different cloud scenes provides a first step for such analysis in the future and are in our opinion worth to be shown here as auxiliary information.

8. Section 4.2. I suggest to report here only the concentration levels of the main indexes of biological productivity (chlorophyll concentrations) and their spatial distribution (sub-section 4.2.2 is ok), while I would postpone the discussion about pigment distribution to section 5.4.

We thank the referee for this suggestion. We understood that the order of presentation may lead to some confusion and have re-structured this part (following also the suggestions of reviewer #1). In the revised version, we mentioned the chl-a concentrations in section 4.2. and shifted the discussion of the pigment results to section 5.4.1. To underline that pigments as well as bacteria were analysed we:

i) renamed section 4.2.1 to “Pigment and bacteria concentration in seawater”

ii) included this information in section 4.2.1 and it now reads “ *Chl-a concentrations varied between 0.11 $\mu\text{g L}^{-1}$ and 0.6 $\mu\text{g L}^{-1}$, and are more thoroughly discussed together with the pigment composition in section 5.4.1. Moreover, as other but phytoplankton organisms can contribute to the OM pool, bacterial abundance was analysed in the SML and bulk water samples and these data are reported in section 5.7.3.*”

iii) concluded section 5.4.1. with: “ First analyses show that the DOC concentrations were not directly linked to the increasing chl-a concentrations, however their relation to single pigments, *to the microbial abundance*, to the background dust concentrations and finally to wind speed and solar radiation will be further resolved to elucidate potential biological and meteorological controls on the concentration and enrichment of DOC.”

Due to the inclusion of the pigment discussion in chapter 5.4.1. (that therefore became very long) we have added another subsection for the comparison of DOC data from the two sampling techniques (5.4.2 DOC concentrations: manual glass plate vs. MarParCat sampling) in 5.4.2.

9. Line 535: satellite fluorescence measurements. Which satellite?

They were achieved from the MODIS-Terra satellite, this is stated in the Figure caption and we also added this information in the text.

10. Section 5.1.1: see major comments.

We included CCN information and discussion as described above in 1).

11. Lines 602-603: “..suggested an ocean influence on cloud water”. Please, be more precise here. The data show a cloud water composition dominated by seasalt with little quantities of other solutes: this is simply the effect of the larger scavenging efficiency of coarse particles with respect to submicron ones, but it is worth reminding that this picture is “mass-based” (all cloud drops coalesce into one single sample inside the CASC) while in terms of number concentrations (how many cloud drops originated from marine sources), it is difficult to tell solely on the basis of the data shown in Fig.11.

We thank the reviewer for these comments and fully agree that the mass vs. number discussion was not clearly addressed. We accordingly included the suggestions in Chapter 5.1.2 and it now reads:

“These compounds were also observed in the coarse fraction of the aerosol particles, suggesting that the coarse mode particles served as efficient CCN and were efficiently transferred to the cloud water. *To emphasize, these chemical analyses are based on mass, but the control of the cloud droplet number concentration comes from CCN number concentrations, including all particles with sizes of roughly above 100 nm. As larger particles contribute more to the total mass, chemical bulk measurements give no information about a direct influence of sea spray particles on cloud droplet concentrations, but it can show that the chemical composition is consistent with an (expected) oceanic influence on cloud water.*”

And at the end of the paragraph:

“In summary, cloud water chemical composition seemed to be *dominated* by coarse mode aerosol particle composition, and the presence of inorganic marine tracers (sodium, methane sulfonic acid) *shows that material from the ocean is transported to the atmosphere where it can become immersed in cloud droplets. More detailed investigations on the chemical composition, including comparison of constituents from submicron aerosol particles and the SML with the cloud water composition are planned.*”

Another aspect that suggests an influence of marine-derived particles on cloud processes is the finding of TEP (particles that are clearly of marine origin), in submicron aerosol particles. While these are first results, the occurrence of these ocean-derived compounds in very small particles might be related to cloud processes that will be investigated in future studies.

We have mentioned this in chapter 5.7.2 and underlined it more clearly:

“Interestingly, a major part of TEP seems to be located in the sub-micron aerosol particles (Fig. 19). Sub-micron aerosol particles represent the longest living aerosol particle fraction and have a high probability to reach cloud level and *to contribute to cloud formation* and the occurrence of TEP in cloud water, which strongly underlines a possible vertical transport of these ocean-derived compounds.”

Finally, we included the particle mass/number issue in the conclusion as follows:

“We clearly see a link between the ocean and the atmosphere as (i) the particles measured at the surface are well mixed within the marine boundary layer up to cloud level and (ii) ocean-derived compounds can be found in the (*submicron*) aerosol particles at mountain height and in the cloud water. The organic measurements will be implemented in a new source function for the oceanic emission of OM. From a perspective of particle number concentrations, the marine contributions to both CCN and INP are, however, rather limited. *These findings underline that further in depth studies differentiating between the aerosol number and aerosol mass are strongly required.*”

12. Line 644: “.. by southern Hemisphere”: it is not clearly shown in Figure S1.

Fig. S1 shows the air mass back trajectories and from the plots, it seems that the air masses partly passed the Southern Hemisphere. However, this is subjective information and from the wind direction plot, this was not confirmed. We agree with the reviewer that “by the southern Hemisphere” could not clearly been shown and therefore we have revised this part. It now reads: “High DMS concentrations on September 19th – 20th occurred when air originated predominantly from the Mauritanian upwelling region (Figure SI1) and on September 26th and 27th. ” As furthermore stated, ”These elevated concentrations will be linked to the phytoplankton composition ... ” to elucidate further biological connections.

13. Section 5.3.2: The results about HONO look so preliminary that I am not sure they deserve a dedicated section. They can be reduced to a short paragraph at the end of the previous section about trace gases.

We agree and included the HONO discussion in the trace gas section 5.3.

14. Line 850: Enrichment factors for DOM are reported. Please specify the concentration unit used for DOM (organic carbon, organic nitrogen or UV absorption?).

The DOM comprises the sum of the single DOM fractions (in $\mu\text{g/L}$) and the EF is the (dimensionless) ratio between the DOC/DOM in the SML and the ULW. We added this information in chapter 5.7.1 and it now reads: “*The DOM concentrations were derived from the sum of the individual compound groups (in $\mu\text{g L}^{-1}$) and the EFs for DOM varied from 0.83 to 1.46, which agreed very well to the DOC measurements described in section 5.4.1.*”

15. Lines 987 – 988: “.. a daily variation of the number of particles formed was observed (but from a very limited set of samples, $n = 3$) probably related to the daily sampling conditions. To explain these observations, two different hypothesis can be postulated...”
Actually, it is not clear at all what observations the Authors are referring to, because no data are shown but simply a “daily variation” is observed and only for three samples. Similarly to the HONO case, the impression is that the state of the analysis of the Go:PAM dataset is just too preliminary to be discussed in a dedicated section.

We thank the reviewer for this comment. In fact, the results were very preliminary at the time of writing the first version of the current paper. However, the data interpretation has strongly proceeded in the meantime and the data are included and discussed in a separate paper by Zabalegui, et al., 2019. In the revised version, we referred to this paper (that is also published within this SI) and shortly summarized the main findings and it reads now: “A subset of the collected SML samples were investigated within the Go:PAM and showed that particles were formed when these samples were exposed to actinic irradiation. These particles resulted most likely from the reaction of ozone with gaseous products that were released from the SML as shown recently (Ciuraru et al. 2015) and the results obtained herein are explained in more detail in a separate paper by Zabalegui, et al., 2019. “

16. Section 5.9.1. Please provide a short description of COSMO.

We included a short description of COSMO and it now reads: “*COSMO is a compressible and non-hydrostatic meteorological model and the current weather forecast model of the German Weather Service. The numerical calculation of the weather forecast is achieved by using information of the underlying orography and land-use, as well as boundary data of all meteorological fields. The needed boundary and initial fields will be derived from re-analysis-data and/or input parameters from coarse-resolved weather model data.*”

17. Section 5.9.2. Please provide a short description of MUSCAT.

We included a short description of MUSCAT and it now reads: “*The new emission scheme will be implemented to the aerosol chemical transport model MUSCAT (Multi-Scale Chemistry Aerosol Transport). MUSCAT is able to treat atmospheric transport and chemical transformation of different traces gases as well as particle properties. In addition to advection and turbulent diffusion, sedimentation, dry and wet deposition through the transport processes are considered, too. MUSCAT is coupled with COSMO that provides MUSCAT with all needed meteorological fields (Wolke, et al. 2004). The multiscale model system COSMO-MUSCAT will be used further to validate the emission scheme of OM via small and meso-scale simulations.* “

18. The graphical quality of the figures must be improved.

The quality of the Figures appears partly poor due to the “copy and paste” procedure to the pdf document. We will provide the graphics in highest resolution and will upload them as a separate file.

19. Figure S4: there is something wrong in this figure

We thank the reviewer for this note and corrected the Figure S4. It now clearly shows the cloud events (according to the method of Gong e al. 2019a).

Additional information: Data availability: We uploaded out data on World Data Centre PANGAEA (<https://www.pangaea.de/>) and included the respective DOIs in the revised manuscript.

Marine organic matter in the remote environment of the Cape Verde Islands – An introduction and overview to the MarParCloud campaign

Manuela van Pinxteren^{1*}, Kanneh Wadinga Fomba¹, Nadja Triesch¹, Christian Stolle^{2,3}, Oliver Wurl³, Enno Bahlmann^{2,4}, Xianda Gong¹, Jens Voigtländer¹, Heike Wex¹, Tiera-Brandy Robinson³, Stefan Barthel¹, Sebastian Zeppenfeld¹, Erik H. Hoffmann¹, Marie Roveretto⁵, Chunlin Li⁵, Benoit Grosselin⁶, Veronique Daële⁶, Fabian Senf¹, Dominik van Pinxteren¹, Malena Manzi⁷, Nicolás Zabalegui⁷, Sanja Frka⁸, Blaženka Gašparović⁸, Ryan Pereira⁹, Tao Li¹⁰, Liang Wen¹⁰, Jiarong Li¹¹, Chao Zhu¹¹, Hui Chen¹¹, Jianmin Chen¹¹, Björn Fiedler¹², Wolf von Tümpling¹³, Katie A. Read¹⁴, Shalini Punjabi^{14,15}, Alastair C. Lewis^{14,15}, James R. Hopkins¹⁴, Lucy J. Carpenter¹⁵, Ilka Peeken¹⁶, Tim Rixen⁴, Detlef Schulz-Bull², María Eugenia Monge⁷, Abdelwahid Mellouki^{6,10}, Christian George⁵, Frank Stratmann¹, Hartmut Herrmann^{1,10*}

*corresponding authors: Manuela van Pinxteren (manuela@tropos.de) and Hartmut Herrmann (herrmann@tropos.de)

¹ Leibniz-Institute for Tropospheric Research (TROPOS), 04318 Leipzig, Germany

² Leibniz-Institute for Baltic Sea Research Warnemuende, 18119 Rostock, Germany

³ Institute for Chemistry and Biology of the Marine Environment, Carl-von-Ossietzky University Oldenburg, 26382 Wilhelmshaven, Germany

⁴ Leibniz Centre for Tropical Marine Research (ZMT), 28359 Bremen, Germany

⁵ Institut de Recherches sur la Catalyse et l'Environnement de Lyon, Lyon, France.

⁶ Institut de Combustion, Aérothermique, Réactivité et Environnement, Centre National de la Recherche Scientifique, Orléans, France.

⁷ Centro de Investigaciones en Bionanociencias (CIBION), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), C1425FQD, Ciudad de Buenos Aires, Argentina

⁸ Division for Marine and Environmental Research, Ruđer Bošković Institute, 10000 Zagreb, Croatia

⁹ Lyell Centre, Heriot-Watt University, EH14 4AP, Edinburgh, United Kingdom

¹⁰ School of Environmental Science and Engineering, Shandong University, Qingdao 266237, China

¹¹ Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention, Institute of Atmospheric Sciences, Fudan University, Shanghai, 200433, China

¹² GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany

¹³ Helmholtz Centre for Environmental Research - UFZ, 39114, Magdeburg, Germany

¹⁴ National Centre for Atmospheric Science (NCAS), University of York, Heslington, York, YO10 5DD

¹⁵ Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, Heslington, York, YO10 5DD

¹⁶ Alfred-Wegener-Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

Abstract

The project MarParCloud (Marine biological production, organic aerosol Particles and marine Clouds: a process chain) aims at improving our understanding of the genesis, modification and impact of marine organic matter (OM), from its biological production, via its export to marine aerosol particles and, finally, towards its ability to act as ice nucleating particles (INP) and cloud condensation nuclei (CCN). A field campaign at the Cape Verde Atmospheric Observatory (CVAO) in the tropics in September/October 2017 formed the core of this project that was jointly performed with the project MARSU (MARine atmospheric Science Unravelled). A suite of chemical, physical, biological and meteorological techniques was applied and comprehensive measurements of bulk water, the sea surface microlayer (SML), cloud water and ambient aerosol particles collected at a ground-based and a mountain station took place.

Key variables comprised the chemical characterization of the atmospherically relevant OM components in the ocean and the atmosphere as well as measurements of INP and CCN. Moreover, bacterial cell counts, mercury species and trace gases were analysed. To interpret the results, the measurements were accompanied by various auxiliary parameters such as air mass back trajectory analysis, vertical atmospheric profile analysis, cloud observations and pigment measurements in seawater. Additional modelling studies supported the experimental analysis.

During the campaign, the CVAO exhibited marine air masses with low and partly moderate dust influences. The marine boundary layer was well mixed as indicated by an almost uniform particle number size distribution within the boundary layer. Lipid biomarkers were present in the aerosol particles in typical concentrations of marine background conditions. Accumulation and coarse mode particles served as CCN and were efficiently transferred to the cloud water. The ascent of ocean-derived compounds, such as sea salt and sugar-like compounds, to the cloud level as derived from chemical analysis and atmospheric transfer modelling results denote an influence of marine emissions on cloud formation. Organic nitrogen compounds (free amino acids) were enriched by several orders of magnitude in submicron aerosol particles and in cloud water compared to seawater. However, INP measurements indicated also a significant contribution of other non-marine sources to the local INP concentration, as (biologically active) INP were mainly present in supermicron aerosol particles that are not suggested to undergo strong enrichment during ocean-atmosphere transfer. ~~or strong enrichment processes during upward transport.~~ In addition, the number of CCN at the supersaturation of 0.30% was about 2.5 times higher during dust periods compared to marine periods. Lipids, sugar-like compounds, UV absorbing humic-like substances and low molecular weight neutral components were important organic compounds in the seawater and highly surface-active lipids were enriched within the SML. The selective enrichment of specific organic compounds in the SML needs to be studied in further detail and implemented in an OM source function for emission modelling to better understand transfer patterns, mechanisms of marine OM transformation in the atmosphere and the role of additional sources.

In summary, when looking at particulate mass, we do see oceanic compounds transferred to the atmospheric aerosol and to the cloud level, while from a perspective of particle number concentrations, marine contributions to both CCN and INP are rather limited.

Keywords

MarParCloud, MARSU, organic matter, seawater, sea surface microlayer, aerosol particles, cloud water, Cape Verde Atmospheric Observatory (CVAO)

1 Introduction and Motivation

The ocean covers around 71% of the Earth's surface and acts as a source and sink for atmospheric gases and particles. However, the complex interactions between the marine boundary layer (MBL) and the ocean surface are still largely unexplored (Cochran, et al. 2017; de Leeuw, et al. 2011; Gantt and Meskhidze 2013; Law, et al. 2013). In particular, the role of marine organic matter (OM) with its sources and contribution to marine aerosol particles, is still ~~poorly understood~~elusive. For example, where this particle fraction might lead to a variety of effects such as impacting health through the generation of reactive oxygen species, OM composition increasing or decreasing the absorption of solar radiation and therefore radiative properties, and impacting marine ecosystems via atmospheric deposition ~~where this particle fraction might lead to a variety of effects such as changing health effects, changing radiative properties, changing effects of marine particles deposited to the ecosystems~~ (e.g. Abbatt, et al. 2019; Brooks and Thornton 2018; Burrows, et al. 2013; Gantt and Meskhidze 2013; Pagnone, et al. 2019; [Patel and Rastogi, 2020](#)). Furthermore, knowledge on the properties of marine organic aerosol particles and their ability to act as cloud condensation nuclei (CCN) or ice nucleating particle (INP) is ~~still elusive~~not fully understood. The fraction of marine CCN made up of sea spray aerosol is still debated and suggested to comprise about 30% on a global scale (excluding the high southern latitudes) (Quinn, et al. 2017) and important pieces of information about marine CCN are still missing (e.g. Bertram, et al. 2018). Ocean-derived INPs were proposed to play a dominating role in determining INP concentrations in near-surface-air over the remote areas such as the Southern Ocean, however their source strength in other oceanic regions as well as knowledge about which physicochemical properties determine the INP efficiency ~~is~~are still largely unknown (Burrows, et al. 2013; McCluskey, et al. 2018a; McCluskey, et al. 2018b). In recent years, it was clearly demonstrated that marine aerosol particles contain a significant organic mass fraction derived from primary and secondary processes (Middlebrook, et al. 1998; Prather, et al. 2013; Putaud, et al. 2000; van Pinxteren, et al. 2017; van Pinxteren, et al. 2015). Although it is known that the main OM groups show similarities to the oceanic composition and comprise carbohydrates, proteins, lipids as well as humic-like and refractory organic matter, a large fraction of OM in the marine environment is still unknown at a molecular level, thereby limiting our ability to constrain interlinked processes (e.g. Gantt and Meskhidze 2013).

The formation of ocean-derived aerosol particles and their precursors is influenced by the uppermost layer of the ocean, the sea surface microlayer (SML) which forms due to different physicochemical properties of air and water (Engel, et al. 2017; Wurl, et al. 2017). Recent investigations suggest that the SML is stable up to wind speeds of $> 10 \text{ m s}^{-1}$ and is therefore

existent at the global average wind speed of 6.6 m s^{-1} and a fixed component influencing the ocean atmosphere interaction on global scales (Wurl, et al. 2011). The SML is involved in the generation of sea-spray (or primary) particles including their organic fraction by ~~either transfer of OM to rising bubbles before they burst out~~ to jet droplets and film droplets (de Leeuw, et al. 2011). ~~or through a more direct transfer of OM from the ocean compartments to the marine particles.~~ A mechanistic and predictable understanding of these complex and interacting processes is still lacking (e.g. Engel, et al. 2017). Moreover, surface films influence air-sea gas exchange and may undergo (photo)chemical reactions leading to a production of unsaturated and functionalized volatile organic compounds (VOCs) acting as precursors for the formation of secondary organic aerosol (SOA) particles (Brueggemann, et al. 2018; Ciuraru, et al. 2015). Thus, dynamics of OM and especially surface-active compounds present at the air-water interface may have global impacts on the air-sea exchange processes necessary to understand oceanic feedbacks on the atmosphere (e.g. Pereira, et al. 2018).

Within the SML, OM is a mixture of different compounds ~~such as including~~ polysaccharides, amino acids, proteins, lipids ~~and it occurs as and particulate and~~ chromophoric dissolved organic matter (CDOM) that are either dissolved or particulate (e.g. Gašparović, et al. 1998a; Gašparović, et al. 2007; Stolle, et al. 2019). In addition, the complex microbial community is assumed to exert a strong control on the concentration and the composition of OM (Cunliffe, et al. 2013). In calm conditions, bacteria accumulate in the SML (Rahlff, et al. 2017) and are an integral part of the biofilm-like habitat forming at the air-sea interface (Stolle, et al. 2010; Wurl, et al. 2016).

A variety of specific organic compounds such as surface-active substances (SAS), volatile organic compounds (VOC), and acidic polysaccharides aggregating to transparent exopolymer particles (TEP), strongly influence the physico-chemical properties of OM in the SML. SAS (or surfactants) are highly enriched in the SML relative to bulk water and contribute to the formation of surface films (Frka, et al. 2009; Frka, et al. 2012; Wurl, et al. 2009). SAS are excreted by phytoplankton, during zooplankton grazing and bacterial activities (e.g. Gašparović, et al. 1998b). The enrichment of SAS in the SML occurs predominantly via advective and diffusive transport at low wind speeds or bubble scavenging at moderate to high wind speeds (Wurl, et al. 2011). When transferred to the atmosphere, OM with surfactant properties, ubiquitously present in atmospheric aerosol particles, has the potential to affect the cloud droplet formation ability of these particles (e.g. Kroflič, et al. 2018).

Sticky and gel-like TEP are secreted by phytoplankton and bacteria and can form via abiotic processes (Wurl, et al. 2009). Depending on their buoyancy they may contribute to sinking particles (marine snow) or can rise and accumulate at the sea surface. Due to their sticky nature TEP is called the “marine glue” and as such it contributes to the formation of hydrophobic films by trapping other particulate and dissolved organic compounds (Wurl, et al. 2016). Additionally, TEP is suspected to play a pivotal role in the release of marine particles into the air via sea spray and bursting bubbles (Bigg and Leck 2008).

Many studies recognize a possible link between marine biological activity and marine-derived organic aerosol particles (Facchini, et al. 2008; O'Dowd, et al. 2004; Ovadnevaite, et al. 2011), and thus to the SML due to the linkages outlined before. Yet, the environmental drivers and mechanisms for the OM enrichment are not very clear (Brooks and Thornton 2018; Gantt and Meskhidze 2013) and individual compound studies can only explain a small part of OM cycling

(e.g. van Pinxteren, et al. 2017; van Pinxteren and Herrmann 2013). The molecular understanding of the occurrence and the processing of OM in all marine compartments is essential for a deeper understanding and for an evidence-based implementation of organic aerosol particles and their relations to the oceans in coupled ocean-atmosphere models. Synergistic measurements in comprehensive interdisciplinary field campaigns in representative areas of the ocean and also laboratory studies under controlled conditions are required to explore the biology, physics and chemistry in all marine compartments (e.g. Quinn, et al. 2015). Accordingly, the project MarParCloud together with contributions from the project MARSU addresses central aspects of ocean atmosphere interactions focusing on the marine OM within an interdisciplinary field campaign at the Cape Verde Islands. Synergistic measurements will deliver an improved understanding of the role of marine organic matter. MarParCloud focuses on the following main research questions:

- To what extent is seawater a source of OM to aerosol particles (regarding number, mass, chemical composition, CCN and INP concentration) and in cloud water?
- What are the important chemically-defined OM groups (proteins, lipids, carbohydrates - as sum parameters and on molecular level) in oceanic surface films, aerosol particles and cloud water (and how are they linked)?
- What are the main biological and physical factors responsible for the occurrence and accumulation of OM ~~Is the occurrence and accumulation of OM~~ in the surface film and in other marine compartments (aerosol particles, cloud water) ~~controlled by biological and meteorological factors?~~
- Which functional role do bacteria play in aerosol particles?
- Does the ~~surface film~~ SML contribute to the formation of ice nuclei, and at what temperatures do these nuclei become ice-active? Are these ice nuclei found in cloud water?
- Does the presence of marine OM in the surface ocean drive the concentration of CCN in the MBL ~~Is the marine OM connected to the CCN concentration in the MBL?~~
- How must an emission parameterization for OM (including individual species) be designed in order to best reflect the concentrations in the aerosol depending on those in seawater or biological productivity under given ambient conditions?

The tropics with a high photochemical activity are of central importance in several aspects of the climate system. Approximately 75% of the tropospheric production and loss of ozone occurs within the tropics, and in particular in the tropical upper troposphere (Horowitz, et al. 2003). The Cape Verde islands are located downwind of the Mauritanian coastal upwelling region off northwest in the islands. In addition, they are in a region of the Atlantic that is regularly impacted by dust deposition from the African Sahara (Carpenter, et al. 2010). The remote station of CVAO is therefore an excellent site for process-oriented campaigns embedded into

the long-term measurements of atmospheric constituents, which are essential for understanding the atmospheric processes and its impact on climate.

2 Strategy of the campaign

The present contribution intends to provide an introduction, overview and first results of the comprehensive MarParCloud field campaign to the MarParCloud Special Issue. We will describe the oceanic and atmospheric ambient conditions at the CVAO site that have not been synthesized elsewhere and are valuable in themselves because of the sparseness of the existing information at such a tropical remote location. Next, we will describe the sampling and analytical strategy during MarParCloud, taking into account all marine compartments i.e. the seawater (SML and bulk water), ambient aerosol particles (at ground-level and the Mt Verde, elevation: 744 m a.s.l.), and cloud water. Detailed aerosol investigations were carried out, both for the chemical composition and for physical properties at both stations. In addition, vertical profiles of meteorological parameters were measured at CVAO using a helikite. These measurements were combined with modelling studies to determine the MBL height. In conjunction, they are an indicator for the mixing state within the MBL providing further confidence for ground-level measured aerosol properties being representative for those at cloud level. The chemical characterization of OM in the aerosol particles as well as in the surface ocean and cloud water included sum parameters (e.g. OM classes like biopolymers and humic-like substances) and molecular analyses (e.g. lipids, sugars and amino acids). Additionally, to address the direct oceanic transfer (bubble bursting), seawater and aerosol particle characterization obtained from a systematic plunging waterfall tank are presented. ~~As an example for trace metals,~~ Ocean surface mercury (Hg) associated with OM was ~~studied~~investigated. Marine pigments and marine microorganisms were ~~captured~~analysed to investigate their relation to OM and to algae produced trace gases. Marine trace gases such as dimethyl sulphide (DMS), other VOCs and oxygenated (O)VOCs were measured and discussed. Furthermore, a series of continuous nitrous acid (HONO) measurements was conducted at the CVAO with the aim of elucidating the possible contribution of marine surfaces at the production of this acid. To explore whether marine air masses exhibit a significant potential to form SOA, an oxidation flow reactor (OFR) was deployed at the CVAO. Finally, modelling studies to describe the vertical transport of selected marine organic compounds from the ocean to the atmosphere up to cloud level taking into account advection and wind conditions will be applied. From the obtained results of organic compound measurements, a new source function for the oceanic emission of OM will be developed. The measurements, first interpretations and conclusions aggregated here will provide a basis for upcoming detailed analysis.

3 Experimental

3.1 General CVAO site and meteorology

The Cape Verde archipelago Islands are situated in the Eastern Tropical North Atlantic (ETNA). The Archipelago experiences strong North-East trade winds that divide the islands into two groups, the Barlavento (windward) and Sotavento (leeward) islands. The North-Western Barlavento Islands of São Vicente and Santo Antão, as well as São Nicolao, are rocky and hilly making them favourable for the formation of orographic clouds.

The CVAO is part of a bilateral initiative between Germany and the UK to conduct long-term studies in the tropical north-east Atlantic Ocean ($16^{\circ} 51.49' \text{ N}$, $-24^{\circ} 52.02' \text{ E}$). The station is located directly at the shoreline at the northeastern tip of the island of São Vicente at 10 m a.s.l. The air temperature varies between 20 and 30 °C with a mean of 23.6 °C. The relative humidity is in average at 79% and precipitation is very low (Carpenter, et al. 2010). Due to the trade winds, this site is free from local island pollution and provides reference conditions for studies of ocean-atmosphere interactions. However, it also lies within the Saharan dust outflow corridor to the Atlantic Ocean and experiences strong seasonal dust outbreaks with peaks between late November and February (Fomba, et al. 2014; Patey, et al. 2015; Schepanski, et al. 2009). Air mass inflow to this region can vary frequently within a day leading to strong inter-day temporal variation in the aerosol mass and chemical composition (Fomba, et al. 2014, Patey, et al. 2015). Despite the predominant NE trade winds, air masses from the USA as well as from Europe are partly observed. However, during autumn, marine air masses are mainly present with few periods of dust outbreaks because at these times the dust is transported at higher altitudes in the Saharan Air Layer (SAL) over the Atlantic to the Americas (Fomba, et al. 2014). During autumn, there is no significant transport of the dust at lower altitudes and only intermittent effects of turbulence in the SAL leads to occasional dust deposition and sedimentation from the SAL to lower altitudes and at ground level. Furthermore, during autumn the mountain site (Mt. Verde) is often covered with clouds as surface temperatures drop after typically very hot summer months. Due to the frequent cloud coverage and less dust influence in autumn, the MarParCloud campaign was scheduled from September 13th to October 13th 2017.

3.2 CVAO equipment during MarParCloud

The setup of the CVAO station is explained in detail in Carpenter, et al. (2010) and Fomba, et al. (2014). During the MarParCloud campaign, the 30 m high tower was equipped with several aerosol particle samplers, including high volume PM_{10} (Digitel, Rieme, Germany), and total suspended particle (TSP, Sieria Anderson, USA) samplers, low volume TSP (homebuilt) and PM_{10} (Comde-Derenda, Germany) samplers and a size-resolved aerosol particle Berner impactor (5 stages). The sampling times were usually set to 24 h (more details in the SI). On-line aerosol instruments included a Cloud Condensation Nuclei counter (CCNC, Droplet Measurement Technologies, Boulder, USA) (Roberts and Nenes 2005) to measure cloud condensation nuclei number concentration (N_{CCN}). A TROPOS-type Scanning Mobility Particle Sizer (SMPS) (Wiedensohler, et al. 2012), and an APS (Aerodynamic Particle Sizer, model 3321, TSI Inc., Paul, MN, USA) were used to measure in the size range from 10 nm to 10 μm . The particles hygroscopicity (expressed as κ (Petters and Kreidenweis 2007)) was derived from combined N_{CCN} and particle number size distributions (PNSDs) measurements from the SMPS and APS. Vertical profiles of meteorological parameters were measured using

a 16 m³ Helikite (Allsopp Helikites Ltd, Hampshire, UK), a combination of a kite and a tethered balloon. Additional equipment at the CVAO station on ground included the plunging waterfall tank, the Long Path Absorption Photometer (LOPAP), and the Gothenburg Potential Aerosol Mass Reactor (Go:PAM) chamber. Further details on the measurements are listed and explained in the SI and all instruments can be found in the Table S1.

3.3 Mt. Verde

Mt. Verde was a twin site for aerosol particle measurements and the only site with cloud water-sampling during the MarParCloud campaign. It is the highest point of the São Vicente Island (744 m) situated in the northeast of the Island (16° 86.95' N, -24° 93.38' E) and northwest to the CVAO. Mt. Verde also experiences direct trade winds from the ocean with no significant influence of anthropogenic activities from the island. Mt. Verde was in clouds during roughly 58% of the time during the campaign. However, the duration of the cloud coverage varied between 2 h and 18 h with longer periods of cloud coverage observed in the nights when surface temperatures dropped.

During the campaign, Mt. Verde was, for the first time, equipped with similar collectors as operated at the CVAO, namely the high volume Digitel sampler for the PM₁ and PM₁₀ bulk aerosol particles, a low volume TSP sampler and a five-stage Berner impactor for the size-resolved aerosol particle sampling. Bulk cloud water was collected using six (4 plastic and 2 stainless steel) compact Caltech Active Strand Cloud water Collectors (CASCC2) (Demoz, et al. 1996). The six samplers were run in parallel for a sampling time between 2.5 and 13 hours collecting between 78 to 544 mL cloud water per sampler in an acid-precleaned plastic bottle. It needs to be pointed out that the aerosol particle samplers run continuously and aerosol particles were also sampled during cloud events. The cloud droplets were efficiently removed due to the pre-conditioning of the aerosol particles sampled with the Berner impactor (more information in the SI) and due to the size cut the PM₁ sampler. However, for aerosol particles sampled with the PM₁₀ sampler, small cloud droplets can be collected as well. In addition, the particles sampled with the low volume TSP sampler can be influenced by cloud droplets to some extent. The cloud liquid water content was measured continuously by a particle volume monitor (PVM-100, Gerber Scientific, USA), which was mounted on a support at the same height with the cloud water samplers. The same suite of on-line aerosol instruments as employed at the CVAO (SMPS, APS, CCNc) was installed at the mountain side. All instruments employed at the Mt. Verde site are listed in the Table S2.

3.4 Oceanographic setting and seawater sampling site

The ETNA around Cape Verde is characterized by a so-called oxygen minimum zone (OMZ) at a water depth of approximately 450 m and by sluggish water velocities (Brandt, et al. 2015). The region is bounded by a highly productive eastern-boundary upwelling system (EBUS) along the African coast, by the Cape Verde Frontal Zone (CVFZ) on its western side, and by zonal current bands towards the equator (Stramma, et al. 2005). Upper water masses towards the archipelago are dominated by North Atlantic Central Water masses (NACW) with enhanced

salinity, whereas the South Atlantic Central Water mass (SACW) is the dominating upper layer water mass in the EBUS region (Pastor, et al. 2008). Filaments and eddies generated in the EBUS region are propagating westwards into the open ocean and usually dissipate before reaching the archipelago. However, observations from the Cape Verde Ocean Observatory (CVOO) 60 nautical miles northeast of the Sao Vicente island (17° 35.00 N', -24° 17.00 E', <http://cvo0.geomar.de>) also revealed the occurrence of water masses originating from the EBUS region which got advected by stable mesoscale eddies (Fiedler, et al. 2016; Karstensen, et al. 2015).

For the MarParCloud campaign, the water samples were taken at Bahia das Gatas, a beach that is situated upwind of the CVAO about 4 km northwest in front of the station. The beach provided shallow access to the ocean that allowed the employment of the fishing boats for manual SML and bulk water sampling and the other equipment. For SML sampling, the glass plate technique as one typical SML sampling strategy was applied (Cunliffe and Wurl 2014). A glass plate with a sampling area of 2000 cm² was vertically immersed into the water and then slowly drawn upwards with a withdrawal rate between 5 and 10 cm s⁻¹. The surface film adheres to the surface of the glass and is removed using framed Teflon wipers (Stolle, et al. 2010; van Pinxteren, et al. 2012). Bulk seawater was collected from a depth of 1 m using a specially designed device consisting of a glass bottle mounted on a telescopic rod used to monitor sampling depth. The bottle was opened underwater at the intended sampling depth with a specifically conceived seal-opener.

In addition, the MarParCat, a remotely controllable catamaran, was applied for SML sampling using the same principle as manual sampling (glass plate). The MarParCat sampled bulk water in a depth of 70 cm. A more detailed description of the MarParCat can be found in the SI. Using the two devices, manual sampling and the MarParCat, between one and six liters of SML were sampled at each sampling event. For the sampling of the SML, great care was taken that all parts that were in contact with the sample (glass plate, bottles, catamaran tubing) underwent an intense cleaning with 10% HCl to avoid contamination and carry over problems.

The sampling sites with the different set up and equipment are illustrated in Figure 1. All obtained SML and bulk water samples and their standard parameters are listed in Table S3.

4 Ambient conditions

4.1 Atmospheric conditions during the campaign

4.1.1 Marine and dust influences

During autumn, marine background air masses are mainly observed at the CVAO, interrupted by a few periods of dust outbreaks (Carpenter, et al. 2010; Fomba, et al. 2014). A 5 years' average dust record showed low concentrations with average values of 25 µg m⁻³ and 17 µg m⁻³ during September and October, respectively (Fomba, et al. 2014). The dust concentrations during the campaign were generally < 30 µg m⁻³ however, strong temporal variation of mineral dust markers were observed (Table 1). According to Fomba, et al. (2013, 2014), a classification into: marine conditions (dust < 5µg/m³, typically Fe < 50 ng m⁻³), low dust (dust < 20 µg/m³) and moderate dust (dust < 60µg/m³) conditions was used to describe the dust influence during

this period. Following this classification, one purely marine period was defined from September 22nd to 24th, which was also evident from the course of the back trajectories (Fig SII). For the other periods, the air masses were classified as mixed with marine and low or moderate dust influences as listed in Table 1. Based on a three-modal parameterization method that regarded the number concentrations in different aerosol particle modes, a similar but much finer classification of the aerosol particles was obtained as discussed in Gong, et al. (2020a).

The classification of the air masses was complemented by air mass backward trajectory analyses. 96 hours back trajectories were calculated on an hourly basis within the sampling intervals, using the HYSPLIT model (HYbrid Single-Particle Lagrangian Integrated Trajectory, <http://www.arl.noaa.gov/ready/hysplit4.html>, 26.07.19) published by the National Oceanic and Atmospheric Administration (NOAA) in the ensemble mode at an arrival height of 500 m \pm 200 m (van Pinxteren, et al. 2010). The back trajectories for the individual days of the entire campaign, based on the sampling interval for aerosol particle sampling, were calculated and are listed in Figure SII. Air parcel residence times over different sectors are plotted in Figure 2. The comparison of dust concentration and the residence time of the back trajectories revealed that in some cases low dust contributions were observed although the air masses travelled almost completely over the ocean (e.g. first days of October). In such cases, entrainment of dust from higher altitudes might explain this finding. The related transport of Saharan dust to the Atlantic during the measurement period can be seen in a visualization based on satellite observations (<https://svs.gsfc.nasa.gov/12772>, last visited on Oct. 1st, 2019). For specific days with a low MBL height, it might be more precise to employ back trajectories that start at a lower height and therefore exclude entrainment effects from the free troposphere for the characterisation of CVAO data. Similarly, for investigating long-lived components, it might be helpful to analyse longer trajectory integration times (e.g. 10 days instead of 4 days). However, the longer the back trajectories, the higher is the level of uncertainty. Regarding aerosol analysis, it is important to notice that dust influences are generally more pronounced on super-micron particles than on sub-micron particles (e.g. Fomba, et al. 2013; Müller, et al. 2009; Müller, et al. 2010) meaning that bigger particles may be affected by dust sources whereas smaller particles may have stronger oceanic and anthropogenic as well as long-range transport influences. Consequently, the herein presented classification represents a first general characterisation of the air mass origins. Depending on the sampling periods of other specific analysis, slight variations may be observed and this will be indicated in the specific analysis and manuscripts.

4.1.2 Meteorological condition

Air temperature, wind direction, wind speed measured between September 15th and October 6th (17.5 m a.s.l.) are shown in Figure 3 together with the mixing ratios of the trace gases ozone, ethane, ethene, acetone, methanol and DMS. During this period the air temperature ranged from 25.6 °C (6:00 UTC) to 28.3 °C (14:00 UTC) with an average diurnal variation of 0.6 °C. The wind direction was north-easterly (30 to 60 °), except for a period between September 19th and 20th and again on September 21st when northerly air, and lower wind speeds, prevailed. The meteorological conditions observed during the campaign were typical for this site (e.g.

Carpenter, et al. 2010, Fomba, et al. 2014). The concentrations of the different trace gases will be more thoroughly discussed in section 5.3.

4.1.3 Measured and modelled marine boundary layer (MBL) height

The characterization of the MBL is important for the interpretation of both the ground-based as well as the vertically-resolved measurements, because the MBL mixing state allows to elucidate the possible connections between ground-based processes (e.g. aerosol formation) and the higher (e.g. mountain and cloud level) altitudes. The Cape Verdes typically exhibit a strong inversion layer with a sharp increase in the potential temperature and a sharp decrease of the humidity (Carpenter, et al. 2010).

The vertical measurements of meteorological parameters were carried out at CVAO with a 16 m³ Helikite. The measurements demonstrate that a Helikite is a reliable and useful instrument that can be deployed under prevailing wind conditions such as at this measurement site. 19 profiles on ten different days could be obtained and Figure 4 shows an exemplary profile, from September 17th. During the campaign, the wind speed varied between 2 and 14 m s⁻¹ and the MBL height was found to be between about 600 and 1100 m (compare to Fig. 5). Based on the measured vertical profiles, the MBL was found to be often well mixed. However, there are indications for a decoupled boundary layer in a few cases that will be further analysed.

As it was not possible to obtain information of the MBL height for the entire campaign from online measurements, the MBL height was also simulated using the Bulk-Richardson number. The simulations showed that the MBL height was situated where the Bulk-Richardson number exceeded the critical value 0.25. Figure 5 shows, that the simulated MBL height was always lower compared to the measured one during the campaign and also compared to previous measurements reported in the literature. Based on long-term measurements, Carpenter, et al. (2010) observed an MBL height of 713 ± 213 m at the Cape Verdes. In the present study a simulated MBL height of 452 ± 184 m was found, however covering solely a period over one month. The differences might be caused by the grid structure of the applied model (more details in the SI). The vertical resolution of 100 to 200 m might lead to a misplacement of the exact position of the MBL-height. Moreover, the model calculations were constructed to identify the lowest inversion layer. Therefore, the modelled MBL height might represent a low, weak internal layer within the MBL and not the actual MBL. These issues will be analysed in further studies.

4.1.4 Cloud conditions

The Cape Verde Islands are dominated by a marine tropical climate and as mentioned above, marine air is constantly supplied from a north-easterly direction which also transports marine boundary-layer clouds towards the islands. Average wind profiles derived from the European Center for Medium-Range Weather Forecasts (ECWMF) model simulations are shown in Figure 6a. On the basis of the wind profiles, different cloud scenes have been selected and quantified (Derrien and Le Gleau 2005) using geostationary Meteosat SEVIRI data with a spatial resolution of 3 km (Schmetz, et al. 2002) and are shown in Figure 6b – f. The island Sao Vicente is located in the middle of each picture. The first scene at 10:00 UTC on September 19th was characterized by low wind speeds throughout the atmospheric column (Fig. 6b). In this calm situation, a compact patch of low-level clouds was located north-west of the Cape Verde Islands. The cloud field was rather spatially homogeneous, i.e. marine stratocumulus, which transitioned to more broken cumulus clouds towards the island. South-eastwards of the islands, high-level ice clouds dominated and possibly mask lower-level clouds. For the second cloud scene at 10:00 UTC on September 22nd (Fig. 6c), wind speed was higher with more than 12 m s⁻¹ in the boundary layer. Similarly, coverage of low- to very low-level clouds was rather high in the region around Cape Verde Islands. A compact stratocumulus cloud field approached the islands from north-easterly direction. The clouds that had formed over the ocean dissolved when the flow traverses the islands. Pronounced lee effects appeared downstream of the islands. Cloud scene three at 10:00 UTC on September 27th was again during a calm phase with wind speed of a few m s⁻¹ only (Fig. 6d). The scene was dominated by fractional clouds (with a significant part of the spatial variability close to or below the sensor resolution). These clouds formed locally and grew. Advection of clouds towards islands was limited. The last two cloud scenes (at 10:00 UTC on October 1st in Fig. 6e and at 10:00 UTC on October 11th in Fig. 6f) were shaped by higher boundary-layer winds and changing wind directions in higher atmospheric levels. The scene in Fig. 6e shows a complex mixture of low-level cloud fields and higher-level cirrus patches. The scene in Fig. 6f was again dominated by low- to very low-level clouds. The eastern part of the islands was embedded in a rather homogeneous stratocumulus field. A transition of the spatial structure of the cloud field happened in the centre of the domain with more cumuliform clouds and cloud clumps west to the Cape Verde Island. Overall, the majority of low-level clouds over the islands were formed over the ocean and ocean-derived aerosol particles, e.g. sea salt and marine biogenic compounds, might be expected to have some influence on cloud formation. Infrequent instances of locally formed clouds influenced by the orography of the islands could be also identified in the satellite data.

The different cloud scenes reflect typical situations observed in conditions with either weaker or stronger winds. The average in-cloud time of an air parcel might depend on cloud type and cloud cover that in turn impacts in-cloud chemical processes (e.g. Lelieveld and Crutzen, 1991), such as the formation of methane-sulfonic acid and other organic acids (Hoffmann et al., 2016 und Chen et al., 2018). Future studies will relate the chemical composition of the aerosol particles and cloud water to the cloud scenes and their respective oxidation capacity. However, the rather coarse horizontal resolution of the satellite sensor and the missing information about time-resolved vertical profiles of thermodynamics and cloud condensate limits a further detailed characterization of these low-level cloud fields and their formation processes. A synergistic combination with ground-based in-situ and remote sensing measurements would be highly beneficial for future

investigations to elucidate how cloud chemistry might be different for the varying cloud scenes depending on horizontal cloud patterns and vertical cloud structures.

4.2 Biological seawater conditions

4.2.1 Pigment and bacteria concentration in seawater

To characterize the biological conditions at CVAO, a variety of pigments including chlorophyll-*a* (chl-*a*) were measured in the samples of Cape Verdean bulk water (data in Table S4 and illustrated in section 5.4.1). Chl-*a* is the most prominently used tracer for biomass in seawater; however information of phytoplankton composition can only be determined by also determining marker pigments. Therefore, each time when a water sample was taken, also several liters of bulk water were collected for pigment analysis (more details in the SI). Chl-concentrations varied between $0.11 \mu\text{g L}^{-1}$ and $0.6 \mu\text{g L}^{-1}$, and are more thoroughly discussed together with the pigment composition in section 5.4.1. Moreover, as other but phytoplankton organisms can contribute to the OM pool, bacterial abundance was analysed in the SML and bulk water samples and these data are reported in section 5.7.3.

~~Phytoplankton biomass expressed in chl-*a* was very low with $0.11 \mu\text{g L}^{-1}$ at the beginning. Throughout the campaign two slight increases of biomass occurred, but were always followed by a biomass depression. The biomass increase occurred towards the end of the study, where pre-bloom conditions were reached with values up to $0.6 \mu\text{g L}^{-1}$. These are above the typical chl-*a* concentration in this area. In contrast, the abundance of chlorophyll degradation products as phaeophorbide *a* and phaeophytin *a* decreased over time. The low concentrations of the chlorophyll degradation products suggested that only moderate grazing took place and the pigment-containing organisms were fresh and in a healthy state. The most prominent pigment throughout the campaign was zeaxanthin, suggesting cyanobacteria being the dominant group in this region. This is in a good agreement with the general low biomass in the waters of the Cape Verde region and in line with previous studies, reporting the dominance of cyanobacteria during the spring and summer seasons (Franklin, et al. 2009; Hepach, et al. 2014; Zindler, et al. 2012). However, once the biomass increased, cyanobacteria were repressed by diatoms as indicated by the relative increase of fucoxanthin. The prymnesiophyte and haptophyte marker 19-hexanoyloxyfucoxanthin and the pelagophyte and haptophytes marker 19-butanoyloxyfucoxanthin were present and also increased when cyanobacteria decreased. In contrast, dinoflagellates and chlorophytes were background communities as indicated by their respective markers peridinin and chlorophyll b. Still, chlorophytes were much more abundant than dinoflagellates. In summary, the pigment composition indicated the presence of cyanobacteria, haptophytes and diatoms with a change in dominating taxa (from cyanobacteria to diatoms). The increasing concentration of chl-*a* and fucoxanthin implied that a bloom started to develop within the campaign dominated by diatoms. The increasing concentrations could also be related to changing water masses, however, since the oceanographic setting was relatively stable, the increasing chl-*a* concentrations suggest that a local bloom had developed, that might be related to the low but permanent presence of atmospheric dust input, which needs further verification. In the course of further data analysis of the campaign, the phytoplankton groups will be related to the abundance of e.g. DMS (produced by haptophytes) or isoprene that~~

~~has been reported to be produced by diatoms or cyanobacteria (Bonsang, et al. 2010), as well as to other VOCs.~~

4.2.2 Wave glider fluorescence measurements

Roughly at the same time as the MarParCloud field campaign took place, an unmanned surface vehicle (SV2 Wave Glider, Liquid Robotics Inc.) equipped with a biogeochemical sensor package, a conductivity-temperature-depth sensor (CTD) and a weather station was operated in the vicinity of the sampling location. The Wave Glider carried out continuous measurements of surface water properties (water intake depth: 0.3 m) along a route near the coast (Fig. 7a), and on October 5th it was sent on a transect from close to the sampling location towards the open ocean in order to measure lateral gradients in oceanographic surface conditions. The glider measurements delivered information on the spatial resolution of several parameters. Fluorescence measurements, which can be seen as a proxy of chl-*a* concentration in surface waters and hence of biological production, indicated some enhanced production leeward of the islands and also at one location upwind of the island of Santa Luzia next to São Vicente. In the vicinity of the MarParCloud sampling site the glider observed a slight enhancement in fluorescence when compared to open-ocean waters. This is in agreement with the measured pigment concentration. The overall pattern of slightly enhanced biological activity was also confirmed by [the MODIS-Terra](#) satellite fluorescence measurements (Fig. 7b). However, both in situ glider and sample data as well as remote sensing data did not show any particular strong coastal bloom events and thus indicate that the MarParCloud sampling site well represented the open-ocean regime during the sampling period.

5 Measurements and selected results

5.1 Vertical resolution measurements

5.1.1 Physical aerosol characterization

Based on aerosol particles measured during the campaign, air masses could be classified into different types, depending on differences in PNSDs. Marine type and dust type air masses could be clearly distinguished, even if the measured dust concentrations were only low to medium, according to the annual mean at the CVAO (Fomba, et al. 2013, 2014). The median of PNSDs during marine conditions is illustrated in Figure 8 and showed three modes, i.e., Aitken, accumulation and coarse mode. There was a minimum between the Aitken- and accumulation-mode of PNSDs (Hoppel minimum; see (Hoppel, et al. 1986) at roughly 70 nm. PNSDs measured during marine type air masses featured the lowest Aitken, accumulation and coarse mode particle number concentrations, with median values of 189, 143 and 7 cm⁻³, respectively. The PNSDs present during times with dust influences featured a single mode in the sub-micron size range (Fig. 8), and no visible Hoppel minimum was found. The dust type air masses

featured the highest total particle number concentration (994 cm^{-3}) and a median coarse-mode particle number concentration of 44 cm^{-3} .

N_{CCN} at different supersaturations were compared during dust and marine periods, as shown in Figure 9. During dust periods, the aerosol particles show a great enhancement in Aitken, accumulation and coarse mode number concentrations, such that overall N_{CCN} increases distinctly. N_{CCN} at a supersaturation of 0.30% (proxy for the supersaturation encountered in clouds present during the campaign) during the strongest observed dust periods is about 2.5 times higher than that during marine periods. As suggested by Modini, et al. (2015), Wex, et al. (2016) and Quinn, et al (2017), the coarse mode aerosol particles can be attributed to sea spray aerosol (SSA) in a marine environment. In these studies, the fraction of sea spray aerosol was determined based on three-modal fits from which the particle number concentrations in the different modes were determined. A similar analysis was done for this study. During marine periods, SSA accounted for about 3.7% of CCN number concentrations at 0.30% supersaturation and for 1.1% to 4.4% of N_{total} (total particle number concentration). The hygroscopicity parameter κ averaged 0.28, suggesting the presence of OM in the particles (see Gong, et al., 2020a). Particle sizes for which κ was determined (i.e., the critical diameters determined during CCN analysis) were roughly 50 to 130 nm. The low value determined for κ is in line with the fact that sodium chloride from sea salt was below detection limit in the size segregated chemical analysis for particles in this size range (Figure 11, while insoluble EC and WSOM made up 30% of the main constituents at CVAO on average.

~~The particle number concentrations for the coarse mode of the aerosol particles that is attributed to sea spray aerosol (SSA) accounted for about 3.7% of $N_{\text{CCN}, 0.30\%}$ (CCN number concentration at 0.30% supersaturation) and for 1.1% to 4.4% of N_{total} (total particle number concentration).~~
A thorough statistical analysis of N_{CCN} and particle hygroscopicity concerning different aerosol types is reported in Gong, et al. (2020a). Figure 9a-10a shows the median of marine type PNSDs for cloud free conditions and cloud events at CVAO and Mt. Verde. Figure 9b-10b shows the scatter plot of N_{CCN} at CVAO versus those on Mt. Verde. For cloud free conditions, all data points are close to the 1:1 line, indicating N_{CCN} being similar at the CVAO and Mt. Verde. However, during cloud events, larger particles, mainly accumulation- and coarse-mode particles, were activated to cloud droplet and were, consequently, removed by the inlet. Therefore, during these times, N_{CCN} at the CVAO was larger than the respective values measured on Mt. Verde. Altogether, these measurements suggested that, for cloud free conditions, the aerosol particles measured at ground level (CVAO) represent the aerosol particles at the cloud level (Mt. Verde).

5.1.2 Chemical composition of aerosol particles and cloud water

Between October 2nd and 9th, size-resolved aerosol particles at the CVAO and the Mt. Verde were collected simultaneously. The relative contribution of their main chemical constituents (inorganic ions, water-soluble organic matter (WSOM), and elemental carbon) at both sites is shown in Figure 10. Sulfate, ammonium, and WSOM dominated the sub-micron particles and the chemical composition aligned well with the κ value from the hygroscopicity measurements (Gong, et al. 2020a). The super-micron particles were mainly composed of sodium and chloride at both stations. These findings agreed well with previous studies at the CVAO (Fomba, et al.

2014; van Pinxteren, et al. 2017). From the chemical composition no indications for anthropogenic influences was found as concentrations of elemental carbon and submicron potassium were low (see Tab. S5). However, according to the dust concentrations (Table 2) and the air mass origins (Fig. S1), as well as the PNSD (Gong et al. 2020a), the air masses during this period experienced low dust influences, that was however not visible from the main chemical constituents studied here. These findings warrant more detailed chemical investigations (like size-resolved dust measurements), a distinction between mass-based and number-based analysis as well as detailed source investigations that are currently ongoing. The absolute concentrations of the aerosol constituents were lower at the Mt. Verde compared to the CVAO site (Table S5); they were reduced by factor of seven (super-micron particle) and by a factor of four (sub-micron particles). This decrease in the aerosol mass concentrations and the differences in chemical composition between the ground-based aerosol particles and the ones at Mt. Verde, could be due to cloud effects as described in the previous section. Different types of clouds consistently formed and disappeared during the sampling period of the aerosol particles at the Mt. Verde (more details about the frequency of the cloud events are available in the SI and in Gong, et al., ~~(202019a)~~ and potentially affected the aerosol chemical composition. These effects will be more thoroughly examined in further analysis.

A first insight in the cloud water composition of a connected cloud water sampling event from October 5th till October 6th is presented in Figure 142. Sea salt, sulfate and nitrate compounds dominated the chemical composition making up more than 90% of the mass of the investigated chemical constituents. These compounds were also observed in the coarse fraction of the aerosol particles, indicating-suggesting that the coarse mode particles served as efficient CCN and were efficiently transferred to the cloud water. To emphasize, these chemical analyses are based on mass, but the control of the cloud droplet number concentration comes from CCN number concentrations, including all particles with sizes of roughly above 100 nm. As larger particles contribute more to the total mass, chemical bulk measurements give no information about a direct influence of sea spray particles on cloud droplet concentrations, but it can show that the chemical composition is consistent with an (expected) oceanic influence on cloud water.

No strong variations were found for the main cloud water constituents over the here reported sampling period. However, the WSOM contributed with maximal 10% to the cloud water composition and with higher contributions in the beginning and at the end of the sampling event, which warrants further analysis. The measured pH values of the cloud water samples ranged between 6.3 and 6.6 and agreed with previous literature data for marine clouds (Herrmann, et al. 2015). In summary, cloud water chemical composition seemed to be ~~controlled-dominated~~ by coarse mode aerosol particle composition, and the presence of inorganic marine tracers (sodium, methane sulfonic acid) shows that material from the ocean is transported to the atmosphere where it can become immersed in cloud droplets. More detailed investigations on the chemical composition, including comparison of constituents from submicron aerosol particles and the SML with the cloud water composition are planned.

5.2 Lipid biomarkers in aerosol particles

Lipids from terrestrial sources such as plant waxes, soils and biomass burning have frequently been observed in the remote marine troposphere (Kawamura, et al. 2003; Simoneit, et al. 1977)

and are common in marine deep-sea sediments. Within MarParCloud, marine-derived lipids were characterized in aerosol particles using lipid biomarkers in conjunction with compound specific stable carbon isotopes. Bulk aerosol filters sampled at the CVAO and PM₁₀ filter sampled at the Mt. Verde (not reported here) were extracted and the lipids were separated into functional groups for molecular and compound specific carbon isotope analysis. The content of identifiable lipids was highly variable and ranged from 4 to 140 ng m⁻³. These concentrations are in the typical range for marine aerosol particles (Mochida, et al. 2002; Simoneit, et al. 2004) but somewhat lower than previously reported for the tropical North East Atlantic (Marty & Saliot, 1979) and 1 to 2 orders of magnitude lower than reported from urban and terrestrial rural sites (Simoneit, 2004). It mainly comprised the homologue series of n-alkanoic acids, n-alkanols and n-alkanes. Among these the c16:0 acid and the c18:0 acids were by far the dominant compounds, each contributing 20 to 40% to the total observed lipids. This result aligns well with the findings of Cochran, et al. (2016) from sea spray tank studies that connected the transfer of lipid-like compounds to their physicochemical properties such as solubility and surface activity. Among the terpenoids, dehydroabietic acid, 7-oxo-dehydroabietic acid and friedelin were in some samples present in remarkable amounts. Other terpenoid biomarker in particular phytosterols were rarely detectable. The total identifiable lipid content was inversely related to dust concentration, as shown exemplary for the fatty acids (Fig. 123) with generally higher lipid concentrations in primary marine air masses. This is consistent with previous studies reporting low lipid yields in Saharan dust samples and higher yields in dust from the more vegetated Savannas and dry tropics (Simoneit, et al. 1977). First measurements of typical stable carbon isotope ratios of the lipid fractions were (-28.1 ± 2.5) ‰ for the fatty acids and (-27.7 ± 0.7) ‰ for the n-alkanes suggesting a mixture of terrestrial c3 and c4, as well as marine sources. In a separate contribution the lipid fraction of the aerosol particles in conjunction with its typical stable carbon isotope ratios will be further resolved.

5.3 Trace gas measurements:

~~5.3.1~~ Dimethyl sulphide, ozone, ~~and~~ (oxygenated) volatile organic compounds and nitrous acid

Trace gases such as dimethyl sulfide (DMS), volatile organic compounds (VOCs) and oxygenated (O)VOCs have been measured during the campaign and the results are presented together with the meteorological data in Figure 3. The atmospheric mixing ratios of DMS during this period ranged between 68 ppt and 460 ppt with a mean of 132 ± 57 ppt (1σ). These levels were higher than the annual average mixing ratio for 2015 of 57 ± 56 ppt, however this may be due to seasonably high and variable DMS levels observed during summer and autumn at Cape Verde (observed mean mixing ratios were 86 ppt and 107 ppt in September and October 2015). High DMS concentrations on September 19th – 20th occurred when air originated predominantly from the Mauritanian upwelling region ~~and on September 26th and 27th when the footprint was influenced by southern hemisphere~~ (Figure SI1) and on September 26th and 27th. These elevated concentrations will be linked to the phytoplankton composition reported in section ~~4.2.15.4.1.~~ to elucidate associations for example between DMS and coccolith (individual plates of calcium carbonate formed by *coccolithophores* phytoplankton) as

observed by Marandino, et al. (2008). Ethene showed similar variability to DMS, with coincident peaks (> 300 ppt DMS and > 40 ppt ethene) on September 20th, 26th and 27th, consistent with an oceanic source for ethene. Ethene can be emitted from phytoplankton (e.g. McKay, et al. 1996) and therefore it is possible that it originated from the same biologically active regions as DMS. In the North Atlantic atmosphere, alkenes such as ethene emitted locally have been shown to exhibit diurnal behaviour with a maximum at solar noon, suggesting photochemical production in seawater (Lewis, et al. 2005). There was only weak evidence of diurnal behaviour at Cape Verde (data not shown), possibly because of the very short atmospheric lifetime of ethene (8 hours assuming $[OH] = 4 \times 10^6$ molecules cm^{-3} , Vaughan, et al. 2012) in this tropical environment, which would mask photochemical production. Mean acetone and methanol mixing ratios were 782 ppt (566 ppt – 1034 ppt) and 664 ppt (551 ppt – 780 ppt), respectively. These are similar to previous measurements at Cape Verde and in the remote Atlantic at this time of year (Lewis, et al. 2005; Read, et al. 2012). Methanol and acetone showed similar broad-scale features, indicating common sources. Highest monthly methanol and acetone concentrations have often been observed in September at Cape Verde, likely as a result of increased biogenic emissions from vegetation or plant matter decay in the Sahel region of Africa (Read, et al. 2012). In addition to biogenic sources, (O)VOCs are anthropogenically produced from fossil fuels and solvent usage in addition to having a secondary source from the oxidation of precursors such as methane. Carpenter, et al. (2010) showed that air masses originating from North America (determined via 10-day back trajectories) could impact (O)VOCs at the CVAO.

The average ozone mixing ratio during the campaign was 28.7 ppb (19.4 ppb – 37.8 ppb). Lower ozone concentrations on September 27th to 28th were associated with influence from southern hemispheric air. Ozone showed daily photochemical loss, as expected in these very low-NO_x conditions, on most days with an average daily (from 9:00 UTC to 17:00 UTC) loss of 4 ppbV. It was previously shown that the photochemical loss of O₃ at Cape Verde and over the remote ocean is attributable to halogen oxides (29% at Cape Verde) as well as ozone photolysis (54%) (e.g. Read, et al. 2008).

Finally, a series of continuous measurements of nitrous acid (HONO) has been conducted, aiming at evaluating the possible contribution of marine surfaces to the production of HONO. The measurements indicated that HONO concentrations exhibited diurnal variations peaking at noontime. The concentrations during daytime (08:00 to 17:00, local time) and night-time (17:30 to 07:00 local time) periods were around 20 ppt and 5 ppt on average, respectively. The fact that the observed data showed higher values during the day compared to the night-time was quite surprising since HONO is expected to be photolyzed during the daytime. If confirmed, the measurements conducted here may indicate that there is an important HONO source in the area of interest. Altogether, for the trace gases, a variety of conditions were observed in this three-week period with influence from ocean-atmosphere exchange and also potential impacts of long-range transport.

5.3.2 Nitrous acid

Nitrous acid (HONO) plays a significant role in the atmospheric chemistry as an important source of hydroxyl radical ($\cdot\text{OH}$). It is well recognized that significant uncertainties remain on its emission sources as well as on its in-situ tropospheric formation processes. During the campaign, a series of continuous measurements of HONO has been conducted, aiming at evaluating the possible contribution of marine surfaces to the production of HONO. The measurements indicated that HONO concentrations exhibited diurnal variations peaking at noontime. The concentrations during daytime (08:00 to 17:00, local time) and nighttime (17:30 to 07:00 local time) periods were around 20 ppt and 5 ppt on average, respectively. The fact that the observed data showed higher values during the day compared to the nighttime was quite surprising since HONO is expected to be photolyzed during the daytime. If confirmed, the measurements conducted here may indicate that there is an important HONO source in the area of interest. In a separate paper, the data obtained will be described and discussed and tentative explanation of the observed phenomena will be developed.

5.4 Organic Matter and related compounds in seawater

5.4.1. Dissolved organic carbon and pigments

Dissolved organic carbon (DOC) comprise a complex mixtures of different compound groups and is diverse in its composition. For a first overview, DOC as a sum parameter was analyzed in all SML and bulk water samples (data in Table S4). DOC concentration varied between 1.8 and 3.2 mg L⁻¹ in the SML and 0.9 and 2.8 mg L⁻¹ in the bulk water and were in general agreement with previous studies at this location (e.g. van Pinxteren, et al. 2017). A slight enrichment in the SML with an enrichment factor (EF) = 1.66 (± 0.65) was found, i.e. SML concentrations contain roughly 70% more DOC than the corresponding bulk water. The concentrations of DOC in the bulk water together with the temporal evolution of biological indicators (pigments and the total bacterial cell numbers) and atmospheric dust concentrations are presented in Figure 134.

Phytoplankton biomass expressed in chl-*a* was very low with 0.11 $\mu\text{g L}^{-1}$ at the beginning of the campaign. Throughout the campaign two slight increases of biomass occurred, but were always followed by a biomass depression. The biomass increase occurred towards the end of the study, where pre-bloom conditions were reached with values up to 0.6 $\mu\text{g L}^{-1}$. These are above the typical chl-*a* concentration in this area. In contrast, the abundance of chlorophyll degradation products as phaeophorbide *a* and phaeophytin *a* decreased over time. The low concentrations of the chlorophyll degradation products suggested that only moderate grazing took place and the pigment-containing organisms were fresh and in a healthy state. The most prominent pigment throughout the campaign was zeaxanthin, suggesting cyanobacteria being the dominant group in this region. This is in a good agreement with the general low biomass in the waters of the Cape Verde region and in line with previous studies, reporting the dominance of cyanobacteria during the spring and summer seasons (Franklin, et al. 2009; Hepach, et al. 2014; Zindler, et al. 2012). However, once the biomass increased, cyanobacteria were repressed

by *diatoms* as indicated by the relative increase of fucoxanthin. The *prymnesiophyte* and *haptophyte* marker 19-hexanoyloxyfucoxanthin and the *pelagophyte* and *haptophytes* marker 19-butanoyloxyfucoxanthin were present and also increased when *cyanobacteria* decreased. In contrast, *dinoflagellates* and *chlorophytes* were background communities as indicated by their respective markers peridinin and chlorophyll b. Still, *chlorophytes* were much more abundant than *dinoflagellates*. In summary, the pigment composition indicated the presence of *cyanobacteria*, *haptophytes* and *diatoms* with a change in dominating taxa (from *cyanobacteria* to *diatoms*). The increasing concentration of chl-*a* and fucoxanthin implied that a bloom started to develop within the campaign dominated by *diatoms*. The increasing concentrations could also be related to changing water masses, however, since the oceanographic setting was relatively stable, the increasing chl-*a* concentrations suggest that a local bloom had developed, that might be related to the low but permanent presence of atmospheric dust input, which needs further verification. In the course of further data analysis of the campaign, the phytoplankton groups will be related to the abundance of e.g. DMS (produced by *haptophytes*) or isoprene that has been reported to be produced by *diatoms* or *cyanobacteria* (Bonsang, et al. 2010), as well as to other VOCs. First analyses show that the DOC concentrations were not directly linked to the increasing chl-*a* concentrations, however their relation to single pigments, to the microbial abundance, to the background dust concentrations and finally to wind speed and solar radiation will be further resolved to elucidate potential biological and meteorological controls on the concentration and enrichment of DOC.

5.4.2 DOC concentrations: Manual glass plate vs. MarParCat sampling

For several dates, both SML sampling devices (glass plate and catamaran) were applied in parallel to compare the efficiency of different sampling approaches: manual glass plate and the catamaran sampling (Fig. 145). As mentioned above both techniques used the same principle, i.e. the collection of the SML on a glass plate and its removal with a Teflon wiper. The deviation between both techniques concerning DOC measurements was below 25% in 17 out of 26 comparisons and therefore within the range of variability of these measurements. However, in roughly 30% of all cases the concentration differences between manual glass plate and catamaran were larger than 25%. The discrepancy for the bulk water results could be related to the slightly different bulk water sampling depths using the MarParCat bulk water sampling system (70 cm) and the manual sampling with the telescopic rods (100 cm). Although the upper meters of the ocean are assumed to be well mixed, recent studies indicate that small scale variabilities can be observed already within the first 100 cm of the ocean (Robinson, et al. 2019a).

The variations within the SML measurements could be due to the patchiness of the SML that has been tackled in previous studies (e.g. Mustafa, et al. 2017, 2018). Small-scale patchiness was recently reported as a common feature of the SML. The concentrations and compositions probably undergo more rapid changes due to a high physical and biological fluctuations. Mustafa, et al. (2017) have recently shown that the enrichment of fluorescence dissolved matter (a part of DOC) showed short time-scale variability, changing by 6% within ten-minute intervals. The processes leading to the enrichment of OM in the SML are probably much more complex than previously assumed (Mustafa, et al. 2018). In addition, the changes in DOC

concentrations between the glass plate and the catamaran could result from the small variations of the sampling location as the catamaran was typically 15 to 30 m apart from the boat where the manual glass plate sampling was carried out.

Given the high complex matrix of seawater and especially the SML, the two devices applied were in quite good agreement considering DOC measurements. However, this is not necessarily the case for the single parameters like specific organic compounds and INP concentrations. Especially low concentrated constituents might be more affected by small changes in the sampling procedure and this remains to be evaluated for the various compound classes.

5.4.32. Surfactants and lipids in seawater

Due to their physicochemical properties, surfactants (SAS) are enriched in the SML relative to the bulk water and form surface films (Frka, et al. 2009; Frka, et al. 2012; Wurl, et al. 2009). During the present campaign, the SAS in the dissolved fraction of the SML samples ranged from 0.037 to 0.125 mg TX-100 eqL⁻¹ (Triton-X-100 equivalents) with a mean of 0.073 ± 0.031 mg TX-100 eqL⁻¹ (n = 7). For bulk water, the dissolved SAS ranged from 0.020 to 0.068 mg TX-100 eqL⁻¹ (mean 0.051 ± 0.019 mg TX-100 eqL⁻¹, n = 12). The SAS enrichment showed EFs from 1.01 to 3.12 (mean EF = 1.76 ± 0.74) (Fig. 156) and was slightly higher than that for the DOC (mean EF = 1.66 ± 0.65) indicating some higher surfactant activity of the overall DOM in the SML in respect to the bulk DOM. An accumulation of the total dissolved lipids (DL) in the SML was observed as well (mean EF = 1.27 ± 0.12). Significant correlation was observed between the SAS and DL concentrations in the SML ($r = 0.845$, n = 7, $p < 0.05$) while no correlation was detected for the bulk water samples. Total DL concentrations ranged from 82.7 to 148 $\mu\text{g L}^{-1}$ (mean $108 \pm 20.6 \mu\text{g L}^{-1}$, n = 8) and from 66.5 to 156 $\mu\text{g L}^{-1}$ (mean $96.9 \pm 21.7 \mu\text{g L}^{-1}$, n = 17) in the SML and the bulk water, respectively. In comparison to the bulk water, the SML samples were enriched with lipid degradation products e.g. free fatty acids and long chain alcohols (DegLip; mean EF = 1.50 ± 0.32), particularly free fatty acids and long-chain alcohols (Fig. 156), pointing to their accumulation from the bulk and/or enhanced OM degradation within the SML. DegLip are strong surface-active compounds (known as dry surfactants), which play an important role in surface film establishment (Garrett 1965). The overall surfactant activity of the SML is the result of the competitive adsorption of highly surface-active lipids and other less surface-active macromolecular compounds (polysaccharides, proteins, humic material) (Ćosović and Vojvodić 1998) dominantly present in seawater. The presence of even low amounts of lipids results in their significant contribution to the overall surface-active character of the SML complex organic mixture (Frka, et al. 2012). The observed biotic and/or abiotic lipid degradation processes within the SML will be further resolved by combining surfactant and lipid results with detailed pigment characterisation and microbial measurements. The same OM classes of the ambient aerosol particles will be investigated and compared with the seawater results. This will help to tackle the questions to what extent the seawater exhibits a source of OM on aerosol particles and which important aerosol precursors are formed or converted in surface films.

5.5 Seawater Untargeted Metabolomics

For a further OM characterization of SML and bulk seawater an ambient MS-based metabolomics method using direct analysis in real time quadrupole time-of-flight mass spectrometry (DART-QTOF-MS) coupled to multivariate statistical analysis was designed (Zabalegui, et al. 2019). A strength of a DART ionization source is that it is less affected by high salt levels than an electrospray ionization source (Kaylor, et al. 2014), allowing the analysis of seawater samples without observing salt deposition at the mass spectrometer inlet, or having additional limitations such as low ionization efficiency due to ion suppression (Tang, et al. 2004). Based on these advantages, paired SML/bulk water samples were analyzed without the need of desalinization by means of a transmission mode (TM) DART-QTOF-MS-based analytical method that was optimized to detect lipophilic compounds (Zabalegui, et al. 2019). An untargeted metabolomics approach, addressed as seaomics, was implemented for sample analysis. SML samples were successfully discriminated from ULW samples based on a panel of ionic species extracted using chemometric tools. The coupling of the DART ion source to high-resolution instrumentation allowed generating elemental formulae for unknown species and tandem MS capability contributed to the identification process. Tentative identification of discriminant species and the analysis of relative compound abundance changes among sample classes (SML and bulk water) suggested that fatty alcohols, halogenated compounds, and oxygenated boron-containing organic compounds may be involved in water-air transfer processes and in photochemical reactions at the water-air interface of the ocean (Zabalegui, et al., 2019). These identifications (e.g. fatty alcohols) agree well with the abundance of lipids in the respective samples. In this context, TM-DART-HR-MS appears to be an attractive strategy to investigate the seawater OM composition without requiring a desalinization step.

5.6 Ocean surface mercury associated with organic matter

Several ~~trace metals~~elements are known to accumulate in the SML. In the case of Hg, the air-sea exchange plays an important role in its global biogeochemical cycle and hence processing of Hg in the SML is of particular interest. Once deposited from the atmosphere to the ocean surface via dry and wet deposition, the divalent mercury (Hg^{II}) can be transported to the deeper ocean by absorbing on sinking OM particles, followed by methylation. On the other hand, Hg^{II} complexed by DOM in the ocean surface can be photo-reduced to Hg^0 , which evades into the gas phase. In both processes, OM, dissolved or particulate, is the dominant factor influencing the complexation and adsorption of Hg. To explore the Hg behaviour with OM, the concentrations of total and dissolved Hg as well as the methylmercury (MeHg) were determined in the SML and in the bulk water using the US EPA method 1631 and 1630, as described in Li, et al. (2018). Figure 167 shows the concentrations of Hg and MeHg associated with DOC and POC in the SML and bulk water. The total Hg concentrations were 3.6 and 4.6 ng L^{-1} in the SML but 3.1 and 1.3 ng L^{-1} in the bulk water on September 26th and 27th, respectively, which were significantly enriched compared to data reported for the deep North Atlantic ($0.18 \pm 0.06 \text{ ng L}^{-1}$) (Bowman, et al. 2015). Atmospheric deposition and more OM adsorbing Hg are supposed to result in the high total Hg at ocean surface. The dissolved Hg concentrations were enriched by 1.7 and 2.7 times in the SML relative to bulk water, consistent with the enrichments of DOC by a factor of 1.4 and 1.9 on September 26th and 27th, respectively. Particulate Hg in

the SML accounted for only 6% of the total Hg concentration on September 26th but 55% on September 27th, in contrast to their similar fractions of ~35% in the bulk water on both days. According to the back trajectories (Figure SII) stronger contribution of African continental sources (e.g., dust) was observed on September 27th that might be linked to the higher concentrations of particulate Hg in the SML on this day. The water-particle partition coefficients ($\log K_d$) for Hg in the SML (6.8 L kg⁻¹) and bulk water (7.0 L kg⁻¹) were similar regarding POC as the sorbent, but one unit higher than the reported $\log K_d$ values in seawater (4.9–6.1 L kg⁻¹) (Batrakova, et al. 2014). MeHg made up lower proportions of the total Hg concentrations in the SML (2.0%) than bulk water (3.4% and 4.2%), probably due to photo-degradation or evaporation of MeHg at the surface water (Blum, et al. 2013). From the first results, it seems that the SML is the major compartment where Hg associated with OM is enriched, while MeHg is more likely concentrated in deeper water. The limited data underlines the importance of SML in Hg enrichment dependent on OM, which needs further studies to understand the air-sea exchange of Hg.

5.7 Ocean-atmosphere transfer of organic matter and related compounds

5.7.1 Dissolved organic matter classes

To investigate the complexity of dissolved organic matter (DOM) compound groups, liquid chromatography, organic carbon detection, organic nitrogen detection, UV absorbance detection (LC-OCD-OND-UVD; Huber, et al. (2011), more details in the SI) was applied to identify five different DOM classes. These classes include (i) biopolymers (likely hydrophobic, high molecular weight $\gg 20,000$ g mol⁻¹, largely non-UV absorbing extracellular polymers); (ii) “humic substances” (higher molecular weight ~ 1000 g mol⁻¹, UV absorbing); (iii) “building blocks” (lower molecular weight 300-500 g mol⁻¹, UV absorbing humics); (iv) low molecular weight “neutrals” (350 g mol⁻¹, hydro- or amphiphilic, non-UV absorbing); and (v) low molecular weight acids (350 g mol⁻¹). These measurements were performed from a first set of samples from all the ambient marine compartments. That comprised three SML samples and the respective bulk water, three aerosol particle filter samples (PM₁₀) from the CVAO and two from the Mt. Verde and finally four cloud water samples collected during the campaign. The SML-EFs for DOM concentrations were derived from the sum of the individual compound groups (in $\mu\text{g L}^{-1}$) and the EFs for DOM-varied from 0.83 to 1.46, which agreed very well to the DOC measurements described in section 5.4.1. A clear compound group that drove this change could not be identified so far. Figure 178 shows the relative composition of the measured DOM groups in the distinct marine compartments as an average of the single measurements (concentrations are listed in Table S6). In the SML and in the bulk water, the low molecular weight neutral (LMWN) compounds generally dominated the overall DOM pool (37 to 51%). Humic-like substances, building blocks, and biopolymeric substances contributed 22 to 32%, 16 to 23%, and 6 to 12%, respectively. Interestingly, low molecular weight acids (LMWA) were predominantly observed in the SML (2 to 8%) with only one bulk water time point showing any traces of LMWA. This finding agreed well with the presence of free amino acids (FAA) in the SML; e.g. the sample with highest LMWA concentration showed highest

FAA concentration (more details in Triesch, et al., 2020~~19~~). Further interconnections between the DOM fractions and single organic markers and groups (e.g. sugars, lipids and surfactants, see section 5.4.3~~2~~) are subject to ongoing work. In contrast, aerosol particles were dominated by building blocks (46 to 66%) and LMWN (34 to 51%) compound groups, with a minor contribution of LMWA (> 6%). Interestingly, higher molecular weight compounds of humic-like substances and biopolymers were not observed. Cloud water samples had a variable contribution of substances in the DOM pool with humic substances and building blocks generally dominating (27 to 63% and 16 to 29%, respectively) and lower contributions of biopolymers (2 to 4%) and LMW acids and neutrals (1 to 20% and 18 to 34%) observed. The first measurements indicate that the composition of the cloud waters is more consistent with the SML and bulk water and different from the aerosol particle's composition. This observation suggests a two-stage process where selective aerolisation mobilises lower molecular weight humics (building blocks) into the aerosol particle phase, which may aggregate in cloud waters to form larger humic substances in cloud waters. These preliminary observations need to be further studied with a larger set of samples and could relate to either different solubilities of the diverse OM groups in water, the interaction between DOM and particulate OM (POM), including TEP formation, as well as indicating the different OM sources and transfer pathways. In addition, the chemical conditions, like pH-value or redox, could preferentially preserve or mobilise DOM fractions within the different types of marine waters. In summary, all investigated compartments showed a dominance of LMW neutrals and building blocks, which suggests a link between the seawater, aerosol particles and cloud water at this location and possible transfer processes. Furthermore, the presence of humic-like substances and biopolymers and partly LMWA in the seawater and cloud water, but not in the aerosol particles, suggests an additional source or formation pathway of these compounds. For a comprehensive picture; however, additional samples need to be analysed and interpreted in future work. It is worth noting that the result presented here are the first for such a diverse set of marine samples and demonstrate the potential usefulness in identifying changes in the flux of DOM between marine compartments.

A more comprehensive set of samples was analysed for FAA on molecular level as important organic nitrogen- containing compounds (Triesch, et al. 2020). The FAA, likely resulting from the ocean, were strongly enriched in the submicron aerosol particles ($EF_{aer (FAA)} 10^2$ - 10^4) and to a lower extent enriched in the supermicron aerosol particles ($EF_{aer (FAA)} 10^1$). The cloud water contained the FAA in significantly higher concentrations compared to their respective seawater concentrations and they were enriched by a factor of $4 \cdot 10^3$ compared to the SML. These high concentrations cannot be currently explained and possible sources such as biogenic formation or enzymatic degradation of proteins, selective enrichment processes or pH dependent chemical reactions are subject to future work. The presence of high concentrations of FAA in submicron aerosol particles and in cloud water together with the presence of inorganic marine tracers (sodium, methane sulfonic acid) point to an influence of oceanic sources on the local clouds (Triesch, et al. 2020).

5.7.2. Transparent exopolymer particles: field and tank measurements

As part of the OM pool, gel particles, such as positive buoyant transparent exopolymer particles (TEP), formed by the aggregation of precursor material released by plankton and bacteria, accumulate at the sea surface. The coastal water in Cape Verde has shown to be oligotrophic with low chl-*a* abundance during the campaign (more details in section 4.2.1). Based on previous work (Wurl, et al. 2011) it is expected that surfactant enrichment, which is closely linked to TEP enrichment, in the SML would be higher in oligotrophic waters but have a lower absolute concentration. This compliments the here achieved findings, which showed low TEP abundance in these nearshore waters; the abundance in the bulk water ranged from 37 to 144 μgXeqL^{-1} (xanthan gum equivalents) and 99 to 337 μgXeqL^{-1} in the SML. However while the SML layer was relatively thin ($\sim 125\ \mu\text{m}$) there was positive enrichment of TEP in the SML with an average EF of 2.0 ± 0.8 (Fig. 189a). The enrichment factor for TEP was furthermore very similar to surfactant enrichment (section 5.4.32).

In addition to the field samples, a tank experiment was run simultaneously using the same source of water (Fig. 198b). Breaking waves were produced via a waterfall system (details in the SI) and samples were collected from the SML and bulk water after a wave simulation time of 3 h. TEP abundance in the tank experiment matched the field samples at the beginning but quickly increased to 1670 μgXeqL^{-1} in the SML with an EF of 13.2 after the first day of bubbling. The enrichment of TEP in the SML during the tank experiment had a cyclical increase and decrease pattern. Interestingly, in the field samples, even on days with moderate wind speeds ($> 5\ \text{m s}^{-1}$) and occasional presence of white caps, TEP abundance or enrichment didn't increase, but it did increase substantially due to the waves in the tank experiment. This suggests that the simulated waves are very effective in enriching TEP in the SML and TEP were more prone to transport or formation by bubbling than by other physical forces, confirming bubble-induced TEP enrichment in recent artificial set-ups (Robinson, et al. 2019b). Besides the detailed investigations of TEP in seawater, first analyses show a clear abundance of TEP in the aerosol particles and in cloud water. Interestingly, a major part of TEP seems to be located in the sub-micron aerosol particles (Fig. 2049). Sub-micron aerosol particles represent the longest living aerosol particle fraction and have a high probability to reach cloud level and to contribute to cloud formation and the occurrence of TEP in cloud water, which strongly underlines a possible vertical transport of these ocean-derived compounds.

5.7.3 Bacterial abundance in distinct marine samples: field and tank measurements

The OM concentration and composition is closely linked with biological and especially microbial processes within the water column. Throughout the sampling period, the temporal variability of bacterial abundance in SML and bulk water was studied (data listed in Tab.SI4). Mean absolute cell numbers were $1.3 \pm 0.2 \times 10^6\ \text{cells mL}^{-1}$ and $1.2 \pm 0.1 \times 10^6\ \text{cells mL}^{-1}$ for SML and bulk water, respectively (Fig. 210a, all data listed in Table S4). While comparable SML data is lacking for this oceanic province, our data is in range with previous reports for surface water of subtropical regions (Zäncker, et al. 2018). A strong day-to-day variability of absolute cell numbers was partly observed (e.g. the decline between September 25th and 26th), but all these changes were found in both, in the SML and bulk water (Fig. 210a). This indicates that the upper water column of the investigated area experienced strong changes, e.g. by inflow of different water masses and/or altered meteorological forcing. As for the absolute abundance,

the enrichment of bacterial cells in the SML was also changing throughout the sampling period, with EFs ranging from 0.88 to 1.21 (Fig. 201b). A detailed investigation of physical factors (e.g. wind speed, solar radiation) driving OM concentration and bacterial abundance in the SML and bulk water will be performed to explain the short-term variability observed. Further ongoing investigations aim to determine the bacterial community composition by 16S sequencing approaches. The resulting comparison of water and aerosol particle samples will help to better understand the specificity of the respective communities and to gain insights into the metabolic potential of abundant bacterial taxa in aerosol particles. During the tank experiment, cell numbers ranged between 0.6 and 2.0×10^6 cells mL⁻¹ (Fig 210c); the only exception being observed on October 3rd, when cell numbers in the SML reached 4.9×10^6 cells mL⁻¹. Both, in the SML and bulk water, bacterial cell numbers decreased during the experiment, which may be attributed to limiting substrate supply in the closed system. Interestingly, SML cell numbers always exceeded those from the bulk water (Fig. 210d), although the SML was permanently disturbed by bursting bubbles throughout the entire experiment. This seems to be in line with the high TEP concentrations observed for the SML in the tank (section 5.7.2). A recent study showed that bubbles are very effective transport vectors for bacteria into the SML, even within minutes after disruption (Robinson, et al. 2019a). The decline of SML bacterial cell numbers (both absolute and relative) during the experiment may be partly caused by permanent bacterial export into the air due to bubble bursting. Although this conclusion remains speculative as cell abundances of air samples are not available for our study, previous studies have shown that aerolisation of cells may be quite substantial (Rastelli, et al. 2017). Bacterial abundance in cloud water samples taken at the Mt. Verde during the MarParCloud campaign ranged between 0.4 and 1.5×10^5 cells mL⁻¹ (Fig 201a). Although only few samples are available, these numbers agree well with previous reports (e.g. Hu, et al. 2018).

5.7.4 Ice-nucleating particles

The properties of ice-nucleating particles (INP) in the SML and in bulk seawater, airborne in the marine boundary layer as well as the contribution of sea-spray aerosol particles to the INP population in clouds were examined during the campaign. The numbers of INP (N_{INP}) at -12, -15 and -18 °C in the PM₁₀ samples from the CVAO varied from 0.000318 to 0.0232, 0.00580 to 0.0533 and 0.0279 to 0.100 std L⁻¹, respectively. INP measurements in the ocean water showed that enrichment as well as depletion of INP in SML compared to the bulk seawater occurred and enrichment factors EF varied from 0.36 to 11.40 and 0.36 to 7.11 at -15 and -20 °C, respectively (details in Gong, et al. 202019b). N_{INP} in PM₁ were generally lower than those in PM₁₀ and, furthermore, N_{INP} in PM₁ at CVAO did not show elevated N_{INP} at warm temperatures, in contrast to N_{INP} in PM₁₀. These elevated concentrations in PM₁₀ decreased upon heating the samples, clearly pointing to a biogenic origin of these INP. Therefore, ice active particles in general and biologically active INP in particular were mainly present in the supermicron particles, and particles in this size range are not suggested to undergo strong enrichment of OM during oceanic transfer via bubble bursting (Quinn et al., 2015 and refs. therein). N_{INP} (per volume of water) of the cloud water was roughly similar or slightly above that of the SML (Fig. 224), while concentrations of sea salt were clearly lower in cloud water compared to ocean water. Assuming sea salt and the INP to be similarly distributed in both, sea and cloud water (i.e., assuming that INP would not be enriched or altered during the

production of supermicron sea spray particles), N_{INP} is at least four orders of magnitude higher than what would be expected if all airborne INP would originated from sea spray. These first measurements indicate that other sources besides the ocean, such as mineral dust or other long ranged transported particles, contributed to the local INP concentration (details in Gong, et al. 2020a).
2020a).

5.8 The SML potential to form secondary organic aerosol particles

To explore if marine air masses exhibit a significant potential to form SOA, a Gothenburg Potential Aerosol Mass Reactor (Go:PAM) was used, that relies on providing a highly oxidizing medium reproducing atmospheric oxidation on timescales ranging from a day to several days in much shorter timescales (i.e., a few minutes). During the campaign, outdoor air and gases produced from a photochemical reactor was flowed through the Go:PAM (Watne, et al. 2018), and exposed to high concentrations of OH radicals formed via the photolysis of ozone and subsequent reaction with water vapour (Zabalegui, et al. 2019 and refs. therein). The aerosol particles produced at the outlet of the OFR were monitored by means of an SMPS i.e., only size distribution and number concentration were monitored. A subset of the collected SML samples were investigated within the Go:PAM and showed that particles were formed when these samples were exposed to actinic irradiation. These particles resulted most likely from the reaction of ozone with gaseous products that were released from the SML as shown recently (Ciuraru et al. 2015) and the results obtained here are explained in more detail in a separate paper by Zabalegui, et al. (2019). ~~and showed varying trends briefly discussed below. Ozone is known to react with iodide anions to produce different iodinated gases acting as aerosol precursors (Carpenter, et al. 2012; Carpenter and Nightingale 2015). In principle, this chemistry is mainly a bulk process and not related to the SML composition. However, a daily variation of the number of particles formed was observed (but from a very limited set of samples, $n = 3$) probably related to the daily sampling conditions. To explain these observations, two different hypothesis can be postulated: (i) the ozone bulk reaction is not efficient enough for our lab-to-the-field approach, (ii) ozone is scavenged away by the organic SML constituents and the products of these reactions are producing, or not, the aerosol particles in the Go:PAM. Zabalegui, et al. (2019) also pointed out Due to the limited number of samples, no firm conclusions can be made, but we observed~~ the clear need to have concentrated SML samples (~~reproduced-achieved~~ here by centrifugation of the authentic samples) as a prerequisite of aerosol formation which is pointing toward a specific “organic-rich” chemistry. Outdoor air masses were also investigated for their secondary mass production potential. During the campaign, northeast wind dominated i.e., predominantly clean marine air masses were collected. Those did not show any distinct diurnal difference for their secondary aerosols formation potential. However, a significant decrease of secondary organic mass was observed on September 30th, which will be analysed in more detail.

5.9 The way to advanced modelling

5.9.1 Modelling of cloud formation and vertical transfer of ocean-derived compounds

Besides for the assessment of the cloud types (section 4.1.4) it is intended to apply modelling approaches to simulate the occurrence and formation of clouds at the Mt. Verde site including advection, wind, effective transport and vertical transport. This will allow to model chemical multiphase processes under the given physical conditions. Furthermore, the potential vertical transfer of ocean-derived compounds to cloud level will be modelled. To this end, the meteorological model data by the Consortium for Small-scale Modelling-Multiscale Chemistry Aerosol Transport Model (COSMO) (Baldauf, et al. 2011) will be used to define a vertical meteorological data field. COSMO is a compressible and non-hydrostatic meteorological model and the current weather forecast model of the German Weather Service. The numerical calculation of the weather forecast is achieved by using information of the underlying orography and land-use, as well as boundary data of all meteorological fields. The needed boundary and initial fields will be derived from re-analysis-data and/or input parameters from coarse-resolved weather model data. First simulations show that clouds frequently occurred at heights of 700 m to 800 m (Fig. 232) in strong agreement with the observations. This demonstrates that clouds at Mt. Verde can form solely due to the local meteorological conditions and not necessarily due to orographic effects. Accordingly, the combination of the ground-based aerosol measurements and the in-cloud measurements at the top of Mt. Verde will be applied to examine important chemical transformations of marine aerosol particles during horizontal and vertical transport within the MBL. From the here presented measurements, a transfer of ocean-derived compounds to cloud level is very likely. To link and understand both measurement sites, in terms of important multiphase chemical pathways, more detailed modelling studies regarding the multiphase chemistry within the marine boundary layer combined with the impact of the horizontal and vertical transport on the aerosol and cloud droplet composition will be performed by using different model approaches (more details in the SI). In general, both projected model studies will focus on (i) determining the oxidation pathways of key marine organics and (ii) the evolution of aerosol and cloud droplet acidity by chemical aging of the sea spray aerosol. The model results will finally be linked to the measurements and compared with the measured aerosol particle concentration and composition and the in-cloud measurements at the top of the Mt. Verde.

5.9.2 Development of a new organic matter emission source function

The link of ocean biota with marine derived organic aerosol particles has been recognized (e.g. O'Dowd, et al. 2004). However, the usage of a single parameter like chl-*a* as indicator for biological processes and its implementation in oceanic emission parameterisations is insufficient as it does not reflect pelagic community structure and associated ecosystem functions. It is strongly suggested to incorporate process-based models for marine biota and OM rather than relying on a simple parameterizations (Burrows, et al. 2014). A major challenge is the high level of complexity of the OM in marine aerosol particles as well as in the bulk water

and the SML as potential sources. Within MarParCloud modelling, a new source function for the oceanic emission of OM will be developed as a combination of the sea spray source function of Salter, et al. (2015) and a new scheme for the enrichment of OM within the emitted sea spray droplets. This new scheme will be based on the Langmuir-Adsorption of organic species at the bubble films. The oceanic emissions will be parameterised following Burrows, et al. (2014), where the OM is partitioned into several classes based on their physicochemical properties. The measured concentration of the species in the ocean surface water and the SML (e.g. lipids, sugars and proteins) will be included in the parameterisation scheme. Finally, size class resolved enrichment functions of the organic species groups within the jet droplets will be implemented in the new scheme. The new emission scheme will be implemented to the aerosol chemical transport model MUSCAT (Multi-Scale Chemistry Aerosol Transport). MUSCAT is able to treat atmospheric transport and chemical transformation of different traces gases as well as particle properties. In addition to advection and turbulent diffusion, sedimentation, dry and wet deposition through the transport processes are considered, too. MUSCAT is coupled with COSMO that provides MUSCAT with all needed meteorological fields (Wolke, et al. 2004). The multiscale model system COSMO-MUSCAT will be used further to validate the emission scheme of OM via small and meso-scale simulations.~~The new emission scheme will be implemented to the aerosol model MUSCAT (Multi-Scale Chemistry Aerosol Transport) and be validated via small and meso-scale simulations using COSMO MUSCAT (Wolke, et al. 2004).~~

6 Summary and Conclusion

Within MarParCloud and with substantial contributions from MARSU, an interdisciplinary campaign in the remote tropical ocean took place in autumn 2017. This paper delivers a description of the measurement objectives including first results and provides an overview for upcoming detailed investigations.

Typical for the measurement site, the wind direction was almost constant from the north-easterly sector (30 – 60 °). The analysis of the air masses and dust measurements showed that dust input was generally low, however, partly moderate dust influences were observed. Based on very similar particle number size distributions at the ground and mountain sites, it was found that the MBL was generally well mixed with a few exceptions and the MBL height ranged from 600 to 1100 m. Differences in the PNSDs arose from the dust influences. The chemical composition of the aerosol particles and the cloud water indicated that the coarse mode particles served as efficient CCN. Furthermore, lipid biomarkers were present in the aerosol particles in typical concentrations of marine background conditions and anti-correlated with dust concentrations.

From the satellite cloud observations and supporting modelling studies, it was suggested that the majority of low-level clouds observed over the islands formed over the ocean and could form solely due to the local meteorological conditions. Therefore, ocean-derived aerosol particles, e.g. sea salt and marine biogenic compounds, might be expected to have some influence on cloud formation. The presence of compounds of marine origin in cloud water samples (e.g. sodium, methane sulfonic acid, FAA, TEP, distinct DOM classes) at the Mt. Verde supported an ocean-cloud link. The transfer of ocean-derived compounds, e.g. TEP, from

the ocean to the atmosphere was confirmed in controlled tank measurements. The DOM composition of the cloud waters was consistent with the SML and bulk water composition and partly different from the aerosol particle's composition. ~~However, INP measurements indicated that other sources besides the ocean and/or atmospheric transformations significantly contribute to the local INP concentration.~~ However, based on the findings that (biologically active) INP were mainly present in supermicron aerosol particles that are not suggested to undergo strong enrichment during ocean-atmosphere transfer as well as the INP abundance in seawater and in cloud water, other non-marine sources most likely significantly contributed to the local INP concentration.

The bulk water and SML analysis comprised a wide spectrum of biological and chemical constituents and consistently showed enrichment in the SML. Especially for the complex OM characterisation, some of the methods presented here have been used for the first time for such diverse sets of marine samples (e.g. DOM fractioning, metabolome studies with DART-HR-MS). Chl-*a* concentrations were typical for oligotrophic regions such as Cape Verde. The pigment composition indicated the presence of cyanobacteria, haptophytes and diatoms with a temporal change in dominating groups (from cyanobacteria to diatoms) suggests the start of the diatom bloom. Possible linkages to the background dust input will be resolved. Concentrations and SML enrichment of DOC were comparable to previous campaigns at the same location. . For the DOC as a sum parameter, the two applied sampling devices (manual and catamaran glass plate) provided very similar results. However, if this is also true for the various compound classes remains to be evaluated. Lipids established an important organic compound group in the SML and a selective enrichment of surface-active lipid classes within the SML was found. Observed enrichments also indicated on biotic and/or abiotic lipid degradation processing within the SML. The temporal variability of bacterial abundance was studied and provided first co-located SML and cloud water measurements for this particular oceanic province. Whether the strong day-to-day variability of absolute cell numbers in the SML and bulk water derived from changing water bodies and/or altered meteorological forcing needs to be further elucidated. Regarding mercury species, results indicate that the SML is the major compartment where (dissolved plus particulate) Hg were enriched, while MeHg was more likely concentrated in the bulk water, underlining the importance of SML in Hg enrichment dependent on OM.

For the trace gases, a variety of conditions were observed showing influences from ocean as well as long-range transport of pollutants. High sunlight and high humidity in this tropical region are key in ensuring that primary and secondary pollutants (e.g. ethene and ozone) are removed effectively, however additional processes need to be regarded. Measurements within the marine boundary layer and at the ocean-atmosphere interface, such as those shown here, are essential to understand the various roles of these short-lived trace gases with respect to atmospheric variability and wider climatic changes. The Cape Verde islands are likely a source region for HONO and the potential of the SML to form secondary particles needs to be further elucidated.

This paper shows the proof of concept of the connection between organic matter emission from the ocean to the atmosphere and up to the cloud level. We clearly see a link between the ocean and the atmosphere as (i) the particles measured at the surface are well mixed within the marine boundary layer up to cloud level and (ii) ocean-derived compounds can be found in the (submicron) aerosol particles at mountain height and in the cloud water. The organic

measurements will be implemented in a new source function for the oceanic emission of OM. From a perspective of particle number concentrations, the marine contributions to both CCN and INP are, however, –rather limited. These findings underline that further in depth studies differentiating between submicron and supermicron particles as well as between aerosol number and aerosol mass are strongly required. ~~A~~However, a clear description of any potential transfer patterns and the quantification of additional important sources must await the complete analysis of all the samples collected. The main current objective is to finalize all measurements and interconnect the meteorological, physical, biological and chemical parameters also to be implemented as key variables in model runs. Finally, we aim to achieve a comprehensive picture of the seawater and atmospheric conditions for the period of the campaign to elucidate ~~in particular~~ the abundance, ~~and~~ cycling and transfer mechanisms of organic matter between the marine environmental compartments.

Data availability. The data are available through the World Data Centre PANGAEA (<https://www.pangaea.de/>) under the following links:
<https://doi.pangaea.de/10.1594/PANGAEA.910693>, van Pinxteren, M (2020): (Table 2) Concentrations of pigments, DOC and microbial parameters in the Sea surface Microlayer (SML) and bulk water during MarParCloud at Cape Verde islands
<https://doi.pangaea.de/10.1594/PANGAEA.910692>, van Pinxteren, M (2020): (Table 1) Seawater chemistry during MarParCloud at Cape Verde islands

Special issue statement.

Appendix A1: List of acronyms

APS – Aerodynamic particle sizer
CCN – Cloud condensation nuclei
CCNc – Cloud condensation nuclei counter
CDOM – Chromophoric dissolved organic matter
chl-*a* – Chlorophyll-*a*
COSMO – Consortium for small-scale modelling-multiscale chemistry aerosol transport model
CTD – Conductivity-temperature-depth sensor
CVAO – Cape Verde atmospheric observatory
CVFZ – Cape Verde frontal zone
CVOO – Cape Verde ocean observatory
DART-QTOF-MS – Direct analysis in real time quadrupole time-of-flight mass spectrometry
DegLip – Lipid degradation products
DL – Dissolved lipids
DMS – Dimethyl sulfide
DOC – Dissolved organic carbon
DOM – Dissolved organic matter
ECWMF – European center for medium-range weather forecasts
EBUS – Eastern-boundary upwelling system

1321 EF – Enrichment factor (analyte concentration in the SML in respect to the analyte concentration in
 1322 the bulk water)
 1323 ETNA – Eastern tropical north Atlantic
 1324 FAA – Free amino acids
 1325 Go:PAM – Gothenburg potential aerosol mass reactor
 1326 HONO – Nitrous acid
 1327 HYSPLIT – Hybrid single-particle lagrangian integrated trajectory
 1328 INP – Ice nucleating particle(s)
 1329 LOPAP – Long path absorption photometer
 1330 LMWA – Low molecular weight acids
 1331 LMWN – Low molecular weight neutrals
 1332 MarParCat – Catamaran with glass plates for SML sampling
 1333 MarParCloud – Marine biological production, organic aerosol Particles and marine Clouds: a process
 1334 chain
 1335 MARSU – MARine atmospheric Science Unravelled
 1336 MBL – Marine boundary layer
 1337 MeHg – Methylmercury (MeHg)
 1338 Mt. Verde – Highest point of the São Vicente island (744 m)
 1339 MUSCAT – Multi-scale chemistry aerosol transport
 1340 NACW – North Atlantic central water masses
 1341 N_{CCN} – Cloud condensation nuclei number concentration
 1342 N_{INP} – Numbers of INP
 1343 OH – Hydroxyl radical
 1344 OFR – Oxidation flow reactor
 1345 OM – Organic matter
 1346 OMZ – Oxygen minimum zone
 1347 (O)VOC – (Oxygenated) volatile organic compounds
 1348 PM₁ – Particulate matter (aerosol particles) smaller than 1 µm
 1349 PM₁₀ – Particulate matter (aerosol particles) smaller than 10 µm
 1350 PNSDs – Particle number size distributions
 1351 POM – Particulate organic matter
 1352 PVM – Particle volume monitor
 1353 SACW – South Atlantic central water mass
 1354 SAL –Saharan air layer
 1355 SAS – Surface-active substances/surfactants
 1356 SML – Sea surface microlayer
 1357 SOA – Secondary organic aerosol
 1358 SSA – Sea spray aerosol
 1359 SMPS – Scanning mobility particle sizer
 1360 TEP – Transparent exopolymer particles
 1361 TSP – Total suspended particle
 1362 TM – Transmission mode
 1363 WSOM – Water-soluble organic matter
 1364
 1365
 1366 Acknowledgement

This work was funded by Leibniz Association SAW in the project “Marine biological production, organic aerosol particles and marine clouds: a Process Chain (MarParCloud)” (SAW-2016-TROPOS-2) and within the Research and Innovation Staff Exchange EU project MARSU (69089). We acknowledge the CVAO site manager Luis Neves and to the Atmospheric Measurement Facility at the National Centre for Atmospheric Science (AMF, NCAS) for the funding of the trace gas measurements. We thank the European Regional Development fund by the European Union under contract no. 100188826. The authors acknowledge Thomas Conrath, Tobias Spranger and Pit Strehl for their support in the fieldwork Kerstin Lerche from the Helmholtz-Zentrum für Umweltforschung GmbH – UFZ in Magdeburg is acknowledged for the pigment measurements. The authors thank Susanne Fuchs, Anett Dietze, Sontje Krupka, René Rabe and Anke Rödger for providing additional data and filter samples and Elisa Berdalet for the discussion about the pigment concentrations. Kay Weinhold, Thomas Müller und Alfred Wiedensohler are acknowledged for their data support. We thank Johannes Lampel for providing the photograph of Figure 1. María Eugenia Monge is a research staff member from CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina). Jianmin Chen thanks for funding from the Ministry of Science and Technology of China (No.2016YFC0202700), and National Natural Science Foundation of China (No. 91843301, 91743202, 21527814). Sanja Frka and Blaženka Gašparović acknowledge the Croatian Science Foundation for the full support under the Croatian Science Foundation project IP-2018-01-3105. In addition, the use of SEVIRI data and NWCSAF processing software distributed by EUMETSAT and obtained from the TROPOS satellite archive is acknowledged. Erik H. Hoffmann thanks the Ph.D. scholarship program of the German Federal Environmental Foundation (Deutsche Bundesstiftung Umwelt, DBU, AZ: 2016/424) for its financial support. Ryan Pereira thanks Juliane Bischoff and Sara Trojahn for technical support. We also thank the Monaco Explorations programme as well as captain and crew of MV YERSIN for supporting the Wave Glider deployment. Finally, the authors thank the anonymous reviewers for their valuable comments and suggestions.

Author contributions. MvP, KWF, NT and HH organized and coordinated the MarParCloud campaign. MvP, KWF, NT, CS, EB, XG, JV, HW, TBR, MR, CL, BG, TL, LW, JL, HC participated in the campaign. All authors were involved in the analysis, data evaluation and discussion of the results. MvP and HH wrote the manuscript with contributions from all co-authors. All co-authors proofread and commented the manuscript.

Competing interest. The authors declare that they have no conflict of interest.

References

Abbatt, J. P. D., Leaitch, W. R., Aliabadi, A. A., Bertram, A. K., Blanchet, J. P., Boivin-Rioux, A., Bozem, H., Burkart, J., Chang, R. Y. W., Charette, J., Chaubey, J. P., Christensen, R. J., Cirisan, A., Collins, D. B., Croft, B., Dionne, J., Evans, G. J., Fletcher, C. G., Gali, M., Ghahremaninezhad, R., Girard, E., Gong, W. M., Gosselin, M., Gourdal, M., Hanna, S. J., Hayashida, H., Herber, A. B., Hesarakis, S., Hoor, P., Huang, L., Husserr, R., Irish, V. E., Keita, S. A., Kodros, J. K., Kollner, F., Kolonjari, F., Kunkel, D., Ladino, L. A., Law, K., Lévassieur, M., Libois, Q., Liggio, J., Lizotte, M., Macdonald, K. M., Mahmood, R., Martin, R. V., Mason, R. H., Miller, L. A., Moravek, A., Mortenson,

- 1411 E., Mungall, E. L., Murphy, J. G., Namazi, M., Norman, A. L., O'Neill, N. T., Pierce, J. R., Russell, L.
1412 M., Schneider, J., Schulz, H., Sharma, S., Si, M., Staebler, R. M., Steiner, N. S., Thomas, J. L., von
1413 Salzen, K., Wentzell, J. J. B., Willis, M. D., Wentworth, G. R., Xu, J. W., and Yakobi-Hancock, J. D.:
1414 Overview paper: New insights into aerosol and climate in the Arctic, *Atmos. Chem. Phys.*, 19, 2527-
1415 2560, 10.5194/acp-19-2527-2019, 2019.
- 1416 Baldauf, M., Seifert, A., Forstner, J., Majewski, D., Raschendorfer, M., and Reinhardt, T.: Operational
1417 Convective-Scale Numerical Weather Prediction with the COSMO Model: Description and
1418 Sensitivities, *Monthly Weather Review*, 139, 3887-3905, 10.1175/mwr-d-10-05013.1, 2011.
- 1419 Batrakova, N., Travnikov, O., and Rozovskaya, O.: Chemical and physical transformations of mercury
1420 in the ocean: a review, *Ocean Science*, 10, 1047-1063, 10.5194/os-10-1047-2014, 2014.
- 1421 Bertram, T. H., Cochran, R. E., Grassian, V. H., and Stone, E. A.: Sea spray aerosol chemical
1422 composition: elemental and molecular mimics for laboratory studies of heterogeneous and multiphase
1423 reactions, *Chemical Society Reviews*, 47, 2374-2400, 10.1039/c7cs00008a, 2018.
1424
- 1425 Bigg, E. K., and Leck, C.: The composition of fragments of bubbles bursting at the ocean surface,
1426 *Journal of Geophysical Research-Atmospheres*, 113, 10.1029/2007jd009078, 2008.
- 1427 Blum, J. D., Popp, B. N., Drazen, J. C., Anela Choy, C., and Johnson, M. W.: Methylmercury production
1428 below the mixed layer in the North Pacific Ocean, *Nature Geoscience*, 6, 879-884, 10.1038/ngeo1918,
1429 2013.
- 1430 Bonsang, B., Gros, V., Peeken, I., Yassaa, N., Bluhm, K., Zoellner, E., Sarda-Estevé, R., and Williams,
1431 J.: Isoprene emission from phytoplankton monocultures, relationship with chlorophyll, cell volume, and
1432 carbon content., *Environmental Chemistry*, 7, 554-563, DOI: 10.1071/EN09156 2010.
- 1433 Bowman, K. L., Hammerschmidt, C. R., Lamborg, C. H., and Swarr, G.: Mercury in the North Atlantic
1434 Ocean: The U.S. GEOTRACES zonal and meridional sections, *Deep Sea Research Part II: Topical*
1435 *Studies in Oceanography*, 116, 251-261, 10.1016/j.dsr2.2014.07.004, 2015.
- 1436 Brandt, P., Bange, H. W., Banyte, D., Dengler, M., Didwischus, S. H., Fischer, T., Greatbatch, R. J.,
1437 Hahn, J., Kanzow, T., Karstensen, J., Krortzinger, A., Krahmann, G., Schmidtko, S., Stramma, L.,
1438 Tanhua, T., and Visbeck, M.: On the role of circulation and mixing in the ventilation of oxygen
1439 minimum zones with a focus on the eastern tropical North Atlantic, *Biogeosciences*, 12, 489-512,
1440 10.5194/bg-12-489-2015, 2015.
- 1441 Brooks, S. D., and Thornton, D. C. O.: Marine Aerosols and Clouds, in: *Annual Review of Marine*
1442 *Science*, Vol 10, edited by: Carlson, C. A., and Giovannoni, S. J., *Annual Review of Marine Science*,
1443 *Annual Reviews*, Palo Alto, 289-313, 2018.
- 1444 Brueggemann, M., Hayeck, N., and George, C.: Interfacial photochemistry at the ocean surface is a
1445 global source of organic vapors and aerosols, *Nature Communications*, 9, 10.1038/s41467-018-04528-
1446 7, 2018.
- 1447 Burrows, S. M., Hoose, C., Poschl, U., and Lawrence, M. G.: Ice nuclei in marine air: biogenic particles
1448 or dust?, *Atmos. Chem. Phys.*, 13, 245-267, 10.5194/acp-13-245-2013, 2013.
- 1449 Burrows, S. M., Ogunro, O., Frossard, A. A., Russell, L. M., Rasch, P. J., and Elliott, S. M.: A physically
1450 based framework for modeling the organic fractionation of sea spray aerosol from bubble film Langmuir
1451 equilibria, *Atmos. Chem. Phys.*, 14, 13601-13629, 10.5194/acp-14-13601-2014, 2014.
- 1452 Carpenter, L. J., Fleming, Z. L., Read, K. A., Lee, J. D., Moller, S. J., Hopkins, J. R., Purvis, R. M.,
1453 Lewis, A. C., Müller, K., Heinold, B., Herrmann, H., Fomba, K. W., van Pinxteren, D., Müller, C.,
1454 Tegen, I., Wiedensohler, A., Müller, T., Niedermeier, N., Achterberg, E. P., Patey, M. D., Kozlova, E.

- 1455 A., Heimann, M., Heard, D. E., Plane, J. M. C., Mahajan, A., Oetjen, H., Ingham, T., Stone, D., Whalley,
1456 L. K., Evans, M. J., Pilling, M. J., Leigh, R. J., Monks, P. S., Karunaharan, A., Vaughan, S., Arnold, S.
1457 R., Tschritter, J., Pöhler, D., Friess, U., Holla, R., Mendes, L. M., Lopez, H., Faria, B., Manning, A. J.,
1458 and Wallace, D. W. R.: Seasonal characteristics of tropical marine boundary layer air measured at the
1459 Cape Verde Atmospheric Observatory, *Journal of Atmospheric Chemistry*, 67, 87-140, 10.1007/s10874-
1460 011-9206-1, 2010.
- 1461 Carpenter, L. J., Archer, S. D., and Beale, R.: Ocean-atmosphere trace gas exchange, *Chemical Society*
1462 *Reviews*, 41, 6473-6506, 10.1039/c2cs35121h, 2012.
- 1463 Carpenter, L. J., and Nightingale, P. D.: Chemistry and Release of Gases from the Surface Ocean,
1464 *Chemical Reviews*, 115, 4015-4034, 10.1021/cr5007123, 2015.
- 1465 Chen, Q., Sherwen, T., Evans, M., and Alexander, B.: DMS oxidation and sulfur aerosol formation in
1466 the marine troposphere: a focus on reactive halogen and multiphase chemistry, *Atmos Chem Phys*, 18,
1467 *13617-13637*, <https://doi.org/10.5194/acp-18-13617-2018>, 2018.
- 1468 Ciuraru, R., Fine, L., van Pinxteren, M., D'Anna, B., Herrmann, H., and George, C.: Unravelling New
1469 Processes at Interfaces: Photochemical Isoprene Production at the Sea Surface, *Environmental Science*
1470 *& Technology*, 49, 13199-13205, 10.1021/acs.est.5b02388, 2015.
- 1471 Cochran, R. E., Jayarathne, T., Stone, E. A., and Grassian, V. H.: Selectivity Across the Interface: A
1472 Test of Surface Activity in the Composition of Organic-Enriched Aerosols from Bubble Bursting,
1473 *Journal of Physical Chemistry Letters*, 7, 1692-1696, 10.1021/acs.jpcllett.6b00489, 2016.
1474
- 1475 Cochran, R. E., Ryder, O. S., Grassian, V. H., and Prather, K. A.: Sea Spray Aerosol: The Chemical
1476 Link between the Oceans, Atmosphere, and Climate, *Accounts Chem. Res.*, 50, 599-604,
1477 10.1021/acs.accounts.6b00603, 2017.
- 1478 Čosović, B., and Vojvodić, V.: Voltammetric analysis of surface active substances in natural seawater,
1479 *Electroanalysis*, 10, 429-434, 10.1002/(sici)1521-4109(199805)10:6<429::Aid-elan429>3.3.Co;2-z,
1480 1998.
- 1481 Cunliffe, M., Engel, A., Frka, S., Gašparović, B., Guitart, C., Murrell, J. C., Salter, M., Stolle, C., Upstill-
1482 Goddard, R., and Wurl, O.: Sea surface microlayers: A unified physicochemical and biological
1483 perspective of the air-ocean interface, *Progress in Oceanography*, 109, 104-116,
1484 10.1016/j.pocean.2012.08.004, 2013.
- 1485 Cunliffe, M., and Wurl, O.: "Guide to best practices to study the ocean's surface", Occasional
1486 Publications of the Marine Biological Association of the United Kingdom (Plymouth, UK), 118 pp. ,
1487 (<http://www.mba.ac.uk/NMBL/>). 2014.
- 1488 de Leeuw, G., Andreas, E. L., Anguelova, M. D., Fairall, C. W., Lewis, E. R., O'Dowd, C., Schulz, M.,
1489 and Schwartz, S. E.: Production Flux of Sea Spray Aerosol, *Reviews of Geophysics*, 49,
1490 10.1029/2010rg000349, 2011.
- 1491 Demoz, B. B., Collett, J. L., and Daube, B. C.: On the Caltech Active Strand Cloudwater Collectors,
1492 *Atmos Res*, 41, 47-62, Doi 10.1016/0169-8095(95)00044-5, 1996.
- 1493 Derrien, M., and Le Gleau, H.: MSG/SEVIRI cloud mask and type from SAFNWC, *International*
1494 *Journal of Remote Sensing*, 26, 4707-4732, 10.1080/01431160500166128, 2005.
- 1495 Engel, A., Bange, H., Cunliffe, M., Burrows, S., Friedrichs, G., Galgani, L., Herrmann, H., Hertkorn,
1496 N., Johnson, M., Liss, P., Quinn, P., Schartau, M., Soloviev, A., Stolle, C., Upstill-Goddard, R., van

- 1497 Pinxteren, M., and Zäncker, B.: The Ocean's Vital Skin: Toward an Integrated Understanding of the Sea
1498 Surface Microlayer, *Front. Mar. Sci.*, 4, doi: 10.3389/fmars.2017.00165, 2017.
- 1499 Facchini, M. C., Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S., Ceburnis, D.,
1500 Flanagan, R., Nilsson, E. D., de Leeuw, G., Martino, M., Woeltjen, J., and O'Dowd, C. D.: Primary
1501 submicron marine aerosol dominated by insoluble organic colloids and aggregates, *Geophysical*
1502 *Research Letters*, 35, 10.1029/2008gl034210, 2008.
- 1503 Fiedler, B., Grundle, D. S., Schutte, F., Karstensen, J., Loscher, C. R., Hauss, H., Wagner, H., Loginova,
1504 A., Kiko, R., Silva, P., Tanhua, T., and Kortzinger, A.: Oxygen utilization and downward carbon flux
1505 in an oxygen-depleted eddy in the eastern tropical North Atlantic, *Biogeosciences*, 13, 5633-5647,
1506 10.5194/bg-13-5633-2016, 2016.
- 1507 Fomba, K. W., Müller, K., van Pinxteren, D., and Herrmann, H.: Aerosol size-resolved trace metal
1508 composition in remote northern tropical Atlantic marine environment: case study Cape Verde islands,
1509 *Atmos. Chem. Phys.*, 13, 4801-4814, 10.5194/acp-13-4801-2013, 2013.
- 1510 Fomba, K. W., Mueller, K., van Pinxteren, D., Poulain, L., van Pinxteren, M., and Herrmann, H.: Long-
1511 term chemical characterization of tropical and marine aerosols at the Cape Verde Atmospheric
1512 Observatory (CVAO) from 2007 to 2011, *Atmos. Chem. Phys.*, 14, 8883-8904, 10.5194/acp-14-8883-
1513 2014, 2014.
- 1514 Franklin, D., Poulton, J. A., Steinke, M., Young, J., Peeken, I., and Malin, G.: Dimethylsulphide, DMSP-
1515 lyase activity and microplankton community structure inside and outside of the Mauritanian upwelling,
1516 *Progress in Oceanography*, 83, 134–142, 2009.
- 1517 Frka, S., Kozarac, Z., and Čosović, B.: Characterization and seasonal variations of surface active
1518 substances in the natural sea surface micro-layers of the coastal Middle Adriatic stations, *Estuarine*
1519 *Coastal and Shelf Science*, 85, 555-564, 10.1016/j.ecss.2009.09.023, 2009.
- 1520 Frka, S., Pogorzelski, S., Kozarac, Z., and Čosović, B.: Physicochemical Signatures of Natural Sea Films
1521 from Middle Adriatic Stations, *Journal of Physical Chemistry A*, 116, 6552-6559, 10.1021/jp212430a,
1522 2012.
- 1523 Gantt, B., and Meskhidze, N.: The physical and chemical characteristics of marine primary organic
1524 aerosol: a review, *Atmos. Chem. Phys.*, 13, 3979-3996, 10.5194/acp-13-3979-2013, 2013.
- 1525 Garrett, W. D.: Collection of slick-forming materials from the sea surface, *Limnol. Oceanogr.*, 10, 602–
1526 605, 1965.
- 1527 Gašparović, B., Kozarac, Z., Saliot, A., Čosović, B., and Möbius, D.: Physicochemical characterization
1528 of natural and ex-situ reconstructed sea-surface microlayers, *Journal of Colloid and Interface Science*,
1529 208, 191-202, 10.1006/jcis.1998.5792, 1998a.
- 1530 Gašparović, B., Vojvodić, V., and Čosović, B.: Excretion of organic matter during an experimental
1531 phytoplankton bloom followed using o-nitrophenol as an electrochemical probe, *Croatica Chemica*
1532 *Acta*, 71, 271-284, 1998b.
- 1533 Gašparović, B., Plavšić, M., Čosović, B., and Saliot, A.: Organic matter characterization in the sea
1534 surface microlayers in the subarctic Norwegian fjords region, *Marine Chemistry*, 105, 1-14,
1535 10.1016/j.marchem.2006.12.010, 2007.
- 1536 Gong, X., Wex, H., van Pinxteren, M., Triesch, N., Fomba, K. W., Lubitz, J., Stolle, C., Robinson, T.-B.,
1537 Müller, T., Herrmann, H., and Stratmann, F.: Characterization of aerosol particles at Cabo Verde close
1538 to sea level and at the cloud level – Part 2: Ice-nucleating particles in air, cloud and seawater, *Atmos.*
1539 *Chem. Phys.*, 20, 1451–1468, <https://doi.org/10.5194/acp-20-1451-2020>, 2020b. Gong, X., Wex, H.,

- van Pinxteren, M., Triesch, N., Fomba, K. W., Lubitz, J., Stolle, C., Robinson, T. B., Müller, T., Herrmann, H., and Stratmann, F.: Characterization of aerosol particles at Cape Verde close to sea and cloud level heights—Part 2: ice nucleating particles in air, cloud and seawater, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2019-729>, 2019b.
- Gong, X., Wex, H., Voigtländer, J., Fomba, K. W., Weinhold, K., van Pinxteren, M., Henning, S., Müller, T., Herrmann, H., and Stratmann, F.: Characterization of aerosol particles at Cabo Verde close to sea level and at the cloud level – Part 1: Particle number size distribution, cloud condensation nuclei and their origins, *Atmos. Chem. Phys.*, 20, 1431–1449, <https://doi.org/10.5194/acp-20-1431-2020>, 2020a. Gong, X., Wex, H., Voigtländer, J., Fomba, K. W., Weinhold, K., van Pinxteren, M., Henning, S., Müller, T., Herrmann, H., and Stratmann, F.: Characterization of aerosol particles at Cape Verde close to sea and cloud level heights—Part 1: particle number size distribution, cloud condensation nuclei and their origins, <https://doi.org/10.5194/acp-2019-585>, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2019-585>, 2019a.
- Hepach, H., Quack, B., Ziska, F., Fuhlbrügge, S., Atlas, E., Peeken, I., Krüger, K., and Wallace, D. W. R.: Drivers of diel and regional variations of halocarbon emissions from the tropical North East Atlantic, *Atmos. Chem. Phys.*, 14, 1255–1275, [10.5194/acp-14-1255-2014](https://doi.org/10.5194/acp-14-1255-2014), 2014.
- Herrmann, H., Schaefer, T., Tilgner, A., Styler, S. A., Weller, C., Teich, M., and Otto, T.: Tropospheric Aqueous-Phase Chemistry: Kinetics, Mechanisms, and Its Coupling to a Changing Gas Phase, *Chemical Reviews*, 115, 4259–4334, [10.1021/cr500447k](https://doi.org/10.1021/cr500447k), 2015.
- Hoffmann, E. H., Tilgner, A., Schrödner, R., Bräuer, P., Wolke, R., Herrmann, H.: An advanced modeling study on the impacts and atmospheric implications of multiphase dimethyl sulfide chemistry, *Proceedings of the National Academy of Sciences*, 113(42), 11776–11781, 2016.
- Hoppel, W. A., Frick, G. M., and Larson, R. E.: Effect of nonprecipitating clouds on the aerosol size distribution in the marine boundary-layer, *Geophysical Research Letters*, 13, 125–128, [10.1029/GL013i002p00125](https://doi.org/10.1029/GL013i002p00125), 1986.
- Horowitz, L. W., Walters, S., Mauzerall, D. L., Emmons, L. K., Rasch, P. J., Granier, C., Tie, X. X., Lamarque, J. F., Schultz, M. G., Tyndall, G. S., Orlando, J. J., and Brasseur, G. P.: A global simulation of tropospheric ozone and related tracers: Description and evaluation of MOZART, version 2, *Journal of Geophysical Research-Atmospheres*, 108, [10.1029/2002jd002853](https://doi.org/10.1029/2002jd002853), 2003.
- Hu, W., Niu, H. Y., Murata, K., Wu, Z. J., Hu, M., Kojima, T., and Zhang, D. Z.: Bacteria in atmospheric waters: Detection, characteristics and implications, *Atmos. Environ.*, 179, 201–221, [10.1016/j.atmosenv.2018.02.026](https://doi.org/10.1016/j.atmosenv.2018.02.026), 2018.
- Huber, S. A., Balz, A., Abert, M., and Pronk, W.: Characterisation of aquatic humic and non-humic matter with size-exclusion chromatography–organic carbon detection–organic nitrogen detection (LC-OCD-OND), *Water research*, 45, 879–885, 2011.
- Karstensen, J., Fiedler, B., Schutte, F., Brandt, P., Kortzinger, A., Fischer, G., Zantopp, R., Hahn, J., Visbeck, M., and Wallace, D.: Open ocean dead zones in the tropical North Atlantic Ocean, *Biogeosciences*, 12, 2597–2605, [10.5194/bg-12-2597-2015](https://doi.org/10.5194/bg-12-2597-2015), 2015.
- Kawamura, K., Ishimura, Y., and Yamazaki, K.: Four years' observations of terrestrial lipid class compounds in marine aerosols from the western North Pacific, *Global Biogeochemical Cycles*, 17, 2003.
- Kaylor, A., Dwivedi, P., Pittman, J. J., Monge, M. E., Cheng, G., Li, S., and Fernandez, F. M.: Plasma-spray ionization (PLASI): a multimodal atmospheric pressure ion source for liquid stream analysis,

1585 Journal of the American Society for Mass Spectrometry, 25, 1788-1793, 10.1007/s13361-014-0948-2,
1586 2014.

1587 Kroflič, A., Frka, S., Simmel, M., Wex, H., and Grgić, I.: Size-Resolved Surface-Active Substances of
1588 Atmospheric Aerosol: Reconsideration of the Impact on Cloud Droplet Formation, Environmental
1589 Science & Technology, 52, 9179-9187, 10.1021/acs.est.8b02381, 2018.

1590 Law, C. S., Breiviere, E., de Leeuw, G., Garcon, V., Guieu, C., Kieber, D. J., Konradowitz, S., Paulmier,
1591 A., Quinn, P. K., Saltzman, E. S., Stefels, J., and von Glasow, R.: Evolving research directions in Surface
1592 Ocean-Lower Atmosphere (SOLAS) science, Environmental Chemistry, 10, 1-16, 10.1071/en12159,
1593 2013.

1594 Lelieveld, J., and Crutzen, P. J.: The role of clouds in tropospheric photochemistry, J. Atmos. Chem.,
1595 12, 229-267, <https://doi.org/10.1007/BF00048075>, 1991.

1596 Lewis, A. C., Hopkins, J. R., Carpenter, L. J., Stanton, J., Read, K. A., and Pilling, M. J.: Sources and
1597 sinks of acetone, methanol, and acetaldehyde in North Atlantic marine air, Atmos. Chem. Phys., 5, 1963-
1598 1974, 10.5194/acp-5-1963-2005, 2005.

1599 Li, T., Wang, Y., Mao, H., Wang, S., Talbot, R. W., Zhou, Y., Wang, Z., Nie, X., and Qie, G.: Insights
1600 on Chemistry of Mercury Species in Clouds over Northern China: Complexation and Adsorption,
1601 Environ Sci Technol, 52, 5125-5134, 10.1021/acs.est.7b06669, 2018.

1602 Marandino, C. A., De Bruyn, W. J., Miller, S. D., and Saltzman, E. S.: DMS air/sea flux and gas transfer
1603 coefficients from the North Atlantic summertime coccolithophore bloom, Geophysical Research Letters,
1604 35, 10.1029/2008gl036370, 2008.

1605 McCluskey, C. S., Hill, T. C. J., Humphries, R. S., Rauker, A. M., Moreau, S., Strutton, P. G.,
1606 Chambers, S. D., Williams, A. G., McRobert, I., Ward, J., Keywood, M. D., Harnwell, J.,
1607 Ponsonby, W., Loh, Z. M., Krummel, P. B., Protat, A., Kreidenweis, S. M., and DeMott, P. J.:
1608 Observations of Ice Nucleating Particles Over Southern Ocean Waters, Geophysical
1609 Research Letters, 45: 11989-97 2018a.

1610
1611 McCluskey, C. S., Ovadnevaite, J., Rinaldi, M., Atkinson, J., Belosi, F., Ceburnis, D., Marullo,
1612 S., Hill, T. C. J., Lohmann, U., Kanji, Z. A., O'Dowd, C., Kreidenweis, S. M., and DeMott, P.
1613 J.: Marine and Terrestrial Organic Ice-Nucleating Particles in Pristine Marine to Continentally
1614 Influenced Northeast Atlantic Air Masses, Journal of Geophysical Research-Atmospheres, 123:
1615 6196-212, 2018b.

1616

1617 McKay, W. A., Turner, M. F., Jones, B. M. R., and Halliwell, C. M.: Emissions of hydrocarbons from
1618 marine phytoplankton - Some results from controlled laboratory experiments, Atmos. Environ., 30,
1619 2583-2593, 10.1016/1352-2310(95)00433-5, 1996.

1620 Middlebrook, A. M., Murphy, D. M., and Thomson, D. S.: Observations of organic material in individual
1621 marine particles at Cape Grim during the First Aerosol Characterization Experiment (ACE 1), Journal
1622 of Geophysical Research: Atmospheres, 103, 16475-16483, 10.1029/97JD03719, 1998.

1623 Mochida, M., Kitamori, Y., Kawamura, K., Nojiri, Y., and K. Suzuki. K.: Fatty acids in the
1624 marine atmosphere: Factors governing their concentrations and evaluation of organic films on
1625 sea-salt particles , Journal of Geophysical Research-Atmospheres, 107, 2001.

1626
1627 Modini, R. L., Frossard, A.A., Ahlm, L., Russell, L. M., Corrigan, C. E., Roberts, G. C., Hawkins, L.N.,
1628 Schroder, J. C., Bertram, A. K., Zhao, R., Lee, A. K. Y. , Abbatt, J. P. D., Lin, J., Nenes, A., Wang, Z.,
1629 Wonaschutz, A., Sorooshian,A., Noone, K. J., Jonsson, H., Seinfeld, J. H., Toom-Sauntry, D.,
1630 Macdonald, A. M., and Leaitch, W. R., Primary marine aerosol-cloud interactions off the coast of

1631 [California, *J. Geophys. Res.-Atmos.*, 120\(9\), 4282-4303, doi:10.1002/2014jd022963, 2015.](#)
1632
1633 Müller, C., Iinuma, Y., Karstensen, J., van Pinxteren, D., Lehmann, S., Gnauk, T., and Herrmann, H.:
1634 Seasonal variation of aliphatic amines in marine sub-micrometer particles at the Cape Verde islands,
1635 *Atmos. Chem. Phys.*, 9, 9587-9597, 2009.

1636 Müller, K., Lehmann, S., van Pinxteren, D., Gnauk, T., Niedermeier, N., Wiedensohler, A.,
1637 Herrmann, H.: Particle characterization at the Cape Verde atmospheric observatory during the
1638 2007 RHaMBLe intensive, *Atmos. Chem. Phys.*, 10, 2709-21, 2010.
1639

1640 Mustaffa, N. I. H., Ribas-Ribas, M., and Wurl, O.: High-resolution variability of the enrichment of
1641 fluorescence dissolved organic matter in the sea surface microlayer of an upwelling region, *Elementa-*
1642 *Science of the Anthropocene*, 5, 10.1525/elementa.242, 2017.

1643 Mustaffa, N. I. H., Badewien, T. H., Ribas-Ribas, M., and Wurl, O.: High-resolution observations on
1644 enrichment processes in the sea-surface microlayer, *Scientific Reports*, 8, 10.1038/s41598-018-31465-
1645 8, 2018.

1646 O'Dowd, C. D., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Yoon, Y.
1647 J., and Putaud, J. P.: Biogenically driven organic contribution to marine aerosol, *Nature*, 431, 676-680,
1648 Doi 10.1038/Nature02959, 2004.

1649 Ovadnevaite, J., O'Dowd, C., Dall'Osto, M., Ceburnis, D., Worsnop, D. R., and Berresheim, H.:
1650 Detecting high contributions of primary organic matter to marine aerosol: A case study, *Geophysical*
1651 *Research Letters*, 38, 10.1029/2010gl046083, 2011.

1652 Pagnone, A., Volker, C., and Ye, Y.: Processes affecting dissolved iron across the Subtropical North
1653 Atlantic: a model study, *Ocean Dyn.*, 69, 989-1007, 10.1007/s10236-019-01288-w, 2019.

1654 Pastor, M. V., Pelegri, J. L., Hernandez-Guerra, A., Font, J., Salat, J., and Emellanov, M.: Water and
1655 nutrient fluxes off Northwest Africa, *Continental Shelf Research*, 28, 915-936,
1656 10.1016/j.csr.2008.01.011, 2008.

1657 [Patel, A., and Rastogi, N.: Chemical Composition and Oxidative Potential of Atmospheric PM10 over](#)
1658 [the Arabian Sea, *ACS Earth Space Chem.*, 4, 112-121, 10.1021/acsearthspacechem.9b00285, 2020.](#)
1659

1660 Patey, M. D., Achterberg, E. P., Rijkenberg, M. J., and Pearce, R.: Aerosol time-series measurements
1661 over the tropical Northeast Atlantic Ocean: Dust sources, elemental composition and mineralogy,
1662 *Marine Chemistry*, 174, 103-119, 10.1016/j.marchem.2015.06.004, 2015.

1663 Pereira, R., Ashton, I., Sabbaghzadeh, B., Shutler, J. D., and Upstill-Goddard, R. C.: Reduced air-sea
1664 CO₂ exchange in the Atlantic Ocean due to biological surfactants, *Nature Geoscience*, 11, 492-+,
1665 10.1038/s41561-018-0136-2, 2018.

1666 Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and
1667 cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, 10.5194/acp-7-1961-2007,
1668 2007.

1669 Prather, K. A., Bertram, T. H., Grassian, V. H., Deane, G. B., Stokes, M. D., DeMott, P. J., Aluwihare,
1670 L. I., Palenik, B. P., Azam, F., Seinfeld, J. H., Moffet, R. C., Molina, M. J., Cappa, C. D., Geiger, F. M.,
1671 Roberts, G. C., Russell, L. M., Ault, A. P., Baltrusaitis, J., Collins, D. B., Corrigan, C. E., Cuadra-
1672 Rodriguez, L. A., Ebben, C. J., Forestieri, S. D., Guasco, T. L., Hersey, S. P., Kim, M. J., Lambert, W.
1673 F., Modini, R. L., Mui, W., Pedler, B. E., Ruppel, M. J., Ryder, O. S., Schoepp, N. G., Sullivan, R. C.,
1674 and Zhao, D.: Bringing the ocean into the laboratory to probe the chemical complexity of sea spray

1675 aerosol, Proceedings of the National Academy of Sciences of the United States of America, 110, 7550-
1676 7555, 10.1073/pnas.1300262110, 2013.

1677 Putaud, J. P., Van Dingenen, R., Mangoni, M., Virkkula, A., Raes, F., Maring, H., Prospero, J. M.,
1678 Swietlicki, E., Berg, O. H., Hillamo, R., and Mäkelä, T.: Chemical mass closure and assessment of the
1679 origin of the submicron aerosol in the marine boundary layer and the free troposphere at Tenerife during
1680 ACE-2, Tellus B, 52, 141-168, 10.1034/j.1600-0889.2000.00056.x, 2000.

1681 Quinn, P. K., Collins, D. B., Grassian, V. H., Prather, K. A., and Bates, T. S.: Chemistry and Related
1682 Properties of Freshly Emitted Sea Spray Aerosol, Chemical Reviews, 115, 4383-4399,
1683 10.1021/cr500713g, 2015.

1684 [Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L. M., and Bates, T. S.: Small fraction of marine
1685 cloud condensation nuclei made up of sea spray aerosol, Nature Geoscience, 10, 674-+,
1686 10.1038/ngeo3003, 2017.](#)

1687

1688 Rahlff, J., Stolle, C., Giebel, H. A., Brinkhoff, T., Ribas-Ribas, M., Hodapp, D., and Wurl, O.: High
1689 wind speeds prevent formation of a distinct bacterioneuston community in the sea-surface microlayer,
1690 Fems Microbiology Ecology, 93, 10.1093/femsec/fix041, 2017.

1691 Rastelli, E., Corinaldesi, C., Dell'Anno, A., Lo Martire, M., Greco, S., Facchini, M. C., Rinaldi, M.,
1692 O'Dowd, C., Ceburnis, D., and Danovaro, R.: Transfer of labile organic matter and microbes from the
1693 ocean surface to the marine aerosol: an experimental approach, Scientific Reports, 7, 10.1038/s41598-
1694 017-10563-z, 2017.

1695 Read, K. A., Mahajan, A. S., Carpenter, L. J., Evans, M. J., Faria, B. V. E., Heard, D. E., Hopkins, J. R.,
1696 Lee, J. D., Moller, S. J., Lewis, A. C., Mendes, L., McQuaid, J. B., Oetjen, H., Saiz-Lopez, A., Pilling,
1697 M. J., and Plane, J. M. C.: Extensive halogen-mediated ozone destruction over the tropical Atlantic
1698 Ocean, Nature, 453, 1232-1235, 10.1038/nature07035, 2008.

1699 Read, K. A., Carpenter, L. J., Arnold, S. R., Beale, R., Nightingale, P. D., Hopkins, J. R., Lewis, A. C.,
1700 Lee, J. D., Mendes, L., and Pickering, S. J.: Multiannual Observations of Acetone, Methanol, and
1701 Acetaldehyde in Remote Tropical Atlantic Air: Implications for Atmospheric OVOC Budgets and
1702 Oxidative Capacity, Environmental Science & Technology, 46, 11028-11039, 10.1021/es302082p,
1703 2012.

1704 Roberts, G. C., and Nenes, A.: A continuous-flow streamwise thermal-gradient CCN chamber for
1705 atmospheric measurements, Aerosol Science and Technology, 39, 206-221, 10.1080/027868290913988,
1706 2005.

1707 Robinson, T.-B., Stolle, C., and Wurl, O.: Depth is Relative: The Importance of Depth on TEP in the
1708 Near Surface Environment, Ocean Sci. Discuss., 2019a.

1709 Robinson, T.-B., Wurl, O., Bahlmann, E., Jürgens, K., and Stolle, C.: 2019 b. 'Rising bubbles
1710 enhance the gelatinous nature of the air-sea interface', Limnology and Oceanography,
1711 10.1002/lno.11188, 2019b.

1712

1713 Salter, M. E., Zieger, P., Navarro, J. C. A., Grythe, H., Kirkevåg, A., Rosati, B., Riipinen, I., and Nilsson,
1714 E. D.: An empirically derived inorganic sea spray source function incorporating sea surface temperature,
1715 Atmos. Chem. Phys., 15, 11047-11066, 10.5194/acp-15-11047-2015, 2015.

1716 Schepanski, K., Tegen, I., and Macke, A.: Saharan dust transport and deposition towards the tropical
1717 northern Atlantic, Atmos. Chem. Phys., 9, 1173-1189, DOI 10.5194/acp-9-1173-2009, 2009.

1718 Schmetz, J., Pili, P., Tjemkes, S., Just, D., Kerkmann, J., Rota, S., and Ratier, A.: An introduction to
1719 Meteosat Second Generation (MSG), Bulletin of the American Meteorological Society, 83, 977-+,
1720 10.1175/BAMS-83-7-Schmetz-1, 2002.

1721 Simoneit, B. R. T., Chester, R., and Eglinton, G.: Biogenic lipids in particulates from lower
1722 atmosphere over eastern Atlantic , Nature, 267: 682-85, 1977.
1723

1724 Simoneit, B. R. T., Kobayashi, M., Mochida, M., Kawamura, K., Lee, M., Lim, H. J., Turpin,
1725 B. J., and Komazaki, Y.: Composition and major sources of organic compounds of aerosol
1726 particulate matter sampled during the ACE-Asia campaign, Journal of Geophysical Research-
1727 Atmospheres, 109, 2004.
1728

1729 Stolle, C., Nagel, K., Labrenz, M., and Jürgens, K.: Succession of the sea-surface microlayer in the
1730 coastal Baltic Sea under natural and experimentally induced low-wind conditions, Biogeosciences, 7,
1731 2975-2988, 2010.

1732 Stolle, C., Ribas-Ribas, M., Badewien, T.H., Barnes, J., Carpenter, L.J., Chance, R., Damgaard,
1733 L.R., Durán Quesada, A.M., Engel, A., Frka, S., Galgani, L., Gašparović, B., Gerriets, M.,
1734 Hamizah Mustaffa, N.I., Herrmann, H., Kallajoki, L., Pereira, R., Radach, F., Revsbech, N.P.,
1735 Rickard, P., Saint, A., Salter, M., Striebel, M., Triesch, N., Uher, G., Upstill-Goddard, R.C., van
1736 Pinxteren, M., Zäncker, B., Zieger, P., and Wurl, O.: The MILAN campaign: Studying diel
1737 light effects on the air-sea interface, accepted for Bulletin of the American Meteorological
1738 Society, 2019.
1739

1740 Stramma, L., Huttel, S., and Schafstall, J.: Water masses and currents in the upper tropical northeast
1741 Atlantic off northwest Africa, J. Geophys. Res.-Oceans, 110, 10.1029/2005jc002939, 2005.

1742 Tang, K., Page, J. S., and Smith, R. D.: Charge competition and the linear dynamic range of detection
1743 in electrospray ionization mass spectrometry, Journal of the American Society for Mass Spectrometry,
1744 15, 1416-1423, 10.1016/j.jasms.2004.04.034, 2004.

1745 Triesch, N., van Pinxteren, M., Engel, A., and Herrmann, H.: Concerted measurements of free amino
1746 acids at the Cape Verde Islands: High enrichments in submicron sea spray aerosol particles and cloud
1747 droplets, Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2019-976>, in review, 2020

1748 van Pinxteren, D., Brüeggemann, E., Gnauk, T., Mueller, K., Thiel, C., and Herrmann, H.: A GIS based
1749 approach to back trajectory analysis for the source apportionment of aerosol constituents and its first
1750 application, Journal of Atmospheric Chemistry, 67, 1-28, 10.1007/s10874-011-9199-9, 2010.

1751 van Pinxteren, M., and Herrmann, H.: Glyoxal and methylglyoxal in Atlantic seawater and marine
1752 aerosol particles: method development and first application during the Polarstern cruise ANT XXVII/4,
1753 Atmos. Chem. Phys., 13, 11791-11802, 10.5194/acp-13-11791-2013, 2013.

1754 van Pinxteren, M., Fiedler, B., van Pinxteren, D., Iinuma, Y., Koertzinger, A., and Herrmann,
1755 H.: Chemical characterization of sub-micrometer aerosol particles in the tropical Atlantic
1756 Ocean: marine and biomass burning influences, Journal of Atmospheric Chemistry, 72: 105-
1757 25, 2015.
1758

1759 van Pinxteren, M., Barthel, S., Fomba, K., Müller, K., von Tümpling, W., and Herrmann, H.: The
1760 influence of environmental drivers on the enrichment of organic carbon in the sea surface microlayer
1761 and in submicron aerosol particles – measurements from the Atlantic Ocean, Elem Sci Anth, 5,
1762 <https://doi.org/10.1525/elementa.225>, 2017.

1763 Vaughan, S., Ingham, T., Whalley, L. K., Stone, D., Evans, M. J., Read, K. A., Lee, J. D., Moller, S. J.,
 1764 Carpenter, L. J., Lewis, A. C., Fleming, Z. L., and Heard, D. E.: Seasonal observations of OH and HO₂
 1765 in the remote tropical marine boundary layer, *Atmos. Chem. Phys.*, 12, 2149-2172, 10.5194/acp-12-
 1766 2149-2012, 2012.

1767 Watne, Å. K., Psichoudaki, M., Ljungström, E., Le Breton, M., Hallquist, M., Jerksjö, M., Fallgren, H.,
 1768 Jutterström, S., and Hallquist, Å. M.: Fresh and Oxidized Emissions from In-Use Transit Buses Running
 1769 on Diesel, Biodiesel, and CNG, *Environmental Science & Technology*, 52, 7720-7728,
 1770 10.1021/acs.est.8b01394, 2018.

1771 Wex, H., Dieckmann, K., Roberts, G. C., Conrath, T., Izaguirre, M. A., Hartmann, S., Herenz, P.,
 1772 Schäfer, M., Ditas, F., Schmeissner, T., Henning, S., Wehner, B., Siebert, H., and Stratmann, F., Aerosol
 1773 arriving on the Caribbean island of Barbados: Physical properties and origin, *Atmos. Chem. Phys.*,
 1774 14107–14130, 2016 doi:10.5194/acp-16-14107-2016.

1775 Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., Tuch,
 1776 T., Pfeifer, S., Fiebig, M., Fjaraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., Laj,
 1777 P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., Hüglin, C., Fierz-
 1778 Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F., Santos, S., Grunig, C., Faloon, K.,
 1779 Beddows, D., Harrison, R. M., Monahan, C., Jennings, S. G., O'Dowd, C. D., Marinoni, A., Horn, H.
 1780 G., Keck, L., Jiang, J., Scheckman, J., McMurry, P. H., Deng, Z., Zhao, C. S., Moerman, M., Henzing,
 1781 B., de Leeuw, G., Loschau, G., and Bastian, S.: Mobility particle size spectrometers: harmonization of
 1782 technical standards and data structure to facilitate high quality long-term observations of atmospheric
 1783 particle number size distributions, *Atmospheric Measurement Techniques*, 5, 657-685, 10.5194/amt-5-
 1784 657-2012, 2012.

1785 Wolke, R., Knuth, O., Hellmuth, O., Schröder, W., and Renner, E.: The parallel model system LM-
 1786 MUSCAT for chemistry-transport simulations: Coupling scheme, parallelization and application, in:
 1787 *Parallel Computing: Software Technology, Algorithms, Architectures, and Applications*, edited by: G.R.
 1788 Joubert, W. E. N., F.J. Peters, and W.V. Walter, Elsevier, Amsterdam, Niederlande, 363-370, 2004.

1789 Wurl, O., Miller, L., Ruttgers, R., and Vagle, S.: The distribution and fate of surface-active substances
 1790 in the sea-surface microlayer and water column, *Marine Chemistry*, 115, 1-9,
 1791 10.1016/j.marchem.2009.04.007, 2009.

1792 Wurl, O., Wurl, E., Miller, L., Johnson, K., and Vagle, S.: Formation and global distribution of sea-
 1793 surface microlayers, *Biogeosciences*, 8, 121-135, 10.5194/bg-8-121-2011, 2011.

1794 Wurl, O., Stolle, C., Van Thuoc, C., The Thu, P., and Mari, X.: Biofilm-like properties of the sea surface
 1795 and predicted effects on air-sea CO₂ exchange, *Progress in Oceanography*, 144, 15-24,
 1796 10.1016/j.pocean.2016.03.002, 2016.

1797 Wurl, O., Ekau, W., Landing, W. M., and Zappa, C. J.: Sea surface microlayer in a changing ocean - A
 1798 perspective, *Elementa-Science of the Anthropocene*, 5, 10.1525/elementa.228, 2017.

1799 Zabalegui, N., Manzi, M., Depoorter, A., Hayeck, N., Roveretto, M., Li, C., van Pinxteren, M.,
 1800 Herrmann, H., George, C., and Monge, M.E.: Seawater Analysis by Ambient Mass
 1801 Spectrometry-Based Seaomics and Implications on Secondary Organic Aerosol Formation,
 1802 submitted to *Atmos. Chem. Phys.*, 2019.

1803
 1804 Zäncker, B., Cunliffe, M., and Engel, A.: Bacterial Community Composition in the Sea Surface
 1805 Microlayer Off the Peruvian Coast. *Front. Microbiol.*, 9:2699. doi: 10.3389/fmicb.2018.02699,
 1806 2018.

1807

1808 Zindler, C., Peeken, I., Marandino, C. A., and Bange, H. W.: Environmental control on the variability
 1809 of DMS and DMSP in the Mauritanian upwelling region, Biogeosciences, 9, 1041-1051, 10.5194/bg-9-
 1810 1041-2012, 2012.
 1811

1812 **Caption of Figures:**

1813 Figure 1: Illustration of the different sampling sites during the campaign.

1814 Figure 2: The residence time of the air masses calculated from 96 h (4 days) back trajectories
 1815 in ensemble mode.

1816 Figure 3: Time-series of air temperature, wind direction, wind speed, ethene, dimethyl sulfide,
 1817 methanol, acetone, ethane and ozone.

1818 Fig. 4: The measured temperature and humidity profiles at the CVAO on September 17th
 1819 using a 16 m³ Helikite. From the measurements the boundary layer height was determined
 1820 (here: ~ 850 m).

1821 Fig. 5: Time series and vertical profiles of the MBL height simulated with COSMO-
 1822 MUSCAT on the N2 domain and measured with the helikite.

1823 Fig. 6: (a) ECMWF wind forecasts and (b – f) cloud scenery derived from Meteosat SEVIRI
 1824 observations for the Cape Verde Islands region using a , a state-of-the-art cloud classification
 1825 algorithm (the cloud retrieval software of the Satellite Application Facility on support to
 1826 Nowcasting and Very Short-Range Forecasting version 2016 (a) Average horizontal winds
 1827 have been derived from a 2.5 x 2.5 degree (250 km x 250 km) domain centered on Cape
 1828 Verde Islands and are plotted for each pressure level from 1000 to 250 hPa against time using
 1829 arrows. The arrow colours refer to the pressure level. Gray vertical lines mark the times of the
 1830 subsequently shown cloud scenes. (b – f) Different cloud scenes observed with Meteosat
 1831 SEVIRI for a domain of size 1500 km x 1000 km centered on the Cape Verde Islands. The
 1832 shadings refer to different cloud types derived with the cloud classification algorithm of the
 1833 NWC-SAF v2016.

1834 Fig. 7: (a) The mission track of a SV2 Wave Glider as color-coded fluorescence data derived
 1835 from a Wetlabs FLNTURT sensor installed on the vehicle (data in arbitrary units) (b).
 1836 Chlorophyll-a surface ocean concentrations derived from the MODIS-Terra satellite (mean
 1837 concentration for October 2017). Please note that logarithmic values are shown.

1838 Fig. 8: (a) The median of PNSDs of marine type (blue) and dust type2 (black), with a linear
 1839 and (b) a logarithmic scaling on the y axis, measured from September 21st 03:30:00 to
 1840 September 21st 20:00:00 (UTC) and from September 28th 09:30:00 to September 30th
 1841 18:30:00 (UTC). The error bar indicates the range between 25% and 75% percentiles.

1842 Fig. 9: N_{CCN} as a function of supersaturation during dust (black line) and marine (blue line)
 1843 periods. The shadows show the 25% to 75% percentiles.

1844 Fig. 910: (a) The median of PNSDs for marine type particle during cloud events and non-
 1845 cloud events at CVAO and MV; (b) Scatter plots of N_{CCN} at CVAO against those at MV at
 1846 supersaturation of ~ 0.30%. Slope and R² are given.

1847 Fig.110: (a) Percentage aerosol composition at the CVAO (mean value of 5 blocks) and (b) at
 1848 the Mt. Verde (mean value of 6 blocks) between October 2nd and October 9th. Aerosol particles

were samples in five different size stages from 0.05-0.14 μm (stage 1), 0.14-0.42 μm (stage 2), 0.42-1.2 μm (stage 3), 1.2-3.5 μm (stage 4) and 3.5-10 μm (stage 5).

Fig. 142: Cloud water composition for one connected sampling event between October 5th 7:45 (start, local time, UTC-1) and October 6th, 08:45 (start, local time, UTC-1).

Fig. 132: Straight chain unsaturated fatty acids ($\Sigma(\text{c12 to c33})$) concentrations on the PM_{10} aerosol particles versus atmospheric dust concentrations.

Fig. 134: Temporal evolution of DOC concentrations in the bulk water samples along the campaign together with the main pigment concentrations (chl-*a*, zeaxanthin and fucoxanthin) concentrations and total cell numbers measured in the bulk water and dust concentrations in the atmosphere (yellow background area).

Fig. 145: (a) Concentrations of DOC in the SML and (b) and in the bulk water sampled for paired glass plate (GP) and the MarParCat (cat) sampling events.

Fig 156: Average enrichments (EF) of surfactants (SAS) and dissolved lipid classes indicating organic matter degradation (DegLip).

Fig. 167: Concentrations of Hg, MeHg, DOC and POC in the sea surface microlayer (SML) and bulk water sampled on September 26th and 27th 2017.

Fig. 187: DOM classes measured in all compartments. The data represent mean values of three SML samples and the respective bulk water, three aerosol particle samples (PM_{10}) from the CVAO and two aerosol samples (PM_{10}) from the Mt. Verde and four cloud water samples, all collected between 26. – 27.09., 01. – 02.10., and 08. – 09.10.2017.

Fig. 198: (a) Total TEP abundance in the SML and the bulk water as well as enrichment factor (SML/ULW) of TEP for field samples taken in nearshore water Cape Verde; (b) together with tank experiment with > 3 h bubbling of water collected from nearshore Cape Verde.


Fig. 2049: Microscopy image of TEP in TSP aerosol particles sampled at the CVAO sampled between September 29th and 30th with a flow rate of 8 L min⁻¹.

Fig. 210: Bacterial abundance of SML and ULW from (a) field and (c) tank water samples as well as from cloud water samples (diamonds, a) taken during the campaign are shown. Additionally, enrichment factors (i.e. SML versus ULW) are presented (b, d). In panel a, please note the different power values between SML/ ULW (10^6 cells mL⁻¹) and cloud water samples (10^4 cells mL⁻¹).

Fig. 224: N_{INP} of SML seawater ($n = 9$) and cloud water ($n = 13$) as a function of temperature.

Fig. 223: Modelled 2D vertical wind field on October 5th after 12 hours of simulation time. The model domain spans 222 km length and 1.5 km height. The black contour lines represent the simulated cloud liquid water content (with a minimum of 0.01 g m⁻³ and a maximum of 0.5 g m⁻³). The more dense the lines, the higher the simulated liquid water content of the clouds.

1907
1908
1909
1910



1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927

Figure 1

1928
1929
1930




Figure 2

1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944




Figure 3

1960




Figure 4

1961

1962

1963

1964

1965

1966

1967

1968

1969

1970

1971

1972

1973

1974

1975




Figure 5

1976

1977

1978

1979

1980

1981

1982




Figure 6

1996
1997
1998





Figure 7

1999
2000
2001
2002
2003
2004

2005
2006
2007



2008
2009
2010
2011
2012

Figure 8




Figure 9




Figure 910




Figure 1011

2067

2068

2069

2070

2071

2072

2073

2074

2075

2076

2077

2078

2079

2080

2081

2082

2083




Figure 112




Figure 132

2102
2103
2104
2105




Figure 143

2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118





Figure 154

2130



2131

2132

Figure 156

2133

2134

2135

2136

2137

2138

2139

2140

2141

2142


2143

2144

2145

2146

2147
2148



2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161

Figure 176






Figure 178




(a)



(b)

Figure 189

2179



2180

2181

2182

2183

2184

2185

2186

2187

2188

2189

2190

2191

2192

2193


2194

2195

2196

Figure [1920](#)


2197
2198



2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211

Figure 219


2212
2213
2214



2215
2216
2217
2218
2219
2220
2221
2222
2223
2224

Figure 212

2225
2226
2227



2228

2229

Figure 232

2230

2231

2232

2233

2234

2235

2236

2237