

Measurement report: Chemical characteristics of PM_{2.5} during typical biomass burning season at an agricultural site of the North China Plain

Linlin Liang¹, Guenter Engling^{2,3}, Chang Liu¹, Wanyun Xu¹, Xuyan Liu⁴, Yuan Cheng⁵, Zhenyu Du⁶, Gen Zhang¹, Junying Sun¹, Xiaoye Zhang¹

¹ State Key Laboratory of Severe Weather & Key Laboratory for Atmospheric Chemistry, Chinese Academy of Meteorological Sciences, Beijing 100081, China

² Division of Atmospheric Sciences, Desert Research Institute, Reno, NV 89512, USA

³ Now at: California Air Resources Board, El Monte, CA 91731, USA

⁴ National Satellite Meteorological Center, Beijing 100081, China

⁵ School of Environment, Harbin Institute of Technology, Harbin 150001, China

⁶ National Research Center for Environmental Analysis and Measurement, Beijing 100029 China

Abstract:

Biomass burning activities are ubiquitous in China, especially in North China, where there is an enormous rural population and winter heating custom. Biomass burning tracers (i.e., levoglucosan, mannosan ~~-(MN)~~ and potassium (K⁺)), as well as other chemical components were quantified at a rural site (Gucheng, GC) in North China from 15 October to 30 November, during a transition heating season, when the field burning of agricultural residues was becoming intense. The measured daily average concentrations of ~~L~~Glevoglucosan, ~~mannosanMN~~ and K⁺ in PM_{2.5} during this study were $0.79 \pm 0.75 \mu\text{g m}^{-3}$, $0.03 \pm 0.03 \mu\text{g m}^{-3}$ and $1.52 \pm 0.62 \mu\text{g m}^{-3}$, respectively. Carbonaceous components and biomass burning tracers showed higher levels at nighttime than daytime, while secondary inorganic ions were enhanced during daytime. An episode with high levels of biomass burning tracers was encountered at the end of October, 2016, with high ~~L~~Glevoglucosan at $4.37 \mu\text{g m}^{-3}$. Based on the comparison of chemical components during different biomass burning pollution periods, it appeared that biomass combustion can obviously elevate carbonaceous components levels, whereas no essentially effect on secondary inorganic aerosols in the ambient air. Moreover, the ~~levoglucosanLG/mannosanMN~~ ratios during different biomass burning pollution periods remained at high values (in the range of 18.3 - 24.9), however, the ~~levoglucosanLG/K⁺~~ ratio was significantly elevated during the intensive biomass burning pollution period (1.67) when air

temperatures decreasing, substantially higher than in other biomass burning periods (averaged at 0.47).

Keywords: Biomass burning; Organic tracers; Levoglucosan; Mannosan; Potassium

1. Introduction

Particulate air pollution is attracting more and more concerns in China because of their obvious adverse impact on visibility reduction, as well as health implication and regional or global climate change (Kanakidou et al., 2009; Pope and Dockery, 2006; Cheng et al., 2016). Carbonaceous species, i.e., organic carbon (OC) and elemental carbon (EC), and water-soluble inorganic ions, e.g., SO_4^{2-} , NO_3^- and NH_4^+ are the major components of ambient aerosols (Liang et al., 2017; Du et al., 2014; Zheng et al., 2015; Tan et al., 2016). Biomass burning (BB) emissions constitute a large source of ambient particulate pollution, especially for carbonaceous components, i.e., primary organic carbon (POC) and black carbon (BC) on global scale (Bond et al., 2004; Tang et al., 2018; Titos et al., 2017). As an important aerosol component, black carbon from industrial and combustion emissions contributes to the enhanced $\text{PM}_{2.5}$ ([particulate matter with aerodynamic diameters less than 2.5 \$\mu\text{m}\$](#)) mass concentrations and influences regional radiative forcing (Chen et al., 2017). Fresh biomass burning aerosol was found to be mainly comprised of carbonaceous species which typically constitutes 50-60% of the total particle mass (Hallquist et al., 2009). Yao et al. (2016) identified approximately half of carbonaceous aerosols being contributed by biomass burning at Yucheng, a rural site in the North China Plain.

Biomass burning emissions also represent a potentially large source of secondary organic aerosol (SOA). The precursors and formation pathways of SOA from biomass burning emissions were investigated by extensive field observations (e.g., Zhu et al., 2015; 2017; Adler et al., 2011; Zhang et al., 2010; 2015). Based on morphological particle analysis, Yao et al. (2016) investigated the smoke emitted from biomass burning impacting SOA production. Sun et al. (2010) found that phenolic compounds, which were emitted in large amounts from wood combustion, can form SOA at high yields in aqueous-phase reactions. In addition, smoke from biomass burning can be transported thousands of kilometers downwind from the source areas. Biomass burning aerosol from Southeast Asia can be transported to China, Singapore and even further to North America (Liang et

al., 2017; Hertwig et al., 2015; Peltier et al., 2008). Based on molecular tracer measurements, synoptic data as well as air mass back trajectory analysis, a fire episode was captured at a background site of East China with smoke advected from Southeast Asia (Liang et al., 2017).

The North China Plain (NCP) is one of the most polluted regions in China. Severe haze–fog of longer duration and more extensive coverage has occurred frequently in the NCP area, especially during the seasons of autumn and winter. NCP covers one quarter of China's cultivated land and yields 35% of the agricultural products in China (Boreddy et al., 2017). The rural population in NCP is also large and dense, and biomass burning activities are common in this region in form of cooking and heating. Intense fire activity typically occurs in October after the corn harvest. Abundant smoke is emitted from agricultural burning, i.e., residential biofuel combustion, open field burns, etc. Various field observations have investigated different aspects of biomass burning, e.g., seasonal variations, chemical and physical properties of smoke particles, spatial distribution, sources, transport, etc., in the NCP region (Cheng et al., 2013; Shen et al., 2018; Sun et al., 2013; 2016; Boreddy et al., 2017; Xu et al., 2019). However, these field investigations of the contribution of biomass burning to ambient aerosols in the NCP region were concentrated on the city of Beijing (Cheng et al., 2013; Zheng et al., 2015; Duan et al., 2004). Little field research about biomass burning was reported for rural areas in the NCP. In fact, biomass burning activities are common in the rural areas of the NCP region, and the resulting smoke aerosol can be transported to urban areas, e.g., the city of Beijing, resulting in haze episodic events.

~~The objective of this study is — gain insights about the abundance of smoke during the typical biomass burning season, i.e., autumn–winter transition season, following the corn harvest. In this paper, we focus on quantifying multiple biomass burning tracers, i.e., levoglucosan, LG, mannosan, MN and K^+ as well as other chemical species in $PM_{2.5}$ in the rural areas of the NCP region during the typical biomass burning season. In order to characterize the biomass burning pollution status within rural areas of the NCP region, multiple biomass burning tracers, i.e., levoglucosan, mannosan and K^+ in $PM_{2.5}$ sampled at a heavily polluted rural site in Hebei province were quantified during a typical biomass burning season, i.e., autumn–winter transition season, following the corn harvest. Combined with the analysis of other chemical components, it reveals different levels of biomass combustion pollution impacting on different types of chemical components in ambient~~

aerosol. Meanwhile, based on the multi-analysis of biomass burning molecular tracers, back trajectory analysis, fire activity data and synoptic conditions, the results of this study demonstrate the biomass burning pollution status, as well as the formation process of severe biomass burning pollution episode in the rural atmosphere of North China. These results can provide valuable information about the biomass burning activities in all of Northern China.

~~The results of this study demonstrated the biomass burning pollution status, as well as chemical properties of ambient aerosols under different biomass burning pollution levels in the rural atmosphere of North China.~~

2. Site description and experimental Methods

2.1 Site description and sampling

Samples were collected at a rural site, Gucheng (GC, 39°09'N, 115°44'E; 15.2 m a.s.l), located on a platform at the China Meteorological Administration farm in the town of Gucheng (GC site), approximately 110 km southwest of Beijing and 35 km north of the city of Baoding (population of about 5 million) in Hebei province, as shown in Figure S1. The station is surrounded by agricultural fields, with major crop species being corn and wheat. The dominant wind direction at GC is southwest and northeast during the study period. This site is upwind of Beijing, when the wind blows from the south or southwest, where heavily polluted cities and regions of Hebei province, i.e., Baoding, Shijiazhuang, Xingtai, Handan, are located. Thus, it is an appropriate station for representing the air pollution situation in the NCP region (Sheng et al., 2018; Chi et al., 2018; Xu et al., 2019; 2020; Kuang et al., 2020).

Daytime and nighttime PM_{2.5} samples were collected from 15 October, 2016 to 23 November, 2016, by using PM_{2.5} High-volume (Hi-Vol) sampler (GUV-15HBL1, Thermo Fisher Scientific CO., LTD), at the nominal flow rate of 1.13 m³ min⁻¹. The daytime samples were collected from 07:00 to 19:00, while nighttime samples were collected from 19:00 to 07:00 local time of the next day. All PM_{2.5} samples were collected on quartz fiber filters, prebaked at 850 °C for at least 5 h to remove carbonaceous material. A total of 33 couples of daytime/nighttime samples and 6 whole-day samples as well as 4 field blank samples were collected during the sampling period. The filters were stored at -20 °C after sample collection.

2.2 Experimental Methods

2.2.1 Anhydrosugar and water-soluble inorganic ion analysis

The quartz filter samples were analyzed for biomass burning anhydrosugar tracers, i.e., levoglucosan and mannosan~~LG and MN~~, using an improved high-performance anion-exchange chromatography (HPAEC) method with pulsed amperometric detection (PAD) on a Dionex ICS-5000+ system. Levoglucosan~~LG~~ and mannosan~~MN~~ were separated by a Dionex Carbopac MA1 analytical column and guard column with an aqueous sodium hydroxide (NaOH, 480 mM) eluent at a flow rate of 0.4 mL min⁻¹. The detection limit of levoglucosan~~LG~~ and mannosan~~MN~~ was 0.002 mg L⁻¹ and 0.005 mg L⁻¹, respectively. More details about the HPAEC-PAD method can be found elsewhere (Iinuma et al., 2009).

The quartz filter samples were also analyzed for water-soluble inorganic ions by a Dionex ICS-5000+ ion chromatograph, including SO₄²⁻, NO₃⁻, NH₄⁺, Cl⁻, Ca²⁺, Na⁺, K⁺ and Mg²⁺, and the method detection limits for the individual ionic species were 0.18 µg L⁻¹, 0.15 µg L⁻¹, 0.03 µg L⁻¹, 0.048 µg L⁻¹, 0.08 µg L⁻¹, 0.01 µg L⁻¹, 0.01 µg L⁻¹, 0.008 µg L⁻¹, respectively. The cations were separated on an Ionpac CS12 analytical column and CG12 guard column with a 20 mM methanesulfonic acid as eluent at a flow rate of 1.0 mL min⁻¹, while the anions were separated on an Ionpac AS11-HC column and AG11-HC guard column with 21.5 mM KOH eluent at a flow rate of 1.0 mL min⁻¹. The water-soluble inorganic ion data were corrected by field blanks.

2.2.2 Organic carbon/elemental carbon analysis

OC and EC were measured on a punch (0.526 cm²) of each quartz sample by a thermal/optical carbon analyzer (DRI Model 2001, Desert Research Institute, USA), using the Interagency Monitoring of Protected Visual Environments (IMPROVE) thermal evolution protocol with reflectance charring correction. The analytical error of OC was within 10%, and one sample of every 10 samples was selected at random for duplicate analysis. The detection limit of OC was 0.82 µgC cm⁻² (Liang et al., 2017).

2.2.3 Gas online monitoring (i.e., NO, NO₂, SO₂, O₃, CO and NH₃)

During this campaign, commercial instruments from Thermo Fisher Scientific Co., LTD were used to measure O₃ (TE 49C), NO/NO₂/Nox (Model 42CTL), CO (TE 48CTL), and SO₂ (TE43CTL), while NH₃ was measured by an ammonia analyzer (DLT-100, Los Gatos Research,

USA) at GC station. All measurement data quality was controlled according to standard gases (Xu et al., 2019; Lin et al., 2011; Meng et al., 2018; Ge et al., 2018).

2.2.4 Meteorological parameters

The meteorological parameters, including air temperature, relative humidity (RH) and wind speed at a 24-h resolution at the GC site are presented in Figure 1. During this campaign, the daily average RH value was observed at $77 \pm 13\%$, with a range from 48% to 99%, while the daily wind speed was observed with an average value of $1.07 \pm 1.14 \text{ m s}^{-1}$, exhibiting moist and stable synoptic conditions at this rural site during the autumn-winter transition season. Moreover, there was rare precipitation during the sampling period at the GC site, except for two days, i.e., 20 and 27 October, 2016 (Figure 1).

2.2.5 Back trajectory and fire spot analysis

To characterize the transport pathways of the aerosol at the Gucheng site, back-trajectories were calculated with the NOAA Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model via NOAA ARL READY Website (<http://ready.arl.noaa.gov/HYSPLIT.php>).

To investigate the influence of biomass burning activities in surrounding areas, fire hot spot counts were obtained from the Fire Information for Resource Management System (FIRMS) (available at <https://firms.modaps.eosdis.nasa.gov/download/>).

2.2.6 Statistical analysis

Statistical analysis of data, i.e., the correlation analysis between the concentrations of levoglucosan, mannosan and K^{\pm} at the Gucheng site during the sampling period were conducted with the linear fitting method.

3. Results and discussion

3.1 Characteristics of chemical components in $\text{PM}_{2.5}$

In this study, the mass concentration of $\text{PM}_{2.5\text{-cal}}$ was reconstituted by the sum of carbonaceous components ($1.6 \times \text{OC} + \text{EC}$) and inorganic ions ($\text{SO}_4^{2-} + \text{NH}_4^+ + \text{NO}_3^- + \text{Cl}^- + \text{Ca}^{2+} + \text{Na}^+ + \text{K}^+ + \text{Mg}^{2+}$). Figure 1 describes the time-series variation obtained for daily $\text{PM}_{2.5\text{-cal}}$, and its major components (OC, EC, SO_4^{2-} , NO_3^- and NH_4^+), biomass burning tracers (levoglucosan ~~LG~~,

mannosan~~MN~~ and K⁺), ratios of levoglucosan/OC and meteorological factors (temperature~~(T)~~,
relative humidity (RH), wind speed (~~WS~~) and planetary boundary layer (PBL) height-rainfall)
during the sampling period. The average daily PM_{2.5-cal} mass concentration in the autumn-winter
transition season at GC reached $137 \pm 72.4 \mu\text{g m}^{-3}$, ranging from $23.3 \mu\text{g m}^{-3}$ to $319 \mu\text{g m}^{-3}$ (Table
1, Figure 1a), which is higher than during the severe winter haze in January, 2013 at an urban site
in Beijing ($121 \mu\text{g m}^{-3}$) (Zheng et al., 2015). The mass concentrations of these chemical species
during the day are distributed as follows (from highest to lowest): $\text{OC} > \text{EC} > \text{NO}_3^- > \text{SO}_4^{2-} >$
 $\text{NH}_4^+ > \text{Cl}^- > \text{Ca}^{2+} > \text{K}^+ > \text{Na}^+ > \text{Mg}^{2+}$. Organic matter (OM), calculated by multiplying OC values
with a coefficient of 1.6, was the most abundant PM component, the daily average value of which
was $70.4 \pm 49.6 \mu\text{g m}^{-3}$, accounting for nearly half (46.7%) of PM_{2.5-cal} mass, indicating obvious
organic pollution at the rural site in the North China Plain during the sampling season.

The measured daily average concentrations of biomass burning tracers, i.e., levoglucosan-LG,
mannosan~~MN~~ and K⁺ in PM_{2.5} during our study were $0.79 \pm 0.75 \mu\text{g m}^{-3}$, $0.03 \pm 0.03 \mu\text{g m}^{-3}$ and
 $1.52 \pm 0.62 \mu\text{g m}^{-3}$, respectively (Table 1). The~~-anhydrosugar ambient concentrations of~~
levoglucosan levels (levoglucosan-LG and mannosan-MN) in this study were higher than those
observed in the city of Beijing during the summer (averaged at $0.23 \pm 0.37 \mu\text{g m}^{-3}$, in the range of
 0.06 to $2.30 \mu\text{g m}^{-3}$) and winter (averaged at $0.59 \pm 0.42 \mu\text{g m}^{-3}$, in the range of 0.06 to $1.94 \mu\text{g m}^{-3}$
³ of 2010-2011) (Cheng et al., 2013). The biomass burning tracer levels and ratios observed in this
study and other field studies are summarized in Table S1. The highest concentrations of
levoglucosan-LG in GC were observed on 31 October, 2016 with $4.37 \mu\text{g m}^{-3}$, which is a sharp
increase (over 30 times) of the minimum concentration ($0.14 \mu\text{g m}^{-3}$) during that period (Figure 1c).
Accordingly, the PM_{2.5-cal} concentration during that period was also elevated (as high as $236 \mu\text{g m}^{-3}$)
³) (Figure 1a). Secondary inorganic aerosol (sulfate, SO_4^{2-} ; nitrate, NO_3^- and ammonium, NH_4^+ ,
SNA) species, were the major water soluble ions, accounting for 82.8% of total water soluble ions,
the daily average values of which were $10.5 \pm 6.87 \mu\text{g m}^{-3}$, $15.9 \pm 9.29 \mu\text{g m}^{-3}$ and $10.9 \pm 5.51 \mu\text{g m}^{-3}$,
respectively (Table 1). SNA species exhibited a synchronous temporal trend (Figure 1c), while
the NO_3^- concentrations exceeded those of SO_4^{2-} at the GC site, in contrast to the results of previous
studies, e.g., Tan et al. (2016), who found SO_4^{2-} to be the dominant species in PM_{2.5} during winter
in 2006 in Beijing. Similarly, Chi et al., (2018) also found NO_3^- concentrations exceeded those of

SO₄²⁻ at both Beijing and GC sites during the winter in 2016, although they observed that NH₄⁺ was the dominant component of SNA (the concentrations of SO₄²⁻, NO₃⁻ and NH₄⁺ were 14.0 μg m⁻³, 14.2 μg m⁻³, and 24.2 μg m⁻³, respectively).

3.2 Day-night variations in the characteristics of PM_{2.5} chemical components

Carbonaceous components and biomass burning tracers exhibited higher levels during nighttime than daytime, while secondary inorganic ions showed the opposite pattern, i.e., higher concentrations during daytime than nighttime (Figure 2 and Figure S2). Besides, the gap of carbonaceous components and anhydrosugars between daytime and nighttime (two-fold) was more significant than for secondary inorganic ions. EC and primary organic components (POC) are not subject to significant differences in chemical reactions in ambient air between daytime and nighttime, and they will be mainly influenced by the variations of the planetary boundary layer (PBL) height. In the night, the PBL height decreases, compressing air pollutants into a shallow layer, and subsequently resulting in faster accumulation and higher concentrations of pollutants (Zheng et al., 2015; Zhong et al., 2018; 2019). The contributions of OM and EC to PM_{2.5-cal} were observed to be higher at nighttime (53.9% and 16.6%) than daytime (43.8% and 13.7%) as well (Figure 3).

Besides the influence from variations of the PBL height, the chemical degradation of levoglucosan may occur due to photochemical reaction in the ambient aerosols during day-time, further enlarging the gap of levoglucosan levels between daytime and nighttime (Sang et al., 2016; Gensch et al., 2018). Consequently, the contribution of levoglucosan to PM_{2.5-cal} during daytime (0.370.45%) was observed to be considerably lower than that during nighttime (0.64%) (Figure 3). However, secondary inorganic ions have an important formation pathway, i.e., photochemical processing, during daytime. Thus, the secondary inorganic species (SO₄²⁻, NO₃⁻ and NH₄⁺) were enhanced during daytime due to photochemical formation (Sun et al., 2013; Zheng et al., 2015; Wu et al., 2018). The mass contributions of SO₄²⁻, NO₃⁻ and NH₄⁺ to PM_{2.5-cal} were decreased from daytime (9.9%, 14.5% and 10.0%) to nighttime (6.5%, 9.6% and 7.1%) (Figure 3). Such an enhancement in secondary transformations during daytime is more evident in terms of the sulfur and nitrogen oxidation ratios (SOR and NOR, molar ratio of sulfate or nitrate to the sum of sulfate and SO₂ or nitrate and NO₂), which have been used previously as indicators of secondary transformations (Sun et al., 2013; Zheng et al., 2015). Both SOR and NOR during daytime were

higher than those during nighttime (Figure S3), further confirming the elevated secondary formations of sulfate and nitrate during daytime.

In addition, the concentrations of other water-soluble inorganic ions, i.e., K^+ and Cl^- during nighttime ($1.78 \pm 0.95 \mu g m^{-3}$ and $6.08 \pm 4.00 \mu g m^{-3}$) were higher than those in daytime ($1.43 \pm 0.54 \mu g m^{-3}$ and $4.33 \pm 2.30 \mu g m^{-3}$), while their contributions to $PM_{2.5-cal}$ were reversed, due to the significant accumulation and higher concentrations of pollutants during nighttime. As Ca^{2+} , Mg^{2+} and Na^+ , mainly emitted from primary natural sources, such as dust, soil resuspension and sea salt, are subject to more activity during the daytime and also influenced by the airflow dynamics, the contribution of those species in nighttime were lower than those during daytime, especially for Ca^{2+} , decreasing from 2.2% in daytime to 0.9% at nighttime (Figure 3).

3.3 Biomass burning episodes and the impacts on chemical $PM_{2.5}$ characteristics

An episode with high biomass burning tracer levels was encountered on 31 October, 2016. The concentrations of levoglucosan_{LG} in $PM_{2.5}$ during this one-day episode ($4.37 \mu g m^{-3}$) were significantly higher than those during typical transition season at the GC site ($0.69 \pm 0.47 \mu g m^{-3}$) (Figure 1d). Meanwhile, there was significant change in the meteorological conditions, i.e., the wind direction changed from southwesterly to northerly winds (Figure S4). Northerly winds advected cold and dry air masses, with the lowest hourly temperature observed at $-5.3^\circ C$ (Figure S5). This notable temperature decline before the commencing of the operation of the central heating systems should have caused intense combustion activities for heating purposes at the rural site. Moreover, the synoptic situation on 31 October, 2016 was under weaker turbulence with low PBL height and small wind speeds (Figure 1f). These worsened meteorological conditions would further enhance aerosol accumulation.

Here, we mainly distinguish four sub-periods based on daily levoglucosan_{LG} concentrations during the time frame from 15 October to 23 November, 2016. The four periods were separated as follows: 15-30 October (Period I: Minor biomass burning), 31 October (Period II: Intensive biomass burning), 1-14 November (Period III: Major biomass burning), 15-23 November (Period IV: Heating season). Table 2 compares the concentrations of $PM_{2.5-cal}$ mass, chemical components and gases at the GC site during these four periods, as well as the ratios between the intensive, major BB periods and heating season to minor BB period. The level of levoglucosan_{LG}

during the intensive BB episode II was about 12 times of that during the minor BB period I. K^+ and Cl^- , the common biomass burning tracers utilized in many studies (Duan et al., 2004; Cheng et al., 2013), were also observed with increased abundance during intensive BB episode II. When entering into November, the weather was becoming cold, and thus combustion activities for heating in the rural areas commenced, resulting in the ambient levels of levoglucosan to increase to $0.92 \pm 0.47 \mu\text{g m}^{-3}$ during period III, about 3 times of those in Period I. The central heating systems in North China cities were operated during period IV, and the ambient level of levoglucosan was observed at $0.96 \pm 0.63 \mu\text{g m}^{-3}$, which was similar to that observed in period III.

The concentrations of OC and EC were also observed to be strongly elevated in period II (Table 2), and especially OC levels increased to $59.996.3 \pm 25.3 \mu\text{g m}^{-3}$ during the intensive BB episode II, nearly 6 times of ~~that those~~ during the minor BB period ($16.2 \pm 7.52 \mu\text{g m}^{-3}$). The levoglucosan/OC ratio was utilized to estimate the effect of biomass burning to ambient organic aerosols. Accordingly, levoglucosan/OC ratios sharply increased to 0.045 during period II, which ~~is~~ was ~~obvious~~ noticeably higher than during other periods in this study (Figure 1e). Moreover, this level is also higher than most of the published field observations, i.e., at urban sites (Zhang et al., 2008; Cheng et al., 2013; Zhang et al., 2014), rural sites (Sang et al., 2013; Ho et al., 2014; Pietrogrande et al., 2015; Mkoma et al., 2013) and agricultural sites (Ho et al., 2014; Jung et al., 2014), yet lower than at an urban site in northern Italy during winter time (in the range of 0.01 to 0.13) (Pietrogrande et al., 2015). This illustrates that biomass combustion played an important role in organic aerosol pollution during the intensive BB episode II. However, During the major BB (period III) and heating season (period IV), due to other emissions of OC enhanced during the major BB episode (period III) and heating season (period IV), i.e., combustion of coal and biofuel for heating, OC increased to a higher level ($55.2 \pm 17.1 \mu\text{gC m}^{-3}$ and $69.4 \pm 24.6 \mu\text{gC m}^{-3}$, respectively). Due to the abundance of organic aerosols, the contribution from biomass burning emission was thereby reduced and the levoglucosan/OC ratios during periods III and IV decreased to 0.016 ± 0.005 and 0.014 ± 0.006 , respectively, even lower than those observed in the minor BB period I (0.025 ± 0.008).

Compared to the carbonaceous components, the concentrations of secondary inorganic aerosol species (SO_4^{2-} , NO_3^- , NH_4^+) exhibited a different pattern, i.e., showing no obvious differences between minor BB period I and other three periods. The ratios of SO_4^{2-} , NO_3^- , NH_4^+ during periods

II, III and IV to period I were all around 1.0 (Table 2), with no increasing trend. Moreover, the relationships between levoglucosan-LG and OC (and EC) were better than those between levoglucosan-LG and SNA during daytime and nighttime (Figure S3). The precursor gases of SNA, i.e., SO₂, NO, NO₂ and NH₃, were observed to have an increasing trend when biomass burning was prevalent during periods III and IV, with the ratios to period I arranged from 1.13 to 1.90 (Table 2). The time-series variations of the gases (SO₂, NO_x, NH₃, CO and O₃) and PBL during the sampling period are shown in Figure S4. The primary emission gases were exhibited negative relationships with PBL, while O₃ exhibited obvious positive relationship with PBL (Figure S5). Combustion from different fossil fuels (coal, gasoline, diesel, etc.) and biomasses (straws, woods, leaves, etc.) can all emit CO into the atmosphere (Streets et al., 2003; Chantara et al., 2019; Merico et al., 2020). Due to the more abundant combustion in the colder weather, the concentrations of CO also increased to 1.65 ± 0.53 ppm and 1.18 ± 0.83 ppm during the major biomass burning period III and the heating season period IV, respectively.

The combustion of biomass, especially of agricultural residues (e.g., wheat and corn straw) is very common in the rural areas in North China during the autumn-winter transition period. During the autumn harvest season in North China, wheat and corn straw burning is common practice, resulting in more abundant fire spots when entering into November than period I (Figure 4). The intense biomass burning event on 31 October, 2016 was also supported by air mass back trajectory analysis (Figure 5), performed with the TrajStat software. Based on the 48 h back trajectories at the GC site at 00:00 (UTC time) on 1 November, 2016, the air mass at the GC site was restricted in the region of Beijing-Tianjing-Hebei, the polluted area where fire spots were numerous. However, on the previous and following day of this episode, i.e., 31 October and 2 November onward, the air masses arriving at GC were advected from the northwest of Mongolia, where mostly desert areas are present, with less farm land and rare biomass burning activities (Figure 5).

Mean percentiles of major components in PM_{2.5} with respect to different BB pollution periods at GC site during the sampling time are shown in Figure 6. With the variation of BB pollution periods, the EC fraction seems to exhibit no obvious change during periods I, II and III, but slightly increased during the heating season (period IV), while the OC fraction increased significantly from 34.0% during the minor BB period I elevated to 65.4% during the intense BB period II. The

contributions of sulfate, nitrate and ammonium to $PM_{2.5-cal}$ all decreased sharply from the minor BB period to the intense period (Figure 6). This suggests that organic aerosol species become more important during BB pollution periods, concerning their contribution to the $PM_{2.5-cal}$, while EC has no such character. The OM percentage during intense BB period II was 65.4%, about double of that during the minor biomass burning period (34.0%), indicating that there was a large fraction of OM in $PM_{2.5-cal}$ originating from BB at the GC site during intensive BB period II. Opposite to OM, contributions of secondary inorganic ions to $PM_{2.5-cal}$ significantly decreased with the BB pollution becoming more severe. The contributions of SO_4^{2-} , NO_3^- and NH_4^+ to $PM_{2.5-cal}$ during the minor BB episode (11.6%, 20.5% and 12.5%) obviously declined during the intense BB episode (1.73%, 7.73% and 4.24%).

3.4 Relationships among tracers during different biomass burning pollution periods

In addition to pollution level information of biomass burning molecular tracers, the ratios between them could also be used to identify the different biomass types or indicate the burning formation processes of atmospheric aerosols. Levoglucosan and mannosan showed a good relationship during the entire sampling period (Figure 7a, $r = 0.97$, $p < 0.01$). The levoglucosan_{LG}/mannosan_{MN} ratios during minor, intense, major biomass pollution and heating season periods were observed at high values, i.e., 24.9, 24.1, 24.8 and 18.3 respectively (Table 2, Figure 7). LG and MN in the four periods showed a good relationship (Figure 7a, $R^2 = 0.94$). Compared to the former three episodes (24.1 to 24.9, averaged at 24.6), the levoglucosan/mannosan ratio during the heating season period (18.3) decreased by 25.6%. Based on source emission studies, the levoglucosan_{LG}/mannosan_{MN} ratios from crop residue burning, i.e., rice straw, wheat straw and corn straw, are similar and are characterized by high values (averaged at 29, in the range of 12 to 55) (Zhang et al., 2007; Engling et al., 2009; Cheng et al., 2013; Jung et al., 2014), yet overlapped-overlapping with those from hardwood (averaged at 28, in the range of 11 to 146) (Bari et al., 2009; Jung et al., 2014) and grass burning (18.2 ± 10.2) (Sullivan et al., 2008), while softwood is characterized by relatively lower levoglucosan_{LG}/mannosan_{MN} ratios (averaged at 4.3, in the range of 2.5 to 4.7) (Engling et

al., 2006; Cheng et al., 2013; Jung et al., 2014). Subsequently, this declining trend of the levoglucosan/mannosan ratios during the heating season period was partly caused by the higher proportion of softwood combustion, which is characterized by relatively lower levoglucosan/mannosan ratios. According to the local habits, soft woods, e.g. China fir and pine are also commonly used as biofuels for stove heating in North China, since they allow sustained long-time heating duration.

The concentrations of levoglucosan and K^+ during minor, major BB episode and heating season were correlated well (Figure 7b, $r = 0.84$, $p < 0.01$), while the red dot of period II being off from the fitted regression line. The levoglucosan/ K^+ ratios during periods III and IV (0.52 and 0.51) were similar to those during a BB episode at an urban site in Beijing during winter time (levoglucosan/ $K^+ = 0.51$) (Cheng et al., 2013). However, the levoglucosan/ K^+ ratio during the intense BB period II increased to 1.67, which was significantly higher than that in typical straw combustion (< 1.0). Correspondingly, there was a significant drop in temperatures at the GC site during period II, with the average daily temperature sharply decreasing from 7.5 °C on 30 Oct to 0.31 °C on 31 October, 2016, and the average temperature at night of 31 October even decreased to -3.4 °C (Figure 1g). Hence, the combustion activities were apparently intense around the sampling site for heating purposes. Compared to K^+ , there is a large enrichment of levoglucosan in wood burning emissions, based on the results from previous biomass source combustion studies (Engling et al., 2006; Chantara et al., 2019). The influence of softwood and/or other materials from softwood, which are commonly used as biofuels for stove heating in North China (Cheng et al., 2013; Zhou et al., 2017), should be larger during this low temperature period. Moreover, levoglucosan/ K^+ ratios also can be influenced by combustion conditions, i.e., smoldering versus flaming burns. Biofuels are typically subject to smoldering combustion condition in residential stoves for heating purposes in the rural areas in North China, which was reflected in relatively higher levoglucosan/ K^+ ratios than during flaming combustion (Schkolnik et al., 2005; Lee et al., 2010).

4. Summary and conclusion

Anhydrosugars, including levoglucosan and mannosan, and water-soluble potassium ion were employed as molecular tracers to investigate the characteristics of biomass burning activities as well as chemical properties of ambient aerosols under different biomass burning pollution levels. The

measured daily average concentrations of ~~levoglucosan~~^{LG}, ~~mannosan~~^{MN} and K^+ in $PM_{2.5}$ during a typical biomass burning season from 15 October to 30 November, 2016 were $0.79 \pm 0.75 \mu g m^{-3}$, $0.03 \pm 0.03 \mu g m^{-3}$ and $1.52 \pm 0.62 \mu g m^{-3}$, respectively. The concentrations of carbonaceous components and biomass burning tracers were observed higher at nighttime than daytime, while the patterns of secondary inorganic ions (SO_4^{2-} , NO_3^- and NH_4^+) were opposite, since they were enhanced by photochemical formation during daytime. An episode with extreme biomass burning tracer levels was encountered on 31 October, 2016, with concentrations of ~~levoglucosan~~^{LG} as high as $4.37 \mu g m^{-3}$. Comparing the chemical compositions between different biomass burning periods, it was apparent that biomass burning can considerably elevate the levels of organic components, while not showing a significant effect on the production of secondary inorganic ions. Compared to the other biomass burning episodes, the levoglucosan/mannosan ratios during the heating season period slightly decreased, ~~Moreover, due to more local soft wood and smoldering combustion taking place for heating under the low temperatures, the~~ while ~~levoglucosan~~^{LG}/ K^+ ratio during the intensive BB period was unusually higher than those in the other three biomass burning periods.

Data availability. The data used in this study can be obtained from this open link: <https://pan.baidu.com/s/11bKUZff1KJbzNVxS3VsLaA> code: jvqx. It is also available from the corresponding author upon request (lianglinlin@cma.gov.cn).

Author contributions. LL designed conducted all observations and drafted the paper. GE revised the paper and improved the English writing. XL drew the Figure 4 and Figure 5. CL, WX, YC, ZD, GZ, JS and XZ interpreted the data and discussed the results. All authors approved the final version for publication.

Competing interests. The authors declare that they have no conflict of interest.

Special issue statement. This article is part of the special issue– “In-depth study of air pollution sources and processes within Beijing and its surrounding region (APHH-Beijing) (ACP/AMT interjournal SI)”. It is not associated with a conference.

Acknowledgements. This research is supported by the Beijing Natural Science Foundation (8192055) and CAMS Fundamental Research Funds (No. 2017Z011). The authors would like to

acknowledge Yingli Yu and Ye Kuang for their help with PM_{2.5} samples collection; Hongbing Cheng for help with chemical analyses.

Financial support. This research has been supported by the Beijing Natural Science Foundation (No. 8192055), State Environmental Protection Key Laboratory of Sources and Control of Air Pollution Complex (No. SCAPC201701) and Chinese Academy of Meteorological Sciences Fundamental Research Funds (No. 2017Z011).

References:

- Bari, M. A., Baumbach, G., Kuch, B., and Scheffknecht, G.: Wood smoke as a source of particle-phase organic compounds in residential areas, *Atmos. Environ.*, 43, 4722-4732, <https://doi.org/10.1016/j.atmosenv.2008.09.006>, 2009.
- Boreddy, S. K. R., Kawamura, K., Okuzawa, K., Kanaya, Y., and Wang, Z.: Temporal and diurnal variations of carbonaceous aerosols and major ions in biomass burning influenced aerosols over Mt. Tai in the North China Plain during MTX2006, *Atmos. Environ.*, 154, 106-117, <http://dx.doi.org/10.1016/j.atmosenv.2017.01.042>, 2017.
- Caseiro, A., Bauer, H., Schmidl, C., Pio, C. A., and Puxbaum, H.: Wood burning impact on PM₁₀ in three Austrian regions, *Atmos. Environ.*, 43, 2186-2195, <https://doi.org/10.1016/j.atmosenv.2009.01.012>, 2009.
- Chantara, S., Thepnuan, D., Wiriy, W., Prawan, S., and Tsai, Y.I.: Emissions of pollutant gases, fine particulate matters and their significant tracers from biomass burning in an open-system combustion chamber, *Chemos.* 224, 407-416, <https://doi.org/10.1016/j.chemosphere.2019.02.153>, 2019.
- Cheng, Y., He, K.B., Du, Z.Y., Engling, G., Liu, J.M., Ma, Y.L., Zheng, M., Weber, R.J.: The characteristics of brown carbon aerosol during winter in Beijing, *Atmos. Environ.*, 127, 355-364, <https://doi.org/10.1016/j.atmosenv.2015.12.035>, 2016.
- Cheng, Y., Engling, G., He, K.B., Duan, F.K., Ma, Y.L., Du, Z.Y., Liu, J.M., Zheng, M., and Weber, R.J.: Biomass burning contribution to Beijing aerosol, *Atmos. Chem. Phys.*, 13, 7765-7781, <https://doi.org/10.5194/acp-13-7765-2013>, 2013.
- Chi, X., He, P., Jiang, Z., Yu, X., Yue, F., Wang, L., Li, B., Kang, H., Liu, C., and Xie, Z.: Acidity of aerosols during winter heavy haze events in Beijing and Gucheng, China, *J. Meteorol. Res.*, 32, 14-25, <https://doi.org/10.1007/s13351-018-7063-4>, 2018.
- Drewnick, F., Hings, S. S., Curtius, J., Eerdekens, G., and Williams, J.: Measurement of fine particulate and gas-phase species during the New Year's fireworks 2005 in Mainz, Germany, *Atmos. Environ.*, 40, 4316-4327, <https://doi.org/10.1016/j.atmosenv.2006.03.040>, 2006.
- Du, Z.Y., He, K.B., Cheng, Y., Duan, F.K., Ma, Y. L., Liu, J.M., Zhang, X.L., Zheng, M., and Weber, R.: A yearlong study of water-soluble organic carbon in Beijing I: Sources and its primary vs. secondary nature, *Atmos. Environ.*, 92, 514-521, <https://doi.org/10.1016/j.atmosenv.2014.04.060>, 2014.
- Duan, F., Liu, X., Yu, T., and Cachier, H.: Identification and estimate of biomass burning contribution to the urban aerosol organic carbon concentrations in Beijing, *Atmos. Environ.*, 38, 1275-1282, <https://doi.org/10.1016/j.atmosenv.2003.11.037>, 2004.
- Engling, G., Carrico, C.M., Kreidenweis, S.M., Collett Jr, J.L., Day, D.E., Malm, W.C., Lincoln, L., Hao, W.M., Iinuma, Y., and Herrmann, H.: Determination of levoglucosan in biomass

444 combustion aerosol by high-performance anion-exchange chromatography with pulsed
 445 amperometric detection, *Atmos. Environ.*, 40, S299-
 446 S311, <https://doi.org/10.1016/j.atmosenv.2005.12.069>, 2006.

447 Engling, G., Lee, J. J., Tsai, Y. W., Lung, S. C. C., Chou, C. C. K., and Chan, C. Y.: Size resolved
 448 anhydrosugar composition in smoke aerosol from controlled field burning of rice straw,
 449 *Aerosol Sci. Tech.*, 43, 662-672, <https://doi.org/10.1080/02786820902825113>, 2009.

450 Ge, B.Z., Wang, Z.F., Lin, W.L., Xu, X.B., Li, J., Ji, D.S., and Ma, Z.Q.: Air pollution over the north
 451 china plain and its implication of regional transport: a new sight from the observed evidences,
 452 *Environ. Pollut.*, 166, 29-38, <https://doi.org/10.1016/j.envpol.2017.10.084>, 2017.

453 Gensch, I, Sang-Arlt, X.F, Laumer, W, Chan, C,Y, Engling, G, Rudolph, J., and Kiendler-Scharr, A.:
 454 Using $\delta^{13}\text{C}$ of levoglucosan as a chemical clock, *Environ. Sci.*
 455 *Tech.* <https://pubs.acs.org/doi/10.1021/acs.est.8b03054>, 2018.

456 He, K.B., Zhao, Q., Ma, Y.L., Duan, F.K., Yang, F.M., Shi, Z.B., and Chen, G.: Spatial and seasonal
 457 variability of PM_{2.5} acidity at two Chinese megacities: insights into the formation of secondary
 458 inorganic aerosols, *Atmos. Chem. Phys.*, 12, 1377-1395, [https://doi.org/10.5194/acp-12-1377-](https://doi.org/10.5194/acp-12-1377-2012)
 459 [2012](https://doi.org/10.5194/acp-12-1377-2012), 2012.

460 Hertwig, D., Burgin, L., Gan, C., Hort, M., Jones, A., Shaw, F., Witham, C., and Zhang, K.:
 461 Development and demonstration of a Lagrangian dispersion modeling system for real-time
 462 prediction of smoke haze pollution from biomass burning in Southeast Asia, *J. Geophys. Res.:*
 463 *Atmos.*, 120, 12605-12630, <https://doi.org/10.1002/2015JD023422>, 2015.

464 Ho, K.F., Engling, G., Sai Hang Ho, S., Huang, R., Lai, S., Cao, J., and Lee, S.C.: Seasonal
 465 variations of anhydrosugars in PM_{2.5} in the Pearl River Delta Region, China, *Tellus B*, 66,
 466 22577, <https://doi.org/10.3402/tellusb.v66.22577>, 2014.

467 Jung, J., Lee, S., Kim, H., Kim, D., Lee, H., and Oh, S.: Quantitative determination of the biomass-
 468 burning contribution to atmospheric carbonaceous aerosols in Daejeon, Korea, during the rice-
 469 harvest period, *Atmos. Environ.*, 89, 642-650, <https://doi.org/10.1016/j.atmosenv.2014.03.010>,
 470 2014.

471 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van
 472 Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y.,
 473 Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati,
 474 E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review,
 475 *Atmos. Chem. Phys.*, 5, 1053-1123, <https://doi.org/10.5194/acp-5-1053-2005>, 2005.

476 Klejnowski, K., Janoszka, K., and Czaplicka, M.: Characterization and seasonal variations of
 477 organic and elemental carbon and levoglucosan in PM₁₀ in Krynica Zdroj, Poland, *Atmos.*,
 478 8(10), 190. <https://doi.org/10.3390/atmos8100190>, 2017.

479 Kuang, Y., Xu, W.Y., Lin, W.L., Meng, Z.Y., Zhao, H. R., Ren, S.X. Zhang, G., Liang, L. L., and
 480 Xu, X. B.: Explosive morning growth phenomena of NH₃ on the North China Plain: Causes
 481 and potential impacts on aerosol formation, *Environ.*
 482 *Pollut.*, <https://doi.org/10.1016/j.envpol.2019.113621>, in press, 2020.

483 Lee, T., Sullivan, A.P., Mack, L., Jimenez, J.L., Kreidenweis, S.M., Onasch, T.B., Worsnop, D.R.,
 484 Malm, W., Wold, C.E., Hao, W.M., and Collett Jr., J.L.: Chemical smoke marker emissions
 485 during flaming and smoldering phases of laboratory open burning of wildland fuels, *Aerosol*
 486 *Science and Technology* 44. <http://dx.doi.org/10.1080/02786826.2010.499884>, 2010.

487 Liang, L., Engling, G., Du, Z., Cheng, Y., Duan, F., Liu, X., and He, K.: Seasonal variations and
 488 source estimation of saccharides in atmospheric particulate matter in Beijing, China,
 489 *Chemosphere*, 150, 365-377, [10.1016/j.chemosphere.2016.02.002](https://doi.org/10.1016/j.chemosphere.2016.02.002), 2016.

490 Liang, L., Engling, G., Zhang, X., Sun, J., Zhang, Y., Xu, W., Liu, C., Zhang, G., Liu, X., and Ma,
 491 Q.: Chemical characteristics of PM 2.5 during summer at a background site of the Yangtze
 492 River Delta in China, *Atmospheric Research*, 198, 163-172, [10.1016/j.atmosres.2017.08.012](https://doi.org/10.1016/j.atmosres.2017.08.012),

493 2017.

494 Liang, L., Engling, G., Cheng, Y., Liu, X., Du, Z., Ma, Q., Zhang, X., Sun, J., Xu, W., Liu, C., Zhang,
 495 G., and Xu, H.: Biomass burning impacts on ambient aerosol at a background site in East China:
 496 Insights from a yearlong study, *Atmospheric Research*, 231, 104660,
 497 10.1016/j.atmosres.2019.104660, 2020.

498 Lin, W.L., Xu, X.B., Sun, J.Y., Liu, X.W., and Wang, Y.: Background concentrations of reactive
 499 gases and the impacts of long-range transport at the jinsha regional atmospheric background
 500 station, *Science China (Earth Sciences)*, 54, 1604-1613, 2011.

501 Liu, L., Liu, Y., Wen, W., Liang, L., Ma, X., Jiao, J., and Guo, K.: Source Identification of Trace
 502 Elements in PM_{2.5} at a Rural Site in the North China Plain, *Atmos.*, 11(2),
 503 179, <https://doi.org/10.3390/atmos11020179>, 2020.

504 Meng, Z.Y., Xu, X.B., Lin, W. L., Ge, B.Z., Xie, Y.L., Song, B., Jia, S.H., Zhang, R., Peng, W.,
 505 Wang, Y., Cheng, H.B., Yang, W., and Zhao, H.R.: Role of ambient ammonia in particulate
 506 ammonium formation at a rural site in the North China Plain, *Atmos. Chem. Phys.*, 18, 167–
 507 184, <https://doi.org/10.5194/acp-18-167-2018>, 2018.

508 Merico, E., Grasso, F.M., Cesari, D., Decesari, S., Belosi, F., Manarini, F., De Nuntis, P., Rinaldi,
 509 M., Gambaro, A., Morabito, E., and Contini, D.: Characterisation of atmospheric pollution near
 510 an industrial site with a biogas production and combustion plant in southern
 511 Italy, <https://doi.org/10.1016/j.scitotenv.2020.137220>, *Sci. Tot. Environ.*, 717, 137220, 2020.

512 Mkoma, S. L., Kawamura, K., and Fu, P. Q.: Contributions of biomass/biofuel burning to organic
 513 aerosols and particulate matter in Tanzania, East Africa, based on analyses of ionic species,
 514 organic and elemental carbon, levoglucosan and mannosan, *Atmos. Chem. Phys.*, 13, 10325–
 515 10338, <https://doi.org/10.5194/acp-13-10325-2013>, 2013.

516 Peltier, R. E., Hecobian, A. H., Weber, R. J., Stohl, A., Atlas, E. L., Riemer, D. D., Blake, D. R.,
 517 Apel, E., Campos, T., and Karl, T.: Investigating the sources and atmospheric processing of
 518 fine particles from Asia and the Northwestern United States measured during INTEX B, *Atmos.*
 519 *Chem. Phys.*, 8, 1835-1853, <https://doi.org/10.5194/acp-8-1835-2008>, 2008.

520 Pietrogrande, M.C., Bacco, D., Ferrari, S., Kaipainen, J., Ricciardelli, I., Riekkola, M.-L., Trentini,
 521 A., and Visentin, M.: Characterization of atmospheric aerosols in the Po valley during the
 522 supersito campaigns — Part 3: Contribution of wood combustion to wintertime atmospheric
 523 aerosols in Emilia Romagna region (Northern Italy), *Atmos. Environ.*, 122, 291-
 524 305, <https://doi.org/10.1016/j.atmosenv.2015.09.059>, 2015

525 Pope, C.A., and Dockery, D.W.: Health effects of fine particulate air pollution: lines that connect, *J.*
 526 *Air Waste Manage.*, 56, 709-742, <https://doi.org/10.1080/10473289.2006.10464485>, 2006.

527 Sang, X.F., Zhang, Z.S., Chan, C.Y., and Engling, G.: Source categories and contribution of biomass
 528 smoke to organic aerosol over the southeastern Tibetan plateau, *Atmos. Environ.*, 78, 113-
 529 123, <https://doi.org/10.1016/j.atmosenv.2012.12.012>, 2013.

530 Sang, X.F., Gensch, I., Kammer, B., Khan, A., Kleist, E., Laumer, W., Schlag, P., Schmitt, S.H.,
 531 Wildt, J., Zhao, R., Mungall, E.L., Abbatt, J.P.D., and Kiendler-Scharr, A.: Chemical stability
 532 of levoglucosan: An isotopic perspective, *J. Geophys. Res.*, 43, 5419-
 533 5424, <https://doi.org/10.1002/2016GL069179>, 2016.

534 Schkolnik, G., Falkovich, A.H., Rudich, Y., Maenhaut, W., and Artaxo, P.: New analytical method
 535 for the determination of levoglucosan, polyhydroxy compounds, and 2-methylerythritol and its
 536 application to smoke and rainwater samples, *Environ. Sci. Technol.*, 39, 2744-
 537 2752, <https://doi.org/10.1021/es048363c>, 2005.

538 Shen, X.J., Sun, J.Y., Zhang, X.Y., Zhang, Y.M., Wang, Y.Q., Tan, K.Y., Wang, P., Zhang, L., Qi,
 539 X.F., Che, H.Z., Zhang, Z., Zhong, J.T., Zhao, H.R., and Ren, S.X.: Comparison of submicron
 540 particles at a rural and an urban site in the North China Plain during the December 2016 heavy
 541 pollution episodes, *J. Meteorol. Res.*, 32, 14-25, <https://doi.org/10.1007/s13351-018-7060-7>,

2018.

Streets, D.G., Bond, T.C., Carmichael, G.R., Fernandes, S.D., Fu, Q., He, D., Klimont, Z., Nelson, S.M., Tsai, N.Y., Wang, M.Q., Woo, J.H., and Yarber, K.F.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *J. Geophys. Res.*, 108 (D21), 8809, <http://dx.doi.org/10.1029/2002JD003093>, 2003.

Sullivan, A.P., Holden, A.S., Patterson, L.A., McMeeking, G.R., Kreidenweis, S.M., Malm, W.C., Hao, W.M., Wold, C.E., and Collett Jr., J.L.: A method for smoke marker measurements and its potential application for determining the contribution of biomass burning from wildfires and prescribed fires to ambient PM_{2.5} organic carbon, *J. Geophys. Res.*, 113, D22302, <https://doi.org/10.1029/2008JD010216>, 2008.

Sun, K., Liu, X.G., Gu, J.W., Li, Y.P., Qu, Y., An, J.L., Wang, J.L., Zhang, Y.H., Hu, M., Zhang, F.: Chemical characterization of size-resolved aerosols in four seasons and hazy days in the megacity Beijing of China, *J. Environ. Sci.*, 32, 155-167, <https://doi.org/10.1016/j.jes.2014.12.020>, 2015.

Sun, Y.L., Wang, Z.F., Fu, P.Q., Yang, T., Jiang, Q., Dong, H.B., Li, J., and Jia, J.J.: Aerosol composition, sources and processes during wintertime in Beijing, China, *Atmos. Chem. Phys.*, 13, 4577-4592, <https://doi.org/10.5194/acp-13-4577-2013>, 2013.

Sun, Y.L., Zhang, Q., Anastasio, C., and Sun, J.: Insights into secondary organic aerosol formed via aqueous-phase reactions of phenolic compounds based on high resolution mass spectrometry, *Atmos. Chem. Phys.*, 10, 4809-4822, <https://doi.org/10.5194/acp-10-4809-2010>, 2010.

Tan, J.H., Duan, J.C., Zhen, N.J., He, K.B., and Hao, J.M.: Chemical characteristics and source of size-fractionated atmospheric particle in haze episode in Beijing, *Atmos. Res.*, 167, 24-33, <https://doi.org/10.1016/j.atmosres.2015.06.015>, 2016.

Tang, R., Wu, Z., Li, X., Wang, Y., Shang, D., Xiao, Y., Li, M., Zeng, L., Wu, Z., Hallquist, M., Hu, M., and Guo, S.: Primary and secondary organic aerosols in summer 2016 in Beijing, *Atmos. Chem. Phys.*, 18, 4055-4068, <https://doi.org/10.5194/acp-18-4055-2018>, 2018.

Urban, R.C., Lima-Souza, M., Caetano-Silva, L., Queiroz, M.E.C., Nogueira, R.F.P., Allen, A.G., Cardoso, A.A., Held, G., and Campos, M.L. A.M.: Use of levoglucosan, potassium, and water-soluble organic carbon to characterize the origins of biomass-burning aerosols, *Atmos. Environ.*, 61, 562-569, <https://doi.org/10.1016/j.atmosenv.2012.07.082>, 2012.

Wang, Z., Wang, T., Guo, J., Gao, R., Xue, L.K., Zhang, J.M., Zhou, Y., Zhou, X.H., Zhang, Q.Z., and Wang, W.X., 2012. Formation of secondary organic carbon and cloud impact on carbonaceous aerosols at Mount Tai, North China, *Atmos. Environ.*, 46, 516-527, <https://doi.org/10.1016/j.atmosenv.2011.08.019>, 2012.

Wu, Y., Ge, X., Wang, J., Shen, Y., Ye, Z., Ge, S., Wu, Y., Yu, H., and Chen, M.: Responses of secondary aerosols to relative humidity and photochemical activities in an industrialized environment during late winter, *Atmospheric Environment*, 193, 66-78, [10.1016/j.atmosenv.2018.09.008](https://doi.org/10.1016/j.atmosenv.2018.09.008), 2018.

Xu, W., Kuang, Y., Zhao, C., Tao, J., Zhao, G., Bian, Y., Yang, W., Yu, Y., Shen, C., Liang, L., Zhang, G., Lin, W., and Xu, X.: NH₃-promoted hydrolysis of NO₂ induces explosive growth in HONO, *Atmos. Chem. Phys.*, 19, 10557-10570, <https://doi.org/10.5194/acp-19-10557-2019>, 2019

Xu, W., Kuang, Y., Liang, L., He, Y., Cheng, H., Bian, Y., Tao, J., Zhang, G., Zhao, P., Ma, N., Zhao, H., Zhou, G., Su, H., Cheng, Y., Xu, X., Shao, M., and Sun, Y.: Dust-Dominated Coarse Particles as a Medium for Rapid Secondary Organic and Inorganic Aerosol Formation in Highly Polluted Air, *Environ Sci Technol*, 54, 15710-15721, [10.1021/acs.est.0c07243](https://doi.org/10.1021/acs.est.0c07243), 2020.

Xu, X., Zhang, H., Lin, W., Wang, Y., Xu, W., and Jia, S.: First simultaneous measurements of peroxyacetyl nitrate (PAN) and ozone at Nam Co in the central Tibetan Plateau: impacts from the PBL evolution and transport processes, *Atmos. Chem. Phys.*, 18, 5199-5217, <https://doi.org/10.5194/acp-18-5199-2018>, 2018.

- 591 Yang, F.M., Tan, J.H., Zhao, Q. Du, Z.Y., He, K.B., Ma, Y.L., Duan, F.K., Chen, G., and Zhao, Q.:
 592 Characteristics of PM_{2.5} speciation in representative megacities and across China, *Atmos.*
 593 *Chem. Phys.*, 11, 5207-5219, <https://doi.org/10.5194/acp-11-5207-2011>, 2011.
- 594 Yao, L., Yang, L.X., Chen, J.M., Wang, X.F., Xue, L.K., Li, W.J., Sui, X., Wen, L., Chi, J.W., Zhu,
 595 Y.H., Zhang, J.M., Xu, C.H., Zhu, T., and Wang, W.X.: Characteristics of carbonaceous
 596 aerosols: Impact of biomass burning and secondary formation in summertime in a rural area of
 597 the North China Plain, *Sci. Tot. Environ.*, 557-
 598 558, <https://doi.org/10.1016/j.scitotenv.2016.03.111>, 2016.
- 599 Yu, J.T., Yan, C.Q., Liu, Y., Li, X.Y., Zhou, T., and Zheng, M.: Potassium: A Tracer for Biomass
 600 Burning in Beijing? *Aerosol Air Qual. Res.*, 18, 2447-2459, doi: 10.4209/aaqr.2017.11.0536,
 601 2018.
- 602 Zhang, T., Cao, J.J., Chow, J.C., Shen, Z.X., Ho, K.F., Ho, S.S.H., Liu, S.X., Han, Y.M., Watson,
 603 J.G., Wang, G.H., and Huang, R.J.: Characterization and seasonal variations of levoglucosan
 604 in fine particulate matter in Xi'an, China, *J. Air Waste Manage.*, 64, 1317-
 605 1327, <https://doi.org/10.1080/10962247.2014.944959>, 2014.
- 606 Zhang, T., Claeyes, M., Cachier, H., Dong, S.P., Wang, W., Maenhaut, W., and Liu, X.D.:
 607 Identification and estimation of the biomass burning contribution to Beijing aerosol using
 608 levoglucosan as a molecular marker, *Atmos. Environ.*, 42, 7013-
 609 7021, <https://doi.org/10.1016/j.atmosenv.2008.04.050>, 2008.
- 610 Zhang, Z., Engling, G., Lin, C.-Y., Chou, C. C. K., Lung, S.-C. C., Chang, S.-Y., Fan, S., Chan, C.-
 611 Y., and Zhang, Y.-H.: Chemical speciation, transport and contribution of biomass burning
 612 smoke to ambient aerosol in Guangzhou, a mega city of China, *Atmospheric Environment*, 44,
 613 3187-3195, 10.1016/j.atmosenv.2010.05.024, 2010.
- 614 Zhang, Z., Gao, J., Engling, G., Tao, J., Chai, F., Zhang, L., Zhang, R., Sang, X., Chan, C.Y., Lin,
 615 Z., and Cao, J.: Characteristics and applications of size-segregated biomass burning tracers in
 616 China's Pearl River Delta region, *Atmos. Environ.*, 102, 290-
 617 301, <https://doi.org/10.1016/j.atmosenv.2014.12.009>, 2015.
- 618 Zheng, G.J., Duan, F.K., Su, H., Ma, Y.L., Cheng, Y., Zheng, B., Zhang, Q., Huang, T., Kimoto, T.,
 619 Chang, D., Pöschl, U., Cheng, Y. F., and He, K. B.: Exploring the severe winter haze in Beijing:
 620 the impact of synoptic weather, regional transport and heterogeneous reactions, *Atmos. Chem.*
 621 *Phys.*, 15, 2969-2983, <https://doi.org/10.5194/acp-15-2969-2015>, 2015.
- 622 Zheng, M., Salmon, L. G., Schauer, J. J., Zeng, L., Kiang, C. S., Zhang, Y., and Cass, G. R.: Seasonal
 623 trends in PM_{2.5} source contributions in Beijing, China, *Atmos. Environ.*, 39, 3967-
 624 3976, <https://doi.org/10.1016/j.atmosenv.2005.03.036>, 2005.
- 625 Zhong, J.T., Zhang, X.Y., and Wang, Y.Q.: Reflections on the threshold for PM_{2.5} explosive growth
 626 in the cumulative stage of winter heavy aerosol pollution episodes (HPEs) in Beijing, *Tellus B*,
 627 71, 1-7, <https://doi.org/10.1080/16000889.2018.1528134>, 2019.
- 628 Zhong, J.T., Zhang, X.Y., Wang, Y.Q., Liu, C., and Dong, Y.S.: Heavy aerosol pollution episodes in
 629 winter Beijing enhanced by radiative cooling effects of aerosols, *Atmos. Res.*, 209, 59-
 630 64, <https://doi.org/10.1016/j.atmosres.2018.03.011>, 2018.
- 631 Zhou, Y., Xing, X., Lang, J., Chen, D., Cheng, S., Wei, L., Wei, X., and Liu, C.: A comprehensive
 632 biomass burning emission inventory with high spatial and temporal resolution in China, *Atmos.*
 633 *Chem. Phys.*, 17, 2839-2864, 10.5194/acp-17-2839-2017, 2017.
- 634 Zhu, C., Kawamura, K., and Kunwar, B.: Effect of biomass burning over the western North Pacific
 635 Rim: wintertime maxima of anhydrosugars in ambient aerosols from Okinawa, *Atmos. Chem.*
 636 *Phys.*, 15, 1959-1973, <https://doi.org/10.5194/acp-15-1959-2015>, 2015.
- 637 Zhu, Y.H., Yang, L.X., Kawamura, K., Chen, J.M., Ono, K., Wang, X.F., Xue, L.K., and Wang, W.X.:
 638 Contributions and source identification of biogenic and anthropogenic hydrocarbons to
 639 secondary organic aerosols at Mt. Tai in 2014, *Environ. Pollut.*, 220, 863-

872, <https://doi.org/10.1016/j.envpol.2016.10.070>, 2017.

Table 1. Average concentrations and the range of PM_{2.5-cal} and its chemical components, biomass burning tracers ($\mu\text{g m}^{-3}$), gaseous species, ratios of OC/EC and levoglucosan LG/OC, as well as meteorological data observed at GC site at daytime, nighttime and whole day, respectively, during the sampling period from 15 Oct to 23 Nov 2016.

Species	Daytime (N = 34)		Nighttime (N = 33)		Whole period (N = 37)*	
	<u>Average</u> concentration	Range	<u>Average</u> concentration	Range	<u>Average</u> concentration	Range
PM _{2.5-cal}	117 \pm 58.8	19.0 - 225	170 \pm 116	21.1 - 465	137 \pm 72.4	23.3 - 319
OC	26.8 \pm 15.7	3.78 - 64.8	61.6 \pm 49.5	2.88 - 175	44.0 \pm 31.0	4.13 - 117
EC	13.4 \pm 8.49	1.44 - 34.0	30.9 \pm 28.5	2.21 - 129	21.7 \pm 15.8	2.46 - 74.9
TC	49.3 \pm 27.6	5.76 - 124	92.5 \pm 73.6	5.10 - 289	65.8 \pm 44.1	7.36 - 192
OC/EC	2.02 \pm 1.26	1.09 - 3.31	2.25 \pm 1.04	1.04 - 6.72	1.95 \pm 0.60	0.83 - 3.10
SO ₄ ²⁻	12.1 \pm 9.31	1.65 - 39.7	9.02 \pm 6.22	1.55 - 23.2	10.5 \pm 6.87	1.66 - 29.5
NO ₃ ⁻	16.9 \pm 9.96	1.85 - 41.2	13.1 \pm 8.52	1.56 - 38.0	15.9 \pm 9.29	2.40 - 45.2
Cl ⁻	4.33 \pm 2.30	0.82 - 9.46	6.08 \pm 4.00	0.62 - 16.0	4.90 \pm 2.46	0.93 - 9.37
NH ₄ ⁺	11.7 \pm 6.76	1.84 - 26.0	10.0 \pm 5.75	1.33 - 22.2	10.9 \pm 5.51	1.99 - 25.4
K ⁺	1.43 \pm 0.54	0.20 - 2.64	1.78 \pm 0.95	0.22 - 4.19	1.52 \pm 0.62	0.50 - 2.96
Mg ²⁺	0.26 \pm 0.14	0.07-0.64	0.19 \pm 0.09	0.06 - 0.38	0.14 \pm 0.12	0.04 - 0.43
Ca ²⁺	2.24 \pm 1.01	1.02-4.75	1.56 \pm 0.08	0.77 - 3.56	1.54 \pm 0.90	0.49 - 3.84

Na ⁺	0.44 ± 0.17	0.10 - 0.79	0.43 ± 0.24	0.10 - 1.31	0.42 ± 0.17	0.11 - 0.88
NO ₃ ⁻ / SO ₄ ²⁻	1.67 ± 0.82	0.75 - 5.52	1.54 ± 0.57	0.74 - 3.50	1.65 ± 0.62	0.78 ± 3.96
Levogluconan	0.57 ± 0.62	0.05 - 3.74	1.10 ± 0.99	0.05 - 4.82	0.79 ± 0.75	0.14 - 4.37
Mannosan	0.024 ± 0.023	0.00 - 0.14	0.05 ± 0.04	0.00 - 0.21	0.03 ± 0.03	0.00 - 0.18
Levogluconan LG/OC	0.018 ± 0.011	0.005 - 0.067	0.020 ± 0.010	0.004 - 0.047	0.020 ± 0.009	0.006 - 0.045
NO (ppb)	23.0 ± 14.7	2.07 - 56.0	45.9 ± 29.5	1.59 - 96.9	31.8 ± 18.3	1.81 - 68.5
NO ₂ (ppb)	25.8 ± 10.4	8.18 - 51.6	29.3 ± 9.37	8.81 - 51.1	26.6 ± 8.74	8.62 - 51.4
SO ₂ (ppb)	9.78 ± 4.96	3.11 - 22.5	9.63 ± 5.67	2.91 - 28.7	8.61 ± 4.04	3.37 - 20.4
CO (ppm)	0.96 ± 0.73	0.03 - 2.49	1.29 ± 1.04	0.02 - 3.26	1.05 ± 0.76	0.12 - 2.48
O ₃ (ppb)	13.0 ± 9.10	1.42 - 41.84	5.00 ± 5.73	1.60 - 24.30	9.25 ± 5.78	1.67 - 24.0
NH ₃ (ppb)	16.4 ± 11.3	1.68 - 46.2	18.3 ± 10.7	1.03 - 42.7	17.1 ± 9.88	1.46 - 44.4
Temperature (°C)	7.71 ± 4.01	- 2.07-15.9	3.30 ± 4.69	- 6.60 - 14.5	6.95 ± 4.58	- 4.33 - 15.4
Relative Humidity (%)	68 ± 17	31 - 98	85 ± 14	34 - 100	77 ± 13	48 - 99
Wind speed (m s ⁻¹)	1.43 ± 1.17	0.09 - 5.65	0.79 ± 1.55	0.03 - 7.19	1.07 ± 1.14	0.04 - 5.02

662 * Six whole-day samples were included used in the data analysis of the “Whole period”.

663

664

665

666

667

668

669 **Table 2.** Concentrations of chemical components in PM_{2.5} aerosols as well as their ratios and gaseous
670 species collected at the GC site, during the four biomass burning periods (i.e., Minor, Intensive, Major
671 and Heating period) from 15 Oct to 23 Nov 2016.

Species	Period I (15-30 Oct) Minor BB	Period II (31 Oct) Intensive BB	Ratio*	Period III (1 -14, Nov) Major BB	Ratio*	Period IV (15 -23, Nov) Heating period	Ratio*
	<u>Average</u> <u>concentration</u>	<u>Average</u> <u>concentration</u>		<u>Average</u> <u>concentration</u>		<u>Average</u> <u>concentration</u>	
PM _{2.5-cal}	81.0 ± 44.5	235	2.91	163 ± 46.7	2.01	189 ± 83.0	2.33
Levogluconan	0.36 ± 0.14	4.37	12.1	0.90 ± 0.37	2.50	0.96 ± 0.63	2.67
Mannosan	0.015 ± 0.005	0.18	12.0	0.038 ± 0.015	2.53	0.050 ± 0.026	3.33
OC	16.2 ± 7.52	96.3	5.93	55.2 ± 17.1	3.41	69.4 ± 24.6	4.28
EC	12.2 ± 5.85	36.0	2.96	25.5 ± 10.1	2.09	36.4 ± 21.5	2.98
TC	28.4 ± 13.1	132	4.66	80.9 ± 34.6	2.85	106 ± 55.3	3.73
SO ₄ ²⁻	10.3 ± 8.96	4.56	0.44	11.8 ± 6.02	1.15	9.08 ± 3.87	0.88
NO ₃ ⁻	16.6 ± 12.9	18.1	1.09	16.5 ± 6.42	0.99	12.6 ± 5.76	0.76
NH ₄ ⁺	10.1 ± 7.40	10.0	0.99	12.0 ± 4.35	1.19	10.3 ± 3.62	1.02
K ⁺	1.16 ± 0.36	2.61	2.25	1.76 ± 0.46	1.52	1.65 ± 0.84	1.42
Cl ⁻	3.46 ± 1.97	7.49	2.16	5.58 ± 2.16	1.61	6.27 ± 2.58	1.81
OC/EC	1.53 ± 0.35	2.67	1.75	2.31 ± 0.59	1.51	2.04 ± 0.31	1.33
NO ₃ ⁻ /SO ₄ ²⁻	1.74 ± 0.60	3.96	2.28	1.50 ± 0.35	0.86	1.42 ± 0.47	0.82

levoglucosan- LG/OC	0.025 ± 0.008	0.045	1.80	0.016 ± 0.005	0.64	0.014 ± 0.006	0.56
levoglucosan LG/EC	0.039 ± 0.019	0.121	3.10	0.038 ± 0.017	0.97	0.028 ± 0.013	0.72
levoglucosan- LG/ mannosan MN	24.9 ± 4.44	24.1	0.97	24.8 ± 6.46	1.00	18.3 ± 4.27	0.73
levoglucosan- G/K ⁺	0.36 ± 0.081	1.67	4.64	0.51 ± 0.16	1.42	0.53 ± 0.15	1.47
NO (ppb)	21.7 ± 12.5	21.7	1.00	39.6 ± 15.4	1.82	39.3 ± 23.6	1.81
NO ₂ (ppb)	21.8 ± 4.95	26.5	1.22	32.7 ± 7.27	1.50	24.6 ± 10.2	1.13
NO _x (ppb)	43.6 ± 16.3	48.2	1.11	72.4 ± 17.8	1.66	64.0 ± 33.4	1.47
SO ₂ (ppb)	5.83 ± 2.46	8.04	1.38	11.1 ± 4.10	1.90	9.75 ± 3.31	1.67
CO (ppm)	0.44 ± 0.33	0.70	1.59	1.65 ± 0.53	3.75	1.18 ± 0.83	2.68
O ₃ (ppb)	9.79 ± 4.88	23.2	2.37	7.51 ± 3.87	0.77	9.59 ± 7.55	0.98
NH ₃ (ppb)	14.3 ± 6.12	11.1	0.78	18.6 ± 8.03	1.30	21.2 ± 14.2	1.48

*: indicates that the ratios of the heating period, intense BB period or major biomass burning period
 were divided by those from the minor BB period.




Figure 1. Time-series variation obtained for $PM_{2.5-cal}$ and its major components, biomass burning tracers as well as meteorological factors at the GC site during the sampling period from 15 Oct to 23 Nov 2016. (a) $PM_{2.5-cal}$ and its major components (b) OC, and EC, (c) secondary inorganic aerosols, i.e., SO_4^{2-} , NO_3^- and NH_4^+ , (d) biomass burning tracers (levoglucosan LG, mannosan MN and K^+), (e) ratios of levoglucosan to OC (LG/OC) and levoglucosan to EC (LG/EC), and meteorological factors (f) PBL and wind speed (WS), (g) temperature (T) and relative humidity (RH), wind speed (WS) and rainfall at the GC site during the sampling period from 15 Oct to 23 Nov 2016.




Figure 2. Day and night distributions of mean concentrations of main chemical components (OC, EC, SO₄²⁻, NO₃⁻ and NH₄⁺) and biomass burning tracers (levoglucosan LG and mannosan MN) in PM_{2.5} observed at GC site during the sampling period.




Figure 3. Percent contributions of individual component mass concentrations to total estimated PM_{2.5-cal} mass in daytime and nighttime during the sampling period.




Figure 4. Fire spots at GC site and the surrounding provinces from (a) 15-30 October, 2016 and (b) 1 - 23, November, 2016, observed by MODIS Terra satellites (blue dot is GC station).




Figure 5. 48 h back trajectories at 500 m at GC site (39°09'N, 115°44'E) at 00:00 (UTC time) from 31 October to 2 November, 2016.




Figure 6. Mean percentiles of major components in PM_{2.5} with respect to different biomass burning pollution periods at GC site during the sampling time.




Figure 7. Scatter plots of (a) levoglucosan versus mannosan, (b) levoglucosan versus K^+ . [Statistical analysis of sampling data was conducted with the linear fitting method.](#)