

## ***Interactive comment on “10–year satellite–constrained fluxes of ammonia improve performance of chemistry transport models” by Nikolaos Evangeliou et al.***

**Nikolaos Evangeliou et al.**

nikolaos.evangeliou@nilu.no

Received and published: 16 January 2021

This manuscript provides a description of an inverse method based on the NH<sub>3</sub> lifetime to estimate NH<sub>3</sub> global emissions from the satellite IASI observations over the 10 yr-period 2008-2017. As NH<sub>3</sub> is a key species for understanding the PM levels, the quantification of its global emissions is important and would be useful to a wide community. The authors cover an important topic, appropriate for ACP. Nevertheless, I have some major comments listed below that should be considered by the authors before publication.

Response: We acknowledge reviewer's effort to improve our manuscript.

C1

Major comments:

1/ The fact that NH<sub>3</sub> columns in the atmosphere depend not only on NH<sub>3</sub> emissions, but is also linked to the abundance of nitric and sulfuric acids (and consequently to NO<sub>x</sub> and SO<sub>2</sub> emissions) is not fully described. To tackle the large variability of the ammonia lifetime, the authors calculated the NH<sub>3</sub> lifetime with a CTM and the spatial variability of ammonia is taken into account. I have more doubt about the temporal variability of ammonia and its main drivers in the atmosphere. If I well understand, the variable lifetime chosen for this study is a gridded average over the 10-yr period. If it is correct, the temporal trend in nitric and sulfuric acids is not fully taken into account, while it could have an importance for the deduced NH<sub>3</sub> emissions over a 10-yr period. This choice should be explained in the text. Would it possible to calculate yearly lifetimes as a sensitivity test to assess the robustness of your study?

Response: We appreciate reviewer's help to clarify this very important issue. As seen in Figure 1d and explained in the legend, the lifetime, as well as the emissions were calculated in monthly timesteps.

However, we admit this is not clear in the text, and therefore we have tried to clarify it further there. Some examples of our corrections are in section 2.3 (second paragraph, see Track Changes), Section 3 (first paragraph, see Track Changes), section 3.2 (first paragraph, see Track Changes). As we show in Figure 1d, the temporal trends of ammonia's reactants are considered and appear to have an effect on the lifetime, which varies from 10.3 to 12.2 hours.

2/ A comprehensive overview about the existing literature is missing. For example, result for SO<sub>2</sub> changes in Figure S2 is not in agreement with Krotkov et al., 2016, ACP, showing strong decrease of SO<sub>2</sub> between 2005 and 2015 at least over Eastern US and over Eastern Europe. Also, different publications have shown NH<sub>3</sub> peak in spring over northwestern European countries, not seen here. At least, discrepancies with previous studies should be discussed. These features could be explained by the choice of the

C2

authors to analyze their results for Europe or for the US as a whole. An analysis done for the hot-spot regions, of interest, where the emissions are high in Figure 4 may help the analysis.

Response: The legend of SO<sub>2</sub> explains that these are not results from our model/setup, but assimilated data from NASA's OMI (Ozone Monitoring Instrument) and MERRA2 (Modern-Era Retrospective Analysis for Research and Applications, Version 2). This is also explained in the manuscript (section 3.2, third paragraph, see Track Changes). About the seasonal variability of the NH<sub>3</sub> emissions, we agree with the reviewer that the spring peaks over northwestern European countries are not seen, because of our choice on the presentation of these results. Since we conduct a global study, we have chosen to study continental emissions rather than focusing only on hot-spot regions. The reason why we did this is because the aim of the paper is not to study the hot-spot emissions of NH<sub>3</sub> as seen from IASI. This has been highlighted already by Van Damme et al. Nature paper (see reference list of the manuscript). We focus on how the prescribed emissions retrieved from IASI can improve modelled concentrations and if models need higher emissions to capture measured concentrations. As a response to if our results are consistent with those of northwestern European countries highlighted in other papers, we plot seasonal emissions of NH<sub>3</sub> for all years, as in Figure 4 of the manuscript (see Fig. 1 below).

Except for years 2013 and 2015 that peak in summertime, all other years peaked in spring, which is in agreement with the reported hot-spot emissions in northwestern Europe.

3/ The impact of the abundance of sulfuric acid on NH<sub>3</sub> columns is detailed, but not the impact of the abundance of nitric acid. Is this impact considered negligible compared to those of sulfuric acids? This should be discussed. The same Figure S2 for NO<sub>2</sub> columns and nitrate concentrations may help analyzing the results.

Response: We agree with the reviewer. Reactions with nitric acid are not negligi-

C3

ble. However, they may have different results in NH<sub>3</sub> concentrations depending on the physicochemical parameters as we explain in the text (neutralization or production of NH<sub>3</sub>). We have retrieved NO<sub>2</sub> from OMI, in consistency with SO<sub>2</sub>, which we now present in Figure 2 of the manuscript and discuss in the text (section 3.2, circa p. 320-340).

Specific comments:

line 87: a comma is missing before "the Tropospheric Emission Spectrometer"

Response: Corrected (see Track Changes, circa L.87).

line 90-95: a verb is missing in this sentence

Response: Corrected (see Track Changes, circa L.91).

line 96-97: Note that Kuenen and Dore, [2019] estimated the uncertainties linked to the agricultural sector at about 100-300% at the European and annual scale. <https://www.eea.europa.eu/publications/emep-eea-guidebook-2019/part-a-general-guidance-chapters/5-uncertainties/view>

Response: We have added this useful information in the manuscript (see Track Changes, circa L.98).

line 98-102 : What is the differences between the different IASI products? The terms NE, VD0.5 and VDgrlf are not intuitive and are not explained at this stage.

Response: In principle, we agree with the reviewer here. However, we cannot add methodological details in the Introduction, and we'd rather prefer to leave only the names of the different datasets used in the analysis. Further down in the Methods section, we explain in detail what each name refers to and how the results were obtained. We have added a sentence explaining this in circa L.106 (see Track Changes).

line 105: please add references of studies using this state-of-the-art inventory.

C4

Response: The sentence we have added in circa L.106 (see Track Changes) refers to all emission datasets used in the present study including the state-of-the-art emissions from ECLIPSE-GFED4-GEIA and EDGAR-GFED4.

Line 124: could a difference of  $2\pm 24\%$  just due to the use of particular vertical profiles be interpreted as “small uncertainties”?

Response: We are not sure if we can judge the reported by Van Damme et al. (2018) values on uncertainties. However, as it is stated in their paper, the calculation does not refer to just particular vertical profiles, but rather to a global average: “Differences between columns derived with a fixed vertical profile (baseline) and columns derived using variable modelled profiles are of the order of  $2\% \pm 24\%$  on a global scale, but may be substantially larger for individual locations linked to regional differences in meteorological mixing and recirculation.”

Line 126-151: the description for CrIS gives more information than for IASI. The analysis of the results may be facilitating with the same information for both the instruments. I encourage you to give more information for IASI (total column uncertainties, peak sensitivity, detection limit, etc).

Response: We have added further details on errors and detection limits for IASI ammonia (see Track Changes in section 2.1.1). Though, we have tried to keep the length of the section consistent with this of CrIS and avoid repetitions, since detailed information of the product is published elsewhere (see references within the manuscript).

Line 152, Section 2.2: could you please provide a map of the interpolated IASI observations? As you performed simulations, it would be great to see the comparison between IASI and the CTM.

Response: We have added this plot in the Supplementary Figure 11, which gives an example of how the gridded results of IASI ammonia compares to the raw data. We believe it is more appropriate to show it there.

C5

Line 155: What is the CTM? As the variable lifetime in section 2.3 is based on this CTM, it should be described before. I would have described LMDZ-OR-INCA before section 2.3.

Response: We agree with the reviewer that the structure was awkward. We have moved the presentation of the CTM first in section 2 of the Methodology (see Track Changes section 2).

Line 160: I would refer to IASI ammonia total columns.

Response: Corrected. Please check at circa L. 214 (Track Changes).

Line 188: Please precise the regions where nitric and sulfuric acids are abundant in the text or at least, refer to Figure 2c and to Figure 2d.

Response: At this point, we discuss the method in general and do not refer to our results. We say that the use of a variable lifetime, and not a constant one, will be able to capture any variability caused by the chemical reactions of ammonia in the atmosphere, where and if they occur.

Line 211: Is the variable lifetime from a CTM for the quantification of VDgrlf emissions similar to the one for the quantification of NE emissions? This is not clear.

Response: We appreciate reviewer's help here. Indeed, this is not clear, and we have now corrected this part (see Track Changes at circa L. 266 of section 2.4).

Line 227-239: Has the NH<sub>3</sub> deposition of LMDz-OR-INCA been already evaluated? Is the bi-directional exchange with surfaces taken into account? This is not discussed. If not, how does it impact your NH<sub>3</sub> emissions?

Response: The total deposition of SO<sub>x</sub>, (SO<sub>2</sub>+SO<sub>4</sub><sup>2+</sup>), NH<sub>x</sub> (NH<sub>3</sub>+NH<sub>4</sub><sup>+</sup>), and NO<sub>y</sub> (NO+NO<sub>2</sub>+NO<sub>3</sub>+HNO<sub>2</sub>+HNO<sub>3</sub>+HNO<sub>4</sub>+N<sub>2</sub>O<sub>5</sub>+organic nitrates+particulate NO<sub>3</sub>-) have been evaluated (see Hauglustaine et al., 2014, in the manuscript). However, we admit we do not account for a compensation parameterization in the CTM, as high-

C6

lighted by the reviewer. We only have the emissions on one side, and the dry deposition ion on the other.

Line 253: you do not focus on hotspot regions but on continents as a whole.

Response: This is true; we agree with the reviewer and we have amended the text at this point (see Track Changes in circa L. 276, first paragraph of section 3).

Line 256-271: the different lifetimes of the literature and your results could be highlighted in a Table.

Response: There is a relevant supplementary Table in Van Damme et al. (2018) Nature paper (see reference within the manuscript), which presents literature values for ammonia lifetimes. We point to this table as "The atmospheric lifetimes of ammonia were summarized in Van Damme et al. (2018)." We do not want to be repetitive and put the same Table here. If the reviewer/editor has a different suggestion, we are willing to correct this in a next stage.

Line 276: As Ammonia lifetime depends on the presence of ammonia's reactants (sulfuric and nitric acid), it also depends on NO<sub>x</sub> and SO<sub>2</sub> emissions, not only NH<sub>3</sub> emissions. I would have written "(sulfuric and nitric acids, through SO<sub>2</sub> and NO<sub>x</sub> emissions)".

Response: We agree with the reviewer. As we have now clarified in the text, NO<sub>2</sub> and SO<sub>2</sub> are precursors of ammonia's atmospheric reactants, hence lifetime is indirectly linked to their concentrations. We have followed reviewer's suggestion to amend this sentence (see Track Changes in circa L. 376, p.10).

Figure 1: space is missing between the legend and Figure 1c and 1d

Response: We are not sure we understand where the problem is in Figure 1. Both the legend and the figure appear to be fine in our version. We have corrected some space problems in the title of Fig. 1c (reactants of NH<sub>3</sub>) that were overplotted by latitudinal values. If the reviewer still thinks there's a space missing somewhere, we could correct

C7

it in a next stage of the reviewing process.

Line 287: "which is in the range of the previously reported values". Your results are far from the results from Dammers et al [2019] for example. How do you explain such differences? Could the simulated NH<sub>3</sub> lifetime by CTM be over-estimated?

Response: We cannot judge the values calculated by Dammers et al. [2019]. As we report in circa L. 367-370 "The majority of ammonia lifetimes reported regionally or globally fall within 10 and 24 hours independently of the different approaches (Hauglustaine et al., 2014; Hertel et al., 2012; Möller and Schieferdecker, 1985; Sutton et al., 1993; Whitburn et al., 2016b),...".

Line 296: Please note in the legend of Figure 1b that the average ammonia emissions are calculated from the 10-year IASI observations and precise with which lifetime. I first thought it was the average ammonia emissions from ECLIPSEv5-GFED4-GEIA. Please also verify the legend of Figure S3.

Response: We thank the reviewer here; We have now clarified that the plot refers to the NE emissions (Track Changes at legend of Figure 1). We have also clarified this in the legend of the supplementary Figure S3.

Line 320-321: The sentence "Although column concentrations of both sulfur dioxide and sulfates present strong interannual variability, they do not show significant changes on an annual basis" is not clear. Please rephrase.

Response: We have amended this sentence to be consistent with what the figures show. Please see Track Changes at circa L. 428-432 (p.12).

Line 331: I do not understand why the anomaly is calculated only after 2015. Please explain.

Response: We initially thought to study anomalies after 2015, as our calculated emissions seem to increase after 2015. We agree with the reviewer that changes are already obvious since 2012 and now provide a more complete reasoning supported by

C8



relevant references. However, the largest reductions were seen after 2015, in agreement with the emissions of NH<sub>3</sub> that we present here, as seen in the attached Fig.2 and that is why we have chosen to restrict anomalies after 2015. Please see Track Changes at p.11-12.

Line 334-337: why the NH<sub>3</sub> emissions based on IASI observations could be impacted by changes in SO<sub>2</sub> and NO<sub>x</sub> emissions only after 2015? In Lachatre et al., 2019, the study you are citing line 337, the changes in SO<sub>2</sub> at least are seen before 2015. This is also the case in your Figure S2. Please strengthen this discussion.

Response: As we now discuss in L. 418-p.11, although the SO<sub>2</sub> and NO<sub>x</sub> reduction is evident since 2012, the largest changes are calculated for the period after 2015, which is in agreement with our suggested NH<sub>3</sub> emissions. This is also evident if we compare anomalies after 2012 with those after 2015 as in the attached Fig.2. Therefore, we present anomalies after 2015. We have tried to explain this in the manuscript (please see Track Changes in p.11-12).

Line 352: please deeply detail why the fact that northern India has been previously identified as a hot-spot region for ammonia explains the differences between the emission datasets.

Response: We believe we do not imply that the fact that N. India has been identified as a hotspot region explains the difference in the emission datasets. We only say that these hotspot emissions in N. India have been highlighted to be due to agricultural activities and we give 2 references to support this. We have now tried to re-write the sentence (see Track Changes at circa L. 476).

Line 335: Please verify the species indices

Response: We have amended this part and the overall discussion in this section as explained in previous comments (see Track Changes at p. 11-12).

Line 356: the ammonia emissions remain mostly constant at the global scale. Is it still

C9

true at continental scale?

Response: Yes, it is actually true that no significant continental changes occurred. For example, the ECLIPSE emissions which are based on the GAINS model are produced for 5-year timesteps. What global models assume is usually a linear interpolation to scale the emissions for each of the years in between. For justification, we plot the annual emissions from ECLIPSEv5-GFED4-GEIA and from EDGARv4.3.1-GFED4 in the attached Fig. 3 and 4.

Line 357: "The total calculated ammonia emissions": which one?

Response: We have amended this sentence (Track Changes at circa L.641-642).

Line 360-363: could you please provide statistics (average and standard deviation) for South American and European emissions as well as for the global budget?

Response: We have amended this part. Numbers have been added everywhere in this paragraph presenting average and sd (see Track Changes at L.491-496, p. 13).

Line 363-364: "Based upon IASI retrievals, Liu et al. (2019) showed an increase of surface NH<sub>3</sub> concentrations trend of more than 0.2 μgNm<sup>-3</sup>yr<sup>-1</sup>": I do not understand the link with the previous sentence.

Response: We consent with this comment and we have removed this sentence thanks to the reviewer (see Track Changes at circa L.496, p. 13).

Line 365: "Ammonia emissions derived over China in this work are among the highest worldwide (Figure S1)": is this already the case in the EDGAR and EGG bottom-up inventories or is this a new feature?

Response: We have made clear that by saying "in this work" we mean the emissions highlighted as NE (see Track Changes at circa L.496, p. 13).

Line 370: please precise "The comparison of the annual ammonia NE emissions. . ." In general, you should specify the inventory or the sensitivity test you are referring to,

C10

it would help for the reading and for the understanding of the study.

Response: The reviewer is again right here. We have modified the sentence as follows: “The comparison of the annual ammonia emissions in the NE dataset to the ...” (see Track Changes, L.509, p.14).

Line 377: I would add “in these regions” at the end of the sentence. Indeed, the impact of the different lifetimes seems to be slight over the other regions of the world.

Response: Corrected as suggested by the reviewer (see Track Changes, L.517, p.14)

Line 385-386: is this contradictory with the sentence “European emissions are practically identical in all datasets” in line 361?

Response: We have modified the sentence as follows “. . . in all datasets except EGG . . .” (see Track Changes, L.493, p.13).

Line 460: consist in?

Response: We have corrected as suggested (see Track Changes, L.601, p.17).

Line 461-470: The description of the different inventories and of the different performed simulations should occur before in the text. I would have placed this paragraph at the end of the introduction.

Response: The sequence of the paper is (a) proof that modelled lifetimes are realistic, (b) presentations of the different emission inventories for NH<sub>3</sub> based on different methodologies, (c) comparison with state of the art datasets (ECLIPSE-GFED-GEIA, EDGAR-GFED) that are frequently used to simulate NH<sub>3</sub> concentrations in global models. Then, we need to prove that the emissions presented in the paper produced more realistic modelled concentrations, and for this reason, we simulate NH<sub>3</sub> using each of the different emissions and compare model concentrations with surface measurements and satellite data. We explain all these in an introductory paragraph in discussions. However, we agree with the reviewer and have moved the part that explains what the

C11

EGG emissions refer to into the place that appear for the first time. Instead, we use abbreviations everywhere in this paragraph.

Figure 4: you should number the different graphs. It would be easier to reference them in the text. Please better describe the NH<sub>3</sub> emission dataset in the legend.

Response: We do not really use the numbering in any part of the text when refer to this figure. This is mainly done because each graph placed in any row shows exactly the same thing for different continental regions. We do not think this is necessary (since it's not used) and if the reviewer/editor insists, we may do so in a next step.

Line 532: there is an empty bracket.

Response: Bracket has now been removed (see Track Changes, p.19).

Section 4.2: Does the evaluation against CrIS done at the global scale? It is not specified. If it is the case, it is not comparable with the surface evaluation done at the regional scale. It would be very interested to do it also at regional scale for the analysis, as in Figure 5, 6 and 7 and particularly over hot-spots as explained in the major comments.

Response: Yes, the comparison with CrIS NH<sub>3</sub> refers to global data, which we now specify in L. 677-678. As we already answered in a previous comment, an evaluation of IASI ammonia for several hotspot regions has been done in Van Damme et al. (2018) Nature paper. What we do here is to use IASI NH<sub>3</sub> to produce emissions and see if a model that participates in CMIP and IPCC simulations can improve its performance, also giving these emissions to be used by anyone interested. We evaluate the modelled concentrations against ground measurements that we trust more, in general for N. America, Europe and Southeastern Asia. As a supplement we compare with another global product (CrIS), to prove that concentrations are better reproduced, not only in N. America, Europe and Southeastern Asia, but in a global scale.

Line 599: the word “already” is misplaced in the sentence.

C12

Response: “already” should be “although” in this sentence. We thank the reviewer for pointing this out. We have amended the sentence (see Track Changes, L.736, p.21).

Figure 9: the colors of the scale should be changed: when the uncertainty is high, the borders on the map are not clearly visible.

Response: We have used another colormap as suggested by the reviewer, in order to have visible coastlines (see manuscript with Track Changes).

Line 612: what are the regions with “changing balance between nitrate and sulfate abundances”? Please detail in the text.

Response: We have amended the sentence to make a more concrete statement as suggested (see Track Changes as L. 751-753, p. 21).

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2020-1008>, 2020.

C13






Fig. 1. Seasonal emissions of  $NH_3$  in northwestern European countries.

C14



**Fig. 2.** Left column: SO<sub>2</sub>, NO<sub>2</sub> and SO<sub>4</sub> anomalies after 2015, as in the manuscript. Right column: same anomalies calculated after 2012, when the first reductions of these precursors were observed.

C15



**Fig. 3.** Annual emissions of NH<sub>3</sub> in ECLIPSEv5-GFED4-GEIA.

C16

EDGARv4.3.1-GFED4




Fig. 4. Annual emissions of NH<sub>3</sub> in EDGARv4.3.1-GFED4.