


## Supplemental Information


### Source Apportionment of Fine Aerosol by application of a Chemical Mass Balance (CMB) Model at an Urban site of Beijing

Jingsha Xu, Di Liu, Xuefang Wu, Tuan V. Vu, Yanli Zhang, Pingqing Fu, Yele Sun, Weiqi Xu, Bo Zheng, Roy M. Harrison, Zongbo Shi




**Figure S1.** Stacked bar plots of the concentrations of PM<sub>2.5</sub> components. **Abbreviations:** OC: organic carbon; EC: elemental carbon; PAH: polycyclic aromatic hydrocarbon; SOA: secondary organic aerosol. “Metals” is the summed concentrations of Al, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, Cd, Sr, Sb, Sn, Ba, Pb; “PAH” is the summed concentrations of phenanthrene, anthracene, fluoranthene, acenaphthylene, acenaphthene, fluorene, pyrene, benzo(a)anthracene, chrysene, benzo(b)fluorene, benzo(b)fluoranthene, benzo(k)fluoranthene, benzo(a)pyrene, benzo(e)pyrene, perylene, indeno(1,2,3-cd)pyrene, dibenzo(a,h)anthracene, benzo(ghi)perylene, coronene, picene and retene; “n-alkane” is the summed concentrations of C24, C25, C26, C27, C28, C29, C30, C31, C32, C33, C34; “cooking markers” is the summed concentrations of palmitic acid, stearic acid, cholesterol; “vehicle markers” is the summed concentrations of 17a(H)-22,29,30-trisnorhopane and 17b(H),21a(H)-norhopane; “SOA” is the summed concentrations of 2-methylthreitol, 2-methylerythritol, 2-methylglyceric acid, cis-2-methyl-1,3,4-trihydroxy-1-butene, 3-methyl-2,3,4-trihydroxy-1-butene, trans-2-methyl-1,3,4-trihydroxy-1-butene, 3-hydroxyglutaric acid(3-HGA), Pinic acid, Pinonic acid, C5-alkene triols, 2-methyltetrols, 3-MBTCA (3-Methyl-1,2,3-butanetricarboxylic Acid), beta-caryophyllinic acid, 3-acetylpentanedioic acid, 3-


acetylhexanedioic acid, 3-isopropylpentanedioic acid, DHOPA (2,3-dihydroxy-4-oxopentanoic acid) and 3-Hydroxy-4,4-dimethylglutaric acid.



**Figure S2.** Regression analysis of gravimetrically measured  $PM_{2.5}$  (offline  $PM_{2.5}$ ) and online  $PM_{2.5}$  in winter and summer at IAP, Beijing



**Figure S3.** Regression results between reconstructed  $PM_{2.5}$  and offline/online  $PM_{2.5}$  by chemical mass closure method in winter excluding outliers.



**Figure S4.** Regression results between inorganics (reconstructed  $PM_{2.5}$  excluding OM) and OM with offline- $PM_{2.5}$  by chemical mass closure method in winter excluding outliers.

**Table S1.** Concentrations (mean±SD (min-max),  $\mu\text{g m}^{-3}$ ) of chemical components in  $\text{PM}_{2.5}$  by applying chemical mass closure method

Component	Winter	Summer
$\text{NO}_3^-$	12.49±9.38 (0.87-34.63)	7.16±4.94 (1.53-26.12)
$\text{SO}_4^{2-}$	8.53±7.05 (1.27-24.21)	6.87±3.96 (1.96-19.48)
$\text{NH}_4^+$	8.24±5.63 (0.50-22.62)	3.62±3.14 (0.08-14.83)
$\text{Cl}^-$	3.80±2.32 (0.00-8.73)	0.47±0.42 (0.12-1.96)
$\text{K}^+$	1.30±1.01 (0.15-3.80)	0.37±0.35 (0.11-2.05)
$\text{Na}^+$	0.42±0.24 (0.09-0.93)	0.20±0.16 (0.03-0.73)
Geological minerals	5.33±3.25 (1.03-12.72)	3.52±1.79 (0.65-6.99)
Total other elements	0.53±0.35 (0.12-1.29)	0.35±0.20 (0.07-1.00)
EC	3.5±2.0 (0.3-6.6)	0.9±0.4 (0.2-1.7)
OM	37.7±21.5 (6.9-85.4)	12.9±4.7 (3.6-25.4)
Bound water-Offline	4.0±3.7 (0.2-13.4)	2.8±1.4 (0.8-7.3)
Reconstructed $\text{PM}_{2.5}$ -Offline	83.4±53.6 (14.0-202.1)	39.4±13.0 (19.6-80.0)
Offline $\text{PM}_{2.5}$	88.6±63.6 (10.3-239.9)	30.0±12.7 (12.2-66.7)
Bound water-Online	4.5±4.1 (0.3-14.9)	3.5±1.8 (0.9-9.1)
Reconstructed $\text{PM}_{2.5}$ -Online	83.8±53.9 (14.0-203.7)	40.1±13.4 (19.8-81.8)
Online $\text{PM}_{2.5}$	99.7±79.1 (8.1-328.7)	36.4±14.9 (13.7-78.1)

Note: 5 samples in winter and 7 samples in summer were not included for the calculation of bound water and reconstructed  $\text{PM}_{2.5}$  due to insufficient ions and  $\text{NH}_3$  data. These samples were excluded for the calculation of average online and offline  $\text{PM}_{2.5}$  in this table as well for better comparison.

**Table S2.** Annual average primary OC emissions (Unit: tonne) in Beijing from the 2016 and 2017 Multi-resolution Emission Inventory for China (MEIC)

Sector	Fuel	2016	2017
Power	Coal	0	0
Power	Oil	0	0
Power	Natural Gas	0	0
Industry	Coal	0	0
Industry	Oil	538	583
Industry	Natural Gas	0	0
Industry	Process	1161	1083
Residential	Coal	6687	4312
Residential	Oil	24	23
Residential	Natural Gas	0	0
Residential	Biofuel	5548	4993
Transportation	Oil	1059	1026
Total		15017	12020

**Table S3.** The ratios of OC/PM<sub>2.5</sub> (or OM/OC) for different sources

	OC/PM <sub>2.5</sub>	Reference
Straw burning	0.546	(Zhang et al., 2007)
Wood burning	0.3	(Wang et al., 2009)
Cooking	1.4 (OM/OC)	(Zhao et al., 2007)
Light-duty gasoline cars	0.317	
Heavy-duty gasoline cars	0.549	(Cai et al., 2017)
Diesel cars	0.569	
Vegetative detritus	2.1 (OM/OC)	(Bae et al., 2006b)
Anthracite	0.446	
Bituminite	0.403	(Zhang et al., 2008)
Coal briquette	0.432	
Secondary organic aerosol	2.17 (OM/OC)	(Bae et al., 2006a)
Oxygenated OA	2.2 (OM/OC)	(Zhang et al., 2005)
Oxygenated OA	1.85 ~ 2.3 (OM/OC)	(Aiken et al., 2008)

**Table S4.** Reconstructed source contributions and their relative abundance in reconstructed PM<sub>2.5</sub> mass in urban Beijing

	Mass concentrations ( $\mu\text{g m}^{-3}$ )			Mass concentrations/ Reconstructed mass (%)	
	Winter	Summer		Winter	Summer
SNA	30.5±21.8	17.7±10.5	SNA/ RM	34.1±9.7	48.5±11.9
Geological minerals	5.3±3.2	3.5±1.8	Geological minerals/ RM	7.0±3.0	10.4±5.5
Vegetative detritus	0.2±0.2	0.2±0.2	Vegetative detritus/ RM	0.3±0.2	0.7±0.4
Biomass burning	8.9±6.2	0.8±0.9	Biomass burning/ RM	11.0±5.9	2.1±1.5
Gasoline vehicles	4.7±3.6	0.7±0.4	Gasoline vehicles/ RM	6.5±4.8	2.2±1.4
Diesel vehicles	0.9±2.0	0.1±0.3	Diesel vehicles/ RM	1.0±2.0	0.4±0.9
Industrial CC	11.6±9.7	4.3±1.7	Industrial CC/ RM	13.8±7.3	13.2±6.1
Residential CC	6.1±7.3	0.4±0.3	Residential CC/ RM	7.6±5.7	1.4±1.1
Cooking	3.1±3.0	0.9±0.6	Cooking/ RM	3.9±2.5	2.8±1.6
Other OM	11.7±10.8	6.5±3.2	Other OM/ RM	14.8±7.9	18.3±5.9
Reconstructed mass (RM)	83.1±49.0	35.1±15.0			
Offline PM <sub>2.5</sub> mass	105.0±77.4	36.5±17.0	Reconstructed mass/ Offline PM <sub>2.5</sub> mass	96.6±17.6	121.7±26.6
Online PM <sub>2.5</sub> mass	94.8±64.4	30.2±14.8	Reconstructed mass/ Offline PM <sub>2.5</sub> mass	91.9±24.1	99.0±19.1

## References

- Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC Ratios of Primary, Secondary, and Ambient Organic Aerosols with High-Resolution Time-of-Flight Aerosol Mass Spectrometry, *Environ. Sci. Technol.*, 42, 4478-4485, 10.1021/es703009q, 2008.
- Bae, M.-S., Demerjian, K. L., and Schwab, J. J.: Seasonal estimation of organic mass to organic carbon in PM<sub>2.5</sub> at rural and urban locations in New York state, *Atmospheric Environment*, 40, 7467-7479, <https://doi.org/10.1016/j.atmosenv.2006.07.008>, 2006a.
- Bae, M.-S., Schauer, J. J., and Turner, J. R.: Estimation of the Monthly Average Ratios of Organic Mass to Organic Carbon for Fine Particulate Matter at an Urban Site, *Aerosol Sci. Technol.*, 40, 1123-1139, 10.1080/02786820601004085, 2006b.
- Cai, T., Zhang, Y., Fang, D., Shang, J., Zhang, Y., and Zhang, Y.: Chinese vehicle emissions characteristic testing with small sample size: Results and comparison, *Atmospheric Pollution Research*, 8, 154-163, <https://doi.org/10.1016/j.apr.2016.08.007>, 2017.
- Wang, Q., Shao, M., Zhang, Y., Wei, Y., Hu, M., and Guo, S.: Source apportionment of fine organic aerosols in Beijing, *Atmos. Chem. Phys.*, 9, 8573-8585, 10.5194/acp-9-8573-2009, 2009.
- Zhang, Q., Worsnop, D. R., Canagaratna, M. R., and Jimenez, J. L.: Hydrocarbon-like and oxygenated organic aerosols in Pittsburgh: insights into sources and processes of organic aerosols, *Atmospheric Chemistry And Physics*, 5, 3289-3311, 10.5194/acp-5-3289-2005, 2005.
- Zhang, Y.-x., Shao, M., Zhang, Y.-h., Zeng, L.-m., He, L.-y., Zhu, B., Wei, Y.-j., and Zhu, X.-l.: Source profiles of particulate organic matters emitted from cereal straw burnings, *Journal of Environmental Sciences*, 19, 167-175, [https://doi.org/10.1016/S1001-0742\(07\)60027-8](https://doi.org/10.1016/S1001-0742(07)60027-8), 2007.
- Zhang, Y., Schauer, J. J., Zhang, Y., Zeng, L., Wei, Y., Liu, Y., and Shao, M.: Characteristics of Particulate Carbon Emissions from Real-World Chinese Coal Combustion, *Environ. Sci. Technol.*, 42, 5068-5073, 10.1021/es7022576, 2008.
- Zhao, Y., Hu, M., Slanina, S., and Zhang, Y.: Chemical Compositions of Fine Particulate Organic Matter Emitted from Chinese Cooking, *Environ. Sci. Technol.*, 41, 99-105, 10.1021/es0614518, 2007.