

Atmospheric conditions and composition that influence PM_{2.5} oxidative potential in Beijing, China

Steven J. Campbell^{1,2*}, Kate Wolfer^{1*}, Battist Uttinger¹, Joe Westwood², Zhi-hui Zhang^{1,2}, Nicolas Bukowiecki¹, Sarah S. Steimer^{2§}, Tuan V. Vu^{3#}, Jingsha Xu³, Nicholas Straw⁴, Steven Thomson³, Atallah Elzein⁵, Yele Sun⁶, Di Liu^{3,6}, Linjie Li⁶, Pingqing Fu⁸, Alastair C. Lewis^{5,7}, Roy M. Harrison^{3†}, William J. Bloss³, Miranda Loh⁹, Mark R. Miller⁴, Zongbo Shi³ and Markus Kalberer^{1,2}

¹Department of Environmental Sciences, University of Basel, Basel, Switzerland

²Department of Chemistry, University of Cambridge, Cambridge, UK

10 ³School of Geography Earth and Environmental Sciences, University of Birmingham, Birmingham, UK

⁴Centre for Cardiovascular Science, Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK

⁵Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York, UK

⁶State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

15 ⁷National Centre for Atmospheric Science, University of York, York, UK

⁸Institute of Surface Earth System Science, Tianjin University, Tianjin, China

⁹Institute of Occupational Medicine, Edinburgh, UK

§ Now at: Department of Environmental Science, Stockholm University, Stockholm, Sweden

20 †Also at: Department of Environmental Sciences / Center of Excellence in Environmental Studies, King Abdulaziz University, PO Box 80203, Jeddah, 21589, Saudi Arabia

#Now at School of Public Health, Imperial College London, London, UK

*Authors contributed equally to the manuscript

25

Correspondence to: stevenjohn.campbell@unibas.ch

Abstract. Epidemiological studies have consistently linked exposure to PM_{2.5} with adverse health effects. The oxidative potential (OP) of aerosol particles has been widely suggested as a measure of their potential toxicity. Several acellular chemical assays are now readily employed to measure OP, however, uncertainty remains regarding the atmospheric conditions and specific chemical components of PM_{2.5} that drive OP. A limited number of studies have simultaneously utilised multiple OP assays with a wide range of concurrent measurements and investigated the seasonality of PM_{2.5} OP. In this work, filter samples were collected in winter 2016 and summer 2017 during the atmospheric pollution and human health in a Chinese megacity (APHH-Beijing) campaign, and PM_{2.5} OP was analysed using four acellular methods; ascorbic acid (AA), dithiothreitol (DTT), 2-7-dichlorofluorescein/hydrogen peroxidase (DCFH) and electron paramagnetic resonance spectroscopy (EPR). Each assay reflects different oxidising properties of PM_{2.5}, including particle-bound ROS (DCFH), superoxide radical production (EPR) and catalytic redox chemistry (DTT/AA), and combination of these four assays provided a detailed overall picture of the oxidising properties of PM_{2.5} at a central site in Beijing. Positive correlations of OP (normalised per volume of air) of all four

30

35

assays with overall PM_{2.5} mass was observed, with stronger correlations in winter compared to summer. In contrast, when OP assay values were normalised for particle mass, days with higher PM_{2.5} mass concentrations ($\mu\text{g m}^{-3}$) were found to have lower mass-normalised OP values as measured by AA and DTT. This finding supports that total PM_{2.5} mass concentrations alone may not always be the best indicator for particle toxicity. Univariate analysis of OP values and an extensive range of additional measurements, 107 in total, including PM_{2.5} composition, gas phase composition and meteorological data, provided detailed insight into the chemical components and atmospheric processes that determine PM_{2.5} OP variability. Multivariate statistical analyses highlighted associations of OP assay responses with varying chemical components in PM_{2.5} for both mass- and volume-normalised data. AA and DTT assays were well predicted by a small set of measurements in multiple linear regression (MLR) models, and indicated fossil fuel combustion, vehicle emissions and biogenic SOA as influential particle sources in the assay response. Mass MLR models of OP associated with compositional source profiles predicted OP almost as well as volume MLR models, illustrating the influence of mass composition on both particle-level OP and total volume OP. Univariate and multivariate analysis showed that different assays cover different chemical spaces, and through comparison of mass- and volume-normalised data we demonstrate that mass-normalised OP provides a more nuanced picture of compositional drivers and sources of OP compared to volume-normalised analysis. This study constitutes one of the most extensive and comprehensive composition datasets currently available, and provides a unique opportunity to explore chemical variations in PM_{2.5} and how they affect both PM_{2.5} OP and the concentrations of particle-bound ROS.

1 Introduction

Large-scale epidemiological studies have consistently linked the exposure of airborne particulate matter (PM) with a range of adverse human health effects (Hart et al., 2015; Laden et al., 2006; Lepeule et al., 2012). A recent study by the World Health Organisation estimated that 1 in 8 deaths globally in 2014 were linked to air pollution exposure (World Health Organisation, 2016) with urban areas in India and China particularly affected (Lelieveld et al., 2020). However, large uncertainty remains regarding the physical and chemical characteristics of PM that result in adverse health outcomes upon exposure (Bates et al., 2019).

Studies have suggested that oxidative stress promoted by PM components *in vivo* could be a key mechanism that results in adverse health outcomes (Donaldson and Tran, 2002; Knaapen et al., 2004; Øvreivik et al., 2015). Oxidative stress occurs when excess concentrations of reactive oxygen species (ROS) overwhelm cellular anti-oxidant defences, resulting in an imbalance of the oxidant-antioxidant ratio in favour of the former, which can subsequently lead to inflammation and disease (Knaapen et al., 2004; Li et al., 2003, 2008). The term ROS typically refers to H₂O₂, in some cases including organic peroxides, the hydroxyl radical ($\cdot\text{OH}$), superoxide ($\text{O}_2^{\cdot-}$) and organic oxygen-centred radicals. Particle-bound ROS is exogenously delivered into the lung through PM inhalation, or can be produced *in vivo via* redox-chemistry initiated by certain particle components, in addition to baseline tissue ROS produced by metabolic processes (Dellinger et al., 2001). The capability of PM to produce ROS with subsequent depletion of anti-oxidants upon inhalation is defined as oxidative potential (OP) (Bates et al., 2019).

70 OP is a fairly simple measure of PM redox activity, but reflects a complex interplay of particle size, composition and chemistries which induce oxidative stress by free radical generation which triggers cellular signal transduction and damage. These effects can be both localised (to lung epithelial surfaces and alveoli, reviewed by Tao et al., 2003) and systemic, through immune system activation and cytokine release (Miyata and van Eeden, 2011), translocation of ultrafine particles into the circulatory system (Oberdorster et al., 1992), increased circulating monocytes (Tan et al., 2000), and propagation to other cells
75 and organs (Laing et al., 2010; Meng and Zhang, 2006). Oxidative stress is implicated in the majority of toxicological effects related to air pollution (Ghio et al., 2012; Kelly, 2003; Pope and Dockery, 2006; Risom et al., 2005). A rapid and simple metric to capture the oxidative exposure burden which can be easily implemented for epidemiological studies will enable greater insight into the mechanisms of PM toxicity beyond total PM mass concentrations alone.

There are now a wide range of acellular chemical methods that attempt to quantify particle-bound ROS and the entire OP of
80 PM, as typically acellular assays allow faster measurement and are less labour intensive compared to cell cultures or *in vivo* methods (Bates et al., 2019). These include, but are not limited to, the dithiothreitol assay (DTT), ascorbic acid assay (AA), 2-7-dichlorofluorescein/hydrogen peroxidase assay (DCFH), electron paramagnetic spectroscopy (EPR), glutathione assay (GSH) and 9-(1,1,3,3-tetramethylisindolin-2-yl)oxyl-5-ethynyl-10-(phenylethynyl)anthracene (BPEAnit). These acellular assays all have differing sensitivities to specific particle components that may contribute to increased particle-bound ROS concentrations
85 and aerosol OP. For instance, DTT has been shown to be sensitive to soluble metals (Shinyashiki et al., 2009), including copper and manganese (Charrier et al., 2015; Charrier and Anastasio, 2012), as well as a range of organic particle components including water soluble organic carbon (WSOC, a mixture of 100's to 1000's of compounds), oxidised polycyclic aromatic hydrocarbons (PAHs) e.g. quinones (Chung et al., 2006; McWhinney et al., 2013a), and humic-like substances (HULIS) (Dou et al., 2015; Verma et al., 2015a). AA is particularly sensitive to redox-active transition metals, most notably Fe (Godri et al.,
90 2011) and Cu (Janssen et al., 2014; Pant et al., 2015), and has demonstrated sensitivity to organic carbon (Calas et al., 2018) including secondary organic aerosol (Campbell et al., 2019a). EPR is applied to speciate and quantify radical species either bound to aerosol particles (Arangio et al., 2016; Campbell et al., 2019b; Gehling and Dellinger, 2013), so-called environmentally persistent free radicals (EPFR), or radicals formed upon suspension of particles into aqueous solution (Gehling et al., 2014; Tong et al., 2016, 2017) or in some cases into synthetic lung lining fluid (Tong et al., 2018) consisting
95 of a mixture of AA, glutathione and uric acid. EPR has the advantage of not being influenced by the dark colour of particulate suspensions (detection is *via* magnetic excitation rather than magnetic absorbance), does not require extraction of the PM from the filter, and that speciation of the free radical generated can be explored using spin-trap reagents that are selective for specific radicals (Miller et al., 2009). The DCFH assay has been shown to be particularly sensitive to hydrogen peroxide (H₂O₂) and organic peroxides (Venkatachari and Hopke, 2008; Wragg et al., 2016), also present in secondary organic aerosol (SOA)
100 particles (Gallimore et al., 2017), and is a particularly useful assay for measuring particle-bound ROS (Wragg et al., 2016). The application of these four commonly used assays simultaneously allows different mechanisms of ROS generation to be assessed; the variability of particle-bound ROS (DCFH), the production of superoxide upon aqueous particle suspension (EPR) and the catalytic generation of ROS via redox-active components (DTT/AA). Therefore, these data provide a broad picture of

the variability of both particle-bound ROS and OP, and comparison to a comprehensive compositional dataset provides a
105 unique opportunity to probe the chemical changes in PM that affect the burden of particle-bound ROS and OP.

Despite several studies utilising the aforementioned assays, further exploratory work is required to determine specifically
which sources, physical properties and chemical components influence aerosol OP variability. A limited number of studies
have explored the role of chemical composition on aerosol OP, and it is often unclear which specific chemical components are
responsible for driving aerosol OP; for example, studies show transition metals such as Cu and Mn dominate DTT activity
110 (Charrier et al., 2015; Charrier and Anastasio, 2012), whereas others highlight the enhanced role of organics, in particular
water soluble organic carbon (WSOC) such as HULIS, and quinones (Cho et al., 2005; Fang et al., 2016). Furthermore, several
studies correlate volume-normalised OP measurements with compositional variability, but given the potential co-linearity of
many aerosol components with overall mass, mass-normalised intrinsic OP values may provide additional insight into the
effect of chemical composition on aerosol OP (Bates et al., 2019; Puthussery et al., 2020). Thus, a comprehensive
115 characterisation of gaseous and particle phase pollution conditions combined with measurements utilising multiple OP assays
simultaneously provides a wide range of information on particle-bound ROS and aerosol OP, allowing the identification of the
most important components that drive aerosol OP. Ultimately, a greater understanding of the specific aerosol characteristics
that influence OP, as well as specific sources that contribute more to aerosol OP, could allow the development of more targeted
and efficient air pollution mitigation strategies. Further details of the selection of OP assays, their scope and biological
120 applicability are described in **Section S2** of the Supplementary Information.

In this work, PM_{2.5} filter samples collected in winter 2016 and summer 2017 during the APHH campaign were analysed using
four acellular methods; AA, DCFH, DTT and EPR, providing a wealth of data on the health-relevant properties of PM_{2.5}
including particle-bound ROS, redox-active components contributing to aerosol OP, and the formation of superoxide radicals
upon sample extraction. As the APHH campaign simultaneously captured one of the most extensive and comprehensive
125 datasets atmospheric composition datasets including a broad range of PM compositional data, we aimed to establish what
individual PM components, meteorological and atmospheric conditions contributed to increased OP assay response, whether
these influences and compositions differed substantially between assays, and if the compositions reflected particular PM
sources. We included 107 different measurements, comprising transition metals, AMS measurements, total elemental and
organic carbon, and a broad panel of organic species relating to biomass and fossil fuel burning, cooking emissions, vehicular
130 markers, secondary organic aerosol compounds, plus gaseous species and general atmospheric conditions. We sought to
investigate the differences between volume-based and mass-based responses, as mass-based analysis facilitates site and
temporal comparisons more readily than volume measurements and provide details on intrinsic particle properties that
influence OP. In order to highlight underlying trends in such a broad and complex dataset, we also applied multivariate
statistical analysis and developed multiple linear regression models to fully characterise the chemical compositional factors
135 driving each assay response.

2 Materials and methods

2.1 Air Pollution and Human Health in a Chinese Megacity Campaign (APHH)

2.1.1 Site description

140 High-volume 24 hr aerosol filter samples were collected at the Institute of Atmospheric Physics (IAP) in Beijing, China (39°58'28" N, 116°22'15" E) (**Figure S1**). Winter PM was collected during the months of Nov-Dec 2016 and summer PM was collected during the months of May-June 2017. $n = 31$ filters for winter 2016 and $n = 34$ filters for summer 2017 were collected. A PM_{2.5} high-volume sampler (RE-6070VFC, TICS, USA) was used at a flow rate of $\sim 1.06 \text{ m}^3/\text{min}$. PM_{2.5} for subsequent OP analysis was collected onto quartz microfiber filters (Whatman, 20.3 x 25.4 cm) with a collection area of 405 cm^2 .

2.1.2 PM_{2.5} composition, gas phase composition and meteorological data

145 Oxidative potential measurements were correlated with a range of additional particle phase composition, gas phase composition and meteorological measurements conducted concurrently during the APHH-Beijing campaign (Shi et al., 2019). Briefly, the following composition data was collated: total organic and elemental carbon (OC, EC), soluble inorganic ions (K^+ , Na^+ , Ca^{2+} , NH_4^+ , NO_3^- , SO_4^{2-} and Cl^-) measured using ion chromatography (IC), low-oxidised organic aerosol and more-oxidised organic aerosol (LOOOA/MOOOA) fractions using aerosol mass spectrometry (AMS), biomass burning markers
150 (galactosan, mannosan and levoglucosan), 16 polycyclic aromatic hydrocarbons (PAHs) (see (Elzein et al., 2019, 2020)), C₂₄-C₃₄ *n*-alkanes, aerosol cooking markers (palmitic acid, stearic acid, cholesterol), vehicle exhaust markers (17a(H)-22, 29,30-trisnorhopane (C27a) and 17b(H)-21a-norhopane (C30ba)), isoprene SOA markers (2-methylglyceric acid, 2-methylerythritol, 2-methylthreitol, 3-hydroxyglutaric acid), C₅-alkene triols (cis-2-methyl-1,3,4-trihydroxy-1-butene, 3-methyl-2,3,4-trihydroxy-1-butene, trans-2-methyl-1,3,4-trihydroxy-1-butene), α -pinene SOA tracers (cis-pinonic acid, pinic acid, 3-methyl-
155 1,2,3-butanetricarboxylic acid (MBTCA), 2,3-dihydroxy-4-oxopentanoic acid, aged α -pinene SOA marker), β -caryophyllene SOA tracer (β -caryophyllinic acid) and an aromatic volatile organic compound (VOC) SOA tracer (3-isopropylpentanedioic acid) (Liu et al., 2020). The following additional data was obtained from the Centre for Environmental Data Analysis (CEDA) archive: concentrations of inorganic elements Al, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, Cd, Sb, Ba and Pb in PM_{2.5} using X-ray fluorescence (XRF) (Xu et al., 2020a), gas phase concentrations of methanol, acetonitrile, acetaldehyde, acrolein, acetone,
160 isoprene, methacrolein, methyl ethyl ketone, benzene, toluene, C₂-benzenes and C₃-benzenes measured using proton transfer reaction time-of-flight mass spectrometry (PTR-ToF-MS) (Acton et al., 2018), gas phase concentrations of O₃, CO, NO, NO₂, NO_y and SO₂ as well as relative humidity (RH) and air temperature measurements (Shi et al., 2019), photolysis rates for J O¹D and J NO₂ (Whalley et al., 2020) and gas phase concentrations of hydroxyl radicals (OH), peroxy radicals (HO₂) and organic peroxy radicals (RO₂) measured using fluorescence assay gas expansion (FAGE) (Whalley et al., 2020).

165 2.2 Oxidative potential measurements

2.2.1 Reagents

Chemicals and gases were obtained from Sigma-Aldrich unless otherwise indicated and were used without further purification: ascorbic acid ($\geq 99.0\%$), ChelexTM 100 sodium form, 0.1 M HCl solution, 0.1 M NaOH solution, dichlorofluorescein-diacetate (DCFH-DA), 1 M potassium phosphate buffer solution, horseradish peroxidase (HRP), methanol (HPLC grade), and *o*-phenylenediamine ($\geq 99.5\%$). H₂O used for the DCFH, HRP and AA solution were obtained from a Milli-Q high purity water unit (resistivity $\geq 18.2\text{ M}\Omega\text{ cm}^{-1}$, Merck Millipore, USA). For DTT analysis, 9,10-phenanthrenequinone (PQN) ($\geq 99\%$), 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB) (99%), DL-dithiothreitol (DTT) ($\geq 98\%$), potassium phosphate dibasic ($\geq 98\%$, Krebs buffer), potassium phosphate monobasic ($\geq 98\%$, Krebs buffer), and methanol ($\geq 99.9\%$) were all obtained from Fisher Chemical. Nitrogen (oxygen free) was obtained from BOC (Cambridge, UK).

175 2.2.2 Acellular oxidative potential assays

Four offline acellular methods for measuring PM_{2.5} oxidative potential and particle-bound ROS were utilised in this work; The DCFH/HRP assay (Fuller et al., 2014), which quantifies the fluorescent product 2,7-dichlorofluorescein, an assay that is particularly sensitive to species which are likely particle-bound ROS, the ascorbic acid (AA) assay (Campbell et al. (2019)) which quantifies the dominant product of AA oxidation, dehydroascorbic acid (DHA) *via* condensation with a dye and
180 fluorescence spectroscopy. This is an AA-only assay and does not contain other components normally present in synthetic lung fluid (SLF); filter extracts are performed at pH 7, whereas the AA reaction with the filter extract is performed at pH 2 to improve assay stability and sensitivity (Campbell et al. (2019)). Electron Paramagnetic Resonance spectroscopy (EPR) (Miller et al., 2009) specifically targets the measurement of superoxide ($\text{O}_2^{\cdot-}$) and the dithiothreitol (DTT) assay (e.g. Cho et al., 2005) quantifies the rate of loss of DTT via absorbance measurements. These acellular methods have been widely applied in the
185 literature to study particle OP and particle-bound ROS (Bates et al., 2019). For detailed descriptions of the assay protocols, see Section S3 in the Supplementary Information. Assessing OP and particle-bound ROS in filters with the aforementioned assays is performed offline. There is potential to underestimate PM OP and particle-bound ROS using offline filter-based analysis, as short-lived components which contribute to particle-bound ROS and OP may undergo degradation prior to analysis. However, using an offline method allows the opportunity to correlate with a wide range of additional composition
190 measurements, allowing a more explicit characterisation of the chemical components of PM that contribute to observed acellular assay responses.

2.3 Statistical analysis

We aimed to analyse the data as thoroughly as possible with respect to characterising the OP measured by each assay, and to attempt to robustly connect assays to both individual measurements and potential PM sources. As data were collated from

195 several different experimental projects, and as analytical uncertainty values were not available for the majority of the data, the use of positive matrix factorisation (PMF) was not undertaken for source apportionment, and will be published subsequently for selected analyses (Xu et al., 2020a). Multiple analytical platforms were used for the acquisition of compositional data, uncertainty estimates for each measurement were not easily estimable, a factor-based chemical mass balance approach was not required specifically, and temperature, relative humidity, actinic flux and other non-mass measurements could also be
200 influential on the OP response, and are factors mainly independent of PM sources. On this basis we considered that PMF would not ultimately give useful models in the specific context of OP. However, these issues are managed adequately by principal components analysis (PCA), which is a useful general unsupervised method for examining underlying variance and latent effects in data, and handles multicollinearity well, although it is not optimal for chemical mass balance source apportionment (Paatero and Tapper, 1994).

205 PCA and partial least squares regression (PLSR) models were produced in SIMCA+ 16.0 (Umetrics, Umeå, Sweden). Missing values were not altered prior to model construction, although measurements with more than 56% missing values per season were discarded from models. R^2 and Q^2 values were used to assess the goodness-of-fit of the model and the goodness-of-prediction of the data through 7-fold cross-validation respectively. Data were unit-variance scaled and mean-centred to remove effects related to absolute data magnitude. Models were allowed to optimise to the maximum number of latent variables (LV)
210 at which the cumulative Q^2 value stabilised, which for most PLSR models was a single LV. PLSR model robustness was assessed through permutation testing, where the classifier (i.e. OP assay response) for all samples was randomly permuted 999 times and the PLSR model constructed for each permutation; the model was considered robust if the real model R^2 and Q^2 values outperformed those from all random permutation models. Negative Q^2 values indicate no predictive power of the data in the model, and LVs with Q^2 significantly lower than the R^2 value (arbitrarily defined for this study as Q^2 at more than 10%
215 below the R^2) can be considered at least partially overfitted.

Spearman rank correlations (R_s) between OP measurements and $PM_{2.5}$ were calculated using OriginPro (2020) or R, and were used to assess the relationships between assay responses and individual measurements, with Mann-Whitney-U tests (in R) used for pairwise testing of the differences in seasonal response for both assays and individual measurements. All other multivariate analyses, multiple linear regression models and selected univariate analyses were produced in R 4.0.2 (R Core
220 Team, Vienna, Austria), implemented in RStudio 1.3.959 (Boston, Massachusetts, USA).

For multiple linear regression models, outlier values were arbitrarily deemed to be those greater than 5 times the standard deviation and replaced with the season median where appropriate for analysis. Measurement subsets manually selected as relevant to source composition were then subjected to a variable selection process, whereby pairwise Spearman correlations for all measurements were calculated, and measurements removed from subsets if they were highly correlated with other
225 measurements but predicted OP more poorly than the other co-correlated measurements, to reduce the number of variables contributing identical information in the final models. Multiple linear regression models were then further optimised from this initial subset using the *regsubsets* function in the *leaps* R package, to allow for between 4-8 variables which best predicted the OP response (models could be constructed with fewer or even more measurements, but the aim was to examine a small panel


of contributors to potential source compositions). The variable selection process precludes the use of linear regression mode
performance indicators such as the Aikake or Bayesian information criteria, as the model component sets are not identical.
The stability of model predictions and features were assessed using bootstrap resampling of data, by randomly splitting one
fifth of the data as a test set and using the remaining samples to construct the model and predict the left-out samples, for 500
random iterations. Stability was also assessed through overall variance in OP predictions, measurement feature coefficients and
model residuals plots, and run order/date bias (not differentiable as samples were analysed in date order) was assessed in
residuals plots. Although not all data distributions were strictly normal when examined in the univariate kernel density plots,
data were not log-transformed for multiple linear regression models, as this creates non-linearity in the model component
response, which can complicate interpretation. Model residuals were plotted for manual examination and were all generally
normally distributed despite the relatively small number of samples, and biases were related to periods of missing
measurements or samples with values below the limit of quantification. Code developed for analysis is publicly available at
<https://github.com/katewolfer/Beijing>.

3 Results and discussion

Both volume-normalised (OP_v , per m^3 air) and particle mass normalised (OP_m , per μg $PM_{2.5}$) values are considered in this
work, where the OP value of the specific assay and sample is normalised by the volume of air collected or by the total $PM_{2.5}$
mass on the filter, respectively. OP_v is useful when considering exposure or epidemiological outcomes, but OP_m is likely a
more informative metric when exploring how chemical composition influences $PM_{2.5}$ OP, and potentially enabling better OP
response, site and composition intercomparisons (Bates et al., 2019). Henceforth, OP_v and OP_m will be used when discussing
the overall response of all four methods; specific discussion of the acellular methods will be referred to as AA_v , DTT_v , $DCFH_v$
and EPR_v for volume-normalised values, and AA_m , DTT_m , $DCFH_m$ and EPR_m for mass-normalised values. For comparison of
mass normalised OP_m values, $PM_{2.5}$ composition measurements were also normalised for total PM mass (*e.g.* $ng/\mu g$ per μg
 $PM_{2.5}$).

3.1 Seasonal variation of OP_m and OP_v

24-hour $PM_{2.5}$ mass concentrations in winter 2016 (08/11/2016-09/12/2016) ranged from $8.1 - 328.7 \mu g m^{-3}$, with an average
 $PM_{2.5}$ mass of $98.7 \pm 75 \mu g m^{-3}$, whereas in summer 2017 (21/05/2017-24/06/2017) $PM_{2.5}$ concentrations ranged of $13.6 - 85$
 $\mu g m^{-3}$ with an average of $36.7 \pm 16 \mu g m^{-3}$ (**Figure S7**) (Shi et al., 2019; Xu et al., 2020a). Average seasonal values for each
assay are summarised in Table S1. An example data set showing 24-hr average data, for AA_v and $PM_{2.5}$ mass in both the winter
and summer campaign, is shown in **Figure 1** (for $DCFH_v$, DTT_v and EPR_v , see **Section S6** “Summary statistics for all
measurements” in the Supplementary Information).



260 **Figure 1.** 24-hour averaged volume-normalised AA_v (red bars) and $PM_{2.5}$ mass (blue dots), analysed from 24-hour high volume filters, for both winter 2016 (08/11/2016 – 08/12/2016) and summer 2017 (21/05/2017 - 24/06/2017) (Shi et al., 2019; Xu et al., 2020a). Substantially higher average $PM_{2.5}$ mass concentrations ($\mu g m^{-3}$) and AA_v were observed in the winter season compared to the summer (see Table S1 for summary). DCFH_v, DTT_v and EPR_v 24-hour averaged datasets can be found in Figures S8-S10 respectively.

For all assays, a higher average $PM_{2.5}$ OP_v was observed in the winter compared to the summer in Beijing (**Table S1**). The average AA_v was 96.7 ± 42.7 nM [DHA] m^{-3} in the winter, whereas a mean value of 24.1 ± 6.1 nM [DHA] m^{-3} was observed in the summer. Given the recent introduction of this AA-based assay, which measures the formation of the AA oxidation product DHA rather than measuring the decay of AA *via* UV absorbance, limited literature values are available for direct comparison (Campbell et al., 2019a). Average DCFH_v in the winter was 0.71 ± 0.52 nmol $H_2O_2 m^{-3}$ compared to 0.17 ± 0.11 nmol $H_2O_2 m^{-3}$ in the summer, which is within the range of DCFH_v values observed in previous studies in Taiwan, the USA and Singapore (OP_{DCFH} 0.02 - 5.7 nmol $H_2O_2 m^{-3}$) (Hasson and Paulson, 2003; Hewitt and Kok, 1991; Hung and Wang, 2001; See et al., 2007; Venkatachari et al., 2005). Mean observed values for DTT_v in the winter and summer were 2.9 ± 0.11 nmol $min^{-1} m^{-3}$ and 0.9 ± 0.40 nmol $min^{-1} m^{-3}$, respectively. The mean values of DTT_v observed in this study are greater than those measured in similar studies in Beijing (Liu et al., 2014) (0.11-0.49, mean = 0.19 nmol $min^{-1} m^{-3}$) with similar mass concentrations of $PM_{2.5}$ (mean = 140 $\mu g m^{-3}$), although they are within the range of DTT_v values observed in a number of

275 previous studies in several locations, including Europe (Jedynska et al., 2017; Yang et al., 2015), the US (Fang et al., 2015; Verma et al., 2014) and Northern China (Liu et al., 2018) ($0.1\text{--}14.7\text{ nmol min}^{-1}\text{ m}^{-3}$). The mean EPR_v values, relating to the specific detection of $\text{O}_2^{\cdot-}$, were $2.4 \times 10^6 \pm 1.6 \times 10^6$ and $5.8 \times 10^5 \pm 4.1 \times 10^6$ counts m^{-3} in the winter and summer campaign, respectively.

Spearman rank correlation coefficients (R_s) of aerosol OP_v with $\text{PM}_{2.5}$ vary between the winter and summer season, and also
280 between OP assays, as illustrated in **Figure 2**. All four assays, when normalised per volume (OP_v), show a stronger correlation with $\text{PM}_{2.5}$ mass concentration in the winter compared to the summer, consistent with results observed in Chamonix, France by Calas *et al.* (Calas et al., 2018) For example, DCFH_v correlates well with 24-hr average total $\text{PM}_{2.5}$ mass concentration ($\mu\text{g m}^{-3}$) in both winter ($R_s = 0.96$) and summer ($R_s = 0.76$) (**Figure 2B**), whereas AA_v correlates well in the winter ($R_s = 0.89$) and poorly in summer ($R_s = 0.21$). Similar correlations of DCFH_v with $\text{PM}_{2.5}$ mass concentrations in both winter and summer
285 suggest that species influencing DCFH_v variability (e.g. H_2O_2 and organic peroxides, likely particle-bound ROS) present in the particles are relatively consistent between both seasons. Similar to AA_v , differences between the seasons are also observed for DTT_v and EPR_v , where correlations of aerosol OP_v vs. $\text{PM}_{2.5}$ are stronger in winter compared to summer (**Figure 2C and 2D**), also generally consistent with previous studies, although in contrast to Calas et al. (2018), who observed no difference in EPR_v between seasons in Chamonix, although in that study the spin trap DMPO was used to study hydroxyl radicals, whereas
290 in this study we focus on the formation of superoxide upon particle suspension in aqueous solution. The differences in the correlation shown in **Figure 2** suggests that the four assays are sensitive to different PM components and that in winter and summer different PM sources or components are important for the assay's responses (Calas et al., 2018; Saffari et al., 2013; Verma et al., 2014). **Figure 2** demonstrates that $\text{PM}_{2.5}$ mass could be a reasonable predictor of total OP_v in winter but the poorer correlations between all OP_v assays and $\text{PM}_{2.5}$ in the summer indicate that a more detailed understanding is necessary
295 to elucidate and ultimately predict aerosol OP. However, the variability in the strength of correlation between OP_v and $\text{PM}_{2.5}$ mass as well as the seasonal difference indicates that compositional differences in $\text{PM}_{2.5}$ or additional atmospheric processes influence $\text{PM}_{2.5}$ OP.




Figure 2. Comparison of PM_{2.5} OP_v during winter 2016 (blue) and summer 2017 (orange) vs. PM_{2.5} mass (μg m⁻³). (a) AA_v, (b) DCFH_v, (c) DTT_v and (d) EPR_v. Each datapoint represents a 24-hour average for OP measurements and PM_{2.5} mass. Corresponding R_s and linear fit equations are included. For AA_v, DCFH_v and DTT_v, error bars represent the standard deviation observed over three repeat measurements for each filter sample, and in some cases the error is smaller than the data point. Uncertainty values are unavailable for EPR_v measurements.

To gain further insights into the potential particle-level compositional differences underlying assay OP response, the OP data for the four assays was normalised to the PM_{2.5} mass in each sample. As shown in **Figure 3**, mass-normalised OP_m values vary up to a factor of ten within a single season. AA_m, DCFH_m, DTT_m and EPR_m for both winter and summer are also displayed in Figure 3, with colour bars indicating the 24-hr average total PM_{2.5} mass (μg m⁻³) for the corresponding OP_m measurement. The average OP_m response observed in this study shows a similar trend to OP_v (**Table S2**), where higher OP_m values are observed for winter compared to summer (Figure 3), as observed previously (Liu et al., 2018; Saffari et al., 2014). This demonstrates that there are specific properties of PM_{2.5} in the winter that result in overall higher intrinsic OP_m compared to the summer.

For AA_m, an inverse relationship between total PM_{2.5} mass concentration and AA_m is observed in both seasons, where days with high PM_{2.5} mass loadings have correspondingly low AA_m values in both the winter and summer, with almost a factor of

6 difference between the AA_m on the highest PM_{2.5} mass day (PM_{2.5} = 328 µg m⁻³, AA_m = 0.6 nM [DHA] µg⁻¹) and lowest PM_{2.5} mass day observed during the winter campaign (PM_{2.5} = 8 µg m⁻³, AA_m = 3.53 nM [DHA] µg⁻¹). A similar trend is observed for DTT_m, where in general days with higher overall PM_{2.5} mass concentrations have correspondingly low DTT_m values, which has also been observed previously (Wang et al., 2020b). The DTT_m response is also not correlated with Cu and Mn concentrations, despite the non-linear but monotonic relationship between these components being demonstrated in other studies (Charrier et al., 2016). These results indicate that on high-pollution days a large fraction of the PM mass might be OP-inactive, resulting in low intrinsic OP_m values. In general, smaller particles have been observed to have higher DTT_m values compared to larger particles (Bates et al., 2019; Janssen et al., 2014), an effect which may also play a role here. Another possibility is that on higher PM_{2.5} mass days, selected chemical species interact with or deactivate redox-active components present in PM_{2.5} (e.g. interaction of organics with metals (Tapparo et al., 2020), therefore reducing the observed OP_m signal. It is also possible that components present in PM_{2.5} on higher PM_{2.5} mass concentration days interfere with the assay response. It is currently unclear which chemical components are responsible for the observed inverse relationship between PM_{2.5} mass with AA_m and DTT_m. However, statistically significant inverse correlations are observed between AA_m and DTT_m in both the winter and summer with the chemically undetermined “unknown” fraction of PM_{2.5} for DTT_m (R_s = -0.81) and AA_m (R_s = -0.75), implying that PM_{2.5} chemical components unaccounted for in this study are likely responsible for the lower intrinsic AA_m and DTT_m values on high PM_{2.5} mass days (See **Section 3.2** “Univariate analysis of PM OP and additional measurements”, **Figure S11** and **Figure S12**).

In contrast, higher DCFH_m responses are observed on days with greater PM_{2.5} mass concentrations in both winter and summer. Increased DCFH_m responses on more polluted days could indicate that the mass fraction of particle-bound ROS (e.g. organic peroxides from SOA) increases with increasing PM_{2.5} mass concentration, or that the capacity of PM components to produce H₂O₂ upon extraction, as measured by DCFH, is enhanced. Despite the significant seasonal difference in EPR_m, no obvious relationship between EPR_m and PM_{2.5} mass was observed in our study. There is potential to underestimate PM OP and particle-bound ROS using offline filter-based analysis, as short-lived components which contribute to particle-bound ROS and OP may undergo degradation prior to analysis. However, using an offline based method allows the opportunity to correlate with a wide range of additional composition measurements, allowing a more explicit characterisation of the chemical components of PM that contribute to observed acellular assay responses.




Figure 3. Summer and winter 24-hour averaged mass-normalised OP_m (a) AA_m (μM DHA μg⁻¹), (b) DCFH_m (nmol H₂O₂ μg⁻¹), (c) EPR_m (counts μg⁻¹) and (d) DTT_m. Box plots indicate the median, 25% and 75% percentiles, and the data range. Data points are colour coded with respect to the 24-hour average PM_{2.5} mass (μg m⁻³), with a separate colour scale for winter and summer PM_{2.5} masses given the difference in total PM_{2.5} masses observed between the seasons.

Spearman rank correlations (R_s) between the four assays, for mass-normalised OP_m and volume-normalised OP_v are presented in **Table 1**. In terms of OP_v, all four assays show significantly strong correlations with each other in the winter season (R_s 0.72 – 0.89), but weaker correlations are observed between assays in the summer (R_s 0.01-0.58), a seasonal difference observed previously by Calas et al. (2018). In contrast, the only statistically significant correlation observed for OP_m is between AA_m and DTT_m in the winter season only (R_s = 0.58).

Seasonality of both OP_v and OP_m observed in the assays could be driven by changes in PM sources influencing overall OP, or a number of physical and chemical factors directly affecting particle composition. For instance, lower ambient temperatures in the winter may increase the partitioning of semi-volatile organic compounds, such as small quinones (e.g. anthracenequinone and 2,3-dimethylantraquinone (Delgado-Saborit et al., 2013)) and nitro-PAHs, which have been shown to influence DTT activity (Ntziachristos et al., 2007; Verma et al., 2011), observations which are supported by lab-based studies showing decreasing aerosol OP at higher temperatures (Biswas et al., 2009; Verma et al., 2011). Changing boundary layer height between the seasons may also contribute to higher concentrations of species which correlate with PM_{2.5} mass responsible for increasing aerosol OP during the winter, compared to summer, especially affecting OP_v seasonality (Wang et al., 2020a). Furthermore, air mass history may play an important role in the observed seasonality of OP. For instance, it was

observed that winter days with high PM_{2.5} mass concentrations typically originate from regional sources south of Beijing, which is widely industrialised, whereas high mass days in the summer typically have more varied air mass histories (Panagi et al., 2020; Steimer et al., 2020). There are likely varying contributions between different sources in different seasons, e.g. more photochemistry in the summer driving oxidation and biogenic sources, and more contributions from residential heating combustion in the winter (Xu et al., 2020a). In order to gain further insight into what causes the observed variability of OP, relationships between particle chemical composition and aerosol OP will be explored in detail below.

Table 1. Correlation of volume-normalised (OP_v, top panel) and mass-normalised (OP_m, bottom panel) assay responses in the winter (blue) and summer (orange) campaign. It should be noted that assay responses expressed as mass-normalised (OP per µg) are correlated with mass-normalised additional particle phase composition measurements (i.e. µg or ng per µg PM_{2.5}).

<u>OP_v R_s</u>	AA _v	DCFH _v	EPR _v	DTT _v
AA _v		0.89***	0.86***	0.83***
DCFH _v	0.35*		0.86***	0.72***
EPR _v	0.19	0.01		0.88***
DTT _v	0.41*	0.58***	0.07	

<u>OP_m R_s</u>	AA _m	DCFH _m	EPR _m	DTT _m
AA _m		-0.29	0.22	0.60**
DCFH _m	-0.20		-0.08	-0.15
EPR _m	-0.26	0.15		0.27
DTT _m	0.2	-0.28	0.14	

Bold indicates R_s ≥ 0.5, *p < 0.05, **p < 0.01, *** p < 0.001.

370

3.2 Univariate analysis of PM OP_m and additional measurements

Spearman rank correlations between OP_m of the four assays and 107 additional measurements conducted during the APHH campaign (see **Section 2.1.2** “PM_{2.5} composition, gas phase composition and meteorological data”), were calculated for both the winter (*n* = 31) and summer (*n* = 33). We focus on OP_m in the forthcoming discussion; as mentioned previously, as we consider it a particularly informative metric when determining the role of chemical composition on OP (Bates et al., 2019; Puthussery et al., 2020) (all univariate statistical summaries are presented in **Section S8**).

The majority of additional particle phase composition, gas phase composition and meteorological measurements differed significantly by season. Exceptions included Al, V, Zn, Pb, Ca²⁺, Na⁺, NH₄⁺, acetaldehyde, acetonitrile, methanol, methyl ethyl ketone, methyl vinyl ketone/methacrolein, trans-2-methyl-1,3,4-trihydroxy-1-butene, β-caryophyllinic acid, 3-hydroxyglutaric

380 acid, C5-alkene triols, cholesterol, LOOOA and MOOOA. Stacked bar plots illustrating the total daily concentrations for both mass-normalised and volume-normalised data are shown in **Figure 4** and **Figure S13**. Total concentrations of individual PM components (excluding all composite measures) account for approximately 0.3-0.8 $\mu\text{g}/\mu\text{g}$, i.e. 30 – 80% of the total PM mass (data not shown). Interestingly there were no marked or characteristic changes in mass composition associated with haze days; however, haze events were generally correlated with increased biomass burning marker concentration and total organic carbon

385 in winter for the mass-normalised data (also observed during recent later winter haze events in Beijing (Li et al., 2019)), and small inorganic ion concentrations in both seasons in the volume-normalised data (Figure S13).

IC measurements (K^+ , Na^+ , Ca^{2+} , NH_4^+ NO_3^- and SO_4^{2-}) account for the greatest proportion of total particle mass in both seasons, all of which are major components of secondary inorganic PM mass (NH_4^+ , NO_3^- , SO_4^{2-}), mineral dust (Ca^{2+} , K^+), and marine aerosols (Na^+ , Cl^-). These species were present at higher daily concentrations in summer than in winter. Summer

390 compositions for each category were generally consistent for the whole sampling period, with a larger total proportion of SOA markers, whereas winter compositions were more variable, with greater contributions from elemental carbon, PAHs, *n*-alkanes and cooking-related compounds than for summer samples. Although PAHs are not redox-active (Charrier and Anastasio, 2012), they are precursors to redox-active oxy-PAHs (quinones) and nitro-PAHs (Atkinson and Arey, 2007), and have well-established intrinsic cellular toxicity (reviewed in Moorthy et al., 2015), mediated by their conversion to hydroxy-PAHs, which

395 exert mutagenic and teratogenic effects, and also inducing transcriptional modifications and oxidative stress. EC and *n*-alkanes are also non-redox-active and the exact mechanism of their toxicities is unclear (Levy et al., 2012); however, SOA derived from the interaction of *n*-alkanes with NO_x with photo-oxidation (Lim and Ziemann, 2005; Presto et al., 2010) is likely both to contribute to the redox activity of samples (Tuet et al., 2017), and to have more toxic properties than its precursors (Xu et al., 2020b). The sample from 22 November 2016 has a particularly high concentration of cooking markers (palmitic acid, stearic acid and cholesterol).

400 This could reflect the fact that the traditional Chinese winter solar term Xiao Xue (小雪, “Light Snow”), begins on this date (Li, 2006), a period associated with the preparation of warm foods as the ambient temperatures in northern China drop; a similar elevation of palmitic acid and stearic acid has been observed around the same week in a more recent study in Shanghai (Wang et al., 2020c).

405





Figure 4. Stacked bar plots of total concentrations for mass-normalised data. **Abbreviations:** OC: organic carbon; EC: elemental carbon; PAH: polycyclic aromatic hydrocarbon; SOA: secondary organic aerosol “Metals” is the summed concentrations of Al, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, Cd, Sb, Ba, Pb; “biomass burning” is the summed concentrations of palmitic acid, stearic acid and cholesterol; “PAH” is the summed concentrations of naphthalene, acenaphthylene, acenaphthene, fluorene, phenanthrene, fluoranthene, pyrene, benzo(a)anthracene, chrysene, benzo(b)fluoranthene, benzo(k)fluoranthene, benzo(a)pyrene, indeno(1,2,3-cd)pyrene, dibenzo(a,h)anthracene and benzo(ghi)perylene; “n-alkane” is the summed concentrations of C24, C25, C26, C27, C28, C29, C30, C31, C32, C33, C34; “cooking markers” is the summed concentrations of palmitic acid, stearic acid, cholesterol; “vehicle markers” is the summed concentrations of 17a(H)-22,29,30-trisnorhopane (C27a) and 17b(H),21a(H)-norhopane (C30ba); “SOA” is the summed concentrations of 2-methylthreitol, 2-methylerythritol, 2-methylglyceric acid, cis-2-methyl-1,3,4-trihydroxy-1-butene, -methyl-2,3,4-trihydroxy-1-butene, trans-2-methyl-1,3,4-trihydroxy-1-butene, C5-alkene triols, 2-methyltetrols, 3-hydroxyglutaric acid, cis-pinonic acid, acid, MBTCA, beta-caryophyllinic acid,

glutaric acid derivative, 3-acetylpentanedioic acid, 3-acetylhexanedioic acid, 3-isopropylpentanedioic acid and 2,3-dihydroxy-4-oxopentanoic acid. Dates marked in red indicate partial or total day haze events as described in Shi et al. (2019). Measurement uncertainty values were unavailable for most data types, and for selected dates in the upper plots, the sum of the total mass measurements is slightly more than 1 (i.e. more than 1 μg per μg); for these dates, the data has been proportionately scaled. It should be noted that the OC measurement in the upper plots incorporates the variety of organic carbon species represented in the lower plots.

R_s calculated for OP_v and OP_m with the individual compositional measurements have strikingly different univariate correlations, as illustrated in correlation heatmaps (**Figure 5**). Cumulative scores, referring to the number of R_s correlations ≥ 0.5 for OP_m and OP_v (Table S3), demonstrate that for all assays, considerably more significant correlations are observed for OP_v in the winter compared to OP_m . For both OP_v and OP_m , all assays show more statistically significant correlations in winter compared to summer, particularly for the AA response (AA_m , $n = 54$ winter, $n = 15$ summer, AA_v , $n = 67$ winter, $n = 4$ summer). Volume-based correlation analysis (**Figure 5A**) indicates that a very large number of the 107 atmospheric components measured in this study correlate statistically significantly with all four assays. The large number of correlations in the volume-normalised data indicate strong collinearity between concentrations of chemical components in $\text{PM}_{2.5}$ and overall $\text{PM}_{2.5}$ mass concentrations likely due to meteorological processes, complicating analysis of the sources and processes contributing to OP variability in particles. However, the mass-based analysis (**Figure 5B**) reveals that the mass fractions of chemical components and sources to which the four assays are sensitive to differ significantly (further illustrated by the weaker inter-assay correlations shown in Table 1), which demonstrates that mass-based analysis of OP data is also important to elucidate atmospheric processes and particle sources responsible for the different OP metrics.

440



445

Figure 5. Heatmaps demonstrating the correlation of OP, expressed as volume-normalised OP_m (A) and mass-normalised OP_v (B) vs a range of additional measurements conducted during the APHH campaign. Red indicates positive correlation; blue indicates inverse correlation. For OP_m , particle-phase components are also mass normalised ($\mu\text{g per } \mu\text{g PM}_{2.5}$ and for OP_v , volume-normalised ($\mu\text{g or ng per m}^3$).

A range of transition metals were all positively correlated with AA_m and DTT_m, including V, Cr, Mn, Fe, Co, Ni, Zn, Cd and Pb (all $R_s \geq 0.5$, $p < 0.05$). This reinforces the importance of their contribution to urban PM_{2.5} and potential to substantially influence PM_{2.5} OP, particularly Fe, Cr, V and Co which are commonly major components of vehicle emissions, and which can undergo redox-cycling reactions producing ROS (Charrier et al., 2014; Shen and Anastasio, 2012; Valko et al., 2005) contributing to higher AA_m and DTT_m in the winter compared to the summer. Stronger correlations between Fe and AA_m are observed in the winter (R_s 0.73) compared to summer (R_s 0.48) despite Fe concentrations ($\mu\text{g}/\mu\text{g}$) being lower in winter samples than summer samples, again highlighting the enhanced role of redox-active transition metals in winter. It is not established whether this seasonal difference is related to the chemical availability (i.e. redox state, solubility, speciation) of Fe, to the variability of emission sources of Fe between the seasons, or to some other important unknown additional contribution of Fe to AA_m in the summer; complexation of Fe in PM may differ between seasons, and the ligands directly influence the redox state and thus the bioavailability of the metal (Ghio et al., 1999). Ultimately, the direct correlation of transition metals measured only by ICP-MS with OP does not adequately reflect the nuances in redox behaviour of these species when they are complexed with organic ligands (Calas et al., 2017), as well as their range of oxidation states; this represents further gaps in the standard chemical (and particularly the transition metal) characterisation of PM. The epidemiological effects related to bioavailability of the metal when complexed (Costa and Dreher, 1997) in humans are also still not fully explored, although it is clear from multiple atmospheric and clinical studies that complexation affects both transition metal uptake in both the atmosphere and in the body. Interestingly, a mild inverse correlation of Fe with DCFH_m is observed (**Table S8**), which may be linked to the destruction of particle-bound organic peroxides by Fe *via* Fenton-type chemistry (Charrier et al., 2014), a process which the DCFH assay is specifically sensitive to (Gallimore et al., 2017; Wragg et al., 2016), and which has been observed in other recent studies (Paulson et al., 2019). No significant positive correlation between any metals measured in this study and DCFH_m and EPR_m was observed. Few EPR studies have looked specifically at superoxide formation, as is the case here, but those conducted so far show that EPR specifically detecting O₂⁻ is less sensitive to transition metal chemistry compared to traditional EPR methods focussing on OH formation.

In the summer, from the measured transition metals, only Fe correlated significantly positively (Spearman correlation p-value < 0.05) with DTT_m and AA_m response ($R_s = 0.48, 0.51$ respectively), whereas in the winter, DTT_m and AA_m correlated with a number of transition metals including V, Cr, Mn, Fe, Co, Ni, Zn, Cd. Of particular note, AA_m is mildly correlated with Cu in winter samples (R_s 0.48), whereas no correlation is observed between DTT_m and Cu in either winter or summer, in agreement with a recent online DTT study also (Puthussery et al., 2020). In contrast, previous reports from other locations have implicated Cu as a dominant contributor to DTT oxidation, considering volume normalised and mass normalised data (Calas et al., 2018; Charrier et al., 2015). Interestingly, in contrast with OP_m, strong correlations ($R_s > 0.6$) are observed in this study between AA_v, EPR_v, DCFH_v and DTT_v and Cu in the winter, but poorer correlations are observed in the summer for all assays ($R_s < 0.39$). Higher average Cu concentrations in winter compared to summer (winter = 17.7 ng m^{-3} , summer = 4.9 ng m^{-3}) may explain the higher R_s observed for Cu vs. OP_v in winter compared to summer, whereas mass normalised concentrations of Cu are more similar between the seasons. Poor correlation of Cu concentrations with AA_m and DTT_m response in winter may hint

at more insoluble Cu complex formation observed at this site in Beijing, as predominantly water-soluble Cu participates in redox reactions, therefore the sensitivity of AA and DTT towards Cu probably depends on the soluble fraction of Cu (Bates et al., 2019; Charrier and Anastasio, 2012; Fang et al., 2016). Furthermore, the presence of organic chelating ligands in PM may reduce the redox-activity of Cu and Fe (Charrier et al., 2014; Charrier and Anastasio, 2011; Shen and Anastasio, 2012).

Correlations between AA_m and DTT_m with total OC are observed in both summer and winter (**Tables S6 and S7**), and with total EC in the winter season, whereas DCFH_m is negatively correlated with total OC (**Table S8**). In contrast, DCFH_m is positively correlated with MOOOA and LOOOA, whereas DTT_m and AA_m show no correlation and even exhibit slight negative correlations with MOOOA and LOOOA in both summer and winter. This potentially indicates that the MOOOA and LOOOA AMS fractions, typically associated with water-soluble organic carbon content (Verma et al., 2015b), may contain higher concentrations of particle-bound ROS (i.e. organic peroxides) as measured by DCFH_m, but on a per-mass basis these species may contribute less significantly to AA_m and DTT_m compared to redox-active transition metals and other organic components. Total OC and EC correlations with AA_m and DTT_m may relate to concentrations of redox-active organic components such as oxidised PAHs and quinones, which may not be represented by MOOOA and LOOOA factors and which have been shown to significantly contribute to DTT_m (Chung et al., 2006; McWhinney et al., 2013b).

Significant correlations are also observed between AA_m and a range of *n*-alkanes and hopanes (17a(H)-22, 29,30-trisnorhopane (C27a) and 17b(H)-21a-norhopane (C30ba), **Table S6**), markers of primary organic aerosol emitted from vehicles (Schauer et al., 1999; Subramanian et al., 2006). Although these species are not redox-active, they are co-emitted with redox-active transition metals such as Fe, V and Cu from vehicle activity, either directly (Bates et al., 2019) or *via* dust resuspension, and other organics contributing to SOA (Platt et al., 2014) and highlight the potential importance of vehicular emissions on AA_m. Vehicular emissions and dust-resuspension have been previously shown to be the dominant sources of Cu and Fe in Beijing (Gao et al., 2014). EPR_m, DTT_m and DCFH_m responses do not show any significant correlations with these organic traffic markers.

Notably, AA_m correlates well with *cis*-pinonic acid, pinic acid and 3-methyl-2,3,4-butanetricarboxylic acid (MBTCA) in both seasons, all of which are biogenic SOA markers and products of α -pinene oxidation, with MBTCA a marker for OH-initiated ageing of first generation α -pinene oxidation products (Müller et al., 2012). AA sensitivity towards α -pinene SOA has been demonstrated previously (Campbell et al., 2019a). Although these three carboxylic acids are also not redox-active, they may correlate with the formation of particle-bound ROS such as peroxides or peroxy acids in SOA (Steimer et al., 2018), or with species that decompose liberate ROS upon extraction (e.g. (Tong et al., 2017)); these processes are highly likely to contribute to AA_m, highlighting the assay's potential sensitivity to redox-active particle phase components and particle-bound ROS. Generally, DTT_m has been previously shown to be relatively insensitive to SOA as observed here (Bates et al., 2015; Verma et al., 2015b), and both DTT_m and DCFH_m correlate poorly with the SOA markers analysed in the present study (**Tables S7 and S8**).

Compared to the three other assays, few significant correlations are observed between EPR_m and additional measurements, despite the much better correlations with the EPR_v data, particularly for the summer samples. However, seasonality in the

EPR_m response is still observed, with substantial variability in the mass-normalised EPR_m response (\approx factor of 10 in the summer, factor of 2 in the winter, **Figure 3**). Therefore, we observe differences in aerosol composition influencing EPR_m, but with the current comprehensive measurements (i.e., 107 parameters) are unable to determine the specific PM_{2.5} components responsible for the observed EPR_m variation.

520 The univariate analysis presented here clearly shows that OP_m enables a more nuanced identification of aerosol components influencing the oxidising properties of PM_{2.5} as compared to OP_v. Many more correlations are observed when considering volume-normalised OP_v, likely related to collinearity of species with overall PM_{2.5} mass concentration due to meteorological effects. Metal and organic tracers of traffic emissions (exhaust and non-exhaust) such as Fe, Cu and hopanes and SOA markers show especially strong correlations with AA_m, whereas the other three OP_m metrics (DTT_m, DCFH_m and EPR_m) provide a less
525 clear picture.

3.3 Multivariate modelling of OP from measured components

To assess potential latent influences from the individual components on assay response and hence on OP, a systematic multivariate analysis was undertaken. Initially, principal components analysis was applied to the whole set of independent measurements excluding the OP assay responses (i.e. the values to be predicted by the models), to investigate which contributed
530 most to the variation in the data, whether there were relationships between measurements which characterised OP, and if the OP_m response could be predicted from the individual component measurements.

In the PCA model, the seasonal variation within the samples was clearly apparent (**Figure 6**). The first four principal components (PC) accounted for 68.2% of the observed variation in the dataset (R^2 or goodness-of-fit), of which 50.5% was stable through 7-fold cross-validation (Q^2 , or model variation accounted for through cross-validation), indicating about half of
535 the variation in the model was robust with respect to sample score prediction. The loadings plot (**Figure 7**) indicated that seasonality in the first principal component was related to increased PAHs (Feng et al., 2019), *n*-alkanes (He et al., 2006) and biomass burning markers (He et al., 2006) in winter, and increased ozone (Zhao et al., 2018), ambient temperature and selected SOA markers (including 2-methylerythritol (Liang et al., 2012) and 2-methylglyceric acid (Ding et al., 2016; Shen et al., 2018)) in summer, findings which are consistent with existing volume-based studies. When scores were coloured by OP,
540 the AA_m (**Figure 6A**), DTT_m (**Figure 6C**) and DCFH_m (**Figure 6D**) assay responses could be observed in the second and sometimes also the first principal components (although the EPR_m response demonstrated no specific trend, **Figure 6B**). When loadings plots were examined by general measurement category (**Figure 7**), it was observed some categories of measurements cluster together (e.g. PAH, *n*-alkanes, NO_x, temperature, relative humidity), but this was related to strong correlation of these species with the OP_m measurement and known compound behaviour rather than to measurement bias, as other categories
545 showed broader variation (e.g. inorganic and small organic ions, gases, metals and SOA markers).





Figure 6. Principal components analysis scores plot of all data. A: coloured by AA_m response; B: coloured by EPR_m response; C: coloured by DTT_m response; D: coloured by DCFH_m response. Both principal component 1 and principal component 2 demonstrate variance associated with AA and DTT response, and there is greater variation associated with the winter response than the summer response (highlighted in A). PC 1 R²X 35.90%, Q² 29.28%; PC 2 R²X 19.34%, Q² 23.73%; the model included six principal components, with a cumulative R²X of 68.2% and Q² of 50.5%.



560

Figure 7. Principal components analysis loading plot for all data points. Points are coloured by measurement category; fully labelled loadings are provided in **Figure S14**. The plot is annotated with the same orientation as the scores plot, to indicate the direction of visualised trends for selected assays and for season from the latent variable origin as shown in **Figure 6**. In PC 1, the winter classification is driven by increased gas radicals, *n*-alkanes, PAH, vehicle markers, biomass burning markers, total OC and selected metals and SOA markers; the summer classification is driven by increased temperature and photolysis, ozone (the single gas species in this section of the plot), selected SOA markers and metals, and selected VOCs. In PC 2, high AA_m and DTT_m response is associated with increased SOA, transition metals, cooking markers, *n*-alkanes and PAH concentrations in samples; low AA_m and DTT_m response associated with low VOCs, gases and selected meteorological parameters (relative humidity).

565

Partial least squares regression (PLSR) is a supervised regression extension of PCA, which models the variation in the data associated with a defined sample classification (Eriksson et al., 2013). PLSR models were constructed for each individual OP

assay and season, to examine the most specific markers associated with assay response. **Table 2** provides the model performances for all PLSR assay models, and example PLSR scores plots for AA_m and DTT_m models (both seasons) are illustrated in **Figures 8** and **9** (analogous plots for other assays provided in **Figures S18** and **S19**). The performance indicators show that while the mass-normalised measurement data can be used to explain and predict a large majority of the variation associated with AA_m summer/winter and DTT_m winter assay response, the other assay responses were less consistent; R² and Q² values for these models indicated that less than 70% of the variance in response can be predicted from the individual component measurements, and the predictions were much less stable through cross-validation. These results could suggest either that assay responses are not as adequately sensitive at µg/µg concentrations as for the total PM per sample, or that a proportion of the OP_m response is contributed to by species not measured directly in this campaign and which cannot also be inferred from total organic carbon measurements. As total OC is estimated from combustion properties of the sample rather than from a sum of individually validated component measurements, and as multiple organic and transition metal-organic complexed species contribute to the total OC measurements with unknown redox properties, these observations reiterate the need for more comprehensive chemical characterisation of PM composition. Similarly to the univariate correlations, the summer samples were less well modelled in both mass-normalised and volume-normalised data, indicating either reduced assay sensitivity (which may also be compounded by the reduced collected filter PM mass in summer) or the influence of unmeasured components.

Table 2. Performance assessment of PLSR models for all assays, for both mass-normalised (left) and volume-normalised (right) data. Models are considered to perform well when both cumulative (i.e. across all latent variables included in the model) R² and Q² values are high, or at a minimum where Q² values are within 10% of the R² value, indicating that the variance is well accounted for in model cross-validation. Permutation tests were rejected for robustness if any single random permutation model performance surpassed the performance of the real cross-validated model; on this basis, the winter DCFH_m and summer DTT_v models were rejected (highlighted with *), although fewer than three random models outperformed the real model, and none of the permuted model Q² values outperformed those of the real model.

assay	season	mass (µg/µg)				volume (µg/m ³)			
		optimal LVs	cumulative R ²	cumulative Q ²	permutation test pass	optimal LVs	cumulative R ²	cumulative Q ²	permutation test pass
EPR	winter	1	43.2	19.3	no	2	83.9	75.2	yes
	summer	1	11.3	-10.0	no	1	52.0	3.7	no
AA	winter	1	81.4	78.2	yes	2	94.1	87.9	yes
	summer	2	79.3	49.7	yes	1	41.8	22.6	no
DTT	winter	2	76.0	62.0	yes	2	86.8	67.0	yes
	summer	1	47.4	31.6	no	1	66.2	50.9	no*
DCFH	winter	2	71.9	50.4	no*	2	67.0	55.2	yes
	summer	1	28.2	-6.6	no	1	86.0	66.7	yes




Figure 8. PLSR scores plot for AA_m assay. Model performance parameters given in Table 2. Left: winter samples; right: summer samples. Points coloured by overall AA assay response for both seasons. Red bar indicates 2 × SD for all scores, orange dotted line indicates 1 × SD for all scores. Models which have only one latent variable have the X-axis replaced by date for easier visualisation.




Figure 9. PLSR scores plot for DTT_m assay. Model performance parameters given in Table 2. Left: winter samples; right: summer samples. Points coloured by overall DTT assay response for both seasons.

Table 3. Characteristic loadings most influential in PLSR models of OP_m as defined by ordered variable importance in projection for each model. Blue upward arrows indicate positive correlation with the assay measurement, red downward arrows for inverse correlation, and * for p < 0.05 in Spearman correlation of the feature with the assay in the univariate analysis.

EPR _m winter		AA _m winter		DTT _m winter		DCFH _m winter	
feature	VIP	feature	VIP	feature	VIP	feature	VIP
indeno(1,2,3-cd)-pyrene *	2.12 ↑	cis-pinonic acid *	1.44 ↑	SO ₂ *	1.46 ↓	NH ₄ ⁺	2.16 ↑
acenaphthylene	2.02 ↑	Cl ⁻ *	1.42 ↑	Ca ²⁺ *	1.40 ↑	chrysene *	1.61 ↓
benzo(ghi)-perylene *	2.01 ↑	total OC *	1.33 ↑	Fe *	1.37 ↑	benzo(b)-fluoranthene *	1.59 ↓
benzo(a)pyrene *	2.01 ↑	MOOOA *	1.30 ↑	fluorene	1.34 ↑	RH8 *	1.59 ↑
fluorene	1.82 ↑	pyrene *	1.30 ↑	acetaldehyde *	1.33 ↓	benzo(a)anthracene*	1.58 ↓
benzo(a)-anthracene *	1.81 ↑	2-methylthreitol	1.29 ↑	phenanthrene *	1.33 ↑	pyrene *	1.58 ↓
dibenzo(a,h)-anthracene *	1.80 ↑	ORG *	1.29 ↑	acetone *	1.33 ↓	LOOOA *	1.57 ↑
phenanthrene *	1.77 ↑	benzo(k)-fluoranthene *	1.29 ↑	Cl ⁻ *	1.31 ↑	fluoranthene *	1.56 ↓
chrysene *	1.66 ↑	3-methyl-2,3,4-trihydroxy-1-butene *	1.28 ↑	benzene *	1.31 ↓	RH120 * / RH240 *	1.55 ↑
naphthalene *	1.62 ↑	fluoranthene *	1.27 ↑	toluene *	1.30 ↓	K ⁺ *	1.51 ↑

EPR _m summer		AA _m summer		DTT _m summer		DCFH _m summer	
feature	VIP	feature	VIP	feature	VIP	feature	VIP
LOOOA	2.59 ↑	ORG *	1.80 ↑	OH	1.58 ↑	cis-pinonic acid *	2.38 ↓
T8 / T120 / T240	2.28/2.15/2.08 ↑	cis-pinonic acid *	1.62 ↑	dibenzo(a,h)-anthracene *	1.51 ↑	C31 *	1.76 ↓
O ₃	2.00 ↑	MOOOA *	1.58 ↑	C26 *	1.48 ↑	pinic acid *	1.74 ↓
RO ₂ *	1.76 ↑	cholesterol	1.58 ↓	benzo(a)-pyrene *	1.48 ↑	acetonitrile *	1.69 ↑
galactosan *	1.74 ↓	naphthalene *	1.57 ↑	total OC *	1.46 ↑	3-methyl-2,3,4-trihydroxy-1-butene	1.65 ↓
K ⁺	1.70 ↑	palmitic acid *	1.49 ↑	C30 *	1.46 ↑	benzo(ghi)-perylene	1.62 ↓
17a(H)-22,29,30-trisnorhopane (C27a)	1.55 ↓	RH8	1.39 ↓	C28 *	1.43 ↑	C32	1.61 ↓
cis-2-methyl-1,3,4-trihydroxy-1-butene	1.55 ↑	stearic acid *	1.39 ↑	benzo(ghi)-perylene *	1.41 ↑	dibenzo(a,h)-anthracene *	1.61 ↓
Ba	1.47 ↓	benzo(ghi)-perylene *	1.36 ↑	C33 *	1.40 ↑	acetaldehyde *	1.61 ↑
RH8	1.46 ↓	benzo(a)-pyrene *	1.34 ↑	C29 *	1.39 ↑	isoprene *	1.61 ↓

Table 3 shows the top ten features in the variable importance in projection (VIP) for the PLSR loadings, which enable a ranking of the features which contribute most to the model (Naes and Martens, 1988). It is evident from these data that the features which best model the OP_m seasonal response are derived from multiple particle sources and atmospheric aging processes. For example, the AA_m and DTT_m responses show similar trends in the multivariate models, but the main contributors to their responses have little overlap, with AA_m responses being more strongly associated with SOA tracers, PAHs and general measures of organic carbon, and the DTT_m more characterised by combustion and vehicle emissions markers (**Figures S19-S22**; Figures S17-S24 list the top 50 contributors to each assay model response). Notably, compounds which are not generally recognised as being redox-active were frequently observed to be important in PLSR classification, and though they do not directly contribute to the OP_m response, they are likely co-emitted with or are secondary products of redox-active particle components.

3.4 Multiple Linear Regression (MLR) modelling to predict OP_m associated with specific sources

While multivariate model loadings highlighted the measurements most associated with assay response, multivariate models are not always amenable to variable selection, which is important to characterise the chemical profiles contributing to each assay OP response. Multiple linear regression modelling has been used in previous studies (Calas et al., 2018) to establish contributors to total OP response, rather than looking at source apportionment in relation to OP, and only simple forward variable selection was used for model refinement. In the present study, relevant measurements were grouped into six categories of known contributors to Beijing PM (biogenic SOA, biomass burning, coal and fossil power generation, cooking, dust and vehicle emissions). The full method description, references, model formulae and performance parameters for the mass-normalised data models are presented in the Methods (**Section 2.3** “Statistical analysis”) and in **Section S10**. Briefly, literature sources (**Table S13**, Section S10) and the SPECIEUROPE database (Pernigotti et al., 2016) were used to establish which individual chemical measurements were likely to be characteristic of each source, with several measurements appearing in multiple categories (e.g. total EC). All proxy and composite measurements (except total EC, as multiple organic carbon species are represented in the dataset, but elemental carbon should be independent of the majority of these), AMS measurements, and general atmospheric measurements including temperature, relative humidity and actinic flux measurements were excluded from models entirely. Composite measures duplicate selected individual measurements; atmospheric measurements complicate model interpretation, and are more likely to be useful as random effects terms in a mixed effects model approach (not pursued in the present study due to the complexity of model parameterisation and measurement uncertainties). Multiple linear regression models were then constructed for each assay and season for each category, using both mass-normalised and volume-normalised data.

MLR models further reinforced that not all putative sources and components of PM_{2.5} contribute equally to OP_m response (

Table 4). OP_m assay response models based on measurements characteristic of vehicle emissions, coal/fossil fuel combustion and biomass burning gave accurate and robust predictions of OP_m, and these are important contributors to PM (reported as

mass per volume) in Beijing urban background sites (Yu et al., 2013; Zheng et al., 2005). As expected, OP_v models also gave very good predictions for these source profiles, but additionally gave improved models of OP_v for biogenic SOA and dust compared with the OP_m data. Although the same base sets of predictors for each source were used for each model (season, OP assay and PM normalisation), there was only partial overlap of the final selected predictors between models from the same source and season, again illustrating the complex dynamic between OP and overall mass/volume composition. As with the PLSR models, the most important contributors to regression models were often not redox-active species, indicating that they are probably influencing or contributing to the oxidation state of the redox-active PM components. As with the univariate and multivariate analyses, the summer samples gave less robust linear regression models (and thus OP predictions) from both mass- and volume-normalised data. However, AA and DTT measurements produced the best models for all source contributions, indicating that these assays might be most optimal for measuring OP in an urban environment, as they appear to reflect the variety and composition of PM sources well.

Vehicle emissions, biogenic SOA and winter biomass burning contributions to AA and DTT response (as measured by the model R² value) were generally comparable across both assays, contrasting with the findings of Fang et al. (2016), who observed greater OP response in positive matrix factorisation-chemical mass balance (PMF-CMB) models associated with traffic emissions for AA_v over DTT_v, and biomass burning for DTT_v over AA_v in multiple locations in the southeastern US. However, a more recent study conducted in the coastal areas adjacent to Beijing (Liu et al., 2018) observed similar seasonality to the present study in the DTT OP_m response. Vehicle emissions (Wang et al., 2016; Yu et al., 2019), coal combustion (Ma et al., 2018; Yu et al., 2019), biomass burning (Ma et al., 2018) and dust (Yu et al., 2019) sources have been shown in other studies using PMF models to contribute to OP_v in Beijing, all using the DTT assay. Cooking markers (palmitic acid, stearic acid and cholesterol) contributed a substantial proportion of the known organic fraction of the PM mass and volume concentrations (see **Figure 4**), but did not contribute robustly to the modelled OP response for either normalisation type, suggesting they are either not strongly contributing to or affected by oxidative conditions in PM, or that their variation over the sampling period cannot be linearly modelled. Similarly, biomass burning markers contribute a comparable number of variables in the model base sets, but appear to contribute much more significantly to the OP_v than to the OP_m response. Biogenic SOA and dust models (which incorporate K⁺, Na⁺, Ca²⁺, Cl⁻, Al, Ti, Mn, Fe and Zn) explained a significant proportion of winter OP_v responses, but were only strongly correlated with winter AA and DTT for mass-normalised models. These observations suggest these sources contribute to PM OP_v by total quantity rather than through their particularly high intrinsic OP, i.e. their mass as a proportion of the PM mass is smaller, but their concentration per volume is high, and the AA and DTT assays have a notable selectivity for these species over the EPR and DCFH assays.

It should be noted that the MLR models represent a sub-optimal prediction of the OP response from measured components, as numerous species which are known source components (e.g. PAHs from combustion processes and those which distinguish gasoline from diesel vehicle emissions, or VOCs relevant to biomass burning such as methanol or acrolein) could not be included in models. Not all measurements which were associated in the literature with a particular assay response passed the stages of variable selection for mass-normalised models, which could reflect a lower limit of detection in either the OP_m assay

670 responses, or in the individual component measurements. Synergistic effects between individual measured components (e.g. transition metals with organic components such as quinones or carboxylic acids (Wang et al., 2018)) cannot be interpreted from linear models when the complexation and oxidation states of the contributing compounds are essentially unknown. MLR models do not fully account for the proportion of each measurement which may originate from multiple emissions sources, and PMF-CMB or mixed effects models can address this issue more adequately. Validation of both the multivariate and MLR
675 models using secondary datasets (both from Beijing and other locations) is also needed prior to their future implementation.

Table 4. R² values for optimised subset multiple linear regression models of relevant source contributions. R² values greater than 0.7 are highlighted in bold. Full model performance indicators are provided in **Section S11** of the SI, including all model terms, residuals, coefficients and p-values.

		EPR R ²		AA R ²		DTT R ²		DCFH R ²	
(µg/µg)	source model	winter	summer	winter	summer	winter	summer	winter	summer
	vehicle emissions	0.88	0.72	0.95	0.73	0.91	0.80	0.89	0.62
	biomass burning	0.41	0.29	0.49	0.47	0.45	0.41	0.58	0.31
	coal/fossil fuel combustion	0.84	0.56	0.88	0.61	0.86	0.68	0.75	0.71
	cooking markers	0.19	0.11	0.66	0.20	0.39	0.36	0.08	0.24
	dust	0.23	0.23	0.88	0.47	0.72	0.46	0.50	0.26
	biogenic SOA	0.55	0.35	0.95	0.74	0.79	0.61	0.55	0.70
(µg/m ³)	vehicle emissions	0.94	0.79	0.97	0.74	0.96	0.87	0.94	0.86
	biomass burning	0.85	0.23	0.89	0.24	0.72	0.62	0.78	0.53
	coal/fossil fuel combustion	0.91	0.69	0.95	0.62	0.88	0.77	0.93	0.91
	cooking markers	0.10	0.08	0.09	0.22	0.10	0.44	0.11	0.49
	dust	0.79	0.21	0.92	0.30	0.78	0.54	0.73	0.63
	biogenic SOA	0.87	0.36	0.84	0.59	0.80	0.63	0.94	0.90

4 Conclusions

This study presents a detailed and comprehensive analysis of PM_{2.5} oxidative potential and particle-bound ROS concentrations measured in winter 2016 and summer 2017 during the APHH-Beijing campaign at a central site in Beijing, China. Four acellular methods for measuring OP were applied, providing a broad assessment of the oxidative properties of particles
685 including particle-bound ROS concentrations, superoxide radical production and catalytic redox activity. We correlated the acellular assay responses with an extensive and comprehensive dataset including 107 additional atmospheric measurements (particle components, trace gases, meteorological parameters) to delineate chemical particle components and atmospheric processes and sources responsible for driving PM_{2.5} OP. Higher volume-normalised and mass-normalised OP values across all assays were observed in the winter compared to the summer. An inverse correlation was observed between AA_m and DTT_m
690 with overall PM_{2.5} mass concentrations, i.e. days with higher PM_{2.5} mass concentrations have lower intrinsic OP values. This

is likely due to an increase in OP-inactive material in high PM_{2.5} mass days, and/or a mass fraction that is at present undetermined and highlights that a focus on total PM exposure only does not necessarily capture accurately the oxidising properties and therefore certain toxicological effects of PM.

Univariate analysis with the additional 107 measurement parameters acquired during the APHH-Beijing campaign highlight significant assay-specific responses to chemical components of PM_{2.5}, as well as a seasonal difference in what components drive aerosol OP. It also highlights the importance of considering both volume-normalised and mass-normalised OP metrics when drawing conclusions on the role of chemical composition on OP, as assay correlations vary significantly between the two metrics. The data presented in this study illustrate that mass-normalised OP_m values provide a more nuanced picture of specific chemical components and sources that influence intrinsic OP, whereas many more correlations with OP_v values are observed, likely due to collinearity of many chemical components with overall PM_{2.5} mass concentrations driven by changes in meteorological conditions. Both metrics, mass-normalised OP_m as well as volume-normalised OP_v, are important to consider with OP_v a more relevant metric with respect to exposure and epidemiological studies, whereas OP_m provides more insight into what sources and what composition drives OP concentrations in particles. Furthermore, OP_m may allow easier study and site inter-comparisons.

The multivariate statistical analyses encapsulated the observations from the univariate analyses into comprehensive single models of OP relating to PM composition, and the inference from the univariate analyses that OP_m measured by each assay is related to different compounds present in the particle was confirmed. It is clear from these differences that assay chemistry must contribute directly to its chemical selectivity, as the independent chemical measurements were given equal analytical weight with respect to each assay. The relationship between each assay and the independent measurements also confirmed that while there may exist a correlative relationship between an assay and non-redox active compounds such as *n*-alkanes or PAHs, the assay is more likely to be measuring either secondary oxidation products of these primary compounds or species co-emitted that contribute to particle OP. This represents a gap in the chemical analysis of these samples, and more detailed redox active compound speciation is required, especially for functionalised organics. Furthermore, variable selection of measurements and evaluation through multiple linear regression models indicated that OP_m is well predicted by measurement panels characteristic of combustion sources, particularly (exhaust and non-exhaust) vehicle emissions, and biogenic SOA. This study demonstrates further that these commonly applied acellular assays are sensitive to a wide and differing range of chemical components, highlighting the advantage of using these assays as they encompass multiple chemical components and sources of aerosol into an integrated measurement. Further comprehensive work is needed to identify the direct links between these OP assays and biological and toxicology data.

Author Contributions. SJC collated data, analysed filters for AA and DCFH, performed data analysis and interpretation and wrote the manuscript. KW performed univariate and multivariate statistical analysis, data interpretation and wrote the manuscript. BU, JW, ST and NS analysed filters for AA, DCFH, DTT and EPR respectively. TV provided XRF and additional

data. AMS data were provided by YS. PAH data was provided by AE and AL. SOA tracer data was provided by DL, LL and
725 PF. All other authors contributed to data analysis, interpretation and writing of the manuscript.

Competing Interests. The authors declare that they have no conflict of interest

Special Issue Statement. This article is part of the special issue “In-depth study of air pollution sources and processes within
730 Beijing and its surrounding region (APHH-Beijing) (ACP/AMT inter-journal SI)

Acknowledgements. This work was funded by the European Research Council (ERC grant 279405), by the Natural
Environment Research Council (NERC) (NE/K008218/1) and the Swiss National Science Foundation (200021_192192 / 1).
This work has also received funding from the European Union’s Horizon 2020 research and innovation programme through
735 the EUROCHAMP-2020 Infrastructure Activity under grant agreement No. 730997. SSS additionally acknowledges support
from the Swiss National Science Foundation (fellowship P2EZP2_162258) and a 2017 LIFE PostDoc fellowship by the AXA
Research Fund. We acknowledge the support from Pingqing Fu, Zifa Wang, Jie Li and Yele Sun from IAP for hosting the
APHH-Beijing campaign at IAP. We thank Di Liu from the University of Birmingham, Siyao Yue, Liangfang Wei, Hong Ren,
Qiaorong Xie, Wanyu Zhao, Linjie Li, Ping Li, Shengjie Hou, Qingqing Wang from IAP, Rachel Dunmore and James Lee
740 from the University of York, Kebin He and Xiaoting Cheng from Tsinghua University, and James Allan and Hugh Coe from
the University of Manchester for providing logistic and scientific support for the field campaigns.

References

- Acton, J., Hewitt, N., Huang, Z. and Wang, X.: APHH: Volatile organic compound (VOC) mixing ratios made at the IAP-
Beijing site during the summer and winter campaigns. Centre for Environmental Data Analysis, 18.06.2020
745 <https://catalogue.ceda.ac.uk/uuid/de37c54e59a548ccb9f168ee724f3769>, 2018.
- Arangio, A. M., Tong, H., Socorro, J., Pöschl, U. and Shiraiwa, M.: Quantification of environmentally persistent free radicals
and reactive oxygen species in atmospheric aerosol particles, *Atmos. Chem. Phys.*, 16(20), 13105–13119, doi:10.5194/acp-
16-13105-2016, 2016.
- Atkinson, R. and Arey, J.: Mechanisms of the gas-phase reactions of aromatic hydrocarbons and pahs with oh and NO3
750 radicals, *Polycycl. Aromat. Compd.*, 27(1), 15–40, doi:10.1080/10406630601134243, 2007.
- Bates, J. T., Weber, R. J., Abrams, J., Verma, V., Fang, T., Klein, M., Strickland, M. J., Sarnat, S. E., Chang, H. H., Mulholland,
J. A., Tolbert, P. E. and Russell, A. G.: Reactive Oxygen Species Generation Linked to Sources of Atmospheric Particulate
Matter and Cardiorespiratory Effects, *Environ. Sci. Technol.*, 49(22), 13605–13612, doi:10.1021/acs.est.5b02967, 2015.
- Bates, J. T., Fang, T., Verma, V., Zeng, L., Weber, R. J., Tolbert, P. E., Abrams, J. Y., Sarnat, S. E., Klein, M., Mulholland, J.
755 A. and Russell, A. G.: Review of Acellular Assays of Ambient Particulate Matter Oxidative Potential: Methods and

- Relationships with Composition, Sources, and Health Effects, *Environ. Sci. Technol.*, 53(8), 4003–4019, doi:10.1021/acs.est.8b03430, 2019.
- 760 Biswas, S., Verma, V., Schauer, J. J., Cassee, F. R., Cho, A. K. and Sioutas, C.: Oxidative potential of semi-volatile and non volatile particulate matter (PM) from heavy-duty vehicles retrofitted with emission control technologies, *Environ. Sci. Technol.*, 43(10), 3905–3912, doi:10.1021/es9000592, 2009.
- Calas, A., Uzu, G., Martins, J. M. F., Voisin, Di., Spadini, L., Lacroix, T. and Jaffrezzo, J. L.: The importance of simulated lung fluid (SLF) extractions for a more relevant evaluation of the oxidative potential of particulate matter, *Sci. Rep.*, 7(1), 1–12, doi:10.1038/s41598-017-11979-3, 2017.
- 765 Calas, A., Uzu, G., Kelly, F. J., Houdier, S., Martins, J. M. F., Thomas, F., Molton, F., Charron, A., Dunster, C., Oliete, A., Jacob, V., Besombes, J. L., Chevrier, F. and Jaffrezzo, J. L.: Comparison between five acellular oxidative potential measurement assays performed with detailed chemistry on PM10 samples from the city of Chamonix (France), *Atmos. Chem. Phys.*, 18(11), 7863–7875, doi:10.5194/acp-18-7863-2018, 2018.
- Campbell, S. J., Uttinger, B., Lienhard, D. M., Paulson, S. E., Shen, J., Griffiths, P. T., Stell, A. C. and Kalberer, M.: Development of a physiologically relevant online chemical assay to quantify aerosol oxidative potential, *Anal. Chem.*, 91, 13088–13095, doi:10.1021/acs.analchem.9b03282, 2019a.
- 770 Campbell, S. J., Stevanovic, S., Miljevic, B., Bottle, S. E., Ristovski, Z. and Kalberer, M.: Quantification of Particle-Bound Organic Radicals in Secondary Organic Aerosol, *Environ. Sci. Technol.*, 53(12), 6729–6737, doi:10.1021/acs.est.9b00825, 2019b.
- Charrier, J. G. and Anastasio, C.: Impacts of antioxidants on hydroxyl radical production from individual and mixed transition metals in a surrogate lung fluid, *Atmos. Environ.*, 45(40), 7555–7562, doi:10.1016/j.atmosenv.2010.12.021, 2011.
- 775 Charrier, J. G. and Anastasio, C.: On dithiothreitol (DTT) as a measure of oxidative potential for ambient particles: evidence for the importance of soluble transition metals, *Atmos. Chem. Phys.*, 12(19), 9321–9333, doi:10.5194/acp-12-9321-2012, 2012.
- Charrier, J. G., McFall, A. S., Richards-Henderson, N. K. and Anastasio, C.: Hydrogen peroxide formation in a surrogate lung fluid by transition metals and quinones present in particulate matter, *Environ. Sci. Technol.*, 48(12), 7010–7017, doi:10.1021/es501011w, 2014.
- 780 Charrier, J. G., Richards-Henderson, N. K., Bein, K. J., McFall, A. S., Wexler, A. S. and Anastasio, C.: Oxidant production from source-oriented particulate matter - Part 1: Oxidative potential using the dithiothreitol (DTT) assay, *Atmos. Chem. Phys.*, 15(5), 2327–2340, doi:10.5194/acp-15-2327-2015, 2015.
- 785 Charrier, J. G., McFall, A. S., Vu, K. K. T., Baroi, J., Olea, C., Hasson, A. and Anastasio, C.: A bias in the “mass-normalized” DTT response – An effect of non-linear concentration-response curves for copper and manganese, *Atmos. Environ.*, 144, 325–334, doi:10.1016/j.atmosenv.2016.08.071, 2016.
- Cho, A. K., Sioutas, C., Miguel, A. H., Kumagai, Y., Schmitz, D. A., Singh, M., Eiguren-Fernandez, A. and Froines, J. R.: Redox activity of airborne particulate matter at different sites in the Los Angeles Basin, *Environ. Res.*, 99(1), 40–47,

- 790 doi:10.1016/j.envres.2005.01.003, 2005.
- Chung, M. Y., Lazaro, R. A., Lim, D., Jackson, J., Lyon, J., Rendulic, D. and Hasson, A. S.: Aerosol-borne quinones and reactive oxygen species generation by particulate matter extracts, *Environ. Sci. Technol.*, 40(16), 4880–4886, doi:10.1021/es0515957, 2006.
- Costa, D. L. and Dreher, K. L.: Bioavailable transition metals in particulate matter mediate cardiopulmonary injury in healthy and compromised animal models., *Environ. Health Perspect.*, 105 Suppl 5(September), 1053–1060, doi:10.1289/ehp.97105s51053, 1997.
- Delgado-Saborit, J. M., Alam, M. S., Godri Pollitt, K. J., Stark, C. and Harrison, R. M.: Analysis of atmospheric concentrations of quinones and polycyclic aromatic hydrocarbons in vapour and particulate phases, *Atmos. Environ.*, 77, 974–982, doi:10.1016/j.atmosenv.2013.05.080, 2013.
- 800 Dellinger, B., Pryor, W. A., Cueto, R., Squadrito, G. L., Hegde, V. and Deutsch, W. A.: Role of free radicals in the toxicity of airborne fine particulate matter, *Chem. Res. Toxicol.*, 14(10), 1371–1377, doi:10.1021/tx010050x, 2001.
- Ding, X., He, Q. F., Shen, R. Q., Yu, Q. Q., Zhang, Y. Q., Xin, J. Y., Wen, T. X. and Wang, X. M.: Spatial and seasonal variations of isoprene secondary organic aerosol in China: Significant impact of biomass burning during winter, *Sci. Rep.*, 6(20411), 1–10, doi:10.1038/srep20411, 2016.
- 805 Donaldson, K. and Tran, C. L.: Inflammation Caused By Particles and Fibers, *Inhal. Toxicol.*, 14(1), 5–27, doi:10.1080/089583701753338613, 2002.
- Dou, J., Lin, P., Kuang, B. Y. and Yu, J. Z.: Reactive oxygen species production mediated by humic-like substances in atmospheric aerosols: Enhancement effects by pyridine, imidazole, and their derivatives, *Environ. Sci. Technol.*, 49(11), 6457–6465, doi:10.1021/es5059378, 2015.
- 810 Elzein, A., Dunmore, R. E., Ward, M. W., Hamilton, J. F. and Lewis, A. C.: Variability of polycyclic aromatic hydrocarbons and their oxidative derivatives in wintertime Beijing, China, *Atmos. Chem. Phys. Discuss.*, 1–28, doi:10.5194/acp-2019-120, 2019.
- Elzein, A., Stewart, G. J., Swift, S. J., Nelson, B. S., Crilley, L. R., Alam, M. S., Reyes-villegas, E., Gadi, R., Harrison, R. M., Hamilton, J. F. and Lewis, A. C.: A comparison of PM_{2.5}-bound polycyclic aromatic hydrocarbons in summer Beijing (China) and Delhi (India), *Atmos. Chem. Phys. Discuss.*, (August), 1–25, doi:https://doi.org/10.5194/acp-2020-770, 2020.
- 815 Eriksson, L., Byrne, T., Johansson, E., Trygg, J. and Vikstrom, C.: Multi-and megavariable data analysis basic principles and applications., 2013.
- Fang, T., Verma, V., Guo, H., King, L. E., Edgerton, E. S. and Weber, R. J.: A semi-automated system for quantifying the oxidative potential of ambient particles in aqueous extracts using the dithiothreitol (DTT) assay: Results from the Southeastern Center for Air Pollution and Epidemiology (SCAPE), *Atmos. Meas. Tech.*, 8(1), 471–482, doi:10.5194/amt-8-471-2015, 2015.
- 820 Fang, T., Verma, V., Bates, J. T., Abrams, J., Klein, M., Strickland, M. J., Sarnat, S. E., Chang, H. H., Mulholland, J. A., Tolbert, P. E., Russell, A. G. and Weber, R. J.: Oxidative potential of ambient water-soluble PM_{2.5} in the southeastern United States : contrasts in sources and health associations between ascorbic acid (AA) and dithiothreitol (DTT) assays, ,

- 3865–3879, doi:10.5194/acp-16-3865-2016, 2016.
- 825 Feng, B., Li, L., Xu, H., Wang, T., Wu, R., Chen, J., Zhang, Y., Liu, S., Ho, S. S. H., Cao, J. and Huang, W.: PM_{2.5}-bound polycyclic aromatic hydrocarbons (PAHs) in Beijing: Seasonal variations, sources, and risk assessment, *J. Environ. Sci. (China)*, 77, 11–19, doi:10.1016/j.jes.2017.12.025, 2019.
- Fuller, S. J., Wragg, F. P. H., Nutter, J. and Kalberer, M.: Comparison of on-line and off-line methods to quantify reactive oxygen species (ROS) in atmospheric aerosols, *Atmos. Environ.*, 92, 97–103, doi:10.1016/j.atmosenv.2014.04.006, 2014.
- 830 Gallimore, P. J., Mahon, B. M., Wragg, F. P. H., Fuller, S. J., Giorio, C., Kourtchev, I. and Kalberer, M.: Multiphase composition changes and reactive oxygen species formation during limonene oxidation in the new Cambridge Atmospheric Simulation Chamber (CASC), *Atmos. Chem. Phys.*, 17, 9853–9868, doi:10.5194/acp-2017-186, 2017.
- Gao, J., Tian, H., Cheng, K., Lu, L., Wang, Y., Wu, Y., Zhu, C., Liu, K., Zhou, J., Liu, X., Chen, J. and Hao, J.: Seasonal and spatial variation of trace elements in multi-size airborne particulate matters of Beijing, China: Mass concentration, enrichment
- 835 characteristics, source apportionment, chemical speciation and bioavailability, *Atmos. Environ.*, 99, 257–265, doi:10.1016/j.atmosenv.2014.08.081, 2014.
- Gehling, W. and Dellinger, B.: Environmentally persistent free radicals and their lifetimes in PM_{2.5}, *Environ. Sci. Technol.*, 47(15), 8172–8178, doi:10.1021/es401767m, 2013.
- Gehling, W., Khachatryan, L. and Dellinger, B.: Hydroxyl radical generation from environmentally persistent free radicals
- 840 (EPFRs) in PM_{2.5}, *Environ. Sci. Technol.*, 48(8), 4266–4272, doi:10.1021/es401770y, 2014.
- Ghio, A. J., Stonehuerner, J., Dailey, L. A. and Carter, J. D.: Metals associated with both the water-soluble and insoluble fractions of an ambient air pollution particle catalyze an oxidative stress, *Inhal. Toxicol.*, 11(1), 37–49, doi:10.1080/089583799197258, 1999.
- Ghio, A. J., Carraway, M. S. and Madden, M. C.: Composition of air pollution particles and oxidative stress in cells, tissues,
- 845 and living systems, *J. Toxicol. Environ. Heal. - Part B Crit. Rev.*, 15(1), 1–21, doi:10.1080/10937404.2012.632359, 2012.
- Godri, K. J., Harrison, R. M., Evans, T., Baker, T., Dunster, C., Mudway, I. S. and Kelly, F. J.: Increased oxidative burden associated with traffic component of ambient particulate matter at roadside and Urban background schools sites in London, *PLoS One*, 6(7), doi:10.1371/journal.pone.0021961, 2011.
- Hart, J. E., Liao, X., Hong, B., Puett, R. C., Yanosky, J. D., Suh, H., Kioumourtzoglou, M. A., Spiegelman, D. and Laden, F.:
- 850 The association of long-term exposure to PM_{2.5} on all-cause mortality in the Nurses' Health Study and the impact of measurement-error correction, *Environ. Heal.*, 14(1), 38, doi:10.1186/s12940-015-0027-6, 2015.
- Hasson, A. S. and Paulson, S. E.: An investigation of the relationship between gas-phase and aerosol-borne hydroperoxides in urban air, *J. Aerosol Sci.*, 34(4), 459–468, doi:10.1016/S0021-8502(03)00002-8, 2003.
- He, L. Y., Hu, M., Huang, X. F., Zhang, Y. H. and Tang, X. Y.: Seasonal pollution characteristics of organic compounds in
- 855 atmospheric fine particles in Beijing, *Sci. Total Environ.*, 359(1–3), 167–176, doi:10.1016/j.scitotenv.2005.05.044, 2006.
- Hewitt, C. N. and Kok, G. L.: Formation and occurrence of organic hydroperoxides in the troposphere: Laboratory and field observations, *J. Atmos. Chem.*, 12(2), 181–194, doi:10.1007/BF00115779, 1991.

- Hung, H. F. and Wang, C.-S.: Experimental determination of reactive oxygen species in Taipei aerosols, *J. Aerosol Sci.*, 32(10), 1201–1211, doi:10.1016/S0021-8502(01)00051-9, 2001.
- 860 Janssen, N. A. H., Yang, A., Strak, M., Steenhof, M., Hellack, B., Gerlofs-Nijland, M. E., Kuhlbusch, T., Kelly, F., Harrison, R., Brunekreef, B., Hoek, G. and Cassee, F.: Oxidative potential of particulate matter collected at sites with different source characteristics, *Sci. Total Environ.*, 472, 572–581, doi:10.1016/j.scitotenv.2013.11.099, 2014.
- Jedynska, A., Hoek, G., Wang, M., Yang, A., Eeftens, M., Cyrus, J., Keuken, M., Ampe, C., Beelen, R., Cesaroni, G., Forastiere, F., Cirach, M., de Hoogh, K., De Nazelle, A., Nystad, W., Akhlaghi, H. M., Declercq, C., Stempfelet, M., Eriksen,
- 865 K. T., Dimakopoulou, K., Lanki, T., Meliefste, K., Nieuwenhuijsen, M., Yli-Tuomi, T., Raaschou-Nielsen, O., Janssen, N. A. H., Brunekreef, B. and Kooter, I. M.: Spatial variations and development of land use regression models of oxidative potential in ten European study areas, *Atmos. Environ.*, 150, 24–32, doi:10.1016/j.atmosenv.2016.11.029, 2017.
- Kelly, F. J.: Oxidative stress: Its role in air pollution and adverse health effects, *Occup. Environ. Med.*, 60(8), 612–616, doi:10.1136/oem.60.8.612, 2003.
- 870 Knaapen, A. M., Borm, P. J. A., Albrecht, C. and Schins, R. P. F.: Inhaled particles and lung cancer. Part A: Mechanisms, *Int. J. Cancer*, 109(6), 799–809, doi:10.1002/ijc.11708, 2004.
- Laden, F., Schwartz, J., Speizer, F. E. and Dockery, D. W.: Reduction in fine particulate air pollution and mortality: Extended follow-up of the Harvard Six Cities Study, *Am. J. Respir. Crit. Care Med.*, 173(6), 667–672, doi:10.1164/rccm.200503-443OC, 2006.
- 875 Laing, S., Wang, G., Briazova, T., Zhang, C., Wang, A., Zheng, Z., Gow, A., Chen, A. F., Rajagopalan, S., Chen, L. C., Sun, Q. and Zhang, K.: Airborne particulate matter selectively activates endoplasmic reticulum stress response in the lung and liver tissues, *Am. J. Physiol. - Cell Physiol.*, 299(4), 736–749, doi:10.1152/ajpcell.00529.2009, 2010.
- Lelieveld, J., Pozzer, A., Pöschl, U., Fnais, M., Haines, A. and Münzel, T.: Loss of life expectancy from air pollution compared to other risk factors: a worldwide perspective, *Cardiovasc. Res.*, 116(11), 1910–1917, doi:10.1093/cvr/cvaa025, 2020.
- 880 Lepeule, J., Laden, F., Dockery, D. and Schwartz, J.: Chronic exposure to fine particles and mortality: An extended follow-up of the Harvard six cities study from 1974 to 2009, *Environ. Health Perspect.*, 120(7), 965–970, doi:10.1289/ehp.1104660, 2012.
- Levy, J. I., Diez, D., Dou, Y., Barr, C. D. and Dominici, F.: A meta-analysis and multisite time-series analysis of the differential toxicity of major fine particulate matter constituents, *Am. J. Epidemiol.*, 175(11), 1091–1099, doi:10.1093/aje/kwr457, 2012.
- 885 Li, N., Hao, M., Phalen, R. F., Hinds, W. C. and Nel, A. E.: Particulate air pollutants and asthma: A paradigm for the role of oxidative stress in PM-induced adverse health effects, *Clin. Immunol.*, 109(3), 250–265, doi:10.1016/j.clim.2003.08.006, 2003.
- Li, N., Xia, T. and Nel, A. E.: The Role of Oxidative Stress in Ambient Particulate Matter Induced Lung Diseases and its Implications in the Toxicity of Engineered Nanoparticles, *Free Radic. Bio. Med.*, 9(44), 1689–1699, doi:10.1038/mp.2011.182, 2008.
- 890 Li, X., Jiang, L., Bai, Y., Yang, Y., Liu, S., Chen, X., Xu, J., Liu, Y., Wang, Y., Guo, X., Wang, Y. and Wang, G.: Wintertime

- aerosol chemistry in Beijing during haze period: Significant contribution from secondary formation and biomass burning emission, *Atmos. Res.*, 218(October 2018), 25–33, doi:10.1016/j.atmosres.2018.10.010, 2019.
- Li, Y.: Observational accuracy of sunrise and sunset times in the sixth century China, *Chinese J. Astron. Astrophys.*, 6(5),
895 doi:10.1088/1009-9271/6/5/16, 2006.
- Liang, L., Engling, G., Duan, F., Cheng, Y. and He, K.: Characteristics of 2-methyltetrols in ambient aerosol in Beijing, China, *Atmos. Environ.*, 59, 376–381, doi:10.1016/j.atmosenv.2012.05.052, 2012.
- Lim, Y. Bin and Ziemann, P. J.: Products and mechanism of secondary organic aerosol formation from reactions of n-alkanes with OH radicals in the presence of NO_x, *Environ. Sci. Technol.*, 39(23), 9229–9236, doi:10.1021/es051447g, 2005.
- 900 Liu, D., Harrison, R. M., Vu, T., Xu, J., Shi, Z., Li, L., Sun, Y. and Fu, P.: Estimation of biogenic and anthropogenic precursor contributions to secondary organic aerosol in Beijing using molecular tracers, *Prep.*, 2020.
- Liu, Q., Baumgartner, J., Zhang, Y., Liu, Y., Sun, Y. and Zhang, M.: Oxidative potential and inflammatory impacts of source apportioned ambient air pollution in Beijing, *Environ. Sci. Technol.*, 48(21), 12920–12929, doi:10.1021/es5029876, 2014.
- Liu, W. J., Xu, Y. S., Liu, W. X., Liu, Q. Y., Yu, S. Y., Liu, Y., Wang, X. and Tao, S.: Oxidative potential of ambient PM_{2.5}
905 in the coastal cities of the Bohai Sea, northern China: Seasonal variation and source apportionment, *Environ. Pollut.*, 236, 514–528, doi:10.1016/j.envpol.2018.01.116, 2018.
- Ma, Y., Cheng, Y., Qiu, X., Cao, G., Fang, Y., Wang, J., Zhu, T., Yu, J. and Hu, D.: Sources and oxidative potential of water-soluble humic-like substances (HULISWS) in fine particulate matter (PM_{2.5}) in Beijing, *Atmos. Chem. Phys.*, 18(8), 5607–5617, doi:10.5194/acp-18-5607-2018, 2018.
- 910 McWhinney, R. D., Badali, K., Liggio, J., Li, S. M. and Abbatt, J. P. D.: Filterable redox cycling activity: A comparison between diesel exhaust particles and secondary organic aerosol constituents, *Environ. Sci. Technol.*, 47(7), 3362–3369, doi:10.1021/es304676x, 2013a.
- McWhinney, R. D., Zhou, S. and Abbatt, J. P. D.: Naphthalene SOA: Redox activity and naphthoquinone gas-particle partitioning, *Atmos. Chem. Phys.*, 13(19), 9731–9744, doi:10.5194/acp-13-9731-2013, 2013b.
- 915 Meng, Z. and Zhang, Q.: Oxidative damage of dust storm fine particles instillation on lungs, hearts and livers of rats, *Environ. Toxicol. Pharmacol.*, 22(3), 277–282, doi:10.1016/j.etap.2006.04.005, 2006.
- Miller, M. R., Borthwick, S. J., Shaw, C. A., McLean, S. G., McClure, D., Mills, N. L., Duffin, R., Donaldson, K., Megson, I. L., Hadoke, P. W. F. and Newby, D. E.: Direct impairment of vascular function by diesel exhaust particulate through reduced bioavailability of endothelium-derived nitric oxide induced by superoxide free radicals, *Environ. Health Perspect.*, 117(4),
920 611–616, doi:10.1289/ehp.0800235, 2009.
- Miyata, R. and van Eeden, S. F.: The innate and adaptive immune response induced by alveolar macrophages exposed to ambient particulate matter, *Toxicol. Appl. Pharmacol.*, 257(2), 209–226, doi:10.1016/j.taap.2011.09.007, 2011.
- Moorthy, B., Chu, C. and Carlin, D. J.: Polycyclic aromatic hydrocarbons: From metabolism to lung cancer, *Toxicol. Sci.*, 145(1), 5–15, doi:10.1093/toxsci/kfv040, 2015.
- 925 Müller, L., Reinnig, M. C., Naumann, K. H., Saathoff, H., Mentel, T. F., Donahue, N. M. and Hoffmann, T.: Formation of 3-

- methyl-1,2,3-butanetricarboxylic acid via gas phase oxidation of pinonic acid - A mass spectrometric study of SOA aging, *Atmos. Chem. Phys.*, 12(3), 1483–1496, doi:10.5194/acp-12-1483-2012, 2012.
- Naes, T. and Martens, H.: Principal component regression in NIR analysis: viewpoints, background details and selection of components, *J. Chemom.*, 2(2), 155–167, 1988.
- 930 Ntziachristos, L., Froines, J. R., Cho, A. K. and Sioutas, C.: Relationship between redox activity and chemical speciation of size-fractionated particulate matter, *Part. Fibre Toxicol.*, 4, 1–12, doi:10.1186/1743-8977-4-5, 2007.
- Oberdorster, G., Ferin, J., Gelein, R., Soderholm, S. C. and Finkelstein, J.: Role of the alveolar macrophage in lung injury: Studies with ultrafine particles, *Environ. Health Perspect.*, 97, 193–199, doi:10.1289/ehp.97-1519541, 1992.
- Øvrevik, J., Refsnes, M., Låg, M., Holme, J. A. and Schwarze, P. E.: Activation of proinflammatory responses in cells of the
935 airway mucosa by particulate matter: Oxidant- and non-oxidant-mediated triggering mechanisms, *Biomolecules*, 5(3), 1399–1440, doi:10.3390/biom5031399, 2015.
- Paatero, P. and Tapper, U.: Postivie Matrix Factorisation: A non-negative factor model with optimal utilization of error estimates of data values, *Environmetrics*, 5(2), 111–126, 1994.
- Panagi, M., Fleming, Z. L., Monks, P. S., Ashfold, M. J., Wild, O., Hollaway, M., Zhang, Q., Squires, F. A. and Vande Hey,
940 J. D.: Investigating the regional contributions to air pollution in Beijing: A dispersion modelling study using CO as a tracer, *Atmos. Chem. Phys.*, 20(5), 2825–2838, doi:10.5194/acp-20-2825-2020, 2020.
- Pant, P., Baker, S. J., Shukla, A., Maikawa, C., Godri Pollitt, K. J. and Harrison, R. M.: The PM10 fraction of road dust in the UK and India: Characterization, source profiles and oxidative potential, *Sci. Total Environ.*, 530–531, 445–452, doi:10.1016/j.scitotenv.2015.05.084, 2015.
- 945 Paulson, S. E., Gallimore, P. J., Kuang, X. M., Chen, J. R., Kalberer, M. and Gonzalez, D. H.: A light-driven burst of hydroxyl radicals dominates oxidation chemistry in newly activated cloud droplets, *Sci. Adv.*, 5(5), 1–8, doi:10.1126/sciadv.aav7689, 2019.
- Pernigotti, D., Belis, C. A. and Spano, L.: SPECIEUROPE: The European data base for PM source profiles., *Atmos. Pollut. Res.*, 7(2), 307–314, 2016.
- 950 Platt, S. M., Haddad, I. El, Pieber, S. M., Huang, R. J., Zardini, A. A., Clairrotte, M., Suarez-Bertoa, R., Barmet, P., Pfaffenberger, L., Wolf, R., Slowik, J. G., Fuller, S. J., Kalberer, M., Chirico, R., Dommen, J., Astorga, C., Zimmermann, R., Marchand, N., Hellebust, S., Temime-Roussel, B., Baltensperger, U. and Prévôt, A. S. H.: Two-stroke scooters are a dominant source of air pollution in many cities, *Nat. Commun.*, 5(May), 1–7, doi:10.1038/ncomms4749, 2014.
- Pope, C. A. and Dockery, D. W.: Health effects of fine particulate air pollution: Lines that connect, *J. Air Waste Manag. Assoc.*, 56(6), 709–742, doi:10.1080/10473289.2006.10464485, 2006.
- 955 Presto, A. A., Miracolo, M. A., Donahue, N. M. and Robinson, A. L.: Secondary organic aerosol formation from high-NO_x Photo-oxidation of low volatility precursors: N-alkanes, *Environ. Sci. Technol.*, 44(6), 2029–2034, doi:10.1021/es903712r, 2010.
- Puthussery, J. V., Singh, A., Rai, P., Bhattu, D., Kumar, V., Vats, P., Furger, M., Rastogi, N., Slowik, J. G., Ganguly, D.,

- 960 Prevot, A. S. H., Tripathi, S. N. and Verma, V.: Real-Time Measurements of PM 2.5 Oxidative Potential Using a Dithiothreitol Assay in Delhi, India , *Environ. Sci. Technol. Lett.*, doi:10.1021/acs.estlett.0c00342, 2020.
- Risom, L., Møller, P. and Loft, S.: Oxidative stress-induced DNA damage by particulate air pollution, *Mutat. Res. - Fundam. Mol. Mech. Mutagen.*, 592(1–2), 119–137, doi:10.1016/j.mrfmmm.2005.06.012, 2005.
- Saffari, A., Daher, N., Shafer, M. M., Schauer, J. J. and Sioutas, C.: Seasonal and spatial variation in reactive oxygen species activity of quasi-ultrafine particles (PM0.25) in the Los Angeles metropolitan area and its association with chemical composition, *Atmos. Environ.*, 79, 566–575, doi:10.1016/j.atmosenv.2013.07.058, 2013.
- 965 Saffari, A., Daher, N., Shafer, M. M., Schauer, J. J. and Sioutas, C.: Seasonal and spatial variation in dithiothreitol (DTT) activity of quasi-ultrafine particles in the Los Angeles Basin and its association with chemical species, *J. Environ. Sci. Heal. - Part A Toxic/Hazardous Subst. Environ. Eng.*, 49(4), 441–451, doi:10.1080/10934529.2014.854677, 2014.
- 970 Schauer, J. J., Kleeman, M. J., Cass, G. R. and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 2. C1 through C30 organic compounds from medium duty diesel trucks, *Environ. Sci. Technol.*, 33(10), 1578–1587, doi:10.1021/es980081n, 1999.
- See, S. W., Wang, Y. H. and Balasubramanian, R.: Contrasting reactive oxygen species and transition metal concentrations in combustion aerosols, *Environ. Res.*, 103(3), 317–324, doi:10.1016/j.envres.2006.08.012, 2007.
- 975 Shen, H. and Anastasio, C.: A comparison of hydroxyl radical and hydrogen peroxide generation in ambient particle extracts and laboratory metal solutions, *Atmos. Environ.*, 46(530), 665–668, doi:10.1016/j.atmosenv.2011.10.006, 2012.
- Shen, R., Liu, Z., Liu, Y., Wang, L., Li, D., Wang, Y., Wang, G., Bai, Y. and Li, X.: Typical polar organic aerosol tracers in PM2.5 over the North China Plain: Spatial distribution, seasonal variations, contribution and sources, *Chemosphere*, 209, 758–766, doi:10.1016/j.chemosphere.2018.06.133, 2018.
- 980 Shi, Z., Vu, T., Kotthaus, S., Harrison, R. M., Grimmond, S., Yue, S., Zhu, T., Lee, J., Han, Y., Demuzere, M., Dunmore, R. E., Ren, L., Liu, D., Wang, Y., Wild, O., Allan, J., Acton, W. J., Barlow, J., Barratt, B., Beddows, D., Bloss, W. J., Calzolari, G., Carruthers, D., Carslaw, D. C., Chan, Q., Chatzidiakou, L., Chen, Y., Crilley, L., Coe, H., Dai, T., Doherty, R., Duan, F., Fu, P., Ge, B., Ge, M., Guan, D., Hamilton, J. F., He, K., Heal, M., Heard, D., Hewitt, C. N., Hollaway, M., Hu, M., Ji, D., Jiang, X., Jones, R., Kalberer, M., Kelly, F. J., Kramer, L., Langford, B., Lin, C., Lewis, A. C., Li, J., Li, W., Liu, H., Liu, J.,
- 985 Loh, M., Lu, K., Lucarelli, F., Mann, G., Mcfiggans, G., Miller, M. R., Mills, G., Monk, P., Nemitz, E., Ouyang, B., Palmer, P. I., Percival, C., Popoola, O., Reeves, C., Rickard, A. R., Shao, L., Shi, G., Spracklen, D., Stevenson, D., Sun, Y., Sun, Z., Tao, S., Tong, S., Wang, Q., Wang, W., Wang, X., Wang, X., Wang, Z., Wei, L., Whalley, L., Wu, X., Wu, Z., Xie, P., Yang, F., Zhang, Q., Zhang, Y., Zhang, Y. and Zheng, M.: Introduction to the special issue “ In-depth study of air pollution sources and processes within Beijing and its surrounding region, , 7519–7546, 2019.
- 990 Shinyashiki, M., Eiguren-Fernandez, A., Schmitz, D. A., Di Stefano, E., Li, N., Linak, W. P., Cho, S. H., Froines, J. R. and Cho, A. K.: Electrophilic and redox properties of diesel exhaust particles, *Environ. Res.*, 109(3), 239–244, doi:10.1016/j.envres.2008.12.008, 2009.
- Steimer, S., Patton, D., Vu, T., Panagi, M., Monks, P., Harrison, R., Fleming, Z., Shi, Z. and Kalberer, M.: Seasonal Differences

- in the Composition of Organic Aerosols in Beijing: a Study by Direct Infusion Ultrahigh Resolution Mass Spectrometry, *Atmos. Chem. Phys. Discuss.*, (January), 1–26, doi:10.5194/acp-2019-1009, 2020.
- Steimer, S. S., Delvaux, A., Campbell, S. J., Gallimore, P. J., Grice, P., Howe, D. J., Pitton, D., Claeys, M., Hoffmann, T. and Kalberer, M.: Synthesis and characterisation of peroxydicarboxylic acids as proxies for highly oxygenated molecules (HOMs) in secondary organic aerosol, *Atmos. Chem. Phys.*, 18(15), 10973–10983, doi:10.5194/acp-18-10973-2018, 2018.
- Subramanian, R., Donahue, N. M., Bernardo-Bricker, A., Rogge, W. F. and Robinson, A. L.: Contribution of motor vehicle emissions to organic carbon and fine particle mass in Pittsburgh, Pennsylvania: Effects of varying source profiles and seasonal trends in ambient marker concentrations, *Atmos. Environ.*, 40(40), 8002–8019, doi:10.1016/j.atmosenv.2006.06.055, 2006.
- Tan, W. C., Qiu, D., Li, B. L., Ng, T. P., Lee, S. H., Van Eeden, S. F., D’Yachkova, Y. and Hogg, J. C.: The human bone marrow response to acute air pollution caused by forest fires, *Am. J. Respir. Crit. Care Med.*, 161(4 Pt 1), 1213–1217, doi:10.1164/ajrccm.161.4.9904084, 2000.
- Tao, F., Gonzalez-Flecha, B. and Kobzik, L.: Reactive oxygen species in pulmonary inflammation by ambient particulates, *Free Radic. Biol. Med.*, 35(4), 327–340, doi:10.1016/S0891-5849(03)00280-6, 2003.
- Tapparo, A., Di Marco, V., Badocco, D., D’Aronco, S., Soldà, L., Pastore, P., Mahon, B. M., Kalberer, M. and Giorio, C.: Formation of metal-organic ligand complexes affects solubility of metals in airborne particles at an urban site in the Po valley, *Chemosphere*, 241, 1–13, doi:10.1016/j.chemosphere.2019.125025, 2020.
- Tong, H., Arangio, A. M., Lakey, P. S. J., Berkemeier, T., Liu, F., Kampf, C. J., Brune, W. H., Pöschl, U. and Shiraiwa, M.: Hydroxyl radicals from secondary organic aerosol decomposition in water, *Atmos. Chem. Phys.*, 16(3), 1761–1771, doi:10.5194/acp-16-1761-2016, 2016.
- Tong, H., Lakey, P. S. J., Arangio, A. M., Socorro, J., Kampf, C. J., Berkemeier, T., Brune, W. H., Pöschl, U. and Shiraiwa, M.: Reactive oxygen species formed in aqueous mixtures of secondary organic aerosols and mineral dust influencing cloud chemistry and public health in the Anthropocene, *Faraday Discuss.*, 200, 251–270, doi:10.1039/c7fd00023e, 2017.
- Tong, H., Lakey, P. S. J., Arangio, A. M., Socorro, J., Shen, F., Lucas, K., Brune, W. H., Pöschl, U. and Shiraiwa, M.: Reactive Oxygen Species Formed by Secondary Organic Aerosols in Water and Surrogate Lung Fluid, *Environ. Sci. Technol.*, 52, 11642–11651, doi:10.1021/acs.est.8b03695, 2018.
- Tuet, W. Y., Chen, Y., Xu, L., Fok, S., Gao, D., Weber, R. J. and Ng, N. L.: Chemical oxidative potential of secondary organic aerosol (SOA) generated from the photooxidation of biogenic and anthropogenic volatile organic compounds, , 839–853, doi:10.5194/acp-17-839-2017, 2017.
- Valko, M., Morris, H. and Cronin, M. T. D.: Metals, Toxicity and Oxidative Stress, *Curr. Med. Chem.*, 12(10), 1161–1208, 2005.
- Venkatachari, P. and Hopke, P. K.: Development and laboratory testing of an automated monitor for the measurement of atmospheric particle-bound reactive oxygen species (ROS), *Aerosol Sci. Technol.*, 42(8), 629–635, 2008.
- Venkatachari, P., Hopke, P. K., Grover, B. D. and Eatough, D. J.: Measurement of particle-bound reactive oxygen species in urban aerosols, *J. Atmos. Chem.*, 50(1), 49–58, doi:10.1007/s10874-005-1662-z, 2005.

- Verma, V., Pakbin, P., Cheung, K. L., Cho, A. K., Schauer, J. J., Shafer, M. M., Kleinman, M. T. and Sioutas, C.: Physicochemical and oxidative characteristics of semi-volatile components of quasi-ultrafine particles in an urban atmosphere, *Atmos. Environ.*, 45(4), 1025–1033, doi:10.1016/j.atmosenv.2010.10.044, 2011.
- 1030 Verma, V., Fang, T., Guo, H., King, L., Bates, J. T., Peltier, R. E., Edgerton, E., Russell, A. G. and Weber, R. J.: Reactive oxygen species associated with water-soluble PM_{2.5} in the southeastern United States: Spatiotemporal trends and source apportionment, *Atmos. Chem. Phys.*, 14(23), 12915–12930, doi:10.5194/acp-14-12915-2014, 2014.
- Verma, V., Wang, Y., El-Afifi, R., Fang, T., Rowland, J., Russell, A. G. and Weber, R. J.: Fractionating ambient humic-like substances (HULIS) for their reactive oxygen species activity - Assessing the importance of quinones and atmospheric aging, *Atmos. Environ.*, 120, 351–359, doi:10.1016/j.atmosenv.2015.09.010, 2015a.
- 1035 Verma, V., Fang, T., Xu, L., Peltier, R. E., Russell, A. G., Ng, N. L. and Weber, R. J.: Organic aerosols associated with the generation of reactive oxygen species (ROS) by water-soluble PM_{2.5}, *Environ. Sci. Technol.*, 49(7), 4646–4656, doi:10.1021/es505577w, 2015b.
- 1040 Wang, G., Cheng, S., Wei, W., Zhou, Y., Yao, S. and Zhang, H.: Characteristics and source apportionment of VOCs in the suburban area of Beijing, China, *Atmos. Pollut. Res.*, 7(4), 711–724, doi:10.1016/j.apr.2016.03.006, 2016.
- Wang, H., Li, Z., Lv, Y., Zhang, Y., Xu, H., Guo, J. and Goloub, P.: Determination and climatology of diurnal cycle of atmospheric mixing layer height over Beijing 2013–2018: Lidar measurements and implication for airpollution, *Atmos. Chem. Phys.*, 1–25, doi:10.5194/acp-2020-175, 2020a.
- 1045 Wang, J., Jiang, H., Jiang, H., Mo, Y., Geng, X., Li, J., Mao, S., Bualert, S., Ma, S., Li, J. and Zhang, G.: Source apportionment of water-soluble oxidative potential in ambient total suspended particulate from Bangkok: Biomass burning versus fossil fuel combustion, *Atmos. Environ.*, 235(May), 117624, doi:10.1016/j.atmosenv.2020.117624, 2020b.
- Wang, Q., He, X., Zhou, M., Huang, D., Qiao, L., Zhu, S., Ma, Y., Wang, H., Li, L., Huang, C., Huang, X. H. H., Xu, W., Worsnop, D. R., Goldstein, A. H., Guo, H. and Yu, J. Z.: Hourly Measurements of Organic Molecular Markers in Urban Shanghai, China: Primary Organic Aerosol Source Identification and Observation of Cooking Aerosol Aging, *ACS Earth Sp. Chem.*, doi:10.1021/acsearthspacechem.0c00205, 2020c.
- 1050 Wang, Y., Plewa, M. J., Mukherjee, U. K. and Verma, V.: Assessing the cytotoxicity of ambient particulate matter (PM) using Chinese hamster ovary (CHO) cells and its relationship with the PM chemical composition and oxidative potential, *Atmos. Environ.*, 179(September 2017), 132–141, doi:10.1016/j.atmosenv.2018.02.025, 2018.
- 1055 Whalley, L. K., Slater, E. J., Woodward-Massey, R., Ye, C., Lee, J. D., Squires, F., Hopkins, J. R., Dunmore, R. E., Shaw, M., Hamilton, J. F., Lewis, A. C., Mehra, A., Worrall, S. D., Bacak, A., Bannan, T. J., Coe, H., Ouyang, B., Jones, R. L., Crilley, L. R., Kramer, L. J., Bloss, W. J., Vu, T., Kotthaus, S., Grimmond, S., Sun, Y., Xu, W., Yue, S., Ren, L., Joe, W., Acton, F., Hewitt, C. N., Wang, X., Fu, P., Heard, D. E. and Whalley, L.: Evaluating the sensitivity of radical chemistry and ozone formation to ambient VOCs and NO_x in Beijing, , (x), 1–41, doi:10.5194/acp-2020-785, 2020.
- 1060 World Health Organisation: Ambient Air Pollution: A Global Assessment of Exposure and Burden of Disease., 2016.
- Wragg, F. P. H., Fuller, S. J., Freshwater, R., Green, D. C., Kelly, F. J. and Kalberer, M.: An automated online instrument to

- quantify aerosol-bound reactive oxygen species (ROS) for ambient measurement and health-relevant aerosol studies, *Atmos. Meas. Tech.*, 9(10), 4891–4900, doi:10.5194/amt-9-4891-2016, 2016.
- Xu, J., Liu, D., Wu, X., Vu, T. V., Zhang, Y. Z., Fu, P., Sun, Y., Xu, W., Ji, D., Harrison, R. M. and Shi, Z.: Source apportionment of Fine Aerosol at an Urban Site of Beijing Using a Chemical Mass Balance Model, *Prep.*, 2020a.
- 1065 Xu, J., Hu, W., Liang, D. and Gao, P.: Technology Photochemical impacts on the toxicity of PM_{2.5}, *Crit. Rev. Environ. Sci. Technol.*, 0(0), 1–27, doi:10.1080/10643389.2020.1816126, 2020b.
- Yang, A., Wang, M., Eeftens, M., Beelen, R., Dons, E., Leseman, D. L. A. C., Brunekreef, B., Cassee, F. R., Janssen, N. A. H. and Hoek, G.: Spatial variation and land use regression modeling of the oxidative potential of fine particles, *Environ. Health Perspect.*, 123(11), 1187–1192, doi:10.1289/ehp.1408916, 2015.
- 1070 Yu, L., Wang, G., Zhang, R., Zhang, L., Song, Y., Wu, B., Li, X., An, K. and Chu, J.: Characterization and source apportionment of PM_{2.5} in an urban environment in Beijing, *Aerosol Air Qual. Res.*, 13(2), 574–583, doi:10.4209/aaqr.2012.07.0192, 2013.
- Yu, S. Y., Liu, W. J., Xu, Y. S., Yi, K., Zhou, M., Tao, S. and Liu, W. X.: Characteristics and oxidative potential of atmospheric PM_{2.5} in Beijing: Source apportionment and seasonal variation, *Sci. Total Environ.*, 650, 277–287, doi:10.1016/j.scitotenv.2018.09.021, 2019.
- 1075 Zhao, H., Zheng, Y. and Li, C.: Spatiotemporal distribution of PM_{2.5} and O₃ and their interaction during the summer and winter seasons in Beijing, China, *Sustain.*, 10(12), doi:10.3390/su10124519, 2018.
- Zheng, M., Salmon, L. G., Schauer, J. J., Zeng, L., Kiang, C. S., Zhang, Y. and Cass, G. R.: Seasonal trends in PM_{2.5} source contributions in Beijing, China, *Atmos. Environ.*, 39(22), 3967–3976, doi:10.1016/j.atmosenv.2005.03.036, 2005.
- 1080