



Oxidation of low-molecular weight organic compounds in cloud droplets: global impact on tropospheric oxidants

Simon Rosanka¹, Rolf Sander², Bruno Franco³, Catherine Wespes³, Andreas Wahner¹, and Domenico Taraborrelli¹

¹Forschungszentrum Jülich GmbH, Institute of Energy and Climate Research, IEK-8: Troposphere, Jülich, Germany

²Atmospheric Chemistry Department, Max-Planck Institute of Chemistry, Mainz, Germany

³Université libre de Bruxelles (ULB), Spectroscopy, Quantum Chemistry and Atmospheric Remote Sensing (SQUARES), Brussels 1050, Belgium

Correspondence: Simon Rosanka (s.rosanka@fz-juelich.de)

Abstract. In liquid cloud droplets, superoxide anion ($O_{2(aq)}^-$) is known to quickly consume ozone ($O_{3(aq)}$), which is relatively insoluble. The significance of this reaction as tropospheric O_3 sink is sensitive to the abundance of $O_{2(aq)}^-$ and therefore to the production of its main precursor, hydroperoxyl radical ($HO_{2(aq)}$). The aqueous-phase oxidation of oxygenated volatile organic compounds (OVOCs) is the major source of $HO_{2(aq)}$ in cloud droplets. Hence, the lack of explicit aqueous-phase chemical kinetics in global atmospheric models leads to a general underestimation of clouds as O_3 sinks. In this study, the importance of in-cloud OVOC oxidation for tropospheric composition is assessed by using the Chemistry As A Boxmodel Application (CAABA) and the global atmospheric model ECHAM/MESSy (EMAC), which are both capable of explicitly representing the relevant chemical transformations. For this analysis, three different in-cloud oxidation mechanisms are employed: (1) one including the basic oxidation of $SO_{2(aq)}$ via $O_{3(aq)}$ and $H_2O_{2(aq)}$, which thus represents the capabilities of most global models, (2) the more advanced standard EMAC mechanism, which includes inorganic chemistry and simplified degradation of methane oxidation products, and (3) the detailed in-cloud OVOC oxidation scheme Jülich Aqueous-phase Mechanism of Organic Chemistry (JAMOC). By using EMAC, the global impact of each mechanism is assessed focusing mainly on tropospheric volatile organic compounds (VOCs), HO_x ($HO_x=OH+HO_2$), and O_3 . This is achieved by performing a detailed HO_x and O_3 budget analysis in the gas- and aqueous-phase. The resulting changes are evaluated against O_3 and methanol (CH_3OH) satellite observations from the Infrared Atmospheric Sounding Interferometer (IASI) for 2015. In general, the explicit in-cloud oxidation leads to an overall reduction of predicted OVOCs levels, and reduces EMAC's overestimation of some OVOCs in the tropics. The in-cloud OVOC oxidation shifts the HO_2 production from the gas- to the aqueous-phase. As a result, the O_3 budget is perturbed with scavenging being enhanced and the gas-phase chemical losses being reduced. With the simplified in-cloud chemistry, about 13 Tg a^{-1} of O_3 are scavenged, which increases to 336 Tg a^{-1} when JAMOC is used. The highest O_3 reduction of 12 % is predicted in the upper troposphere/lower stratosphere (UTLS). These changes in the free troposphere significantly reduce the modelled tropospheric ozone columns, which are known to be generally overestimated by EMAC and other global atmospheric models.



1 Introduction

Aqueous-phase chemistry in cloud droplets differs significantly from gas-phase chemistry, mainly due to photolysis enhanced
25 by scattering effects within cloud droplets (Bott and Zdunkowski, 1987; Mayer and Madronich, 2004), faster reaction rates,
and chemical reactions that do not occur in the gas-phase (Herrmann, 2003; Epstein and Nizkorodov, 2012). Moreover, the
conversion of nitrogen monoxide (NO) to nitrogen dioxide (NO₂) by peroxy radicals (RO₂) essentially does not take place
in liquid droplets because NO is very insoluble. Compared to gas-phase chemistry, models of aqueous-phase chemistry still
suffer from large uncertainties and most global models only include rudimentary implementations (Ervens, 2015). In general,
30 warm (liquid) clouds can act as a sink for ozone (O₃) and its precursors in the troposphere. Figure 1 gives an overview of the
inorganic aqueous-phase chemistry for O_{3(aq)} according to the mechanism by Staehelin et al. (1984). When O₃ is taken up
into cloud droplets, it is mainly destroyed via:



The superoxide anion O_{2(aq)}⁻ is in equilibrium with its conjugate acid, the hydroperoxyl radical (HO_{2(aq)}):



Here, HO_{2(aq)} is either scavenged from the gas phase or produced by photo-oxidation inside the cloud droplet. The realistic
representation of clouds as O₃ sinks is thus sensitive to a proper representation of HO_{2(aq)} in cloud droplets.

The importance of aqueous-phase chemistry for tropospheric O₃ has already been the topic of many earlier studies. Lelieveld
and Crutzen (1990) proposed that clouds strongly influence O₃, HO_x (HO_x=HO₂+OH), and NO_x (NO_x=NO+NO₂). They
40 concluded that under high-NO_x conditions, the net-O₃ production is decreased by as much as 40 %. However, Liang and Jacob
(1997) suggested that Lelieveld and Crutzen (1990) grossly overestimated the impact of clouds on O₃ because they made the
arduous assumption that the methyl peroxy radical (CH₃O₂) could have the same solubility as HO₂. They predicted that clouds
reduce tropospheric O₃ by less than 3 % in summer. A major aqueous-phase source of HO_{2(aq)} is the oxidation of water soluble
oxygenated volatile organic compounds (OVOCs). By not considering additional in-cloud HO_{2(aq)} sources, Liang and Jacob
45 (1997) underestimated O_{2(aq)}⁻ concentrations dampening the in-cloud destruction of O_{3(aq)}. An extensive and explicit in-cloud
OVOC oxidation scheme for global models, including the global ECHAM/MESSy Atmospheric Chemistry (EMAC) model, is
currently not available. Most global models only include basic sulfur dioxide (SO_{2(aq)}) oxidation as the only in-cloud O_{3(aq)}
destruction pathway in the aqueous-phase (Ervens, 2015). By neglecting in-cloud OVOC oxidation, aqueous-phase HO_{2(aq)}
concentrations are very likely underestimated in EMAC and other global models. Thus, it is expected that global atmospheric
50 models underestimate clouds as O₃ sinks. To date no global estimates of O₃ loss by scavenging has been reported.

Within this study, the global importance of in-cloud OVOC oxidation on tropospheric volatile organic compounds (VOCs),
HO_x, and O₃ is addressed. For the detailed in-cloud OVOC oxidation scheme, the Jülich Aqueous-phase Mechanism of Organic
Chemistry (JAMOC) suitable for global model applications is developed and implemented into the atmospheric chemistry
mechanism Module Efficiently Calculating the Chemistry of the Atmosphere (MECCA) in our companion paper by Rosanka
55 et al. (2020a). In JAMOC, the phase transfer of species containing up to ten carbon atoms is taken into account and a selection of



species containing up to four carbon atoms is considered to react in the aqueous-phase. Isoprene (C₅H₈), the most abundantly emitted VOC, is not explicitly dissolved in cloud droplets but many of its oxidation products explicitly react inside cloud droplets. In the aqueous phase, OVOCs mainly react with hydroxyl radicals (OH_(aq)) during daytime and with nitrate radicals (NO_{3(aq)}) during nighttime.

60 In this study, JAMOC is implemented into the global model EMAC (Sect. 2). The performance of JAMOC is compared to the performance of an aqueous-phase mechanism including only minimal aqueous-phase chemistry and of the standard mechanism of EMAC (each presented in Sect. 2.1). In order to understand the mechanistic behind the impact of in-cloud OVOC oxidation on a single air parcel, a box-model study is performed in Sect. 3. Afterwards, the impact on a global scale is analysed (Sect. 4). The analysis focuses on a selection of VOCs, HO_x, and O₃. The multiphase chemistry of JAMOC is expected to impact
65 tropospheric organic acids, which will be the topic of a further study. When considering the global O₃ budget, odd oxygen is also analysed to account for rapid cycling between species of the O_x family. In the scope of this study, O_x is defined as:

$$\begin{aligned} O_x \equiv & O + O_3 + NO_2 + 2 \times NO_3 + 3 \times N_2O_5 + HNO_3 + HNO_4 + ClO + HOCl + ClNO_2 + ClNO_3 + BrO + \\ & HOBr + BrNO_2 + 2 \times BrNO_3 + PANs + PNs + ANs + NPs \end{aligned} \quad (1)$$

where PANs are peroxyacyl nitrates, PNs are alkyl peroxy nitrates, ANs are alkyl nitrates, and NPs are nitrophenols. In Sect. 4, all performed EMAC simulations are evaluated against satellite observations of O₃ and methanol (CH₃OH) obtained
70 from the Infrared Atmospheric Sounding Interferometer (IASI). Model uncertainties are discussed in Sect. 5, followed by a general conclusion (Sect. 6).

2 Modelling approach

The aqueous- and gas-phase mechanisms are presented in Sect. 2.1. They are used within two different modelling frameworks: a box-model and a global atmospheric model. The box-model, used to investigate the local impact on an air parcel, is presented
75 in Sect. 2.2, and the global chemical atmospheric model in Sect. 2.3. Section 2.4 provides an overview of all simulations performed in this study.

2.1 The chemical mechanism

The study is based on the comparison of three different aqueous-phase mechanisms (Sect. 2.1.1). While they are characterised by different complexity, especially in the species and reactions taken into account, they are all coupled to the same gas-phase
80 mechanism (Sect. 2.1.2).

2.1.1 Aqueous-phase

The first aqueous-phase mechanism includes the uptake of a few soluble compounds, their acid-base equilibria, and the oxidation of SO_{2(aq)} via O_{3(aq)} and H₂O_{2(aq)}. This mechanism was applied by Jöckel et al. (2006) and is considered to represent the capabilities of most global models (Ervens, 2015). The second aqueous-phase mechanism includes an advanced scheme,



85 representing more than 150 reactions (Tost et al., 2007; Jöckel et al., 2016). It includes in-cloud $\text{HO}_{x(\text{aq})}$ chemistry and the
destruction of $\text{O}_{3(\text{aq})}$ by $\text{O}_{2(\text{aq})}^-$, but misses a detailed in-cloud OVOC oxidation scheme. This mechanism can be considered
to be the current standard mechanism used in EMAC. The last aqueous-phase mechanism is the complex OVOC oxidation
90 scheme JAMOC developed in our companion paper by Rosanka et al. (2020a). This mechanism is based on the box-model
mechanism CLOUD Explicit Physico-chemical Scheme (CLEPS 1.0, Mouchel-Vallon et al. (2017)). In order to make it appli-
cable for global models, Rosanka et al. (2020a) reduced the number of species (i.e. only a selection of species containing up
to four-carbon atoms), which react within the aqueous-phase. Still, the phase transfer of soluble species containing up to ten
carbon atoms is represented in JAMOC. In addition to CLEPS, Rosanka et al. (2020a) extended JAMOC by: (1) simulating
hydration and dehydration explicitly, (2) taking the oligomerisation of formaldehyde, glyoxal and methylglyoxal into account,
(3) adding further aqueous-phase photolysis reactions, and (4) considering the gas-phase photo-oxidation of new outgassed
95 species. A complete description of JAMOC, including a list of all reactions, is available in Rosanka et al. (2020a). Even though
Fenton's chemistry is an in-cloud source of $\text{OH}_{(\text{aq})}$, this chemistry is not considered in this study (switched off in JAMOC),
due to missing global iron (Fe) distributions and emissions in EMAC. The associated uncertainties for excluding this $\text{OH}_{(\text{aq})}$
sources are discussed in Sect. 5.

2.1.2 Gas-phase

100 The Mainz Organic Mechanism (MOM, Sander et al., 2019) is used to model gas-phase chemistry, containing an extensive
oxidation scheme for isoprene (Taraborrelli et al., 2009, 2012; Nölscher et al., 2014), monoterpenes (Hens et al., 2014), and
aromatics (Cabrera-Perez et al., 2016). In addition, comprehensive reactions schemes are considered for the modelling of the
chemistry of NO_x , HO_x , CH_4 , and anthropogenic linear hydrocarbons. VOCs are oxidised by OH , O_3 , and NO_3 , whereas
 RO_2 reacts with HO_2 , NO_x , NO_3 , and undergoes self- and cross-reactions (Sander et al., 2019). When the complex in-cloud
105 OVOC oxidation scheme JAMOC is coupled to MOM, MOM is modified following the gas-phase additions as described in
Rosanka et al. (2020a).

2.2 Chemistry box-model CAABA

Each of the three mechanisms is implemented in the Chemistry As A Boxmodel Application (CAABA, Sander et al., 2019), in
order to investigate their implications on a single air-parcel under predefined atmospheric conditions. It is capable to numeri-
110 cally integrate the multiphase chemical mechanism as one single system of Ordinary Differential Equations (ODEs) with ap-
propriate phase-transfer reactions (Sander, 1999; Kerkweg et al., 2007). The Kinetic Pre-Processor (KPP version 2.2.3, Sandu
and Sander (2006)) is used in MECCA to integrate these ODE systems. Further, photolysis, emissions and dry deposition of
chemical species, and the exchange with other air masses outside the box (entrainment) are represented in a simplified manner.

In this study, an air parcel during summer is simulated at mid-latitude with a constant temperature of 278 K and a relative
115 humidity of 70 %. The same initial conditions are used as proposed in Rosanka et al. (2020a, see their Table 2), but the NO
emissions are neglected in this study. In order to represent a realistic atmospheric cloud event and investigate the impact of
the newly developed aqueous-phase mechanism, three atmospheric conditions are modelled during the simulated day. First,



CAABA is initialised at 0 UTC and no cloud droplets are present until 12 UTC. At 12 UTC a cloud is formed with droplet radii of 20 μm and a liquid water content of $3.0 \times 10^{-1} \text{ g L}^{-1}$. After one hour the cloud evaporates and all species outgas. The rest of the day is simulated using the same conditions as before the cloud event.

2.3 Global model EMAC

The ECHAM/MESSy Atmospheric Chemistry (EMAC) model is a numerical chemistry and climate simulation system that includes submodels describing tropospheric and middle atmospheric processes and their interaction with oceans, land, and human influences (Jöckel et al., 2010). It uses the second version of the Modular Earth Submodel System (MESSy2) to link multi-institutional computer codes. The core atmospheric model is the 5th generation European Centre Hamburg general circulation Model (ECHAM5, Roeckner et al., 2003). For the present study, EMAC (ECHAM5 version 5.3.02, MESSy version 2.54.0) is used at T63L90MA resolution, i.e. with a spherical truncation of T63 (corresponding to a quadratic Gaussian grid of approximately 1.875° by 1.875° degrees in latitude and longitude) with 90 vertical hybrid pressure levels up to 0.01 hPa.

In contrast to CAABA, gas- and aqueous-phase chemistry are calculated separately. In order to model the gas-phase mechanism MOM in the troposphere and stratosphere, the submodel MECCA is used. The SCAVenging submodel (SCAV, Tost et al., 2006) is used to simulate the removal of trace gases and aerosol particles by clouds and precipitation. SCAV calculates the transfer of species into and out of rain and cloud droplets using the Henry's law equilibrium, acid dissociation equilibria, oxidation-reduction reactions, heterogeneous reactions on droplet surfaces, and aqueous-phase photolysis reactions (Tost et al., 2006). In this study, SCAV is used to calculate the three aqueous-phase mechanisms presented in Sect. 2.1.1. Like MECCA, SCAV treats the aqueous-phase mechanism as an ODE system and uses KPP (version 1) to solve it. This operator splitting is necessary because the ODE systems resulting from the combination of gas-phase and in-cloud aqueous-phase mechanisms would suffer from (1) a higher stiffness due to fast acid-base equilibria and phase-transfer reactions, and (2) load imbalances on High-Performance Computing (HPC) systems due to the sparsity of clouds. In both MECCA and to some degree SCAV, tagging systems are used to calculate detailed gas- and aqueous-phase O_x and HO_x budgets. These systems allow to estimate the full implications of the aqueous-phase mechanism on atmospheric chemistry. The tagging system of MECCA is more sophisticated and allows to obtain reaction rates from multiple reactions and combine them into a single tracer (Gromov et al., 2010). For the tropospheric O_x budget, the gas-phase chemical production and loss, and the scavenging and wet deposition are taken into account by using MECCA and SCAV, respectively. Additionally, the dry deposition of O_x and many MOM species is calculated by the submodel Dry DEPosition (DDEP, Kerkweg et al., 2006) using its default scheme.

The MESSy submodel Model of Emissions of Gases and Aerosols from Nature (MEGAN) is used to model biogenic VOC emissions (Guenther et al., 2006). Global isoprene emissions are scaled to the best estimate of Sindelarova et al. (2014), which is 595 Tg a^{-1} . Biomass burning emission fluxes are calculated using the MESSy submodel BIOBURN, which calculates these fluxes based on biomass burning emission factors and dry matter combustion rates. For the latter, Global Fire Assimilation System (GFAS) data are used, which are based on satellite observations of fire radiative power from the Moderate Resolution Imaging Spectroradiometer (MODIS) satellite instruments (Kaiser et al., 2012). The biomass burning emission factors for VOCs are based on Akagi et al. (2011).



The submodel SORBIT (Jöckel et al., 2010) is used to sample the model state along sun-synchronous satellite orbits, at the time of the satellite overpass, and to compare the model outputs to satellite observations obtained from the Infrared Atmospheric Sounding Interferometer (IASI, Clerbaux et al., 2009) onboard the Metop-A (IASI-A) and Metop-B (IASI-B) satellites. In particular, the Fast Optimal Retrievals on Layers for IASI Ozone (FORLI-O₃, version 20151001; see Hurtmans et al. (2012), for a description of the retrievals) are used for the comparison of tropospheric O₃ columns. In general, when analysing tropospheric burdens and budgets, the standard EMAC tropopause definition is used. Here, the tropopause is defined in the extra tropics using potential vorticity whereas temperature lapse rates are used in the tropics (Jöckel et al., 2006). However, when comparing modelled tropospheric O₃ columns to IASI-FORLI measurements, the troposphere is defined as ranging from the ground to 300 hPa in order to limit the influences of the stratospheric O₃, but to include the altitude of maximum sensitivity of IASI in the troposphere (Wespes et al., 2017). Moreover, this allows to avoid larger errors that affect the O₃ retrievals in the upper troposphere/lower stratosphere (UTLS) and that result in a positive column bias (Boynard et al., 2016). The evaluation of simulation results against global observational datasets of VOC abundance can be performed for only a few species. Daily global distributions of methanol total columns are available from IASI-A and IASI-B observations, using a neural network-based retrieval approach (Franco et al., 2018). Due to the limited vertical information on methanol that is contained in the IASI spectra, only total columns have been retrieved. Since the neural network-based retrievals do not rely on scene-dependent a priori information, no averaging kernels are produced and the retrieved total columns are meant to be compared at face value with model data (see Franco et al., 2018, and references therein). For this purpose, the IASI methanol measurements have been daily averaged on the EMAC T63 spatial grid. The comparisons with IASI O₃ and methanol data are associated with some observational uncertainties. IASI retrievals are obtained in the thermal infrared range, resulting in an especially high sensitivity to clouds. Appropriate filters are applied in order to account for cloud-contaminated IASI scenes observations. These filters are based on defined cloud cover thresholds, using information from the Eumetcast operational processing system (August et al., 2012). The fractional cloud cover threshold depends on the species observed. For O₃ and methanol, all observations with a fractional cloud cover above 13 % (Wespes et al., 2017) and 25 % (Franco et al., 2018) have been excluded, respectively. The IASI methanol retrievals are less sensitive to the presence of residual clouds since no radiative transfer model is used, resulting in a higher threshold for methanol. Of course, it cannot be completely ruled out that individual IASI measurements are locally affected by residual clouds that passed the filtering. However, due to the huge dataset used for the seasonal averages, it is considered that such an effect is diluted and is globally negligible.

2.4 Simulations performed

In both modelling frameworks, multiple simulations are performed. In CAABA, the impact of each aqueous-phase mechanism on a single air-parcel is investigated. For comparison, the same day is simulated in CAABA using the same initial conditions but excluding the specific cloud event at 12 UTC. The global impact is investigated by performing a reference and two sensitivity simulations with EMAC. Global simulations without any in-cloud aqueous-phase chemistry lead to unrealistic concentrations of O₃ and other chemical species (Tost et al., 2007). Therefore, the reference simulation includes the minimal scavenging mechanism (in the following called Scm). The two sensitivity simulations use the standard EMAC (in the following called



ScSta) and the detailed OVOC oxidation aqueous-phase mechanism (in the following called ScJAMOC). For consistency, the same simulation names are used for the CAABA simulations. In EMAC, the years 2014 and 2015 are simulated, where 2014 is discarded as a spin-up. All simulations were performed at the Jülich Supercomputing Center with the JURECA/JUWELS clusters (Jülich Supercomputing Centre, 2018, 2019).

190 3 Box model results

Figure 2 shows the time evolution of selected gas-phase species for the different aqueous-phase mechanisms Scm, ScSta and ScJAMOC for the cloud scenario of CAABA (see Sect. 2.2). For comparison, the results of the no-cloud scenario are also shown. Both Scm and ScSta have only little impact on most of the OVOCs explicitly treated in JAMOC. For some OVOCs, the phase transfer considered in Scm and ScSta leads to reduced gas-phase concentrations during the cloud event. After the cloud
195 evaporates, gas-phase concentrations are slightly higher compared to the no-cloud scenario, since the OVOCs transferred into the cloud droplet generally do not oxidise. Within ScSta, a subset of these OVOCs (containing one carbon atom) are oxidised leading to a slight reduction compared to Scm. In contrast, ScJAMOC efficiently removes OVOCs, leading to overall reduced OVOC concentrations. Glyoxal, one of the OVOC examples presented in Fig. 2, is completely removed from the gas-phase and quickly hydrated within the cloud droplet. The irreversible oxidation of its hydrated forms and oligomers leads to a reduction
200 of in-cloud glyoxal concentrations. In the gas-phase, glyoxal itself is produced by the oxidation of hydrocarbons. Due to low aqueous-phase HO_x concentrations during the cloud event, the oxidation of these hydrocarbons is reduced. After the cloud evaporates, the higher hydrocarbon concentrations lead to some glyoxal being produced.

Each mechanism leads to changes in most gas-phase radical concentrations. As soon as the cloud droplets form, gas-phase HO_x is reduced due to the uptake of radicals and radical precursors within the first minutes. This becomes evident when
205 inspecting the results of Scm: in this mechanism, the uptake of HO_x is not taken into account. Here, the gas-phase HO₂ concentration is still reduced due to the uptake of a few HO₂ sources (e.g. formaldehyde). In the case of the other mechanisms, the uptake of HO_x is explicitly considered and leads to an additional reduction in gas-phase concentrations when the cloud forms. In the case of ScJAMOC and, to some extent, of ScSta, the additional partitioning of OVOCs into the cloud droplet leads to a further decrease of gas-phase HO_x concentrations. The reduction of OH is in line with other modelling studies for
210 cloud events (Tilgner et al., 2013). When the cloud evaporates, radicals and radical sources are transferred to the gas phase. For ScJAMOC, the efficient in-cloud oxidation of radical sources induces significantly lower HO_x concentrations after the cloud evaporates. The photolysis of OVOCs and their oxidation within cloud droplets cause an increase of HO_{x(aq)} of about 50 %.

When the cloud forms, gas-phase O₃ is reduced in comparison to the no-cloud scenario because of its reactive uptake into the cloud droplet. Within Scm, O_{3(aq)} only reacts with SO_{2(aq)}, leading to only a little reduction in gas-phase O₃. This
215 reduction is more pronounced for ScSta and ScJAMOC due to additional aqueous-phase sinks and the uptake of HO₂ into the cloud droplet. For ScJAMOC, the reduction in O₃ is larger due to the additional aqueous-phase HO_{2(aq)} sources from OVOC oxidation. In the gas phase, the significantly reduced HO₂ concentrations cause NO_x to increase (HO₂ being the major sink of NO_x). However, it mostly dampens the production of O₃ after the cloud event.



4 Global impact on atmospheric composition

220 This section evaluates the importance of in-cloud OVOC oxidation on a global scale by focusing on VOCs (Sect. 4.1), and HO_x (Sect. 4.2). The importance for tropospheric O_3 is discussed in Sect. 4.3.

4.1 Impact on tropospheric VOCs

The extensive aqueous-phase OVOC oxidation scheme JAMOC considers many VOC sinks. These significantly influence the concentrations of tropospheric VOCs. In general, VOCs can be split into primarily emitted VOCs and OVOCs mostly formed from secondary production (e.g., oxidation of primarily emitted VOCs). The main global source of primarily emitted VOCs are biogenic processes. The largest biogenic emissions take place in the equatorial region (e.g. Amazon Basin, Central Africa) with additional emissions in the Northern (NH) and Southern Hemispheric (SH) extratropics. Isoprene, the most abundant biogenic VOC, is only slightly influenced by ScJAMOC. The yearly mean tropospheric burden increases from 204 Gg (Scm) to 213 Gg (ScJAMOC). This increase is caused by changes in OH concentrations, the main isoprene oxidant (see Sect. 4.2). Primarily emitted VOCs are quickly oxidised in the lower troposphere, leading to low concentrations in the free troposphere. The top panel of Fig. 3 shows the zonal mean of the sum of all OVOCs that are explicitly treated in JAMOC for Scm. High OVOC concentrations are predicted in the lower troposphere and at lower latitudes, consistent with strong terrestrial biogenic emissions at the Earth surface. By the general upward transport in the equatorial region, OVOCs are transported into the free troposphere. Due to deep convection events in the same region, OVOCs are even transported into the dry tropical upper troposphere. The lower panel of Fig. 3 shows the changes of the sum of OVOCs explicitly treated in JAMOC induced in ScJAMOC. Overall, the tropospheric OVOC burden is reduced with the largest change in the tropical free troposphere. The frequent occurrence of clouds in this region and the high OVOC concentrations lead to an efficient removal of gas-phase OVOCs. The ubiquity of clouds in the NH extratropics allows for additional removal of OVOCs from the gas-phase. These results are in line with the box-model results presented above (see Fig. 2). The efficient removal of OVOCs in warm clouds significantly affect the OVOC levels in the dry tropical upper troposphere. Here, these OVOCs act as an important HO_x source, potentially influencing the production of O_3 (Jaeglé et al., 2001).

Table 1 provides an overview of the annual tropospheric burden for a selection of VOCs explicitly treated in JAMOC. As shown in Fig. 3, the global burden of most VOCs is reduced due to the uptake and oxidation processes implemented in ScJAMOC. Because of the low number of VOCs containing one carbon atom treated in ScSta, changes between Scm and ScSta are only minor. The burden of some VOCs even increases in ScSta, which is caused by reduced HO_x concentrations (see Sect. 4.2). The impact in ScJAMOC differs for each VOC, with some VOCs in terms of absolute changes being efficiently removed whereas others are only slightly impacted. The varying efficiency of the VOCs removal by clouds is explained by differences in their Henry's law constants, accommodation coefficients, and aqueous-phase reactivities. The burden of methanol, the OVOC containing one carbon atom for which the highest absolute change is predicted, is reduced by about 1000 Gg. For methyl hydroperoxide the total change is lower but the relative reduction is higher, which is due to a slightly higher solubility and overall higher reaction rate constants for the oxidation via $\text{OH}_{(\text{aq})}$ and $\text{NO}_{3(\text{aq})}$. Formaldehyde is reduced by about 16 %.



Even though ethanol has a Henry's law constant similar to methanol, the relative reduction is still significantly smaller, due to a slower aqueous-phase oxidation. Ethylene glycol has a slow aqueous-phase oxidation, but a very high solubility, which results in a substantial reduction of its tropospheric burden. The opposite holds for ethyl hydroperoxide, which is four times
255 less soluble but undergoes a fast aqueous-phase oxidation. This leads to a relative change that is similar to the one of ethylene glycol. Acetaldehyde is the only OVOC for which an enhanced burden is predicted. This is partially due to newly implemented in-cloud sources, but in particular to the aqueous-phase oxidation of methylglyoxal yielding pyruvic acid, which is a known source of acetaldehyde (Berges and Warneck, 1992).

Figure 4 shows the seasonal mean methanol column for the IASI observations. In addition, the differences of Sc_m vs IASI
260 and Sc_{JAMOC} vs Sc_m are shown. The highest methanol columns occur close to its major biogenic sources (e.g., Amazon Basin, boreal forests). When using Sc_m , EMAC underestimates methanol at mid-latitudes and overestimates it close to its main tropical biogenic sources (see center column Fig. 4). Both these model inconsistencies are caused by an incorrect spatial distribution of biogenic emissions. The submodel MEGAN, used to simulate biogenic methanol emissions (see Sect. 2.3), estimates yearly biogenic methanol emissions of 104 Tg a^{-1} , which is close to the 103 Tg a^{-1} estimated by Millet et al. (2008,
265 their Table 2). However, the spatial distribution of biogenic emissions from MEGAN is different to their predictions. Compared to Millet et al. (2008), MEGAN significantly overestimates biogenic emissions in the Amazon Basin, but underestimates emissions at mid- and high-latitudes. EMAC simulates the Amazon basin too dry in the dry season (September–November, SON) and consequently too hot (Hagemann and Stacke, 2015). The biogenic emissions in MEGAN are temperature-dependent and generally higher temperatures induce higher emissions. Thus, the positive bias of surface temperatures in EMAC leads to an
270 overestimation in the Amazon basin. Additionally, uncertainties for all coefficients used in MEGAN, related to the emissions of methanol and primarily emitted VOCs (e.g. isoprene) further influence the incorrect emission distribution. EMAC also underestimates methanol over the oceans. In the current simulation setup, the ocean is represented to only act as a methanol sink but should be considered to be a source as well over certain oceans (e.g. over the Pacific, see Millet et al., 2008). However, EMAC models the ocean as a net sink with an uptake of about 2.1 Tg a^{-1} , which is smaller than the predicted net sink from
275 Millet et al. (2008) of 16 Tg a^{-1} . It is thus expected that there is an additional deficiency in the representation of the gas-phase chemistry of methanol in MOM. Still, when using Sc_{JAMOC} , the model bias for methanol is partially resolved (see right column of Fig. 4). In areas where the sources are expected to be modelled correctly (i.e. Central Africa, East Asia), the additional in-cloud OVOC oxidation leads to a reduction of methanol partially resolving the model bias in these regions. However, Sc_{JAMOC} is not able to completely resolve the model bias over the Amazon basin. The positive model bias away
280 from its major sources (i.e. over oceans) is reduced and partially resolved. Especially during NH autumn (SON), the strong model bias over the East Pacific and the South Atlantic Ocean is reduced. At the same time, a high overestimation for Sc_m is observed southeast of India over the Indian Ocean. The strong El Niño event in 2015–2016 led to droughts, draining the already dry Indonesian peatland. This drying, in combination with widespread deforestation, led to strong Indonesian fires, emitting large amounts of VOCs (Parker et al., 2016). This positive model bias is strongly reduced when in-cloud methanol oxidation
285 is taken into account (Sc_{JAMOC}).



To the best of our knowledge, glyoxal satellite retrievals from the Ozone Monitoring Instrument (OMI, Levelt et al., 2006) are only available up to 2014, while the TROPOspheric Monitoring Instrument (TROPOMI) started its operations in late 2017. Levelt et al. (2018) report that this is due to detector degradation and the challenging nature of glyoxal retrievals. A detailed analysis for the year 2007 is performed by Alvarado et al. (2014). Figure 5 gives the yearly mean integrated glyoxal column for Scm and the changes introduced by ScJAMOC. In the gas-phase, glyoxal is an oxidation product of hydrocarbons. Therefore, high glyoxal concentrations are predicted by EMAC close to strong biogenic hydrocarbon sources (e.g. Amazon Basin). As found with the CAABA box model, atmospheric glyoxal levels are significantly reduced by the chemical loss in cloud droplets with ScJAMOC (see Table 1). When comparing these results to satellite retrievals from Alvarado et al. (2014, their Fig. 9), it can be concluded that the spatial distribution is reasonably well captured by Scm. However, glyoxal levels are generally overestimated in regions where biogenic emissions dominate. The additional sink introduced in ScJAMOC leads to a significant reduction of the model bias, especially in the Amazon Basin and over Central Africa. However, the model bias is not fully resolved yet in the Amazon Basin. Here, the too high biogenic hydrocarbon emissions from MEGAN are the cause of an overestimated production of glyoxal. It is important to keep in mind that the comparability with these satellite retrievals is limited due to a different year simulated. It is still expected that the yearly mean spatial distribution of biogenic emissions is comparable for both years and mainly varies in their magnitude. To conclude, when using JAMOC (ScJAMOC) the representation of methanol and glyoxal gas-phase concentrations is significantly improved within EMAC.

4.2 Impact on tropospheric HO_x

VOCs play an important role in the production and loss of OH and HO₂. Thus, the additional uptake of VOCs will influence the tropospheric OH budget. In the troposphere, OH is primarily produced by the reaction of O(¹D) with H₂O. Here, the main source of O(¹D) is the photolysis of O₃. Figure 6 gives the zonal mean of the total OH production of Scm and the changes predicted by ScJAMOC. OH is mainly produced in the lower troposphere by both its primary and secondary sources, whereas in the upper troposphere secondary sources dominate. Table 2 gives an overview on the tropospheric gas-phase OH sources and sinks. With ScJAMOC, the gross OH formation decreases by about 7.3 % from 280.2 Tmol a⁻¹ to 259.8 Tmol a⁻¹. This finding is consistent with the box model results (Fig. 2). The uptake and oxidation of VOCs in the aqueous-phase reduce the contribution of VOCs to the OH production. However, the major reduction in the OH production is caused by overall reduced tropospheric O₃ concentrations. Specifically, the two largest O₃ sinks, namely the OH production induced by O₃ photolysis and the reaction of O₃ with HO₂, are reduced by 8.5 %. O₃ has a long atmospheric lifetime, leading to a low spatial variability in the reduction in tropospheric O₃. However, the reduction in VOC concentrations has a high spatial variability (see Fig. 3), largely determining the spatial distribution of the reduction in the total OH formation by ScJAMOC (lower panel Fig. 6). The removal of VOCs containing one carbon atom presents the largest contribution to the reduction. The reduction in HO_x leads to an additional reduction in the destruction of OH from HO_x cross-reactions (HO₂ + OH and OH + OH). The OH budget presented in this study compares well with earlier EMAC studies by Lelieveld et al. (2016), which used the standard in-cloud EMAC mechanism (ScSta). The relative contributions of each OH source and sink in ScSta are comparable with their reported budgets. However, they report a lower tropospheric gross OH formation of 251.2 Tmol a⁻¹ while using the same tropopause



320 definition. This difference is mainly related to different years simulated (leading to different emissions), and a lower model resolution used (T42L31, approximately 2.8 by 2.8 degrees in latitude and longitude with 31 vertical layers). Specifically, the lower number of tropospheric levels is expected to influence tropospheric budgets.

Figure 7 shows the zonal HO₂ production for Scm and the changes predicted in ScJAMOC. Due to the fast interconversion within the HO_x family, the spatial distribution and magnitude of the HO₂ production are similar to the production of OH. Table 3 gives the gas-phase HO₂ budget for each simulation. The HO₂ production changes from about 315 Tmol a⁻¹ to 290 Tmol a⁻¹ for Scm and ScJAMOC, respectively. Lower VOC concentrations lead to a reduction in the HO₂ production. Here, the influence of VOCs containing one carbon atom is the highest (see Table 1). Thus, VOCs become less important as a HO₂ sink. The highest reduction is caused by the reduced availability of HO₂, significantly reducing radical-radical reactions as a HO₂ sink.

330 Table 2 and 3 also provide the in-cloud budgets for OH_(aq) and HO_{2(aq)}. The representation of the aqueous-phase chemistry of OH_(aq) in clouds strongly affects the HO_{2(aq)} production. The aqueous-phase budget of OH_(aq) differs significantly between ScSta and ScJAMOC, which explicitly treat in-cloud HO_{x(aq)} kinetics. ScJAMOC has the highest total OH_(aq) production with more than 12 Tmol a⁻¹, which is about four times higher than in ScSta. The higher increase, compared to the box-model (Sect. 3), is attributed to the specific box-model scenario (Sect. 2.2 and Rosanka et al., 2020a, their Table 2). In both ScSta and ScJAMOC, most OH_(aq) is formed via the destruction of O_{3(aq)}. In ScJAMOC, the photolysis of OVOCs leads to the second highest formation of OH_(aq). Here, OVOCs containing one carbon atom contribute the most, while most OH_(aq) is formed from methyl hydroperoxide. Due to higher radical concentrations, the reactions of OH_(aq) with O_{3(aq)} and radical-radical reactions in ScJAMOC contribute about four times as much to the loss of HO_{x(aq)} compared to ScSta. The oxidation of OVOCs is the major OH_(aq) sink, with OVOCs containing one carbon atom contributing the most. This oxidation leads to the most significant production of HO_{2(aq)}, followed by OVOC photolysis. Due to increased aqueous-phase OH_(aq) and H₂O_{2(aq)} concentrations, the oxidation of H₂O_{2(aq)} increases by a factor of four in ScJAMOC. The destruction of O_{3(aq)} leads to a reduction in O_{2(aq)}⁻. This equilibrium is therefore the dominant HO_{2(aq)} sink for both ScSta and ScJAMOC, since HO_{2(aq)} is in equilibrium with O_{2(aq)}⁻ (R2). In the literature no in-cloud HO_{x(aq)} budget on a global scale has been presented so far. The novel in-cloud aqueous-phase budgets can thus not be compared to earlier studies.

345 4.3 Impact on tropospheric O₃

The efficient oxidation of OVOCs by cloud droplets leads to elevated aqueous-phase HO_{2(aq)} concentrations accelerating the in-cloud O_{3(aq)} destruction. This has a significant impact on tropospheric O₃ levels predicted by EMAC. Table 4 gives the O_x budget for the three simulations. The chemical production increases for ScSta compared to Scm. Slightly elevated NO_x concentrations lead to an increased contribution of methylperoxy radicals and RO₂ reactions with NO, compensating the reduced production from HO₂. For ScJAMOC, the chemical production decreases by about 150 Tg a⁻¹ (2.6 %), mainly caused by an overall reduction in HO₂ (see Sect. 4.2) and in RO₂ radicals due to the uptake and explicit oxidation of VOCs. The chemical loss on the other hand is reduced by about 90 Tg a⁻¹ (1.7 %) and about 420 Tg a⁻¹ (8.0 %) for ScSta and ScJAMOC, respectively. This reduction is mainly attributed to an overall reduction in tropospheric levels of O₃ and HO_x. The



loss by dry deposition reduces by about 50 Tg a^{-1} (5.6 %) for ScJAMOC, due to generally reduced surface O_3 concentrations.
355 The largest change in the O_x budget is related to scavenging processes. O_x scavenging increases from about 150 Tg a^{-1} (Scm)
to about 260 Tg a^{-1} (73.3 %) and 480 Tg a^{-1} (220.0 %) for ScSta and ScJAMOC, respectively. Here, the biggest increase
occurs for O_3 scavenging, due to the accelerated $\text{O}_{3(\text{aq})}$ destruction by enhanced $\text{HO}_{2(\text{aq})}$ (R1), which in turn enhances the
 O_3 uptake. These changes in the O_x budget terms lead to a reduced O_3 burden. Compared to the literature, the O_3 burden
from ScJAMOC is closer to the observational estimate from satellite retrievals for the same time period of 287-311 Tg in the
360 60°S - 60°N latitudinal band and closer to the global tropospheric burden of 324 Tg derived from the IASI-FORLI observations
(Gaudel et al., 2018, their Table 5). However, it is important to take into account that different tropopause definitions are used in
the extra tropics. In Gaudel et al. (2018), the tropopause definition for IASI-FORLI is the WMO tropopause altitude definition,
based on the temperature lapse rate (WMO, 1957). In this study, potential vorticity is used as tropopause definition in the extra
tropics (see Sect. 2.3). All three O_x budgets (Table 4) compare well with a recent multi-model comparison of Young et al.
365 (2018, see their Fig. 3). The chemical loss and chemical production get closer to the multi-model mean of about 4500 Tg a^{-1}
and about 4950 Tg a^{-1} , respectively. The tropospheric O_3 burden in ScJAMOC is now lower than the multi-model mean of
about 340 Tg but closer to the observational estimate from Ziemke et al. (2011). The increased Stratospheric-Tropospheric
Exchange (STE) is still lower than the multi-model mean (about 540 Tg a^{-1}) and the observational estimate of 489 Tg a^{-1}
by Olsen et al. (2013). The tropospheric O_3 lifetime is reduced by one day, due to higher relative changes in the O_x loss than
370 in the tropospheric O_3 burden.

Figure 8 gives the zonal net- O_x production for Scm and the changes in ScJAMOC. In general, O_x is produced where
 NO_x concentrations are high (close to the surface and in the upper troposphere). In the free troposphere, above the planetary
boundary layer (PBL), the increased destruction of O_3 over the ocean leads to an overall net O_x loss in the zonal mean.
The changes in the chemical production and in the loss of O_x , and the increase of scavenging lead to changes in the net- O_x
375 production in ScJAMOC. At the surface, the net O_x production increases. Here, the efficient uptake of O_3 sink precursors
overcompensates the reduction in the chemical production and leads to a reduced chemical loss. This increase mainly occurs
over continental regions. In the free troposphere above the PBL, the net- O_x change is reduced leading to an increased O_x
destruction. This is directly caused by the efficient uptake of HO_2 , VOCs, and O_3 precursors in this cloud dominated region in
ScJAMOC. In the tropical UTLS, VOCs are an important HO_2 source. The efficient removal of VOCs in the lower troposphere
380 reduces the total VOC mass transported into this region (see Fig. 3). The chemical production of O_x is therefore reduced in the
tropical UTLS, due to a limited availability in HO_2 .

Figure 9 and 10 give the yearly mean surface concentration and the zonal mean of O_3 for Scm and the changes of ScJAMOC.
In general, O_3 concentrations are higher in the NH with the highest values found over continental areas. Overall, surface O_3
slightly decreases for ScJAMOC with the maximum mean reduction of about 4 nmol/mol. The decrease in surface O_3 is
385 very low where the net- O_x production increases. The highest reduction in O_3 is predicted in the UTLS, where tropospheric O_3
concentrations are the highest. Here, O_3 is reduced by more than 12 % for ScJAMOC. Even though the total lower tropospheric
change is similar in both hemispheres, the relative reduction is higher in the SH (NH: about 4 %, SH: about 10 %).



Figure 11 shows the seasonal tropospheric integrated O_3 columns from IASI-FORLI O_3 retrievals. In addition, the differences of Scm with respect to IASI-FORLI and ScJAMOC with respect to Scm are shown. As explained previously, the comparison is performed here by using tropospheric O_3 column integrated between the Earth surface and 300 hPa (see Sect. 2.3). To meaningfully compare the model profile to the IASI observation, the non-uniform sensitivity of the IASI-FORLI retrievals to the O_3 vertical distribution was accounted for by applying the averaging kernels. It provided the model vertical distribution of O_3 as would be seen by IASI. For this purpose, the model profiles sampled at the place and time of the IASI overpasses (see Sect. 2.3) were first vertically interpolated to the IASI pressure levels. Then the smoothing of the model profiles to the lower vertical resolution of IASI was performed following Rodgers (2000). In order to take the specific scene of each IASI observation into account, the averaging kernels of the different observations contained in the model grid cell have all been considered to smooth the gridded model profile. The smoothed model profiles are finally averaged to derive the smoothed gridded model profile. In Scm, EMAC generally overestimates tropospheric O_3 in the tropics and at mid-latitudes regionally by more than 10 DU. This general overestimation is lower but consistent with an earlier EMAC study by Jöckel et al. (2016). They report an overestimation of up to 15 DU (see their Fig. 29), based on a comparison of a nudged simulation with OMI O_3 retrievals using EMACs standard aqueous-phase mechanism (here ScSta). These differences can be attributed to a much simplified gas-phase chemical mechanism, a lower spatial resolution (inducing artificial dilution of NO_x point sources, Fiore et al., 2003), and different emission data sets. At higher latitudes, especially during the NH winter (December-February, DJF) and spring (March-May, MAM), EMAC slightly underestimates tropospheric O_3 . In ScJAMOC, the overall modelled O_3 bias compared to IASI-FORLI is reduced by 1-2 DU, improving the representation of O_3 in EMAC. Here, due to the long lifetime of O_3 , the reduction in tropospheric O_3 is not limited to the typical cloud dominated and precipitation regions. This demonstrates the importance of a proper representation of in-cloud $O_{3(aq)}$ and OVOC oxidation chemistry in global models. By not taking these processes into account, as done in most global models (Ervens, 2015), tropospheric O_3 is overestimated. It is expected that the bias reduction is even more pronounced for the complete troposphere (when using the standard EMAC definition, see Sect. 2.3), since the highest relative reduction in O_3 is predicted in the UTLS above 300 hPa (Fig. 10). Similar to methanol, Scm strongly overestimates the tropospheric O_3 column west of Indonesia over the Indian Ocean in the NH autumn. This overestimation is also linked to the strong Indonesian peatland fires (Parker et al., 2016). Due to the ongoing Asian monsoon, the emitted VOCs are quickly transported to higher altitudes, where they act as O_3 precursors. The efficient upward transport of the biomass burning tracers isocyanic acid (HNCO) and hydrogen cyanide (HCN) during the summer monsoon phase, has already been investigated in earlier EMAC simulations by Rosanka et al. (2020b). In the same region, surface O_3 is also substantially reduced in ScJAMOC (Fig. 9). These results indicate that soluble OVOCs are efficiently removed by clouds. As a consequence, the reactive uptake of O_3 is enhanced and O_3 production dampens. This leads to a reduction of the modelled bias for this region and period when using JAMOC.



5 Model uncertainties

420 In our companion paper (Rosanka et al., 2020a), uncertainties related to the kinetic data used in JAMOC are already discussed. The global model simulations performed in this study suffer from additional uncertainties mainly attributed to (1) the representation of VOC emissions, and (2) missing sources of key oxidants. Each uncertainty will be shortly discussed in this section.

As demonstrated for methanol (see Sect. 4.1), a satisfactory reproduction of tropospheric VOC concentrations strongly depends on the realistic representation of VOC emissions. As pointed out earlier, the highest uncertainty is introduced by the biogenic emission submodel MEGAN. For instance, isoprene emissions are very sensitive to temperature and light. These uncertainties are not well quantified. Drought stress also affects isoprene emissions and it is estimated to reduce the emissions by 17-50 % globally (Jiang et al., 2018; Sindelarova et al., 2014). Additionally, biomass burning emissions in Indonesia are potentially underestimated. Parker et al. (2016) pointed out that in the monsoon period of 2015, a high fraction of the Indonesian fire emissions originates from peatland, which is known to produce significantly high VOC emissions (Akagi et al., 2011). In the GFAS retrievals used for biomass burning, the dominant fire type in Indonesia is assigned to tropical forest fires with the exceptions of a few grid points. The strength of VOC emissions for the Indonesian fire period in 2015 is therefore underestimated. It is thus expected that when using JAMOC and a realistic combination of peatland and tropical forest fire types the overestimation of tropospheric O_3 in this region and time period will be further reduced (see Sect. 4.3 and Fig. 11).

435 Fenton chemistry is a major source of in-cloud $OH_{(aq)}$ (Deguillaume et al., 2004). Even though these reactions are available in JAMOC, Fenton chemistry is not taken into account in this study, due to missing global iron (Fe) distributions and emissions in EMAC. However, Scanza et al. (2018) present an approach to implement these into a global model. Realising this approach in EMAC would make Fenton chemistry feasible in the future. From the literature, no global modelling study is known that couples this $OH_{(aq)}$ source to a detailed in-cloud OVOC oxidation scheme, making it difficult to estimate its impact on a global scale. In the highly idealised box modelling study of Mouchel-Vallon et al. (2017), most $OH_{(aq)}$ (63 %) is produced from Fenton chemistry (see their supplemental material SM5). This indicates the importance of Fenton chemistry in areas with high iron concentrations. The major source of atmospheric iron is mineral dust. Fossil fuel and biomass burning also emit some iron. Thus, iron concentrations are high close to deserts with the highest concentrations in the Sahara, Lut, Thar, and Arabian desert (Wang et al., 2015, their Fig. 6). Not considering this $OH_{(aq)}$ source catalysed by iron might lead to an underestimation of OVOC oxidation rates in the aqueous-phase. In particular Central Africa, a region with high biogenic VOC emissions, might be influenced by Fe being transported from the Sahara. In addition, mineral dust will be transported over the tropical Atlantic to the Amazon basin. Here, the missing $OH_{(aq)}$ source could be responsible for the underestimation of in-cloud OVOC oxidation and thus the destruction of $O_3(aq)$.

450 To conclude, the impact of the in-cloud OVOC chemistry on the tropospheric composition estimated in this study, is influenced by some model and observational uncertainties. However, the findings of the simulations performed in this study are still consistent with earlier studies and improve the representation of a selection of OVOCs and the EMAC bias towards high O_3 concentrations. Due to their complexity, reducing the model uncertainties introduced by biogenic and biomass burning



emissions, and missing aqueous-phase Fenton chemistry, is outside the scope of this study. Model representation of the latter is expected to substantially increase the oxidation rate of OVOCs in the cloud droplets and aerosols. Additional global modelling
455 studies need to be performed to address these issues.

6 Conclusions

In this study, the influence of in-cloud oxidation of soluble OVOCs on the tropospheric gas-phase composition was studied. This was achieved by implementing the extensive aqueous-phase OVOC oxidation scheme JAMOC, initially presented by Rosanka et al. (2020a), into the global model EMAC. The mechanism considers a selection of VOCs containing up to four
460 carbon atoms, their acid/base and/or hydration/dehydration equilibria, and their reactions with $\text{OH}_{(\text{aq})}$, $\text{NO}_{3(\text{aq})}$, and other oxidants (if available). Additionally, the phase transfer of species containing up to ten carbon atoms is taken into account. In addition to the EMAC simulations, a representative cloud droplet was simulated in the box-model CAABA in order to understand all processes involved.

When in-cloud OVOC oxidation is taken into account, VOCs are efficiently removed from the gas-phase leading to generally
465 reduced tropospheric VOC burdens. The reduction in modelled methanol and glyoxal concentrations is in line with satellite retrievals. The overall reduction in VOC concentrations leads to lower formation rates of HO_x in the gas-phase. Higher in-cloud $\text{HO}_{2(\text{aq})}$ concentrations, formed from OVOC oxidation, lead to an accelerated destruction of $\text{O}_{3(\text{aq})}$ in clouds. In addition, the chemical production and loss of O_3 in the gas-phase are reduced due to lower VOC and HO_x concentrations. This results in a reduced O_3 burden and decreases EMAC's bias towards too high O_3 concentrations. In ScJAMOC, many secondary
470 organic aerosol (SOA) precursors are explicitly treated, impacting the formation of SOA (Blando and Turpin, 2000; Ervens et al., 2011; Ervens, 2015). The potentially enhanced SOA formation will further influence tropospheric HO_x chemistry and NO_2 photolysis, resulting in a higher reduction of tropospheric O_3 and EMAC's O_3 bias. However, studying the influence of in-cloud OVOC oxidation on SOA formation is outside the scope of this study.

The findings in this study demonstrate the importance of in-cloud chemistry on tropospheric O_3 . Most atmospheric global
475 models do not take detailed aqueous-phase chemistry into account (Ervens, 2015). With the minimal oxidation of $\text{SO}_{2(\text{aq})}$ by $\text{O}_{3(\text{aq})}$, which is representative for most global models, only about 13 Tg a^{-1} of O_3 are scavenged by clouds. With explicit in-cloud OVOC oxidation considered, O_3 scavenging increases to about 336 Tg a^{-1} . This estimate neglects the O_3 sink in deliquescent aerosols, which might turn out to be significant as well. The predicted O_3 loss by clouds is significantly higher than the results suggested by Liang and Jacob (1997). Even though regional changes might be in the order of the results
480 presented by Lelieveld and Crutzen (1990), the global reduction does not get close to their reduction of 40 %. To conclude, global models, which neglect explicit in-cloud OVOC oxidation, significantly underestimate clouds as O_3 sinks and show a general tendency to overestimate tropospheric O_3 .



Data availability. The simulation results are archived at the Jülich Supercomputing Centre (JSC) and are available on request. The IASI O₃ data processed with FORLI-O₃ v0151001 can be downloaded from the AERIS portal at <http://iasi.aeris-data.fr/O3/> (last access: 2 August 2020). The IASI methanol columns retrieved with the ANNI framework are available upon request.

Author contributions. SR and DT designed the study. SR performed the simulations and analysed the data with contributions from DT. BF and CW acted as IASI data providers and analysts. SR and DT discussed the results with contributions from RS, BF, and AW. The manuscript was prepared by SR with the help of all co-authors.

Competing interests. The authors declare that they have no competing of interest.

Acknowledgements. The work described in this paper has received funding from the Initiative and Networking Fund of the Helmholtz Association through the project “Advanced Earth System Modelling Capacity (ESM)”. The content of this paper is the sole responsibility of the author(s) and it does not represent the opinion of the Helmholtz Association, and the Helmholtz Association is not responsible for any use that might be made of the information contained. The authors gratefully acknowledge the Earth System Modelling Project (ESM) for funding this work by providing computing time on the ESM partition of the supercomputer JUWELS at the Jülich Supercomputing Centre (JSC). The authors gratefully acknowledge the computing time granted through JARA on the supercomputer JURECA at Forschungszentrum Jülich. IASI is a joint mission of EUMETSAT and the Centre National d’Etudes Spatiales (CNES, France). The authors acknowledge the AERIS data infrastructure for providing access to the IASI data in this study and ULB-LATMOS, in particular Daniel Hurtmans, for the development of the retrieval algorithms. The research in Belgium is funded by the Belgian State Federal Office for Scientific, Technical and Cultural Affairs, and the European Space Agency (ESA Prodex IASI Flow and B-AC SAF).



Table 1. Mean gas-phase tropospheric burden in 2015 for a selection of VOCs for Scm and the changes induced by ScSta and ScJAMOC. Burden values are given in Gg.

	Scm	Δ ScSta	Δ ScJAMOC
C₁-VOCs			
Formaldehyde	1212.3	- 46.6	- 204.2
Methanol	3279.3	- 341.0	- 998.8
Methyl hydroperoxide	1914.5	- 32.9	- 849.9
Hydroxymethyl hydroperoxide	67.8	+ 0.2	- 16.0
C₂-VOCs			
Ethanol	110.9	+ 0.4	- 16.6
Ethylene glycol	3.1	+ 0.1	- 1.4
Acetaldehyde	147.1	+ 1.7	+ 12.1
Glycolaldehyde	278.8	- 0.9	- 101.2
Glyoxal	44.6	0.0	- 12.7
Ethyl hydroperoxide	62.9	- 0.9	- 28.3
C₃-VOCs			
Methylglyoxal	181.8	- 0.6	- 35.3
Isopropyl hydro peroxide	13.0	- 0.2	- 4.6



Table 2. Global tropospheric mean gas- and aqueous-phase source and sink fluxes of OH for Scm and the changes induced by ScSta and ScJAMOC. All values are given in Tmol a^{-1} . The aqueous-phase budget is only based on cloud droplets. Rain droplets are not taken into account.

	Scm	Δ ScSta	Δ ScJAMOC
Gas-phase sources			
$\text{O}(^1\text{D}) + \text{H}_2\text{O}$	96.67	- 1.39	- 7.11
$\text{NO} + \text{HO}_2$	84.53	- 0.25	- 2.10
$\text{O}_3 + \text{HO}_2$	32.36	- 0.95	- 3.93
$\text{H}_2\text{O}_2 + h\nu$	26.70	- 0.85	- 1.39
OVOCs	30.40	- 0.30	- 5.82
Other	9.54	+ 0.01	- 0.02
Total	280.20	- 3.73	- 20.37
Gas-phase sinks			
$\text{OH} + \text{HO}_{y\text{g}}^{\text{a}}$	49.88	+ 0.06	- 1.90
$\text{OH} + \text{NO}_y^{\text{b}}$	4.73	+ 0.01	+ 0.11
$\text{OH} + \text{CH}_4$	32.85	- 0.02	- 0.35
$\text{OH} + \text{C}_1^{\text{c}}$	150.90	- 2.73	- 16.20
$\text{OH} + \text{C}_n\text{-VOCs}$	39.75	- 0.15	- 2.70
Other	2.09	0.00	0.00
Total	280.20	- 3.73	- 20.37
Aqueous-phase sources			
$\text{O}_3 + \text{O}_2^-$	-	+ 1.94	+ 6.30
$\text{H}_2\text{O}_2 + h\nu$	-	+ 0.95	+ 1.08
$\text{C}_1\text{-VOCs} + h\nu$	-	-	+ 4.71
$\text{C}_n\text{-VOCs} + h\nu$	-	-	+ 0.32
Other	-	+ 0.02	+ 0.02
Total	-	+ 2.91	+ 12.43
Aqueous-phase sinks			
$\text{OH} + \text{HO}_{y\text{aq}}^{\text{d}}$	-	+ 0.42	+ 2.20
$\text{C}_1\text{-VOCs}$	-	+ 2.40	+ 8.98
$\text{C}_n\text{-VOCs}$	-	-	+ 0.91
Other	-	+ 0.09	+ 0.34
Total	-	+ 2.91	+ 12.43

^a $\text{HO}_{y\text{g}} \equiv \text{H}_2, \text{O}_3, \text{H}_2\text{O}_2$, radical–radical reactions

^b $\text{NO}_y \equiv \text{NO}, \text{NO}_2, \text{HNO}_2, \text{HNO}_3, \text{HNO}_4, \text{NH}_3$, N-reaction products

^c $\text{C}_1 \equiv \text{CO}$, VOCs with one C-atom

^d $\text{HO}_{y\text{aq}} \equiv \text{O}_2^-, \text{H}_2\text{O}_2$, radical–radical reactions



Table 3. Global tropospheric mean gas- and aqueous-phase source and sink fluxes of HO₂ for Scm and the changes induced by ScSta and ScJAMOC. All values are given in Tmol a⁻¹. The aqueous-phase budget is only based on cloud droplets. Rain droplets are not taken into account.

	Scm	Δ ScSta	Δ ScJAMOC
Gas-phase sources			
OH + O ₃	12.51	- 0.18	- 0.71
H ₂ O ₂ + OH	13.86	- 0.44	- 0.56
HNO ₄ ^a	26.38	- 0.52	- 1.59
C ₁ -VOCs	214.71	- 3.58	- 17.76
C _n -VOCs	22.33	+ 0.01	- 0.64
Photolysis	24.64	- 0.47	- 3.88
Other	1.26	- 0.01	- 0.01
Total	315.69	- 5.19	- 25.15
Gas-phase sinks			
HO ₂ + O ₃	32.36	- 0.95	- 3.93
HO ₂ + OH	12.86	- 0.21	- 0.69
HO ₂ + HO ₂	77.34	- 2.33	- 8.37
HO ₂ + NO	84.53	- 0.25	- 2.10
HO ₂ + NO ₂ & NO ₃	27.31	- 0.44	- 1.58
C ₁ -VOCs + HO ₂	47.63	- 1.34	- 6.74
C _n -VOCs + HO ₂	26.85	- 0.22	- 2.08
Other	6.81	+ 0.55	+ 0.34
Total	315.69	- 5.19	- 25.15
Aqueous-phase sources			
Mass transfer	-	+ 0.60	+ 0.51
H ₂ O ₂ + OH	-	+ 0.38	+ 1.61
C ₁ -VOCs	-	+ 2.39	+ 10.80
C ₂ -VOCs	-	-	+ 0.92
Other	-	+ 0.01	+ 0.09
Total	-	+ 3.38	+ 13.93
Aqueous-phase sinks			
HO ₂ ⇌ O ₂ ⁻ + H ⁺	-	+ 2.68	+ 8.69
HO ₂ + HO _{y, aq} ^b	-	+ 0.69	+ 5.22
Other	-	+ 0.01	+ 0.02
Total	-	+ 3.38	+ 13.93

^a HNO₄ → NO₂ + HO₂

^b HO_{y, aq} ≡ O₂⁻, radical–radical reactions



Table 4. Detailed tropospheric O_x budget for Scm and the changes induced by ScSta and ScJAMOC. The gross terms as well as the relative contributions of the major contributors are given.

	Scm	Δ ScSta	Δ ScJAMOC
Sources [Tg a⁻¹]			
Chemical production	5895.6	+ 7.1	- 155.8
HO ₂ + NO	4050.3	- 12.8	- 101.3
CH ₃ O ₂ + NO	1084.8	+ 13.1	- 22.9
RO ₂ + NO	731.1	+ 6.7	- 30.8
Other	29.4	+ 0.1	+ 0.1
STE ^a	355.2	+ 5.6	+ 15.3
Sinks [Tg a⁻¹]			
Chemical loss	5254.7	- 91.2	- 423.2
O(¹ D) + H ₂ O	2317.3	- 35.0	- 167.3
HO ₂ + O ₃	1550.1	- 42.4	- 187.6
OH + O ₃	599.0	- 1.4	- 0.6
HOBr + hv	341.6	- 0.8	- 54.6
PhO + O ₃ ^b	215.4	+ 1.5	- 31.8
Other	231.3	- 4.1	- 81.5
Dry deposition	846.5	- 9.1	- 47.3
O ₃	801.6	- 9.4	- 47.1
Other	44.9	+ 0.3	- 0.2
Scavenging	149.7	+ 112.9	+ 329.7
O ₃	13.2	+ 104.4	+ 323.1
N ₂ O ₅	25.0	- 2.3	- 2.7
HNO ₃	111.5	- 0.3	- 1.0
Other	-	+ 11.2	+ 10.3
O₃ burden [Tg]	348.2	- 5.0	- 25.0
O₃ lifetime [days]	20.3	- 0.3	- 1.0

^a Stratospheric-Tropospheric Exchange

^b O₃ loss due reaction with phenoxy radicals from oxidation of aromatics
 (Taraborrelli et al., 2020)




Figure 1. Graphical representation of inorganic aqueous-phase ozone chemistry based on Staehelin et al. (1984).




Figure 2. Time evolution for gas-phase mixing ratios of the sum of all the OVOCs explicitly reacting in JAMOC (\sum VOCs = methanol + formaldehyde + methyl hydroperoxide + hydroxymethylhydroperoxide + ethanol + ethylene glycol + acetaldehyde + glycolaldehyde + glyoxal + hydroperoxide + methylglyoxal + isopropanol + isopropyl hydro peroxide + methacrolein + methyl vinyl ketone), glyoxal, glycolaldehyde, methylglyoxal, HO_2 , OH , NO_x , and O_3 within the boxmodel CAABA. The time when the cloud is present (between 12 and 13 UTC) is indicated by a blue background shading. Nighttime is indicated by a grey background shading. Mixing ratios are provided for no cloud event (black line), Scm (green line), ScSta (purple line) and ScJAMOC (red line). Note that lines may overlap.




Figure 3. Yearly zonal mean of the sum of all the OVOCs explicitly reacting in JAMOC (\sum VOCs = methanol + formaldehyde + methyl hydroperoxide + hydroxymethylhydroperoxide + ethanol + ethylene glycol + acetaldehyde + glycolaldehyde + glyoxal + hydroperoxide + methylglyoxal + isopropanol + isopropyl hydro peroxide + methacrolein + methyl vinyl ketone): for Scm (top) and in comparison to ScJAMOC (bottom). The yearly mean tropopause is depicted by a black line.




Figure 4. Seasonal (December-February, DJF; March-May, MAM; June-August, JJA; September-November, SON) mean integrated methanol column obtained from IASI satellite observations (left), of the Scm simulation in comparison to IASI observations (center), and of ScJAMOC in comparison to Scm (right).




Figure 5. Mean integrated tropospheric glyoxal column for Scm (top) and in comparison to ScJAMOC (bottom).




Figure 6. Zonal mean gross OH formation for Scm (top) and in comparison to ScJAMOC (bottom). The yearly mean tropopause is depicted by a black line.




Figure 7. Zonal mean gross HO₂ formation for Scm (top) and in comparison to ScJAMOC (bottom). The yearly mean tropopause is depicted by a black line.




Figure 8. Mean zonal net- O_x change for Scm (top) and in comparison to ScJAMOC (bottom). The yearly mean tropopause is depicted by a black line. Deposition in the lowest model layer is not taken into account.




Figure 9. Mean surface O_3 concentration for Scm (top) and in comparison to ScJAMOC (bottom).




Figure 10. Mean zonal O₃ concentration for Scm (top) and in comparison to ScJAMOC (bottom). The yearly mean tropopause is depicted by a red solid line. In addition, the 300 hPa tropopause layer used for the O₃ IASI-FORLI comparison (see Fig. 11) is depicted by a red dash-dotted line.




Figure 11. Seasonal (December-February, DJF; March-May, MAM; June-August, JJA; September-November, SON) tropospheric O₃ column comparison between IASI-FORLI satellite observations and EMAC. IASI-FORLI satellite observations (left), Scm simulation in comparison to IASI-FORLI observations (center), and ScJAMOC in comparison to Scm (right). For this comparison, the tropopause is defined at 300 hPa.



500 References

- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmospheric Chemistry and Physics*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- Alvarado, L. M. A., Richter, A., Vrekoussis, M., Wittrock, F., Hilboll, A., Schreier, S. F., and Burrows, J. P.: An improved glyoxal retrieval from OMI measurements, *Atmospheric Measurement Techniques*, 7, 4133–4150, <https://doi.org/10.5194/amt-7-4133-2014>, 2014.
- 505 August, T., Klaes, D., Schlüssel, P., Hultberg, T., Crapeau, M., Arriaga, A., O’Carroll, A., Coppens, D., Munro, R., and Calbet, X.: IASI on Metop-A: Operational Level 2 retrievals after five years in orbit, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 113, 1340–1371, <https://doi.org/10.1016/j.jqsrt.2012.02.028>, *three Leaders in Spectroscopy*, 2012.
- Berges, M. G. M. and Warneck, P.: Product Quantum Yields for the 350 nm Photodecomposition of Pyruvic Acid in Air, *Berichte der Bunsengesellschaft für physikalische Chemie*, 96, 413–416, <https://doi.org/10.1002/bbpc.19920960334>, 1992.
- 510 Blando, J. D. and Turpin, B. J.: Secondary organic aerosol formation in cloud and fog droplets: a literature evaluation of plausibility, *Atmospheric Environment*, 34, 1623–1632, [https://doi.org/10.1016/S1352-2310\(99\)00392-1](https://doi.org/10.1016/S1352-2310(99)00392-1), 2000.
- Bott, A. and Zdunkowski, W.: Electromagnetic energy within dielectric spheres, *J. Opt. Soc. Am. A*, 4, 1361–1365, <https://doi.org/10.1364/JOSAA.4.001361>, 1987.
- 515 Boynard, A., Hurtmans, D., Koukouli, M. E., Goutail, F., Bureau, J., Safieddine, S., Lerot, C., Hadji-Lazaro, J., Wespes, C., Pommereau, J.-P., Pazmino, A., Zyrichidou, I., Balis, D., Barbe, A., Mikhailenko, S. N., Loyola, D., Valks, P., Van Roozendael, M., Coheur, P.-F., and Clerbaux, C.: Seven years of IASI ozone retrievals from FORLI: validation with independent total column and vertical profile measurements, *Atmospheric Measurement Techniques*, 9, 4327–4353, <https://doi.org/10.5194/amt-9-4327-2016>, 2016.
- Cabrera-Perez, D., Taraborrelli, D., Sander, R., and Pozzer, A.: Global atmospheric budget of simple monocyclic aromatic compounds, *Atmospheric Chemistry and Physics*, 16, 6931–6947, <https://doi.org/10.5194/acp-16-6931-2016>, 2016.
- 520 Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., Wespes, C., and Coheur, P.-F.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmospheric Chemistry and Physics*, 9, 6041–6054, <https://doi.org/10.5194/acp-9-6041-2009>, 2009.
- Deguillaume, L., Leriche, M., Monod, A., and Chaumerliac, N.: The role of transition metal ions on HO_x radicals in clouds: a numerical evaluation of its impact on multiphase chemistry, *Atmospheric Chemistry and Physics*, 4, 95–110, <https://doi.org/10.5194/acp-4-95-2004>, 2004.
- 525 Epstein, S. A. and Nizkorodov, S. A.: A comparison of the chemical sinks of atmospheric organics in the gas and aqueous phase, *Atmospheric Chemistry and Physics*, 12, 8205–8222, <https://doi.org/10.5194/acp-12-8205-2012>, 2012.
- Ervens, B.: Modeling the Processing of Aerosol and Trace Gases in Clouds and Fogs, *Chemical Reviews*, 115, 4157–4198, <https://doi.org/10.1021/cr5005887>, 2015.
- 530 Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmospheric Chemistry and Physics*, 11, 11 069–11 102, <https://doi.org/10.5194/acp-11-11069-2011>, 2011.
- Fiore, A. M., Jacob, D. J., Mathur, R., and Martin, R. V.: Application of empirical orthogonal functions to evaluate ozone simulations with regional and global models, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD003151>, 2003.
- 535



- Franco, B., Clarisse, L., Stavrou, T., Müller, J.-F., Van Damme, M., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Taraborrelli, D., Clerbaux, C., and Coheur, P.-F.: A General Framework for Global Retrievals of Trace Gases From IASI: Application to Methanol, Formic Acid, and PAN, *Journal of Geophysical Research: Atmospheres*, 123, 13,963–13,984, <https://doi.org/10.1029/2018JD029633>, 2018.
- 540 Gaudel, A., Cooper, O. R., Ancellet, G., Barret, B., Boynard, A., Burrows, J. P., Clerbaux, C., Coheur, P. F., Cuesta, J., Cuevas, E., Doniki, S., Dufour, G., Ebojie, F., Foret, G., Garcia, O., Granados Muños, M. J., Hannigan, J. W., Hase, F., Huang, G., Hassler, B., Hurtmans, D., Jaffe, D., Jones, N., Kalabokas, P., Kerridge, B., Kulawik, S. S., Latter, B., Leblanc, T., Le Flochmoën, E., Lin, W., Liu, J., Liu, X., Mahieu, E., McClure-Begley, A., Neu, J. L., Osman, M., Palm, M., Petetin, H., Petropavlovskikh, I., Querel, R., Rahpoe, N., Rozanov, A., Schultz, M. G., Schwab, J., Siddans, R., Smale, D., Steinbacher, M., Tanimoto, H., Tarasick, D. W., Thouret, V., Thompson, A. M., Trickl, T., Weatherhead, E., Wespes, C., Worden, H. M., Vigouroux, C., Xu, X., Zeng, G., and Ziemke, J.: Tropospheric Ozone Assessment
- 545 Report: Present-day distribution and trends of tropospheric ozone relevant to climate and global atmospheric chemistry model evaluation, *Elementa*, 6, 39, <https://doi.org/10.1525/elementa.291>, 12.03.01; LK 01, 2018.
- Gromov, S., Jöckel, P., Sander, R., and Brenninkmeijer, C. A. M.: A kinetic chemistry tagging technique and its application to modelling the stable isotopic composition of atmospheric trace gases, *Geoscientific Model Development*, 3, 337–364, <https://doi.org/10.5194/gmd-3-337-2010>, 2010.
- 550 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, <https://doi.org/10.5194/acp-6-3181-2006>, 2006.
- Hagemann, S. and Stacke, T.: Impact of the soil hydrology scheme on simulated soil moisture memory, *Climate Dynamics*, 44, 1731–1750, <https://doi.org/10.1007/s00382-014-2221-6>, 2015.
- 555 Hens, K., Novelli, A., Martinez, M., Auld, J., Axinte, R., Bohn, B., Fischer, H., Keronen, P., Kubistin, D., Nölscher, A. C., Oswald, R., Paasonen, P., Petäjä, T., Regelin, E., Sander, R., Sinha, V., Sipilä, M., Taraborrelli, D., Tatum Ernest, C., Williams, J., Lelieveld, J., and Harder, H.: Observation and modelling of HO_x radicals in a boreal forest, *Atmospheric Chemistry and Physics*, 14, 8723–8747, <https://doi.org/10.5194/ACP-14-8723-2014>, 2014.
- Herrmann, H.: Kinetics of Aqueous Phase Reactions Relevant for Atmospheric Chemistry, *Chemical Reviews*, 103, 4691–4716, <https://doi.org/10.1021/cr020658q>, 2003.
- 560 Hurtmans, D., Coheur, P.-F., Wespes, C., Clarisse, L., Scharf, O., Clerbaux, C., Hadji-Lazaro, J., George, M., and Turquety, S.: FORLI radiative transfer and retrieval code for IASI, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 113, 1391 – 1408, <https://doi.org/10.1016/j.jqsrt.2012.02.036>, three Leaders in Spectroscopy, 2012.
- Jaeglé, L., Jacob, D. J., Brune, W. H., and Wennberg, P. O.: Chemistry of HO_x radicals in the upper troposphere, *Atmospheric Environment*, 35, 469 – 489, [https://doi.org/10.1016/S1352-2310\(00\)00376-9](https://doi.org/10.1016/S1352-2310(00)00376-9), 2001.
- 565 Jiang, X., Guenther, A., Potosnak, M., Geron, C., Seco, R., Karl, T., Kim, S., Gu, L., and Pallardy, S.: Isoprene emission response to drought and the impact on global atmospheric chemistry, *Atmospheric Environment*, 183, 69 – 83, <https://doi.org/10.1016/j.atmosenv.2018.01.026>, 2018.
- Jöckel, P., Tost, H., Pozzer, A., Brühl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M. G., Sander, R., Steil, B., Stiller, G., Tanarhte, M., Taraborrelli, D., van Aardenne, J., and Lelieveld, J.: The atmospheric chemistry general circulation model ECHAM5/MESSy1: consistent simulation of ozone from the surface to the mesosphere, *Atmospheric Chemistry and Physics*, 6, 5067–5104, <https://doi.org/10.5194/acp-6-5067-2006>, 2006.



- Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geoscientific Model Development*, 3, 717–752, <https://doi.org/10.5194/gmd-3-717-2010>, 2010.
- 575 Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C. A. M., Brinkop, S., Cai, D. S., Dyroff, C., Eckstein, J., Frank, F., Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, M., Meul, S., Neumaier, M., Nützel, M., Oberländer-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth System Chemistry integrated Modelling (ESCiMo) with the Modular Earth Submodel System (MESSy) version 2.51, *Geoscientific Model Development*, 9, 1153–1200, <https://doi.org/10.5194/gmd-9-1153-2016>, 2016.
- 580 Jülich Supercomputing Centre: JURECA: Modular supercomputer at Jülich Supercomputing Centre, *Journal of large-scale research facilities*, 4, <https://doi.org/10.17815/jlsrf-4-121-1>, 2018.
- Jülich Supercomputing Centre: JUWELS: Modular Tier-0/1 Supercomputer at the Jülich Supercomputing Centre, *Journal of large-scale research facilities*, 5, <https://doi.org/10.17815/jlsrf-5-171>, 2019.
- 585 Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöckel, P.: Technical Note: An implementation of the dry removal processes DRY DEPosition and SEDimentation in the Modular Earth Submodel System (MESSy), *Atmospheric Chemistry and Physics*, 6, 4617–4632, <https://doi.org/10.5194/acp-6-4617-2006>, 2006.
- 590 Kerkweg, A., Sander, R., Tost, H., Jöckel, P., and Lelieveld, J.: Technical Note: Simulation of detailed aerosol chemistry on the global scale using MECCA-AERO, *Atmospheric Chemistry and Physics*, 7, 2973–2985, <https://doi.org/10.5194/acp-7-2973-2007>, 2007.
- Lelieveld, J. and Crutzen, P. J.: Influences of cloud photochemical processes on tropospheric ozone, *Nature*, 343, 227–233, <https://doi.org/10.1038/343227a0>, 1990.
- 595 Lelieveld, J., Gromov, S., Pozzer, A., and Taraborrelli, D.: Global tropospheric hydroxyl distribution, budget and reactivity, *Atmospheric Chemistry and Physics*, 16, 12477–12493, <https://doi.org/10.5194/acp-16-12477-2016>, 2016.
- Levelt, P. F., van den Oord, G. H. J., Dobber, M. R., Malkki, A., Huib Visser, Johan de Vries, Stammes, P., Lundell, J. O. V., and Saari, H.: The ozone monitoring instrument, *IEEE Transactions on Geoscience and Remote Sensing*, 44, 1093–1101, <https://doi.org/10.1109/TGRS.2006.872333>, 2006.
- 600 Levelt, P. F., Joiner, J., Tamminen, J., Veefkind, J. P., Bhartia, P. K., Stein Zweers, D. C., Duncan, B. N., Streets, D. G., Eskes, H., van der A, R., McLinden, C., Fioletov, V., Carn, S., de Laat, J., DeLand, M., Marchenko, S., McPeters, R., Ziemke, J., Fu, D., Liu, X., Pickering, K., Apituley, A., González Abad, G., Arola, A., Boersma, F., Chan Miller, C., Chance, K., de Graaf, M., Hakkarainen, J., Hassinen, S., Ialongo, I., Kleipool, Q., Krotkov, N., Li, C., Lamsal, L., Newman, P., Nowlan, C., Suleiman, R., Tilstra, L. G., Torres, O., Wang, H., and Wargan, K.: The Ozone Monitoring Instrument: overview of 14 years in space, *Atmospheric Chemistry and Physics*, 18, 5699–5745, <https://doi.org/10.5194/acp-18-5699-2018>, 2018.
- 605 Liang, J. and Jacob, D. J.: Effect of aqueous phase cloud chemistry on tropospheric ozone, *Journal of Geophysical Research: Atmospheres*, 102, 5993–6001, <https://doi.org/10.1029/96JD02957>, 1997.
- Mayer, B. and Madronich, S.: Actinic flux and photolysis in water droplets: Mie calculations and geometrical optics limit, *Atmospheric Chemistry and Physics*, 4, 2241–2250, <https://doi.org/10.5194/acp-4-2241-2004>, 2004.



- 610 Millet, D. B., Jacob, D. J., Custer, T. G., de Gouw, J. A., Goldstein, A. H., Karl, T., Singh, H. B., Sive, B. C., Talbot, R. W., Warneke, C., and Williams, J.: New constraints on terrestrial and oceanic sources of atmospheric methanol, *Atmospheric Chemistry and Physics*, 8, 6887–6905, <https://doi.org/10.5194/acp-8-6887-2008>, 2008.
- Mouchel-Vallon, C., Deguillaume, L., Monod, A., Perroux, H., Rose, C., Ghigo, G., Long, Y., Leriche, M., Aumont, B., Patryl, L., Armand, P., and Chaumerliac, N.: CLEPS 1.0: A new protocol for cloud aqueous phase oxidation of VOC mechanisms, *Geoscientific Model Development*, 10, 1339–1362, <https://doi.org/10.5194/gmd-10-1339-2017>, 2017.
- 615 Nölscher, A., Butler, T., Auld, J., Veres, P., Muñoz, A., Taraborrelli, D., Vereecken, L., Lelieveld, J., and Williams, J.: Using total OH reactivity to assess isoprene photooxidation via measurement and model, *Atmos. Environ.*, 89, 453–463, <https://doi.org/10.1016/j.atmosenv.2014.02.024>, 2014.
- Olsen, M. A., Douglass, A. R., and Kaplan, T. B.: Variability of extratropical ozone stratosphere–troposphere exchange using microwave limb sounder observations, *Journal of Geophysical Research: Atmospheres*, 118, 1090–1099, <https://doi.org/10.1029/2012JD018465>, 2013.
- 620 Parker, R. J., Boesch, H., Wooster, M. J., Moore, D. P., Webb, A. J., Gaveau, D., and Murdiyarto, D.: Atmospheric CH₄ and CO₂ enhancements and biomass burning emission ratios derived from satellite observations of the 2015 Indonesian fire plumes, *Atmospheric Chemistry and Physics*, 16, 10 111–10 131, <https://doi.org/10.5194/acp-16-10111-2016>, 2016.
- Rodgers, C. D.: *Inverse Methods for Atmospheric Sounding*, WORLD SCIENTIFIC, <https://doi.org/10.1142/3171>, 2000.
- 625 Roeckner, E., Bäuml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornblueh, L., Manzini, E., Rhodin, A., Schlese, U., Schulzweida, U., and Tompkins, A.: The atmospheric general circulation model ECHAM 5. Part I: Model description, Tech. Rep. 349, Max-Planck-Institute for Meteorology, Hamburg, 2003.
- Rosanka, S., Sander, R., Wahner, A., and Taraborrelli, D.: Oxidation of low-molecular weight organic compounds in cloud droplets: development of the JAMOC chemical mechanism in CAABA/MECCA (version 4.5.0gmd), *Geoscientific Model Development Discussions*, companion paper, submitted, 2020a.
- 630 Rosanka, S., Vu, G. H. T., Nguyen, H. M. T., Pham, T. V., Javed, U., Taraborrelli, D., and Vereecken, L.: Atmospheric chemical loss processes of isocyanic acid (HNCO): a combined theoretical kinetic and global modelling study, *Atmospheric Chemistry and Physics*, 20, 6671–6686, <https://doi.org/10.5194/acp-20-6671-2020>, 2020b.
- Sander, R.: *Modeling Atmospheric Chemistry: Interactions between Gas-Phase Species and Liquid Cloud/Aerosol Particles*, *Surveys in Geophysics*, 20, 1–31, <https://doi.org/10.1023/A:1006501706704>, 1999.
- 635 Sander, R., Baumgaertner, A., Cabrera-Perez, D., Frank, F., Gromov, S., Grooß, J.-U., Harder, H., Huijnen, V., Jöckel, P., Karydis, V. A., Niemeyer, K. E., Pozzer, A., Riede, H., Schultz, M. G., Taraborrelli, D., and Tauer, S.: The community atmospheric chemistry box model CAABA/MECCA-4.0, *Geoscientific Model Development*, 12, 1365–1385, <https://doi.org/10.5194/gmd-12-1365-2019>, 2019.
- Sandu, A. and Sander, R.: Technical note: Simulating chemical systems in Fortran90 and Matlab with the Kinetic PreProcessor KPP-2.1, *Atmospheric Chemistry and Physics*, 6, 187–195, <https://doi.org/10.5194/acp-6-187-2006>, 2006.
- 640 Scanza, R. A., Hamilton, D. S., Perez Garcia-Pando, C., Buck, C., Baker, A., and Mahowald, N. M.: Atmospheric processing of iron in mineral and combustion aerosols: development of an intermediate-complexity mechanism suitable for Earth system models, *Atmospheric Chemistry and Physics*, 18, 14 175–14 196, <https://doi.org/10.5194/acp-18-14175-2018>, 2018.
- Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavroukou, T., Müller, J.-F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmospheric Chemistry and Physics*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014.



- Stahelin, J., Buehler, R. E., and Hoigné, J.: Ozone decomposition in water studied by pulse radiolysis. 2. Hydroxyl and hydrogen tetroxide (HO₄) as chain intermediates, *The Journal of Physical Chemistry*, 88, 5999–6004, <https://doi.org/10.1021/j150668a051>, 1984.
- 650 Taraborrelli, D., Lawrence, M. G., Butler, T. M., Sander, R., and Lelieveld, J.: Mainz Isoprene Mechanism 2 (MIM2): an isoprene oxidation mechanism for regional and global atmospheric modelling, *Atmospheric Chemistry and Physics*, 9, 2751–2777, <https://doi.org/10.5194/ACP-9-2751-2009>, 2009.
- Taraborrelli, D., Lawrence, M. G., Crowley, J. N., Dillon, T. J., Gromov, S., Groß, C. B. M., Vereecken, L., and Lelieveld, J.: Hydroxyl radical buffered by isoprene oxidation over tropical forests, *Nature Geoscience*, 5, 190–193, <https://doi.org/10.1038/ngeo1405>, 2012.
- 655 Taraborrelli, D., Cabrera-Perez, D., Bacer, S., Gromov, S., Lelieveld, J., Sander, R., and Pozzer, A.: Influence of aromatics on tropospheric gas-phase composition, *Atmospheric Chemistry and Physics Discussions*, 2020, 1–26, <https://doi.org/10.5194/acp-2020-461>, 2020.
- Tilgner, A., Brüner, P., Wolke, R., and Herrmann, H.: Modelling multiphase chemistry in deliquescent aerosols and clouds using CAPRAM3.0i, *Journal of Atmospheric Chemistry*, 70, 221–256, <https://doi.org/10.1007/s10874-013-9267-4>, 2013.
- Tost, H., Jöckel, P., Kerkweg, A., Sander, R., and Lelieveld, J.: Technical note: A new comprehensive SCAVenging submodel for global atmospheric chemistry modelling, *Atmospheric Chemistry and Physics*, 6, 565–574, <https://doi.org/10.5194/acp-6-565-2006>, 2006.
- 660 Tost, H., Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., and Lelieveld, J.: Global cloud and precipitation chemistry and wet deposition: tropospheric model simulations with ECHAM5/MESSy1, *Atmospheric Chemistry and Physics*, 7, 2733–2757, <https://doi.org/10.5194/acp-7-2733-2007>, 2007.
- Wang, R., Balkanski, Y., Boucher, O., Bopp, L., Chappell, A., Ciais, P., Hauglustaine, D., Peñuelas, J., and Tao, S.: Sources, transport and deposition of iron in the global atmosphere, *Atmospheric Chemistry and Physics*, 15, 6247–6270, [https://doi.org/10.5194/acp-15-6247-](https://doi.org/10.5194/acp-15-6247-2015)
665 2015, 2015.
- Wespes, C., Hurtmans, D., Clerbaux, C., and Coheur, P.-F.: O₃ variability in the troposphere as observed by IASI over 2008–2016: Contribution of atmospheric chemistry and dynamics, *Journal of Geophysical Research: Atmospheres*, 122, 2429–2451, <https://doi.org/10.1002/2016JD025875>, 2017.
- WMO: Meteorology-A three-dimensional science: Second session of the commission for aerology, *WMO Bull.*, p. 134–138, 1957.
- 670 Young, P. J., Naik, V., Fiore, A. M., Gaudel, A., Guo, J., Lin, M., Neu, J., Parrish, D., Rieder, H., Schnell, J., et al.: Tropospheric Ozone Assessment Report: Assessment of global-scale model performance for global and regional ozone distributions, variability, and trends, *Elem Sci Anth*, 6, <https://doi.org/10.1525/elementa.265>, 2018.
- Ziemke, J. R., Chandra, S., Labow, G. J., Bhartia, P. K., Froidevaux, L., and Witte, J. C.: A global climatology of tropospheric and stratospheric ozone derived from Aura OMI and MLS measurements, *Atmospheric Chemistry and Physics*, 11, 9237–9251,
675 <https://doi.org/10.5194/acp-11-9237-2011>, 2011.