

**Convergent ~~Estimates of Evidence for Pervasive but~~  
~~Limited Contribution of Biomass Burning to Biomass~~  
~~Burning-Derived~~ Atmospheric Ammonia in Peninsular  
Southeast Asia**

5 Yunhua Chang<sup>1</sup>, Yan-Lin Zhang<sup>1</sup>, Sawaeng Kawichai<sup>2</sup>, Qian Wang<sup>1</sup>, Martin  
Van Damme<sup>3</sup>, Lieven Clarisse<sup>3</sup>, Tippawan Prapamontol<sup>2</sup>, and Moritz F.  
Lehmann<sup>4</sup>.

<sup>1</sup>Yale-NUIST Center on Atmospheric Environment, Nanjing University of Information Science & Technology,  
Nanjing 210044, China

10 <sup>2</sup>Research Institute for Health Sciences (RIHES), Chiang Mai University, Chiang Mai 50200, Thailand

<sup>3</sup>Université libre de Bruxelles (ULB), Spectroscopy, Quantum Chemistry and Atmospheric Remote Sensing  
(SQUARES), Brussels B-1050, Belgium

<sup>4</sup>Department of Environmental Sciences, University of Basel, Basel 4056, Switzerland

*Correspondence to:* Yan-Lin Zhang (dryanlinzhang@outlook.com)

15 **Abstract.** Ammonia (NH<sub>3</sub>) is an important agent involved in atmospheric chemistry and nitrogen cycling. Current  
estimates of NH<sub>3</sub> emissions from biomass burning (BB) differ by more than a factor of two, impeding a reliable

assessment of their environmental consequences. Combining high-resolution satellite observations of  $\text{NH}_3$  columns with network measurements of the concentration and stable nitrogen isotope composition ( $\delta^{15}\text{N}$ ) of  $\text{NH}_3$ , we present coherent estimates on the amount of  $\text{NH}_3$  derived from BB in the heartland of Southeast Asia, a tropical monsoon environment. Our results reveal a strong variability of atmospheric  $\text{NH}_3$  levels in time and space across different landscapes. All evidence in hand suggests that anthropogenic activities are the most important modulating control with regards to the observed patterns of  $\text{NH}_3$  distribution in the study area. N-isotope balance considerations revealed that during the intensive fire period, the atmospheric input from BB accounts for not more than  $21 \pm 5\%$  ( $1\sigma$ ) of the ambient  $\text{NH}_3$ , even at the rural sites and in the proximity of burning areas. Our N-isotope based assessment of the variation of the relative contribution of BB-derived  $\text{NH}_3$  is further validated independently through the measurements of particulate  $\text{K}^+$ , a chemical tracer of BB. Our findings underscore that BB-induced  $\text{NH}_3$  emissions in the tropical monsoon environments can be much lower than previously anticipated, with important implications for future modeling studies to better constrain the climate and air quality effects of wildfires.

## 1 Introduction

Biomass burning (BB) in tropical vegetation regions due to wildfires has been recognized as a globally important source of trace gases (including  $\text{CO}_2$ , CO, and ozone precursors) and aerosols (mostly black and organic carbon) (Crutzen and Andreae, 1990; Andreae and Merlet, 2001; Shi et al., 2015; Andreae, 2019; Crutzen et al., 1979). Most BB hotspots occur in West Africa and South America (Crutzen and Andreae, 1990; van der Werf et al., 2006; Shi et al., 2015), but recent studies have highlighted the importance also of Southeast (SE) Asia in this regard, mainly because of the much higher population densities near intensive fire burning areas (Huang et al., 2013; Marlier et al., 2013; Lee et al., 2017; Betha et al., 2014). The climate over large parts of SE Asia is governed by a wet (typically May, June, July) and dry (typically February, March, April) season caused by seasonal shifts in the monsoon winds. During the dry season, dry plant materials (e.g., forest, peatland, banana leaf) readily ignite,

resulting in large wildfires that can markedly modify the atmospheric composition in the tropics, while the tropical  
40 rain belt causes plentiful rainfall during summer, preventing such fires during the rainy season (Lee et al., 2017;  
Chu et al., 2018).

Besides carbon soot, BB also emits large amounts of reactive nitrogen compounds (Lobert et al., 1990; Bauters et  
al., 2018), in particular ammonia ( $\text{NH}_3$ ), which is believed to represent the major source of  $\text{NH}_3$  during intensive  
fires period (Akagi et al., 2011; Whitburn et al., 2015). However, these emissions are subject to large uncertainties  
45 (differences by a factor of two or greater). (Bray et al., 2018; Whitburn et al., 2015; Whitburn et al., 2016b; Van  
Damme et al., 2015b) For example, BB is probably the second most important  $\text{NH}_3$  source after agriculture,  
contributing 11-23% of the global burden (Paulot et al., 2017; Bouwman et al., 1997). Minor  $\text{NH}_3$  sources include  
fossil fuel burning and biogenic activity (Chang et al., 2012; Chang et al., 2016b; Chang et al., 2019b; Chang et al.,  
2020). A recent paper also highlighted the underestimated importance of industrial emissions (Van Damme et al.,  
50 2018). Once emitted in the atmosphere,  $\text{NH}_3$  is rapidly removed by dry or wet deposition (Asman et al., 1998).  
Excess  $\text{NH}_3$  is known to be responsible for several environmental issues: eutrophication of terrestrial and aquatic  
ecosystem, soil acidification, and loss of plant diversity (Sutton et al., 2008; Aneja et al., 2008; Sutton et al., 2011).  
In the atmosphere,  $\text{NH}_3$  can neutralize acid gases (mostly sulfuric acid, nitric acid or hydrochloric acid), resulting  
in the formation of secondary aerosols that in turn negatively affect climate and human health (Wang et al., 2011;  
55 Wang et al., 2013; Paulot and Jacob, 2014; Souri et al., 2017).

To assess the environmental impacts of BB (e.g., air quality and climate change), atmospheric chemistry models  
incorporating BB-related emissions have widely been used over the past decades (Huang et al., 2013; Aouizerats  
et al., 2015; Wang et al., 2013; Wang et al., 2011; Souri et al., 2017) but are afflicted with a relatively large  
uncertainty regarding the input parameters used in the models (Hantson et al., 2016; Whitburn et al., 2015; Paulot  
60 et al., 2017). The uncertainties, for example, for carbon emissions and for other trace gases (including  $\text{NH}_3$ ), can

be over 200% (Whitburn et al., 2015; Paulot et al., 2017; Zhu et al., 2013; Pan et al., 2019). In recent years, hyperspectral sounders on board satellites have demonstrated their capabilities to directly measure tropospheric column concentrations of NH<sub>3</sub> gas (Van Damme et al., 2018; Van Damme et al., 2014; Van Damme et al., 2015b; Clarisse et al., 2009). Satellite observations therefore offer a “top-down” alternative to the bottom-up estimates. 65 Still, the biggest challenge of satellite-based NH<sub>3</sub> assessments is the requirement for the atmosphere to be cloud-free during observations, and the need for a sizeable temperature difference between land or sea surface and the atmosphere (Van Damme et al., 2015a; Whitburn et al., 2015; Martin, 2008; Streets et al., 2013; Clarisse et al., 2010).


Large uncertainties remain regarding global or regional atmospheric budgets of NH<sub>3</sub>, and the attribution of 70 emissions to specific sources, emphasizing the need for independent verification methods. An impressive body of previous work has studied the BB influence on the concentration and composition of aerosols in SE Asia (Betha et al., 2014; Aouizerats et al., 2015; Lee et al., 2017; Bikina et al., 2019). However, to our knowledge, there are no reports on the detailed spatiotemporal patterns of atmospheric NH<sub>3</sub> concentration and nitrogen isotopic composition ( $\delta^{15}\text{N-NH}_3$ ) associated with BB in this region. Due to isotopic fractionation associated with NH<sub>3</sub> production, 75 pyrogenic NH<sub>3</sub> displays a distinctly higher  $\delta^{15}\text{N-NH}_3$  ( $\delta^{15}\text{N}$  defined as  $(R_{\text{sample}}/R_{\text{standard}} - 1) \times 1000$ , where R refers to the <sup>15</sup>N/<sup>14</sup>N ratio in a sample or a standard) than temperature-dependent volatilized sources (Felix et al., 2013; Chang et al., 2016a). The N isotopic analysis of ambient NH<sub>3</sub> has been proven a useful tool to constrain sources of NH<sub>3</sub> emissions in the atmosphere, where both natural and anthropogenic activities are relevant (Chang et al., 2019b; Chang et al., 2019a; Elliott et al., 2019). Here, we integrate high-resolution satellite observations with discrete NH<sub>3</sub> 80 concentration measurements and  $\delta^{15}\text{N-NH}_3$  data obtained from a regional passive monitoring network during and after the dry season of large-scale forest fires in the mountain areas of northern Thailand, SE Asia.

## 2 Methods

### 2.1 Site description

Surrounded by the mountain ranges of the northern Thailand highlands, the Chiang Mai Province covers an area of approximately 20107 km<sup>2</sup>, with a total population of over 1.7 million. Chiang Mai is characterized by a tropical monsoon climate, tempered by the low latitude and moderate elevation, with warm to hot weather year-round. Some 70% of the area is covered by forests, and 13.4% of the area is under agriculture. A continuing environmental issue in Chiang Mai is the smoke pollution from wildfires that primarily occurs every year towards the end of the dry season between February and April (Tsai et al., 2013) before the relatively cool and rainy season from May on.

During the period from March to July 2018, ambient NH<sub>3</sub> concentrations and  $\delta^{15}\text{N-NH}_3$  values were determined at 9 monitoring stations across the Chiang Mai Province. Figure 1 illustrates the location of sampling sites (with the different land use regimes indicated), while Fig. S1 reports meteorological data for Chiang Mai and Table S1 details the information of each station.



95 Figure 1. Location of sampling sites. Land-cover map (left; revised from Chang et al., [2016b](#)) with zoom-in in  
lower panels (right). The chosen sampling sites are representative for gradient in land use from urban to rural.  
Images in the right are obtained from © Google Maps.

## 2.2 Sampling and laboratory analysis

In order to obtain information regarding the spatial and temporal variability of  $\text{NH}_3$  concentrations over Chiang  
100 Mai, ambient gas-phase  $\text{NH}_3$  concentrations at each site were collected weekly using passive sampling devices  
(PSD) (ALPHA - Adapted Low-cost, Passive High Absorption; Centre for Ecology and Hydrology, Edinburg, UK)  
(Chang et al., 2016a). The ALPHA PSD is a circular polyethylene vial (26 mm height, 27 mm diameter) with one  
open end. The vial holds a 25 mm phosphorous acid-impregnated filter and a PTFE membrane for gaseous  $\text{NH}_3$   
diffusion. These PSDs have been widely used in Europe, China, and the US, and are capable of detecting  $\text{NH}_3$   
105 concentrations as low as  $0.03 \mu\text{g m}^{-3}$  (Chang et al., 2016a; Puchalski et al., 2011; Liu et al., 2013; Tang et al., 2018).  
In the laboratory, the ALPHA filter samples were soaked in 10 mL deionized water ( $18 \text{ M}\Omega \cdot \text{cm}$ ) in a 15 mL vial  
for 30 min with occasional shaking. Concentrations of  $\text{NH}_3$ -derived  $\text{NH}_4^+$  in extracts were determined using a  
Dionex<sup>TM</sup> ICS-5000<sup>+</sup> system (Thermo Fisher Scientific, Sunnyvale, USA). The IC system was equipped with an  
automated sampler (AS-DV), an IonPac CG12A guard column, and a CS12A separation column. Aqueous  
110 methanesulfonic acid (MSA,  $30 \text{ mM L}^{-1}$ ) served as eluent at a flow rate of  $1 \text{ mL min}^{-1}$ . The isotopic analysis of the  
extracted  $\text{NH}_4^+$  was based on the isotopic analysis of nitrous oxides ( $\text{N}_2\text{O}$ ) after chemical conversion (Liu et al.,  
2014). More precisely, dissolved  $\text{NH}_4^+$  in DI extracts was oxidized to  $\text{NO}_2^-$  by alkaline hypobromite ( $\text{BrO}^-$ ), and  
then reduced to  $\text{N}_2\text{O}$  by hydroxylamine hydrochloride ( $\text{NH}_2\text{OH} \cdot \text{HCl}$ ). The produced  $\text{N}_2\text{O}$  was analyzed using a  
purge and cryogenic trap system (Gilson GX-271, IsoPrime Ltd., Cheadle Hulme, UK), coupled to an isotope ratio  
115 mass spectrometer (PT-IRMS) (IsoPrime 100, IsoPrime Ltd., Cheadle Hulme, UK) (Liu et al., 2014). In order to  
correct for any machine drift and procedural blank contribution, international  $\text{NH}_4^+$  (IAEA N1, USGS 25, and

USGS 26) standards were processed in the same way as samples (Liu et al., 2014). The analytical precision for N isotope analyses was better than 0.5‰ ( $n = 5$ ).

设置了格式: 字体: 倾斜

### 2.3 Isotope-based source apportionment

120 Isotopic mixing models represent valuable tools to estimate the fractional contributions of multiple sources (emission sources of  $\text{NH}_3$  in this study) within a mixture (the ambient  $\text{NH}_3$  in this study) (Layman et al., 2012). By explicitly taking into account the uncertainties associated with the isotopic signatures of single sources and the N isotope fractionation during transformations, the application of Bayesian methods to stable isotope mixing models yields robust probability estimates on source apportionments, and its application to natural systems is more  
125 appropriate than the application of simple linear mixing models (Parnell et al., 2010). Here, a novel Bayesian approach using a mixing model, implemented in the software package SIAR (Stable Isotope Analysis in R), was used to resolve multiple  $\text{NH}_3$  source categories by generating potential solutions of source apportionment as true probability distributions of the single source contribution to the total  $\text{NH}_3$  pool. The generation of such source contribution probability distributions allows estimating likelihood ranges of source contributions even at under-  
130 constrained conditions (i.e., the number of potential sources exceeds the number of different isotope system parameters + 1). The SIAR package is available to download from the packages section of the Comprehensive R Archive Network site (CRAN; <http://cran.r-project.org/>), which has been widely applied in a number of fields (Chang et al., 2019a; Chang et al., 2019b). Model frame and computing methods are detailed in Text S1.

### 2.4 Satellite observations of ammonia and fires

135  $\text{NH}_3$  total columns ( $\text{molec cm}^{-2}$ ) are retrieved from the Infrared Atmospheric Sounding Interferometer (IASI) observations. IASI instruments are onboard the Metop satellite series; in this work, we use IASI/Metop-A (launched in 2006) and IASI/Metop-B (launched in 2012) data. Both instruments have an overpass time around 9:30 am and

pm (local solar time when crossing the equator) and therefore provide in total a global coverage four times a day. The retrieval strategy, based on artificial neural networks, is fully detailed in previous work (Whitburn et al., 2016a; Van Damme et al., 2017). Here we only consider morning observations, as they are more sensitive to the lower layer of the atmosphere. Fire Radiative Power (FRP) from the Moderate Resolution Imaging Spectroradiometer (MODIS) and fire counts derived from the 375-m Visible Infrared Imaging Radiometer Suite (VIIRS) are also used (Li et al., 2020).

### 3 Results and discussion

#### 3.1 Satellite-observed NH<sub>3</sub> distributions ~~probably linked with biomass burning~~

Figure 2a illustrates the monthly spatial distribution of NH<sub>3</sub> columns obtained from IASI in 2018 at a spatial resolution of  $0.25^\circ \times 0.5^\circ$  cells. Our study area is set within a large domain of  $5.00^\circ \times 3.25^\circ$  (red and black rectangles in Fig. 2a and Fig. 2b, respectively), in which totally 260 gridded pixels ( $0.25^\circ \times 0.25^\circ$  per cell) are used for dividing active fire points (Fig. 2b). Intriguingly, from this plot, one is tempted to conclude that fires do play a very important role in NH<sub>3</sub> emissions, as the NH<sub>3</sub> columns are much higher in March and April (dry season), which is coincident with high number of monthly fire activities (indicated by the number of fire points). Further, using 11 years (2008-2018) of IASI satellite data, Figure S2 presents a climatology of monthly NH<sub>3</sub> columns over Chiang Mai at a much finer spatial resolution, which also support the pervasive contribution of BB during dry season (March and April). Based on the average observed temporal distribution of satellite-constrained wildfires, the sampling period in this study can be divided into two contrasting fire-regime periods, i.e., BB season (March and April) and non-BB season (May and June). Interestingly, however, although the number of fire points in March (43613 points) is significantly ( $p < 0.01$ ) higher than that in April (27905 points) (Fig. 2b), the average NH<sub>3</sub> column in March is nearly the same as that in April (Fig. 2a). This implies that there is not a one-to-one relationship between BB and NH<sub>3</sub> emissions,

and in turn that other sources or factors (e.g., soil dryness, agricultural emissions, precipitation, temperature  
 160 dependence, etc.) must also play a significant role.




Figure 2. (a) Monthly (March, April, May, June) spatial distributions of the NH<sub>3</sub> total columns (molecules cm<sup>-2</sup>) in  
 2018 obtained from the satellite measurements by the Infrared Atmospheric Sounding Interferometer  
 (IASI)/MetOp-A instrument in 0.25° × 0.5° cells. (b) Monthly distributions of gridded counts of active fire pixels  
 165 (0.25° × 0.25° per cell) derived from the 375-m Visible Infrared Imaging Radiometer Suite (VIIRS). The red and  
 black squares indicate our study area in Chiang Mai. (c) Daily variations of (°C) and rainfall (mm) in Chiang Mai  
 City.

Given that the average monthly temperature varies only little, in contrast to the drastic change of rainfall during  
 our study period (Fig. 2c), it is reasonable to assume that temperature-dependent NH<sub>3</sub> volatilization is not the main  
 170 driver of changes in the NH<sub>3</sub> columns. The amount of rainfall, on the other hand, can have multi-faceted impact on  
 NH<sub>3</sub> emissions. First of all, there is an obvious link between precipitation rates and the number of wildfires, and, if

BB is a major  $\text{NH}_3$  emission source, we can also expect a relationship between the  $\text{NH}_3$  columns and monthly rainfall rates. Secondly, and maybe more importantly, rain will dissolve atmospheric particulate  $\text{NH}_4^+$ , and will act to clean the air from  $\text{NH}_3$ , which may partly explain the low  $\text{NH}_3$  levels during May and June. On the other hand, comparison between March and April reveals higher  $\text{NH}_3$  levels in April despite higher rain rates, suggesting that other processes than BB and rain-scavenging of BB-derived  $\text{NH}_3$  must be relevant factors. In Fig. 3a and 3b, we superimposed the orography at the scale of the study area (Chiang Mai and surrounding mountains) onto the images of year-long averaged MODIS FRP (Fire Radiative Power) and IASI- $\text{NH}_3$  for 2018, respectively. Just by visual evaluation it seems obvious that there is no strong correlation between fire intensity/number of fires and the observed IASI- $\text{NH}_3$ , suggesting only limited influence of BB on  $\text{NH}_3$ . Yet more strikingly, the IASI- $\text{NH}_3$  distribution matches that of the population density quite well (Fig. 3c). More precisely, hot spots of atmospheric  $\text{NH}_3$  (Fig. 3b) appear to be concentrated in urban areas with dense population. Hence, our satellite remote sensing observations suggest a significant influence of non-BB emissions on  $\text{NH}_3$  concentrations, seemingly related to urban anthropogenic activities.





Figure 3. (a) The MODIS FRP (fire radiative power) (the size of red dots is proportional to [arbitrary FRP values](#)) and (b) the IASI Metop-A and Metop-B averaged NH<sub>3</sub> distribution (molec cm<sup>-2</sup>) for 2018. (c) Number of people per grid-cell in the Chiang Mai area in 2018 at a resolution of 3 arc minute.

### 190 3.2 Discrete concentration measurements confirm urban areas as hot spots of NH<sub>3</sub> emissions

A total of ~~153~~ [more than 150](#) samples were collected in this study for analyzing NH<sub>3</sub> concentrations (~~and reported in Fig. 4~~). ~~Considering all weekly samples, the~~ [The range of](#) atmospheric NH<sub>3</sub> concentrations over Chiang Mai ~~ranged~~ [was](#) from 2.5 to 46.4 μg m<sup>-3</sup>, with ~~a~~ mean (± 1σ) and median values of 14.5 (± 9.2) and 11.4 μg m<sup>-3</sup>, respectively. Consistent with the IASI satellite-based NH<sub>3</sub> assessment, the weighted average NH<sub>3</sub> concentration

195 (mean<sub>min</sub><sup>max</sup> ± 1σ) ~~during the dry season (i.e., when wildfires were markedly more abundant)~~ [was](#) significantly (p < 0.01) higher (~~20.6<sub>6.8</sub><sup>46.4</sup> ± 9.8 μg m<sup>-3</sup>~~ [during the dry season when there were significantly \(p < 0.01\) more wild fires \(20.6<sub>6.8</sub><sup>46.4</sup> ± 9.8 μg m<sup>-3</sup>\)](#)) than ~~that~~ during the rainy season (10.2<sub>2.5</sub><sup>31.9</sup> ± 5.7 μg m<sup>-3</sup>). Again, it is tempting to conclude that there is a direct link between higher atmospheric NH<sub>3</sub> levels and the higher number of BB events.

However, there are several aspects that appear to ~~preclude~~ [speak against](#) BB ~~at least~~ as the main, or only, driver of ambient NH<sub>3</sub> concentrations. Firstly, from a global perspective, the ambient NH<sub>3</sub> concentrations we measured in northern Thailand are generally lower than in tropical regions with dense population or intensive agricultural production (also see Fig. 2a) (Carmichael et al., 2003; Chang et al., 2016b). Secondly, within the study area, large spatial differences in NH<sub>3</sub> concentrations were found (Fig. 4). Yet, despite their proximity to wildfires at the time, the three rural sites always displayed the lowest NH<sub>3</sub> concentrations (8.3<sub>2.5</sub><sup>26.8</sup> ± 4.6 μg m<sup>-3</sup>; Fig. 4; see detailed discussion in the next section).

205

城代码已更改




Figure 4. Temporal variations of measured  $\text{NH}_3$  concentrations ( $\mu\text{g m}^{-3}$ ) between sites of different land-use regimes. The error bar indicates one standard deviation.

Although afflicted with ~~Despite relatively~~ large uncertainties, it is well accepted that, globally, atmospheric  $\text{NH}_3$  is primarily emitted by agricultural activities and biomass burning (Asman et al., 1998; Bouwman et al., 1997). As a consequence, one would expect the  $\text{NH}_3$  concentrations in the atmosphere over rural environments with lush vegetation and agricultural land-use to be higher than those in (sub)urban areas, where agricultural activities are mostly absent. In our study, the average  $\text{NH}_3$  concentrations at the nine sites, respectively, are:  $19.5^{39.4}_{6.5} \pm 9.5$  (S1; suburban),  $11.9^{19.7}_{4.5} \pm 4.6$  (S2; suburban),  $8.8^{16.6}_{4.4} \pm 3.9$  (S3; rural),  $9.0^{26.8}_{2.8} \pm 5.8$  (S4; rural),  $7.0^{13.7}_{2.5} \pm 3.8$  (S5; rural),  $20.2^{40.5}_{9.1} \pm 8.6$  (S6; urban traffic),  $18.1^{46.1}_{7.2} \pm 12.1$  (S7; urban traffic),  $19.6^{46.4}_{4.2} \pm 10.1$  (S8; urban), and  $16.6^{30.6}_{6.7} \pm 8.0$  (S9; urban)  $\mu\text{g m}^{-3}$  (see also compilation in Fig. 4). Thus, against current paradigms, the observed  $\text{NH}_3$  concentrations, from high to low, clearly display clearly reflect an urban ( $18.6^{46.4}_{4.2} \pm 9.7$ ,  $n = 68$ ) to suburban ( $15.6^{39.4}_{4.5} \pm 8.3$ ,  $n = 34$ ) to rural ( $8.3^{26.8}_{2.5} \pm 4.6$ ,  $n = 51$ ) gradient. Such a concentration gradient can be taken as evidence that non-agricultural activities (including on-road traffic), at least in some regions, can overtake-outweigh agriculture and/or BB as the dominant  $\text{NH}_3$  source in urban areas.

Indeed, a growing body of studies confirm that the urban atmosphere can be a hot spot of NH<sub>3</sub> ~~releaser~~release. Non-agricultural activities, ~~such as like~~wastewater treatment, coal combustion, solid garbage ~~handling, as well as~~ vehicular exhaust, and urban green space ~~also~~ contribute ~~strongly to urban~~ to NH<sub>3</sub> emissions (Chang et al., 2016a; Chang et al., 2019b; Teng et al., 2017; Chang et al., 2015; Li et al., 2016; Sun et al., 2017). For example, high  
225 vehicular NH<sub>3</sub> emissions from ~~noble metal based~~ three-way catalysts (TWCs) have been demonstrated in chassis dynamometer vehicle experiments, road tunnel tests, and through ambient air measurements (Huang et al., 2018; Chang et al., 2016b; Chang et al., 2019b).

### 3.3 N isotopic constraints on the sources of natural and anthropogenic NH<sub>3</sub>

The correlative analysis of spatiotemporal concentration patterns with variations in land use effects provides first  
230 qualitative constraints with regards to the relative importance of natural/BB and anthropogenic NH<sub>3</sub> emissions, but it is insufficient when a more quantitative assessment is required. The N-isotopic composition of NH<sub>3</sub> (i.e., δ<sup>15</sup>N-NH<sub>3</sub>-) can provide help in this regard, as it is sensitive to changes of NH<sub>3</sub> sources with distinct isotopic composition (Elliott et al., 2019; Felix et al., 2013). δ<sup>15</sup>N-NH<sub>3</sub> values determined in this study ( $-27.04_{-46.28}^{-12.35} \pm 7.22\text{‰}$ , n = 145) show a relatively large variability in time and space (Fig. 5). NH<sub>3</sub> emitted from the five major NH<sub>3</sub> sources displays  
235 distinct isotopic signatures (N-fertilizer application,  $-50.0 \pm 1.8\text{‰}$ ; urban waste volatilized sources,  $-37.8 \pm 3.6\text{‰}$ ; livestock breeding,  $-29.1 \pm 1.7\text{‰}$ ; on-road traffic,  $-12.0 \pm 1.8\text{‰}$ ; biomass burning, 12‰) (see ~~colored~~coloured bars in Fig. 4) (Chang et al., 2016a; Kawashima and Kurahashi, 2011; Chang and Ma, 2016). Thus, the measurement of δ<sup>15</sup>N-NH<sub>3</sub> can be used to distinguish between specific sources, and to quantify their contribution to the measured total NH<sub>3</sub> pool (~~n~~Note that the isotopic signature determined by Kawashima and Kurahashi (2011) is for particulate  
240 NH<sub>4</sub><sup>+</sup>, not NH<sub>3</sub>). As a first step, we examine the spatiotemporal characteristics of the measured δ<sup>15</sup>N-NH<sub>3</sub> relative to the N isotopic source signatures to infer seasonal changes of NH<sub>3</sub> sources.

设置了格式: 下标

设置了格式: 上标

设置了格式: 下标




Figure 5. (left) Weekly-based variations of the  $\delta^{15}\text{N}$  values (%) of ambient  $\text{NH}_3$  measured in urban, suburban, and rural environments (setting 0 as the breaking point). The error bars indicate two standard deviations. (right) Box

245 plots of the distribution of  $\delta^{15}\text{N-NH}_3$  during BB season and non-BB season for each type of sampling sites.

The lowest  $\delta^{15}\text{N-NH}_3$  values were observed at the rural sites (S3-S5) during the dry season ( $-32.72_{-46.28}^{-20.38} \pm 6.46\%$ ,  $n = 21$ ). These  $\delta^{15}\text{N}$  values are much lower than the  $\delta^{15}\text{N}$  of BB-related  $\text{NH}_3$  and indicate the pervasive influence of agricultural  $\text{NH}_3$  emissions in rural environments, rather than BB. During the rainy season, a drastic increase of  $\delta^{15}\text{N-NH}_3$  ( $-23.97_{-37.99}^{-12.35} \pm 7.50\%$ ,  $n = 22$ ) at the rural sites was observed. Again, if

250 BB was the dominating modulator of  $\text{NH}_3$  levels, an increased contribution from of BB-derived  $\text{NH}_3$  during the dry versus the wet season in rural areas should have resulted in higher, and not lower  $\delta^{15}\text{N-NH}_3$  values. The increased  $\delta^{15}\text{N-NH}_3$  during the non-BB (i.e., rainy) period can probably be explained by the fact that agricultural  $\text{NH}_3$  emissions with low  $\delta^{15}\text{N-NH}_3$  can be dramatically lowered by continuous and heavy rainfall (Zheng et al., 2018; Chang et al., 2019a) so that at low levels, local sources can become more important (e.g., residential kitchens,

255 nearby burning of biofuels for cooking).

As for the urban sites, the mean  $\delta^{15}\text{N-NH}_3$  values at S6, S7, S8, and S9 were  $-23.95_{-34.00}^{-16.35} \pm 4.61\%$  ,  
 $-25.53_{-35.01}^{-14.10} \pm 6.11\%$  ,  $-24.47_{-33.08}^{-16.86} \pm 4.20\%$  , and  $-25.32_{-39.44}^{-17.13} \pm 7.75\%$  , respectively. The overall  
average  $\delta^{15}\text{N-NH}_3$  value at the four urban sites ( $-24.82_{-39.44}^{-14.10} \pm 5.74\%$  , n = 68) was significantly (p < 0.01; one-  
way analysis of variance and paired-sample t test; similarly hereinafter) higher than that at the rural


260 ( $-28.24_{-46.28}^{-12.35} \pm 8.22\%$  , n = 43) and suburban ( $-29.94_{-45.62}^{-18.78} \pm 7.35\%$  , n = 34) sites, respectively, indicating a  
greater contribution of  $\text{NH}_3$  emissions from pyrogenic (e.g., on road traffic) sources. The average value of urban  
 $\delta^{15}\text{N-NH}_3$  during the dry season ( $-24.21_{-34.47}^{-14.10} \pm 4.82\%$  , n = 28) was very similar to the average value observed  
during the rainy season ( $-25.25_{-39.44}^{-15.31} \pm 6.33\%$  , n = 40), after the pronounced decrease of  $\text{NH}_3$  concentrations  
due to wet removal. This rather minor difference can hardly be ascribed to the influence of BB emissions, given  
265 the large seasonal fluctuation of wildfire intensity mentioned above. Based on the absolute  $\delta^{15}\text{N-NH}_3$  values in the  
urban settings, and their rather invariant temporal trends, we argue that vehicle/transport-related is a more important  
and apparently steady source of pyrogenic  $\text{NH}_3$  in the studied urban areas.

The two suburban sites (S1 and S2) are located geographically within the transition zone between the urban and  
rural environments, and this transitional character seems also indicated by their intermediate  $\delta^{15}\text{N-NH}_3$  values  
270 ( $-29.94_{-45.62}^{-18.78} \pm 7.35\%$  , n = 34). However, interestingly, in comparison to the urban and rural sites, the overall  
 $\delta^{15}\text{N-NH}_3$  value for the two suburban sites was significantly (p < 0.01) higher during the BB season  
( $-24.84_{-28.56}^{-21.49} \pm 2.29\%$  , n = 14) than that during non-BB season ( $-33.52_{-45.62}^{-18.78} \pm 7.59\%$  , n = 20) (Fig. 5). In  
fact, among the three different land-use regimes, during the BB season, the average  $\delta^{15}\text{N-NH}_3$  was highest for the  
suburban sites, and closer the  $\text{NH}_3$  isotopic signatures of pyrogenic sources, raising questions as to how important  
275 the contribution of BB versus road traffic in the suburban areas during BB season really is.

设置了格式: 字体: 倾斜

### 3.4 Isotope-based quantification of BB contribution to ambient NH<sub>3</sub>

There are several challenges that need to be overcome when trying to more accurately quantify the contribution of BB emissions to ambient NH<sub>3</sub> based on N isotope data. Firstly, given the use of only one isotope parameter ( $\delta^{15}\text{N}$ ; in contrast to NO<sub>x</sub> where also the  $\delta^{18}\text{O}$  can be analyzed), more than three potential NH<sub>3</sub> sources (e.g., urban and suburban sites) will introduce large uncertainties in isotopic endmember mixing models in terms of quantifying their relative contributions to the ambient NH<sub>3</sub> (Chang et al., 2015). Secondly, atmospheric wet scavenging could further compromise/alter the primary NH<sub>3</sub> N isotopic signatures (Elliott et al., 2019; Zheng et al., 2018; Chang et al., 2019a). For these reasons, we focus here on the samples collected at the three rural sites during the dry BB season (lasting seven weeks) to isotopically examine the contribution of BB emissions to ambient NH<sub>3</sub>. We separated these samples into seven groups based on the week of their sampling, and we integrated the measured  $\delta^{15}\text{N-NH}_3$  values as well as the N isotopic signatures of potential NH<sub>3</sub> sources (i.e., biomass burning, livestock breeding, fertilizer application) into the Bayesian isotopic mixing model (see Text S1 for details). The results of NH<sub>3</sub> source apportionment are reported in Fig. 6. With a certain degree of variability, the contribution of BB to the ambient NH<sub>3</sub> in the rural areas during the seven weeks of sampling in the dry season was only 21.0% ( $\pm$  4.7%). Hence, NH<sub>3</sub> emission from BB significantly less important than from livestock breeding (37.1  $\pm$  7.1%) and fertilizer application (41.8  $\pm$  5.9%). This comes at a surprise, given the fact that the study area belongs to one of the most important BB regions in SE Asia, or even in the world, and the samples used for isotopic source apportionment were collected during the season of intensive BB.




295 Figure 6. Source apportionment results of ambient NH<sub>3</sub> in rural areas during the dry season based on Bayesian isotopic mixing modelling and the isotopic source signatures. The error bars indicate two standard deviations.

During the dry season, we also analyzed particulate potassium (K<sup>+</sup>), a chemical tracer of biomass combustion, at two rural sites (S4, S5; in 39 daily fine-particle (PM<sub>2.5</sub>) samples). The particulate K<sup>+</sup> data offer a valuable opportunity to validate our isotope-based source apportionment results. Again, we divided the dry-season data set into seven groups based on the week of NH<sub>3</sub> passive sampling. The correlation between the particulate K<sup>+</sup> concentration and the total NH<sub>3</sub> concentration at the rural sites was rather poor ( $r^2 = 0.43$ ; blue symbols in Fig. 7a). Such a weak correlation supports our conclusion regarding the isotope-based source apportionment results (see above), providing additional independent evidence that BB can hardly be the dominant source of NH<sub>3</sub> during the sampling period at the studied rural areas. In contrast, the correlation between the particulate K<sup>+</sup> concentration and the estimated BB-derived NH<sub>3</sub> concentration (instead of total NH<sub>3</sub>) is much better ( $r^2 = 0.76$ ; Fig. 7a), and thus further validates our modeling approach. While the independent particulate K<sup>+</sup> data further increase our confidence in the N-isotope based assessment, still some uncertainty remains with regards to the robustness of the endmember source  $\delta^{15}\text{N}$  values, potential source-altering effects, and in turn our estimates on the BB-associated NH<sub>3</sub> contribution. In other words, the latter is probably sensitive to the considered range in the  $\delta^{15}\text{N}$  of potential

310 NH<sub>3</sub> emission sources, and this range may be quite large/uncertain at least for some of the sources. The δ<sup>15</sup>N-NH<sub>3</sub> from BB, in particular, is only poorly constrained, with hardly any reports from the literature (e.g., Kawashima and Kurahashi, 2011). In recent chamber experiments, we found that the δ<sup>15</sup>N-NH<sub>3</sub> produced by combustion of a variety of biomass types (subtropical trees and agricultural residues) ranged between -11.8‰ and -4.6‰ (Chang et al., 2016, 2019a), which is distinctly lower than the N isotopic signature of BB-emitted NH<sub>3</sub>-NH<sub>4</sub><sup>+</sup> (12‰) determined previously (Kawashima and Kurahashi, 2011), and adopted in this study. Assuming that the true N isotopic signatures of BB-emitted NH<sub>3</sub> in the study area falls somewhere within the range of -12‰ and 12‰ (based on our published data (Chang et al., 2016,2019a) and the value reported in (Kawashima and Kurahashi, 2011)), we re-calculated the source apportionment estimates as function of the different δ<sup>15</sup>N values for BB-emitted NH<sub>3</sub> (Fig. 7b). The estimates are not sensitive to the choice of the N isotopic composition of the BB-associated NH<sub>3</sub> source. Specifically, independent of the chosen δ<sup>15</sup>N-NH<sub>3</sub> value, BB is always the least important of the three main NH<sub>3</sub> sources in rural areas, contributing not more than 29.6%. This is because that although the isotopic signatures of BB-emitted NH<sub>3</sub> have a wide range of δ<sup>15</sup>N values, their δ<sup>15</sup>N-NH<sub>3</sub> values are still significantly (p < 0.01) higher (i.e., without overlap as shown in Fig. 4) than the measured δ<sup>15</sup>N values of ambient NH<sub>3</sub> at the rural sites.

设置了格式: 上标



325 Figure 7. (a) Scatter plots of the aerosol K<sup>+</sup> concentrations versus total NH<sub>3</sub> concentrations, as well as the NH<sub>3</sub> concentrations from BB emissions, at the rural sites during the dry season. (b) Bayesian isotope modelling-based

source apportionment results of ambient  $\text{NH}_3$  at the rural sites during dry season, as function of the assumed N isotopic signatures of BB-emitted  $\text{NH}_3$ .

As illustrated by the pie chart in Fig. 6, the average contribution of BB to ambient  $\text{NH}_3$  at the rural sites during the season of intensive fire events is  $2.4 \mu\text{g m}^{-3}$ , which can be regarded as the maximum possible concentration of BB-emitted  $\text{NH}_3$  for the urban and rural sites, which are much further away from the fire areas. Based on the total  $\text{NH}_3$  concentrations measured at the other sites, we calculate, in turn, that the contribution of BB to the ambient  $\text{NH}_3$  in the urban and suburban areas are on the order of 9.6% (ranging from 5.2% to 14.8%) and 12.3% (ranging from 6.1% to 19.9%), respectively.

#### 335 **4 Conclusion**

In this study, we integrated satellite constraints on atmospheric  $\text{NH}_3$  levels and fire intensity, discrete  $\text{NH}_3$  concentration measurement, and N isotopic analysis of  $\text{NH}_3$  in order to assess the regional-scale contribution of BB to ambient  $\text{NH}_3$  in the heartland of Southeast Asia. The combined approach provides a cross-validation framework for source apportioning of  $\text{NH}_3$  in the lower atmosphere and will thus help to ameliorate predictions of BB emissions beyond the tropics, particularly in areas of high vegetation fire risk. Our results suggest that during the dry wildfire season, BB emissions represent a ubiquitous but comparatively small  $\text{NH}_3$  source, which accounts for 9.6%, 12.3%, and 21.0% of ambient  $\text{NH}_3$  in urban, suburban, and rural environments, respectively. While we do not claim that our results necessarily apply also at the global scale, and we do not question that globally BB is one of the most important  $\text{NH}_3$  sources, we find that at least in the heartland of SE Asia, BB related  $\text{NH}_3$  emissions to the atmosphere are rather moderate, and vary significantly in time and space. Both satellite observations and field/ground-based measurements capture these variations. Our findings underscore that BB-induced  $\text{NH}_3$  emissions in tropical monsoon environments can be much lower than previously anticipated. Existing atmospheric transport models may overestimate current, and likely future,  $\text{NH}_3$  emissions under changing climate conditions.

While the full implications of our results remain to be explored, they promise to provide important guidance for  
350 revising NH<sub>3</sub> emissions from BB in atmospheric transport models to assess on air quality, human health and climate  
change.

### **Acknowledgements**

This study was supported by the International (regional) Cooperation and Exchange Project (NSFC-TRF project;  
Grant no. 41761144056), the National Natural Science Foundation of China (Grant nos. 41975166, 41977305,  
355 41761144056, 41705100), the Provincial Natural Science Foundation of Jiangsu (Grant no. BK20180040,  
BK20170946), the special fund of State Key Joint Laboratory of Environment Simulation and Pollution Control  
(Grant no. 19K01ESPCT). Lieven Clarisse and Martin Van Damme are respectively a research associate and a  
postdoctoral researcher supported by the F.R.S.-FNRS.

### **Conflict of interest**

360 The authors declare that they have no competing interests.

### **Data availability statement**

Time series of data used in this paper can be found online (at <https://doi.org/10.5281/zenodo.4025673>).

### **References**

- 365 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D., and  
Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric  
models, *Atmos. Chem. Phys.*, 11, 4039-4072, 10.5194/acp-11-4039-2011, 2011.  
Andreae, M. O., and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global  
Biogeochem Cy.*, 15, 955-966, 10.1029/2000gb001382, 2001.  
Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment,  
370 *Atmos. Chem. Phys.*, 19, 8523-8546, 10.5194/acp-19-8523-2019, 2019.

- Aneja, V. P., Schlesinger, W. H., and Erisman, J. W.: Farming pollution, *Nat Geosci*, 1, 409-411, Doi 10.1038/Ngeo236, 2008.
- Aouizerats, B., van der Werf, G. R., Balasubramanian, R., and Betha, R.: Importance of transboundary transport of biomass burning emissions to regional air quality in Southeast Asia during a high fire event, *Atmos. Chem. Phys.*, 15, 363-373, 10.5194/acp-15-363-2015, 2015.
- 375 Asman, W. A., Sutton, M. A., and Schjørring, J. K.: Ammonia: emission, atmospheric transport and deposition, *New Phytol*, 139, 27-48, 1998.
- Bauters, M., Drake, T. W., Verbeeck, H., Bodé, S., Hervé-Fernández, P., Zito, P., Podgorski, D. C., Boyemba, F., Makelele, I., Cizungu Ntaboba, L., Spencer, R. G. M., and Boeckx, P.: High fire-derived nitrogen deposition on central African forests, *Proceedings of the National Academy of Sciences*, 115, 549-554, 10.1073/pnas.1714597115, 2018.
- 380 Betha, R., Behera, S. N., and Balasubramanian, R.: 2013 Southeast Asian Smoke Haze: Fractionation of Particulate-Bound Elements and Associated Health Risk, *Environmental Science & Technology*, 48, 4327-4335, 10.1021/es405533d, 2014.
- 385 Bikkina, S., Haque, M. M., Sarin, M., and Kawamura, K.: Tracing the Relative Significance of Primary versus Secondary Organic Aerosols from Biomass Burning Plumes over Coastal Ocean Using Sugar Compounds and Stable Carbon Isotopes, *ACS Earth and Space Chemistry*, 3, 1471-1484, 10.1021/acsearthspacechem.9b00140, 2019.
- Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., VanderHoek, K. W., and Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, *Global Biogeochem Cy*, 11, 561-587, 1997.
- 390 Bray, C. D., Battye, W., Aneja, V. P., Tong, D. Q., Lee, P., and Tang, Y.: Ammonia emissions from biomass burning in the continental United States, *Atmospheric Environment*, 187, 50-61, <https://doi.org/10.1016/j.atmosenv.2018.05.052>, 2018.
- Carmichael, G. R., Ferm, M., Thongboonchoo, N., Woo, J.-H., Chan, L. Y., Murano, K., Viet, P. H., Mossberg, C., Bala, R., Boonjawat, J., Upatum, P., Mohan, M., Adhikary, S. P., Shrestha, A. B., Pienaar, J. J., Brunke, E. B., Chen, T., Jie, T., Guoan, D., Peng, L. C., Dhiharto, S., Harjanto, H., Jose, A. M., Kimani, W., Kirouane, A., Lacaux, J.-P., Richard, S., Barturen, O., Cerda, J. C., Athayde, A., Tavares, T., Cotrina, J. S., and Bilici, E.: Measurements of sulfur dioxide, ozone and ammonia concentrations in Asia, Africa, and South America using passive samplers, *Atmospheric Environment*, 37, 1293-1308, [https://doi.org/10.1016/S1352-2310\(02\)01009-9](https://doi.org/10.1016/S1352-2310(02)01009-9), 2003.
- 400 Chang, Y., Deng, C., Dore, A. J., and Zhuang, G.: Human Excreta as a Stable and Important Source of Atmospheric Ammonia in the Megacity of Shanghai, *Plos One*, 10, e0144661, 2015.
- Chang, Y., Liu, X., Deng, C., Dore, A. J., and Zhuang, G.: Source apportionment of atmospheric ammonia before, during, and after the 2014 APEC summit in Beijing using stable nitrogen isotope signatures, *Atmospheric Chemistry and Physics*, 16, 11635-11647, 10.5194/acp-16-11635-2016, 2016a.
- 405 Chang, Y., and Ma, H.: Comment on "Fossil Fuel Combustion-Related Emissions Dominate Atmospheric Ammonia Sources during Severe Haze Episodes: Evidence from 15N-Stable Isotope in Size-Resolved Aerosol Ammonium", *Environ Sci Technol*, 50, 10765-10766, 10.1021/acs.est.6b03458, 2016.
- 410 Chang, Y., Zou, Z., Deng, C., Huang, K., Collett, J. L., Lin, J., and Zhuang, G.: The importance of vehicle emissions as a source of atmospheric ammonia in the megacity of Shanghai, *Atmospheric Chemistry and Physics*, 16, 3577-3594, 10.5194/acp-16-3577-2016, 2016b.

- Chang, Y., Zhang, Y. L., Li, J., Tian, C., Song, L., Zhai, X., Zhang, W., Huang, T., Lin, Y. C., Zhu, C., Fang, Y., Lehmann, M. F., and Chen, J.: Isotopic constraints on the atmospheric sources and formation of nitrogenous species in clouds influenced by biomass burning, *Atmos. Chem. Phys.*, 19, 12221-12234, 10.5194/acp-19-12221-2019, 2019a.
- 415 Chang, Y., Zou, Z., Zhang, Y., Deng, C., Hu, J., Shi, Z., Dore, A. J., and Collett, J.: Assessing contributions of agricultural and non-agricultural emissions to atmospheric ammonia in a Chinese megacity, *Environmental Science & Technology*, 10.1021/acs.est.8b05984, 2019b.
- Chang, Y., Clarisse, L., Van Damme, M., Tao, Y., Zou, Z., Dore, A. J., and Collett, J. L.: Ammonia  
420 Emissions from Mudflats of River, Lake, and Sea, *ACS Earth and Space Chemistry*, 10.1021/acsearthspacechem.0c00017, 2020.
- Chang, Y. H., Liu, X. J., Dore, A. J., and Li, K.: Stemming PM<sub>2.5</sub> pollution in China: Re-evaluating the role of ammonia, aviation and non-exhaust road traffic emissions, *Environmental science & technology*, 46, 13035-13036, 10.1021/es304806k, 2012.
- 425 Chu, J.-E., Kim, K.-M., Lau, W. K. M., and Ha, K.-J.: How Light-Absorbing Properties of Organic Aerosol Modify the Asian Summer Monsoon Rainfall?, *Journal of Geophysical Research: Atmospheres*, 123, 2244-2255, 10.1002/2017jd027642, 2018.
- Clarisse, L., Clerbaux, C., Dentener, F., Hurtmans, D., and Coheur, P.-F.: Global ammonia distribution derived from infrared satellite observations, *Nat Geosci*, 2, 479-483, 10.1038/ngeo551, 2009.
- 430 Clarisse, L., Shephard, M. W., Dentener, F., Hurtmans, D., Cady-Pereira, K., Karagulian, F., Van Damme, M., Clerbaux, C., and Coheur, P.-F.: Satellite monitoring of ammonia: A case study of the San Joaquin Valley, *Journal of Geophysical Research: Atmospheres*, 115, 10.1029/2009JD013291, 2010.
- Crutzen, P. J., Heidt, L. E., Krasnec, J. P., Pollock, W. H., and Seiler, W.: Biomass burning as a source of atmospheric gases CO, H<sub>2</sub>, N<sub>2</sub>O, NO, CH<sub>3</sub>Cl and COS, *Nature*, 282, 253, 10.1038/282253a0, 1979.
- 435 Crutzen, P. J., and Andreae, M. O.: Biomass Burning in the Tropics: Impact on Atmospheric Chemistry and Biogeochemical Cycles, *Science*, 250, 1669-1678, 10.1126/science.250.4988.1669, 1990.
- Elliott, E. M., Yu, Z., Cole, A. S., and Coughlin, J. G.: Isotopic advances in understanding reactive nitrogen deposition and atmospheric processing, *Science of The Total Environment*, 662, 393-403, <https://doi.org/10.1016/j.scitotenv.2018.12.177>, 2019.
- 440 Felix, J. D., Elliott, E. M., Gish, T. J., McConnell, L. L., and Shaw, S. L.: Characterizing the isotopic composition of atmospheric ammonia emission sources using passive samplers and a combined oxidation-bacterial denitrifier approach, *Rapid communications in mass spectrometry : RCM*, 27, 2239-2246, 10.1002/rcm.6679, 2013.
- Hantson, S., Arneeth, A., Harrison, S. P., Kelley, D. I., Prentice, I. C., Rabin, S. S., Archibald, S., Mouillot, F., Arnold, S. R., Artaxo, P., Bachelet, D., Ciais, P., Forrest, M., Friedlingstein, P., Hickler, T., Kaplan, J. O., Kloster, S., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Meyn, A., Sitch, S., Spessa, A., van der Werf, G. R., Voulgarakis, A., and Yue, C.: The status and challenge of global fire modelling, *Biogeosciences*, 13, 3359-3375, 10.5194/bg-13-3359-2016, 2016.
- 445 Huang, C., Hu, Q., Lou, S., Tian, J., Wang, R., Xu, C., An, J., Ren, H., Ma, D., Quan, Y., Zhang, Y., and Li, L.: Ammonia Emission Measurements for Light-Duty Gasoline Vehicles in China and Implications for Emission Modeling, *Environmental Science & Technology*, 52, 11223-11231, 10.1021/acs.est.8b03984, 2018.
- 450

- Huang, K., Fu, J. S., Hsu, N. C., Gao, Y., Dong, X., Tsay, S.-C., and Lam, Y. F.: Impact assessment of biomass burning on air quality in Southeast and East Asia during BASE-ASIA, *Atmospheric Environment*, 78, 291-302, <https://doi.org/10.1016/j.atmosenv.2012.03.048>, 2013.
- 455 Kawashima, H., and Kurahashi, T.: Inorganic ion and nitrogen isotopic compositions of atmospheric aerosols at Yurihonjo, Japan: Implications for nitrogen sources, *Atmospheric environment*, 45, 6309-6316, 2011.
- Layman, C. A., Araujo, M. S., Boucek, R., Hammerschlag-Peyer, C. M., Harrison, E., Jud, Z. R., Matich, P., Rosenblatt, A. E., Vaudo, J. J., Yeager, L. A., Post, D. M., and Bearhop, S.: Applying stable isotopes to examine food-web structure: an overview of analytical tools, *Biological Reviews*, 87, 545-562, [10.1111/j.1469-185X.2011.00208.x](https://doi.org/10.1111/j.1469-185X.2011.00208.x), 2012.
- 460 Lee, H. H., Bar-Or, R. Z., and Wang, C.: Biomass burning aerosols and the low-visibility events in Southeast Asia, *Atmos. Chem. Phys.*, 17, 965-980, [10.5194/acp-17-965-2017](https://doi.org/10.5194/acp-17-965-2017), 2017.
- 465 Li, F., Zhang, X., and Kondragunta, S.: Biomass Burning in Africa: An Investigation of Fire Radiative Power Missed by MODIS Using the 375 m VIIRS Active Fire Product, *Remote Sensing*, 12, [10.3390/rs12101561](https://doi.org/10.3390/rs12101561), 2020.
- Li, Q., Jiang, J., Cai, S., Zhou, W., Wang, S., Duan, L., and Hao, J.: Gaseous Ammonia Emissions from Coal and Biomass Combustion in Household Stoves with Different Combustion Efficiencies, *Environmental Science & Technology Letters*, 3, 98-103, [10.1021/acs.estlett.6b00013](https://doi.org/10.1021/acs.estlett.6b00013), 2016.
- 470 Liu, D., Fang, Y., Tu, Y., and Pan, Y.: Chemical method for nitrogen isotopic analysis of ammonium at natural abundance, *Anal Chem*, 86, 3787-3792, [10.1021/ac403756u](https://doi.org/10.1021/ac403756u), 2014.
- Liu, X., Zhang, Y., Han, W., Tang, A., Shen, J., Cui, Z., Vitousek, P., Erisman, J. W., Goulding, K., Christie, P., Fangmeier, A., and Zhang, F.: Enhanced nitrogen deposition over China, *Nature*, 494, 459-462, [10.1038/nature11917](https://doi.org/10.1038/nature11917), 2013.
- 475 Lobert, J. M., Scharffe, D. H., Hao, W. M., and Crutzen, P. J.: Importance of biomass burning in the atmospheric budgets of nitrogen-containing gases, *Nature*, 346, 552-554, [10.1038/346552a0](https://doi.org/10.1038/346552a0), 1990.
- Marlier, M. E., DeFries, R. S., Voulgarakis, A., Kinney, P. L., Randerson, J. T., Shindell, D. T., Chen, Y., and Faluvegi, G.: El Niño and health risks from landscape fire emissions in southeast Asia, *Nat Clim Chang*, 3, 131-136, [10.1038/nclimate1658](https://doi.org/10.1038/nclimate1658), 2013.
- 480 Martin, R. V.: Satellite remote sensing of surface air quality, *Atmospheric Environment*, 42, 7823-7843, <https://doi.org/10.1016/j.atmosenv.2008.07.018>, 2008.
- Pan, X., Ichoku, C., Chin, M., Bian, H., Darmenov, A., Colarco, P., Ellison, L., Kucsera, T., da Silva, A., Wang, J., Oda, T., and Cui, G.: Six Global Biomass Burning Emission Datasets: Inter-comparison and Application in one Global Aerosol Model, *Atmos. Chem. Phys. Discuss.*, 2019, 1-39, [10.5194/acp-2019-475](https://doi.org/10.5194/acp-2019-475), 2019.
- 485 Parnell, A. C., Inger, R., Bearhop, S., and Jackson, A. L.: Source partitioning using stable isotopes: coping with too much variation, *Plos One*, 5, e9672, 2010.
- Paulot, F., and Jacob, D. J.: Hidden cost of US agricultural exports: particulate matter from ammonia emissions, *Environmental science & technology*, 48, 903-908, 2014.
- 490 Paulot, F., Paynter, D., Ginoux, P., Naik, V., Whitburn, S., Van Damme, M., Clarisse, L., Coheur, P.-F., and Horowitz, L. W.: Gas-aerosol partitioning of ammonia in biomass burning plumes: Implications for the interpretation of spaceborne observations of ammonia and the radiative forcing of ammonium nitrate, *Geophysical Research Letters*, 44, 8084-8093, [10.1002/2017gl074215](https://doi.org/10.1002/2017gl074215), 2017.

- 495 Puchalski, M. A., Sather, M. E., Walker, J. T., Lehmann, C. M., Gay, D. A., Mathew, J., and Robarge,  
W. P.: Passive ammonia monitoring in the United States: Comparing three different sampling devices,  
Journal of Environmental Monitoring, 13, 3156-3167, 2011.
- Shi, Y., Matsunaga, T., and Yamaguchi, Y.: High-Resolution Mapping of Biomass Burning Emissions in  
Three Tropical Regions, Environmental Science & Technology, 49, 10806-10814,  
10.1021/acs.est.5b01598, 2015.
- 500 Souri, A. H., Choi, Y., Jeon, W., Kochanski, A. K., Diao, L., Mandel, J., Bhave, P. V., and Pan, S.:  
Quantifying the Impact of Biomass Burning Emissions on Major Inorganic Aerosols and Their  
Precursors in the U.S, Journal of Geophysical Research: Atmospheres, 122, 12,020-012,041,  
10.1002/2017jd026788, 2017.
- 505 Streets, D. G., Canty, T., Carmichael, G. R., de Foy, B., Dickerson, R. R., Duncan, B. N., Edwards, D. P.,  
Haynes, J. A., Henze, D. K., Houyoux, M. R., Jacob, D. J., Krotkov, N. A., Lamsal, L. N., Liu, Y., Lu,  
Z., Martin, R. V., Pfister, G. G., Pinder, R. W., Salawitch, R. J., and Wecht, K. J.: Emissions estimation  
from satellite retrievals: A review of current capability, Atmospheric Environment, 77, 1011-1042,  
<https://doi.org/10.1016/j.atmosenv.2013.05.051>, 2013.
- 510 Sun, K., Tao, L., Miller, D. J., Pan, D., Golston, L. M., Zondlo, M. A., Griffin, R. J., Wallace, H. W.,  
Leong, Y. J., Yang, M. M., Zhang, Y., Mauzerall, D. L., and Zhu, T.: Vehicle Emissions as an Important  
Urban Ammonia Source in the United States and China, Environmental Science & Technology, 51,  
2472-2481, 10.1021/acs.est.6b02805, 2017.
- Sutton, M. A., Erisman, J. W., Dentener, F., and Moller, D.: Ammonia in the environment: From ancient  
515 times to the present, Environmental Pollution, 156, 583-604, DOI 10.1016/j.envpol.2008.03.013, 2008.
- Sutton, M. A., Oenema, O., Erisman, J. W., Leip, A., van Grinsven, H., and Winiwarter, W.: Too much  
of a good thing, Nature, 472, 159-161, Doi 10.1038/472159a, 2011.
- Tang, Y. S., Braban, C. F., Dragosits, U., Dore, A. J., Simmons, I., van Dijk, N., Poskitt, J., Dos Santos  
Pereira, G., Keenan, P. O., Conolly, C., Vincent, K., Smith, R. I., Heal, M. R., and Sutton, M. A.:  
520 Drivers for spatial, temporal and long-term trends in atmospheric ammonia and ammonium in the UK,  
Atmos. Chem. Phys., 18, 705-733, 10.5194/acp-18-705-2018, 2018.
- Teng, X., Hu, Q., Zhang, L., Qi, J., Shi, J., Xie, H., Gao, H., and Yao, X.: Identification of Major Sources  
of Atmospheric NH<sub>3</sub> in an Urban Environment in Northern China During Wintertime, Environmental  
Science & Technology, 51, 6839-6848, 10.1021/acs.est.7b00328, 2017.
- 525 Tsai, Y. I., Sopajaree, K., Chotruksa, A., Wu, H.-C., and Kuo, S.-C.: Source indicators of biomass burning  
associated with inorganic salts and carboxylates in dry season ambient aerosol in Chiang Mai Basin,  
Thailand, Atmospheric Environment, 78, 93-104, <https://doi.org/10.1016/j.atmosenv.2012.09.040>,  
2013.
- Van Damme, M., Clarisse, L., Heald, C. L., Hurtmans, D., Ngadi, Y., Clerbaux, C., Dolman, A. J.,  
530 Erisman, J. W., and Coheur, P. F.: Global distributions, time series and error characterization of  
atmospheric ammonia (NH<sub>3</sub>) from IASI satellite observations, Atmos. Chem. Phys., 14,  
2905-2922, 10.5194/acp-14-2905-2014, 2014.
- Van Damme, M., Clarisse, L., Dammers, E., Liu, X., Nowak, J. B., Clerbaux, C., Flechard, C. R., Galy-  
Lacaux, C., Xu, W., Neuman, J. A., Tang, Y. S., Sutton, M. A., Erisman, J. W., and Coheur, P. F.:  
535 Towards validation of ammonia (NH<sub>3</sub>) measurements from the IASI satellite, Atmos.  
Meas. Tech., 8, 1575-1591, 10.5194/amt-8-1575-2015, 2015a.

- Van Damme, M., Erismann, J. W., Clarisse, L., Dammers, E., Whitburn, S., Clerbaux, C., Dolman, A. J., and Coheur, P.-F.: Worldwide spatiotemporal atmospheric ammonia (NH<sub>3</sub>) columns variability revealed by satellite, *Geophysical Research Letters*, 42, 8660-8668, 10.1002/2015gl065496, 2015b.
- 540 Van Damme, M., Whitburn, S., Clarisse, L., Clerbaux, C., Hurtmans, D., and Coheur, P. F.: Version 2 of the IASI NH<sub>3</sub> neural network retrieval algorithm: near-real-time and reanalysed datasets, *Atmos. Meas. Tech.*, 10, 4905-4914, 10.5194/amt-10-4905-2017, 2017.
- Van Damme, M., Clarisse, L., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Clerbaux, C., and Coheur, P.-F.: Industrial and agricultural ammonia point sources exposed, *Nature*, 564, 99-103, 10.1038/s41586-018-0747-1, 2018.
- 545 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr, A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423-3441, 10.5194/acp-6-3423-2006, 2006.
- Wang, S. X., Xing, J., Jang, C. R., Zhu, Y., Fu, J. S., and Hao, J. M.: Impact Assessment of Ammonia Emissions on Inorganic Aerosols in East China Using Response Surface Modeling Technique, *Environmental Science & Technology*, 45, 9293-9300, 2011.
- 550 Wang, Y., Zhang, Q. Q., He, K., Zhang, Q., and Chai, L.: Sulfate-nitrate-ammonium aerosols over China: response to 2000–2015 emission changes of sulfur dioxide, nitrogen oxides, and ammonia, *Atmospheric Chemistry and Physics*, 13, 2635-2652, 10.5194/acp-13-2635-2013, 2013.
- 555 Whitburn, S., Van Damme, M., Kaiser, J. W., van der Werf, G. R., Turquety, S., Hurtmans, D., Clarisse, L., Clerbaux, C., and Coheur, P. F.: Ammonia emissions in tropical biomass burning regions: Comparison between satellite-derived emissions and bottom-up fire inventories, *Atmospheric Environment*, 121, 42-54, <https://doi.org/10.1016/j.atmosenv.2015.03.015>, 2015.
- Whitburn, S., Van Damme, M., Clarisse, L., Bauduin, S., Heald, C. L., Hadji-Lazaro, J., Hurtmans, D., Zondlo, M. A., Clerbaux, C., and Coheur, P. F.: A flexible and robust neural network IASI-NH<sub>3</sub> retrieval algorithm, *Journal of Geophysical Research: Atmospheres*, 121, 6581-6599, 10.1002/2016JD024828, 2016a.
- 560 Whitburn, S., Van Damme, M., Clarisse, L., Turquety, S., Clerbaux, C., and Coheur, P.-F.: Doubling of annual ammonia emissions from the peat fires in Indonesia during the 2015 El Niño, *Geophysical Research Letters*, 43, 11,007-011,014, 10.1002/2016gl070620, 2016b.
- Zheng, X.-D., Liu, X.-Y., Song, W., Sun, X.-C., and Liu, C.-Q.: Nitrogen isotope variations of ammonium across rain events: Implications for different scavenging between ammonia and particulate ammonium, *Environmental Pollution*, 239, 392-398, <https://doi.org/10.1016/j.envpol.2018.04.015>, 2018.
- 570 Zhu, L., Henze, D. K., Cady-Pereira, K. E., Shephard, M. W., Luo, M., Pinder, R. W., Bash, J. O., and Jeong, G.-R.: Constraining U.S. ammonia emissions using TES remote sensing observations and the GEOS-Chem adjoint model, *Journal of Geophysical Research: Atmospheres*, 118, 3355-3368, 10.1002/jgrd.50166, 2013.