

Compositions and mixing states of aerosol particles by aircraft observations in the Arctic springtime, 2018

Kouji Adachi¹, Naga Oshima¹, Sho Ohata^{2,3,4}, Atsushi Yoshida², Nobuhiro Moteki², and Makoto Koike²

5 ¹ Department of Atmosphere, Ocean, and Earth System Modeling Research, Meteorological Research Institute, Tsukuba, Japan

² Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, Tokyo, Japan

³ Institute for Space–Earth Environmental Research, Nagoya University, Nagoya, Japan

10 ⁴ Institute for Advanced Research, Nagoya University, Nagoya, Japan

Correspondence to: Kouji Adachi (adachik@mri-jma.go.jp)

15 **Abstract.** Aerosol particles were collected at various altitudes in the Arctic during the Polar Airborne Measurements and Arctic Regional Climate Model Simulation Project (PAMARCMiP 2018) conducted in the early spring of 2018. The composition, size, number fraction, and mixing state of individual aerosol particles were analyzed using transmission electron microscopy (TEM), and their sources and transport were evaluated by numerical model simulations. We found that sulfate, sea-salt, mineral-dust, K-bearing, and carbonaceous particles were the major aerosol constituents. Many particles were composed of two or more compositions that had coagulated and were coated with sulfate, organic materials, or both, and were internally mixed. The number fraction of mineral-dust and sea-salt particles decreased with increasing altitude. The K-bearing particles increased within a biomass burning (BB) plume at altitudes > 3900 m, which originated from Siberia. Chlorine in sea-salt particles was replaced with sulfate at high altitudes.

25 These results suggest that the sources, transport, and aging of Arctic aerosols largely vary depending on the altitude and air mass history. We also provide the occurrences of solid-particle inclusions (soot, fly-ash, and Fe-aggregate particles), some of which are light-absorbing ~~and potential ice-nucleating~~ particles. They were mainly emitted from anthropogenic and biomass burning sources and were embedded within other relatively large host particles. Our TEM measurements revealed, ~~for the first time,~~ the detailed mixing state of individual particles at various altitudes in the Arctic. This information facilitates the accurate evaluation of the aerosol influences on Arctic haze, radiation balance, cloud formation, and snow/ice albedo when deposited.

30

1 Introduction

35 The Arctic is sensitive to climate change. The surface-air temperature in this region is rising more rapidly than the global average, resulting in decreases in seasonal sea ice coverage and thickness (Stroeve et al., 2012). Aerosol particles are an important factor influencing the Arctic climate (Abbatt et al., 2019). They scatter and absorb solar radiation and affect cloud formation. Light-absorbing aerosol particles decrease the snow albedo when deposited on snow surfaces (Hansen and Nazarenko et al., 2004). Hence, 40 understanding their chemical and physical properties, such as size, abundance, composition, and mixing state, is required to accurately evaluate the influence of aerosol particles on the climate in this region.

The sources of the aerosol particles within the Arctic region include anthropogenic activities, the ocean, glaciers, and ground surfaces. In general, the local anthropogenic sources within the Arctic are limited because of the limited human activities. However, air pollution over the Arctic, named “Arctic haze”, due 45 to long-range transport (LRT) aerosols from mid-low latitudes has been observed in winter and spring (Stone et al., 2014; Shaw et al., 1995; Law et al., 2014; Law and Stohl, 2007). These LRT aerosol particles originate from, for example, anthropogenic emissions from urban cities in East Asia, North America, and Europe, biomass burning (BB) from Siberia and North America, and desert areas (Abbatt et al., 2019; Latham et al., 2013; Hecobian et al., 2011).

50 In the Arctic, the near-surface air temperature is low at lower altitudes, resulting in a temperature inversion, thereby forming a polar dome with little vertical mixing and removal of aerosols (Law et al., 2014; Arnold et al., 2016; Shaw et al., 1995). LRT particles travel along several pathways from their source regions ~~and to altitudes above each beyond~~ the polar dome (Law et al., 2014; Schulz et al., 2019). As a result, LRT particles are difficult to monitor via ground measurements. Thus, aircraft observations 55 have been conducted over the Arctic during field observation campaigns such as the Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS), Aerosol, Radiation, and Cloud Processes affecting Arctic Climate (ARCPAC), Arctic Study of Tropospheric Aerosol, Clouds and Radiation (ASTAR), and Network on Climate and Aerosols: Addressing Key Uncertainties in Remote Canadian Environments (NETCARE) (Law et al., 2014; Arnold et al., 2016; Brock, et al., 2011; 60 Yamanouchi et al., 2005; McNaughton et al., 2011; Willis et al., 2019; Wendisch et al., 2019; Jacob et al., 2010).

LRT particles are subject to mixing processes with other aerosol particles or gas condensation processes during transport. The composition and mixing state of individual particles largely differ from their original ones. These differences influence their optical and hygroscopic properties, thus altering their 65 climate effects. For example, soot particles coated with sulfate or organic matter exhibit an enhanced light absorption by the focusing of incoming light on the soot core, and the coatings may change them into efficient cloud condensation nuclei (CCN) (Bond et al., 2013). Information on the soot-mixing state is

also essential in climate models to accurately evaluate their atmospheric lifetime (i.e., their removal or transport efficiency) and, in turn, their direct radiative effects (Matsui and Moteki, 2020; Moteki et al., 2019).

These mixing processes occur on the surfaces of individual particles with size ranges of hundreds of nanometers. Single-particle mass spectrometry (PALMS; Murphy et al., 2006; Brock et al., 2011) and microscopy measurements have been performed to examine the composition and mixing state of individual aerosol particles in the Arctic. Brock et al. (2011) investigated the individual aerosol composition in the Arctic during the ARCPAC aircraft campaign using PALMS. In addition, various microscopy measurements of Arctic aerosol particles have been conducted using samples collected during ground observations (e.g., Anderson et al., 1992; Geng et al., 2010; Chi et al., 2015). However, electron microscopy measurements at various altitudes using an airplane have been absent during recent campaigns except that performed by Hara et al. (2003) during the ASTAR 2000 campaign. The concentration of black carbon (BC) or sulfate in the Arctic atmosphere has changed over the past decades (Gong et al., 2010), and sea ice melting is accelerating (Stroeve et al., 2012). Thus, knowledge of the individual particles in the current Arctic atmosphere is essential to better understand the influence of aerosol particles on the Arctic climate.

We used transmission electron microscopy (TEM) for single-particle analysis of aerosol particles at various altitudes in the Arctic. TEM is an off-line technique and requires aerosol particle samples collected with TEM grids, thus resulting in a low time resolution and loss of volatile materials. On the other hand, TEM directly measures the shape, composition, mixing state, and inclusions within host particles. TEM also directly identifies solid particles consisting of soot, metals, or dust, which potentially become ice-nucleating particles (INPs; Krämer et al., 2017).

The current study was conducted as part of the Polar Airborne Measurements and Arctic Regional Climate Model Simulation Project 2018 (PAMARCMiP 2018) in the early spring of 2018. The overall goal of this campaign is to better understand and quantify the interaction between atmospheric aerosols, surface optical properties, and clouds in the central Arctic. In this campaign, Yoshida et al. (2020) measured the size, abundance, and mixing state of BC and Fe-oxide (FeOx) particles using a single-particle soot photometer (SP2). Hartmann et al. (2020) also reported that particles from marine sources exhibit enhanced INP activities and characterized the aerosol particles within high-INP activity samples using TEM. The current study focused on TEM analysis and measured the composition and mixing state of aerosol particles at various altitudes. The goal of this study is to investigate the detailed individual-particle composition and mixing state to better understand aerosol climate influences, aging, and LRT in the Arctic.

2.1 Campaign and TEM sampling

The PAMARCMiP 2018 campaign was conducted from 23 March to 4 April 2018 using the research aircraft Polar 5 (Wesche et al., 2016) to observe aerosols, sea ice, and clouds. The aircraft was based at the Villum Research Station (81°36 N, 16°40 W) at Station Nord in Greenland. Fourteen flights were
 105 executed during the campaign above the Arctic Ocean and the Fram Strait at various altitudes up to ~5200 m (Fig. 1).

TEM samples were collected using an aerosol sampler (AS-16W, Arios Inc., Tokyo, Japan) with two impactor stages. The airflow was 1.0 L/min with 0.5- and 1.0-mm nozzle sizes for the fine and coarse modes, respectively. The sampled particle sizes are ~0.1-0.7 μm in aerodynamic diameter (50% cutoff
 110 size) at the fine stage and $>0.7 \mu\text{m}$ at the coarse stage. The sampler mounted 16 TEM grids under both the fine and coarse modes during each research flight. TEM 200-mesh Cu grids with a formvar carbon substrate (U1007, EM-Japan, Tokyo, Japan) were applied for sample collection. The sampling time was 19 min during regular flights and 29 min during long flights (April 3 and 4) with a 1-min pause time between each sampling. In total, we collected 139 TEM samples during the campaign. The samples
 115 were sealed after each flight and stored under dry conditions at room temperature.

2.2 TEM measurements

We recorded ~30 TEM images per each grid from all TEM samples under the fine mode and selected 26 TEM samples from eight research flights to cover the various altitudes and periods of interest for detailed composition and mixing state analysis (Table S1). A transmission electron microscope (JEM-
 120 1400, JEOL, Tokyo, Japan) equipped with an energy-dispersive X-ray spectrometer (EDS; X-Max 80 mm, Oxford Instruments, Tokyo, Japan) was used for both TEM- and scanning TEM (STEM)-mode analyses. A semiautomatic particle measurement system using the STEM-EDS measured the particle size and composition at an acceleration voltage of 120 kV and a 20-sec acquisition time. More than 200 particles around the center parts under the jet nozzle were analyzed for each sample. As the particle
 125 number concentration in the sampled air was low, the collected particles were dispersed and hardly overlapped on the substrates. The particle size was measured based on the area-equivalent diameter (AED), which was obtained from the particle area identified by making binary images using appropriate thresholds to distinguish them from the substrate in the STEM image-determined from the STEM images. The AED may be larger than the aerodynamic diameter for particles with a flat shape or low
 130 density. ~~The AED was obtained for each particle by assuming that it only consists of its representative composition with the recognition that most particles are internally mixed and contain several components.~~ Elemental mapping images of representative particles were also acquired using STEM-EDS. Details of the STEM-EDS analysis have been provided elsewhere (Adachi et al., 2019, 2020).

The STEM-EDS analysis normalizes the elemental composition to 100% among the selected elements (C, N, O, Na, Mg, Al, Si, P, S, Cl, K, Ca, V, Cr, Mn, Fe, Zn, and Pb). The detection limits of each element are determined based on one sigma of the peak intensity and calculated with EDS software (INCA Energy version 5.02, Oxford Instruments, Tokyo, Japan). The EDS technique has a limitation for the quantifications of light elements because of their relatively high self-absorption of outgoing X-ray, and the uncertainties of the EDS measurements for C, N, O, and S were evaluated within ~5 weight % (Adachi et al., 2019). This study considers only nonvolatile materials such as mineral-dust, sea-salt, K-bearing, sulfate, and non- or low-volatile carbonaceous particles (e.g., soot and tarballs). Volatile materials such as volatile organic compounds, nitrate, and water are lost after sampling and in the TEM chamber and are thus not considered.

2.3 Particle classifications

Aerosol particles and their inclusions, if any, are classified into each particle type. First, we classified each major aerosol particle type based on its composition and measured their AED. Then, if the particles embed or attach inclusions (soot, fly-ash, and Fe-oxide aggregate), we classified them based on their compositions and shapes.

2.3.1 Major types

We categorize the analyzed particles into six particle types based on their composition, i.e., 1. mineral-dust particles (both Al and Fe > 0.2 wt.%), 2. sea-salt particles (Na > 1 wt.%), 3. K-bearing particles (K > 2 wt.%), 4. sulfate particles (S > 2 wt.%), 5. carbonaceous particles (C + O > 90 wt.%), and 6. other particles. The workflow of particle classification is shown in Fig. S1. When a particle agrees with multiple particle type definitions, the particle is classified as the upper categories. This classification may underestimate the secondary particles in the lower categories in the flow chart, such as sulfate and carbonaceous particles, which partially coat or are attached to other particles. We thus consider the occurrences of secondary particles using elemental mapping images. The classification is the same as that used by Adachi et al. (2020) except for P-bearing particles, a proxy of primary biological aerosol (PBA) particles. The P-bearing particles are not included in the current study because of the lack of PBA particles.

2.3.2 Inclusions

In addition to the major particle-type category based on the composition, we identified the inclusions embedded within the particles. Soot, fly-ash, and Fe-oxide aggregate (hereafter Fe-aggregate) particles were identified as inclusions based on their shape and composition. Soot particles consist of aggregates of ~40-nm nanospheres possessing structures of concentrically wrapped, graphene-like layers. They have been defined as nanosphere-soot when analyzed using TEM (Buseck et al., 2014). We adopt the

definition of nanosphere-soot for our soot particles and employ the term soot for simplicity. We also use the term BC instead of soot in the model simulations and for the results from those instruments that measure BC by assuming that soot and BC are qualitatively the same material. An individual Fe-aggregate particle usually consists of more than two spherical ~~iron~~Fe-oxide (magnetite) particles (Moteki et al., 2017; Yoshida et al., 2018; Ohata et al., 2018; Kurisu et al., 2019; Li et al., 2017). Fe-aggregate particles mainly originate from anthropogenic sources (Moteki et al., 2017). Fly-ash particles exhibit spherical shapes and occur as either single particles or aggregates. They mainly consist of Si, Al, and Fe and originate from anthropogenic combustion sources. Nonaggregated Fe particles were categorized as fly-ash particles.

2.3.4 Model simulations

We applied the Meteorological Research Institute Earth System Model version 2 (MRI-ESM2; Yukimoto et al., 2019; Oshima et al., 2020) to estimate the possible source regions and transport pathways of the mineral-dust, sea-salt, and soot particles sampled by the aircraft. The model configurations are basically the same as those in Adachi et al. (2020) except for the simulation period and the classification of the BC sources. In the current study, the simulation period was extended to 2018 (from January 2008 to December 2018), and the BC concentrations were further divided into those originating from anthropogenic and BB sources. We used an atmospheric general circulation model (AGCM) with land processes (MRI-AGCM3.5) and the Model of Aerosol Species in the Global Atmosphere mark-2 (MASINGAR mk-2) as atmospheric and aerosol component models, respectively. The model calculates mass mixing ratios of non-sea-salt sulfate, BC, organic carbon, sea salt, mineral dust, and aerosol precursor gases. The size distributions of sea salt and mineral dust are divided into 10 discrete bins, while the other aerosol components are represented by lognormal size distributions. The model treats non-sea-salt sulfate, BC, organic carbon, sea salt, mineral dust, and aerosol precursor ~~gases.~~ The model uses a horizontal resolution with an approximately 120-km grid (TL159) and 80 vertical layers from the surface to the model top at 0.01 hPa in a hybrid sigma-pressure coordinate system. The horizontal wind fields were nudged toward the 6-hourly Japanese 55-year Reanalysis data (Kobayashi et al. 2015) to reproduce realistic meteorological fields in the simulations. We used the daily BB emissions from the global fire assimilation system dataset of Kaiser et al. (2012) and the monthly anthropogenic emissions dataset of Lamarque et al. (2010). In addition to the baseline simulation, we performed a model sensitivity simulation without BC emissions from BB sources (i.e., anthropogenic BC). The BC concentration from BB sources (hereafter referred to as BB BC) was estimated by subtracting the anthropogenic BC concentration (the sensitivity simulation) from the total BC concentration (the baseline simulation). The emissions of mineral dust and sea salt were calculated based on the meteorological conditions in the simulations (Tanaka and Chiba, 2005; Yumimoto et al., 2017).

3 Results and discussion

3.1 Number fraction and size of the major aerosol types

205 The aerosol particles in our samples mainly consist of sulfate, sea-salt, mineral-dust, K-bearing, and carbonaceous particles. They are mixed at the single-particle scale and sometimes contain inclusions. We measured the particle-size distribution and particle number fraction of each aerosol type (Figs. 2 and 3). We classified the samples based on the sampling altitude below 1000 m and above 1000 m. The aerosol particles above 1000 m (above the polar dome) were more influenced by LRT than those below 1000 m. The former was mainly influenced by local emissions, and the latter included LRT aerosol
210 particles.

The median diameter of all particles is 470 nm ($\sigma = 410$) in the AED (Fig. 2). The sulfate, K-bearing, and carbonaceous particles exhibit similar size distributions (the median AEDs are 470, 450, and 400 nm, respectively), whereas the carbonaceous particles demonstrate a sharper peak than do the others. The mineral-dust and sea-salt particles attain larger and broader size distributions than do the others
215 with median diameters of 680 and 740 nm, respectively. Samples above 1000 m had smaller size distributions for all particle types except mineral dust particles than those below 1000 m (Fig. 2). This result is consistent with that by Yamanouchi et al. (2005), who showed decreases of larger particles (200-1000 nm) at a higher altitude, whereas small particles (100-200 nm) did not largely show an altitude dependency. An explanation is that large particles are neither transported for long distances nor
220 uplifted toward high altitudes.

Sulfate particles are the most abundant in all samples except the sample collected on 31 March, 14:40, which predominantly contains sea-salt particles. The sulfate number fraction ranges from 29% to 86% (59% on average) (Table S1; Fig. 3). The second most dominant particle types are sea-salt and carbonaceous particles for the samples collected at < 1000 m and > 1000 m, respectively. Potassium-
225 bearing particles are relatively abundant (accounting for > 10% of the number fraction) in three samples collected at > 3900 m on 2, 3, and 4 April (Figs. 4 and 5). Based on the TEM observations and model simulations, we found that the above K-bearing particle-rich samples were collected from air masses influenced by BB, and we denoted them as BB samples (please refer to section 3.2 for the transport of the BB plume).

230 The size-dependent number fractions (Fig. 3) indicate that sulfate particles dominate the < 1 μm particles in the samples collected below 1000 m. In the samples collected above 1000 m, sulfate particles dominate all bin sizes except the largest size bin (> 2 μm), which only contains 12 particles and attains a low statistical significance (Fig. 3 (b)). The fraction of mineral-dust particles increases with increasing particle size. The number fractions of the sea-salt and carbonaceous particles increase
235 with increasing and decreasing particle size, respectively. In contrast to the negligible contributions of

the K-bearing particles in the samples collected below 1000 m, their number fraction in the samples collected above 1000 m is ~10% in all size bins except the largest one.

The number fraction of the aerosol types varies depending on the sampling altitude. The mineral-dust and sea-salt number fractions decrease with increasing sampling altitude (Fig. 4). The K-bearing particle fraction is high in the BB samples collected above 3900 m. The number fractions of the sulfate and carbonaceous particles did not show a clear altitude dependency in this study, although other studies have demonstrated that their absolute number concentrations depend on the altitude (e.g., Brock et al., 2011).

3.2 K-bearing particles and biomass burning plume

K-bearing particles are the second dominant type (11-37% in number fraction) in the BB samples (Fig. 5 and Table S1). The K-bearing particles mainly occur as potassium sulfate mixed with organic matter, soot, or both (Fig. 6). Other than in the K-bearing particles, K is detected as a minor component in approximately half of all particles and 76% of the soot-bearing particles.

BB smoke contains a large amount of K in addition to other aerosols, such as organic matter and soot particles (Reid et al., 2005), and K can be used as a tracer element of BB. The occurrences of the K-bearing particles in the BB samples (Fig. 6) are similar to those from BB in other areas (e.g., Li et al., 2003).

We used the modeled BB BC to identify the transport pathways of the BB samples (Fig. 7). The BB BC emissions in the model simulations show that the BB source is southeast Siberia (Fig. S2), where forest BB frequently occurs (Brock et al., 2011; Warneke et al., 2009; Schulz et al., 2019). During the campaign, BB started on approximately 20 March in the area and persisted for several months. On 28 March, the emitted BB BC plume was driven toward the northeast by a low-pressure system. Thereafter, the BB plume was lifted and subsequently transported toward the North Pole until 1 April (Fig. S2). On April 2, 3, and 4, a part of the BB plume approached the Fram Strait at ~600 hPa (~4000 m), where we conducted the sampling. Although the model simulations estimated the BB contributions only for the BC concentrations, we interpret that the K-bearing particles and other BB emissions were also transported along with BC as seen in the TEM results (Figs. 5 and 6). In terms of the BB samples, the sampling points were > 4600 km away from the BB sources, and the samples had aged for a week or more (Figs. 7 and S2).

3.3 Sea-salt particles

Sea-salt particles are globally abundant, especially over oceans, and considerably influence the climate as CCN and via sunlight scattering (Lewi and Schwartz, 2004). They are formed from seawater film droplets over the open ocean and local leads in the Arctic. When the sea freezes, sea-salt particles form

on frost flowers over sea ice (Hara et al., 2017; Xu et al., 2016) and from blowing snow (Huang and
270 Jaeglé, 2017). Our samples could originate from these sources as our research flights flew over both
open water (near Svalbard islands) and sea ice (near Greenland) (Fig. 1). Original sea-salt particles
mainly consist of sodium chloride as well as other inorganic salts (e.g., Mg, Ca, and K as chlorides or
sulfates). In the atmosphere, the composition of sea-salt particles is altered through the reactions with
acidic gases, thus forming sodium sulfate or nitrate (Adachi and Buseck, 2015; Gard et al., 1998;
275 Yoshizue et al., 2019), [and with organic matters \(Laskin et al., 2012; Chi et al., 2015\)](#).

In our samples, the sea-salt particles are complex mixtures of, for example, sodium chloride, sodium
sulfate, magnesium sulfate, Mg-C-O, calcium sulfate, and others (Fig. 8 and Fig. S3). The sea-salt
occurrences are similar to those found at a ground site in Svalbard (Chi et al., 2015). Although nitrate
can react with sea-salt particles, N was rarely detected in our sea-salt particles, possibly because the
280 nitrate fraction relative to that of sulfate is limited in spring (Brock et al., 2011; Fenger et al., 2013). It is
also possible that the measured particle sizes are too small for nitrate to retain as a particle phase and
that nitrates are lost from the TEM samples after sampling because of their high volatility.

Some Mg occurs around NaCl cores (Fig. 8a-b). These mixtures may form either in the atmosphere or
on the substrate when liquid particles change into solid phases after sampling. Such Mg occurs with C
285 and O as an amorphous phase, suggesting that they constitute organic matter. Similar organic Mg-C
coatings on sea-salt particles have been observed [over California by Laskin et al. \(2012\)](#) and in the
Arctic (Chi et al., 2015). As Mg salts exhibit a high hygroscopicity, it is possible that they are liquid in
the atmosphere and absorb organic matter either from anthropogenic or marine sources (Shaw et al.,
2010). Some sea-salt particles demonstrate homogeneous distributions of Mg, S, and Na (Fig. 8c and
290 Fig. S3e-f). These particles commonly attach soot particles, and Cl is replaced with sulfate, suggesting
that they are well aged in the atmosphere.

Most Cl in the sea-salt particles was replaced with sulfate in the samples collected above 1000 m (Fig.
S4). Such Cl loss in NaCl at high altitudes in the Arctic has also been observed by Hara et al. (2002).
Mg and Ca are correlated with Na in all samples, although their weight percent is lower in the samples
295 collected above 1000 m than those collected below 1000 m (Fig. 9). This result suggests that most Mg,
Ca, and Na originates from sea-salt particles and that their weight percent within the individual particles
decreases with increasing sampling altitudes because of the condensation and coagulation of sulfate or
other material on the sea-salt particles.

In the model simulations, the sea-salt concentrations near the sea surface are high (Fig. S5). On 30 and
300 31 March, the sea-salt particle concentrations are higher than those on the other sampling days based on
the TEM analysis (Table S1), and the model simulation indicates that the sea-salt particles are
transported from the north.

3.4 Mineral-dust particles

Certain types of mineral-dust particles (e.g., feldspar) act as INPs (Kanji et al., 2017) and have a climatological impact on the Arctic (Fan, 2013). In our samples, the number fraction of the mineral-dust particles increases with increasing particle size (Fig. 3) and decreasing altitude (Fig. 4). They mainly consist of Si, Al, or both and include Na, Mg, K, Ca, and Fe as minor components or small grains (e.g., Fe) (Fig. 10). They exhibit irregular shapes with crystalline structures. The mineral-dust particles are commonly mixed with sea-salt (Fig. 10a and S6) and sulfate particles (Fig. 10c), both of which change the mineral-dust particles from hydrophobic to hydrophilic. Such mixtures have also been observed in low-latitude regions (e.g., the Amazon basin) in other studies (Adachi et al., 2020), suggesting that mixtures of mineral dust and hygroscopic particles (e.g., sea salt and sulfate) occur globally. The Na and Cl weight percent within the mineral-dust particles increases in the samples collected below 1000 m (Fig. S6), indicating that mineral-dust and sea-salt particle mixing occurs near the surface. These results suggest that the mineral-dust particles originate from local ground surfaces. However, although mineral-dust particles may have local sources in summer, such as glacial outwash planes (Tobo et al., 2019) and bare ground surfaces, the ground surfaces in Greenland or the Svalbard islands were mostly, if not completely, covered by snow and ice during the sampling period (March and April). Our model simulation does not include mineral-dust emissions in the Arctic during the sampling period because of the snow coverage and yields only a small amount of LRT mineral dust above ~700 hPa in the sampling area (Fig. S7). McNaughton et al. (2011) measured the dust mass concentration over the Western Arctic in spring during the ARCTAS/ARCPAC 2008 campaigns and demonstrated that the mineral-dust concentration was high at > 4 km in altitude, where an Asian dust plume possibly influenced. They also showed that mineral-dust particles decrease toward the surface, although the sea-salt mass vertical distributions were consistent with our results. On the one hand, long-term ground observations of the aerosol composition at station Nord have indicated small contributions from soil particles in the early spring, although they exhibit peaks in summer (Nguyen et al., 2013; Heidam et al., 1999). Groot Zwaafink et al. (2016) also showed a simulation result of substantial contributions of mineral dust from Eurasia and north America >60°N to the surface dust concentrations at the Arctic region during early spring. Their studies agree with our observations, suggesting that the mineral-dust particles originate from the Arctic sources. ~~In contrast to these previous studies, our observation suggests that mineral-dust particles could be emitted from local ground surfaces, but the source of the mineral-dust particles in our samples cannot be confirmed.~~ As the number fractions of the mineral-dust particles are nonnegligible (4% on average), further observations are necessary to identify their exact sources.

3.5 Sulfate particles

Sulfate particles are one of the most dominant aerosol species in the Arctic (Brock et al., 2011; Hara et al., 2003; Matsui et al., 2011). This study also showed that sulfate particles were the most dominant

species (~60%) in their number fraction (Fig. 3). The sulfate particle number fractions are not well correlated with the altitude ($R^2=0.16$), suggesting that the detected sulfate particles and their precursors originate from both marine surface sources (Willis et al., 2017) and LRT at high altitudes. Sulfate commonly mixes with other species, and S is detected in almost all sampled particles. Sulfate particles generally exhibit various chemical forms, including ammonium sulfate, sodium sulfate, potassium sulfate, and calcium sulfate (Figs. 6 and 10; Fig. S3). The sulfate particles commonly have satellite structures around their rims in the TEM images (Fig. 11), suggesting that sulfuric acid and ammonium bisulfate contribute to the sulfate particles (Kojima et al., 2004). This result is consistent with other measurements in the Arctic that show contributions of acidic sulfate (Brock et al., 2011; Hara et al., 2003; Fisher et al., 2011). In addition, Yu et al. (2019) reported similar satellite structures from samples collected at an Arctic ground site (Ny-Ålesund, Svalbard Islands) during late summer and showed sulfate ($^{32}\text{S}^-$ and $^{16}\text{O}^-$) as well as $^{12}\text{C}^{14}\text{N}^-$ signals as a proxy of organic matter on the satellites. Although our TEM measurements did not detect C and N on the satellites, a trace amount of organic matter may be possible in the Arctic sulfates.

3.6 Carbonaceous particles

Carbonaceous particles primarily consist of C and O and include secondary and primary organic aerosols, soot particles, and tarballs. Secondary organic aerosols coat or embed other species and are mostly classified into the categories of their host species. Volatile organic compounds are likely lost before the analysis, and thus, their number fraction may be underestimated over the mass fraction measured using online instruments. Tarballs or tarball-like particles are spherical organic particles originating from BB (Pósfai et al., 2004) (Fig. 12) and are a major aerosol type originating from BB as brown carbon (Chakrabarty et al., 2010; Sedlacek et al., 2018). As a result, they potentially influence the climate, although their detailed occurrences, including their removal processes, remain unknown. In the BB samples, we encountered a small number of tarball particles (<1 % in number fractions). These tarballs mainly consist of C and O and include some N and K (Fig. 12), ~~and~~ their main composition is similar to that of particles from young BB smoke plumes (e.g., Pósfai et al., 2004; Adachi and Buseck, 2011), but, different from the young ones, some had sulfate coatings (Fig. 12). Moroni et al. (2017, 2020) also found tarballs with K-bearing particles using scanning electron microscopy in ground observations on the Svalbard Islands in the summer of 2015. Tarballs from Siberia BB were also found over the northwestern Pacific in 2016, showing a transport of tarballs toward the Arctic region (Yoshizue et al., 2020). In contrast to the smooth spherical shapes of fresh and several days aged tarballs (e.g., Li et al., 2003; Yoshizue et al., 2020), the surfaces of our tarballs are not smooth and contain sulfates (Fig. 12), ~~suggestingsuggesting~~ that they reacted with ~~other species and had aged~~ sulfate during LRT. The tarballs having sulfate are removed more efficiently from the atmosphere by precipitation than the original ones as the sulfate absorbs water. As the mass fraction of tarballs in fresh

BB smoke may reach ~40% (Sedlaeck et al., 2018), which is much higher than that in our samples, most tarballs could have been removed from the atmosphere during LRT. This tarball removal process has been proposed by Pósfai et al. (2004) when they first characterized tarballs, and our study provides evidence of the above-hypothesized processes.

Soot particles were also commonly found in our samples. However, only 13% of the carbonaceous particles consisted of soot with thin or no coatings, i.e., external mixtures, ~~and~~ ~~†~~ They were mainly embedded within or attached to other particles (internal mixtures) and were classified as a part of their host species.

3.7 Soot, fly-ash, and Fe-aggregate inclusions

Inclusions are particles embedded within or attached to host particles. Independent from the classifications of their host particles, ~~†~~ inclusions ~~are~~ ~~were~~ identified based on both their composition and shape determined from the TEM images after the removal of beam-sensitive materials (e.g., sulfate) by exposure to an electron beam during the STEM-EDS analysis (Fig. 13 and Fig. S8). When they overlap with non-beam-sensitive materials (e.g., mineral dust and sea salt), it is difficult to detect inclusions in the TEM images. Thus, the actual number fraction of the inclusions may be larger than that reported in this study. On the other hand, this technique has the advantage of identifying small inclusions (e.g., < 50 nm), which are difficult to detect using other methods. Although we observed several metal particles as inclusions (e.g., Zn and Pb), we focused on the occurrences of soot, fly-ash, and Fe-aggregate particles in this study.

3.7.1 Soot inclusions

Soot particles absorb solar radiation and exert a positive radiative forcing (Bond et al., 2013). When they are coated by or embedded within other species (e.g., sulfate and organic matter), their light absorption is enhanced by the focusing of light on the soot core (Bond et al., 2006; Oshima et al., 2009). Studies involving Arctic BC observations have shown a 54% increase in BC light absorption due to its coating (Zanatta et al., 2018) or an approximately 20% increase in the direct radiative effect due to the different BC mixing state assumptions (Kodros et al., 2018). In addition, although soot particles are initially hydrophobic, such coatings commonly make them hydrophilic. The degrees of the soot mixing states with hygroscopic materials influence their atmospheric lifetime in models, resulting in uncertainty in the simulated atmospheric BC concentrations in the Arctic (Eckhardt et al., 2015). Thus, the mixing states of soot particles are of interest regarding their climate influences (Fierce et al., 2020).

In our samples, soot particles were found in ~17% of all measured particles (Fig. S9 and Table S1).

~~Most soot particles were internally mixed with other particles (Fig. 13 and Fig. S8).~~ The number fraction of the externally mixed soot particles, which are those without any apparent coatings, was only

~1% of all soot particles. The relatively low number fractions of the externally mixed soot particles are similar to the background conditions observed by Hara et al. (2003) and indicate that most LRT soot particles were internally mixed in the Arctic.

Soot particles were observed within all aerosol types, and the sampling altitude was not clearly
410 correlated with the soot particle fraction ($R^2=0.13$). On the other hand, more soot particles were found in large host particles than in small ones, i.e., the size distributions of the soot-bearing particles (host particles) were larger than those of all particles (Fig. S10). This result is consistent with those in other areas (Adachi and Buseck, 2008; Adachi et al., 2014). A possible explanation is that the coagulation process of soot particles with large host particles occurs more efficiently than that with small particles.

415 BB is one of the major sources of BC in the Arctic atmosphere and snow (Spackman et al., 2010; Hegg et al., 2010). In our BB samples, ~93% of the soot-bearing particles contained K, showing substantial BC contributions from BB. In comparison, the ratio was 74% for the non-BB samples. However, there was no apparent enhancement of the soot particle number fraction measured by TEM in the BB samples (Table S1). This inconsistency may be explained by the fact that we measured the number fraction and
420 that the other species, such as the K-bearing salts, sulfates, and organic matters, also increased within the BB plume, resulting in relatively lower soot number fractions in the BB samples. A comparison with number concentrations of BC by SP2 will be discussed in sect. 3.7.3.

3.7.2 Fly-ash and Fe-aggregate inclusions

The fly-ash particles and primary particles of Fe aggregates commonly exhibited spherical shapes,
425 which indicated that they formed through rapid cooling after melting or evaporation at high temperatures during emission. The shape and composition of the Fe aggregates are consistent with those observed near sources in East Asia (Moteki et al., 2017). Moteki et al. showed that Fe-aggregate particles consist of magnetite, which absorbs light and imposes warming effects on the climate. Yoshida et al. (2020) also demonstrated light-absorbing FeOx ~~Fe-oxide~~ particles using SP2 during this
430 campaign. Thus, we suggest that our Fe-aggregate particles also exhibit light-absorbing properties.

The number fractions of the particles containing fly-ash or Fe-aggregate particles are 1.4% and 0.5%, respectively (Fig. S9 and Table S1). Although these particles are commonly found in polluted areas from anthropogenic sources such as stationary combustion sources and vehicles (Li et al. 2016), there are almost no such local anthropogenic sources in the Arctic area. Instead, the fly-ash and Fe-aggregate
435 particles attain better relations with the soot number fraction ($R^2=0.35$ and 0.24 for the fly-ash and Fe aggregates, respectively; Fig. S11), and soot, fly-ash, and Fe-aggregate particles are often found in the same particles (Fig. 13 and Fig. S8). The positive relations between soot, fly-ash, and Fe-aggregate particles measured by TEM ~~Our TEM measurements~~ agree with the SP2 observations of Yoshida et al. (2020), who also found correlations among BC and FeOx ~~Fe-oxide~~ during this campaign. These results

440 indicate that the soot, fly-ash, and Fe-aggregate particles originated from anthropogenic sources through LRT.

3.7.3 Comparison between TEM and SP2 data for soot/BC and Fe-bearing particles/FeOx

445 We further compared the TEM results with SP2 data reported by Yoshida et al. (2020). The TEM and SP2 measurements for soot/BC and Fe-bearing particle/FeOx reasonably correlate for non-BB samples (Fig. S12). The R^2 values of non-BB samples for soot/BC and Fe-bearing particle/FeOx are 0.45 and 0.33, respectively. The latter has a weaker correlation possibly because of their small number in the Arctic atmosphere. Although TEM and SP2 utilize different particle properties, i.e., composition/shapes and optical properties, respectively, their correlations assure these techniques. The BB samples, on the other hand, showed outliers from the relations, especially for the soot/BC. A possible reason is that BB
450 samples have higher total particle number concentrations than non-BB samples, resulting in smaller number fractions in the TEM samples than SP2.

3.8 Implications for particle aging and the climate

In the Arctic, although there are few anthropogenic sources and vegetation, we found that many aerosol particles originated from anthropogenic and BB sources. The soot and Fe-aggregate particles have light-
455 absorbing properties, and the mineral-dust particles demonstrate water-insoluble, crystalline structures with hygroscopic attachments. These optical and ~~physical~~~~structural~~ properties influence their ~~direct~~ radiative forcing by scattering and absorbing solar radiation and their ~~indirect radiative forcing through~~ cloud modification by serving as CCN and INPs. When these light-absorbing particles are deposited on snow or ice surfaces, they reduce the surface albedo, resulting in snow/ice melting acceleration (Law
460 and Stohl, 2007). ~~Soot particles (or BC)~~ ~~has~~~~ve~~ commonly been detected at various concentrations in the snow (Mori et al., 2019; Kinease et al., 2019) and ice cores (McConnell et al., 2007) in polar regions. Their deposition process from the atmosphere onto snow/ice is important to better understand their contributions to snowmelt and albedo changes. As most soot particles are embedded with or attached to other species, mostly hygroscopic ones, they will be more easily removed from the atmosphere than
465 those without coatings.

Soot, tarballs, and Fe-aggregate particles have been intensively investigated near their sources, such as urban areas and BB smoke. This study shows the composition and shape of aged particles ~~traveled~~ (> 1 week) ~~based on our model calculations~~. We observed that ~~T~~the soot and Fe-aggregate particles still exhibit fractal structures, especially the small particles ($<$ several tens of primary particles; Fig. 13 and
470 Fig. S8), similar to those found in source regions such as East Asia (Adachi et al., 2016). This observation is inconsistent with the result showing highly compacted soot particles at a remote marine free-troposphere site (China et al., 2015). Whether soot particles become core-shell structures as they age in the atmosphere is of interest for the accurate evaluation of soot climate effects (Cappa et al.,

2012; Adachi et al., 2010). Our observation implies that the soot particles in the Arctic cannot be simply assumed to be spherical particles but should be treated as fractal particles when calculating their optical properties ~~of aged samples~~. One explanation is that soot particles consisting of several tens or fewer primary particles hardly attain highly compact shapes because of the insufficient monomer numbers to turn them into spherical aggregates. On the other hand, most soot particles were coated with sulfate or other materials, suggesting that they possess enhanced light absorption ~~and CCN activity~~. Overall effects of soot shapes and coatings observed in this study are mixtures of both positive and negative on its radiative forcing evaluations.

~~We also observed the aging of tarball particles in a BB plume, i.e., the surfaces of certain tarball particles had reacted with sulfate, making them hygroscopic. Aged tarballs are removed more efficiently from the atmosphere by precipitation than the original ones. As the mass fraction of tarballs in fresh BB smoke may reach ~40% (Sedlaeck et al., 2018), which is much higher than that in our samples, most tarballs could have been removed from the atmosphere during LRT. This tarball removal process has been proposed by Pósfai et al. (2004) when they first characterized tarballs, and our study provides evidence of the above hypothesized processes.~~

4. Summary

This study reveals that the aerosol particles in the Arctic troposphere exhibit various composition, shape, and mixing state ~~ranges~~ depending on the sampling altitude and air mass history. Sulfate is the dominant aerosol type, and sea-salt, mineral-dust, K-bearing, and carbonaceous particles are also observed as major aerosol species. The aerosol particles are commonly mixtures of several components, resulting in different optical and hygroscopic properties than the original particles. In addition to the main components, they also include soot, fly-ash, and Fe-aggregate particles, all of which are solid, primary particles originating from anthropogenic or BB sources. The aerosol particles that include soot and Fe-aggregate particles may exhibit light-absorbing properties and contribute to Arctic warming and snow/ice melting when deposited on the surface. BB is an important source of ~~the~~ aerosol particles in the Arctic area. In the BB samples, we find K-bearing and tarball particles, both of which are tracers of BB emissions. Our model simulations indicated the BB contributions from Siberia. Tarballs could have aged-traveled for a week or more after their emission and have reacted with sulfate. Many mineral-dust particles are mixed with sea salt and are relatively abundant in the samples collected near the surface. The mineral-dust occurrence suggests that they originated from the Arctic local sources but not from LRT ~~contradicts previous studies that show that they mainly originate from LRT in spring, and further studies are required to confirm their sources~~. The ability of solid particles, such as mineral-dust, soot, fly-ash, and Fe-aggregate particles, to function as INPs should be considered to evaluate their contributions to cloud formation. This study highlights a wide range of mixing states, and the mixing

states of aerosol particles after LRT should be accounted for to accurately evaluate their climate influences.

510 **Data and code availability**

The TEM and simulation data used in this publication are available upon request (adachik@mri-jma.go.jp). Access to the MRI-ESM2 code is available under a collaboration framework with the MRI.

Author contributions

515 KA conducted the TEM analysis and data processing. SO, AY, and MK executed the TEM sampling and field observations. KA and NM set up the TEM sampler. MK supervised the TEM sampling. NO performed the model simulations and analyses. KA prepared the manuscript with contributions from all coauthors.

Competing interests

The authors declare that they have no conflicts of interest.

520 **Acknowledgments**

We are indebted to all the PAMARCMiP 2018 participants for their cooperation and support. The authors also acknowledge the Alfred Wegener Institute (AWI) for both the support to conduct the PAMARCMiP 2018 campaign and the use of the Polar5 research aircraft and the skill and safety exemplified by the pilots and flight staff. We thank the financial support of the Environment Research
525 and Technology Development Fund (JPMEERF20205001, JPMEERF20202003, JPMEERF20172003, and JPMEERF20165005) of the Environmental Restoration and Conservation Agency of Japan, the Global Environmental Research Coordination System of the Ministry of the Environment of Japan, and the Arctic Challenge for Sustainability (ArCS) project ~~the ArCS~~ (JPMXD1300000000) and ArCS II (JPMXD1420318865) projects of the Ministry of Education, Culture, Sports, Science, and Technology
530 (MEXT) of Japan. KA and NO acknowledge the financial support provided by the Japan Society for the Promotion of Science (JSPS) KAKENHI (grant numbers JP26701004, JP16K16188, JP16H01772, JP18H04134, JP18H03363, JP18H05292, JP19H01972, JP19H04236, JP19K21905, and JP19H04259).

References

- 535 Abbatt, J. P. D., Leaitch, W. R., Aliabadi, A. A., Bertram, A. K., Blanchet, J.-P., Boivin-Rioux, A.,
Bozem, H., Burkart, J., Chang, R. Y. W., Charette, J., Chaubey, J. P., Christensen, R. J.,
Cirisan, A., Collins, D. B., Croft, B., Dionne, J., Evans, G. J., Fletcher, C. G., Galí, M.,
Ghahremaninezhad, R., Girard, E., Gong, W., Gosselin, M., Gourdal, M., Hanna, S. J.,
Hayashida, H., Herber, A. B., Hesaraki, S., Hoor, P., Huang, L., Hussherr, R., Irish, V. E.,
540 Keita, S. A., Kodros, J. K., Köllner, F., Kolonjari, F., Kunkel, D., Ladino, L. A., Law, K.,
Levasseur, M., Libois, Q., Liggio, J., Lizotte, M., Macdonald, K. M., Mahmood, R., Martin, R.
V., Mason, R. H., Miller, L. A., Moravek, A., Mortenson, E., Mungall, E. L., Murphy, J. G.,
Namazi, M., Norman, A.-L., amp, apos, Neill, N. T., Pierce, J. R., Russell, L. M., Schneider, J.,
Schulz, H., Sharma, S., Si, M., Staebler, R. M., Steiner, N. S., Thomas, J. L., von Salzen, K.,
545 Wentzell, J. J. B., Willis, M. D., Wentworth, G. R., Xu, J.-W., and Yakobi-Hancock, J. D.:
Overview paper: New insights into aerosol and climate in the Arctic, ~~Atmospheric Chemistry
and Physics~~*Atmos. Chem. Phys.*, 19, 2527-2560, 10.5194/acp-19-2527-2019, 2019.
- Adachi, K. and Buseck, P. R.: Internally mixed soot, sulfates, and organic matter in aerosol particles
from Mexico City, ~~Atmospheric Chemistry and Physics~~*Atmos. Chem. Phys.*, 8, 6469–6481,
550 <https://doi.org/10.5194/acp-8-6469-2008>, 2008.
- Adachi, K., and Buseck, P. R.: Atmospheric tar balls from biomass burning in Mexico, ~~Journal of
Geophysical Research~~*J. Geophys. Res.*, 116, 10.1029/2010jd015102, 2011.
- Adachi, K., and Buseck, P. R.: Changes in shape and composition of sea-salt particles upon aging in an
urban atmosphere, ~~Atmospheric Atmos. Environment~~*Environ.*, 100, 1-9,
555 [10.1016/j.atmosenv.2014.10.036](https://doi.org/10.1016/j.atmosenv.2014.10.036), 2015.
- Adachi, K., Chung, S. H., and Buseck, P. R.: Shapes of soot aerosol particles and implications for their
effects on climate, ~~Journal of Geophysical Research~~*J. Geophys. Res.*, 115,
10.1029/2009jd012868, 2010.
- Adachi, K., Oshima, N., Gong, Z., de Sá, S., Bateman, A. P., Martin, S. T., de Brito, J. F., Artaxo, P.,
560 Cirino, G. G., Sedlacek III, A. J., and Buseck, P. R.: Mixing states of Amazon-basin aerosol
particles transported over long distances using transmission electron microscopy, ~~Atmospheric
Chemistry and Physics~~*Atmos. Chem. Phys.*, 20, 11923–11939, [https://doi.org/10.5194/acp-20-
11923-2020](https://doi.org/10.5194/acp-20-11923-2020), 2020.<https://doi.org/10.5194/acp-2020-452>, 2020. (accepted)
- Adachi, K., Zaizen, Y., Kajino, M., and Igarashi, Y.: Mixing state of regionally transported soot
565 particles and the coating effect on their size and shape at a mountain site in Japan, ~~Journal of
Geophysical Research~~*J. Geophys. Res.:-Atmospheres*, 119, 5386-5396,
[10.1002/2013jd020880](https://doi.org/10.1002/2013jd020880), 2014.

- Adachi, K., Moteki, N., Kondo, Y., and Igarashi, Y.: Mixing states of light-absorbing particles measured using a transmission electron microscope and a single-particle soot photometer in Tokyo, Japan, ~~Journal of Geophysical Research~~[J. Geophys. Res.: Atmospheres](#), 121, 9153-9164, 10.1002/2016jd025153, 2016.
- Adachi, K., Sedlacek, A. J., III, Kleinman, L., Springston, S. R., Wang, J., Chand, D., Hubbe, J. M., Shilling, J. E., Onasch, T. B., Kinase, T., Sakata, K., Takahashi, Y., and Buseck, P. R.: Spherical tarball particles form through rapid chemical and physical changes of organic matter in biomass-burning smoke, *Proc. Natl. Acad. Sci. USA*, 116, 19336-19341, 10.1073/pnas.1900129116, 2019.
- Anderson, J. R., Buseck, P. R., Saucy, D. A., Pacyna, J. M.: Characterization of individual fine-fraction particles from the Arctic aerosol at Spitsbergen, May–June 1987, ~~Atmospheric~~[Atmos. Environment](#)~~Environ.~~, 26, 1747-1762, 1992.
- Arnold, S. R., Law, K. S., Brock, C. A., Thomas, J. L., Starkweather, S. M., von Salzen, K., Stohl, A., Sharma, S., Lund, M. T., Flanner, M. G., Petäjä, T., Tanimoto, H., Gamble, J., Dibb, J. E., Melamed, M., Johnson, N., Fidel, M., Tynkkynen, V. P., Baklanov, A., Eckhardt, S., Monks, S. A., Browse, J., and Bozem, H.: Arctic air pollution: Challenges and opportunities for the next decade, ~~Elementa: Science of the Anthropocene~~, 4, 000104, 10.12952/journal.elementa.000104, 2016.
- Bond, T. C., Habib, G., and Bergstrom, R. W.: Limitations in the enhancement of visible light absorption due to mixing state, ~~Journal of Geophysical Research~~[J. Geophys. Res.](#), 111, 10.1029/2006jd007315, 2006.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, ~~Journal of Geophysical Research~~[J. Geophys. Res.: Atmospheres](#), 118, 5380-5552, 10.1002/jgrd.50171, 2013.
- Brock, C. A., Cozic, J., Bahreini, R., Froyd, K. D., Middlebrook, A. M., McComiskey, A., Brioude, J., Cooper, O. R., Stohl, A., Aikin, K. C., de Gouw, J. A., Fahey, D. W., Ferrare, R. A., Gao, R. S., Gore, W., Holloway, J. S., Hübler, G., Jefferson, A., Lack, D. A., Lance, S., Moore, R. H., Murphy, D. M., Nenes, A., Novelli, P. C., Nowak, J. B., Ogren, J. A., Peischl, J., Pierce, R. B., Pilewskie, P., Quinn, P. K., Ryerson, T. B., Schmidt, K. S., Schwarz, J. P., Sodemann, H., Spackman, J. R., Stark, H., Thomson, D. S., Thornberry, T., Veres, P., Watts, L. A., Warneke,

C., and Wollny, A. G.: Characteristics, sources, and transport of aerosols measured in spring 2008 during the aerosol, radiation, and cloud processes affecting Arctic Climate (ARCPAC) Project, ~~Atmospheric Chemistry and Physics~~[Atmos. Chem. Phys.](#), 11, 2423-2453, 10.5194/acp-11-2423-2011, 2011.

Buseck, P. R., Adachi, K., Gelencsér, A., Tompa, É., and Pósfai, M.: Ns-Soot: A ~~Material~~[material-Based-based Term-term](#) for ~~Strongly-strongly Lightlight-Absorbing-absorbing Carbonaceous carbonaceous Particlesparticles~~, ~~Aerosol Science and Technology~~[Aerosol Sci.ence and Technol.ogy](#), 48, 777-788, 10.1080/02786826.2014.919374, 2014.

610 Cappa, C. D., Onasch, T. B., Massoli, P., Worsnop, D. R., Bates, T. S., Cross, E. S., Davidovits, P., Hakala, J., Hayden, K. L., Jobson, B. T., Kolesar, K. R., Lack, D. A., Lerner, B. M., Li, S. M., Mellon, D., Nuaaman, I., Olfert, J. S., Petaja, T., Quinn, P. K., Song, C., Subramanian, R., Williams, E. J., and Zaveri, R. A.: Radiative absorption enhancements due to the mixing state of atmospheric black carbon, *Science*, 337, 1078-1081, 10.1126/science.1223447, 2012.

615 Chakrabarty, R. K., Moosmüller, H., Chen, L.-W. A., Lewis, K., Arnott, W. P., Mazzoleni, C., Dubey, M. K., Wold, C. E., Hao, W. M., and Kreidenweis, S. M.: Brown carbon in tar balls from smoldering biomass combustion, ~~Atmospheric Chemistry and Physics~~[Atmos. Chem. Phys.](#), 10, 6363–6370, <https://doi.org/10.5194/acp-10-6363-2010>, 2010.

Chi, J. W., Li, W. J., Zhang, D. Z., Zhang, J. C., Lin, Y. T., Shen, X. J., Sun, J. Y., Chen, J. M., Zhang, X. Y., Zhang, Y. M., and Wang, W. X.: Sea salt aerosols as a reactive surface for inorganic and organic acidic gases in the Arctic troposphere, ~~Atmospheric Chemistry and Physics~~[Atmos. Chem. Phys.](#), 15, 11341-11353, 10.5194/acp-15-11341-2015, 2015.

China, S., Scarnato, B., Owen, R. C., Zhang, B., Ampadu, M. T., Kumar, S., Dzepina, K., Dziobak, M. P., Fialho, P., Perlinger, J. A., Hueber, J., Helmig, D., Mazzoleni, L. R., and Mazzoleni, C.: Morphology and mixing state of aged soot particles at a remote marine free troposphere site: Implications for optical properties, ~~Geophysical Research Letters~~[Geophys. Research Res. LettersLett.](#), 42, 1243-1250, 10.1002/2014gl062404, 2015.

Eckhardt, S., Quennehen, B., Olivié, D. J. L., Berntsen, T. K., Cherian, R., Christensen, J. H., Collins, W., Crepinsek, S., Daskalakis, N., Flanner, M., Herber, A., Heyes, C., Hodnebrog, Ø., Huang, L., Kanakidou, M., Klimont, Z., Langner, J., Law, K. S., Lund, M. T., Mahmood, R., Massling, A., Myriokefalitakis, S., Nielsen, I. E., Nøjgaard, J. K., Quaas, J., Quinn, P. K., Raut, J. C., Rumbold, S. T., Schulz, M., Sharma, S., Skeie, R. B., Skov, H., Uttal, T., von Salzen, K., and Stohl, A.: Current model capabilities for simulating black carbon and sulfate concentrations in the Arctic atmosphere: a multi-model evaluation using a comprehensive measurement data set,

- 635 ~~Atmospheric Chemistry and Physics~~[Atmos. Chem. Phys.](#), 15, 9413-9433, 10.5194/acp-15-9413-2015, 2015.
- Fan, S.-M.: Modeling of observed mineral dust aerosols in the arctic and the impact on winter season low-level clouds, ~~Journal of Geophysical Research~~[J. Geophys. Res.: Atmospheres](#), 118, 11,161-111,174, 10.1002/jgrd.50842, 2013.
- 640 Fenger, M., Sørensen, L. L., Kristensen, K., Jensen, B., Nguyen, Q. T., Nøjgaard, J. K., Massling, A., Skov, H., Becker, T., and Glasius, M.: Sources of anions in aerosols in northeast Greenland during late winter, ~~Atmospheric Chemistry and Physics~~[Atmos. Chem. Phys.](#), 13, 1569-1578, 10.5194/acp-13-1569-2013, 2013.
- Fierce, L., Onasch, T. B., Cappa, C. D., Mazzoleni, C., China, S., Bhandari, J., Davidovits, P., Fischer, D. A., Helgestad, T., Lambe, A. T., Sedlacek, A. J., III, Smith, G. D., and Wolff, L.: Radiative absorption enhancements by black carbon controlled by particle-to-particle heterogeneity in composition, *Proc. Natl. Acad. Sci. USA*, 117, 5196-5203, 10.1073/pnas.1919723117, 2020.
- 645 Fisher, J. A., Jacob, D. J., Wang, Q., Bahreini, R., Carouge, C. C., Cubison, M. J., Dibb, J. E., Diehl, T., Jimenez, J. L., Leibensperger, E. M., Lu, Z., Meinders, M. B. J., Pye, H. O. T., Quinn, P. K., Sharma, S., Streets, D. G., van Donkelaar, A., and Yantosca, R. M.: Sources, distribution, and acidity of sulfate–ammonium aerosol in the Arctic in winter–spring, ~~Atmospheric~~[Atmos. Environment](#)[Environ.](#), 45, 7301-7318, 10.1016/j.atmosenv.2011.08.030, 2011.
- 650 Gard, E. E., Kleeman, M. J., Gross, D. S., Hughes, L. S., Allen, J. O., Morrical, B. D., Fergenson, D. P., Dienes, T., E. Gälli, M., Johnson, R. J., Cass, G. R., and Prather, K. A.: Direct ~~Observation~~[observation](#) of ~~Heterogeneous~~[heterogeneous](#) ~~Chemistry~~[chemistry](#) in the ~~Atmosphere~~[atmosphere](#), *Science*, 279, 1184-1187, 10.1126/science.279.5354.1184, 1998.
- 655 Geng, H., Ryu, J. Y., Jung, H.-J., Chung, H., Ahn, K.-H., and Ro, C.-U.: Single-particle characterization of summertime arctic aerosols collected at Ny-Alesund, Svalbard, ~~Environmental~~[Environ. Science and Technology](#)[Technol.](#), 44, 2348–2353, <https://doi.org/10.1021/es903268j>, 2010.
- 660 Hansen, J., and Nazarenko, L.: Soot climate forcing via snow and ice albedos, *Proc. Natl. Acad. Sci. USA*, 101, 423-428, 10.1073/pnas.2237157100, 2004.
- Hara, K.: Mixing states of individual aerosol particles in spring Arctic troposphere during ASTAR 2000 campaign, ~~Journal of Geophysical Research~~[J. Geophys. Res.](#), 108, 10.1029/2002jd002513, 2003.
- 665 Hara, K., Osada, K., Nishita, C., Yamagata, S., Yamanocuchi, T., Herber, A., Matsunaga, K., Iwasaka, Y., Nagatani, M., and Nakata, H.: Vertical variations of sea-salt modification in the boundary

layer of spring Arctic during the ASTAR 2000 campaign, ~~Tellus, 54B~~~~Tellus B: Chemical and Physical Meteorology, 54~~, 361-376, 10.3402/tellusb.v54i4.16671, 2002.

670 Hara, K., Matoba, S., Hirabayashi, M., and Yamasaki, T.: Frost flowers and sea-salt aerosols over seasonal sea-ice areas in northwestern Greenland during winter–spring, ~~Atmospheric Chemistry and Physics~~~~Atmos. Chem. Phys.~~, 17, 8577-8598, 10.5194/acp-17-8577-2017, 2017.

Hartmann, M., Adachi, K., Eppers, O., Haas, C., Herber, A., Holzinger, R., Hünnerbein, A., Jäkel, E., Jentzsch, C., Pinxteren, M., Wex, H., Willmes, S., and Stratmann, F.: Wintertime ~~Airborne~~~~airborne Measurements~~~~measurements~~ of ~~Ice-ice~~~~Nucleating~~~~nucleating~~ ~~Particles~~~~particles~~ in the ~~High-high~~ Arctic: A ~~Hint-hint~~ to a ~~Marine~~~~marine~~, ~~Biogenic~~~~biogenic~~ ~~Source~~~~source~~ for ~~Ice-ice~~~~Nucleating~~~~nucleating~~ ~~Particles~~~~particles~~, ~~Geophysical Research Letters~~~~Geophys. Res. Lett.~~, 47, 10.1029/2020gl087770, 2020.

680 Hecobian, A., Liu, Z., Hennigan, C. J., Huey, L. G., Jimenez, J. L., Cubison, M. J., Vay, S., Diskin, G. S., Sachse, G. W., Wisthaler, A., Mikoviny, T., Weinheimer, A. J., Liao, J., Knapp, D. J., Wennberg, P. O., Kürten, A., Crounse, J. D., Clair, J. S., Wang, Y., and Weber, R. J.: Comparison of chemical characteristics of 495 biomass burning plumes intercepted by the NASA DC-8 aircraft during the ARCTAS/CARB-2008 field campaign, ~~Atmospheric Chemistry and Physics~~~~Atmos. Chem. Phys.~~, 11, 13325-13337, 10.5194/acp-11-13325-2011, 2011.

685 Hegg, D. A., Warren, S. G., Grenfell, T. C., and Clarke, A. D.: Sources of light-absorbing aerosol in arctic snow and their seasonal variation, ~~Atmospheric Chemistry and Physics~~~~Atmos. Chem. Phys.~~, 10, 10923-10938, 10.5194/acp-10-10923-2010, 2010.

Heidam, N. Z., Wählin, P., and Christensen, J. H.: Tropospheric ~~Gases-gases~~ and ~~Aerosols-aerosols~~ in ~~Northeast-northeast~~ Greenland, ~~Journal of the Atmospheric Sciences~~~~Sci.~~, 56, 261-278, 690 1999.

Huang, J., and Jaeglé, L.: Wintertime enhancements of sea salt aerosol in polar regions consistent with a sea ice source from blowing snow, ~~Atmospheric Chemistry and Physics~~~~Atmos. Chem. Phys.~~, 17, 3699-3712, 10.5194/acp-17-3699-2017, 2017.

Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Emmons, L. K., Ferrare, R. A., 695 Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E., Pederson, J. R., and Fisher, J. A.: The Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) mission: design, execution, and first results, ~~Atmospheric Chemistry and Physics~~~~Atmos. Chem. Phys.~~, 10, 5191-5212, 10.5194/acp-10-5191-2010, 2010.

- 700 Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J.,
Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions
estimated with a global fire assimilation system based on observed fire radiative power,
Biogeosciences, 9, 527-554, 10.5194/bg-9-527-2012, 2012.
- Kinase, T., Adachi, K., Oshima, N., Goto-Azuma, K., Ogawa-Tsukagawa, Y., Kondo, Y., Moteki, N.,
705 Ohata, S., Mori, T., Hayashi, M., Hara, K., Kawashima, H., and Kita, K.: Concentrations and
~~Size-size Distributions-distributions~~ of ~~Black-black Carbon-carbon~~ in the ~~Surface-surface Snow~~
~~snow~~ of ~~Eastern-eastern~~ Antarctica in 2011, ~~Journal of Geophysical ResearchJ. Geophys. Res.:~~
~~Atmospheres~~, 125, 10.1029/2019jd030737, 2020.
- Kobayashi, S., Ota, Y., Harada, Y., Ebita, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H.,
710 Kobayashi, C., Endo, H., Miyaoka, K., and Takahashi, K.: The JRA-55 ~~Reanalysisreanalysis~~:
General ~~Specifications-specifications~~ and ~~Basie-basic~~ ~~Characteristicscharacteristics~~, ~~Journal of~~
~~the. Meteorological-Meteorol. Society of.~~ Japan, 93, 5–48, [https://doi.org/10.2151/jmsj.2015-](https://doi.org/10.2151/jmsj.2015-001)
001, 2015.
- Kodros, J. K., Hanna, S. J., Bertram, A. K., Leaitch, W. R., Schulz, H., Herber, A. B., Zannata, M.,
715 Burkart, J., Willis, M. D., Abbatt, J. P. D., and Pierce, J. R.: Size-resolved mixing state of
black carbon in the Canadian high Arctic and implications for simulated direct radiative effect,
~~Atmospheric Chemistry and PhysicsAtmos. Chem. Phys.~~, 18, 11345-11361, 10.5194/acp-18-
11345-2018, 2018.
- Kojima, T. and Buseck, P. R.: Aerosol particles from tropical convective systems: 2. Cloud bases,
720 ~~Journal of Geophysical ResearchJ. Geophys. Res.~~, 110, 10.1029/2004jd005173, 2005.
- Krämer, M., Cziczo, D. J., Burkert-Kohn, M., Boose, Y., Wex, H., Ladino, L. A., and Kanji, Z. A.:
Overview of ~~Ice-ice Nucleating-nucleating Particlesparticles~~, Meteorol.ogical Monogr.aphs, 58,
1.1-1.33, 10.1175/amsmonographs-d-16-0006.1, 2017.
- Kurisu, M., Adachi, K., Sakata, K., and Takahashi, Y.: Stable ~~Isotope-isotope Ratios-ratios~~ of
725 ~~Combustion-combustion Iron-iron Produced-produced~~ by ~~Evaporation-evaporation~~ in a ~~Steel~~
~~steel Plantplant~~, ACS Earth and Space Chem.istry, 3, 588-598,
10.1021/acsearthspacechem.8b00171, 2019.
- Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Lioussé, C.,
Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne,
730 J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van
Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions
of reactive gases and aerosols: methodology and application, ~~Atmospheric Chemistry and~~
~~PhysicsAtmos. Chem. Phys.~~, 10, 7017-7039, 10.5194/acp-10-7017-2010, 2010.

- 735 [Laskin, A., Moffet, R. C., Gilles, M. K., Fast, J. D., Zaveri, R. A., Wang, B., Nigge, P., and Shutthanandan, J.: Tropospheric chemistry of internally mixed sea salt and organic particles: Surprising reactivity of NaCl with weak organic acids, *J. Geophys. Res.*, **117**, 10.1029/2012jd017743, 2012.](#)
- 740 Latham, T. L., Beyersdorf, A. J., Thornhill, K. L., Winstead, E. L., Cubison, M. J., Hecobian, A., Jimenez, J. L., Weber, R. J., Anderson, B. E., and Nenes, A.: Analysis of CCN activity of Arctic aerosol and Canadian biomass burning during summer 2008, [Atmospheric Chemistry and Physics](#) *Atmos. Chem. Phys.*, **13**, 2735-2756, 10.5194/acp-13-2735-2013, 2013.
- 745 Law, K. S., Stohl, A., Quinn, P. K., Brock, C. A., Burkhardt, J. F., Paris, J.-D., Ancellet, G., Singh, H. B., Roiger, A., Schlager, H., Dibb, J., Jacob, D. J., Arnold, S. R., Pelon, J., and Thomas, J. L.: Arctic [Air-air](#) Pollution: New [Insights-insights](#) from POLARCAT-IPY, [Bulletin of the American Meteorological Society](#) *Am. Meteorol. Soc.*, **95**, 1873-1895, 10.1175/bams-d-13-00017.1, 2014.
- Law, K. S., and Stohl, A.: Arctic [Air-air Pollutionpollution](#): Origins and [Impactsimpacts](#), *Science*, **315**, 1537-1540, 10.1126/science.1137695, 2007.
- 750 Lewis, E.R., Schwartz, S.E., 2004. Sea [Salt-salt Aerosol-aerosol Productionproduction](#): Mechanisms, [Methodsmethods](#), [Measurement-measurement](#) and [Model-models](#) A Critical Review. In: *Geophys. Monogr.*, vol. 152. AGU, Washington, D. C, p. 413.
- Li, J., Pósfai, M., Hobbs, P. V., and Buseck, P. R.: Individual aerosol particles from biomass burning in southern Africa: 2, Compositions and aging of inorganic particles, [Journal of Geophysical Research](#) *J. Geophys. Res.: Atmospheres*, **108**, 10.1029/2002jd002310, 2003.
- 755 Li, W., Shao, L., Zhang, D., Ro, C.-U., Hu, M., Bi, X., Geng, H., Matsuki, A., Niu, H., and Chen, J.: A review of single aerosol particle studies in the atmosphere of East Asia: morphology, mixing state, source, and heterogeneous reactions, [Journal of Cleaner Clean. ProductionProd., **112**, 1330-1349, 10.1016/j.jclepro.2015.04.050, 2016.](#)
- 760 Li, W., Xu, L., Liu, X., Zhang, J., Lin, Y., Yao, X., Gao, H., Zhang, D., Chen, J., Wang, W., Harrison, R. M., Zhang, X., Shao, L., Fu, P., Nenes, A., and Shi, Z.: Air pollution–aerosol interactions produce more bioavailable iron for ocean ecosystems, [Science-Sci. AdvancesAdv., **3**, e1601749, 10.1126/sciadv.1601749, 2017.](#)
- 765 Matsui, H., Kondo, Y., Moteki, N., Takegawa, N., Sahu, L. K., Koike, M., Zhao, Y., Fuelberg, H. E., Sessions, W. R., Diskin, G., Anderson, B. E., Blake, D. R., Wisthaler, A., Cubison, M. J., and Jimenez, J. L.: Accumulation-mode aerosol number concentrations in the Arctic during the ARCTAS aircraft campaign: Long-range transport of polluted and clean air from the Asian

continent, ~~Journal of Geophysical Research~~J. Geophys. Res., 116, 10.1029/2011jd016189, 2011.

770 Matsui, H., and Moteki, N.: High ~~Sensitivity~~sensitivity of Arctic ~~Black-black Carbon-carbon Radiative~~
~~radiative Effects-effects~~ to ~~Subgrid-subgrid Vertical-vertical Velocity-velocity~~ in ~~Aerosol~~
~~aerosol Activationactivation~~, ~~Geophysical Research Letters~~Geophys. Res. Lett., 47,
10.1029/2020gl088978, 2020.

775 McConnell, J. R., Edwards, R., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J. R.,
Pasteris, D. R., Carter, M. M., and Kahl, J. D. W.: 20th-Century-century ~~Industrial-industrial~~
~~Black-black Carbon-carbon Emissions-emissions~~ ~~Altered-altered~~ Arctic ~~Climate-climate~~
~~Forcingforcing~~, *Science*, 317, 1381-1384, 10.1126/science.1144856, 2007.

780 McNaughton, C. S., Clarke, A. D., Freitag, S., Kapustin, V. N., Kondo, Y., Moteki, N., Sahu, L.,
Takegawa, N., Schwarz, J. P., Spackman, J. R., Watts, L., Diskin, G., Podolske, J., Holloway,
J. S., Wisthaler, A., Mikoviny, T., de Gouw, J., Warneke, C., Jimenez, J., Cubison, M., Howell,
S. G., Middlebrook, A., Bahreini, R., Anderson, B. E., Winstead, E., Thornhill, K. L., Lack, D.,
Cozic, J., and Brock, C. A.: Absorbing aerosol in the troposphere of the Western Arctic during
the 2008 ARCTAS/ARCPAC airborne field campaigns, ~~Atmospheric-Chemistry-and~~
~~Physies~~Atmos. Chem. Phys., 11, 7561-7582, 10.5194/acp-11-7561-2011, 2011.

785 Mori, T., Goto-Azuma, K., Kondo, Y., Ogawa-Tsukagawa, Y., Miura, K., Hirabayashi, M., Oshima, N.,
Koike, M., Kupiainen, K., Moteki, N., Ohata, S., Sinha, P. R., Sugiura, K., Aoki, T.,
Schneebeli, M., Steffen, K., Sato, A., Tsushima, A., Makarov, V., Omiya, S., Sugimoto, A.,
Takano, S., and Nagatsuka, N.: Black ~~Carbon-carbon~~ and ~~Inorganic-inorganic Aerosols~~
~~aerosols~~ in Arctic ~~Snowpacksnowpack~~, ~~Journal of Geophysical Research~~J. Geophys. Res.:
~~Atmospheres~~, 124, 13325-13356, 10.1029/2019jd030623, 2019.

790 Moroni, B., Cappelletti, D., Crocchianti, S., Becagli, S., Caiazzo, L., Traversi, R., Udisti, R., Mazzola,
M., Markowicz, K., Ritter, C., and Zielinski, T.: Morphochemical characteristics and mixing
state of long range transported wildfire particles at Ny-Ålesund (Svalbard Islands),
~~Atmospheric-Atmos. Environment~~Environ., 156, 135-145, 10.1016/j.atmosenv.2017.02.037,
2017.

795 Moroni, B., Ritter, C., Crocchianti, S., Markowicz, K., Mazzola, M., Becagli, S., Traversi, R., Krejci,
R., Tunved, P., and Cappelletti, D.: Individual ~~Partiele-particle~~ ~~Characteristiescharacteristics~~,
~~Optical-optical Properties-properties~~ and ~~Evolution-evolution~~ of an ~~Extreme-extreme~~ ~~Longlong-~~
~~Range-range Transported-transported Biomass-biomass Burning-burning Event-event~~ in the
European Arctic (Ny-Ålesund, Svalbard Islands), ~~Journal of Geophysical Research~~J. Geophys.
800 ~~Res.:-Atmospheres~~, 125, 10.1029/2019jd031535, 2020.

- Moteki, N., Adachi, K., Ohata, S., Yoshida, A., Harigaya, T., Koike, M., and Kondo, Y.: Anthropogenic iron oxide aerosols enhance atmospheric heating, *Nat. Commun.*, 8, 15329, 10.1038/ncomms15329, 2017.
- 805 Moteki, N., Mori, T., Matsui, H., and Ohata, S.: Observational constraint of in-cloud supersaturation for simulations of aerosol rainout in atmospheric models, *npj Climate and Atmospheric Science*, 2, 10.1038/s41612-019-0063-y, 2019.
- Murphy, D. M., Cziczo, D. J., Froyd, K. D., Hudson, P. K., Matthew, B. M., Middlebrook, A. M., Peltier, R. E., Sullivan, A., Thomson, D. S., and Weber, R. J.: Single-particle mass spectrometry of tropospheric aerosol particles, *Journal of Geophysical Research*, 111, 10.1029/2006jd007340, 2006.
- 810 *J. Geophys. Res.: Atmospheres*, 111, 10.1029/2006jd007340, 2006.
- Nguyen, Q. T., Skov, H., Sørensen, L. L., Jensen, B. J., Grube, A. G., Massling, A., Glasius, M., and Nøjgaard, J. K.: Source apportionment of particles at Station Nord, North East Greenland during 2008–2010 using COPREM and PMF analysis, *Atmospheric Chemistry and Physics*, 13, 35-49, 10.5194/acp-13-35-2013, 2013.
- 815 Ohata, S., Yoshida, A., Moteki, N., Adachi, K., Takahashi, Y., Kurisu, M., and Koike, M.: Abundance of ~~Light~~~~light~~-~~Absorbing~~~~absorbing~~ ~~Anthropogenic~~~~anthropogenic~~ ~~Iron~~~~iron~~ ~~Oxide~~~~oxide~~ ~~Aerosols~~~~aerosols~~ in the ~~Urban~~~~urban~~ ~~Atmosphere~~~~atmosphere~~ and ~~Their~~~~their~~ ~~Emission~~~~emission~~ ~~Source~~~~sources~~, *Journal of Geophysical Research*, 10.1029/2018jd028363, 2018.
- 820 Oshima, N., Koike, M., Zhang, Y., and Kondo Y.: Aging of black carbon in outflow from anthropogenic sources using a mixing state resolved model: 2. Aerosol optical properties and cloud condensation nuclei activities, *Journal of Geophysical Research*, 114, D18202, 10.1029/2008JD011681, 2009.
- Oshima, N., Yukimoto, S., Deushi, M., Koshiro, T., Kawai, H., Tanaka, T. Y., and Yoshida, K.: Global and Arctic effective radiative forcing of anthropogenic gases and aerosols in MRI-ESM2.0, *Progress in Earth and Planetary Science*, 7, 38, <https://doi.org/10.1186/s40645-020-00348-w>, 2020.
- 825 Pósfai, M., Gelencsér, A., Simonics, R., Arató, K., Li, J., Hobbs, P. V., and Buseck, P. R.: Atmospheric tar balls: Particles from biomass and biofuel burning, *Journal of Geophysical Research*, 109, 10.1029/2003jd004169, 2004.
- 830 *J. Geophys. Res.: Atmospheres*, 109, 10.1029/2003jd004169, 2004.
- Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmospheric Chemistry and Physics*, 5, 799-825, 10.5194/acp-5-799-2005, 2005.

- 835 Schulz, H., Zanatta, M., Bozem, H., Leaitch, W. R., Herber, A. B., Burkart, J., Willis, M. D., Kunkel, D., Hoor, P. M., Abbatt, J. P. D., and Gerdes, R.: High Arctic aircraft measurements characterising black carbon vertical variability in spring and summer, ~~Atmospheric Chemistry and Physics~~Atmos. Chem. Phys., 19, 2361-2384, 10.5194/acp-19-2361-2019, 2019.
- 840 Sedlacek III, A. J., Buseck, P. R., Adachi, K., Onasch, T. B., Springston, S. R., and Kleinman, L.: Formation and evolution of tar balls from northwestern US wildfires, ~~Atmospheric Chemistry and Physics~~Atmos. Chem. Phys., 18, 11289–11301, [https://doi.org/10.5194/acp-18-11289-](https://doi.org/10.5194/acp-18-11289-2018) 2018, 2018.
- Shaw, G. E.: The Arctic ~~Haze-haze Phenomenon~~phenomenon, ~~Bull. Am. Meteorol. Soc.~~Bulletin of the American Meteorological Society, 76, 2403-2414, 1995.
- 845 Shaw, P. M., Russell, L. M., Jefferson, A., and Quinn, P. K.: Arctic organic aerosol measurements show particles from mixed combustion in spring haze and from frost flowers in winter, ~~Geophysical Research Letters~~Geophys. Res. Lett., 37, 10.1029/2010gl042831, 2010.
- 850 Spackman, J. R., Gao, R. S., Neff, W. D., Schwarz, J. P., Watts, L. A., Fahey, D. W., Holloway, J. S., Ryerson, T. B., Peischl, J., and Brock, C. A.: Aircraft observations of enhancement and depletion of black carbon mass in the springtime Arctic, ~~Atmospheric Chemistry and Physics~~Atmos. Chem. Phys., 10, 9667-9680, 10.5194/acp-10-9667-2010, 2010.
- Stone, R. S., Sharma, S., Herber, A., Eleftheriadis, K., and Nelson, D. W.: A characterization of Arctic aerosols on the basis of aerosol optical depth and black carbon measurements, ~~Elementa: Science of the Anthropocene~~, 2, 000027, 10.12952/journal.elementa.000027, 2014.
- 855 Stroeve, J. C., Serreze, M. C., Holland, M. M., Kay, J. E., Malanik, J., and Barrett, A. P.: The Arctic's rapidly shrinking sea ice cover: a research synthesis, ~~Climatic Clim.~~Change, 110, 1005-1027, 10.1007/s10584-011-0101-1, 2011.
- Tanaka, T. Y. and Chiba, M.: Global simulation of dust aerosol with a chemical transport model, MASINGAR. ~~Journal of the Meteorological Meteorol. Society Soc. of Japan~~, 83A, 255–278, 2005.
- 860 Tobo, Y., Adachi, K., DeMott, P. J., Hill, T. C. J., Hamilton, D. S., Mahowald, N. M., Nagatsuka, N., Ohata, S., Uetake, J., Kondo, Y., and Koike, M.: Glacially sourced dust as a potentially significant source of ice nucleating particles, ~~Nature Geoscience~~Geosci., 12, 253-258, 10.1038/s41561-019-0314-x, 2019.
- 865 Warneke, C., Bahreini, R., Brioude, J., Brock, C. A., de Gouw, J. A., Fahey, D. W., Froyd, K. D., Holloway, J. S., Middlebrook, A., Miller, L., Montzka, S., Murphy, D. M., Peischl, J., Ryerson, T. B., Schwarz, J. P., Spackman, J. R., and Veres, P.: Biomass burning in Siberia and

Kazakhstan as an important source for haze over the Alaskan Arctic in April 2008,
~~Geophysical Research Letters~~Geophys. Res. Lett., 36, 10.1029/2008gl036194, 2009.

Wendisch, M., Macke, A., Ehrlich, A., Lüpkes, C., Mech, M., Chechin, D., Dethloff, K., Velasco, C. B.,
870 Bozem, H., Brückner, M., Clemen, H.-C., Crewell, S., Donth, T., Dupuy, R., Ebell, K., Egerer,
U., Engelmann, R., Engler, C., Eppers, O., Gehrmann, M., Gong, X., Gottschalk, M.,
Gourbeyre, C., Griesche, H., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann,
H., Heygster, G., Hoor, P., Jafariserajehlou, S., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U.,
Kecorius, S., Knudsen, E. M., Köllner, F., Kretzschmar, J., Lelli, L., Leroy, D., Maturilli, M.,
875 Mei, L., Mertes, S., Mioche, G., Neuber, R., Nicolaus, M., Nomokonova, T., Notholt, J., Palm,
M., van Pinxteren, M., Quaas, J., Richter, P., Ruiz-Donoso, E., Schäfer, M., Schmieder, K.,
Schnaiter, M., Schneider, J., Schwarzenböck, A., Seifert, P., Shupe, M. D., Siebert, H., Spreen,
G., Stapf, J., Stratmann, F., Vogl, T., Welti, A., Wex, H., Wiedensohler, A., Zanatta, M., and
Zeppenfeld, S.: The Arctic ~~Cloud-cloud Puzzle~~puzzle: Using ACLOUD/PASCAL
880 ~~Multiplatform-multiplatform Observations-observations~~ to ~~Unravel-unravel~~ the ~~Role-role~~ of
~~Clouds-clouds~~ and ~~Aerosol-aerosol Particles-particles~~ in Arctic ~~Amplificationamplification~~,
~~Bulletin of the American Meteorological Society~~Bull. Am. Meteorol. Soc., 100, 841-871,
10.1175/bams-d-18-0072.1, 2019.

Wesche, C., Steinhage, D., and Nixdorf, U.: Polar aircraft Polar5 and Polar6 operated by the Alfred
885 Wegener Institute, Journal of large-scale research facilities JLSRF, 2, 10.17815/jlsrf-2-153,
2016.

Willis, M. D., Köllner, F., Burkart, J., Bozem, H., Thomas, J. L., Schneider, J., Aliabadi, A. A., Hoor, P.
M., Schulz, H., Herber, A. B., Leaitch, W. R., and Abbatt, J. P. D.: Evidence for marine
biogenic influence on summertime Arctic aerosol, ~~Geophysical Research Letters~~Geophys. Res.
890 Lett., 44, 6460-6470, 10.1002/2017gl073359, 2017.

Willis, M. D., Bozem, H., Kunkel, D., Lee, A. K. Y., Schulz, H., Burkart, J., Aliabadi, A. A., Herber, A.
B., Leaitch, W. R., and Abbatt, J. P. D.: Aircraft-based measurements of High Arctic
springtime aerosol show evidence for vertically varying sources, transport and composition,
~~Atmospheric Chemistry and Physics~~Atmos. Chem. Phys., 19, 57-76, 10.5194/acp-19-57-2019,
895 2019.

Xu, L., Russell, L. M., and Burrows, S. M.: Potential sea salt aerosol sources from frost flowers in the
pan-Arctic region, ~~Journal of Geophysical Research~~J. Geophys. Res.: Atmospheres, 121,
10,840-810,856, 10.1002/2015jd024713, 2016.

Yamanouchi, T., Treffeisen, R., Herber, A., Shiobara, M., Yamagata, S., Hara, K., Sato, K., Yabuki, M.,
900 Tomikawa, Y., Rinke, A., Neuber, R., Schumacher, R., Kriews, M., Ström, J., Schrems, O.,

and Gernandt, H.: Arctic Study of Tropospheric Aerosol and Radiation (ASTAR) 2000: Arctic haze case study, *Tellus-B: Chemical and Physical Meteorology*, 57B, 141-152, 10.3402/tellusb.v57i2.16784, 2017.

905 Yoshida, A., Ohata, S., Moteki, N., Adachi, K., Mori, T., Koike, M., and Takami, A.: Abundance and ~~Emission emission Flux-flux~~ of the ~~Anthropogenic-anthropogenic Iron-iron Oxide-oxide Aerosols-aerosols From-from~~ the ~~East-east~~ Asian ~~Continental-continental Outflowoutflow~~, *Journal of Geophysical ResearchJ. Geophys. Res.: Atmospheres*, 123, 11,194-111,209, 10.1029/2018jd028665, 2018.

910 Yoshida, A., Moteki, N., Ohata, S., Mori, T., Koike, M., Kondo, Y., Matsui, H., Oshima, N., Takami, A., and Kita, K.: Abundances and microphysical properties of light-absorbing iron oxide and black carbon aerosols over East Asia and the Arctic, *Journal of Geophysical ResearchJ. Geophys. Res.: Atmospheres*, 10.1029/2019jd032301, 2020.

915 Yoshizue, M., Iwamoto, Y., Adachi, K., Kato, S., Sun, S., Miura, K., and Uematsu, M.: Individual particle analysis of marine aerosols collected during the North–South transect cruise in the Pacific Ocean and its marginal seas, *Journal of OceanographyOceanogr.*, 75, 513-524, 10.1007/s10872-019-00519-4, 2019.

920 Yu, H., Li, W., Zhang, Y., Tunved, P., Dall'Osto, M., Shen, X., Sun, J., Zhang, X., Zhang, J., and Shi, Z.: Organic coating on sulfate and soot particles during late summer in the Svalbard Archipelago, *Atmos. Chem. Phys.*, 19, 10433–10446, <https://doi.org/10.5194/acp-19-10433-2019>, 2019.

925 Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yabu, S., Yoshimura, H., Shindo, E., Mizuta, R., Obata, A., Adachi, Y., and Ishii, M.: The Meteorological Research Institute ~~Earth-earth System-system Model model Version-version~~ 2.0, MRI-ESM2.0: Description and ~~Basic-basic Evaluation-evaluation~~ of the ~~Physical-physical Componentcomponent~~, *Journal of the Meteorological Meteorol. Society of Japan*. Ser. II, 97, 931-965, 10.2151/jmsj.2019-051, 2019.

Yumimoto, K., Tanaka, T. Y., Oshima, N., and Maki, T.: JRAero: the Japanese Reanalysis for Aerosol v1.0, *Geoscientific-Geosci. Model DevelopmentDev.*, 10, 3225-3253, 10.5194/gmd-10-3225-2017, 2017.

930 Zanatta, M., Laj, P., Gysel, M., Baltensperger, U., Vratolis, S., Eleftheriadis, K., Kondo, Y., Dubuisson, P., Winiarek, V., Kazadzis, S., Tunved, P., and Jacobi, H.-W.: Effects of mixing state on optical and radiative properties of black carbon in the European Arctic, *Atmospheric Chemistry and PhysicsAtmos. Chem. Phys.*, 18, 14037-14057, 10.5194/acp-18-14037-2018, 2018.





Figure 1. Flight tracks and locations of the TEM samplings during the PAMARCMiP 2018. The blue star indicates the Villum Research Station in Greenland. The yellow points indicate the sampling locations below 1000 m. The red points indicate those above 1000 m. The sampling points are the midpoints of the sampling time; 19 min for all samples except during the 3 and 4 April flights, which had a sampling time of 29 min.



945 **Figure 2.** Normalized size distributions of each particle type for all samples (left) and for samples at altitudes above and below 1000 m (right). The particle sizes were determined from the area-equivalent

diameters obtained from the STEM images. The size bins are shown on a log scale and are < 0.25, 0.25-0.35, 0.35-0.5, 0.5-0.71, 0.71-1.00, 1.00-1.41, and >1.41 μm . The number fraction values indicate the percentage of particles in each bin size among each particle type. The particle numbers used for the mineral-dust, sea-salt, K-bearing, sulfate, and carbonaceous particles are 7844, 337, 1205, 421, 4672, and 809, respectively.




Figure 3. Size-dependent number fractions in the samples (a) < 1000 m and (b) > 1000 m and number fractions of the total particles of the samples (c) < 1000 m and (d) > 1000 m. The size bins in (a) and (b) are shown on a log-scale and are <0.35, 0.35-0.5, 0.5-0.71, 0.71-1.00, 1.00-1.41, 1.41-2.00, and > 2.00 μm . The particle numbers used in (c) and (d) are 4400 and 3444, respectively.






Figure 4. Vertical changes in the number fractions (%) of (a) mineral-dust, (b) sea-salt, and (c) K-

bearing particles. Each plot indicates the number fraction of a given sample. The number of particles in the samples ranges from 216 to 442. The red solid-circle points in the K-bearing particle plots indicate the samples originating from the biomass burning plume (BB samples).



965 **Figure 5.** Number fractions of the aerosol particles in the BB samples. (a) Size-dependent and (b) total number fractions. Three BB samples were collected on 2, 3, and 4 April. The size bins are shown on a log-scale and are 0.35-0.5, 0.5-0.71, 0.71-1.00, 1.00-1.41, 1.41-2.00, and $> 2.00 \mu\text{m}$. (b) Number fractions of the total particles of the BB samples. The particle number of the BB samples is 1041.




970

975


Figure 6. TEM and elemental mapping images of a BB sample (10:40-10:59, 2 April 2018). (a) TEM image of K-bearing particles. (b) The same area of the TEM image (a) but after the STEM-EDS analysis. Beam-sensitive materials are removed, and soot inclusions are more apparent. (c) and (d): Examples of K-bearing particles containing soot inclusions and organic coatings, which appear in gray in the TEM image. (e) STEM and elemental mapping images for C, N, O, S, and K in the TEM image. The distributions of K, S, and O correspond to the K-bearing particles, and that of C represents the soot and organic particles.

(a) Horizontal distributions of BB BC at 600 hPa


9:00, April 2



9:00, April 3




9:00, April 4




(b) Vertical distributions of BB BC at 0° E longitude


9:00, April 2



9:00, April 3



9:00, April 4



5 10 20 40 60 80 100 150 200 500 (ng/m³)
BB BC concentration

5 10 20 40 60 80 100 150 200 500 (ng/m³)
BB BC concentration

Figure 7. Model simulations of the (a) horizontal and (b) vertical distributions of the mass

980 concentration of BC originating from biomass burning (BB) during the sampling periods of the BB samples. The white circles indicate the sampling locations. The transport pathway of the BB plume from its source area is shown in Fig. S2.




Figure 8. TEM and elemental mapping images of the sea-salt particles in a sample (10:00-10:19, 30 March 2018). Top images (a)-(c): TEM images of the sea-salt particles mixed with various grains. Bottom images (d)-(f): Elemental mapping images of the TEM areas for C, O, Na, Mg, S, Cl, K, and Ca. Soot particles are attached to the sea-salt particles (b) and (c). C-O-Mg in the particles (a) and (b) occurs around NaCl grains.




Figure 9. Relations of Na with Mg, Cl, and Ca among all particles. The plots show the average atomicweight percent values of all particles within each sample. The black and red plots are the samples < 1000 m and > 1000 m, respectively. R^2 values are 0.84, 0.61, and 0.89 for (a), (b), and (c), respectively. Approximated curves are shown in the gray dashed lines. The error bars indicate the 95% confidential intervals.




Figure 10. TEM and elemental mapping images of the mineral-dust particles. Top (a)-(c): TEM images of the mineral-dust particles. Bottom (d)-(f): Elemental mapping images of the TEM areas for the detected elements. All particles were found in the samples collected from 10:00-10:19 on 30 March 2018. Soot, NaCl, and Ca-S particles are attached to these mineral-dust particles in (a). Fe occurs as grains in (b). Sulfate is attached to the mineral-dust particle in (c).





Figure 11. TEM and elemental mapping images of the sulfate particles. Top (a)-(c): TEM images.


Bottom (d)-(f): Elemental mapping images of the TEM areas for C, N, Na, and S. All particles were collected from 14:40-14:59, 31 March 2018. Most particles mainly consist of sulfate. Some particles also include N and Na, suggesting that they are mixtures of ammonium sulfate and sodium sulfate. The carbon grains correspond to the internally mixed soot particles. Small droplets occur around relatively large sulfate particles, forming a satellite structure, as indicated in (a).



1010

Figure 12. TEM and elemental mapping images of the tarball and tarball-like particles in a BB sample (10:30-10:59, 2 April 2018). (a) TEM image of the tarball-like particles containing sulfate. As they exhibit deformed spherical shapes and the characteristic elements of tarballs (e.g., C, N, O, and K), we label them tarball-like particles. Other examples of tarballs are shown in (b) and (c). (d)-(f): Elemental mapping images of the TEM areas.

1015






Figure 13. TEM and elemental mapping images of the sulfate particles with inclusions. (a) TEM image before the mapping analysis and (b) after the mapping analysis. Electron-beam exposure removed sulfate, and inclusions became more apparent. (c) Elemental mapping images of the TEM areas for the detected elements. (d) A fly-ash particle. (e) Mineral-dust and fly-ash particles. (f) An Fe-aggregate particle. (g) Mineral-dust and soot particles. The white squares in (b) and (c) indicate the areas of (d)-(g). The sample was collected from 14:40-14:59, 31 March 2018.