



Compositions and mixing states of aerosol particles by aircraft observations in the Arctic springtime, 2018

Kouji Adachi¹, Naga Oshima¹, Sho Ohata^{2,3,4}, Atsushi Yoshida², Nobuhiro Moteki², and Makoto Koike²

5 ¹ Department of Atmosphere, Ocean, and Earth System Modeling Research, Meteorological Research Institute, Tsukuba, Japan

² Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, Tokyo, Japan

³ Institute for Space–Earth Environmental Research, Nagoya University, Nagoya, Japan

10 ⁴ Institute for Advanced Research, Nagoya University, Nagoya, Japan

Correspondence to: Kouji Adachi (adachik@mri-jma.go.jp)

15 **Abstract.** Aerosol particles were collected at various altitudes in the Arctic during the Polar Airborne Measurements and Arctic Regional Climate Model Simulation Project (PAMARCMiP 2018) conducted in the early spring of 2018. The composition, size, number fraction, and mixing state of individual aerosol particles were analyzed using transmission electron microscopy (TEM), and their sources and transport were evaluated by numerical model simulations. We found that sulfate, sea-salt, mineral-dust, K-bearing, and carbonaceous particles were the major aerosol constituents and were internally mixed. The number fraction of mineral-dust and sea-salt particles decreased with increasing altitude. The K-bearing particles increased within a biomass burning (BB) plume at altitudes > 3900 m, which originated from Siberia. Chlorine in sea-salt particles was replaced with sulfate at high altitudes. These results suggest that the sources, transport, and aging of Arctic aerosols largely vary depending on the altitude and air mass history.

20 We also provide the occurrences of solid-particle inclusions (soot, fly-ash, and Fe-aggregate particles), some of which are light-absorbing and potential ice-nucleating particles. Our TEM measurements revealed, for the first time, the detailed mixing state of individual particles at various altitudes in the Arctic. This information facilitates the accurate evaluation of the aerosol influences on Arctic haze, radiation balance, cloud formation, and snow/ice albedo when deposited.

30



1 Introduction

The Arctic is sensitive to climate change. The surface-air temperature in this region is rising more rapidly than the global average, resulting in decreases in seasonal sea ice coverage and thickness (Stroeve et al., 2012). Aerosol particles are an important factor influencing the Arctic climate (Abbatt et al., 2019). They scatter and absorb solar radiation and affect cloud formation. Light-absorbing aerosol particles decrease the snow albedo when deposited on snow surfaces (Hansen and Nazarenko et al., 2004). Hence, understanding their chemical and physical properties, such as size, abundance, composition, and mixing state, is required to accurately evaluate the influence of aerosol particles on the climate in this region.

The sources of the aerosol particles within the Arctic region include anthropogenic activities, the ocean, glaciers, and ground surfaces. In general, the local anthropogenic sources within the Arctic are limited because of the limited human activities. However, air pollution over the Arctic named Arctic haze due to long-range transport (LRT) aerosols from mid-low latitudes has been observed in winter and spring (Stone et al., 2014; Shaw et al., 1995; Law et al., 2014; Law and Stohl, 2007). These LRT aerosol particles originate from, for example, anthropogenic emissions from urban cities in East Asia, North America, and Europe, biomass burning (BB) from Siberia and North America, and desert areas (Abbatt et al., 2019; Latham et al., 2013; Hecobian et al., 2011).

In the Arctic, the near-surface air temperature is low at lower altitudes, resulting in a temperature inversion, thereby forming a polar dome with little vertical mixing and removal of aerosols (Law et al., 2014; Arnold et al., 2016; Shaw et al., 1995). LRT particles travel along several pathways from their source regions and reach beyond the polar dome (Law et al., 2014; Schulz et al., 2019). As a result, LRT particles are difficult to monitor via ground measurements. Thus, aircraft observations have been conducted over the Arctic during field observation campaigns such as the Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS), Aerosol, Radiation, and Cloud Processes affecting Arctic Climate (ARCPAC), Arctic Study of Tropospheric Aerosol, Clouds and Radiation (ASTAR), and Network on Climate and Aerosols: Addressing Key Uncertainties in Remote Canadian Environments (NETCARE) (Law et al., 2014; Arnold et al., 2016; Brock, et al., 2011; Yamanouchi et al., 2005; McNaughton et al., 2011; Willis et al., 2019; Wendisch et al., 2019; Jacob et al., 2010).

LRT particles are subject to mixing processes with other aerosol particles or gas condensation processes during transport. The composition and mixing state of individual particles largely differ from their original ones. These differences influence their optical and hygroscopic properties, thus altering their climate effects. For example, soot particles coated with sulfate or organic matter exhibit an enhanced light absorption by the focusing of incoming light on the soot core, and the coatings may change them into efficient cloud condensation nuclei (CCN) (Bond et al., 2013). Information on the soot-mixing state is



65 also essential in climate models to accurately evaluate their atmospheric lifetime (i.e., their removal or
transport efficiency) and, in turn, their direct radiative effects (Matsui and Moteki, 2020; Moteki et al.,
2019).

These mixing processes occur on the surfaces of individual particles with size ranges of hundreds of
nanometers. Single-particle mass spectrometry (PALMS; Murphy et al., 2006; Brock et al., 2011) and
70 microscopy measurements have been performed to examine the composition and mixing state of
individual aerosol particles in the Arctic. Brock et al. (2011) investigated the individual aerosol
composition in the Arctic during the ARCPAC aircraft campaign using PALMS. In addition, various
microscopy measurements of Arctic aerosol particles have been conducted using samples collected during
ground observations (e.g., Anderson et al., 1992; Geng et al., 2010; Chi et al., 2015). However, electron
75 microscopy measurements at various altitudes using an airplane have been absent during recent
campaigns except that performed by Hara et al. (2003) during the ASTAR 2000 campaign. The
concentration of black carbon (BC) or sulfate in the Arctic atmosphere has changed over the past decades
(Gong et al., 2010), and sea ice melting is accelerating (Stroeve et al., 2012). Thus, knowledge of the
individual particles in the current Arctic atmosphere is essential to better understand the influence of
80 aerosol particles on the Arctic climate.

We used transmission electron microscopy (TEM) for single-particle analysis of aerosol particles at
various altitudes in the Arctic. TEM is an off-line technique and requires aerosol particle samples
collected with TEM grids, thus resulting in a low time resolution and loss of volatile materials. On the
other hand, TEM directly measures the shape, composition, mixing state, and inclusions within host
85 particles. TEM also directly identifies solid particles consisting of soot, metals, or dust, which potentially
become ice-nucleating particles (INPs; Krämer et al., 2017).

The current study was conducted as part of the Polar Airborne Measurements and Arctic Regional Climate
Model Simulation Project 2018 (PAMARCMiP 2018) in the early spring of 2018. The overall goal of this
campaign is to better understand and quantify the interaction between atmospheric aerosols, surface
90 optical properties, and clouds in the central Arctic. In this campaign, Yoshida et al. (2020) measured the
size, abundance, and mixing state of BC and Fe-oxide particles using a single-particle soot photometer
(SP2). Hartmann et al. (2020) also reported that particles from marine sources exhibit enhanced INP
activities and characterized the aerosol particles within high-INP activity samples using TEM. The current
study focused on TEM analysis and measured the composition and mixing state of aerosol particles at
95 various altitudes. The goal of this study is to investigate the detailed individual-particle composition and
mixing state to better understand aerosol climate influences, aging, and LRT in the Arctic.



2 Methods

2.1 Campaign and TEM sampling

The PAMARCMiP 2018 campaign was conducted from 23 March to 4 April 2018 using the research
100 aircraft Polar 5 (Wesche et al., 2016) to observe aerosols, sea ice, and clouds. The aircraft was based at
the Villum Research Station (81°36 N, 16°40 W) at Station Nord in Greenland. Fourteen flights were
executed during the campaign above the Arctic Ocean and the Fram Strait at various altitudes up to
~5200 m (Fig. 1).

TEM samples were collected using an aerosol sampler (AS-16W, Arios Inc., Tokyo, Japan) with two
105 impactor stages. The airflow was 1.0 L/min with 0.5- and 1.0-mm nozzle sizes for the fine and coarse
modes, respectively. The sampled particle sizes are ~0.1-0.7 μm in aerodynamic diameter (50% cutoff
size) at the fine stage and $>0.7 \mu\text{m}$ at the coarse stage. The sampler mounted 16 TEM grids under both
the fine and coarse modes during each research flight. TEM 200-mesh Cu grids with a formvar carbon
substrate (U1007, EM-Japan, Tokyo, Japan) were applied for sample collection. The sampling time was
110 19 min during regular flights and 29 min during long flights (April 3 and 4) with a 1-min pause time
between each sampling. In total, we collected 139 TEM samples during the campaign. The samples
were sealed after each flight and stored under dry conditions at room temperature.

2.2 TEM measurements

We recorded ~30 TEM images per each grid from all TEM samples under the fine mode and selected 26
115 TEM samples from eight research flights to cover the various altitudes and periods of interest for
detailed composition and mixing state analysis (Table S1). A transmission electron microscope (JEM-
1400, JEOL, Tokyo, Japan) equipped with an energy-dispersive X-ray spectrometer (EDS; X-Max 80
mm, Oxford Instruments, Tokyo, Japan) was used for both TEM- and scanning TEM (STEM)-mode
analyses. A semiautomatic particle measurement system using the STEM-EDS measured the particle
120 size and composition at an acceleration voltage of 120 kV and a 20-sec acquisition time. More than 200
particles around the center parts under the jet nozzle were analyzed for each sample. As the particle
number concentration in the sampled air was low, the collected particles were dispersed and hardly
overlapped on the substrates. The particle size was measured based on the area-equivalent diameter
(AED) determined from the STEM images. The AED may be larger than the aerodynamic diameter for
125 particles with a flat shape or low density. The AED was obtained for each particle by assuming that it
only consists of its representative composition with the recognition that most particles are internally
mixed and contain several components. Elemental mapping images of representative particles were also
acquired using STEM-EDS. Details of the STEM-EDS analysis have been provided elsewhere (Adachi
et al., 2019, 2020).



130 The STEM-EDS analysis normalizes the element composition to 100% among the selected elements (C,
N, O, Na, Mg, Al, Si, P, S, Cl, K, Ca, V, Cr, Mn, Fe, Zn, and Pb). The detection limits of each element
are determined based on one sigma of the peak intensity and calculated with EDS software (INCA
Energy version 5.02, Oxford Instruments, Tokyo, Japan). This study considers only nonvolatile
135 materials such as mineral-dust, sea-salt, K-bearing, sulfate, and non- or low-volatile carbonaceous
particles (e.g., soot and tarballs). Volatile materials such as volatile organic compounds, nitrate, and
water are lost after sampling and in the TEM chamber and are thus not considered. We categorize the
analyzed particles into six particle types based on their composition, i.e., mineral-dust particles (both Al
and Fe > 0.2 wt.%), sea-salt particles (Na > 1 wt.%), K-bearing particles (K > 2 wt.%), sulfate particles
(S > 2 wt.%), carbonaceous particles (C + O > 90 wt.%), and other particles. The workflow of particle
140 classification is shown in Fig. S1. This classification may underestimate the particles in the lower
categories in the flow chart, such as sulfate and carbonaceous particles, which partially coat or are
attached to other particles. The classification is the same as that used by Adachi et al. (2020) except for
P-bearing particles, a proxy of primary biological aerosol (PBA) particles. The P-bearing particles are
not included in the current study because of the lack of PBA particles.

145 In addition to the particle-type category based on the composition, we identified the inclusions
embedded within the particles. Soot, fly-ash, and Fe-oxide aggregate (hereafter Fe-aggregate) particles
were identified as inclusions based on their shape and composition. Soot particles consist of aggregates
of ~40-nm nanospheres possessing structures of concentrically wrapped, graphene-like layers. They
have been defined as nanosphere-soot when analyzed using TEM (Buseck et al., 2014). We adopt the
150 definition of nanosphere-soot for our soot particles and employ the term soot for simplicity. We also use
the term BC instead of soot in the model simulations and for the results from those instruments that
measure BC by assuming that soot and BC are qualitatively the same material. An individual Fe-
aggregate particle usually consists of more than two spherical iron-oxide (magnetite) particles (Moteki
et al., 2017; Yoshida et al., 2018; Ohata et al., 2018; Kurisu et al., 2019; Li et al., 2017). Fe-aggregate
155 particles mainly originate from anthropogenic sources (Moteki et al., 2017). Fly-ash particles exhibit
spherical shapes and occur as either single particles or aggregates. They mainly consist of Si, Al, and Fe
and originate from anthropogenic combustion sources. Nonaggregated Fe particles were categorized as
fly-ash particles.

2.3 Model simulations

160 We applied the Meteorological Research Institute Earth System Model version 2 (MRI-ESM2;
Yukimoto et al., 2019; Oshima et al., 2020) to estimate the possible source regions and transport
pathways of the mineral-dust, sea-salt, and soot particles sampled by the aircraft. The model
configurations are basically the same as those in Adachi et al. (2020) except for the simulation period



and the classification of the BC sources. In the current study, the simulation period was extended to
165 2018 (from January 2008 to December 2018), and the BC concentrations were further divided into those
originating from anthropogenic and BB sources. We used an atmospheric general circulation model
(AGCM) with land processes (MRI-AGCM3.5) and the Model of Aerosol Species in the Global
Atmosphere mark-2 (MASINGAR mk-2) as atmospheric and aerosol component models, respectively.
The model treats non-sea-salt sulfate, BC, organic carbon, sea salt, mineral dust, and aerosol precursor
170 gases. The model uses a horizontal resolution with an approximately 120-km grid (TL159) and 80
vertical layers from the surface to the model top at 0.01 hPa in a hybrid sigma-pressure coordinate
system. The horizontal wind fields were nudged toward the 6-hourly Japanese 55-year Reanalysis data
(Kobayashi et al. 2015) to reproduce realistic meteorological fields in the simulations. We used the
daily BB emissions from the global fire assimilation system dataset of Kaiser et al. (2012) and the
175 monthly anthropogenic emissions dataset of Lamarque et al. (2010). In addition to the baseline
simulation, we performed a model sensitivity simulation without BC emissions from BB sources (i.e.,
anthropogenic BC). The BC concentration from BB sources (hereafter referred to as BB BC) was
estimated by subtracting the anthropogenic BC concentration (the sensitivity simulation) from the total
BC concentration (the baseline simulation). The emissions of mineral dust and sea salt were calculated
180 based on the meteorological conditions in the simulations (Tanaka and Chiba, 2005; Yumimoto et al.,
2017).

3 Results and discussion

3.1 Number fraction and size of the major aerosol types

The aerosol particles in our samples mainly consist of sulfate, sea-salt, mineral-dust, K-bearing, and
185 carbonaceous particles. They are mixed at the single-particle scale and sometimes contain inclusions.
We measured the particle-size distribution and particle number fraction of each aerosol type (Figs. 2
and 3). We classified the samples based on the sampling altitude below 1000 m and above 1000 m. The
former was mainly influenced by local emissions, and the latter included LRT aerosol particles.

The median diameter of all particles is 470 nm ($\sigma = 410$) in the AED (Fig. 2). The sulfate, K-bearing,
190 and carbonaceous particles exhibit similar size distributions (the median AEDs are 470, 450, and 400
nm, respectively), whereas the carbonaceous particles demonstrate a sharper peak than do the others.
The mineral-dust and sea-salt particles attain larger and broader size distributions than do the others
with median diameters of 680 and 740 nm, respectively.

Sulfate particles are the most abundant in all samples except the sample collected on 31 March, 14:40,
195 which predominantly contains sea-salt particles. The sulfate number fraction ranges from 29% to 86%
(59% on average) (Table S1; Fig. 3). The second most dominant particle types are sea-salt and



carbonaceous particles for the samples collected at < 1000 m and > 1000 m, respectively. Potassium-bearing particles are relatively abundant (accounting for $> 10\%$ of the number fraction) in three samples collected at > 3900 m on 2, 3, and 4 April (Figs. 4 and 5). Based on the TEM observations and model
200 simulations, we found that the above K-bearing particle-rich samples were collected from air masses influenced by BB, and we denoted them as BB samples (please refer to section 3.2 for the transport of the BB plume).

The size-dependent number fractions (Fig. 3) indicate that sulfate particles dominate the $< 1 \mu\text{m}$ particles in the samples collected below 1000 m. In the samples collected above 1000 m, sulfate
205 particles dominate all bin sizes except the largest size bin ($> 2 \mu\text{m}$), which only contains 12 particles and attains a low statistical significance (Fig. 3 (b)). The fraction of mineral-dust particles increases with increasing particle size. The number fractions of the sea-salt and carbonaceous particles increase with increasing and decreasing particle size, respectively. In contrast to the negligible contributions of the K-bearing particles in the samples collected below 1000 m, their number fraction in the samples
210 collected above 1000 m is $\sim 10\%$ in all size bins except the largest one.

The number fraction of the aerosol types varies depending on the sampling altitude. The mineral-dust and sea-salt number fractions decrease with increasing sampling altitude (Fig. 4). The K-bearing particle fraction is high in the BB samples collected above 3900 m. The number fractions of the sulfate and carbonaceous particles did not show a clear altitude dependency in this study, although other
215 studies have demonstrated that their absolute number concentrations depend on the altitude (e.g., Brock et al., 2011).

3.2 K-bearing particles and biomass burning plume

K-bearing particles are the second dominant type (11-37% in number fraction) in the BB samples (Fig. 5 and Table S1). The K-bearing particles mainly occur as potassium sulfate mixed with organic matter, soot, or both (Fig. 6). Other than in the K-bearing particles, K is detected as a minor component in
220 approximately half of all particles and 76% of the soot-bearing particles.

BB smoke contains a large amount of K in addition to other aerosols, such as organic matter and soot particles (Reid et al., 2005), and K can be used as a tracer element of BB. The occurrences of the K-bearing particles in the BB samples (Fig. 6) are similar to those from BB in other areas (e.g., Li et al.,
225 2003).

We used the modeled BB BC to identify the transport pathways of the BB samples (Fig. 7). The BB BC emissions in the model simulations show that the BB source is southeast Siberia (Fig. S2), where forest BB frequently occurs (Brock et al., 2011; Warneke et al., 2009; Schulz et al., 2019). During the campaign, BB started on approximately 20 March in the area and persisted for several months. On 28
230 March, the emitted BB BC plume was driven toward the northeast by a low-pressure system.



Thereafter, the BB plume was lifted and subsequently transported toward the North Pole until 1 April (Fig. S2). On April 2, 3, and 4, a part of the BB plume approached the Fram Strait at ~600 hPa (~4000 m), where we conducted the sampling. Although the model simulations estimated the BB contributions only for the BC concentrations, we interpret that the K-bearing particles and other BB emissions were also transported along with BC as seen in the TEM results. In terms of the BB samples, the sampling points were > 4600 km away from the BB sources, and the samples had aged for a week or more.

3.3 Sea-salt particles

Sea-salt particles are globally abundant, especially over oceans, and considerably influence the climate as CCN and via sunlight scattering (Lewi and Schwartz, 2004). They are formed from seawater film droplets over the open ocean and local leads in the Arctic. When the sea freezes, sea-salt particles form on frost flowers over sea ice (Hara et al., 2017; Xu et al., 2016) and from blowing snow (Huang and Jaeglé, 2017). Our samples could originate from these sources as our research flights flew over both open water (near Svalbard islands) and sea ice (near Greenland) (Fig. 1). Original sea-salt particles mainly consist of sodium chloride as well as other inorganic salts (e.g., Mg, Ca, and K as chlorides or sulfates). In the atmosphere, the composition of sea-salt particles is altered through the reactions with acidic gases, thus forming sodium sulfate or nitrate (Adachi and Buseck, 2015; Gard et al., 1998; Yoshizue et al., 2019).

In our samples, the sea-salt particles are complex mixtures of, for example, sodium chloride, sodium sulfate, magnesium sulfate, Mg-C-O, calcium sulfate, and others (Fig. 8 and Fig. S3). The sea-salt occurrences are similar to those found at a ground site in Svalbard (Chi et al., 2015). Although nitrate can react with sea-salt particles, N was rarely detected in our sea-salt particles, possibly because the nitrate fraction relative to that of sulfate is limited in spring (Brock et al., 2011; Fenger et al., 2013). It is also possible that the measured particle sizes are too small for nitrate to retain as a particle phase and that nitrates are lost from the TEM samples after sampling because of their high volatility.

Some Mg occurs around NaCl cores (Fig. 8a-b). These mixtures may form either in the atmosphere or on the substrate when liquid particles change into solid phases after sampling. Such Mg occurs with C and O as an amorphous phase, suggesting that they constitute organic matter. Similar organic Mg-C coatings on sea-salt particles have been observed in the Arctic (Chi et al., 2015). As Mg salts exhibit a high hygroscopicity, it is possible that they are liquid in the atmosphere and absorb organic matter either from anthropogenic or marine sources (Shaw et al., 2010). Some sea-salt particles demonstrate homogeneous distributions of Mg, S, and Na (Fig. 8c and Fig. S3e-f). These particles commonly attach soot particles, and Cl is replaced with sulfate, suggesting that they are well aged in the atmosphere.

Most Cl in the sea-salt particles was replaced with sulfate in the samples collected above 1000 m (Fig. S4). Such Cl loss in NaCl at high altitudes in the Arctic has also been observed by Hara et al. (2002).



265 Mg and Ca are correlated with Na in all samples, although their weight percent is lower in the samples collected above 1000 m than those collected below 1000 m (Fig. 9). This result suggests that most Mg, Ca, and Na originates from sea-salt particles and that their weight percent within the individual particles decreases with increasing sampling altitudes because of the condensation and coagulation of sulfate or other material on the sea-salt particles.

270 In the model simulations, the sea-salt concentrations near the sea surface are high (Fig. S5). On 30 and 31 March, the sea-salt particle concentrations are higher than those on the other sampling days based on the TEM analysis (Table S1), and the model simulation indicates that the sea-salt particles are transported from the north.

3.4 Mineral-dust particles

275 Certain types of mineral-dust particles (e.g., feldspar) act as INPs (Kanji et al., 2017) and have a climatological impact on the Arctic (Fan, 2013). In our samples, the number fraction of the mineral-dust particles increases with increasing particle size (Fig. 3) and decreasing altitude (Fig. 4). They mainly consist of Si, Al, or both and include Na, Mg, K, Ca, and Fe as minor components or small grains (e.g., Fe) (Fig. 10). They exhibit irregular shapes with crystalline structures. The mineral-dust particles are

280 commonly mixed with sea-salt and sulfate particles, both of which change the mineral-dust particles from hydrophobic to hydrophilic. Such mixtures have also been observed in low-latitude regions (e.g., the Amazon basin) in other studies (Adachi et al., 2020), suggesting that mixtures of mineral dust and hygroscopic particles (e.g., sea salt and sulfate) occur globally. The Na and Cl weight percent within the mineral-dust particles increases in the samples collected below 1000 m (Fig. S6), indicating that

285 mineral-dust and sea-salt particle mixing occurs near the surface. These results suggest that the mineral-dust particles originate from local ground surfaces. However, although mineral-dust particles may have local sources in summer, such as glacial outwash planes (Tobo et al., 2019) and bare ground surfaces, the ground surfaces in Greenland or the Svalbard islands were mostly, if not completely, covered by snow and ice during the sampling period (March and April). Our model simulation does not include

290 mineral-dust emissions in the Arctic during the sampling period because of the snow coverage and yields only a small amount of LRT mineral dust above ~700 hPa in the sampling area (Fig. S7). Long-term ground observations of the aerosol composition at station Nord have indicated small contributions from soil particles in the early spring, although they exhibit peaks in summer (Nguyen et al., 2013; Heidam et al., 1999). McNaughton et al. (2011) measured the dust mass concentration over the Western

295 Arctic in spring during the ARCTAS/ARCPAC 2008 campaigns and demonstrated that the mineral-dust concentration was high at > 4 km in altitude, where an Asian dust plume possibly influenced. They also showed that mineral-dust particles decrease toward the surface, although the sea-salt mass vertical distributions were consistent with our results. In contrast to these previous studies, our observation



suggests that mineral-dust particles could be emitted from local ground surfaces, but the source of the
300 mineral-dust particles in our samples cannot be confirmed. As the number fractions of the mineral-dust
particles are nonnegligible (4% on average), further observations are necessary to identify their sources.

3.5 Sulfate particles

Sulfate particles are one of the most dominant aerosol species in the Arctic (Brock et al., 2011; Hara et
al., 2003; Matsui et al., 2011). This study also showed that sulfate particles were the most dominant
305 species (~60%) in their number fraction (Fig. 3). The sulfate particle number fractions are not well
correlated with the altitude ($R^2=0.16$), suggesting that the detected sulfate particles and their precursors
originate from both marine surface sources (Willis et al., 2017) and LRT at high altitudes. Sulfate
commonly mixes with other species, and S is detected in almost all sampled particles. Sulfate particles
generally exhibit various chemical forms, including ammonium sulfate, sodium sulfate, potassium
310 sulfate, and calcium sulfate (Figs. 6 and 10; Fig. S3). The sulfate particles commonly have satellite
structures around their rims in the TEM images (Fig. 11), suggesting that sulfuric acid and ammonium
bisulfate contribute to the sulfate particles (Kojima et al., 2004). This result is consistent with other
measurements in the Arctic that show contributions of acidic sulfate (Brock et al., 2011; Hara et al.,
2003; Fisher et al., 2011).

315 3.6 Carbonaceous particles

Carbonaceous particles primarily consist of C and O and include secondary and primary organic
aerosols, soot particles, and tarballs. Secondary organic aerosols coat or embed other species and are
mostly classified into the categories of their host species. Volatile organic compounds are likely lost
before the analysis, and thus, their number fraction may be underestimated over the mass fraction
320 measured using online instruments. Tarballs or tarball-like particles are spherical organic particles
originating from BB (Pósfai et al., 2004) (Fig. 12) and are a major aerosol type originating from BB as
brown carbon (Chakrabarty et al., 2010; Sedlacek et al., 2018). As a result, they potentially influence
the climate, although their detailed occurrences, including their removal processes, remain unknown. In
the BB samples, we encountered a small number of tarball particles (<1 % in number fractions). These
325 tarballs mainly consist of C and O and include some N and K (Fig. 12), and their composition is similar
to that of particles from young BB smoke plumes (e.g., Pósfai et al., 2004; Adachi and Buseck, 2011).
Moroni et al. (2017, 2020) also found tarballs with K-bearing particles using scanning electron
microscopy in ground observations on the Svalbard Islands in the summer of 2015. In contrast to the
smooth spherical shapes of fresh tarballs (e.g., Li et al., 2003), the surfaces of our tarballs are not
330 smooth and contain sulfates (Fig. 12), suggesting that they reacted with other species and had aged
during LRT.



Soot particles were also commonly found in our samples. However, only 13% of the carbonaceous particles consisted of soot with thin or no coatings, i.e., external mixtures, and they were mainly embedded within or attached to other particles (internal mixtures) and were classified as a part of their host species.

3.7 Soot, fly-ash, and Fe-aggregate inclusions

Inclusions are particles embedded within or attached to host particles. Inclusions are identified based on both their composition and shape determined from the TEM images after the removal of beam-sensitive materials (e.g., sulfate) by exposure to an electron beam during the STEM-EDS analysis (Fig. 13 and Fig. S8). When they overlap with non-beam-sensitive materials (e.g., mineral dust and sea salt), it is difficult to detect inclusions in the TEM images. Thus, the actual number fraction of the inclusions may be larger than that reported in this study. On the other hand, this technique has the advantage of identifying small inclusions (e.g., < 50 nm), which are difficult to detect using other methods. Although we observed several metal particles as inclusions (e.g., Zn and Pb), we focused on the occurrences of soot, fly-ash, and Fe-aggregate particles in this study.

3.7.1 Soot inclusions

Soot particles absorb solar radiation and exert a positive radiative forcing (Bond et al., 2013). When they are coated by or embedded within other species (e.g., sulfate and organic matter), their light absorption is enhanced by the focusing of light on the soot core (Bond et al., 2006; Oshima et al., 2009). Studies involving Arctic BC observations have shown a 54% increase in BC light absorption due to its coating (Zanatta et al., 2018) or an approximately 20% increase in the direct radiative effect due to the different BC mixing state assumptions (Kodros et al., 2018). In addition, although soot particles are initially hydrophobic, such coatings commonly make them hydrophilic. The degrees of the soot mixing states with hygroscopic materials influence their atmospheric lifetime in models, resulting in uncertainty in the simulated atmospheric BC concentrations in the Arctic (Eckhardt et al., 2015). Thus, the mixing states of soot particles are of interest regarding their climate influences (Fierce et al., 2020).

In our samples, soot particles were found in ~17% of all measured particles (Fig. S9 and Table S1). Most soot particles were internally mixed with other particles (Fig. 13 and Fig. S8). The number fraction of the externally mixed soot particles, which are those without any apparent coatings, was only ~1% of all soot particles. The relatively low number fractions of the externally mixed soot particles are similar to the background conditions observed by Hara et al. (2003) and indicate that most LRT soot particles were internally mixed in the Arctic.

Soot particles were observed within all aerosol types, and the sampling altitude was not clearly correlated with the soot particle fraction ($R^2=0.13$). On the other hand, more soot particles were found



365 in large host particles than in small ones, i.e., the size distributions of the soot-bearing particles (host particles) were larger than those of all particles (Fig. S10). This result is consistent with those in other areas (Adachi and Buseck, 2008; Adachi et al., 2014). A possible explanation is that the coagulation process of soot particles with large host particles occurs more efficiently than that with small particles.

BB is one of the major sources of BC in the Arctic atmosphere and snow (Spackman et al., 2010; Hegg et al., 2010). In our BB samples, ~93% of the soot-bearing particles contained K, showing substantial BC contributions from BB. In comparison, the ratio was 74% for the non-BB samples. However, there was no apparent enhancement of the soot particle number fraction in the BB samples (Table S1). This inconsistency may be explained by the fact that we measured the number fraction and that the other species, such as the K-bearing salts, sulfates, and organic matters, also increased within the BB plume, 375 resulting in relatively lower soot number fractions in the BB samples.

3.7.2 Fly-ash and Fe-aggregate inclusions

The fly-ash particles and primary particles of Fe aggregates commonly exhibited spherical shapes, which indicated that they formed through rapid cooling after melting or evaporation at high temperatures during emission. The shape and composition of the Fe aggregates are consistent with those 380 observed near sources in East Asia (Moteki et al., 2017). Moteki et al. showed that Fe-aggregate particles consist of magnetite, which absorbs light and imposes warming effects on the climate. Yoshida et al. (2020) also demonstrated light-absorbing Fe-oxide particles using SP2 during this campaign. Thus, we suggest that our Fe-aggregate particles also exhibit light-absorbing properties.

The number fractions of the particles containing fly-ash or Fe-aggregate particles are 1.4% and 0.5%, 385 respectively (Fig. S9 and Table S1). Although these particles are commonly found in polluted areas from anthropogenic sources such as stationary combustion sources and vehicles (Li et al. 2016), there are almost no such local anthropogenic sources in the Arctic area. Instead, the fly-ash and Fe-aggregate particles attain better relations with the soot number fraction ($R^2=0.35$ and 0.24 for the fly-ash and Fe aggregates, respectively; Fig. S11), and soot, fly-ash, and Fe-aggregate particles are often found in the 390 same particles (Fig. 13 and Fig. S8). Our TEM measurements agree with the SP2 observations of Yoshida et al. (2020), who also found correlations among BC and Fe oxide during this campaign. These results indicate that the soot, fly-ash, and Fe-aggregate particles originated from anthropogenic sources through LRT.

3.8 Implications for particle aging and the climate

395 In the Arctic, although there are few anthropogenic sources and vegetation, we found that many aerosol particles originated from anthropogenic and BB sources. The soot and Fe-aggregate particles have light-absorbing properties, and the mineral-dust particles demonstrate water-insoluble, crystalline structures



with hygroscopic attachments. These optical and structural properties influence their direct radiative forcing by scattering and absorbing solar radiation and their indirect radiative forcing through cloud
400 modification by serving as CCN and INPs. When these light-absorbing particles are deposited on snow or ice surfaces, they reduce the surface albedo, resulting in snow/ice melting acceleration (Law and Stohl, 2007). Soot particles (or BC) have commonly been detected at various concentrations in the snow (Mori et al., 2019; Kinease et al., 2019) and ice cores (McConnell et al., 2007) in polar regions. Their deposition process from the atmosphere onto snow/ice is important to better understand their
405 contributions to snowmelt and albedo changes.

Soot, tarballs, and Fe-aggregate particles have been intensively investigated near their sources, such as urban areas and BB smoke. This study shows the composition and shape of aged particles (> 1 week). The soot and Fe-aggregate particles still exhibit fractal structures, especially the small particles (< several tens of primary particles; Fig. 13 and Fig. S8), similar to those found in source regions such as
410 East Asia (Adachi et al., 2016). This observation is inconsistent with the result showing highly compacted soot particles at a remote marine free-troposphere site (China et al., 2015). Whether soot particles become core-shell structures as they age in the atmosphere is of interest for the accurate evaluation of soot climate effects (Cappa et al., 2012; Adachi et al., 2010). Our observation implies that the soot particles in the Arctic cannot be assumed to be spherical particles but should be treated as
415 fractal particles when calculating the optical properties of aged samples. One explanation is that soot particles consisting of several tens or fewer primary particles hardly attain highly compact shapes because of the insufficient monomer numbers to turn them into spherical aggregates. On the other hand, most soot particles were coated with sulfate or other materials, suggesting that they possess enhanced light absorption and CCN activity. We also observed the aging of tarball particles in a BB plume, i.e.,
420 the surfaces of certain tarball particles had reacted with sulfate, making them hygroscopic. Aged tarballs are removed more efficiently from the atmosphere by precipitation than the original ones. As the mass fraction of tarballs in fresh BB smoke may reach ~40% (Sedlaeck et al., 2018), which is much higher than that in our samples, most tarballs could have been removed from the atmosphere during LRT. This tarball removal process has been proposed by Pósfai et al. (2004) when they first characterized tarballs,
425 and our study provides evidence of the above hypothesized processes.

4. Summary

This study reveals that the aerosol particles in the Arctic troposphere exhibit various composition, shape, and mixing state ranges depending on the sampling altitude and air mass history. Sulfate is the dominant aerosol type, and sea-salt, mineral-dust, K-bearing, and carbonaceous particles are also
430 observed as major aerosol species. The aerosol particles are commonly mixtures of several components, resulting in different optical and hygroscopic properties than the original particles. In addition to the



main components, they also include soot, fly-ash, and Fe-aggregate particles, all of which are solid, primary particles originating from anthropogenic sources. The aerosol particles that include soot and Fe-aggregate particles may exhibit light-absorbing properties and contribute to Arctic warming and
435 snow/ice melting when deposited on the surface. BB is an important source of the aerosol particles in the Arctic area. In the BB samples, we find K-bearing and tarball particles, both of which are tracers of BB emissions. Our model simulations indicated the BB contributions from Siberia. Tarballs could have aged for a week or more after their emission and have reacted with sulfate. Many mineral-dust particles are mixed with sea salt and are relatively abundant in the samples collected near the surface. The
440 mineral-dust occurrence contradicts previous studies that show that they mainly originate from LRT in spring, and further studies are required to confirm their sources. The ability of solid particles, such as mineral-dust, soot, fly-ash, and Fe-aggregate particles, to function as INPs should be considered to evaluate their contributions to cloud formation. This study highlights a wide range of mixing states, and the mixing states of aerosol particles after LRT should be accounted for to accurately evaluate their
445 climate influences.

Data and code availability

The TEM and simulation data used in this publication are available upon request (adachik@mri-jma.go.jp). Access to the MRI-ESM2 code is available under a collaboration framework with the MRI.

Author contributions

450 KA conducted the TEM analysis and data processing. SO, AY, and MK executed the TEM sampling and field observations. KA and NM set up the TEM sampler. MK supervised the TEM sampling. NO performed the model simulations and analyses. KA prepared the manuscript with contributions from all coauthors.

Competing interests

455 The authors declare that they have no conflicts of interest.

Acknowledgments

We are indebted to all the PAMARCMiP 2018 participants for their cooperation and support. The authors also acknowledge the Alfred Wegener Institute (AWI) for both the support to conduct the PAMARCMiP 2018 campaign and the use of the Polar5 research aircraft and the skill and safety
460 exemplified by the pilots and flight staff. We thank the financial support of the Environment Research and Technology Development Fund (JPMEERF20205001, JPMEERF20202003, JPMEERF20172003,



and JPMEERF20165005) of the Environmental Restoration and Conservation Agency of Japan, the Global Environmental Research Coordination System of the Ministry of the Environment of Japan, and the ArCS (JPMXD130000000) and ArCS II (JPMXD1420318865) projects of the Ministry of
465 Education, Culture, Sports, Science, and Technology (MEXT) of Japan. KA and NO acknowledge the financial support provided by the Japan Society for the Promotion of Science (JSPS) KAKENHI (grant numbers JP26701004, JP16K16188, JP16H01772, JP18H04134, JP18H03363, JP18H05292, JP19H01972, JP19H04236, JP19K21905, and JP19H04259).



470 **References**

- Abbatt, J. P. D., Leaitch, W. R., Aliabadi, A. A., Bertram, A. K., Blanchet, J.-P., Boivin-Rioux, A.,
Bozem, H., Burkart, J., Chang, R. Y. W., Charette, J., Chaubey, J. P., Christensen, R. J.,
Cirisan, A., Collins, D. B., Croft, B., Dionne, J., Evans, G. J., Fletcher, C. G., Galí, M.,
Ghahremaninezhad, R., Girard, E., Gong, W., Gosselin, M., Gourdal, M., Hanna, S. J.,
475 Hayashida, H., Herber, A. B., Hesaraki, S., Hoor, P., Huang, L., Husserr, R., Irish, V. E.,
Keita, S. A., Kodros, J. K., Köllner, F., Kolonjari, F., Kunkel, D., Ladino, L. A., Law, K.,
Levasseur, M., Libois, Q., Liggio, J., Lizotte, M., Macdonald, K. M., Mahmood, R., Martin, R.
V., Mason, R. H., Miller, L. A., Moravek, A., Mortenson, E., Mungall, E. L., Murphy, J. G.,
Namazi, M., Norman, A.-L., amp, apos, Neill, N. T., Pierce, J. R., Russell, L. M., Schneider, J.,
480 Schulz, H., Sharma, S., Si, M., Staebler, R. M., Steiner, N. S., Thomas, J. L., von Salzen, K.,
Wentzell, J. J. B., Willis, M. D., Wentworth, G. R., Xu, J.-W., and Yakobi-Hancock, J. D.:
Overview paper: New insights into aerosol and climate in the Arctic, *Atmospheric Chemistry
and Physics*, 19, 2527-2560, 10.5194/acp-19-2527-2019, 2019.
- Adachi, K. and Buseck, P. R.: Internally mixed soot, sulfates, and organic matter in aerosol particles
485 from Mexico City, *Atmospheric Chemistry and Physics*, 8, 6469–6481,
<https://doi.org/10.5194/acp-8-6469-2008>, 2008.
- Adachi, K., and Buseck, P. R.: Atmospheric tar balls from biomass burning in Mexico, *Journal of
Geophysical Research*, 116, 10.1029/2010jd015102, 2011.
- Adachi, K., and Buseck, P. R.: Changes in shape and composition of sea-salt particles upon aging in an
490 urban atmosphere, *Atmospheric Environment*, 100, 1-9, 10.1016/j.atmosenv.2014.10.036,
2015.
- Adachi, K., Chung, S. H., and Buseck, P. R.: Shapes of soot aerosol particles and implications for their
effects on climate, *Journal of Geophysical Research*, 115, 10.1029/2009jd012868, 2010.
- Adachi, K., Oshima, N., Gong, Z., de Sá, S., Bateman, A. P., Martin, S. T., de Brito, J. F., Artaxo, P.,
495 Cirino, G. G., Sedlacek III, A. J., and Buseck, P. R.: Mixing states of Amazon-basin aerosol
particles transported over long distances using transmission electron microscopy, *Atmospheric
Chemistry and Physics*, <https://doi.org/10.5194/acp-2020-452>, 2020. (accepted)
- Adachi, K., Zaizen, Y., Kajino, M., and Igarashi, Y.: Mixing state of regionally transported soot
particles and the coating effect on their size and shape at a mountain site in Japan, *Journal of
500 Geophysical Research: Atmospheres*, 119, 5386-5396, 10.1002/2013jd020880, 2014.
- Adachi, K., Moteki, N., Kondo, Y., and Igarashi, Y.: Mixing states of light-absorbing particles
measured using a transmission electron microscope and a single-particle soot photometer in



- Tokyo, Japan, *Journal of Geophysical Research: Atmospheres*, 121, 9153-9164, 10.1002/2016jd025153, 2016.
- 505 Adachi, K., Sedlacek, A. J., III, Kleinman, L., Springston, S. R., Wang, J., Chand, D., Hubbe, J. M., Shilling, J. E., Onasch, T. B., Kinase, T., Sakata, K., Takahashi, Y., and Buseck, P. R.: Spherical tarball particles form through rapid chemical and physical changes of organic matter in biomass-burning smoke, *Proc. Natl. Acad. Sci. USA*, 116, 19336-19341, 10.1073/pnas.1900129116, 2019.
- 510 Anderson, J. R., Buseck, P. R., Saucy, D. A., Pacyna, J. M.: Characterization of individual fine-fraction particles from the Arctic aerosol at Spitsbergen, May–June 1987, *Atmospheric Environment*, 26, 1747-1762, 1992.
- Arnold, S. R., Law, K. S., Brock, C. A., Thomas, J. L., Starkweather, S. M., von Salzen, K., Stohl, A., Sharma, S., Lund, M. T., Flanner, M. G., Petäjä, T., Tanimoto, H., Gamble, J., Dibb, J. E.,
515 Melamed, M., Johnson, N., Fidel, M., Tynkkynen, V. P., Baklanov, A., Eckhardt, S., Monks, S. A., Browse, J., and Bozem, H.: Arctic air pollution: Challenges and opportunities for the next decade, *Elementa: Science of the Anthropocene*, 4, 000104, 10.12952/journal.elementa.000104, 2016.
- Bond, T. C., Habib, G., and Bergstrom, R. W.: Limitations in the enhancement of visible light
520 absorption due to mixing state, *Journal of Geophysical Research*, 111, 10.1029/2006jd007315, 2006.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K.,
525 Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *Journal of Geophysical Research: Atmospheres*, 118, 5380-5552, 10.1002/jgrd.50171, 2013.
- Brock, C. A., Cozic, J., Bahreini, R., Froyd, K. D., Middlebrook, A. M., McComiskey, A., Brioude, J.,
530 Cooper, O. R., Stohl, A., Aikin, K. C., de Gouw, J. A., Fahey, D. W., Ferrare, R. A., Gao, R. S., Gore, W., Holloway, J. S., Hübler, G., Jefferson, A., Lack, D. A., Lance, S., Moore, R. H., Murphy, D. M., Nenes, A., Novelli, P. C., Nowak, J. B., Ogren, J. A., Peischl, J., Pierce, R. B., Pilewskie, P., Quinn, P. K., Ryerson, T. B., Schmidt, K. S., Schwarz, J. P., Sodemann, H., Spackman, J. R., Stark, H., Thomson, D. S., Thornberry, T., Veres, P., Watts, L. A., Warneke,
535 C., and Wollny, A. G.: Characteristics, sources, and transport of aerosols measured in spring 2008 during the aerosol, radiation, and cloud processes affecting Arctic Climate (ARCPAC)



- Project, Atmospheric Chemistry and Physics, 11, 2423-2453, 10.5194/acp-11-2423-2011, 2011.
- 540 Buseck, P. R., Adachi, K., Gelencsér, A., Tompa, É., and Pósfai, M.: Ns-Soot: A Material-Based Term for Strongly Light-Absorbing Carbonaceous Particles, *Aerosol Science and Technology*, 48, 777-788, 10.1080/02786826.2014.919374, 2014.
- 545 Cappa, C. D., Onasch, T. B., Massoli, P., Worsnop, D. R., Bates, T. S., Cross, E. S., Davidovits, P., Hakala, J., Hayden, K. L., Jobson, B. T., Kolesar, K. R., Lack, D. A., Lerner, B. M., Li, S. M., Mellon, D., Nuaaman, I., Olfert, J. S., Petaja, T., Quinn, P. K., Song, C., Subramanian, R., Williams, E. J., and Zaveri, R. A.: Radiative absorption enhancements due to the mixing state of atmospheric black carbon, *Science*, 337, 1078-1081, 10.1126/science.1223447, 2012.
- 550 Chakrabarty, R. K., Moosmüller, H., Chen, L.-W. A., Lewis, K., Arnott, W. P., Mazzoleni, C., Dubey, M. K., Wold, C. E., Hao, W. M., and Kreidenweis, S. M.: Brown carbon in tar balls from smoldering biomass combustion, *Atmospheric Chemistry and Physics*, 10, 6363–6370, <https://doi.org/10.5194/acp-10-6363-2010>, 2010.
- Chi, J. W., Li, W. J., Zhang, D. Z., Zhang, J. C., Lin, Y. T., Shen, X. J., Sun, J. Y., Chen, J. M., Zhang, X. Y., Zhang, Y. M., and Wang, W. X.: Sea salt aerosols as a reactive surface for inorganic and organic acidic gases in the Arctic troposphere, *Atmospheric Chemistry and Physics*, 15, 11341-11353, 10.5194/acp-15-11341-2015, 2015.
- 555 China, S., Scarnato, B., Owen, R. C., Zhang, B., Ampadu, M. T., Kumar, S., Dzepina, K., Dziobak, M. P., Fialho, P., Perlinger, J. A., Hueber, J., Helmig, D., Mazzoleni, L. R., and Mazzoleni, C.: Morphology and mixing state of aged soot particles at a remote marine free troposphere site: Implications for optical properties, *Geophysical Research Letters*, 42, 1243-1250, 10.1002/2014gl062404, 2015.
- 560 Eckhardt, S., Quennehen, B., Olivié, D. J. L., Berntsen, T. K., Cherian, R., Christensen, J. H., Collins, W., Crepinsek, S., Daskalakis, N., Flanner, M., Herber, A., Heyes, C., Hodnebrog, Ø., Huang, L., Kanakidou, M., Klimont, Z., Langner, J., Law, K. S., Lund, M. T., Mahmood, R., Massling, A., Myriokefalitakis, S., Nielsen, I. E., Nøjgaard, J. K., Quaas, J., Quinn, P. K., Raut, J. C., Rumbold, S. T., Schulz, M., Sharma, S., Skeie, R. B., Skov, H., Uttal, T., von Salzen, K., and
565 Stohl, A.: Current model capabilities for simulating black carbon and sulfate concentrations in the Arctic atmosphere: a multi-model evaluation using a comprehensive measurement data set, *Atmospheric Chemistry and Physics*, 15, 9413-9433, 10.5194/acp-15-9413-2015, 2015.
- Fan, S.-M.: Modeling of observed mineral dust aerosols in the arctic and the impact on winter season low-level clouds, *Journal of Geophysical Research: Atmospheres*, 118, 11,161-111,174,
570 10.1002/jgrd.50842, 2013.



- Fenger, M., Sørensen, L. L., Kristensen, K., Jensen, B., Nguyen, Q. T., Nøjgaard, J. K., Massling, A., Skov, H., Becker, T., and Glasius, M.: Sources of anions in aerosols in northeast Greenland during late winter, *Atmospheric Chemistry and Physics*, 13, 1569-1578, 10.5194/acp-13-1569-2013, 2013.
- 575 Fierce, L., Onasch, T. B., Cappa, C. D., Mazzoleni, C., China, S., Bhandari, J., Davidovits, P., Fischer, D. A., Helgestad, T., Lambe, A. T., Sedlacek, A. J., III, Smith, G. D., and Wolff, L.: Radiative absorption enhancements by black carbon controlled by particle-to-particle heterogeneity in composition, *Proc. Natl. Acad. Sci. USA*, 117, 5196-5203, 10.1073/pnas.1919723117, 2020.
- Fisher, J. A., Jacob, D. J., Wang, Q., Bahreini, R., Carouge, C. C., Cubison, M. J., Dibb, J. E., Diehl, T.,
580 Jimenez, J. L., Leibensperger, E. M., Lu, Z., Meinders, M. B. J., Pye, H. O. T., Quinn, P. K., Sharma, S., Streets, D. G., van Donkelaar, A., and Yantosca, R. M.: Sources, distribution, and acidity of sulfate–ammonium aerosol in the Arctic in winter–spring, *Atmospheric Environment*, 45, 7301-7318, 10.1016/j.atmosenv.2011.08.030, 2011.
- Gard, E. E., Kleeman, M. J., Gross, D. S., Hughes, L. S., Allen, J. O., Morrical, B. D., Fergenson, D. P.,
585 Dienes, T., E. Gälli, M., Johnson, R. J., Cass, G. R., and Prather, K. A.: Direct Observation of Heterogeneous Chemistry in the Atmosphere, *Science*, 279, 1184-1187, 10.1126/science.279.5354.1184, 1998.
- Geng, H., Ryu, J. Y., Jung, H.-J., Chung, H., Ahn, K.-H., and Ro, C.-U.: Single-particle characterization of summertime arctic aerosols collected at Ny-Alesund, Svalbard, *Environmental Science and Technology*, 44, 2348–2353, <https://doi.org/10.1021/es903268j>, 2010.
590
- Hansen, J., and Nazarenko, L.: Soot climate forcing via snow and ice albedos, *Proc. Natl. Acad. Sci. USA*, 101, 423-428, 10.1073/pnas.2237157100, 2004.
- Hara, K.: Mixing states of individual aerosol particles in spring Arctic troposphere during ASTAR 2000 campaign, *Journal of Geophysical Research*, 108, 10.1029/2002jd002513, 2003.
- 595 Hara, K., Osada, K., Nishita, C., Yamagata, S., Yamanocuchi, T., Herber, A., Matsunaga, K., Iwasaka, Y., Nagatani, M., and Nakata, H.: Vertical variations of sea-salt modification in the boundary layer of spring Arctic during the ASTAR 2000 campaign, *Tellus B: Chemical and Physical Meteorology*, 54, 361-376, 10.3402/tellusb.v54i4.16671, 2002.
- Hara, K., Matoba, S., Hirabayashi, M., and Yamasaki, T.: Frost flowers and sea-salt aerosols over
600 seasonal sea-ice areas in northwestern Greenland during winter–spring, *Atmospheric Chemistry and Physics*, 17, 8577-8598, 10.5194/acp-17-8577-2017, 2017.
- Hartmann, M., Adachi, K., Eppers, O., Haas, C., Herber, A., Holzinger, R., Hünerbein, A., Jäkel, E., Jentsch, C., Pinxteren, M., Wex, H., Willmes, S., and Stratmann, F.: Wintertime Airborne



- 605 Measurements of Ice Nucleating Particles in the High Arctic: A Hint to a Marine, Biogenic
Source for Ice Nucleating Particles, *Geophysical Research Letters*, 47, 10.1029/2020gl087770,
2020.
- Hecobian, A., Liu, Z., Hennigan, C. J., Huey, L. G., Jimenez, J. L., Cubison, M. J., Vay, S., Diskin, G.
S., Sachse, G. W., Wisthaler, A., Mikoviny, T., Weinheimer, A. J., Liao, J., Knapp, D. J.,
Wennberg, P. O., Kürten, A., Crouse, J. D., Clair, J. S., Wang, Y., and Weber, R. J.:
610 Comparison of chemical characteristics of 495 biomass burning plumes intercepted by the
NASA DC-8 aircraft during the ARCTAS/CARB-2008 field campaign, *Atmospheric
Chemistry and Physics*, 11, 13325-13337, 10.5194/acp-11-13325-2011, 2011.
- Hegg, D. A., Warren, S. G., Grenfell, T. C., and Clarke, A. D.: Sources of light-absorbing aerosol in
arctic snow and their seasonal variation, *Atmospheric Chemistry and Physics*, 10, 10923-
615 10938, 10.5194/acp-10-10923-2010, 2010.
- Heidam, N. Z., Wählin, P., and Christensen, J. H.: Tropospheric Gases and Aerosols in Northeast
Greenland, *Journal of the Atmospheric Sciences*, 56, 261-278, 1999.
- Huang, J., and Jaeglé, L.: Wintertime enhancements of sea salt aerosol in polar regions consistent with a
sea ice source from blowing snow, *Atmospheric Chemistry and Physics*, 17, 3699-3712,
620 10.5194/acp-17-3699-2017, 2017.
- Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Emmons, L. K., Ferrare, R. A.,
Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E.,
Pederson, J. R., and Fisher, J. A.: The Arctic Research of the Composition of the Troposphere
from Aircraft and Satellites (ARCTAS) mission: design, execution, and first results,
625 *Atmospheric Chemistry and Physics*, 10, 5191-5212, 10.5194/acp-10-5191-2010, 2010.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J.,
Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions
estimated with a global fire assimilation system based on observed fire radiative power,
Biogeosciences, 9, 527-554, 10.5194/bg-9-527-2012, 2012.
- 630 Kinase, T., Adachi, K., Oshima, N., Goto-Azuma, K., Ogawa-Tsukagawa, Y., Kondo, Y., Moteki, N.,
Ohata, S., Mori, T., Hayashi, M., Hara, K., Kawashima, H., and Kita, K.: Concentrations and
Size Distributions of Black Carbon in the Surface Snow of Eastern Antarctica in 2011, *Journal
of Geophysical Research: Atmospheres*, 125, 10.1029/2019jd030737, 2020.
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H.,
635 Kobayashi, C., Endo, H., Miyaoka, K., and Takahashi, K.: The JRA-55 Reanalysis: General



- Specifications and Basic Characteristics, *Journal of the Meteorological Society of Japan*, 93, 5–48, <https://doi.org/10.2151/jmsj.2015-001>, 2015.
- Kodros, J. K., Hanna, S. J., Bertram, A. K., Leaitch, W. R., Schulz, H., Herber, A. B., Zanatta, M., Burkart, J., Willis, M. D., Abbatt, J. P. D., and Pierce, J. R.: Size-resolved mixing state of
640 black carbon in the Canadian high Arctic and implications for simulated direct radiative effect, *Atmospheric Chemistry and Physics*, 18, 11345–11361, [10.5194/acp-18-11345-2018](https://doi.org/10.5194/acp-18-11345-2018), 2018.
- Kojima, T. and Buseck, P. R.: Aerosol particles from tropical convective systems: 2. Cloud bases, *Journal of Geophysical Research*, 110, [10.1029/2004jd005173](https://doi.org/10.1029/2004jd005173), 2005.
- Krämer, M., Cziczo, D. J., Burkert-Kohn, M., Boose, Y., Wex, H., Ladino, L. A., and Kanji, Z. A.:
645 Overview of Ice Nucleating Particles, *Meteorological Monographs*, 58, 1.1–1.33, [10.1175/amsmonographs-d-16-0006.1](https://doi.org/10.1175/amsmonographs-d-16-0006.1), 2017.
- Kurusu, M., Adachi, K., Sakata, K., and Takahashi, Y.: Stable Isotope Ratios of Combustion Iron Produced by Evaporation in a Steel Plant, *ACS Earth and Space Chemistry*, 3, 588–598, [10.1021/acsearthspacechem.8b00171](https://doi.org/10.1021/acsearthspacechem.8b00171), 2019.
- 650 Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmospheric Chemistry and Physics*, 10, 7017–7039, [10.5194/acp-10-7017-2010](https://doi.org/10.5194/acp-10-7017-2010), 2010.
- 655 Latham, T. L., Beyersdorf, A. J., Thornhill, K. L., Winstead, E. L., Cubison, M. J., Hecobian, A., Jimenez, J. L., Weber, R. J., Anderson, B. E., and Nenes, A.: Analysis of CCN activity of Arctic aerosol and Canadian biomass burning during summer 2008, *Atmospheric Chemistry and Physics*, 13, 2735–2756, [10.5194/acp-13-2735-2013](https://doi.org/10.5194/acp-13-2735-2013), 2013.
- 660 Law, K. S., Stohl, A., Quinn, P. K., Brock, C. A., Burkhart, J. F., Paris, J.-D., Ancellet, G., Singh, H. B., Roiger, A., Schlager, H., Dibb, J., Jacob, D. J., Arnold, S. R., Pelon, J., and Thomas, J. L.: Arctic Air Pollution: New Insights from POLARCAT-IPY, *Bulletin of the American Meteorological Society*, 95, 1873–1895, [10.1175/bams-d-13-00017.1](https://doi.org/10.1175/bams-d-13-00017.1), 2014.
- Law, K. S., and Stohl, A.: Arctic Air Pollution: Origins and Impacts, *Science*, 315, 1537–1540, [10.1126/science.1137695](https://doi.org/10.1126/science.1137695), 2007.
- 665 Lewis, E.R., Schwartz, S.E., 2004. Sea Salt Aerosol Production: Mechanisms, Methods, Measurement and Model e A Critical Review. In: *Geophys. Monogr.*, vol. 152. AGU, Washington, D. C, p. 413.



- Li, J., Pósfai, M., Hobbs, P. V., and Buseck, P. R.: Individual aerosol particles from biomass burning in
670 southern Africa: 2, Compositions and aging of inorganic particles, *Journal of Geophysical
Research: Atmospheres*, 108, 10.1029/2002jd002310, 2003.
- Li, W., Shao, L., Zhang, D., Ro, C.-U., Hu, M., Bi, X., Geng, H., Matsuki, A., Niu, H., and Chen, J.: A
review of single aerosol particle studies in the atmosphere of East Asia: morphology, mixing
675 state, source, and heterogeneous reactions, *Journal of Cleaner Production*, 112, 1330-1349,
10.1016/j.jclepro.2015.04.050, 2016.
- Li, W., Xu, L., Liu, X., Zhang, J., Lin, Y., Yao, X., Gao, H., Zhang, D., Chen, J., Wang, W., Harrison,
R. M., Zhang, X., Shao, L., Fu, P., Nenes, A., and Shi, Z.: Air pollution–aerosol interactions
produce more bioavailable iron for ocean ecosystems, *Science Advances*, 3, e1601749,
10.1126/sciadv.1601749, 2017.
- 680 Matsui, H., Kondo, Y., Moteki, N., Takegawa, N., Sahu, L. K., Koike, M., Zhao, Y., Fuelberg, H. E.,
Sessions, W. R., Diskin, G., Anderson, B. E., Blake, D. R., Wisthaler, A., Cubison, M. J., and
Jimenez, J. L.: Accumulation-mode aerosol number concentrations in the Arctic during the
ARCTAS aircraft campaign: Long-range transport of polluted and clean air from the Asian
continent, *Journal of Geophysical Research*, 116, 10.1029/2011jd016189, 2011.
- 685 Matsui, H., and Moteki, N.: High Sensitivity of Arctic Black Carbon Radiative Effects to Subgrid
Vertical Velocity in Aerosol Activation, *Geophysical Research Letters*, 47,
10.1029/2020gl088978, 2020.
- McConnell, J. R., Edwards, R., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J. R.,
Pasteris, D. R., Carter, M. M., and Kahl, J. D. W.: 20th-Century Industrial Black Carbon
690 Emissions Altered Arctic Climate Forcing, *Science*, 317, 1381-1384,
10.1126/science.1144856, 2007.
- McNaughton, C. S., Clarke, A. D., Freitag, S., Kapustin, V. N., Kondo, Y., Moteki, N., Sahu, L.,
Takegawa, N., Schwarz, J. P., Spackman, J. R., Watts, L., Diskin, G., Podolske, J., Holloway,
J. S., Wisthaler, A., Mikoviny, T., de Gouw, J., Warneke, C., Jimenez, J., Cubison, M., Howell,
695 S. G., Middlebrook, A., Bahreini, R., Anderson, B. E., Winstead, E., Thornhill, K. L., Lack, D.,
Cozic, J., and Brock, C. A.: Absorbing aerosol in the troposphere of the Western Arctic during
the 2008 ARCTAS/ARCPAC airborne field campaigns, *Atmospheric Chemistry and Physics*,
11, 7561-7582, 10.5194/acp-11-7561-2011, 2011.
- Mori, T., Goto-Azuma, K., Kondo, Y., Ogawa-Tsukagawa, Y., Miura, K., Hirabayashi, M., Oshima, N.,
700 Koike, M., Kupiainen, K., Moteki, N., Ohata, S., Sinha, P. R., Sugiura, K., Aoki, T.,
Schneebeil, M., Steffen, K., Sato, A., Tsushima, A., Makarov, V., Omiya, S., Sugimoto, A.,
Takano, S., and Nagatsuka, N.: Black Carbon and Inorganic Aerosols in Arctic Snowpack,



- Journal of Geophysical Research: Atmospheres, 124, 13325-13356, 10.1029/2019jd030623, 2019.
- 705 Moroni, B., Cappelletti, D., Crocchianti, S., Becagli, S., Caiazzo, L., Traversi, R., Udisti, R., Mazzola, M., Markowicz, K., Ritter, C., and Zielinski, T.: Morphochemical characteristics and mixing state of long range transported wildfire particles at Ny-Ålesund (Svalbard Islands), *Atmospheric Environment*, 156, 135-145, 10.1016/j.atmosenv.2017.02.037, 2017.
- Moroni, B., Ritter, C., Crocchianti, S., Markowicz, K., Mazzola, M., Becagli, S., Traversi, R., Krejci, R., Tunved, P., and Cappelletti, D.: Individual Particle Characteristics, Optical Properties and Evolution of an Extreme Long-Range Transported Biomass Burning Event in the European Arctic (Ny-Ålesund, Svalbard Islands), *Journal of Geophysical Research: Atmospheres*, 125, 10.1029/2019jd031535, 2020.
- 710
- Moteki, N., Adachi, K., Ohata, S., Yoshida, A., Harigaya, T., Koike, M., and Kondo, Y.: Anthropogenic iron oxide aerosols enhance atmospheric heating, *Nat. Commun.*, 8, 15329, 10.1038/ncomms15329, 2017.
- 715
- Moteki, N., Mori, T., Matsui, H., and Ohata, S.: Observational constraint of in-cloud supersaturation for simulations of aerosol rainout in atmospheric models, *npj Climate and Atmospheric Science*, 2, 10.1038/s41612-019-0063-y, 2019.
- 720
- Murphy, D. M., Cziczo, D. J., Froyd, K. D., Hudson, P. K., Matthew, B. M., Middlebrook, A. M., Peltier, R. E., Sullivan, A., Thomson, D. S., and Weber, R. J.: Single-particle mass spectrometry of tropospheric aerosol particles, *Journal of Geophysical Research: Atmospheres*, 111, 10.1029/2006jd007340, 2006.
- Nguyen, Q. T., Skov, H., Sørensen, L. L., Jensen, B. J., Grube, A. G., Massling, A., Glasius, M., and Nøjgaard, J. K.: Source apportionment of particles at Station Nord, North East Greenland during 2008–2010 using COPREM and PMF analysis, *Atmospheric Chemistry and Physics*, 13, 35-49, 10.5194/acp-13-35-2013, 2013.
- 725
- Ohata, S., Yoshida, A., Moteki, N., Adachi, K., Takahashi, Y., Kurisu, M., and Koike, M.: Abundance of Light-Absorbing Anthropogenic Iron Oxide Aerosols in the Urban Atmosphere and Their Emission Sources, *Journal of Geophysical Research: Atmospheres*, 10.1029/2018jd028363, 2018.
- 730
- Oshima, N., Koike, M., Zhang, Y., and Kondo Y.: Aging of black carbon in outflow from anthropogenic sources using a mixing state resolved model: 2. Aerosol optical properties and cloud condensation nuclei activities, *Journal of Geophysical Research*, 114, D18202, 10.1029/2008JD011681, 2009.
- 735



- Oshima, N., Yukimoto, S., Deushi, M., Koshiro, T., Kawai, H., Tanaka, T. Y., and Yoshida, K.: Global and Arctic effective radiative forcing of anthropogenic gases and aerosols in MRI-ESM2.0, *Progress in Earth and Planetary Science*, 7, 38, <https://doi.org/10.1186/s40645-020-00348-w>, 2020.
- 740 Pósfai, M., Gelencsér, A., Simonics, R., Arató, K., Li, J., Hobbs, P. V., and Buseck, P. R.: Atmospheric tar balls: Particles from biomass and biofuel burning, *Journal of Geophysical Research: Atmospheres*, 109, 10.1029/2003jd004169, 2004.
- Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmospheric Chemistry and*
745 *Physics*, 5, 799-825, 10.5194/acp-5-799-2005, 2005.
- Schulz, H., Zanatta, M., Bozem, H., Leaitch, W. R., Herber, A. B., Burkart, J., Willis, M. D., Kunkel, D., Hoor, P. M., Abbatt, J. P. D., and Gerdes, R.: High Arctic aircraft measurements characterising black carbon vertical variability in spring and summer, *Atmospheric Chemistry and Physics*, 19, 2361-2384, 10.5194/acp-19-2361-2019, 2019.
- 750 Sedlacek III, A. J., Buseck, P. R., Adachi, K., Onasch, T. B., Springston, S. R., and Kleinman, L.: Formation and evolution of tar balls from northwestern US wildfires, *Atmospheric Chemistry and Physics*, 18, 11289–11301, <https://doi.org/10.5194/acp-18-11289-2018>, 2018.
- Shaw, G. E.: The Arctic Haze Phenomenon, *Bulletin of the American Meteorological Society*, 76, 2403-2414, 1995.
- 755 Shaw, P. M., Russell, L. M., Jefferson, A., and Quinn, P. K.: Arctic organic aerosol measurements show particles from mixed combustion in spring haze and from frost flowers in winter, *Geophysical Research Letters*, 37, 10.1029/2010gl042831, 2010.
- Spackman, J. R., Gao, R. S., Neff, W. D., Schwarz, J. P., Watts, L. A., Fahey, D. W., Holloway, J. S., Ryerson, T. B., Peischl, J., and Brock, C. A.: Aircraft observations of enhancement and
760 depletion of black carbon mass in the springtime Arctic, *Atmospheric Chemistry and Physics*, 10, 9667-9680, 10.5194/acp-10-9667-2010, 2010.
- Stone, R. S., Sharma, S., Herber, A., Eleftheriadis, K., and Nelson, D. W.: A characterization of Arctic aerosols on the basis of aerosol optical depth and black carbon measurements, *Elementa: Science of the Anthropocene*, 2, 000027, 10.12952/journal.elementa.000027, 2014.
- 765 Stroeve, J. C., Serreze, M. C., Holland, M. M., Kay, J. E., Malanik, J., and Barrett, A. P.: The Arctic's rapidly shrinking sea ice cover: a research synthesis, *Climatic Change*, 110, 1005-1027, 10.1007/s10584-011-0101-1, 2011.



- Tanaka, T. Y. and Chiba, M.: Global simulation of dust aerosol with a chemical transport model, MASINGAR. *Journal of the Meteorological Society of Japan*, 83A, 255–278, 2005.
- 770 Tobo, Y., Adachi, K., DeMott, P. J., Hill, T. C. J., Hamilton, D. S., Mahowald, N. M., Nagatsuka, N., Ohata, S., Uetake, J., Kondo, Y., and Koike, M.: Glacially sourced dust as a potentially significant source of ice nucleating particles, *Nature Geoscience*, 12, 253-258, 10.1038/s41561-019-0314-x, 2019.
- Warneke, C., Bahreini, R., Brioude, J., Brock, C. A., de Gouw, J. A., Fahey, D. W., Froyd, K. D.,
775 Holloway, J. S., Middlebrook, A., Miller, L., Montzka, S., Murphy, D. M., Peischl, J., Ryerson, T. B., Schwarz, J. P., Spackman, J. R., and Veres, P.: Biomass burning in Siberia and Kazakhstan as an important source for haze over the Alaskan Arctic in April 2008, *Geophysical Research Letters*, 36, 10.1029/2008gl036194, 2009.
- Wendisch, M., Macke, A., Ehrlich, A., Lüpkes, C., Mech, M., Chechin, D., Dethloff, K., Velasco, C. B.,
780 Bozem, H., Brückner, M., Clemen, H.-C., Crewell, S., Donth, T., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Engler, C., Eppers, O., Gehrman, M., Gong, X., Gottschalk, M., Gourbeyre, C., Griesche, H., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Hoor, P., Jafariserajehlou, S., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U., Kecorius, S., Knudsen, E. M., Köllner, F., Kretzschmar, J., Lelli, L., Leroy, D., Maturilli, M.,
785 Mei, L., Mertes, S., Mioche, G., Neuber, R., Nicolaus, M., Nomokonova, T., Notholt, J., Palm, M., van Pinxteren, M., Quaas, J., Richter, P., Ruiz-Donoso, E., Schäfer, M., Schmieder, K., Schnaiter, M., Schneider, J., Schwarzenböck, A., Seifert, P., Shupe, M. D., Siebert, H., Spreen, G., Stapf, J., Stratmann, F., Vogl, T., Welti, A., Wex, H., Wiedensohler, A., Zanatta, M., and Zeppenfeld, S.: The Arctic Cloud Puzzle: Using ACLOUD/PASCAL Multiplatform
790 Observations to Unravel the Role of Clouds and Aerosol Particles in Arctic Amplification, *Bulletin of the American Meteorological Society*, 100, 841-871, 10.1175/bams-d-18-0072.1, 2019.
- Wesche, C., Steinhage, D., and Nixdorf, U.: Polar aircraft Polar5 and Polar6 operated by the Alfred Wegener Institute, *Journal of large-scale research facilities JLSRF*, 2, 10.17815/jlsrf-2-153,
795 2016.
- Willis, M. D., Köllner, F., Burkart, J., Bozem, H., Thomas, J. L., Schneider, J., Aliabadi, A. A., Hoor, P. M., Schulz, H., Herber, A. B., Leaitch, W. R., and Abbatt, J. P. D.: Evidence for marine biogenic influence on summertime Arctic aerosol, *Geophysical Research Letters*, 44, 6460-6470, 10.1002/2017gl073359, 2017.
- 800 Willis, M. D., Bozem, H., Kunkel, D., Lee, A. K. Y., Schulz, H., Burkart, J., Aliabadi, A. A., Herber, A. B., Leaitch, W. R., and Abbatt, J. P. D.: Aircraft-based measurements of High Arctic



- springtime aerosol show evidence for vertically varying sources, transport and composition, *Atmospheric Chemistry and Physics*, 19, 57-76, 10.5194/acp-19-57-2019, 2019.
- 805 Xu, L., Russell, L. M., and Burrows, S. M.: Potential sea salt aerosol sources from frost flowers in the pan-Arctic region, *Journal of Geophysical Research: Atmospheres*, 121, 10,840-810,856, 10.1002/2015jd024713, 2016.
- Yamanouchi, T., Treffeisen, R., Herber, A., Shiobara, M., Yamagata, S., Hara, K., Sato, K., Yabuki, M., Tomikawa, Y., Rinke, A., Neuber, R., Schumacher, R., Kriews, M., Ström, J., Schrems, O., and Gernandt, H.: Arctic Study of Tropospheric Aerosol and Radiation (ASTAR) 2000: Arctic
810 haze case study, *Tellus B: Chemical and Physical Meteorology*, 57, 141-152, 10.3402/tellusb.v57i2.16784, 2017.
- Yoshida, A., Ohata, S., Moteki, N., Adachi, K., Mori, T., Koike, M., and Takami, A.: Abundance and Emission Flux of the Anthropogenic Iron Oxide Aerosols From the East Asian Continental Outflow, *Journal of Geophysical Research: Atmospheres*, 123, 11,194-111,209,
815 10.1029/2018jd028665, 2018.
- Yoshida, A., Moteki, N., Ohata, S., Mori, T., Koike, M., Kondo, Y., Matsui, H., Oshima, N., Takami, A., and Kita, K.: Abundances and microphysical properties of light-absorbing iron oxide and black carbon aerosols over East Asia and the Arctic, *Journal of Geophysical Research: Atmospheres*, 10.1029/2019jd032301, 2020.
- 820 Yoshizue, M., Iwamoto, Y., Adachi, K., Kato, S., Sun, S., Miura, K., and Uematsu, M.: Individual particle analysis of marine aerosols collected during the North–South transect cruise in the Pacific Ocean and its marginal seas, *Journal of Oceanography*, 75, 513-524, 10.1007/s10872-019-00519-4, 2019.
- Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M.,
825 Tanaka, T., Hosaka, M., Yabu, S., Yoshimura, H., Shindo, E., Mizuta, R., Obata, A., Adachi, Y., and Ishii, M.: The Meteorological Research Institute Earth System Model Version 2.0, MRI-ESM2.0: Description and Basic Evaluation of the Physical Component, *Journal of the Meteorological Society of Japan. Ser. II*, 97, 931-965, 10.2151/jmsj.2019-051, 2019.
- Yumimoto, K., Tanaka, T. Y., Oshima, N., and Maki, T.: JRAero: the Japanese Reanalysis for Aerosol
830 v1.0, *Geoscientific Model Development*, 10, 3225-3253, 10.5194/gmd-10-3225-2017, 2017.
- Zanatta, M., Laj, P., Gysel, M., Baltensperger, U., Vratolis, S., Eleftheriadis, K., Kondo, Y., Dubuisson, P., Winiarek, V., Kazadzis, S., Tunved, P., and Jacobi, H.-W.: Effects of mixing state on optical and radiative properties of black carbon in the European Arctic, *Atmospheric Chemistry and Physics*, 18, 14037-14057, 10.5194/acp-18-14037-2018, 2018.



835 **Figures**




Figure 1. Flight tracks and locations of the TEM samplings during the PAMARCMiP 2018. The blue star indicates the Villum Research Station in Greenland. The yellow points indicate the sampling locations below 1000 m. The red points indicate those above 1000 m. The sampling points are the midpoints of the sampling time; 19 min for all samples except during the 3 and 4 April flights, which had a sampling time of 29 min.






Figure 2. Size distributions of each particle type. The particle sizes were determined from the area-equivalent diameters obtained from the STEM images. The size bins are shown on a log scale and are < 0.25, 0.25-0.35, 0.35-0.5, 0.5-0.71, 0.71-1.00, 1.00-1.41, and >1.41 μm. The particle numbers used for the mineral-dust, sea-salt, K-bearing, sulfate, and carbonaceous particles are 7844, 337, 1205, 421, 4672, and 809, respectively.



850 **Figure 3.** Size-dependent number fractions in the samples (a) < 1000 m and (b) > 1000 m and number fractions of the total particles of the samples (c) < 1000 m and (d) > 1000 m. The size bins in (a) and (b) are shown on a log-scale and are <0.35, 0.35-0.5, 0.5-0.71, 0.71-1.00, 1.00-1.41, 1.41-2.00, and > 2.00 μm . The particle numbers used in (c) and (d) are 4400 and 3444, respectively.



855

Figure 4. Vertical changes in the number fractions (%) of (a) mineral-dust, (b) sea-salt, and (c) K-bearing particles. Each plot indicates the number fraction of a given sample. The number of particles in the samples ranges from 216 to 442. The red solid-circle points in the K-bearing particle plots indicate the samples originating from the biomass burning plume (BB samples).

860




Figure 5. Number fractions of the aerosol particles in the BB samples. (a) Size-dependent and (b) total number fractions. Three BB samples were collected on 2, 3, and 4 April. The size bins are shown on a log-scale and are 0.35-0.5, 0.5-0.71, 0.71-1.00, 1.00-1.41, 1.41-2.00, and > 2.00 μm. (b) Number fractions of the total particles of the BB samples. The particle number of the BB samples is 1041.






Figure 6. TEM and elemental mapping images of a BB sample (10:40-10:59, 2 April 2018). (a) TEM
image of K-bearing particles. (b) The same area of the TEM image (a) but after the STEM-EDS
870 analysis. Beam-sensitive materials are removed, and soot inclusions are more apparent. (c) and (d):
Examples of K-bearing particles containing soot inclusions and organic coatings, which appear in gray
in the TEM image. (e) STEM and elemental mapping images for C, N, O, S, and K in the TEM image.
The distributions of K, S, and O correspond to the K-bearing particles, and that of C represents the soot
and organic particles.



875

Figure 7. Model simulations of the (a) horizontal and (b) vertical distributions of the mass concentration of BC originating from biomass burning (BB) during the sampling periods of the BB samples. The white circles indicate the sampling locations. The transport pathway of the BB plume from its source area is shown in Fig. S2.



880

Figure 8. TEM and elemental mapping images of the sea-salt particles in a sample (10:00-10:19, 30 March 2018). Top images (a)-(c): TEM images of the sea-salt particles mixed with various grains. Bottom images (d)-(f): Elemental mapping images of the TEM areas for C, O, Na, Mg, S, Cl, K, and Ca. Soot particles are attached to the sea-salt particles (b) and (c). C-O-Mg in the particles (a) and (b) occurs around NaCl grains.

885




Figure 9. Relations of Na with Mg, Cl, and Ca among all particles. The plots show the average weight percent values of all particles within each sample. The black and red plots are the samples < 1000 m and > 1000 m, respectively. The error bars indicate the 95% confidential intervals.




Figure 10. TEM and elemental mapping images of the mineral-dust particles. Top (a)-(c): TEM images of the mineral-dust particles. Bottom (d)-(f): Elemental mapping images of the TEM areas for the detected elements. All particles were found in the samples collected from 10:00-10:19 on 30 March 895 2018. Soot, NaCl, and Ca-S particles are attached to these mineral-dust particles in (a). Fe occurs as grains in (b). Sulfate is attached to the mineral-dust particle in (c).






Figure 11. TEM and elemental mapping images of the sulfate particles. Top (a)-(c): TEM images. Bottom (d)-(f): Elemental mapping images of the TEM areas for C, N, Na, and S. All particles were
900 collected from 14:40-14:59, 31 March 2018. Most particles mainly consist of sulfate. Some particles
also include N and Na, suggesting that they are mixtures of ammonium sulfate and sodium sulfate. The
carbon grains correspond to the internally mixed soot particles. Small droplets occur around relatively
large sulfate particles, forming a satellite structure, as indicated in (a).



905 **Figure 12.** TEM and elemental mapping images of the tarball and tarball-like particles in a BB sample (10:30-10:59, 2 April 2018). (a) TEM image of the tarball-like particles containing sulfate. As they exhibit deformed spherical shapes and the characteristic elements of tarballs (e.g., C, N, O, and K), we label them tarball-like particles. Other examples of tarballs are shown in (b) and (c). (d)-(f): Elemental mapping images of the TEM areas.



910

Figure 13. TEM and elemental mapping images of the sulfate particles with inclusions. (a) TEM image before the mapping analysis and (b) after the mapping analysis. Electron-beam exposure removed sulfate, and inclusions became more apparent. (c) Elemental mapping images of the TEM areas for the detected elements. (d) A fly-ash particle. (e) Mineral-dust and fly-ash particles. (f) An Fe-aggregate particle. (g) Mineral-dust and soot particles. The white squares in (b) and (c) indicate the areas of (d)-
915 (g). The sample was collected from 14:40-14:59, 31 March 2018.