

Measurement report: Regional trends of stratospheric ozone evaluated using the Merged GRidded Dataset of Ozone Profiles (MEGRIDOP)

5 Viktoria F. Sofieva¹, Monika Szlag¹, Johanna Tamminen¹, Erkki Kyrölä¹, Doug Degenstein², Chris Roth², Daniel Zawada², Alexei Rozanov³, Carlo Arosio³, John P. Burrows³, Mark Weber³, Alexandra Laeng⁴, Gabriele Stiller⁴, Thomas von Clarmann⁴, Lucien Froidevaux⁵, Nathaniel Livesey⁵, Michel van Roozendael⁶, Christian Retscher⁷

10 ¹ Finnish Meteorological Institute, Helsinki, Finland

² Institute of Space and Atmospheric Studies, University of Saskatchewan, Saskatoon, Canada

³ Institute of Environmental Physics, University of Bremen, Bremen, Germany

⁴ Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research, Karlsruhe, Germany

⁵ Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA

15 ⁶ Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

⁷ ESA/ESRIN, Frascati, Italy

Correspondence to: Viktoria F. Sofieva (viktoria.sofieva@fmi.fi)

Abstract

20 In this paper, we present the Merged GRidded Dataset of Ozone Profiles (MEGRIDOP) in the stratosphere with a resolved longitudinal structure, which is derived from data by six limb and occultation satellite instruments: GOMOS, SCIAMACHY and MIPAS on Envisat, OSIRIS on Odin, OMPS on Suomi-NPP, and MLS on Aura. The merged dataset was generated as a contribution to the European Space Agency Climate Change Initiative Ozone project (Ozone_cci). The period of this merged time series of ozone profiles is from late 2001 until the end of 2018.

25 The monthly mean gridded ozone profile dataset is provided in the altitude range from 10 to 50 km in bins of 10° latitude x 20° longitude. The merging is performed using deseasonalized anomalies. The created MEGRIDOP dataset can be used for analyses which probe our understanding of stratospheric chemistry and dynamics. To illustrate some possible applications, we created a climatology of ozone profiles with resolved longitudinal structure. We found zonal asymmetry/structures in the

30 climatological ozone profiles at middle and high latitudes associated with the polar vortex. At Northern high latitudes, the amplitude of the seasonal cycle also has a longitudinal dependence.

The MEGRIDOP dataset has been also used to evaluate regional vertically-resolved ozone trends in the stratosphere, including polar regions. It is found that stratospheric ozone trends exhibit longitudinal structures at Northern Hemisphere middle and high latitudes, with enhanced trends over Scandinavia and the Atlantic region. This agrees well with previous analyses and might be due to changes in dynamical processes related to the Brewer-Dobson circulation.

35 **1 Introduction**

Nowadays, the importance of protecting the ozone layer and monitoring its recovery from the effect of ozone depleting substances is well recognized (e.g., Petropavlovskikh et al., 2019; WMO, 2014, 2018). Past analyses have demonstrated that ozone is recovering in the upper stratosphere (e.g., Arosio et al., 2019; Bourassa et al., 2014; Kyrölä et al., 2013; Petropavlovskikh et al., 2019; Sofieva et al., 2017; Steinbrecht et al., 2017; WMO, 2018). The ozone recovery in the lower
40 stratosphere is not yet observed, and the lower stratospheric ozone trends are the subject of recent controversial discussions (Ball et al., 2018, 2019; Chipperfield et al., 2018).

In the majority of studies of ozone profile trends using satellite observations made in limb-viewing geometry, analyses are performed on zonal mean data. This representation allows ozone trends to be estimated globally. At the same time, such representation provides a sufficiently large amount of experimental data in spatio-temporal bins (usually 10° latitude and one
45 month) to enable robust estimation of trends. This is especially important for the period before 2001, when long data records are available only from solar occultation instruments having relatively sparse data coverage.

A recent study by Arosio et al. (2019) using the merged SCIAMACHY-OMPS dataset has shown that ozone trends for the period 2003-2018 have a significant dependence on longitude. Also, total ozone column trends (WMO, 2018 and references therein) have a pronounced zonal structure.

50 This paper is focused on a new longitudinally resolved merged dataset of ozone profiles in the stratosphere based on several limb and occultation instruments. This new merged dataset is a contribution to the European Space Agency Climate Change Initiative ozone project (Ozone_cci). It can be used in different applications, including the evaluation of regional ozone trends in the stratosphere.

The paper is organized as follows. In Section 2, we briefly discuss the satellite data used for creating the merged dataset.
55 Section 3 is dedicated to the methodological aspects of data merging. Examples of ozone distributions are shown in Section 4. Section 5 is dedicated to regional trend analysis. A discussion and summary (Section 6) conclude the paper.

2 Data

The MEGRIDOP dataset is a merged and gridded dataset generated using ozone profiles retrieved from several limb and occultation instruments, viz. MIPAS (Michelson Interferometer for Passive Atmospheric Sounding), SCIAMACHY (SCanning Imaging Spectrometer for Atmospheric CHartographY) and GOMOS (Global Ozone Monitoring by occultation of Stars), all on Envisat, OSIRIS (Optical Spectrograph and InfraRed Imaging System) on Odin, OMPS-LP (Ozone Mapping and Profiles Suite - Limb Profiler) on Suomi-NPP, and MLS (Microwave Limb Sounder) on Aura.

These instruments provide high-quality ozone profiles with a good vertical resolution of 2-4 km and a relatively dense spatio-temporal coverage (100-3500 ozone profiles per day with fairly uniform sampling in longitude). The important information about the datasets is collected in Table 1. More information about the datasets from the individual satellite instruments is found in Petropavlovskikh et al., (2019), Sofieva et al., (2017), and references therein.

Table 1. General information about the datasets.

Instrument/ satellite	Level 2 processor, references	Years	Vertical range/retrieval coordinate	Local time of Level 2 data	Number of profiles per day
MIPAS/Envisat	KIT/IAA V7R_O3_240 von Clarmann et al. (2003; 2009)	2005-2012	6-70 km, Altitude	10 a.m. and p.m.	~1000
SCIAMACHY/Envisat	UBr v3.5 (Jia et al., 2015)	2002-2012	8-65 km, Altitude	10 a.m.	~1300
GOMOS/Envisat	ALGOM2s v1 (Kyrölä et al., 2010; Sofieva et al., 2016)	2002-2011	10-105 km, altitude	10 p.m.	~110
OSIRIS/Odin	USask v5.10 (Bourassa et al., 2017; Degenstein et al., 2009)	2001-present	10-59 km, altitude	6 a.m. and p.m.	~250
OMLS-LP /SUOMI-NPP	USask 2D v 1.1.0 (Zawada et al., 2017)	2012-present	6- 59 km, altitude	1:30 p.m.	~1600
MLS/Aura	NASA v4.2 (Livesey et al., 2013)	2004 - present	261-0.02 hPa (~8-75 km), pressure	1:30 a.m. and p.m.	~3000

For all instruments except MLS, the original ozone profile retrievals are performed on an altitude grid. GOMOS, OSIRIS, SCIAMACHY and OMPS-LP - provide number density ozone profiles; therefore this representation (number density on an altitude grid) is used for the merged dataset. For MIPAS, the retrievals are performed in volume mixing ratio vs. altitude grid. The conversion to number density profiles is performed using temperature profiles retrieved by MIPAS and the pressure profiles provided with the MIPAS ozone data ; the latter are constructed from altitude and temperature using one (z,p,T) data point from the ERA-Interim reanalysis (<https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era-interim>; Dee et al., 2011).

For MLS, retrievals are performed in mixing ratio on a pressure grid. Similarly to the conversion procedure of MIPAS data, we performed the conversion to number density using the retrieved MLS temperatures, but for altitude-pressure conversion, we used the ERA-Interim reanalysis data. Such conversion might introduce some uncertainty in the MLS data. For studies of long-term changes, this uncertainty is associated with a potentially imperfect representation of temperature trends in ERA-Interim, which might influence ozone trends. However, since current stratospheric temperature trends (after 2000) are small (Maycock et al., 2018; Steiner et al., 2020), this uncertainty is expected to be small. The MLS ozone profiles data record is stable (Hubert et al., 2016), therefore including MLS data into the merged dataset is advantageous, especially for the merging method applied in our work (see also below).

For all the instruments, we use the ozone profiles from the updated HARMONized dataset of Ozone profiles (HARMOZ_ALT) developed in the ESA Ozone_cci project (Sofieva et al., 2013), <https://climate.esa.int/en/projects/ozone/>. HARMOZ consists of the original retrieved ozone profiles from each instrument, which are screened for invalid data by the instrument experts and are presented on a vertical grid (altitude-gridded profiles are used in our paper) and in a common netCDF4 format. Detailed information about the original datasets can be found in (Sofieva et al., 2013), and references to the corresponding publications are also collected in Table 1 of our paper.

3 Merging method

The method used for creating the MEGRIDOP dataset is similar to that used for the creation of the merged SAGE-CCI-OMPS dataset (Sofieva et al., 2017). Below we describe and illustrate the merging process.

3.1 Gridded monthly means from individual instruments

First, gridded ozone profile data $\rho_i(z, b, t)$ in each $10^\circ \times 20^\circ$ latitude-longitude bin b and at altitude z were created for each individual dataset i and each month t . The mean number density profile in each spatio-temporal bin is $\rho_i(z, b, t)$. For each instrument, we required more than 10 measurements in each spatio-temporal bin. The uncertainty of the averaged data $\sigma_i(z, b, t)$ is approximated by the standard error of the mean (see discussion in Toohey and von Clarmann (2013) on possible influence of correlations caused by orbital sampling on the standard error of the mean).

The non-uniformity of the sampling pattern can be characterized by the inhomogeneity measure, which is defined as the linear combination of two classical inhomogeneity measures, asymmetry A and entropy E : $H = \frac{1}{2}(A + (1 - E))$ (Sofieva et al., 2014). The unitless inhomogeneity measure H ranges from 0 to 1 (the more homogeneous, the smaller H is). For our application, we considered the inhomogeneity in time (H_{time}) as the main contribution to sampling uncertainty.

Examples of gridded datasets at 30 km altitude for individual satellite instruments are shown in Figures 1 and 2. All instruments show a similar morphology, although biases between individual datasets exist. The coverage is instrument-specific

and to some extent time-dependent; the most complete coverage is achieved by MIPAS and MLS. The spatial bins are covered rather uniformly by the data. Examples of the inhomogeneity measure H_{time} are presented in the Supplement, Figure S1. H_{time} is very close to zero for the instruments with dense sampling (MIPAS, SCIAMACHY, MLS, OMPS). For OSIRIS and GOMOS, H is usually below 0.1 (good homogeneity of the data) with a few exceptions for some months and locations. In this work, the inhomogeneity measure H_{time} is used for detection of spatial bins with high inhomogeneity of data (see below).

110





Figure 1. Examples of gridded monthly mean ozone number density (cm^{-3}) at 30 km for individual satellite instruments in January 2008.



115

Figure 2. Examples of gridded monthly mean ozone number density (cm^{-3}) at 30 km for individual satellite instruments in January 2018.

3.2 Seasonal cycle and deseasonalized anomalies

120 For each instrument i , latitude-longitude bin b , and altitude level z , deseasonalized anomalies are computed as:

$$\Delta_i(z, b, t) = \frac{\rho_i(z, b, t) - \rho_{m,i}(z, b)}{\rho_{m,i}(z, b)}, \quad (1)$$

where $\rho_i(z, b, t)$ is the monthly mean in this spatial bin and $\rho_{m,i}(z, b)$ is the climatological mean value for the month m . In other words, from each January we removed mean January values, from each February – the mean February value, and so on.

125 In our computations, we removed values for spatial bins with less than 10 profiles and inhomogeneity H_{time} larger than 0.9. For all instruments except for OMPS, the seasonal cycle is estimated using the years 2005-2011. For OMPS, the seasonal cycle is evaluated using data from 2012-2018. Figure 3 illustrates the seasonal cycle at 40 km for all instruments except GOMOS, as the GOMOS data do not cover all months for the considered spatial bins. Although biases are visible, the overall behavior of the seasonal cycle is similar for the different datasets. In the tropics (left panel), small differences in seasonal cycle between two longitude regions, 0-20° E and 120-140°E are observed, while at mid-latitudes, all satellite instruments show consistently different seasonal cycles in two different longitude regions.





Figure 3. Examples of seasonal cycles in the tropics (left) and NH upper stratosphere (right) at 40 km. Solid lines: longitudes 0-20°E, dashed lines: longitudes 120-140°E. In the tropics, a semi-annual cycle is observed.

135

For two instruments - MIPAS and MLS - which measure during day and during night, and thus provide data at all latitudes in all seasons, we compared the relative amplitude of the seasonal cycle $\frac{\max(\rho_m) - \min(\rho_m)}{\text{mean}(\rho_m)}$ at several altitude levels (Figure

4). As seen from Figure 4, longitudinal structures in the relative amplitude of the seasonal cycle are observed, to be largest in the Northern middle and high latitudes, particularly in the middle and upper stratosphere.



140 **Figure 4. Relative amplitude of seasonal cycle at 25 km (left), 35 km(center) and 45 km(right) for MIPAS (top) and MLS (bottom).**

The merging of individual datasets was performed on deseasonalized anomalies. The main advantage of using deseasonalized anomalies is that various biases between the individual datasets - e.g., instrumental-specific, or those due to the difference in
145 local time - are automatically removed. The deseasonalization also removes spatial sampling biases if the sampling patterns do not change over time. Details of the applied merging method are presented in the next section.

3.3 Merging the data

The merging method used for creating MERGRIDOP is similar to that used in creating the merged SAGE-CCI-OMPS dataset (Sofieva et al., 2017). The deseasonalized anomalies of all instruments except OMPS are aligned, as the seasonal cycle
150 was estimated using the same period. First, we offset the OMPS deseasonalized anomalies to the median of the deseasonalized anomalies from all other instruments. These additive offsets are computed using the data from years 2012-2018, and the offsetting procedure is illustrated in Figure 5. In this figure, we selected a spatial bin where the effect of the offsetting is clearly visible. In many other bins, the offsets are small or negligible. As observed in Figure 5 (and also below in Figure 6), the deseasonalized anomalies from individual datasets are in good agreement.

155




Figure 5. Illustration of offsetting the OMPS deseasonalized anomalies. The data are shown for altitude 35 km and the latitude/longitude bin 0-10° N/0-20° E.

160 After offsetting OMPS, the merged ozone profiles in each spatiotemporal bin and at each altitude level is obtained from the median of the deseasonalized anomalies corresponding to individual instruments:

$$\Delta_{merged}(z, b, t) = median(\Delta_i(z, b, t)) \quad (2)$$

The advantage of using the median estimate is that the merged anomaly follows the majority of the data, and it is not very sensitive to exclusion/addition of an individual data record, in cases where there are several (and consistent) anomaly datasets available. The sensitivity of the dataset and the evaluated trends to the number of instruments was studied in detail for SAGE-CCI-OMPS dataset, which is created with the same merging algorithm (Sofieva et al., 2017), and this is valid also for MEGRIDOP.

165 The uncertainties of the merged deseasonalized anomalies are computed similarly to those used for the merged SAGE-CCI-OMPS dataset (Sofieva et al., 2017). For each instrument, the uncertainty of the deseasonalized anomalies, σ_{Δ_i} , is estimated via Gaussian error propagation; it is given by

$$\sigma_{\Delta_i} = \Delta_i \sqrt{\frac{\sigma_i^2}{\rho_i^2} + \frac{\sigma_{m,i}^2}{\rho_{m,i}^2}} \quad (3)$$

where σ_i is the uncertainty of the gridded ozone profiles (see Sect. 3.1.) and $\sigma_{m,i}$ is the uncertainty of the seasonal cycle $\rho_{m,i}$, which can be estimated via propagation of random uncertainties to the mean value:

$$\sigma_{m,i}^2 = \frac{1}{N_m^2} \sum_{j=1}^{N_m} \sigma_i^2(z, b, t_j) \quad (4)$$


175 Analogously to (Sofieva et al., 2017), the uncertainties of the merged deseasonalized anomalies are estimated as:

$$\sigma_{\Delta,merged} = \min \left(\sigma_{\Delta,i_{med}}, \sqrt{\frac{1}{N} \sum_{i=1}^N \sigma_{\Delta,i}^2 + \frac{1}{N^2} \sum_{i=1}^N (\Delta_i - \Delta_{merged})^2} \right), \quad (5)$$

180 where $\sigma_{\Delta,i_{med}}$ is the anomaly uncertainty of the instrument corresponding to the median value. In cases where there are even number of measurements, the mean of two neighbors to the median is used. Analogously to uncertainty estimates in the merged SAGE-CCI-OMPS dataset (Sofieva et al., 2017), the uncertainties given by Eq. (5) can be interpreted as follows. If individual anomalies are significantly different, the uncertainty of the merged anomaly is the uncertainty corresponding to the median value. In cases where several instruments report a similar anomaly (intersecting error bars), this provides more confidence in this anomaly value, and the resulting uncertainty of the merged anomaly is approximated by the second term in Eq. (5).

185 The deseasonalized anomalies from individual datasets are usually very close to each other, so that several values can be typically found within the uncertainty interval of the merged anomaly $\Delta_{merged} \pm \sigma_{\Delta,merged}$. This is similar to the approach taken with the SAGE_CCI-OMPS dataset (Sofieva et al. (2017), Figure S8).

Examples of deseasonalized anomalies and their estimated uncertainties are displayed in Figures 6 and 7, respectively.



190 **Figure 6 . An example of deseasonalized anomalies (in %) for individual instruments and the merged dataset in the spatial bin 0-10° N, 0-20° E.**




Figure 7. An example of uncertainties in deseasonalized anomalies (in %) for individual instruments and the merged dataset in the spatial bin 0-10° N, 0-20° E

195

The average estimated uncertainty of the merged ozone is usually less than 2% before 2012 and below 1% after 2012. In the UTLS, uncertainties are larger than in the stratosphere; they are typically in the range of 2-12 % before 2012 and 2-6 % after 2012.

4 The merged dataset and selected examples

200


The merged deseasonalized anomalies can be directly used for evaluation of ozone trends in the stratosphere. The evaluation of regional ozone trends is discussed in Section 5 of our paper. We also created a version of MEGRIDOP in number density through restoration of the seasonal cycle. This was achieved in a manner similar to that applied in creating the merged SAGE-CCI-OMPS dataset (Sofieva et al., 2017). The best estimates of the amplitude and morphology of the seasonal cycle are provided by MIPAS and MLS, as these two instruments provide global coverage in all seasons. The ozone profiles from OSIRIS and MLS have the smallest biases with respect to ozone soundings (Hubert et al., 2016). For the seasonal cycle of the merged dataset, we computed the mean of MIPAS and MLS seasonal cycles and offset it to the mean of OSIRIS and MLS values (this offset does not depend on season). By this procedure, the seasonal cycle in the merged dataset has absolute values,

205

which have the smallest biases with respect to the ground-based instruments, and a realistic amplitude. An example of a number density MERGRIDOP dataset is shown in Figure 8.


210

The merged dataset allows us to provide a gridded climatology of ozone profiles, i.e., the collection of ozone profiles categorized by calendar month, latitude, longitude, and altitude. Figure 9 shows these climatological ozone values, for four months and at four altitude levels. The polar projections of these distributions are presented in the Supplement (Figures S2 and S3). As observed in these figures, there is zonal asymmetry associated with the polar vortex, in both hemispheres. In other locations, the ozone distributions are rather uniform in longitude.



215

Figure 8. An example of number density ozone profiles (in cm^{-3}) for individual instruments and the merged dataset in the spatial bin 0-10° N, 0-20° E.



220 **Figure 9. Climatological ozone distributions (in DU/km), for January, April, July, and October, for selected altitude levels (15, 20, 30, and 40 km).**

5 Evaluation of regional ozone trends

For evaluation of the regional ozone trends, we exploited the standard approach of multiple linear regression and applied it to the deseasonalized anomalies:

$$\Delta_{merged}(t) = at + b + q_1 QBO_{30}(t) + q_2 QBO_{50}(t) + s F_{10.7}(t) + d ENSO(t), \quad (6),$$

225 where we model the trend with a simple linear term, $QBO_{30}(t)$ and $QBO_{50}(t)$ are the equatorial winds at 30 hPa and 50 hPa, respectively (<http://www.cpc.ncep.noaa.gov/data/indices/>), $F_{10.7}(t)$ is the monthly average solar 10.7 cm radio flux (ftp://ftp.geolab.nrcan.gc.ca/data/solar_flux/monthly_averages/), and $ENSO(t)$ is the 2 month lagged ENSO proxy (<http://www.esrl.noaa.gov/psd/enso/mei/table.html>). The evaluation of trends has been performed for each latitude-longitude bin and for each altitude level separately. Autocorrelations are removed using the Cochrane–Orcutt transformation (Cochrane and Orcutt, 1949).

230

In our analysis, we consider long-term trends over the years covered by MEGRIDOP, and approximate them by a linear function (which describes bulk changes). However, real changes in the atmosphere can be non-linear (Laine et al., 2014): if variations are analyzed on a shorter time scale, they can be different from long-term trends (e.g., Arosio et al., 2019; Chipperfield et al., 2018; Galytska et al., 2019; Strahan et al., 2020). We selected the years after 2003, in order to avoid the


235 influence of a major sudden stratospheric warming in September 2002 on ozone trends at Southern Hemisphere middle and high latitudes (see also a discussion below).

Ozone trends (expressed in percent per decade) estimated at several altitude levels for years 2003-2018 are shown in Figure 10. Figure 11 displays the trends at these altitudes in absolute units, DU/(km decade). In Figures 10 and 11, black stars indicate the statistically significant trends, i.e., trends different from zero at a 95% or greater confidence level. The morphology of ozone trends presented in absolute and in relative units looks similar. As shown in Figures 10 and 11, statistically significant trends are observed in the upper stratosphere. A longitudinal structure is clearly visible in the NH mid-latitude trends above 40 km: the trends are significantly larger over Scandinavia/Atlantic Ocean (5-6 % decade⁻¹) than over Siberia (~1 % decade⁻¹). The same feature was also observed by Arosio et al. (2019). Enhanced ozone trends over the mid-latitude Atlantic sector are seen in both absolute and relative units, and also at lower altitudes (but the ozone trends are not statistically significant below 40 km).

We also compared the trends in late 2004 – 2018, the common measurement period, using MEGRIDOP, only MLS data and the merged SCIAMACHY-OMPS dataset by Arosio et al. (2019). We found that the spatial distributions of ozone trends are similar for the considered datasets (Figure 12, top). The MEGRIDOP and pure MLS ozone trends in 2004-2018 are similar (as expected, MLS data are used in MEGRIDOP). SCIAMACHY-OMPS trends are somewhat larger, which might be related to the OMPS drift (Kramarova et al., 2018), but within error limits, and the morphology of ozone trends is similar. Specifically interesting is a two-core structure of ozone trends in the NH polar region, and this is seen nearly at all altitude levels (Figure 12, bottom) for all datasets.

There are several analyses showing that the residual circulation has a pronounced longitudinal two-core structure at Northern Hemisphere high and middle latitudes (e.g., Demirhan Bari et al., 2013; Kozubek et al., 2015). Kozubek et al. (2015) also performed a trend analysis and showed a weakening of the two-core structure, which possibly affects the ozone distribution in the region. Arosio et al. (2019) suggested that this longitudinal structure in the NH mid-latitude ozone trends is due to changes in dynamical processes related to the 3D structure of the Brewer Dobson circulation. However, the origin of the longitudinal structure of ozone trends requires more detailed investigations, including simulations with chemistry-transport models, in the future.

Statistically significant (at 95% confidence level) positive trends (1-2 % decade⁻¹) are observed also at SH mid-latitudes (~40°-50°S) at 25 km. This is in agreement with other studies of zonally averaged ozone trends (e.g., Arosio et al., 2019; Petropavlovskikh et al., 2019; Sofieva et al., 2017). In our analysis, there is a zonal asymmetry with larger trends in the sector 50°W - 10°E. At altitudes 20-25 km, the trends patterns are different in the Northern and Southern hemispheres.



265

Figure 10. Ozone trends ($\% \text{ decade}^{-1}$) in 2003-2018, for several altitudes. Statistically significant trends are indicated by stars.





Figure 11. Same as Figure 10, but trends in $\text{DU km}^{-1} \text{ decade}^{-1}$.



270 **Figure 12. Top: ozone trends in late 2004- 2018 (% decade⁻¹) at 35 km, bottom: longitude –altitude cross section of the ozone trends at ~ 65 °N (the latitude is indicated by dashed line on top panels). Ozone trends are estimated using the MEGRIDOP (left), MLS (center), and SCIAMACHY-OMPS datasets (right). For the SCIAMACHY-OMPS dataset, ozone trends in the Southern Atlantic Anomaly (SAA) region are not shown, because SCIAMACHY data are flagged in this region.**

275 Comparisons of MEGRIDOP ozone trends at 35 km in Figures 10 and 12 show larger positive ozone trends in the tropics in the period 2004 – 2018 compared to the period 2003 – 2018. A pronounced sensitivity of tropical ozone trends at ~35 km to the selection of the period for evaluation of ozone trends has been reported in several papers (e.g., Laine et al., 2014, Arosio et al., 2019, Galytska et al., 2019). As a hypothesis, this might be related to a decadal-scale ozone oscillation resulting from changes in Brewer-Dobson Circulation.

280 In previous studies (e.g., Petropavlovskikh et al., 2019; Steinbrecht et al., 2017; WMO, 2018), ozone trends have been evaluated at latitudes 60° S- 60° N, e.g., excluding polar regions. In this study, we have made an attempt to evaluate ozone trends also in polar regions. The ozone trends in polar projections are shown in the Supplement.

285 We found statistically significant positive trends in the NH polar middle stratosphere (25-30 km). In the SH polar regions, the estimated ozone trends are mostly positive, but they are not statistically significant. We found that the estimated trends in the SH polar regions are sensitive to the inclusion of 2002 data into the trend analysis. Quite exceptional (larger) ozone values in 2002 due to a SH major sudden stratospheric warming result in negative, although not statistically significant, ozone trends in the SH polar stratosphere, as expected since 2002 is at the beginning of the time period. If data from 2002 are excluded from the analysis, the estimated trends over Antarctica are not sensitive to the selection of the starting point for the trend analysis. This can be observed, for example, by comparison of ozone trends at 35 km in Figure 10 (trends for 2003-2018) and
 290 Figure 12 (trends for late 2004 to 2018).

Since natural variability is high in polar regions and the observational period is relatively short, it is quite expected that a simple multiple regression will lead to trend estimates that are not statistically significant. Other methods for trend analysis in polar regions, such as considering seasonal trends (Solomon et al., 2016; Szelađ et al., 2020; Galytska et al., 2019) can be explored in future works. In addition, the relation of winter-spring trends with respect to the position of the polar vortex would be an interesting subject in future studies.

The satellite data quality typically degrades in the UTLS compared to higher levels in the stratosphere. Our merging principle seems to be particularly optimal for the UTLS datasets, as it automatically removes biases, which can be significant in this altitude region. The very large natural ozone variability results in the MEGRIDOP the trend estimates below 20 km being not statistically significant in most locations.

6 Summary

In this paper, we presented the merged gridded dataset of ozone profiles (MEGRIDOP), which combines ozone data from six limb-viewing satellite instruments. The merged gridded ozone profiles are the monthly means in $10^{\circ} \times 20^{\circ}$ latitude-longitude bins, and they cover altitudes from 10 to 50 km. This dataset covers the years 2001-2018 and will be extended regularly in the future.

The merging was performed using aligned deseasonalized anomalies: the merged dataset represents the median of the deseasonalized anomalies from the individual instruments. The merged deseasonalized anomalies can be used directly for evaluation of ozone trends. For other applications, the MEGRIDOP is also available in the form of ozone number density profiles. The dataset is available through open access at <https://climate.esa.int/en/projects/ozone/data/> and at ftp://cci_web@ftp-ae.oma.be/esacci.

The MEGRIDOP dataset can be used in different analyses. As an illustration of one of the possible applications, a climatology of ozone profiles with resolved longitudinal structure has been created. We found zonal asymmetry/structures in the climatological ozone profiles at middle and high latitudes associated with the polar vortex. At Northern high latitudes, the amplitude of the seasonal cycle also has a longitudinal dependence.

We evaluated regional ozone trends over the years 2001-2018 using a multiple linear regression method. Overall, the estimated trends agree well with the trends derived from zonal mean ozone profiles. We found a zonal asymmetry in the upper stratospheric ozone trends at middle and high latitudes in the Northern Hemisphere: the trends are larger over Scandinavia/Atlantic Ocean than over Siberia. This feature agrees well with previous analyses and might be due to changes in dynamical processes related to the Brewer-Dobson circulation.

We estimated regional and vertically resolved ozone trends also in the polar regions. As far as we know, this is the first such analysis using limb satellite measurements. We found statistically significant positive trends in the NH polar middle

stratosphere (25-30 km). In the SH polar regions, the estimated ozone trends are mostly positive, but they are not statistically significant.

325 The MEGRIDOP dataset can be used in different analyses. In particular, it can be used for validation/intercomparison of climate data records from ground-based and other satellite measurements and chemistry-transport models in the future.

Acknowledgements

330 The work is performed in the framework of the ESA Ozone_cci project. The GOMOS ALGOM2s dataset was created in the framework of ESA ALGOM project. The KIT team would like to thank the European Space Agency (ESA) for giving access to MIPAS level-1 data. The SCIAMACHY ozone retrieval was funded in parts by ESA, the German Academic Exchange Service (DAAD) the German Aerospace Agency (DLR), and the University and State of Bremen. The data set was calculated with resources provided by the North-German Supercomputing Alliance (HLRN). The FMI team thanks the Academy of
335 Finland (Project SECTIC, and the Centre of Excellence of Inverse Modelling and Imaging). The authors thank the Canadian Space Agency. Work at the Jet Propulsion Laboratory, California Institute of Technology, was performed under contract with the National Aeronautics and Space Administration (NASA).

7 References

340 Arosio, C., Rozanov, A., Malinina, E., Weber, M. and Burrows, J. P.: Merging of ozone profiles from SCIAMACHY, OMPS and SAGE II observations to study stratospheric ozone changes, *Atmos. Meas. Tech.*, 12(4), 2423–2444, doi:10.5194/amt-12-2423-2019, 2019.

Ball, W. T., Alsing, J., Mortlock, D. J., Staehelin, J., Haigh, J. D., Peter, T., Tummon, F., Stübi, R., Stenke, A., Anderson, J., Bourassa, A., Davis, S. M., Degenstein, D., Frith, S., Froidevaux, L., Roth, C., Sofieva, V., Wang, R., Wild, J., Yu, P., Ziemke, J. R. and Rozanov, E. V.: Evidence for a continuous decline in lower stratospheric ozone offsetting ozone layer recovery, *Atmos. Chem. Phys.*, 18(2), 1379–1394, doi:10.5194/acp-18-1379-2018, 2018.

Ball, W. T., Alsing, J., Staehelin, J., Davis, S. M., Froidevaux, L. and Peter, T.: Stratospheric ozone trends for 1985-2018: sensitivity to recent large variability, *Atmos. Chem. Phys. Discuss.*, (March), 1–27, doi:10.5194/acp-2019-243, 2019.

Bourassa, A. E., Degenstein, D. A., Randel, W. J., Zawodny, J. M., Kyrölä, E., McLinden, C. A., Sioris, C. E. and Roth, C. Z.: Trends in stratospheric ozone derived from merged SAGE II and Odin-OSIRIS satellite observations, *Atmos. Chem. Phys.*, 14(13), 6983–6994, doi:10.5194/acp-14-6983-2014, 2014.

Bourassa, A. E., Roth, C. Z., Zawada, D. J., Rieger, L. A., McLinden, C. A. and Degenstein, D. A.: Drift corrected Odin-OSIRIS ozone product: algorithm and updated stratospheric ozone trends, *Atmos. Meas. Tech. Discuss.*, 2017, 1–16,

doi:10.5194/amt-2017-229, 2017.

355 Chipperfield, M. P., Dhomse, S., Hossaini, R., Feng, W., Santee, M. L., Weber, M., Burrows, J. P., Wild, J. D., Loyola, D. and Coldewey-Egbers, M.: On the Cause of Recent Variations in Lower Stratospheric Ozone, *Geophys. Res. Lett.*, 45(11), 5718–5726, doi:10.1029/2018GL078071, 2018.

von Clarmann, T., Glatthor, N., Grabowski, U., Höpfner, M., Kellmann, S., Kiefer, M., Linden, A., Tsidu, G. M., Milz, M., Steck, T., Stiller, G. P., Wang, D. Y., Fischer, H., Funke, B., Gil-López, S., López-Puertas, M., Mengistu Tsidu, G., Milz, M.,
360 Steck, T., Stiller, G. P., Wang, D. Y., Fischer, H., Funke, B., Gil-López, S. and López-Puertas, M.: Retrieval of temperature and tangent altitude pointing from limb emission spectra recorded from space by the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS), *J. Geophys. Res.*, 108(D23), 4736, doi:10.1029/2003JD003602, 2003.

von Clarmann, T., Höpfner, M., Kellmann, S., Linden, A., Chauhan, S., Funke, B., Grabowski, U., Glatthor, N., Kiefer, M., Schieferdecker, T., Stiller, G. P. and Versick, S.: Retrieval of temperature, H₂O, O₃, HNO₃, CH₄, N₂O, ClONO₂ and ClO from
365 MIPAS reduced resolution nominal mode limb emission measurements, *Atmos. Meas. Tech.*, 2(1), 159–175, doi:10.5194/amt-2-159-2009, 2009.

Cochrane, D. and Orcutt, G. H.: Application of Least Squares Regression to Relationships Containing Auto-Correlated Error Terms, *J. Am. Stat. Assoc.*, 44(245), 32–61, doi:10.1080/01621459.1949.10483290, 1949.

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G.,
370 Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597, doi:10.1002/qj.828, 2011.

375 Degenstein, D. A., Bourassa, A. E., Roth, C. Z. and Llewellyn, E. J.: Limb scatter ozone retrieval from 10 to 60 km using a multiplicative algebraic reconstruction technique, *Atmos. Chem. Phys.*, 9(17), 6521–6529, doi:10.5194/acp-9-6521-2009, 2009.

Demirhan Bari, D., Gabriel, A., Körnich, H. and Peters, D. W. H.: The effect of zonal asymmetries in the Brewer-Dobson circulation on ozone and water vapor distributions in the northern middle atmosphere, *J. Geophys. Res. Atmos.*, 118(9), 3447–
380 3466, doi:10.1029/2012JD017709, 2013.

Galytska, E., Rozanov, A., Chipperfield, M., Dhomse, S., Weber, M., Arosio, C., Wuhu, F. and Burrows, J.: Dynamically controlled ozone decline in the tropical mid-stratosphere observed by SCIAMACHY, *Atmos. Chem. Phys.*, 19, 767–783, doi:10.5194/acp-19-767-2019, 2019.

Hubert, D., Lambert, J.-C., Verhoelst, T., Granville, J., Keppens, A., Baray, J.-L., Bourassa, A. E., Cortesi, U., Degenstein, D.
385 A., Froidevaux, L., Godin-Beekmann, S., Hoppel, K. W., Johnson, B. J., Kyrölä, E., Leblanc, T., Lichtenberg, G., Marchand,

- M., McElroy, C. T., Murtagh, D., Nakane, H., Portafaix, T., Querel, R., Russell III, J. M., Salvador, J., Smit, H. G. J., Stebel, K., Steinbrecht, W., Strawbridge, K. B., Stübi, R., Swart, D. P. J., Taha, G., Tarasick, D. W., Thompson, A. M., Urban, J., van Gijssel, J. A. E., Van Malderen, R., von der Gathen, P., Walker, K. A., Wolfram, E. and Zawodny, J. M.: Ground-based assessment of the bias and long-term stability of 14 limb and occultation ozone profile data records, *Atmos. Meas. Tech.*, 9(6), 2497–2534, doi:10.5194/amt-9-2497-2016, 2016.
- 390
- Jia, J., Rozanov, A., Ladstätter-Weissenmayer, A. and P. Burrows, J.: Global validation of improved SCIAMACHY scientific ozone limb data using ozonesonde measurements, *Atmos. Meas. Tech. Discuss.*, 8(5), 4817–4858, doi:10.5194/amtd-8-4817-2015, 2015.
- Kozubek, M., Krizan, P. and Lastovicka, J.: Northern Hemisphere stratospheric winds in higher midlatitudes: longitudinal distribution and long-term trends, *Atmos. Chem. Phys.*, 15, 2203–2213, doi:https://doi.org/10.5194/acp-15-2203-2015, 2015.
- 395
- Kramarova, N. A., Bhartia, P. K., Jaross, G., Moy, L., Xu, P., Chen, Z., DeLand, M., Froidevaux, L., Livesey, N., Degenstein, D., Bourassa, A., Walker, K. A. and Sheese, P.: Validation of ozone profile retrievals derived from the OMPS LP version~2.5 algorithm against correlative satellite measurements, *Atmos. Meas. Tech.*, 11(5), 2837–2861, doi:10.5194/amt-11-2837-2018, 2018.
- 400
- Kyrölä, E., Tamminen, J., Sofieva, V. F., Bertaux, J.-L., Hauchecorne, A., Dalaudier, F., Fussen, D., Vanhellemont, F., Fanton D’Andon, O., Barrot, G., Guirlet, M., Mangin, A., Blanot, L., Fehr, T., de Miguel, L. and Fraisse, R.: Retrieval of atmospheric parameters from GOMOS data, *Atmos. Chem. Phys.*, 10(23), 11881–11903, doi:10.5194/acp-10-11881-2010, 2010.
- Kyrölä, E., Laine, M., Sofieva, V. F., Tamminen, J., Päivärinta, S.-M., Tukiainen, S., Zawodny, J. and Thomason, L.: Combined SAGE II–GOMOS ozone profile data set for 1984–2011 and trend analysis of the vertical distribution of ozone, *Atmos. Chem. Phys.*, 13(21), 10645–10658, doi:10.5194/acp-13-10645-2013, 2013.
- 405
- Laine, M., Latva-Pukkila, N. and Kyrölä, E.: Analysing time-varying trends in stratospheric ozone time series using the state space approach, *Atmos. Chem. Phys.*, 14(18), 9707–9725, doi:10.5194/acp-14-9707-2014, 2014.
- Livesey, N. J., Read, W. G., Froidevaux, L., Lambert, A., Manney, G. L., Pumphrey, H. C., Santee, M. L., Schwartz, M. J., Wang, S., Cofield, R. E., Cuddy, D. T., Fuller, R. A., Jarnot, R. F., Jiang, J. H., Knosp, B. W., Stek, P. C., Wagner, P. A., Wu, D. L. and L., D.: EOS MLS Version 3.3 and 3.4 Level 2 data quality and description document. [online] Available from: http://mls.jpl.nasa.gov/data/v3_data_quality_document.pdf, 2013.
- 410
- Maycock, A. C., Randel, W. J., Steiner, A. K., Karpechko, A. Y., Christy, J., Saunders, R., Thompson, D. W. J., Zou, C.-Z., Chrysanthou, A., Luke Abraham, N., Akiyoshi, H., Archibald, A. T., Butchart, N., Chipperfield, M., Dameris, M., Deushi, M., Dhomse, S., Di Genova, G., Jöckel, P., Kinnison, D. E., Kirner, O., Ladstädter, F., Michou, M., Morgenstern, O., O’Connor, F., Oman, L., Pitari, G., Plummer, D. A., Revell, L. E., Rozanov, E., Stenke, A., Visioni, D., Yamashita, Y. and Zeng, G.: Revisiting the Mystery of Recent Stratospheric Temperature Trends, *Geophys. Res. Lett.*, 45(18), 9919–9933, doi:10.1029/2018GL078035, 2018.
- 415

- Petropavlovskikh, I., Godin-Beekmann, S., Hubert, D., Damadeo, R., Hassler, B. and Sofieva, V.: SPARC/IO3C/GAW Report on Long-term Ozone Trends and Uncertainties in the Stratosphere, edited by M. Kenntner and B. Ziegele, [online] Available from: <https://elib.dlr.de/126666/>, 2019.
- Sofieva, V. F., Rahpoe, N., Tamminen, J., Kyrölä, E., Kalakoski, N., Weber, M., Rozanov, A., von Savigny, C., Laeng, A., von Clarmann, T., Stiller, G., Lossow, S., Degenstein, D., Bourassa, A., Adams, C., Roth, C., Lloyd, N., Bernath, P., Hargreaves, R. J., Urban, J., Murtagh, D., Hauchecorne, A., Dalaudier, F., van Roozendaal, M., Kalb, N. and Zehner, C.: Harmonized dataset of ozone profiles from satellite limb and occultation measurements, *Earth Syst. Sci. Data*, 5(2), 349–363, doi:10.5194/essd-5-349-2013, 2013.
- Sofieva, V. F., Kalakoski, N., Päivärinta, S.-M., Tamminen, J., Laine, M. and Froidevaux, L.: On sampling uncertainty of satellite ozone profile measurements, *Atmos. Meas. Tech.*, 7(6), 1891–1900, doi:10.5194/amt-7-1891-2014, 2014.
- Sofieva, V. F., Ialongo, I., Hakkarainen, J., Kyrölä, E., Tamminen, J., Laine, M., Hauchecorne, A., Dalaudier, F., Bertaux, J.-L., Fussen, D., Blanot, L., Barrot, G. and Dehn, A.: Improved GOMOS/Envisat ozone retrievals in the upper troposphere and the lower stratosphere, *Atmos. Meas. Tech. Discuss.*, 2016, 1–26, doi:10.5194/amt-2016-219, 2016.
- Sofieva, V. F., Kyrölä, E., Laine, M., Tamminen, J., Degenstein, D., Bourassa, A., Roth, C., Zawada, D., Weber, M., Rozanov, A., Rahpoe, N., Stiller, G., Laeng, A., von Clarmann, T., Walker, K. A., Sheese, P., Hubert, D., van Roozendaal, M., Zehner, C., Damadeo, R., Zawodny, J., Kramarova, N. and Bhartia, P. K.: Merged SAGE II, Ozone_cci and OMPS ozone profile dataset and evaluation of ozone trends in the stratosphere, *Atmos. Chem. Phys.*, 17(20), 12533–12552, doi:10.5194/acp-17-12533-2017, 2017.
- Solomon, S., Ivy, D. J., Kinnison, D., Mills, M. J., Neely, R. R. and Schmidt, A.: Emergence of healing in the Antarctic ozone layer, *Science* (80-.), 353(6296), 269–274, doi:10.1126/science.aae0061, 2016.
- Steinbrecht, W., Froidevaux, L., Fuller, R., Wang, R., Anderson, J., Roth, C., Bourassa, A., Degenstein, D., Damadeo, R., Zawodny, J., Frith, S., McPeters, R., Bhartia, P., Wild, J., Long, C., Davis, S., Rosenlof, K., Sofieva, V. F., Walker, K., Rahpoe, N., Rozanov, A., Weber, M., Laeng, A., von Clarmann, T., Stiller, G., Kramarova, N., Godin-Beekmann, S., Leblanc, T., Querel, R., Swart, D., Boyd, I., Hocke, K., Kämpfer, N., Maillard Barras, E., Moreira, L., Nedoluha, G., Vigouroux, C., Blumenstock, T., Schneider, M., Garcia, O., Jones, N., Mahieu, E., Smale, D., Kotkamp, M., Robinson, J., Petropavlovskikh, I., Harris, N., Hassler, B., Hubert, D. and Tummou, F.: An update on ozone profile trends for the period 2000 to 2016, *Atmos. Chem. Phys.*, 17(17), 10675–10690, doi:10.5194/acp-17-10675-2017, 2017.
- Steiner, A. K., Ladstädter, F., Randel, W. J., Maycock, A. C., Fu, Q., Claud, C., Gleisner, H., Haimberger, L., Ho, S.-P., Keckhut, P., Leblanc, T., Mears, C., Polvani, L. M., Santer, B. D., Schmidt, T., Sofieva, V., Wing, R. and Zou, C.-Z.: Observed temperature changes in the troposphere and stratosphere from 1979 to 2018, *J. Clim.*, 1–72, doi:10.1175/JCLI-D-19-0998.1, 2020.
- Strahan, S. E., Smale, D., Douglass, A. R., Blumenstock, T., Hannigan, J. W., Hase, F., Jones, N. B., Mahieu, E., Notholt, J.,

- 450 Oman, L. D., Ortega, I., Palm, M., Prignon, M., Robinson, J., Schneider, M., Sussmann, R. and Velazco, V. A.: Observed Hemispheric Asymmetry in Stratospheric Transport Trends From 1994 to 2018, *Geophys. Res. Lett.*, 47(17), e2020GL088567, doi:<https://doi.org/10.1029/2020GL088567>, 2020.
- Szeląg, M. E., Sofieva, V. F., Degenstein, D., Roth, C., Davis, S. and Froidevaux, L.: Seasonal stratospheric ozone trends over 2000–2018 derived from several merged data sets, *Atmos. Chem. Phys. Discuss.*, 2020, 1–16, doi:10.5194/acp-2019-1144, 455 2020.
- Toohey, M. and von Clarmann, T.: Climatologies from satellite measurements: the impact of orbital sampling on the standard error of the mean, *Atmos. Meas. Tech.*, 6(4), 937–948, doi:10.5194/amt-6-937-2013, 2013.
- WMO: Scientific assessment of ozone depletion, Global Ozone Research and Monitoring Project-Report No. 52, Geneva, Switzerland. [online] Available from: <https://www.esrl.noaa.gov/csd/assessments/ozone/>, 2014.
- 460 WMO: Scientific Assessment of Ozone Depletion: 2018., 2018.
- Zawada, D. J., Rieger, L. A., Bourassa, A. E. and Degenstein, D. A.: Tomographic retrievals of ozone with the OMPS Limb Profiler: algorithm description and preliminary results, *Atmos. Meas. Tech. Discuss.*, 2017, 1–26, doi:10.5194/amt-2017-236, 2017.