

Observation and modeling of high-⁷Be concentration events at the surface in Northern Europe associated with the instability of the Arctic polar vortex in early 2003

5 Erika Brattich¹, Hongyu Liu², Bo Zhang², Miguel Ángel Hernández-Ceballos³, Jussi Paatero⁴, Darko Sarvan⁵, Vladimir Djurdjevic⁶, Laura Tositti⁷, and Jelena Ajtić⁵

¹ Department of Physics and Astronomy “Augusto Righi” DIFA, Alma Mater Studiorum University of Bologna, via Irnerio 46, 40126 Bologna (BO), Italy

² National Institute of Aerospace, 100 Exploration Way, Hampton, VA 23666, USA

³ Department of Physics, University of Cordoba, Rabanales Campus, 14071 Cordoba, Spain

10 ⁴ Finnish Meteorological Institute, P.O. Box 503, FI-00101, Helsinki, Finland

⁵ Faculty of Veterinary Medicine, University of Belgrade, Bulevar oslobođenja 18, 11000 Belgrade, Serbia

⁶ Institute of Meteorology, Faculty of Physics, University of Belgrade, Studentski trg 18, 11000 Belgrade, Serbia

⁷ Department of Chemistry “G. Ciamician”, Alma Mater Studiorum University of Bologna, via Selmi 2, 40126 Bologna (BO), Italy

15 *Correspondence to:* Erika Brattich (erika.brattich@unibo.it)

Abstract. Events of very high concentrations of ⁷Be cosmogenic radionuclide have been recorded at low-elevation surface stations in the subpolar regions of Europe during the cold season. With an aim to investigate the mechanisms responsible for those peak ⁷Be events, and in particular to verify if they are associated with the fast descent of stratospheric air masses occurring during sudden stratospheric warming (SSWs), we analyse ⁷Be observations at
20 six sampling sites in Fennoscandia during January-March 2003 when very high ⁷Be concentrations were observed and the Arctic vortex was relatively unstable as a consequence of several SSWs. We use the GEOS-Chem chemistry and transport model driven by the MERRA-2 meteorological reanalysis to simulate tropospheric ⁷Be over Northern Europe. We show that the model reasonably reproduces the temporal evolution of surface ⁷Be concentrations observed at the six sampling sites. Our analysis of model simulations, surface ⁷Be observations, atmospheric
25 soundings of ozone and temperature, as well as surface ozone measurements indicates that the ⁷Be peak observed in late February 2003 (between 20 and 28 February 2003) at the six sampling sites in Fennoscandia was associated with downward transport of stratospheric vortex air originated during an SSW that occurred a few days earlier (between 18 and 21 February 2003).

Beryllium-7 (^7Be) is a cosmogenic radionuclide widely monitored and analysed around the world (e.g., Tositti et al., 2004, 2014; Gourdin et al., 2014; Sýkora et al., 2017). Due to its relatively long radioactive half-life (53.22 days) and its cosmogenic origin in the upper troposphere and lower stratosphere (UT-LS) (Lal and Peters, 1967), ^7Be is considered a tracer for intrusion of stratospheric air to the troposphere and large-scale subsidence (e.g., Liu et al., 2016; Chae and Kim, 2019; Heikkila et al., 2008). The variability of the ^7Be activity concentration in surface layers is driven by both static and dynamic factors, e.g., geographical location of the monitoring sites (e.g., Hernández-Ceballos et al., 2015), seasonal atmospheric processes driving transport of carrier aerosols (Lal and Peters, 1967), stratosphere-troposphere air mass exchange (Cristofanelli et al., 2003; Cristofanelli et al., 2009; Putero et al., 2016; Brattich et al., 2017a), synoptic influences (Usoskin et al., 2009), downward transport in the troposphere (Lee et al., 2007), solar activity and dry and wet deposition (e.g., Hernández-Ceballos et al., 2015, 2016a; Ioannidou and Papastefanou, 2006).

The spatial and temporal variability of the ^7Be surface concentrations in Europe and their relationship with meteorological variables was previously analysed in many studies (e.g., Piñero García et al., 2012; Błażej and Mieltski, 2014). The impact of the 11-year solar modulation on the ^7Be concentrations in the air is well established (e.g., Leppänen et al., 2010). The distinctive spring/summer maximum of ^7Be concentrations is widely described and mainly linked with the increased downward transport from the upper troposphere resulting from the intense convection and higher tropopause height typical of the warm season (Cristofanelli et al., 2006; Gerasopoulos et al., 2001, 2003). In addition, cases of high ^7Be surface concentration, some of which occurring over the autumn/winter season, have been analysed in Europe, e.g., over the Iberian Peninsula (Hernández-Ceballos et al., 2017) and at high-altitude stations in the Alps and the Apennines (Brattich et al., 2017; Cristofanelli et al., 2006, 2009). The spring/summer maximum was originally observed with fission products injected into the stratosphere during atmospheric nuclear tests (Dutkiewicz and Husain, 1985; Cristofanelli et al., 2018). A recent study by Salminen-Paatero et al. (2019) who used potential vorticity analysis to gain insights into stratosphere-to-troposphere transport of radionuclides at Rovaniemi (Finnish Lapland) indicated that the transfer of stratospheric air into the upper troposphere was at its maximum in March followed by descending to the ground-level during late spring and early summer.

Further, over the last decade many studies have investigated the ^7Be records in Northern Europe (Leppänen et al., 2010, 2012; Leppänen and Paatero, 2013; Sarvan et al., 2017; Leppänen, 2019), one of the three regions in Europe identified with a distinct ^7Be behaviour in the surface air (Ajtić et al., 2016, 2017; Hernández-Ceballos et

60 al. 2015, 2016b). Among these studies, Ajtić et al. (2016) analysed the ^7Be concentration measured in Helsinki, Finland, over 25 years (1987-2011), and pointed out a relatively high number of ^7Be extremes occurring over autumn and winter: more specifically, 10 % of the highest ^7Be concentrations (above 90th percentile) were observed in the cold season (October-March). Furthermore, recent studies have also indicated that the polar vortices can have a notable influence on the wintertime ^7Be surface concentrations in both the Northern (Ajtić et al., 2018; Bianchi
65 et al., 2019; Terzi and Kalinowski, 2017) and the Southern Hemisphere (Pacini et al., 2015).

In particular, Ajtić et al. (2018) and Bianchi et al. (2019) employed two different methodologies to identify episodes of extremely high ^7Be surface concentrations in autumn and winter, pointing out a large number of cases over the October-March period of the years investigated. The comparison of the dates identified in both analyses showed an overlap with the events of the so-called Sudden Stratospheric Warming (SSW) of the Arctic vortex, i.e.,
70 a sudden rise in the polar temperatures that leads to a highly irregular shape of the vortex and its misalignment from the pole. Ajtić et al. (2018) also noted cases of extremely high ^7Be concentrations occurring right after a very low ^7Be concentration over the Fennoscandian Peninsula during autumn and winter. Overall, this relationship between the SSW of the Arctic vortex and high ^7Be surface concentrations is likely linked to the perturbed stratosphere-troposphere interactions associated with SSWs, which could favor a fast descent of: 1) midlatitude air
75 rich in ^7Be , thus increasing this radionuclide's surface abundance, and 2) aged vortex air wherein ^7Be is subjected to radioactive decay and not transported from outside the vortex, thus decreasing the ^7Be surface abundance.

The atmospheric circulation in the Arctic is dominated by the presence of two distinct polar vortices, one in the troposphere and one in the stratosphere (Waugh et al., 2017). The two vortices present well distinct features: first, the vortex in the troposphere is much more extended than the stratospheric one; and second, while the
80 tropospheric vortex is present all-year round, the stratospheric polar vortex exists only from fall to spring (Waugh et al., 2017). The stratospheric polar vortex in winter stems from the large-scale temperature gradients between the midlatitudes and the poles. Therefore, the stratospheric polar vortex begins to form in autumn as a result of the decreasing solar heating in the polar regions; it strengthens during winter and then breaks down in spring when solar radiation returns to the polar region. Larger topographic and land-sea contrasts and the resulting stronger
85 upward-propagating waves in the Northern Hemisphere, make the northern stratospheric vortex, or the Arctic vortex, weaker and more distorted than its Southern Hemisphere counterpart, the Antarctic vortex. SSW, a major mode of the temporal variability of the Arctic vortex, is the strongest manifestation of the coupling of the stratosphere-troposphere system, with influence on the tropospheric flow lasting for many weeks (Charlton and Polvani, 2007) and with significant effects on chemical composition in the middle atmosphere (Sofieva et al., 2011;
90 Tao et al., 2015). While major SSWs, the so-called vortex split (Charlton and Polvani, 2007; Charlton et al., 2007),

can even cause the stratospheric vortex to break down during midwinter (Vaugh et al., 2017), vortex displacements are instead characterized by a shift of the polar vortex off the pole and its subsequent distortion into a “comma shape” during the extrusion of a vortex filament (Charlton and Polvani, 2007; Charlton et al., 2007). Previous works suggested that the occurrence of SSWs is capable to perturb greatly the polar vortex and hence the stratospheric potential vorticity (PV) distribution (e.g., Matthewman et al., 2009), and the vertical distribution of ozone (e.g., 95 Sonneman et al., 2006; Madhu, 2016). Additionally, the meteorological conditions associated with SSWs in the Arctic have been linked with the occurrences of ^7Be winter extremes, especially in the presence of a very high Scandinavian teleconnection index (Ajtic et al., 2018).

While the initial scientific interest over the stratospheric polar vortex was especially linked to the stratospheric ozone loss over the poles, it is now recognized that the vortices might affect the processes in the 100 troposphere and surface weather (e.g., Mitchell et al., 2013). The present work aims at investigating in detail the atmospheric processes responsible for high ^7Be activities recorded in the cold season over the Fennoscandian Peninsula and its relationship with the Arctic polar vortex through model simulations. For this purpose, we conduct ^7Be simulations for the period of January-March 2003 using the GEOS-Chem global 3-D chemical and transport (CTM) model. The period was selected because of the large number of events with extremely high ^7Be 105 concentrations at surface in Fennoscandia; some of these events were preceded by very low surface concentrations in the lower troposphere ($< 10^{\text{th}}$ percentile). This period thus offers the opportunity to test the hypothesis that SSWs facilitate a fast descent of not only the midlatitude but also vortex air (Ajtić et al., 2018). To achieve this goal, our analysis will therefore focus on:

- 110 • Investigating the processes responsible for the variability of ^7Be concentrations in surface air in Northern Europe;
- Better understanding whether and how SSW and the Arctic vortex winter-time instability influence the surface concentrations of ^7Be in Northern Europe;
- Quantifying the rate of air subsidence on the inner and outer side of the vortex during the period of its 115 instability.

To analyse the influence of SSW and of the Arctic polar vortex on ^7Be concentrations, we first assess the performance of the GEOS-Chem model in reproducing the observed ^7Be variability. We then use model simulations together with other supporting measurements from soundings and meteorological datasets to examine the processes responsible for the variability in the ^7Be concentrations over the period of January-March 2003.

120 As opposite to the cosmogenic origin of ^7Be , ^{210}Pb (half-life 22.3 years) is a nuclide of crustal origin derived from decay of ^{222}Rn (half-life 3.8 days), which is emitted from soils by decay of ^{226}Ra . Owing to the contrasting

natural origins of the two nuclides, the ${}^7\text{Be}/{}^{210}\text{Pb}$ ratio is often regarded as indicative of vertical transport processes and convective activity in the atmosphere (e.g., Koch et al., 1996; Tositti et al., 2004; Brattich et al., 2017a,b). After being produced by contrasting physical mechanisms, both ${}^7\text{Be}$ and ${}^{210}\text{Pb}$ rapidly attach to ambient submicron-
125 sized particles (e.g., Gaffney et al., 2004) and are removed by wet (mainly) and dry (secondarily) deposition processes of their carrier aerosol. The bias in the simulated ${}^7\text{Be}/{}^{210}\text{Pb}$ ratio due to uncertainties in the model deposition schemes is thus reduced. For this reason, besides ${}^7\text{Be}$, the ${}^7\text{Be}/{}^{210}\text{Pb}$ ratio was also analysed to gain further insights into vertical transport processes during the study period.

The rest of this paper is organized as follows. Section 2 describes the radioactivity (${}^7\text{Be}$ and ${}^{210}\text{Pb}$) and
130 meteorological data used. Section 3 provides a brief description of GEOS-Chem, the HYSPLIT trajectory model, and statistical parameters used to assess the model's performance in reproducing the observations. Section 4 presents an overview of the ${}^7\text{Be}$ observations made in Northern Europe in the boreal winter 2003. Section 4 presents and discusses the main results of this work. Specifically, Section 4.1 analyses the precipitation and transport pattern in the study region, while Section 4.2 assesses how well the GEOS-Chem model performs in reproducing the
135 observed variability in the monthly mean surface ${}^7\text{Be}$ concentrations during the study period. Section 4.3 further evaluates the performance of the model in reproducing the short-time variability of ${}^7\text{Be}$ in Northern Europe, followed by an interpretation of the observed variability using model simulations and additional meteorological and ozone observations in section 4.4. Finally, summary and conclusions are given in Section 5.

2 Data

140 In this section we briefly describe the ${}^7\text{Be}$ and ${}^{210}\text{Pb}$ radioactivity data as well as the meteorological datasets analysed in this work.

2.1 ${}^7\text{Be}$ data

Since 1988, the Radioactivity Environmental Monitoring data bank (REMdb) (<https://rem.jrc.ec.europa.eu/RemWeb/>) has brought together and stored in a harmonised way environmental
145 radioactivity data (air, water, milk and mixed diet) measured by the European Member States (Sangiorgi et al., 2019). Among the set of sample types and measurements recommended in 2000/473/Euratom (European Commission, 2000), measurements of natural radioelements, such as ${}^7\text{Be}$ in surface air, are required, and hence, it is very closely monitored and widely stored in REMdb (De Cort et al., 2007).

150 Within the REMdb, the activity concentration of ^7Be in the surface air in Northern Europe (latitude north of 55°N) is available for six surface sampling sites (Hernández-Ceballos et al., 2015), all located at elevations ranging from 9 to 130 m a.s.l.: Ivalo, Umeå, Helsinki, Kista, Harku and Risø, however, with varying start dates and sampling frequencies (Figure 1a). The largest dataset is for Helsinki where, since 1999, the sampling has been performed daily or once every two days. Datasets for Ivalo, Umeå, Kista and Risø also span more than two decades and have a good temporal coverage (roughly once a week since 1995). In particular, ^7Be activity concentrations
165 were obtained by gamma-spectrometry analysis performed by the European Union Competent Authorities. Aerosol samples were collected on filter papers using air samplers with a flow rate of several hundred cubic meters per day and then their radioactivity concentrations were analysed in laboratories.

2.2 ^{210}Pb data

Daily aerosol samples were collected in Helsinki on the roof of the Finnish Meteorological Institute's main
160 building ($60^\circ 10'\text{N}$, $24^\circ 57'\text{E}$). Filters (Munktell MGA, diameter $\varnothing = 240$ mm) were changed every day at 06 UTC. The air volume was about $3500\text{ m}^3/\text{day}$. The filters were assayed for ^{210}Pb by alpha counting of the in-grown daughter nuclide ^{210}Po (Mattsson et al., 1996).

2.3 Meteorological and ozone data

The ^7Be variability is tightly linked to horizontal and vertical transport of the carrier aerosol, and to
165 precipitation that leads to the radionuclide's removal from the atmosphere. Here we used the Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2) meteorological reanalysis (Gelaro et al., 2017) to assist in the data analysis and to drive the GEOS-Chem model simulations. MERRA-2 is produced with version 5.12.4 of the Goddard Earth Observing System (GEOS) atmospheric data assimilation system. It assimilates modern observations of the atmosphere, ocean, land, and chemistry, and includes assimilation of aerosol
170 remote sensing data.

Vertical soundings of air temperature from the Finnish Meteorological Institute's (FMI) Arctic Space Centre (<http://fmiarc.fmi.fi>) at Sodankylä, northern Finland (67.37°N , 26.63°E) were obtained from the University of Wyoming (<http://weather.uwyo.edu/upperair/sounding.html>). To study the effect of downward transport of stratospheric air masses into the troposphere, potential vorticity (PV) values (Holton et al., 1995) were calculated
175 from ERA-Interim wind, temperature, and surface pressure fields (Dee et al., 2011) obtained from the European Centre for Medium-Range Weather Forecasts (ECMWF), Reading, UK.

Ozone sounding data (Kivi et al., 2007; Denton et al., 2019) was retrieved from the database of the FMI's Arctic Space Centre (<http://litdb.fmi.fi>).

3 Methods

180 In this section we give a brief description of the GEOS-Chem and HYSPLIT models and the statistical parameters used to indicate the model performances.

3.1 GEOS-Chem model

185 GEOS-Chem (<http://www.geos-chem.org>) is a global 3-D CTM that has been widely used to study atmospheric composition and processes (e.g., Bey et al., 2001; Park et al., 2004; Eastham et al., 2014). In this study, we use the GEOS-Chem v11-01f to simulate ^7Be and ^{210}Pb and assist in interpreting the observations. GEOS-Chem includes a radionuclide simulation option (^{222}Rn - ^{210}Pb - ^7Be), which simulates the emission, transport (advection, convection, boundary layer mixing), deposition and decay of the radionuclide tracers (Jacob et al., 1997; Liu et al., 2001; Yu et al., 2018).

190 We used the ^7Be production rates recommended by Lal and Peters (1967) for a maximum solar activity year (1958), which has been shown to produce the best results compared to aircraft ^7Be observations in the stratosphere where ^7Be concentrations mainly result from a balance between production and radioactive decay and their observations can be used as a constraint on the ^7Be source (Koch et al., 1996; Liu et al., 2001). In this work, the production rates of Lal and Peters (1967) are formulated as a function of latitude and pressure without seasonal variation (Koch et al., 1996). About two thirds of atmospheric ^7Be is generated in the stratosphere. While there are 195 more accurate modern models (e.g., Masarik and Beer, 1999; Webber et al., 2007; Usoskin and Kovaltsov, 2008; Poluianov et al., 2016), the use of the Lal and Peters production rates is sufficient for this study where we mainly focus on atmospheric transport features. Although the Lal and Peters sources are used here for temporal variability studies, they are not applicable for quantitative studies of the ^7Be activities (e.g., Golubenko et al., 2021)

200 ^{222}Rn emission follows a recent work by Zhang et al. (2020), in which a customized emission map was built upon a few previously published emission scenarios and evaluated against global ^{222}Rn surface observations and aircraft profiles. ^{222}Rn emission flux rate is a function of latitude, longitude, and month. ^7Be and ^{210}Pb are assumed to behave like aerosols once formed in the atmosphere and subject to dry and wet deposition (Liu et al., 2001). Both wet and dry deposition for ^{222}Rn are neglected due to its inert nature.

205 GEOS-Chem simulations in this work are driven by the MERRA-2 meteorological reanalysis. The native resolution of MERRA-2 is 0.667° longitude by 0.5° latitude, with 72 vertical layers (top at 0.01hPa). The meteorological fields are regridded into 2.5° longitude by 2° latitude for the GEOS-Chem simulations in this work. GEOS-Chem uses the TPCORE advection algorithm of Lin and Rood (1996). Convective transport is calculated using archived convective mass fluxes (Wu et al., 2007). Boundary-layer mixing is based on the non-local scheme implemented by Lin and McElroy (2010). The wet deposition scheme follows that of Liu et al. (2001) and includes 210 rainout (in-cloud scavenging) due to stratiform and anvil precipitation, scavenging in convective updrafts (Mari et al., 2000), and washout (below-cloud scavenging) by precipitation (Wang et al., 2011). Precipitation formation and evaporation fields are archived in MERRA-2 and used directly by the model wet deposition scheme. Dry deposition is based on the resistance-in-series scheme of Wesely (1989).

215 In addition to the standard model simulations of ^7Be and ^{210}Pb , we separately transport ^7Be produced in the model layers above the MERRA-2 thermal tropopause (i.e., stratospheric ^7Be tracer) to quantify the stratospheric contribution to ^7Be in the troposphere. This approach was previously used by Liu et al. (2001, 2016). Stratospheric fraction of ^7Be is defined as the ratio of the stratospheric ^7Be tracer concentration to the ^7Be concentration from the standard simulation. All model simulations are conducted for the period of January 2002 – March 2003 with initial conditions from a previous five-year simulation. Hourly and monthly mean outputs for January-March 2003 are 220 used for analysis.

3.2 HYSPLIT

The Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPLIT) model, developed by the NOAA's Air Resources Laboratory (ARL) (Stein et al., 2015), was used to calculate a set of backward trajectories during the study period. To compute the 96 h 3D backward trajectories at 00, 06, 12, and 18 UTC and with different ending 225 heights: 100, 500, 1000 and 1500 m above ground level, the NCEP (National Centers for Environmental Prediction) FNL Operational Global Analysis (NCEP/NWS/NOAA/U.S. Department of Commerce, 2000) meteorological files were used. Computation used the vertical velocity (w , m/s) field contained in the meteorological input file. While 96 h was considered a sufficiently long period to represent the synoptic air flows, the heights were selected to help us to understand the behaviour of the airflows circulating in the Atmospheric Boundary Layer (ABL), just above 230 the ABL, and in the free troposphere. We used the cluster methodology implemented in the HYSPLIT model to group the calculated trajectories according to their length and curvature, and thus identify the airflow patterns over the whole period of the analysis (Hernández-Ceballos et al., 2013; Brattich et al., 2016). It is worth mentioning that

clusters, as well as trajectories, indicate an estimation of the general airflow rather than the exact pathway of an air parcel (e.g., Jorba et al., 2004; Salvador et al., 2008).

235 3.3 Evaluation of the model output

The performance of the model in reproducing observed activity concentrations was evaluated by calculating some basic statistical parameters, such as the mean and standard deviation and other indicators, according to the methodology developed by Hanna (1993) and summarized later by Chang and Hanna (2004). Specifically, the performance of the CTM was evaluated using the following set of indicators, proposed by
 240 Carruthers et al. (2004):

- The mean bias (*MB*), a measure of the mean difference between the modelled and observed concentrations:

$$MB = \overline{C_m - C_o} \quad (1)$$

where C_m is modelled concentration and C_o is observed concentration.

- 245 • The normalized mean square error (*NMSE*), a measure of the mean difference between matched pairs of modelled and observed concentrations:

$$NMSE = \frac{\overline{(C_m - C_o)^2}}{C_m C_o} \quad (2)$$

- The fraction of modelled concentrations within a factor of 2 of observations (*FA2*), i.e., for which $0.5 < C_m/C_o < 2$

- 250 • The Pearson's correlation coefficient (*R*), a measure of the extent of a linear relationship between the modelled and observed concentrations:

$$R = \frac{\sum_{i=1}^n (C_{o,i} - \overline{C_o})(C_{m,i} - \overline{C_m})}{\sqrt{\sum_{i=1}^n (C_{o,i} - \overline{C_o})^2 \sum_{i=1}^n (C_{m,i} - \overline{C_m})^2}} \quad (3)$$

Additionally, the statistical significance of the Pearson's correlation coefficient and the Z-test were also evaluated to identify the presence of statistically significant differences between observed and simulated
 255 concentrations. A perfect model has *MB* and *NMSE* values equal to 0 and *FA2* value equal to 1, while the *R* results range from -1 (perfect negative relationship) to +1 (perfect positive relationship), where 0 implies no relationship between the variables. To better understand the quantitative differences between observations and simulations, scatter plots were used.

4 Results and discussion

260 4.1 Boreal winter 2002/2003

As indicated by Ajtić et al. (2018) and Bianchi et al. (2019), the winter of 2003 offers a good opportunity to investigate a possible link between SSWs and extreme surface concentrations of ^7Be detected in Northern Europe. This period is sufficiently covered by the ^7Be activity concentration measurements at all six monitoring sites.

265 In particular, very high ^7Be activity concentrations, above the 90th percentile (calculated over the 1995-2011 period) simultaneously at most of the Fennoscandian Peninsula sampling sites, were recorded around 23-24 February 2003 (Ajtić et al., 2016, 2018) (Figure 1b). During the 2002/2003 boreal winter, the Arctic vortex was relatively unstable, with six SSWs taking place over the whole season (Peters et al., 2010). Two very pronounced episodes, which were both associated with the vortex splitting and fast SSW recovery, occurred in January and
270 February, respectively (Günther et al., 2008). The evolution of the vortex caused vortex filamentation and vigorous mixing of the vortex and midlatitude stratospheric air (Günther et al., 2008; Müller et al., 2003). Several balloon flights inside the Arctic polar vortex in early 2003 observed unusual trace gas distributions connected to an intrusion of mesospheric air down to altitudes of about 25 km (Engel et al., 2006; Huret et al., 2006; Müller et al., 2007). Such disturbances around the pole are expected to affect the troposphere, i.e., weather conditions (Baldwin and
275 Dunkerton, 2001), and air chemical composition (Hsu, 1980; Limpasuvan et al., 2004). Hence, the high ^7Be concentrations that were measured in Fennoscandia around 24 February 2003 could be a result of downward motion of midlatitude stratospheric air. Interestingly, prior to this episode, very low (below the 10th percentile for each site) surface concentrations of ^7Be were measured in Risø, Kista and Ivalo on 3, 10 and 16 February 2003, respectively (Figure 1). These low values were tentatively linked by Ajtić et al. (2018) with the transport of aged stratospheric
280 vortex air poor in ^7Be , even though they are more likely related to precipitation scavenging that occurred the days before, as shown by the ECA&D (European Climate Assessment & Dataset, <https://www.ecad.eu/>) records. The reader is referred to Ajtić et al. (2018) for more details.

4.2 Analysis of winter precipitation and transport in the Fennoscandian Peninsula: observations vs. model simulations

285 Before analysing the temporal pattern of simulated ^7Be concentrations, we analysed the precipitation and transport pattern in the MERRA-2 meteorological dataset that drives the GEOS-Chem simulations. In particular,

the MERRA-2 precipitation was evaluated against the data from Global Precipitation Climatology Project (GPCP) v2.2 (<https://rda.ucar.edu/datasets/ds728.2/>) satellite and surface observations in winter 2003 (Adler et al., 2003).

290 Figure 2 shows the MERRA-2 and GPCP monthly precipitation in winter for the region within 0-90°N and 90°W – 90°E. Good agreement is found between the MERRA-2 and the GPCP precipitations averaged over the region. Specifically, the geographical distribution of precipitation in MERRA-2 shows some important features that are consistent with the observed climatology precipitations: the desert climate in North Africa with very low precipitation throughout the period, high precipitation over the North Atlantic region during winter, and Europe where the seasonal pattern of precipitation is similar to that in the North Atlantic region.

295 To assess the capability of the model to correctly capture the trend in precipitation during the observation period at the sampling sites, we examined the normalized differences between the MERRA-2 and the observed precipitation, calculated as a difference between the MERRA-2 and the observed values, normalized over the observed value (Table 1).

Overall, the MERRA-2 precipitation tends to be higher than that of GPCP at all sampling sites (Table 1) 300 except for Harku and Helsinki, and especially in the February-March period. This result is in agreement with the findings of Gelaro et al. (2017) who compared the global precipitation of MERRA-2 and GPCP, and reported a general positive bias over northern high latitudes. However, the agreement between MERRA-2 and GPCP precipitation seasonality is reasonable, as indicated by the correlation coefficient values, higher than 0.85 at all sites except for Ivalo (-0.32), and the low *NMSE* values, in the range of 0-0.42 (Table 1). The low negative 305 correlation at Ivalo is due to the fact that while the GPCP-observed precipitation at this site is similar between January and February with a general tendency towards lower values from January to March 2003, the model simulates a decrease from January to February with a small increase in March. However, the statistical parameters reported in Table 1 indicate an overall small discrepancy between the GPCP and MERRA-2 precipitation at all sites.

310 Figure 3 shows that winter circulation in the Fennoscandian Peninsula is dominated by SW and W winds (Chen, 2000; Linderson, 2001). The analysis of the main circulation in the three months in Figure 3 reveals low wind speeds from S-SW in the study area and period. A region of strong wind speeds, possibly corresponding to the Arctic vortex, is clearly visible at surface level to the west of the study area in all the three months. In addition, there appears to be a convergence area (opposite wind directions) between 60 and 75°N. Model-simulated $^7\text{Be}/^{210}\text{Pb}$ 315 ratios and fraction of stratospheric ^7Be increased over the three months period and peaked in March, suggesting increasing stratospheric influence, subsidence, or convective mixing in the study region.

4.3 Variations of the monthly mean surface ^7Be concentrations in the Arctic region: model simulations vs. observations

Figure 4a shows a scatter plot comparing the simulated and observed monthly mean ^7Be concentrations at the six sampling sites. Table 2 reports the statistical parameters and the normalized differences that indicate the performance of the GEOS-Chem model in reproducing the observed ^7Be monthly means.

In general, the model well simulates the month-to-month trend in ^7Be concentrations measured at the sampling sites, as indicated by the fact that all the values fall within the 95% confidence levels (Figure 4a) and the high positive correlation coefficients (> 0.7) except for Ivalo and the low *MB* and *NMSE* values (Table 2). In fact, the normalized differences are not very high (generally < 1), except at Risø. The bias between the model and the observations is partly attributed to the coarse horizontal resolution of the model. Overall, the simulations underestimate the observed values, likely due to uncertainties associated with the deposition schemes and/or precipitation as discussed earlier.

The use of the ^7Be production rate of Lal and Peters (1967) for a solar maximum year (1958) may also partly explain the tendency of simulated ^7Be to be lower than observed. The sunspot number in 2003 (99.3) was rather low (slowly decreasing from 2000, a solar maximum year, and reaching minimum in 2008) compared with the value of 184.8 in 1958. As known, the galactic cosmic-ray intensity, largely responsible for the production of cosmogenic radionuclides, at the Earth's orbit is inversely related to solar activity (Potgieter, 2013), leading to the well-known phase opposition between sunspot number and ^7Be concentration (e.g., Hernández-Ceballos et al., 2015). Sunspot number data herein used were extracted from the World Data Center for the production, preservation and dissemination of the international sunspot number (Sunspot Index and Long-term Solar Observation, SILSO, Royal Observatory of Belgium, Brussels, <http://www.sidc.be/silso/datafiles> - total).

4.4 Variations of the ^7Be weekly and daily mean surface concentrations in the Arctic region: observations vs. model simulations

After analysing the model's performance in reproducing ^7Be monthly mean observations in the previous Section, here we compare the simulated and observed weekly (daily in the case of Helsinki) ^7Be activity concentrations at the six sampling sites (Figure 4b). Table 3 shows the corresponding parameters that indicate the performance of the GEOS-Chem model in reproducing observations. The weekly evolution of simulated versus observed ^7Be concentrations at these sites is shown in Figure 5.

As with the monthly means, the model generally represents adequately the temporal pattern but not the magnitude of weekly mean concentrations, which tend to be lower than those observed (Table 3, Figures 4b and

5). This bias can arise from the higher precipitation in the model than in the observations and/or from errors in the deposition schemes. However, the correct reproduction of the ^7Be temporal pattern, as indicated by the high correlation values at all sampling sites with the exception of Risø, suggests that the model captures the transport processes leading to the peak in ^7Be concentrations at the end of February 2003 and the preceding very low concentration values. In addition, the low *MB* and *NMSE* values calculated at all sites and especially at Ivalo and Umeå suggest that the model reproduces adequately the observed values (Figure 5). The significance of the Z-test for the differences between mean observed and simulated values indicates the presence of statistically significant differences, reflecting the model's general tendency to underestimate observations.

355 As for the $^7\text{Be}/^{210}\text{Pb}$ ratio in Helsinki (Table 3), the model tends to underestimate the observed ratio, which could be due to the model underestimating ^7Be and overestimating ^{210}Pb . Nevertheless, the relatively high correlation between the simulated and observed ratios suggests a reasonable simulation of the temporal pattern of this tracer.

4.5 Understanding the ^7Be variations during the 2002/2003 boreal winter

360 As mentioned earlier (Section 4.1), an SSW event occurred at the end of February 2003. We concentrated our analysis on two different periods during the month: early in the month, between 3 and 16 February when very low ^7Be concentration values were recorded, and at the end of the month between 20 and 28 February characterized by extremely high ^7Be concentrations. To gain further insights into the ^7Be variations during the 2002/2003 boreal winter (Section 4.1), we analysed the simulated $^7\text{Be}/^{210}\text{Pb}$ ratio, maps of surface winds and relative humidity, ozone soundings, vertical cross sections of simulated ^7Be activity concentrations and calculated potential vorticity, and simulated and observed vertical profiles of air temperature. The results were further supported with the analysis of the clusters of back-trajectories during the two different periods of low and high ^7Be concentrations.

370 Risø Figure 6 presents the temporal (weekly mean) pattern of $^7\text{Be}/^{210}\text{Pb}$ and of the stratospheric fraction of ^7Be (calculated as the ratio of the stratospheric ^7Be tracer concentration to the total ^7Be concentration in the troposphere) at the six sampling sites, while daily observations of the $^7\text{Be}/^{210}\text{Pb}$ ratio at Helsinki and Sodankylä (67.367°N, 26.629°E; 160 km south of Ivalo) are presented in the Supplementary Information (hereafter SI). At the beginning of February, the ratio was generally quite low at all the sites. In contrast, the week of 19-26 February 2003 was marked by an evident peak in the $^7\text{Be}/^{210}\text{Pb}$ ratio and a simultaneous increase in the fraction of ^7Be originating in the stratosphere at all sites, which together could be the first indication of a prominent vertical transport from the UT-LS region. The low $^7\text{Be}/^{210}\text{Pb}$ ratio at Risø results from the simultaneous increase of both

^7Be and ^{210}Pb concentrations, together with the delayed stratospheric influence at this site as evidenced by the pattern of the stratospheric ^7Be fraction (Figure 6b).

Hence, we further examined the vertical profiles of temperature with an aim to identify differences in vertical transport near the beginning and end of February. The soundings from the Sodankylä station in the Arctic offer three sets of measurements for each of the investigated periods: on 10 and 16 February, that fall into the period when very low ^7Be concentrations were recorded in Kista and Ivalo, respectively; and 22 and 24 February, the days marked by extremely high ^7Be concentrations over the Fennoscandian Peninsula; 20 and 21 February in the period of transition to high ^7Be concentration over the Fennoscandian Peninsula. Figure 7 shows air temperature profiles in the MERRA-2 dataset and atmospheric soundings at the Sodankylä station. Besides the very good agreement between the MERRA-2 and observed temperatures, a warming of the stratosphere (20-60 km) and a different vertical temperature structure of the lower stratosphere around 20-24 February as compared to the 10 and 16 February profiles are also evident. The observation of the increase in stratospheric temperatures suggests the link between the SSW and the ^7Be peak observed at the six sampling sites located in Northern Europe. In addition, the ozone soundings at the Sodankylä station reveal an ozone mixing ratio peak in the lower troposphere (~1.5-3km) on 19th February 2003 as compared to those observed during 12, 26 and 28 February 2003 (Figure 8a), consistent with downward transport from higher altitudes around that day. Despite the chemical ozone loss in the Arctic vortex in the stratosphere in 2003 as observed by ozone soundings (Tilmes et al., 2006), obviously lower-stratospheric ozone was still enhanced relative to tropospheric ozone. In addition, average O_3 values recorded at surface air quality stations located in Denmark, Finland, and Sweden, which are available through the saqgetr R package (Grange, 2019), show enhanced O_3 concentrations in late February 2003, consistent with the aforementioned peaks in the $^7\text{Be}/^{210}\text{Pb}$ ratio as well as stratospheric ^7Be fraction. This further suggests the transport of stratospheric air masses to the surface.

Simultaneously, the analysis of maps of surface transport and relative humidity (Figure 9) highlights the different winds and relative humidity values in the two periods, with low relative humidity values (~40-50%) suggesting subsidence from 18 to 21 February, and the transition from a clockwise circulation to the fast and complex wind system typical of the second period corresponding to the high ^7Be peak and the SSW. Together with the increase in stratospheric temperatures observed in Figure 7, the reversal of zonal winds confirms the link between the ^7Be peak and the SSW event, in agreement with Sofieva et al. (2012). Interestingly, the parcel of the lowest relative humidity values occurs during the 18-21 February period, i.e., a couple of days before the dates of the ^7Be peaks in the measurements and those peaks in the simulated $^7\text{Be}/^{210}\text{Pb}$ ratio and stratospheric ^7Be fraction (Figure 1b and Figure 6). Together with enhanced ozone concentrations observed in the lower troposphere on 19th

February, this suggests that the downward transport from the UT-LS was triggered by the SSW occurring a few days before.

To better constrain the stratospheric origin of the air masses arriving at the sampling sites during the two
410 periods, we further analyzed the potential vorticity data from ECMWF during the month of February 2003 at three
latitudes (63, 64.5 and 66°N) along the 21°E meridian (Figure 10), while vertical cross sections of MERRA-2 PV
are provided in Figure S3. The data reveal clearly a bubble of high potential vorticity down to the surface at the
three latitudes from 18 till 22 February 2003, particularly at the northernmost latitude where values higher than 1.6
PVU, a value considered as a threshold for stratospheric air in the lower troposphere especially when in conjunction
415 with low relative humidity, high $^7\text{Be}/^{210}\text{Pb}$ ratios and ozone (Cristofanelli et al., 2006), were observed. Indeed, PV
is considered as a quasi-passive tracer, because of its conservation properties under adiabatic conditions (Hoor et
al., 2010; Gettelman et al., 2011), and the tropopause level can be identified as regions of strong enhancements in
gradients of PV, essentially indicated by distinct values in the troposphere and stratosphere. Recent studies of PV
cutoff lows in the Northern Hemisphere extratropics (Pinheiro et al., 2017; Portmann et al., 2020; Muñoz et al.,
420 2020) suggest that the majority of the events are relatively short-lived, persisting for about 2-3 days, a value in
good agreement with the vertical motion subject of this study.

The low relative humidity (Figure 9) and high potential vorticity (Figure 10) corresponded to high ^7Be
descending to lower atmospheric levels, as simulated by the model (Figure 11). The descending vertical motion
from the upper vertical levels during the period is clearly visible in the MERRA-2 vertical pressure velocity
425 (omega) fields sampled at the six sampling sites for the month of February 2003, especially for the northernmost
sites (Figure 12) where omega is largely positive with near-surface values up to $0.3\text{-}0.4 \text{ Pa s}^{-1}$ around 18-19
February. Further evidence of this is seen from the maps of vertical pressure velocity (Figure 13) and of the
stratospheric fraction of ^7Be originated in the stratosphere (Figure 14) at 940 hPa in the boundary layer. Note that
positive vertical pressure velocity as seen over both Fennoscandia and Greenland (Figure 13) during 18-25
430 February 2003 indicates descending motions that can facilitate the transport of stratospherically influenced air, if
present, to the ground-level. This explains why Fennoscandia saw increased stratospheric influence on surface ^7Be
concentrations during this period (Figure 14), but Greenland did not see much. However, on monthly average,
Greenland is a region with significant stratospheric influences in February 2003 (middle right panel, Figure 3).

Analysis and comparison of 4-day back-trajectories at each sampling station allowed a reconstruction of
435 two distinct atmospheric circulation patterns in the two periods. Figure 15 shows the clustering results for three
sampling sites (Ivalo, Harku, and Risø) during both periods. The stations are ordered as a function of decreasing
latitude from high (upper panels) to low (lower panels). Only results at 1000 m (well above the winter PBL height)

are shown and cluster results for other altitudes in the lower troposphere are similar. While the first period (with low ^7Be values) presents the dominance of westerly winds (air masses flowing eastward), as typical of these latitudes (Zanis et al., 1999), the second period is characterized by a clockwise displacement of airflows with origins at higher altitudes (Figure 15). This pattern in the second period is better established at lower latitude stations (Helsinki, Harku, Kista and Risø) compared to higher latitude ones (Ivalo, Umeå). It likely results from the aged vortex (Günther et al., 2008) and the SSW at the end of February, corresponding to a decrease in the MERRA-2 daily average height of the thermal tropopause on 21st-22nd February at the Sodankylä station in Finland (Figure S4) during the SSW (Peethani et al., 2014; Wargan and Coy, 2016). Associated with these processes is the downward transport of stratospheric air previously identified with an independent approach.

5 Summary and conclusions

We have used a global 3-D model (GEOS-Chem) driven by the MERRA-2 meteorological reanalysis to simulate atmospheric concentrations of ^7Be of cosmogenic origin for the period of January-March 2003. The aim was to verify the mechanisms responsible for the surface ^7Be variabilities in Northern Europe, and to test the hypothesis that SSWs may facilitate fast descent of UT-LS vortex air to the surface. The period was selected as it involves two intense SSWs and observations of extremely high ^7Be concentrations at six sampling sites in Fennoscandia.

Before using the model's output to investigate the processes responsible for ^7Be variability in Northern Europe over the period, we evaluated the MERRA-2 precipitation fields against the GPCP satellite and surface observations. A generally good agreement was found both at regional scale and at the six sampling sites. Analysis of the wind fields in the study period indicated low wind speeds from S-SW in agreement with the major circulation patterns over the Fennoscandian Peninsula in winter, and the presence of a region of strong wind speeds to the west of the study area, likely in connection with the Arctic polar vortex.

The model reproduced efficiently the ^7Be and $^7\text{Be}/^{210}\text{Pb}$ temporal (i.e., monthly and weekly) patterns at the six sampling sites in the study period, even though it presented a tendency to underestimate the observed surface ^7Be concentrations. The lower modelled values are likely due to its coarse horizontal resolution (2.5° longitude by 2° latitude), lack of year-to-year variation in ^7Be production rates, and uncertainties associated with precipitation scavenging.

In order to investigate the processes responsible for ^7Be variability at the six sampling sites during the study period, and in particular to test whether the peak ^7Be concentrations measured in Fennoscandia around 24 February

2003 originated from fast descent of stratospheric vortex air facilitated by SSW, we analysed time-height cross sections of simulated ^7Be and potential vorticity, vertical profiles of air temperature, maps of surface winds and relative humidity, and ozone soundings. The analysis of the temporal variations of simulated $^7\text{Be}/^{210}\text{Pb}$ ratio and
470 fraction of ^7Be originated in the stratosphere indicates a peak during the week of 19-26 February 2003, suggesting downward transport from the UT-LS region. The latter was corroborated by a layer of ozone mixing ratio enhancements in the lower troposphere recorded by the soundings at the Sodankylä station on 19th February. Furthermore, the vertical profiles of air temperature indicated a warming of the stratosphere and a change in shape in the vicinity of the tropopause region during the period, suggesting the link between the downward transport of
475 the vortex air and SSW.

Our analysis of time-height cross sections of simulated ^7Be concentrations, calculated potential vorticity and MERRA-2 vertical pressure velocity (omega) revealed the vertical downward transport to the surface of a stratospheric air parcel characterized by high potential vorticity, high vertical velocity (in particular at Ivalo on 19th February) and high ^7Be concentrations, further supporting the stratospheric origin of the air masses during the
480 investigated period.

Additionally, low relative humidity and a change in the circulation pattern from slow, clockwise to fast, swirling winds occurred over the study area. The change in the circulation pattern and the downward transport of stratospheric air was verified by the analysis of the clusters of back trajectories during the periods of low and high ^7Be concentrations, which showed a change from westerlies to airflows from upper vertical levels.

485 Altogether, these analyses confirm the link between the SSW and transport of stratospheric air to the surface, resulting in high surface ^7Be concentrations observed in February 2003 in Fennoscandia. Since more frequent SSWs are expected in a warmer climate (Kang and Tziperman, 2017, 2018; Simpkins, 2017), this link has important implications for the impact of climate change on atmospheric transport, tropospheric composition, and air quality in northern high-latitude regions.

490 **Data availability**

^7Be activity concentration data are available in the Radioactivity Environmental Monitoring (REM) database (<https://data.jrc.ec.europa.eu/collection/id-0117>). All model output, and ^{210}Pb daily observational data at Helsinki and Sodankylä for January-March 2003 are available online (<http://doi.org/10.5281/zenodo.4117521>).

Author contributions

495 JA, MAHC and EB designed the study. HL and BZ conducted the GEOS-Chem model simulations. MAHC led the calculation and analysis of HYSPLIT back-trajectories. EB developed the analysis methodology and led the analysis of observational data and model output, with contributions from all coauthors. JP contributed ²¹⁰Pb and meteorological observational datasets. EB wrote the manuscript with contributions from all coauthors.

Competing interests

500 The authors declare that they have no conflict of interest.

Acknowledgements

The study was partially supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Contract numbers 451-03-9/2021-14/200143 and 451-03-9/2021-14/200162). HL and BZ acknowledge funding support from the NASA Modeling, Analysis and Prediction Program (grant
505 80NSSC17K0221) and Atmospheric Composition Campaign Data Analysis and Modeling program (grant NNX14AR07G). NASA Center for Climate Simulation (NCCS) provided supercomputing resources. The GEOS-Chem model is managed by the Atmospheric Chemistry Modeling Group at Harvard University with support from NASA ACPMAP and MAP programs. The University of Wyoming, ECMWF, Dr. Rigel Kivi from the Finnish Meteorological Institute Arctic Space Center and Dr. Laura Thölix from the Finnish Meteorological Institute are
510 gratefully acknowledged for providing and helping with temperature profiles, ozone soundings and potential vorticity data used to support the detection of air masses of stratospheric origin in the troposphere. The World Data Center is gratefully acknowledged for the production, preservation and dissemination of the international sunspot number (Sunspot Index and Long-term Solar Observation, SILSO, Royal Observatory of Belgium, Brussels, <http://www.sidc.be/silso/datafiles#total>). Support for the Twentieth Century Reanalysis Project version 3 dataset is
515 provided by the U.S. Department of Energy, Office of Science Biological and Environmental Research (BER; <http://science.energy.gov/ber/>), by the National Oceanic and Atmospheric Administration Climate Program Office, and by the NOAA Physical Sciences Laboratory. NASA/GSFC/SED/ESD/LA/MAPB and contributors are gratefully acknowledged for developing and computing the GPCP (Global Precipitation Climatology Project) as part of the Global Energy and Water Cycle Exchanges (GEWEX) project. Stuart Grange is gratefully acknowledged
520 for providing access to European air quality data through the saqgetr R package.

References

- Adler, R.F., Huffmann, G.J., Chang, A., Ferraro, R., Xie, P.-P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D., Gruber, A., Susskind, J. Arkin, P, and Nelkin, E., 2003: The version-2 Global Precipitation Climatology Project (GPCP) monthly precipitation analysis (1979-present). *J. Hydrometeorol.*, 4(6), 1147-1167, doi:10.1175/1525-7541(2003)004<1147:TVGPCP>2.0.CO;2
- 525 Ajtić, J., Djurdjevic, V., Sarvan, D., Brattich, E., and Hernández-Ceballos, M.A.: Analysis of extreme beryllium-7 specific activities in surface air, *Radiation & Applications*, 1(3), 216–221, doi:10.21175/RadJ.2016.03.040, 2016.
- Ajtić, J. V., Sarvan D., Djurdjevic, V. S., Hernández-Ceballos, M. A., and Brattich, E.: Beryllium-7 surface concentration extremes in Europe, *Facta Universitatis – Series: Physics, Chemistry and Technology*, 15 (1), 45–
- 530 55, 2017.
- Ajtić, J., Brattich, E., Sarvan, D., Djurdjevic, V., and Hernández-Ceballos, M. A.: Factors affecting the ^7Be surface concentration and its extremely high occurrences over the Scandinavian Peninsula during autumn and winter, *Chemosphere*, 199, 278–285, doi:10.1016/j.chemosphere.2018.02.052, 2018.
- Albers, J.R., Kiladis, G.N., Birner, T., and Dias, J.: Tropical upper-tropospheric potential vorticity intrusions during sudden stratospheric warmings. *J. Atmos. Sci.*, 73 (6), 2361-2384, doi:10.1175/JAS-D-15-0238.1
- 535 Baldwin, M.P., and Dunkerton, T. J.: Stratospheric Harbingers of Anomalous Weather Regimes, *Science*, 294, 581–584, doi:10.1126/science.1063315, 2001.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q. B., Liu, H. G. Y., Mickley, L. J. and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res. Atmos.*, 106(D19), 23073–23095, doi:10.1029/2001JD000807, 2001.
- 540 Bianchi, S., Plastino, W., Brattich, E., Djurdjevic, V., Longo, A., Hernández-Ceballos, M. A., Sarvan, D., and Ajtić, J.: Analysis of trends, periodicities, and correlations in the beryllium-7 time series in Northern Europe, *Appl. Radiat. Isot.* 148, 160-167, doi:10.1016/j.apradiso.2019.03.038, 2019.
- Błażej, S., and Mietelski, J. W.: Cosmogenic ^{22}Na , ^7Be and terrestrial ^{137}Cs , ^{40}K radionuclides in ground level air samples collected weekly in Kraków (Poland) over years 2003–2006, *J. Radioanal. Nucl.*, 300, 747–756, doi:10.1007/s10967-014-3049-6, 2014.
- Brattich, E., Hernández-Ceballos, M.A., Orza, J.A.G., Bolívar, J.P., and Tositti, L.: The western Mediterranean basin as an aged aerosols reservoir. Insights from an old-fashioned but efficient radiotracer. *Atmos. Environ.* 141, 481-493, doi:10.1016/j.atmosenv.2016.07.022, 2016

- 550 Brattich, E., Orza, J.A.G., Cristofanelli, P., Bonasoni, P., and Tositti, L.: Influence of stratospheric air masses on radiotracers and ozone over the central Mediterranean, *J. Geophys. Res.*, 122, doi:10.1002/2017JD027036, 2017a.
- Brattich, E., Liu, H., Tositti, L., Considine, D. B., and Crawford, J. H.: Processes controlling the seasonal variations in ^{210}Pb and ^7Be at the Mt. Cimone WMO-GAW global station, Italy: a model analysis, *Atmos. Chem. Phys.*, 17(2), 1061-1080, doi:10.5194/acp-17-1061-2017, 2017b.
- 555 Butler, A.H., Seidel, D. J., Hardiman, S. C., Butchart, N., Birner, T., and Match, A.: Defining Sudden Stratospheric Warmings, *Bull. Am. Meteorol. Soc.*, 96, 1913–1928, doi:10.1175/BAMS-D-13-00173.1, 2015.
- Carruthers, D.J., Edmunds, H.A., Lester, A.E., McHugh, C.A., and Singles, R.J.: Use and validation of ADMS-Urban in contrasting urban and industrial locations, *Int. J. Environ. Pollut.*, 14, 363-374, doi:10.1504/IJEP.2000.000558, 2000.
- 560 Chae, J-S., and Kim, G.: Large seasonal variations in fine aerosol precipitation rates revealed using cosmogenic ^7Be as a tracer, *Sci. Total Environ.*, 673, 1-6, doi:10.1016/j.scitotenv.2019.03.482, 2019.
- Chang, J.C., and Hanna, S.R.: Air quality model performance evaluation, *Meteorol. Atmos. Phys.*, 87, 167-196, doi:10.1007/s00703-003-0070-7, 2004.
- Charlton, A.J., and Polvani, L.M.: A new look at Stratospheric Sudden Warmings. Part I: climatology and
565 modelling benchmarks. *J. Clim.*, 20(3), 449-469, doi:10.1175/JCLI3996.1, 2007.
- Charlton, A.J., Polvani, L.M., Perlwitz, J., Sassi, F., Manzini, E., Shibata, K., Pawson, S., Nielsen, J.E., and Rind, D.: A new look at Stratospheric Sudden Warmings. Part II: evaluation of numerical simulations. *J. Clim.*, 20(3), 470-488, doi:10.1175/JCLI3994.1, 2007.
- Chen, D.: A monthly circulation climatology for Sweden and its application to a winter temperature case study,
570 *Int. J. Climatol.*, 20, 1067-1076, doi:10.1002/1097-0088(200008)20:10<1067::AID-JOC528>3.0.CO;2-Q, 2000.
- Cristofanelli, P., Bonasoni, P., Collins, W., Feichter, J., Forster, C., James, P., Kentarchos, A., Kubik, P.W., Land, C., Meloen, J., Roelofs, G.J., Siegmund, P., Sprenger, M., Schnabel, C., Stohl, A., Tobler, L., Tositti, L., Trickl, T., and Zanis, P.: Stratosphere-to-troposphere transport: A model and method evaluation, *J. Geophys. Res. Atmos.*, 108 (12), doi:10.1029/2002JD002600, 2003.
- 575 Cristofanelli, P., Bonasoni, P., Tositti, L., Bonafè, U., Calzolari, F., Evangelisti, F., Sandrini, S., and Stohl, A.: A 6-year analysis of stratospheric intrusions and their influence on ozone at Mt. Cimone (2165 m above sea level), *J. Geophys. Res. Atmos.*, 111, doi: 10.1029/2005JD006553, 2006
- Cristofanelli, P., Calzolari, F., Bonafé, U., Duchi, R., Marinoni, A., Roccatò, F., Tositti, L., and Bonasoni, P.: Stratospheric intrusion index (SI2) from baseline measurement data, *Theor. Appl. Climatol.*, 97, 317-325, doi:
580 10.1007/s00704-008-0073-x, 2009.

- Cristofanelli, P., Brattich, E., Decesari, S., Landi, T.C., Maione, M., Putero, D., Tositti, L., and Bonasoni, P.: Studies on environmental radionuclides at Mt. Cimone, Chapter in: High Mountain Atmospheric Research. The Italian Mt. Cimone WMO/GAW Global Station (2165 m a.s.l.), SpringerBriefs in Meteorology, Springer International Publishing, ISBN: 978-3-319-61126-6, doi: 10.1007/978-3-319-61127-3, 2018.
- 585 De Cort, M., Sangiorgi, M. Hernandez Ceballos, M. A., Vanzo, S., Nweke, E., Tognoli, P. V., Tollefsen, T.: REM data bank - Years 1984-2006. European Commission, Joint Research Centre (JRC) [Dataset] doi:10.2905/jrc-10117-10024 PID: <http://data.europa.eu/89h/jrc-10117-10024>, 2007
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N.,
590 Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137, 553-597, <https://doi.org/10.1002/qj.828>, 2011.
- 595 Denton, M.H., Kivi, R., Ulich, T., Rodger, C.J., Clilverd, M.A., Denton, J.S., and Lester, M.: Observed response of stratospheric and mesospheric composition to sudden stratospheric warmings. *J. Atmos. Sol.*, 191, 105054, doi: 10.1016/j.jastp.2019.06.001, 2019.
- Dutkiewicz, V.A., and Husain, L.: Stratospheric and tropospheric components of ^7Be in surface air, *J. Geophys. Res. Atmos.*, 90, D3, 5783-5788, doi:10.1029/JD090iD03p05783, 1985.
- 600 Eastham, S. D., Weisenstein, D. K. and Barrett, S. R. H.: Development and evaluation of the unified tropospheric-stratospheric chemistry extension (UCX) for the global chemistry-transport model GEOS-Chem, *Atmos. Environ.*, 89, 52–63, doi:10.1016/j.atmosenv.2014.02.001, 2014.
- Engel, A., Möbius, T., Haase, H.-P., Bönisch, H., Wetter, T., Schmidt, U., Levin, I., Reddman, T., Oelhaf, H., Wetzell, G., Grunow, K., Huret, N., Pirre, M.: Observation of mesospheric air inside the arctic stratosphere polar
605 vortex in early 2003. *Atmos. Chem. Phys.*, 6, 267-282, doi:10.5194/acp-6-267-2006, 2006.
- European Commission, 2000. Commission recommendation of 8 June 2000 on the application of Article 36 of the Euratom Treaty concerning the monitoring of the levels of radioactivity in the environment for the purpose of assessing the exposure of the population as a whole. Available at <https://op.europa.eu/it/publication-detail/-/publication/8116b329-eb85-4bda-8d4b-e8c9e0d152c2/language-en>, last accessed on 2 July 2021

- 610 European Union (EU): Treaty Establishing the European Atomic Energy Community (E.A.E.C. – EURATOM), Consolidated Version, Title Two - Provisions for the Encouragement of the Progress in the Field of Nuclear Energy, Chapter III: Health and Safety, 2010.
- Feely, H.W., Larsen, R. J., and Sanderson, C. G.: Factors that cause seasonal variations in beryllium-7 concentrations in surface air, *J. Environ. Radioact.*, 9, 223–249, doi:10.1016/0265-931X(89)90046-5, 1989.
- 615 Gaffney, J.S., Marley, N., and Cunningham, M.M.: Natural radionuclides in fine aerosols in the Pittsburgh area. *Atmos. Environ.*, 38, 3191-3200, doi:10.1016/j.atmosenv.2004.03.015, 2004.
- Gelaro, R., McCarthy, W., Suárez, M.J., Todling, R., Molod, A., Takacs, L., Randles, C.A., Darmenov, A., Bosilovich, M.G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A.M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J.E., Partyka, G., Pawson, S.,
- 620 Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Clim.*, 30(14), 5419-5454, doi:10.1175/JCLI-D-16-0758.1, 2017.
- Gerasopoulos, E., Zanis, P., Stohl, A., Zerefos, C. S., Papastefanou, C., Ringer, W., Tobler, L., Hübener, S., Gäggeler, H. W., Kanter, H. J., Tositti, L., and Sandrini, S.: A climatology of ⁷Be at four high-altitude stations at
- 625 the Alps and the Northern Apennines, *Atmos. Environ.*, 35, 6347–6360, doi :10.1016/S1352-2310(01)00400-9, 2001.
- Gerasopoulos, E., Zerefos, C. S., Papastefanou, C., Zanis, P., and O’Brien, K.: Low-frequency variability of beryllium-7 surface concentrations over the Eastern Mediterranean, *Atmos. Environ.*, 37, 1745–1756, doi:10-1016/S13552-2310(03)00068-2, 2003.
- 630 Gettelman, A., Hoor, P., Pan, L.L., Randel, W.J., Hegglin, M.I., and Birner, T. : The extratropical upper troposphere and lower stratosphere. *Rev. Geophys.*, 49, RG3003, doi :10.1029/2011RG000355, 2011.
- Golubenko, K., Rozanov, E., Kovaltsov, G., Leppänen, A.-P., Sukholodov, T., and Usoskin, I.: Chemistry-climate model SOCOL-AERv2-BE v1 with the cosmogenic Beryllium-7 isotope cycle. *Geosci. Model Dev. Discuss.*, doi: 10.5194/gmd-2021-56, in review, 2021.
- 635 Gourdin, E., Evrard, O., Huon, S., Reyss, J.-L., Ribolzi, O., Bariac, T., Sengtaheuanghoung, O., and Ayrault, S.: Spatial and temporal variability of ⁷Be and ²¹⁰Pb wet deposition during four successive monsoon storms in a catchment of northern Laos, *J. Environ. Radioact.*, 136, 195–205, doi:10.1016/j.jenvrad.2014.06.008, 2014.
- Grange, S.K.: Technical note: saqgetr R package. Available at <https://drive.google.com/open?id=1IgDODHqBHewCTKLdAAxRyR7ml8ht6Ods>. Last accessed on 2 July 2021,
- 640 2019.

- Günther, G., Müller, R., von Hobe, M., Stroh, F., Konopka, P., and Volk, C. M.: Quantification of transport across the boundary of the lower stratospheric vortex during Arctic winter 2002/2003, *Atmos. Chem. Phys.*, 8, 3655–3670, doi:10.5194/acp-8-3655-2008, 2008.
- Hanna, S.R.: Uncertainties in air quality model predictions, *Bound.-Layer Meteorol.*, 62, 3-20, 1993.
- 645 Hernández-Ceballos, M.A., Cinelli, G., Marín Ferrer, M., Tollefsen, T., De Felice, L., Nweke, E., Tognoli, P. V., Vanzo, S., and De Cort, M.: A climatology of ^7Be in surface air in European Union, *J. Environ. Radioactiv.*, 141, 62–70, doi:10.1016/j.jenvrad.2014.12.003, 2015.
- Heikkilä, U., Beer, J., and Feichter, J.: Modeling cosmogenic radionuclides ^{10}Be and ^7Be during the Maunder Minimum using the ECHAM5-HAM General Circulation Model., *Atmos. Chem. Phys.*, 8, 1797-2809, doi: 10.5194/acp-8-2797-2008, 2008
- 650 Hernández-Ceballos, M.A., Adame, J.A., Bolívar, J.P., De La Morena, P.: Vertical behaviour and meteorologica properties of air masses in the southwest of the Iberian Peninsula (1997-2007). *Meteorol. Atmos. Phys.* 119, 163-175, doi:10.1007/s00703-012-0225-5, 2013.
- Hernández-Ceballos, M.A., Brattich, E., Cinelli, G., Ajtić, J., and Djurdjevic, V. Seasonality of ^7Be concentrations in Europe and influence of tropopause height, *Tellus B*, 68, 29534, doi:10.3402/tellusb.v68.29534, 2016.
- 655 Hernández-Ceballos, M.A., Brattich, E., Lozano, R. L., and Cinelli, G.: ^7Be behaviour and meteorological conditions associated with ^7Be peak events in Spain, *J. Environ. Radioactiv.*, 166, 17–26, doi:10.1016/j.jenvrad.2016.03.019, 2017
- Hoor, P., Wernli, H., Hegglin, M.I., and Bönisch, H.: Transport timescales and tracer properties in the extratropical UTLS. *Atmos. Chem. Phys.*, 10, 7929-7944, doi:10.5194/acp-10-7929-2010, 2010.
- 660 Hsu, C.-P.F.: Air Parcel Motions during a Numerically Simulated Sudden Stratospheric Warming, *J. Atmos. Sci.*, 37, 2768–2792, doi:10.1175/1520-0469(1980)037<2768:APMDAN>2.0.CO;2, 1980.
- Huret, N., Pirre, M., Hauchecorne, A., Robert, C., Catoire, V., 2006. On the vertical structure of the stratosphere at midlatitudes during the first stage of the polar vortex formation and in the polar region in the presence of a large mesospheric descent. *J. Geophys. Res.* 111(D6), doi:10.1029/2005JD006102
- 665 Ioannidou, A., and Papastefanou, C.: Precipitation scavenging of ^7Be and ^{137}Cs radionuclides in air, *J. Environ. Radioactiv.*, 85, 121–136, doi:10.1016/j.jenvrad.2005.06.005, 2006.
- Jacob, D. J. D., Prather, M. J. M., Rasch, P. P. J., Shia, R. R.-L., Balkanski, Y. J., Beagley, S. R., Bergmann, D. J., Blackshear, W. T., Brown, M., Chiba, M., Chipperfield, M.P., de Grandpré, J., Dignon, J.E., Feichter, J., Genthon, C., Grose, W.L., Kasibhatla, P.S., Köhler, I., Kritz, M.A., Law, K., Penner, J.E., Ramonet, M., Reeves, C.E.,
- 670

- Rotman, D.A., Stockwell, D.Z., Van Velthoven, P.F.J., Verver, G., Wild, O., Yang H., and Zimmermann, P.: Evaluation and intercomparison of global atmospheric transport models using ^{222}Rn and other short-lived tracers, *J. Geophys. Res. Atmos.*, 102(D5), 5953–5970, doi:10.1029/96JD02955, 1997.
- Jorba, O., Pérez, C., Rocadenbosch, F., and Baldasano, J.M. Cluster analysis of 4-day back trajectories arriving in the Barcelona area, Spain, from 1997 to 2002, *J. Appl. Meteorol. Climatol.*, 43,887–901, doi:10.1175/1520-0450(2004)043<0887:CAODBT>2.0.CO;2, 2004.
- Kang, W., and Tziperman, E.: More frequent sudden stratospheric warming events due to enhanced MJO forcing expected in a warmer climate. *J. Clim.*, 30(21), 8727-8743, doi:10.1175/JCLI-D-17-0044.1, 2017.
- Kang, W., and Tziperman, E.: The MJO-SSW teleconnection: interaction between MJO-forced waves and the midlatitude jet. *Geophys. Res. Lett.*, 45(9), 4400-4409, doi:10.1029/2018GL077937, 2018.
- Kivi, R., Kyrö, E., Turunen, T., Harris, N.R.P., von der Gathen, P., Rex, M., Anderson, S.B., and Wohltmann, I.: Ozone observations in the Arctic during 1989-2003: ozone variability and trends in the lower stratosphere and free troposphere, *J. Geophys. Res.*, 112, p. D08306, doi:10.1029/2006JD007271, 2007.
- Koch, D.M., Jacob, D.J., and Graustein, W.C.: Vertical transport in tropospheric aerosols as indicated by ^7Be and ^{210}Pb in a chemical tracer model, *J. Geophys. Res. Atmos.*, 101, 18651-18666, doi:10.1029/96JD01176, 1996.
- Lal, D., and Peters, B.: Cosmic ray produced radioactivity on the earth, in: Sitte, K. (Ed.), *Cosmic Rays II*. Springer Berlin Heidelberg, pp. 551–612, 1967.
- Lee, H.N., Tositti, L., Zheng, X., and Bonasoni, P.: Analyses and comparisons of ^7Be , ^{210}Pb and activity ratio $^7\text{Be}/^{210}\text{Pb}$ with ozone observations at two GAW stations from high mountains, *J. Geophys. Res. Atmos.*, 112, 1-11, doi: 10.1029/2006JD007421, 2007.
- Leppänen, A.-P.: Deposition of naturally occurring ^7Be and ^{210}Pb in Northern Finland, *J. Environ. Radioactiv.*, 208-209, 105995, doi:10.1016/j.jenvrad.2019.105995, 2019.
- Leppänen, A.-P., Pacini, A.A., Usoskin, I.G., Aldahan, A., Echer, E., Evangelista, H., Klemola, S., Kovaltsov, G.A., Mursula, K., and Possnert, G.: Cosmogenic ^7Be in air: a complex mixture of production and transport, *J. Atmos. Sol.*, 72, 1036-1043, doi:10.1016/j.jastp.2010.06.006, 2010.
- Limpasuvan, V., Thompson, D. W. J., Hartmann, D. L.: The Life Cycle of the Northern Hemisphere Sudden Stratospheric Warmings. *J. Clim.*, 17, 2584–2596, doi:10.1175/1520-0442(2004)017<2584:TLCOTN>2.0.CO;2, 2004
- Lin, J. T., and McElroy, M. B.: Impacts of boundary layer mixing on pollutant vertical profiles in the lower troposphere: Implications to satellite remote sensing, *Atmos. Environ.*, 44(14), 1726–1739, doi:10.1016/j.atmosenv.2010.02.009, 2010.

- Lin, S.-J. and Rood, R. B.: Multidimensional Flux-Form Semi-Lagrangian Transport Schemes, *Mon. Weather Rev.*, 124(9), 2046–2070, doi:10.1175/1520-0493(1996)124<2046:MFFSLT>2.0.CO;2, 1996.
- Linderson, M.-L.: Objective classification of atmospheric circulation over Southern Scandinavia, *Int. J. Climatol.*, 21, 155-169, doi:10.1002/joc.604, 2001.
- 705 Liu, H., Jacob, D. J., Bey, I. and Yantosca, R. M.: Constraints from ^{210}Pb and ^7Be on wet deposition and transport in a global three-dimensional chemical tracer model driven by assimilated meteorological fields, *J. Geophys. Res. Atmos.*, 106(D11), 12109–12128, doi:10.1029/2000JD900839, 2001.
- Liu, H., Considine, D. B., Horowitz, L. W., Crawford, J. H., Rodriguez, J. M., Strahan, S. E., Damon, M. R., 710 Steenrod, S. D., Xu, X., Kouatchou, J., Carouge, C., and Yantosca, R. M.: Using beryllium-7 to assess cross-tropopause transport in global models, *Atmos. Chem. Phys.*, 16, 4641–4659, doi:10.5194/acp-16-4641-2016, 2016.
- Madhu, V.: Effects of sudden stratospheric warming events on the distribution of total column ozone over polar and middle latitude regions. *Open J. Mar. Sci.* 6(2), doi:10.4236/ojms.2016.62025
- Mari, C., Jacob, D. J. and Bechtold, P.: Transport and scavenging of soluble gases in a deep convective cloud, *J. Geophys. Res. Atmos.*, 105(D17), 22255–22267, doi:10.1029/2000JD900211, 2000.
- 715 Masarik, J. and Beer, J.: Simulation of particle fluxes and cosmogenic nuclide production in the Earth's atmosphere. *J. Geophys. Res.*, 104, D10, 12099-12111, doi:10.1029/1998JD200091, 1999.
- Mattsson, R., Paatero, J., and Hatakka, J.: Automatic Alpha/Beta Analyser for Air Filter Samples - Absolute Determination of Radon Progeny by Pseudo-coincidence Techniques, *Radiat. Prot. Dosim.*, 63, 133-139, 720 doi:10.1093/oxfordjournals.rpd.a031520, 1996.
- Matthewmann, N.J., Esler, J.G., Charlton-Perez, A.J., and Polvani, L.: A new look at stratospheric sudden warmings. Part III: polar vortex evolution and vertical structure. *J. Clim.*, 22, 1566-1585, doi:10.1175/2008JCLI2365.1
- Mitchell, D.M., Gray, L.J., and Anstey, J: The influence of stratospheric vortex displacements and splits on surface 725 climate, *J. Clim.*, 26, 2668-2682, doi:10.1175/JCLI-D-12-00030.1, 2013.
- Müller, M., Neuber, R., Fierli, F., Hauchecorne, A., Vömel, H., and Oltmans, S. J.: Stratospheric water vapour as tracer for vortex filamentation in the Arctic winter 2002/2003, *Atmos. Chem. Phys.*, 3, 4393–4410, doi:10.5194/acp-3-1991-2003, 2003.
- Müller, R., Tilmes, S., Groöf, J.-U., Engel, A., Oelhaf, H., Wetzell, G., Huret, N., Pirre, M., Catoire, V., Toon, G., 730 Nakjima, H., 2007. Impact of mesospheric intrusions on ozone-tracer relations in the stratospheric polar vortex. *J. Geophys. Res.*, 112(D23), doi:10.1029/2006JD008315

- Muñoz, C., Schultz, D., and Vaughan, G.: A midlatitude climatology and interannual variability of 200- and 500-hPa cutoff lows. *J. Clim.*, 33, 2201-2222, doi:10.1175/JCLI-D-19-0497, 2020.
- NCEP/NWS/NOAA/U.S. Department of Commerce: NCEP FNL Operational Model Global Tropospheric
735 Analyses, continuing from July 1999. Research Data Archive at the National Center for Atmospheric Research. Computational and Information Systems Laboratory, doi:10.5065/D6M043C6, Updated daily, Accessed July 2021, 2000.
- Paatero, J., Hatakka, J., and Viisanen, Y.: Trajectory Analysis of ^{210}Pb and ^7Be in Ground-Level Air in Southern Finland. *Radiochemistry*, 43, 475–481, doi:10.1023/A:1013069206962, 2001.
- 740 Pacini, A.A., Usoskin, I. G., Mursula, K., Echer, E., and Evangelista, H.: Signature of a sudden stratospheric warming in the near-ground ^7Be flux, *Atmos. Environ.*, 113, 27–31, doi:10.1016/j.atmosenv.2015.04.065, 2015.
- Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M. and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: Implications for policy, *J. Geophys. Res. Atmos.*, 109(15), doi:10.1029/2003JD00473, 2004.
- 745 Peethani, S., Sharma, N., and Pathakoti, M.: Effect of tropospheric and stratospheric temperatures on tropopause height, *Remote Sens. Lett.*, 5, 11, 933-940, doi:10.1080/2150704X.2014.973078, 2014.
- Peters, D.H.W., Vargin, P., Gabriel, A., Tsvetkova, N., and Yushkov, V.: Tropospheric forcing of the boreal polar vortex splitting in January 2003, *Annales Geophysicae*, 28, 2133–2148, doi:10.5194/angeo-28-2133-2010, 2010.
- Piñero García, F., Ferro García, M. A., and Azahra, M.: ^7Be behaviour in the atmosphere of the city of Granada
750 January 2005 to December 2009, *Atmos. Environ.*, 47, 84–91, doi:10.1016/j.atmosenv.2011.11.034, 2012.
- Pinheiro, H.R., Hodges, K.I., Gan, M.A., Ferreira, N.J.: A new perspective of the climatological features of upper-level cut-off lows in the Southern Hemisphere. *Clim. Dyn.*, 48, 541-559, doi:10.1007/s00382-016-3093-8
- Poluianov, S.V., Kovaltsov, G.A., Mishev, A.L., Usoskin, I.G.: Production of cosmogenic isotopes ^7Be , ^{10}Be , ^{14}C , ^{22}Na , and ^{36}Cl in the atmosphere: altitudinal profiles of yield functions, *J. Geophys. Res.*, 121(13), 8125-8136,
755 doi:10.1002/2016JD025034, 2016.
- Portmann, R., Sprenger, M., and Wernli, H.: The three-dimensional life cycle of potential vorticity cutoffs: a global ERA-interim climatology (1979-2017). *Weather Clim. Dynam. Discuss.*, doi:10.5194/wcd-2020-030, in review, 2020.
- Potgieter, M.S.: Solar modulation of cosmic rays, *Living Rev. Sol. Phys.*, 10(3), doi:10.12942/lrsp-2013-3, 2013.
- 760 Putero, D., Cristofanelli, P., Sprenger, M., Škerlak, B., Tositti, L., and Bonasoni, P.: STEFLUX, a tool for investigating stratospheric intrusions: application to two WMO/GAW global stations, *Atmos. Chem. Phys.*, 16, 14203-14217, doi: 10.5194/acp-16-14203-2016, 2016

- Salminen-Paatero, S., Thölix, L., Kivi, R., and Paatero, J.: Nuclear contamination sources in surface air of Finnish Lapland in 1965–2011 studied by means of ^{137}Cs , ^{90}Sr , and total beta activity, *Environ. Sci. Pollut. Res.*, 26, 21511–21523, doi:10.1007/s11356-019-05451-0, 2019.
- 765
- Salvador, P., Artíñano, B., Querol, X., Alastuey, A.: A combined analysis of backward trajectories and aerosol chemistry to characterise long-range transport episodes of particulate matter: The Madrid air basin, a case study, *Sci Tot Environ.*, 390, 495-506, doi:10.1016/j.scitotenv.2007.10.052, 2008.
- Sangiorgi, M., Hernández Ceballos, M.A., Iurlaro, G., Cinelli, G., and De Cort, M.: 30 years of European Commission Radioactivity Environmental Monitoring data bank (REMdb) – an open door to boost environmental radioactivity research. *Earth Sys. Sci. Data*, 11, 589-601, doi:10.5194/essd-11-589-2019, 2019.
- 770
- Simpkins, G.: Tropics to stratosphere. *Nat. Clim. Change* 7, 624, doi:10.1038/nclimate3385, 2017.
- Sofieva, V.F., Kalakoski, N., Verronen, P.T., Päivärinta, S.-M., Kyrölä, E., Backman, L., and Tamminen, J.: Polar-night O_3 , NO_2 , and NO_3 distributions during sudden stratospheric warmings in 2003-2008 as seen by GOMOS/Envisat. *Atmos. Chem. Phys.* 12, 1051-1066, doi:10.5194/acp-12-1051-2012, 2012.
- 775
- Sonnemann, G.R., Grygalashvyly, M., and Berger, U.: Impact of a stratospheric warming event in January 2001 on the minor constituents in the MLT region calculated on the basis of a new 3D-model LIMA of the dynamics and chemistry of the middle atmosphere. *J. Atmos. Sol.-Terr. Phys.* 68(17), 2012-2025, doi:10-1016/j.jastp.2006.04.005, 2006.
- 780
- Stein, A.F., Draxler, R.R., Rolph, G.D., Stunder, B.J.B., Cohen, M.D., and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bull. Am. Meteorol. Soc.*, 96, 2059-2077, doi:10.1175/BAMS-D-14-00110.1, 2015.
- Sýkora, I., Holý, K., Jeskovský, M., Müllerov, M., Bulko, M., and Povinec, P. P.: Long-term variations of radionuclides in the Bratislava air, *J. Environ. Radioactiv.*, 166, 27–35, doi:10.1016/j.jenvrad.2016.03.004, 2017.
- 785
- Tao, M., Konopka, Ploeger, F., Grooß, J.-U., Müller, R., Volk, C.M., Walker, K.A., and Riese, M.: Impact of the 2009 major sudden stratospheric warming on the composition of the stratosphere. *Atmos. Chem. Phys.*, 15, 8695-8715, doi:10.5194/acp-15-8695-2015, 2015.
- Terzi, L., and Kalinowski, M. B.: World-wide seasonal variation of ^7Be related to large-scale atmospheric circulation dynamics, *J. Environ. Radioactiv.*, 178-179, 1–15, doi:10.1016/j.jenvrad.2017.06.031, 2017.
- 790
- Tilmes, S., Müller, R., Engel, A., Rex, M., Russell III, J.M., 2006. Chemical ozone loss in the Arctic and Antarctic stratosphere between 1992 and 2005. *Geophys. Res. Lett.* 33(20), doi:10.1029/2006GL026925

- Tositti, L., Hübener, S., Kanter, H.J., Ringer, W., Sandrini, S., and Tobler, L.: Intercomparison of sampling and measurement of ^7Be in air at four high-altitude locations in Europe, *Appl. Rad. Isot.*, 61, 1497-1502, doi:10.1016/j.apradiso.2004.04.003, 2004.
- 795 Tositti, L., Hübener, S., Kanter, H.J., Ringer, W., Sandrini, S., and Tobler, L.: Intercomparison of sampling and measurement of ^7Be in air at four high-altitude locations in Europe. *Appl. Radiat. Isot.*, 61(6), 1497-1502, doi:10.1016/j.apradiso.2004.04.003
- Tositti, L., Brattich, E., Cinelli, G., and Baldacci, D.: 12 years of ^7Be and ^{210}Pb in Mt. Cimone and their correlation with meteorological parameters, *Atmos. Environ.*, 87, 108-122, doi: 10.1016/j.atmosenv.2014.01.014, 2014.
- 800 Usoskin, I.G., and Kovaltsov, G.A.: Production of cosmogenic ^7Be isotope in the atmosphere: Full 3-D modeling, *J. Geophys. Res.*, 113, D12107, doi:10.1029/2007JD009725, 2008.
- Usoskin, I.G, Field, C.V., Schmidt, G.A., Leppänen, A.-P., Aldahan, A., Kovaltsov, G.A., Possnert, G., and Ungar, R.K.: Short-term production and synoptic influences on atmospheric ^7Be concentrations, *J. Geophys. Res. Atmos.*, 114(D6), doi:10.1029/2008JD9011333, 2009.
- 805 Wang, Q., Jacob, D. J., Fisher, J. A., Mao, J., Leibensperger, E. M., Carouge, C. C., Le Sager, P., Kondo, Y., Jimenez, J. L., Cubison, M. J., Doherty, S. J., Sager, P. Le, Kondo, Y., Jimenez, J. L., Cubison, M. J., and Doherty, S.J.: Sources of carbonaceous aerosols and deposited black carbon in the Arctic in winter-spring: implications for radiative forcing, *Atmos. Chem. Phys.*, 11(23), 12453–12473, doi:10.5194/acp-11-12453-2011, 2011, 2011.
- Wargan, K., and Coy, L.: Strengthening of the tropopause inversion layer during the 2009 Sudden Stratospheric
- 810 Warming: a MERRA-2 study. *J. Atmos. Sci.*, 73(5), 1871-1887, doi:10.1175/JAS-D-15-0333.1
- Waugh, D.W., Sobel, A.H., Polvani, L.M.: What is the polar vortex and how does it influence weather? *Bull. Am. Meteorol. Soc.*, 98, 37-44, doi:10.1175/BAMS-D-15-00212.1, 2017.
- Webber, W.R., Higbie, P.R., and McCracken, K.G.: Production of the cosmogenic isotopes ^3H , ^7Be , ^{10}Be , and ^{36}Cl in the Earth's atmosphere by solar and galactic cosmic rays. *J. Geophys. Res.*, 112, A10, doi:
- 815 10.1029/2007JA012499, 2007.
- Wesley, M. L.: Parameterization of surface resistance to gaseous dry deposition in regional numerical models, *Atmos. Environ.*, 16, 1293-1304, 1989.
- Wu, S., Mickley, L. J., Jacob, D. J., Logan, J. A., Yantosca, R. M. and Rind, D. Why are there large differences between models in global budgets of tropospheric ozone? *J. Geophys. Res. Atmos.*, 112(D5), 1–18,
- 820 doi:10.1029/2006JD007801, 2007.

Yu, K., Keller, C. A., Jacob, D. J., Molod, A. M., Eastham, S. D. and Long, M. S.: Errors and improvements in the use of archived meteorological data for chemical transport modeling: an analysis using GEOS-Chem v11-01 driven by GEOS-5 meteorology, *Geosci. Model Dev.*, 11, 305–319, doi:10.5194/gmd-2017-125, 2018.

825 Zanis, P., Schuepbach, E., Gäggeler, H.W., Hübener, S., and Tobler, L.: Factors controlling beryllium-7 at Jungfrauoch in Switzerland, *Tellus*, 51 (4), 789e805, doi:10.1034/j.1600-0889.1999.t01-3-00004.x, 1999.

Zhang, B., H. Liu, J.H. Crawford, G. Chen, T.D. Fairlie, S. Chambers, C.-H. Kang, A.G. Williams, K. Zhang, D.B. Considine, M. P. Sulprizio, and R.M. Yantosca: Simulation of radon-222 with the GEOS-Chem global model: Emissions, seasonality, and convective transport, *Atmos. Chem. Phys. Discuss.*, in review, doi:10.5194/acp-2020-804, 2020.

830

Tables

835 **Table 1.** (left) Normalized differences between the MERRA-2 and observed precipitation, calculated as differences between the MERRA-2 and the observed values, normalized over the observed value, at each sampling site. Positive values indicate that the model tends to overestimate observations while the negative ones indicate underestimation. (right) Statistical parameters (mean \pm *SD* = mean \pm standard deviation; *MB* = mean bias; *NMSE* = normalized mean square error; *R* = correlation coefficient; *FA2* = factor of 2x) indicating the model performance in reproducing GPCP monthly accumulated precipitation at the six sampling sites in Northern Europe.

Sampling site	Normalized differences			Statistical parameters					
	Jan-03	Feb-03	Mar-03	Mean \pm <i>SD</i> (mm)		<i>MB</i> (mm)	<i>NMSE</i>	<i>R</i>	<i>FA2</i>
				MERRA-2	GPCP				
Ivalo	-0.07	0.63	1.25	2.23 \pm 0.67	1.48 \pm 0.43	0.75	0.34	-0.32	1
Umeå	-0.03	0.07	0.08	1.53 \pm 0.86	1.53 \pm 0.94	0.02	0	1.00	1
Helsinki	-0.09	-0.21	0.17	1.23 \pm 0.92	1.32 \pm 1.03	-0.09	0.01	0.99	1
Kista	0.96	-0.03	1.73	1.12 \pm 0.92	0.67 \pm 0.46	0.46	0.42	0.88	0.67
Harku	-0.2	0.18	-0.07	1.23 \pm 0.92	1.42 \pm 1.28	-0.18	0.05	1.00	1
Risø	-0.06	2.45	1.52	1.63 \pm 0.37	1.05 \pm 0.98	0.58	0.3	0.99	0.33

840

845 **Table 2. (left)** Normalized differences between the simulated and observed ^7Be monthly means, calculated as differences between the simulated and the observed values, normalized over the observed value, at each sampling site. (right) Statistical parameters (mean \pm SD = mean \pm standard deviation; MB = mean bias; $NMSE$ = Normalized Mean Square Error; $FA2$ = Factor of 2) indicating the model performance in reproducing observed ^7Be monthly means at the six sampling sites in Northern Europe and $^7\text{Be}/^{210}\text{Pb}$ and ^{210}Pb monthly means in Helsinki.

Sampling Site	Tracer	Normalized differences			Statistical parameters				
		Jan-03	Feb-03	Mar-03	Mean \pm SD		MB	$NMSE$	$FA2$
					Modelled	Observed			
Ivalo	^7Be	0.15	0.08	-0.07	(1.82 \pm 0.49) mBq m ⁻³	(1.85 \pm 0.78) mBq m ⁻³	-0.03 mBq m ⁻³	0.07	1
Umeå		-0.01	-0.14	-0.05	(1.69 \pm 0.72) mBq m ⁻³	(1.88 \pm 0.96) mBq m ⁻³	1.69 mBq m ⁻³	0.89	1
Helsinki		-0.22	-0.31	-0.2	(1.58 \pm 0.80) mBq m ⁻³	(2.30 \pm 0.60) mBq m ⁻³	1.57 mBq m ⁻³	0.76	0.74
Kista		-0.28	-0.35	-0.25	(1.68 \pm 0.69) mBq m ⁻³	(2.41 \pm 0.89) mBq m ⁻³	-0.73 mBq m ⁻³	0.16	0.92
Harku		-0.21	-0.13	-0.28	(1.61 \pm 0.61) mBq m ⁻³	(2.16 \pm 0.81) mBq m ⁻³	-0.54 mBq m ⁻³	0.17	0.93
Risø		-0.32	-0.4	-0.08	(2.08 \pm 0.83) mBq m ⁻³	(3.31 \pm 1.52) mBq m ⁻³	2.08 mBq m ⁻³	0.7	0.62
Helsinki	$^7\text{Be}/^{210}\text{Pb}$	-0.28	-0.5	-0.5	4.89 \pm 3.57	10.4 \pm 7.5	-5.04	0.78	0.45
Helsinki	^{210}Pb	0.73	0.5	0.66	(0.48 \pm 0.29) mBq m ⁻³	(0.36 \pm 0.32) mBq m ⁻³	0.12 mBq m ⁻³	0.3	0.78


855 **Table 3.** Statistical parameters indicating the model performance in reproducing observed ^7Be weekly (daily in the case of Helsinki) means at the six sampling sites in Northern Europe. The asterisk aside the value of the correlation coefficient and of the Z indicates that the value is statistically significant at the 0.05 significance level.

Sampling site	Tracer	Mean \pm SD		<i>MB</i>	<i>NMSE</i>	<i>R</i>	<i>FA2</i>	<i>Z</i>
		Modelled	Observed					
Ivalo	^7Be	(1.52 \pm 0.44) mBq m ⁻³	(1.85 \pm 0.78) mBq m ⁻³	-0.33 mBq m ⁻³	0.12	0.73*	1.00	-0.13
Umeå		(1.43 \pm 0.72) mBq m ⁻³	(1.88 \pm 0.96) mBq m ⁻³	-0.45 mBq m ⁻³	0.11	0.88*	0.92	-0.61
Helsinki		(1.35 \pm 0.83) mBq m ⁻³	(2.30 \pm 1.15) mBq m ⁻³	-0.88 mBq m ⁻³	0.47	0.65*	0.57	-5.69*
Kista		(1.43 \pm 0.62) mBq m ⁻³	(2.41 \pm 0.89) mBq m ⁻³	-0.98 mBq m ⁻³	0.30	0.85*	0.62	-2.44*
Harku		(1.36 \pm 0.56) mBq m ⁻³	(2.16 \pm 0.81) mBq m ⁻³	-0.79 mBq m ⁻³	-0.45	0.68*	0.86	-2.09*
Risø		(1.84 \pm 0.89) mBq m ⁻³	(3.31 \pm 1.52) mBq m ⁻³	-1.47 mBq m ⁻³	0.77	0.19	0.38	-2.68*
Helsinki	$^7\text{Be}/^{210}\text{Pb}$	4.86 \pm 3.96	10.3 \pm 7.5	-5.06	0.85	0.77*	0.43	-16.1

860

865


Figures




870

Figure 1. a) Location of the ⁷Be sampling sites in Northern Europe (source: <https://mapamundiparaimprimir.com/europa/>); b) ⁷Be concentrations measured at six surface sampling sites in Northern Europe during the 2002/2003 boreal winter. Dashed lines indicate the 90th percentile reference line for each sampling site.


Total Precipitation, 2003



875 **Figure 2.** Comparison of the MERRA-2 total precipitation during January-March 2003 with the GPCP observations. The black dots indicate the locations of the sampling sites: 1=Ivalo, 2=Umeå, 3= Helsinki, 4=Kista, 5=Harku, 6=Risø.



880 **Figure 3.** Simulated monthly mean ${}^7\text{Be}$ surface concentrations (mBq m^{-3}), ${}^7\text{Be}/{}^{210}\text{Pb}$ ratio and fraction of stratospheric ${}^7\text{Be}$ for January-March 2003. Arrows represent winds in the MERRA-2 reanalysis. The dots indicate the locations of the sampling sites: 1=Ivalo, 2=Umeå, 3=Helsinki, 4=Kista, 5=Harku, 6=Risø.



885 **Figure 4.** Scatter plots of: a) simulated vs. observed ^7Be monthly means at the six sampling sites; b) simulated vs. observed ^7Be weekly (daily in the case of Helsinki) means at the six sampling sites. Also shown are the linear regression line and the 95% confidence and prediction bands around the linear fit.

890




Figure 5. Temporal evolution of simulated and observed ^7Be surface concentrations at the six sampling sites. Values are weekly (daily in the case of Helsinki) means.





Figure 6. Temporal evolution of the simulated a) ${}^7\text{Be}/{}^{210}\text{Pb}$ and b) fraction of stratospheric ${}^7\text{Be}$ (calculated as the ratio between the stratospheric ${}^7\text{Be}$ and the total ${}^7\text{Be}$ concentrations, in percentage) at the six sampling sites during January-March 2003. Values are weekly means.



905

Figure 7. Vertical profiles of air temperature in the MERRA-2 reanalysis (dotted line) and in the soundings at the Sodankylä station in Finland on selected days of low ^7Be values (top panels: 10 and 16 Feb 2003, 00 UTC), transition to high ^7Be values (middle panels: 20 Feb 2003, 00 UTC and 21 Feb 2003, 12 UTC) and high ^7Be values (bottom panels: 22 and 24 Feb 2003, 00 UTC) at the six sampling sites in Northern Europe.

910





Figure 8. a) Vertical profiles of ozone mixing ratios (ppbv) obtained by ozone soundings at the Sodankylä Arctic station during 4 different days in February 2003: 12, 19, 16 and 28 February 2003; b) Daily mean O₃ concentrations recorded at ground-based air quality stations located in Denmark (DK), Finland (FI), Sweden (SE) during January-March 2003..



920 **Figure 9.** MERRA-2 daily mean relative humidity (colors) and winds (arrows) at ground level during February 18-25, 2003. The dots indicate the locations of the sampling sites: 1=Ivalo, 2=Umeå, 3=Helsinki, 4=Kista, 5=Harku, 6=Risø.




Figure 10. Time-height cross-sections of calculated daily potential vorticity during the month of February 2003 at three latitudes: a) 63°N, b) 64.5°N, and c) 66°N along the 21°E meridian.





Figure 11. Time-height cross-sections of simulated hourly ^7Be concentrations (mBq SCM^{-1} where SCM stands for Standard Cubic Meter) during the month of February 2003 at the six sampling sites.



935 **Figure 12.** Time-height cross-sections of MERRA-2 3-hourly average vertical pressure velocity (ω , in Pa s^{-1}) during the month of February 2003 sampled at the six sampling sites. Red color indicates large positive ω values, corresponding to a descending motion, whereas blue color stands for large negative ω values, corresponding to a rising motion. Black lines are contours of ω values.

940





Figure 13. MERRA-2 daily mean vertical pressure velocity (omega) at 940 hPa during February 18-25, 2003. The dots indicate the locations of the sampling sites: 1=Ivalo, 2=Umeå, 3=Helsinki, 4=Kista, 5=Harku, 6=Risø.



945

Figure 14. Simulated daily mean fraction of ^7Be originating in the stratosphere (%) at 940 hPa. The dots indicate the locations of the sampling sites: 1=Ivalo, 2=Umeå, 3=Helsinki, 4=Kista, 5=Harku, 6=Risø.




Figure 15. Average trajectory cluster results (centroids) arriving in Ivalo, Harku and Risø at 1000 m for low ^7Be values (left: 3-16 February 2003) and high ^7Be values (right: 20-28 February 2003), respectively. The stations are ordered by latitude from top to bottom (coordinates of the receptor site are provided on the left of each plot). The right numbers between brackets in the centroids are the percentage of complete trajectories occurring in that cluster, and the left numbers are an identification number of the centroid. In each panel, the upper plot shows the average latitude-longitude plot of each trajectory cluster, while the bottom panel presents the time vs. trajectory average altitude plot.