

~~Organic~~ The impact of organic pollutants from tropical Indonesian peatland fires : ~~regional influences and its impact on the tropospheric and lower stratospheric~~ ozone composition

Simon Rosanka¹, Bruno Franco², Lieven Clarisse², Pierre-François Coheur², Andrea Pozzer³, Andreas Wahner¹, and Domenico Taraborrelli¹

¹Institute of Energy and Climate Research, IEK-8: Troposphere, Forschungszentrum Jülich GmbH, 52428 Jülich, Germany

²Spectroscopy, Quantum Chemistry and Atmospheric Remote Sensing (SQUARES), Université libre de Bruxelles (ULB), Brussels 1050, Belgium

³Atmospheric Chemistry Department, Max-Planck-Institute for Chemistry, 55128 Mainz, Germany

Correspondence: Simon Rosanka (s.rosanka@fz-juelich.de)

Abstract. The particularly strong dry season in Indonesia in 2015, caused by an exceptional strong El Niño, led to severe peatland fires resulting in high volatile organic compound (VOC) biomass burning emissions. At the same time, the developing Asian monsoon anticyclone (ASMA) and the general upward transport in the intertropical convergence zone (ITCZ) efficiently transported the resulting primary and secondary pollutants to the upper troposphere/lower stratosphere (UTLS). In this study, we assess the importance of these VOC emissions for the composition of the lower troposphere and the UTLS, and we investigate the effect of in-cloud oxygenated VOC (OVOC) oxidation during such a strong pollution event. This is achieved by performing multiple chemistry simulations using the global atmospheric model ECHAM/MESy (EMAC). By comparing modelled columns of the biomass burning marker hydrogen cyanide (HCN) and carbon monoxide (CO) to spaceborne measurements from the Infrared Atmospheric Sounding Interferometer (IASI), we find that EMAC properly captures the exceptional strength of the Indonesian fires.

In the lower troposphere, the increase in VOC levels is higher in Indonesia compared to other biomass burning regions. This has a direct impact on the oxidation capacity, resulting in the largest regional reduction in ~~hydroxyl radicals~~ the hydroxyl radical (OH) and nitrogen oxides (NO_x). ~~Even though~~ While an increase in ozone (O₃) is predicted close to the peatland fires, ~~particular high concentrations of phenols lead to an~~ simulated O₃ ~~depletion~~ decreases in eastern Indonesia ~~due to particularly high phenol concentrations~~. ~~By employing the detailed in-cloud OVOC oxidation scheme Jülich Aqueous-phase Mechanism of Organic Chemistry (JAMOC), we find that the predicted changes are dampened and that by ignoring these processes, global models tend to overestimate the impact of such extreme pollution events.~~ In the ASMA and the ITCZ, the upward transport leads to elevated VOC concentrations in the UTLS region/lower stratosphere, which results in ~~a depletion~~ the reduction of OH and NO_x and the increase of hydroperoxyl radical (HO₂). In addition, the degradation of VOC emissions from the Indonesian fires becomes a major source of lower stratospheric nitrate radicals (NO₃), which increase by up to 20 %. Enhanced phenol levels in the upper troposphere result in a 20 % increase in the contribution of O₃. ~~We find that this is caused by a high destruction of~~ destruction by phenoxy radicals ~~and by the increased formation of reservoir species, which~~

~~dampen the chemical production of~~ to the chemical destruction of O₃, which is predicted to be as large as 40 % of the total chemical O₃ loss in the UTLS. In the months following the fires, this loss propagates into the lower stratosphere and potentially contributes to the variability of lower stratospheric O₃ observed by satellite retrievals. The Indonesian peatland fires regularly occur during El Niño years and ~~contribute to the depletion of~~. ~~In the time period from 2001 to 2016~~the largest perturbations of radicals concentrations in the lower stratosphere are predicted for particular strong El Niño years. By activating the detailed in-cloud OVOC oxidation scheme Jülich Aqueous-phase Mechanism of Organic Chemistry (JAMOC), we find that the ~~lower stratospheric is reduced by about 0.38 and contributes to about 25 to the lower stratospheric reduction observed by remote sensing~~. ~~By not considering these processes, global models might not be able to reproduce this variability in lower stratospheric predicted changes are dampened~~. Global models that neglect in-cloud OVOC oxidation tend to overestimate the impact of such extreme pollution events on the atmospheric composition.

1 Introduction

Particularly strong Indonesian wildfires during the El Niño in 2015 led to severe air pollution and reduced visibility (Kim et al., 2015; Lee et al., 2017), resulting in increased morbidity and mortality (Marlier et al., 2013; Reddington et al., 2014; Crippa et al., 2016) in South-East Asia (SEA). In general, El Niño is a large-scale climate anomaly, which is characterised by significantly warmer eastern equatorial Pacific Ocean sea surface temperatures (Trenberth, 1997), resulting in a dry season in SEA (Weng et al., 2007). The very strong El Niño phase in 2015-2016, which is the third strongest on record (after 1997–1998 and 1982–1983, NOAA, 2020), led to a particularly strong dry season in Indonesia (Jiménez-Muñoz et al., 2016), ~~which started~~
in mid July and lasted until November (Field et al., 2016). In the past, much of the originally forested and moist peatland in Kalimantan and Sumatra has been drained and cleared during agricultural land management. In order to clear these forests, landscape fires are commonly used. Even small local fires in these regions during ~~non-El~~~~non-El~~ Niño years may induce particular strong biomass burning emissions. Gaveau et al. (2014) ~~estimate~~~~estimated~~ that a local one-week Indonesian biomass burning event in 2013 contributed to about 5-10 % of Indonesian's total greenhouse gas emissions in that year. The additional drying during El Niño years favours fires that burn deep down into the peat and can last for multiple weeks. Due to their long lifetimes, these fires spread and ignite new areas, which are not necessarily prone to biomass burning. Compared to ~~non-El~~~~non-El~~ Niño years, this results in strong biomass burning emissions from Indonesia (van der Werf et al., 2017). The underground conditions inherently determine smouldering fires, which are characterised by low combustion temperatures. In combination with the high carbon content of peat, ~~this~~ smouldering fires emit much larger amounts of non-CO₂ emissions from peatlands than from other fuels (Christian et al., 2003; Rein et al., 2009; Yu et al., 2010). A major fraction of these non-CO₂ emissions ~~are~~~~is~~ volatile organic compounds (VOCs), which comprise a large variety of species ~~influencing and can influence~~ atmospheric chemistry on a regional and global scale. In the atmosphere, VOCs mainly react with the hydroxyl radical (OH), ozone (O₃), and the nitrate radical (NO₃), or photodissociate. Their atmospheric lifetimes range from minutes to years. Figure 1 shows the dry matter burned (DMB) during the 2015 Indonesian fires along the distribution of the peatlands (indicated in blue). It

55 becomes evident that most of the areas influenced by biomass burning (e.g. Sumatra, Kalimantan) are covered with peatland, indicating that the 2015 Indonesian fires are characterised by high VOC emissions.

During the Indonesian biomass burning season, usually the Asian monsoon is ongoing such that a large anticyclone spanning from tropical to temperate regions (from about 10° N to 40° N) evolves. This ~~almost-stationary-globally-prevailing~~ semi-stationary large-scale meteorological pattern typically ~~extends-extents~~ from the Middle East to Asia in the upper tropo-
60 sphere and lower stratosphere (UTLS) (Basha et al., 2020). As a convective system, the Asian monsoon anticyclone (ASMA) acts as pollution pump facilitating a fast transport of surface emissions to the UTLS (Park et al., 2008; Randel et al., 2010; Lelieveld et al., 2018). Vogel et al. (2015) analysed the impact of different regions in Asia on the chemical composition of the 2012 ASMA by using a chemical Lagrangian model. They found that air masses from SEA contribute significantly to the com-
65 position of the anticyclone in the UTLS. In addition, the vertically convective transport in the Inter Tropical Convergence Zone (ITCZ) and in the south-eastern flank of the anticyclone carries air masses from SEA into the UTLS. Thus, even short-lived VOCs from Indonesian fires are transported into the UTLS and potentially affect the lower stratospheric composition.

The Asian monsoon is characterised by the frequent occurrence of clouds and precipitation, and it has been demonstrated that the ASMA has a higher water vapour content than ~~usual-other~~ meteorological systems (Fu et al., 2006). At the same time, the Madden-Julian Oscillation (MJO) leads to enhanced water vapour concentrations and precipitation over the Indian Ocean
70 and Indonesia (Zhang, 2013). Many oxygenated VOCs (OVOCs) have a high solubility and quickly partition and react in cloud droplets influencing radical concentrations and the atmospheric composition in general (Herrmann et al., 2015). Rosanka et al. (2020b) showed that the in-cloud OVOC oxidation has a significant impact on the predicted concentrations of VOCs, key oxidants, and O₃. In the past, global atmospheric chemistry models were not capable to represent this process explicitly nor in its full complexity (Ervens, 2015). However, the recently developed Jülich Aqueous-phase Mechanism of Organic
75 Chemistry (JAMOC, Rosanka et al., 2020c,b) comprises an advanced in-cloud OVOC oxidation scheme suitable to be used in the ECHAM/MESSy Atmospheric Chemistry (EMAC, Jöckel et al., 2010) model. This allows us to assess the importance of this in-cloud oxidation process during the VOC-dominated Indonesian peatland fires.

In this study, we therefore investigate the importance of biomass burning VOC emissions from the strong 2015 Indonesian peatland fires on ~~the~~ (1) the lower tropospheric composition, (2) the UTLS, and (3) the importance of in-cloud OVOC oxidation
80 in such an extreme pollution event, ~~and (3) the influence of the emitted VOCs on the UTLS. This is addressed by performing multiple global chemistry simulations using the ECHAM/MESSy Atmospheric Chemistry (EMAC, Sect. 2) model.~~ In addition to the 2015 fires, strong peatland fires frequently occur in Indonesia. Especially during El Niño years (in 2002–2003, 2004–2005, 2006–2007, 2009–2010, and 2014–2016), high emissions have been observed (van der Werf et al., 2017). There-
85 fore, the long-term impact of these periodically occurring events is additionally addressed. Globally, biomass burning is not limited to Indonesia and many regions are frequently affected. In each region, biomass burning varies in strength, frequency, the characteristics of the biomass burned, and the chemical background conditions. ~~Peatland only covers about 2.84 of the Earth's land mass (Xu et al., 2018) making equatorial Asia the region where most peatland is burned. Since non-peatland biomass burning emissions result in lower VOC emissions (Akagi et al., 2011), Indonesia is characterised by a unique emission footprint. In order to address the resulting differences to other biomass burning dominated regions, we compare the influence~~

90 to seven regions with high biomass burning emissions. Figure 2 and Table 2 provide an overview of each region. This approach also allows us to analyse the global impact of all biomass burning VOC emissions on the atmospheric composition. Figure 3 shows the relative contribution of each region to the 2015 total biomass burning emissions of VOCs and aromatics, a subgroup of VOCs. These emissions are calculated based on dry matter combustion rates from the Global Fire Assimilation System (GFAS) and emission factors from Akagi et al. (2011) (for further details see Sect. 2.1.2). Due to the unique emission footprint from the Indonesian fires, SEA contributes 30% of the total biomass burning VOC emissions and more than to the total aromatic emissions. The two northern regions Alaska (ALA) and North Asia (NAS), which are characterised by extratropical forest with organic soil, add significantly to the global VOC emissions from biomass burning, even though their contribution to the total burned mass is low. The highest total emissions of almost 40% originate from Central Africa (CAF) and South Africa (SAF). This is twice as much as from SEA but since mainly tropical forest and savanna are burned, the contribution to the total biomass burning attributed to aromatic emissions is only half, when compared to SEA. The two regions dominated by savanna, Central South America (CSA) and North Australia (NAU), contribute the least to the VOC emissions.

The importance of biomass burning VOC emissions is addressed by performing multiple global chemistry simulations using the ECHAM/MESy Atmospheric Chemistry (EMAC) model. In a first step, we therefore compare the Indonesian peatland fires to other biomass burning regions, focusing on their specific emission footprint (Sect. 2) model. In a first step, the ability of EMAC to represent biomass burning events is evaluated using hydrogen cyanide (HCN) and carbon monoxide (CO) satellite retrievals (Sect. 4). Afterwards the impact of the 2015 Indonesian peatland fires on the lower troposphere (Sect. 5) troposphere and the UTLS is analysed, focusing on VOCs hydrocarbons, oxygenated organics, nitrogen containing compounds, and key radicals. Some VOCs are known to be toxic of which some are emitted or formed from biomass burning. They might have a significant influence on the Indonesia's population, which is the World's fourth highest (United Nations, 2019). Therefore, a selection of toxic species is additionally analysed in this section in Sect. 7-5 and Sect. 6 discuss, respectively. In Sect. 7, the importance of in-cloud OVOC oxidation during this pollution event and the influence on the UTLS, respectively is addressed. Modelling uncertainties related to this study are discussed in Sect. 8 before drawing final conclusions (Sect. 9).

2 Modelling approach

This section provides an overview on the global model used in this study. The main focus is placed on the representation of atmospheric gas- and aqueous-phase chemistry, biogenic and biomass burning emissions, and the strategy to compare EMAC's prediction to simulated results with satellite retrievals (Sect. 2.1). Sect. 2.2 provides an overview of each simulation performed in this study.

2.1 EMAC

The ECHAM/MESy Atmospheric Chemistry (EMAC) model is a numerical chemistry and climate simulation system that includes submodels describing tropospheric and middle-atmosphere processes and their interaction with oceans, land, and human influences (Jöckel et al., 2010). It uses the second version of the Modular Earth Submodel System (MESSy2) to

link multi-institutional computer codes. The core atmospheric model is the 5th-generation European Centre Hamburg general circulation model (ECHAM5, Roeckner et al., 2006). Jöckel et al. (2010) ~~provide~~provided an update on all modelling components used. For the present study, we applied EMAC (ECHAM5 version 5.3.02, MESSy version 2.54.0) in the T106L90MA and T42L90MA resolution, i.e. with a spherical truncation of T106 and T42 (corresponding to a quadratic Gaussian grid of approximately 1.1° by 1.1° and 2.8° by 2.8°, respectively). By using this horizontal resolution, addressing the short-term implications for 2015–2016 as well as the long-term impact (2001–2016) on a global scale is still feasible while at the same time, the computational costs are affordable. For both resolutions, 90 (~~L90~~) vertical hybrid pressure levels up to 0.01 hPa ~~(are used focusing on the lower and middle atmosphere)~~are used(MA), representing tropospheric and stratospheric transport processes reasonably well (Jöckel et al., 2010). Thus, the impact on the troposphere and the UTLS can be addressed. A detailed discussion on the comparability of both resolutions is performed in Sect. 8.

2.1.1 Atmospheric chemistry

~~Within~~For this study, the gas- and aqueous-phase ~~chemistry is modelled~~chemical kinetics is integrated by two separate submodels. For the atmospheric gas-phase chemistry, the applied model setup comprised the submodel Module Efficiently Calculating the Chemistry of the Atmosphere (MECCA, Sander et al., 2019) using the gas-phase Mainz Organic Mechanism (MOM). MOM contains an extensive oxidation scheme for isoprene (Taraborrelli et al., 2009, 2012; Nölscher et al., 2014), monoterpenes (Hens et al., 2014), and aromatics (Cabrera-Perez et al., 2016) and is therefore capable ~~to represent of~~representing all the biomass burning VOCs considered in EMAC. In addition, comprehensive reactions schemes are considered for the modelling of the chemistry of NO_x (NO+NO₂), HO_x (OH+HO₂), CH₄ and anthropogenic ~~linear-aliphatic and~~aromatic hydrocarbons. VOCs are oxidised by OH, O₃, and NO₃, whereas peroxy radicals (RO₂) react with HO₂, NO_x, and NO₃, and undergo self- and cross-reactions (Sander et al., 2019). Isocyanic acid (HNCO) is a chemical constituent that is ~~heavily-primarily~~ emitted by biomass burning and potentially harmful to humans (Wang et al., 2007; Roberts et al., 2011; Leslie et al., 2019). In order to properly represent this toxic constituent within EMAC, MOM has been extended to represent the atmospheric chemistry of HNCO. For this, the mechanism proposed by Rosanka et al. (2020d) is implemented into MOM. Their mechanism includes formamide as an additional chemical source of HNCO and chemical mechanisms for nitromethane, methylamine, dimethylamine, and trimethylamine.

The atmospheric aqueous-phase chemistry is modelled using the SCAVenging submodel (SCAV, Tost et al., 2006). It simulates the removal of trace gases and aerosol particles by clouds and precipitation. SCAV calculates the transfer of species into and out of rain and cloud droplets using the Henry's law equilibrium, acid dissociation equilibria, oxidation-reduction reactions, heterogeneous reactions on droplet surfaces, and aqueous-phase photolysis reactions (Tost et al., 2006). As mentioned earlier and as demonstrated by Rosanka et al. (2020b), in-cloud OVOC oxidation significantly influences the atmospheric composition. However, the ordinary differential equations (ODE) systems resulting from the combination of gas-phase and in-cloud aqueous-phase suffer from (1) a higher stiffness due to fast acid-base equilibria and phase-transfer reactions, and (2) load imbalance on High-Performance Computing (HPC) systems due to the sparsity of clouds. This leads to a significant increase in computational costs when using larger chemical mechanisms like the Jülich Aqueous-phase Mechanism of Organic Chemistry

(JAMOC), i.e. larger ODE systems (Rosanka et al., 2020c). Using JAMOC in each simulation performed in this study is thus not feasible. As a trade-off, JAMOC is used in a simulation subset in order to address and estimate its implications on the other simulations. Thus, two different aqueous-phase mechanisms are used within this study: (1) the standard aqueous-phase mechanism of EMAC (in the following called ScSta), which includes a detailed oxidation scheme and represents more than 150 reactions (Jöckel et al., 2016), and (2) JAMOC (Rosanka et al., 2020c), which includes a complex in-cloud OVOC oxidation scheme. In JAMOC, the phase transfer of species containing up to ten carbon atoms and the oxidation of species containing up to four carbon atoms are represented. Similar to MOM, both aqueous-phase mechanisms are modified to include the changes proposed by Rosanka et al. (2020d) to properly represent HNCO.

2.1.2 Biogenic and Biomass Burning VOC Emissions

165 In the atmosphere, biogenic and biomass burning emissions are ~~the dominant two major~~ sources of VOCs. The largest biogenic emissions take place in the equatorial region (e.g. Amazon Basin, Central Africa) with additional emissions in the Northern (NH) and Southern Hemisphere (SH) extratropics. The MESSy submodel ~~uses the~~ Model of Emissions of Gases and Aerosols from Nature (MEGAN, Guenther et al., 2006) ~~is used~~ to calculate biogenic VOC emissions. The global emissions of isoprene, the most abundant biogenic VOC, are scaled to 595 Tg a^{-1} , the best estimate of Sindelarova et al. (2014).

170 Biomass burning emission fluxes are calculated using the MESSy submodel BIOBURN, which determines these fluxes based on biomass burning emission factors and dry matter combustion rates. For the latter, data from the ~~GFAS Global Fire Assimilation System (GFAS)~~ are used that are based on satellite observations of the fire radiative power obtained from the Moderate Resolution Imaging Spectroradiometer (MODIS) satellite instruments (Kaiser et al., 2012). In BIOBURN, the emission strength depends on the dominant ~~fire-fuel~~ type in the respective area. From the GFAS dataset used in EMAC, in 2015, 175 the dominant ~~fire-fuel~~ type over Indonesia is tropical forest fire. However, as discussed earlier, peatland fires contribute substantially to the Indonesian fires. The GFAS dataset of EMAC is changed such that the dominant ~~fire-fuel~~ type over Indonesia is a combination of peat and tropical forest fires with equal contributions (following van der Werf et al., 2017). In general, biomass burning emission factors for VOCs are based on Akagi et al. (2011). Biomass burning emissions for HNCO, formamide, nitromethane, methylamine, dimethylamine, and trimethylamine are implemented following Rosanka et al. (2020d) 180 using emission factors from Koss et al. (2018) for HNCO and formamide.

2.1.3 Observational comparison

The evaluation of model simulation results against global observational datasets of VOC abundance can be performed for only a few species, mainly because of the limited availability in spaceborne measurements of such compounds. Among them, several VOCs are retrieved globally from the observations made by the nadir-viewing hyperspectral Infrared Atmospheric Sounding Interferometer (IASI, Clerbaux et al., 2009). Embarked on the Metop platforms on sun-synchronous polar orbits, IASI crosses the equator at 9:30 and 21:30 local solar time and achieves a global coverage twice daily with a fairly dense spatial sampling. ~~Although significant enhancements of carbon monoxide (CO) and ammonia (NH₃) have already been captured by the IASI measurements in the 2015 Indonesian fires (Whitburn et al., 2016b; Neechita-Banda et al., 2018), here~~ ~~Here~~, we make use

of the HCN abundance retrieved from the IASI/Metop–A and –B observations to assess the ability of EMAC to represent such an important biomass burning event. ~~This choice is explained in Sect. 4.~~ In addition, IASI methanol (CH_3OH) data are used to assess the impact of in-cloud OVOC oxidation in the model simulations (Sect. 7).

The retrieval method used to obtain the HCN measurements from the IASI observations follows closely the version 3 of the Artificial Neural Network for IASI (ANNI), which already allowed the retrieval of a suite of VOCs, including CH_3OH (Franco et al., 2018). ANNI is a general retrieval framework that consists in quantifying, for each IASI observation, the spectral signature of the target gas with a sensitive hyperspectral metric, and in converting this metric into gas total column via an artificial feedforward neural network (NN). Details on the ANNI retrieval approach, the HCN retrieval specificities, and the HCN product itself are provided in Appendix A. We refer to Franco et al. (2018) for a description of the IASI methanol retrievals. The satellite datasets exploited in this study consist of daily global distributions of HCN and CH_3OH total columns derived from the daytime observations (approximately 9:30 *a.m.*, local time) of the IASI/Metop–A and –B overpasses. These offer a better measurement sensitivity than the evening overpasses (Franco et al., 2018). Scenes affected by clouds or poor retrieval performance are removed from the final dataset by specific filters. Examples of daily regional distributions of HCN columns in the 2015 Indonesian fires as well as the seasonal global distributions of HCN as retrieved from IASI are presented in Appendix A. Those highlight the ability of IASI to capture the enhancements of HCN during biomass burning events as well as its downwind transport over long distances.

Significant enhancements of carbon monoxide (CO) have already been captured by IASI in the 2015 Indonesian fires (e.g., Whitburn et al., 2016b; Nechita-Banda et al., 2018). Therefore, we also evaluate the ability of EMAC to reproduce the CO columns observed from space during this event. The vertical profile and column abundance of CO are obtained in near real time from the IASI/Metop–A and –B spectra with the Fast Optimal Retrievals on Layers for IASI (FORLI) algorithm (Hurtmans et al., 2012). Several quality flags ensure that IASI observations affected by clouds, unstable retrieval and too weak measurement sensitivity are excluded from the final CO dataset. The FORLI algorithm, characterization of the retrieved CO product, and validation against independent measurements are reported in Hurtmans et al. (2012) and George et al. (2015). Following the formalism of Rodgers (2000), the IASI averaging kernels are applied to the CO model profiles to account for the inhomogeneous vertical sensitivity of the IASI measurements and to compute modelled CO columns as would be seen by the satellite instrument (see, e.g., Sect. 5.1 in Schultz et al., 2018).

2.2 Simulations performed

Within this study, seven simulations are performed, which can be summarised in three simulation sets. Each simulation differs either in the biomass burning emissions, the aqueous-phase mechanism used, or the modelled time period. Table 1 provides an overview of all simulations and their characteristics. For each simulation set, ~~one simulation exists, in which~~ in one simulation all VOC emissions from biomass burning are switched off (named REF and REF_{LONG}). A second simulation includes biomass burning VOC emissions as described in Sect. 2.1.2 (named FIR and FIR_{LONG}). Performing high resolution simulations with the highest complexity in the chemical mechanisms in EMAC comes with high computational costs. The strong Indonesian peatland fires of 2015 and the following year are selected as a specific case study (named REF and FIR). For both simulations,

the year 2014 is simulated as spin up, which is not considered for the analysis. For this case study, high resolution simulations are performed at T106L90MA. In order to isolate the impact of the Indonesian peatland fires in 2015, an additional simulation (named FIR_{NOINDO}) is performed, for which all biomass burning VOC emissions from Indonesia are switched off. In order to address the impact of in-cloud OVOC oxidation on such VOC-dominated pollution event, two simulations including JAMOC are performed (named REF_{JAMOC} and FIR_{JAMOC}). However, to reduce the computational demand (see Sect. 2.1.1), these simulations focus only on the second half of 2015 at T106L90MA resolution. The long-term effect of reoccurring Indonesian peatland fires are addressed by performing two long simulations for the time period of 2001–2016 (named REF_{LONG} and FIR_{LONG}). Here, the year 2000 is simulated for spin up, which is not used for the analysis. Performing these simulations at T106L90MA and using JAMOC is computationally not feasible. Therefore, the EMAC's standard aqueous-phase mechanism is used and the resolution is reduced to T42L90MA. All simulations are performed using the Chemistry-Transport Model mode (QCTM mode, Deckert et al., 2011), meaning that chemistry and dynamics are decoupled, e.g. fixed tracer mixing ratios are used as input for the radiation scheme instead of the prognostic chemical tracers. In this way, the meteorology is the same for all simulations and all changes in the atmospheric chemical composition predicted by EMAC are either due to the additional VOC emissions from biomass burning (when comparing REF with FIR or REF_{LONG} with FIR_{LONG}) or the in-cloud OVOC oxidation (when comparing REF, FIR, REF_{JAMOC} and FIR_{JAMOC}).

3 The representation of Peatland fires in Indonesia compared to biomass burning events in EMAC other regions

Globally, biomass burning frequently occurs in seven regions for which Fig. 2 and Table 2 provide an overview. In each region, biomass burning varies in strength, frequency, the characteristics of the biomass burned, and the chemical background conditions. Only about 2.84 % of the Earth's land mass is covered by peatland (Xu et al., 2018), making equatorial Asia the region where the most of peatland is burned. Since non-peatland biomass burning fuels have lower VOC (and higher NO_x) emission factors (Akagi et al., 2011), Indonesia is characterised by a unique emission footprint. Figure 3 shows the total, the VOC, and the aromatic emissions for each region in non-El Niño years, El Niño years, and in 2015 predicted by EMAC (based on REF_{LONG} and FIR_{LONG}). In non-El Niño years, the highest total emissions originate from Central Africa (CAF) and South Africa (SAF), whereas the lowest emissions occur in Alaska (ALA). Compared to SAF, SEA contributes only about one third. However, in El Niño years this almost doubles and in the exceptional strong year 2015 the emissions from SEA are almost the same as SAF. In CAF and SAF, mainly tropical forest and savanna are burned resulting in a significant lower contribution to the VOC emissions and aromatic emissions than SEA. The two northern regions ALA and North Asia (NAS), which are characterised by extratropical forest with organic soil, add significantly to the global VOC emissions from biomass burning, even though their contribution to the total emissions is low. The contribution of NAS to the total biomass burning is less than half of the contribution by SAF but its contribution to the aromatic emissions is almost the same. The contribution of ALA strongly depends on the El Niño. In non-El Niño years, the emissions are very low but increase in El Niño years. In the exceptional strong year 2015, the contribution of ALA to the aromatic emissions is of similar strength as from CAF even

255 though its total contribution is only one third when compared to CAF. The two regions dominated by savanna, Central South America (CSA) and North Australia (NAU), contribute little to the VOC emissions.

4 The representation of the Indonesian peatland fires in EMAC

In order to analyse the ability of EMAC to represent the Indonesian peatland fires, we compare predicted EMAC total columns of HCN and CO to observations obtained from IASI retrievals.

260 4.1 Comparison to IASI HCN retrievals

HCN mainly originates from combustion processes and is therefore largely emitted by biomass burning (Shim et al., 2007). Other emission sources including industrial activities, automobile exhaust, and domestic biofuel are assumed to be very weak (Lobert et al., 1990; Li et al., 2009). Reactions involving acetonitrile (CH_3CN) are the only gas-phase source of HCN, but those are estimated to be a minor contribution to the atmospheric HCN burden (Li et al., 2009). The slow oxidation
265 of HCN by OH and $\text{O}(^1\text{D})$ is considered to be the most important atmospheric gas-phase sink, leading to long chemical lifetimes (Cicerone and Zellner, 1983). However, due to a strong ocean uptake, the overall atmospheric lifetime is reduced to a few months (Li et al., 2000, 2009). The almost exclusive biomass burning source, combined with a long atmospheric residence time that allows for long-range transport, makes HCN a widely used primary tracer of biomass burning emissions and fire plumes (Li et al., 2009). ~~Other typical fire tracers such as and either have several other sources or are too~~
270 ~~short-lived to track fire plumes over long distance~~Moreover, substantial emissions of HCN are expected from strong peatland fires (e.g., Akagi et al., 2011; Andreae, 2019). Therefore, HCN satellite data from IASI are used here to evaluate the performance of EMAC in representing the 2015 Indonesian peatland fires.

~~Figure 4 shows the comparison of modelled total columns to IASI satellite retrievals for the three months with strong peatland emissions in Indonesia.~~ At the beginning of the Indonesian fires, the emitted HCN is transported westwards leading
275 to high HCN column values over the Indian Ocean (see Fig. 19). While the fires are ongoing throughout October, the strong westward transport of HCN results in the complete covering of the Indian ocean. Some HCN is also transported eastwards over Australia and the Pacific Ocean. In November, the air masses from Indonesia mix with emissions from Africa and the eastward transported air masses reach South America. Figure 4 shows the comparison of modelled HCN total columns to IASI satellite retrievals for the three months mean with strong peatland emissions in Indonesia. In general, EMAC strongly underestimates
280 HCN when its ~~biomass burning source~~ main source from biomass burning is not taken into account (simulation REF). Once the HCN biomass burning emissions are taken into account, the overall underprediction in EMAC is mostly resolved. ~~However in the simulation, but EMAC partially overpredicts HCN in SEA. Figure 5 gives the frequency of the global HCN is partially overpredicted in SEA during the main fire period (September and October). Additionally, EMAC total column bias in relation to the IASI retrievals during the Indonesian peatland fires, once including biomass burning emissions in the simulations and~~
285 ~~once not. This comparison clearly shows that HCN columns are slightly overestimated in CSA (September and October) and CAF (September). Interestingly, the relative model bias (not shown) is similar in all three regions. Due to the particular strength~~

of the Indonesian fires, the absolute bias is more pronounced in SEA. is strongly underestimated when its main source is not represented in EMAC. With HCN from biomass burning, the mean column bias reduces from -5.32×10^{15} molecules cm^{-2} to -1.06×10^{15} molecules cm^{-2} and its variance reduces from 1.75×10^{31} molecules² cm^{-4} to 2.57×10^{30} molecules² cm^{-4} , significantly improving the representation of HCN in EMAC.

In general, EMAC's representation of HCN is associated with some uncertainties. The occurrence of biomass burning events is very low in the Amazon Basin, suggesting that the high column values are not caused by fires or transport from CSA. Especially in NH autumn (SON), the Amazon basin is known to be highly influenced by biogenic emissions. Shim et al. (2007) already suggested that biogenic. Another important source of HCN emissions is terrestrial vegetation which may contribute to atmospheric concentrations by up to 18 % .In (Shim et al., 2007). In EMAC, the submodule MEGAN , biogenic emissions are taken into account and calculates that biogenic emissions contribute about 15 % to the total HCN emissions, suggesting that EMAC's overprediction is not caused by a misrepresentation of other sources. Additionally, Considering the particular high atmospheric concentrations in 2015, this slightly lower contribution suggests that this source strength may be well represented in EMAC. Overall, it is expected that the atmospheric lifetime of is reasonably well represented HCN atmospheric lifetime is realistically modelled, since globally HCN columns are well reproduced. Moreover, the ocean uptake accounts for 1.2 Tg(N) a^{-1} , which is well in the range of 1.1 to 2.6 Tg(N) a^{-1} proposed by Li et al. (2000) and very close to the Singh et al. (2003) estimate of 1.0 Tg(N) a^{-1} . The representation of biomass burning within EMAC depends on satellite observations (Sect. 2.1.2), which retrieve the fire radiative power and are thus sensitive to clouds. This introduces some uncertainties in regions that are characterised by the frequent occurrence of clouds, like equatorial Asia. Focusing on Indonesia, Liu et al. (2020) compared five different global fire inventories and found that GFAS, the inventory used in this study, represents the strength of these fires best. Still, GFAS even tends to slightly underestimate the strength, when compared to regional observations in Singapore, Malaysia, and Indonesia. This suggests that the magnitude of the Indonesian fires is well represented in EMAC. In this study However, from the literature a high uncertainty in the emission factors for HCN are reported. Here, we use the emission factors optimised for atmospheric models by Akagi et al. (2011), which suggest 5.0 g kg^{-1} for HCN from peatland fires. However, from the literature a high uncertainty in the emission factors for are reported. From recent field measurements in Indonesia and Malaysia, Stockwell et al. (2016) and Smith et al. (2018) report values ranging from 0.34 g kg^{-1} to 8.21 g kg^{-1} , whereas lab measurements for Indonesian peatland by Stockwell et al. (2015) suggest values between 3.30 g kg^{-1} and 3.83 g kg^{-1} . Overall, this results in a mean emission factor of 4.40 g kg^{-1} across all studies (Andreae, 2019), suggesting that some of EMAC's overestimation is caused by a the HCN emission factor used in EMAC is slightly too high, influencing EMAC's overprediction of HCN emission factor. Due to its long lifetime, is transported over long distances. West of Indonesia, EMAC also predicts higher columns than observed by IASI, suggesting columns. Lastely, the EMAC's overprediction west of Indonesia, suggests that some of the overprediction is caused by the deviation of horizontal transport (further discussed in Sect. 8). Figure 5 gives the frequency

4.2 Comparison to IASI CO retrievals

320 At the surface, CO is primarily emitted by natural and anthropogenic combustion processes like biomass and fossil fuel burning
and to a lesser extent by biogenic and oceanic sources. The degradation of methane (CH₄) and non-methane hydrocarbons
(NMHC) in the atmosphere accounts for almost half of the global EMAC total column bias in relation CO sources (Zheng et al., 2019)
. In the atmosphere, CO mainly reacts with OH and the EMAC estimates by Lelieveld et al. (2016) and more recently by
Rosanka et al. (2020b) show that CO largely determines the atmospheric oxidation capacity. To a lesser extent, CO is deposited
325 (Stein et al., 2014).

Figure 6a shows the total CO columns observed by IASI for the three months mean with strong peatland emissions
in Indonesia. Similar to HCN, high CO columns up to about 6.0×10^{18} molecules cm⁻² are observed over Indonesia.
Additionally, high CO columns are also observed in Africa and South America. Compared to HCN, CO is characterised
by a shorter lifetime. Therefore, less CO from Indonesia is transported towards Africa at the end of the peatland fire period.
330 Figure 6b shows the comparison of modelled total CO columns for FIR to the IASI retrievals during the Indonesian peatland
fires, once including biomass burning emissions in the simulations and once not. This comparison confirms that is strongly
underestimated when its main source is not represented in EMAC. With from biomass burning, the mean column bias reduces
from for the same period. Overall EMAC captures the spatial CO pattern with overprediction of 1.0 to and its variance reduces
from 1.75×10^{18} molecules cm⁻² in South America. Rosanka et al. (2020b) showed that EMAC predicts total methanol columns
335 too high in this region, which is related to EMAC's tendency to simulate the Amazon basin too dry in the dry season
(September-November) and consequently too hot (Hagemann and Stacke, 2015). This results in an overestimation of biogenic
VOC emissions in South America. Since VOC degradation is the main atmospheric CO source, their overprediction explains
EMAC's high bias for CO total columns in this region and its outflows.

As seen in Fig. 6b and 6c, EMAC constantly underestimate total CO columns over Indonesia during the main peatland fire
340 period. However, overall the model bias stays within a factor of two (dashed lines in Fig. 6c). This underprediction can be
explained by the emission factors used by EMAC. Stockwell et al. (2016) and Smith et al. (2018) report CO emission factors
ranging from 216 to 314 g kg⁻¹ obtained from observation in Indonesia and Malaysia during the Indonesian peatland fires of
2015. In addition, in the recent assessment of Andreae (2019) a mean emission factor of 260 to 2.57, significantly improving
the representation of in EMAC. g kg⁻¹ for peatland is reported across multiple studies, which is higher than the emission factor
345 used by EMAC (182 g kg⁻¹, Akagi et al., 2011).

From this analysis we conclude that even though EMAC does not reproduce HCN and CO columns perfectly, the Indonesian fires are reasonably well represented, especially when considering the exceptional strength of the 2015 Indonesian fires (for further discussion see Appendix A and Fig. 19). This also holds true considering all global biomass burning emission events.

5 The influence impact of biomass burning on the lower troposphere

350 In the following subsections, the impact of the 2015 Indonesian peatland fires on the lower tropospheric composition is analysed. In addition, substantial differences to the other six biomass burning dominated regions are discussed. All results are

based on the simulations REF and FIR. [Figure 7 depicts how the Indonesian peatland fires affect the atmospheric gas-phase composition.](#) Table 3 provides an overview on the global and regional changes (between simulation REF and FIR) in the tropospheric burden of each species discussed in the following subsections. The regional changes reported in Table 3 are calculated for the respective main biomass burning season defined in Table 2.

5.1 ~~Impact on VOCs~~ [Hydrocarbons](#)

Many VOCs are characterised by short lifetimes resulting in highly location-dependent changes within the troposphere. ~~Direct emissions are the only source of atmospheric hydrocarbons.~~ Globally, biomass burning emissions of VOCs significantly increase the atmospheric concentration of many hydrocarbons, ~~with acetylene (C₂H₂) and ethane (C₂H₆).~~ [In general, hydrocarbons can be separated into the aliphatic hydrocarbons and aromatic hydrocarbons. For both, direct emissions are the only atmospheric source.](#)

[Aliphatic hydrocarbons are further grouped into alkanes \(only single covalent bonds\), alkenes \(containing at least one C–C double bond\), and alkynes \(containing at least one C–C triple bond\). Ethane \(C₂H₆\) being the two hydrocarbons is globally the most abundant alkane and is impacted the most. Their tropospheric burden increases by 20.5% by biomass burning. Its global burden is increased by 32.6%, whereas the burden of less abundant alkanes like propane \(C₃H₈\) and *n*-butane \(C₄H₁₀\) only increases by 6.3% and 32.62.4%, respectively, with the highest regional change is.](#) Overall, the global change in the burden of alkenes is lower than of alkanes. Here, ethylene (C₂H₄) has the highest absolute change of 13.5 Gg (12.5%) followed by propene (C₃H₆) with 3.3 Gg (11.3%). Even though its abundance is the lowest, the highest global relative change of 20.2% is predicted for isobutene (C₄H₈). In addition, EMAC predicts an increase of 20.5% due to biomass burning emissions for the alkyne acetylene (C₂H₂). In general, the highest absolute change is predicted for SEA, except for propane since its burden increase is by 0.3 Gg higher in NAS than in SEA. In both cases, the relative change is very similar. For many aliphatic hydrocarbons, the lowest absolute changes are predicted in ALA. However, due to the generally low background concentrations in this area, the relative changes are the highest, making biomass burning in this region the major source of these hydrocarbons.

The two most abundant ~~aromatics~~ [aromatic hydrocarbons](#), benzene (C₆H₆) and toluene (C₆H₅CH₃), are strongly emitted by biomass burning events. In the FIR simulation, the tropospheric burden of [benzene](#) increases by 27.3%. Toluene has a slightly lower increase of only 15.3%. ~~The tropospheric concentration of oxygenated aromatics also increases strongly. The most dominant change is a higher relative impact is predicted for phenol (C₆H₅O), whose tropospheric burden is more than doubled and increases to 2.3. Even though phenol is directly emitted by biomass burning, the overall high aromatic emissions lead to an enhanced chemical production of phenol from benzene oxidation. The highest absolute change is observed in SEA.~~ However, due to low aromatic background concentrations, the relative increase is higher in ALA, CSA, and NAU ~~less abundant aromatics like ethylbenzene (C₈H₁₀) and styrene (C₈H₈). Here, the global burden changes by 65.9% and 29.8%, respectively. As it is for the aliphatic hydrocarbons, the highest absolute changes for benzene and toluene are predicted in SEA during the Indonesian peatland fires. Opposite to this, EMAC predicts the lowest change in SEA for ethylbenzene and styrene, which is related to the fact that EMAC's uses significant lower emissions for both aromatic hydrocarbons for peatland, when compared to the recent values reported by Andreae (2019).~~

~~Methanol (CH₃OH) is directly~~

5.2 Oxygenated organics

The degradation of aliphatic and aromatic hydrocarbons leads to the formation of oxygenated organic compounds. Additionally, they are emitted by biomass burning ~~emissions but also chemically produced from VOC oxidation~~ such as the Indonesian peatland fires. Globally, ~~methanol increases by 7.9%~~ biomass burning has only a little impact on formaldehyde (HCHO), the simplest aldehyde (R-CHO). However, regional changes are predicted to be higher and range from 2.2 to 35.1 %. The global and regional changes are higher for more complex aldehydes. The global burden of acetaldehyde (CH₃CHO) and glycolaldehyde (HOCH₂CHO) ~~increases by 11.4 % when biomass burning VOC emissions are taken into account. The high VOC emissions during the Indonesian peatland fires result in the highest changes and 8.0 %, respectively. In all cases, the highest absolute and relative change is predicted~~ in SEA. The two α -dicarbonyls glyoxal (OCHCHO) and methyl glyoxal (CH₃C(O)CHO) are primarily produced from VOC oxidation. ~~The global burden of glyoxal and methyl glyoxal~~ Their global burden increases by 9.3 % and 1.3 %, respectively. Again, the highest absolute changes are predicted in SEA. However, the highest relative change occurs in ALA due to generally low background VOC concentrations. Globally, methanol (CH₃OH) increases by 7.9 % when biomass burning VOC emissions are taken into account. Here, the Indonesian peatland fires contribute by far the most. A significant higher impact is predicted for 2,3-butanedione ((CH₃CO)₂). Its global burden is tripled due to biomass burning and the absolute changes predicted regionally are the highest in NSA and SEA.

In the atmosphere, organic acids are mainly produced from the photo-oxidation of biogenic and anthropogenic VOCs but may also be emitted from biomass burning. Formic acid (HCOOH) is slightly impacted by biomass burning VOC emissions and globally increases by 4.9 % with the highest changes in SEA and Africa (CAF and SAF). The acid impacted the most by biomass burning is acetic acid (CH₃COOH), which globally gains 23.3 % with the highest changes in SEA, CAF, CSA, and SAF. Interestingly, the high increase predicted in CSA only leads to a low relative rise. This is due to generally high background concentrations in this region from high biogenic VOC emissions.

The largest change in oxygenated aromatics is predicted for phenol (C₆H₅OH), whose tropospheric burden is more than doubled and increases to 2.3 Gg. Even though phenol is directly emitted by biomass burning, the overall high aromatic emissions lead to an enhanced chemical production of phenol from benzene oxidation. The highest absolute change is observed in SEA. However, due to low aromatic background concentrations, the relative increase is higher in ALA, CSA, and NAU. The increase in benzaldehyde (C₆H₅CHO) is significantly lower (globally by 14.6 %) with similar absolute changes in ALA, CAF, NAS, SAF, and SEA.

The oxidation of VOCs leads to the formation of CO (see Sect. 4.2). Overall, the VOC emissions from biomass burning only result in a global CO increase of 2.4 %, with regional changes between 2.4 % and 6.5 %.

5.3 ~~Impact on hydroxyl radicals (OH)~~ Nitrogen containing compounds

Besides looking at HCN, we also analysed the impact of the Indonesian peatland fires on two nitrogen-bearing compounds that are toxic for humans (isocyanic acid) and for vegetation (nitrophenols). Isocyanic acid (HNCO) is known to be a toxic

420 constituent of biomass burning emissions. It is linked to protein carbamylation, which causes adverse health effects such as rheumatoid arthritis, cardiovascular diseases, and cataracts (Wang et al., 2007; Roberts et al., 2011; Leslie et al., 2019). It is expected that the protein carbamylation potentially starts if humans are exposed to ambient concentrations above 1 ppb (Roberts et al., 2011). Rosanka et al. (2020d) already reported that HNC0 concentrations are high in regions characterised by strong biomass burning events. Globally, similar high concentrations are predicted in this study. However, we predict higher concentrations in Indonesia than Rosanka et al. (2020d), who reported that ambient HNC0 conditions of 1 ppb are exceeded 425 for less than 30 days in Indonesia in 2011. The year 2011 is known to have low biomass burning emissions in this region (van der Werf et al., 2017). Figure 8 shows the number of days, in which this threshold is exceeded during the 2015 Indonesian peatland fires between August and October. Here, 1 ppb of HNC0 is regularly exceeded and some regions are affected during the complete fire period. This causes potentially severe health effects for the population of Indonesia, which is the World's fourth highest (United Nations, 2019).

430 In the atmosphere, nitrophenols are mainly formed from the oxidation of the aromatic compounds benzene, toluene, phenols, and cresols (Nojima et al., 1975; Atkinson et al., 1980; Grosjean, 1984), of which the first three are emitted by biomass burning (see Sect. 5.1 and 5.2). Without biomass burning emissions of aromatics, the modelled nitrophenol concentrations are only high in regions with high anthropogenic activities (Fig. 9a). When biomass burning emissions of benzene, toluene, and phenols are included, nitrophenol concentrations significantly increase in areas affected by biomass burning. The strongest changes occur 435 in SEA, CAF, and SAF (Fig. 9b). Many biomass burning regions frequently exceed nitrophenol thresholds that are determined for regions where anthropogenic aromatic emissions dominate. On a global scale, biomass burning becomes the main source of nitrophenols. Nitrophenols are known to have a high phytotoxic activity that is prolonged given their photochemical stability (Grosjean, 1991). Rippen et al. (1987) and Natangelo et al. (1999) suggested that nitrophenols could have contributed to the forest decline in Northern and Central Europe in the 1980's but also in other parts of the world. Therefore, the overall increase 440 of nitrophenols in biomass burning areas is a potential danger for plants in these regions where plants are already under stressed conditions due to the biomass burning itself. At the same time, nitrophenols are known to absorb solar radiation (Hems and Abbatt, 2018) and therefore enhance hazy conditions in those areas (Lee et al., 2017), contributing to increased morbidity and mortality (Crippa et al., 2016).

5.4 Radicals


445 In general, organic molecules react with OH by either H-abstraction or addition to double bonds, making OH the most important daytime VOC oxidant. Figure 10a gives the mean tropospheric surface OH concentration in 2015 and Fig. 10b presents the changes due to biomass burning VOC emissions. OH concentrations are significantly reduced in most regions with frequent biomass burning events. This reduction is caused by the direct reaction of OH with VOCs, and the enhanced formation of CO from VOC degradation. The reduction in OH is not uniformly distributed and depends on the local chemical regime. In 450 Indonesia, the high VOC emissions lead to the highest absolute and relative OH reduction. The enhanced oxidation of VOCs by OH leads to an overall increase in HO₂. In ALA and NAS, the most northern areas of interest, the absolute change in OH is low (see Table 3). Within the biomass burning plume, the enhanced HO₂ concentrations react with NO producing OH and

compensating the OH reduction by VOC degradation, resulting in a regional surface OH increase. Still, outside the biomass burning plume, an overall decrease in OH is predicted in ALA and NAS. Here, VOCs from biomass burning become the highest OH sink resulting in strong relative changes in OH reactivity. In general, OH reactivity is the highest in the Amazon Basin (100 s⁻¹) and the lowest in Antarctica (0.5 s⁻¹). The additional VOC emissions in Indonesia result in a significant increase of about 50 % in the OH reactivity, which is similar to the increases predicted in ALA and NAS.

5.5 Impact on nitrogen oxides (–) and nitrate radicals (–)

Figures 11a and 11b show the mean surface NO_x concentrations and the changes induced by the VOC biomass burning emissions, respectively. The additional VOC emissions significantly reduce the regional concentrations in tropospheric NO_x. In SEA, the absolute changes are large but small in relative (about 8 %), whereas the highest absolute and relative NO_x changes are predicted in ALA. These reductions are caused by enhanced reactions of RO₂ with NO_x resulting in an increased formation of NO_x reservoir species (i.e. alkyl and acyl peroxy nitrates) and nitrogen-containing aromatics (e.g. nitrophenols).

NO₃ is the most important nighttime oxidant, which is globally increased by about 5 % when the biomass burning emissions of VOCs are included (see Table 3). On the one hand, the formation of NO₃ is enhanced from aromatic RO₂ reacting with NO₂, but on the other hand the loss of NO₃ by reactions with RO₂ and aldehydes is increased. In the two northern regions (ALA and NAS), the elevated O₃ and regionally increased NO₂ concentrations induce an enhanced formation from inorganic reactions, resulting in an additional rise of NO₃. The absolute increase in NO₃ is high in SEA, especially in Indonesia. Here, the particularly large increase in phenols results in enhanced concentrations of phenyl peroxy radicals (C₆H₅O₂), which form NO₃ when reacting with NO₂ following Jagiella and Zabel (2007):



(R1)

Taraborrelli et al. (2021) recently studied the importance of aromatics on the atmospheric composition on a global scale. They also demonstrated the importance of this reaction but in opposition to this study our findings, predicted a reduction of NO₃ in Indonesia. Taraborrelli et al. (2021) used a different resolution and analysed 2010, a year with little biomass burning emissions in Indonesia (van der Werf et al., 2017), reducing the importance of this production channel.

5.5 Impact on ozone (–)

The perturbed NO_x–HO_x relation consequently leads to changes in tropospheric O₃. Figure ??a shows the mean tropospheric column and Fig. ??b illustrates the changes induced by VOC biomass burning emissions. Overall, EMAC predicts an enhanced formation of O₃ when the biomass burning emissions are included (see Table 3). The increase in HO₂ leads to an enhanced chemical O₃ production by reacting with NO. Due to high NO_x emissions from biomass burning, the O₃ production is to a large extent VOC-limited. In the two northern regions, the background VOC concentrations are low resulting in the highest

relative changes of more than 10 %. ~~The same can be observed in Fig. ??b, which shows the zonal-mean changes in the tropospheric column from 2001 onward. The~~ Based on the long term simulations (REF_{LONG} and FIR_{LONG}) we find that the largest changes are predicted in the NH high latitudes in 2003, a year with intense biomass burning in Boreal Asia (van der Werf et al., 2017). However, compared to the averaged tropospheric background O₃ concentrations, these changes are negligible on a global scale.

As described in Sect. 4, most VOC emissions from Indonesia are transported towards the Indian Ocean. Therefore, O₃ is predicted to increase in Sumatra and west of it (Fig. ??b). Interestingly, away from biomass burning emissions in Kalimantan and in east Indonesia, O₃ concentrations are slightly reduced, even though the chemical O₃ production still increases in this area. The ~~decrease can also be observed in Fig. ??b during strong El Niño years, especially in 2006, 2009, 2014, and 2015.~~ The particularly strong emissions of aromatics lead to enhanced concentrations of phenoxy radicals (C₆H₅O), which directly ~~destruet~~ destroy O₃ (Tao and Li, 1999) in lower NO_x regions:



This O₃ sink increases by 780 % resulting in a net loss of O₃ in these areas. Globally, this O₃ destruction channel ~~gains~~ increases from 144.9 to 200.1 Tg a⁻¹ in the troposphere. Also, Taraborrelli et al. (2021) reported a similar strength of this destruction channel of about 200 Tg a⁻¹. Therefore, biomass burning emissions regionally control the importance of this destruction channel.

5.5 Pollution and toxic conditions

~~The direct emission and degradation of primarily emitted VOCs lead to the formation of toxic compounds that are of special interest in highly populated areas. One prominent example is nitrophenols, which are known to have a high phytotoxic activity that is enhanced by a high photochemical stability (Grosjean, 1991). Rippen et al. (1987) and Natangelo et al. (1999) suggested that nitrophenols could have contributed to the forest decline in Northern and Central Europe in-~~

6 The influence of Indonesian peatland fires on the UTLS

As illustrated in Fig. 7, some of the biomass burning VOC emissions from Indonesia are quickly transported, by the ASMA and the ~~80's but also in other parts of~~ general tropical updraft, into the UTLS (see Sect. 1 and Vogel et al., 2015). In the following, we define the lower stratosphere between 147-32 hPa (about 13–24 km) above 30° in latitudinal direction and between 100-32 hPa (about 17–24 km) below 30° latitude. In Table 4, the lower stratospheric burden in November for each discussed species is presented, including the changes induced by biomass burning and the contribution from the Indonesian peatland fires.

510 6.1 Hydrocarbons

Even though their atmospheric lifetime is generally short, the upward transport in the tropics leads to an increase of hydrocarbons in the lower stratosphere due to biomass burning. Similar to the changes in the troposphere, ~~the world. In the atmosphere, nitrophenols are mainly formed from the oxidation of the aromatic compounds benzene, toluene, phenols, and cresols (Nojima et al., 1975; Atkinson et al., 1980; Grosjean, 1984), of which the first three are emitted by biomass burning. Without aromatic biomass burning emissions, nitrophenol concentrations are only high in regions with high anthropogenic emissions (Fig. 9a)~~ aliphatic hydrocarbon with the highest absolute change is ethane. The lower stratospheric burden in November increases by about 41 % to 43.56 Gg. The lower stratospheric burden of other aliphatic hydrocarbons like propane and *n*-butane changes by around 30 %. The lower stratospheric burden of benzene is tripled whereas toluene is doubled, which is consistent with the difference between their chemical lifetimes. The contribution of the Indonesian peatland fires for most hydrocarbons ranges between 69 to 87 %, except ethane. We expect that additional non-Indonesian fires from SEA contribute the rest, since the Indonesian peatland fires contribute about 76 % to the total biomass burning emissions from SEA in 2015. In the case of ethane, the contribution from the Indonesian peatland fires is only about 24 %. Compared to other hydrocarbons, ethane has an atmospheric lifetime of about two months (Hodnebrog et al., 2018). Thus, we expect that its long lifetime allows ethane emitted from other biomass burning regions to be transported into the lower stratosphere. At the same time, the recent biomass burning inventory by Andreae (2019) indicates that EMAC underestimates ethane emissions from the Indonesian peatland fires by a factor of three.

6.2 Oxygenated organics

In addition to the upward transport of the directly emitted OVOCs, the elevated hydrocarbon concentrations also form OVOCs in the lower stratosphere. In the lower stratosphere, methanol is one of the most abundant OVOCs and its burden increases by 34 %. EMAC predicts a higher relative increase for less abundant OVOCs like glyoxal (about 49 %) and acetic acid (about 64 %). The high increase in benzene, due to the strong aromatic emissions from the Indonesian peatland fires (see Sect. 3), results in the particular large production of phenol. Here, the lower stratospheric burden increases by about 167 %. The contribution from the Indonesian peatland fires to the lower stratospheric burden of all OVOCs is in a similar range as for the hydrocarbons, namely from about 72 to 86 %. ~~When biomass burning emissions of benzene, toluene, and phenols are included, nitrophenol concentrations significantly increase in areas affected by biomass burning. The strongest changes occur in SEA, CAF, and SAF~~

6.3 Nitrogen containing compounds

The Indonesian peatland fires resulted in substantial HCN emissions, as seen in Fig. 9b). ~~Many biomass burning regions frequently exceed nitrophenol thresholds that are determined for regions, where anthropogenic aromatic emissions dominate. On a global scale, biomass burning becomes the main source of nitrophenols, which have been measured in rain droplets (Leuenberger et al., 1985; Schummer et al., 2009). JAMOC represents the phase transfer of some nitrophenols. However, their~~

reduction due to this additional sink is calculated to be below 14 and as discussed in Sect. 4.1, which results in a strong increase of HCN in the lower stratosphere. Here, EMAC predicts an increase of more than 36 Gg (about 220 %). This insignificant reduction results from the missing sink of these nitrophenols (Hems and Abbatt, 2018) in JAMOC). Sheese et al. (2017) report that the highest increase on record in lower stratospheric HCN was observed by the Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS) instrument on the SCISAT satellite following the 2015 Indonesian peatland fires. Therefore, the predicted nitrophenol concentrations are expected to be slightly overestimated. Still, the overall increase of nitrophenols in biomass burning areas is a potential danger for plants in these regions where plants are already under stressed conditions due to the biomass burning itself based on the long term simulations (simulation REF_{LONG} and FIR_{LONG}). EMAC reproduces their findings. The elevated lower stratospheric benzene and toluene concentrations lead to an increase in lower stratospheric nitrophenol concentrations of about 60 %.

6.4 Radicals

The oxidation of VOCs transported into the lower stratosphere influences the lower stratospheric oxidation capacity. Overall, the lower stratospheric OH burden is reduced by about 3 %, whereas the burden of HO₂ increases by 2.4 % (see Table 4). The enhanced formation of NO_x reservoir species results in a 6 % and 4.2 % reduction of NO and NO₂, respectively. At the same time, nitrophenols are known to absorb solar radiation (Hems and Abbatt, 2018) and therefore enhance hazy conditions in those areas (Lee et al., 2017), contributing to increased morbidity and mortality (Crippa et al., 2016).

Isoeyanic acid (i) is also known to be a toxic constituent of the enhanced reactions of NO₂ with aromatic RO₂ results in an increase of NO₃ of more than 9 %. Figure 12 provides the mean longitudinal relative change in lower stratospheric OH, HO₂, NO_x, and NO₃ between 2001 and 2016 based on the long-term simulations (simulation REF_{LONG} and FIR_{LONG}). After each Indonesian peatland fire period, the lower stratospheric oxidants are influenced. With decreasing lower stratospheric VOC concentrations over time, the influence on the oxidants vanishes in the second half of the following year. Particularly strong influences are observed during El Niño periods, caused by enhanced VOC emissions from peatland fires. For example, intense fires in 2006 and 2015 led to a significant change in lower stratospheric oxidants in early 2007 and 2016, respectively. In 2010, almost no fires occurred in Indonesia (van der Werf et al., 2017), resulting in only a little change in oxidant concentrations in 2011. Even in non-El Niño years, EMAC predicts changes in lower stratospheric radicals due to VOC biomass burning emissions. It is linked to protein carbamylation, which causes adverse health effects such as rheumatoid arthritis, cardiovascular diseases, and cataracts (Wang et al., 2007; Roberts et al., 2011; Leslie et al., 2019). It is expected that the protein carbamylation potentially starts if humans are exposed to ambient concentrations above 1 (Roberts et al., 2011). Rosanka et al. (2020d) already reported that concentrations are high in regions characterised by strong biomass burning events. Globally, similar high concentrations are predicted in this study. However, we predict higher concentrations in Indonesia than Rosanka et al. (2020d) who reported that ambient conditions of 1. We expect that these originate from other biomass burning events in SEA (see Sect. 6.1). In each year, the highest influence is predicted for NO₃. Following the intense fires of 2002, 2006, 2009, 2014, and 2015 the zonal mean NO₃ concentration changes by more than 20 are exceeded for less than 30 in Indonesia in 2011. The year 2011 is known to have low biomass burning emissions in % in the tropical and subtropical lower stratosphere at the end of the

same year and at the beginning of the next year. Thus, our findings indicate that VOC emissions from the Indonesian peatland fires quickly transported into the lower stratosphere become a major source of lower stratospheric NO_3 .

The elevated phenol and consequently phenoxy radicals in the UTLS influence the importance of the O_3 loss due to reaction with phenoxy radicals (Reaction R2). In the upper tropical troposphere this loss process contributes significantly to the total chemical O_3 loss. Figure 13a shows the zonal mean in the relative contribution of this O_3 loss pathway to the total chemical O_3 loss without VOC biomass burning emissions in November 2015. Especially in the upper northern tropical troposphere, this loss process contributes up to 40 % to the total chemical O_3 loss. Following the benzene emissions from biomass burning, elevated phenoxy radicals in the UTLS double the contribution of this region (van der Werf et al., 2017) O_3 loss processes in the upper southern tropical troposphere (i.e. an increase of more than 20 %) and increase the contribution in the upper northern tropical troposphere by about 10 % in November 2015 (see Fig. 13b). A similar impact can be observed following other intense Indonesian biomass burning seasons. Figure 13c shows the zonal mean change in the relative contribution of this O_3 loss pathway in the UTLS between 2001 and 2016. Here, we define the UTLS from 250 hPa to 50 hPa above the tropopause calculated by EMAC. Especially after the intense Indonesian peatland fires during strong El Niño periods, a change in the upper southern tropical UTLS of more than 10 % is predicted at the end of each year. In all years, the increase in the upper northern tropical UTLS is lower. Figure 8 shows the number of days, in which this threshold is exceeded during the 2015–14 shows the zonal mean relative change in the phenoxy radical O_3 loss pathway due to VOC biomass burning in April 2016. Following the increase of benzene in the lower stratosphere (Sect. 6.1), EMAC predicts an increase in this O_3 loss process by more than 400 % which propagates into the upper tropical lower stratosphere. These findings suggest that the frequently re-occurrence of strong Indonesian peatland fires. Here, 1 of is regularly exceeded and some regions are affected during the complete fire period. This causes potentially severe health effects for the population of Indonesia could contribute to the variability in lower stratospheric O_3 which is observed by remote sensing measurements (Kyrölä et al., 2013; Nair et al., 2015; Vigouroux et al., 2015; Chipperfield et al., 20

In the UTLS, aviation is the only direct anthropogenic activity and contributes about 3.5 % to the total anthropogenic climate change (Lee et al., 2021). Here, aviation NO_x emissions lead to a formation of O_3 and a depletion of methane (CH_4). Recently, Rosanka et al. (2020a) showed that the enhancement in O_3 is limited by the background concentrations of NO_x and HO_x . If enough HO_x is available, a lower background NO_x concentration results in a higher O_3 gain. In general, low background HO_x concentrations limit the O_3 gain in winter. In our study, we find (not shown) that in the North Atlantic flight sector (between 400–100 hPa), the NO_x burden is reduced due to VOC emissions from SEA fires by about 6 % with regional changes of more than 20 % in 2015. At the same time, HO_x increases regionally by 10 %. Even though NO_x emissions from the frequently occurring Indonesian peatland fires are expected to result in an increase in UTLS NO_x , substantial VOC emissions from the same fires potentially compensate the impact of the NO_x increase and favour the formation of O_3 from aviation activities. In the simulation setup used, EMAC neglects VOC emissions reported from aviation activity (e.g., Wilkerson et al., 2010). Our findings indicate that the direct emissions of benzene, toluene, and phenol in the UTLS potentially enhance the loss of O_3 by phenoxy radicals and consequently affect the impact of aviation on O_3 in that region.

The influence Recently, Rosanka et al. (2020b) showed that the in-cloud oxidation of OVOCs significantly influences tropospheric VOCs and oxidants. By using JAMOC (Rosanka et al., 2020c,b), we investigate the importance of in-cloud OVOC oxidation is addressed by applying JAMOC during the Indonesian fire period peatland fires on the troposphere and lower stratosphere (simulations REF_{JAMOC} and FIR_{JAMOC}). In order to isolate the influence of the Indonesian peatland fires from the back-ground changes induced by JAMOC, the changes from the Indonesian fires due to the in-cloud OVOC oxidation are calculated following:

$$\Delta\text{JAMOC} = (\text{FIR}_{\text{JAMOC}} - \text{FIR}) - (\text{REF}_{\text{JAMOC}} - \text{REF}) \quad (1)$$

7.1 On the lower troposphere

Figure 15 shows the changes in the zonal mean concentration over Indonesia and the Indian Ocean of all OVOCs explicitly reacting in JAMOC (\sum OVOCs, see Equation B1 in Appendix B) for the simulations without JAMOC (Fig. 15a) and the predicted changes due to JAMOC (Fig. 15b; calculated using Reaction 1), focusing on the Indonesian fire period (SON). Due to the high solubility of many OVOCs and their in-cloud oxidation, their concentration is strongly reduced at altitudes that are characterised by frequent cloud events. The Table 5 provides the SEA burden changes for a selection of species that are represented in JAMOC. Overall, the additional in-cloud sink for methanol, glyoxal, and methyl glyoxal leads to a lower increase in their burden ranging between 23 and 32 results in a more limited increase of their predicted burden (e.g., only about 87.9 Gg instead of 112.3 Gg for methanol). However, their predicted burden in the reference simulation (REF_{JAMOC}) is also significantly lower, which results in a relative change which is in a similar order as if no in-cloud OVOC oxidation is taken into account.

Figure 16 shows the Probability Density Function (PDF) for EMAC's methanol column bias when compared to IASI satellite retrievals (Franco et al., 2018) in SEA during the Indonesian peatland fires. Without VOC emissions from biomass burning, methanol is slightly underestimated by simulation REF. This underestimation is more pronounced when the in-cloud oxidation of OVOCs is taken into account (simulation REF_{JAMOC}). In both cases, EMAC tends to strongly underestimate methanol in some regions. When VOC biomass burning emissions are taken into account (simulation FIR), these underpredictions are resolved. However, now EMAC tends to strongly overestimate methanol mainly close to biomass burning sources (not shown). These overpredictions are reduced once in-cloud OVOC oxidation is implemented (simulation FIR_{JAMOC}). A high fraction of SEA is covered by oceans. Millet et al. (2008) suggested that some regions of the Pacific and Indian Ocean are a net source of methanol. As discussed by Rosanka et al. (2020b), EMAC represents the ocean as a net methanol sink. Therefore, when comparing the predictions of methanol from EMAC to satellite observations, a certain underestimation is expected. Thus, simulation FIR_{JAMOC} compares the best with IASI retrievals, since it has overall the lowest relative biases.

Changes in hydrocarbons are minimal due to their low solubility, whereas strong changes are predicted for the relative relatively insoluble O₃. Due to in-cloud OVOC oxidation, the initially predicted increase in O₃ in western Indonesia and over the Indian Ocean (Sect. 2.5.4) is dampened by more than 60 % once JAMOC is implemented. This reduced limited increase is

caused by the ~~increasing~~ importance of clouds as ~~an~~ O_3 ~~sinks~~~~sink~~. This process is globally analysed by Rosanka et al. (2020b) and is based on the enhanced HO_2 formation in cloud droplets by OVOC oxidation. Within clouds, HO_2 is in acid equilibrium with the superoxide anion (O_2^-), which actively destroys O_3 .

7.2 On the lower stratosphere

As seen in Fig. 15b, the in-cloud OVOC oxidation leads to the reduction of their concentrations in the UTLS. Table 6 presents the lower stratospheric burden changes in November due to JAMOC. Overall, the in-cloud oxidation of the OVOCs leads to a more limited increase of their concentration in the lower stratospheric burden induced by the biomass burning emissions. For example, the increase in the methanol burden is limited to 2.6 Gg (instead of 5.8 Gg). In the case of ethylene glycol ($HOCH_2CH_2OH$), the lower stratospheric burden decreases even by about 85 % when JAMOC is used, with and without VOC biomass burning emissions taken into account. Similarly to the changes in the lower stratosphere, the relative change for the simulation using JAMOC is in a similar order as for the simulation without VOC biomass burning emissions but tends to be slightly lower for some OVOCs. This is especially the case for isopropyl hydro peroxide ($(CH_3)_2CHOOH$), which is by about 23 % lower. The increase in the lower stratospheric phenol concentrations is only slightly impacted and decrease by about 13 % (from original 167 %, see Table 4) and thus has only little impact on the destruction of O_3 by phenoxy radicals. This is consistent with the main source of phenol being the oxidation of benzene, which has a lifetime in the order of 1-2 weeks.

To conclude, in-cloud OVOC oxidation is important to properly represent the resulting impacts from strong pollution events especially during the monsoon season. Overall, the predicted impact on VOCs, radicals, and O_3 is dampened by the in-cloud oxidation and models neglecting this process probably tend to overestimate the impact of such an event. It is widely recognised that clouds may act as a source of secondary organic aerosols (SOA) ~~,-which even enhances by due to~~ in-cloud oxidation processes (Blando and Turpin, 2000; Ervens et al., 2011; Ervens, 2015). Ervens et al. (2011) suggested that cloud processes might contribute ~~in the same order~~ to SOA formation on the same order as gas-phase processes. Within this study, SOA formation from cloud processes are not explicitly represented. However, it is expected that the enhanced VOC concentrations from biomass burning will lead to an increased SOA formation from aqueous-phase processes due to the enhanced formation of oligomers (e.g. from glyoxal and methyl glyoxal) within clouds.

8 ~~The influence of Indonesian peatland fires on the UTLS~~

~~Some of the biomass burning VOC emissions from SEA are quickly transported by the ASMA and the general tropical updraft into the UTLS (see Sect. 1 and Vogel et al., 2015), which significantly increases lower stratospheric VOC concentrations. The effect of this transport process can be observed in Fig. 15. For example, the highest increase in glyoxal, methanol, and phenol concentrations in the lower stratosphere (48.5, 33.1, and 149.4 , respectively) is predicted in November. In the following months, these VOCs actively react with and change the lower stratospheric radical chemistry. Overall, this results in a reduction in lower stratospheric peaking in April 2016, as shown in Fig. ??.~~ Especially in the tropics, lower stratospheric diminishes by

more than 12. We find that the substantial increase in phenols, caused by the high aromatic emissions from Indonesia, favours
675 the formation of phenoxy radicals that contribute the most to this depletion via Reaction R2. Under high conditions, many
VOCs form reservoirs (i.e. alkyl and acyl peroxy nitrates), which flattens the peak of the burden in the lower stratosphere by
7.5 and increases the burden by 3.3.

Figure ?? shows the zonal mean reduction in lower stratospheric based on the long-term simulations (simulation and) from
2001 onward. After each Indonesian peatland fire period, stratospheric is depleted. With decreasing lower stratospheric VOC
680 concentrations over time, slightly recovers in the second half of the following year. Particularly strong decreases are observed
during El Niño periods, caused by enhanced VOC emissions from peatland fires. For example, intense fires in 2006 led to a
significant decrease in lower stratospheric in early 2007. In 2010, almost no fires occurred in Indonesia (van der Werf et al., 2017)
, resulting in the partial recovery of lower stratospheric in 2011. Even in non El Niño years, VOC emissions from SEA biomass
burning events contribute to the depletion of lower stratospheric. Though of low intensity, the 2011 Indonesian peatland
685 fires led to a reduction in , which is comparable to the reduction during the 2009 El Niño year. In 2011, a strong La Niña
occurred, which is the colder counterpart of El Niño (NOAA, 2020). In general, La Niña strengthens whereas El Niño weakens
the ASMA. Additionally, the ASMA strengthens over time with a particular increase in 2011-2016, compared to 2001-2010
(Basha et al., 2020). Yuan et al. (2019) reported that the 2011 ASMA was stronger with a higher spatial extend. Additionally,
the 2011 Indonesian fires occurred earlier during the strongest phase of the Asian monsoon. The combination of earlier fires and
690 an enhanced updraft results in a higher than usual transport of VOCs to the UTLS, leading to a more important stratospheric
destruction than in other non El Niño years. A higher than usual increase in UTLS concentrations from surface sources in 2011
was also reported by Yuan et al. (2019).

Figure 15 shows that the enhanced OVOC concentrations in the UTLS are reduced when in-cloud OVOC oxidation is
taken into account. For example, EMAC predicts that the increase in the lower stratospheric methanol burden in November is
695 reduced by about 25.5 with the implementation of in-cloud OVOC oxidation. The increase of phenol, the aromatic that strongly
contributes to the lower stratospheric destruction, is only dampened by 8.7. Overall, this results in a 25.2 reduced destruction
of lower stratospheric and a 19.3 less reduction in lower stratospheric. In Fig. 3, it becomes obvious that other non-Indonesian
biomass burning events occur on the Indochinese Peninsula (i.e. northwest SEA) during the Asian monsoon. VOC emissions
from these fires are also transported by the ASMA into the UTLS, which contributes to the lower stratospheric depletion.
700 Without VOC emissions from the Indonesian peatland fires (simulation), the lower stratospheric decrease is dampened by
70.3. This difference is almost equal to the relative contribution of Indonesia's contribution to the total SEA VOC biomass
burning emissions in 2015, which accounts for 71.4.

In the second half of the 20th-century, stratospheric declined mainly due to halogen-containing substances from anthropogenic
activities (Molina and Rowland, 1974). After the Montreal Protocol has been implemented in 1989, a slowdown of the anthropogenic
705 stratospheric depletion was observed (Strahan and Douglass, 2018). Even though recovers in the upper- and mid-stratosphere, a
decline in lower stratospheric is observed by remote sensing measurements (Kyrölä et al., 2013; Nair et al., 2015; Vigouroux et al., 2015)
. Recently, a lower stratospheric decline of about 1.5 has been reported between 2001 and 2016 by Ball et al. (2018, their Fig. 3)
. This reduction is mainly attributed to meteorological variability's in dynamical processes (Chipperfield et al., 2018b; Ball et al., 2019)

710 ~~Between 2001 and 2016, we predict a lower stratospheric decrease of about 0.38, when using the lower stratosphere definition of Ball et al. (2018). They define the lower stratosphere between 147-32 (about 13-24) above 30° in latitudinal direction and between 100-32 (about 17-24) below 30° latitude. Based on our results, we therefore appoint that biomass burning VOC emissions from SEA (in particular the Indonesian peatland fires) contribute to this observed decline by about 25, which is robust against the influence of in-cloud OVOC oxidation (about 20). To our knowledge, most global stratospheric models do not consider this kind of VOC emissions and their chemistry, and are thus not able to capture this variability. However, it is~~
715 ~~important to keep in mind that our simulations are to some degree idealised in order to be able to isolate the impact of these emissions. For example, meteorological variations induced by changes in the chemical composition are neglected.~~

~~Another interesting aspect is the reduction in lower stratospheric. In the UTLS, aviation is the only direct anthropogenic activity and contributes about 3-5 to the total anthropogenic climate change (Lee et al., 2010). Here, aviation emissions lead to a formation of and a depletion of methane (CH_4). Recently, Rosanka et al. (2020a) showed that the enhancement in is limited by~~
720 ~~the background concentrations of and. If enough is available, a lower background concentration results in a higher gain. In general, low background concentrations limit the gain in winter. In our study, we find that in the North Atlantic flight sector (between 400-100), the burden is reduced due to SEA fires by about 6 with regional changes of more than 20 in 2015. At the same time, increases regionally by 10. Therefore, VOC emissions from frequently occurring Indonesian peatland fires potentially favour the formation of from aviation activities.~~

725 **8 Model uncertainties**

The most important aspects that influence our results are the representation of the transport processes, using different model resolutions, and the chemical kinetics. Each aspect is associated with some uncertainties of which all are shortly discussed in this section.

730 The magnitude of the depletion in lower stratospheric O_3 depends closely on the representation of the vertical transport that conveys the emitted VOCs into the UTLS. In order to evaluate the vertical transport processes of global models, ^{222}Rn (^{222}Rn , radioactive decay half-lifetime of 3.8 days) is typically used (Mahowald et al., 1997; Zhang et al., 2008; Jöckel et al., 2010). Jöckel et al. (2010) and more recently Brinkop and Jöckel (2019) analysed the ability of EMAC to capture the ^{222}Rn surface concentrations and vertical profiles. Their findings indicate that the vertical transport is well represented in EMAC (using the T42L90MA resolution) and that they are comparable to the earlier analysis with ECHAM5 (the base model of

735 EMAC) by Zhang et al. (2008). Figure 4 shows that the horizontal transport is also an important aspect that influences the distribution of the emitted VOCs from Indonesian peatland fires. Evaluating the horizontal transport using observations (like ^{222}Rn) is currently not possible. Recently however, Orbe et al. (2018) compared transport time scales of various global models, including EMAC. They found that the horizontal transport from NH mid-latitudes to the tropics differs by 30 %. Based on this comparison, it can be assumed that the horizontal transport is reasonably well represented in EMAC.

740 In this study, we transfer our process understanding from the fine- (T106L90MA) to the coarse-resolution (T42L90MA) simulations. It is therefore important to understand how well transport processes agree between both resolutions. Currently,

no direct analysis has been performed that focuses on the impact of different resolutions on transport processes in EMAC. However, Aghedo et al. (2010) analysed the influence of different horizontal and vertical resolutions in ECHAM5. Since EMAC uses the same horizontal and vertical transport scheme as ECHAM5, we assume that their findings also apply to EMAC. They
745 find that the vertical transport mainly depends on the number of levels used. By increasing the number of layers from 19 to 31 levels, the mass transported into the stratosphere reduces globally by about 36 %, whereas increasing the resolution from T42 to T106 only decreases the vertically transported mass globally by about 10 %. Here, the influence is the lowest (about 7 %) at high latitudes and [the](#) highest in the tropics (about 17 %). Aghedo et al. (2010) suggested that the higher impact in the tropics is probably related to tropical convection processes. Increasing the resolution changes the meridional transport in most regions
750 by less than 2 % and is thus negligible. For our purposes, differences in the inter-hemispheric transport are also negligible. The mean transport time from the NH to SH decreases from 11.9 to 11.8 months and for the SH to NH transport from 11.4 to 11.5 months when increasing the horizontal resolution from T42 to T106. By using the same vertical resolution (90 levels), the highest uncertainty introduced by using different resolutions is eliminated. It is therefore expected that the important transport processes are comparable and properly represented in both resolutions.

755 We find that the reaction of phenoxy radicals with O_3 (Reaction R2) has a significant influence at the surface, in the troposphere, and the lower stratosphere. As discussed by Taraborrelli et al. (2021), the chemical kinetics used in MOM to represent this O_3 loss is associated with some uncertainties. Currently, only the measured reaction rate constant for C_6H_5O is available and this is used for all phenoxy radicals. Yet, no experimental evidence has been found for the formation of phenyl peroxy radical ($C_6H_5O_2$), which might influence the cycling nature of this O_3 loss by Reactions R1 and R2. However, this product is still
760 to be expected. Even with different products, a significant ~~deletion~~ [depletion](#) of O_3 is anticipated by Reaction R2. At the same time, the reaction rate from Tao and Li (1999) is reported to be at the lower end, whereas a higher reaction rate would increase the depleted O_3 . Additionally, Taraborrelli et al. (2021) report that MOM neglects the non-HONO formation channel from nitrophenol photolysis, which does not destroy the aromatic ring and reforms phenoxy radicals (Cheng et al., 2009; Vereecken et al., 2016). It is therefore expected that, due to increasing nitrophenol concentrations in the lower troposphere (Sect. 5.3) as
765 well as in the UTLS, the importance of Reaction R2 as an O_3 sink is potentially underestimated.

9 Conclusions

In this study, the influence of VOC emissions from reoccurring Indonesian peatland fires is analysed with the main focus on 2015, a particularly strong year. This is achieved by performing multiple global simulations using EMAC. By comparing EMAC's prediction of HCN [and](#) CO columns to IASI satellite retrievals, we show that EMAC properly represents the emissions
770 from the Indonesian peatland fires and global biomass burning events.

Our results indicate that VOC emissions from biomass burning are important to reproduce hydrocarbons and secondary OVOCs in the atmosphere. Compared to other biomass burning regions, a particularly strong increase is modelled in SEA region, due to the unique emission footprint from the Indonesian peatland fires. [The enhanced formation of nitrophenols and strong HNCO emissions create toxic conditions in most parts of Indonesia, directly influencing its population.](#) Regionally,

775 significant changes in radical concentrations (HO_x and NO_x) are predicted. In general, O_3 increases in the lower troposphere with the highest changes in the NH high latitudes due to strong fires in Boreal Asia. However, on a global scale, tropospheric changes in O_3 are negligible. High aromatic emissions from peatland fires lead to a depletion of O_3 in eastern Indonesia. ~~The enhanced formation of nitrophenols and strong emissions create toxic conditions in most parts of Indonesia, directly influencing its population. The overall impact in the lower troposphere is reduced when in-cloud OVOC oxidation is taken into account.~~
780 ~~Especially, the increase initially predicted is reduced due to its enhanced destruction within clouds. However, the increased formation of oligomers in cloud droplets potentially leads to enhanced SOA concentrations.~~

The ongoing ASMA and the general tropical upward transport during the Indonesian fires, lift the emitted VOCs and their oxidation products quickly to the UTLS lower stratosphere. Here, ~~the enhanced VOC concentrations contribute to the depletion especially large increases are predicted for levels of the aromatic compounds benzene and toluene. The oxidation of VOCs~~
785 results in the reduction of OH and NO_x and the increase of HO_2 . Additionally, the Indonesian fires becomes a major source of lower stratospheric ~~. In particular, the NO_3 . Indonesian fires enhance the O_3 destruction by phenoxy radicals plays a key role. The predicted depletion is in line with remote sensing measurements and supported by our results that these VOC emissions contribute by about 25 by up to 20 % . Although high VOC emissions from biomass burning events in ALA and NAS have a large regional impact, their impact on the UTLS is negligible due to missing fast upward transport at higher latitudes.~~
790 southern tropical UTLS. This chemical loss propagates into the lower stratosphere and potentially influences the variability of O_3 retrieved from satellite observations. Overall, the highest changes in lower stratospheric radicals during the period between 2001 to 2016 is predicted for particular strong El Niño years, due to strong Indonesian peatland fires.

The overall impact of Indonesian fires on the composition of the troposphere and lower stratosphere is reduced when in-cloud OVOC oxidation is taken into account. Especially, the predicted O_3 increase in the troposphere is dampened due to enhanced
795 destruction of O_3 within clouds. This suggests that models neglecting the in-cloud oxidation of OVOCs probably tend to overestimate the impact of such an event like the Indonesian peatland fires.

Appendix A: HCN retrievals from IASI observations

The spaceborne data of HCN columns used in this study are obtained from the IASI radiance spectra by applying the version 3 of the Artificial Neural Network for IASI (ANNI) retrieval framework. Initially developed for the retrieval of NH_3 and dust
800 from the IASI observations (Whitburn et al., 2016a; Clarisse et al., 2019), ANNI v3 incorporates updates and modifications to allow the retrieval of a suite of VOCs. Until now, it has been used to retrieve methanol, formic acid, and PAN (Franco et al., 2018), then acetone (Franco et al., 2019) and acetic acid (Franco et al., 2020). Here, we perform the HCN retrieval by applying the full ANNI v3 procedure. As this approach has already been described in detail (see Franco et al., 2018, and references therein), we limit ourselves here to a summary of the main retrieval steps, and to the elements specific to the retrieval of HCN.
805 Examples of HCN columns from IASI single overpasses in the 2015 Indonesian fire plumes and averaged distributions are also presented.

As mentioned in Sect. 2.1.3, the ANNI retrieval method proceeds in two major steps. First, in each individual IASI radiance spectrum, the target species is detected and the strength of its absorption is quantified by a metric called the Hyperspectral Range Index (HRI). Then, the HRI is converted into a gas total column by means of an artificial feedforward neural network (NN), which also provides an uncertainty on the retrieved column.

The HRI is a dimensionless metric of the magnitude of the spectral signature of a target species in a given IASI spectrum, relative to the spectral variability of a "background" atmosphere in absence of the target gas, i.e. a variability resulting from all other parameters that contribute to the spectral radiance, such as other atmospheric gases (see Walker et al., 2011). The HRI is calculated over the main spectral range, in which the target species absorbs. The HCN absorption band (ν_2 branch) included in the IASI spectrum is situated close to a strong Q branch of CO_2 near 720 cm^{-1} . Therefore, the whole $700\text{--}800 \text{ cm}^{-1}$ spectral range covering many HCN features is used to calculate the HRI. The CO_2 line mixing in that range is accounted for as described by Duflot et al. (2013). A first HRI of HCN was already set up for the IASI observations by Duflot et al. (2015), but here we set up a new more sensitive one following the iterative procedure presented by Franco et al. (2018).

In contrast to Duflot et al. (2015), who used pre-calculated coefficients to link the HRI to the HCN total column, the ANNI v3 procedure implements an artificial feedforward NN for this purpose. Such a NN is set up to mimic in a comprehensive way the complex connections that exist between the HRI, the state of the atmosphere and Earth's surface, and the gas abundance. Setting up a NN requires a training phase, in which the NN learns from the presentation of an extensive dataset including all the necessary input and output variables. In ANNI v3, the NN inputs are the HRI, a spectral baseline temperature, the H_2O columns, the temperature profile, the surface pressure and emissivity, and the IASI viewing angle, whereas the output is the HCN column. Here, we built this training set from over 250,000 synthetic IASI spectra simulated by a line-by-line radiative transfer model. The advantage of such a synthetic training set is that it is free of the noise and/or scarcity of real measurements and that the spectra can be generated in large amounts in order to make the training set – and hence the NN – representative of all possible conditions. For example, the NN set up for HCN is trained to retrieve gas column from 1×10^{14} to $15 \times 10^{16} \text{ molecules cm}^{-2}$. Actually, two separate synthetic datasets are assembled per target species, one being representative of conditions close to emission sources, the other of mixing/transport conditions (see Whitburn et al., 2016a; Franco et al., 2018, for the rationale). Each training set leads to the setup of a specific NN that is used to globally retrieve the target species in emission or transport regimes, successively. The training performances are similar to those of the other VOCs retrieved with ANNI v3 and are reached with a NN made of two computational layers, each layer deploying eight nodes.

In addition to the total column, the NN returns an associated error that is calculated via a perturbation method of the input variables (see Whitburn et al., 2016a). A pre-filter prevents the retrieval on cloudy scenes (cloud coverage $> 10 \%$) or for observations with missing ancillary data. Consistent with the other ANNI VOCs products, a post-filter discards the individual retrievals affected by too large uncertainties or poor measurement sensitivity to HCN, specifically when

$$|\text{column}_{(\text{HCN})} / \text{HRI}_{(\text{HCN})}| > 8 \times 10^{15} \text{ molecules cm}^{-2} \quad (\text{A1})$$

or spectral baseline temperatures $< 268 \text{ K}$. This post-filter is not (directly) driven by the gas abundance, but rather by the thermal contrast (Franco et al., 2020). Finally, the constant climatological background of target gas abundance that is not

accounted for by the HRI has been estimated as 1.85×10^{15} molecules cm^{-2} for HCN (see Franco et al., 2018); this offset is thus added to the individual retrieved columns. Once set up, the NN is fed for each individual IASI observation with the appropriate input data. Here, we chose to use the ERA-5 reanalysis dataset (Hersbach et al., 2020) for the meteorological input data in the network. In the framework of the evaluation of EMAC in the 2015 Indonesian fires (see Sect. 4), only the HCN
 845 product obtained with the NN in transport/mixing regime has been exploited. Indeed, the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) onboard CALIPSO indicates fire plume located in the free troposphere during this massive biomass burning event.

Figure 17 presents the daily distributions of HCN total columns from IASI/Metop-A and -B observations in South-East Asia, for six successive days taken during the 2015 Indonesian fires. Whereas background areas are characterized by HCN
 850 total columns generally lower than 0.5×10^{16} molecules cm^{-2} , on the first day (September 29, 2015), strong HCN enhancements ($> 4 \times 10^{16}$ molecules cm^{-2}) are detected by IASI in the vicinity of Sumatra, indicating a massive fire plume. After six days (October 4, 2015), we can observe that the plume has grown progressively and that the bulk of HCN has been transported to the west across the Indian Ocean. The retrieved column uncertainties in the area generally fall in the range of $2\text{--}5 \times 10^{15}$ molecules cm^{-2} . Note that these uncertainties are reduced significantly by averaging numerous IASI measurements
 855 to build monthly or seasonal mean distributions of HCN columns. The typical seasonal distributions of IASI/Metop-A HCN columns are presented in Fig. 18 for the 2011–2014 time period, i.e. for years without massive fire events, such as the 2010 Russian fires or the 2015 Indonesian fires. These distributions highlight the dominant contribution of biomass burning to the atmospheric HCN burden, with HCN enhancements detected in Africa throughout the year, in South-East Asia in March–April–May, in India, eastern China and North Hemisphere mid- and high latitudes during the boreal summer, and within the
 860 tropics in September–October–November. Important outflows from these source regions are also noticeable, especially over the oceans. Figure 19 presents the monthly mean HCN columns during the 2015 Indonesian fires (from September to December) along with the corresponding distributions over the 2011–2014 time period. It illustrates the exceptional intensity of the 2015 fires compared to the previous years, with important HCN enhancements detected throughout the entire intertropical band.

Appendix B: Definition of \sum OVOCs

865 In Fig. 15, the mixing ratios of the sum of all the OVOCs explicitly reacting in JAMOC (\sum OVOCs) are shown. In this case, \sum OVOCs is defined as:

$$\begin{aligned} \sum \text{OVOCs} = & \text{methanol} + \text{formaldehyde} + \text{methyl hydroperoxide} + \text{hydroxymethylhydroperoxide} + \text{ethanol} + \\ & \text{ethylene glycol} + \text{acetaldehyde} + \text{glycolaldehyde} + \text{glyoxal} + \text{1-hydroperoxyacetone} + \\ & \text{methylglyoxal} + \text{isopropanol} + \text{isopropyl hydro peroxide} + \text{methacrolein} + \text{methyl vinyl ketone} \end{aligned}$$

(B1)

Data availability. The simulation results are archived at the Jülich Supercomputing Centre (JSC) and are available on request. The IASI VOC columns retrieved with the ANNI framework are available upon request. The IASI CO data processed with FORLI-CO v0151001 can be downloaded from the AERIS portal at <http://iasi.aeris-data.fr/CO/> (last access: 22 February 2021).

Author contributions. The study was designed by SR and DT. AP implemented the algorithms for VOC emissions from biomass burning and terrestrial vegetation. SR adjusted the biomass burning emission factors. AP and DT implemented HCN deposition to the ocean. SR performed the simulations and analysed the data with contributions from DT. BF, LC, and PFC developed the IASI VOC products and contributed to the analyses. SR and DT discussed the results with contributions from BF and AW. The manuscript was prepared by SR with the help of all co-authors.

Competing interests. The authors declare that they have no competing of interest.

Acknowledgements. The work described in this paper has received funding from the Initiative and Networking Fund of the Helmholtz Association through the project “Advanced Earth System Modelling Capacity (ESM)”. The content of this paper is the sole responsibility of the author(s) and it does not represent the opinion of the Helmholtz Association, and the Helmholtz Association is not responsible for any use that might be made of the information contained. The authors gratefully acknowledge the Earth System Modelling Project (ESM) for funding this work by providing computing time on the ESM partition of the supercomputer JUWELS at the Jülich Supercomputing Centre (JSC). IASI is a joint mission of EUMETSAT and the Centre National d’Etudes Spatiales (CNES, France). The [authors acknowledge the AERIS data infrastructure for providing access to the IASI data, Daniel Hurtmans for the development of the CO retrievals, and Eumetsat/AC SAF for CO data production.](#) The research at ULB has been supported by the project OCTAVE (Oxygenated Compounds in the Tropical Atmosphere: Variability and Exchanges (<http://octave.aeronomie.be/>)) of the Belgian Research Action through Interdisciplinary Networks (BRAIN-be; 2017–2021; Research project BR/175/A2/OCTAVE) and by the IASI. Flow Prodex arrangement (ESA–BELSPO). LC is a research associate supported by the F.R.S.–FNRS.

Table 1. Characteristics List of the different biomass burning regions focusing on the dominant fire type, the main biomass burning season, EMAC simulations performed in this study. Here, ScSta indicates EMAC's standard aqueous-phase mechanism (Jöckel et al., 2016) and JAMOC indicates the Dry-Matter-Burned-complex in-cloud OVOC oxidation scheme by Rosanka et al. (2020c,b) (DMB for further details see Sect. 2.1.1). The global DMB is 4985 Tg a⁻¹. The naming of each region is as follows: ALA = Alaska, CAF = Central Africa, CSA = Central South America, NAS = North Asia, NAU = North Australia, SAF = South Africa, and SEA = South East Asia. Each region is graphically illustrated

Name	Analysed Period	VOC Bioburn Emissions	Aqueous-phase mechanism	Resolution
REF	2015-2016	no	ScSta	T106L90MA
FIR	2015-2016	yes	ScSta	T106L90MA
FIR _{NOINDO}	SOND^a in 2015	yes^b	ScSta	T106L90MA
REF _{LONG}	2001-2016	no	ScSta	T42L90MA
FIR _{LONG}	2001-2016	yes	ScSta	T42L90MA
REF _{JAMOC}	SOND^a in Fig-2, 2015	no	JAMOC	T106L90MA
FIR _{JAMOC}	SOND^a in 2015	yes	JAMOC	T106L90MA

^aFocus on Indonesia in September, October, November, and December

^bNo VOC biomass burning emissions from Indonesian peatland fires

Table 2. Characteristics of the different biomass burning regions focusing on the dominant fuel type, the main biomass burning season, and the Dry Matter Burned (DMB). The global DMB by GFAS (Kaiser et al., 2012) for the year 2015 is 4985 Tg a⁻¹. The naming of each region is as follows: ALA = Alaska, CAF = Central Africa, CSA = Central South America, NAS = North Asia, NAU = North Australia, SAF = South Africa, and SEA = South East Asia. Each region is graphically illustrated in Fig. 2.

Region	Dominant fuel type	Main biomass burning season	2015 DMB [Tg a ⁻¹]
ALA	Extratropical forest with organic soil	JJA	295
CAF	Tropical forest & Savanna	DJF	778
CSA	Savanna	SON	439
NAS	Extratropical forest with organic soil	MMA & JJA	363
NAU	Savanna	SON	260
SAF	Tropical forest & Savanna	JJA	1036
SEA	Tropical forest ^a	SON	1237 ^b

^aIn this study a combination of tropical forest (50 %) & peatland (50 %) is assumed in Indonesia (Sect. 2.1.2)

^bOf which 949 Tg a⁻¹ are from Indonesian peatland fires

Table 3. Absolute (Abs.) and relative (Rel.) changes in the tropospheric burden for each region and each species discussed. Regional differences are calculated for the main biomass burning season (see Table 2) and the global changes are calculated for the complete year of 2015. If not stated otherwise, The unit for the absolute differences are explicitly given in and, whereas relative changes are provided in parenthesis always given in %. Most radical burdens are presented in mol. The differences are calculated between simulation REF and FIR.

Species	Unit ^a	Global		ALA		CAF		CSA		NAS		NAU		SAF		SEA	
		Abs.	Rel.	Abs.	Rel.	Abs.	Rel.	Abs.	Rel.	Abs.	Rel.	Abs.	Rel.	Abs.	Rel.	Abs.	Rel.
Aliphatic hydrocarbons																	
Ethane	Gg	422.9	32.6	20.4	140.1	18.4	43.6	23.9	144.9	31.8	47.3	16.7	128.3	26.5	123.8	47.6	48.2
Propane	Gg	19.4	6.3	2.3	290.8	1.5	12.8	1.7	51.9	3.1	18.7	0.7	79.8	2.4	84.3	2.8	18.1
<i>n</i> -Butane	Mg	3727.4	2.4	470.4	242.7	256.1	6.0	336.3	16.9	671.3	8.6	86.3	33.2	324.9	34.9	779.6	9.4
Ethylene	Gg	13.5	12.5	2.0	103.5	2.9	59.9	3.6	9.2	2.2	46.3	0.8	26.4	4.5	81.5	8.3	85.3
Propene	Mg	3283.5	11.3	590.7	116.2	1021.5	83.4	863.5	5.4	560.7	54.3	279.5	34.5	1420.7	100.9	2304.0	83.6
Isobutene	Mg	107.1	20.2	36.2	7299.4	21.7	103.8	41.0	379.8	29.3	293.8	4.1	380.5	27.0	161.0	53.8	70.9
Acetylene	Gg	45.7	20.5	1.4	119.8	4.4	56.0	6.0	305.0	2.5	20.8	2.8	258.8	6.6	184.5	13.2	64.2
Aromatic hydrocarbons																	
Benzene	Gg	38.8	27.3	4.4	1312.7	2.7	46.4	4.3	368.2	5.9	84.2	1.9	498.1	4.6	228.7	20.0	207.7
Toluene	Mg	6655.3	15.3	802.6	1308.7	595.5	62.8	1007.7	85.0	968.2	62.0	175.9	198.2	950.0	199.0	10169.8	366.8
Styrene	Mg	56.2	29.8	42.8	11375.3	8.0	270.4	10.2	151.1	31.7	627.7	2.1	181.9	13.2	582.3	10.0	35.6
Ethylbenzene	Mg	2431.9	65.9	774.2	27408.4	421.5	958.0	351.8	470.8	820.3	592.5	139.2	1377.7	731.0	3203.8	237.3	75.4
OVOCs																	
Formaldehyde	Gg	25.4	2.2	3.6	35.1	5.4	11.7	5.3	2.2	3.4	12.6	1.6	3.9	8.4	17.7	17.9	15.9
Acetaldehyde	Gg	15.5	11.4	1.0	79.0	2.6	47.7	4.2	6.9	1.3	31.2	0.9	22.5	3.7	74.3	8.8	80.7
Glycolaldehyde	Gg	21.3	8.0	1.7	65.6	3.7	27.8	6.9	5.1	2.4	36.5	1.2	9.5	5.8	34.7	17.2	67.1
Methanol	Gg	223.3	7.9	11.7	31.6	17.0	16.8	31.5	6.6	18.8	16.2	14.4	17.5	27.4	28.5	112.3	60.7
Glyoxal	Mg	3872.3	9.3	388.7	126.2	633.0	24.5	591.5	3.9	397.3	38.6	240.1	12.7	1048.3	36.7	3186.1	62.2
Methyl glyoxal	Mg	2481.7	1.3	208.1	17.4	438.9	4.1	206.5	0.2	212.4	6.9	144.4	1.3	743.4	5.8	2195.8	10.2
2,3-butanedione	Mg	487.1	205.5	0.4	180.5	125.7	812.6	319.7	8060.4	1.0	21.0	6.4	960.4	94.2	2588.8	304.2	807.0
Phenol	Mg	1167.8	105.7	171.8	4282.2	179.1	316.4	275.7	1305.1	155.7	323.5	63.9	1353.8	277.1	1339.7	1400.3	1226.3
Benzaldehyde	Mg	282.1	14.6	102.9	4472.0	76.1	196.9	48.6	92.4	83.2	221.3	19.4	284.9	108.2	574.7	125.2	49.9
CO	Gg	8341.1	2.4	153.3	3.9	253.4	2.4	446.4	2.8	307.5	2.6	503.2	6.2	273.4	3.4	1908.0	6.5
Acids																	
Formic acid	Gg	32.9	4.9	1.5	24.0	7.0	25.2	5.8	4.7	2.1	12.9	3.2	13.0	11.4	49.9	11.3	20.7
Acetic acid	Gg	119.3	23.3	8.7	441.7	29.0	124.3	34.1	15.3	9.8	128.5	10.4	45.3	48.7	238.9	37.7	117.4
Oxidants																	
O ₃	Gg	1115.3	0.3	92.0	1.9	61.6	0.7	-28.3	-0.3	104.7	0.7	2.0	0.0	60.4	0.8	83.2	0.4
OH	Kmol	-240.3	-1.7	-6.4	-4.9	-8.5	-2.4	-15.9	-4.7	-10.8	-3.4	-21.6	-6.8	-10.3	-3.5	-58.2	-4.5
HO ₂	Kmol	3537.2	0.4	354.6	3.7	410.0	1.7	367.6	0.8	409.6	1.8	157.7	0.7	533.4	2.8	945.4	1.3
NO	Mmol	-82.3	-2.8	-38.4	-52.7	-5.9	-6.5	-5.7	-6.6	-26.0	-26.6	-6.8	-10.6	-6.4	-7.5	-22.0	-7.7
NO ₂	Mmol	-178.5	-2.2	-73.5	-36.7	-16.3	-4.7	-14.2	-4.1	-60.1	-19.3	-13.0	-6.5	-18.8	-5.5	-70.5	-8.3
NO ₃	Kmol	6926.5	5.3	151.3	35.5	385.0	6.0	193.8	6.6	407.2	15.2	497.8	16.9	432.3	9.3	2368.5	15.5

^aUnit for absolute values only. Relative values given in %.

Table 4. Stratospheric burden in November 2015 and changes induced by VOC biomass burning emissions. In addition the relative difference (Rel.) and the Indonesian contribution (Indo. contr.). The latter is calculated based on the difference of FIR and FIR_{NOINDO} .

Species	Unit	REF	Δ FIR	Rel. [%]	Indo. contr. [%]
Ethane	Gg	30.8	43.6	41.6	24.1
Propane	Mg	666.3	855.9	28.5	69.1
<i>n</i> -Butane	Mg	116.2	152.2	31.0	80.0
Benzene	Mg	242.1	724.2	199.2	87.2
Methanol	Gg	17.1	22.9	34.0	76.0
Glyoxal	Mg	9.8	14.7	48.9	69.4
Phenol	Kg	629.8	1685.6	167.7	85.8
Acetic acid	Mg	289.4	473.6	63.7	72.6
HCN	Gg	16.7	53.5	220.4	62.6
Nitrophenols	Mg	14.7	23.6	60.2	69.5
OH	Mmol	3.8	-0.1	-3.2	70.2
NO	Gmol	1.9	-0.1	-6.1	67.7
NO ₂	Gmol	3.6	-0.2	-4.2	66.2
NO ₃	Mmol	10.3	1.0	9.5	74.7
HO ₂	Mmol	24.1	0.6	2.4	63.9

Table 5. The SEA tropospheric burden during the Indonesian fire period with and without VOC biomass burning emissions of OVOCs explicitly reacting in JAMOC. The tropospheric burden is given in Gg.

Species	REF	Δ FIR	Rel. [%]	REF _{JAMOC}	Δ FIR _{JAMOC}	Rel. [%]
Methanol	185.2	112.3	60.7	133.8	87.9	65.7
Methyl hydroperoxide	140.8	10.7	7.6	71.7	4.0	5.6
Hydroxymethylhydroperoxide	5.0	1.0	20.2	3.4	0.6	17.0
Ethanol	9.1	1.0	10.5	7.1	0.8	11.5
Ethylene glycol	0.2	0.2	73.6	0.1	0.1	70.1
Glycolaldehyde	25.6	17.2	67.1	13.2	10.6	80.8
1-Hydroperoxyacetone	4.7	0.7	14.5	2.4	0.3	13.2
Methyl glyoxal	21.5	2.2	10.2	15.5	1.7	10.9
Isopropyl hydro peroxide	0.8	0.1	18.1	0.4	0.1	17.3

Table 6. The lower stratospheric burden in November with and without VOC biomass burning emissions of OVOCs explicitly reacting in JAMOC. The stratospheric burden is given in Mg.

Species	REF	Δ FIR	Rel. [%]	REF _{JAMOC}	Δ FIR _{JAMOC}	Rel. [%]
<u>Methanol</u>	<u>17097.87</u>	<u>5821.01</u>	<u>34.05</u>	<u>10102.29</u>	<u>2634.34</u>	<u>26.08</u>
<u>Methyl hydroperoxide</u>	<u>637.88</u>	<u>55.45</u>	<u>8.69</u>	<u>562.11</u>	<u>28.3</u>	<u>5.03</u>
<u>Hydroxy methyl hydroperoxide</u>	<u>60.56</u>	<u>12.14</u>	<u>20.04</u>	<u>49.16</u>	<u>6.17</u>	<u>12.55</u>
<u>Ethanol</u>	<u>45.35</u>	<u>15.5</u>	<u>34.18</u>	<u>24.15</u>	<u>5.45</u>	<u>22.55</u>
<u>Ethylene glycol</u>	<u>0.64</u>	<u>0.24</u>	<u>38.47</u>	<u>0.09</u>	<u>0.03</u>	<u>31.04</u>
<u>Glycolaldehyde</u>	<u>84.23</u>	<u>16.41</u>	<u>19.48</u>	<u>54.31</u>	<u>11.73</u>	<u>21.61</u>
<u>1-Hydroperoxyacetone</u>	<u>10.83</u>	<u>3.86</u>	<u>35.59</u>	<u>5.96</u>	<u>1.76</u>	<u>29.57</u>
<u>Methyl glyoxal</u>	<u>6.58</u>	<u>0.34</u>	<u>5.11</u>	<u>3.5</u>	<u>0.2</u>	<u>5.6</u>
<u>Isopropyl hydro peroxide</u>	<u>2.61</u>	<u>1.9</u>	<u>72.5</u>	<u>2.1</u>	<u>1.03</u>	<u>49.06</u>




Figure 1. Accumulated dry matter burned (DMB) during the Indonesian peatland fires of 2015. The distribution of Indonesian peatland is indicated in blue. The data for the peatland distribution are obtained from Xu et al. (2017, 2018).




Figure 2. Mean dry matter burned (DMB) in 2015. The naming of each region is as follows: ALA = Alaska, CAF = Central Africa, CSA = Central South America, NAS = North Asia, NAU = North Australia, SAF = South Africa, and SEA = South East Asia. Further details about each region are presented in Table 2.




Figure 3. Relative contribution of each defined emission region to the The total, the VOC, and the aromatic emissions from biomass burning for each region in 2015, non-El Niño years, El Niño years, and in 2015 predicted by EMAC (based on REF_{LONG} and FIR_{LONG}). Further details about each region are presented in Table 2 and Fig. 2.




Figure 4. HCN column-comparison between IASI satellite-observations and EMAC for September, October-REF and November-2015. IASI satellite-observations (left), the simulation without VOC biomass burning emissions-FIR (simulation a) in comparison to Mean global observed IASI observations HCN columns for September to November. (centre) - Mean global HCN column comparison between REF and the simulation including VOC biomass burning emissions IASI for September to November. (simulation c) in Mean global HCN column comparison to between FIR and IASI observations (right) for September to November.




Figure 5. Global HCN column bias between EMAC simulations and IASI satellite data. The column bias is calculated based on monthly mean data during the Indonesian peatland fires in 2015.




Figure 6. CO comparison between IASI and FIR. (a) Mean global observed IASI CO columns for September to November. (b) Mean global CO column comparison between FIR and IASI for September to November. (c) Scatter plot for direct comparison between FIR and IASI over Indonesia for September to November.




Figure 7. Illustration of the impact of VOC emissions from the Indonesian peatland fires on the atmospheric composition.




Figure 8. Number of days in which ambient concentrations of 1 ppb of HCHO are exceeded during the Indonesian peatland fires in 2015 between August and October.




Figure 9. (a) Yearly mean tropospheric nitrophenol (NPs) column without biomass burning VOC emissions. (b) Changes in the yearly mean tropospheric nitrophenol (NPs) due to VOC biomass burning.




Figure 10. (a) Yearly mean surface OH concentration without biomass burning VOC emissions. (b) Changes in the yearly mean surface OH concentration due to VOC biomass burning.




Figure 11. (a) Yearly mean surface NO_x concentration without biomass burning VOC emissions. (b) Changes in the yearly mean surface NO_x concentration due to VOC biomass burning.




Figure 12. The mean longitudinal relative change in lower stratospheric (a) Yearly-OH, (b) HO₂, (c) NO_x, and (d) NO₃ between 2001 and 2016. Results are based on both long-term simulations (simulation REF_{LONG} and FIR_{LONG}). The lower stratosphere is defined between 147-32 hPa (about 13–24 km) above 30° in latitudinal direction and between 100-32 hPa (about 17–24 km) below 30° latitude.




Figure 13. Zonal mean contribution of the destruction of O₃ by phenoxy radicals (Reaction R2) to the total chemical O₃ loss. (a) Relative contribution if no VOC biomass burning emissions are taken into account (REF) in November 2015. (b) Changes in the relative contribution due to VOC biomass burning emissions (FIR) in November 2015. (c) Changes in the relative contribution due to VOC biomass burning emissions in the UTLS. Here, we define the UTLS from 250 hPa to 50 hPa above the tropopause calculated by EMAC. The tropopause pressure level in (a) and (b) is indicated in black.




Figure 14. Zonal mean relative change in the destruction of O₃ by phenoxy radicals (Reaction R2) due to VOC biomass burning emissions (FIR vs REF) in April 2016. The tropopause pressure level is indicated in black.




Figure 15. Mean zonal change in the sum of all OVOCs explicitly reacting in JAMOC (\sum OVOCs, see Equation B1 in Appendix B) over Indonesia and the Indian Ocean during the 2015 Indonesian fire period (SON). (a) Changes due to VOC biomass burning emissions (difference between simulation FIR and REF) and (b) changes due to JAMOC (Δ JAMOC).




Figure 16. Probability density function of EMAC's methanol column bias to the IASI satellite measurements for simulation REF, FIR, REF_{JAMOC}, and FIR_{JAMOC} in SEA during the 2015 Indonesian peatland fires.




Figure 17. Daily regional distributions of HCN total column (in molecules cm^{-2}) derived from the IASI spectra recorded in the morning overpasses of Metop-A and -B, for six successive days during the 2015 Indonesian fires. These distributions take into account the actual footprint on the Earth's surface of each individual IASI measurement, i.e. a small circle at nadir and an elongated ellipse at the limit of the across-track swath of the satellite. Note the complementarity of the IASI/Metop-A and -B flight tracks that avoid gaps between the successive overpasses in the tropics. The white areas correspond to data filtered out because of unsatisfactory retrieval quality or the presence of clouds.




Figure 18. Seasonal means (on a $1 \times 1^\circ$ grid) of the HCN total columns (in molecules cm^{-2}) retrieved from the IASI/Metop-A measurements over the 2011–2014 time period. The HCN columns over the continents have been retrieved with the NN in emission regime, whereas the NN in transport/mixing regime has been used over the oceans.




Figure 19. Monthly means (on a $1 \times 1^\circ$ grid) of the HCN total columns (in molecules cm^{-2}) retrieved from the IASI/Metop-A measurements over the 2011–2014 time period (left plots) and over the year 2015 (right plots).

References

- Aghedo, A. M., Rast, S., and Schultz, M. G.: Sensitivity of tracer transport to model resolution, prescribed meteorology and tracer lifetime in the general circulation model ECHAM5, *Atmospheric Chemistry and Physics*, 10, 3385–3396, <https://doi.org/10.5194/acp-10-3385-2010>, 2010.
- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmospheric Chemistry and Physics*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmospheric Chemistry and Physics*, 19, 8523–8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.
- Atkinson, R., Carter, W. P. L., Darnall, K. R., Winer, A. M., and Pitts Jr., J. N.: A smog chamber and modeling study of the gas phase NO_x–air photooxidation of toluene and the cresols, *International Journal of Chemical Kinetics*, 12, 779–836, <https://doi.org/10.1002/kin.550121102>, 1980.
- Ball, W. T., Alsing, J., Mortlock, D. J., Staehelin, J., Haigh, J. D., Peter, T., Tummon, F., Stübi, R., Stenke, A., Anderson, J., Bourassa, A., Davis, S. M., Degenstein, D., Frith, S., Froidevaux, L., Roth, C., Sofieva, V., Wang, R., Wild, J., Yu, P., Ziemke, J. R., and Rozanov, E. V.: Evidence for a continuous decline in lower stratospheric ozone offsetting ozone layer recovery, *Atmospheric Chemistry and Physics*, 18, 1379–1394, <https://doi.org/10.5194/acp-18-1379-2018>, 2018.
- Ball, W. T., Alsing, J., Staehelin, J., Davis, S. M., Froidevaux, L., and Peter, T.: Stratospheric ozone trends for 1985–2018: sensitivity to recent large variability, *Atmospheric Chemistry and Physics*, 19, 12 731–12 748, <https://doi.org/10.5194/acp-19-12731-2019>, 2019.
- Basha, G., Ratnam, M. V., and Kishore, P.: Asian summer monsoon anticyclone: trends and variability, *Atmospheric Chemistry and Physics*, 20, 6789–6801, <https://doi.org/10.5194/acp-20-6789-2020>, 2020.
- Blando, J. D. and Turpin, B. J.: Secondary organic aerosol formation in cloud and fog droplets: a literature evaluation of plausibility, *Atmospheric Environment*, 34, 1623 – 1632, [https://doi.org/10.1016/S1352-2310\(99\)00392-1](https://doi.org/10.1016/S1352-2310(99)00392-1), 2000.
- Brinkop, S. and Jöckel, P.: ATTLA 4.0: Lagrangian advective and convective transport of passive tracers within the ECHAM5/MESSy (2.53.0) chemistry–climate model, *Geoscientific Model Development*, 12, 1991–2008, <https://doi.org/10.5194/gmd-12-1991-2019>, 2019.
- Cabrera-Perez, D., Taraborrelli, D., Sander, R., and Pozzer, A.: Global atmospheric budget of simple monocyclic aromatic compounds, *Atmospheric Chemistry and Physics*, 16, 6931–6947, <https://doi.org/10.5194/acp-16-6931-2016>, 2016.
- Cheng, S.-B., Zhou, C.-H., Yin, H.-M., Sun, J.-L., and Han, K.-L.: OH produced from o-nitrophenol photolysis: A combined experimental and theoretical investigation, *The Journal of Chemical Physics*, 130, 234 311, <https://doi.org/10.1063/1.3152635>, 2009.
- Chipperfield, M. P., Dhomse, S., Hossaini, R., Feng, W., Santee, M. L., Weber, M., Burrows, J. P., Wild, J. D., Loyola, D., and Coldewey-Egbers, M.: On the Cause of Recent Variations in Lower Stratospheric Ozone, *Geophysical Research Letters*, 45, 5718–5726, <https://doi.org/https://doi.org/10.1029/2018GL078071>, 2018a.
- Chipperfield, M. P., Dhomse, S., Hossaini, R., Feng, W., Santee, M. L., Weber, M., Burrows, J. P., Wild, J. D., Loyola, D., and Coldewey-Egbers, M.: On the Cause of Recent Variations in Lower Stratospheric Ozone, *Geophysical Research Letters*, 45, 5718–5726, <https://doi.org/10.1029/2018GL078071>, 2018b.

- Christian, T. J., Kleiss, B., Yokelson, R. J., Holzinger, R., Crutzen, P. J., Hao, W. M., Saharjo, B. H., and Ward, D. E.: Comprehensive laboratory measurements of biomass-burning emissions: 1. Emissions from Indonesian, African, and other fuels, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2003JD003704>, 2003.
- 925 Cicerone, R. J. and Zellner, R.: The atmospheric chemistry of hydrogen cyanide (HCN), *Journal of Geophysical Research: Oceans*, 88, 10 689–10 696, <https://doi.org/10.1029/JC088iC15p10689>, 1983.
- Clarisse, L., Clerbaux, C., Franco, B., Hadji-Lazaro, J., Whitburn, S., Kopp, A. K., Hurtmans, D., and Coheur, P.-F.: A Decadal Data Set of Global Atmospheric Dust Retrieved From IASI Satellite Measurements, *Journal of Geophysical Research: Atmospheres*, 124, 1618–1647, 930 <https://doi.org/10.1029/2018jd029701>, 2019.
- Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., Wespes, C., and Coheur, P.-F.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmospheric Chemistry and Physics*, 9, 6041–6054, <https://doi.org/10.5194/acp-9-6041-2009>, 2009.
- Crippa, P., Castruccio, S., Archer-Nicholls, S., Lebron, G. B., Kuwata, M., Thota, A., Sumin, S., Butt, E., Wiedinmyer, C., and 935 Spracklen, D. V.: Population exposure to hazardous air quality due to the 2015 fires in Equatorial Asia, *Scientific Reports*, 6, 37 074, <https://doi.org/10.1038/srep37074>, 2016.
- Deckert, R., Jöckel, P., Grewe, V., Gottschaldt, K.-D., and Hoor, P.: A quasi chemistry-transport model mode for EMAC, *Geoscientific Model Development*, 4, 195–206, <https://doi.org/10.5194/gmd-4-195-2011>, 2011.
- Duflot, V., Hurtmans, D., Clarisse, L., R'honi, Y., Vigouroux, C., Mazière, M. D., Mahieu, E., Servais, C., Clerbaux, C., and Coheur, P.- 940 F.: Measurements of hydrogen cyanide (HCN) and acetylene (C₂H₂) from the Infrared Atmospheric Sounding Interferometer (IASI), *Atmospheric Measurement Techniques*, 6, 917–925, <https://doi.org/10.5194/amt-6-917-2013>, 2013.
- Duflot, V., Wespes, C., Clarisse, L., Hurtmans, D., Ngadi, Y., Jones, N., Paton-Walsh, C., Hadji-Lazaro, J., Vigouroux, C., Mazière, M. D., Metzger, J.-M., Mahieu, E., Servais, C., Hase, F., Schneider, M., Clerbaux, C., and Coheur, P.-F.: Acetylene (C₂H₂) and hydrogen cyanide (HCN) from IASI satellite observations: global distributions, validation, and comparison with model, *Atmospheric Chemistry and Physics*, 945 15, 10 509–10 527, <https://doi.org/10.5194/acp-15-10509-2015>, 2015.
- Ervens, B.: Modeling the Processing of Aerosol and Trace Gases in Clouds and Fogs, *Chemical Reviews*, 115, 4157–4198, <https://doi.org/10.1021/cr5005887>, 2015.
- Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmospheric Chemistry and Physics*, 11, 11 069–11 102, <https://doi.org/10.5194/acp-11-11069-2011>, 950 2011.
- Field, R. D., van der Werf, G. R., Fanin, T., Fetzer, E. J., Fuller, R., Jethva, H., Levy, R., Livesey, N. J., Luo, M., Torres, O., and Worden, H. M.: Indonesian fire activity and smoke pollution in 2015 show persistent nonlinear sensitivity to El Niño-induced drought, *Proceedings of the National Academy of Sciences*, 113, 9204–9209, <https://doi.org/10.1073/pnas.1524888113>, 2016.
- Franco, B., Clarisse, L., Stavroukou, T., Müller, J.-F., Van Damme, M., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Taraborrelli, D., Cler- 955 baux, C., and Coheur, P.-F.: A General Framework for Global Retrievals of Trace Gases From IASI: Application to Methanol, Formic Acid, and PAN, *Journal of Geophysical Research: Atmospheres*, 123, 13,963–13,984, <https://doi.org/10.1029/2018JD029633>, 2018.
- Franco, B., Clarisse, L., Stavroukou, T., Müller, J.-F., Pozzer, A., Hadji-Lazaro, J., Hurtmans, D., Clerbaux, C., and Coheur, P.-F.: Acetone Atmospheric Distribution Retrieved From Space, *Geophysical Research Letters*, 46, 2884–2893, <https://doi.org/10.1029/2019gl082052>, 2019.

- 960 Franco, B., Clarisse, L., Stavrou, T., Müller, J.-F., Taraborrelli, D., Hadji-Lazaro, J., Hannigan, J. W., Hase, F., Hurtmans, D., Jones, N., Lutsch, E., Mahieu, E., Ortega, I., Schneider, M., Strong, K., Vigouroux, C., Clerbaux, C., and Coheur, P.-F.: Spaceborne Measurements of Formic and Acetic Acids: A Global View of the Regional Sources, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2019gl086239>, 2020.
- Fu, R., Hu, Y., Wright, J. S., Jiang, J. H., Dickinson, R. E., Chen, M., Filipiak, M., Read, W. G., Waters, J. W., and Wu, D. L.: Short circuit
965 of water vapor and polluted air to the global stratosphere by convective transport over the Tibetan Plateau, *Proceedings of the National Academy of Sciences*, 103, 5664–5669, <https://doi.org/10.1073/pnas.0601584103>, 2006.
- Gaveau, D. L. A., Salim, M. A., Hergoualc'h, K., Locatelli, B., Sloan, S., Wooster, M., Marlier, M. E., Molidena, E., Yaen, H., DeFries, R., Verchot, L., Murdiyarso, D., Nasi, R., Holmgren, P., and Sheil, D.: Major atmospheric emissions from peat fires in Southeast Asia during non-drought years: evidence from the 2013 Sumatran fires, *Scientific Reports*, 4, 6112, <https://doi.org/10.1038/srep06112>, 2014.
- 970 George, M., Clerbaux, C., Bouarar, I., Coheur, P.-F., Deeter, M. N., Edwards, D. P., Francis, G., Gille, J. C., Hadji-Lazaro, J., Hurtmans, D., Inness, A., Mao, D., and Worden, H. M.: An examination of the long-term CO records from MOPITT and IASI: comparison of retrieval methodology, *Atmospheric Measurement Techniques*, 8, 4313–4328, <https://doi.org/10.5194/amt-8-4313-2015>, 2015.
- Grosjean, D.: Atmospheric reactions of ortho cresol: Gas phase and aerosol products, *Atmospheric Environment (1967)*, 18, 1641 – 1652, [https://doi.org/10.1016/0004-6981\(84\)90386-X](https://doi.org/10.1016/0004-6981(84)90386-X), 1984.
- 975 Grosjean, D.: Atmospheric fate of toxic aromatic compounds, *Science of The Total Environment*, 100, 367 – 414, [https://doi.org/10.1016/0048-9697\(91\)90386-S](https://doi.org/10.1016/0048-9697(91)90386-S), an Honour Volume For Eric I. Hamilton, 1991.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, <https://doi.org/10.5194/acp-6-3181-2006>, 2006.
- 980 Hagemann, S. and Stacke, T.: Impact of the soil hydrology scheme on simulated soil moisture memory, *Climate Dynamics*, 44, 1731–1750, <https://doi.org/10.1007/s00382-014-2221-6>, 2015.
- Hems, R. F. and Abbatt, J. P. D.: Aqueous Phase Photo-oxidation of Brown Carbon Nitrophenols: Reaction Kinetics, Mechanism, and Evolution of Light Absorption, *ACS Earth and Space Chemistry*, 2, 225–234, <https://doi.org/10.1021/acsearthspacechem.7b00123>, 2018.
- Hens, K., Novelli, A., Martinez, M., Auld, J., Axinte, R., Bohn, B., Fischer, H., Keronen, P., Kubistin, D., Nölscher, A. C., Oswald, R.,
985 Paasonen, P., Petäjä, T., Regelin, E., Sander, R., Sinha, V., Sipilä, M., Taraborrelli, D., Tatum Ernest, C., Williams, J., Lelieveld, J., and Harder, H.: Observation and modelling of HO_x radicals in a boreal forest, *Atmospheric Chemistry and Physics*, 14, 8723–8747, <https://doi.org/10.5194/ACP-14-8723-2014>, 2014.
- Herrmann, H., Schaefer, T., Tilgner, A., Styler, S. A., Weller, C., Teich, M., and Otto, T.: Tropospheric Aqueous-Phase Chemistry: Kinetics, Mechanisms, and Its Coupling to a Changing Gas Phase, *Chemical Reviews*, 115, 4259–4334, <https://doi.org/10.1021/cr500447k>, pMID: 25950643, 2015.
- 990 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

- Hodnebrog, Ø., Dalsøren, S. B., and Myhre, G.: Lifetimes, direct and indirect radiative forcing, and global warming potentials of ethane (C₂H₆), propane (C₃H₈), and butane (C₄H₁₀), *Atmospheric Science Letters*, 19, e804, <https://doi.org/https://doi.org/10.1002/asl.804>, 2018.
- 1000 Hurtmans, D., Coheur, P.-F., Wespes, C., Clarisse, L., Scharf, O., Clerbaux, C., Hadji-Lazaro, J., George, M., and Turquety, S.: FORLI radiative transfer and retrieval code for IASI, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 113, 1391–1408, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2012.02.036>, three Leaders in Spectroscopy, 2012.
- Jagiella, S. and Zabel, F.: Reaction of phenylperoxy radicals with NO₂ at 298 K, *Phys. Chem. Chem. Phys.*, 9, 5036–5051, <https://doi.org/10.1039/B705193J>, 2007.
- 1005 Jiménez-Muñoz, J. C., Mattar, C., Barichivich, J., Santamaría-Artigas, A., Takahashi, K., Malhi, Y., Sobrino, J. A., and Schrier, G. v. d.: Record-breaking warming and extreme drought in the Amazon rainforest during the course of El Niño 2015–2016, *Scientific Reports*, 6, 33 130, <https://doi.org/10.1038/srep33130>, 2016.
- Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geoscientific Model Development*, 3, 717–752, <https://doi.org/10.5194/gmd-3-717-2010>, 1010 2010.
- Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C. A. M., Brinkop, S., Cai, D. S., Dyroff, C., Eckstein, J., Frank, F., Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, M., Meul, S., Neumaier, M., Nützel, M., Oberländer-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth System Chemistry integrated Modelling (ESCiMo) with the Modular Earth Submodel System (MESSy) version 2.51, *Geoscientific Model Development*, 9, 1153–1200, 1015 <https://doi.org/10.5194/gmd-9-1153-2016>, 2016.
- Jülich Supercomputing Centre: JUWELS: Modular Tier-0/1 Supercomputer at the Jülich Supercomputing Centre, *Journal of large-scale research facilities*, 5, <https://doi.org/10.17815/jlsrf-5-171>, 2019.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- 1020 Kim, P. S., Jacob, D. J., Mickley, L. J., Koplitz, S. N., Marlier, M. E., DeFries, R. S., Myers, S. S., Chew, B. N., and Mao, Y. H.: Sensitivity of population smoke exposure to fire locations in Equatorial Asia, *Atmospheric Environment*, 102, 11 – 17, <https://doi.org/10.1016/j.atmosenv.2014.09.045>, 2015.
- Koss, A. R., Sekimoto, K., Gilman, J. B., Selimovic, V., Coggon, M. M., Zarzana, K. J., Yuan, B., Lerner, B. M., Brown, S. S., Jimenez, J. L., 1025 Krechmer, J., Roberts, J. M., Warneke, C., Yokelson, R. J., and de Gouw, J.: Non-methane organic gas emissions from biomass burning: identification, quantification, and emission factors from PTR-ToF during the FIREX 2016 laboratory experiment, *Atmospheric Chemistry and Physics*, 18, 3299–3319, <https://doi.org/10.5194/acp-18-3299-2018>, 2018.
- Kyrölä, E., Laine, M., Sofieva, V., Tamminen, J., Päivärinta, S.-M., Tukiainen, S., Zawodny, J., and Thomason, L.: Combined SAGE II—GOMOS ozone profile data set for 1984–2011 and trend analysis of the vertical distribution of ozone, *Atmospheric Chemistry and Physics*, 1030 13, 10 645–10 658, <https://doi.org/10.5194/acp-13-10645-2013>, 2013.
- Lee, D., Pitari, G., Grewe, V., Gierens, K., Penner, J., Petzold, A., Prather, M., Schumann, U., Bais, A., Berntsen, T., Iachetti, D., Lim, L., and Sausen, R.: Transport impacts on atmosphere and climate: Aviation, *Atmospheric Environment*, 44, 4678 – 4734, <https://doi.org/10.1016/j.atmosenv.2009.06.005>, transport Impacts on Atmosphere and Climate: The ATTICA Assessment Report, 2010.

- Lee, D., Fahey, D., Skowron, A., Allen, M., Burkhardt, U., Chen, Q., Doherty, S., Freeman, S., Forster, P., Fuglestedt, J., Gettel-
1035 man, A., De León, R., Lim, L., Lund, M., Millar, R., Owen, B., Penner, J., Pitari, G., Prather, M., Sausen, R., and Wilcox, L.:
The contribution of global aviation to anthropogenic climate forcing for 2000 to 2018, *Atmospheric Environment*, 244, 117 834,
<https://doi.org/https://doi.org/10.1016/j.atmosenv.2020.117834>, 2021.
- Lee, H.-H., Bar-Or, R. Z., and Wang, C.: Biomass burning aerosols and the low-visibility events in Southeast Asia, *Atmospheric Chemistry
and Physics*, 17, 965–980, <https://doi.org/10.5194/acp-17-965-2017>, 2017.
- 1040 Lelieveld, J., Gromov, S., Pozzer, A., and Taraborrelli, D.: Global tropospheric hydroxyl distribution, budget and reactivity, *Atmo-
spheric Chemistry and Physics*, 16, 12 477–12 493, <https://doi.org/10.5194/acp-16-12477-2016>, [https://acp.copernicus.org/articles/16/
12477/2016/](https://acp.copernicus.org/articles/16/12477/2016/), 2016.
- Lelieveld, J., Bourtsoukidis, E., Brühl, C., Fischer, H., Fuchs, H., Harder, H., Hofzumahaus, A., Holland, F., Marno, D., Neumaier, M.,
Pozzer, A., Schlager, H., Williams, J., Zahn, A., and Ziereis, H.: The South Asian monsoon—pollution pump and purifier, *Science*, 361,
1045 270–273, <https://doi.org/10.1126/science.aar2501>, 2018.
- Leslie, M. D., Ridoli, M., Murphy, J. G., and Borduas-Dedekind, N.: Isocyanic acid (HNCO) and its fate in the atmosphere: a review, *Environ.
Sci.: Processes Impacts*, 21, 793–808, <https://doi.org/10.1039/C9EM00003H>, 2019.
- Leuenberger, C., Ligocki, M. P., and Pankow, J. F.: Trace organic compounds in rain. 4. Identities, concentrations, and scavenging mech-
anisms for phenols in urban air and rain, *Environmental Science & Technology*, 19, 1053–1058, <https://doi.org/10.1021/es00141a005>,
1050 pMID: 22288749, 1985.
- Li, Q., Jacob, D. J., Bey, I., Yantosca, R. M., Zhao, Y., Kondo, Y., and Notholt, J.: Atmospheric hydrogen cyanide (HCN): Biomass burning
source, ocean sink?, *Geophysical Research Letters*, 27, 357–360, <https://doi.org/10.1029/1999GL010935>, 2000.
- Li, Q., Palmer, P. I., Pumphrey, H. C., Bernath, P., and Mahieu, E.: What drives the observed variability of HCN in the troposphere and lower
stratosphere?, *Atmospheric Chemistry and Physics*, 9, 8531–8543, <https://doi.org/10.5194/acp-9-8531-2009>, 2009.
- 1055 Liu, T., Mickley, L. J., Marlier, M. E., DeFries, R. S., Khan, M. F., Latif, M. T., and Karambelas, A.: Diagnosing spatial biases and
uncertainties in global fire emissions inventories: Indonesia as regional case study, *Remote Sensing of Environment*, 237, 111 557,
<https://doi.org/10.1016/j.rse.2019.111557>, 2020.
- Lobert, J. M., Scharffe, D. H., Hao, W. M., and Crutzen, P. J.: Importance of biomass burning in the atmospheric budgets of nitrogen-
containing gases, *Nature*, 346, 552–554, <https://doi.org/10.1038/346552a0>, 1990.
- 1060 Mahowald, N. M., Rasch, P. J., Eaton, B. E., Whittlestone, S., and Prinn, R. G.: Transport of 222radon to the remote troposphere using
the Model of Atmospheric Transport and Chemistry and assimilated winds from ECMWF and the National Center for Environmental
Prediction/NCAR, *Journal of Geophysical Research: Atmospheres*, 102, 28 139–28 151, <https://doi.org/10.1029/97JD02084>, 1997.
- Marlier, M. E., DeFries, R. S., Voulgarakis, A., Kinney, P. L., Randerson, J. T., Shindell, D. T., Chen, Y., and Faluvegi, G.: El Niño and health
risks from landscape fire emissions in southeast Asia, *Nature Climate Change*, 3, 131–136, <https://doi.org/10.1038/nclimate1658>, 2013.
- 1065 Millet, D. B., Jacob, D. J., Custer, T. G., de Gouw, J. A., Goldstein, A. H., Karl, T., Singh, H. B., Sive, B. C., Talbot, R. W., Warneke, C.,
and Williams, J.: New constraints on terrestrial and oceanic sources of atmospheric methanol, *Atmospheric Chemistry and Physics*, 8,
6887–6905, <https://doi.org/10.5194/acp-8-6887-2008>, 2008.
- Molina, M. J. and Rowland, F. S.: Stratospheric sink for chlorofluoromethanes: chlorine atom-catalysed destruction of ozone, *Nature*, 249,
810–812, <https://doi.org/10.1038/249810a0>, 1974.
- 1070 Nair, P. J., Froidevaux, L., Kuttippurath, J., Zawodny, J. M., Russell III, J. M., Steinbrecht, W., Claude, H., Leblanc, T., van Gi-
jssel, J. A. E., Johnson, B., Swart, D. P. J., Thomas, A., Querel, R., Wang, R., and Anderson, J.: Subtropical and midlati-

- tude ozone trends in the stratosphere: Implications for recovery, *Journal of Geophysical Research: Atmospheres*, 120, 7247–7257, <https://doi.org/10.1002/2014JD022371>, 2015.
- 1075 Natangelo, M., Mangiapan, S., Bagnati, R., Benfenati, E., and Fanelli, R.: Increased concentrations of nitrophenols in leaves from a damaged forestal site, *Chemosphere*, 38, 1495 – 1503, [https://doi.org/10.1016/S0045-6535\(98\)00370-1](https://doi.org/10.1016/S0045-6535(98)00370-1), 1999.
- Nechita-Banda, N., Krol, M., van der Werf, G. R., Kaiser, J. W., Pandey, S., Huijnen, V., Clerbaux, C., Coheur, P., Deeter, M. N., and Röckmann, T.: Monitoring emissions from the 2015 Indonesian fires using CO satellite data, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373, 20170 307, <https://doi.org/10.1098/rstb.2017.0307>, 2018.
- NOAA: Multivariate ENSO Index Version 2 (MEI.v2), <https://psl.noaa.gov/enso/mei/>, (last accessed 19 September 2020), 2020.
- 1080 Nojima, K., Fukaya, K., Fukui, S., and Kanno, S.: Studies on photochemistry of aromatic hydrocarbons II: The formation of nitrophenols and nitrobenzene by the photochemical reaction of benzene in the presence of nitrogen monoxide, *Chemosphere*, 4, 77–82, 1975.
- Nölscher, A., Butler, T., Auld, J., Veres, P., Muñoz, A., Taraborrelli, D., Vereecken, L., Lelieveld, J., and Williams, J.: Using total OH reactivity to assess isoprene photooxidation via measurement and model, *Atmos. Environ.*, 89, 453–463, <https://doi.org/10.1016/j.atmosenv.2014.02.024>, 2014.
- 1085 Orbe, C., Yang, H., Waugh, D. W., Zeng, G., Morgenstern, O., Kinnison, D. E., Lamarque, J.-F., Tilmes, S., Plummer, D. A., Scinocca, J. F., Josse, B., Marecal, V., Jöckel, P., Oman, L. D., Strahan, S. E., Deushi, M., Tanaka, T. Y., Yoshida, K., Akiyoshi, H., Yamashita, Y., Stenke, A., Revell, L., Sukhodolov, T., Rozanov, E., Pitari, G., Visioni, D., Stone, K. A., Schofield, R., and Banerjee, A.: Large-scale tropospheric transport in the Chemistry–Climate Model Initiative (CCMI) simulations, *Atmospheric Chemistry and Physics*, 18, 7217–7235, <https://doi.org/10.5194/acp-18-7217-2018>, 2018.
- 1090 Park, M., Randel, W. J., Emmons, L. K., Bernath, P. F., Walker, K. A., and Boone, C. D.: Chemical isolation in the Asian monsoon anticyclone observed in Atmospheric Chemistry Experiment (ACE-FTS) data, *Atmospheric Chemistry and Physics*, 8, 757–764, <https://doi.org/10.5194/acp-8-757-2008>, 2008.
- Randel, W. J., Park, M., Emmons, L., Kinnison, D., Bernath, P., Walker, K. A., Boone, C., and Pumphrey, H.: Asian Monsoon Transport of Pollution to the Stratosphere, *Science*, 328, 611–613, <https://doi.org/10.1126/science.1182274>, 2010.
- 1095 Reddington, C. L., Yoshioka, M., Balasubramanian, R., Ridley, D., Toh, Y. Y., Arnold, S. R., and Spracklen, D. V.: Contribution of vegetation and peat fires to particulate air pollution in Southeast Asia, *Environmental Research Letters*, 9, 094 006, <https://doi.org/10.1088/1748-9326/9/9/094006>, 2014.
- Rein, G., Cohen, S., and Simeoni, A.: Carbon emissions from smouldering peat in shallow and strong fronts, *Proceedings of the Combustion Institute*, 32, 2489–2496, <https://doi.org/10.1016/j.proci.2008.07.008>, 2009.
- 1100 Rippen, G., Zietz, E., Frank, R., Knacker, T., and Klöpffer, W.: Do airborne nitrophenols contribute to forest decline?, *Environmental Technology Letters*, 8, 475–482, <https://doi.org/10.1080/09593338709384508>, 1987.
- Roberts, J. M., Veres, P. R., Cochran, A. K., Warneke, C., Burling, I. R., Yokelson, R. J., Lerner, B., Gilman, J. B., Kuster, W. C., Fall, R., and de Gouw, J.: Isocyanic acid in the atmosphere and its possible link to smoke-related health effects, *Proceedings of the National Academy of Sciences*, 108, 8966–8971, <https://doi.org/10.1073/pnas.1103352108>, 2011.
- 1105 Rodgers, C. D.: *Inverse Methods for Atmospheric Sounding*, WORLD SCIENTIFIC, <https://doi.org/10.1142/3171>, 2000.
- Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornbluh, L., Manzini, E., Schlese, U., and Schulzweida, U.: Sensitivity of Simulated Climate to Horizontal and Vertical Resolution in the ECHAM5 Atmosphere Model, *Journal of Climate*, 19, 3771–3791, <https://doi.org/10.1175/JCLI3824.1>, 2006.

- Rosanka, S., Frömming, C., and Grewe, V.: The impact of weather patterns and related transport processes on aviation's contribution to ozone and methane concentrations from NO_x emissions, *Atmospheric Chemistry and Physics*, 20, 12 347–12 361, <https://doi.org/10.5194/acp-20-12347-2020>, 2020a.
- Rosanka, S., Sander, R., Franco, B., Wespes, C., Wahner, A., and Taraborrelli, D.: Oxidation of low-molecular weight organic compounds in cloud droplets: global impact on tropospheric oxidants, *Atmospheric Chemistry and Physics Discussions*, 2020, 1–33, <https://doi.org/10.5194/acp-2020-1041>, 2020b.
- 1115 Rosanka, S., Sander, R., Wahner, A., and Taraborrelli, D.: Oxidation of low-molecular weight organic compounds in cloud droplets: development of the JAMOC chemical mechanism in CAABA/MECCA (version 4.5.0gmd), *Geoscientific Model Development Discussions*, 2020, 1–18, <https://doi.org/10.5194/gmd-2020-337>, 2020c.
- Rosanka, S., Vu, G. H. T., Nguyen, H. M. T., Pham, T. V., Javed, U., Taraborrelli, D., and Vereecken, L.: Atmospheric chemical loss processes of isocyanic acid (HNCO): a combined theoretical kinetic and global modelling study, *Atmospheric Chemistry and Physics*, 20, 6671–
- 1120 6686, <https://doi.org/10.5194/acp-20-6671-2020>, 2020d.
- Sander, R., Baumgaertner, A., Cabrera-Perez, D., Frank, F., Gromov, S., Grooß, J.-U., Harder, H., Huijnen, V., Jöckel, P., Karydis, V. A., Niemeyer, K. E., Pozzer, A., Riede, H., Schultz, M. G., Taraborrelli, D., and Tauer, S.: The community atmospheric chemistry box model CAABA/MECCA-4.0, *Geoscientific Model Development*, 12, 1365–1385, <https://doi.org/10.5194/gmd-12-1365-2019>, 2019.
- Schultz, M. G., Stadler, S., Schröder, S., Taraborrelli, D., Franco, B., Krefting, J., Henrot, A., Ferrachat, S., Lohmann, U., Neubauer, D., Siegenthaler-Le Drian, C., Wahl, S., Kokkola, H., Kühn, T., Rast, S., Schmidt, H., Stier, P., Kinnison, D., Tyndall, G. S., Orlando, J. J., and Wespes, C.: The chemistry–climate model ECHAM6.3-HAM2.3-MOZ1.0, *Geoscientific Model Development*, 11, 1695–1723, <https://doi.org/10.5194/gmd-11-1695-2018>, 2018.
- Schummer, C., Groff, C., Al Chami, J., Jaber, F., and Millet, M.: Analysis of phenols and nitrophenols in rainwater collected simultaneously on an urban and rural site in east of France, *Science of The Total Environment*, 407, 5637 – 5643, <https://doi.org/10.1016/j.scitotenv.2009.06.051>, 2009.
- 1130 Sheese, P. E., Walker, K. A., and Boone, C. D.: A global enhancement of hydrogen cyanide in the lower stratosphere throughout 2016, *Geophysical Research Letters*, 44, 5791–5797, <https://doi.org/10.1002/2017GL073519>, 2017.
- Shim, C., Wang, Y., Singh, H. B., Blake, D. R., and Guenther, A. B.: Source characteristics of oxygenated volatile organic compounds and hydrogen cyanide, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD007543>, 2007.
- 1135 Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrakou, T., Müller, J.-F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmospheric Chemistry and Physics*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014.
- Singh, H. B., Salas, L., Herlth, D., Kolyer, R., Czech, E., Viezee, W., Li, Q., Jacob, D. J., Blake, D., Sachse, G., Harward, C. N., Fuelberg, H., Kiley, C. M., Zhao, Y., and Kondo, Y.: In situ measurements of HCN and CH₃CN over the Pacific Ocean: Sources, sinks, and budgets, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD003006>, 2003.
- 1140 Smith, T. E. L., Evers, S., Yule, C. M., and Gan, J. Y.: In Situ Tropical Peatland Fire Emission Factors and Their Variability, as Determined by Field Measurements in Peninsula Malaysia, *Global Biogeochemical Cycles*, 32, 18–31, <https://doi.org/10.1002/2017GB005709>, 2018.
- Stein, O., Schultz, M. G., Bouarar, I., Clark, H., Huijnen, V., Gaudel, A., George, M., and Clerbaux, C.: On the wintertime low bias of Northern Hemisphere carbon monoxide found in global model simulations, *Atmospheric Chemistry and Physics*, 14, 9295–9316, <https://doi.org/10.5194/acp-14-9295-2014>, <https://acp.copernicus.org/articles/14/9295/2014/>, 2014.
- 1145

- Stockwell, C. E., Veres, P. R., Williams, J., and Yokelson, R. J.: Characterization of biomass burning emissions from cooking fires, peat, crop residue, and other fuels with high-resolution proton-transfer-reaction time-of-flight mass spectrometry, *Atmospheric Chemistry and Physics*, 15, 845–865, <https://doi.org/10.5194/acp-15-845-2015>, 2015.
- 1150 Stockwell, C. E., Jayarathne, T., Cochrane, M. A., Ryan, K. C., Putra, E. I., Saharjo, B. H., Nurhayati, A. D., Albar, I., Blake, D. R., Simpson, I. J., Stone, E. A., and Yokelson, R. J.: Field measurements of trace gases and aerosols emitted by peat fires in Central Kalimantan, Indonesia, during the 2015 El Niño, *Atmospheric Chemistry and Physics*, 16, 11 711–11 732, <https://doi.org/10.5194/acp-16-11711-2016>, 2016.
- Strahan, S. E. and Douglass, A. R.: Decline in Antarctic Ozone Depletion and Lower Stratospheric Chlorine Determined From Aura Microwave Limb Sounder Observations, *Geophysical Research Letters*, 45, 382–390, <https://doi.org/10.1002/2017GL074830>, 2018.
- 1155 Tao, Z. and Li, Z.: A kinetics study on reactions of C₆H₅O with C₆H₅O and O₃ at 298 K, *International Journal of Chemical Kinetics*, 31, 65–72, [https://doi.org/10.1002/\(SICI\)1097-4601\(1999\)31:1<65::AID-KIN8>3.0.CO;2-J](https://doi.org/10.1002/(SICI)1097-4601(1999)31:1<65::AID-KIN8>3.0.CO;2-J), 1999.
- Taraborrelli, D., Lawrence, M. G., Butler, T. M., Sander, R., and Lelieveld, J.: Mainz Isoprene Mechanism 2 (MIM2): an isoprene oxidation mechanism for regional and global atmospheric modelling, *Atmospheric Chemistry and Physics*, 9, 2751–2777, <https://doi.org/10.5194/ACP-9-2751-2009>, 2009.
- 1160 Taraborrelli, D., Lawrence, M. G., Crowley, J. N., Dillon, T. J., Gromov, S., Groß, C. B. M., Vereecken, L., and Lelieveld, J.: Hydroxyl radical buffered by isoprene oxidation over tropical forests, *Nature Geoscience*, 5, 190–193, <https://doi.org/10.1038/ngeo1405>, 2012.
- Taraborrelli, D., Cabrera-Perez, D., Bacer, S., Gromov, S., Lelieveld, J., Sander, R., and Pozzer, A.: Influence of aromatics on tropospheric gas-phase composition, *Atmospheric Chemistry and Physics*, 21, 2615–2636, <https://doi.org/10.5194/acp-21-2615-2021>, 2021.
- Tost, H., Jöckel, P., Kerkweg, A., Sander, R., and Lelieveld, J.: Technical note: A new comprehensive SCAVENGING submodel for global atmospheric chemistry modelling, *Atmospheric Chemistry and Physics*, 6, 565–574, <https://doi.org/10.5194/acp-6-565-2006>, 2006.
- 1165 Trenberth, K. E.: The Definition of El Niño, *Bulletin of the American Meteorological Society*, 78, 2771–2778, [https://doi.org/10.1175/1520-0477\(1997\)078<2771:TDOENO>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<2771:TDOENO>2.0.CO;2), 1997.
- United Nations, Department of Economic and Social Affairs, Population Division: World Population Prospects 2019: Data Booklet, https://population.un.org/wpp/Publications/Files/WPP2019_DataBooklet.pdf, ST/ESA/SER.A/424, 2019.
- 1170 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth System Science Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- Vereecken, L., Chakravarty, H. K., Bohn, B., and Lelieveld, J.: Theoretical Study on the Formation of H- and O-Atoms, HONO, OH, NO, and NO₂ from the Lowest Lying Singlet and Triplet States in Ortho-Nitrophenol Photolysis, *International Journal of Chemical Kinetics*, 48, 785–795, <https://doi.org/10.1002/kin.21033>, 2016.
- 1175 Vigouroux, C., Blumenstock, T., Coffey, M., Errera, Q., García, O., Jones, N. B., Hannigan, J. W., Hase, F., Liley, B., Mahieu, E., Mellqvist, J., Notholt, J., Palm, M., Persson, G., Schneider, M., Servais, C., Smale, D., Thölix, L., and De Mazière, M.: Trends of ozone total columns and vertical distribution from FTIR observations at eight NDACC stations around the globe, *Atmospheric Chemistry and Physics*, 15, 2915–2933, <https://doi.org/10.5194/acp-15-2915-2015>, 2015.
- 1180 Vogel, B., Günther, G., Müller, R., Grooß, J.-U., and Riese, M.: Impact of different Asian source regions on the composition of the Asian monsoon anticyclone and of the extratropical lowermost stratosphere, *Atmospheric Chemistry and Physics*, 15, 13 699–13 716, <https://doi.org/10.5194/acp-15-13699-2015>, 2015.

- Walker, J. C., Dudhia, A., and Carboni, E.: An effective method for the detection of trace species demonstrated using the MetOp Infrared Atmospheric Sounding Interferometer, *Atmospheric Measurement Techniques*, 4, 1567–1580, <https://doi.org/10.5194/amt-4-1567-2011>, 2011.
- 1185
- Wang, Z., Nicholls, S. J., Rodriguez, E. R., Kummu, O., Hörkkö, S., Barnard, J., Reynolds, W. F., Topol, E. J., DiDonato, J. A., and Hazen, S. L.: Protein carbamylation links inflammation, smoking, uremia and atherogenesis, *Nature Medicine*, 13, 1176–1184, <https://doi.org/10.1038/nm1637>, 2007.
- Weng, H., Ashok, K., Behera, S. K., Rao, S. A., and Yamagata, T.: Impacts of recent El Niño Modoki on dry/wet conditions in the Pacific rim during boreal summer, *Climate Dynamics*, 29, 113–129, <https://doi.org/10.1007/s00382-007-0234-0>, 2007.
- 1190
- Whitburn, S., Van Damme, M., Clarisse, L., Bauduin, S., Heald, C. L., Hadji-Lazarou, J., Hurtmans, D., Zondlo, M. A., Clerbaux, C., and Coheur, P.-F.: A flexible and robust neural network IASI-NH₃ retrieval algorithm, *Journal of Geophysical Research: Atmospheres*, 121, 6581–6599, <https://doi.org/10.1002/2016jd024828>, 2016a.
- Whitburn, S., Van Damme, M., Clarisse, L., Turquety, S., Clerbaux, C., and Coheur, P.-F.: Doubling of annual ammonia emissions from the peat fires in Indonesia during the 2015 El Niño, *Geophysical Research Letters*, 43, 11,007–11,014, <https://doi.org/10.1002/2016gl070620>, 2016b.
- 1195
- Wilkerson, J. T., Jacobson, M. Z., Malwitz, A., Balasubramanian, S., Wayson, R., Fleming, G., Naiman, A. D., and Lele, S. K.: Analysis of emission data from global commercial aviation: 2004 and 2006, *Atmospheric Chemistry and Physics*, 10, 6391–6408, <https://doi.org/10.5194/acp-10-6391-2010>, <https://acp.copernicus.org/articles/10/6391/2010/>, 2010.
- 1200
- Xu, J., Morris, P. J., Liu, J., and Holden, J.: PEATMAP: Refining estimates of global peatland distribution based on a meta-analysis, <https://doi.org/10.5518/252>, (Dataset), 2017.
- Xu, J., Morris, P. J., Liu, J., and Holden, J.: PEATMAP: Refining estimates of global peatland distribution based on a meta-analysis, *CATENA*, 160, 134 – 140, <https://doi.org/10.1016/j.catena.2017.09.010>, 2018.
- Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the Last Glacial Maximum, *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2010GL043584>, 2010.
- 1205
- Yuan, C., Lau, W. K. M., Li, Z., and Cribb, M.: Relationship between Asian monsoon strength and transport of surface aerosols to the Asian Tropopause Aerosol Layer (ATAL): interannual variability and decadal changes, *Atmospheric Chemistry and Physics*, 19, 1901–1913, <https://doi.org/10.5194/acp-19-1901-2019>, 2019.
- Zhang, C.: Madden–Julian Oscillation: Bridging Weather and Climate, *Bulletin of the American Meteorological Society*, 94, 1849–1870, <https://doi.org/10.1175/BAMS-D-12-00026.1>, 2013.
- 1210
- Zhang, K., Wan, H., Zhang, M., and Wang, B.: Evaluation of the atmospheric transport in a GCM using radon measurements: sensitivity to cumulus convection parameterization, *Atmospheric Chemistry and Physics*, 8, 2811–2832, <https://doi.org/10.5194/acp-8-2811-2008>, 2008.
- Zheng, B., Chevallier, F., Yin, Y., Ciais, P., Fortems-Cheiney, A., Deeter, M. N., Parker, R. J., Wang, Y., Worden, H. M., and Zhao, Y.: Global atmospheric carbon monoxide budget 2000–2017 inferred from multi-species atmospheric inversions, *Earth System Science Data*, 11, 1411–1436, <https://doi.org/10.5194/essd-11-1411-2019>, 2019.
- 1215