

Mountain-wave-induced polar stratospheric clouds and their representation in the global chemistry model ICON-ART

Michael Weimer^{1,2,*}, Jennifer Buchmüller^{2,†}, Lars Hoffmann³, Ole Kirner¹, Beiping Luo⁴, Roland Ruhnke², Michael Steiner⁵, Ines Tritscher⁶, and Peter Braesicke²

¹Steinbuch Centre for Computing, Karlsruhe Institute of Technology (KIT), Eggenstein-Leopoldshafen, Germany

²Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology (KIT), Eggenstein-Leopoldshafen, Germany

³Jülich Supercomputing Centre, Forschungszentrum Jülich, Jülich, Germany

⁴Institute for Atmospheric and Climate Science, ETH Zurich, Switzerland

⁵Laboratory for Air Pollution / Environmental Technology, EMPA, Switzerland

⁶Institute of Energy and Climate Research: Stratosphere (IEK-7), Forschungszentrum Jülich, Jülich, Germany

* now at: Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA, USA

† now at: Steinbuch Centre for Computing, Karlsruhe Institute of Technology (KIT), Eggenstein-Leopoldshafen, Germany

Correspondence: Michael Weimer (michael.weimer@kit.edu)

Abstract. Polar stratospheric clouds (PSCs) are a driver for ozone depletion in the lower polar stratosphere. They provide surface for heterogeneous reactions activating chlorine and bromine reservoir species during the polar night. The large-scale effects of PSCs are represented by means of parametrisations in current global chemistry-climate models, but one process is still a challenge: the representation of PSCs formed locally in conjunction with unresolved mountain waves. In this study, we

5 investigate direct simulations of PSCs formed by mountain waves with the ICOSahedral Nonhydrostatic modelling framework (ICON) with its extension for Aerosols and Reactive Trace gases (ART) including local grid refinements (nesting) with two-way interaction. Here, the nesting is set up around the Antarctic Peninsula which is a well-known hot spot for the generation of mountain waves in the southern hemisphere. We compare our model results with satellite measurements of PSCs from the

10 Cloud-Aerosol Lidar with Orthogonal Polarisation (CALIOP) and gravity wave observations of the Atmospheric InfraRed

Sounder (AIRS). For a mountain wave event from 19 to 29 July 2008 we find similar structures of PSCs as well as a fairly realistic development of the mountain wave between the satellite data and the ICON-ART simulations in the Antarctic Peninsula nest. We compare a global simulation without nesting with the nested configuration to show the benefits of adding the nesting. Although the mountain waves cannot be resolved explicitly in the used global resolution (about 160 km), their effect from the nested regions (about 80 and 40 km) on the global domain is represented. Thus, we show in this study that the ICON-ART

15 model has the potential to bridge the gap between directly resolved mountain-wave-induced PSCs and their representation and effect on chemistry in coarse global resolutions.

1 Introduction

Polar stratospheric clouds (PSCs) play a key role in explaining the rapid ozone loss in the polar stratosphere during local spring (e.g., Solomon et al., 1986; Solomon, 1999; Braesicke et al., 2018). Three different types of PSCs or mixtures thereof can be found in the lower stratosphere: solid nitric acid trihydrate particles (NAT), liquid supercooled ternary solution droplets (STS) and ice particles (e.g., Peter and GroöB, 2012; Tritscher et al., 2021, and references therein). Heterogeneous reactions on the surface of PSCs lead to activation of chlorine and bromine species during the polar night, thus enhancing the catalytic ozone depletion cycles as soon as the sun rises (e.g., Solomon, 1999). In addition, PSCs can irreversibly remove nitrogen-containing species by sedimentation, a process known as denitrification, thus extending the period of low ozone concentrations during polar spring (e.g., Waibel et al., 1999). PSCs may form below specific threshold temperatures (e.g., Hanson and Mauersberger, 1988; Marti and Mauersberger, 1993; Carslaw et al., 1994), indicating that temperature is a crucial parameter for PSC formation.

Mountain waves (orographic gravity waves) are stationary waves in the lee of a mountain which can develop in a stably stratified atmosphere (e.g., Fritts and Alexander, 2003) when a sizeable component of the large scale flow is perpendicular to the mountain range. Mountain waves can propagate upwards into the stratosphere and higher (Wright et al., 2017) and may perturb the synoptic temperature field with local amplitudes of up to ± 15 K or even more (Carslaw et al., 1998b; Eckermann et al., 2009; Dörnbrack et al., 2020). In the polar regions, mountain waves are of particular interest for the formation of PSCs because mountain-wave-induced temperature fluctuations can lead to localised cooling that triggers PSC formation even if synoptic-scale temperatures are above the PSC formation temperatures (Carslaw et al., 1998b).

Mountain-wave-induced PSCs have a significant influence on the ozone depletion over both Antarctica and the Arctic (e.g., Höpfner et al., 2006a; McDonald et al., 2009; Alexander et al., 2011; Hoffmann et al., 2017; Langematz et al., 2018). Various regions have been identified from observations and models as being hot spots for mountain-wave-induced PSCs, such as Greenland and Scandinavia in the northern hemisphere and the Southern Andes, the Antarctic Peninsula or the Transantarctic Mountains in the southern hemisphere (Dörnbrack et al., 2002; Plougonven et al., 2008; Eckermann et al., 2009; Noel et al., 2009; Hoffmann et al., 2013, 2017). McDonald et al. (2009) estimated that up to 40 % of Antarctic PSC formation are associated with mountain waves in the early Antarctic winter when temperatures are close to the NAT formation threshold. Alexander et al. (2013) concluded that about 5 % of Antarctic and 12 % of Arctic PSCs are related to mountain wave activity.

Mountain waves are mesoscale features of the atmospheric circulation, which poses a challenge for simulating them explicitly with global general circulation models because these models have limited spatial resolution. The horizontal wavelengths of mountain waves most relevant for the middle atmosphere are in the range of tens to hundreds of kilometres (e.g., Fritts and Alexander, 2003; Eckermann et al., 2006; Orr et al., 2020). Several studies suggest that in a discrete numerical model eight grid points are needed to represent gravity waves and their dynamics adequately (Geller et al., 2013; Preusse et al., 2014; Kang et al., 2017). However, current global chemistry climate models (CCMs) with horizontal resolutions in the order of a few hundreds of kilometres are not able to resolve the full range of mountain waves adequately (Lamarque et al., 2013; Orr et al., 2015; Morgenstern et al., 2017). Mesoscale models with resolutions as high as 7 km have been developed in the past to calculate the

local effect of mountain-wave-induced PSCs (e.g., Fueglistaler et al., 2003; Eckermann et al., 2006; Plougonven et al., 2008; Noel and Pitts, 2012) but they need input of a previous simulation of a global model or a reanalysis to provide the boundary conditions (e.g., Weimer et al., 2016). Thus, mountain waves and mountain-wave-induced PSCs either have to be parametrised (Orr et al., 2015; Zhu et al., 2017; Orr et al., 2020) or have to be calculated in a post-processing step via Lagrangian models (e.g., Mann et al., 2005) or via mesoscale models. An approach for interactive two-way coupling between the high-resolution simulations and the global models, in particular CCMs, is missing so far.

In this study, mountain-wave-induced PSCs are simulated seamlessly with the ICOSahedral Nonhydrostatic modelling framework (ICON, Zängl et al., 2015) and its extension for Aerosols and Reactive Trace gases (ART, Rieger et al., 2015; Weimer et al., 2017; Schröter et al., 2018). ICON-ART provides the possibility of local grid refinement (nesting) with two-way interaction (see Reinert et al., 2019, for details). Thus, a global low-resolution simulation provides boundary conditions for a region with refined grid, similar to mesoscale models. But additionally, the refined grid also feeds back to the low resolution which is a novel approach in atmospheric chemistry modelling.

We perform a comprehensive case study for a well-observed mountain wave event at the Antarctic Peninsula in July 2008 (Noel and Pitts, 2012). We start at a global resolution of about 160 km which is comparable to other global CCMs. We then apply the two-way nesting at the Antarctic Peninsula with a resolution of 40 km in the grid refinement. This resolution still misses directly resolving gravity waves with horizontal wavelengths lower than about 300 km (cf., Geller et al., 2013), but we chose this configuration (1) for a balance between accuracy and computational expense and (2) to show how CCMs could already benefit from modest higher resolutions. We show how the higher resolution in the refinement impacts the gravity wave dynamics, PSCs and finally the ozone chemistry in the global model. Thus, this study is a first step to close the gap between direct simulations of mountain-wave-induced PSC formation and their treatment in coarse global resolutions.

We selected the Antarctic Peninsula for this study with the ICON-ART model because it is a well-known hot spot of mountain-wave-induced PSCs (Bacmeister, 1993; Bacmeister et al., 1994; McDonald et al., 2009; Alexander et al., 2011; Hoffmann et al., 2017). Given the strong incident flow at low levels and the Antarctic Peninsula protruding a distance of 1300 km with peak heights of up to 2800 m posing a substantial barrier to this flow, mountain waves with large amplitudes, and mesoscale horizontal (> 300 km) and vertical wavelengths (> 10 km) are typically excited (Alexander and Teitelbaum, 2007; Plougonven et al., 2008; Hoffmann et al., 2013; Orr et al., 2020). Thus, our ICON-ART model configuration is expected to be able to capture such waves.

Despite the Antarctic Peninsula being in a remote location, it is frequently covered by gravity wave and PSC satellite observations that allows to evaluate the ICON-ART simulations (e.g., Hoffmann et al., 2016; Spang et al., 2018; Höpfner et al., 2018; Pitts et al., 2018). Here, we specifically use Atmospheric InfraRed Sounder (AIRS, Aumann et al., 2003; Chahine et al., 2006) and Cloud-Aerosol Lidar with Orthogonal Polarisation (CALIOP, Pitts et al., 2009; Höpfner et al., 2009; Pitts et al., 2018) data to evaluate our simulations. AIRS is a cross-track scanning nadir instrument that is able to detect stratospheric temperature fluctuations with high horizontal resolution (up to 14 km at nadir). Therefore, it is particularly suited to detect the horizontal structures of mountain waves in the lower and mid stratosphere (e.g., Hoffmann et al., 2013, 2017). CALIOP, a nadir lidar instrument with high resolution in both horizontal (5 km) and vertical (180 m in the lower stratosphere) direction, can detect

and discriminate the different types of PSCs (Pitts et al., 2018). Therefore, CALIOP is able to identify PSCs formed in mountain waves and can be used for the evaluation of PSC schemes in models. Sampling the model results on the measurement grid of the satellite instruments and converting the data from model quantities to measured quantities (e.g., by means of radiative transfer calculations) is a precise way to evaluate the ICON-ART simulations with AIRS and CALIOP observations (Mishchenko et al., 1996; Grimsdell et al., 2010; Orr et al., 2015).

This study is organised as follows: Section 2 briefly describes the ICON-ART model and its PSC scheme. In Sect. 3, the simulation setup is pointed out that is used to examine the mountain wave event at the Antarctic Peninsula. This is followed by a description of the AIRS and CALIOP instruments in Sect. 4. The model results are compared with CALIOP and AIRS measurements and the impact of the two-way nesting on the chemistry is investigated in Sect. 5. Finally, conclusions and outlook follow in Sect. 6.

2 The ICON-ART model

The ICON model is the operational model for numerical weather prediction at the German Weather Service (DWD, Zängl et al., 2015). In addition, it can be applied to large eddy simulations (Dipankar et al., 2015) and fully coupled with an ocean and a land surface model for climate integrations with the climate physics configuration (Giorgetta et al., 2018). The ART extension has been developed to incorporate aerosols and the atmospheric chemistry into ICON. It can be coupled to ICON in configurations for numerical weather prediction (Rieger et al., 2015) and allows flexible configurations for weather and climate integrations (Schröter et al., 2018).

The PSC scheme in ICON-ART calculates ice PSCs based on the microphysics of the operational configuration at DWD (Doms et al., 2011). For stratospheric temperatures, the microphysics assumes an ice number concentration of 0.25 cm^{-3} . Liquid (sulfate and STS) particles are formed by the module of Carslaw et al. (1995), with some adaption of the used size distributions. NAT particles are formed by a non-equilibrium approach based on Carslaw et al. (2002) and van den Broek et al. (2004). The NAT size distribution can be flexibly selected by the model user via XML files (see Schröter et al., 2018). A maximum NAT number concentration of $2.3 \times 10^{-4} \text{ cm}^{-3}$ is assumed (van den Broek et al., 2004). The PSC particles are treated separately and externally mixed in each grid box. A detailed description of the PSC scheme can be found in Appendix A.

While STS particles are calculated diagnostically, NAT particles are separate tracers for each size bin. Both NAT and ice include prognostic equations that allow the advection of these particles into regions with temperatures too large for PSC formation, which has been observed in mountain waves (Eckermann et al., 2009). The other parts of the scheme are designed similar to global models like ECHAM/MESSy Atmospheric Chemistry model (EMAC, Jöckel et al., 2010; Kirner et al., 2011) which have been shown to reflect the main properties of PSCs on a global scale. This study focuses on the benefit of increasing the resolution and showing the impact of the two-way nesting.

The chemistry in ICON-ART is based on the Module Efficiently Calculating the Chemistry of the Atmosphere (MECCA, Sander et al., 2011a) which uses the Kinetic PreProcessor (KPP) to generate Fortran files for solving the specified chemical




Figure 1. Simulation setup in this study: A free-running simulation from 01 March 2008 until 30 April 2008 is followed by a period until 18 July 2008 where the meteorology is reinitialised every second day. During the mountain wave event until 29 July 2008, two simulations are performed: one global simulation without nests (with 160 km resolution) and one simulation with the nests (with 160, 80 and 40 km resolution) as visualised in Fig. 2 including two-way nesting.

mechanism (Sandu and Sander, 2006). Photolysis rates are calculated with the CloudJ module (Prather, 2015; Weimer et al., 2017). A system of 142 chemical reactions including 38 photolytic reactions and 11 heterogeneous reactions on the surface of PSCs is used. It covers the chemical families O_x , HO_x , NO_x , ClO_x , BrO_x , a basic hydrocarbon chemistry and the oxidation of SO_2 . The reaction system is similar to other studies (e.g., Stone et al., 2019; Zambri et al., 2019; Nakajima et al., 2020) and can be found in the supplement. Trace gas emissions at the Earth’s surface are included by a module described in Weimer et al. (2017).

The model equations of ICON-ART are discretised horizontally on an icosahedral-triangular C grid (e.g., Staniforth and Thuburn, 2012; Zängl et al., 2015). The global resolution can be refined by root divisions and bisections of the original icosahedron, resulting in the horizontal resolution description $RnBk$, as defined by e.g. Zängl et al. (2015). Vertical discretisation is performed on generalised smooth-level coordinates (Leuenberger et al., 2010).

For the purpose of detailed simulations around a specific region, the grid can be refined for the area of interest by further bisections. Here, the parent domain provides boundary conditions for the nested domain. The simulated values in the nested domain are interpolated to the parent grid with a relaxation-based method (Reinert et al., 2019). Thus, the global domain is nudged towards the values in the nests, which will be further investigated in Sect. 5.

3 Simulation with nests around the Antarctic Peninsula

In 2008, a mountain-wave event took place between 19 and 29 July around the Antarctic Peninsula (Noel and Pitts, 2012) which is further investigated in this study. A three-step simulation is conducted, see Fig. 1.

In the first step, the simulation starts on 01 March 2008 with a global resolution of R2B04 (Δx of about 160 km) as a free-running simulation until 01 May 2008. The first of March is chosen because almost no PSCs are formed at this time either in the northern or in the southern hemisphere (Tully et al., 2011). This period is used as a spin-up period for the chemistry until the southern hemispheric polar vortex intensifies (Schoeberl and Newman, 2003) and PSCs formed in 2008 (Tully et al., 2011).




Figure 2. Visualisation of the nested domains used in the simulation with the nests: A global resolution of R2B04 ($\Delta x \approx 160$ km) is used, with first circular refined grid around the Antarctic Continent (R2B05, $\Delta x \approx 80$ km) and a second rectangular refinement around the Antarctic Peninsula (R2B06, $\Delta x \approx 40$ km). The white line shows the location of the cross section analysed in Sect. 5.3.

Table 1. Overview of the simulation setup for the investigation of the mountain wave event in July 2008. Details see text.

Time period (in 2008)	Resolution	Output interval (h)	Remark
01 March – 30 April	R2B04	24	Spin-up for chemistry, free-running
01 May – 18 July	R2B04	24	Dynamics reinitialised every second day
19 July – 29 July	R2B04	1	Free-running
	see Fig. 2	see text	Including two-way nesting

140 In the second step, the meteorological variables are reinitialised every second day by the reanalysis product of ECMWF, ERA-Interim (Dee et al., 2011), in the period between 01 May and 18 July 2008, but the chemical tracers are free-running. This ensures a comparable evolution of the polar vortex in the model with respect to the reanalysis. This method was already introduced e.g. in Schröter et al. (2018).

145 In the third step, we conducted two simulations covering the mountain wave event from 19 to 29 July 2008: one simulation without any nests and one simulation with two-way nesting around the Antarctic Continent (R2B05, Δx of about 80 km) and around the Antarctic Peninsula (R2B06, Δx of about 40 km), see Fig. 2 and Table 1. The chemistry, including PSCs, photolysis and transport of tracers, is calculated in all domains. Since we are interested in the interaction between the model domains, we decided to avoid every influence by other models during this part of the simulation and hence they are free-running. Output in this third step of the simulation is given (1) at the specific Antarctic Peninsula overpasses of AIRS for the Antarctic Peninsula
150 nest and (2) hourly during the whole period of the mountain wave event.

On 01 March 2008, the meteorological variables are initialised with ERA-Interim. The chemical tracers are initialised by an EMAC simulation which included tropospheric as well as stratospheric chemistry similar to Jöckel et al. (2016). Sea surface temperature and sea ice cover are based on monthly varying values of the climatology by Taylor et al. (2000), linearly interpolated to the simulation date. The advective model time step is set to 360 s. Vertically, the same 90 levels are used as in

Table 2. Emission datasets used in this study.

Species	GEIA ^a	MACCcity ^b	MEGAN-MACC ^c	GFED3 ^d	EDGARv4.2 ^e
CH ₄	–	✓	✓	✓	–
CO	–	✓	✓	–	✓
CO ₂	–	–	–	✓	✓
N ₂ O	–	–	–	✓	✓
SO ₂	–	✓	–	✓	–
CFCl ₃	✓	–	–	–	–

^a Cunnold et al. (1994)

^b van der Werf et al. (2006); Lamarque et al. (2010); Granier et al. (2011); Diehl et al. (2012)

^c Sindelarova et al. (2014)

^d van der Werf et al. (2010)

^e Janssens-Maenhout et al. (2011, 2013)

155 the operational setup of DWD weather forecasts, covering the altitude range from the surface up to 75 km (see e.g., Weimer et al., 2017, Fig. 1). In the lower stratosphere, the vertical grid spacing increases from 400 m at an altitude of 12 km up to about 1200 m at 30 km.

The emission datasets used in this study are summarised in Table 2. Emissions of CH₄, CO, CO₂, N₂O, SO₂ and CFCl₃ are considered. The GEIA dataset for chlorofluorocarbons (CFCs) provided by the Emissions of atmospheric Compounds and Compilation of Ancillary Data base (ECCAD) includes only the year 1986 and should eventually be adapted by the online
160 emission tool by Jähn et al. (2020) for the simulated year. Since the emission rates of CFCl₃ have decreased by more than 40 % since 1986 (Montzka et al., 2018), we neglect the emissions of other CFCs for the less than 1-year simulation. In combination with the emission tool by Jähn et al. (2020) and in the context of the recently found source of CFCs (Montzka et al., 2018; Lickley et al., 2020), further CFCs should be considered in future simulations.

165 NAT PSCs are simulated using a size distribution based on van den Broek et al. (2004), shown in Fig. 3. We prescribe H₂SO₄ in the lower stratosphere (Thomason et al., 2008; SPARC, 2013) in the global domain. In the nested domains, H₂SO₄ runs freely after initialised by the parent domain.

4 Satellite datasets

4.1 CALIOP

170 The Cloud-Aerosol Lidar with Orthogonal Polarisation (CALIOP, Pitts et al., 2009; Höpfner et al., 2009; Pitts et al., 2018) onboard the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) was launched on 28 April 2006 (Winker et al., 2007). In 2008, the satellite flew as part of NASA’s A-train constellation in an orbit with 98° inclination at an




Figure 3. Size distribution of NAT particles used in this study. Based on van den Broek et al. (2004).

altitude of 705 km (Stephens et al., 2002; Pitts et al., 2018). Its orbit was sun-synchronous with equator crossings at about 01:30 and 13:30 LT. It measured down to 82°S with a repeat cycle of 16 days. In February 2018, it was moved to a lower orbit.

175 CALIOP is a light detecting and ranging (lidar) instrument, set up in nadir geometry. It scans the atmosphere at wavelengths of 532 and 1064 nm with parallel and orthogonal polarisations for the 532-nm channel (Winker et al., 2007). In altitudes from 8.4 to 30 km, the vertical resolution is 180 m or higher. Horizontal averaging of 5 km is applied for the detection of PSCs with CALIOP (Pitts et al., 2018). At the Earth’s surface, the light beam has a diameter of about 100 m (Höpfner et al., 2009; Pitts et al., 2018).

180 For the detection of PSCs, a combination of two values is used (Pitts et al., 2018): (1) the ratio of total and molecular backscatter coefficient R_{532} and (2) the backscatter coefficient at perpendicular polarisation β_{\perp} , both at a wavelength of 532 nm. Discrimination of the PSC types can be seen in diagrams of R_{532} vs. β_{\perp} . Different regions in this diagram refer to the different PSC categories: STS, NAT mixtures, enhanced NAT mixtures, Ice and Wave-ice (see Pitts et al., 2018). The thresholds to distinguish PSCs from background noise are calculated as daily median plus one absolute standard deviation and depend on potential temperature ($\beta_{\perp, \text{thres}}$ and $R_{532, \text{thres}}$). Particles with $\beta_{\perp} < \beta_{\perp, \text{thres}}$ and $R_{532} > R_{532, \text{thres}}$ are attributed to the STS category. If $\beta_{\perp} > \beta_{\perp, \text{thres}}$, non-spherical particles are assumed. Pitts et al. (2018) estimated that 10 to 15 % of particles classified as NAT mixtures and STS could be misclassified and may lead to enhancements in the respective other class. The boundary between the NAT mixtures and ice ($R_{\text{NAT|ice}}$) is calculated dynamically depending on the state of dehydration and denitrification (Pitts et al., 2018). The category of enhanced NAT mixtures represents NAT particles nucleated heterogeneously on wave ice PSCs. Both the enhanced NAT mixtures and Wave-ice categories depend on empirically set thresholds of β_{\perp} and R_{532} and therefore a not “all-inclusive” (Pitts et al., 2018). Overall, the classification scheme has been shown to be applicable to ground-based lidars with comparable results (Snels et al., 2019) and has been shown to be in the expected thermodynamic existence regimes (Pitts et al., 2018).

185

190

We use the PSC climatology of Pitts et al. (2018), which is the version 2 level 1B data. It is restricted to night-time southern
195 hemispheric data during the mountain wave event (data is available within the time frame we are interested in from 22 to 29
July 2008). We examine the altitude levels between 15 and 30 km where (1) no tropospheric clouds contaminate the results
and (2) most of the PSCs can be found (see, e.g., Fig. 13 of Pitts et al., 2018). In addition, the dataset includes temperature and
pressure data of Modern-Era Retrospective analysis for Research and Applications version 2 (MERRA2, Gelaro et al., 2017),
which originally has a horizontal resolution of $0.5^\circ \times 0.625^\circ$, interpolated on the CALIOP paths.

200 4.2 AIRS

The Atmospheric InfraRed Sounder (AIRS, Aumann et al., 2003; Chahine et al., 2006) is one of the instruments onboard the
Aqua satellite, which was launched in May 2002 and which is part of the A-train constellation right ahead of CALIPSO. Thus,
its orbit is the same as CALIPSO in 2008 but about 1 to 2 min ahead of CALIPSO¹.

The AIRS instrument is a nadir sounder with across-track scanning capabilities that scans the atmosphere by 90 footprints
205 per scan with a ground coverage of 1780 km and a size of $13.5 \times 13.5 \text{ km}^2$ (nadir) to $41 \times 21.4 \text{ km}^2$ (scan edge) per footprint
(e.g., Orr et al., 2015; Hoffmann et al., 2017). It measures the spectrally resolved radiances in wavelengths between 3.74
and $15.4 \mu\text{m}$. Brightness temperatures in the $15 \mu\text{m}$ band can be used to derive information about gravity waves in the lower
polar stratosphere (Hoffmann et al., 2017). In this study we use a data product averaging over 21 channels around $15 \mu\text{m}$ to
improve the signal-to-noise ratio (Hoffmann et al., 2017). The temperature weighting function in this band peaks at an altitude
210 of around 23 km with a full width at half maximum of 15 km and with information from the altitude range between 17 and
32 km. Therefore, it is well suited to derive information about gravity waves in the altitude region where PSCs are expected to
exist.

This band is used to examine the mountain wave event at the Antarctic Peninsula mentioned above. The same algorithm as
in previous studies is used to compare the model data with specific Antarctic Peninsula overpasses of AIRS (Hoffmann and
215 Alexander, 2010; Hoffmann et al., 2016, 2017). In particular, the ICON-ART data are resampled on the AIRS measurement
grid and a radiative transfer model is used to simulate AIRS measurements based on the ICON-ART data to allow for a direct
comparison with the real observations.

5 Mountain-wave-induced PSCs with ICON-ART

Here, we investigate a mountain wave event during July 2008 with ICON-ART in a configuration with interactive chemistry
220 and local grid refinement around the Antarctic Peninsula. This section comprises an evaluation of the dynamical structure of
the mountain wave with AIRS (Sect. 5.1) and comparisons of the model results with CALIOP measurements (Sect. 5.2). In
addition, it discusses the impact of the direct simulation of mountain-wave-induced PSCs on the polar ozone chemistry (Sect.
5.3).

¹<https://www-calipso.larc.nasa.gov/about/atrain.php>, last access 19 March 2021




Figure 4. Meteorological situation at the Antarctic Peninsula for a) 19 and b) 22 July 2008 at 00 UTC from ERA-Interim. The colours and the wind barbs depict the surface wind in knots. The white lines correspond to the mean sea level pressure in hPa. The black lines are geopotential heights at 500 hPa in gpdam. The shaded regions show the air masses within the polar vortex, determined according to Nash et al. (1996) on $\theta = 475$ K. Panels covering the whole period of the mountain wave event can be found in the supplement.

5.1 Near-surface meteorological conditions and stratospheric dynamical structure of the mountain wave

225 The evolution of the ozone loss and the Antarctic polar vortex in 2008 was comparable to the previous years although the so-called ozone hole lasted rather long into December (Tully et al., 2011). It was a year with increased gravity wave activity at the Antarctic Peninsula (Hoffmann et al., 2016). We chose a gravity wave event lasting ten days from 19 July until 29 July 2008 with lowest temperatures during the whole winter season (Noel and Pitts, 2012). By end of July 2008, the polar vortex was close to its maximum extension whereas the stratospheric ozone concentration started to decrease (Tully et al., 2011).

230 Figure 4 summarises the large-scale meteorological conditions around the Peninsula, based on the ERA-Interim reanalysis. As indicated e.g. by Alexander and Teitelbaum (2007), the ability of ERA-Interim to capture mesoscale mountain wave events is limited. Therefore, we use the reanalysis to show the meteorological background conditions and will evaluate the lower stratospheric temperature perturbations with AIRS later in this section. Figure 4 shows several variables at the beginning of the mountain wave event on 19 July 2008 (panel a) and around its peak dynamics on 22 July 2008 (panel b). Panels covering the
235 whole event can be found in the supplement (Figure S1).

The Antarctic Peninsula was located at the vortex edge during the whole mountain wave event, as depicted by the shaded regions in the panels and determined by the method by Nash et al. (1996). Starting from 19 July 2008, an approaching high pressure system led to an increase of the mean sea level pressure gradient (white lines) at the Antarctic Peninsula with a corresponding increase of easterly winds at the mountain range (wind barbs and colour-coded). The gradient on the geopotential
240 height at 500 hPa (black lines) was also increased showing the large-scale easterly flow at this altitude on 22 July.

These conditions led to a mountain wave event lasting for ten days which is now compared to AIRS measurements. For this, temperature and pressure of ICON-ART in the Antarctic Peninsula nest is saved at the time step closest to each of the AIRS overpasses during 20 and 21 July 2008. These data are then convolved with the same temperature weighting functions that apply for the AIRS observations (see e.g., Hoffmann et al., 2017). The same methodology was applied to model simulations




Figure 5. Comparison of AIRS and ICON-ART brightness temperature (BT) perturbations at wavelengths of 15 μm for all Antarctic Peninsula overpasses of AIRS during 20 July 2008. The first column shows the BT perturbation observed by AIRS, the second column the simulated perturbation based on ICON-ART in the Antarctic Peninsula nest. The third column shows the perturbation for AIRS (black) and ICON-ART (orange) at a latitude of 70°S. The rows show different overpasses at (a-c) 03:49 UTC, (d-f) 05:28 UTC, (g-i) 18:55 UTC and (j-l) 20:33 UTC.




Figure 6. Same as Fig. 5 but for 21 July 2008. The rows show different overpasses at (a-c) 04:32 UTC and (d-f) 19:38 UTC.

245 by Orr et al. (2015). The resulting brightness temperature (BT) perturbations can be found in Fig. 5 on 20 July and in Fig. 6 on 21 July 2008 for AIRS and ICON-ART.

Horizontal structures of the BT perturbations are shown in the first and second columns for AIRS and ICON-ART, respectively. Largest perturbations are present directly above the Antarctic Peninsula for both AIRS and ICON-ART which demonstrates that the perturbations originate from mountain waves propagating into the lower stratosphere. In addition, the mountain wave has an angle with respect to the Antarctic Peninsula mountains of about 45° , represented in both AIRS and
 250 ICON-ART. The horizontal wavelength of the simulated mountain wave is in the order of 300 km, which is a medium-large wavelength compared to other events (see e.g., Alexander and Teitelbaum, 2007; Plougonven et al., 2008; Hoffmann et al., 2014). Therefore, the ICON-ART simulation with lower stratospheric vertical resolution in the order of 500 m and horizontal resolution of 40 km in the Antarctic Peninsula nest can capture the main features of the mountain wave.

255 Fine structures as e.g. in panel j of Fig. 5 cannot be simulated in this simulation setup of ICON-ART (e.g. panel k) since the resolution of 40 km is still too coarse to predict them, which was already indicated by the comparison to CALIOP. This might be a resolution issue which would perhaps be improved by going to higher grid spacings, as shown by Orr et al. (2015). The chemistry non-linearly depends on temperature which means that small temperature variations could have a measurable effect on the chemistry (e.g., Murphy and Ravishankara, 1994). As pointed out previously, the microphysics of PSCs is one example
 260 for this. If the amplitude is underestimated, like in Fig. 5b and c, a higher resolved model would most probably generate more

ice PSCs, thus improving the CALIOP comparison at the lowest temperatures, cf. e.g. Figs. 5 to 7 of Orr et al. (2020) who used a higher resolution simulation to parametrise mountain-wave-induced PSCs.

These results are also stressed by the comparison of the perturbations at the latitude of 70°S that are shown in the third column of Figs. 5 and 6. The largest BT perturbations can be found at the longitude of the Antarctic Peninsula in all the panels. Some fine structures are missing in ICON-ART. For instance, at some overpasses the amplitude of the wave is underestimated compared to AIRS (e.g. panel c of Fig. 6). An analogous study by Orr et al. (2015) using a 4 km resolution seems to suggest a better match between model and observations. At other overpasses, the phase of the wave is shifted with respect to AIRS, as e.g. in panels c of Fig. 5 and f in Fig. 6. This is most probably a result of the free-running simulation where the wave cannot be expected to be located at exactly the same location as in the measurements.

In total, the comparison with AIRS showed that, apart from some missing fine structures, the mountain wave event taking place in the end of July 2008 can be represented with the resolution of 40 km in comparison with AIRS. In the following, we will analyse how this directly simulated mountain wave event impacts the dynamics in the coarser resolution domains in ICON-ART.

Figure 7 shows cross sections of the mountain wave dynamics along the line shown in Fig. 2 for the example of 22 July 2008 at 04 UTC in the altitude range between 12 and 30 km. The model data is interpolated to the path by an inverse distance method including the three neighboured grid points. The left column shows the simulation without nests whereas the three other columns illustrate the dynamics in the different domains shown in Fig. 2 of the simulation with the nests.

The bottom row of Fig. 7 shows the resolution of the Antarctic Peninsula in the different domains. As can be seen, the higher the horizontal resolution the better the Antarctic Peninsula can be represented as mountain range. Thus, this indicates that the interaction between the flow and the detailed orography in the Antarctic Peninsula nest is improved with respect to the global resolution of 160 km.

This is reflected in the variables shown in Fig. 7. Temperatures close to 175 K only occur in the Antarctic Peninsula nest with a 40 km resolution in the lee of the mountain. The temperature in the Antarctic Peninsula nest also shows the characteristic high and low temperature patterns as calculated by theory (e.g., Queney, 1947; Smith, 1989) and seen by measurements (e.g., Wright et al., 2017). As already indicated by Figs. 5 and 6, the temperature perturbation in the order of 10 K compares well to the AIRS measurements and is also consistent with previous studies of mountain waves (Meilinger et al., 1995; Carslaw et al., 1998a; Eckermann et al., 2009).

The relatively large temperature gradients in the Antarctic Peninsula nest are also in agreement with gradients in the wind velocities which are shown in the second and third row of Fig. 7. In the lee of the Antarctic Peninsula, the vertical wind velocity changes signs in altitudes where the temperature increases or decreases and has maximum values above 0.3 m s^{-1} . Largest horizontal wind velocities in the order of 75 m s^{-1} occur in altitudes above 27 km.

The mountains also cause perturbations in the potential temperature (fourth row in Fig. 7) in the lee. If diabatic processes are negligible, the flow follows the contours of potential temperature which will be shown in the PSC precursors in Sect. 5.3.2.

Up to now, it was only demonstrated that this mountain wave event can be directly simulated in the Antarctic Peninsula nest (right column), compares well to measurements and is consistent with the theory of mountain waves (Queney, 1947; Smith,




Figure 7. Cross sections on 22 July 2008 at 04 UTC along the white line in Fig. 2. Each row represents a different variable in the model. The bottom row shows the surface altitude with the Antarctic Peninsula at around 65°W. The columns represent the different simulations and domains: The first column is the simulation without nests ($\Delta x \approx 160$ km), the second one is the global domain ($\Delta x \approx 160$ km) including two-way nesting and the two right columns represent the Antarctica ($\Delta x \approx 80$ km) and Antarctic Peninsula nests ($\Delta x \approx 40$ km), respectively. The dynamical variables temperature, vertical wind velocity, horizontal wind velocity and potential temperature are shown in this figure.

1989). In addition, Figure 7 also shows the impact of the two-way nesting in ICON-ART. The nest around the Antarctica with resolution of about 80 km (third column) interacts with both nests and shows the transition between these resolutions. The mountain wave at the Antarctic Peninsula cannot be represented adequately in the resolution of 160 km in the simulation without the nests (left column in Fig. 7). The characteristic wave patterns do not occur in this simulation. In contrast to this, the global domain in the simulation with the nests (second column in Fig. 7) shows a decrease in temperature of about 2 K in the lee of the mountains. This is a result of the two-way nesting and the lower temperature due to the directly simulated mountain wave in the Antarctic Peninsula nest. The amplitude is lower than it is expected by mountain waves but the effect of the mountain wave is still remarkable also in the global grid of 160 km resolution.

This is also visible in the other variables of Fig. 7. Especially for the vertical wind velocity (second row), one can see that wave-like structures occur in the global domain where they cannot be represented without the nests. Therefore, we can expect that mountain-wave-induced PSCs can also be represented in this relatively low global resolution because they are directly simulated in the locally refined regions. In the following section, we will compare PSCs to CALIOP measurements in the Antarctic Peninsula nest before we analyse the impact of further model variables on the chemistry.

5.2 Comparison of simulated PSCs with CALIOP measurements

For a comparison with CALIOP, the ICON-ART PSC volume concentrations are interpolated (1) horizontally by an inverse distance method, (2) linearly in time and (3) linearly in geometric altitude to all CALIOP paths from 22 to 29 July 2008 where CALIOP's orbit was within the region of Antarctic Peninsula nest. Details about the interpolation can be found in Weimer (2019, Sect. 4.7.4).

As pointed out by previous studies, an adequate comparison of CALIOP with model data can only be set up if the model data is transferred into the optical space measured by CALIOP at 532 nm. We apply the method by Engel et al. (2013), Tritscher et al. (2019) and Steiner et al. (2021). It is based on T-matrix and Mie calculations (e.g., Mishchenko et al., 1996) with particle number densities and particle radii of the PSC types as input. The external PSC mixtures of ICON-ART are combined to optical properties (R_{532} and β_{\perp}) for each grid point. In accordance with Tritscher et al. (2019) and Steiner et al. (2021) we use aspect ratios of ice and NAT of 0.9 and refractive indices of 1.44 for STS (Krieger et al., 2000), 1.31 for ice and 1.48 for NAT (Middlebrook et al., 1994).

The three thresholds to determine the boundaries between the PSC categories, as mentioned in Sect. 4.1, are taken from the measurement data and averaged daily for each CALIOP height level ($\bar{\beta}_{\perp, \text{thres}}$, $\bar{R}_{532, \text{thres}}$ and $\bar{R}_{\text{NAT}|\text{ice}}$) (Tritscher et al., 2019; Steiner et al., 2021). As pointed out by Engel et al. (2013), it is important to account for the measurement uncertainties σ_R and $\sigma_{\beta, \perp}$ to compare the simulated R_{532} and β_{\perp} with CALIOP. These uncertainties are calculated according to Eqs. (5) and (6) of Tritscher et al. (2019), respectively. The simulated R_{532} and β_{\perp} are scaled by a normal distribution with mean at the simulated value and standard deviation as the respective σ . To summarise, the condition to determine if a PSC is detected is then

$$\beta_{\text{scal}} > \bar{\beta}_{\perp, \text{thres}} + \sigma_{\beta, \perp}, \quad \beta_{\text{scal}} \sim \mathcal{N}(\beta_{\perp}, \sigma_{\beta, \perp}^2) \quad (1)$$




Figure 8. Statistical analysis of PSC occurrence for a) the ICON-ART simulation without nests, b) the ICON-ART simulation with nests in the Antarctic Peninsula nest and c) CALIOP measurements. The data is restricted to the region of the Antarctic Peninsula nest for all panels to get comparable results. The ICON-ART data is interpolated to the CALIOP paths in the altitude range from 15 to 30 km in the nest in the time range between 22 and 29 July 2008 where CALIOP data is available. The colours correspond to the numbers in the circles and show the fraction of the PSC category that is present in the temperature bins with a width of 3.5 K. The sizes of the circles correspond to the total number of grid points with PSCs in the temperature bins relative to the maximum in each panel, also denoted at the right hand side of the panels as N_{PSC} , following the nomenclature by Spang et al. (2016). The “k” behind the numbers abbreviate thousand, i.e. 179.7k = 179700 grid points with PSCs. The temperature data for CALIOP originates from MERRA2, which is part of the CALIOP product.

or

$$R_{532,scal} > \overline{R}_{532,thres} + \sigma_R, \quad R_{532,scal} \sim \mathcal{N}(R_{532}, \sigma_R^2) \quad (2)$$

Since the ICON-ART simulations are free-running we cannot expect the PSCs to occur at exactly the same locations as in the measurements. Therefore, we compare the evolution of PSCs with respect to the temperature in both CALIOP and ICON-ART, which is a crucial parameter in the formation of PSCs (e.g., Solomon, 1999). We count the occurrence of the different PSC categories by the method described above in temperature bins and compare them between ICON-ART and CALIOP. The "No" PSC category is neglected in this comparison to emphasise the occurrence of PSCs in the temperature bins. The data is restricted to the region of the Antarctic Peninsula nest resulting in a total number of about one million grid points with PSCs for CALIOP and about 0.5 million grid points with PSCs in ICON-ART. This large difference could either be a result of horizontal shifted PSCs especially in the later part of the free-running simulation or a problem with the constant number concentrations used in the model. It should be further analysed in the future.

Figure 8 shows the relative number of grid points (in %) on the CALIOP paths where the different PSC categories occur. The size of the circles correspond to the total number of grid points with PSCs and the colour of the circles show the relative number of this PSC category occurrence in the respective temperature bin. The panels show the PSC occurrence in a) the simulation without the nests, b) in the Antarctic Peninsula nest and c) in the CALIOP measurements. For ICON-ART (panels

a and b), the modelled temperatures are used directly. The temperature for CALIOP (panel c) is interpolated from MERRA2 and provided as part of the original dataset (Pitts et al., 2018).

345 As can be seen, the relative distribution of grid points with PSCs coincides in all panels down to a temperature of about 180 K. Temperatures lower than that, however, are underrepresented in the simulation without the nest, compared to CALIOP. This is expected from Fig. 7 since temperatures are higher than 180 K in the simulation without nests in the lee of the mountains. In addition, the simulation without nests overestimates the fraction of the ice category at temperatures lower than 183 K compared to CALIOP.

350 Both deficiencies are improved in the Antarctic Peninsula nest with resolution of 40 km (panel b). PSCs in the ice category occur with a similar fraction like in CALIOP in the 181 K bin and the relative number of grid points with PSCs in the lowest temperature bin coincides with the measurements. The fractions of the NAT mixtures and NAT enhanced categories are also comparable to CALIOP over the whole range of temperatures. For medium temperatures (188 and 184.5 K) the fraction of the STS category is slightly overestimated compared to CALIOP. A reason for this could be missing fine structures in the gravity
355 wave, as already indicated by Figs. 5 and 6. The fraction of the ice category is 9 % larger in the lowest temperature bin than in CALIOP.

Although Fig. 7 suggested that main parts of the mountain wave can be captured in the Antarctic Peninsula nest, no PSCs in the “Wave-ice” category are simulated by ICON-ART. As mentioned in Sect. 2, the ice particle number concentration is essentially set to the constant value of 0.25 cm^{-3} in the stratosphere. The ice number concentration in mountain waves can
360 increase to values of a few cm^{-3} and then lead to larger backscatter ratios (e.g., Engel et al., 2013). Therefore, this too low ice number concentration could explain why the backscatter ratio to determine the PSC categories does not get as large as 50 which is needed for the “Wave-ice” category (Pitts et al., 2018). On the other hand, as mentioned in Sect. 4.1, the Wave-ice category is based on an empirical threshold of $R_{532} > 50$, so that parts of the Wave-ice clouds could also be attributed to the ice category (Pitts et al., 2018).

365 Comparing the three panels of Fig. 8, there seems to be a resolution-dependent shift of overestimated fraction in the ice category. At the resolution of 160 km, the overestimation starts at the 181 K bin, for 40 km it begins at 177.5 K. Thus, this figure gives first hints that using an even higher resolution could improve the comparison of the PSC distributions at low temperatures with CALIOP since the amplitude of the mountain wave may be better represented, as already mentioned in the comparison with AIRS in Sect. 5.1. Similar findings by Orr et al. (2020) corroborate this. It is also consistent to other previous
370 studies with mesoscale models that used a higher resolution to study mountain waves (e.g., Noel and Pitts, 2012) and should be one focus of future simulations.

This shift in the fraction of the ice category is emphasised in Figs. 9 and 10. They show the evolution of the PSC categories relative to the NAT and ice formation temperatures, T_{NAT} and T_{ice} . We calculated these temperatures with $X_{\text{H}_2\text{O}} = 2.5 \text{ ppmv}$ and $X_{\text{HNO}_3} = 2 \text{ ppbv}$ based on the formulas by Hanson and Mauersberger (1988) and Marti and Mauersberger (1993), respec-
375 tively, to get a comparable pressure-dependent reference temperature for both the model and the measurements. These constant volume mixing ratios are used only to calculate the PSC existence temperatures. They are based on satellite measurements shown by Tritscher et al. (2019) for late July, accounting for denitrification and dehydration.




Figure 9. Same as Fig. 8, but relative to T_{NAT} . Here, a common T_{NAT} for both ICON-ART and CALIOP y-axes is derived from Hanson and Mauersberger (1988) with input of $X_{\text{H}_2\text{O}} = 2.5$ ppmv and $X_{\text{HNO}_3} = 2$ ppbv based on Tritscher et al. (2019). The data are binned in terms of temperature difference with a bin width of 1 K.

As can be seen in Fig. 9, the number of grid points with PSCs grows when the temperature gets lower than T_{NAT} in all three panels. Similar patterns as in Fig. 8 can be seen: Overestimation of the STS category at temperatures around 1 to 3 K lower than T_{NAT} in the Antarctic Peninsula nest and differences in the ice category. In case of the simulation without nests, the fractions of ice are larger than in CALIOP for $\Delta T_{\text{NAT}} < 0$ K. In the Antarctic Peninsula nest, this overestimation can be seen at temperatures $\Delta T_{\text{NAT}} < -2$ K. These deficiencies should be analysed in future simulations.

The comparison with respect to T_{ice} in Fig. 10 emphasises the previous findings. In the simulation without the nests, temperatures lower than $\Delta T_{\text{ice}} = -3$ K are underrepresented compared to CALIOP. The fraction of the ice category peaks with 69 % at $\Delta T_{\text{ice}} = -1$ K. The resolution of 40 km in the Antarctic Peninsula nest is able to reproduce the general evolution of PSCs for $\Delta T_{\text{ice}} < -3$ K, but the fraction peaks with 74 % at $\Delta T_{\text{ice}} = -2$ K. In contrast, the CALIOP measurements suggest that the fraction of the ice category should be larger for $\Delta T_{\text{ice}} < -2$ K. An even higher resolution might help to reflect ice PSCs at these temperatures, as discussed in the comparison with AIRS (compare also with Orr et al., 2020).




Figure 10. Same as Fig. 8, but relative to T_{ice} . Here, a common T_{ice} for ICON-ART and CALIOP y-axes is derived from Marti and Mauersberger (1993) with input of $X_{H_2O} = 2.5$ ppmv based on Tritscher et al. (2019). The data are binned in terms of temperature difference with a bin width of 1 K.

In total, this section demonstrated that the PSC scheme in ICON-ART is able to generate PSCs similar to CALIOP. The peak fraction of the ice category seems to move to lower temperatures if the resolution is increased, which might indicate that an even higher resolution is needed to capture the ice formation at the lowest temperatures (cf., Orr et al., 2020). On the other hand, this analysis demonstrated that the evolution of the PSCs is clearly improved in the Antarctic Peninsula nest with respect to CALIOP. Some differences in the STS category with respect to CALIOP also indicate that some fine structures of the mountain wave are missing, consistent to the previous findings in the comparison with AIRS.

5.3 Impact of mountain-wave-induced PSCs on the chemistry

In the previous sections, it was demonstrated that both the PSC formation and the formation of the mountain wave are in relatively good agreement with measurements considering the limits in measurements and simulation setup. In this section, we investigate the impact of directly simulated mountain-wave-induced PSCs on the interactively calculated chemistry in ICON-ART. The analysis is focused on the mountain wave event at its peak dynamics on 22 July 2008 (see Figs. 4 and S1).




Figure 11. Same as Fig. 7 but for the volume concentrations of the different PSCs: liquid aerosol, NAT and ice. Please note the different colour bars.

400 5.3.1 The formation of PSCs in the mountain wave

Figure 11 demonstrates that the ICON-ART model has the potential to close the gap between direct simulations of mountain-wave-induced PSCs and their treatment in relatively coarse global resolutions. The volume concentration of liquid particles (first row) in the global domain with nests is influenced by the mountain wave, especially at altitudes higher than 20 km where particle volume concentrations close to zero occur in the global domain which do not exist without the nesting technique. The influence of the mountain wave on liquid aerosol is amplified within the nests. The liquid particles are assumed to freeze at temperatures 3 K below the frost point (Carslaw et al., 1995; Koop et al., 2000). Thus, liquid particles are only computed for higher temperatures so that they are formed in the mountain wave where the temperature is higher than this threshold.

STS and NAT compete with each other in taking up HNO_3 . This is why in this case study distinct layers exist: NAT PSCs at altitudes higher than 20 km and STS PSCs at lower altitudes where also H_2SO_4 is enhanced, see Sect. 5.3.2. The NAT volume concentrations are increased in the Antarctic Peninsula nest (second row, right column) which is why they are also increased in the global domain comparing the simulations with and without nests.

In contrast to the literature (e.g., Carslaw et al., 1999; Svendsen et al., 2005), the NAT volume concentration decreases when the air masses approach the mountain wave. Since the NAT size bins are advected with the general air masses, the wave-like

patterns occur in both nests. As a result of the operator splitting used in ICON-ART, the largest fraction of gaseous H_2O leads
415 to ice formation at temperatures lower than about 180 K and is not available for NAT PSCs and liquid particles in the mountain
wave anymore. Therefore, the largest signal of mountain-wave-induced PSCs can be found in ice PSCs in Fig. 11. This is an
issue of further investigation in the future. In addition, NAT PSCs are formed by freezing STS particles in the mountain wave,
as shown e.g. by Bertram et al. (2000) and Salcedo et al. (2001). This is not integrated in the model so far and should be
considered in the future.

420 The best example of the formation of mountain-wave-induced PSCs are ice PSCs (third row in Fig. 11). In the lee of the
Antarctic Peninsula ice PSCs occur with volume concentrations as large as $700 \mu\text{m}^3 \text{cm}^{-3}$. These relatively high values might
be overestimated because of the assumptions in the hydrometeor microphysics of the meteorological model (Doms et al.,
2011). As discussed above, the ice number concentration is too low for mountain-wave conditions compared to measurements.
A more realistic number concentration would lead to smaller ice particles and could reduce the ice volume concentration to
425 more realistic values.

On the other hand, the ice PSCs are clearly connected to the regions where temperature is decreased and show similar wave-
like patterns as the temperature. These increased volume concentrations are also present in the global domain where wave-like
patterns can be simulated with the nests in contrast to the simulation without the nests where these structures do not exist.

Therefore, this figure shows that mountain-wave-induced PSCs can be directly simulated with ICON-ART for this specific
430 event. Their effect can also be treated in the global domain where mountain-wave-induced PSCs cannot be represented without
the nests. The PSCs also interact with the gaseous species which are analysed in the next section.

5.3.2 Influence of the mountain wave on PSC precursors

As mentioned above, long-lived tracers closely follow the potential temperature. Therefore, the wave perturbations in potential
temperature can be seen in all tracers shown in Fig. 12 in the lee of the mountain for the simulation with the nests. The wave-like
435 structures in sulfuric acid (H_2SO_4 , first row), nitric acid (HNO_3 , second row) and water vapour (H_2O , third row) correspond
to the structures seen in the potential temperature (see Fig. 7).

H_2SO_4 is prescribed by a climatology in the global domain, but free-running in the nests, and only shows a minor signal of
the two-way nesting. Due to the missing sink of H_2SO_4 by sedimentation of aerosols (see Sect. 3) and advection processes,
the mixing ratio accumulates in the nested domains (two right columns).

440 HNO_3 , shown in the second row of Fig. 12, is taken up by STS and NAT PSCs so that it is lower in the simulation with nests
than in the simulation without the nests. This is shown not only in both nested domains, but can also be returned to the global
domain as a result of the two-way nesting.

This property especially occurs for H_2O . In the lee at altitudes between 22 and 25 km, temperatures lower than about 179 K
lead to volume mixing ratios lower than 1 ppmv that are connected with the uptake in ice PSCs (cf. Fig. 11). The resolution
445 of 80 km in the Antarctica nest (third column) misses a larger part of the wave but the wave-like pattern can still be seen for
 H_2O . It cannot be represented in the simulation without the nests (left column) where the H_2O volume mixing ratio does




Figure 12. Same as Fig. 7 but for the tracers that are relevant for the formation of PSCs: H_2SO_4 , HNO_3 and H_2O . Please note the different colour bars.

not decrease to values lower than 2.5 ppmv. In the global domain of the simulation with the nests, however, values as low as 2 ppmv occur in the lee of the mountain as a result of the two-way nesting.

5.3.3 Impact of mountain-wave-induced polar stratospheric clouds on chlorine activation

Both the directly simulated mountain-wave-induced PSCs and the lower wave-driven temperature in the simulation with the nests also affect other species. The chlorine species are summarised in Fig. 13 in the same way as in the previous figures. The reservoir species hydrochloric acid (HCl , first row) and chlorine nitrate (ClONO_2 , second row) are shown together with the active chlorine species (third row), summarised as ClO_x :

$$X_{\text{ClO}_x} = X_{\text{ClONO}_2} + 2X_{\text{Cl}_2\text{O}_2} + X_{\text{OCIO}} + 2X_{\text{Cl}_2} + X_{\text{BrCl}} + X_{\text{HOCl}} + X_{\text{Cl}} + X_{\text{ClO}} \quad (3)$$

During July, which is a relatively late stage of the southern polar winter, most of the chlorine species in the altitude range between 15 and 25 km have been already activated (cf., Tully et al., 2011). The broad band of volume mixing ratios of ClO_x with values up to about 2.2 ppbv that are present in all panels of the third row suggests this. Therefore, additional chlorine activation can only be expected in altitude regions above 25 km. In the previous sections, it was shown that the directly simulated mountain-wave-induced ice PSCs reaches altitudes of about 26 km (see Fig. 11).




Figure 13. Same as Fig. 7 but for the chlorine species: HCl, ClONO₂ and the sum of all other (active) chlorine-containing species (ClO_x). The colour bars are equal in this figure.

460 Low temperature and mountain-wave-induced PSCs lead to increased chlorine activation in the lee of the Antarctic Peninsula that is not present in the simulation without the nests. At altitudes around 26 km, the additional ice PSCs activate both ClONO₂ and HCl in the lee of the mountains. This is shown by values of the reservoir species around zero and values of ClO_x up to 3 ppbv in this region that cannot be simulated without the nests. These increased ClO_x mixing ratios also have an effect on ozone in the model which will be shown in the next section.

465 5.3.4 Impact of mountain-wave-induced polar stratospheric clouds on ozone

A summary of the impact of the two-way nesting on mountain-wave-induced PSCs and the chemistry can be found in Fig. 14. It shows the differences of the two global domains between with and without the nests for various variables of the previous sections. Panel a shows the difference in the NAT volume concentration and it can be seen that more NAT particles are produced in the simulation with the nests as a result of the two-way nesting. These enhanced NAT volume concentrations lead to (1) a
 470 decreased volume concentration of liquid particles (panel b) as a result of the used operator splitting and (2) to decreased values of HNO₃ (panel d) with differences lower than -1.4 ppbv in this region since HNO₃ is taken up by NAT particles. In regions where no additional NAT particles exist, the liquid volume concentration is enhanced due to the two-way nesting (panel b). In the ice volume concentration in panel c, the wave-like structure is clearly present.




Figure 14. Difference in global domain between with and without nest around the Antarctic Peninsula along the cross section shown in Fig. 2 at the same date (22 July 2008, 04 UTC) and using the same algorithm as in the previous figures. The shown variables are differences in (a-c) the NAT, liquid and ice volume concentrations, respectively, (d) the HNO_3 volume mixing ratio, (e) the volume mixing ratio of the reservoir species ClONO_2 and HCl and (f) the ClO_x as defined in Eq. (3). Minimum values of HNO_3 are around 2.3 ppbv.

The mountain-wave-induced (ice) PSCs and low temperatures lead to additional chlorine activation, in this example in an altitude of about 26 km. The negative difference in the sum of ClONO_2 and HCl (panel e) is closely connected to the positive differences in ClO_x in the lee of the mountain (panel f). HNO_3 is also a product of heterogeneous reactions on the surface of PSCs and therefore is enhanced in this region with respect to the simulation without the nests (panel d).

Due to transport of the activated chlorine species and PSCs, ozone depletion might take place downstream of the Antarctic Peninsula (e.g., Höpfner et al., 2006b). Therefore, ozone is affected by the mountain wave event (1) far in the lee of the mountain and (2) later in the year. This is why the time series of the southern hemispheric mean total ozone column for the whole third period of the simulation is shown in Fig. 15. The ozone columns are averaged for latitudes south of 60°S . Two lines are shown in panel a: the solid line is the mean total ozone column in the global domain of the simulation with the nests, whereas the dashed line illustrates the simulation without the nests.

As can be seen, differences in the simulations are below 1 DU during the whole day of 19 July (see also panel b). This first day is a spin-up period for the nests that are initialised with the global domain to form the mountain wave. From 20 July until end of the simulation, however, the mean ozone column in the simulation with the nests is smaller than in the simulation without the nests. While ozone is generally decreasing during this period in both simulations, the absolute value of the difference peaks at around 8 DU on 25 July and decreases afterwards. This order of magnitude has also been simulated by the parametrisation




Figure 15. Time series of the mean total ozone column during the third period of the simulation in the global domains for latitudes south of 60°S (panel a). The dashed line is the simulation without the nests and the solid line shows the ozone development in the simulation with the nests. The difference between these lines is illustrated in panel b.

of mountain-wave-induced PSCs of Orr et al. (2020). The higher resolution around the Antarctica and the Antarctic Peninsula
 490 seems to lead to generally lower ozone columns. This additionally highlights the need of higher resolutions in atmospheric chemistry models.

Figure 15 demonstrates that the higher load of active chlorine species leads to a by up to 8 DU larger decrease of ozone in the southern hemisphere. The prediction skill after 10 days is fairly low so that an analysis after the shown period is not possible with a free-running simulation. Future investigations could extend this period with specified dynamics in both simulations to
 495 be able to examine the impact of mountain-wave-induced PSCs on the ozone hole during September and October.

6 Conclusions and outlook

Seamless modelling of chemistry-climate interactions is challenging. Not many modelling systems can do this in a consistent way. In the past, it was impossible to directly simulate mountain-wave-induced PSCs in global chemistry models due to the coarse resolution in that kind of simulations. In this study, we investigated this problem with the scheme for PSCs in the
 500 ICOSahedral Non-hydrostatic modelling framework with its extension for Aerosols and Reactive Trace gases (ICON-ART). The scheme forms ice PSCs based on the microphysics of the meteorological model, liquid (sulfate and STS) particles by the analytic expression of Carslaw et al. (1995) with some improvements with respect to the constant particle number concentration and NAT particles by a kinetic non-equilibrium approach with a flexibly selectable size distribution.

We performed a three-step simulation to investigate the impact of mountain-wave-induced PSCs in ICON-ART on the chem-
 505 istry: First, a free-running simulation was conducted from 01 March to 30 April 2008. Second, the dynamics were reinitialised

every second day by ERA-Interim data to ensure a realistic development of the polar vortex until 18 July 2008. Third, two free-running simulations followed that covered the investigated mountain wave event from 19 July to 29 July 2008: a simulation including two-way nesting with nests around the Antarctica ($\Delta x \approx 80$ km) and the Antarctic Peninsula ($\Delta x \approx 40$ km) and a simulation without these nests.

510 The results were compared with measurements by CALIOP and AIRS. The CALIOP PSC categories of ICON-ART were derived from an algorithm that transfers the PSC volume concentrations to the spectral space of CALIOP so that a statistical comparison between both datasets could be established. The total number of grid points with PSCs in ICON-ART was half of that measured by CALIOP although the temperature distribution was similar. This should be further investigated in future simulations. The CALIOP “Wave-ice” category could not be simulated with the model most probably as a result of the constant
515 ice number concentration set to 0.25 cm^{-3} . The analyses also showed the need of an interactive calculation of PSCs to treat the different PSC types competing with the available gaseous HNO_3 and H_2O . The comparison with all CALIOP measurements within the Antarctic Peninsula nest demonstrated that the general formation of most of the PSCs in ICON-ART is similar with respect to temperature. We found a resolution-dependent overestimation of the fraction of ice clouds in comparison to CALIOP at high temperatures. This suggested that an even higher resolution is needed to capture PSCs in the lowest temperatures.
520 However, the resolution of 40 km clearly improved the PSC formation compared to 160 km and CALIOP. These findings were also pronounced by evaluations with respect to the NAT and ice formation temperatures.

The comparison to AIRS demonstrated for all Antarctic Peninsula overpasses of 20 and 21 July 2008 that the main features of the mountain wave could be represented in the resolution of 40 km. The measured angle of the mountain wave to the mountain range could be reflected by the model and the brightness temperature perturbation was in the correct order of magnitude for
525 all overpasses. The investigated mountain wave had a horizontal wavelength of about 300 km that could be captured by the chosen resolution. For mountain waves with smaller wavelengths, an even higher resolution would be needed to resolve the wave adequately.

By introducing the two-way nesting around the Antarctic Peninsula with resolutions down to 40 km, we were able to directly simulate the main features of the mountain wave and transfer its effect back to the global domain. Thus, additional mountain-
530 wave-induced PSCs together with lower temperatures led to enhanced chlorine activation in the global resolution of 160 km. An up to 8 DU larger ozone depletion above the Antarctic Continent was simulated.

Thus, this study shows the need of treating mountain-wave-induced PSCs in CCMs. The study also demonstrated that (1) dynamics, tracers, PSCs and chemistry are interactively and consistently integrated in ICON-ART. In addition, ICON-ART showed the potential to bridge the gap between direct simulations of mountain-wave-induced PSCs and their treatment in
535 coarse global resolutions.

Future simulations should exploit the nesting technique further and use it for other known mountain wave hot spots (e.g., Hoffmann et al., 2013, 2017). Waves of non-orographic origin, such as southern hemispheric storm tracks, can also induce mesoscale temperature perturbation that have not been considered in our simulations (e.g., Tritscher et al., 2021). In addition, the impact on ozone depletion should be evaluated with measurements and simulations of longer periods. The northern hemi-
540 sphere should be also a focus in the future where PSC formation and ozone depletion has been shown to be highly sensitive to

the existence of mountain waves (e.g., Tabazadeh et al., 2000; Eckermann et al., 2006; Dörnbrack et al., 2012; Khosrawi et al., 2018).

Code and data availability. Licences of the ICON code are currently managed by the Max-Planck-Institute for Meteorology (MPI-M) and the German Weather Service (DWD). Please visit https://code.mpimet.mpg.de/projects/iconpublic/wiki/How_to_obtain_the_model_code (last access on 19 March 2021) for further information. For ART, please contact Bernhard Vogel (bernhard.vogel@kit.edu). The version 2 data of CALIOP PSCs first published in Pitts et al. (2018) can be directly obtained by contacting Michael Pitts (michael.c.pitts@nasa.gov) and will be made available soon at CALIPSO Science Team (2015). The AIRS data is distributed by the NASA Goddard Earth Sciences Data Information and Services Center (AIRS project, 2007). The AIRS gravity wave datasets used in this study can be accessed at https://datapub.fz-juelich.de/slcs/airs/gravity_waves (last access on 19 March 2021). The code to transfer model PSC data to the optical space of CALIOP has been recently published as supplement by Steiner et al. (2021).

Appendix A: The PSC scheme in ICON-ART

In ICON-ART, the three types of PSCs (ice, NAT and STS) are treated separately. Sensitivity simulations showed that the ICON microphysics for ice clouds, operationally computed up to an altitude of 22.5 km, can be extended to the lower stratosphere up to 30 km (Weimer, 2019). The hydrometeor microphysics in ICON include heterogeneous nucleation of cloud ice, nucleation of cloud ice due to homogeneous freezing of cloud water and depositional growth and sublimation of cloud ice interacting with the other hydrometeors as well as sedimentation of the ice particles (Doms et al., 2011). Thus, both nucleation of ice PSCs and dehydration of the lower stratosphere can be captured by the ICON microphysics. A similar approach is used in the Whole Atmosphere Community Climate Model (WACCM, Wegner et al., 2013). Ice particle radius (r_{ice} in m) and particle number concentration (N_{ice} in m^{-3}) correspond to the assumed size distributions in the microphysics (Doms et al., 2011):

$$N_{\text{ice}}(T) = \min \left(5 \exp [0.304 (273.15 - T)], 250 \times 10^3 \right) \quad (\text{A1})$$

$$r_{\text{ice}} = \frac{1}{2} \sqrt[3]{\frac{\rho q_{\text{ice}}}{130 N_{\text{ice}}}} \quad (\text{A2})$$

where ρ is the air density in kg m^{-3} , T is the temperature in K and q_{ice} is the mass mixing ratio in kg kg^{-1} of water in ice. Equation A1 means that the ice particle number concentration is assumed to be 0.25 cm^{-3} for temperatures lower than 239 K.

Two parametrisations for the microphysics of NAT particles are integrated in ICON-ART. The thermodynamic NAT parametrisation is diagnostic and therefore computes the number of moles of NAT particles in thermodynamic equilibrium, calculates its sedimentation and evaporates the particles again within the same model time step. The volume mixing ratio of HNO_3 condensed in NAT ($X_{\text{HNO}_3(\text{NAT})}$ in mol mol^{-1}) is calculated on the basis of the difference between vapour pressure of HNO_3 (p_{HNO_3} in Pa) and the saturation vapour pressure over NAT ($p_{\text{sat,NAT}}$ in Pa):

$$X_{\text{HNO}_3(\text{NAT})} = \frac{p_{\text{HNO}_3} - p_{\text{sat,NAT}}}{p} \quad (\text{A3})$$

570 The saturation vapour pressure over NAT is calculated according to Hanson and Mauersberger (1988) and p is the air pressure (in Pa). Particle number concentration (N_{NAT}) and radius (r_{NAT}) in the thermodynamic NAT parametrisation are calculated using a threshold in the number concentration of $N_{\text{NAT,max}} = 2.3 \times 10^{-4} \text{ cm}^{-3}$ which is based on van den Broek et al. (2004). Below this threshold, the radius of the NAT particles is set to $0.1 \mu\text{m}$. Above this threshold, the particle number concentration is set to $N_{\text{NAT,max}}$ and the radius of the particles is increased accordingly. This method has already been used in a similar way
575 for solid particles by Kirner et al. (2011).

The kinetic NAT parametrisation is a non-equilibrium approach based on prognostic equations for the particle mass (see e.g., Seinfeld and Pandis, 2006, p. 539). Carslaw et al. (2002) applied this approach to NAT particles in a Lagrangian model and van den Broek et al. (2004) extended it to Eulerian models. In the Lagrangian description by Carslaw et al. (2002), the change in NAT particle radius ($r_{\text{NAT},b}$) is calculated prognostically by a diffusive growth

$$580 \quad \frac{dr_{\text{NAT},b}}{dt} = \frac{G_b}{r_{\text{NAT},b}} \quad (\text{A4})$$

where G_b (in $\text{m}^2 \text{s}^{-1}$) is a growth factor which depends on the diffusion coefficient of HNO_3 in air (D_{HNO_3} , in $\text{m}^2 \text{s}^{-1}$), the air temperature (T , in K) and the saturation difference of the HNO_3 vapour pressure (in Pa):

$$G_b = \frac{D_{\text{HNO}_3,b}^* M_{\text{NAT}}}{\rho_{\text{NAT}} R^* T} (p_{\text{HNO}_3} - p_{\text{sat,NAT}}) \quad (\text{A5})$$

with

$$585 \quad D_{\text{HNO}_3,b}^* = \frac{D_{\text{HNO}_3}}{1 + 4 D_{\text{HNO}_3} / (\bar{v}_{\text{HNO}_3} r_{\text{NAT},b})} \quad (\text{A6})$$

In these equations, \bar{v}_{HNO_3} is the mean thermal velocity of air molecules (in m s^{-1}), R^* stands for the universal constant of an ideal gas in $\text{J mol}^{-1} \text{K}^{-1}$, ρ_{NAT} is the crystal density of NAT ($1.626 \times 10^6 \text{ g m}^{-3}$, Drdla et al., 1993; van den Broek et al., 2004) and M_{NAT} is the NAT molar mass of 117 g mol^{-1} . In one of the Eulerian formulations by van den Broek et al. (2004), this radius change is applied to particles in size bins, but directly converted into a change in the particle number concentration
590 (“FixedRad” approach). This approach is used in this study with a size distribution based on van den Broek et al. (2004) as shown in Fig. 3. That is why we added a b subscript to all variables in Eqs. (A4), (A5) and (A6) that depend on the size bin b .

The size distribution of NAT particles can be flexibly specified by the user to investigate its impact on denitrification without any change in the Fortran code but rather changing the respective XML control file (cf. Schröter et al., 2018). Each size bin is defined by radius limits and a maximum particle number concentration, which are kept constant during the simulation. If the
595 calculated particle number concentration in a bin exceeds the maximum, the excess mass is transferred to the next larger size bin. The sum of the maximum particle number concentrations of the bins has to equal the value by van den Broek et al. (2004), which is $2.3 \times 10^{-4} \text{ cm}^{-3}$ and based on measurements of large NAT particles. Sensitivity studies by van den Broek et al. (2004) showed that this value in combination with the used size distribution leads to denitrification comparable to measurements. The size bins are transported as passive tracers in ICON-ART.

600 Since the calculation of $p_{\text{sat,NAT}}$ by Hanson and Mauersberger (1988) has specific temperature limits, NAT particles are evaporated automatically at temperatures higher than 220 K. For temperatures below 180 K, we calculate p_{HNO_3} with a constant

temperature of 180 K. These two NAT parametrisations are also implemented in the ECHAM/MESSy Atmospheric Chemistry model (EMAC, Jöckel et al., 2010; Kirner et al., 2011).

Sedimentation of NAT particles, either formed by thermodynamic or kinetic NAT parametrisation, is calculated by a simple upwind method, using the Stokes velocity of assumed spherical particles (Stokes, 1851).

The microphysics of liquid (sulfate and STS) particles in the module are calculated by the scheme first published by Carslaw et al. (1995), with one exception: in the original code by Carslaw et al. (1995) the particle number concentration is set to the constant value of 10 cm^{-3} . We improved this fixed value by applying the mean of all balloon-borne STS measurements by Hervig and Deshler (1998) in order to derive the particle surface area concentration S_{STS} and the radius r_{STS} of STS particles from the internally calculated particle volume concentration V_{STS} :

$$S_{\text{STS}} = 6.068 (V_{\text{STS}})^{0.671} \quad (\text{A7})$$

$$r_{\text{STS}} = \frac{3 V_{\text{STS}}}{S_{\text{STS}}} \quad (\text{A8})$$

In these equations, V_{STS} has to be given in $\mu\text{m}^3 \text{ cm}^{-3}$ to get S_{STS} and r_{STS} in $\mu\text{m}^2 \text{ cm}^{-3}$ and μm , respectively. Sedimentation is neglected for liquid particles since they are too small to result in relevant redistribution of the major constituents H_2O , HNO_3 and H_2SO_4 (Tabazadeh et al., 2000; Considine et al., 2000).

Since the PSCs in the current version of the model do not interact with each other like in a fully coupled PSC scheme (e.g., Zhu et al., 2015), there are essentially two approaches to calculate PSCs: either they are computed with the total (gaseous + liquid + solid) concentrations of HNO_3 and H_2O as input for all PSC types (e.g., Kirner et al., 2011) or the PSCs are calculated subsequently with the gaseous fraction that remains after formation of the previously calculated PSC types (operator splitting). Both approaches have their advantages and disadvantages. We use the second approach because the maximum of HNO_3 and H_2O taken up by PSCs cannot exceed the gaseous concentrations in this case. First, ice PSCs are calculated, then NAT PSCs and finally liquid particles.

Particle radius, particle number concentration and particle surface area concentration are used to calculate the heterogeneous reaction rate constants on the surface of PSCs. For NAT and ice, which can grow to relatively large sizes in the order of tens of μm , the following equation is used to calculate the heterogeneous reaction rate constant, assuming spherical particles (Drdla et al., 1993):

$$k_{\text{het},h,c} = \frac{\gamma_{h,c} \pi r_c^2 \bar{v}_{i(h)} N_c}{N_{j(h)} \left(1 + \frac{3\gamma_{h,c}}{4\text{Kn}} \right)}, \quad c \in \{\text{NAT, ice}\} \quad (\text{A9})$$

where $\gamma_{h,c}$ is the uptake coefficient (i.e. reaction probability) of heterogeneous reaction h and PSC type c , $\bar{v}_{i(h)}$ is the mean thermal velocity of the gaseous reactant $i(h)$ (in m s^{-1}), N_c is the PSC number concentration and $N_{j(h)}$ is the number concentration of the reactant $j(h)$ adsorbed on the particle (both in m^{-3}). The Knudsen number Kn is calculated by λ_{mfp}/r_c with the mean free path λ_{mfp} of air molecules (in m), calculated according to Kennard (1938). In case of the kinetic NAT parametrisation, Eq. (A9) is evaluated for each size bin separately and summed up subsequently.

Liquid particles are considerably smaller than ice and NAT particles (e.g., Considine et al., 2000) and hence $Kn \gg 1$ so that the slip flow correction term in Eq. (A9) can be neglected, which yields:

$$635 \quad k_{\text{het},h,\text{liq}} = \frac{\gamma_{h,\text{liq}} \bar{v}_{i(h)} S_{\text{liq}}}{4N_{j(h)}} \quad (\text{A10})$$

The total reaction rate constant is the sum of the reaction rate constants on all three types of PSCs. The uptake coefficients in ICON-ART are used either from Carslaw et al. (1995) or Sander et al. (2011b). A summary of the γ values can be found in Table S3 of the paper’s supplement. They also compare well to other models (e.g., Solomon et al., 2015).

Author contributions. This paper is part of MW’s thesis supervised by PB, RR and OK. MW developed the module for PSCs in ICON-ART and performed the simulations with contributions by JB, OK, RR and PB. LH performed the comparison between AIRS and ICON-ART. MW, BL, IT and MS performed the comparison between ICON-ART and CALIOP. MW prepared the manuscript with contributions by all authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. Parts of the simulations were performed on bwUniCluster for which the authors acknowledge support by the state of Baden-Württemberg through bwHPC. Other parts of the work as well as evaluations were performed on the supercomputer ForHLR funded by the Ministry of Science, Research and the Arts Baden-Württemberg and by the Federal Ministry of Education and Research. This work was performed with the help of the Large Scale Data Facility at the Karlsruhe Institute of Technology funded by the Ministry of Science, Research and the Arts Baden-Württemberg and by the Federal Ministry of Education and Research. We acknowledge ECCAD for archiving and distributing the emission data. We acknowledge funding from the Initiative and Networking Fund of the Helmholtz Association through the projects “Digital Earth” and “Advanced Earth System Modelling Capacity”. I. Tritscher was funded by the Deutsche Forschungsgemeinschaft (DFG) under project number 310479827.

References

- AIRS project: AIRS/Aqua L1B Infrared (IR) geolocated and calibrated radiances V005, <https://doi.org/10.5067/YZEXEVN4JGGJ>, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), Accessed: 31 December 2015, 2007.
- 655 Alexander, M. J. and Teitelbaum, H.: Observation and analysis of a large amplitude mountain wave event over the Antarctic peninsula, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD008368>, 2007.
- Alexander, S. P., Klekociuk, A. R., Pitts, M. C., McDonald, A. J., and Arevalo-Torres, A.: The effect of orographic gravity waves on Antarctic polar stratospheric cloud occurrence and composition, *J. Geophys. Res.: Atmos.*, 116, <https://doi.org/10.1029/2010JD015184>, 2011.
- Alexander, S. P., Klekociuk, A. R., McDonald, A. J., and Pitts, M. C.: Quantifying the role of orographic gravity waves on polar stratospheric cloud occurrence in the Antarctic and the Arctic, *J. Geophys. Res.: Atmos.*, 118, 11,493–11,507, <https://doi.org/10.1002/2013JD020122>, 2013.
- 660 Aumann, H. H., Chahine, M. T., Gautier, C., Goldberg, M. D., Kalnay, E., McMillin, L. M., Revercomb, H., Rosenkranz, P. W., Smith, W. L., Staelin, D. H., Strow, L. L., and Susskind, J.: AIRS/AMSU/HSB on the Aqua mission: design, science objectives, data products, and processing systems, *IEEE Trans. Geosci. Remote Sens.*, 41, 253–264, <https://doi.org/10.1109/TGRS.2002.808356>, 2003.
- 665 Bacmeister, J.: Mountain-Wave Drag in the Stratosphere and Mesosphere Inferred from Observed Winds and a Simple Mountain-Wave Parameterization Scheme, *J. Atmos. Sci.*, 50, 377–399, [https://doi.org/10.1175/1520-0469\(1993\)050<0377:MWDITS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050<0377:MWDITS>2.0.CO;2), 1993.
- Bacmeister, J. T., Newman, P. A., Gary, B. L., and Chan, K. R.: An Algorithm for Forecasting Mountain Wave-Related Turbulence in the Stratosphere, *Weather Forecast.*, 9, 241–253, [https://doi.org/10.1175/1520-0434\(1994\)009<0241:AAFFMW>2.0.CO;2](https://doi.org/10.1175/1520-0434(1994)009<0241:AAFFMW>2.0.CO;2), 1994.
- Bertram, A. K., Dickens, D. B., and Sloan, J. J.: Supercooling of type 1 polar stratospheric clouds: The freezing of submicron nitric acid aerosols having HNO_3 mol fractions less than 0.5, *J. Geophys. Res.: Atmos.*, 105, 9283–9290, <https://doi.org/10.1029/1999JD901174>, 2000.
- 670 Braesicke, P., J. Neu (Lead Authors), Fioletov, V., Godin-Beekmann, S., Hubert, D., Petropavlovskikh, I., Shiotani, M., and Sinnhuber, B.-M.: Chapter 3: Update on Global Ozone: Past, Present, and Future, in: *Scientific Assessment of Ozone Depletion: 2018*, edited by Fahey, D. W., Newman, P. A., Pyle, J. A., and Safari, B., Global Ozone Research and Monitoring Project – Report No. 58, World Meteorological Organization, Geneva, Switzerland, 2018.
- 675 CALIPSO Science Team: CALIPSO Lidar Level 2 Polar Stratospheric Clouds (PSC) Data Product Ensemble, Provisional V1-10, https://doi.org/10.5067/CALIOP/CALIPSO/CAL_LID_L2_PSCMASK-PROV-V1-00_L2-001.00, 2015.
- Carlsaw, K. S., Luo, B. P., Clegg, S. L., Peter, T., Brimblecombe, P., and Crutzen, P. J.: Stratospheric aerosol growth and HNO_3 gas phase depletion from coupled HNO_3 and water uptake by liquid particles, *Geophys. Res. Lett.*, 21, 2479–2482, <https://doi.org/10.1029/94GL02799>, 1994.
- 680 Carlsaw, K. S., Luo, B., and Peter, T.: An analytic expression for the composition of aqueous HNO_3 - H_2SO_4 stratospheric aerosols including gas phase removal of HNO_3 , *Geophys. Res. Lett.*, 22, 1877–1880, <https://doi.org/10.1029/95GL01668>, 1995.
- Carlsaw, K. S., Wirth, M., Tsias, A., Luo, B. P., Dörnbrack, A., Leutbecher, M., Volkert, H., Renger, W., Bacmeister, J. T., and Peter, T.: Particle microphysics and chemistry in remotely observed mountain polar stratospheric clouds, *J. Geophys. Res.: Atmos.*, 103, 5785–5796, <https://doi.org/10.1029/97JD03626>, 1998a.
- 685 Carlsaw, K. S., Wirth, M., Tsias, A., Luo, B. P., Dörnbrack, A., Leutbecher, M., Volkert, H., Renger, W., Bacmeister, J. T., Reimer, E., and Peter, T.: Increased stratospheric ozone depletion due to mountain-induced atmospheric waves, *Nature*, 391, 675–678, <https://doi.org/10.1038/35589>, 1998b.

- Carslaw, K. S., Peter, T., Bacmeister, J. T., and Eckermann, S. D.: Widespread solid particle formation by mountain waves in the Arctic stratosphere, *Journal of Geophysical Research: Atmospheres*, 104, 1827–1836, <https://doi.org/10.1029/1998JD100033>, 1999.
- 690 Carslaw, K. S., Kettleborough, J. A., Northway, M. J., Davies, S., Gao, R.-S., Fahey, D. W., Baumgardner, D. G., Chipperfield, M. P., and Kleinböhl, A.: A vortex-scale simulation of the growth and sedimentation of large nitric acid hydrate particles, *J. Geophys. Res.: Atmos.*, 107, SOL 43–1–SOL 43–16, <https://doi.org/10.1029/2001JD000467>, 8300, 2002.
- Chahine, M. T., Pagano, T. S., Aumann, H. H., Atlas, R., Barnet, C., Blaisdell, J., Chen, L., Divakarla, M., Fetzer, E. J., Goldberg, M.,
 695 Gautier, C., Granger, S., Hannon, S., Irion, F. W., Kakar, R., Kalnay, E., Lambrigtsen, B. H., Lee, S.-Y., Le Marshall, J., Mcmillan, W. W., McMillin, L., Olsen, E. T., Revercomb, H., Rosenkranz, P., Smith, W. L., Staelin, D., Strow, L. L., Susskind, J., Tobin, D., Wolf, W., and Zhou, L.: AIRS: Improving Weather Forecasting and Providing New Data on Greenhouse Gases, *B. Am. Meteorol. Soc.*, 87, 911–926, <https://doi.org/10.1175/BAMS-87-7-911>, 2006.
- Considine, D. B., Douglass, A. R., Connell, P. S., Kinnison, D. E., and Rotman, D. A.: A polar stratospheric cloud parameterization for the
 700 global modeling initiative three-dimensional model and its response to stratospheric aircraft, *J. Geophys. Res.: Atmos.*, 105, 3955–3973, <https://doi.org/10.1029/1999JD900932>, 2000.
- Cunnold, D. M., Fraser, P. J., Weiss, R. F., Prinn, R. G., Simmonds, P. G., Miller, B. R., Alyea, F. N., and Crawford, A. J.: Global trends and annual releases of CCl_3F and CCl_2F_2 estimated from ALE/GAGE and other measurements from July 1978 to June 1991, *J. Geophys. Res.: Atmos.*, 99, 1107–1126, <https://doi.org/10.1029/93JD02715>, 1994.
- 705 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- 710 Diehl, T., Heil, A., Chin, M., Pan, X., Streets, D., Schultz, M., and Kinne, S.: Anthropogenic, biomass burning, and volcanic emissions of black carbon, organic carbon, and SO_2 from 1980 to 2010 for hindcast model experiments, *Atmos. Chem. Phys. Discuss.*, 12, 24 895–24 954, <https://doi.org/10.5194/acpd-12-24895-2012>, 2012.
- Dipankar, A., Stevens, B., Heinze, R., Moseley, C., Zängl, G., Giorgetta, M., and Brdar, S.: Large eddy simulation using the general circulation model ICON, *J. Adv. Model. Earth Sy.*, 7, 963–986, <https://doi.org/10.1002/2015MS000431>, 2015.
- 715 Doms, G., Förstner, J., Heise, E., Herzog, H.-J., Mironov, D., Raschendorfer, M., Reinhardt, T., Ritter, B., Schrodin, R., Schulz, J.-P., and Vogel, G.: A Description of the Nonhydrostatic Regional COSMO Model. Part II: Physical Parametrization, Tech. rep., Deutscher Wetterdienst, Offenbach, 2011.
- Dörnbrack, A., Birner, T., Fix, A., Flentje, H., Meister, A., Schmid, H., Browell, E. V., and Mahoney, M. J.: Evidence for inertia gravity waves forming polar stratospheric clouds over Scandinavia, *J. Geophys. Res.: Atmospheres*, 107, SOL 30–1–SOL 30–18, <https://doi.org/10.1029/2001JD000452>, 2002.
- 720 Dörnbrack, A., Pitts, M. C., Poole, L. R., Orsolini, Y. J., Nishii, K., and Nakamura, H.: The 2009–2010 Arctic stratospheric winter – general evolution, mountain waves and predictability of an operational weather forecast model, *Atmos. Chem. Phys.*, 12, 3659–3675, <https://doi.org/10.5194/acp-12-3659-2012>, 2012.
- Dörnbrack, A., Kaifler, B., Kaifler, N., Rapp, M., Wildmann, N., Garhammer, M., Ohlmann, K., Payne, J. M., Sandercock, M., and
 725 Austin, E. J.: Unusual appearance of mother-of-pearl clouds above El Calafate, Argentina ($50^{\circ}21'S$, $72^{\circ}16'W$), *Weather*, 75, 378–388, <https://doi.org/10.1002/wea.3863>, 2020.

- Drdla, K., Turco, R. P., and Elliott, S.: Heterogeneous chemistry on Antarctic polar stratospheric clouds: A microphysical estimate of the extent of chemical processing, *J. Geophys. Res.: Atmos.*, 98, 8965–8981, <https://doi.org/10.1029/93JD00164>, 1993.
- Eckermann, S. D., Dörnbrack, A., Flentje, H., Vosper, S. B., Mahoney, M. J., Bui, T. P., and Carslaw, K. S.: Mountain Wave-
730 Induced Polar Stratospheric Cloud Forecasts for Aircraft Science Flights during SOLVE/THESEO 2000, *Weather Forecast.*, 21, 42–68, <https://doi.org/10.1175/WAF901.1>, 2006.
- Eckermann, S. D., Hoffmann, L., Höpfner, M., Wu, D. L., and Alexander, M. J.: Antarctic NAT PSC belt of June 2003: Observational validation of the mountain wave seeding hypothesis, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2008GL036629>, 2009.
- Engel, I., Luo, B. P., Pitts, M. C., Poole, L. R., Hoyle, C. R., Grooß, J.-U., Dörnbrack, A., and Peter, T.: Heterogeneous formation of polar
735 stratospheric clouds – Part 2: Nucleation of ice on synoptic scales, *Atmos. Chem. Phys.*, 13, 10 769–10 785, <https://doi.org/10.5194/acp-13-10769-2013>, 2013.
- Fritts, D. C. and Alexander, M. J.: Gravity wave dynamics and effects in the middle atmosphere, *Rev. Geophys.*, 41, <https://doi.org/10.1029/2001RG000106>, 2003.
- Fueglistaler, S., Buss, S., Luo, B. P., Wernli, H., Flentje, H., Hostetler, C. A., Poole, L. R., Carslaw, K. S., and Peter, T.: Detailed modeling
740 of mountain wave PSCs, *Atmos. Chem. Phys.*, 3, 697–712, <https://doi.org/10.5194/acp-3-697-2003>, 2003.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454,
745 <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Geller, M. A., Alexander, M. J., Love, P. T., Bacmeister, J., Ern, M., Hertzog, A., Manzini, E., Preusse, P., Sato, K., Scaife, A. A., and Zhou, T.: A Comparison between Gravity Wave Momentum Fluxes in Observations and Climate Models, *J. Climate*, 26, 6383 – 6405, <https://doi.org/10.1175/JCLI-D-12-00545.1>, 2013.
- Giorgetta, M. A., Brokopf, R., Crueger, T., Esch, M., Fiedler, S., Helmert, J., Hohenegger, C., Kornblueh, L., Köhler, M., Manzini, E.,
750 Mauritsen, T., Nam, C., Raddatz, T., Rast, S., Reinert, D., Sakradzija, M., Schmidt, H., Schneck, R., Schnur, R., Silvers, L., Wan, H., Zängl, G., and Stevens, B.: ICON-A, the Atmosphere Component of the ICON Earth System Model: I. Model Description, *J. Adv. Model. Earth Sy.*, 18, <https://doi.org/10.1029/2017MS001242>, 2018.
- Granier, C., Bessagnet, B., Bond, T., D’Angiola, A., Denier van der Gon, H., Frost, G., Heil, A., Kaiser, J., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J.-F., Lioussé, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J.-C., Riahi, K., Schultz, M., Smith, S., Thompson, A., van Aardenne, J., van der Werf, G., and van Vuuren, D.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980-2010 period, *Climatic Change*, 109, 163–190, <https://doi.org/10.1007/s10584-011-0154-1>, 2011.
- Grimsdell, A. W., Alexander, M. J., May, P. T., and Hoffmann, L.: Model Study of Waves Generated by Convection with Direct Validation via Satellite, *J. Atmos. Sci.*, 67, 1617 – 1631, <https://doi.org/10.1175/2009JAS3197.1>, 2010.
- 760 Hanson, D. R. and Mauersberger, K.: Laboratory studies of the nitric acid trihydrate: Implications for the south polar stratosphere, *Geophys. Res. Lett.*, 15, 855–858, <https://doi.org/10.1029/GL015i008p00855>, 1988.
- Hervig, M. E. and Deshler, T.: Stratospheric aerosol surface area and volume inferred from HALOE, CLAES, and ILAS measurements, *J. Geophys. Res.: Atmos.*, 103, 25 345–25 352, <https://doi.org/10.1029/98JD01962>, 1998.

Hoffmann, L. and Alexander, M. J.: Occurrence frequency of convective gravity waves during the North American thunderstorm season, *J. Geophys. Res.: Atmos.*, 115, <https://doi.org/10.1029/2010JD014401>, 2010.

Hoffmann, L., Xue, X., and Alexander, M. J.: A global view of stratospheric gravity wave hotspots located with Atmospheric Infrared Sounder observations, *Journal of Geophysical Research: Atmospheres*, 118, 416–434, <https://doi.org/10.1029/2012JD018658>, 2013.

Hoffmann, L., Alexander, M. J., Clerbaux, C., Grimsdell, A. W., Meyer, C. I., Rößler, T., and Tournier, B.: Intercomparison of stratospheric gravity wave observations with AIRS and IASI, *Atmospheric Measurement Techniques*, 7, 4517–4537, <https://doi.org/10.5194/amt-7-4517-2014>, 2014.

Hoffmann, L., Grimsdell, A. W., and Alexander, M. J.: Stratospheric gravity waves at Southern Hemisphere orographic hotspots: 2003–2014 AIRS/Aqua observations, *Atmos. Chem. Phys.*, 16, 9381–9397, <https://doi.org/10.5194/acp-16-9381-2016>, 2016.

Hoffmann, L., Spang, R., Orr, A., Alexander, M. J., Holt, L. A., and Stein, O.: A decadal satellite record of gravity wave activity in the lower stratosphere to study polar stratospheric cloud formation, *Atmos. Chem. Phys.*, 17, 2901–2920, <https://doi.org/10.5194/acp-17-2901-2017>, 2017.

Höpfner, M., Larsen, N., Spang, R., Luo, B. P., Ma, J., Svendsen, S. H., Eckermann, S. D., Knudsen, B., Massoli, P., Cairo, F., Stiller, G., v. Clarmann, T., and Fischer, H.: MIPAS detects Antarctic stratospheric belt of NAT PSCs caused by mountain waves, *Atmos. Chem. Phys.*, 6, 1221–1230, <https://doi.org/10.5194/acp-6-1221-2006>, 2006a.

Höpfner, M., Luo, B. P., Massoli, P., Cairo, F., Spang, R., Snels, M., Di Donfrancesco, G., Stiller, G., von Clarmann, T., Fischer, H., and Biermann, U.: Spectroscopic evidence for NAT, STS, and ice in MIPAS infrared limb emission measurements of polar stratospheric clouds, *Atmos. Chem. Phys.*, 6, 1201–1219, <https://doi.org/10.5194/acp-6-1201-2006>, 2006b.

Höpfner, M., Pitts, M. C., and Poole, L. R.: Comparison between CALIPSO and MIPAS observations of polar stratospheric clouds, *J. Geophys. Res.: Atmos.*, 114, <https://doi.org/10.1029/2009JD012114>, d00H05, 2009.

Höpfner, M., Deshler, T., Pitts, M., Poole, L., Spang, R., Stiller, G., and von Clarmann, T.: The MIPAS/Envisat climatology (2002–2012) of polar stratospheric cloud volume density profiles, *Atmos. Meas. Tech.*, 11, 5901–5923, <https://doi.org/10.5194/amt-11-5901-2018>, 2018.

Jähn, M., Kuhlmann, G., Mu, Q., Haussaire, J.-M., Ochsner, D., Osterried, K., Clément, V., and Brunner, D.: An online emission module for atmospheric chemistry transport models: implementation in COSMO-GHG v5.6a and COSMO-ART v5.1-3.1, *Geosci. Model Dev.*, 13, 2379–2392, <https://doi.org/10.5194/gmd-13-2379-2020>, 2020.

Janssens-Maenhout, G., Petrescu, A. M., Muntean, M., and Blujdea, V.: Verifying Greenhouse Gas Emissions: Methods to Support International Climate Agreements, *Greenhouse Gas Measurement and Management*, 1, 132–133, <https://doi.org/10.1080/20430779.2011.579358>, 2011.

Janssens-Maenhout, G., Diego, V., and Marilena Muntean, G.: Global emission inventories in the Emission Database for Global Atmospheric Research (EDGAR)—Manual (I), Gridding: EDGAR emissions distribution on global gridmaps, Publications Office of the European Union, Luxembourg, 2013.

Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geosci. Model Dev.*, 3, 717–752, <https://doi.org/10.5194/gmd-3-717-2010>, 2010.

Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C. A. M., Brinkop, S., Cai, D. S., Dyroff, C., Eckstein, J., Frank, F., Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, M., Meul, S., Neumaier, M., Nützel, M., Oberländer-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth System Chemistry integrated Modelling (ES-CiMo) with the Modular Earth Submodel System (MESSy) version 2.51, *Geosci. Model Dev.*, 9, 1153–1200, <https://doi.org/10.5194/gmd-9-1153-2016>, 2016.

- Kang, M.-J., Chun, H.-Y., and Kim, Y.-H.: Momentum Flux of Convective Gravity Waves Derived from an Offline Gravity Wave Parameterization. Part I: Spatiotemporal Variations at Source Level, *J. Atmos. Sci.*, 74, 3167 – 3189, <https://doi.org/10.1175/JAS-D-17-0053.1>, 2017.
- 805 Kennard, E. H.: Kinetic theory of gases, with an introduction to statistical mechanics, McGraw-Hill Book Company, New York and London, <https://archive.org/details/in.ernet.dli.2015.1789>, 1938.
- Khosrawi, F., Kirner, O., Stiller, G., Höpfner, M., Santee, M. L., Kellmann, S., and Braesicke, P.: Comparison of ECHAM5/MESy Atmospheric Chemistry (EMAC) simulations of the Arctic winter 2009/2010 and 2010/2011 with Envisat/MIPAS and Aura/MLS observations, *Atmos. Chem. Phys.*, 18, 8873–8892, <https://doi.org/10.5194/acp-18-8873-2018>, 2018.
- 810 Kirner, O., Ruhnke, R., Buchholz-Dietsch, J., Jöckel, P., Bruehl, C., and Steil, B.: Simulation of polar stratospheric clouds in the chemistry-climate-model EMAC via the submodel PSC, *Geosci. Model Dev.*, 4, 169–182, <https://doi.org/10.5194/gmd-4-169-2011>, 2011.
- Koop, T., Luo, B., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, <https://doi.org/10.1038/35020537>, 2000.
- Krieger, U. K., Mössinger, J. C., Luo, B., Weers, U., and Peter, T.: Measurement of the refractive indices of H₂SO₄–HNO₃–H₂O solutions to stratospheric temperatures, *Appl. Opt.*, 39, 3691–3703, <https://doi.org/10.1364/AO.39.003691>, 2000.
- 815 Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liou, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, <https://doi.org/10.5194/acp-10-7017-2010>, 2010.
- 820 Lamarque, J.-F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, A., and Zeng, G.: The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): overview and description of models, simulations and climate diagnostics, *Geosci. Model Dev.*, 6, 179–206, <https://doi.org/10.5194/gmd-6-179-2013>, 2013.
- 825 Langematz, U., M. Tully (Lead Authors), Calvo, N., Dameris, M., de Laat, A. T. J., Klekociuk, A., Müller, R., and Young, P.: Chapter 4: Polar Stratospheric Ozone: Past, Present, and Future, in: *Scientific Assessment of Ozone Depletion: 2018*, edited by Fahey, D. W., Newman, P. A., Pyle, J. A., and Safari, B., Global Ozone Research and Monitoring Project – Report No. 58, World Meteorological Organization, Geneva, Switzerland, 2018.
- Leuenberger, D., Koller, M., Fuhrer, O., and Schär, C.: A Generalization of the SLEVE Vertical Coordinate, *Mon. Weather Rev.*, 138, 3683–3689, <https://doi.org/10.1175/2010MWR3307.1>, 2010.
- 830 Lickley, M., Solomon, S., Fletcher, S., Velders, G. J. M., Daniel, J., Rigby, M., Montzka, S. A., Kuijpers, L. J. M., and Stone, K.: Quantifying contributions of chlorofluorocarbon banks to emissions and impacts on the ozone layer and climate, *Nature Comm.*, 11, 1380, <https://doi.org/10.1038/s41467-020-15162-7>, 2020.
- Mann, G. W., Carslaw, K. S., Chipperfield, M. P., Davies, S., and Eckermann, S. D.: Large nitric acid trihydrate particles and denitrification caused by mountain waves in the Arctic stratosphere, *J. Geophys. Res.: Atmos.*, 110, <https://doi.org/10.1029/2004JD005271>, d08202, 2005.
- 835 Marti, J. and Mauersberger, K.: A survey and new measurements of ice vapor pressure at temperatures between 170 and 250K, *Geophys. Res. Lett.*, 20, 363–366, <https://doi.org/10.1029/93GL00105>, 1993.

- McDonald, A. J., George, S. E., and Woollands, R. M.: Can gravity waves significantly impact PSC occurrence in the Antarctic?, *Atmos. Chem. Phys.*, 9, 8825–8840, <https://doi.org/10.5194/acp-9-8825-2009>, 2009.
- Meilinger, S. K., Koop, T., Luo, B. P., Huthwelker, T., Carslaw, K. S., Krieger, U., Crutzen, P. J., and Peter, T.: Size-dependent stratospheric droplet composition in Lee wave temperature fluctuations and their potential role in PSC freezing, *Geophys. Res. Lett.*, 22, 3031–3034, <https://doi.org/10.1029/95GL03056>, 1995.
- Middlebrook, A. M., Berland, B. S., George, S. M., Tolbert, M. A., and Toon, O. B.: Real refractive indices of infrared-characterized nitric-acid/ice films: Implications for optical measurements of polar stratospheric clouds, *J. Geophys. Res.: Atmospheres*, 99, 25 655–25 666, <https://doi.org/10.1029/94JD02391>, 1994.
- Mishchenko, M. I., Travis, L. D., and Mackowski, D. W.: T-matrix computations of light scattering by nonspherical particles: A review, *J. Quant. Spectrosc. Radiat. Transf.*, 55, 535 – 575, [https://doi.org/10.1016/0022-4073\(96\)00002-7](https://doi.org/10.1016/0022-4073(96)00002-7), light Scattering by Non-Spherical Particles, 1996.
- Montzka, S. A., Dutton, G. S., Yu, P., Ray, E., Portmann, R. W., Daniel, J. S., Kuijpers, L., Hall, B. D., Mondeel, D., Siso, C., Nance, J. D., Rigby, M., Manning, A. J., Hu, L., Moore, F., Miller, B. R., and Elkins, J. W.: An unexpected and persistent increase in global emissions of ozone-depleting CFC-11, *Nature*, 557, 413–417, <https://doi.org/10.1038/s41586-018-0106-2>, 2018.
- Morgenstern, O., Hegglin, M. I., Rozanov, E., O'Connor, F. M., Abraham, N. L., Akiyoshi, H., Archibald, A. T., Bekki, S., Butchart, N., Chipperfield, M. P., Deushi, M., Dhomse, S. S., Garcia, R. R., Hardiman, S. C., Horowitz, L. W., Jöckel, P., Josse, B., Kinnison, D., Lin, M., Mancini, E., Manyin, M. E., Marchand, M., Marécal, V., Michou, M., Oman, L. D., Pitari, G., Plummer, D. A., Revell, L. E., Saint-Martin, D., Schofield, R., Stenke, A., Stone, K., Sudo, K., Tanaka, T. Y., Tilmes, S., Yamashita, Y., Yoshida, K., and Zeng, G.: Review of the global models used within phase 1 of the Chemistry–Climate Model Initiative (CCMI), *Geoscientific Model Development*, 10, 639–671, <https://doi.org/10.5194/gmd-10-639-2017>, 2017.
- Murphy, D. M. and Ravishankara, A. R.: Temperature averages and rates of stratospheric reactions, *Geophysical Research Letters*, 21, 2471–2474, <https://doi.org/10.1029/94GL02287>, 1994.
- Nakajima, H., Murata, I., Nagahama, Y., Akiyoshi, H., Saeki, K., Kinase, T., Takeda, M., Tomikawa, Y., Dupuy, E., and Jones, N. B.: Chlorine partitioning near the polar vortex edge observed with ground-based FTIR and satellites at Syowa Station, Antarctica, in 2007 and 2011, *Atmos. Chem. Phys.*, 20, 1043–1074, <https://doi.org/10.5194/acp-20-1043-2020>, 2020.
- Nash, E. R., Newman, P. A., Rosenfield, J. E., and Schoeberl, M. R.: An objective determination of the polar vortex using Ertel's potential vorticity, *J. Geophys. Res.: Atmos.*, 101, 9471–9478, <https://doi.org/10.1029/96JD00066>, 1996.
- Noel, V. and Pitts, M.: Gravity wave events from mesoscale simulations, compared to polar stratospheric clouds observed from spaceborne lidar over the Antarctic Peninsula, *J. Geophys. Res.: Atmos.*, 117, <https://doi.org/10.1029/2011JD017318>, 2012.
- Noel, V., Hertzog, A., and Chepfer, H.: CALIPSO observations of wave-induced PSCs with near-unity optical depth over Antarctica in 2006–2007, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2008JD010604>, 2009.
- Orr, A., Hosking, J. S., Hoffmann, L., Keeble, J., Dean, S. M., Roscoe, H. K., Abraham, N. L., Vosper, S., and Braesicke, P.: Inclusion of mountain-wave-induced cooling for the formation of PSCs over the Antarctic Peninsula in a chemistry–climate model, *Atmos. Chem. Phys.*, 15, 1071–1086, <https://doi.org/10.5194/acp-15-1071-2015>, 2015.
- Orr, A., Hosking, J. S., Delon, A., Hoffmann, L., Spang, R., Moffat-Griffin, T., Keeble, J., Abraham, N. L., and Braesicke, P.: Polar stratospheric clouds initiated by mountain waves in a global chemistry–climate model: a missing piece in fully modelling polar stratospheric ozone depletion, *Atmos. Chem. Phys.*, 20, 12 483–12 497, <https://doi.org/10.5194/acp-20-12483-2020>, 2020.

- Peter, T. and Grooß, J.-U.: Chapter 4 Polar Stratospheric Clouds and Sulfate Aerosol Particles: Microphysics, Denitrification and Heterogeneous Chemistry, in: *Stratospheric Ozone Depletion and Climate Change*, edited by Müller, R., pp. 108–144, The Royal Society of Chemistry, <https://doi.org/10.1039/9781849733182-00108>, 2012.
- Pitts, M. C., Poole, L. R., and Thomason, L. W.: CALIPSO polar stratospheric cloud observations: second-generation detection algorithm and composition discrimination, *Atmos. Chem. Phys.*, 9, 7577–7589, <https://doi.org/10.5194/acp-9-7577-2009>, 2009.
- Pitts, M. C., Poole, L. R., and Gonzalez, R.: Polar stratospheric cloud climatology based on CALIPSO spaceborne lidar measurements from 2006 to 2017, *Atmos. Chem. Phys.*, 18, 10 881–10 913, <https://doi.org/10.5194/acp-18-10881-2018>, 2018.
- Plougonven, R., Hertzog, A., and Teitelbaum, H.: Observations and simulations of a large-amplitude mountain wave breaking over the Antarctic Peninsula, *J. Geophys. Res.: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009739>, 2008.
- 885 Prather, M. J.: Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3c, *Geosci. Model Dev.*, 8, 2587–2595, <https://doi.org/10.5194/gmd-8-2587-2015>, 2015.
- Preusse, P., Ern, M., Bechtold, P., Eckermann, S. D., Kalisch, S., Trinh, Q. T., and Riese, M.: Characteristics of gravity waves resolved by ECMWF, *Atmospheric Chemistry and Physics*, 14, 10 483–10 508, <https://doi.org/10.5194/acp-14-10483-2014>, 2014.
- Queney, P.: *Theory of perturbations in stratified currents with applications to air flow over mountain barriers*, University of Chicago Press, 890 Chicago, Ill, 1947.
- Reinert, D., Prill, F., Zängl, G., Rieger, D., Schröter, J., Förstner, J., Werchner, S., Weimer, M., Ruhnke, R., and Vogel, B.: Working with the ICON Model, Tech. rep., Deutscher Wetterdienst, Offenbach, https://code.mpimet.mpg.de/attachments/download/19568/ICON_tutorial_2019.pdf, 2019.
- Rieger, D., Bangert, M., Bischoff-Gauss, I., Förstner, J., Lundgren, K., Reinert, D., Schröter, J., Vogel, H., Zängl, G., Ruhnke, R., and 895 Vogel, B.: ICON-ART 1.0 - a new online-coupled model system from the global to regional scale, *Geosci. Model Dev.*, 8, 1659–1676, <https://doi.org/10.5194/gmd-8-1659-2015>, 2015.
- Salcedo, D., Molina, L. T., and Molina, M. J.: Homogeneous Freezing of Concentrated Aqueous Nitric Acid Solutions at Polar Stratospheric Temperatures, *J. Phys. Chem.*, 105, 1433–1439, <https://doi.org/10.1021/jp001639s>, 2001.
- Sander, R., Baumgaertner, A., Gromov, S., Harder, H., Jöckel, P., Kerkweg, A., Kubistin, D., Regelin, E., Riede, H., Sandu, A., Taraborrelli, D., Tost, H., and Xie, Z.-Q.: The atmospheric chemistry box model CAABA/MECCA-3.0, *Geosci. Model Dev.*, 4, 373–380, 900 <https://doi.org/10.5194/gmd-4-373-2011>, 2011a.
- Sander, S., Abbatt, J., Barker, J., Burkholder, J., Friedl, R., Golden, D., Huie, R., Kolb, C., Kurylo, M., Moortgat, K., Orkin, V., and Wine, P.: *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies*, Evaluation No. 17, JPL Publication 10-6, 2011b.
- Sandu, A. and Sander, R.: Technical note: Simulating chemical systems in Fortran90 and Matlab with the Kinetic PreProcessor KPP-2.1, 905 *Atmos. Chem. Phys.*, 6, 187–195, <https://doi.org/10.5194/acp-6-187-2006>, 2006.
- Schoeberl, M. and Newman, P.: MIDDLE ATMOSPHERE, Polar Vortex, in: *Encyclopedia of Atmospheric Sciences*, edited by Holton, J. R., pp. 1321–1328, Academic Press, Oxford, <https://doi.org/10.1016/B0-12-227090-8/00228-1>, 2003.
- Schröter, J., Rieger, D., Stassen, C., Vogel, H., Weimer, M., Werchner, S., Förstner, J., Prill, F., Reinert, D., Zängl, G., Giorgetta, M., Ruhnke, R., Vogel, B., and Braesicke, P.: ICON-ART 2.1: a flexible tracer framework and its application for composition studies in numerical 910 weather forecasting and climate simulations, *Geosci. Model Dev.*, 11, 4043–4068, <https://doi.org/10.5194/gmd-11-4043-2018>, 2018.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, Wiley, Hoboken, NJ, 2. ed. edn., previous ed.: 1998, 2006.

- Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrakou, T., Müller, J.-F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmos. Chem. Phys.*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014.
- Smith, R. B.: Hydrostatic Airflow over Mountains, vol. 31 of *Adv. Geophys.*, pp. 1–41, Elsevier, [https://doi.org/10.1016/S0065-2687\(08\)60052-7](https://doi.org/10.1016/S0065-2687(08)60052-7), 1989.
- Snels, M., Scoccione, A., Di Liberto, L., Colao, F., Pitts, M., Poole, L., Deshler, T., Cairo, F., Cagnazzo, C., and Fierli, F.: Comparison of Antarctic polar stratospheric cloud observations by ground-based and space-borne lidar and relevance for chemistry–climate models, *Atmos. Chem. Phys.*, 19, 955–972, <https://doi.org/10.5194/acp-19-955-2019>, 2019.
- Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, *Rev. Geophys.*, 37, 275–316, <https://doi.org/10.1029/1999RG900008>, 1999.
- Solomon, S., Garcia, R., Rowland, F., and Wuebbles: On the depletion of Antarctic ozone, *Nature*, 321, 755–758, <https://doi.org/10.1038/321755a0>, 1986.
- Solomon, S., Kinnison, D., Bandoro, J., and Garcia, R.: Simulation of polar ozone depletion: An update, *J. Geophys. Res.: Atmos.*, 120, 7958–7974, <https://doi.org/10.1002/2015JD023365>, 2015.
- Spang, R., Hoffmann, L., Höpfner, M., Griessbach, S., Müller, R., Pitts, M. C., Orr, A. M. W., and Riese, M.: A multi-wavelength classification method for polar stratospheric cloud types using infrared limb spectra, *Atmos. Meas. Tech.*, 9, 3619–3639, <https://doi.org/10.5194/amt-9-3619-2016>, 2016.
- Spang, R., Hoffmann, L., Müller, R., Grooß, J.-U., Tritscher, I., Höpfner, M., Pitts, M., Orr, A., and Riese, M.: A climatology of polar stratospheric cloud composition between 2002 and 2012 based on MIPAS/Envisat observations, *Atmos. Chem. Phys.*, 18, 5089–5113, <https://doi.org/10.5194/acp-18-5089-2018>, 2018.
- SPARC: SPARC Newsletter No. 40, available at <http://www.sparc-climate.org/publications/newsletter>, 2013.
- Staniforth, A. and Thuburn, J.: Horizontal grids for global weather and climate prediction models: a review, *Q. J. Roy. Meteor. Soc.*, 138, 1–26, <https://doi.org/10.1002/qj.958>, 2012.
- Steiner, M., Luo, B., Peter, T., Pitts, M. C., and Stenke, A.: Evaluation of polar stratospheric clouds in the global chemistry–climate model SOCOLv3.1 by comparison with CALIPSO spaceborne lidar measurements, *Geosci. Model Dev.*, 14, 935–959, <https://doi.org/10.5194/gmd-14-935-2021>, 2021.
- Stephens, G. L., Vane, D. G., Boain, R. J., Mace, G. G., Sassen, K., Wang, Z., Illingworth, A. J., O’connor, E. J., Rossow, W. B., Durden, S. L., Miller, S. D., Austin, R. T., Benedetti, A., and Mitrescu, C. a.: The CloudSat Mission and the A-Train, *B. Am. Meteorol. Soc.*, 83, 1771–1790, <https://doi.org/10.1175/BAMS-83-12-1771>, 2002.
- Stokes, G. G.: On the Effect of the Internal Friction of Fluids on the Motion of Pendulums, *Transactions of the Cambridge Philosophical Society, Part II*, 8 – 106, 1851.
- Stone, K. A., Solomon, S., Kinnison, D. E., Baggett, C. F., and Barnes, E. A.: Prediction of Northern Hemisphere Regional Surface Temperatures Using Stratospheric Ozone Information, *J. Geophys. Res.: Atmos.*, 124, 5922–5933, <https://doi.org/10.1029/2018JD029626>, 2019.
- Svendsen, S. H., Larsen, N., Knudsen, B., Eckermann, S. D., and Browell, E. V.: Influence of mountain waves and NAT nucleation mechanisms on polar stratospheric cloud formation at local and synoptic scales during the 1999–2000 Arctic winter, *Atmos. Chem. Phys.*, 5, 739–753, <https://doi.org/10.5194/acp-5-739-2005>, 2005.

- 950 Tabazadeh, A., Santee, M. L., Danilin, M. Y., Pumphrey, H. C., Newman, P. A., Hamill, P. J., and Mergenthaler, J. L.: Quantifying Denitrification and Its Effect on Ozone Recovery, *Science*, 288, 1407–1411, <https://doi.org/10.1126/science.288.5470.1407>, 2000.
- Taylor, K., Williamson, D., and Zwiers, F.: The Sea Surface Temperature and Sea-ice Concentration Boundary Conditions for AMIP II Simulations, PCMDI report, Program for Climate Model Diagnosis and Intercomparison, Lawrence Livermore National Laboratory, 2000.
- Thomason, L. W., Burton, S. P., Luo, B.-P., and Peter, T.: SAGE II measurements of stratospheric aerosol properties at non-volcanic levels, 955 *Atmos. Chem. Phys.*, 8, 983–995, <https://doi.org/10.5194/acp-8-983-2008>, 2008.
- Tritscher, I., Grooß, J.-U., Spang, R., Pitts, M. C., Poole, L. R., Müller, R., and Riese, M.: Lagrangian simulation of ice particles and resulting dehydration in the polar winter stratosphere, *Atmos. Chem. Phys.*, 19, 543–563, <https://doi.org/10.5194/acp-19-543-2019>, 2019.
- Tritscher, I., Pitts, M. C., Poole, L. R., Alexander, S. P., Cairo, F., Chipperfield, M. P., Grooß, J.-U., Höpfner, M., Lambert, A., Luo, B. P., Molleker, S., Orr, A., Salawitch, R., Snels, M., Spang, R., Woiwode, W., and Peter, T.: Polar Stratospheric Clouds Satellite Observations, 960 *Processes, and Role in Ozone Depletion*, *Rev. Geophys.*, 59, e2020RG000702, <https://doi.org/10.1029/2020RG000702>, e2020RG000702, 2021.
- Tully, M., Klekociuk, A., Alexander, S., Dargaville, R., Deschamps, L., Fraser, P., Gies, H., Henderson, S., Javorniczky, J., Krummel, P., Petelina, S., Shanklin, J., Siddaway, J., and Stone, K.: The Antarctic ozone hole during 2008 and 2009, *J. So. Hemisph. Earth*, 61, 77–90, <https://doi.org/10.22499/2.6101.007>, 2011.
- 965 van den Broek, M. M. P., Williams, J. E., and Bregman, A.: Implementing growth and sedimentation of NAT particles in a global Eulerian model, *Atmos. Chem. Phys.*, 4, 1869–1883, <https://doi.org/10.5194/acp-4-1869-2004>, 2004.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, <https://doi.org/10.5194/acp-6-3423-2006>, 2006.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van 970 Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11 707–11 735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.
- Waibel, A. E., Peter, T., Carslaw, K. S., Oelhaf, H., Wetzel, G., Crutzen, P. J., Pöschl, U., Tsias, A., Reimer, E., and Fischer, H.: Arctic Ozone Loss Due to Denitrification, *Science*, 283, 2064–2069, <https://doi.org/10.1126/science.283.5410.2064>, 1999.
- Wegner, T., Kinnison, D. E., Garcia, R. R., and Solomon, S.: Simulation of polar stratospheric clouds in the specified dynamics version of 975 the whole atmosphere community climate model, *J. Geophys. Res.: Atmospheres*, 118, 4991–5002, <https://doi.org/10.1002/jgrd.50415>, 2013.
- Weimer, M.: Towards Seamless Simulations of Polar Stratospheric Clouds and Ozone in the Polar Stratosphere with ICON-ART, Ph.D. thesis, Karlsruher Institut für Technologie (KIT), <https://doi.org/10.5445/IR/1000100338>, 2019.
- Weimer, M., Mieruch, S., Schädler, G., and Kottmeier, C.: A new estimator of heat periods for decadal climate predictions – a complex 980 network approach, *Nonlin. Processes Geophys.*, 23, 307–317, <https://doi.org/10.5194/npg-23-307-2016>, 2016.
- Weimer, M., Schröter, J., Eckstein, J., Deetz, K., Neumaier, M., Fischbeck, G., Hu, L., Millet, D. B., Rieger, D., Vogel, H., Vogel, B., Reddmann, T., Kirner, O., Ruhnke, R., and Braesicke, P.: An emission module for ICON-ART 2.0: implementation and simulations of acetone, *Geosci. Model Dev.*, 10, 2471–2494, <https://doi.org/10.5194/gmd-10-2471-2017>, 2017.
- Winker, D. M., Hunt, W. H., and McGill, M. J.: Initial performance assessment of CALIOP, *Geophys. Res. Lett.*, 34, 985 <https://doi.org/10.1029/2007GL030135>, 2007.

- Wright, C. J., Hindley, N. P., Hoffmann, L., Alexander, M. J., and Mitchell, N. J.: Exploring gravity wave characteristics in 3-D using a novel S-transform technique: AIRS/Aqua measurements over the Southern Andes and Drake Passage, *Atmos. Chem. Phys.*, 17, 8553–8575, <https://doi.org/10.5194/acp-17-8553-2017>, 2017.
- 990 Zambri, B., Solomon, S., Kinnison, D. E., Mills, M. J., Schmidt, A., Neely III, R. R., Bourassa, A. E., Degenstein, D. A., and Roth, C. Z.: Modeled and Observed Volcanic Aerosol Control on Stratospheric NO_y and Cly, *J. Geophys. Res.: Atmos.*, 124, 10 283–10 303, <https://doi.org/10.1029/2019JD031111>, 2019.
- Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOsahedral Non-hydrostatic) modelling framework of DWD and MPI-M: Description of the non-hydrostatic dynamical core, *Q. J. R. Meteorol. Soc.*, 141, 563–579, <https://doi.org/10.1002/qj.2378>, 2015.
- 995 Zhu, Y., Toon, O. B., Lambert, A., Kinnison, D. E., Brakebusch, M., Bardeen, C. G., Mills, M. J., and English, J. M.: Development of a Polar Stratospheric Cloud Model within the Community Earth System Model using constraints on Type I PSCs from the 2010-2011 Arctic winter, *J. Adv. Model. Earth Sy.*, 7, 551–585, <https://doi.org/10.1002/2015MS000427>, 2015.
- Zhu, Y., Toon, O. B., Pitts, M. C., Lambert, A., Bardeen, C., and Kinnison, D. E.: Comparing simulated PSC optical properties with CALIPSO observations during the 2010 Antarctic winter, *J. Geophys. Res.: Atmos.*, 122, 1175–1202, <https://doi.org/10.1002/2016JD025191>, 2017.