

Elevated 3D structures of PM_{2.5} and impact of complex terrain-forcing circulations on heavy haze pollution over Sichuan Basin, China

Zhuozhi Shu^{1,4}, Yubao Liu^{1,4}, Tianliang Zhao^{1,4,*}, Junrong Xia¹, Chenggang Wang¹,
5 Le Cao¹, Haoliang Wang¹, Lei Zhang², Yu Zheng², Lijuan Shen¹, Lei Luo³, Yueqing
Li³

¹Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Key
Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Nanjing
10 University of Information Science and Technology, Nanjing, 210044, China

²State Key Laboratory of Disastrous Weather, China Academy of Meteorological Sciences, Beijing,
100081, China

³Institute of Plateau Meteorology, China Meteorological Administration, Chengdu, 610072, China

⁴Precision Regional Earth Modeling and Information Center, Nanjing University of Information
15 Science and Technology, Nanjing, 210044, China

Correspondence: Tianliang Zhao (tlzhao@nuist.edu.cn)

Abstract. Deep basins create a uniquely favorable ~~condition for the formation of~~ air pollution ~~causing~~
~~condition~~, and the Sichuan Basin (SCB) in Southwest China ~~is such a basin featuring features~~ frequent
heavy pollution. A wintertime heavy haze pollution event in ~~the~~ SCB was studied with conventional
20 and intensive observation data and the WRF-~~Chem~~ model to explore the three-dimensional distribution
of PM_{2.5} ~~for to~~ understanding the impact of regional pollutant emissions, basin circulations associated
with plateaus, and downwind transport to the adjacent areas. It was found that the vertical structure of
PM_{2.5} over SCB was characterized ~~with by~~ a remarkable hollow sandwiched by high PM_{2.5} layers at
heights of 1.5–3 km and ~~athe~~ highly polluted near-surface layer. The southwesterlies ~~passed~~ over the
25 Tibetan Plateau (TP) and Yunan-Guizhou Plateau (YGP) resulted in a lee vortex over ~~the~~ SCB, which
helped form and maintain heavy PM_{2.5} pollution. The basin PM_{2.5} was lifted into the free troposphere
and transported outside of ~~the~~ SCB. At the bottom of ~~the~~ SCB, high PM_{2.5} concentrations were mostly
located in the northwest~~ern~~ and southern regions. Due to the blocking effect of the plateau terrain on
the northeasterly winds, PM_{2.5} gradually increased from northeast to southwest in the basin. In the
30 lower free troposphere, the high PM_{2.5} centers were distributed over the northwestern and southwestern
SCB areas, as well as the central SCB region. For this event, ~~the~~ regional emissions ~~from SCB~~
contributed 75.4–94.6 % to the surface PM_{2.5} concentrations in ~~the~~ SCB. The SCB emissions ~~export~~

~~was~~ were the major source of ~~the~~ PM_{2.5} over the eastern regions of ~~the~~ TP and the northern regions of YGP, with contribution rates of 72.7 % and 70.5 %, respectively, during the dissipation stage of heavy air pollution over SCB, which was regarded as the major pollutant source affecting atmospheric environmental changes in Southwest China.

1 Introduction

Haze pollution has ~~brought~~ caused serious environmental problems, especially in the densely populated and economically developed regions in China, ~~which have with~~ high levels of fine ~~particular~~ particulate matters (PM_{2.5}) (particulate matter with an aerodynamical diameter equal to or less than 2.5 μm) (Guo et al., 2014; Li et al., 2015; Gu and Yim, 2016; Lin et al., 2018). Owing to the significant adverse effects on human health and climate change (Dawson et al., 2007; Langrish et al., 2012; Megaritis et al., 2014; Guo et al., 2016), understanding PM_{2.5} pollution distributions and mechanisms is of high interest in environmental and climate studies.

Anthropogenic pollutant emissions and stagnant meteorological conditions are commonly regarded as two key factors ~~for~~ influencing haze pollution with excessive concentrations of PM_{2.5} (Yim et al., 2014; Zhang et al., 2015; Cai et al., 2017). With strong anthropogenic emissions and favorable meteorological conditions, four main regions with frequent heavy haze pollution ~~are determined~~ have been identified, centered over the North China Plain (NCP) (Tao et al., 2012; Ye et al., 2016; Zhang et al., 2016; Huang et al., 2017), the Yangtze River Delta (YRD) in East China (Wang et al., 2012; Li et al., 2015; Tang et al., 2015; Ming et al., 2017), the Pearl River Delta (PRD) in South China (Wu et al., 2013; Zhang et al., 2013; Zhang et al., 2014; Guo et al., 2016), and the Sichuan Basin (SCB) in Southwest China (Tao et al., 2013; Chen and Xie, 2014; Zhou et al., 2019), ~~respectively~~. Haze pollution over ~~the~~ NCP, YRD and PRD, the main economic centers, with ~~the~~ large flatlands has been extensively studied. However, air pollution in the SCB region with highly frequent heavy PM_{2.5} pollution has ~~not~~ been ~~insufficiently~~ completely understood owing to the complex deep basin terrain, ~~in~~ particularly the effect of the immediately adjoining Tibetan Plateau (TP).

The TP's "harbor effect" on the tropospheric westerlies favors a stable atmospheric stratification and low wind speeds in the boundary layer over the downstream SCB (Xu et al., 2015; Xu et al., 2016), which is conducive to air pollutant accumulation in SCB (Yim et al., 2014; Xu et al., 2016; Wang et al.,

2018). The downslope flows at the lee side of the plateau ~~could-can~~ induce ~~the-a~~ special stagnation meteorological condition in the lower troposphere (Wang et al., 2015; Ning et al., 2018a). ~~The-Air~~ ~~air-~~stagnation days account for 76.6 % of the total days in winter over ~~the~~ SCB (Liao et al., 2018), where near-surface ~~weakened~~ wind, strong vertical air temperature inversion and shallow boundary layer ~~could-~~significantly restrain ~~the~~ atmospheric diffusion capacity (Ning et al., 2018a; Wang et al., 2018; Tian et al., 2019), resulting in the occurrence of heavy air pollution in ~~the~~ SCB.

The SCB, covering 260,000 km² of ~~the~~ Sichuan-Chongqing plain with a dense population of more than 100 million people, is a deep basin ~~in Southwest China~~ surrounded by plateaus and mountains ~~in Southwest China~~. It ~~is-lies~~ immediately to the east of ~~the~~ TP₁ with a large elevation drop exceeding 3000 m over a short horizontal distance. The unique terrain effect generates the asymmetries of meteorological and air pollutant distribution (Zhang et al., 2019), with a remarkable difference ~~of-in~~ PM_{2.5} concentrations between the eastern and western regions over ~~the~~ SCB (Chen and Xie, 2012; Ning et al., 2018b). The weak vertical diffusion in the atmospheric boundary layer is one of the main causes ~~for-of~~ air pollution in winter (Ye et al., 2013; Hu et al., 2014; Tian et al., 2017; Zhao et al., 2018). Many ~~evidences-studies have~~ suggested that air pollution over ~~the~~ SCB is mostly caused by the accumulation of air pollutants originating from ~~the~~ local emissions (Chen et al., 2014; Liao et al., 2017; Wang et al., 2018; Qiao et al., 2019). However, ~~due-to~~~~because of~~ the complex flows in SCB, it is important to study how ~~the~~ PM_{2.5} is circulated three-dimensionally ~~in-order-to~~ ~~more-accurately~~ estimate the roles of local emissions and exchanges with outside regions ~~more accurately~~.

In this ~~paperstudy~~, ~~the~~ observation data analysis and numerical experiments were ~~carried-outconducted~~ to analyze the three-dimensional distribution of PM_{2.5} concentrations in SCB during a heavy haze pollution episode in January 2017. The contributions of the SCB pollutant emissions and ~~the~~ PM_{2.5} transport to the surrounding plateaus and mountains were estimated. Section 2 introduces the observation data and ~~the~~ modeling methods ~~for-used in~~ this study. Section 3 characterizes the horizontal and vertical distributions of PM_{2.5} concentrations, during the formation, maintenance and dissipation stages of the heavy haze pollution episode. We also assessed the contribution of local emissions to the heavy PM_{2.5} pollution within ~~the~~ SCB, and the impact of external transport of the ~~SCB-PM_{2.5}~~ ~~in the SCB on to~~ the surrounding areas in Southwest China. ~~The sSummaryies~~ and conclusions are provided in Section 4.

90 2. Data and model

2.1 Observation data

The surface air pollutant concentrations and meteorological elements observed in 18 cities (Fig. 1; Table 1) over the SCB were used to investigate the distribution of PM_{2.5}, weather circulations ~~as well as~~ the ~~and~~ modeling performance. The hourly meteorological observational data, ~~containing including~~ surface air temperature, relative humidity, wind speed and wind direction, were obtained from the Chinese meteorological monitoring network, and the hourly observational PM_{2.5} concentrations were obtained from the China National Environmental Monitoring Center (<http://www.cnemc.cn>).

~~Besides In addition to~~ the above-~~mentioned~~ conventional observations, sounding observations were conducted every ~~three hours~~ 3 h-~~by~~ using a kite balloon with the sounding system TT12 DigiCORA (Vaisala, Finland) at the Meteorological Observatory of Chengdu (Site 1 in Fig. 1) during 1–20 January, ~~January~~-2017. The vertical sounding data of air temperature, wind speed, wind direction, and relative humidity were observed at time intervals of 1 ~~second~~. In addition, a Micro Pulse Lidar Type 4 System-(MPL-4B-IDS, Sigma Space, America) was operated at the observational site Ya'an (Site 15 in Fig. 1) in the western SCB edge to retrieve the vertical PM_{2.5} structures at 532 nm (laser emission wavelength), 2500 Hz (laser repetition rate) and 6–8 μJ (optimal laser output range).

2.3 Model configuration and simulation experiments

The Weather Research and Forecasting with Chemistry (WRF-Chem, version 3.8.1) model was employed to simulate ~~the~~ severe haze pollution events over 2–8 January 2017 in the SCB (Fig. 2). ~~A~~ The spin-up time of modeling for the first 24 ~~hours~~ starting on 1 January 2017, was dropped. The ERA-Interim meteorological reanalysis data of the European Center for Medium-Range Weather Forecasts (<https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/>) ~~were~~ served as the initial and boundary conditions of the WRF-Chem simulation. The model domains and ~~the~~ topography ~~were~~ are ~~shown~~ presented in Fig. 1. There were three nesting domains with ~~aa~~ domain 1 (D1) covering ~~the~~ most areas of China, ~~a~~ domain 2 (D2) covering Southwest China, and an inner domain 3 (D3) covering the SCB and the surrounding areas, at grid intervals of 48, 12 and 3 km, respectively (Fig. 1). Considering

120 the complex terrain underlying the SCB's deep basin and surrounding plateaus and mountains in Southwest China, we adopted a grid ratio of 1:4 for simulation experiments with a precisely defined horizontal resolution. It should be noted that the even grid ratio may cause interpolation errors at the nested-domain boundary conditions owing to the nature of Arakawa C-grid staggering. An adequate vertical resolution is fundamental for evaluating thermal stratification over a complex terrain. Therefore, 35 vertical layers were set with fine resolutions of 30–120 m in the boundary layer. The physical schemes for the WRF-Chem simulations ~~were~~are listed in Table 2.

125 The Regional Acid Deposition Model, version 2 (RADM2) (Stockwell et al., 1990) was selected for the atmospheric chemistry mechanism, including the main inorganic ions, elemental carbon, primary and secondary organic aerosols, ~~as well as~~and other aerosol species. (Tuccella et al. 2012). The Multi-resolution Emission Inventory for China (MEIC) from 2012 (<http://www.meicmodel.org>) with a horizontal resolution of $0.25 \times 0.25^\circ$ was used ~~for modeling to model the~~ anthropogenic emissions of air pollutants. The Model of Emissions of Gases and Aerosols from Nature (v2.1) was applied to the natural emission sources in the simulation with dust emission parameterization.

130 High PM_{2.5} levels in the atmosphere could significantly reduce the near-ground solar radiation for stable atmospheric stratification, which decreases the vertical turbulent diffusion in the boundary layer (Wang et al., 2019). This is an important mechanism in the formation of severe haze pollution with the exponential growth of PM_{2.5} (Zhong et al., 2018). The overestimated vertical diffusion capacity under poor air quality conditions (Ren et al., 2019) causes deviations in air pollutant concentrations simulated in air quality models (Wang et al., 2018). In this study, the vertical turbulent diffusion coefficient of the atmospheric boundary layer was cut halfway for better simulation of the 3D structures of PM_{2.5}, during the heavy air pollution event over the SCB region.

140 Two simulation experiments were conducted: 1) a baseline simulation (Emi-Real), with the MEIC anthropogenic emission inventory over all three domains, and 2) a sensitivity simulation (Emi-Non~~;-~~), ~~same as~~similar to the baseline simulation–Emi-Real but involved shutting down the anthropogenic emission sources in the SCB (Fig. 1). By comparing the PM_{2.5} concentrations between ~~the experiments~~ Emi-Real and Emi-Non, we ~~could~~quantified the contribution of local emission sources to the heavy haze pollution over the SCB and estimated the transport from the polluted SCB to the adjoined areas over the ~~Eastern~~eastern TP, ~~the~~ northern YGP and ~~the~~ Daba Mountain (DBM) region (Fig. 1). The definite ranges of the three regions were defined with the altitudes of 750–3500 m over 30.5–33.0° N,

145

102.7–105.3 °E (the eastern TP edge), 750–3000 m over 27.8–29 °N, 103.5–108.5 °E (the northern YGP edge), and above 750 m over 31.5–33.0 °N, 106.0–109.4 °E (the DBM region), as shown in Fig. S5.

2.4 Case description

150 A severe haze pollution event occurred during 2–8 January 2017 in the SCB. As shown in Fig. 2, high and low PM_{2.5} concentrations were centered ~~respectively~~ in the western and eastern regions during the episode, respectively, presenting a generally asymmetric horizontal distribution.

Based on the National Ambient Air Quality Standards of China by the Ministry of Ecology and Environment in 2012 (<http://www.mee.gov.cn/>), light and heavy air pollution levels of PM_{2.5} ~~are-were~~ 155 categorized with daily average PM_{2.5} concentrations exceeding 75 and 150 µg m⁻³ in ambient air, respectively. The most heavily polluted regions ~~was-were~~ mainly concentrated in the northwestern city cluster of the SCB, including Chengdu, Deyang and Meishan, with ~~the~~ daily mean ~~concentrations of~~ PM_{2.5} concentrations exceeding 150 µg m⁻³ (Figs. 1 and 2a). An hourly PM_{2.5} peak of 345.0 µg m⁻³ was observed in Chengdu, a representative megacity in Southwest China. According to the hourly PM_{2.5} 160 variations in the city cluster over the northwestern SCB region, we divided the heavy haze episode into three periods, P1, P2 and P3, ~~respectively for~~ corresponding to formation (from 12:00 p.m. on 2 January to 0:00 on 5 January 2017), maintenance (from 0:00 a.m. on 5 January to 12:00 p.m. on 6 January 2017) and dissipation (from 12:00 p.m. on 6 January 0:00 a.m. on 6 January 2017) stages, respectively (local time was used in this study). As shown in Fig. 2b, during P1, the surface PM_{2.5} concentrations sharply 165 increased to the heavy haze pollution level, and then ~~it~~ fluctuated at the heavy pollution level in P2. Finally in P3, the concentrations of PM_{2.5} dropped below 75 µg m⁻³ and the event ended on 8 January 2017 (Fig. 2b).

The meteorological overview of the haze event was characterized by the 700 hPa fields of geopotential heights and wind vectors (Fig. 3). A trough in the mid-latitude westerlies moved eastward 170 from the eastern edge of the TP to the western SCB margin during P1, the trough of low pressure evolved over the SCB region during P2, and the westerly trough shifted out the SCB region with the low-pressure system disappearing during P3 (Fig. 3). The changes in atmospheric circulations in the three stages reflected the meteorological modulation of heavy haze development over the SCB in

association with the effect of TP topography on the westerlies.

175 Analysis of the observations ~~found-revealed~~ noteworthy patterns of spatial distribution of surface PM_{2.5} concentrations over SCB in the ~~formation, maintenance and dissipation~~three periods (Fig. 4). During P1, the surface PM_{2.5} concentrations were distributed relatively even over SCB, but ~~in the maintenance period~~during P2, the PM_{2.5} concentrations exhibited a northeast-southwest gradient and a dramatic increase in the western SCB area. For example, the surface PM_{2.5} concentrations increased
180 from 202.1 to 276.6 $\mu\text{g m}^{-3}$, from 148.6 to 181.0 $\mu\text{g m}^{-3}$, from 104.9 to 205.7 $\mu\text{g m}^{-3}$ and from 145.6 to 168.4 $\mu\text{g m}^{-3}$ ~~respectively~~, at sites 1, 3, 6 and 15, respectively (Fig. 1; Table 1). In contrast, during the dissipation period P3, strong northeasterly winds developed and the air quality was improved from the northeast to the southwest regions, with the reduction in PM_{2.5} cleaned-up concentrations in the northeastern SCB (Fig. 4). The northeast-southwest gradients of the surface PM_{2.5} concentrations in the
185 SCB ~~were~~ mostly resulted from the near-surface northeasterly winds that were blocked by plateaus and mountains located to the southwest of the SCB, which will be further discussed in the following sections.

3. Results and discussion

3.1 Model evaluation

190 ~~We first~~, we validated the WRF-Chem simulation performance. The simulation results ~~are were~~ compared with the meteorological and PM_{2.5} observations in the SCB, including the intensive vertical soundings, for verifying the vertical structures of the simulated boundary layer. The simulated vertical PM_{2.5} distribution in the lower troposphere was evaluated ~~with the~~using ground-based MPL detection at site 15 ~~at in~~ the western SCB (Fig. 1, Table 1).

195 ~~The A~~ reasonable simulation of meteorology is crucial for modeling variations in air pollutants ~~changes~~ (Hanna et al., 2001). The meteorological simulation was validated by comparing with hourly surface meteorological observations of 2 m air temperature (T2), 10 m wind speed (WS10) and relative humidity (RH). The statistical metrics of comparisons between simulated and observed meteorological variables ~~were are~~ given in Table 3, including the mean bias ~~(MB)~~, ~~the~~ mean error, ~~(ME)~~ and ~~the~~ root mean squared error ~~(RMSE)~~. The verification metrics showed a reasonably good model performance
200 with a reference to ~~the~~ previous ~~works studies~~ (Emery et al., 2001; Chang and Hanna, 2004), although

RH was ~~a bit~~slightly underestimated and wind speed was slightly overestimated. ~~Furthermore,~~ ~~the~~ statistical verification of the simulated surface PM_{2.5} concentrations ~~were~~ ~~are~~ shown in Table 4 with the normalized mean bias (~~NMB~~), ~~the~~ normalized mean error (~~NME~~), ~~the~~ mean fractional bias (~~MFB~~) and ~~the~~ mean fractional error (~~MFE~~) in two levels of light PM_{2.5} pollution (75–150 µg m⁻³) and heavy PM_{2.5} pollution (> 150 µg m⁻³). In general, the verification suggested that the WRF-Chem simulations reasonably reproduced the meteorological conditions and the evolution of PM_{2.5} concentrations over ~~the~~ SCB, within the criteria for regulatory applications (Emery et al., 2017).

The vertical structure of the atmospheric boundary layer directly affects the vertical diffusion of atmospheric pollutants. Therefore, we compared the vertical profiles of the model simulation with the intensive sounding observations in terms of ~~the~~ variation range and average profiles during the heavy haze episode. ~~Compared with the observed air temperature, the WRF-Chem simulations were evaluated to reasonably capture the vertical temperature profiles for understanding atmospheric stability in the vertical thermodynamic structures of the boundary layer over the SCB (Fig. S3).~~ The potential temperature, wind speed and ~~relative humidity~~RH of the simulation were ~~also~~ validated for both daytime and nighttime, ~~as shown~~ in Fig. 5. The simulated vertical profiles of ~~the~~ meteorological variables were generally acceptable in the lower troposphere (Fig. 5). It should be ~~pointed out~~noted that the significant underestimation of RH above 1 km, where the observed RH reached nearly 100%, was caused by the clouds due to the abundant moisture at night, ~~that which~~ the model failed to reproduce.

The MPL-4B lidar, located at site 15 (Fig. 1) ~~in on~~ the western edge of ~~the~~ SCB to the east of ~~the~~ TP, continuously detected aerosol extinction ratios in the troposphere. The vertical distribution of ~~the~~ PM_{2.5} mass concentrations ~~were~~ ~~was~~ derived from the extinction ratio (Ansmann et al., 2012; Córdoba-Jabonero et al., 2016). The height-time cross-section of ~~the~~ derived and simulated PM_{2.5} mass concentrations from 7:00 a.m. to 2:00 p.m. on 5 January 2017 ~~were~~ ~~are~~ presented in Fig. 6. It can be seen that a good agreement between the lidar observation and the WRF-Chem simulation was achieved. One of the significant features is that ~~besides in addition to~~ the occurrence of near-surface high PM_{2.5}, which is typical for most heavy haze pollution ~~events~~ over ~~the~~ areas with ~~a~~ relatively flat terrain, a layer of high PM_{2.5} ~~concentrations was~~ developed between 1 and 2 km above ground level (Fig. 6a), leaving a hollow layer between the two heavily polluted layers. The upper high PM_{2.5} layer was built ~~with due~~ ~~to the~~ uplifting and then overturning of the air flows associated with the blocking effect of the TP terrain, which ~~will be~~is addressed in the next section.

3.2 Surface PM_{2.5} concentrations

Figure 7 shows the simulated surface PM_{2.5} concentrations and near-surface wind fields during the formation, maintenance and dissipation periods of 2–8 January 2017. The high PM_{2.5} concentrations were mostly centered in the northwest and southern SCB regions, featuring the Chengdu-Chongqing urban agglomeration (Fig. 1). ~~From the formation to the maintenance and the dissipation periods,~~ The prevailing northeasterly winds strengthened gradually over ~~the SCB~~ from the P1 to P2 and P3 periods (Fig. 7). The high plateaus and mountains, especially YGP and TP to the west of ~~the SCB~~, blocked the upcoming northeasterly winds. The spatial distribution of surface PM_{2.5} concentrations (Fig. 7) clearly reflects the combined effect of the urban anthropogenic air pollutant emissions and the PM_{2.5} accumulations by the flow convergence forced by the TP and the YGP blocking ~~to the~~ prevailing winds. During the formation and maintenance stage, the surface winds were weak (1.4–1.7 m s⁻¹) over ~~the SCB~~, ~~which was and were~~ insufficient to dispel the air pollutants, but ~~led to~~ continuously an accumulation of PM_{2.5}, locally from light to heavy pollution conditions (Fig. 7a, Fig. 7b). ~~In the maintenance period~~ During P2, heavy air pollution blanketed a large area in ~~the SCB~~ with excessive PM_{2.5} concentrations (mostly >150.0 μg m⁻³). ~~By the dissipation period~~ During P3, the northeasterly winds intensified and removed PM_{2.5} from ~~the SCB~~ (Fig. 7c).

3.3 Vertical structures of PM_{2.5} concentrations

The high terrain of ~~the~~ YGP and TP blocked the northeastern airflows over ~~the SCB~~ by lifting the airflow along with air pollutants, altering the vertical PM_{2.5} distribution. Therefore, it was of great interest to analyze the vertical distribution and ~~the~~ transport structures of PM_{2.5} over ~~the SCB~~ and the surrounding regions.

The terrain effect of TP, the “world roof” on the mid-latitude westerlies could modulate haze pollution in the downstream region over China (Xu et al., 2016). The SCB is immediately to the east of the TP, with a large elevation drop exceeding 3000 m over a short horizontal distance. The unique terrain effect generates the asymmetries of in meteorological and air pollutant distributions over the SCB (Zhang et al., 2019). Chengdu (site 1), situated on in the far west side of the SCB, we was selected to better understand the elevated 3D structures of PM_{2.5} with the impact of TP terrain-forcing circulations on the haze pollution episode event over the SCB. Chengdu is a metropolis in the SCB with high anthropogenic pollutant emissions and the has the highest most polluted environment pollution

levels in Southwest China (Ning et al., 2018b). It is important to investigate how the urban surface-high PM_{2.5} levels evolved vertically in the atmosphere over urban surfaces with the combination of between the high urban emissions and TP's terrain-forcing lifting over SCB.

To examine the vertical structures of PM_{2.5} concentrations over SCB, we selected the urban site 1 (104.02° E; 30.67° N) in Chengdu (cf. Fig. 1) as a reference point to investigate the distributions of PM_{2.5} and the atmospheric circulations respectively in the vertical-meridional and vertical-zonal cross-sections, respectively, over the SCB. The evolution of circulation evolutions from a clean environment (Figs. 8a and 9a), and the formation (Figs. 8b and 9b), maintenance (Figs. 8c and 9c) and dissipation periods (Figs. 8d and 9d) of the heavy haze pollution episode were plotted. A remarkable feature in the vertical distributions of PM_{2.5} was the unique hollows over the SCB, between the high surface concentration and a high PM_{2.5} layers at heights of 1.5–3 km. The PM_{2.5} distribution was developed by the interaction of atmospheric circulations in the free troposphere and topographic effects on the air flows in the boundary layer over the SCB (Figs. 8 and 9). Leeward vortices often occur over the SCB owing due to the effect of the large TP topography on the mid-latitude westerlies in the free troposphere (Zhang et al., 2019). The lee vortex with a strong temperature inversion can act as a lid covering air pollutants within the atmospheric boundary layer over the SCB region (Ning et al. 2018a). In the current case, the lee vortex circulation, working together with the basin near-surface flows, drove a 3D PM_{2.5} transport and its temporal changes over the SCB (Figs. 8 and 9).

Comparing the vertical structures of PM_{2.5} and the circulations in different periods, the so-called lid of vortex circulation with the underlying high PM_{2.5} layers in the uphill near-surface airflows was elevated to the free troposphere in the clean environment and the dissipation periods of the heavy air pollution (Figs. 8a, 8d, 9a and 9d), while whereas the lid with a southwesterly wind in vortex circulation was pressed down in the formation and maintenance periods with while confining the strong vertical sub-circulations along the eastern TP upslope to the atmospheric boundary layer (Figs. 8b, 8c, 9b and 9c).

Driven by the near-surface northeasterly winds (Fig. 7), the near-ground airflows with high PM_{2.5} concentrations over the SCB were uplifted respectively over the windward slopes of TP and YGP. And the uphill airflows were restrained and overturned at heights of 1.5–3 km (a.s.l.), forming a vertical sub-circulation over the SCB region, especially the well-structured vertical circulations in the formation and maintenance P1 and P2 stages of the heavy air pollution (Figs. 8–9). Governed by the

vertical sub-circulations, the downward transport from the high PM_{2.5} layers could replenish the surface PM_{2.5} concentrations in the northwest SCB with the addition of near-surface accumulation of air pollutants (Figs. [8b–8c](#), [9b–9c](#)).

The TP and YGP lee vortices over the SCB also modify the vertical thermo-dynamical structures in the atmosphere (Xu et al., 2016), altering the height and intensity of the lid in stable stratification and covering air pollutants (Ning et al., 2018a). The potential temperature vertical gradients (Fig. 5), which are used for assessing atmospheric stability, were estimated with 4.0 K/km, 7.8 K/km and 5.2 K/km in the boundary layer during the three periods of haze pollution with near-surface strong temperature inversion (Fig. S3), presenting a thermodynamic structure with stable stratification in the atmospheric boundary layer, weakening the air pollutant dispersion. From the formation to maintenance periods of heavy air pollution, accompanying the lowering of the stable layer in the free troposphere, the uplifted airflows along the windward slopes of TP and YGP were enhanced, and the high PM_{2.5} layers were restricted at the lower altitudes of 1.5–3.0 km, where the vertical structure of PM_{2.5} over the SCB was characterized with a remarkable hollow sandwiched by a high PM_{2.5} layers at heights of 1.5–3 km and a highly polluted near-surface layer (Figs. [8b–8c](#), [9b–9c](#)). In-During the dissipation period, the aloft high PM_{2.5} concentrations over the SCB were transported to the downwind regions following the airflows in the lower troposphere, as the lid in the southwest region was weakening-weakened and elevated into the free troposphere (Figs. [8d](#) and [9d](#)).

3.4 Distribution of PM_{2.5} in upper high concentration layer

This section examined-describes the characteristics of the upper-layer high PM_{2.5} concentrations. The PM_{2.5} concentrations were averaged between the heights of 1.5–2.5 km, and-as shown in Fig. 10. Comparing-Compared with the surface PM_{2.5} concentrations, the PM_{2.5} concentrations decreased significantly in the lower free troposphere (Figs. 7 and 10), reflecting an-the important role of surface air pollutant emissions in the atmospheric environment over the SCB. During the formation period of heavy air pollution events, the PM_{2.5} particles in the free troposphere were concentrated in the northwestern SCB (Fig. 10a);-. In-During the-maintenance-periodP2,-the high PM_{2.5} centers were developed in the northwestern SCB edge, and PM_{2.5} concentrations increased obviously-in the southwestern and central SCB regions (Fig. 10b), reflecting the strong vertical diffusion of PM_{2.5} in the lower troposphere during the heavy air pollution periods (Figs. [8c](#) and [9c](#)). Driven by strong

320 northeasterly winds ~~in the dissipation period during P3~~ (Fig. 7c), the high PM_{2.5} concentrations in the lower free troposphere were centered in the narrow southwestern and southern SCB areas (Fig. 10c), where ~~the~~ PM_{2.5} from the polluted SCB region ~~were was~~ transported out ~~at from~~ the gap between the eastern TP and northern YGP edge.

3.5 Contribution of local emission and outflow transport

325 Local emissions and regional transport of air pollutants are the two key factors that affecting air quality. The SCB region in the northeastern part of Southwest China, characterized by a deep-bowl structure, is isolated by plateaus (TP in the west and YGP in the south) and mountains with a clean atmospheric environment. Haze pollution events with extremely high PM_{2.5} concentrations over the SCB are ascribed to the accumulation of local anthropogenic pollutants and air pollutant transport over
330 the basin (Wang et al., 2018; Qiao et al., 2019; Zhao et al., 2019). High local anthropogenic emissions in the SCB dominate regional air pollution over the SCB (Liao et al., 2017). The transport of air pollutants from neighboring countries in South Asia is mostly concentrated in the neighboring regions of the southern TP and southern YGP (Wang et al., 2018; Zhao et al., 2019; Yin et al., 2020). Therefore, the anthropogenic emission data of South Asian neighboring countries of China are not included in the
335 WRF-Chem simulation on haze pollution over the SCB during January 2–8 2017, considering the less effects of northward cross-border transport of air pollutants from South Asian neighboring countries on air pollution in SCB with prevailing northeasterly wind during Asian winter monsoon season with a negligible contribution to the wintertime heavy haze pollution over the SCB region.~~considering the lower effects of air pollutant cross-border transport on wintertime air pollution in the SCB with a~~
340 ~~negligible contribution to the wintertime heavy haze pollution over the SCB region.~~ Here, the differences ~~of in the~~ PM_{2.5} concentrations between the numerical experiments, Emi-Real and Emi-Non, were analyzed to assess the contribution of regional air pollutant emissions to surface PM_{2.5} concentrations in the SCB and the impact of PM_{2.5} transport from the SCB to the surrounding plateaus and mountains.

345 Figure 11 shows the PM_{2.5} concentrations originating from local emissions of primary PM_{2.5} and gaseous precursors of PM_{2.5} over the SCB, and the relative contribution rates to air pollution changes. The SCB's regional air pollutant emissions provided surface PM_{2.5} from 40.6 to 136.2 μg m⁻³, contributing 75.4–94.6 % of total concentrations for the heavy pollution episode over the SCB. This

350 indicates the dominant role of local air pollutant emissions on air quality changes over this isolated deep basin in Southwest China. The surface PM_{2.5} concentrations sourced from the regional air pollutant emissions over the SCB were averaged, with 88.64, 91.04, and 65.96 μg m⁻³ for P1, P2, and P3, respectively. However, ~~it was interesting interestingly, the average to point out that the averaged~~ contribution rates of regional air pollutant emissions to surface PM_{2.5} concentrations in the SCB were actually ~~dropped down decreased~~ from 90.7 % ~~in the formation period~~ in P1 to; 85.6 % ~~in the maintenance period~~ in P2 to and 83.3 % ~~in the dissipation period~~ in P3 (Fig. 11). This could be attributed to the exchanges between the PM_{2.5}-rich air mass over the SCB and PM_{2.5}-poor air mass in the surrounding plateaus and mountains over Southwest China (Figs. 8 and 9).

To assess the impact of the PM_{2.5} transport from the SCB on the air quality over the surrounding areas in Southwest China, we calculated the ~~PM_{2.5}-contribution amounts~~ and rates of SCB's regional air pollutant emissions to the PM_{2.5} concentrations in the adjoining regions in the plateaus and mountains, based on the differences ~~of in the~~ PM_{2.5} concentrations between ~~Emi-real~~ Real and Emi-Non (Table 5). The near-surface prevailing northeasterly winds of the SCB brought PM_{2.5} from the SCB to the eastern TP edge, ~~the northern YGP edge,~~ and ~~the DBM region~~ (Fig. 7), resulting in ~~importing an increase in the~~ concentrations of surface PM_{2.5} during the heavy haze pollution event, with averaged averages of ~~respectively with~~ 18.0, 31.3, and 10.4 μg m⁻³ ~~during the heavy haze pollution~~ respectively (Table 5). The TP and YGP, as the cleaner regions in China (Song et al., 2017; Zhan et al., 2018), ~~were received significant remarkably polluted~~ pollution by due to the PM_{2.5} transport from the SCB. During the dissipation period of the heavy air pollution episode, the eastern TP edge and northern YGP regions gained peak imports of PM_{2.5} at 22.9 and 41.9 μg m⁻³ (Table 5). Thus, in this case, the downwind adjoining TP and YGP regions ~~is were~~ the main receptor area of the SCB emissions.

375 Finally, the PM_{2.5} contribution rates, i.e., the percentages ~~of between~~ the PM_{2.5} concentrations transported from the basin to those in the adjacent regions of plateaus and mountains were calculated for different periods of the heavy PM_{2.5} pollution event over the SCB. The surface PM_{2.5} in the eastern TP edge ~~were~~ mostly originated from the source region of the SCB, with ~~the~~ dominant contribution rates ~~respectively~~ of 63.6, 67.4 and 72.7 % in the formation, maintenance, and dispersion periods, respectively. The PM_{2.5} import from the SCB pollutant emissions also contributed to the majority of surface PM_{2.5} concentrations in the northern YGP, with contribution rates of 58.3, 52.8 and 70.5 % during the three different periods, with an overall contribution rate of 58.5 % (averaged for the whole

~~entire~~ SCB heavy air pollution ~~event~~). In contrast, the DBM region was less influenced by the SCB's
380 emission sources with a contribution rate of 31.0 % (~~averaged during the heavy air pollution event~~).

4. Conclusions

By using the multiple ground observations, meteorological sounding data and ~~Micro-Pulse-Lidar~~MPL retrievals as well as conducting modeling experiments with the WRF-Chem model, this study investigated the three-dimensional structures and the development mechanisms of the PM_{2.5} for a
385 wintertime heavy haze pollution ~~episode-event~~ over ~~the~~ SCB, an isolated deep basin in Southwest China. The roles of the basin pollutant emissions and the unique basin circulations were evaluated for their contributions to the 3D distribution of PM_{2.5} over ~~the~~ SCB and to the neighboring YGP, TP, and DBM regions.

The vertical structure of ~~the~~ PM_{2.5} in the lower troposphere over ~~the~~ SCB was characterized ~~with~~
390 ~~by~~ unique hollows located between a high PM_{2.5} layer at ~~a~~ heights of 1.5–3 km and ~~the-a~~ high PM_{2.5} surface layer. ~~It is found that t~~The hollow was developed by the interaction of the upper-level free tropospheric circulations and ~~the~~ lower-level topographic boundary layer. The southwesterlies ~~passed-passing~~ over the TP and YGP resulted in a lee vortex over ~~the~~ SCB, which helped form and maintain ~~heavy-high~~ PM_{2.5} ~~pollutionconcentrations~~, with well-developed vertical secondary circulations along
395 the eastern TP upslope, ~~while-whereas~~ the southwesterlies with the underlying high PM_{2.5} layers were elevated in the dissipation of heavy PM_{2.5} pollution over the SCB.

Due to ~~the-effect-of~~ the joint impact of the urban anthropogenic air pollutant emissions and the large terrain blocking flow at the eastern TP slope and YGP, high surface PM_{2.5} concentrations were mostly distributed in the northwest~~ern~~ and southern SCB regions. The tropospheric circulations, ~~which~~
400 ~~altered with-altering~~ the vertical diffusion of PM_{2.5} exerted a strong impact on PM_{2.5} distribution in the lower free troposphere. The high PM_{2.5} centers in the lower free troposphere were distributed over the northwestern and southwestern SCB edges, as well as the central SCB regions. Driven by strong northeasterly winds in the dissipation period, ~~the~~ PM_{2.5} in the lower free troposphere ~~were~~ converged to the west boundary of ~~the~~ SCB and then transported to the eastern TP edge and the northern YGP edge
405 areas.

The regional emissions of air pollutants in ~~the~~ SCB played a dominant role in the formation of ~~the~~ heavy air pollution, contributing 75.4–94.6 % to surface PM_{2.5} concentrations over the basin for the

heavy pollution ~~episode-event~~ studied herein. Furthermore, the surface PM_{2.5} concentrations in the eastern TP were mostly transported from the SCB's emission sources, with contribution rates of 63.6, 410 67.4 and 72.7 % for ~~the formation, maintenance and dispersion periods~~P1, P2, and P3, respectively. Similarly, the SCB also contributed the majority of surface PM_{2.5} concentrations in the adjacent northern YGP, with an average contribution rate of 58.5 % for the whole SCB pollution period and a very high contribution of 70.5 % during the dissipation period. Therefore, the SCB region is the major source of air pollutants ~~source~~ for the downwind receptor areas over the adjoining TP and YGP regions and affects the atmospheric environmental changes in Southwest China. 415

This ~~work-study highlight~~exposed the unique and important three-dimensional structures of PM_{2.5} and investigated their formation mechanisms and downwind outflow transport over the SCB. The deep basin terrain along with the TP and YGP forcing effect creates very complex PM_{2.5} pollution conditions over the SCB region, which is ~~dramatically-significantly~~ different from those over relatively flat 420 regions. To generalize our findings, further work with more case studies and regional climatic analyses with long-term observation data and numerical modeling with data assimilation and refined physical and chemical schemes are ~~desired~~required. MEIC 2017 was not available for the WRF-Chem model. The SCB is located in Southwest China with larger uncertainties in the anthropogenic emission inventory compared to Eastern China. An accurate emission inventory could improve air pollution simulations and air quality change assessments in future studies. Furthermore, as pointed out in this 425 study, the PM_{2.5} emission sources in the SCB greatly influence the regional environmental changes ~~over~~ in Southwest China. Thus, the regional transport modeling of air pollutants with careful consideration of the thermal and dynamic forcing of the underlying complex plateaus terrain should be further investigated.

430

Data availability. Data used in this paper can be provided by Zhuozhi Shu (shuzhuozhi@foxmail.com) upon request.

Author contributions. ZS and TZ conducted the study design. ZS, TZ, JX, CW and LC conducted the 435 vertical observational experiment. ZS wrote the manuscript with the help of TZ and YL. LZ and YZ assisted with data processing. HL and LS were involved in the scientific interpretation and discussion.

LL and YL provided the surface meteorological data. All of the authors provided commentary on the paper.

440 *Competing interests.* The authors declare that they have no conflicts of interest.

Acknowledgements. This research was supported by the Second Tibetan Plateau Scientific Expedition and Research (STEP) program (2019QZKK0105), the National Natural Science Foundation of China (91744209, 91644223 and 91544109), the National Key R & D Program Pilot Projects of China
445 (2016YFC0203304) and Graduate Research and Innovation Projects of Jiangsu Province (SJKE19_0941).

References

Ansmann, A., Seifert, P., Tesche, M., and Wandinger, U.: Profiling of fine and coarse particle mass: case studies of Saharan dust and Eyjafjallajökull/Grimsvötn volcanic plumes, *Atmos. Chem. Phys.*, 12,
450 9399–9415, <https://doi.org/10.5194/acp-12-9399-2012>, 2012.

Cai, W., Li, K., Liao, H., Wang, H. and Wu, L.: Weather conditions conducive to Beijing severe haze more frequent under climate change, *Nat. Clim. Change*, 7, 257–262, <https://doi.org/10.1038/nclimate3249>, 2017.

Chang, J. C. and Hanna, S. R.: Air quality model performance evaluation, *Meteorol. Atmos. Phys.*, 87,
455 167–196, <https://doi.org/10.1007/s00703-003-0070-7>, 2004.

Chen, Y. and Xie, S.: Temporal and spatial visibility trends in the Sichuan Basin, China, 1973 to 2010, *Atmos. Res.*, 112, 25–34, <https://doi.org/10.1016/j.atmosres.2012.04.009>, 2012.

Chen, Y., Xie, S., Luo, B. and Zhai, C.: Characteristics and origins of carbonaceous aerosol in the Sichuan Basin, China, *Atmos. Environ.*, 94, 215–223, <https://doi.org/10.1016/j.atmosenv.2014.05.037>,
460 2014.

Córdoba-Jabonero, C., Andrey-Andrés, J., Gómez, L., Adame, J.A., Sorribas, M., Navarro-Comas, M., Puentedura, O., Cuevas, E., and Gil-Ojeda, M.: Vertical mass impact and features of Saharan dust intrusions derived from ground-based remote sensing in synergy with airborne in-situ measurements, *Atmos. Environ.*, 142, 420–429, <https://doi.org/10.1016/j.atmosenv.2016.08.003>, 2016.

- 465 Dawson, J. P., Adams, P. J., and Pandis, S. N.: Sensitivity of PM_{2.5} to climate in the Eastern US: a modeling case study, *Atmos. Chem. Phys.*, 7, 4295–4309, <https://doi.org/10.5194/acp-7-4295-2007>, 2007.
- Emery, C. A., Tai, E., and Yarwood, G.: Enhanced meteorological modeling and performance evaluation for two Texas ozone episodes, Project Report prepared for the Texas Natural Resource
- 470 Conservation Commissions, ENVIRON International Corporation, Novato, CA, 2001.
- Emery, C., Liu, Z., Russell, A. G., Odman, M. T., Yarwood, G. and Kumar, N.: Recommendations on statistics and benchmarks to assess photochemical model performance, *J. Air. Waste. Manage.*, 67, 582–598, <https://doi.org/10.1080/10962247.2016.1265027>, 2017.
- Gu, Y. and Yim, S. H.: The air quality and health impacts of domestic trans-boundary pollution in
- 475 various regions of China, *Environ. Int.*, 97, 117–124, <https://doi.org/10.1016/j.envint.2016.08.004>, 2016.
- Guo, S., Hu, M., Zamora, M. L., Peng, J., Shang, D., Zheng, J., Du, Z., Wu, Z., Shao, M., Zeng, L. Molina, M. J. and Zhang, R.: Elucidating severe urban haze formation in China, *P. Natl. Acad. Sci. USA.*, 111, 17373–17378, <https://doi.org/10.1073/pnas.1419604111>, 2014.
- 480 Guo, Y., Zeng, H., Zheng, R., Li, S., Barnett, A. G., Zhang, S., Zou, X., Huxley, R., Chen, W. and Williams, G.: The association between lung cancer incidence and ambient air pollution in China: a spatiotemporal analysis, *Environ. Res.*, 144, 60–65, <https://doi.org/10.1016/j.envres.2015.11.004>, 2016.
- Hanna, S. R. and Yang, R.: Evaluations of mesoscale models' simulations of near-surface winds, temperature gradients, and mixing depths, *J. Appl. Meteorol.*, 40, 1095–1104,
- 485 [https://doi.org/10.1175/1520-0450\(2001\)040<1095:EOMMSO>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<1095:EOMMSO>2.0.CO;2), 2001.
- Huang, X., Liu, Z., Liu, J., Hu, B., Wen, T., Tang, G., Zhang, J., Wu, F., Ji, D., Wang, L., and Wang, Y.: Chemical characterization and source identification of PM_{2.5} at multiple sites in the Beijing–Tianjin–Hebei region, China, *Atmos. Chem. Phys.*, 17, 12941–12962, <https://doi.org/10.5194/acp-17-12941-2017>, 2017.
- 490 Hu, X., Ma, Z., Lin, W., Zhang, H., Hu, J., Wang, Y., Xu, X., Fuentes, J. D. and Xue, M.: Impact of the Loess Plateau on the atmospheric boundary layer structure and air quality in the North China Plain: A case study. *Sci. Total. Environ.*, 499, 228–237, <https://doi.org/10.1016/j.scitotenv.2014.08.053>, 2014.
- Langrish, J. P., Li, X., Wang, S., Lee, M. M. Y., Barnes, G. D., Miller, M. R., Cassee, F. R., Boon, N. A., Donaldson, K., Li, J., Li, L., Mills, N. L., Newby, D. E. and Jiang, L.: Reducing Personal Exposure to

- 495 Particular Air Pollution Improves Cardiovascular Health in Patients with Coronary Heart Disease, *Environ. Health Persp.*, 120, 367–372, <https://doi.org/10.1289/ehp.1103898>, 2012.
- Li, B., Zhang, J., Zhao, Y., Yuan, S., Zhao, Q., Shen, G. and Wu, H.: Seasonal variation of urban carbonaceous aerosols in a typical city Nanjing in Yangtze River Delta, China, *Atmos. Environ.*, 106, 223–231, <https://doi.org/10.1016/j.atmosenv.2015.01.064>, 2015.
- 500 Li, P., Xin, J., Wang, Y., Li, G., Pan, X., Wang, S., Cheng, M., Wen, T., Wang, Guang. And Liu, Z.: Association between particulate matter and its chemical constituents of urban air pollution and daily mortality or morbidity in Beijing city, *Environ. Sci. Pollut. R.*, 22, 358–368, <https://doi.org/10.1007/s11356-014-3301-1>, 2015.
- Lin, Y., Zou, J., Yang, W. and Li, C.: A review of recent advances in research on PM_{2.5} in China, *Int. J. Environ. Res. Public Health*, 15, 438, <https://doi.org/10.3390/ijerph15030438>, 2018.
- 505 Liao, T., Wang, S., Ai, J., Gui, K., Duan, B., Zhao, Q., Zhang, X., Jiang, W. and Sun, Y.: Heavy pollution episodes, transport pathways and potential sources of PM_{2.5}, during the winter of 2013 in Chengdu (China), *Sci. Total. Environ.*, 584, 1056–1065, <https://doi.org/10.1016/j.scitotenv.2017.01.160>, 2017.
- 510 Liao, T., Gui, K., Jiang, W., Wang, S., Wang, B., Zeng, Z., Che, H., Wang, Y. and Sun, Y.: Air stagnation and its impact on air quality during winter in Sichuan and Chongqing, southwestern China, *Sci. Total. Environ.*, 635, 576–585, <https://doi.org/10.1016/j.scitotenv.2018.04.122>, 2018.
- Megaritis, A. G., Fountoukis, C., Charalampidis, P. E., Denier van der Gon, H. A. C., Pilinis, C., and Pandis, S. N.: Linking climate and air quality over Europe: effects of meteorology on PM_{2.5} concentrations, *Atmos. Chem. Phys.*, 14, 10283–10298, <https://doi.org/10.5194/acp-14-10283-2014>, 2014.
- 515 Ming, L., Jin, L., Li, J., Fu, P., Yang, W., Liu, D., Zhang, G., Wang, Z. and Li, X.: PM_{2.5} in the Yangtze River Delta, China: Chemical compositions, seasonal variations, and regional pollution events, *Environ. Pollut.*, 223, 200–212, <https://doi.org/10.1016/j.envpol.2017.01.013>, 2017.
- 520 Ning, G., Wang, S., Yim, S. H. L., Li, J., Hu, Y., Shang, Z., Wang, J., and Wang, J.: Impact of low-pressure systems on winter heavy air pollution in the northwest Sichuan Basin, China, *Atmos. Chem. Phys.*, 18, 13601–13615, <https://doi.org/10.5194/acp-18-13601-2018>, 2018a.
- Ning, G., Wang, S., Ma, M., Ni, C., Shang, Z., Wang, J. and Li, J.: Characteristics of air pollution in different zones of Sichuan Basin, China, *Sci. Total. Environ.*, 612, 975–984,

- 525 <https://doi.org/10.1016/j.scitotenv.2017.08.205>, 2018b.
- Qiao, X., Guo, H., Tang, Y., Wang, P., Deng, W., Zhao, X., Hu, J., Ying, Q., and Zhang, H.: Local and regional contributions to fine particulate matter in the 18 cities of Sichuan Basin, southwestern China, *Atmos. Chem. Phys.*, 19, 5791–5803, <https://doi.org/10.5194/acp-19-5791-2019>, 2019.
- Stockwell, W. R., Middleton, P., Chang, J. S. and Tang, X.: The second generation regional acid
530 deposition model chemical mechanism for regional air quality modeling, *J. Geophys. Res. Atmos.*, 95, 16343–16367, <https://doi.org/10.1029/JD095iD10p16343>, 1990.
- [Ren, Y., Zhang, H., Wei, W., Wu, B., Liu, J., Cai, X., and Song, Y.: Comparison of the turbulence structure during light and heavy haze pollution episodes, *Atmospheric Research*, 230, 0169-8095, <https://doi.org/10.1016/j.atmosres.2019.104645>, 2019.](https://doi.org/10.1016/j.atmosres.2019.104645)
- 535 Song, C., Wu, L., Xie, Y., He, J., Chen, X., Wang, T., Lin, Y., Jin, T., Wang, A., Liu, Y., Dai, Q., Liu, B., Wang, Y. and Mao, H.: Air pollution in China: status and spatiotemporal variations, *Environ. Pollut.*, 227, 334–347, <https://doi.org/10.1016/j.envpol.2017.04.075>, 2017.
- Tang, L., Yu, H., Ding, A., Zhang, Y., Qin, W., Zhuang, W., Chen, W., Hua, Y. and Yang, X.: Regional contribution to PM₁ pollution during winter haze in Yangtze River Delta, China, *Sci. Total. Environ.*,
540 541, 161–166, <https://doi.org/10.1016/j.scitotenv.2015.05.058>, 2015.
- Tian, P., Cao, X., Zhang, L., Sun, N., Sun, L., Logan, T., Shi, J., Wang, Y., Ji, Y., Lin, Y., Huang, Z., Zhou, T., Shi, Y., and Zhang, R.: Aerosol vertical distribution and optical properties over China from long-term satellite and ground-based remote sensing, *Atmos. Chem. Phys.*, 17, 2509–2523, <https://doi.org/10.5194/acp-17-2509-2017>, 2017.
- 545 Tian, M., Liu, Y., Yang, F., Zhang, L., Peng, C., Chen, Y., Shi, G., Wang, H., Luo, B., Jiang, C., Li, B., Takeda, N. and Koizumi, K.: Increasing importance of nitrate formation for heavy aerosol pollution in two megacities in Sichuan Basin, southwest China, *Environ. Pollut.*, 250, 898–905, <https://doi.org/10.1016/j.envpol.2019.04.098>, 2019.
- Tao, J., Zhang, L., Engling, G., Zhang, R., Yang, Y., Cao, J., Zhu, C., Wang, Q. and Luo L.: Chemical
550 composition of PM_{2.5} in an urban environment in Chengdu, China: Importance of springtime dust storms and biomass burning, *Atmos. Res.*, 122, 270–283, <https://doi.org/10.1016/j.atmosres.2012.11.004>, 2013.
- Tao, M., Chen, L., Su, L. and Tao, J.: Satellite observation of regional haze pollution over the North China Plain, *J. Geophys. Res. Atmos.*, 117, D12203, <https://doi.org/10.1029/2012JD017915>, 2012.

- 555 Tuccella, P., Curci, G., Visconti, G., Bessagnet, B., and Menut L., and Park, R. J.: Modeling of gas and aerosol with WRF-Chem over Europe: Evaluation and sensitivity study, *J. Geophys. Res. Atmos.*, 117, D03303, <https://doi.org/10.1029/2011JD016302>, 2012.
- ~~Wang, H., Peng, Y., Zhang, X., Liu, H., Zhang, M., Che, H., Cheng, Y., and Zheng, Y.: Contributions to the explosive growth of PM_{2.5} mass due to aerosol-radiation feedback and decrease in turbulent~~
- 560 ~~diffusion during a red alert heavy haze in Beijing-Tianjin-Hebei, China, *Atmos. Chem. Phys.*, 18, 17717–17733, <https://doi.org/10.5194/acp-18-17717-2018>, 2018.~~
- Wang, T., Jiang, F., Deng, J., Shen, Y., Fu, Q., Wang, Q., Fu, Y., Xu, J. and Zhang, D.: Urban air quality and regional haze weather forecast for Yangtze River Delta region, *Atmos. Environ.*, 58, 70–83, <https://doi.org/10.1016/j.atmosenv.2012.01.014>, 2012.
- 565 Wang, Q. and Tan, Z.: Multi-scale topographic control of southwest vortex formation in Tibetan Plateau region in an idealized simulation, *J. Geophys. Res. Atmos.*, 119, 543–561, <https://doi.org/10.1002/2014JD021898>, 2014.
- Wang, X., Dickinson, R. E., Su, L., Zhou, C. and Wang K.: PM_{2.5} pollution in China and how it has been exacerbated by terrain and meteorological conditions, *Bull. Amer. Meteor. Soc.*, 99, 105–119,
- 570 <https://doi.org/10.1175/BAMS-D-16-0301.1>, 2018.
- Wang, H., Tian, M., Chen, Y., Shi, G., Liu, Y., Yang, F., Zhang, L., Deng, L., Yu, J., Peng, C., and Cao, X.: Seasonal characteristics, formation mechanisms and source origins of PM_{2.5} in two megacities in Sichuan Basin, China, *Atmos. Chem. Phys.*, 18, 865–881, <https://doi.org/10.5194/acp-18-865-2018>, 2018.
- 575 ~~Wang, H., Peng, Y., Zhang, X., Liu, H., Zhang, M., Che, H., Cheng, Y., and Zheng, Y.: Contributions to the explosive growth of PM_{2.5} mass due to aerosol-radiation feedback and decrease in turbulent diffusion during a red alert heavy haze in Beijing-Tianjin-Hebei, China, *Atmos. Chem. Phys.*, 18, 17717–17733, <https://doi.org/10.5194/acp-18-17717-2018>, 2018.~~
- ~~Wang, L., Liu, J., Gao, Z., Li, Y., Huang, M., Fan, S., Zhang, X., Yang, Y., Miao, S., Zou, H., Sun, Y.,~~
- 580 ~~Chen, Y., and Yang, T.: Vertical observations of the atmospheric boundary layer structure over Beijing urban area during air pollution episodes, *Atmos. Chem. Phys.*, 19, 6949–6967, <https://doi.org/10.5194/acp-19-6949-2019>, 2019.~~
- Wu, M., Wu, D., Fan, Q., Wang, B. M., Li, H. W., and Fan, S. J.: Observational studies of the meteorological characteristics associated with poor air quality over the Pearl River Delta in China,

- 585 Atmos. Chem. Phys., 13, 10755–10766, <https://doi.org/10.5194/acp-13-10755-2013>, 2013.
- Xu, R., Tang, G., Wang, Y. and Tie, X.: Analysis of a long-term measurement of air pollutants (2007–2011) in North China Plain (NCP); Impact of emission reduction during the Beijing Olympic Games, Chemosphere, 159, 647–658, <https://doi.org/10.1016/j.chemosphere.2016.06.025>, 2016.
- Xu, X., Wang, Y., Zhao, T., Cheng, X., Meng, Y. and Ding, G.: "Harbor" effect of large topography on
590 haze distribution in eastern China and its climate modulation on decadal variations in haze, Chinese Sci. Bull., 60, 1132–1143, <https://doi.org/10.1360/N972014-00101>, 2015.
- Xu, X., Zhao, T., Liu, F., Gong, S. L., Kristovich, D., Lu, C., Guo, Y., Cheng, X., Wang, Y., and Ding, G.: Climate modulation of the Tibetan Plateau on haze in China, Atmos. Chem. Phys., 16, 1365–1375, <https://doi.org/10.5194/acp-16-1365-2016>, 2016.
- 595 Ye, X., Song, Y., Cai, X. and Zhang, H.: Study on the synoptic flow patterns and boundary layer process of the severe haze events over the North China Plain in January 2013, Atmos. Environ., 124, 129–145, <https://doi.org/10.1016/j.atmosenv.2015.06.011>, 2016.
- Yim, S. H. L., Fung, J. C. H. and Ng, E. Y. Y.: An assessment indicator for air ventilation and pollutant dispersion potential in an urban canopy with complex natural terrain and significant wind variations,
600 Atmos. Environ., 94, 297–306, <https://doi.org/10.1016/j.atmosenv.2014.05.044>, 2014.
- [Yin, D., Zhao, S., Qu, J., Yu, Y., Kang, S., Ren, X., Zhang, J., Zou, Y., Dong, L., Li, J., He, J., Li, P., and Qin, D.: The vertical profiles of carbonaceous aerosols and key influencing factors during wintertime over western Sichuan Basin, China, Atmospheric Environment, 223, 1352-2310, https://doi.org/10.1016/j.atmosenv.2020.117269, 2020.](https://doi.org/10.1016/j.atmosenv.2020.117269)
- 605 Zhang, Y., Shao, M., Lin, Y., Luan, S., Mao, Ning., Chen, W. and Wang, M.: Emission inventory of carbonaceous pollutants from biomass burning in the Pearl River Delta Region, China, Atmos. Environ., 76, 189–199, <https://doi.org/10.1016/j.atmosenv.2012.05.055>, 2013.
- Zhang, Q., Quan, J., Tie, X., Li, X., Liu, Q., Gao, Y. and Zhao, D.: Effects of meteorology and secondary particle formation on visibility during heavy haze events in Beijing, China, Sci. Total.
610 Environ., 502, 578–584, <https://doi.org/10.1016/j.scitotenv.2014.09.079>, 2015.
- Zhang, Z., Zhang, X., Gong, D., Kim, S.-J., Mao, R., and Zhao, X.: Possible influence of atmospheric circulations on winter haze pollution in the Beijing–Tianjin–Hebei region, northern China, Atmos. Chem. Phys., 16, 561–571, <https://doi.org/10.5194/acp-16-561-2016>, 2016.
- Zhang, L., Guo, X., Zhao, T., Gong, S., Xu, X., Li, Y., Luo, L., Gui, K., Wang, H., Zheng, Y. and Yin,

- 615 X.: A modelling study of the terrain effects on haze pollution in the Sichuan Basin, *Atmos. Environ.*, 196, 77–85, <https://doi.org/10.1016/j.atmosenv.2018.10.007>, 2019.
- Zhao, S., Yu, Y., Yin, D., Qin, D., He, J. and Dong, L.: Spatial patterns and temporal variations of six criteria air pollutants during 2015 to 2017 in the city clusters of Sichuan Basin, China, *Sci. Total Environ.*, 624, 540–557, <https://doi.org/10.1016/j.scitotenv.2017.12.172>, 2018.
- 620 Zhao, S., Yu, Y., Qin, D., Yin, D., Dong, L. and He, J.: Analyses of regional pollution and transportation of PM_{2.5} and ozone in the city clusters of Sichuan Basin, China, *Atmos. Pollut. Res.*, 10, 374–385, <https://doi.org/10.1016/j.apr.2018.08.014>, 2019.
- Zhan, D., Kwan, M. P., Zhang, W., Yu, X., Meng, B. and Liu, Q.: The driving factors of air quality index in China, *J. Clean Prod.*, 197, 1342–1351, <https://doi.org/10.1016/j.jclepro.2018.06.108>, 2018.
- 625 [Zhong, J., Zhang, X., Dong, Y., Wang, Y., Liu, C., Wang, J., Zhang, Y., and Che, H.: Feedback effects of boundary-layer meteorological factors on cumulative explosive growth of PM_{2.5} during winter heavy pollution episodes in Beijing from 2013 to 2016, *Atmos. Chem. Phys.*, 18, 247–258, <https://doi.org/10.5194/acp-18-247-2018>, 2018.](https://doi.org/10.5194/acp-18-247-2018)
- Zhou, Y., Luo, B., Li, J., Hao, Y., Yang, W., Shi, F., Chen, Y., Simayi, M. and Xie, S.: Characteristics of
630 six criteria air pollutants before, during, and after a severe air pollution episode caused by biomass burning in the southern Sichuan Basin, China, *Atmos. Environ.*, 215, 116840, <https://doi.org/10.1016/j.atmosenv.2019.116840>, 2019.

Table 1. Names of 18 observation sites with the corresponding site number (Fig. 1b) in [the SCB](#).

Number	1	2	3	4	5	6
Names	Chengdu	Chongqing	Deyang	Guang'an	Leshan	Meishan
Number	7	8	9	10	11	12
Names	Mianyang	Nanchong	Neijiang	Suining	Yibin	Ziyang
Number	13	14	15	16	17	18
Names	Zigong	Luzhou	Ya'an	Bazhong	Dazhou	Guangyuan

Table 2. Setting of physical schemes in the WRF-Chem simulations

Microphysics	Morrison 2-mom
Boundary layer	MYJ
Longwave radiation	RRTMG
Shortwave radiation	RRTM
Land surface	Noah

Table 3. ~~The~~ Statistical metrics of comparisons between simulated and observed meteorological elements during 2–8 January 2017.

	Obs.	Sim.	R	MB	ME	RMSE
T2	9.9	9.2	0.78**	-0.7	1.7	2.1
RH	85.1	77.7	0.67**	-7.4	11.2	13.4
WS10	1.2	1.5	0.41*	0.3	0.8	1.1

Note: MB, ME, RMSE were calculated as following: $MB = \frac{1}{N} \sum_{i=1}^N (M_i - O_i)$; $ME = \frac{1}{N} \sum_{i=1}^N |M_i - O_i|$;

$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (M_i - O_i)^2}$; (M and O represented the results from simulation and observation). The

** and * respectively indicated passing the 99% and 95% significant test.

655

Table 4. Statistical metrics of comparisons between simulated and observed surface PM_{2.5} concentrations in ~~the~~ two levels of light and heavy PM_{2.5} pollution during 2–8 January 2017.


	NMB (%)	NME (%)	MFB (%)	MFE (%)
Light pollution	-4.3	25.4	-7.7	30.0
Heavy pollution	-13.5	33.4	-16.3	37.4

Note: NMB, NME, MFB and MFE were calculated as following: $NMB = \frac{\sum_{i=1}^N (M_i - O_i)}{\sum_{i=1}^N O_i} \cdot 100\%$; $NME =$

660 $\frac{\sum_{i=1}^N |M_i - O_i|}{\sum_{i=1}^N O_i} \cdot 100\%$; $MFB = \frac{1}{N} (2 \cdot \frac{M_i - O_i}{M_i + O_i}) \cdot 100\%$; $MFE = \sum_{i=1}^N \left[2 \cdot \frac{M_i - O_i}{M_i + O_i} \right] \cdot 100\%$.

Table 5. ~~The a~~Amounts and contribution rates of PM_{2.5} trans-boundary transport from the SCB to surface PM_{2.5} concentrations averaged over the eastern TP edge (ETP), northern YGP edge (YGP) and ~~the~~ DBM region during the formation (P1), maintenance (P2) and dissipation (P3) periods of the heavy haze pollution events over the SCB region.

	Region	P1	P2	P3	Averages
Transport amount ($\mu\text{g m}^{-3}$)	ETP	15.4	18.8	22.5	18.0
	YGP	30.1	27.5	41.9	31.3
	DBM	8.6	13.5	8.4	10.4
Contribution rates (%)	ETP	63.6	67.4	72.7	66.6
	YGP	58.3	52.8	70.5	58.5
	DBM	26.7	36.6	30.1	31.0



675 **Figure 1.** (Left panel) three nesting domains D1, D2 and D3 of WRF-Chem simulation with the terrain heights (m in a.s.l.) and (right panel) the location of 18 urban observation sites (black dots, Table 1) including site 1 (Chengdu) with the intensive sounding observations and site 15 (Ya'an) with the ground-based MPL detection in the SCB with the surrounding Tibetan Plateau (TP), Yunnan-Guizhou Plateau (YGP), Mountains Daba (Mt. Daba) and Wu (Mt. Wu) in Southwest China. The red dash lines

680 indicate the location of the cross sections respectively along 30.67° N and 104.02° E.




Figure 2. (a) Surface $PM_{2.5}$ concentrations over 18 urban sites averaged during the heavy haze pollution over 2–8 January 2017, (b) hourly variations of $PM_{2.5}$ concentrations observed at the three heaviest polluted cities Chengdu (CD), Deyang (DY), and Meishan (MS) (Fig. 1; Table 1) over 1–10 January 2017. The P1, P2 and P3 indicated respectively the formation, maintenance and dissipation periods of heavy haze pollution with the light pollution level of $75 \mu\text{g m}^{-3}$ (black dashed line) and heavy pollution level of $150 \mu\text{g m}^{-3}$ (red dashed line).

690




Figure 3. 700 hPa geopotential height fields and wind vectors averaged during (a) P1, (b) P2 and (c) P3 stages with the trough line (brown line) and low-pressure center (L). The SCB was outlined with the red solid lines.

695




Figure 4. The spatial distributions of observed surface $PM_{2.5}$ concentrations in the SCB averaged in the formation, maintenance and dissipation periods, P1, P2 and P3 (Fig. 2b), respectively, of the heavy

700 haze pollution episode over 2–8 January 2017.





Figure 5. Comparisons of observed and simulated vertical profiles of potential temperature, wind speed, and relative humidity in the daytime (a, b, c) and nighttime (d, e, f) at Chengdu (site 1 in Fig. 1) during 2–8 January 2017. The red and gray shaded areas represented the variation range of simulation and observation in all vertical profiles with averaged values (lines), respectively.



710 | **Figure 6.** Vertical and time cross-sections of PM_{2.5} mass concentrations ($\mu\text{g m}^{-3}$) from (a) MPL-4B retrievals products and (b) simulation results at site 15 (Fig. 1; Table 1) ~~in-on~~ the western SCB edge during 7:00 a.m.–2:00 p.m. on 5 January 2017.





Figure 7. Horizontal distribution of surface PM_{2.5} concentrations ($\mu\text{g m}^{-3}$; color contours) and wind vectors at 10 m averaged in the periods (a) P1, (b) P2 and (c) P3, respectively. The SCB was outlined with an altitude contour line of 750 m (a.s.l.; black lines).



720 **Figure 8.** Height-longitude cross-sections of $\text{PM}_{2.5}$ concentrations (color contours: $\mu\text{g m}^{-3}$) and wind
 vectors along 30.67° N in the (a) clean environment at 12:00 a.m. on 2 January 2017 (b) heavy air
 pollution formation stage at 12:00 a.m. on 3 January 2017 (c) maintenance stage at 8:00 a.m. on 6
 January 2017, and (d) dissipation stage at 8:00 a.m. on 7 January 2017. The brown arrows highlighted
 the major air flows (red arrows) associated with the terrain of TP, SCB and Mt. Wu (filled brown
 725 color).




Figure 9. Same as Fig. 7, but for height-latitude cross-sections of PM_{2.5} concentrations and wind vectors.

730





Figure 10. Horizontal distribution of $PM_{2.5}$ concentrations (color contours: $\mu g\ m^{-3}$) averaged between 1.5 and 2.5 km heights (in the lower free troposphere) for (a) formation, (b) maintenance and (c) dissipation periods of heavy haze pollution episode-event over the SCB. The SCB was outlined with an altitude contours of 750 m in a.s.l. (dark black lines).



740 **Figure 11.** Hourly variations of surface PM_{2.5} concentrations originating from the SCB's anthropogenic emissions (blue filled areas) and the contribution proportions to the basin surface PM_{2.5} levels (red curve) during 1–8 January 2017 based on the regional averages over the SCB.