

Responses of Arctic Black Carbon and Surface Temperature to Multi-Region Emission Reductions: an HTAP2 ensemble modeling study

Na Zhao¹, Xinyi Dong², Kan Huang^{1*}, Joshua S. Fu^{3,4*}, Marianne Tronstad Lund⁵, Kengo Sudo⁶, Daven Henze⁷, Tom Kucsera⁸, Yun Fat Lam⁹, Mian Chin¹⁰, Simone Tilmes¹¹

¹ Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention (LAP3), Department of Environmental Science and Engineering, Fudan University, Shanghai, China

² School of Atmospheric Science, Nanjing University, Nanjing, China

³ Department of Civil and Environmental Engineering, The University of Tennessee, Knoxville, Tennessee, USA.

⁴ Computational Earth Science Group, Computational Sciences and Engineering Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

⁵ CICERO Center for International Climate and Environmental Research, Oslo, Norway

⁶ Nagoya University, Furo-cho, Chigusa-ku, Nagoya, Japan

⁷ Department of Mechanical Engineering, University of Colorado, Boulder, CO, USA

⁸ Universities Space Research Association, Greenbelt, MD, USA

⁹ Department of Geography, The University of Hong Kong, HKSAR, China

¹⁰ Earth Sciences Division, NASA Goddard Space Flight Center, Greenbelt, MD, USA

¹¹ Atmospheric Chemistry Observations and Modeling Laboratory, National Center for Atmospheric Research, Boulder, Colorado, USA

Correspondence: huangkan@fudan.edu.cn; jifu@utk.edu

ABSTRACT

Black carbon (BC) emissions play an important role in regional climate change of the Arctic. It is necessary to pay attention to the impact of long-range transport from regions outside the Arctic as BC emissions from local sources in the Arctic were relatively small. The Task Force Hemispheric Transport of Air Pollution Phase2 (HTAP2) set up a series of simulation scenarios to investigate the response of BC in a given region to different source regions. This study investigated the responses of Arctic BC concentrations and surface temperature to 20% anthropogenic emission reductions from six regions in 2010 within the framework of HTAP2 based on ensemble modeling results. Emission reductions from East Asia (EAS) had most (monthly contributions: 0.2 - 1.5 ng m⁻³) significant impact on the Arctic near surface BC concentrations while the monthly contributions from Europe

37 (EUR), Middle East (MDE), North America (NAM), Russia-Belarus-Ukraine (RBU), and South Asia
38 (SAS) were $0.2\text{--}1.0\text{ ng m}^{-3}$, $0.001\text{--}0.01\text{ ng m}^{-3}$, $0.1\text{--}0.3\text{ ng m}^{-3}$, $0.1\text{--}0.7\text{ ng m}^{-3}$, $0.0\text{--}0.2\text{ ng m}^{-3}$,
39 respectively. The responses of the vertical profiles of the Arctic BC to the six regions were found to
40 be different due to multiple transport pathways. Emission reductions from NAM, RBU, EUR, and
41 EAS mainly influenced the BC concentrations in low troposphere of the Arctic, while most of the BC
42 in the upper troposphere of the Arctic derived from SAS. The response of the Arctic BC to emission
43 reductions of six source regions became less significant with the increase of the latitude. The benefit
44 of BC emission reductions in terms of slowing down surface warming in the Arctic was evaluated by
45 using Absolute Regional Temperature-change Potential (ARTP). Compared to the response of global
46 temperature to BC emission reductions, the response of Arctic temperature was substantially more
47 sensitive, highlighting the need for curbing global BC emissions.

48 **1. Introduction**

49 Black carbon (BC) is one of the short-lived climate forcers (SLCFs, AMAP, 2015) and was regarded
50 as the second largest contributor to global warming, only inferior to carbon dioxide (Bond et al.
51 2013). BC over the Arctic can perturb the radiation balance in a number of ways. Direct aerosol
52 forcing occurred through absorption or scattering of solar (shortwave) radiation. BC is the most
53 efficient atmospheric particulate species at absorbing visible light (Bond et al., 2013), the added
54 atmospheric heating will subsequently increase the downward longwave radiation to the surface and
55 warm the surface (AMAP, 2011). Radiative forcing by BC can also result from aerosol-cloud
56 interactions that affected cloud microphysical properties, albedo, extent, lifetime, and longwave
57 emissivity (Twomey 1977; Garrett and Zhao 2006). BC has an additional forcing mechanism after
58 depositing onto snow and ice surfaces (Clarke and Noone, 1985). The surface albedo of snow and ice
59 could be reduced and further enhanced the absorption of solar radiation at the surface. In the Arctic,
60 surface temperature responses were strongly linked to surface radiative forcing as the stable
61 atmosphere of the region prevented rapid heat exchange with the upper troposphere (Hansen and
62 Nazarenko, 2004).

63 The Arctic has been warming twice as rapidly as the world in the past fifty years, and has
64 experienced significant changes in its ice and snow covers as well as permafrost (AMAP, 2017).

65 Reductions of carbon dioxide emissions are the backbone of any meaningful effort to mitigate
66 climate forcing. But even if significant reductions of carbon dioxide are made, slow down of the
67 temperature rise in the Arctic and the sea level rise caused by the melting of glaciers may not be
68 achieved in time. Hence, the goal of slowing down the deterioration of the Arctic may best be
69 achieved by also targeting at shorter-lived climate forcing agents, especially those that could impose
70 appreciable surface forcing and trigger regional-scale climate feedbacks pertaining to the melting of
71 sea ice and snow. Modelling studies by UNEP/WMO (2011) and Stohl et al. (2015) suggested that
72 the climate response of SLCFs mitigation was strongest in the Arctic region. AMAP (2011 and 2015)
73 as well as Sand et al. (2016) demonstrated that per unit of emission reductions of SLCFs in the
74 Northern areas had the largest temperature response on the Arctic, with the Nordic countries
75 (Denmark, Finland, Iceland, Norway, and Sweden) and Russia having the largest impact compared to
76 other Arctic countries such as the United States and Canada.

77 The few studies that investigated specific regional aerosol forcing (Shindell and Faluvegi, 2009;
78 Shindell et al., 2012; Teng et al., 2012) typically used a single climate model at a time to investigate
79 the climate response to idealized, historical, or projected forcing. However, different models varied
80 considerably in the representation of aerosols and radiative properties, resulting in large uncertainties
81 in simulating the aerosol radiative forcing (Myhre et al., 2013b; Shindell et al., 2013). When
82 investigating the climate response to regional emissions, such uncertainties were likely to be
83 confounded even further by the variability between models in regional climate and circulation
84 patterns and variation in the global and regional climate sensitivity (the amount of simulated
85 warming per unit radiative forcing). Hence, the Task Force Hemispheric Transport of Air pollution
86 Phase2 (HTAP2, <http://www.htap.org/>) incorporating multiple global models can avoid the great
87 uncertainty of single model to a certain degree, with the aim to improve model estimates of the
88 impacts of intercontinental transport of air pollutants on climate, ecosystems, and human health
89 (Galmarini et al., 2017). To date, the HTAP2 results have been explored from a variety of scientific
90 and policy-relevant perspectives. For instance, by comparing against observations, sulfur and
91 nitrogen depositions during HTAP2 had been significantly improved compared to HTAP1. From
92 2001 to 2010, the global nitrogen deposition increased 7% while the global sulfur deposition
93 decreased 3% (Tan et al., 2018a). The significant impacts of hemispheric transport on the deposition
94 were specifically focused and the deposition over the coastal regions was more sensitive to

95 hemispheric transport than the non-coastal continental regions (Tan et al., 2018b). Jonson et al. (2018)
96 assessed the contributions from different world regions to European ozone levels and contributions
97 from the non-European regions were mostly from North. America and eastern Asia, larger than those
98 from European emissions. Hogrefe et al. (2018) found that the simulated ozone over the continental
99 US varied very differently by digesting boundary conditions from four hemispheric or global models.
100 The impact of emission changes from six major source regions on global aerosol direct radiative
101 forcing was estimated (Stjern et al., 2016). In the local source regions, the radiative forcing
102 associated with SO_4^{2-} was strengthened (25%) while that from BC was weakened (37%) due to a
103 20% emission reduction. Liang et al. (2018) estimated global air-pollution-related premature
104 mortality from exposure to $\text{PM}_{2.5}$ and ozone and the interregional transport lead to more deaths
105 through changes in $\text{PM}_{2.5}$ than in O_3 . However, the source region contributions to Arctic BC and the
106 spread among multi-model results have been rarely explored from the perspective of HTAP2
107 initiative.

108 This study aims to investigate the responses of Arctic BC concentrations and surface temperature
109 to 20% anthropogenic emission reductions from different regions in the Northern Hemisphere (NH).
110 A comparison of six global modeling works within the framework of HTAP2 experiments for the
111 Arctic region in 2010 was presented. The ensemble modeling results were used to apportion the
112 contribution from different source regions to the near-surface and vertical black carbon in the Arctic.
113 In addition, the Arctic surface temperature responses to the emission reductions were estimated.

114 **2. Methodology**


115 **2.1 HTAP2 experiments**

116 HTAP2 developed a harmonized emissions database covering all countries and the major sectors for
117 global and regional modeling from 2008 to 2010. The emissions database was obtained from the
118 nationally reported emissions (e.g., National Emission Inventory for the United States), the regional
119 scientific inventories (e.g., European Monitoring and Evaluation Programme (EMEP), Netherlands
120 Organisation for Applied Scientific Research (TNO) for Europe, Model Inter-Comparison Study for
121 Asia, MICS–Asia III), and the Emissions Database for Global Atmospheric Research data
122 (EDGARv4.3) for the rest of the world (mainly South America, Africa, Russia, and Oceania).

123 Biomass burning emissions were not prescribed in HTAP2. Temporal resolution of data sources was
 124 monthly, and thus the HTAP2 emission inventory provided harmonized emission data with monthly
 125 resolution for all the air pollutants including BC. It should be noted that the emissions of
 126 international shipping and international aviation in HTAP2 were considered constant over the year. It
 127 was recommended that modeling groups used the Global Fire Emissions Database (GFED4,
 128 <http://globalfiredata.org/>) with a temporal resolution of daily or 3-hour intervals. The detailed
 129 information of different regional inventories can be found in Janssens–Maenhout et al. (2015).

130 Emission perturbations were conducted in sensitivity simulations to investigate the response of
 131 various air pollutants in a given region to different source regions. In this study, the Arctic region was
 132 the targeted receptor region of interest. Six source regions in HTAP2 experiments, namely, East Asia
 133 (EAS), Europe (EUR), Middle East (MDE), North America (NAM), Russia–Belarus–Ukraine (RBU),
 134 and South Asia (SAS) were selected to demonstrate their influences on the BC concentrations over
 135 the Arctic region (Figure 1a). Two emission scenarios were designed for the HTAP2 simulation to
 136 explore the source/receptor relationships, i.e. the base scenario (BASE) with no emission reduction,
 137 and the control scenario (EASALL, EURALL, MDEALL, NAMALL, RBUALL, and SASALL)
 138 with 20% reduction of all anthropogenic emissions in six regions respectively.

139



140

141 **Figure 1.** (a) The sketch map of receptor and source regions. (b)–(g) Spatial distributions of 20% reduction of
 142 annual BC emission in the six source regions in 2010. MDE: Middle East; EUR: Europe; RBU:
 143 Russia–Belarus–Ukraine; NAM: North America; EAS: East Asia; SAS: South Asia. The unit legends from (b) to (g)
 144 are the same of $10^{-11} \text{ kg m}^{-2} \text{ s}^{-1}$.

2.1.1 Anthropogenic emission reductions of BC in HTAP2

Anthropogenic BC emission sectors included power plants, industries, transportation, shipping, aviation, agriculture, and residential sectors. The emission inventory had a monthly temporal resolution and a spatial resolution of $0.1^{\circ} \times 0.1^{\circ}$. The total anthropogenic emissions and 20% emission reductions of BC in six source regions of HTAP2 in 2010 are presented in Table 1. The higher BC emission reductions were found in the EAS and SAS with the values of 355.6 and 232.5 Gg yr⁻¹, respectively, while were much lower in the MDE and RBU with the values of 5.3 and 18.6 Gg yr⁻¹, respectively. The BC emission reductions in the EAS, EUR, and RBU showed significant monthly variations with higher values from November to March, while the monthly variations were not obvious in the MDE, NAM, and SAS.

Table 1. 20% emission reductions and total anthropogenic emissions of BC in different regions of HTAP2 in 2010. (Unit: Gg yr⁻¹).

Regions	Total anthropogenic emissions	20% Emission Reductions												
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2010
EAS ^a	1778.1	46.4	35.6	33.0	24.1	23.9	24.0	24.0	23.6	23.5	25.0	31.4	41.0	355.6
EUR ^b	326.3	6.7	6.4	7.2	6.5	5.3	4.9	4.0	3.7	4.4	5.2	5.3	5.7	65.3
MDE ^c	26.7	0.4	0.4	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.5	5.3
NAM ^d	310.8	5.2	5.1	5.3	5.1	5.1	5.2	5.3	5.3	5.1	5.1	5.1	5.2	62.2
RBU ^e	93.0	2.0	1.9	1.9	1.7	1.4	1.3	1.0	1.1	1.2	1.6	1.7	1.8	18.6
SAS ^f	1162.7	20.4	19.1	19.7	18.9	19.2	18.9	19.2	19.2	19.0	19.3	19.2	20.4	232.5
All	3697.6	81.2	68.5	67.6	56.8	55.4	54.7	53.9	53.3	53.6	56.7	63.2	74.6	739.5
Global	5492.9	110.6	86.2	92.8	103.9	98.7	97.2	86.8	85.4	85.4	84.2	83.3	83.9	1098.6

^a East Asia. ^b Europe. ^c Middle East. ^d North America. ^e Russia–Belarus–Ukraine. ^f South Asia.

Figure 1b–g illustrates the spatial distribution of the 20% reductions of annual BC emissions in six source regions in 2010. It can be found that the most intense reductions of BC emissions in EAS and SAS were concentrated in East China and India, respectively, which were mainly attributed to emissions from residential sectors, followed by transportation and industries. The BC emission reductions of EUR were widely distributed with high values in central Europe, with residential and transportation sectors accounting for the largest proportion. The reductions near the Arctic circle could be found in the north of EUR, NAM, and RBU. For MDE, most BC was emitted from Iran, which located in the northeast of this region. Overall, the spatial pattern of BC emission reductions in six regions was closely related to the spatial distribution of the human population.

168 2.1.2 Model description

169 Considering that the simulations should cover all months of 2010 and all emission source regions,
 170 five global models (i.e. CAMchem, CHASER_re1, GEOS-Chem, GOCART-v5, and Oslo CTM3-v2)
 171 were incorporated to simulate the responses of BC concentrations in the Arctic to 20% BC emission
 172 reductions from EAS, EUR, MDE, NAM, RBU, and SAS, respectively. The brief information of
 173 model configurations is listed in Table 2. As required by HTAP2, all simulations should include a
 174 spin-up time of 6 months prior to the period of interest. The outputs from all models are available
 175 upon request from <http://aerocom.met.no>. The time resolution of the outputs used in this study is
 176 monthly for all models, although models were run at a finer resolution (e.g., daily or hourly). The
 177 model outputs for air pollutants were originally provided in the unit of mass mixing ratio (MMR, kg
 178 kg⁻¹). To facilitate comparison between model and observation and further data analysis, we
 179 converted the original units into ng m⁻³ based on the ideal gas law (Aamaas et al., 2017).

180

181 **Table 2.** Configurations of models used in this study

Models	Meteorological field	Horizontal resolution	Vertical layers	Convection	Reference
CAMchem	GEOS5 v5.2	1.9° × 2.5°	56	Zhang–McFarlane approach for deep convection	Lamarque et al., 2012; Tilmes et al., 2016
CHASER_re1	ERA-Interim and HadISST	2.8° × 2.8°	32	CCSR/NIES AGCM for advection, convection, and other subgrid-scale mixing	Sudo et al., 2002; Takashi et al., 2018
GEOS-Chem	GEOS-5 (MERRA)	2.0° × 2.5°	47	Convective transport is computed from the convective mass fluxes in the meteorological archive	Henze et al., 2007
GOCART-v5	MERRA	1.3° × 1.0°	72	MERRA for moist convection, Arakawa–Schubert (RAS) algorithm for GCTM	Chin et al., 2000
Oslo CTM3-v2	ECMWF-IFS	2.8° × 2.8°	60	Tiedke mass flux scheme for deep convection	Søvde et al., 2012; Lund et al., 2018

182 2.2 Calculation of the temperature response to BC emissions reduction

183 The climate effects of air pollutants have been the focus of climate change research since the last
 184 century (IPCC, 1990; IPCC, 2001). In the last few years, the metrics for estimating this kind of effect
 185 have been constantly improving (Shindell et al., 2012; Bond et al., 2013; Smith and Mizrahi, 2013;

186 Stohl et al., 2015). The Intergovernmental Panel on Climate Change (IPCC) used the Global
187 Warming Potential (GWP) as a method for comparing the potential climate impact of emissions of
188 different greenhouse gases (IPCC, 1990). GWP is the time-integrated radiative forcing due to a pulse
189 emission of a given species, over some given time horizon (commonly 20, 100, or 500 years) relative
190 to a pulse emission of carbon dioxide. GWP does not purport to represent the impact of air pollutant
191 emissions on temperature. Although a short-lived climate pollutant (SLCP) could have the same
192 GWP as a long-lived climate pollutant, identical (in mass terms) pulse emissions could cause a
193 different temperature change at a given time, because long-lived climate pollutants accumulate in the
194 climate system while short-lived climate pollutants can be broken down by various processes.
195 Consequently, warming caused by long-lived climate pollutants is determined by total cumulative
196 emissions to date, while the warming due to short-lived climate pollutants is determined more by the
197 current rate of emissions in any given decade and depends much less on historical emissions. This
198 means the importance of SLCP emissions is often overstated based on GWP. Shine et al. (2005)
199 proposed the Global Temperature Change Potential (GTP) as a replacement for GWP to represent the
200 global-mean surface temperature change for both a pulse emission (GTP_P) and a sustained change in
201 emissions (GTP_S) of a given air pollutant. The distinction between GTP_P and GTP_S avoids the
202 overestimation of GWP for the short-lived climate pollutants. Even for a uniform forcing, there will
203 be differences of spatial patterns in the temperature response. Regional Temperature-change Potential
204 (RTP) (Shindell and Faluvegi, 2010) was applied to analyze the temperature response on the regional
205 scale, since both GWP and GTP focused on the global scale. The GWP, GTP, and RTP were
206 normalized to the corresponding effect of CO_2 as the Absolute Global Warming Potential (AGWP),
207 Absolute Global Temperature Change Potential (AGTP), and Absolute Regional Temperature-change
208 Potential (ARTP), respectively. AGWP represented the absolute forms of radiative forcing. AGTP
209 and ARTP represented the absolute forms of temperature perturbation. The ARTP provide additional
210 insight into the spatial pattern of temperature response to inhomogeneous forcings beyond that
211 available from traditional global metrics. Very few metrics have attempted to examine sub-global
212 scales thus far, though some have used local information with non-linear global damage metrics
213 (Shine et al., 2005a; Lund et al., 2012). Shindell et al. (2012) indicated that the forcing/response
214 portion of the ARTP appeared to be relatively robust across models.

215 ARTPs is more suitable for this study to calculate the temperature response, considering that the

research object is BC with short lifetime and focus on regional impact of the BC emission reductions on temperature changes in the Arctic. For SLCFs with atmospheric lifetimes much shorter than both the time horizon of the ARTP and the response time of the climate system, the general expression for the ARTP following a pulse emission of BC (E) in region r which leads to a response in latitude band m is as follows (Fuglestad et al., 2010; Collins et al., 2013; Aamaas et al., 2017):

$$\text{ARTP}_{r, m, s}(H) = \sum_l \frac{F_{l, r, s}}{E_{r, s}} \times \text{RCS}_{l, m} \times R_T(H) \quad (1)$$

$F_{l, r, s}$ (in W m^{-2}) is the radiative forcing in latitude band l due to emission in region r in season s as a function after the pulse emission $E_{r, s}$ (in Tg). Even though our estimates are based on seasonal emissions, the temperature responses calculated are annual means. Shindell and Faluvegi (2009) analyzed BC climate effect in four different latitudes: southern mid-high latitudes (90°S – 28°S), tropics (28°S – 28°N), northern mid-latitudes (28°N – 60°N), and the Arctic (60°N – 90°N), which gives a better estimate of the global temperature response as it accounts for varying efficacies with latitude. The $\text{RCS}_{l, m}$ is a matrix of regional response coefficients based on the RTP concept (unitless; Collins et al., 2013). As these response coefficients are normalized here, they contain no information on climate sensitivity, only the relative regional responses in the different latitude bands. The global climate sensitivity is included in the impulse response function R_T , which is a temporal temperature response to an instantaneous unit pulse of RF (in $\text{K m}^2 \text{W}^{-1}$). This paper refers to the ARTP values in Aamaas et al. (2017). Aamaas et al. (2017) applies two refinements of the forcing-response coefficients for radiative forcing occurring in the Arctic: one for the aerosol effects in the atmosphere (Shindell and Faluvegi, 2010; Lund et al., 2014) and another for the effects due to BC on snow (Flanner, 2013). The ARTP in this study estimated of the direct effect in the Arctic included both the direct radiative forcing from outside the Arctic and within the Arctic, while the ARTP of the semi-direct effect in the Arctic was due to the semi-direct radiative forcing from outside the Arctic. The contribution by radiative forcing within the Arctic to Arctic temperature changes considered the vertical profile of BC concentrations as both $F_{\text{Arctic}, r, s}$ and $\text{RCS}_{\text{Arctic}, \text{Arctic}}$ have a dependence on the height of the BC (Lund et al., 2014; Lund et al., 2017). The total response in the Arctic was the sum of the contributions from BC forcing outside of the Arctic and inside of the Arctic.

Regional temperature responses at time t of an emission $E(t)$ can be calculated with these ARTP values by a convolution (Aamaas et al., 2016). The temperature response is as follows:

$$\Delta T_{r, m, s, t}(t) = \int_0^t E_{r, s, t}(t') \times ARTP_{r, m, s, t}(t-t') dt \quad (2)$$

$\Delta T_{r, m, s, t}$ refers to the decrease of the Arctic or global surface temperature after 20, 100, or 500 years to 20% BC emission reductions of six regions (namely EAS, EUR, MDE, NAM, RBU, and SAS) in the framework of HTAP2 either during summer or winter in this paper.

3. Results and Discussion

3.1 Model evaluation

To evaluate the model performance from all five models, the monthly simulated surface BC concentrations of the BASE scenario were compared with the observations at four monitoring sites in the Arctic Circle in 2010. The locations of the four sites, including Alert (82.5°N, 62.3°W) in Canada, Barrow (71.3°N, 156.6°W) in Alaska, Tiksi (71.59°N, 128.92°E) in Russia, and Zeppelin (78.9°N, 11.9°E) in Norway, are plotted in Figure S1 in the Supporting Information.

Metrics (Text S1) including correlative coefficient (COR), normalized mean bias (NMB), normalized mean error (NME), mean bias (MB), and mean absolute error (MAE) were selected for evaluating the model performance in this study (U.S. EPA, 2007). In addition to the evaluation for each single model, the multi-model ensemble mean (calculated as the average of all participating models) was also evaluated. The statistical results are listed in Table 3 and Table S1. A comparison between the monthly variations of simulated and observed BC concentrations is shown in Figure S2 (a).

The correlations of the simulated BC concentrations among different models were moderate to high with CORs ranging from 0.33 to 0.98 (Table S1), suggesting the temporal variations of different models were relatively consistent. Overall, CAMchem, GEOS-Chem, GOCART-v5, and Oslo CTM3-v2 underestimated the near-surface BC (Figure S2a), which may be attributed to an underestimation of BC emissions, e.g., gas flaring (Huang et al., 2014, 2015; Stohl et al., 2013) and shipping emissions (Marelle et al., 2016). Also, appropriate temporal allocation of BC emissions from residential combustion was another important factor governing the model performance (Stohl et al., 2013). However, the simulated BC surface concentrations from CHASER_re1 were higher than

271 those of the other four models and observations (Figure S2a), which mainly due to their slow BC
272 aging-rate in remote/polar regions (Sudo et al., 2015).

273 Table 3 shows the model performances at the four Arctic sites. No single model could reproduce
274 the BC concentrations in the Arctic well, and models performed differently at different monitoring
275 sites. Relatively good agreement between the observation and models was found at Zeppelin, with
276 CORs, NME, MB, and MAE of 0.59–0.83, 38.59%–142.64%, –13.53–14.97 ng m⁻³, and 5.40–14.97
277 ng m⁻³ among the five models, respectively. The best correlation (0.83) was found at Zeppelin from
278 Oslo CTM3, while the smallest NMB (38.59%) and MAE (5.40 ng m⁻³) were found at Zeppelin from
279 GOCART. On the contrary, the simulated BC concentrations didn't agree so well with observations
280 at the other three sites with even negative COR values in some models (e.g., CAMchem, and
281 CHASER_re1), which may be explained by the uncertainties in emission inventory, the bias in the
282 meteorological simulations, and chemical mechanisms (Miao et al., 2017; Zhang et al., 2019). All
283 models, except Oslo CTM3, overestimated the BC concentrations in Barrow in July (Figure S2a),
284 mainly due to the large contributions of biomass burning from Siberia in the simulations caused by
285 overestimations of emissions and/or too little removal during transport (Sobhani et al., 2018).

286 The vertical profiles of simulated BC concentrations of the BASE simulation were also compared
287 with aircraft measurements from HIAPER Pole-to-Pole Observations (HIPPO) during 24 March–16
288 April 2010 (Figure S2b). Different from comparison between observed and simulated BC
289 concentrations near the surface, the vertical profiles of BC concentrations were overestimated by
290 most models. As the aircraft ascended and descended along each flight track, BC concentrations from
291 HIPPO varied with time, latitude, longitude, and altitude. However, most of the simulation results of
292 HTAP2 were provided in the temporal resolution of monthly, simulation and observation results
293 cannot be exactly matched. This partly explained the difference between the simulations and
294 observations. Overall, currently no single model could reproduce the BC concentrations over
295 different regions of the Arctic well. There is a number of reasons responsible for this. First, the BC
296 emission inventory in the Arctic is not well understood due to lacking of local activity data and
297 emission factors, e.g. gas flaring in the oil and gas production fields, biofuel combustion, non-road
298 transportation, etc. Secondly, the lifetime of BC in the atmosphere is sensitive to its wet deposition
299 rates. However, different models have divergent treatment of wet scavenging parameterizations,
300 which may be not representative in the Arctic region and could result in the simulated BC

concentrations ranging between several magnitudes. The mechanism of BC sinks is still not well understood in the Arctic. Last but not the least, almost all the global models used the latitude/longitude projection which has very large distortions over the polar regions and this may also affect the ability of global models simulating the air pollutants over the Arctic region.

Although the single model didn't reproduce the BC concentrations in the Arctic well, the consistency of the model ensemble mean with the observation was improved to some extent. The NME and MAE of model ensemble mean was closer to zero compared with the single model. Therefore, to reduce the bias from one single model, the multi-model ensemble mean was used for further analysis.

Table 3 Comparison of the simulations and observations of monthly surface BC concentrations at Alert, Barrow, Tiksi, and Zeppelin in 2010.

Parameters	Sites	CAMchem	CHASER_re1	GEOS-Chem	GOCART-v5	Oslo CTM3-v2	Model ensemble mean
COR ^a	Alert	-0.24	-0.22	0.35	0.20	-0.24	-0.10
	Barrow	-0.28	-0.08	0.06	0.00	0.01	-0.06
	Tiksi	-0.19	0.05	0.50	0.48	0.41	0.11
	Zeppelin	0.72	0.59	0.80	0.76	0.83	0.73
NMB ^b (%)	Alert	-86.75	115.06	-57.81	-34.31	-92.38	-9.21
	Barrow	-38.43	104.10	-38.95	-8.18	-75.58	4.37
	Tiksi	-82.03	10.31	-69.76	-67.34	-84.82	-46.79
	Zeppelin	-79.93	142.64	-45.57	-9.81	-75.98	8.63
NME ^c (%)	Alert	86.75	151.30	66.37	70.77	92.38	74.69
	Barrow	72.07	124.44	69.50	84.20	75.58	72.12
	Tiksi	82.03	64.55	70.16	68.82	84.82	60.81
	Zeppelin	79.93	142.64	45.57	38.59	75.98	42.06
MB ^d (ng m ⁻³)	Alert	-29.03	11.08	-21.41	-16.07	-30.16	-12.81
	Barrow	-22.13	15.35	-19.10	-11.12	-30.40	-9.44
	Tiksi	-55.99	-17.28	-48.51	-48.16	-56.26	-40.64
	Zeppelin	-13.53	14.97	-8.19	-3.59	-12.45	-1.44
MAE ^e (ng m ⁻³)	Alert	29.03	31.05	22.85	22.23	30.16	23.56
	Barrow	29.01	28.91	26.71	30.30	30.40	25.22
	Tiksi	55.99	37.06	48.60	48.49	56.26	43.73
	Zeppelin	13.53	14.97	8.19	5.40	12.45	4.95

^a Correlative coefficient. ^b Normalized mean bias. ^c Normalized mean error. ^d Mean bias. ^e Mean absolute error.

314 **3.2 Near-surface BC concentrations in the Arctic**

315 Before analyzing the responses of Arctic BC to emission reductions, it is necessary to understand the
316 spatial-temporal distribution of BC concentrations in the Arctic region. In this study, months from
317 May to October were defined as summer and November to April were defined as winter due to the
318 special geographical location of the Arctic (Aamaas et al., 2017).

319 Spatial distributions of Arctic near surface BC concentrations in summer and winter simulated
320 from each model are showed in Figure S4. BC simulated by CHASER_re1 showed relatively high
321 concentrations over the whole Arctic, followed by GEOS-chem and GOCART-v5, while those
322 simulated by Oslo CTM3-v2 and CAMchem were lower. The difference of simulated BC
323 concentrations between land and ocean was more obvious in summer than that in winter, especially
324 for GEOS-chem and GOCART-v5. The mean BC concentrations from the ensemble models near the
325 surface Arctic (66–90°N) were 18.6 ng m⁻³ in summer and 16.6 ng m⁻³ in winter in 2010,
326 respectively. Figure 2 shows that the BC concentrations over the polar sea ice region in winter were
327 higher than that in summer. The coverage of the polar dome expanded more southward in winter
328 (Bozem et al., 2019; Law and Stohl, 2007), allowing more BC from lower latitudinal regions to be
329 transported into the Arctic. Turbulent exchange and deposition were reduced during winter as the
330 meteorological conditions in the Arctic were stable and dry (Bradley et al., 1992; Bozem et al., 2019;
331 Law and Stohl, 2007). In addition, BC emissions in EAS, EUR, and RBU regions showed obvious
332 monthly changes with higher emissions from November to March as mentioned earlier (Section
333 2.1.1), leading to the relatively high BC concentrations over the polar sea ice region in winter. Over
334 the terrestrial areas within the Arctic Circle, summer BC concentrations were higher than winter,
335 especially in Siberia and Alaska, which were attributed to intense BC emissions from biomass
336 burning over these areas from Jun to Aug (Figure S3).

337





Figure 2. Spatial distribution of near-surface BC concentrations in **(a)** summer and **(b)** winter in the Arctic in 2010.


3.3 Response of Arctic BC to 20% emission reductions

3.3.1 Contributions of regional emission reductions to the Arctic near-surface BC


The response of the Arctic near-surface BC to 20% emission reductions from different source regions was analyzed through emission perturbation simulations. Figure 3 shows the spatial distribution of the referred response above in summer and in winter of 2010 based on multi-model ensemble mean results. The source region contributions to the surface BC concentrations exhibited significant seasonal variability with higher values in winter. The BC emission reductions in EAS almost affected the whole Arctic, especially in winter, indicating the significance of the intercontinental transport of BC. The spatial distribution of the Arctic near-surface BC response to SAS emission reductions was similar to that of EAS, but the extent was much weaker. The emission reductions from EUR, NAM, and RBU mainly affected the local and nearby areas, which was generally consistent with the spatial pattern of emissions (Figure 1). The contribution from MDE emission reductions was very little.




(a₁) EASALL–summer




(a₂) EASALL–winter




(b₁) EURALL–summer




(b₂) EURALL–winter




(c₁) MDEALL–summer




(c₂) MDEALL–winter




(d₁) NAMALL–summer




(d₂) NAMALL–winter



(e₁) RBUALL–summer



(e₂) RBUALL–winter







Figure 3. Spatial distribution of contribution of 20% emission reductions of different source regions to Arctic near-surface BC in summer and in winter in 2010.

The monthly variations of the response of the Arctic near-surface BC concentrations to 20% emission reductions from six source regions are presented in Figure 4. Results from the ensemble simulations are averaged over the Arctic covering latitudinal areas from 66°N to the north pole. The emission reductions from the total six source regions were 329.6 Gg during May to October, lower than that of 411.9 Gg during November to April (Table 1). Correspondingly, the contributions of 20% BC emission reductions from all six regions to Arctic **monthly** near-surface BC concentrations were 0.8–1.4 ng m⁻³ during May to October and 1.5–3.2 ng m⁻³ during November to April. Arctic sensitivities (Arctic concentration change per unit source region emission change) for BC typically maximized from December to February for EUR and RBU and from March to May for EAS and NAM (Shindell et al., 2008; AMAP, 2008). The enhanced sensitivity from December to May resulted from faster transport and slower removal during winter as the meteorological conditions in the Arctic were stable and dry (Law and Stohl, 2007). The results of deposition changes also proved this result well (Figure S5). The wet deposition in summer was higher than that in winter, which was 7–13 times of dry depositions. Sharma et al. (2013) found that the Arctic region (north of 70°N) was very dry during winter with an average daily precipitation rate between 0 and 1 mm day⁻¹. Precipitation rates over some of the BC source regions such as Eurasia were at the same order of magnitude as the Arctic. Less wet deposition and a shallow boundary layer resulted in higher BC concentrations near the surface during winter. In the summertime, the Arctic region experienced 2 to 3 times higher precipitation rates as well as wet depositions of BC relative to wintertime, thus resulting in lower contributions to the near-surface BC concentrations.

The annual contribution of 20% emission reductions from EAS, EUR, MDE, NAM, RBU, and SAS to the Arctic near-surface BC concentrations reached 0.70, 0.54, 0.01, 0.20, 0.29, and 0.09 ng

377 m^{-3} in 2010, respectively, totaling about 1.83 ng m^{-3} . A simple linear interpolation suggested the
378 contribution of 100% BC emissions from six regions to the Arctic near-surface BC concentrations
379 was about 9.15 ng m^{-3} (five times of 1.83 ng m^{-3}). The annual mean Arctic near-surface BC
380 concentration from the BASE simulation was about 18 ng m^{-3} in 2010. Thus, the impact of emissions
381 from six regions on the Arctic near-surface BC was outstanding. It should be noted that the
382 contributions from six regions only considered anthropogenic emissions while the contribution from
383 biomass burning was not included in the sensitivity experiments of HTAP2. It is known that
384 wildfires in East of Russia and U.S. Alaska are important sources of BC in the Arctic region,
385 especially in summer. Thus, the contributions from six regions to the Arctic BC should be even more
386 dominate over the other regions by including biomass burning in RBU and NAM. The response of
387 Arctic near-surface BC concentration was found strongest to the 20% emission reductions from EAS
388 with the monthly contribution of $0.2\text{--}1.5 \text{ ng m}^{-3}$, accounting for 16.8%–49.0% of the total reduced
389 BC concentrations resulting from all six source regions (Figure 4). On one hand, the BC emission
390 reductions in EAS were the largest among the six source regions (Table 1). On the other hand, BC
391 emission reductions in EAS can influence the Arctic lower troposphere via two pathways (Bozem et
392 al., 2019; Stohl, 2006). BC from northern regions of EAS can enter into the polar dome of the Arctic
393 in winter, as the air masses have cooled during transport. BC from eastern regions of EAS fast
394 uplifted due to convection and then followed by high altitude transport in northerly directions.
395 Radiative cooling eventually led to a slow descent into the polar dome area after air masses arrived
396 in the high Arctic. It occurred both in summer and winter. In addition to EAS, BC emission reduction
397 from EUR also showed significant impacts on the Arctic near-surface BC concentration with the
398 monthly contribution of $0.2\text{--}1.0 \text{ ng m}^{-3}$, accounting for 20.1%–49.0% of the total reduced BC
399 concentrations resulting from all six source regions (Figure 4). Among the three regions in the Arctic
400 Circle (i.e. EUR, NAM, and RBU), EUR region had the largest BC emission reductions. Also, the
401 relatively short distance between EUR and the Arctic made EUR the second most important source
402 region to the Arctic. As for NAM and RBU, their 20% emission reductions induced moderate
403 reductions of the monthly Arctic near-surface BC concentrations by $0.1\text{--}0.3$ and $0.1\text{--}0.7 \text{ ng m}^{-3}$,
404 respectively. The contribution of 20% emission reductions from SAS to the Arctic near-surface BC
405 concentrations was much lower of monthly contributions of $0.0\text{--}0.2 \text{ ng m}^{-3}$ as a significant portion of
406 BC originating from SAS accumulated in the upper troposphere (Section 3.3.2). Compared to the

407 five source regions discussed above, the response of Arctic BC concentrations to emission reductions
 408 from MDE was negligible, owing largely to the low emissions there and long distance from the
 409 Arctic.
 410



411 **Figure 4.** Monthly mean reduced concentrations of the near-surface Arctic BC due to 20% emission reductions
 412 from six source regions in 2010.

413 Figure S6 compares the contributions of 20% emission reductions to Arctic near surface BC
 414 concentrations simulated by different models. All five models showed similar monthly variations, of
 415 which CHASER_re1 simulated high BC concentrations compared to the other models due to slow
 416 aging-speed (Sudo et al., 2015). All models showed the major source regions of Arctic BC from EAS,
 417 EUR, and RUB. NAM and SAS contributed moderately while the contribution from MDE was
 418 negligible.

419

420 3.3.2 Contributions of regional emission reductions to the vertical BC profiles

421 To assess the contributions from various source regions to the BC profiles based on the model
 422 ensemble mean, the vertical stratification needed to be unified as most participating models had
 423 different vertical settings. Since CHASER had a relatively coarse vertical resolution of 32 layers, the
 424 other models were unified to the same vertical stratification, as detailed in Table S2.

425 As shown in Figure 5, the contributions of regional emission reductions to BC exhibited strong
 426 vertical gradients over the Arctic. In general, the BC profiles displayed a bimodal pattern in summer,
 427 showing peaks at around 1.0–1.6 km a.s.l. (4th and 5th layers) and 8.0–8.9 km a.s.l. (13th and 14th

layers). While in winter, the BC profiles showed a unimodal pattern with peaks around 0.6–1.6 km a.s.l. (3rd – 5th layers). Long-range transport of air pollutions may occur near the planetary boundary layer (Eckhardt et al., 2003; Stohl et al., 2002). High contributions in the low layers (e.g., 3rd – 5th layers) were consistent with the height of the planetary boundary layer in the Arctic (Zhang, et al., 2018; Cheng, 2011).

It has been summarized that there were several major transport pathways for BC into the Arctic troposphere (Stohl, 2006). i) BC transported rapidly at low-level, followed by uplifting at the Arctic front when it is located far north. Significant deposition of BC in the Arctic occurs mostly north of 70°N for this transport route. This transport route derived often from the high BC emission areas in northern EUR but seldom from the NAM and RBU. That was mainly due to that the BC emissions exist at high enough latitudes in EUR, which can be north of the polar front. However, the BC emissions in NAM and RBU were concentrated south of the polar front (Figure 1), thus BC emitted from these two regions can't be easily transported into the Arctic through this pathway. ii) Cold air masses into the polar dome transport at low-level. This pathway derived mainly from EUR and high-latitude areas of EAS during winter. The contribution of 20% emission reductions from EUR to the Arctic BC concentrations peaked at around 1.0 km a.s.l. with the multi-model ensemble mean value of 0.4 ng m⁻³ in summer, while peaked at lower altitude of around 1.6 km a.s.l. with the value of 0.8 ng m⁻³ in winter. iii) BC could also ascend south of the Arctic followed either by high-altitude transport or by several cycles of upward and downward transport, and finally slowly descended into the polar dome due to radiative cooling. This was the frequent transport route from source regions such as NAM, RBU, and east EAS. The contribution from NAM and RBU to the Arctic BC peaked at about 1.6 km a.s.l. (0.2 ng m⁻³) and 1.0 km a.s.l. (0.2 ng m⁻³) in summer, and peaked at about 1.0 km a.s.l. (0.3 ng m⁻³) and 0.4 km a.s.l. (0.5 ng m⁻³) in winter. The contribution from EAS including pathways ii and iii, to the Arctic BC peaked at about 1.6 km a.s.l. (0.6 ng m⁻³) in summer and peaked at about 2.4 km a.s.l. (1.6 ng m⁻³) in winter. Matsui et al. (2011) pointed out that Asian anthropogenic air masses were measured most frequently in the upper troposphere, with median values of 20 ng m⁻³ (410hPa) in April 2008 and 5 ng m⁻³ (353hPa) in June–July 2008. In our analysis, the contribution of 20% emission from EAS and SAS to BC in the Arctic was 1.4 ng m⁻³ (432hPa) in April 2010 and 0.7 ng m⁻³ (375hPa) in June–July 2010. If the contribution was linearly interpolated, the contribution of 100% emission from EAS and SAS to BC in the Arctic would be

about 7 ng m^{-3} (432hPa) in April and 3.5 ng m^{-3} (375hPa) in June–July in 2020. In general, our results were at the same magnitude with Matsui et al. (2011). The contribution from MDE was negligible.

As shown in Figure 5, BC can also be transported into the upper troposphere of the Arctic. Air masses preferably kept their potential temperature almost constant during transport as the atmospheric circulation can be well described by adiabatic motions in the absence of diabatic processes related to clouds, radiation, and turbulence. The potential temperature was low within the polar dome area, and thus only air masses experienced diabatic cooling were able to enter the polar dome (Stohl, 2006). That is to say, the air masses from SAS and low-latitude regions of EAS were not easy to penetrate the polar dome but can be lifted and transported to the Arctic in the middle and upper troposphere along the isentropes (AMAP, 2011; Barrie, 1986; Law and Stohl, 2007; Stohl, 2006). This agreed well with the previous study of Koch and Hansen (2005) and Stohl (2006). The contribution from SAS to the Arctic BC concentrations peaked at about 9.7 km a.s.l. (0.4 ng m^{-3}) in summer and 9.7 km a.s.l. (0.5 ng m^{-3}) winter. This was also consistent with the vertical profiles of BC shown in Stjern et al. (2016). The polar dome boundary was variable in time and space and was not zonally symmetric. The range of polar dome expanded southward to about 40°N over Eurasia in winter as the temperature difference of different latitudes became smaller (Bozem et al., 2019; Law and Stohl, 2007), resulting in the contribution of EAS to the Arctic BC concentrations in upper troposphere only peak in summer at 13th layer (8.0 km a.s.l.) with the value of 0.6 ng m^{-3} .




Figure 5. Contribution of 20% emission reductions from six source regions to BC concentrations in different vertical layers (a) in summer and (b) in winter in the Arctic in 2010.

480 3.3.3 Contributions of emission reductions to BC in different latitudinal bands

481 To further analyze the response of the Arctic BC concentrations to emission reductions of six source
482 regions in HTAP2, the contribution of 20% emission reductions to BC concentrations at different
483 latitudes of the Arctic were calculated (Figures 6 and 7). In regard to the different horizontal
484 resolution of participating models, the Arctic region (66–90°N) was divided into eight latitudinal
485 bands with a 3-degree interval, which was based on the coarsest resolution of all models.

486 The response of the Arctic BC concentrations to emission reductions of six source regions became
487 weaker with the increase of the latitude due to the continuous loss of BC during transport (e.g., dry
488 and wet depositions) (Figure 6). The difference of contributions between two adjacent latitudinal
489 bands became smaller as closer to the north pole. The contributions of 20% emission reductions to
490 the Arctic BC concentrations near surface were the highest between 66–69°N both in summer (1.9 ng
491 m⁻³) and winter (4.1 ng m⁻³), which were 1.4–2.8 times higher than the other latitudinal bands.

492 The contributions from EAS and EUR were higher than those from the other four regions in each
493 latitudinal band. In detail, the contributions from EUR (0.8 ng m⁻³ in summer and 1.5 ng m⁻³ in
494 winter) were higher than those from EAS (0.6 ng m⁻³ in summer and 1.2 ng m⁻³ in winter) in the
495 latitudinal band of 66–69°N as the BC concentrations near surface there were more sensitive to the
496 local emission sources. In contrast, the latitudinal contributions from EAS (0.3–0.4 ng m⁻³ in
497 summer and 0.9–1.1 ng m⁻³ in winter) were higher than those from EUR (0.2–0.4 ng m⁻³ in summer
498 and 0.4–0.9 ng m⁻³ in winter) in the other high latitudinal bands where long-range transport played
499 the dominant role.

500 The downward trends of the response of the Arctic near surface BC to emission reductions with
501 the increase of latitude from EUR and RBU were more obvious than that of other regions (Figure 6).
502 Dry and wet depositions of BC decreased with the increase of transport distance, and the decreasing
503 rates became slower (Figure S7). The changes of dry and wet depositions caused by emission
504 reductions from EUR and RBU were still obvious in the Arctic region (66°N–90°N), while
505 depositions caused by emission reductions from the other regions tended to be gentle (Figure S7).
506 This explains why the contribution from EAS to BC at different latitudes remained almost constant
507 while that from EUR and RBU decreased obviously from lower latitudes to the Arctic pol
508




Figure 6. Contributions of 20% emission reductions of different regions to near-surface BC concentrations in each latitudinal band of the Arctic. The results of summer and winter correspond to the left and right panel in the figure.

Figure 7 further depicts the response of the vertical Arctic BC profiles in different latitudinal bands to 20% emission reductions. The contributions of eight latitudinal bands showed a typical bimodal pattern in summer with peaks at 0.6–1.6 km a.s.l. (3rd – 5th layers) and 8.0–8.9 km a.s.l. (13th and 14th layers), while the contribution displayed a single peak at the 0.4–1.0 km a.s.l. (2nd – 4th layers) in winter. Similar to section 3.3.2, the peak value of the contribution at the low layers was due to the transport of EAS, EUR, NAM, and RBU emission reductions to the Arctic through different pathways both in summer and winter. The peak value in the high layers in summer was due to the transport of EAS and SAS. However, a high contribution of 20% emission reductions to BC concentrations in SAS was found in the high layers, while the contribution was low in other regions, leading to a single peak in winter. The statistical results of SAS indicated that the contribution in vertical appeared one peak at the 15th layer (9.7 km a.s.l.) with values of 0.45 and 0.48 ng m⁻³ in summer and winter, respectively (Figure S8).




Figure 7. Contributions of 20% emission reductions from all six source regions to the vertical BC concentrations of the Arctic in different latitude bands varies with vertical layers in (a) summer and (b) winter in 2010.


As same as the whole Arctic region (Section 3.3.1 & 3.3.2), the contributions of 20% emission reductions to BC concentrations in eight latitude bands were higher in winter than in summer, whether near surface or in vertical. The contribution of 20% emission reductions from all six source regions to BC concentrations in eight latitude bands of the Arctic near surface was 0.7–1.9 ng m^{-3} in summer and 1.8–4.1 ng m^{-3} in winter, respectively (Figure 6). The high BC peak at around 0.6–1.6 km a.s.l. (3rd – 5th layers) was 1.1–2.1 ng m^{-3} in summer, and 2.9–4.2 ng m^{-3} in winter (Figure 7).

3.4 Benefit of BC emission reductions on the decrease of Arctic temperature

The impact of BC emission reductions on decreasing the Arctic (60–90°N) surface temperature was assessed by using ARTP (See methods in Section 2.2). Aerosol effects, BC deposition on snow, and BC semi-direct were considered in the calculation of ARTP (Aamaas et al., 2017). As shown in Figure 8, the response of Arctic surface temperature to emission reductions was the most significant at the time scale of 10 years and then gradually decreased with the passage of time. For each source region, the Arctic temperature response was significantly higher in winter than in summer as ARTP was seasonal dependent with higher values in the colder seasons. Obviously, the Arctic surface temperature benefited the most from BC emission reductions from EAS with more than 300 and 660 μK decreases in summer and winter after 10 years, respectively. The influences of EUR and NAM emission reductions on the temperature decrease were similar in summer, reaching about 3–90 μK

544 after 10, 20, 50, and 100 years. However, in winter, the influence of emission reductions from NAM
 545 on temperature decrease (8–200 μK) was weaker than that from EUR (14–370 μK). This was mainly
 546 due to the difference of ARTPs between EUR and NAM was not obvious compared with the
 547 difference of emission reductions from NAM and EUR in summer and winter. The responses of the
 548 temperature decrease to emission reductions from RBU were 9–20 μK in summer and 4–100 μK in
 549 winter after 10–100 years, respectively, which were smaller than that from EUR and NAM. This can
 550 be explained by the low BC emission reductions from RBU (Table 1). The response of the
 551 temperature decrease to emission reductions from SAS in winter (10–320 μK) was similar to that
 552 from EUR, while this response in summer (8–230 μK) was more than twice that of EUR. Although
 553 the ARTP of EUR was higher than that of SAS, the BC emission reductions from SAS were much
 554 higher than that from EUR and the difference between emission reductions from the two regions was
 555 more obvious in summer (Table 1). In spite of the higher Arctic temperature response to EAS than
 556 SAS in the target year of this study, a number of studies have shown that BC emissions in South Asia
 557 were increasing in recent years (Sahu et al., 2008; Paliwal et al., 2016; Sharma et al., 2019) while the
 558 emissions of East Asia were exhibiting a downward trend especially from China (Chen et. al., 2016),
 559 thus it should be given more attention to the impact assessment of South Asia on the Arctic in the
 560 future. The minimum temperature response was found from MDE due to the least emission
 561 reductions and small ARTP.


562



563 **Figure 8.** Arctic surface temperature response to 20% regional BC emission reductions in (a) summer and (b)
 564 winter after 10, 20, 50, and 100 years.

565 In addition, the impacts of BC emission reductions from six source regions on the Arctic and

566 global surface temperature were compared in this study (Figure 9). Due to the BC emission
 567 reductions from the six source regions, the surface temperature in the Arctic decreased 27–780 μK in
 568 summer and 61–1675 μK in winter after 10, 20, 50, and 100 years, which were higher than that of
 569 the global with the values of 10–290 μK in summer and 16–470 μK in winter. It can be seen that the
 570 difference of the temperature response between the Arctic and the globe was more obvious in winter.
 571 Overall, the response of the Arctic surface temperature was more sensitive to emissions perturbation
 572 than that of the globe surface temperature.
 573



574 **Figure 9.** Global and Arctic surface temperature responses to 20% regional BC emission reductions in **(a)** summer
 575 and **(b)** winter after 10, 20, 50, and 100 years.

576 It should be noted the estimation of temperature response was subject to large uncertainties for the
 577 following reasons. On the one hand, even though the HTAP2 emissions database were all constructed
 578 by bottom-up methods, the different inventories and spatiotemporal distributions were constructed
 579 with sub-regional (country, state, county or province level) activity data and emission factors, which
 580 lead to inconsistencies at the borders between two adjacent inventories. The version 5 of Evaluating
 581 the Climate and Air Quality Impacts of Short-Lived Pollutants (ECLIPSEv5, <http://eclipse.nilu.no>)
 582 estimated a 2010 emission inventory, that serves also as a reference point for all projections
 583 (Janssens-Maenhout et al., 2015). At the global level, a relatively good agreement was found with
 584 small relative emission differences compared with the ECLIPSEv5 emission inventory for the
 585 aggregated sectors in 2010. However, larger differences of 29% between HTAP2 and ECLIPSEv5
 586 emissions was present for BC since ECLIPSEv5 relied on provincial statistics for China which
 587 resulted from higher coal consumption than reported national statistics. Hoesly et al. (2018) provided

588 a sectoral and gridded historical (1750–2014) anthropogenic emission inventory for use in the
589 Coupled Model Intercomparison Project phase 6 (CMIP6). The amount of global BC anthropogenic
590 emissions was 7.7 Tg/year in 2010 from the CMIP6 emissions, which was larger than that from
591 HTAP2 emissions (5.5 Tg/year). This was mainly due to the energy, transportation, and international
592 shipping sectors of CMIP6 were higher than those of HTAP2.

593 On the other hand, the time evolution of R_T , a parameter in the calculation of ARTP was also one
594 factor causing the uncertainty of temperature response calculation. This impulse response function
595 was only based on one coupled atmosphere-ocean climate model GISS-ER in this study, while Olivié
596 and Peters (2013) have found a spread in the GTP (20) value of BC of about –60 to +80% due to
597 variability of R_T among various models. However, the uncertainty in R_T was less relevant for the
598 regional patterns. Forcing-response coefficients didn't exist on a seasonal basis since emissions
599 occurring during Northern Hemisphere summer and winter season were differentiated (Aamaas et al,
600 2017). Hence, the seasonal differences presented here in the ARTP values were not due to potential
601 differences in the response sensitivities, but due to differences in the RF. The temperature response
602 will vary by species and location, such as between land surface and ocean surface. These differences
603 are not accounted for in this study, but the increased efficacy in the RCS matrix towards the NH can
604 be partly attributed to a larger land area fraction in the NH (Shindell et al., 2015). Besides, recent
605 studies have found that the positive radiation budget of BC being largely compensated for by rapid
606 atmospheric adjustment, this means that the responses of surface temperatures to BC tends to be
607 weaker than expected (Stjern et al., 2017; Takemura and Suzuki, 2019).

608 Although the HTAP2 emissions database contain uncertainties and ARTP calculations are
609 simplifications, these emission metrics are useful, simple, and quick approximations for calculating
610 the temperature response in the different latitude bands for emissions of BC. It should be noted that
611 the estimated responses of Arctic surface temperature to 20% emission reductions were only valid for
612 the comparison among different source regions but cannot be used to reflect the actual change of
613 temperature. On the one hand, in reality, not all emissions sectors of a specific source region cannot
614 be reduced by 20% at the same time. On the other hand, there were many other factors (e.g.
615 greenhouse gases, sea ice coverage) that can affect the temperature change in the Arctic besides BC.

616 4. Conclusions

617 The CAMchem, CHASER_re1, GEOS-Chem, GOCART, and Oslo CTM3 in HTAP2 experiment
618 were used in this study to estimate the responses of Arctic BC to multi-region emission reductions in
619 2010. Six regions (e.g., EAS, EUR, MDE, RBU, NAM, and SAS) were selected as the source
620 regions and the Arctic was the receptor region. HTAP2 set up the base scenario with all BC
621 emissions, and also simulated BC concentrations with 20% reduction of anthropogenic emissions.
622 The AGPT was further used to calculate the benefit of BC emission reductions on the decrease of
623 Arctic temperature.

624 The statistical results of 20% BC emission reductions showed that emission reductions in EAS
625 were the largest with the values of 355.6 Gg yr⁻¹, followed by SAS (232.5 Gg yr⁻¹), EUR (65.3 Gg
626 yr⁻¹), NAM (62.2 Gg yr⁻¹), RBU (18.6 Gg yr⁻¹), and MDE (5.3 Gg yr⁻¹). The BC emission
627 reductions in the EAS, EUR, and RBU were higher from November to March.

628 The temporal variations of simulations from different models were relatively consistent as the
629 correlations of the simulated BC concentrations among different models ranged from 0.33 to 0.98.
630 However, the simulated BC concentrations didn't agree so well with observations at monitoring sites
631 except Zeppelin. In order to reduce the difference of simulation performance of each model in
632 different areas of the Arctic, the model ensemble mean was used for analysis.

633 The contribution of 20% BC emission reductions from EAS, EUR, MDE, NAM, RBU, and SAS
634 to the Arctic near-surface BC concentrations reached 0.70, 0.54, 0.01, 0.20, 0.29, and 0.09 ng m⁻³,
635 respectively. Correspondingly, the reduced column BC loadings from the six regions above over the
636 Arctic was 8521.7, 2789.1, 28.8, 1762.1, 998.6, and 3640.2 ng m⁻², respectively.

637 The response of Arctic near-surface BC concentrations to 20% emission reductions from EAS and
638 EUR was larger than other four source regions, with the monthly value of 0.2–1.5 ng m⁻³ and 0.2–1.0
639 ng m⁻³, accounting for 16.8%–49.0% and 20.1%–49.0% of the total contributions from all six
640 regions, respectively. The BC profiles displayed a bimodal pattern in summer with peaks at around
641 1.0–1.6 km a.s.l. (4th and 5th layers) and 8.0–8.9 km a.s.l. (13th and 14th layers). While the BC profiles
642 showed a unimodal pattern with peaks around 0.6–1.6 km a.s.l. (3rd – 5th layers) in winter.

643 The response of Arctic BC to emission reductions from source regions in winter was higher than

644 that in summer. The contributions of 20% emission reductions to the Arctic BC concentrations near
645 surface were the highest between 66–69°N both in summer (1.9 ng m^{-3}) and winter (4.1 ng m^{-3}), and
646 became weaker with the increase of the latitude.

647 The response of Arctic temperature to BC emission reductions was the most significant at the time
648 scale of 10 years and then gradually decreased with the passage of time. The Arctic had benefited the
649 most from emission reduction in EAS with more than 300 and 660 μK decreases in summer and
650 winter after 10 years, respectively. The Arctic temperature response was more sensitive to the whole
651 globe in regard of the same emissions perturbation. The estimation of temperature response was
652 subject to large uncertainties due to the uncertainties in the calculation of ARTP and emissions of BC
653 in source regions.

654 Overall, this study provided insights on the source regions and seasonal contributions of Arctic BC
655 from the most recent international ensemble modeling efforts. The discrepancy between model
656 results and observations and the spread among different HTAP models may be attributed to various
657 factors such as emissions in the remote Arctic, physical parameterizations, and convection and
658 deposition processes. This would subsequently result in large uncertainties of the climatic effects of
659 air pollutants. More observation sites on the typical transport pathways from sources regions to the
660 Arctic should be planned to improve the model capability of simulating the transport behavior of
661 black carbon.

662 **Data availability**

663 All data used in this paper can be obtained through the AeroCom servers and web interfaces,
664 accessible at <http://aerocom.met.no>.

666 **Author contributions**

667 KH and JSF designed this study. ML, KS, DH, TK, MC, and ST performed modeling. NZ analyzed
668 data and wrote the paper. All have commented on and reviewed the paper.

669

670 **Competing interests.**

671 The authors declare that they have no conflict of interest.

672

673 **Acknowledgements**

674 We sincerely thank for the HTAPv2 international initiative. This work was partially supported by
675 the National Key R&D Program of China (2018YFC0213105), the National Natural Science
676 Foundation of Shanghai (18230722600), and the National Natural Science Foundation of China
677 (91644105).

678

679 **References**

- 680 Aamaas, B., Berntsen, T. K., Fuglestedt, J. S., Shine, K. P., and Bellouin, N.: Regional emission metrics for
681 short-lived climate forcers from multiple models, *Atmos. Chem. Phys.*, 16, 7451– 7468,
682 <https://doi.org/10.5194/acp-16-7451-2016>, 2016.
- 683 Aamaas, B., Berntsen, T. K., Fuglestedt, J. S., Shine, K. P., and Collins, W. J.: Regional temperature change
684 potentials for short-lived climate forcers based on radiative forcing from multiple models, *Atmos. Chem. Phys.*,
685 17, 10795-10809, 10.5194/acp-17-10795-2017, 2017.
- 686 AMAP: Black carbon and ozone as Arctic climate forcers. Arctic Monitoring and Assessment Programme (AMAP),
687 Oslo, Norway. vii + 116 pp, 2015.
- 688 AMAP: Snow, Water, Ice and Permafrost in the Arctic (SWIPA) 2017. Arctic Monitoring and Assessment
689 Programme (AMAP), Oslo, Norway. xiv + 269 pp. ISBN 978–82–7971–101–8, 2017.
- 690 AMAP: The Impact of Black Carbon on Arctic Climate. By: Quinn, P. K., Stohl, A., Arneth, A., Berntsen, T.,
691 Burkhardt, J. F., Christensen, J., Flanner, M., Kupiainen, K., Lihavainen, H., Shepherd, M., Shevchenko, V.,
692 Skov, H., and Vestreng, V., Arctic Monitoring and Assessment Programme (AMAP), Oslo. 72 pp, 2011.
- 693 AMAP: The Impact of Short-Lived Pollutants on Arctic Climate. By: Quinn, P. K., Bates, T. S., Baum, E., Bond, T.,
694 Burkhardt, J. F., Fiore, A. M., Flanner, M. G., Garrett, T., Koch, D., McConnell, J. R., Shindell, D., and Stohl, A.,
695 Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway, 2008.
- 696 Barrie, L. A.: Arctic air pollution: an overview of current knowl- edge, *Atmos. Environ.*, 20, 643–663, 1986.
- 697 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S.,
698 Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M.,
699 Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z.,
700 Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender,
701 C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys.*
702 *Res.–Atmos.*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.
- 703 Bozem, H., Hoor, P., Kunkel, D., Köllner, F., Schneider, J., Herber, A., Schulz, H., Leaitch, W. R., Aliabadi, A. A.,
704 Willis, M. D., Burkart, J., and Abbatt, J. P. D.: Characterization of transport regimes and the polar dome during
705 Arctic spring and summer using in situ aircraft measurements, *Atmos. Chem. Phys.*, 19, 15049-15071,
706 10.5194/acp-19-15049-2019, 2019.
- 707 Bradley, R. S., Keimig, F. T., and Diaz, H. F.: Climatology of surface-based inversions in the North American
708 Arctic, *J. Geophys. Res.*, 97, 15 699, <https://doi.org/10.1029/92JD01451>, 1992.
- 709 Chen, D. S., Zhao, Y. H., Nelson, P., Li, Y., Wang, X. T., Zhou, Y., Lang, J. L., and Guo, X. R.: Estimating ship
710 emissions based on AIS data for port of Tianjin, China. *Atmos. Environ.* 145: 10–18.
711 <http://dx.doi.org/10.1016/j.atmosenv.2016>.
- 712 Cheng, G.: Analysis of observational data of atmospheric boundary layer characteristics in the Arctic (in Chinese),
713 Nanjing University of information engineering. 2011.
- 714 Chin, M., Rood, R. B., Lin, S.-J., Müller, J.-F., and Thompson, A. M.: Atmospheric sulfur cycle simulated in the
715 global model GOCART: Model description and global properties, *J. Geophys. Res.–Atmos.*, 105,
716 24671–24687, doi:10.1029/2000JD900384, 2000.
- 717 Clarke, A. D. and Noone, K. J.: Soot in the Arctic snowpack: a cause for perturbations in radiative transfer, *Atmos.*
718 *Environ.*, 19, 2045–2053, [https://doi.org/10.1016/0004-6981\(85\)90113-1](https://doi.org/10.1016/0004-6981(85)90113-1), 1985.
- 719 Collins, W. J., Fry, M. M., Yu, H., Fuglestedt, J. S., Shindell, D. T., and West, J. J.: Global and regional

720 temperature-change potentials for near-term climate forcers, *Atmos. Chem. Phys.*, 13, 2471–2485,
 721 10.5194/acp-13-2471-2013, 2013.

722 Eckhardt, S., Stohl, A., Beirle, S., Spichtinger, N., James, P., Forster, C., Junker, C., Wagner, T., Platt, U., and
 723 Jennings, S. G.: The North Atlantic Oscillation controls air pollution transport to the Arctic, *Atmos. Chem.*
 724 *Phys.*, 3, 1769–1778, <https://doi.org/10.5194/acp-3-1769-2003>, 2003.

725 Flanner, M. G.: Arctic climate sensitivity to local black carbon, *J. Geophys. Res.-Atmos.*, 118, 1840–1851,
 726 <https://doi.org/10.1002/jgrd.50176>, 2013.

727 Fuglestad, J. S., Shine, K. P., Berntsen, T., Cook, J., Lee, D. S., Stenke, A., Skeie, R. B., Velders, G. J. M., and
 728 Waitz, I. A.: Transport impacts on atmosphere and climate: metrics, *Atmos. Environ.*, 44, 4648–4677,
 729 <https://doi.org/10.1016/j.atmosenv.2009.04.044>, 2010.

730 Galmarini, S., Koffi, B., Solazzo, E., Keating, T., Hogrefe, C., Schulz, M., Benedictow, A., Griesfeller, J. J.,
 731 Janssens-Maenhout, G., Carmichael, G., Fu, J., and Dentener, F.: Technical note: Coordination and
 732 harmonization of the multi-scale, multi-model activities HTAP2, AQMEII3, and MICS-Asia3: simulations,
 733 emission inventories, boundary conditions, and model output formats, *Atmos. Chem. Phys.*, 17, 1543–1555,
 734 2017.

735 Garrett, T. J. and Zhao, C.: Increased Arctic cloud longwave emissivity associated with pollution from mid-latitudes,
 736 *Nature*, 440, 787–789, 2006.

737 Hansen, J. and Nazarenko, L.: Soot climate forcing via snow and ice albedos, *P. Natl Acad. Sci. USA*, 101,
 738 423–428, <https://doi.org/10.1073/pnas.2237157100>, 2004.

739 Henze, D. K., Hakami, A., and Seinfeld, J. H.: Development of the adjoint of GEOS–Chem, *Atmos. Chem. Phys.*, 7,
 740 2413–2433, [doi:10.5194/acp-7-2413-2007](https://doi.org/10.5194/acp-7-2413-2007), 2007.

741 Hoesly R. M., Smith S. J., Feng L. Y., Klimont Z., Janssens-Maenhout G., Pitkanen T., Seibert J. J., Vu L., Andres R.
 742 J., Bolt R. M., Bond T. C., Dawidowski L., Kholod N., Kurokawa J., Li M., Liu L., Lu Z. F., Moura M. C. P.,
 743 O’Rourke P. R., Zhang Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols
 744 from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369–408,
 745 10.5194/gmd-11-369-2018, 2018.

746 Hogrefe, C., Liu, P., Pouliot, G., Mathur, R., Roselle, S., Flemming, J., Lin, M., and Park, R. J.: Impacts of different
 747 characterizations of large-scale background on simulated regional-scale ozone over the continental United
 748 States, *Atmos. Chem. Phys.*, 18, 3839–3864, 2018.

749 Huang, K., Fu, J. S., Hodson, E. L., Dong, X., Cresko, J., Prikhodko, V. Y., Storey, J. M., and Cheng, M.-D.:
 750 Identification of missing anthropogenic emission sources in Russia: Implication for modeling Arctic haze,
 751 *Aerosol Air Qual. Res.*, 14, 1799–1811, 2014.

752 Huang, K., Fu, J. S., Prikhodko, V. Y., Storey, J. M., Ro- manov, A., Hodson, E. L., Cresko, J., Morozova, I., Ig-
 753 natieva, Y., and Cabaniss, J.: Russian anthropogenic black carbon: Emission reconstruction and Arctic black
 754 carbon simulation, *J. Geophys. Res.-Atmos.*, 120, 11306–11333, <https://doi.org/10.1002/2015JD023358>, 2015.

755 IPCC: Climate Change: The Intergovernmental Panel on Climate Change Scientific Assessment. The
 756 Intergovernmental Panel on Climate Change (IPCC), Cambridge University Press, Cambridge, UK, 1990.

757 IPCC: Climate Change 2001: The Scientific Basis. Intergovernmental Panel on Climate Change. The
 758 Intergovernmental Panel on Climate Change (IPCC), Cambridge University Press, Cambridge, UK, 2001.

759 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q.,
 760 Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S.,
 761 Koffi, B., and Li, M.: HTAP_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to
 762 study hemispheric transport of air pollution, *Atmos. Chem. Phys.*, 15, 11411–11432,
 763 10.5194/acp-15-11411-2015, 2015.

764 Jonson, J. E., Schulz, M., Emmons, L., Flemming, J., Henze, D., Sudo, K., Tronstad Lund, M., Lin, M., Benedictow,

765 A., Koffi, B., Dentener, F., Keating, T., Kivi, R., and Davila, Y.: The effects of intercontinental emission
 766 sources on European air pollution levels, *Atmos. Chem. Phys.*, 18, 13655–13672, 2018.

767 Koch, D. and Hansen, J.: Distant origins of Arctic black carbon: A Goddard Institute for Space Studies ModelE
 768 experiment, *J. Geophys. Res.–Atmos.*, 110, D04204, <https://doi.org/10.1029/2004JD005296>, 2005.

769 Lamarque, J. F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A.,
 770 Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and
 771 evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*,
 772 5, 369–411, 10.5194/gmd-5-369-2012, 2012.

773 Law, K. S. and Stohl, A.: Arctic air pollution: origins and impacts, *Science*, 315, 1537–1540,
 774 <https://doi.org/10.1126/science.1137695>, 2007.

775 Liang, C.-K., West, J. J., Silva, R. A., Bian, H., Chin, M., Davila, Y., Dentener, F. J., Emmons, L., Flemming, J.,
 776 Folberth, G., Henze, D., Im, U., Jonson, J. E., Keating, T. J., Kucsera, T., Lenzen, A., Lin, M., Lund, M. T., Pan,
 777 X., Park, R. J., Pierce, R. B., Sekiya, T., Sudo, K., and Takemura, T.: HTAP2 multi-model estimates of
 778 premature human mortality due to intercontinental transport of air pollution and emission sectors, *Atmos.*
 779 *Chem. Phys.*, 18, 10497–10520, 2018.

780 Lund, M. T., Berntsen, T. K., Heyes, C., Klimont, Z., and Samset, B. H.: Global and regional climate impacts of
 781 black carbon and co-emitted species from the on-road diesel sector, *Atmos. Environ.*, 98, 50–58,
 782 <https://doi.org/10.1016/j.atmosenv.2014.08.033>, 2014.

783 Lund, M. T., Aamaas, B., Berntsen, T., Bock, L., Burkhardt, U., Fuglestad, J. S., and Shine, K. P.: Emission
 784 metrics for quantifying regional climate impacts of aviation, *Earth Syst. Dynam.*, 8, 547–563,
 785 <https://doi.org/10.5194/esd-8-547-2017>, 2017.

786 Lund, M. T., Myhre, G., Haslerud, A. S., Skeie, R. B., Griesfeller, J., Platt, S. M., Kumar, R., Myhre, C. L., and
 787 Schulz, M.: Concentrations and radiative forcing of anthropogenic aerosols from 1750 to 2014 simulated with
 788 the Oslo CTM3 and CEDS emission inventory, *Geosci. Model Dev.*, 11, 4909–4931,
 789 <https://doi.org/10.5194/gmd-11-4909-2018>, 2018.

790 Marelle, L., Thomas, J. L., Raut, J.-C., Law, K. S., Jalkanen, J.-P., Johansson, L., Roiger, A., Schlager, H., Kim, J.,
 791 Reiter, A., and Weinzierl, B.: Air quality and radiative impacts of Arctic shipping emissions in the summertime
 792 in northern Norway: from the local to the regional scale, *Atmos. Chem. Phys.*, 16, 2359–2379,
 793 <https://doi.org/10.5194/acp-16-2359-2016>, 2016.

794 Matsui, H., Kondo, Y., Moteiki, N., Takegawa, N., Sahu, L. K., Zhao, Y., Fuelberg, H. E., Sessions, W. R., Diskin, G.,
 795 Blake, D. R., Wisthaler, A., Koike, M.: Seasonal variation of the transport of black carbon aerosol from the
 796 Asian continent to the Arctic during the ARCTAS aircraft campaign, *J. Geophys. Res.–Atmos.*, 116, D05202,
 797 [doi:10.1029/2010JD015067](https://doi.org/10.1029/2010JD015067).

798 Miao, Y. C., Guo, J. P., Liu, S. H., Liu, H., Zhang, G., Yan, Y., He, J.: Relay transport of aerosols to
 799 Beijing–Tianjin–Hebei region by multiscale atmospheric circulations, *Atmos. Environ.*, 165, 35–45,
 800 <https://doi.org/10.1016/j.atmosenv.2017.06.032>, 2017.

801 Myhre, G., Shindell, D., Breon, F.-M., Collins, W., Fuglestad, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D.,
 802 Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and
 803 Natural Radiative Forcing, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working*
 804 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by:
 805 Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and
 806 Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 2013.

807 Olivie, D. J. L. and Peters, G. P.: Variation in emission metrics due to variation in CO₂ and temperature impulse
 808 response functions, *Earth Syst. Dynam.*, 4, 267–286, <https://doi.org/10.5194/esd-4-267-2013>, 2013.

809 Paliwal, U., Sharma, M., and Burkhardt, J. F.: Monthly and spatially resolved black carbon emission inventory of

810 India: uncertainty analysis, *Atmos. Chem. Phys.*, 16, 12457–12476, 10.5194/acp-16-12457-2016, 2016.

811 Sahu, S. K., Beig, G., and Sharma, C.: Decadal growth of black carbon emissions in India, *Geophysical Research*
812 *Letters*, 35, 10.1029/2007gl032333, 2008.

813 Samset, B. H., Myhre, G., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Diehl, T.,
814 Easter, R. C., Ghan, S. J., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J. F., Lin, G., Liu, X., Penner, J. E.,
815 Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Black carbon vertical profiles
816 strongly affect its radiative forcing uncertainty, *Atmos. Chem. Phys.*, 13, 2423–2434,
817 10.5194/acp-13-2423-2013, 2013.

818 Sand, M., Bernsten, T. K., von Salzen, K., Flanner, M. G., Langner, J., and Victor, D. G.: Response of Arctic
819 temperature to changes in emissions of short-lived climate forcers, *Nature Clim. Change*, 6, 286–289,
820 10.1038/nclimate2880, 2016.

821 Sekiya, T., Miyazaki, K., Ogochi, K., Sudo, K., and Takigawa, M.: Global high-resolution simulations of
822 tropospheric nitrogen dioxide using CHASER V4.0, *Geosci. Model Dev.*, 11, 959–988,
823 10.5194/gmd-11-959-2018, 2018.

824 Sharma, S., Ishizawa, M., Chan, D., Lavoué, D., Andrews, E., Eleftheriadis, K., and Maksyutov, S.: 16-year
825 simulation of Arctic black carbon: Transport, source contribution, and sensitivity analysis on deposition,
826 *Journal of Geophysical Research: Atmospheres*, 118, 943–964, 10.1029/2012jd017774, 2013.

827 Sharma, G., Sinha, B., Pallavi, Hakkim, H., Chandra, B. P., Kumar, A., and Sinha, V.: Gridded Emissions of CO,
828 NO_x, SO₂, CO₂, NH₃, HCl, CH₄, PM_{2.5}, PM₁₀, BC, and NMVOC from Open Municipal Waste Burning in
829 India, *Environ Sci Technol*, 53, 4765–4774, 10.1021/acs.est.8b07076, 2019.

830 Shine, K. P., Fuglestad, J. S., Hailemariam, K., and Stuber, N.: Alternatives to the Global Warming Potential for
831 Comparing Climate Impacts of Emissions of Greenhouse Gases, *Climatic Change*, 68, 281–302,
832 10.1007/s10584-005-1146-9, 2005

833 Shindell, D. and Faluvegi, G.: Climate response to regional radiative forcing during the 20th century, *Nat. Geosci.*,
834 2, 294–300, 2009.

835 Shindell, D. and Faluvegi, G.: The net climate impact of coal-fired power plant emissions, *Atmos. Chem. Phys.*, 10,
836 3247–3260, <https://doi.org/10.5194/acp-10-3247-2010>, 2010.

837 Shindell D., Kuylenstierna J.C.I., Vignati E., van Dingenen R., Amann M., Klimont Z., Anenberg S.C., Muller N.,
838 JanssensMaenhaut G., Raes F., Schwartz J., Faluvegi G., Pozzoli L., Kupiainen K., Höglund–Isaksson L.,
839 Emberson L., Streets D., Ramanathan V., Hicks K., Kim Oanh N.T., Milly G., Williams M., Demkine W.,
840 Fowler D.: Simultaneously Mitigating Near-Term Climate Change and Improving Human Health and Food
841 Security. *Science*, Vol. 335, 183–189, 2012.

842 Shindell, D. T., Chin, M., Dentener, F., Doherty, R. M., Faluvegi, G., Fiore, A. M., Hess, P., Koch, D. M.,
843 MacKenzie, I. A., Sanderson, M. G., Schultz, M. G., Schulz, M., Stevenson, D. S., Teich, H., Textor, C., Wild,
844 O., Bergmann, D. J., Bey, I., Bian, H., Cuvelier, C., Duncan, B. N., Folberth, G., Horowitz, L. W., Jonson, J.,
845 Kaminski, J. W., Marmer, E., Park, R., Pringle, K. J., Schroeder, S., Szopa, S., Takemura, T., Zeng, G., Keating,
846 T. J., Zuber, A.: A multi-model assessment of pollution transport to the Arctic, *Atmos. Chem. Phys. Discuss.*,
847 2008.

848 Shindell, D. T., Faluvegi, G., Rotstain, L., and Milly, G.: Spatial patterns of radiative forcing and surface
849 temperature response, *J. Geophys. Res.-Atmos.*, 120, 5385–5403, <https://doi.org/10.1002/2014JD022752>,
850 2015.

851 Shindell, D. T., Lamarque, J.–F., Schulz, M., Flanner, M., Jiao, C., Chin, M., Young, P. J., Lee, Y. H., Rotstain, L.,
852 Mahowald, N., Milly, G., Faluvegi, G., Balkanski, Y., Collins, W. J., Conley, A. J., Dalsoren, S., Easter, R.,
853 Ghan, S., Horowitz, L., Liu, X., Myhre, G., Nagashima, T., Naik, V., Rumbold, S. T., Skeie, R., Sudo, K.,
854 Szopa, S., Takemura, T., Voulgarakis, A., Yoon, J.–H., and Lo, F.: Radiative forcing in the ACCMIP historical

and future climate simulations, *Atmos. Chem. Phys.*, 13, 2939–2974, 2013.

Shindell, D. T., Voulgarakis, A., Faluvegi, G., and Milly, G.: Precipitation response to regional radiative forcing, *Atmos. Chem. Phys.*, 12, 6969–6982, doi:10.5194/acp-12-6969-2012, 2012.

Smith, S.J. and Mizrahi, A.: Near-term climate mitigation by short-lived forcers. *PNAS* 14202–14206, doi: 10.1073/pnas.1308470110, 2013.

Sobhani, N., Kulkarni, S., Carmichael, G. R.: Source sector and region contributions to black carbon and PM_{2.5} in the Arctic. *Atmos. Chem. Phys.*, 18, 18123–18148, 2018.

Stjern, C. W., Samset, B. H., Myhre, G., Bian, H., Chin, M., Davila, Y., Dentener, F., Emmons, L., Flemming, J., Haslerud, A. S., Henze, D., Jonson, J. E., Kucsera, T., Lund, M. T., Schulz, M., Sudo, K., Takemura, T., and Tilmes, S.: Global and regional radiative forcing from 20 % reductions in BC, OC and SO₄ – an HTAP2 multi-model study, *Atmos. Chem. Phys.*, 16, 13579–13599, <https://doi.org/10.5194/acp-16-13579-13599>, 2016.

Stjern, C. W. et al. Rapid adjustments cause weak surface temperature response to increased black carbon concentrations. *Geophys. Res.* 122, 11462–11481, 2017.

Stohl, A.: Characteristics of atmospheric transport into the Arctic troposphere, *J. Geophys. Res.*, 111, D11306, <https://doi.org/10.1029/2005JD006888>, 2006.

Stohl A., Aamaas B., Amann M., Baker L. H., Bellouin N., Bernsten T. K., Boucher O., Cherian R., Collins W., Daskalakis N., Dusinska M., Eckhardt S., Fuglestedt J. S., Harju M., Heyes C., Hodnebrog Ø., Hao J., Im U., Kanakidou M., Klimont Z., Kupiainen K., Law K. S., Lund M. T., Maas R., MacIntosh C. R., Myhre G., Myriokefalitakis S., Olivie D., Quaas J., Quennehen B., Raut J.-C., Rumbold S. T., Samset B. H., Schulz M., Seland Ø., Shine K. P., Skeie R. B., Wang S., Yttri K. E., and Zhu T.: Evaluating the climate and air quality impacts of short-lived pollutants. *Atmos. Chem. Phys.*, 15, 10529 – 10566, 2015.

Stohl, A., Eckhardt, S., Forster, C., James, P., and Spichtinger, N.: On the pathways and timescales of intercontinental air pollution transport, *J. Geophys. Res.–Atmos.*, 107, ACH 6–1–ACH 6–17, <https://doi.org/10.1029/2001JD001396>, 2002.

Stohl, A., Klimont, Z., Eckhardt, S., Kupiainen, K., Shevchenko, V. P., Kopeikin, V. M., and Novigatsky, A. N.: Black carbon in the Arctic: the underestimated role of gas flaring and residential combustion emissions, *Atmos. Chem. Phys.*, 13, 8833–8855, <https://doi.org/10.5194/acp-13-8833-2013>, 2013

Sudo, K., Sekiya, T., Nagashima, T.: CHASER/MIROC-ESM in HTAP2 status reports, HTAP2 Global and Regional Model Evaluation Workshop, Nagoya University, JAMSTEC, NIES, 2015.

Sudo, K., Takahashi, M., Kurokawa, J.-I., and Akimoto, H.: CHASER: A global chemical model of the troposphere 1. Model description, *J. Geophys. Res.–Atmos.*, 107, ACH 7–1–ACH 7–20, doi:10.1029/2001JD001113, 2002.

Søvde, O. A., Prather, M. J., Isaksen, I. S. A., Bernsten, T. K., Stordal, F., Zhu, X., Holmes, C. D., and Hsu, J.: The chemical transport model Oslo CTM3, *Geosci. Model Dev.*, 5, 1441–1469, doi:10.5194/gmd-5-1441-2012, 2012.

Takemura, T., and Suzuki, K.: Weak global warming mitigation by reducing black carbon emissions, *Scientific Reports*, 9, 10.1038/s41598-019-41181-6, 2019.

Tan, J., Fu, J. S., Dentener, F., Sun, J., Emmons, L., Tilmes, S., Sudo, K., Flemming, J., Jonson, J. E., Gravel, S., Bian, H., Davila, Y., Henze, D. K., Lund, M. T., Kucsera, T., Takemura, T., and Keating, T.: Multi-model study of HTAP II on sulfur and nitrogen deposition, *Atmos. Chem. Phys.*, 18, 6847–6866, 2018a.

Tan, J., Fu, J. S., Dentener, F., Sun, J., Emmons, L., Tilmes, S., Flemming, J., Takemura, T., Bian, H., Zhu, Q., Yang, C.-E., and Keating, T.: Source contributions to sulfur and nitrogen deposition – an HTAP II multi-model study on hemispheric transport, *Atmos. Chem. Phys.*, 18, 12223–12240, 2018b.

Teng, H., Washington, W. M., Branstator, G., Meehl, G. A., and Lamarque, J.-F.: Potential impacts of Asian carbon

900 aerosols on future US warming, *Geophys. Res. Lett.*, 39, L11703, doi:10.1029/2012GL051723, 2012.

901 Tilmes, S., Lamarque, J.-F., Emmons, L. K., Kinnison, D. E., Marsh, D., Garcia, R. R., Smith, A. K., Neely, R. R.,
 902 Conley, A., Vitt, F., Val Martin, M., Tanimoto, H., Simpson, I., Blake, D. R., and Blake, N.: Representation of
 903 the Community Earth System Model (CESM1) CAM4-chem within the Chemistry- Climate Model Initiative
 904 (CCMI), *Geosci. Model Dev.*, 9, 1853– 1890, doi:10.5194/gmd-9-1853-2016, 2016.

905 Twomey, S.: The influence of pollution on the shortwave albedo of clouds, *J. Atmos. Sci.*, 34, 1149–1152, 1977.

906 UNEP/WMO: Integrated Assessment of Black Carbon and Tropospheric Ozone, Nairobi, Kenya, available at:
 907 <http://wedocs.unep.org/handle/20.500.11822/8028> (last access: 13 May 2020), 2011.

908 US EPA: Guidance on the use of models and other analyses for demonstrating attainment of air quality goals for
 909 ozone, PM_{2.5}, and regional haze. U.S. Environmental Protection Agency Office of Air Quality Planning and
 910 Standards Air Quality Analysis Division Air Quality Modeling Group Research Triangle Park, NC, 2007.

911 Zhang, H. Y., Cheng, S. Y., Yao, S., Wang, X. Q., and Zhang, J. F.: Multiple perspectives for modeling regional
 912 PM_{2.5} transport across cities in the Beijing–Tianjin–Hebei region during haze episodes, *Atmos. Environ.*, 212,
 913 22–35, <https://doi.org/10.1016/j.atmosenv.2019.05.031>, 2019.

914 Zhang, X., Zhang, Y. H., Han, J. B., Zhang, L., and Shi, W. R.: Analysis of the climatic characteristics of the
 915 atmospheric boundary layer height in yanmayan Island, Arctic (in Chinese), *Polar studies*, 30(02):132–139,
 916 2018.