



Analysis of Secondary Organic Aerosol Simulation Bias in the Community Earth System Model (CESM2.1)

Yaman Liu^{1,2}, Xinyi Dong^{1,2}, Minghuai Wang^{1,2}, Louisa K. Emmons³, Yawen Liu^{1,2}, Yuan Liang^{1,2}, Xiao Li^{1,2}, Manish Shrivastava⁴

5 ¹School of Atmospheric Science, Nanjing University, Nanjing, China

²Joint International Research Laboratory of Atmospheric and Earth System Sciences & Institute for Climate and Global Change Research, Nanjing University, China

³National Center for Atmospheric Research, Boulder, CO, USA

⁴Pacific Northwest National Laboratory, Richland, Washington, USA

10 *Correspondence to:* Xinyi Dong (dongxy@nju.edu.cn)

Abstract. Organic aerosol (OA) has been considered as one of the most important uncertainties in climate modeling due to the complexity in presenting its chemical production and depletion mechanisms. To better understand the capability of climate models and probe into the associated uncertainties in simulating OA, we evaluate the Community Earth System Model version 2.1 (CESM2.1) configured with the Community Atmosphere Model version 6 (CAM6) with comprehensive tropospheric and stratospheric chemistry representation (CAM6-Chem), through a long-term simulation (1988–2019) with observations collected from multiple datasets in the United States. We find that CESM generally reproduces the inter-annual variation and seasonal cycle of OA mass concentration at surface layer with correlation of 0.40 as compared to ground observations, and systematically overestimates (69 %) in summer and underestimates (-19 %) in winter. Through a series of sensitivity simulations, we reveal that modeling bias is primarily related to the dominant fraction of monoterpene-formed secondary organic aerosol (SOA), and a strong positive correlation of 0.67 is found between monoterpene emission and modeling bias in eastern US during summer. In terms of vertical profile, the model prominently underestimates OA and monoterpene concentrations by 37–99 % and 82–99 % respectively in the upper air (>500 m) as validated against aircraft observations. Our study suggests that the current Volatility Basis Set (VBS) scheme applied in CESM might be parameterized with too high monoterpene SOA yields which subsequently result in strong SOA production near emission source area. We also find that the model has difficulty in reproducing the decreasing trend of surface OA in southeast US, probably because of employing pure gas VBS to represent isoprene SOA which is in reality mainly formed through multiphase chemistry, thus the influence of aerosol acidity and sulfate particle change on isoprene SOA formation has not been fully considered in the model. This study reveals the urgent need to improve the SOA modeling in climate models.

1 Introduction

30 As one of the most important contributors (20 %–90 %) to total fine atmospheric particles (Kanakidou et al., 2004), organic aerosol (OA) plays an important role in the climate system by affecting the radiation budget (Ghan et al., 2012). OA consists



of primary organic aerosol (POA, also called primary organic matter POM) emitted directly from biomass burning, fossil fuels combustion, biological compounds, etc., and secondary organic aerosol (SOA) formed via oxidation of volatile organic compounds (VOCs) (Hallquist et al., 2009;Tsigaridis et al., 2014;Shrivastava et al., 2017). Chamber studies have revealed the important role of biogenic VOCs such as monoterpenes (Docherty and Ziemann, 2003;Kristensen et al., 2016;Signorell and Bertram, 2009) and isoprene (Kroll et al., 2005;Nguyen et al., 2010;Nguyen et al., 2014;Paulot et al., 2009) in SOA production, and the contribution of anthropogenic VOCs such as aromatic compounds, emitted from vehicle emissions and solvents, also has important influence in urban areas (Nakao et al., 2012;Sato et al., 2010).

Understanding the atmospheric burden and spatiotemporal distributions of OA is one of the key priorities in atmospheric research because of the central roles it played in regulating both climate and air quality, but our current modeling knowledge of OA retains significant uncertainty. The radiative forcing effect of OA has been assessed with climate models through tremendous efforts during the past decades, yet the limited capability of climate models in terms of simulating the productions and depletions of OA induce large uncertainties. Substantial divergences were reported for models employed in the framework of Aerosol Comparisons between Observations and Models (AeroCom) phase II project even with the same set of emissions input. The burden of OA varied greatly for 28 models in the range of 0.6 ~ 3.8 Tg, with OA lifetime ranging from 3.8 to 9.6 days (Tsigaridis et al., 2014). As OA loading and properties of aerosols varied, the estimated radiative forcing of OA ranged from -0.06 to -0.01 W/m² among the 16 participating models (Myhre et al., 2013), revealing the fundamental uncertainty of OA simulation.

Modeling discrepancies largely come from the lack of a consensus in the representation of chemical composition and formation processes of OA among different models. Despite the fact that chamber studies have been continuously revealing new reactions and identifying new species, model representation of SOA lags far behind in terms of the chemical production, depletion, deposition velocity, aging, and other processes. The reactions observed in chambers are condensed into a few functions, and the various products are lumped to fewer species in the models for unclear SOA formation processes and to save computational cost. Most of OA undergoes chemical aging via SOA formation (Zhang et al., 2007) with volatility and hygroscopicity changing, but such processing is poorly understood due to inadequate relevant observations. In the remote areas of United States (U.S.), OA and organic carbon (OC) concentrations show opposite bias in half of participating models (Tsigaridis et al., 2014), indicating the models lack a consensus representation of SOA production and depletion.

To reveal the uncertainties associated with OA simulation in climate models, we evaluate a recent version of Community Earth System Model version 2.1 (CESM2.1) in this study with multiple observational datasets in the U.S. The model has been widely applied for OA climate effect assessment purpose and a significant portion of improvements have been implemented in the latest version regarding the chemical mechanisms (Tilmes et al., 2019), but evaluation of simulated OA concentration hasn't been well documented or thoroughly discussed. We focus on the validation over the U.S. because it has long-term surface measurements and flight campaigns that provide solid observation data. We first evaluate the spatiotemporal characteristics of the simulation bias, and then probe into the chemical mechanism to identify the origins through a series of sensitivity runs,



65 and finally demonstrate the urgent need to both improve the current parameterization of the SOA production scheme and
implement a more comprehensive production mechanism in climate model.

2 Methods

2.1 Model

70 Community Earth System Model is a coupled Earth System model composed of atmosphere, ocean, land, sea-ice, land-ice,
river, and wave models (Danabasoglu et al., 2020). CESM2 (versions 2.0 and 2.1) includes updated atmospheric chemistry in
the Community Atmosphere Model version 6 with chemistry (CAM6-Chem) over previous versions. The Model for Ozone
and Related chemical Tracers (MOZART) chemical mechanism covering the troposphere and stratosphere (referred to as
MOZART-TS1) is used in CAM6-Chem. Emmons et al. (2020) reported the updates of MOZART-TS1 in CESM2.1, including
75 the oxidation of isoprene and terpenes, organic nitrate speciation, and aromatic speciation and oxidation, and thus improved
representation of ozone (O₃) and SOA precursors. The most recently released CESM2.2 includes a new version (TS2) of
MOZART tropospheric chemical mechanism with updates for isoprene and terpene chemistry (Schwantes et al., 2020) aiming
at further improving O₃ simulation. In this study we also briefly compare the results between TS1 and TS2 as will be discussed
in Sect. 3.3.


Both biogenic and anthropogenic VOCs are considered in CESM2.1 with improved gas-phase chemical mechanisms (Emmons
80 et al., 2020) and new SOA representation (Tilmes et al., 2019). CESM2.1 applies the Volatility Basis Set (VBS) scheme
(Bergström et al., 2012; Hodzic et al., 2016; Shrivastava et al., 2011; Shrivastava et al., 2013) by lumping SOA precursors based
on their volatility bins to simulate SOA production. In CAM6-Chem, SOA and the gas-phase condensable sources (SOAG)
are categorized into five bins with the saturation concentration (C*) of 0.01, 0.1, 1.0, 10.0, and 100.0 µg/m³ (Tilmes et al.,
2019), respectively. Compared with the simple SOA scheme which proportionally calculated SOAG based on emissions of
85 precursors (Liu et al., 2012), this VBS scheme was demonstrated to improve CESM performance with smaller bias of OA
concentration over remote regions when evaluated against aircraft observations (Shrivastava et al., 2015; Tilmes et al., 2019).
VBS approach relies on empirical parameterizations fitting to chamber experiments thus the parameters vary between models.
The current CAM6-Chem configures VBS scheme and parameters by following the work of Hodzic et al. (2016) with GEOS-
Chem as the host model which differs significantly from CAM-Chem in terms of SOA-related modules such as gas-phase
90 chemistry, aerosol dynamics, dry and wet depositions. Consequently, the same scheme and parameterization employed by
Hodzic et al. (2016) may result in different performance within CAM-Chem. Tilmes et al. (2019) provided a comprehensive
comparison between simple SOA scheme and the CESM2.1 VBS scheme in terms of simulated SOA burden and radiative
forcing, and the simulations were validated against two flight campaigns, yet the evaluation against ground surface
measurements at different temporal scale (e.g., annual, seasonal) over different geophysical areas hasn't been thoroughly
95 discussed.



2.2 Observations

Surface measurements from the Interagency Monitoring of Protected Visual Environments (IMPROVE) network and multiple aircraft campaigns in northern hemisphere are incorporated in this study to validate model performance and also facilitate the analysis of SOA trend. IMPROVE is a long-existing program currently managed by the U.S. Environmental Protection Agency (EPA), and is designed to measure chemical composition of ambient fine particles and its spatial and temporal information (Solomon et al., 2014). We used 1988–2019 daily data for seasonal cycle and long-term trend evaluation. As IMPROVE measures organic carbon (OC) mass instead of OA, the concentration of OA observation is derived with the ratio of OA to OC (OA/OC) which is determined by the aging process. IMPROVE data has been widely employed in many studies for model evaluation purpose (Hodzic et al., 2016; Shrivastava et al., 2015; Tsigaridis et al., 2014). The OA/OC of IMPROVE data varies between sites and the mean value used in this study is 1.8 (1.79 ~ 2.02) as recommended by Malm and Hand (2007). It should be noticed that not all sites have observations for the whole study period, and thus only those (140 sites, locations shown in Fig.1) with more than 10-years continuous data are used in this study to avoid measurement bias.

1988–2019 surface OA simulation bias



110 **Figure 1: 1998–2019 CAM-Chem-SD surface OA concentration mean bias (top figure) and normalized mean bias (bottom figure) compared to IMPROVE data. CONUS is divided into two subdomains, EUS (red box) and WUS (blue box).**



Free troposphere measurements from a total of five aircraft field campaigns are employed to validate simulated vertical OA profiles including CalNex (California Nexus, Ryerson et al. (2013)), DC3 (Deep Convective Clouds and Chemistry, Barth et al. (2015)), SENEX (Southeast Nexus, (Warneke et al., 2016)), SEAC4RS (Studies of Emissions and Atmospheric
115 Composition, Clouds and Climate Coupling by Regional Surveys, (Toon et al., 2016)), FRAPPE (Front Range Air Pollution and Photochemistry Experiment (Flocke et al., 2020). Flights were located in the area of contiguous United States (CONUS) and took place between 2010 and 2014 with more details presented in Table 1 and Fig. 3.

Table 1: Aircraft measurements used in this study

Campaign	Dates	Region
CalNex (Ryerson et al., 2013)	2010.4.30~ 6.22	California and the eastern Pacific coastal region
DC3 (Barth et al., 2015)	2012.5.1~ 6.30	northeastern Colorado, west Texas to central Oklahoma, and northern Alabama
SENEX (Warneke et al., 2016)	2013.6.01~ 7.10	southeast U.S.
SEAC4RS (Toon et al., 2016)	2013.8.6~ 9.24	southeast and west US
FRAPPE (Flocke et al., 2020;Dingle et al., 2016)	2014.7.26~ 8.19	northern front range metropolitan area (central U.S.)

120

2.3 Simulation configurations

To reproduce the observed meteorological conditions and allow direct comparison with OA measurements, a specified dynamic simulation (SD) is conducted from 1987 to 2019 with a spin-up time of one year thus our discussion can focus on
125 chemical mechanism performance of CESM2.1. This experiment (referred as CAM-Chem-SD) uses FCSD component set. Temperature, horizontal winds, and surface fluxes are nudged to Modern-Era Retrospective analysis for Research and Applications (MERRA2) fields. The horizontal resolution is $0.9^\circ \times 1.25^\circ$ and vertical resolution is 32 levels with model top at ~45 km (Emmons et al., 2020). Prescribed historical sea surface temperatures (SSTs) are used in the FCSD component set. Anthropogenic and biomass burning emissions from 1987 to 2014 are from the standard Coupled Model Intercomparison
130 Project round 6 (CMIP6) (Eyring et al., 2016) simulations, and emissions after 2014 are from SSP585 scenario which is based on the shared socioeconomic pathway 5 (SSP5) (O'Neill et al., 2017) and forcing levels of Representative Concentration Pathways 8.5 (RCP8.5). Biogenic emissions are calculated with the Model of Emissions of Gases and Aerosol from Nature (MEGAN) in CESM (Guenther et al., 2012;Emmons et al., 2020).



135 The current VBS scheme in CESM2.1 represents SOAG production from nine precursors through 15 reactions. The 15
reactions and chemical formulas of related species from Emmons et al. (2020) are shown at Table S1 and S2. To identify their
contributions to total SOA mass and associated simulation uncertainties, we conducted 15 sensitivity experiments with one of
the reactions turned off in each experiment. The 15 sensitivity simulations and a 14-month baseline simulation are set up from
January 2010 to evaluate the contribution of each reaction to total SOA production in VBS scheme. To exclude the influence
of potential extreme meteorology condition or emission inputs, these sensitivity runs are configured with FC2010climo
140 component set and Newtonian relaxation time of three hours, and the FC2010climo component set is as same as FCSD
component set except that the emissions are a 10-year average used for each year of the simulation. As one reaction is turned
off for each sensitivity experiment, the difference between base and the sensitivity experiment represents the influence of this
specific reaction (Table 2). For instance, the no ISOP-OH-SOAG experiment excludes SOAG produced by ISOP + OH
reaction (reaction 1 in Table S2) with other configurations same as the base run (CAM-Chem-climo) experiment.

145

Table 2: CESM experiments used in this study

Index	Experiment ID
B1	CAM-Chem-SD
B2	CAM-Chem-climo
E1	no ISOP+OH
E2	no ISOP+O3
E3	no ISOP+NO3
E4	no MTERP+OH
E5	no MTERP+O3
E6	no MTERP+NO3
E7	no BCARY+OH
E8	no BCARY+O3
E9	no BCARY+NO3
E10	no BENZENE+OH
E11	no TOLUENE+OH
E12	no XYLENES+OH
E13	no IVOC+OH
E14	no SVOC+OH
E15	no GLYOXAL



3 Results and Discussion

3.1 Evaluation against IMPROVE

150 We first validate the surface OA simulation with IMIPROVE data for the CAM-Chem-SD scenario and find an overall significant overestimation, with the results and statistics presented in Fig. 1 and Table 3. As demonstrated in Fig. 1, CESM2.1 shows critical simulation bias with normalized mean bias (NMB) varying from -73.87 % to 176.47 % at different sites with daily data pairs against IMPROVE measurements. The model noticeably overestimates annual average surface OA concentration over continental U.S. (CONUS) with mean bias (MB) of $0.41 \mu\text{g}/\text{m}^3$ and NMB of 20.27 %. It also shows large regional difference as demonstrated by the sharp contrast between the two subdomains, eastern US (EUS, $65^\circ\text{--}95^\circ\text{W}$, $25^\circ\text{--}49^\circ\text{N}$) and western US (WUS, $95^\circ\text{--}125^\circ\text{W}$, $25^\circ\text{--}49^\circ\text{N}$) as presented in Fig. 1. Prominent overestimation is found over EUS with NMB more than 100 % at nine sites, while moderate underestimation is shown over WUS. Our validation suggests that the updated SOA representation (by applying VBS) helps improve the performance of CESM as compared with its earlier versions. CAM4-Chem was reported to underestimate OC by $\sim 36\%$ compared with IMPROVE dataset with correlation coefficient of 0.41 (Lamarque et al., 2012), and CAM5-Chem was found to overestimate with NMB by 24 % at urban sites and 217 % at remote sites (Tsigaridis et al., 2014). The AeroCom phase II multi-model ensemble mean showed NMB of -48 % (range -85 % to 24 %) at urban locations and -70 % (range -38 % to 217 %) at remote locations on global scale, suggesting that the performance of CESM2.1 is consistent with other climate models.

Annual and seasonal variations of surface OA from the model and IMPROVE are presented in Fig. 2 to examine the model's capability of reproducing the temporal variation. The model generally reproduces the annual decreasing trend of OA concentration for the whole CONUS domain, and is in good agreement with observation in spring and fall, but substantially overestimates in summer and slightly underestimates in winter as shown in Fig. 2(a). Monthly average of the simulation is found to have larger overestimation in warmer months (May–Sep.) than cooler months as shown in Fig. 2(b). In EUS, the simulation prominently overestimates in summer by $4.26 \mu\text{g}/\text{m}^3$ (131.15 %) but successfully reproduces the temporal change with a strong correlation with observations of 0.60 as shown in Fig. 2(c). As compared with CONUS domain, simulation at EUS shows an even greater overestimation during warmer months as shown in Fig. 2(d). In WUS, simulated OA shows a slow descending trend in summer ($-0.02 \mu\text{g}/\text{m}^3$ per decade) while the observations indicate an ascending trend ($0.23 \mu\text{g}/\text{m}^3$ per decade) with large inter-annual variation from 1999 to 2019 mainly due to the influence of wildfires (Malm et al., 2017). The model shows smaller bias in WUS but also a poor correlation of 0.36 in summer as shown in Fig. 2(e) and 2(f).


175 The large modeling bias of OA in summer over EUS may be attributed to the discrepancy in simulating SOA, because POA is proportionally determined by emissions from fossil fuel and biofuel that have smaller seasonal diversity (see supplementary material Fig. S1). Biomass burning also contributes POA, but wildfire or prescribed fire emission is minimal in EUS (van der Werf et al., 2017). Production of SOA is represented by the MOZART-TS1 and VBS module as mentioned in Sect. 2. MOZART-TS1 has been demonstrated in a few studies (Emmons et al., 2020) to overestimate summertime surface O_3 over southeast US, thus it may subsequently induce uncertainties while simulating the VOCs chemistry. Besides, the performance

180



and uncertainty of the VBS scheme in CESM2 remain largely undocumented regarding whether the VBS scheme tends to overestimate or underestimate. With the same VBS scheme but implemented in GEOS-Chem, Hodzic et al. (2016) reported simulation bias of 34.8 % through validation against IMPROVE data, and the bias was explained by evaporation of OC from IMPROVE samples (Kim et al., 2015) and uncertainties in the boundary layer parameterization. The performance of CESM2.1 seems better as our simulation bias for CONUS domain is 20.27 %. But with the larger biases in simulated OA spatial and temporal trends shown in Fig. 1 and Fig. 2, it is critical to realize that the overall evaluation statistic is averaged from large overestimations and underestimations from sites across the CONUS domain.

surface OA concentration ($\mu\text{g}/\text{m}^3$)



190 **Figure 2:** The left column is 1988–2019 annual mean (black lines) and seasonal average (blue lines for winter, green lines for spring, red lines for summer, and yellow lines for autumn) surface OA concentration of IMPROVE (solid dots) and CAM-Chem-SD (solid lines) over US continent (a), eastern U.S.(c), and western U.S.(e). The right column is seasonal cycle of 1988–2019 average surface OA concentration of IMPROVE (blue dots) and CAM-Chem-SD (red dots) over CONUS (b), EUS (d) and WUS (f). Every blue box denotes the 10th, the 25th, the 75th and the 90th percentiles of the observations for all selected sites in each month.



195 **Table 3: The correlation coefficient (CC), mean bias (MB) and normalized mean bias (NMB) between observations (five field campaigns and IMPROVE surface measurements) and CAM-Chem-SD.**


Observations		CC	Mean Obs.	Mean Sim.	MB	NMB (%)
IMPROVE						
CONUS	Annual	0.40	2.07	2.48	0.41	20.27
	Spring	0.67	1.65	1.57	-0.08	-4.81
	Summer	0.37	2.90	4.87	1.97	68.78
	Fall	0.34	2.23	2.23	-0.01	0.13
	Winter	0.70	1.49	1.22	-0.27	-19.05
EUS	Annual	0.79	2.64	3.72	1.08	40.82
	Spring	0.64	2.49	2.66	0.17	6.71
	Summer	0.60	3.26	7.52	4.26	131.15
	Fall	0.69	2.63	2.88	0.25	9.70
	Winter	0.82	2.25	1.82	-0.43	-19.11
WUS	Annual	0.36	1.78	1.89	0.11	10.49
	Spring	0.77	1.23	1.04	-0.19	-15.36
	Summer	0.48	2.72	3.66	0.94	34.83
	Fall	0.35	2.03	1.91	-0.12	-5.85
	Winter	0.73	1.12	0.91	-0.21	-18.00
Aircraft						
CalNex		0.43	2.06	1.46	-0.60	-29.01
DC3		0.12	2.99	0.72	-2.17	-72.75
SETEX		0.33	7.09	4.22	-2.87	-40.54
SEAC4RS		0.10	6.90	1.93	-4.97	-71.97
FRAPPE		0.27	3.05	2.42	-0.63	-20.64

3.2 Evaluation against flight campaigns

200 We find systematic underestimation in the upper air through the model evaluation against Aerosol Mass Spectrometry (AMS) measurements from flight campaigns, with the profiles shown in Fig. 3 and evaluation statistics summarized in Table 1. These campaigns were conducted over land area of North America during 2010–2014 and most of the measurements were collected



in summer with rest of the data in late spring and early fall. Simulations are paired with measurements by choosing the nearest model grid at the corresponding time at hourly scale. OA concentrations are underestimated in upper air (>500 m) by -20 % – -70 % as compared with the aircraft measurements. The model shows larger bias when compared with SENEX, DC3 and SEAC4RS that mostly cover EUS, while the vertical profiles measured in WUS (CalNex and SENEX) are reproduced better with stronger correlation coefficient and smaller bias. The most prominent bias is found in the comparison with DC3 campaign at ~7 km (geometric altitude above mean sea level) due to the aircraft samples of smoke plume from the High Park fire west of Ft. Collins (Barth et al., 2015), suggesting that fire emission plume structure is not properly represented because of the coarse model grid resolution. In general, the model is found to notably underestimate upper air OA for all flight campaigns with larger discrepancy over EUS. We also examine the simulated and observed isoprene and monoterpene profiles during the SEAC4RS campaign, as shown in Fig. 3(g). These two biogenic VOCs (BVOCs) are reproduced well between surface and ~2 km altitude, but are also substantially underestimated at >2 km height, suggesting too fast oxidation of these VOCs at high altitudes or uncertainties in parameterizations of convective transport and vertical mixing. Validation against the recent ATom (Atmospheric TOMography aircraft campaign) measurements from Wofsy et al. (2018) also suggested underestimation in the upper air over marine areas (see supplementary material Fig. S2 and Table S3), consistent with the recent multi-model evaluation results reported in Hodzic et al. (2020). The photolytic depletion of SOA is represented in VBS scheme of CESM by following Hodzic et al. (2016) which demonstrated this stronger removal process can help lower the GEOS-Chem model overestimation of upper air OA over remote areas (mostly over ocean). But the underestimation of high-altitude SOA within our CESM2.1 model over CONUS land area are different from GEOS-Chem, and are likely related to model-to-model differences in physics, aerosol chemistry, and wet removal.



225 **Figure 3: Flight tracks of five aircraft campaigns, and vertical profiles of average OA (solid lines, panels b-f), ISOP (dashed lines, panel g) and MTERP (dotted lines, panel g) concentration from CAM-Chem-SD (black), CalNex (blue), SENEX (purple), FRAPPE (green), DC3 (orange) and SEAC4RS (red) flight campaigns. The range of OA concentration at each layer is showed as the shaded area or dashed lines.**



3.3 Uncertainties in VBS schemes


This section reveals the contributions from different SOA precursors and probes into the uncertainties within the current VBS
230 scheme in CESM. We first investigate the seasonality and diagnose the components of simulated OA to narrow down the most
important candidates causing the modeling bias. Figure 4(a)–(c) present the CAM-Chem-climo simulated surface OA
concentration with the contributions from POA and five bins of SOA (SOA1–SOA5) over CONUS, EUS, and WUS. On
annual average scale POA consists 35.6 % of the total OA with larger contribution in winter (~50 %), while SOA dominates
in summer (~77.1 %) over CONUS. Our CESM2.1 simulation suggests that the overall ambient OA concentration is
235 determined by SOA for CONUS domain, which is also consistent with observation-based studies (Yu et al., 2004; Zhang et al.,
2018). Figure 4(a) shows that the absolute contribution from POA is relatively less dynamic than SOA through the year
although both show higher values in summer, suggesting that the seasonality of surface OA is primarily determined by SOA.
The five volatility bins show consistent seasonality with high concentrations in warmer months but the two least volatile bins
SOA1 and SOA2 show more prominent seasonal variations, and these two bins also contribute most to total SOA. The
240 dominant influences of SOA1 and SOA2 on total OA presented by CESM is consistent with other VBS-based models (Farina
et al., 2010).

We then examine the contribution from each of the 15 VBS reactions (listed in Table S1) to identify the most influential species
over CONUS, EUS, and WUS as shown in Fig. 4(d)–(f). The SOA formed by each reaction is calculated as the difference
between CAM-Chem-climo and the corresponding sensitivity experiment described in Table 2. For instance, $SOA_{MTERP+O_3}$ is
245 calculated by the difference between SOA from CAM-Chem-climo scenario and SOA from “no MTERP+O₃” scenario. The
sensitivity simulations demonstrate dominant contributions over CONUS and EUS is from 3 BVOC-formed SOA, lumped
monoterpenes (MTERP), followed by isoprene (ISOP) and sesquiterpenes (BCARY), in summer, while the intermediate
volatility organic compounds (IVOCs) and aromatics (BENZENE, TOLUENE, and XYLENES) are mainly emitted by
anthropogenic sources and play a more important role in winter. The contribution from the 15 reactions over WUS show
250 similar results except that SOA formed by ISOP show less contribution compared to SOA formed by BCARY due to less
 $SOA_{ISOP+OH}$ in summer. SOA formed from aromatics also show much smaller seasonal variation as compared with BVOC-
formed SOA because anthropogenic emissions are relatively stable throughout the year, while biogenic emissions are
substantially more intensive during the growing season than other periods. Besides, SOA formed by MTERP dominates total
SOA vertical distribution (~34.5%) as shown in Fig. S3. The sensitivity experiments demonstrate that monoterpene-formed
255 SOA plays the most important role in determining both surface and vertical simulated OA concentration in summer on average
over the CONUS domain.

We further diagnose the SOA concentration in July formed from these reactions because of the prominent seasonality (larger
bias in summer) and regional difference (larger bias in EUS) revealed in earlier sections. Figure 4(g)–(i) present the
contributions from each pathway to the five SOA bins in July over CONUS, EUS, and WUS. Among the 15 pathways to form
260 SOA, the reaction of MTERP+O₃ forms most SOA, followed by MTERP+OH and ISOP+OH over CONUS and EUS, while



SOA formed by MTERP+OH contributes most over WUS, which is consistent with the over estimation of O₃ over EUS in MOZARY-TS1 as mentioned in Emmons et al. (2020). We also find that MTERP-formed SOA is mostly in the second VBS bin SOA2 with low volatility, but most of SOA form by ISOP+OH is in the fourth VBS bin SOA4 with high volatility.



265

270

Figure 4: (a)–(c) Seasonal variation of surface OA concentration from CAM-Chem-climo simulation at all IMPROVE sites located over CONUS (a), EUS (b) and WUS (c). The contribution of POA and SOA in five VBS bins are shown in filled areas. The 14-month average contribution of POA and SOA are shown as the number in the legend. (d)–(f) Seasonal variation of SOA formed by 15 different pathways over CONUS (d), EUS (e) and WUS (f). The contribution of 15 reactions are shown in filled areas and 14-month average contribution is shown as the number in the legend. (g)–(i) the contributions from each pathway to the five SOA bins over CONUS (g), EUS (h) and WUS (i) in 2010 July. SOA concentration in 5 bins are shown in filled bars.

The contribution of monoterpene-derived SOA is consistent with other measurement studies (Xu et al., 2018; Zhang et al., 2018) which demonstrated that a significant fraction of observed OA was monoterpene generated SOA over southeast US. Zhang et al. (2018) indicated monoterpene derived SOA contributed to ~42 % of total AMS PM₁ OA during SOAS (Southern Oxidant and Aerosol Study) field campaign. As Sect. 3.1 reveals substantial overestimation in summer and minor bias in

275



winter, it is thus very likely that monoterpene-related VBS reactions may be primarily responsible for the modeling uncertainty. We have found a strong correlation between monoterpene emission and modeling bias as shown in Fig. 5. The CAM-Chem-SD simulation bias increases along biogenic monoterpene emission with a correlation of 0.41 in summer and -0.12 in winter over the CONUS domain. This relationship is particularly strong (correlation of 0.67) for summer over EUS (Fig. 5(g)) with higher biogenic VOCs, and almost negligible in winter over WUS (correlation of -0.05) with lower biogenic emission. Moreover, we also find biogenic VOCs flux dominates OA bias with higher correlation in summer over EUS, while anthropogenic VOCs flux dominates OA bias over WUS as shown in Fig. S4. Despite the fact that isoprene biogenic emission flux shows the same correlation as monoterpene biogenic emission flux over EUS, the prominent seasonality of modeling bias and the strong correlation with summer time emission suggests that monoterpene SOA dominates the performance of CESM over abundant biogenic emission region.

relationship between OA bias and MTERP emission of CAM-Chem-SD




Figure 5: The relationship between simulated surface OA bias (Y axis) and surface MTERP emissions (X axis) in DJF (a, e, i), MAM (b, f, j), JJA (c, g, k) and SON (d, h, l) over CONUS (a–d), EUS (e–h) and WUS (i–l). The correlation coefficient of OA bias and MTERP emission is shown as the number in the top right of each plot.



The mass yields of VBS in CESM was parameterized following Hodzic et al. (2016) which adjusted the values used in GEOS-Chem (Jo et al., 2013) to account for the wall-loss. These parameters are shown in Table 4. The adjustment by Hodzic et al. (2016) resulted in stronger SOA production and led to overestimation in surface OA concentration compared with AMS global network and biogenic sources contributing to the total SOA production. It is certainly reasonable to take the wall-loss effect into account while applying the chamber measurements, but it also should be noticed those measurements were conducted under artificial environment with predefined chemical species that may vary significantly from the real meteorology condition and atmospheric chemistry regime, thus chamber study reported parameters need to be carefully interpreted and adjusted while being applied in atmospheric model. The current VBS scheme in CESM2.1 differs from Hodzic et al. (2016) by considering the differences of oxidants but doesn't consider the NO_x dependence of monoterpene oxidation. The mass yields applied in CESM2.1 are generally in the middle of low and high NO_x conditions used by Hodzic et al. (2016), but still show substantial overestimation due to monoterpene SOA as mentioned above, suggesting that the mass yields in CESM2.1 require further adjustment.

Table 4: Mass yield coefficient of monoterpene in Jo et al. (2013), Hodzic et al. (2016) and CAM6-Chem in VBS bins. C* is the saturation concentration (µg/m³).

Log(C*)	Condition Oxidant	Jo et al. (2013)		Hodzic et al. (2016)		CAM6-Chem	
		Low NO _x OH, O ₃	NO ₃	low NO _x OH, O ₃ , NO ₃	high NO _x	- OH, O ₃	NO ₃
≤-2				0.093	0.045	0.0508	
-1				0.211	0.015	0.1149	
0		0.01	0.07	0.064	0.142	0.0348	
1		0	0.06	0.102	0.061	0.0554	0.17493
2		0.54	0.24	0.110	0.074	0.1278	0.59019
3		0	0.41	0.125	0.165		

This simplified representation of VBS parameterization also affects simulation bias because BVOCs have been demonstrated to be closely influenced by NO_x concentration (Shrivastava et al., 2017). Since high NO_x usually suppresses the formation of monoterpene SOA, observed OA would be lower at places with high NO_x concentration than those with low NO_x. Subsequently, the overestimation due to monoterpene SOA would be more severe at places with high NO_x level. Fig. 6 (a)–(d) presents the simulation bias at two typical IMPROVE sites representing the high (Agua Tibia: 33.5 °N, 117.0 °W) and low (Lake Sugema: 40.7 °N, 92.0 °W) NO_x conditions respectively. With comparable levels of monoterpene emissions, simulation bias at Agua



315 Tibia is substantially higher than that at the Lake Sugema especially during Sep. and Oct. when NO_x concentrations are maximum. We further investigate the relationship between OA bias and NO_x concentration at low MTERP biogenic emission as shown in Fig. 6(e). OA bias increases with higher NO_x concentration at same MTERP biogenic emission in summer at 53 sites. A recent study (Jo et al., 2020) reported the development of CESM for implementing the NO_x -dependent yields, along with several other updates including isoprene epoxydiols (IEPOX) derived SOA through heterogeneous chemistry, isoprene
320 emission adjustment, biogenic VOCs deposition, and more detailed gas-phase chemistry of isoprene. We also evaluated simulation from CESM2.2 (configured with MOZART-TS2 which includes the above-mentioned updates except for heterogeneous chemistry of IEPOX-SOA) against IMPROVE observations through a full year simulation (2013.03 - 2014.02). We find that these updates moderately lower the simulated OA concentration, but surface OA is still substantially overestimated in summer. The observed OA was $2.4 \mu\text{g}/\text{m}^3$ at IMPROVE sites over the CONUS domain in July, 2013, while
325 simulated OA by CESM2.1 and CESM2.2 were 5.1 and $4.2 \mu\text{g}/\text{m}^3$, respectively (Fig. S5). Thus, we suggest that the monoterpene SOA yield in current CESM2.1 and CESM2.2 might be configured too high and lead to surface OA overestimation, and the VBS parameterization for other BVOCs in CESM may also need further adjustment for reducing OA bias.

In addition to the monoterpene parameterization, representation of isoprene SOA by pure gas-phase VBS may also induce
330 critical uncertainty to the model simulation, since over certain regions like the southeast USA, isoprene SOA is mostly formed from IEPOX via irreversible multiphase chemistry that is closely affected by gas-aqueous phase transfer and acid catalysed reactive uptake, for which neither has been considered in CESM2.1. A few recent studies have revealed that IEPOX-SOA is influenced by anthropogenic emission because of the changes in inorganic aerosol acidity and sulfate particles (Shrivastava et al., 2019;Zheng et al., 2020). Both modeling and measurement evidence have suggested that the steadily decreasing
335 anthropogenic NO_x and SO_2 emission have led to the reduction of biogenic SOA over western and southeastern US during the past decade, yet CESM has difficulty to reproduce this trend as shown in Fig. 7. The IMPROVE data shows a decreasing trend of $0.527 \mu\text{g}/\text{m}^3$ per decade at Cape Romain NWR, but CAM-Chem-SD simulated trend is $0.098 \mu\text{g}/\text{m}^3$ per decade. Ridley et al. (2018) indicated the reduction of anthropogenic emissions, due to EPA regulations on vehicular sources and power generation, can explain more than two-thirds of decline in OA over CONUS from 1990–2012. Marais et al. (2017) showed
340 SO_2 emissions controlled biogenic OA concentration by sulfate-BVOCs interaction in summer over EUS.





Figure 6: 2010 observed surface OA concentration (red dots) and simulated POA (dark filled area) and SOA (gray filled area) at Agua Tiba site (33.5 °N, 117.0 °W) (a) and Lake Sugema site (40.7 °N, 92.0 °W) (b); simulated surface NO_x concentration (blue lines) and MTERP emission from MEGAN (green lines) at Agua Tiba site (c) and Lake Sugema site (d). 1988–2019 summertime surface OA bias (colorful dots) at specific NO_x concentration and MTERP emission flux (e) over 53 sites where MTERP emission flux are between 3×10^{-10} and 6×10^{-10} .

345



350 **Figure 7: 1988–2019 JJA (a) simulated surface OA decade trend and IMPROVE surface OA decade trend at selected IMPROVE sites (filled circles). Black circles indicate sites with statistically significant trends with 95 % confidence according to the student's T-test; (b) simulated (red lines) and observed (red dots) surface OA concentration at Cape Romain NWR site (shown as purple circle in Fig. 7(a)). Simulated and observed OA decade trends in summer at the site are shown as the numbers in the legend.**

4 Summary and Conclusions

The goal of this study is to understand the performance of CESM2.1 and reveal the remaining biases for the simulation of SOA. Through validation against surface measurements and flight campaigns over the US, we have found that CESM2.1 is able to capture inter-annual and seasonal variation of surface OA concentration with a correlation coefficient by 0.41, but systematically overestimates surface OA concentration in summer by 68.78 %. Larger summer time bias (overestimates by 131.15 %) is found over eastern US (overestimates by 40.82 %) where BVOCs emissions are more intensive than western US. Opposite to the overestimation near the surface, consistent underestimations by -20 % – -70 % of OA in the upper air are found in the validation against all five flight campaigns.

360 Our analysis suggests that it is likely simulated monoterpene SOA production is parameterized with too high yields and may be the main factor that affects the modeling bias for three reasons: first, monoterpene SOA contributes most (nearly half) to the total SOA in summertime, while other BVOCs or anthropogenic POA cannot explain the severe overestimation. The large contribution of monoterpene SOA simulated by CESM2.1 is consistent with other measurement and modeling studies, but the current VBS configuration adopted from GEOS-Chem may require further adjustment. Second, the simulation bias showed a strong spatiotemporal correlation with monoterpene emission as demonstrated by the large overestimation in summer over eastern US, and larger overestimation of OA is found at places with higher NO_x condition under same monoterpene emissions level. Third, overestimation of OA at surface layer and underestimation of OA and monoterpene in the free troposphere suggests that both the production and photolytic removal processes might be parameterized too strong.

This study highlights the simulated bias of OA concentration over US and the significance of VBS mass yield parameters in simulating SOA concentration with some limitations. The five aircraft campaigns discussed in this study were all conducted in warm months. The simulated performance in winter and the discrepancy between near and above surface OA bias needs further investigation. The updated VBS scheme under different NO_x condition and multiphase isoprene-derived SOA



chemistry have been recently considered in other global and regional models which showed better performance in SOA
simulation (Hodzic et al., 2016; Shrivastava et al., 2019; Jo et al., 2020), revealing the urgent need for improving process-level
375 representations of SOA formation and removal in models.

Code and data availability. The CESM model code is available at
https://www.cesm.ucar.edu/models/cesm2/release_download.html. Observational data for DC3
(<https://doi.org/10.5067/Aircraft/DC3/DC8/Aerosol-TraceGas>), FRAPPE ([https://doi.org/10.5067/Aircraft/DISCOVER-
380 AQ/Aerosol-TraceGas](https://doi.org/10.5067/Aircraft/DISCOVER-AQ/Aerosol-TraceGas)), SEAC4RS (<https://doi.org/10.5067/AIRCRAFT/SEAC4RS/AEROSOL-TRACEGAS-CLOUD>), and
ATom (<https://doi.org/10.3334/ORNLDAAAC/1581>; Wofsy et al., 2018) can be obtained from the NASA LaRC data archive.
CalNex (<https://csl.noaa.gov/groups/csl7/measurements/2010calnex/P3/DataDownload/>) and SENEX
(<https://esrl.noaa.gov/csd/groups/csd7/measurements/>) data are available via the NOAA ESRL data archive. IMPROVE data
are available at <http://vista.cira.colostate.edu/Improve/improve-data/>.

385

Author contributions. XD and Yaman L conceived the idea and designed the model experiments. Yaman L performed the
simulations, conducted the analysis, and wrote the paper. MW, XD, Yawen L and XL provided conceptualization for the paper.
Yuan L, Yawen L and MS provided guidelines for the simulations in CESM v2.1. Yawen L helped in IMPROVE data
downloading and processing. XD, Yaman L, LEK, MS and MW edited the paper.

390

Competing interests: The authors declare that they have no conflict of interest.

Acknowledgements. This work is supported by the National Natural Science Foundation of China [grant numbers 91744208,
41925023, 41575073, and 41621005] and the Ministry of Science and Technology of the People's Republic of China [grant
395 number 2016YFC0200503]. This research was also supported by the Collaborative Innovation Center of Climate Change,
Jiangsu Province. We greatly thank the High Performance Computing Center of Nanjing University for providing the
computational resources used in this work. We acknowledge Pedro Campuzano-Jost and Jose-Luis Jimenez of the University
of Colorado-Boulder for the DC3 AMS aerosol composition data. We acknowledge Anna Middlebrook and Roya Bahreini for
the CalNex dataset. We acknowledge the use of observations from the FRAPPE campaign, which was funded by the National
400 Science Foundation (NSF) and the State of Colorado. We acknowledge Joost de Gouw of the University of Colorado-Boulder
for help with the SENEX AMS data. We acknowledge NASA for providing the SEAC4RS and ATom AMS data.

ReferencesUncategorized References

Barth, M. C., Cantrell, C. A., Brune, W. H., Rutledge, S. A., Crawford, J. H., Huntrieser, H., Carey, L. D., MacGorman, D.,
405 Weisman, M., Pickering, K. E., Bruning, E., Anderson, B., Apel, E., Biggerstaff, M., Campos, T., Campuzano-Jost, P., Cohen,
R., Crounse, J., Day, D. A., Diskin, G., Flocke, F., Fried, A., Garland, C., Heikes, B., Honomichl, S., Hornbrook, R., Huey, L.



- 410 G., Jimenez, J. L., Lang, T., Lichtenstern, M., Mikoviny, T., Nault, B., O'Sullivan, D., Pan, L. L., Peischl, J., Pollack, I.,
Richter, D., Riemer, D., Ryerson, T., Schlager, H., St. Clair, J., Walega, J., Weibring, P., Weinheimer, A., Wennberg, P.,
Wisthaler, A., Wooldridge, P. J., and Ziegler, C.: The Deep Convective Clouds and Chemistry (DC3) Field Campaign, *Bulletin*
of the American Meteorological Society, 96, 1281-1309, 10.1175/bams-d-13-00290.1, 2015.
- Bergström, R., Denier van der Gon, H. A. C., Prévôt, A. S. H., Yttri, K. E., and Simpson, D.: Modelling of organic aerosols
over Europe (2002-2007) using a volatility basis set (VBS) framework: application of different assumptions regarding the
formation of secondary organic aerosol, *Atmospheric Chemistry and Physics*, 12, 8499-8527, 10.5194/acp-12-8499-2012,
2012.
- 415 Danabasoglu, G., Lamarque, J. F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J.,
Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M.,
Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S.,
Kamphenhout, L., Versteine, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D.,
Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand,
420 W. G.: The Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12,
10.1029/2019ms001916, 2020.
- Dingle, J. H., Vu, K., Bahreini, R., Apel, E. C., Campos, T. L., Flocke, F., Fried, A., Herndon, S., Hills, A. J., Hornbrook, R.
S., Huey, G., Kaser, L., Montzka, D. D., Nowak, J. B., Reeves, M., Richter, D., Roscioli, J. R., Shertz, S., Stell, M., Tanner,
D., Tyndall, G., Walega, J., Weibring, P., and Weinheimer, A.: Aerosol optical extinction during the Front Range Air Pollution
and Photochemistry Experiment (FRAPPÉ) 2014 summertime field campaign, Colorado, USA, *Atmospheric Chemistry and*
425 *Physics*, 16, 11207-11217, 10.5194/acp-16-11207-2016, 2016.
- Docherty, K. S., and Ziemann, P. J.: Effects of Stabilized Criegee Intermediate and OH Radical Scavengers on Aerosol
Formation from Reactions of β -Pinene with O₃, *Aerosol Science and Technology*, 37, 877-891, 10.1080/02786820300930,
2003.
- 430 Emmons, L. K., Schwantes, R. H., Orlando, J. J., Tyndall, G., Kinnison, D., Lamarque, J. F., Marsh, D., Mills, M. J., Tilmes,
S., Bardeen, C., Buchholz, R. R., Conley, A., Gettelman, A., Garcia, R., Simpson, I., Blake, D. R., Meinardi, S., and Péron,
G.: The Chemistry Mechanism in the Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling*
Earth Systems, 12, 10.1029/2019ms001882, 2020.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled
435 Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*,
9, 1937-1958, 10.5194/gmd-9-1937-2016, 2016.
- Farina, S. C., Adams, P. J., and Pandis, S. N.: Modeling global secondary organic aerosol formation and processing with the
volatility basis set: Implications for anthropogenic secondary organic aerosol, *Journal of Geophysical Research*, 115,
10.1029/2009jd013046, 2010.
- 440 Flocke, F., Pfister, G., Crawford, J. H., Pickering, K. E., Pierce, G., Bon, D., and Reddy, P.: Air Quality in the Northern
Colorado Front Range Metro Area: The Front Range Air Pollution and Photochemistry Experiment (FRAPPÉ), *Journal of*
Geophysical Research: Atmospheres, 125, 10.1029/2019jd031197, 2020.
- Ghan, S. J., Liu, X., Easter, R. C., Zaveri, R., Rasch, P. J., Yoon, J. H., and Eaton, B.: Toward a Minimal Representation of
Aerosols in Climate Models: Comparative Decomposition of Aerosol Direct, Semidirect, and Indirect Radiative Forcing,
445 *Journal of Climate*, 25, 6461-6476, 10.1175/jcli-d-11-00650.1, 2012.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of
Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling
biogenic emissions, *Geoscientific Model Development*, 5, 1471-1492, 10.5194/gmd-5-1471-2012, 2012.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George,
C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M., Jimenez, J. L., Kiendler-
Schar, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski,
R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmospheric*
Chemistry and Physics Discussions, 9, 3555-3762, 10.5194/acpd-9-3555-2009, 2009.
- 450 Hodzic, A., Kasibhatla, P. S., Jo, D. S., Cappa, C. D., Jimenez, J. L., Madronich, S., and Park, R. J.: Rethinking the global
secondary organic aerosol (SOA) budget: stronger production, faster removal, shorter lifetime, *Atmospheric Chemistry and*
Physics, 16, 7917-7941, 10.5194/acp-16-7917-2016, 2016.



- 460 Hodzic, A., Campuzano-Jost, P., Bian, H., Chin, M., Colarco, P. R., Day, D. A., Froyd, K. D., Heinold, B., Jo, D. S., Katich, J. M., Kodros, J. K., Nault, B. A., Pierce, J. R., Ray, E., Schacht, J., Schill, G. P., Schroder, J. C., Schwarz, J. P., Sueper, D. T., Tegen, I., Tilmes, S., Tsigaridis, K., Yu, P., and Jimenez, J. L.: Characterization of organic aerosol across the global remote troposphere: a comparison of ATom measurements and global chemistry models, *Atmospheric Chemistry and Physics*, 20, 4607-4635, 10.5194/acp-20-4607-2020, 2020.
- Jo, D. S., Park, R. J., Kim, M. J., and Spracklen, D. V.: Effects of chemical aging on global secondary organic aerosol using the volatility basis set approach, *Atmospheric Environment*, 81, 230-244, 10.1016/j.atmosenv.2013.08.055, 2013.
- 465 Jo, D. S., Hodzic, A., Emmons, L. K., Tilmes, S., Schwantes, R. H., Mills, M. J., Campuzano-Jost, P., Hu, W., Zaveri, R. A., Easter, R. C., Singh, B., Lu, Z., Schulz, C., Schneider, J., Shilling, J. E., Wisthaler, A., and Jimenez, J. L.: Future changes in isoprene-epoxydiol-derived secondary organic aerosol (IEPOX-SOA) under the shared socioeconomic pathways: the importance of explicit chemistry, *Atmos. Chem. Phys. Discuss.*, 2020, 1-57, 10.5194/acp-2020-543, 2020.
- 470 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmospheric Chemistry and Physics Discussions*, 4, 5855-6024, 10.5194/acpd-4-5855-2004, 2004.
- Kim, P. S., Jacob, D. J., Fisher, J. A., Travis, K., Yu, K., Zhu, L., Yantosca, R. M., Sulprizio, M. P., Jimenez, J. L., Campuzano-Jost, P., Froyd, K. D., Liao, J., Hair, J. W., Fenn, M. A., Butler, C. F., Wagner, N. L., Gordon, T. D., Welti, A., Wennberg, P. O., Crouse, J. D., St. Clair, J. M., Teng, A. P., Millet, D. B., Schwarz, J. P., Markovic, M. Z., and Perring, A. E.: Sources, 475 seasonality, and trends of southeast US aerosol: an integrated analysis of surface, aircraft, and satellite observations with the GEOS-Chem chemical transport model, *Atmospheric Chemistry and Physics*, 15, 10411-10433, 10.5194/acp-15-10411-2015, 2015.
- Kristensen, K., Watne, Å. K., Hammes, J., Lutz, A., Pettersen, T., Hallquist, M., Bilde, M., and Glasius, M.: High-Molecular Weight Dimer Esters Are Major Products in Aerosols from α -Pinene Ozonolysis and the Boreal Forest, *Environmental Science & Technology Letters*, 3, 280-285, 10.1021/acs.estlett.6b00152, 2016.
- 480 Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from isoprene photooxidation under high-NO_x conditions, *Geophysical Research Letters*, 32, n/a-n/a, 10.1029/2005gl023637, 2005.
- Lamarque, J. F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric 485 chemistry in the Community Earth System Model, *Geoscientific Model Development*, 5, 369-411, 10.5194/gmd-5-369-2012, 2012.
- Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J. F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., 490 Flanner, M. G., and Mitchell, D.: Toward a minimal representation of aerosols in climate models: description and evaluation in the Community Atmosphere Model CAM5, *Geoscientific Model Development*, 5, 709-739, 10.5194/gmd-5-709-2012, 2012.
- Malm, W. C., and Hand, J. L.: An examination of the physical and optical properties of aerosols collected in the IMPROVE program, *Atmospheric Environment*, 41, 3407-3427, 10.1016/j.atmosenv.2006.12.012, 2007.
- 495 Malm, W. C., Schichtel, B. A., Hand, J. L., and Collett, J. L.: Concurrent Temporal and Spatial Trends in Sulfate and Organic Mass Concentrations Measured in the IMPROVE Monitoring Program, *Journal of Geophysical Research: Atmospheres*, 122, 10,462-410,476, 10.1002/2017jd026865, 2017.
- Marais, E., Jacob, D., Turner, J., and Mickley, L.: Evidence of 1991-2013 decrease of biogenic secondary organic aerosol in response to SO₂ emission controls, *Environmental Research Letters*, 12, 10.1088/1748-9326/aa69c8, 2017.
- 500 Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J. F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J. H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmospheric Chemistry and Physics*, 13, 1853-1877, 10.5194/acp-13-1853-2013, 2013.



- 505 Nakao, S., Liu, Y., Tang, P., Chen, C. L., Zhang, J., and Cocker Iii, D. R.: Chamber studies of SOA formation from aromatic hydrocarbons: observation of limited glyoxal uptake, *Atmospheric Chemistry and Physics*, 12, 3927-3937, 10.5194/acp-12-3927-2012, 2012.
- Nguyen, T. B., Bateman, A. P., Bones, D. L., Nizkorodov, S. A., Laskin, J., and Laskin, A.: High-resolution mass spectrometry analysis of secondary organic aerosol generated by ozonolysis of isoprene, *Atmospheric Environment*, 44, 1032-1042, 10.1016/j.atmosenv.2009.12.019, 2010.
- 510 Nguyen, T. B., Coggon, M. M., Bates, K. H., Zhang, X., Schwantes, R. H., Schilling, K. A., Loza, C. L., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Organic aerosol formation from the reactive uptake of isoprene epoxydiols (IEPOX) onto non-acidified inorganic seeds, *Atmospheric Chemistry and Physics*, 14, 3497-3510, 10.5194/acp-14-3497-2014, 2014.
- O'Neill, B. C., Krieglner, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., van Ruijven, B. J., van Vuuren, D. P., Birkmann, J., Kok, K., Levy, M., and Solecki, W.: The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century, *Global Environmental Change*, 42, 169-180, 10.1016/j.gloenvcha.2015.01.004, 2017.
- 515 Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kroll, J. H., Seinfeld, J. H., and Wennberg, P. O.: Isoprene photooxidation: new insights into the production of acids and organic nitrates, *Atmospheric Chemistry and Physics*, 9, 1479-1501, 10.5194/acp-9-1479-2009, 2009.
- 520 Ridley, D. A., Heald, C. L., Ridley, K. J., and Kroll, J. H.: Causes and consequences of decreasing atmospheric organic aerosol in the United States, *Proc Natl Acad Sci U S A*, 115, 290-295, 10.1073/pnas.1700387115, 2018.
- Ryerson, T. B., Andrews, A. E., Angevine, W. M., Bates, T. S., Brock, C. A., Cairns, B., Cohen, R. C., Cooper, O. R., de Gouw, J. A., Fehsenfeld, F. C., Ferrare, R. A., Fischer, M. L., Flagan, R. C., Goldstein, A. H., Hair, J. W., Hardesty, R. M., Hostetler, C. A., Jimenez, J. L., Langford, A. O., McCauley, E., McKeen, S. A., Molina, L. T., Nenes, A., Oltmans, S. J., Parrish, D. D., Pederson, J. R., Pierce, R. B., Prather, K., Quinn, P. K., Seinfeld, J. H., Senff, C. J., Sorooshian, A., Stutz, J., Surratt, J. D., Trainer, M., Volkamer, R., Williams, E. J., and Wofsy, S. C.: The 2010 California Research at the Nexus of Air Quality and Climate Change (CalNex) field study, *Journal of Geophysical Research: Atmospheres*, 118, 5830-5866, 10.1002/jgrd.50331, 2013.
- 525 Sato, K., Takami, A., Iozaki, T., Hikida, T., Shimono, A., and Imamura, T.: Mass spectrometric study of secondary organic aerosol formed from the photo-oxidation of aromatic hydrocarbons, *Atmospheric Environment*, 44, 1080-1087, 10.1016/j.atmosenv.2009.12.013, 2010.
- Schwantes, R. H., Emmons, L. K., Orlando, J. J., Barth, M. C., Tyndall, G. S., Hall, S. R., Ullmann, K., St. Clair, J. M., Blake, D. R., Wisthaler, A., and Bui, T. P. V.: Comprehensive isoprene and terpene gas-phase chemistry improves simulated surface ozone in the southeastern US, *Atmospheric Chemistry and Physics*, 20, 3739-3776, 10.5194/acp-20-3739-2020, 2020.
- 535 Shrivastava, M., Fast, J., Easter, R., Gustafson, W. I., Zaveri, R. A., Jimenez, J. L., Saide, P., and Hodzic, A.: Modeling organic aerosols in a megacity: comparison of simple and complex representations of the volatility basis set approach, *Atmospheric Chemistry and Physics*, 11, 6639-6662, 10.5194/acp-11-6639-2011, 2011.
- Shrivastava, M., Zelenyuk, A., Imre, D., Easter, R., Beranek, J., Zaveri, R. A., and Fast, J.: Implications of low volatility SOA and gas-phase fragmentation reactions on SOA loadings and their spatial and temporal evolution in the atmosphere, *Journal of Geophysical Research: Atmospheres*, 118, 3328-3342, 10.1002/jgrd.50160, 2013.
- 540 Shrivastava, M., Easter, R. C., Liu, X., Zelenyuk, A., Singh, B., Zhang, K., Ma, P.-L., Chand, D., Ghan, S., Jimenez, J. L., Zhang, Q., Fast, J., Rasch, P. J., and Tiitta, P.: Global transformation and fate of SOA: Implications of low-volatility SOA and gas-phase fragmentation reactions, *Journal of Geophysical Research: Atmospheres*, 120, 4169-4195, 10.1002/2014jd022563, 2015.
- 545 Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., Ng, N. L., Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A., Volkamer, R., Wang, J., Worsnop, D. R., Zaveri, R. A., Zelenyuk, A., and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Reviews of Geophysics*, 55, 509-559, 10.1002/2016rg000540, 2017.
- Shrivastava, M., Andreae, M. O., Artaxo, P., Barbosa, H. M. J., Berg, L. K., Brito, J., Ching, J., Easter, R. C., Fan, J., Fast, J. D., Feng, Z., Fuentes, J. D., Glasius, M., Goldstein, A. H., Alves, E. G., Gomes, H., Gu, D., Guenther, A., Jathar, S. H., Kim, S., Liu, Y., Lou, S., Martin, S. T., McNeill, V. F., Medeiros, A., de Sa, S. S., Shilling, J. E., Springston, S. R., Souza, R. A. F., Thornton, J. A., Isaacman-VanWertz, G., Yee, L. D., Ynoue, R., Zaveri, R. A., Zelenyuk, A., and Zhao, C.: Urban pollution greatly enhances formation of natural aerosols over the Amazon rainforest, *Nat Commun*, 10, 1046, 10.1038/s41467-019-08909-4, 2019.



- 555 Signorell, R., and Bertram, A.: Physical chemistry of aerosols, *Phys Chem Chem Phys*, 11, 7759, 10.1039/b916865f, 2009.
Solomon, P. A., Crumpler, D., Flanagan, J. B., Jayanty, R. K., Rickman, E. E., and McDade, C. E.: U.S. national PM_{2.5} Chemical Speciation Monitoring Networks-CSN and IMPROVE: description of networks, *J Air Waste Manag Assoc*, 64, 1410-1438, 10.1080/10962247.2014.956904, 2014.
- 560 Tilmes, S., Hodzic, A., Emmons, L. K., Mills, M. J., Gettelman, A., Kinnison, D. E., Park, M., Lamarque, J. F., Vitt, F., Shrivastava, M., Campuzano-Jost, P., Jimenez, J. L., and Liu, X.: Climate Forcing and Trends of Organic Aerosols in the Community Earth System Model (CESM2), *Journal of Advances in Modeling Earth Systems*, 10.1029/2019ms001827, 2019.
- Toon, O. B., Maring, H., Dibb, J., Ferrare, R., Jacob, D. J., Jensen, E. J., Luo, Z. J., Mace, G. G., Pan, L. L., Pfister, L., Rosenlof, K. H., Redemann, J., Reid, J. S., Singh, H. B., Thompson, A. M., Yokelson, R., Minnis, P., Chen, G., Jucks, K. W., and Pszenny, A.: Planning, implementation, and scientific goals of the Studies of Emissions and Atmospheric Composition, Clouds and Climate Coupling by Regional Surveys (SEAC4RS) field mission, *Journal of Geophysical Research: Atmospheres*, 121, 4967-5009, 10.1002/2015jd024297, 2016.
- 565 Tsigaridis, K., Daskalakis, N., Kanakidou, M., Adams, P. J., Artaxo, P., Bahadur, R., Balkanski, Y., Bauer, S. E., Bellouin, N., Benedetti, A., Bergman, T., Bernsten, T. K., Beukes, J. P., Bian, H., Carslaw, K. S., Chin, M., Curci, G., Diehl, T., Easter, R. C., Ghan, S. J., Gong, S. L., Hodzic, A., Hoyle, C. R., Iversen, T., Jathar, S., Jimenez, J. L., Kaiser, J. W., Kirkevåg, A., Koch, D., Kokkola, H., Lee, Y. H., Lin, G., Liu, X., Luo, G., Ma, X., Mann, G. W., Mihalopoulos, N., Morcrette, J. J., Müller, J. F., Myhre, G., Myriokefalitakis, S., Ng, N. L., and others, Donnell, D., Penner, J. E., Pozzoli, L., Pringle, K. J., Russell, L. M., Schulz, M., Sciare, J., Seland, Ø., Shindell, D. T., Sillman, S., Skeie, R. B., Spracklen, D., Stavrou, T., Steenrod, S. D., Takemura, T., Tiitta, P., Tilmes, S., Tost, H., van Noije, T., van Zyl, P. G., von Salzen, K., Yu, F., Wang, Z., Wang, Z., Zaveri, R. A., Zhang, H., Zhang, K., Zhang, Q., and Zhang, X.: The AeroCom evaluation and intercomparison of organic aerosol in global models, *Atmospheric Chemistry and Physics*, 14, 10845-10895, 10.5194/acp-14-10845-2014, 2014.
- 570 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth System Science Data*, 9, 697-720, 10.5194/essd-9-697-2017, 2017.
- Warneke, C., Trainer, M., de Gouw, J. A., Parrish, D. D., Fahey, D. W., Ravishankara, A. R., Middlebrook, A. M., Brock, C. A., Roberts, J. M., Brown, S. S., Neuman, J. A., Lerner, B. M., Lack, D., Law, D., Hubler, G., Pollack, I., Sjostedt, S., Ryerson, T. B., Gilman, J. B., Liao, J., Holloway, J., Peischl, J., Nowak, J. B., Aikin, K., Min, K. E., Washenfelder, R. A., Graus, M. G., Richardson, M., Markovic, M. Z., Wagner, N. L., Welti, A., Veres, P. R., Edwards, P., Schwarz, J. P., Gordon, T., Dube, W. P., McKeen, S., Brioude, J., Ahmadov, R., Bougiatioti, A., Lin, J. J., Nenes, A., Wolfe, G. M., Hanisco, T. F., Lee, B. H., Lopez-Hilfiker, F. D., Thornton, J. A., Keutsch, F. N., Kaiser, J., Mao, J., and Hatch, C.: Instrumentation and Measurement Strategy for the NOAA SENEX Aircraft Campaign as Part of the Southeast Atmosphere Study 2013, *Atmos Meas Tech*, 9, 3063-3093, 10.5194/amt-9-3063-2016, 2016.
- 585 Wofsy, S. C., Afshar, S., Allen, H. M., Apel, E. C., Asher, E. C., Barletta, B., Bent, J., Bian, H., Biggs, B. C., Blake, D. R., Blake, N., Bourgeois, I., Brock, C. A., Brune, W. H., Budney, J. W., Bui, T. P., Butler, A., Campuzano-Jost, P., Chang, C. S., Chin, M., Commane, R., Correa, G., Crouse, J. D., Cullis, P. D., Daube, B. C., Day, D. A., Dean-Day, J. M., Dibb, J. E., DiGangi, J. P., Diskin, G. S., Dollner, M., Elkins, J. W., Erdesz, F., Fiore, A. M., Flynn, C. M., Froyd, K. D., Gesler, D. W., Hall, S. R., Hanisco, T. F., Hannun, R. A., Hills, A. J., Hints, E. J., Hoffman, A., Hornbrook, R. S., Huey, L. G., Hughes, S., Jimenez, J. L., Johnson, B. J., Katich, J. M., Keeling, R. F., Kim, M. J., Kupc, A., Lait, L. R., Lamarque, J. F., Liu, J., McKain, K., McLaughlin, R. J., Meinardi, S., Miller, D. O., Montzka, S. A., Moore, F. L., Morgan, E. J., Murphy, D. M., Murray, L. T., Nault, B. A., Neuman, J. A., Newman, P. A., Nicely, J. M., Pan, X., Pappalardo, W., Peischl, J., Prather, M. J., Price, D. J., Ray, E. A., Reeves, J. M., Richardson, M., Rollins, A. W., Rosenlof, K. H., Ryerson, T. B., Scheuer, E., Schill, G. P., Schroder, J. C., Schwarz, J. P., St.Clair, J. M., Steenrod, S. D., Stephens, B. B., Strode, S. A., Sweeney, C., Tanner, D., Teng, A. P., Thames, A. B., Thompson, C. R., Ullmann, K., Veres, P. R., Vieznor, N., Wagner, N. L., Watt, A., Weber, R., Weinzierl, B., Wennberg, P. O., Williamson, C. J., Wilson, J. C., Wolfe, G. M., Woods, C. T., and Zeng, L. H.: ATom: Merged Atmospheric Chemistry, Trace Gases, and Aerosols, in, ORNL Distributed Active Archive Center, 2018.
- 600 Xu, L., Pye, H. O. T., He, J., Chen, Y., Murphy, B. N., and Ng, L. N.: Experimental and model estimates of the contributions from biogenic monoterpenes and sesquiterpenes to secondary organic aerosol in the southeastern United States, *Atmos Chem Phys*, 18, 12613-12637, 10.5194/acp-18-12613-2018, 2018.



- 605 Yu, S., Dennis, R. L., Bhave, P. V., and Eder, B. K.: Primary and secondary organic aerosols over the United States: estimates on the basis of observed organic carbon (OC) and elemental carbon (EC), and air quality modeled primary OC/EC ratios, *Atmospheric Environment*, 38, 5257-5268, 10.1016/j.atmosenv.2004.02.064, 2004.
- Zhang, H., Yee, L. D., Lee, B. H., Curtis, M. P., Worton, D. R., Isaacman-VanWertz, G., Offenberg, J. H., Lewandowski, M., Kleindienst, T. E., Beaver, M. R., Holder, A. L., Lonneman, W. A., Docherty, K. S., Jaoui, M., Pye, H. O. T., Hu, W., Day, D. A., Campuzano-Jost, P., Jimenez, J. L., Guo, H., Weber, R. J., de Gouw, J., Koss, A. R., Edgerton, E. S., Brune, W., Mohr, C., Lopez-Hilfiker, F. D., Lutz, A., Kreisberg, N. M., Spielman, S. R., Hering, S. V., Wilson, K. R., Thornton, J. A., and Goldstein, A. H.: Monoterpenes are the largest source of summertime organic aerosol in the southeastern United States, *Proc Natl Acad Sci U S A*, 115, 2038-2043, 10.1073/pnas.1717513115, 2018.
- 610 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophysical Research Letters*, 34, n/a-n/a, 10.1029/2007gl029979, 2007.
- 615 Zheng, Y., Thornton, J. A., Ng, N. L., Cao, H., Henze, D. K., McDuffie, E. E., Hu, W., Jimenez, J. L., Marais, E. A., Edgerton, E., and Mao, J.: Long-term observational constraints of organic aerosol dependence on inorganic species in the southeast US, *Atmos. Chem. Phys. Discuss.*, 2020, 1-39, 10.5194/acp-2020-575, 2020.
- 620