

Experimental investigation of the stable water isotopes distribution in an Alpine lake environment (L-WAIVE)

Patrick Chazette¹, Cyrille Flamant², Harald Sodemann³, Julien Totems¹, Anne Monod⁴, Elsa Dieudonné⁵, Alexandre Baron¹, Andrew Seidl³, Hans Christian Steen-Larsen³, Pascal Doira¹, Amandine Durand⁴, and Sylvain Ravier⁴

¹Université Paris-Saclay, Laboratoire des Sciences du Climat et de l'Environnement (LSCE), CEA-CNRS-UVSQ, UMR CNRS 8212, CEA Saclay, 91191 Gif-sur-Yvette, France

²LATMOS/IPSL, Sorbonne Université, UVSQ, CNRS UMR 8190, Paris, France

Correspondence to: Patrick Chazette (patrick.chazette@lsce.ipsl.fr)

³University of Bergen, Bergen and Bjerknes Centre for Climate Research, Bergen, Norway

⁴Aix Marseille Univ, CNRS, LCE, Marseille, France

⁵Université du Littoral Côte d'Opale, Laboratoire de Physico-Chimie de l'Atmosphère (ULCO/LPCA), France

Correspondence to: Patrick Chazette (patrick.chazette@lsce.ipsl.fr)

Abstract. In order to gain understanding on the vertical structure of atmospheric water vapor above mountain lakes and to assess its link with the isotopic composition of the lake water and with small-scale dynamics (i.e. valley winds, thermal convection above complex terrain), the L-WAIVE (Lacustrine-Water vApor Isotope inVentory Experiment) field campaign was conducted in the Annecy valley in the French Alps during 10 days in June 2019. This field campaign was based on an original experimental synergy between a suite of ground-based, boat-borne, and two ultra-light aircraft (ULA) measuring platforms implemented to characterise the thermodynamic and isotopic composition above and in the lake. A Cavity Ring-down Spectrometer and an in-cloud liquid water collector were deployed onboard one of the ULA to characterize the vertical distribution of the main stable water isotopes (H_2^{16}O , H_2^{18}O and $\text{H}^2\text{H}^{16}\text{O}$) both in the air and in shallow cumulus clouds. The temporal evolution of the meteorological structures of the low troposphere was derived from an airborne Rayleigh-Mie lidar (embarked on a second ULA), a ground-based Raman lidar and a wind lidar. ULA flight patterns were repeated several times per day to capture the diurnal evolution as well as the variability associated with the different weather events encountered during the field campaign, which influenced the humidity field, cloud conditions and slope wind regimes in the valley. In parallel, throughout the campaign, liquid water samples of rain, at the air-lake water interface, and at 2 m depth in the lake were taken. A significant variability of the isotopic composition was observed along time, depending on weather conditions, linked to the transition from the valley boundary layer towards the free troposphere, the valley wind intensity, and the vertical thermal stability. Thus, significant gradients of isotopic content have been revealed at the transition to the free troposphere, at altitudes between 2.5 and 3.5 km. The influence of the lake on the atmosphere isotopic composition is difficult to isolate from other contributions, especially in the presence of thermal instabilities and valley winds. Nevertheless, such an effect appears to be detectable in a layer of about 300 m thickness above the lake in light wind conditions. We also noted similar isotopic compositions in cloud drops and rainwater.

Keywords: L-WAIVE, valley, lake, Annecy, stable water isotopes, aerosol, lidar, vertical profiles, airborne

1 Introduction

Why are the vertical structures of the stable isotope of the water vapor field in the lower troposphere only sparsely documented above Alpine lakes? This is in part due to the complexity and fast-evolving nature of the low-level atmospheric circulation in Alpine-type valleys which is intimately linked to the surrounding orography interacting with the synoptic scale circulation. Thermally driven wind systems may be induced by hilly terrain, such as slope, mountain, and plateau winds (Kottmeier et al., 2008). Such winds result in mountain venting phenomena that control the variability of the water vapor field in mountain catchment on very small-time scales. Furthermore, small-scale inhomogeneity in soil properties across a mountain valley, as well as lake breezes resulting from land-lake temperature contrasts, may also induce the development of thermal circulations, particularly on clear-sky days, modifying the wind, humidity, and temperature fields on small spatial scales. The interaction of slope-driven and secondary circulations can furthermore influence the thermodynamical environment in the valley by the formation of convergence lines. Such convergence lines may favour the formation of shallow clouds, and in some cases even deep convection (Barthlott et al., 2006). Interaction with the synoptic scale circulation can lead to the formation of strong, gusty down-valley winds such as foehn events (Drobinski et al., 2007) and gap flows (Flamant et al., 2002; Mayr et al., 2007) that can also contribute to rapid modifications of the water vapor field in mountain catchment areas.

Stable water isotopes have long been used as a tool to study processes in lacustrine and hydrologic systems (see review by Gat, 2010), as well as evapotranspiration (e.g. Berkelhammer et al., 2016). For instance, an early study based on water stable isotope measurements conducted in the US Great Lakes region suggested that in the summer up to 15% of the atmospheric water content in the atmosphere downwind of the lakes is derived from lake evaporation (Gat et al., 1994). Isotopic measurements of lake water have shown the relative roles of evaporation from the lake surface and transpiration from surrounding vegetation (Jasechko et al., 2013). The link between hydrology and evaporation has mainly been investigated using vapor and liquid water isotopes measurements gathered just above the Earth's surface and samples from lake water and precipitation (e.g. Cui et al., 2016).

Unexploited potential remains in using stable water isotopes to increase our understanding of the influence of evaporation, boundary-layer processes, and the free troposphere for local and regional climate conditions in Alpine lakes. For example, the depth of the atmospheric layer over which the influence of evaporation from the lake surface is detectable, and how different factors control the depth of this layer are still largely unknown. Detailed and comprehensive analysis of small-scale factors, such as winds in a valley, and how they are related to local and larger-scale dynamics, such as synoptic scale subsidence in complex terrain, are therefore needed.

65 In order to get insights into such aspects, the L-WAIVE (Lacustrine-Water vApor Isotope inVentory Experiment) field campaign was conducted in the Annecy valley (45°47' N, 6°12' E, in Haute-Savoie in the French Alps) around the Annecy lake between 12 and 23 June 2019. Being the second largest natural, glacial lake in France, the Annecy lake is expected to play a substantial role for the regional hydrometeorology.

70 The overarching scientific objective of L-WAIVE field campaign is to study the vertical distribution of the water vapor contents and its heterogeneity over the Annecy lake. For this purpose, we have implemented an original multi-platform experimental approach based on continuous high-resolution vertical profiling of tropospheric water vapor, temperature and wind as well as scattering layers (aerosols, clouds) in the valley. Continuous atmospheric sampling was carried out by lidars together with airborne and boat-borne measurements of stable water isotopes (H_2^{16}O , H^2HO and H_2^{18}O) in vapor, as well as liquid water sampling in the lake, clouds, and precipitation. A consolidated vision of water isotopes across the air/water compartments in the lake area is proposed. Special emphases are made on the interface between the lake and the atmosphere, as well as at the boundary with the free troposphere.

80 It is worth noting that we have defined such an observational strategy including ultra-light aircraft (ULA) means because measurements tall towers only provide incomplete information on the link between the different compartments in the free troposphere where the processes of mixing and distillation occur (e.g. Griffis et al., 2016; Steen-Larsen et al., 2013). In the past, He and Smith (1999) combined airborne measurements and surface sampling of the water isotope composition to study the evaporation process over the forests of New England, before the advent of high-resolution laser-based spectrometers. More recent available airborne measurements of the isotope composition in the boundary layer either focused on areas above the sea (Sodemann et al., 2017), or did not include measurements of the surface isotope composition (Salmon et al., 2019). The use of ULAs makes it possible to sample the atmosphere very close to the surface up to altitudes of more than 4000 m and to fly in deep valleys (Chazette et al., 2005).

90 The rest of the article is divided into 6 sections. The experimental strategy is discussed in Section 2 whereas the instrumental platforms operations and environmental variables monitoring are presented in section 3. In Section 4 we give the synoptic conditions of atmospheric circulation that help to understand the temporal evolution of the measurements in the valley, and more generally the meteorological conditions encountered during the field campaign. Atmospheric water vapor and liquid water isotopes, as well as lake water isotope observations made across the valley are described in Section 5. In Section 6, the vertical distribution of water isotopes is discussed taking into account the vertical structures of the lower troposphere and the lake/atmosphere interface. We conclude in Section 7.

2 L-WAIVE experimental strategy

95 To achieve the scientific and methodological objectives of the L-WAIVE project, the field campaign was implemented in the southern part of the Annecy lake (the so called “Petit Lac”), in the vicinity of the city of Lathuile (Fig. 1). The Annecy lake is

bordered by the city of Annecy to the north, the Massif des Bauges to the west (2217 m above mean sea level – a.m.s.l.), the Massif des Bornes to the East (2438 m a.m.s.l.) and the depression de Faverges to the south (where Lathuile is located). Lathuile is located east of the foothill of the Roc des Boeufs (1774 m a.m.s.l.) to the west of the “Petit Lac” and the Tournette summit (2350 m a.m.s.l.) to the east. The Annecy lake, at a mean altitude of 446.7 m a.m.s.l., covers an area of roughly 27.5 km², and has a mean (maximum) depth of 41.5 m (82 m).

100




Figure 1. Geographical location of L-WAIVE. The different pictures give a view of the environment where the measurements were performed and of the instrumented platforms used: a) location of the experiment, b) lake water sampling from a boat, c) instrumented van, d) instrumented ultra-light aircrafts, and e) panoramic view from UAV showing the location of Lathuile, as well as the Roc des Boeufs to the left, the Tournette summit to the right and the “Petit Lac” in between.

105

2.1 Measurement platforms

Two airborne, one boat-borne and one ground-based instrumented platforms were deployed in the vicinity of Lathuile in order to monitor humidity, temperature, wind, clouds and aerosols in the lower troposphere over the Annecy lake and the surrounding valley environment, as well as to conduct measurements at the interface between the atmosphere and the lake, and in the lake.

110

A brief overview of the platforms is given below, whereas the details on the instrumental payloads are given in Section 3:

- i) Airborne platforms:

- 115
- a. One ultra-light aircraft (ULA) was mainly dedicated to remote sensing measurements. It allowed exploring the two or three-dimensional structure of the lower troposphere thanks to a polarized Rayleigh-Mie lidar. It also carried a Global Positioning System (GPS) device and a meteorological probe (pressure, temperature, relative humidity). This will be referred to as “aerosol ULA” (ULA-A) in the following.
 - b. A second ULA carried both a Cavity Ring-down Spectrometer (CRDS) water vapor isotope analyser, a meteorological probe for pressure, air temperature, GPS and relative humidity, and a cloud water collector. The platform offers the opportunity to measure the vertical profiles of temperature, relative humidity, H^2HO , H_2^{18}O and H_2^{16}O and to collect cloud water samples. The latter were collected during specific cloud flights (when meteorological conditions were favourable) and the cloud collector was opened only in clouds. We will refer to this platform as “isotope and cloud ULA” (ULA-IC) in the following.
- ii) Ground-based platform: Simultaneous high-resolution vertical profiles of water vapor, temperature, aerosols and winds were acquired continuously from two co-located ground-based lidars.
 - iii) Boat-borne platform: An instrumented boat allowed sampling the lake water by vials at the surface film and underneath (~2 m deep) for the assessment of H^2HO and H_2^{18}O . A CRDS water vapor isotope analyser performed measurements during one day at the end of the experiment just above the lake surface in parallel with the lake water sampling.
- 120
- 125

These different platforms are presented in Fig. 1 together with a view of the experiment site.

2.2 Deployment



130 Depending on the weather conditions, airborne platforms were deployed several times a day to document the temporal evolution of the atmospheric boundary layer over the lake. The days of operation of all platforms are summarized in Table A1 of Appendix A. The ground-based water vapor, temperature and aerosol lidar operated continuously between 12 and 21 June in the morning (gathering over 220 h of data), while the ground-based wind lidar (WL) operated continuously between 14 and 23 June in the morning (acquiring also over 220 h of data). A total of 22 successful flights were conducted with ULA-A
135 between 13 and 19 June, while ULA-IC performed 15 flights between 14 and 20 June. Valid cloud water samples were only obtained during the last 3 ULA-IC flights.

The CRDS previously installed on ULA-IC was mounted on the boat from 21 to 22 June. Boat-borne CRDS observations were made at the surface of the lake on the last outings of the boat. Nevertheless, lake water samples were taken on 14 occasions during the field campaign (see Appendix A). Finally, 28 samples of precipitation were taken during the campaign (7 of which
140 on 15 June in the afternoon) between 11 and 22 June 2019.

On days when both ULA flew in coordinated patterns (13, 16-20 June), flights typically began with a profiling sequence between the surface and ~4 km a.m.s.l. which was carried out in the vicinity of the 2 ground-based vertically pointing lidars

(see Fig. 2). Soundings with levelled legs (see blue dotted line in Fig. 2a) were performed at a relatively slow ascent rate (~ 60 m min⁻¹) to ensure that the instruments were as close as possible to equilibrium with the environment. Upon reaching 4 km a.m.s.l., the flight route of the 2 ULA could differ, ULA-A performing a high-altitude survey above the Annecy lake (see red dotted line in Fig. 2a), while ULA-IC was aiming for shallow cumulus clouds to sample cloud water droplets as illustrated in Fig. 2b. Liquid water sampling was performed via multiple passes through the clouds to accumulate enough material to conduct isotope analysis. At the end of the flight, both ULA performed race-track descents around the ground-based lidars on their way back to the airfield.

During ascent and descent, the airborne lidar ALiAS onboard ULA-A was pointing sideways to directly derive the aerosol extinction coefficient (Chazette et al., 2007; Chazette and Totems, 2017). For the exploration of the valley at a cruising altitude between 3.5 and 4.5 km a.m.s.l., ALiAS was pointing to the nadir. The combination of both flight sequences thus allowed to survey the 3-dimensional structure of the lower troposphere over the lake and its surroundings. The individual flight characteristics (time, maximum altitude, type of exploration) are presented in Appendix B for the two ULAs (Tables B1 and B2 for ULA-A and ULA-IC, respectively).



160 **Figure 2. Example of a typical ULA flight plan performed during L-WAIVE (performed on 17 June 2019. a) The flight track adopted during the flight (blue dots for vertical profiling and red dots for horizontal exploration above the lake). DEM is for digital elevation model. b) Schematic representation of the measurement strategy adopted during L-WAIVE. Two types of sampling strategy are illustrated, red for atmospheric sampling and green for cloud sampling. The purple arrows illustrate that the atmospheric sampling is performed during both ascent and descent.**

3 Instrumental set up on each platform

This section provides a detailed description of the payloads on all platforms deployed during L-WAIVE. The periods of operation of each measurement platform are given in Appendix A.

165 3.1 Airborne payloads

We used two Tanarg 912 XS ULA from the company Air Création (Chazette and Totems, 2017). For each ULA (ULA-A and ULA-IC), the maximum total payload is of approximately 250 kg including the pilot. Flight durations were between ~1 and 2 hours, depending on flight conditions, with a cruise speed around 85-100 km h⁻¹. The ULA location was provided by a GPS and an Attitude and Heading Reference System, which are part of the MTi-G components sold by XSens.

170 3.1.1 ULA-A

Rayleigh-Mie lidar. The Airborne Lidar for Atmospheric Studies (ALiAS) was especially developed by LSCE as an airborne payload dedicated to sample the atmospheric scattering layers and then the vertical structures of the atmosphere (Chazette et al., 2012, 2020). It emits a pulse energy of 30 mJ in the ultraviolet at 355 nm with a 20 Hz pulsed Nd:YAG laser (ULTRA) manufactured by QUANTEL™ (<https://www.quantel-laser.com/>). The acquisition system is based on a PXI (PCI eXtensions for Instrumentation) technology. The receiver contains two channels for the detection of the elastic backscatter from the atmosphere in the parallel and perpendicular polarization planes relative to the linear polarization of the emitted radiation. The native resolution along the line-of-sight is 0.75 m, it is degraded here to 100 m during the data treatment to improve the signal to noise ratio. The wide field-of-view of ~2.3 mrad ensures a full-overlap of the transmit and receive paths close to 200-300 m from the emitter.

180 **Meteorological probe.** Part of ULA-A payload was a shielded meteorological probe VAISALA PTU-300 for measuring temperature, pressure and relative humidity. This probe measures the atmospheric pressure, with a 1-minute sampling time, within an uncertainty of 0.25 hPa, the air temperature within an uncertainty of 0.2 K and relative humidity (RH) within a relative uncertainty of 2.5%.

3.1.2 ULA-IC

185 **CRDS water vapor isotope analyser.** ULA-IC carried a CRDS water vapor isotope analyser (L2130-i, Picarro Inc., Sunnyvale, USA; Ser. No. HIDS2254) for the in-situ measurement at about 5 Hz of the H₂¹⁶O mixing ratio, and the isotope

ratios $\delta^{18}\text{O}$ and $\delta^2\text{H}$ for H_2^{18}O and H^2HO , respectively. Water vapor was drawn into the spectrometer through an unheated inlet of 68 cm length (1/4 inch O.D. stainless steel with Silconert coating), pointing backward on the left side of the aircraft at a distance of 38 cm from the CRDS. Pointing forward next to the vapor inlet, a fast-response temperature and humidity probe (iMet XQ-2, InterMet systems, USA; Ser. No. 61124) measured thermodynamic properties (T, RH, p with uncertainties of 0.3 K, 5 %, and 1.5 hPa, respectively) and GPS location at 1 Hz. The CRDS analyser was operated in flight mode, with a flow of about 150 sccm through the inlet maintained by a membrane pump (Part No. S2003, Picarro Inc.). Pressure and water vapor mixing ratio were corrected using calibration functions established at the FARLAB laboratory, University of Bergen, Norway. Raw measurements of the isotope parameters, expressed as δ -values (see Appendix C) were corrected for the mixing ratio-isotope composition dependency and calibrated across the campaign period following recommendations from the International Atomic Energy Agency (IAEA, see Appendix D for details). Tests with a small bubbler system during the campaign indicated an anomaly in the CRDS measurements during 18-20 June, partially affecting two flights in the morning of 19 June, and the first flight on 20 June. The anomaly was due to a saturated inlet system from condensate forming on the aircraft during a cloud sampling flight on 18 June. Flight periods affected by inlet saturation effects were excluded from further analysis (see Appendix D).

A time resolution better than 0.1 Hz was obtained for specific humidity from the CRDS and iMet probe, providing a spatial resolution of 200-300 m in horizontal direction, and 10-50 m in the vertical, assuming a typical horizontal speed of 85-100 m s^{-1} and ascent rate of the aircraft of about 1-5 m s^{-1} . Due to more complex memory effects, the isotope composition has lower effective time resolution (Sodemann et al., 2017; Steen-Larsen et al., 2014). We use here 10 s average data for all parameters on ULA-IC from upward profiles, filtered for rapid elevation changes, defined as exceeding 1.0 hPa ascend or 1.0 hPa descend within 10 s.

Cloud water collector. A pre-cleaned Caltech Active Strand Cloud Water Collector (CASCC, Demoz et al., 1996) was mounted on ULA-IC, modified to efficiently collect cloud water at the relative cruising speed of the ULA (85 to 100 km h^{-1}). The CASCC was modified to efficiently collect cloud liquid droplets from the ULA. In order to sample droplets under the same conditions as those obtained on the ground, the CASCC fan was removed, and its inlet and outlet were prolonged with convergent and divergent High-Density PolyEthylene (HDPE) cones to ensure an isokinetic air sampling. All parts necessary to adapt the HDPE cones on the CASCC were made of plastic. The flow through the probe must be steady and as turbulence-free as possible. The design of these modified inlet and outlet was calculated to get a constant mass of water droplets per time unit through the probe onboard the ULA. The resulting probe was installed on the side of the ULA where the flow was assumed laminar and allowed air sampling with a flow rate of 35 to 47 $\text{m}^3 \text{min}^{-1}$, ahead of the motor exhaust. The CASCC strings and inlet were pre-cleaned with deionized water prior each flight and covered with a clean plastic bag when not in-cloud (especially during take-off and landing). For a flight of 10 minutes inside a shallow cumulus cloud, the probe typically collected 41 to 48 g of cloud liquid droplets corresponding to a liquid water content of 0.10 to 0.16 g m^{-3} , typical for such cloud type (Herrmann, 2003).

220 3.2 Ground-based instrumentation

The ground-based scientific facility hosted by the technical department of the city of Lathuile was mainly composed of the Raman lidar WALI and of a scanning Doppler lidar. WALI was embedded in the ground-based MAS (mobile atmospheric station; Raut and Chazette, 2009). The UAV was also operated from this site, close to the lidar.

3.2.1 Raman lidar

225 WALI has been developed at LSCE (Chazette et al., 2014) based on the same technology as its precursor instruments LESAA (Lidar pour l'Etude et le Suivi de l'Aérosol Atmosphérique; Chazette et al., 2005) and LAUVA (Lidar Aérosol UltraViolet Aéroporté, Chazette et al., 2007; Lesouëf et al., 2013; Raut and Chazette, 2009). It is a custom-made instrument dedicated to atmospheric research activities.

230 The receiver of the lidar is composed of 2 distinct detection paths with a low full-overlap distance (~150-200 m). The first path is dedicated to the detection of the elastic molecular, aerosols and cloud backscatter from the atmosphere (Rayleigh-Mie lidar path). Two different channels are implemented on that path to detect i) the total (co-polarized and cross polarized with respect to the laser emission) and ii) the cross-polarized backscatter coefficients of the atmosphere. The second path, a fibered achromatic reflector, is dedicated to the measurements of the atmospheric Raman scattering, namely the vibrational signal for nitrogen (N₂-channel) and water vapor (H₂O-channel) and the rotational signal to derive the temperature (T-Channel).

235 During the entire experiment, the acquisition was performed for mean profiles of 1000 laser shots leading to a native temporal sampling close to 1 minute. Rayleigh-Mie lidars are very efficient tools to detect scattering layers and then inform on the vertical structures of the atmosphere (e.g. Platt, 1977; Berthier et al., 2004; Chazette et al., 2020). The water vapor mixing ratio (WVMR) is retrieved with an absolute error less than 0.4 g kg⁻¹ in the first 2 km above the ground level (a.g.l.) (Chazette et al., 2014; Totems et al., 2019). The calibration of the T-channel is derived from the methodology presented by Behrendt (2006) and leads to an absolute error on the temperature lower than 1 °C within the first 2 km a.g.l. The final vertical resolution is set to 15 m below 1 km a.g.l. and 30 m above, and the temporal resolution is 0.5 h. In the following, a temporal resolution of 1 h is used.

3.2.2 Wind lidar

245 Wind profiles were measured using a scanning Doppler lidar (Leosphere Windcube WLS100). It operates in the infrared (1.543 μm) with a low pulse energy (0.25 mJ) but a high pulse repetition frequency (20 kHz). The Doppler shift due to the particles' motion along the beam direction (radial wind speed) is determined through heterodyne detection followed by fast Fourier transform analysis. The acquisition time was set to 1 s during the campaign. The pulse duration is 200 ns, corresponding to an axial resolution of 50 m given the pulse shape, with a minimal and maximal range of 100 m and 7.2 km, respectively. The axial resolution can be lowered to 25 m by reducing the pulse duration (100 ns) while increasing the pulse repetition frequency (40 kHz) which in turn reduces the minimal and maximal range to 50 m and 3.3 km, respectively. In practice, the maximum

250

range is limited by the signal level. A minimum Carrier to Noise Ratio (i.e. signal to background noise ratio) of -27 dB is required to keep the radial wind uncertainty (determined from the spectrum peak width) below 0.5 ms^{-1} . Therefore, observations in the free troposphere are possible only when elevated layers of aerosols are present. Even in the boundary layer, several days of very low aerosol load occurred during the campaign, in which case the Carrier to Noise Ratio threshold was lowered to -30 dB. With such a low Carrier to Noise Ratio, the measurements must be considered with caution.

Profiles of the three components of the wind vector were determined using the Doppler Beam Swinging technique originally proposed for Doppler radar (e.g. Koscielny et al., 1984). Here, the measurement cycle includes one vertical beam, for which the radial wind is the vertical component of the wind vector, and four slanted beams (15° from zenith) in the cardinal directions to derive the two horizontal components of the wind. The uncertainty on the horizontal / vertical wind components is determined using the variability inside averaging periods, possibly gathering several measurement cycles. During the campaign, the Doppler lidar was positioned in a wide valley (~ 3 km at the observation site) with a rather flat bottom, and the closest distance existing between one of the slanted beams and an obstacle was ~ 1 km. Therefore, the influence of the orography on the measurement are assumed to be negligible here.

3.3 Boat-borne payload: Lake and atmospheric sampling

The lake water surface and subsurface were sampled at the middle of the “Petit Lac” to measure water isotopes and chemicals from the boat:

- i) The lake water thermal stratification was monitored using an EXO sonde, equipped with temperature, pressure, pH, dissolved dioxygen, ion conductivity and chlorophyll sensors. Profiles recorded in the middle of the “Petit Lac” (Fig. 1) once or twice per day (Table A1) and showed that the thermocline was typically at a depth of about 7 to 13 m, in good agreement with previous studies of the Lake (Danis et al., 2003).
- ii) Subsurface water samples were collected at a depth of 2 m in HDPE capped flasks. Surface water samples were collected using a 30 x 30 cm silica glass plate immersed into water for a minute, then gently removed from water vertically (Cunliffe et al., 2013). The water falling from the plate in a continuous flow was not sampled, then, the dropwise water was collected in HDPE capped flasks. The surface microlayer samples were then collected by scraping the remaining water on the glass plates (using a rubber scraper) in amber glassware capped vials. All liquid water samples were measured for isotopic composition (H_2^{18}O , $\text{H}^2\text{H}^{16}\text{O}$) at FARLAB, University of Bergen according to established laboratory procedures (Appendix D).
- iii) On 22 June, the CRDS isotope analyser was taken on board the boat to sample a cross-section in the atmospheric layer just above the “Petit Lac” through the location used for in situ sampling on the other days. This made it possible to capture the isotope concentrations as a function of the distance from the shore and the depth of the lake on that day. Water vapor isotope measurements were taken from an inlet at either ~ 20 cm or ~ 2 m above the

water surface, while the iMet sonde measured temperature, relative humidity and location. We used 2.5 m of 1/4" PTFE tubing and a 40 cm 1/4" stainless steel tip on the first inlet, and about 1.5 m of tubing with a 50 cm stainless steel tip. A flow rate of about 10 lpm was provided by an external manifold pump (N022AN, KNF, Germany) to either of the selected inlet lines. Post-processing and calibration of water vapor measurements were done as for the aircraft data (Sec. 3.1.2).

3.4 Precipitation sampling

Precipitation samples were taken throughout the campaign period. The sampling device consisted in a pre-cleaned HDPE funnel directly connected to a pre-cleaned HDPE sampling bottle. The precipitation samples were manually operated: after each precipitation event, the sample was aliquoted into 1.5 ml glass vials with rubber/PTFE septa and stored at 4°C prior to isotopic analysis, while the rest of the sample was stored at -18°C. Precipitation sampling times lasted from 20 min to several hours, depending on rainfall rate.

4 Meteorological conditions during L-WAIVE

The synoptic and local weather conditions are analysed in the following, helping with the subsequent interpretation of the water vapour measurements in the lower troposphere over the Annecy valley.

4.1 Synoptic conditions

During L-WAIVE, France was under the influence of two main synoptic features, namely a pronounced trough over Britany and the British Isles and a high-pressure ridge over north Africa, sometimes extending across the Mediterranean and all the way into eastern Europe. This weather situation is illustrated in Fig. 3 using the Fifth European Centre for Medium-Range Weather Forecasts Reanalysis (ERA5) at 0.75° horizontal resolution. This configuration caused a particularly strong pressure gradient over the western Mediterranean for the period 12-15 June (Fig. 3a), flanked in the east by an intense Libyan anticyclone, transporting warm air from sub-tropics to mid-latitudes. The ridge weakened and broadened over the following days (16-19 June, Fig. 3b) before strengthening again (20-23 June) as the Libyan high intensified (Fig. 3c).

During the course of the campaign, the area of interest was under the influence of warmer temperatures in the free troposphere linked with the high-pressure ridge at the beginning and the end of the field campaign, and colder temperature from 20 to 22 June. Since the experiment was conducted in a valley, where ERA5 reanalyses are generally considered not to be very accurate below the average altitude of the mountains (~2 km a.m.s.l. in our case), it is more informative to use the measurements acquired during the field campaign from the lidar and ULAs to describe the evolution of meteorological variables (wind, humidity, temperature, pressure).






Figure 3. ECMWF ERA5 temperature at 700 hPa (K, shading), geopotential height at 700 hPa (gpm/1000, contours) and horizontal winds at 700 hPa (black vectors) on (a) 14 June 2019 at 12 UTC, (b) 17 June 2019 at 12 UTC, and (c) 22 June 2019 at 12 UTC. The location of Lathuile is indicated with a grey dot surrounded by white.

4.2 Insight on the local atmospheric conditions derived from lidars

315 During L-WAIVE, the temporal evolution of the scattering layers, including clouds and aerosols, has been monitored using both the ground-based lidar WALI and the airborne lidar ALiAS. They provide insight into air masses transport, weather situation and local convection. For instance, such a capability was used to improve our understanding of atmospheric circulation in complex situations such as extreme heat wave phenomena (Chazette et al., 2017).

320 Fig. 4 shows the temporal evolution of the vertical profile of the aerosol (or cloud) apparent scattering ratio corrected from the molecular transmission (ASR, Fig. 4a) and of the linear volume depolarization ratio (VDR, Fig. 4b) (Chazette et al., 2012). The vertical extension of aerosols in the valley is highly variable over time as is the presence and nature of clouds during the campaign. The strong diurnal cycle of ASR in the valley is tightly related to the slope winds and vertical stability associated to surface heating and the forcing due to the presence of clouds. The depth of the aerosol layer is the largest at the end of the afternoon when the convection is the strongest in the valley, while it is exhibiting a minimum around 1100-1200 LT when the valley winds reverse. The period was rather cloudy, with cloud types and the rainy periods including thunderstorms being indicated in Fig. 4a. In Fig. 4, we can clearly see the vertical updraft associated with clouds, which drives scatterers and water vapour towards higher altitudes, where there is a transition to the free troposphere. The transition is between the yellow and blue colours in Fig. 4a.

330 Significant day-to-day variations of the aerosol layer depth are thus observed during the field campaign which highlights local, regional, and synoptic scale transport of air masses over the Annecy valley. At the beginning of the campaign (mainly on 14 June) large VDR values are detected up to altitudes exceeding 7 km a.m.s.l. (Fig. 4b). These large VDR values are related to a large-scale Saharan dust transport episode over the Lac d'Annecy valley favoured by the synoptic conditions on that day. These aerosols are progressively mixed downward by subsidence, reaching the valley floor around 0000 LT on 15 June. The local wind in the valley is obviously disturbed during stormy periods and during the episode of long-range dust transport on 335 14 June. The presence of these aerosols at higher altitudes (up to ~5 km a.m.s.l.) allowed the wind to be retrieved with the wind lidar above 2 km a.m.s.l. (Fig. 5a-b).

340 Strong southerly winds (in excess of 20 m s^{-1} above 3.5 km a.m.s.l.) were observed on 14 June in agreement with the synoptic meteorological fields. Weaker winds, less than 5 m s^{-1} , are observed below 2 km a.m.s.l. for the entire period. The wind intensity does not show repetitive patterns from one day to the next. Nevertheless, it is rather in the course of the afternoon, when the wind come from the north to the south, that we have the strongest velocities in the planetary boundary layer. Wind direction shows regular variations (Fig. 5b), with winds directed towards the south during the most of the afternoons. On the contrary, during the night and the first part of the morning, winds are directed towards the city of Annecy (north).

The vertical profiles of WVMR derived from the WALI ground-based lidar are also shown in Fig. 5c. They highlight highly variable water vapor in the first 2 km of the atmosphere, as is the case with wind. Generally higher WVMRs are often observed at night, associated with nighttime downslope winds, but may persist during days associated with heavy precipitation such as on 15 June 2019. The boundary between high and low humidity of the free troposphere is also highlighted, especially at night when the lidar exhibits enhanced sensing capabilities at higher altitude. For example, high humidity is seen up to ~4 km a.m.s.l. on the night of 17-18 June in agreement with what is highlighted in Fig. 4a.

The wind at the surface of the lake controls evaporation, turbulent surface fluxes and influences the thermal convection, as well as cloud forcing influences the depth of the valley boundary layer, allowing the mixing of water vapour in a layer of variable thickness over time. The influence of the lake can therefore be seen up to altitudes of ~4 km a.m.s.l., bearing in mind that the transition altitude to the free troposphere is generally of the order of 2.5 km a.m.s.l. here. The altitude of this sharp transition is also linked to the average altitude of the mountains surrounding the measurement site. There are no big differences between day and night on this boundary. It should be noted, however, that there are substructures that can be characterised by wind shears at different altitudes (Fig. 5b). A 0.5-1 km thick layer over the lake is thus highlighted which can be linked to the wetter structures of Fig. 5c (in dark blue).








Figure 4. Temporal evolution of a) the aerosol scattering ratio (ASR) where the shaded areas correspond to the presence of clouds and b) the linear volume depolarization ratio (VDR) from 13 to 21 June 2019. The cloud type and rain location are indicated, as is the type of the main aerosol structures identified by the VDR (orange/pink for dust on 14 June and other for pollution aerosols, clouds are in pink). Ns, Ac and Cu indicate nimbostratus, altocumulus and cumulus clouds, respectively.



365 **Figure 5. Temporal evolution of a) wind speed (m s^{-1}), b) wind direction (0° is for North and 90° for east) obtained from wind lidar, and the water vapor mixing ratio (WVMR) derived from the ground-based Raman lidar. White areas correspond to missing data caused by detection limitations of the lidar (signal to noise ratio, presence of clouds).**

5 Water vapor isotope during L-WAIVE

Water in vapor phase was sampled throughout the campaign in terms of mixing ratio of the main isotope and abundance of isotopes H_2^{18}O and $\text{H}^2\text{H}^{16}\text{O}$. At the same time, liquid water samples were taken from the surface of the lake, in the clouds, and surface precipitation to obtain their isotopic contents, and to place them in the context of the atmospheric water vapor of the lower troposphere. Water vapor was sampled in-situ using a CRDS, whereas liquid samples, including precipitation, cloud water and lake water were analysed in the laboratory. Here we report the isotope composition (see interpretative framework in Appendix C) as delta-values relative to a standard (e.g. Gat, 1996).

5.1 Atmospheric in situ sampling

370 In total, 15 flights with ULA-IC have been performed (see Table B2) including 13 flights where the CRDS allowed a representative sampling of $\delta^{2\text{H}}$ and $\delta^{18\text{O}}$ (Appendix D). The isotope content $\delta^{2\text{H}}$ observed during the flights ranged between about -340 and -80‰ for flight altitudes up to ~ 3.5 km a.m.s.l. as shown in Fig. 6a. For this figure, we considered ascending and level flight segments, and excluded samplings with pressure increases larger than 1 hPa/10s. As observed in earlier studies, the dataset is clustered along a typical mixing line in $\delta^{2\text{H}} - q$ space (Noone, 2012; Salmon et al., 2019; Sodemann et al., 2017). The end-members of the mixing lines show substantial day-to-day variations, from $\delta^{2\text{H}} = -110$ to -80‰ ($\sim [-16, -12]$ ‰ for $\delta^{18\text{O}}$, not shown) for the more humid end member ($q > 8 \text{ g kg}^{-1}$), and from $\delta^{2\text{H}} = -340$ to -230‰ ($\sim [-30, -20]$ ‰ for $\delta^{18\text{O}}$, not shown) for the drier end member ($q < 3 \text{ g kg}^{-1}$). It is important to keep in mind that both, the variation in maximum flight altitudes and the meteorological situation can contribute to variability.

380 As shown in Fig. 6b-c, the vertical profiles of the isotope content $\delta^{2\text{H}}$ shows small variability with altitude below 2 km a.m.s.l., the greatest variability is observed in the first few hundred metres. In some cases, the profiles exhibit strong vertical gradients at higher elevations, as seen between 2.5 and 3.6 km a.m.s.l. during 5 flights (flights 5, 6, 7, 9 and 10). The same patterns are observed on the vertical profiles of $\delta^{18\text{O}}$ (not shown) and even on the WVMR profiles derived from the meteorological probes onboard the 2 ULAs. The sharp gradients are not observed systematically, as in the case of flight 15 (20 June), or may be present at a higher altitude, not sampled by the flight.

390 Indeed, because of the cloud cover and depending on the wind conditions, ULAs have not been able to fly in the free troposphere on most days. However, this was possible for flight 10 where a marked reduction of $\delta^{2\text{H}}$ occurred just above 3 km a.m.s.l., while higher values of isotopic content are seen at higher altitudes that match those below the transition. The same behaviour was observed on independent measurements of water vapor mixing ratio, which may suggest filamentation

associated with differential transport of drier air layer in a wind shear zone. We will discuss this vertical distribution of isotopes in more detail in Subsection 5.2.5.







Figure 6. Overview of airborne water vapor isotope measurements. (a) Isotope content δ^2H (permil) vs. specific humidity ($g\ kg^{-1}$). The mean mixing lines is computed following Noone (2012) with an end point $q = 1.37\ g\ kg^{-1}$ and $\delta^2H = -335\ ‰$, and reported using a black dotted line. (b) and (c) Vertical profiles of δ^2H (‰). Color indicates flight number. Panel (c) was added to separately plot the profiles with strong vertical gradients. Isotope data are averaged at 10s interval. The red cross associated with Flight 11 indicates that no data was relevant during that flight.

5.2 Cloud liquid water sampling

Four relevant cloud water samples have been taken with the CASCC on 18, 20 and 21 June (Table A1). The isotope contents of the cloud liquid water are shown in Fig. 7a (coloured circles). They are close to the global meteoric water line (GMWL) with corresponding d-excess values ranging from 12.1 to 14.8 ‰. Equilibrium condensates were estimated from relevant water vapor isotope measurements during the time when the cloud samples were taken (Fig. 7a, coloured squares) using the fractionation factors of Majoube (1971), and air temperature measurements at cloud level, ranging between -4 and $1^\circ C$. It is worth noting that given potential sources of uncertainty, the equilibrium condensate values agree remarkably well with the cloud liquid water during 18 June (green symbols), and the afternoon of 20 June (violet symbols). Overall, results here confirm that the cloud water droplets formed from equilibrium fractionation from ambient vapor. It is worth mentioning that such agreement between the completely independent sampling and measurement of vapor and cloud water supports the overall validity of i) the cloud water sampling protocol, ii) the airborne vapor isotope measurements and iii) the consistency of the calibrated water isotope dataset derived from the L-WAIVE campaign.



415 Figure 7. $\delta^2H - \delta^{18}O$ plots of liquid samples compared to the vapor isotope estimates. (a) Cloud water samples (circles), equilibrium
condensate (squares), and precipitation samples (grey triangles). (b) Equilibrium vapor from lake water samples at various depths
(dots and diamonds) compared to vapor isotope measurements from ULA-IC (upward triangle) and boat (downward triangle).
420 Colour denotes matching dates (in the blue boxes are the colour legends in dot representation). Gray colours show data where the
vapor samplings are not available. The times of the lake water sampling are indicated next to the dots with the same colours. Black
line denotes the Global Meteoric Water Line (GMWL).

5.3 Precipitation liquid water isotope composition

In total, 28 precipitation samples, of which 22 are unique, have been taken during the campaign, with sampling times lasting
from 20 min to several hours, depending on rainfall rate (Table A1). The isotope composition in rainfall ranges from -11.2 to
2.2‰ in $\delta^{18}O$ and -73.5 to 9.6 ‰ in δ^2H (Fig. 7a, grey triangles), with corresponding values of d-excess between 3.5 and
425 20.8‰ (not shown). Even if evaporation effects from the sampling setup cannot be fully excluded, in particular for the samples
with a sampling duration of more than 3h, the consistency of duplicate samples indicate that the influence of sampling artefacts
can overall be considered as limited. Notably, there is an overall correspondence between the isotope range observed in cloud
water samples and in a majority of precipitation samples. Deviations of precipitation samples from the GMWL indicate
potential below-cloud exchange processes between the falling raindrops and the ambient water vapor, leading to an enrichment
430 of the precipitation in ^{18}O (Graf et al., 2019; Worden et al., 2007). We have noticed that samples taken between 12 to 15 June,
and on 21 June are most enriched in ^{18}O , exhibiting the higher values in $\delta^{18}O$ and δ^2H and a smaller d-excess. These samples
are from rainfall events associated with local thunderstorms. The corresponding low d-excess (even negative, ~-8‰) of these
samples may point to the partial evaporation of rain droplets during their fall.

5.4 Lake liquid water isotope composition

435 Evaporation from the Annecy lake is expected to be an important source for the water vapor in the low troposphere above the
Annecy valley. In order to link the atmospheric profiles of water vapor isotopes to the lake as a moisture source, 20 lake water
samples (Table A1) were taken throughout the campaign within the lake-atmosphere interface layer (6 samples), as well as
close to 2 m depth (14 samples), and analysed for their stable water isotope composition. The average isotope contents were
found to be -8.3 ± 1.5 ‰ for $\delta^{18}O$, and -63.0 ± 6.0 ‰ for δ^2H . To assess the isotopic balance between the lake water and the
440 water vapor isotope composition measured by ULA and boat, we calculated the equilibrium vapor for the lake surface
temperatures measured during respective days (Fig. 7b) using the equilibrium fractionation factors of Majoube (1971).

Lake water samples taken close to the surface are expected to be most affected by evaporation, causing deviations to the right
from the GMWL along evaporation lines (enrichment in ^{18}O), and corresponding to a negative d-excess. As a matter of fact,
lake water samples clearly cluster on the right-hand side of the GMWL (Fig. 7b, diamonds), whereas vapor data from the
445 ULA-IC (triangles) and the boat (downward triangle) are to the left, with a relatively consistent range of isotope values. The
corresponding d-excess underline this result for the 6 lake water samples from the lake-atmosphere interface layer, with values
of the d-excess ranging between 2.2 and -19.6‰ and an average value of -6.3‰. For the samples taken at a depth of 2 m

(Fig. 7b, diamond symbols), the derived d-excess showed less evaporation influence, with a mean of 0.6‰, and a standard deviation of 8.2‰. While some sampling artifacts cannot be fully excluded, the fact that there are differences between the isotope contents at the lake surface and 2 m depth points to the impact of ongoing evaporation, that decreases with depth. We note that temperature profiles taken within the lake show a typically strong thermocline below about 7 m depth during summer (Danis et al., 2003). Therefore, freshwater input from runoff and precipitation, and loss to evaporation are expected to primarily affect the warmer, upper mixed layer within the timeframe of the campaign.

The range of equilibrium vapor for δ^2H observed in lake water for 16 and 17 June (Fig. 7b, blue and yellow dots) matches to first order with the range of δ^2H from ULA-IC. However, the $\delta^{18}O$ in equilibrium vapor is substantially higher than the measured vapour value. This pleads for a kinetic (non-equilibrium) fractionation during lake evaporation. During 18 June (Fig. 7b, green dot), equilibrium vapor of the liquid samples has a lower heavy isotope content than observed by ULA-IC, pointing to the influence of other factors on either atmospheric or lake water composition on these days. However, one should be rather cautious about this interpretation, as the time of the lake water sampling (~9h LT) was significantly different from the time of the flight (flight 9 of ULA-IC, ~12h LT), while for 16 and 17 June, the measurements were performed at the same time. It is worth noting that Lake Annecy is fed by a catchment whose surface is 10 times larger than that of the lake via ten main tributaries located on the lake periphery. The flows of its tributaries increase significantly in situations of heavy rain, which can substantially influence the isotopic content of the lake during the course of a year; however on 17 and 18 June no rain was observed.

6 Synthesis and discussion

During the two main sunny days of the field campaign (17 and 18 June), the evolution of isotope content profiles in the low troposphere (up to 3.5 km a.m.s.l.) is analysed together with the other observations including lidar measurements. In a second step, we provide a statistical overview of all the available data to compare the various samples from the lake water to the low troposphere, including clouds, and precipitation.


6.1 Vertical profiles in the low troposphere as observed from ULAs

Fig. 8 shows the vertical profiles corresponding to 4 coordinated flights between the 2 ULAs. These flights were carried out in the late morning and afternoon during the two sunny days of the campaign, i.e. 17 and 18 June. The profiles were constructed by averaging the data over 100 m bins. This operation makes it possible to improve the signal-to-noise ratio and to evaluate the dispersion of the measurements as a function of altitude (coloured areas in the figure).

The vertical structures of the lower troposphere are well reproduced across all data types, showing the consistency of the isotopic profiles with the independent observations. The strong gradients on δ^2H above 2.5 km a.m.s.l. are related to the transition to the free troposphere and are confirmed by the lidar observations and meteorological measurements on the two

ULAs. Discrepancies appear between the specific humidity measured from the 2 ULAs on 18 June above 2.7 km a.m.s.l. (Fig. 8c). This can be explained by the presence of a fractional cloud cover. The ULA-IC has crossed part of the cloud base which is at the interface with the free troposphere.

In the morning, the atmosphere near the lake shows near-surface stability over 200 to 300 m which deteriorates/mixes away in the afternoon according to the potential temperature profiles. A few hundred meters above the lake, the atmosphere is very stable on 17 June in the morning (Fig. 8a); it then loses stability in the afternoon due to thermal convection (Fig. 8b). On 18 June morning, the stability is strong below 1.5 km a.m.s.l. (Fig. 8c) and approaches the neutrality above this altitude. In contrast to 17 June, an atmosphere close to neutrality is observed on 18 June afternoon (Fig. 8d), which may favour vertical exchanges from the lake surface to the free troposphere. An important difference between the two days is also related to the surface wind speed. Winds of the order of 5 ms^{-1} were present on 17 June, whereas they were close to null on 18 June. Therefore, significant differences in water exchange between the lake and the atmosphere can be expected between the two days. These differences have already been noted when comparing the samples in Fig. 7b. On 17 June, wind-forced evaporation phenomena may therefore intervene which may explain at least part of the assumed isotopic fractionation. On the other hand, on 18 June, we have seen that the water samples are hardly representative for ULA measurements as they were taken early in the morning, before the sun lit up the valley. Around noon, the structure of about 300 m observed at the bottom of the $\delta \text{}^2\text{H}$ profile (Fig. 8d) and found in the other data is certainly strongly influenced by evaporation from the lake. It is significantly attenuated in the afternoon due to vertical mixing linked to thermal convection. We observe an enrichment in $\text{}^2\text{H}$ in this layer compared to the rest of the vertical profile in the morning that may be due to the lake evaporation into a depleted air.



500 **Figure 8. Vertical profiles of the isotope composition δ^2H from CRDS (ULA-IC), specific humidity (from WALI and meteorological probes onboard ULAs), potential temperature (from meteorological probe of ULA-A), wind speed components from the wind lidar, and apparent scattering ratio (ASR) derived from the airborne lidar ALiAS (ULA-A) on a) 17 June morning (~10:30-12:00), b) 17 June afternoon (~16:00-18:30), c) 18 June morning (~11:00-12:30), and 18 June afternoon (~16:00-18:00). The dispersion on the data is represented by the colored areas.**

6.2 Statistical overview of isotopic content from the lake to the atmosphere

505 The main atmospheric vertical structures are derived from the profiles of potential temperature, aerosols, wind and water vapor mixing ratio. We have therefore defined 3 layers in the low troposphere: i) between the lake surface to 1 km a.m.s.l., ii) between 1 and 2.5 km a.m.s.l., and iii) between 2.5 and 3.5 km a.m.s.l. The uppermost value represents our altitude sampling limit. The first layer can be identified as the lake boundary layer, the second layer is associated to the region of influence of the lake, below the altitude of the surrounding mountains, and the third layer is the transition towards the free troposphere influenced by the regional circulation. The location in altitude of these different layers obviously depends on the weather conditions and the time of day.

510 Using the vertical layering introduced above, the overall results for stable isotopes in water are summarized in Fig. 9 on a statistical basis selecting the highest quality data. This synthesis is presented in the form of whisker boxes for d-excess (Fig. 9a), δ^2H (Fig. 9b) and $\delta^{18}O$ (Fig. 9c). For each sub-figure, the upper part depicted the statistical content of isotope in water vapor for each sampled altitude range in the atmosphere, while the lower part describes the same thing for liquid water samples collected from clouds, precipitations and in the lake, at the interface and sub-surface. For the atmospheric water vapor, 515 the sample number n is significant (numbers given at the right of the whisker boxes in Fig. 9b) and it makes sense to accurately assess the confidence intervals at 95%. As the notches do not overlap from an atmospheric layer to another, we can conclude, with 95% confidence, that the true medians do differ. Hence, we highlight a significant variability in the stable isotope content of water vapor depending on the atmospheric layers previously identified.


520 Between the surface of the lake and 2.5 km a.m.s.l., we find $\delta^{18}O$ values are in the lower range to those recorded by Craig and Gordon (1965) for evaporation over the Mediterranean [-15,-10] ‰. On the other hand, they report much more dispersed d-excess values ranging from 5 to 35‰. Gat (2000) gives narrower values for marine and European air masses with a d-excess interval of [7-11] ‰. Our observations are mainly outside this interval for the first two layers, they overlap it for the 2.5-3.5 km a.m.s.l. layer more influenced by the synoptic scale. Measurements from the boat on the last day of the campaign within 1 m above the lake are overall consistent with the vapor isotope measurements made by ULA-IC on previous days in the lake 525 boundary layer.

Based only on the statistical information content provided by the whisker boxes profiles, it is difficult to see precisely whether the evaporation of lake water significantly influences the first layers of atmosphere just above the lake. Indeed, different factors can influence the relation between water vapor measured by the ULA above the lake, and the evaporating water vapor. In addition to the immediate evaporate, water vapor can be advected and mixed in from other locations horizontally, but also

530 vertically. In particular for a lake with coastal, vegetated areas within less than about 2 km distance anywhere on the lake, influences from evapotranspiration are likely within the lower 1 km above the lake.

Our liquid water samples are not numerous enough to define a confidence interval. As explained previously, we nevertheless note an enrichment in heavy isotope of the surface layer of the lake which is directly in contact with the atmosphere and therefore directly subject to evaporation. The 2 m depth layer is significantly less enriched as evidenced by their isotope
535 content. We highlight therefore a significant enrichment in heavy isotopes of surface water associated with low values of d-excess, which is characteristic of evaporation as described by Gibson and Edwards (2002). Measurements of $\delta^{18}O$ profiles in the Annecy lake have already been carried out between the years 2000 and 2002. They have shown $\delta^{18}O$ values of the order of -9.5‰ for water above the thermocline of "Petit Lac" during the summer months (Danis, 2009), in agreement with our results. This indicates a year-to-year stability of the isotopic content of the lake water below the surface microlayer.

540 From a first analysis of our successful airborne in-situ sampling of water vapor isotopes and cloud water, we conclude that cloud water formed in an equilibrium fractionation process for the samples collected here, and the cloud water isotopes were relatively similar to local precipitation (taken on other days). Since 1992, the Global Network of Isotopes in Precipitation (GNIP, <https://nucleus.iaea.org/wiser>) regularly sample rainwater at the Thonon-les-Bains city (60 km northeast from Annecy) for isotope analyses. Based on data available until 2018, typical summer $\delta^{18}O$ ($\delta^{2}H$) values fall within the interval [-16, -1]
545 ‰ ([-120, -20] ‰), encompassing our precipitation and cloud water measurements. It should be noted that the lake water is poorer in heavy isotopes than rain and cloud water. This difference is probably due to snowmelt during spring and the contribution of precipitation falling at high altitude. Note that GNIP data show median values of -10.5‰ (-78.6‰) for $\delta^{18}O$ ($\delta^{2}H$) during winter versus -6.5‰ (-43.5‰) during summer which agree with the studies of Dutton et al. (2005) and, Kendall and Coplen (2001).



550

Figure 9. Whisker boxes for a) d-excess, b) δ^2H and c) $\delta^{18}O$. For each sub-figure, the upper part depicted the statistical content of isotope in water vapor and the bottom part in the liquid water. For each whisker box, the central continuous vertical line indicates

555 the median (50th percentile, P_{50}), and the bottom and top edges of the box indicate the 25th (P_{25}) and 75th (P_{75}) percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers, which are values outside the interval bounded by the 5th and 95th percentiles. For the atmospheric water vapor, the number of samples n is indicated close to the boxes. The confidence interval at 95% defined by $[(P_{50} - 1.57 \cdot (P_{75} - P_{25})/\sqrt{n}), (P_{50} + 1.57 \cdot (P_{75} - P_{25})/\sqrt{n})]$ is represented by the notches in the upper part of the sub-figure.

7 Conclusion

560 The L-WAIVE field campaign documented for the first-time vertical profiles of stable water isotopes over an alpine lake valley, in combination with lidar and meteorological measurements. Within L-WAIVE, we have developed an innovative approach to retrieve the vertical profiles and their heterogeneity over the Annecy lake in several meteorological situations. The field campaign took place between 12 and 23 June 2019 and used the complementary and/or synergistic measurements from different means of sampling gaseous and liquid water, atmospheric scattering layers and meteorological parameters (temperature, moisture, and wind). Continuous high vertical resolution profiles of tropospheric water vapor, temperature, and 565 wind as well as aerosols in the Annecy valley have been performed using ground-based lidars. They were acquired together with boat-borne and airborne measurements of stable water isotopes (H_2O , H^2HO and H_2^{18}O) in the lake, in the lower atmosphere as well as in precipitation and in-cloud using an original sampling method from ULAs. The atmospheric structure over the Annecy lake has been simultaneously measured using an nadir-looking airborne lidar coupled with a meteorological probe on a second ULA.

570 Strong temporal variability of several measured parameters has been observed during the field campaign, with a marked diurnal cycle influenced by the valley winds and the vertical thermal stability. The valley winds particularly influence the water vapour composition of the atmospheric layers near the lake, over a thickness of about 300 m. Thermal instabilities which evolve strongly during the day, and which can be associated with cloud forcing at the base of the free troposphere (between 2.5 and 3.5 km a.m.s.l.), will rapidly mix water vapour throughout the air column between the lake and the base of the free troposphere. 575 This process makes it difficult to observe the influence of evaporation from the surface waters of the lake. We also note strong vertical gradients in the abundance of isotopes at elevations higher than 2.5 km a.m.s.l., with a marked decrease at the interface with the free troposphere. Such vertical structures are confirmed by the other types of measurements, including lidar profiles of aerosol backscatter. The altitude at which these gradients were observed exhibited a day-to-day variation throughout the measurement campaign in relationship with stratification in the valley, weather conditions, and topography.

580 We note that the cloud water samples are close to the GMWL, (d-excess between 12.1 and 14.8 ‰), which indicates that the cloud water formed from equilibrium fractionation of ambient atmospheric water vapor. There is an overall correspondence between the isotope range observed in cloud water samples and a majority of precipitation samples (d-excess between -8.6 and 20.8/‰). The deviations observed on precipitation samples that are below the GMWL indicate potential below-cloud exchange, or post-condensational exchange processes. It is also worth noting that the average isotope composition of the lake 585 water, taken at 2 m depth, appears different from the one for the surface microlayer sample; this may be due to evaporation

processes. The $\delta^{18}\text{O}$ in equilibrium condensate above the lake is generally substantially more depleted, confirming the existence of non-equilibrium fractionation that could be due to both lake evaporation and air mass advection.

During the two sunny days of 17 and 18 June, it was noted that the effect of the lake on the water vapour content of the valley is difficult to isolate because it involves different processes, including the variability of thermal convection and horizontal advection processes associated with valley winds. Nevertheless, this effect appears to be detectable on a layer of about 300 m thickness above the lake in light wind conditions. The field experiment that was carried out is complex, however it would need to be repeated several times in order to have representative statistics on favourable cases such as the one on 18 June, i.e. with negligible surface winds.

Data availability. Data can be downloaded from <https://metclim-lidars.aeris-data.fr/en/homepage/> (last access: 17 June 2021) upon request to the first author of the paper. ERA5 data were downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store (<https://cds.climate.copernicus.eu#!/search?text=era5&type=dataset>, last access: 17 June 2016).

Author contributions. Patrick Chazette coordinated the field campaign and wrote the paper, with contributions from Cyrille Flamant, Harald Sodemann, Anne Monod, Elsa Dieudonné and Julien Totems. Harald Sodemann and Andrew Seidl sampled and processed the stable water isotope data. Julien Totems calibrated WALI and derived the water vapor profiles. Patrick Chazette, Harald Sodemann, Julien Totems, Anne Monod, Elsa Dieudonné, Alexandre Baron, Andrew Seidl, Pascal Doira, Amandine Durand, and Sylvain Ravier participated in the field campaign. Hans Christian Steen-Larsen and all the authors contributed to proofreading of the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgments. Friendly acknowledgements to local authorities of the town of Lathuile R. Aumaître, H. Bourne, F. Lambert; instrument providers F. Arthoud (USMB); D. Crucciani for its welcome at the Delta Evasion airfield; ULA pilots F., L., F. Toussaint; technical assistance F. Maignan and C. Diana. This research was funded by the WaVIL project (Water Vapor and Isotope Lidar, Agence Nationale de la Recherche grant n° ANR-16-CE01-0009). HS and AS received funding from the Research Council of Norway with the FARLAB project (Contract no. 245907) and the ERC-2017-CoG ISLAS (Grant Agreement Nr. 773245). The IAEA/WMO Global Network of Isotopes in Precipitation (GNIP) is acknowledged for the access to its database accessible at: <https://nucleus.iaea.org/wiser> (last access: 17 June 2016).

References

Barthlott, C., Corsmeier, U., Meißner, C., Braun, F. and Kottmeier, C.: The influence of mesoscale circulation systems on triggering convective cells over complex terrain, *Atmos. Res.*, 81(2), 150–175, doi:10.1016/j.atmosres.2005.11.010, 2006.
Behrendt, A.: Temperature Measurements with Lidar, in *Lidar: Range-Resolved Optical Remote Sensing of the Atmosphere*, pp. 273–305., doi: 10.1007/0-387-25101-4_10, 2006.

- Berkelhammer, M., Noone, D. C., Wong, T. E., Burns, S. P., Knowles, J. F., Kaushik, A., Blanken, P. D. and Williams, M. W.: Convergent approaches to determine an ecosystem's transpiration fraction, *Global Biogeochem. Cycles*, 30(6), 933–951, doi:10.1002/2016GB005392, 2016.
- 620 Berthier, S., Pelon, J., Chazette, P., Couvert, P., Sèze, G., Bréon, F.-M., Laland, M., Winker, D. and Pain, T.: Cloud statistics from spaceborne backscatter lidar data analysis, in European Space Agency, (Special Publication) ESA SP, vol. 2., 2004.
- Chazette, P. and Totems, J.: Mini N2-Raman Lidar onboard ultra-light aircraft for aerosol measurements: Demonstration and extrapolation, *Remote Sens.*, 9(12), doi:10.3390/rs9121226, 2017.
- 625 Chazette, P., Couvert, P., Randriamiarisoa, H., Sanak, J., Bonsang, B., Moral, P., Berthier, S. S., Salanave, S. and Toussaint, F.: Three-dimensional survey of pollution during winter in French Alps valleys, *Atmos. Environ.*, 39(6), 1035–1047, doi:10.1016/j.atmosenv.2004.10.014, 2005.
- Chazette, P., Sanak, J. and Dulac, F.: New approach for aerosol profiling with a lidar onboard an ultralight aircraft: application to the African Monsoon Multidisciplinary Analysis., *Environ. Sci. Technol.*, 41(24), 8335–8341, doi:10.1021/es070343y, 2007.
- 630 Chazette, P., Dabas, a., Sanak, J., Lardier, M. and Royer, P.: French airborne lidar measurements for Eyjafjallajökull ash plume survey, *Atmos. Chem. Phys.*, 12(15), 7059–7072, doi:10.5194/acp-12-7059-2012, 2012.
- Chazette, P., Marnas, F. and Totems, J.: The mobile Water vapor Aerosol Raman Lidar and its implication in the framework of the HyMeX and ChArMEx programs: application to a dust transport process, *Atmos. Meas. Tech.*, 7(6), 1629–1647, doi:10.5194/amt-7-1629-2014, 2014.
- 635 Chazette, P., Totems, J. and Shang, X.: Atmospheric aerosol variability above the Paris Area during the 2015 heat wave - Comparison with the 2003 and 2006 heat waves, *Atmos. Environ.*, 170, 216–233, doi:10.1016/j.atmosenv.2017.09.055, 2017.
- Chazette, P., Totems, J., Baron, A., Flamant, C. and Bony, S.: Trade-wind clouds and aerosols characterized by airborne horizontal lidar measurements during the EUREC4A field campaign, *Earth Syst. Sci. Data*, 12(4), 2919–2936, doi:10.5194/essd-12-2919-2020, 2020.
- 640 Craig, H. and Gordon, L. I.: Deuterium and oxygen 18 variations in the ocean and the marine atmosphere. In *Stable Isotopes in Oceanographic Studies and Paleotemperatures*, edited by E. Tongiorgi, Laboratorio di Geologia Nucleare, Pisa., 1965.
- Cui, B. L., Li, X. Y. and Wei, X. H.: Isotope and hydrochemistry reveal evolutionary processes of lake water in Qinghai Lake, *J. Great Lakes Res.*, 42(3), 580–587, doi:10.1016/j.jglr.2016.02.007, 2016.
- 645 Cunliffe, M., Engel, A., Frka, S., Gašparović, B. Ž., Guitart, C., Murrell, J. C., Salter, M., Stolle, C., Upstill-Goddard, R. and Wurl, O.: Sea surface microlayers: A unified physicochemical and biological perspective of the air-ocean interface, *Prog. Oceanogr.*, 109, 104–116, doi:10.1016/j.pocan.2012.08.004, 2013.
- Danis, P.-A.: Modélisation du fonctionnement thermique , hydrologique et isotopique de systèmes lacustres : sensibilité aux changements climatiques et amélioration des reconstructions paléoclimatiques To cite this version : HAL Id : tel-00360067 Mod ´ elisation du fonc, [online] Available from: <https://tel.archives-ouvertes.fr/tel-00360067> (Accessed 5 November 2020), 2009.
- 650 Danis, P. A., Von Grafenstein, U. and Masson-Delmotte, V.: Sensitivity of deep lake temperature to past and future climatic changes: A modeling study for Lac d'Annecy, France, and Ammersee, Germany, *J. Geophys. Res. Atmos.*, 108(19), doi:10.1029/2003jd003595, 2003.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16(4), 436–468, doi:10.3402/tellusa.v16i4.8993, 1964.
- 655 Drobinski, P., Steinacker, R., Richner, H., Baumann-Stanzer, K., Beffrey, G., Benech, B., Berger, H., Chimani, B., Dabas, A., Dorninger, M., Dürr, B., Flamant, C., Frioud, M., Furger, M., Gröhn, I., Gubser, S., Gutermann, T., Häberli, C., Häller-Scharnhost, E., Jaubert, G., Lothon, M., Mitev, V., Pechinger, U., Piringer, M., Ratheiser, M., Ruffieux, D., Seiz, G., Spatzierer, M., Tschannett, S., Vogt, S., Werner, R. and Zängl, G.: Föhn in the Rhine Valley during MAP: A review of its multiscale dynamics in complex valley geometry, *Q. J. R. Meteorol. Soc.*, 133(625), 897–916, doi:10.1002/qj.70, 2007.
- 660 Dutton, A., Wilkinson, B. H., Welker, J. M., Bowen, G. J. and Lohmann, K. C.: Spatial distribution and seasonal variation in 18O/16O of modern precipitation and river water across the conterminous USA, *Hydrol. Process.*, 19(20), 4121–4146, doi:10.1002/hyp.5876, 2005.
- Flamant, C., Drobinski, P., Nance, L., Banta, R., Darby, L., Dusek, J., Hardesty, M., Pelon, J. and Richard, E.: Gap flow in an Alpine valley during a shallow south fo`hn event: Observations, numerical simulations and hydraulic analogue, *Q. J. R. Meteorol. Soc.*, 128(582), 1173–1210, doi:10.1256/003590002320373256, 2002.
- 665 Gat, J. R.: Oxygen and hydrogen isotopes in the hydrologic cycle, *Annu. Rev. Earth Planet. Sci.*, 24(1), 225–262,

doi:10.1146/annurev.earth.24.1.225, 1996.

Gat, J. R.: Atmospheric water balance-the isotopic perspective, in *Hydrological Processes*, vol. 14, pp. 1357–1369, John Wiley & Sons Ltd., 2000.

670 Gat, J. R.: *Isotope Hydrology - A study of the water cycle*, edited by T. K. Wei, Published by Imperial College Press and distributed by World Scientific Publishing Co., Singapore., 2010.

Gat, J. R., Bowser, C. J. and Kendall, C.: The contribution of evaporation from the Great Lakes to the continental atmosphere: estimate based on stable isotope data, *Geophys. Res. Lett.*, 21(7), 557–560, doi:10.1029/94GL00069, 1994.

675 Gibson, J. J. and Edwards, T. W. D.: Regional water balance trends and evaporation-transpiration partitioning from a stable isotope survey of lakes in northern Canada, *Global Biogeochem. Cycles*, 16(2), 10-1-10–14, doi:10.1029/2001gb001839, 2002.

Graf, P., Wernli, H., Pfahl, S. and Sodemann, H.: A new interpretative framework for below-cloud effects on stable water isotopes in vapour and rain, *Atmos. Chem. Phys.*, 19(2), 747–765, doi:10.5194/acp-19-747-2019, 2019.

680 Griffis, T. J., Wood, J. D., Baker, J. M., Lee, X., Xiao, K., Chen, Z., Welp, L. R., Schultz, N. M., Gorski, G., Chen, M. and Nieber, J.: Investigating the source, transport, and isotope composition of water vapor in the planetary boundary layer, *Atmos. Chem. Phys.*, 16(8), 5139–5157, doi:10.5194/acp-16-5139-2016, 2016.

He, H. and Smith, R. B.: Stable isotope composition of water vapor in the atmospheric boundary layer above the forests of New England, *J. Geophys. Res. Atmos.*, 104(D9), 11657–11673, doi:10.1029/1999JD900080, 1999.

685 Herrmann, H.: Kinetics of Aqueous Phase Reactions Relevant for Atmospheric Chemistry, *Chem. Rev.*, 103(12), 4691–4716, doi:10.1021/cr020658q, 2003.

Jasechko, S., Sharp, Z. D., Gibson, J. J., Birks, S. J., Yi, Y. and Fawcett, P. J.: Terrestrial water fluxes dominated by transpiration, *Nature*, 496(7445), 347–350, doi:10.1038/nature11983, 2013.

Kendall, C. and Coplen, T. B.: Distribution of oxygen-18 and deuterium in river waters across the United States, *Hydrol. Process.*, 15(7), 1363–1393, doi:10.1002/hyp.217, 2001.

690 Koscielny, A. J., Doviak, R. J. and Zrnica, D. S.: An Evaluation of the Accuracy of Some Radar Wind Profiling Techniques, *J. Atmos. Ocean. Technol.*, 1(4), 309–320, doi:10.1175/1520-0426(1984)001<0309:AEOTAO>2.0.CO;2, 1984.

Kottmeier, C., Kalthoff, N., Barthlott, C., Corsmeier, U., Van Baelen, J., Behrendt, A., Behrendt, R., Blyth, A., Coulter, R., Crewell, S., Di Girolamo, P., Dorninger, M., Flamant, C., Foken, T., Hagen, M., Hauck, C., Höller, H., Konow, H., Kunz, M., Mahlike, H., Mobbs, S., Richard, E., Steinacker, R., Weckwerth, T., Wieser, A. and Wulfmeyer, V.: Mechanisms initiating deep convection over complex terrain during COPS, *Meteorol. Zeitschrift*, 17(6), 931–948, doi:10.1127/0941-2948/2008/0348, 2008.

695 Lesouëf, D., Gheusi, F., Chazette, P., Delmas, R. and Sanak, J.: Low Tropospheric Layers Over Reunion Island in Lidar-Derived Observations and a High-Resolution Model, *Boundary-Layer Meteorol.*, 149(3), 425–453, doi:10.1007/s10546-013-9851-9, 2013.

700 Majoube, M.: Fractionnement en oxygène 18 et en deutérium entre l'eau et sa vapeur, *J. Chim. Phys.*, 68, 1423–1436, doi:10.1051/jcp/1971681423, 1971.

Mayr, G. J., Armi, L., Gohm, A., Zängl, G., Durran, D. R., Flamant, C., Gaberšek, S., Mobbs, S., Ross, A. and Weissmann, M.: Gap flows: Results from the Mesoscale Alpine Programme, *Q. J. R. Meteorol. Soc.*, 133(625), 881–896, doi:10.1002/qj.66, 2007.

705 Noone, D.: Pairing measurements of the water vapor isotope ratio with humidity to deduce atmospheric moistening and dehydration in the tropical midtroposphere, *J. Clim.*, doi:10.1175/JCLI-D-11-00582.1, 2012.

Raut, J.-C. C. and Chazette, P.: Assessment of vertically-resolved PM10 from mobile lidar observations, *Atmos. Chem. Phys.*, 9(21), 8617–8638, doi:10.5194/acp-9-8617-2009, 2009.

710 Salmon, O., Welp, L. R., Baldwin, M., Hajny, K., Stirm, B. and Shepson, P.: Vertical profile observations of water vapor deuterium excess in the lower troposphere, *Atmos. Chem. Phys.*, 19(17), 11525–11543, doi:10.5194/acp-19-11525-2019, 2019.

Sodemann, H., Aemisegger, F., Pfahl, S., Bitter, M., Corsmeier, U., Feuerle, T., Graf, P., Hankers, R., Hsiao, G., Schulz, H., Wieser, A. and Wernli, H.: The stable isotopic composition of water vapour above Corsica during the HyMeX SOP1 campaign: Insight into vertical mixing processes from lower-tropospheric survey flights, *Atmos. Chem. Phys.*, 17(9), 6125–6151, doi:10.5194/acp-17-6125-2017, 2017.

715 Steen-Larsen, H. C., Johnsen, S. J., Masson-Delmotte, V., Stenni, B., Risi, C., Sodemann, H., Balslev-Clausen, D., Blunier,

- 720 T., Dahl-Jensen, D., Ellehøj, M. D., Falourd, S., Grindsted, A., Gkinis, V., Jouzel, J., Popp, T., Sheldon, S., Simonsen, S. B., Sjolte, J., Steffensen, J. P., Sperlich, P., Sveinbjörnsdóttir, Á. E., Vinther, B. M. and White, J. W. C.: Continuous monitoring of summer surface water vapor isotopic composition above the Greenland Ice Sheet, *Atmos. Chem. Phys.*, 13(9), 4815–4828, doi:10.3929/ethz-b-000067919, 2013.
- Steen-Larsen, H. C., Sveinbjörnsdóttir, A. E., Peters, A. J., Masson-Delmotte, V., Guishard, M. P., Hsiao, G., Jouzel, J., Noone, D., Warren, J. K. and White, J. W. C.: Climatic controls on water vapor deuterium excess in the marine boundary layer of the North Atlantic based on 500 days of in situ, continuous measurements, *Atmos. Chem. Phys.*, 14(15), 7741–7756, doi:10.5194/acp-14-7741-2014, 2014.
- 725 Totems, J., Chazette, P. and Raut, J.: Accuracy of current Arctic springtime water vapour estimates, assessed by Raman lidar, *Q. J. R. Meteorol. Soc.*, 145(720), 1234–1249, doi:10.1002/qj.3492, 2019.
- Weng, Y., Touzeau, A. and Sodemann, H.: Correcting the impact of the isotope composition on the mixing ratio dependency of water vapour isotope measurements with cavity ring-down spectrometers, *Atmos. Meas. Tech.*, 13(6), 3167–3190, doi:10.5194/amt-13-3167-2020, 2020.
- 730 Worden, J., Noone, D., Bowman, K., Beer, R., Eldering, A., Fisher, B., Gunson, M., Goldman, A., Herman, R., Kulawik, S. S., Lampel, M., Osterman, G., Rinsland, C., Rodgers, C., Sander, S., Shephard, M., Webster, C. R. and Worden, H.: Importance of rain evaporation and continental convection in the tropical water cycle, *Nature*, doi:10.1038/nature05508, 2007.

Appendix A: L-WAIVE data availability

Table A1. Data available during the period of the field experiment between June 11 and 23, 2020. The cross "X" indicates that the ULA or instrument was operated on a half-day basis. The malfunctions are denoted by "M". The number of flights is indicated per ULM for each half-day. For in situ samplings, the number of samples per half-day is indicated. During the first two flights of the ULA-IC, the iMet sonde was not operating (denoted by bold italic numbers). A sequence of ULA-IC flights experienced saturation in the inlet, that requires data filtering (denoted by bold numbers).

Instrument	11	12	13	14	15	16	17	18	19	20	21	22	23											
Ground-based measurements																								
Lidar WALI		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	M				
Wind lidar WLS100					X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Precipitation sampling	1	3	1			1	1	1	10											2	2			
Airborne measurements																								
ULA-A			M	X	X		X		X	X	X	X	X	X	X	X	X	X	X			X	X	X
	ALiAS				1	1		2		1	1	2	2	2	1	2	1	1	1	1		1	1	1
ULA-IC			X	X	X		X		X	X	X	X	X	X	X	X	X	X	X					
	CRDS		<i>I</i>	<i>I</i>	1		1			1	1	1	2	1	1	2	1	1						
	CASCC				M					M				1			1	1	1					
UAV				1	1				1			1	1	1		1								
Boat measurements in the lake																								
Temperature sounding					1					1		2	2	2	1	1		1		1				
Isotope sampling	Micro-layer				1					1		1	1		1			M		1				
	2 m deep				1					1		1	1		1			1		1				

Boat measurements in the atmosphere

CRDS

X

Appendix B: Ultra-light aircraft flights description

Table B1. Flights characteristics for the remote sensing payload (ULA-A).

Flight	Date & time (LT*) dd/mm HHMM-HHMM	Maximum flight altitude during slow spiral ascent (km a.m.s.l.)	Comment
F01	12/06 1845-1850	-	Aborted
F02	13/06 1000-1200	3.5	Profile
F03	13/06 1800-1930	3.8	Profile & transverse
F04	14/06 1630-1700	2.5	Profile
F05	14/06 1730-1830	4.1	Profile
F06	16/06 1030-1100	3.5	Profile
F07	16/06 1700-1800	3.5	Profile & lake axis
F08	17/06 0900-0940	3.4	Profile
F09	17/06 1030-1150	3.7	Profile & transverse
F10	17/06 1545-1700	3.7	Profile & transverse
F11	17/06 1745-1900	3.7	Profile & transverse
F12	18/06 0830-0945	3.7	Profile
F13	18/06 1100-1230	3.7	Profile & transverse
F14	18/06 1550-1715	4.1	Profile & transverse
F15	19/06 0845-1005	3.5	Profile & transverse
F16	19/06 1115-1250	4.5	Profile
F17	19/06 1535-1720	3.7	Profile & other valley
F18	20/06 1210-1335	4.3	Profile & lake axis
F19	20/06 1620-1725	4.0	Profile & lake axis
F20	21/06 1130-1240	3.5	Profile & lake axis
F21	22/06 1100-1200	4.1	Profile
F22	22/06 1600-1650	3.6	Profile
F23	23/06 1120-1330	4.2	Profile & other valleys

745

*Local Time (LT)

Table B2. Flights characteristics for the in situ payload (ULA-IC).

Flight	Date & time (LT*) dd/mm HHMM-HHMM	Maximum flight altitude during slow spiral ascent (km a.m.s.l.)	Comment
F01	12/06 19:23-19:50	1.0	Local survey No iMet data
F02	13/06 10:25-12:50	1.6	Lake survey No iMet data
F03	13/06 18:22-19:49	2.2	Lake survey
F04	14/06 16:53-18:49	2.8	Lake survey
F05	16/06 16:30-18:44	2.6	Lake survey
F06	17/06 10:13-11:20	3.1	Lake survey
F07	17/06 16:35-18:20	3.6	Profiles
F08	18/06 10:30-11:31	2.4	Profiles
F09	18/06 11:56-12:51	3.5	Lake survey
F10	18/06 16:41-18:10	3.8	Clouds

			Partial isotope inlet saturation
F11	19/06 10:34-11:45	2.9	Profiles No isotope data due to isotope inlet saturation
F12	19/06 15:39-16:58	3.1	Lake survey Partial isotope inlet saturation
F13	19/06 17:49-19:18	4.7	Clouds Partial isotope inlet saturation
F14	20/06 11:04-12:39	3.5	Profiles Partial isotope inlet saturation
F15	20/06 14:49-17:16	3.4	Lake survey

*Local Time (LT)

750 **Appendix C: Water vapor stable isotopes interpretive framework**

The isotope composition δ^{2H} and δ^{18O} are given by the relationships:

$$\begin{cases} \delta^{2H} = R^{2H} / R^{2H}_S - 1 \\ \delta^{18O} = R^{18O} / R^{18O}_S - 1 \end{cases} \quad (C1)$$

Where R^S_X is the isotope ratio of Vienna Standard Mean Ocean Water (VSMOW) for the compound X and R_X the isotopic ratio given by

$$\begin{cases} R^{2H} = H^2HO / H_2O \\ R^{18O} = H_2^{18O} / H_2O \end{cases} \quad (C2)$$

The d-excess (d) is therefore defined as

$$d = \delta^{2H} - 8 \cdot \delta^{18O} \quad (C3)$$

755 to measure the deviation from equilibrium fractionation, noting the global average value for precipitation $d = 10\text{‰}$ (Dansgaard, 1964).

Appendix D: Water isotope data processing


760 D.1 Processing of liquid samples for stable water isotope analysis

Liquid water samples were processed at FARLAB, University of Bergen, according to established laboratory routines. For isotopic analysis, liquid water samples were transferred to 1.5 ml glass vials with rubber/PTFE septa (part #548-0907, VWR, USA). An autosampler (A0325, Picarro Inc) injected ca. 2 μ l per injection into a high precision vaporizer (A0211, Picarro Inc, USA) heated to 110°C. After blending with dry N₂ (< 5 ppm H₂O) the gas mixture was directed into the measurement cavity of a Cavity-Ring Down Spectrometer (L2140-i, Picarro Inc, Sunnyvale, USA) for about 7 min with a typical water concentration of 20 000 ppm. Three secondary laboratory standards were measured at the beginning and end of each batch for calibration purposes. Batches consisted typically of 20 samples, with laboratory drift standard DI, measured every 5 samples. For calibration according to International Atomic Energy Agency recommendations (IAEA, 2009), 16 injections of the laboratory standards EVAP ($\delta^2\text{H} = 4.75 \pm 0.11\text{‰}$, $\delta^{18}\text{O} = 5.03 \pm 0.02\text{‰}$) and GSM1, were used, and averaged over the beginning and end of each batch for calibration. Typical short-term reproducibility is 0.3‰ for $\delta^2\text{H}$ and 0.04‰ for $\delta^{18}\text{O}$. Long-term measurement accuracy is better than 0.66‰ for $\delta^2\text{H}$, 0.15‰ for $\delta^{18}\text{O}$, and 0.83‰ for the d-excess, evaluated from the 1-sigma standard deviation of the analysis of an internal laboratory standard over one year.

775 D.2 Processing of stable isotope measurements in water vapor

Raw measurements of pressure and water vapor mixing ratio of the CRDS (Picarro L2130-i, Ser. No. HIDS2254) were corrected using calibration functions established at the FARLAB laboratory, University of Bergen, Norway. Raw measurements of the isotope parameters, expressed as δ -values relative to VSMOW2 (Vienna Standard Mean Ocean Water 2), were corrected for the mixing ratio-isotope composition dependency using time-constant correction functions obtained for this analyzer according to the method of Weng et al. (2020). For calibration, vapor isotope data were transferred onto the VSMOW2-SLAP2 scale by routine measurements of secondary laboratory standards using a Picarro Standard Delivery Module (SDM, Part No. A0101, Picarro Inc.) before and after the campaign. Hereby, secondary laboratory standards GSM1 ($\delta^2\text{H} = -262.95 \pm 0.30\text{‰}$, $\delta^{18}\text{O} = -33.07 \pm 0.05\text{‰}$), and DI ($\delta^2\text{H} = -50.38 \pm 0.37\text{‰}$, $\delta^{18}\text{O} = -7.78 \pm 0.07\text{‰}$) were measured repeatedly for 20 min in the period 1 June 2019 to 19 July 2019 (Fig. D1). Calibrations of the CRDS analyser in the laboratory before and after the field deployment showed minor drift during the measurement period (GSM1 $\delta^{18}\text{O}$: 0.002 ‰/month, DI $\delta^{18}\text{O}$: -0.233 ‰/month, GSM1 $\delta^2\text{H}$: 1.045 ‰/month, DI $\delta^2\text{H}$: -0.853 ‰/month). From 2-sigma standard deviations of all valid SDM calibrations, the combined calibration uncertainty is estimated to be on the order of 0.2‰ for $\delta^{18}\text{O}$ and 2‰ for $\delta^2\text{H}$. A small bubbler system was used in the field for testing instrument drift before and after flight operations in the field. These tests indicated an anomaly in the CRDS measurements 18-20 June, partially affecting two flights in the morning of 19 June, and the first flight on 20 June. The anomaly was due to a saturated inlet system from condensate forming on the aircraft during a

cloud sampling flight on 18 June. Flight periods affected by the saturated inlet were excluded from further analysis, as well as the entire flight 11 of ULA-IC, and are indicated with a corresponding quality flag in the data files (Table B2).




795 Figure D1: Bias between nominal and raw delta-values obtained with the Standards Delivery Module with working standards
 DI and GSM1 on analyzer HIDS2254 before and after the L-WAIVE campaign period. Grey area denotes campaign period.
 Red lines denote the mean and 1-sigma standard deviation of all valid SDM calibrations in the period 1 May 2019 to 19 July
 2019. Black line is a linear regression line.

D.3 Sensitivity of CRDS instrument parameters to flight conditions

800 The CRDS analyser was mounted in the ULA frame without additional thermal insulation. Since the infrared spectrometry is
 sensitive to both, the pressure and temperature conditions in the measurement cavity, and the associated wavelength monitor
 (Gupta et al., 2009), we document here the potential impact of the exposed conditions on the ULA-IC measurement data at 2s
 time averaging. The cavity pressure in the CRDS is narrowly regulated to 50 Torr (66.67 hPa) on ground (Fig. D2a, black
 area). During flight, the spread of the distribution widens due to turbulence and vertical motion of the aircraft, albeit without
 805 inducing a bias to the cavity pressure (grey shading). The cavity temperature is regulated to 80.00 ± 0.02 °C on ground (Fig.
 D2b). During flight, a tail of lower temperatures develops, likely due to the constant cooling effect of airflow around the
 analyser, shifting the maximum of the distribution to slightly below 80°C (grey shading). Data points with cavity temperatures
 below 79.98°C can potentially be compromised, and have been flagged with a corresponding quality flag in the data set.
 810 Temperature of the warm box, housing the wavelength monitor of the CRDS, is maintained to 45.00 ± 0.02 °C on ground, with

815

some deviations to warmer temperatures (Fig. D2c). During flight, the maximum of the distribution shifts to 0.01 K lower temperatures, while retaining an overall similar variability compared to ground condition (grey shading). The temperature within the analyser (Data acquisition system temperature, Fig. D2d) is not regulated, and therefore most strongly affected by ambient conditions. While the overall influence of in-flight conditions appears relatively limited, with the large majority of data points remaining within specification range, insulation to the analyser housing could probably further increase stability of the instrument during flight. This will in particular be important for future operation of the ULA in colder climate zones, and other seasons.



820

Figure D2: CRDS instrument parameters on ground (black) and during flight (grey) with ULA-IC. (a) Distribution of cavity pressure (Torr), (b) cavity temperature (°C), (c) warm-box temperature (°C), and (d) data acquisition system temperature (°C) on ground (black) and during flight segments with level or slowly changing altitude (sequence flag 1 or 2). Data are averaged to a 2 s time interval.