

Interactive comment on “Smoke-charged vortices in the stratosphere generated by wildfires and their behaviour in both hemispheres: comparing Australia 2020 to Canada 2017” by Hugo Lestrelin et al.

Hugo Lestrelin et al.

legras@lmd.ens.fr

Received and published: 2 March 2021

[acpd,discussion]copernicus

bernard.legras@lmd.ipsl.fr Lestrelin, Legras, Podglajen, Salihoglu

Printer-friendly version

Discussion paper



Answer to reviewer #1

General comments

This paper details the dynamics of a newly-discovered phenomenon associated with large pyroCb smoke plumes in the stratosphere, namely the self-organized anticyclonic vortices that are formed due to absorption of solar radiation by black carbon within the plumes. A comparison is made between the 2019-2020 Australian plumes and several plumes associated with large Canadian fires in 2017. Detailed analysis of the Canadian plumes use the Lait Potential Vorticity (PV) from ERA5 to track the vortex evolution and to examine the composite dynamical structures of the vortex. ERA5 ozone is also used to successfully track the vortex locations. Composites of PV, temperature, and ozone help to further understand these features. The paper also details how these features are maintained in the analyses by assimilation of temperature and ozone data, and it examines various dynamical indices to test whether the vortices are in balance and/or inertially unstable. This paper provides an excellent addition to the study of smoke-induced dynamics.

We appreciate the evaluation of the reviewer and his numerous detailed comments.

Specific comments

- *Line 10: You use the term “low absolute potential vorticity” here. Just to be clear, does this mean low magnitude (i.e., absolute value) of the potential vorticity?*

We mean small absolute vorticity and small potential vorticity as well.

- *Line 83: I don't think “g is the free-fall acceleration” is necessary, since “g” isn't in Eq. (1).*

[Interactive comment](#)

[Printer-friendly version](#)

[Discussion paper](#)



We had omitted the required constant $-g$ factor in the definition. It is now restored.

- *Lines 112-114: Quoting from the paper, “we used both Π and the ozone anomaly defined as the deviation with respect to the mean at the same latitude and altitude.” So are both Π and ozone defined using the anomaly with respect to the zonal mean, or is just ozone calculated as the anomaly, while Π is the raw value from Eq. (2)? This sentence could be read either way.*

Only ozone is defined using the anomaly with respect to the zonal mean. The sentence has been modified.

- *Lines 112-114: As mentioned by the authors, the commonly used PV has a disadvantage of large background vertical gradient. While Khaykin et al. (2020) used relative vorticity, Kablick et al. (2020) used the “regular” PV to analyze the 2019-2020 Australian smoke plumes. They used the PV anomaly relative to the zonal mean in units of percent of the absolute value of the zonal mean PV. This alternate approach also reduces the influence of the large background gradient.*

We agree there are several ways in order to remove the background gradient in the display of the vertical structure. The Lait PV has the advantage to remain an adiabatic invariant of motion.

- *Line 139-142: You talk about the “kernel of almost zero PV and low ozone...” To visualize this statement, it would be interesting to see latitude/altitude cross-sections of PV and ozone along the CALIPSO track to compare with Figures 1 and 2.*

The latitude/altitude and longitude/altitude sections are displayed in the animation of vortex O in the supplement and are indeed very instructive. They are centered on the IFS vortex to provide continuity. Doing the same with vortices A, B1 and B2 brings less additional information. Figure 1 and 2 show the location of the vortex

[Printer-friendly version](#)[Discussion paper](#)

center according to the PV and ozone tracking. As these centers are minima in the 3D domain, surrounding contours can be plotted but this would totally obscure the aerosol pattern on many panels.

- *Line 147 (also Lines 158 and 394) : I am unfamiliar with the term “thalweg”. Are you referring to a trough? Could you provide some dynamical field on the maps to indicate where this is occurring to help visualize the point you’re making?*

We mean trough and this has been corrected. Montgomery potential contours have been added to all the isentropic charts of the animations.

- *Line 194-195: Is the statement “it differs from the 2020 case where such effect is not observed for any of the three vortices” referring to results in Khaykin et al. (2020)?*

Yes. The sentence has been improved.

- *Section 3.5. This section provides a helpful reference of other papers that have studied this event. The PV anomaly associated with smoke during the Canadian event was also examined in a recent paper by Allen et al. (2020). While that paper focused primarily on the “Koobor” vortex, they also included a PV anomaly map for 28 August 2017 associated with what this paper calls “Vortex A” (see Figure 16 of the following reference). Allen, D. R., M. D. Fromm, G. P. Kablick III, and G. E. Nedoluha, 2020: Smoke With Induced Rotation and Lofting (SWIRL) in the Stratosphere, *J. Atmos. Sci.*, 77, 4297-4316, <https://doi.org/10.1175/JAS-D-20-0131.1>.*

We missed this paper which was published at a time close to our submission. It is now referred and commented. Indeed the structure shown in the figure 16 of this work displays the vortex A on 28 August but the structure provided by MERRA-2 is apparently much less compact than in the ERA5, perhaps due to a lower spatial resolution in the vertical and horizontal directions. The elongated vortex O is also visible in several panels of this figure.

- *Line 243-245: It is interesting that the 2017 case doesn't show the temperature dipole. Is that simply due to the contours chosen (i.e., the warm anomaly is really there, but it is less than 1.0 K)?*

There is indeed a warm anomaly of weaker amplitude which is actually responsible for the cooling pattern. A new version of the figure which shows the warm anomaly contours is now provided.

- *Line 252: It is unclear exactly how you calculate the horizontal length scale L_h . The text says it is defined as "the diameter of the ring of wind speed maximum". Do you calculate the diameter from the wind speed explicitly for each case? If so, does the wind speed calculation involve removing the background wind in order to focus on the wind associated with the anticyclone?*

We estimate the length explicitly for each case along the East-West direction (i.e. between local extrema of the meridional velocity). The background wind is not removed in the calculation, since by definition its variations at the scale of the vortex are small so that they do not affect local extrema.

- *Line 256: Similarly, L_z is "the vertical extent of the vorticity contour at maximum wind speed". Is this calculated explicitly? Would it be possible to include some more details on this calculation?*

Yes. This is now explained with more details.

- *Line 257: You say "the 2017 vortex A being about 8 times smaller in volume than its gigantic 2020 counterpart". This difference seems too large. If the cloud is considered as a cylinder, then volume is $V = \pi(L_h/2)^2 L_z$. Using numbers from Table 1 we get $V_A = \pi(686/2)^2 3.5 = 1.3 \times 10^6 \text{ km}^3$, and $V_{KooBOR} = \pi(784/2)^2 6.1 = 2.9 \times 10^6 \text{ km}^3$. So KooBOR is only $2.9/1.3=2.2$ times larger in volume than vortex A. Are these estimates correct, or do you use another method to estimate volume?*

Printer-friendly version

Discussion paper



You are correct, thank you for pointing this out. We had previously estimated the volume of the vortices using the same value of vorticity threshold for both cases, but it is more consistent to adapt the value to each case. This has now been corrected.

- *Line 263: You may want to define the condition for inertial instability here.*

The criterion of vanishing PV is actually formally related to symmetric instability (Hoskins, 1974) but, here, we see that the absolute vorticity (not shown) also vanishes, therefore the symmetric instability is an inertial instability.

- *Line 328-330: It looks like the dates used for the Australian vortex are 2-27 February in Figure 8 (mean structure), Figure 9 (heating rate, temperature and vorticity tendencies), and Figure 11 (ozone tendencies), but for Figure 10 (PV increments) the dates are 7-19 January. Is there a particular reason that different dates are chosen for PV? Are the mean composites and increments of PV much different if you calculate them for the different periods? Also, are the green lines on Figure 10 from 7-19 January or 2-27 February?*

There was a mistake here, thank you for spotting it. We have replaced the figure in the main body of the paper by the correct time period, while the 7-19 January period figure has been moved to the appendix. Note that the green contours always correspond to the vorticity in the time period indicated.

The reason for showing the January periods is to emphasize that the LPV increment dipole does not tilt along the direction of the background wind (which corresponds to the drift of the vortex) but rather along the direction of the wind shear, a property which might be found counter intuitive. We follow the reviewers advice and now show this feature in the appendix.

The composite anomaly and increment structures are qualitatively consistent between different periods, but there can be quantitative differences, some of those

[Printer-friendly version](#)[Discussion paper](#)

related to the composite are reported in Tab. 1 of the paper. Regarding the increments, there are differences associated for instance with changes in the ascent rate of the vortex in altitude coordinate which is larger, for Koobor, in January (250 m/day) than in February (150 m/day). This is mentioned in the revised paper together with the new figure in the appendix. The choice of the February time period for the main body of the paper is motivated by the relative steadiness of the structure during that month (see Fig. 7b of Khaykin et al. (2020), which shows a rather constant relative vorticity maximum).

- *Line 335: The northwest-southeast tilt of the PV increments for the Koobor vortex, shown in Figure 10a, is interesting. In the recent analysis by Allen et al. (2020), they examined how Koobor tilts with height and found a NW-SE tilt of the vortex in January. They used a dynamical argument to show how this tilt may develop from internal vortex dynamics in a shear flow. The PV increments shown in this paper appear to support this observed dynamical structure. Also, would the same argument you make in Appendix B apply to the ozone structure seen in Figure 11?*

The argument of Allen et al. (2020) concerns the structure of the vortex and does not apply to the increments, although they are both related to the vertical shear. We now mention their explanation. Vacillations of vortices in a shear flow constitute a fairly standard topic which has been studied a lot, included by the authors in 2D (Legras et al., 2001) and, in 3D, by David Dritschel and collaborators (e.g., Tsang and Dritschel, 2015). There are also a large number of works in the oceanographic literature. Therefore Koobor and its siblings were expected to follow the general laws and to undergo vacillation, erosion and splitting when submitted to a shear. Such vacillations are obvious in Fig. 6b of Khaykin et al. (2020). Koobor went through a first splitting by 25 February 2020, which was actually well predicted by the ECMWF operational model a few days in advance, and a second one by 30 March. For the Canadian case, the deformation of vortex

[Printer-friendly version](#)[Discussion paper](#)

O in the trough led to its splitting in at least three parts that we followed.

Our reasoning in appendix B applies to the ozone increment (or the increment or any quantity organized as a monopole).

- *Line 344: Does “low absolute PV” mean low magnitude (i.e., low absolute value of PV)?*

Yes. The model provides PV and absolute vorticity values close to 0 or even slightly negative. The contrast is such with the ambient PV that the practical situation is that of a ball of zero PV immersed into the stratosphere. Such contrast is made possible by the vertical motion. The familiar formation of cut-off lows or cut-off highs by horizontal isentropic motion never reaches such a situation.

- *Line 408: May want to define terms explicitly in the text here, particularly W and Δ . I assume these are vertical wind and vertical shear of the zonal wind. They are indirectly defined in the Table B1, but not in the text.*

Done.

- *Line 416: How is the wind shear estimate calculated here (i.e., what time range is used)?*

The time range of the composite, this is now specified.

- *Figure 3 caption: Could you include in the caption what time of day was used for the PV analyses?*

Done

- *Figure S4: Is this calculated with normal PV or with the Lait PV? Also, as a reference, it would be useful to include the zonal mean PV on this plot. This should become increasingly negative with time as the parcel ascends. Do the Canadian plumes show a similar behavior?*

Printer-friendly version

Discussion paper



This is the standard PV. The anomaly is defined with respect to the zonal mean PV at same altitude. Therefore this later is the difference between the blue and the orange curves. The initial stage of vortex O exhibits a similar behaviour in 2017. After its split in 3 parts, the offspring exhibit non zero absolute vorticity at the core. As such events usually generate only a small amount of in-mixing, it is unclear whether the effect is real or due to the limitation of assimilation.

- *I enjoyed the animations in the supplementary material. I assume the PV used in the animations is the normal PV, not the Lait PV, right?*

Yes

- *Lastly, there are quite a few different names used in this paper for different aspects of this new phenomenon. For example the terms “smoke-charged vortex”, “smoke charged pancake vortex”, “smoke vortex”, “smoke plume”, “smoke bubble”, and just “bubble” could possibly be condensed into fewer descriptions. The Australian plumes are called “Koobor”, “2nd Vortex”, and “3rd Vortex”, while the Canadian plumes are “Vortex O” (also called “mother vortex”), “Vortex A”, “Vortex B1”, and “Vortex B2”. Different terms are also used for “Koobor”, such as “main vortex”, and “major vortex”. Given this is such a new discovery, to avoid potential confusion, terms could be consolidated and defined (e.g., how does the term “bubble” differ from the terms “plume” and “vortex”). Looking forward, do you have any recommendations for a general scheme as to how these events can be categorized, in order to separate them from stratospheric smoke plumes that do not show a dynamical signature? Allen et al. (2020) coined a new term for this phenomenon, “Smoke With Induced Rotation and Lofting (SWIRL)”, an acronym that accounts for the aerosol source as well as for two of the obvious dynamical aspects of the phenomenon.*

We agree that too much gratuitous variations is undesirable and should be blamed. We have reduced them in the revised version. However, we still find

Printer-friendly version

Discussion paper



useful to have different words to name the smoke bubble or the unshaped smoke plume as seen purely by CALIOP and to name the vortex identified from the reanalysis. We are not very fond of SWIRL as it suggests that heating can generate by itself mean PV or rotation, which is wrong. Swirl is also used already in the fluid dynamics literature, often associated with the word vortex (see, e.g. Mitrofanova et al., 2013), which might be confusing, and has unpleasant meanings in slang. We offer our own acronym, ASTuS (Ascending Smoke Turbo in the Stratosphere) where turbo is a latin word for vortex, which does not have these drawbacks.

Noticeable technical corrections

- *Line 286: What is “beta drift”?*

The beta-drift is a old concept of vortex dynamics on a rotating planet which is standard in the literature on tropical cyclones (see, e.g., Wang and Li, 1992). The beta-drift occurs for a vortex immersed within a planetary PV gradient and results, for a cyclone, into a poleward and westward motion with respect to the mean flow. This is changed to an equatorward motion for an anticyclone. The ingredient of the beta-drift is the wrapping of planetary vorticity around the vortex (Sai-Lap Lam and Dritschel, 2001). This effect is visible in Fig.10 of Allen et al. (2020) but the beta-drift is ignored by these authors who privilege the effect of the vertical tilt.

- *Figure 5 Caption: I think that “orange” and “green” in the figure caption aren’t consistent with the lines on the figure.*

There are actually two curves on each panel of the figure but because the PV tracking and the ozone tracking are very close, they are most often superimposed.

[Printer-friendly version](#)[Discussion paper](#)

- All other corrections have been applied

References

- Allen, D. R., Fromm, M. D., Kablick III, G. P., and Nedoluha, G. E.: Smoke with Induced Rotation and Lofting (SWIRL) in the Stratosphere, *Journal of the Atmospheric Sciences*, 77, 4297–4316, <https://doi.org/10.1175/JAS-D-20-0131.1>, 2020.
- Hoskins, B. J.: The Role of Potential Vorticity in Symmetric Stability and Instability, *Quarterly Journal of the Royal Meteorological Society*, 100, 480–482, <https://doi.org/10.1002/qj.49710042520>, 1974.
- Khaykin, S., Legras, B., Bucci, S., Sellitto, P., Isaksen, L., Tence, F., Bekki, S., Bourassa, A., Rieger, L., Zawada, D., Jumelet, J., and Godin-Beekmann, S.: The 2019-2020 Australian wildfires generated a persistent smoke-charged vortex rising up to 35 km altitude, *Communications Earth and Environment*, 1, 22, <https://doi.org/10.1038/s43247-020-00022-5>, 2020.
- Legras, B., Dritschel, D. G., and Caillol, P.: The Erosion of a Distributed Two-Dimensional Vortex in a Background Straining Flow, *Journal of Fluid Mechanics*, 441, 1–16, <https://doi.org/10.1017/S002211200100502X>, 2001.
- Mitrofanova, O. V., Podzorov, G. D., and Pozdeeva, I. G.: Vortex Structure of Swirl Flows, *International Journal of Heat and Mass Transfer*, 65, 225–234, <https://doi.org/10.1016/j.ijheatmasstransfer.2013.06.007>, 2013.
- Sai-Lap Lam, J. and Dritschel, D. G.: On the Beta-Drift of an Initially Circular Vortex Patch, *Journal of Fluid Mechanics*, 436, 107–129, <https://doi.org/10.1017/S0022112001003974>, 2001.
- Tsang, Y.-K. and Dritschel, D. G.: Ellipsoidal Vortices in Rotating Stratified Fluids: Beyond the Quasi-Geostrophic Approximation, *Journal of Fluid Mechanics*, 762, 196–231, <https://doi.org/10.1017/jfm.2014.630>, 2015.
- Wang, B. and Li, X.: The Beta Drift of Three-Dimensional Vortices: A Numerical Study, *Monthly Weather Review*, 120, 579–593, [https://doi.org/10.1175/1520-0493\(1992\)120<0579:TBDOTD>2.0.CO;2](https://doi.org/10.1175/1520-0493(1992)120<0579:TBDOTD>2.0.CO;2), 1992.

[Printer-friendly version](#)[Discussion paper](#)