

Author's response file

In the following, we include the answers to the referees together with the complete revised manuscript. Changes in the text of the manuscript are marked with red colour.

Response to Referee #1:

First of all we thank the referee for the effort to carefully reading the manuscript and for all comments.

General comments:

This manuscript presents observations of multiple trace gases from the GLORIA instrument over the North Atlantic. It describes the GLORIA instrument and data analysis method, and then uses model simulations to interpret the observations and relate them to the Asian Monsoon and to identify underestimates in the emission inventory. This is an interesting and useful multi-species dataset and the modeling provides a robust tool for analysis. However, the combination of retrieval description and model analysis left the main focus of the manuscript a bit unclear to me. If the primary objective is to present the GLORIA retrieval and data analysis, I would like to see this section expanded and more validation included. If the goal is instead to interpret the observations, the discussion of the driving scientific questions and what new insights are found should be clarified/strengthened.

The objective is both, to present the GLORIA data retrieval of pollutant species together with an interpretation of the data with the help of model simulations. A detailed description of the general retrieval process is given in a previous paper by Johansson et al. (2018) which is cited in the manuscript. We now mention this more clearly in Section 2.2. We also included some more sentences in the conclusion part to better strengthen the messages of this study.

Specific comments:

Line 60: This lifetime is still long enough to allow long-range transport.

We included a corresponding clause to make this clear.

Line 132: Is radiative transport the same as radiative transfer?

Yes, we replaced “transport” by “transfer”.

Section 2.2: Is this the first description of this method, or are there other papers that could be referenced for the GLORIA retrievals and validation?

A detailed description of the general GLORIA retrieval process is given in the paper by Johansson et al. (2018) that is already cited later in the text. We included a clarifying sentence on this issue at the end of the first paragraph in Section 2.2. This paper also contains the

validation of major species observed by GLORIA (e.g. O₃, HNO₃, and ClONO₂). For species discussed by this paper, we lack of in-situ data to compare our GLORIA measurements with.

Line 216: All biomass burning or just “anthropogenic” biomass burning?

We deleted “anthropogenic” in the corresponding sentence.

Line 275: Specify the instrument/satellite.

The instruments are the Atmospheric Chemistry Experiment (ACE) Fourier transform spectrometer on SCISAT-1 and the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) aboard the Envisat satellite. We added a corresponding clause in the text.

Conclusions: Last paragraph: The importance of emission inventories is well-known. Please make the conclusions more specific and emphasize what is new from this study.

We included some more information in the conclusions to better emphasize what is new and important in our study. For instance, we emphasized that enhanced amounts of pollutant species were measured in the upper troposphere with high temporal and spatial resolution. Furthermore, it is important to state that these enhancements were detected far away from the emission sources of these species.

Figure 6: Is this figure the same as the first column of Figure 7?

It is not exactly the same. Figure 6 contains the unsmoothed GLORIA data while the first column of Figure 7 shows the temporally smoothed GLORIA data (as noted now more clearly in the figure caption).

Fig. 9 caption: Does “Coloured encircled areas” mean the cyan dots/circles?

Not only cyan, but also black, blue and green. We added this in the figure caption text for better clarity.

Editorial comments:

Lines 46-50: please reword this sentence for better clarity.

We split the sentence into two parts for better clarity.

References

Johansson, S., Woiwode, W., Höpfner, M., Friedl-Vallon, F., Kleinert, A., Kretschmer, E., Latzko, T., Orphal, J., Preusse, P., Ungermann, J., Santee, M. L., Jurkat-Witschas, T., Marsing, A., Voigt, C., Giez, A., Krämer, M., Rolf, C., Zahn, A., Engel, A., Sinnhuber, B.-M., and Oelhaf, H.: Airborne limb-imaging measurements of temperature, HNO₃, O₃, ClONO₂, H₂O and CFC-12 during the Arctic winter 2015/2016: characterization, in situ validation and comparison to Aura/MLS, Atmos. Meas. Tech., 11, 4737–4756, <https://doi.org/10.5194/amt-11-4737-2018>, 2018.

Response to Referee #2:

First of all we thank the referee for the effort to carefully reading the manuscript and for all comments.

General comments:

My only general comment is that it would have been good to include some discussion of how these observations compare to past observations of these species. Currently this is limited to comparisons to limited-resolution spaceborne remote sounding observations (e.g., the many citations to the Rinsland, Glatthor and Wiegele papers in the manuscript). However, there are a wealth of airborne in-situ observations of many these species in past campaigns (for example the NASA ATOM campaign, among many others from the US, Europe and Asia). Given that GLORIA is a relatively new (but very welcome) addition to the worldwide portfolio of airborne instruments measuring atmospheric composition, and that it is one of the few employing remote-sounding (particularly for such a wide range of species), and further, given the general skepticism some in the community have toward remote sounding observations, some additional statements as to how the GLORIA findings compare to available in situ observations of the same species at similar altitudes/latitudes/seasons, etc. would help cement the value of the GLORIA dataset in the community mind set.

We included some discussion on the comparison to in-situ aircraft measurements in Section 3.1. and added corresponding references. We do not have a direct comparison available (in the sense of a validation) of the pollutant species retrieved by GORIA with in-situ observations. However, retrieved GLORIA amounts of these trace gases are within the spread of values measured by in-situ instruments.

Specific comments:

Sentence spanning lines 24/25: Reword to "Elevated quantities of PAN were measured even in the lowermost stratosphere (locally up to 14 km), likely reflecting the fact that this molecule has the longest lifetime of the four species discussed herein."

We changed the text according to the reviewer's suggestion.

Sentence spanning lines 43-46: Better to split into two sentences along the lines of "... conditions. In particular, rapid vertical transport by deep convection followed by strong horizontal transport associated with the upper troposphere subtropical jet stream ([citations]) is a particularly efficient means by which surface pollutants can be transported long distances.

We changed the text accordingly.

Line 47: Think that the "that" would be better as ", which" in this case.

We split the sentence into two parts for better clarity.

Line 56: "such that C₂H₆ may be" -> "enabling it to be"

We changed this part accordingly.

Line 64: "are important contributors to the tropospheric abundances of this molecule" or something similar.

We changed the sentence accordingly.

Line 66: "like" -> "such as the"

Changed.

Line 105: "using" -> "observing" (to avoid having "using" twice in quick succession)

Changed.

Line 154: I think something like "Test retrievals were used to identify microwindows that combine limited overlap of spectral signatures of disturbing gases with a high sensitivity to changes in the abundance of target gases." would be better wording.

We modified the text accordingly.

Line 252: comma needed after "that"

Changed.

Lines 257-287: As discussed above, it would be good to compare a small number of the wealth finding from airborne in-situ observations of these species.

We included text and citations of airborne in-situ observations for all pollutant species.

Line 262: "stronger enhanced" -> "strong enhancements of"

Changed.

Line 263: "what" -> ", which"

Changed.

Line 265: "picture" -> "behavior"

Changed.

Line 291: "with respect to" -> "given the", also "in" -> "of"

Changed.

Line 292: "Concerning" -> "For"

Changed.

Line 293: "principally" -> "generally"

Changed.

Line 302: "The comparison of" -> "Comparisons for"

Changed.

Line 311: "For C2H2 we note that EMAC predicts elevated concentrations in much the same region where GLORIA reports enhancements (see ...)"

Changed.

Line 311-317: Again, this would be a good place to mention in situ comparisons.

We now mention the in-situ airborne measurements in Section 3.1.

Line 430-434: The sentence "However, the real...". I'm afraid I don't understand what this sentence is trying to say. Does CLAMS not have emissions for these specific species on some kind of fine spatial resolution (EDGAR, MEGAN, etc.?) Please clarify what is meant by "the real regions".

"Real" means where emissions really occur. The original text was misleading, so we rewrote this clause for better understanding.

Line 461: Remove "(primarily CAMS)" and add ", particularly for CAMS" at the end of the sentence.

Changed.

Line 525: Some weird cut and paste typo in citation.

Okay, corrected in bibliography.

Line 679: Extra space between "O" and "3" in citation.

Okay, corrected in bibliography.

Line 746: "n/a-n/a" in citation.

Okay, corrected in bibliography.

Figures 2-5 are nicely put together.

Thanks.

Figure 6 (and 7): The grey line is hard to see, make it thicker. The dashed magenta line is very hard to see. I suggest you make it white and thicker (and possibly not dashed?). "...mark regions

with enhanced VMR levels" - not for O3, perhaps clarify "primary pollutant VMRs" or something like that?

We made the grey lines and the magenta lines thicker so they are clearly visible now (white instead of magenta is not a good option for these lines because they are also included in the following Figure 8 with difference plots containing large white areas). We changed Figures 6 to 8 and the Figure caption 6 according to the reviewer's suggestion.

Figure 9: This is very hard to see given the colored continents/oceans. As pretty as they are I'd suggest a grey-scale version of the background image, or ideally just white oceans and very pale grey continents (single color, no mountains or things like that).

It is a difficult task to find a common optimal background for all individual pictures displayed in Figure 9. We already tested a lot of background and color scale combinations before submitting this manuscript. We again tested many combinations and found that white oceans is not the best solution because yellowish colors are hardly visible. In the end we omitted the surface relief and took for continents a light grey and for oceans a light blue color and changed Figure 9 accordingly.

Response to Referee #3:

First of all we thank the referee for the effort to carefully reading the manuscript and for all comments. Citations mentioned below are included in the manuscript.

General comments:

The presentation of these new observations, along with the description of the measurement technique, is worthwhile. However, the goals of the paper should be made clearer. The uniqueness of the observations could be more strongly emphasized. It would be helpful to have some sort of validation of the observations through comparison to aircraft data from established measurement techniques.

We included additional motivation in the introduction and in the conclusions to better emphasize the goals of this paper. Since no co-incident in-situ observations enabling a direct comparison (in the sense of a validation) are available, we have checked in detail published airborne datasets concerning the pollution trace gases derived from GLORIA. We can conclude that retrieved GLORIA amounts of these trace gases are within the spread of values measured by in-situ instruments. Related citations are now provided in Section 3.1 individually for each species.

The introduction seems rather awkward, with the discussion of the measured compounds seeming rather disjointed. Perhaps more discussion of the measurement technique and its uniqueness would be more appealing to readers, and then an explanation of why these species are discussed - driven by the capability to measure them. The explanation of their role in atmospheric chemistry could be saved for the analysis discussion. At l. 41, PAN is a 'secondary pollutant', not 'secondary order'. At l. 51, in what sense is ethane 'most important'?

We include and modified sentences in the introduction to make the text and the goal of these measurements clearer. A comprehensive description of the technical issues of GLORIA is given in the papers by Friedl-Vallon et al. (2014) and Riese et al. (2014) and references therein. This is written now more clearly in the text. We find that the description of the pollutant species is better suited for the introduction than for the data analysis section (of course, this approach may be a matter of taste). In line 41 we omitted "order". Line 51: It is the most important non-methane hydrocarbon constituent of natural gas. We modified this clause accordingly.

The purpose of the model results in the paper should be made clearer. Are they being used to provide validation of the observations? It would be more appropriate to just use the model to explain the distributions and identify the sources of high mixing ratios.

Of course, the use of the models is not intended to validate the measurements. It should be understood as a kind of intercomparison. One goal was to see if the models EMAC and CAMS are capable to reproduce the locations of the enhanced amounts of pollutants. Another goal was to quantify the differences between measured and simulated data, especially in the case of EMAC where we performed different emission scenarios. Concerning the origin of the detected enhancements, we used backward trajectories and artificial tracers of air mass origin calculations. We tried to make this issue clearer in the revised text.

Using 60-day back trajectories seems rather a stretch. I would not think they are reliable that far back. The forward CLAMS simulations of various regional tracers seem more reliable, so the back trajectories seem unnecessary.

We agree with the referee, that in general, trajectory calculations have limitations due to trajectory dispersion depending on the trajectory length. Therefore the uncertainty of a single trajectory is increasing with the trajectory length, however the variability of a cluster of trajectories starting in the same region, reflects the impact of mixing processes. Therefore, in this study a plenty of back-trajectories are started in the marked regions with enhanced VMR levels.

Frequently employed trajectory lengths to study transport processes in particular in the Asian monsoon region range from a couple of weeks to a few months depending on the transport times from Earth's surface to atmospheric altitudes (Chen et al., 2012; Bergman et al., 2013; Garny and Randel, 2016; Müller et al., 2016; Li et al., 2018; Vogel et al., 2019; Legras and Bucci, 2020; Hanumanthu et al., 2020). In particular the CLaMS backward calculations to analyse the regions with enhanced PAN between 13 and 14.5 km (about 400 K) demonstrate that the transport times from the planetary boundary layer in Asia to the extratropical UTLS over the Atlantic are between 40 and 60 days. Therefore, trajectories up to a lengths of 60 days are necessary in our study to infer the possible source regions of PAN. Further, the endpoints of the trajectories in the planetary boundary layer show a good overall agreement to the results of forward CLaMS simulations for various regional tracers demonstrating that trajectories of a length up to 60 days are suitable for our study.

Moreover, back trajectory calculations have an added valued to the 3-dimensional forward calculations with CLaMS because they demonstrate the detailed transport pathways from the boundary source to the locations of the measurement and its transport times. The trajectory calculations show that the air parcels with enhanced PAN are uplifted by diabatic heating in the upward spiralling range (Vogel et al., 2019) of the Asian monsoon anticyclone up to about 400 K within about 40 days and subsequent transport (within about 20 days) occurred along the subtropical jet to the extratropical UTLS over the Atlantic. This detailed transport pathway and its transport time cannot be inferred from CLaMS 3-dimensional forward calculations, therefore the back-trajectory analysis is an added value to our study.

The conclusions seem to discuss more the model evaluation aspects of the observation-model comparisons, which I do not find fully justified by the presentation of the results.

The conclusion consists of two main parts. First the findings of the measurements and second, the findings connected with the comparison to the models EMAC and CAMS. We included some text to make this issue more clearly.

References

- Bergman, J. W., Fierli, F., Jensen, E. J., Honomichl, S., and Pan, L. L.: Boundary layer sources for the Asian anticyclone: Regional contributions to a vertical conduit, J. Geophys. Res. Atmos., 118, 2560–2575, <https://doi.org/10.1002/jgrd.50142>, 2013.
- Chen, B., Xu, X. D., Yang, S., and Zhao, T. L.: Climatological perspectives of air transport from atmospheric boundary layer to tropopause layer over Asian monsoon regions during boreal summer inferred from Lagrangian approach, Atmos. Chem. Phys., 12, 5827–5839, <https://doi.org/10.5194/acp-12-5827-2012>, 2012.

- Friedl-Vallon, F., Gulde, T., Hase, F., Kleinert, A., Kulesa, T., Maucher, G., Neubert, T., Olschewski, F., Piesch, C., Preusse, P., Rongen, H., Sartorius, C., Schneider, H., Schönfeld, A., Tan, V., Bayer, N., Blank, J., Dapp, R., Ebersoldt, A., Fischer, H., Graf, F., Guggenmoser, T., Höpfner, M., Kaufmann, M., Kretschmer, E., Latzko, T., Nordmeyer, H., Oelhaf, H., Orphal, J., Riese, M., Schardt, G., Schillings, J., Sha, M. K., Suminska-Ebersoldt, O., and Ungermann, J.: Instrument concept of the imaging Fourier transform spectrometer GLORIA, *Atmos. Meas. Tech.*, 7, 3565–3577, <https://doi.org/10.5194/amt-7-3565-2014>, 2014.
- Garny, H. and Randel, W. J.: Transport pathways from the Asian monsoon anticyclone to the stratosphere, *Atmos. Chem. Phys.*, 16, 2703–2718, <https://doi.org/10.5194/acp-16-2703-2016>, 2016.
- Hanumanthu, S., Vogel, B., Müller, R., Brunamonti, S., Fadnavis, S., Li, D., Ölsner, P., Naja, M., Singh, B. B., Kumar, K. R., Sonbawne, S., Jauhiainen, H., Vömel, H., Luo, B., Jorge, T., Wienhold, F. G., Dirksen, R., and Peter, T.: Strong day-to-day variability of the Asian Tropopause Aerosol Layer (ATAL) in August 2016 at the Himalayan foothills, *Atmos. Chem. Phys.*, 20, 14273–14302, <https://doi.org/10.5194/acp-20-14273-2020>, 2020.
- Johansson, S., Woiwode, W., Höpfner, M., Friedl-Vallon, F., Kleinert, A., Kretschmer, E., Latzko, T., Orphal, J., Preusse, P., Ungermann, J., Santee, M. L., Jurkat-Witschas, T., Marsing, A., Voigt, C., Giez, A., Krämer, M., Rolf, C., Zahn, A., Engel, A., Sinnhuber, B.-M., and Oelhaf, H.: Airborne limb-imaging measurements of temperature, HNO₃, O₃, ClONO₂, H₂O and CFC-12 during the Arctic winter 2015/2016: characterization, in situ validation and comparison to Aura/MLS, *Atmos. Meas. Tech.*, 11, 4737–4756, <https://doi.org/10.5194/amt-11-4737-2018>, 2018.
- Legras, B. and Bucci, S.: Confinement of air in the Asian monsoon anticyclone and pathways of convective air to the stratosphere during the summer season, *Atmos. Chem. Phys.*, 20, 11045–11064, <https://doi.org/10.5194/acp-20-11045-2020>, 2020.
- Li, D., Vogel, B., Müller, R., Bian, J., Günther, G., Li, Q., Zhang, J., Bai, Z., Vömel, H., and Riese, M.: High tropospheric ozone in Lhasa within the Asian summer monsoon anticyclone in 2013: influence of convective transport and stratospheric intrusions, *Atmos. Chem. Phys.*, 18, 17979–17994, <https://doi.org/10.5194/acp-18-17979-2018>, 2018.
- Müller, S., Hoor, P., Bozem, H., Gute, E., Vogel, B., Zahn, A., Bönisch, H., Keber, T., Krämer, M., Rolf, C., Riese, M., Schlager, H., and Engel, A.: Impact of the Asian monsoon on the extratropical lower stratosphere: trace gas observations during TACTS over Europe 2012, *Atmos. Chem. Phys.*, 16, 10573–10589, <https://doi.org/10.5194/acp-16-10573-2016>, 2016.
- Riese, M., Oelhaf, H., Preusse, P., Blank, J., Ern, M., Friedl-Vallon, F., Fischer, H., Guggenmoser, T., Höpfner, M., Hoor, P., Kaufmann, M., Orphal, J., Plöger, F., Spang, R., Suminska-Ebersoldt, O., Ungermann, J., Vogel, B., and Woiwode, W.: Gimbalbed Limb Observer for Radiance Imaging of the Atmosphere (GLORIA) scientific objectives, *Atmos. Meas. Tech.*, 7, 1915–1928, <https://doi.org/10.5194/amt-7-1915-2014>, 2014.
- Vogel, B., Müller, R., Günther, G., Spang, R., Hanumanthu, S., Li, D., Riese, M., and Stiller, G. P.: Lagrangian simulations of the transport of young air masses to the top of the Asian monsoon anticyclone and into the tropical pipe, *Atmos. Chem. Phys.*, 19, 6007–6034, <https://doi.org/10.5194/acp-19-6007-2019>, 2019.

Pollution trace gases C₂H₆, C₂H₂, HCOOH, and PAN in the North Atlantic UTLS: observations and simulations

Gerald Wetzel¹, Felix Friedl-Vallon¹, Norbert Glatthor¹, Jens-Uwe Grooß², Thomas Gulde¹, Michael Höpfner¹, Sören Johansson¹, Farahnaz Khosrawi¹,
5 Oliver Kirner³, Anne Kleinert¹, Erik Kretschmer¹, Guido Maucher¹, Hans Nordmeyer¹, Hermann Oelhaf¹, Johannes Orphal¹, Christof Piesch¹, Björn-Martin Sinnhuber¹, Jörn Ungermann², and Bärbel Vogel²

¹Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research, Karlsruhe, Germany

²Research centre Jülich, Institute of Energy and Climate Research – Stratosphere (IEK-7), Jülich, Germany

10 ³Karlsruhe Institute of Technology, Steinbuch Centre for Computing, Karlsruhe, Germany

Correspondence to: Gerald Wetzel (gerald.wetzel@kit.edu)

Abstract

Measurements of the pollution trace gases ethane (C₂H₆), ethyne (C₂H₂), formic acid
15 (HCOOH), and peroxyacetyl nitrate (PAN) were performed in the North Atlantic upper troposphere and lowermost stratosphere (UTLS) region with the airborne limb imager GLORIA (Gimballed Limb Observer for Radiance Imaging of the Atmosphere) with high spatial resolution down to cloud top. Observations were made during flights with the German research aircraft HALO (High Altitude and Long Range Research Aircraft) in the frame of the WISE
20 (Wave-driven ISentropic Exchange) campaign, which was carried out in autumn 2017 from Shannon (Ireland) and Oberpfaffenhofen (Germany). Enhanced volume mixing ratios (VMR) of up to 2.2 ppbv C₂H₆, 0.2 ppbv C₂H₂, 0.9 ppbv HCOOH, and 0.4 ppbv PAN were detected during the flight on 13 September 2017 in the upper troposphere and around the tropopause above the British Isles. **Elevated quantities of PAN were measured** even in the lowermost
25 stratosphere (locally up to 14 km), **likely reflecting the fact that this molecule has the longest lifetime of the four species discussed herein**. Backward trajectory calculations as well as global three-dimensional CLaMS simulations with artificial tracers of air mass origin have shown that the main sources of the observed pollutant species are forest fires in North America and anthropogenic pollution in South and Southeast Asia uplifted and moved within the Asian
30 monsoon anticyclone (AMA) circulation system. After release from the AMA, these species or their precursor substances are transported by strong tropospheric winds over large distances, depending on their particular atmospheric lifetime of up to months. Observations are compared

to simulations with the atmospheric models EMAC (ECHAM5/MESSy Atmospheric Chemistry) and CAMS (Copernicus Atmosphere Monitoring Service). These models are qualitatively able to reproduce the measured VMR enhancements but underestimate the absolute amount of the increase. Increasing the emissions in EMAC by a factor of 2 reduces the disagreement between simulated and measured results and illustrates the importance of the quality of emission databases used in chemical models.

1 Introduction

Organic compounds in the troposphere like ethane, ethyne, formic acid, and secondary (not directly emitted) pollutants like peroxyacetyl nitrate are involved in many atmospheric processes. These pollutants can be transported into remote regions due to their long lifetime under appropriate atmospheric conditions. In particular, rapid vertical transport by deep convection followed by strong horizontal transport associated with the upper troposphere subtropical jet stream is a particularly efficient means by which surface pollutants can be transported long distances (e.g., Lu et al., 2019; Alvarado et al., 2020). At these altitudes, such hydrocarbons and nitrogen-containing substances (like peroxyacetyl nitrate) may also influence the amount of ozone. This is important because ozone is known to be an effective greenhouse gas in the upper troposphere and lowermost stratosphere (UTLS) region since it largely influences the radiation budget around the tropopause (Forster and Shine, 1997; Hansen et al., 1997; Xie et al., 2008; Riese et al., 2012).

Tropospheric emissions of ethane (C_2H_6), the most important non-methane hydrocarbon (NMHC) constituent of natural gas, are connected with biomass burning and natural gas losses (Rudolph, 1995; Singh et al., 2001). The production of fossil fuels together with biofuel use also are important sources of this molecule (Xiao et al., 2008). It is removed from the atmosphere by reaction with the hydroxyl (OH) radical (Xiao et al., 2008). The mean lifetime of C_2H_6 is about two months (Rudolph, 1995) enabling it to be transported far away from its source regions.

The trace gas ethyne (C_2H_2) is emitted into the troposphere by combustion of biofuels and fossil fuels, as well as biomass burning (Xiao et al., 2007). As in the case of C_2H_6 , the reaction with the OH radical is also responsible for the loss of C_2H_2 in the atmosphere. The mean lifetime of C_2H_2 is shorter compared to the one of C_2H_6 and amounts to about two weeks (Xiao et al., 2007) but still allows long-range transport of this species.

65 Formic acid (HCOOH) has many different sources. Direct emissions from plants, biomass
burning, and fossil fuel combustion are important contributors to the tropospheric abundances
of this molecule (Mungall et al., 2018). A secondary photochemical formation takes place from
anthropogenic and biogenic precursors (Yuan et al., 2015) such as the oxidation of volatile
organic compounds (Khare et al., 1999). The loss of HCOOH is possible due to wet and dry
70 deposition as well as oxidation with the OH radical (Paulot et al., 2011). The atmospheric mean
lifetime of HCOOH is very variable and ranges from one or two days in the boundary layer up
to a few weeks in the free troposphere (Millet et al., 2015).

The molecule peroxyacetyl nitrate (CH₃COO₂NO₂), commonly named as PAN, is formed via
a three-body-reaction of peroxyacetyl (CH₃COO₂) with nitrogen dioxide (NO₂) and a third
75 partner M (mainly N₂ or O₂):



The reverse reaction of (R1) is the thermal decomposition of PAN and defines the main loss of
this molecule in the atmosphere (Fischer et al., 2014) while photolysis becomes dominant in
the UTLS region (Fadnavis et al., 2015). Two minor loss processes of PAN are reactions with
80 OH and dry deposition (Fischer et al., 2014). The mean atmospheric lifetime of PAN is very
variable since it is strongly dependent on the ambient temperature. While the mean lifetime
amounts to only one hour at temperatures of 298 K, it rises up to a few months under cold upper
tropospheric conditions (Singh, 1987; Glatthor et al., 2007a; Fischer et al., 2014; Glatthor et al.,
2007b)). Hence, enhanced PAN amounts (which serve as an NO₂ reservoir) can be transported
85 over wide geographical regions in the middle and upper troposphere. Thus, NO₂ released by
the backward reaction of (R1) may contribute to an increase of tropospheric ozone far away
from the PAN sources (Glatthor et al., 2007b; Singh, 1987; Glatthor et al., 2007b; Fadnavis et
al., 2014; Ungermann et al., 2016).

Scientific flights of the airborne limb imager GLORIA (Gimballed Limb Observer for Radiance
90 Imaging of the Atmosphere; Friedl-Vallon et al., 2014; Riese et al., 2014) were carried out
aboard HALO (High Altitude and Long Range Research Aircraft) during the WISE (Wave-
driven Isentropic Exchange) campaign in autumn 2017 above the North Atlantic. This
atmospheric region is characterized by intense dynamical activity reflected in complex
structures like tropopause folds and air masses of different origin. Tropospheric pollutants like
95 C₂H₆, C₂H₂, HCOOH, and PAN were observed together with stratospheric trace gases like O₃
in the UTLS region with high temporal and spatial resolution. An important goal of these

measurements was to identify possible atmospheric regions with enhanced amounts of these pollutant trace gases far away from the emission locations. A description of the GLORIA instrument, data analysis and chemical modelling is given in Sect. 2. A discussion of observed vertical volume mixing ratio (VMR) profiles of trace species follows in Sect. 3 together with a comparison of the measured data to simulations of the chemistry climate model EMAC (ECHAM5/MESSy Atmospheric Chemistry) and to assimilated data of CAMS (Copernicus Atmosphere Monitoring Service). A discussion on the possible origin of air masses at the Earth's boundary layer detected by GLORIA is also included in this section.

2 GLORIA instrument, data analysis and modelling

In the following subsections, we give an overview of the GLORIA instrument and the flights with the HALO aircraft, together with the corresponding data analysis and a description of atmospheric modelling performed for this study.

2.1 GLORIA instrument and HALO flights

The cryogenic Fourier Transform limb emission spectrometer GLORIA operates in the thermal infrared spectral region between about 7 and 13 μm using a two-dimensional detector array observing 128 vertical and 48 horizontal interferograms per measurement (Friedl-Vallon et al., 2014; Riese et al., 2014). The interferograms are Fourier transformed into the spectral domain and radiometrically calibrated using in-flight measurements of two blackbodies (Kleinert et al., 2014). Furthermore, spectra in the horizontal direction were averaged to improve the signal to noise ratio such that the noise equivalent spectral radiance (NESR) finally amounts to about $1.5 \times 10^{-8} \text{ W}(\text{cm}^2 \text{ sr cm}^{-1})^{-1}$. Spectra recorded with maximum optical path difference of 8.0 cm, which corresponds to an unapodized spectral resolution of 0.0625 cm^{-1} , were used for this study. These so-called chemistry mode spectra, are apodized with the Norton and Beer (1976) “strong” function. Due to the high spectral resolution, these measurements allow the retrieval of many species with minor contribution to the spectra by the separation of individual spectral lines from continuum-like emissions. A comprehensive description of technical details of the GLORIA instrument is given by Friedl-Vallon et al. (2014) and Riese et al. (2014).

In this study, we report results from the WISE aircraft campaign, which was dedicated mainly to the investigation of dynamical processes and the evolution of air masses within the context of stratosphere-troposphere exchange. Sixteen flights with HALO were performed from

Shannon (Ireland) and Oberpfaffenhofen (Germany) between 31 August 2017 and 21 October 2017, in a region within about 40° W and 15° E longitude and 40° N and 75° N latitude. In the following sections, we focus on results of the flight on 13 September 2017 where strong enhancements in the VMR of the pollutant species C₂H₆, C₂H₂, HCOOH, and PAN were observed by GLORIA. Figure 1 shows the flight path consisting of two main legs together with tangent points of GLORIA where the instrument was operated in the chemistry mode.

2.2 Data analysis of measured spectra

Radiances are calculated with the Karlsruhe Optimized and Precise Radiative transfer Algorithm (KOPRA; Stiller et al., 2002) based on spectroscopic parameters from the high-resolution transmission molecular absorption database (HITRAN; Gordon et al., 2017). The retrieval itself is performed with the procedure KOPRAFIT (Höpfner et al., 2002) using derivatives of the radiance spectrum with respect to atmospheric state and instrument parameters (Jacobians) calculated by KOPRA. The inverse problem of radiative transfer is solved by KOPRAFIT with a Gauss-Newton iterative method (Rodgers, 2000) in combination with a Tikhonov-Phillips regularization approach (Phillips, 1962; Tikhonov, 1963) using a constraint with respect to a first derivative of the a priori profile of the target species. A detailed description of the general retrieval process together with the validation of major species (e.g. O₃, HNO₃, and ClONO₂) observed by GLORIA is given by Johansson et al. (2018).

In a first step cloud affected spectra are filtered out using a cloud index as described in Spang et al. (2004). Spectra with a colour ratio of the mean radiance in two spectral windows (788.20-796.25 cm⁻¹ and 832.30-834.40 cm⁻¹) larger than 2 were chosen for the retrieval process. Prior to the trace gas retrievals, the pointing elevation angle was retrieved to compensate for systematic misalignment of the line of sight (LOS) of the instrument according to the method described in Johansson et al. (2018). The final pointing (LOS) error was estimated to 0.01°. Vertical profiles of atmospheric species were taken from a climatological atmosphere (Remedios et al., 2007), updated with surface concentration data from NOAA ESRL GMD (National Oceanic and Atmospheric Administration, Earth System Research Laboratory, Global Monitoring Division; Montzka et al., 1999). Subsequently, a temperature retrieval was carried out using pressure-temperature a priori data from the European Centre for Medium-Range Weather Forecasts (ECMWF) which was interpolated to the GLORIA vertical retrieval grid. Spectral windows around 811 and 957 cm⁻¹ containing appropriate CO₂ transitions were

used for this analysis. The total error of the temperature retrieval calculations is estimated to about 1.5 K (Johansson et al., 2018).

The spectral analysis of the target species C_2H_6 , C_2H_2 , $HCOOH$, and PAN is impeded by overlapping features of so-called disturbing gases in the corresponding spectral region. Hence, a careful selection of appropriate microwindows is essential to perform retrievals with good accuracy. Test retrievals were used to identify microwindows that combine limited overlap of spectral signatures of disturbing gases with a high sensitivity to changes in the abundance of target gases. Main interfering species were either adjusted simultaneously together with the target molecule or pre-fitted in a different spectral interval.

The analysis of C_2H_6 was performed in four microwindows within the ν_9 band between 819 and 833 cm^{-1} (see Fig. 2a). Many overlapping features of so-called disturbing gases are visible in this spectral region, first of all H_2O and CO_2 . The error budget of C_2H_6 is shown in Fig. 2b. At higher altitudes with low C_2H_6 values, the total error of this molecule is dominated by the temperature error. Lower down in the VMR maximum region, the systematic spectroscopic error of C_2H_6 (as given in HITRAN) governs the error budget. The total error of C_2H_6 remains within about 15 % in the altitude region of the upper troposphere.

Retrieval calculations of C_2H_2 were carried out in the R-branch region of the ν_5 band. Figure 3a shows spectral contributions of relevant species in four microwindows between 759 and 781 cm^{-1} that have been found best appropriate to derive C_2H_2 from GLORIA spectra. The corresponding retrieval error budget of C_2H_2 is given in Fig. 3b. The random noise error is dominating the budget over nearly the complete altitude range. The total error of C_2H_2 stays within 10-15 % in the region of the VMR maximum in the upper troposphere.

The analysis of the molecule $HCOOH$ was performed in the spectral range between 1086 and 1117 cm^{-1} (see Fig. 4a). Three microwindows were chosen including the strong Q-branch of the $HCOOH$ ν_6 vibrational band. The spectral region is dominated by spectral features due to O_3 , CO_2 , CFC-12, and HCFC-22 transitions. The spectroscopic part of the total error is dominant in the altitude region of the $HCOOH$ VMR maximum in the upper troposphere (see Fig. 4b). Here, the total $HCOOH$ error stays within 10 %. Apart from this altitude region, the error budget is governed by the random noise part and the total error increases significantly (mainly in the upper part of the profile with low $HCOOH$ values).

The retrieval of PAN was conducted in two broad microwindows of the ν_{16} band between 780 and 805 cm^{-1} (see Fig. 5a). The spectral gap between these microwindows was chosen such that

the strong CO₂ Q-branch at 792 cm⁻¹ was excluded from the data analysis. Beside CO₂, the molecules H₂O and CCl₄ are the main interfering species in the spectral region of PAN. The retrieval error budget is depicted in Fig. 5b. Beside random noise, further error sources like inaccuracies in the FOV and the retrieved temperature profile contribute to the total PAN error that remains within 10 % in altitude regions with enhanced PAN amounts.

Besides the retrieval of the above-mentioned pollutant gases, the tracer species ozone was also inferred from the recorded spectra. Many spectral ozone lines are available in the mid-infrared spectral region. Transitions between 780 and 788 cm⁻¹ within the ν_2 band were chosen for the retrieval process similar to the method described in Johansson et al. (2018). The total ozone error is within 10 % with a vertical resolution of 0.3 to 1.5 km.

The altitude resolution of all retrievals, calculated from the full width at half maximum of the columns of the averaging kernel matrix, was used as an a posteriori quality filtering of the retrieved data. Only vertical profile parts with an altitude resolution of better than 2 km were finally used for the data interpretation.

2.3 Model simulations

2.3.1 EMAC

Retrieved vertical profiles of trace species are compared to a multi-year simulation of the chemistry climate model ECHAM5/MESSy Atmospheric Chemistry (EMAC). This Eulerian model includes submodels describing tropospheric and middle atmosphere processes (Jöckel et al., 2010). The core model is the 5th generation European Centre Hamburg general circulation model (ECHAM5; Roeckner et al., 2006) that is connected to the submodels using the interface Modular Earth Submodel System (MESSy). For the present study we applied EMAC (ECHAM5 version 5.3.02, MESSy version 2.53) with a spherical truncation of T106 (corresponding to a resolution of approximately 1.125 by 1.125 degrees in latitude and longitude) with 90 hybrid pressure levels from the ground up to 0.01 hPa. Meteorological data fields are specified using a Newtonian relaxation technique of the surface pressure and prognostic variables below 1 hPa with the ECMWF reanalysis ERA-Interim (Dee et al., 2011). The simulation was initialized on 1 May 2017 and includes a comprehensive chemistry setup from the troposphere to the lower mesosphere. Rate constants of gas-phase reactions originate from Atkinson et al. (2007) and Sander et al. (2011). Photochemical reactions of precursor substances important for the build-up of the species PAN (Fischer et al., 2014) were integrated

into the model setup. For surface emissions of non-methane volatile organic compounds (NMVOCs), a data set of the MACCity emission inventory (MACC/CityZEN; Granier et al., 2011) and ACCMIP (Atmospheric Chemistry and Climate Model Intercomparison Project; Lamarque et al., 2013) was used. **Emission** sources from biomass burning, agricultural waste burning, fossil fuels, ship, road and aircraft, as well as biogenic emissions are considered. For the simulated year 2017, most recent available emissions of 2010 are repeated. In addition to this EMAC standard run, a second model simulation (called EMAC_2) was performed using NMVOC emissions enhanced by a factor of 2 as recommended by Monks et al. (2018). The model output data were saved every 5 hours during the time period of the GLORIA observations. The model output to the GLORIA measurements was interpolated in time and space to the observation geolocations.

2.3.2 CAMS

The Copernicus Atmosphere Monitoring Service (CAMS) produced by ECMWF is a reanalysis dataset that produces continuous data on atmospheric composition (Inness et al., 2019). The Integrated Forecast System (IFS) of ECMWF was integrated to allow for the data assimilation and modelling of aerosols, chemically reactive species and greenhouse gases. Apart from assimilated ozone, no stratospheric chemistry is simulated by the model system. In this study, CAMS reanalyses were used with a horizontal resolution of about 80 km. The vertical resolution consists of 60 pressure levels up to 0.1 hPa. Three-dimensional model output fields are available every 3 hours. Detailed information on the CAMS model architecture is given by Inness et al. (2019). An evaluation study of CAMS using aircraft observations was carried out by Wang et al. (2020). Biases of assimilated species like ozone are found to be less than 20% whereas discrepancies for gases like C₂H₆ and PAN are generally larger.

3 Results and discussion

In this section, vertical profiles retrieved from GLORIA measurements during the WISE campaign on 13 September 2017 over the North Atlantic region are shown. Observed GLORIA chemistry mode data are compared to EMAC and CAMS simulation results. The possible origin of air masses detected by GLORIA is also discussed.

3.1 GLORIA measurements

Retrieved volume mixing ratios of C₂H₆, C₂H₂, HCOOH, and PAN together with O₃ inferred from limb emission spectra during the WISE flight on 13 September 2017 are displayed in Fig. 6. Ozone is a molecule with highest concentrations in the stratosphere (Brasseur and Solomon, 2005). Hence, it can be used as a tracer to diagnose detected air masses whether they are of stratospheric or tropospheric origin. The general shape of O₃ VMR is strongly correlated with the tropopause as shown in Fig. 6a. There are two regions where stratospheric air comes down to about 7 km (around 14:45 UTC and around 16:50 UTC). Here, the tropopause layer reaches these low altitudes in the form of a stratospheric intrusion, while at the beginning and the end of the measurement phase the troposphere extends up to about 12 km. Trajectory calculations have shown that, due to a west-southwesterly mid- and upper tropospheric air flow in the region of the flight path shown in Fig. 1, GLORIA has sounded virtually the same air mass twice yielding to a kind of symmetry in the horizontal trace gas distribution before and after 16:10 UTC. Measured stratospheric ozone volume mixing ratios are within 0.1 and 0.8 ppmv. These values are in line with other mid-latitude remote sensing and in-situ observations performed in this altitude range (e.g., Cortesi et al., 2007; Livesey et al., 2008; Bourgeois et al., 2020).

Measured concentrations of the species C₂H₆ are shown in Fig. 6b. The VMR distribution of C₂H₆ is in parts anti-correlated to the one of ozone. Two regions of stronger enhanced VMR up to about 2.2 ppbv can be seen in the upper troposphere at the beginning and at the end of the measurement period over the region south of Ireland and near the coastline of The Netherlands and Belgium. In the stratosphere, no stronger enhanced C₂H₆ VMR levels are visible and values remain below about 0.6 ppbv, which can be confirmed by mid-latitude satellite (Rinsland et al., 2005; Glatthor et al., 2009; Wiecele et al., 2012) and in-situ aircraft measurements (Rinsland et al., 2005; Glatthor et al., 2009; Wiecele et al., 2012; Rudolph, 1995; Monks et al., 2018).

The same behaviour is present in the vertical and horizontal distribution of C₂H₂ amounts (see Fig. 6c). Elevated volume mixing ratios of up to 0.2 ppbv in the upper troposphere are clearly visible at the beginning and close to the end of the observation period. In the stratosphere, measured C₂H₂ values appear noisy and stay clearly below 0.1 ppbv most of the time. C₂H₂ VMR measured by GLORIA lies within the range of satellite (Rinsland et al., 2005; Wiecele et al., 2012) and aircraft in-situ data (Rinsland et al., 2005; Wiecele et al., 2012; Xiao et al., 2007) obtained in the same altitude region at mid-latitudes.

Measured HCOOH volume mixing ratios are depicted in Fig. 6d. As in the case of the previously mentioned species C₂H₆ and C₂H₂, large amounts of HCOOH are also visible at high altitudes in the troposphere during early and late time of the GLORIA observations with values up to 0.9 ppbv. Comparable amounts were seen by airborne in-situ measurements in the mid-latitude upper troposphere (Reiner et al., 1999). In contrast, stratospheric HCOOH values are low and not higher than 0.1 ppbv. These values are in accordance with space-borne mid-latitude observations by the Atmospheric Chemistry Experiment (ACE) Fourier transform spectrometer and the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) instrument in the altitude regime considered here (Rinsland et al., 2006; Grutter et al., 2010).

The two-dimensional cross section of PAN is shown in Fig. 6e. The distribution of VMR maxima and minima is more structured compared to the one of the previously regarded pollution trace gases. Increased amounts of PAN up to 0.4 ppbv are not only visible at the beginning and end of the observation period in the upper troposphere but also in the lowermost stratosphere around 15:00 UTC, at altitudes of 7 to 8 km. Somewhat less enhanced quantities are noticeable near 14 km around 15:30 UTC. Elevated PAN amounts of comparable magnitude have also been detected in the UTLS region by space-borne instruments (Coheur et al., 2007; Glatthor et al., 2007b; Wiegele et al., 2012; Ungermann et al., 2016; Glatthor et al., 2007b, 2007b) as well as by airborne in-situ measurements (Singh et al., 2000; Singh et al., 2001). This different shape of VMR distribution might be explained by very long lifetimes of PAN under cold UTLS conditions and by the fact that PAN is not emitted directly but dependent on the availability of precursor substances as described in Sect. 1.

3.2 Comparison to model simulations

The comparison of measured species to model simulations is presented in Figs. 7 and 8. The observed data have been temporally smoothed with a 39-point adjacent averaging routine to permit a more realistic comparison given the different horizontal resolutions of the measurement and the EMAC and CAMS simulations. For O₃, both simulations generally reproduce the tropospheric and stratospheric concentrations seen by GLORIA (see Fig. 7a-c) but with coarser spatial structure. The chemistry climate model EMAC is able to simulate finer structures while CAMS only produces a smooth distribution of assimilated ozone. Both models tend to slightly overestimate the amount of ozone in the troposphere. This is also visible in Fig. 8a-c where differences between both EMAC runs (with and without enhanced NMVOC emissions) and GLORIA observations are shown. The amount of simulated ozone in the

EMAC_2 run is only slightly higher (less than 10 ppbv) compared to the EMAC simulation without enhanced NMVOC emissions (VMR differences in Fig. 8b and Fig. 8c are therefore nearly the same).

Comparisons for C_2H_6 are displayed in Fig. 7d-f. Both models are able to qualitatively reproduce the temporal and spatial region of enhanced upper tropospheric C_2H_6 as observed by GLORIA. As in the case of ozone, EMAC again is able to display finer structures in the vertical and horizontal distribution of C_2H_6 compared to CAMS. However, deficits in the simulated absolute C_2H_6 quantities are clearly visible in both models, especially in the case of CAMS. A considerable underestimation of CAMS C_2H_6 with respect to airborne observations was already reported by Wang et al. (2020). The EMAC_2 simulation with increased NMVOC emissions at least reduces the difference to the GLORIA observations compared to the EMAC run without these stronger NMVOC emissions (see Fig. 8d-f).

For C_2H_2 , we note that EMAC predicts elevated concentrations in much the same region where GLORIA reports enhancements (see Fig. 7 g-h). In addition, the measured VMRs in the upper troposphere are only little underestimated in terms of their absolute amount by the EMAC_2 simulation using raised NMVOC emissions. In the stratosphere, simulated C_2H_2 amounts are too low compared to the measurement. Using standard NMVOC emissions in EMAC leads to an increased underestimation of C_2H_2 amounts compared to GLORIA (see Fig. 8g-i).

The comparison of the species HCOOH is shown in Fig. 7i-k and Fig. 8j-l. Elevated HCOOH concentrations, as recorded by GLORIA in the upper troposphere, are clearly underestimated by both models, especially in the CAMS simulation, although the atmospheric region of the (too weak) enhanced HCOOH amounts in the models agrees with the measured one. However, the EMAC_2 simulation at least reduces differences with respect to the GLORIA observations.

Looking at the temporal and spatial distribution of PAN the situation appears somewhat different to the comparisons discussed above (see Fig. 7l-n and Fig. 8m-o). The principal behaviour of enhanced PAN values in the upper troposphere is captured by both atmospheric models. EMAC produces slightly finer structures in the stratosphere compared to CAMS. However, the measured small scale variations in the amount of PAN especially near 14:30 UTC between 6 and 8 km are not reproduced by the EMAC_2 simulation with enhanced NMVOC emissions while the observed elevated PAN values around 15:00 UTC in the lowermost stratosphere from 7 to 8 km are also visible in the model output. Apart from the regions with the highest measured PAN amounts, EMAC_2 tends to overestimate the concentration of PAN

below about 13 km (what is not the case in the standard EMAC run). Interesting VMR variations are also seen by GLORIA in the stratosphere above 13 km. The PAN VMR maximum detected around 15:40 UTC near 14 km is not visible in the model simulations. The PAN VMR minima near 15:00 UTC and between 16:00 UTC and 16:15 UTC at about 13.5 km are reflected as a VMR minimum in EMAC_2, although with lower absolute quantities. The different shape of the horizontal and vertical distribution of PAN VMR is most probably caused by long-range atmospheric transport. Compared to the species discussed before, sources and sinks are different, and atmospheric lifetime of PAN is considerably longer. In the following subsection, we will focus on the origin of the polluted air masses, which have been detected by GLORIA.

3.3 Origin of polluted air masses

To estimate the geographical region (within the Earth's upper planetary boundary layer) of the origin of the measured enhanced amounts of the pollutants **and their pathways** we performed backward trajectory calculations as well as global three-dimensional CLaMS simulations with artificial tracers of air mass origin as described in the following subsections.

3.3.1 CLaMS backward trajectory calculations

To obtain a more detailed insight into the origin and transport pathways of air masses, backward trajectories with the three-dimensional Chemical Lagrangian Model of the Stratosphere (CLaMS; McKenna et al., 2002b; McKenna et al., 2002a; Pommrich et al., 2014) were performed starting from the GLORIA measurements. Although pure trajectories do not include mixing processes, they are well suited to analyse the history of transport pathways of air parcels in the tropics and in the region of the Asian monsoon into the UTLS (Vogel et al., 2014; Li et al., 2018; Ploeger et al., 2012). In this study, 20-, 40-, and 60-days diabatic backward trajectories with a horizontal resolution of 1 x 1 degrees were calculated using ERA-interim reanalysis wind data (Dee et al., 2011). **Frequently employed trajectory lengths to study transport processes in particular in the Asian monsoon region rang from a couple of weeks to a few months depending on the transport times from Earth's surface to atmospheric altitudes (Chen et al., 2012; Bergman et al., 2013; Garny and Randel, 2016; Müller et al., 2016; Li et al., 2018; Vogel et al., 2019; Legras and Bucci, 2020; Hanumanthu et al., 2020). The transport of air from the planetary boundary layer to regions with enhanced PAN between 13 and 14.5 km (about 400 K) needs up to 60 days, therefore a maximum trajectory length of 60 days was chosen for our analysis.**

These trajectories were generated for defined areas, where enhanced or low VMRs of pollutants have been detected by GLORIA. Selected regions are displayed as coloured boxes in Fig. 6. High amounts of pollutants are recorded within the cyan and blue boxes mainly in the upper troposphere. Air masses marked in these two boxes are located near the stratospheric intrusion. This region was probed twice, first at the beginning and second at the end of the flight (see Fig. 1). In addition, the green box marks enhanced quantities of observed PAN found in the stratosphere between 13 and 14.5 km (Fig. 6e). In contrast, the black box stands for an air mass where low pollution VMRs have been observed.

Fig. 9 gives an overview of the trajectory calculations. It is obvious that air masses were transported by westerly winds to the place of GLORIA observations. If we first regard the black box (as defined in Fig. 6) in the lowermost stratosphere, where low mixing ratios for all pollutant species discussed here were observed, we find that only few trajectories penetrate the upper planetary boundary layer (PBL) limit of 800 hPa on their way back from the GLORIA observation points within 20-, 40-, and 60 days (Fig. 9a-c). Further, most of these areas are located over the southern part of the North Pacific where we would not expect much pollution in the PBL. In contrast, looking at the cyan and blues boxes (of Fig. 6) marking mainly air masses in the upper troposphere near the flank of the stratospheric intrusion with generally high amounts of the pollutants, we find lots of trajectories going into the PBL not only over the North Pacific region but also over densely populated regions in Southeast Asia (Fig. 9d-i) where we expect direct anthropogenic emissions or precursors of the considered species (Lelieveld et al., 2001). Furthermore, marked areas of PBL penetration are also visible over the North American continent, especially in Canada where forest fires were frequent and intense in August and September 2017 (Pumphrey et al., in review, 2020; Torres et al., 2020; Hooghiem et al., 2020; Khaykin et al., 2018). These widespread Canadian fires correlate well with the marked PBL areas. Finally, we look at the green region in the stratosphere around 14 km (see Fig. 6) where enhanced amounts of PAN are visible in contrast to the non-elevated values of C_2H_6 , C_2H_2 , and $HCOOH$. The corresponding trajectory calculations (Fig. 9j-l) exhibit no PBL penetration areas in the case of 20- and 40-days backward trajectories. However, the 60-days backward calculations clearly show areas over densely populated Southeast Asia where trajectories entered the PBL. The ascending air masses are clearly visible in the changing colour of the potential temperature along the trajectories (Fig. 9l). This potential source region is located well within the Asian summer monsoon pollution pump (Lelieveld et al., 2018; Randel et al., 2010).

The trajectory calculations show that the air parcels with enhanced PAN are slowly uplifted by

410 diabatic heating superimposed on the anticyclonic motion within about 40 days, which Vogel
et al. (2019) referred to as an “upward spiralling range”. Subsequent transport (within about
20 days) occurred along the subtropical jet to the extratropical UTLS over the Atlantic in
agreement to previous aircraft measurements of long-lived trace gases (like CH₄, H₂O, CO,
N₂O, and SF₆) in the extratropical UTLS over the Atlantic (Müller et al., 2016; Vogel et al.,
415 2016). In this study, it is demonstrated that also for the pollutant PAN with a lifetime of up to
a few months this transport pathway from the Asian monsoon region to the extratropical UTLS
is important. Since PAN has this long lifetime in the free and upper troposphere which is longer
than the one of C₂H₆, C₂H₂, and HCOOH, it is likely that some amounts of PAN are still existing
in the stratospheric region of the green box while concentrations of the three other pollutant
420 species are already depleted.

Backward trajectory calculations are very useful to identify both the origin of an air parcel in
the PBL and its detailed transport pathways and transport times, however mixing processes
between different air parcels are neglected. Therefore, we use in addition global three-
dimensional CLaMS simulations considering mixing of air parcels to characterize the origin of
425 air masses.

3.3.2 Artificial tracers of air mass origin calculations

The Lagrangian three-dimensional chemistry transport model CLaMS (Pommrich et al., 2014)
was used to calculate artificial tracers of air mass origin (Vogel et al., 2016; Vogel et al., 2019).
These artificial tracers refer to marked geographical regions in the boundary layer of the global
430 atmosphere. An overview of these regions is given in Fig. 10 which is an updated configuration
compared to previous studies using artificial tracers of air mass origin in CLaMS. The upper
limit of the model boundary layer follows the orography and extends to about 2-3 km above the
Earth’s surface. In the currently used simulation, the model dynamics is driven by horizontal
winds from the ERA-interim reanalysis (Dee et al., 2011) provided by ECMWF. Transport of
435 air masses from the model boundary into the free troposphere and above is considered from 1
May 2017, the starting time of the simulation. Every 24 h (time step for mixing in CLaMS), air
masses in the model boundary layer are marked by the different tracers of air mass origin and
can be transported like a chemical tracer to other regions of the free troposphere or stratosphere
and subsequent mixing processes between different air masses can occur. Therefore, the value
440 of the individual artificial tracer of air mass origin counts the percentage of an air mass that

originated in the specific model boundary layer region since 1 May 2017 considering advection and mixing processes.

The results of the CLaMS simulation are displayed in Fig. 11. The origin of air masses seen inside the cyan and blue boxes (which contain the largest values of C_2H_6 , C_2H_2 , $HCOOH$, and PAN, see Fig. 6) below the tropical side of the stratospheric intrusion stem to a large part from North America, the Northeast and Northwest Pacific and the Tropical Eastern Pacific; to a smaller part also from Central America. Up to about 40 % in these boxes originate from the South Asian region which includes Tibetan Plateau, Eastern China, Northern India, Indian Ocean, Bay of Bengal, Indian subcontinent, Southeast Asia, and Warm pool (see Fig. 10). This is consistent with the findings of the trajectory calculations as shown in the previous section. Air masses in the stratosphere within the green box where enhanced PAN mixing ratios were detected by GLORIA come mainly from the South Asian region. This is also in agreement with the PBL penetration region of the backward trajectories as discussed before. Concerning the black box in the stratosphere with low concentrations of C_2H_6 , C_2H_2 , $HCOOH$, and PAN, it is obvious that only fragments of air originate from the North American, Northeast and Northwest Pacific region. Some patches of South Asian air masses are visible in the black box zone, obviously from regions without enhanced amounts of the pollutants discussed here.

The artificial tracers of air mass origin mark specific geographical regions in the model boundary layer and are therefore very useful to identify the origin of observed air masses including atmospheric mixing processes. However, some regions on the Earth's surface where high emissions of chemical tracers such as C_2H_6 , C_2H_2 , $HCOOH$, and PAN (or their precursors) really occur are not included in the CLaMS simulations. Nevertheless, CLaMS simulations are useful to show that the enhanced PAN mixing ratios in the lower stratosphere (green box) are mainly from South Asia in agreement to the trajectory calculations.

4 Conclusions

GLORIA observations of pollutant trace gases shown in this work were performed during a flight of the WISE aircraft campaign around the British Isles on 13 September 2017. One of the special characteristics of these remote sensing measurements are the high temporal and spatial resolution of the data. As discussed in Sect. 3.1, the amounts of these gases are in line with data from previously performed satellite and in-situ aircraft measurements. The pollutants can be transported into remote regions due to their long lifetime under appropriate atmospheric

conditions, like convective processes combined with strong wind regimes of the upper troposphere. The following main results can be stated:

475 First, enhanced volume mixing ratios of the pollutant gases C_2H_6 , C_2H_2 , $HCOOH$, and PAN were recorded by the GLORIA instrument in the upper troposphere with high temporal and spatial resolution. It is important to emphasize that these enhancements were detected far away from the emission sources of these species. This is possible due to their long atmospheric lifetimes in the order of weeks to months under free tropospheric conditions. Since PAN has
480 the longest lifetime of this foursome, elevated quantities of this molecule could be measured even in the lowermost stratosphere. Backward trajectories and artificial tracers of air mass origin calculations indicate that the main sources of the emitted species are on the one hand biomass forest fires in North America which reached their maximum a couple of weeks before the GLORIA flight (Pumphrey et al., in review, 2020; Torres et al., 2020) when air masses
485 detected by GLORIA passed this region. On the other hand, another important source region is located in the vast region of South and Southeast Asia where the Asian monsoon anticyclone governs the circulation regime during the summer months. Here, huge amounts of pollutants are lifted upwards into the upper troposphere and further transported to northern mid-latitudes via strong wind fields like the subtropical jet stream (Ungermann et al., 2016; Vogel et al.,
490 2016; Lelieveld et al., 2018; Legras and Bucci, 2020). Indeed, another GLORIA measurement carried out about 6 weeks earlier on 31 July 2017 during the StratoClim (Stratospheric and upper tropospheric processes for better climate predictions) campaign over India and Nepal also shows elevated amounts of more than 0.2 ppbv of C_2H_2 , more than 0.2 ppbv of $HCOOH$, and more than 0.5 ppbv of PAN (Johansson et al., 2020). These values are of comparable magnitude
495 to the ones observed during the WISE campaign. Our study demonstrates that PAN, with a lifetime of up to a few months, is transported from sources in Asia to the northern extratropical UTLS within about 2 months caused by the Asian monsoon which has implications for ozone and thus possibly for the radiative budget in that part of the atmosphere.

Second, the chemistry climate model EMAC and the CAMS assimilation system are able to
500 simulate tropospheric and stratospheric dynamical ozone VMR structures as seen by GLORIA although with coarser spatial resolution compared to the measurement (particularly for CAMS). In addition, both models reproduce the temporal and spatial region of enhanced upper tropospheric VMR levels of the measured pollutant species (C_2H_2 not available in CAMS). However, CAMS clearly underestimates the amount of elevated C_2H_6 , $HCOOH$, and to a lesser

505 extent, PAN. The EMAC_2 simulation using NMVOC emissions enhanced by a factor of 2 (as recommended by Monks et al., 2018) only slightly underestimates the C₂H₂ concentrations while simulated values for C₂H₆ and HCOOH are also too low. Compared to the EMAC standard emission run, the EMAC_2 simulation reduces differences to the GLORIA observations for these gases. In contrast, the size of elevated PAN values is overestimated by
510 the EMAC_2 model run. However, this does not hold for the local PAN VMR enhancements detected near 14 km, which are neither captured by EMAC_2 nor by CAMS.

This study has shown that observations of pollutant species are further needed since biomass burning and wildfires will still occur in the future and seem to have increased in the last years (Witze, 2020). There is still potential to improve chemical models with regard to reproduce the
515 measured VMR enhancements of the pollutant gases in more detail. **The rather coarse model resolution together with deficits in the vertical transport from convection in the meteorological fields used to drive the model impede the reproduction of measured structures of the pollutant trace gas concentrations.** A further aspect is to improve the emission databases in the models because the simulated amount of pollution species is strongly dependent on the local emission
520 place and the intensity of the emissions.

Data availability. GLORIA measurements are available in the database HALO-DB (<https://halo-db.pa.op.dlr.de/mission/96>) and will be available on the KITopen repository. The CAMS model data is available from ECMWF ([https://apps.ecmwf.int/data-catalogues/cams-](https://apps.ecmwf.int/data-catalogues/cams-reanalysis)
525 [reanalysis](https://apps.ecmwf.int/data-catalogues/cams-reanalysis)). EMAC and CLaMS data are available upon request.

Author contributions. GW wrote the paper and performed the bulk of the data analysis, with input from all co-authors. SJ, AK, JU, MH, and NG performed the GLORIA data processing. FFV, TG, EK, GM, HN, and CP operated GLORIA during the WISE campaign in Shannon and
530 Oberpfaffenhofen. OK and FK performed the EMAC simulations and designed the sensitivity studies. BV and JUG performed the CLaMS trajectory- and artificial tracers of air mass origin calculations. BMS, HO, and JO **directed the research and flight planning.** All authors commented on and improved the manuscript.

535 *Competing interests.* The authors declare that they have no conflict of interest.

Special issue statement. This article is part of the special issue “WISE: Wave-driven isentropic exchange in the extratropical upper stratosphere and lower stratosphere (ACP/AMT/WCD inter-journal SI), 2019”. It is not associated with a conference.

540

Acknowledgements. We acknowledge support by the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG Priority Program SPP 1294). We are grateful to the WISE coordination team for excellently conducting the aircraft campaign. Results are based on the efforts of all members of the GLORIA team, including the technology institutes ZEA-1 and
545 ZEA-2 at Forschungszentrum Jülich and the Institute for Data Processing and Electronics at the Karlsruhe Institute of Technology. We would also like to thank the pilots and ground-support team at the Flight Experiments facility of the Deutsches Zentrum für Luft- und Raumfahrt (DLR-FX). We thank ECMWF for providing CAMS data. The EMAC simulations were performed on the supercomputer ForHLR funded by the Ministry of Science, Research and the
550 Arts Baden-Württemberg and by the Federal Ministry of Education and Research. The CLaMS activities contribute to the DFG project AMOS (HALO-SPP 1294/VO 1276/5-1) funded by the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG). The authors gratefully acknowledge the computing time for the CLaMS simulations granted on the supercomputer JURECA at Jülich Supercomputing Centre (JSC) under the VSR project ID
555 JICG11.

Financial support. We acknowledge support by Deutsche Forschungsgemeinschaft and Open Access Publishing Fund of Karlsruhe Institute of Technology.

- Alvarado, L. M. A., Richter, A., Vrekoussis, M., Hilboll, A., Kalisz Hedegaard, A. B., Schneising, O., and Burrows, J. P.: Unexpected long-range transport of glyoxal and formaldehyde observed from the Copernicus Sentinel-5 Precursor satellite during the 2018 Canadian wildfires, *Atmos. Chem. Phys.*, 20, 2057–2072, <https://doi.org/10.5194/acp-20-2057-2020>, 2020.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume III - gas phase reactions of inorganic halogens, *Atmos. Chem. Phys.*, 7, 981–1191, <https://doi.org/10.5194/acp-7-981-2007>, 2007.
- Bergman, J. W., Fierli, F., Jensen, E. J., Honomichl, S., and Pan, L. L.: Boundary layer sources for the Asian anticyclone: Regional contributions to a vertical conduit, *J. Geophys. Res. Atmos.*, 118, 2560–2575, <https://doi.org/10.1002/jgrd.50142>, 2013.
- Bourgeois, I., Peischl, J., Thompson, C. R., Aikin, K. C., Campos, T., Clark, H., Commane, R., Daube, B., Diskin, G. W., Elkins, J. W., Gao, R.-S., Gaudel, A., Hints, E. J., Johnson, B. J., Kivi, R., McKain, K., Moore, F. L., Parrish, D. D., Querel, R., Ray, E., Sánchez, R., Sweeney, C., Tarasick, D. W., Thompson, A. M., Thouret, V., Witte, J. C., Wofsy, S. C., and Ryerson, T. B.: Global-scale distribution of ozone in the remote troposphere from the ATom and HIPPO airborne field missions, *Atmos. Chem. Phys.*, 20, 10611–10635, <https://doi.org/10.5194/acp-20-10611-2020>, 2020.
- Brasseur, G. P. and Solomon, S.: *Aeronomy of the Middle Atmosphere: Chemistry and Physics of the Stratosphere and Mesosphere*, Third revised and enlarged edition, Atmospheric and Oceanographic Sciences Library, 32, Springer, Dordrecht, 2005.
- Chen, B., Xu, X. D., Yang, S., and Zhao, T. L.: Climatological perspectives of air transport from atmospheric boundary layer to tropopause layer over Asian monsoon regions during boreal summer inferred from Lagrangian approach, *Atmos. Chem. Phys.*, 12, 5827–5839, <https://doi.org/10.5194/acp-12-5827-2012>, 2012.
- Coheur, P.-F., Herbin, H., Clerbaux, C., Hurtmans, D., Wespes, C., Carleer, M., Turquety, S., Rinsland, C. P., Remedios, J., Hauglustaine, D., Boone, C. D., and Bernath, P. F.: ACE-FTS observation of a young biomass burning plume: first reported measurements of C₂H₄,

- 590 C₃H₆O, H₂CO and PAN by infrared occultation from space, *Atmos. Chem. Phys.*, 7,
5437–5446, <https://doi.org/10.5194/acp-7-5437-2007>, 2007.
- Cortesi, U., Lambert, J. C., Clercq, C. de, Bianchini, G., Blumenstock, T., Bracher, A.,
Castelli, E., Catoire, V., Chance, K. V., Mazière, M. de, Demoulin, P., Godin-Beekmann,
S., Jones, N., Jucks, K., Keim, C., Kerzenmacher, T., Kuellmann, H., Kuttippurath, J.,
595 Iarlori, M., Liu, G. Y., Liu, Y., McDermid, I. S., Meijer, Y. J., Mencaraglia, F., Mikuteit,
S., Oelhaf, H., Piccolo, C., Pirre, M., Raspollini, P., Ravegnani, F., Reburn, W. J.,
Redaelli, G., Remedios, J. J., Sembhi, H., Smale, D., Steck, T., Taddei, A., Varotsos, C.,
Vigouroux, C., Waterfall, A., Wetzel, G., and Wood, S.: Geophysical validation of
MIPAS-ENVISAT operational ozone data, *Atmos. Chem. Phys.*, 7, 4807–4867,
600 <https://doi.org/10.5194/acp-7-4807-2007>, 2007.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,
Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg,
L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger,
L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M.,
605 Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey,
C., Rosnay, P. de, Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim
reanalysis: configuration and performance of the data assimilation system, *Q.J.R.
Meteorol. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- Fadnavis, S., Semeniuk, K., Schultz, M. G., Kiefer, M., Mahajan, A., Pozzoli, L., and
610 Sonbawane, S.: Transport pathways of peroxyacetyl nitrate in the upper troposphere and
lower stratosphere from different monsoon systems during the summer monsoon season,
Atmos. Chem. Phys., 15, 11477–11499, <https://doi.org/10.5194/acp-15-11477-2015>, 2015.
- Fadnavis, S., Schultz, M. G., Semeniuk, K., Mahajan, A. S., Pozzoli, L., Sonbawane, S.,
Ghude, S. D., Kiefer, M., and Eckert, E.: Trends in peroxyacetyl nitrate (PAN) in the
615 upper troposphere and lower stratosphere over southern Asia during the summer monsoon
season: regional impacts, *Atmos. Chem. Phys.*, 14, 12725–12743,
<https://doi.org/10.5194/acp-14-12725-2014>, 2014.
- Fischer, E. V., Jacob, D. J., Yantosca, R. M., Sulprizio, M. P., Millet, D. B., Mao, J., Paulot,
F., Singh, H. B., Roiger, A., Ries, L., Talbot, R. W., Dzepina, K., and Pandey Deolal, S.:
620 Atmospheric peroxyacetyl nitrate (PAN): a global budget and source attribution, *Atmos.
Chem. Phys.*, 14, 2679–2698, <https://doi.org/10.5194/acp-14-2679-2014>, 2014.

Forster, P. M. d. F. and Shine, K. P.: Radiative forcing and temperature trends from stratospheric ozone changes, *J. Geophys. Res.*, 102, 10841–10855, <https://doi.org/10.1029/96JD03510>, 1997.

625 Friedl-Vallon, F., Gulde, T., Hase, F., Kleinert, A., Kulesa, T., Maucher, G., Neubert, T., Olschewski, F., Piesch, C., Preusse, P., Rongen, H., Sartorius, C., Schneider, H., Schönfeld, A., Tan, V., Bayer, N., Blank, J., Dapp, R., Ebersoldt, A., Fischer, H., Graf, F., Guggenmoser, T., Höpfner, M., Kaufmann, M., Kretschmer, E., Latzko, T., Nordmeyer, H., Oelhaf, H., Orphal, J., Riese, M., Schardt, G., Schillings, J., Sha, M. K., Suminska-
630 Ebersoldt, O., and Ungermann, J.: Instrument concept of the imaging Fourier transform spectrometer GLORIA, *Atmos. Meas. Tech.*, 7, 3565–3577, <https://doi.org/10.5194/amt-7-3565-2014>, 2014.

Garny, H. and Randel, W. J.: Transport pathways from the Asian monsoon anticyclone to the stratosphere, *Atmos. Chem. Phys.*, 16, 2703–2718, [https://doi.org/10.5194/acp-16-2703-](https://doi.org/10.5194/acp-16-2703-2016)
635 2016, 2016.

Glatthor, N., Clarmann, T. von, Stiller, G. P., Funke, B., Koukouli, M. E., Fischer, H., Grabowski, U., Höpfner, M., Kellmann, S., and Linden, A.: Large-scale upper tropospheric pollution observed by MIPAS HCN and C₂H₆ global distributions, *Atmos. Chem. Phys.*, 9, 9619–9634, <https://doi.org/10.5194/acp-9-9619-2009>, 2009.

640 Gordon, I. E., Rothman, L. S., Hill, C., Kochanov, R. V., Tan, Y., Bernath, P. F., Birk, M., Boudon, V., Campargue, A., Chance, K. V., Drouin, B. J., Flaud, J.-M., Gamache, R. R., Hodges, J. T., Jacquemart, D., Perevalov, V. I., Perrin, A., Shine, K. P., Smith, M.-A., Tennyson, J., Toon, G. C., Tran, H., Tyuterev, V. G., Barbe, A., Császár, A. G., Devi, V. M., Furtenbacher, T., Harrison, J. J., Hartmann, J.-M., Jolly, A., Johnson, T. J., Karman, T., Kleiner, I., Kyuberis, A. A., Loos, J., Lyulin, O. M., Massie, S. T., Mikhailenko, S. N.,
645 Moazzen-Ahmadi, N., Müller, H., Naumenko, O. V., Nikitin, A. V., Polyansky, O. L., Rey, M., Rotger, M., Sharpe, S. W., Sung, K., Starikova, E., Tashkun, S. A., Auwera, J. V., Wagner, G., Wilzewski, J., Wcisło, P., Yu, S., and Zak, E. J.: The HITRAN2016 molecular spectroscopic database, *Journal of Quantitative Spectroscopy and Radiative*
650 *Transfer*, 203, 3–69, <https://doi.org/10.1016/j.jqsrt.2017.06.038>, 2017.

Granier, C., Bessagnet, B., Bond, T., D’Angiola, A., van der Denier Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J.-F., Liousse, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J.-C., Riahi, K., Schultz, M. G.,

- Smith, S. J., Thompson, A., van Aardenne, J., van der Werf, G. R., and van Vuuren, D. P.:
 655 Evolution of anthropogenic and biomass burning emissions of air pollutants at global and
 regional scales during the 1980–2010 period, *Climatic Change*, 109, 163–190,
<https://doi.org/10.1007/s10584-011-0154-1>, 2011.
- Grutter, M., Glatthor, N., Stiller, G. P., Fischer, H., Grabowski, U., Höpfner, M., Kellmann,
 S., Linden, A., and Clarmann, T. von: Global distribution and variability of formic acid as
 660 observed by MIPAS-ENVISAT, *J. Geophys. Res.*, 115, 955,
<https://doi.org/10.1029/2009JD012980>, 2010.
- Hansen, J., Sato, M., and Ruedy, R.: Radiative forcing and climate response, *J. Geophys.*
Res., 102, 6831–6864, <https://doi.org/10.1029/96JD03436>, 1997.
- Hanumanthu, S., Vogel, B., Müller, R., Brunamonti, S., Fadnavis, S., Li, D., Ölsner, P., Naja,
 665 M., Singh, B. B., Kumar, K. R., Sonbawne, S., Jauhiainen, H., Vömel, H., Luo, B., Jorge,
 T., Wienhold, F. G., Dirksen, R., and Peter, T.: Strong day-to-day variability of the Asian
 Tropopause Aerosol Layer (ATAL) in August 2016 at the Himalayan foothills, *Atmos.*
Chem. Phys., 20, 14273–14302, <https://doi.org/10.5194/acp-20-14273-2020>, 2020.
- Hooghiem, J. J. D., Popa, M. E., Röckmann, T., Groöß, J.-U., Tritscher, I., Müller, R., Kivi,
 670 R., and Chen, H.: Wildfire smoke in the lower stratosphere identified by in situ CO
 observations, *Atmos. Chem. Phys. Discuss.*, 2020, 1–27, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-2020-65)
[2020-65](https://doi.org/10.5194/acp-2020-65), available at: <https://acp.copernicus.org/preprints/acp-2020-65/>, 2020.
- Höpfner, M., Oelhaf, H., Wetzel, G., Friedl-Vallon, F., Kleinert, A., Lengel, A., Maucher, G.,
 Nordmeyer, H., Glatthor, N., Stiller, G., Clarmann, T. v., Fischer, H., Kröger, C., and
 675 Deshler, T.: Evidence of scattering of tropospheric radiation by PSCs in mid-IR limb
 emission spectra: MIPAS-B observations and KOPRA simulations, *Geophys. Res. Lett.*,
 29, 119-1-119-4, <https://doi.org/10.1029/2001GL014443>, 2002.
- Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M.,
 Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling,
 680 Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy, S., Schulz, M., and
 Suttie, M.: The CAMS reanalysis of atmospheric composition, *Atmos. Chem. Phys.*, 19,
 3515–3556, <https://doi.org/10.5194/acp-19-3515-2019>, 2019.
- Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A.,
 Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System

- 685 (MESSy2), *Geosci. Model Dev.*, 3, 717–752, <https://doi.org/10.5194/gmd-3-717-2010>, 2010.
- Johansson, S., Höpfner, M., Kirner, O., Wohltmann, I., Bucci, S., Legras, B., Friedl-Vallon, F., Glatthor, N., Kretschmer, E., Ungermann, J., and Wetzel, G.: Pollution trace gas distributions and their transport in the Asian monsoon upper troposphere and lowermost
690 stratosphere during the StratoClim campaign 2017, *Atmos. Chem. Phys.*, 20, 14695–14715, <https://doi.org/10.5194/acp-20-14695-2020>, 2020.
- Johansson, S., Woiwode, W., Höpfner, M., Friedl-Vallon, F., Kleinert, A., Kretschmer, E., Latzko, T., Orphal, J., Preusse, P., Ungermann, J., Santee, M. L., Jurkat-Witschas, T., Marsing, A., Voigt, C., Giez, A., Krämer, M., Rolf, C., Zahn, A., Engel, A., Sinnhuber,
695 B.-M., and Oelhaf, H.: Airborne limb-imaging measurements of temperature, HNO₃, O₃, ClONO₂, H₂O and CFC-12 during the Arctic winter 2015/2016: characterization, in situ validation and comparison to Aura/MLS, *Atmos. Meas. Tech.*, 11, 4737–4756, <https://doi.org/10.5194/amt-11-4737-2018>, 2018.
- Khare, P., Kumar, N., Kumari, K. M., and Srivastava, S. S.: Atmospheric formic and acetic
700 acids: An overview, *Rev. Geophys.*, 37, 227–248, <https://doi.org/10.1029/1998RG900005>, 1999.
- Khaykin, S. M., Godin-Beekmann, S., Hauchecorne, A., Pelon, J., Ravetta, F., and Keckhut, P.: Stratospheric smoke with unprecedentedly high backscatter observed by lidars above southern France, *Geophys. Res. Lett.*, 45, 1639–1646,
705 <https://doi.org/10.1002/2017GL076763>, 2018.
- Kleinert, A., Friedl-Vallon, F., Guggenmoser, T., Höpfner, M., Neubert, T., Ribalda, R., Sha, M. K., Ungermann, J., Blank, J., Ebersoldt, A., Kretschmer, E., Latzko, T., Oelhaf, H., Olschewski, F., and Preusse, P.: Level 0 to 1 processing of the imaging Fourier transform spectrometer GLORIA: generation of radiometrically and spectrally calibrated spectra,
710 *Atmos. Meas. Tech.*, 7, 4167–4184, <https://doi.org/10.5194/amt-7-4167-2014>, 2014.
- Lamarque, J.-F., Dentener, F., McConnell, J., Ro, C.-U., Shaw, M., Vet, R., Bergmann, D., Cameron-Smith, P., Dalsoren, S., Doherty, R., Faluvegi, G., Ghan, S. J., Josse, B., Lee, Y. H., MacKenzie, I. A., Plummer, D., Shindell, D. T., Skeie, R. B., Stevenson, D. S., Strode, S., Zeng, G., Curran, M., Dahl-Jensen, D., Das, S., Fritzsche, D., and Nolan, M.: Multi-
715 model mean nitrogen and sulfur deposition from the Atmospheric Chemistry and Climate

Model Intercomparison Project (ACCMIP): evaluation of historical and projected future changes, *Atmos. Chem. Phys.*, 13, 7997–8018, <https://doi.org/10.5194/acp-13-7997-2013>, 2013.

Legras, B. and Bucci, S.: Confinement of air in the Asian monsoon anticyclone and pathways of convective air to the stratosphere during the summer season, *Atmos. Chem. Phys.*, 20, 11045–11064, <https://doi.org/10.5194/acp-20-11045-2020>, 2020.

Lelieveld, J., Bourtsoukidis, E., Brühl, C., Fischer, H., Fuchs, H., Harder, H., Hofzumahaus, A., Holland, F., Marno, D., Neumaier, M., Pozzer, A., Schlager, H., Williams, J., Zahn, A., and Ziereis, H.: The South Asian monsoon-pollution pump and purifier, *Science* (New York, N.Y.), 361, 270–273, <https://doi.org/10.1126/science.aar2501>, 2018.

Lelieveld, J., Crutzen, P. J., Ramanathan, V., Andreae, M. O., Brenninkmeijer, C. M., Campos, T., Cass, G. R., Dickerson, R. R., Fischer, H., Gouw, J. A. de, Hansel, A., Jefferson, A., Kley, D., Laat, A. T. de, Lal, S., Lawrence, M. G., Lobert, J. M., Mayol-Bracero, O. L., Mitra, A. P., Novakov, T., Oltmans, S. J., Prather, K. A., Reiner, T., Rodhe, H., Scheeren, H. A., Sikka, D., and Williams, J.: The Indian Ocean experiment: widespread air pollution from South and Southeast Asia, *Science*, 291, 1031–1036, <https://doi.org/10.1126/science.1057103>, available at: <https://science.sciencemag.org/content/291/5506/1031/tab-pdf>, 2001.

Li, D., Vogel, B., Müller, R., Bian, J., Günther, G., Li, Q., Zhang, J., Bai, Z., Vömel, H., and Riese, M.: High tropospheric ozone in Lhasa within the Asian summer monsoon anticyclone in 2013: influence of convective transport and stratospheric intrusions, *Atmos. Chem. Phys.*, 18, 17979–17994, <https://doi.org/10.5194/acp-18-17979-2018>, 2018.

Livesey, N. J., Filipiak, M. J., Froidevaux, L., Read, W. G., Lambert, A., Santee, M. L., Jiang, J. H., Pumphrey, H. C., Waters, J. W., Cofield, R. E., Cuddy, D. T., Daffer, W. H., Drouin, B. J., Fuller, R. A., Jarnot, R. F., Jiang, Y. B., Knosp, B. W., Li, Q. B., Perun, V. S., Schwartz, M. J., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. A., Avery, M., Browell, E. V., Cammas, J.-P., Christensen, L. E., Diskin, G. S., Gao, R.-S., Jost, H.-J., Loewenstein, M., Lopez, J. D., Nedelec, P., Osterman, G. B., Sachse, G. W., and Webster, C. R.: Validation of Aura Microwave Limb Sounder O₃ and CO observations in the upper troposphere and lower stratosphere, *J. Geophys. Res.*, 113, <https://doi.org/10.1029/2007JD008805>, 2008.

- Lu, X., Zhang, L., and Shen, L.: Meteorology and Climate Influences on Tropospheric Ozone: a Review of Natural Sources, Chemistry, and Transport Patterns, *Curr Pollution Rep*, 5, 238–260, <https://doi.org/10.1007/S40726-019-00118-3>, 2019.
- 750 McKenna, D. S., Konopka, P., Grooß, J.-U., Günther, G., Müller, R., Spang, R., Offermann, D., and Orsolini, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS) 1. Formulation of advection and mixing, *J. Geophys. Res.*, 107, <https://doi.org/10.1029/2000JD000114>, 2002a.
- 755 McKenna, D. S., Grooß, J.-U., Günther, G., Konopka, P., Müller, R., Carver, G., and Sasano, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS) 2. Formulation of chemistry scheme and initialization, *J. Geophys. Res.*, 107, <https://doi.org/10.1029/2000JD000113>, 2002b.
- 760 Millet, D. B., Baasandorj, M., Farmer, D. K., Thornton, J. A., Baumann, K., Brophy, P., Chaliyakunnel, S., Gouw, J. A. de, Graus, M., Hu, L., Koss, A., Lee, B. H., Lopez-Hilfiker, F. D., Neuman, J. A., Paulot, F., Peischl, J., Pollack, I. B., Ryerson, T. B., Warneke, C., Williams, B. J., and Xu, J.: A large and ubiquitous source of atmospheric formic acid, *Atmos. Chem. Phys.*, 15, 6283–6304, <https://doi.org/10.5194/acp-15-6283-2015>, 2015.
- 765 Monks, S. A., Wilson, C., Emmons, L. K., Hannigan, J. W., Helmig, D., Blake, N. J., and Blake, D. R.: Using an Inverse Model to Reconcile Differences in Simulated and Observed Global Ethane Concentrations and Trends Between 2008 and 2014, *J. Geophys. Res. Atmos.*, 123, 11,262–11,282, <https://doi.org/10.1029/2017JD028112>, 2018.
- 770 Montzka, S. A., Butler, J. H., Elkins, J. W., Thompson, T. M., Clarke, A. D., and Lock, L. T.: Present and future trends in the atmospheric burden of ozone-depleting halogens, *Nature*, 398, 690–694, <https://doi.org/10.1038/19499>, 1999.
- 775 Müller, S., Hoor, P., Bozem, H., Gute, E., Vogel, B., Zahn, A., Bönisch, H., Keber, T., Krämer, M., Rolf, C., Riese, M., Schlager, H., and Engel, A.: Impact of the Asian monsoon on the extratropical lower stratosphere: trace gas observations during TACTS over Europe 2012, *Atmos. Chem. Phys.*, 16, 10573–10589, <https://doi.org/10.5194/acp-16-10573-2016>, 2016.
- Mungall, E. L., Abbatt, J. P. D., Wentzell, J. J. B., Wentworth, G. R., Murphy, J. G., Kunkel, D., Gute, E., Tarasick, D. W., Sharma, S., Cox, C. J., Uttal, T., and Liggio, J.: High gas-

- phase mixing ratios of formic and acetic acid in the High Arctic, *Atmos. Chem. Phys.*, 18, 10237–10254, <https://doi.org/10.5194/acp-18-10237-2018>, 2018.
- 780 Norton, R. H. and Beer, R.: New apodizing functions for Fourier spectrometry, *J. Opt. Soc. Am.*, 66, 259, <https://doi.org/10.1364/JOSA.66.000259>, 1976.
- Paulot, F., Wunch, D., Crounse, J. D., Toon, G. C., Millet, D. B., DeCarlo, P. F., Vigouroux, C., Deutscher, N. M., González Abad, G., Notholt, J., Warneke, T., Hannigan, J. W., Warneke, C., Gouw, J. A. de, Dunlea, E. J., Mazière, M. de, Griffith, D. W. T., Bernath, P., Jimenez, J. L., and Wennberg, P. O.: Importance of secondary sources in the atmospheric budgets of formic and acetic acids, *Atmos. Chem. Phys.*, 11, 1989–2013, <https://doi.org/10.5194/acp-11-1989-2011>, 2011.
- 785 Phillips, D. L.: A technique for the numerical solution of certain integral equations of the first kind, *J. Assoc. Comput. Math.*, 9, 84–97, <https://doi.org/10.1145/321105.321114>, 1962.
- 790 Ploeger, F., Konopka, P., Müller, R., Fueglistaler, S., Schmidt, T., Manners, J. C., Grooß, J.-U., Günther, G., Forster, P. M., and Riese, M.: Horizontal transport affecting trace gas seasonality in the Tropical Tropopause Layer (TTL), *J. Geophys. Res.*, 117, n/a-n/a, <https://doi.org/10.1029/2011JD017267>, 2012.
- Pommrich, R., Müller, R., Grooß, J.-U., Konopka, P., Ploeger, F., Vogel, B., Tao, M., Hoppe, C. M., Günther, G., Spelten, N., Hoffmann, L., Pumphrey, H.-C., Viciani, S., D'Amato, F., Volk, C. M., Hoor, P., Schlager, H., and Riese, M.: Tropical troposphere to stratosphere transport of carbon monoxide and long-lived trace species in the Chemical Lagrangian Model of the Stratosphere (CLaMS), *Geosci. Model Dev.*, 7, 2895–2916, <https://doi.org/10.5194/gmd-7-2895-2014>, 2014.
- 795 Pumphrey, H. C., Schwartz, M. J., Santee, M. L., Kablick III, G. P., Fromm, M. D., and Livesey, N. J.: Stratospheric pollution from Canadian forest fires, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-840>, in review, 2020.
- Randel, W. J., Park, M., Emmons, L., Kinnison, D., Bernath, P., Walker, K. A., Boone, C., and Pumphrey, H.: Asian monsoon transport of pollution to the stratosphere, *Science* (New York, N.Y.), 328, 611–613, <https://doi.org/10.1126/science.1182274>, 2010.
- 805 Reiner, T., Möhler, O., and Arnold, F.: Measurements of acetone, acetic acid, and formic acid in the northern midlatitude upper troposphere and lower stratosphere, *J. Geophys. Res.*, 104, 13943–13952, <https://doi.org/10.1029/1999JD900030>, 1999.

- Remedios, J. J., Leigh, R. J., Waterfall, A. M., Moore, D. P., Sembhi, H., Parkes, I.,
810 Greenhough, J., Chipperfield, M. P., and Hauglustaine, D.: MIPAS reference atmospheres
and comparisons to V4.61/V4.62 MIPAS level 2 geophysical data sets, *Atmos. Chem.*
Phys. Discuss., 7, 9973–10017, <https://doi.org/10.5194/acpd-7-9973-2007>, 2007.
- Riese, M., Oelhaf, H., Preusse, P., Blank, J., Ern, M., Friedl-Vallon, F., Fischer, H.,
Guggenmoser, T., Höpfner, M., Hoor, P., Kaufmann, M., Orphal, J., Plöger, F., Spang, R.,
815 Suminska-Ebersoldt, O., Ungermann, J., Vogel, B., and Woiwode, W.: Gimballing Limb
Observer for Radiance Imaging of the Atmosphere (GLORIA) scientific objectives,
Atmos. Meas. Tech., 7, 1915–1928, <https://doi.org/10.5194/amt-7-1915-2014>, 2014.
- Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., and Forster, P.: Impact
of uncertainties in atmospheric mixing on simulated UTLS composition and related
820 radiative effects, *J. Geophys. Res.*, 117, D16305, <https://doi.org/10.1029/2012JD017751>,
2012.
- Rinsland, C. P., Boone, C. D., Bernath, P. F., Mahieu, E., Zander, R., Dufour, G., Clerbaux,
C., Turquety, S., Chiou, L., McConnell, J. C., Neary, L., and Kaminski, J. W.: First space-
based observations of formic acid (HCOOH): Atmospheric Chemistry Experiment austral
825 spring 2004 and 2005 Southern Hemisphere tropical-mid-latitude upper tropospheric
measurements, *Geophys. Res. Lett.*, 33, 955, <https://doi.org/10.1029/2006GL027128>,
2006.
- Rinsland, C. P., Dufour, G., Boone, C. D., Bernath, P. F., and Chiou, L.: Atmospheric
Chemistry Experiment (ACE) measurements of elevated Southern Hemisphere upper
830 tropospheric CO, C₂H₆, HCN, and C₂H₂ mixing ratios from biomass burning emissions
and long-range transport, *Geophys. Res. Lett.*, 32, 24043,
<https://doi.org/10.1029/2005GL024214>, 2005.
- Rodgers, C. D.: *Inverse methods for atmospheric sounding*, 2, World Sci., 2000.
- Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornblueh, L., Manzini,
835 E., Schlese, U., and Schulzweida, U.: Sensitivity of Simulated Climate to Horizontal and
Vertical Resolution in the ECHAM5 Atmosphere Model, *J. Climate*, 19, 3771–3791,
<https://doi.org/10.1175/JCLI3824.1>, 2006.
- Rudolph, J.: The tropospheric distribution and budget of ethane, *J. Geophys. Res.*, 100,
11369, <https://doi.org/10.1029/95JD00693>, 1995.

- 840 Sander, S. P., Friedl, R. R., Barker, J. R., Golden, D. M., Kurylo, M. J., Wine, P. H., Abbatt,
J. P. D., Burkholder, J. B., Kolb, C. E., Moortgat, G. K., Huie, R. E., and Orkin, V. L.:
Chemical kinetics and photochemical data for use in atmospheric studies, Evaluation no.
17, JPL Publ. 10-6, Jet Propulsion Laboratory, Pasadena, CA, 2011.
- Singh, H., Chen, Y., Staudt, A., Jacob, D., Blake, D., Heikes, B., and Snow, J.: Evidence from
845 the Pacific troposphere for large global sources of oxygenated organic compounds,
Nature, 410, 1078–1081, <https://doi.org/10.1038/35074067>, 2001.
- Singh, H., Chen, Y., Tabazadeh, A., Fukui, Y., Bey, I., Yantosca, R., Jacob, D., Arnold, F.,
Wohlfrom, K., Atlas, E., Flocke, F., Blake, D., Blake, N., Heikes, B., Snow, J., Talbot, R.,
Gregory, G., Sachse, G., Vay, S., and Kondo, Y.: Distribution and fate of selected
850 oxygenated organic species in the troposphere and lower stratosphere over the Atlantic, J.
Geophys. Res., 105, 3795–3805, 2000.
- Singh, H. B.: Reactive nitrogen in the troposphere, Environ. Sci. Technol., 21, 320–327,
<https://doi.org/10.1021/es00158a001>, 1987.
- Spang, R., Remedios, J., and Barkley, M.: Colour indices for the detection and differentiation
855 of cloud types in infra-red limb emission spectra, Advances in Space Research, 33, 1041–
1047, [https://doi.org/10.1016/S0273-1177\(03\)00585-4](https://doi.org/10.1016/S0273-1177(03)00585-4), 2004.
- Stiller, G. P., Clarmann, T. von, Funke, B., Glatthor, N., Hase, F., Höpfner, M., and Linden,
A.: Sensitivity of trace gas abundances retrievals from infrared limb emission spectra to
simplifying approximations in radiative transfer modelling, Journal of Quantitative
860 Spectroscopy and Radiative Transfer, 72, 249–280, [https://doi.org/10.1016/S0022-4073\(01\)00123-6](https://doi.org/10.1016/S0022-4073(01)00123-6), 2002.
- Tikhonov, A. N.: On the solution of ill-posed problems and the method of regularization,
Dokl. Acad. Nauk SSSR, 151, 501–504, 1963.
- Torres, O., Bhartia, P. K., Taha, G., Jethva, H., Das, S., Colarco, P., Krotkov, N., Omar, A.,
865 and Ahn, C.: Stratospheric Injection of Massive Smoke Plume From Canadian Boreal
Fires in 2017 as Seen by DSCOVR-EPIC, CALIOP, and OMPS-LP Observations, J.
Geophys. Res. Atmos., 125, <https://doi.org/10.1029/2020JD032579>, 2020.
- Ungermann, J., Ern, M., Kaufmann, M., Müller, R., Spang, R., Ploeger, F., Vogel, B., and
Riese, M.: Observations of PAN and its confinement in the Asian summer monsoon

- 870 anticyclone in high spatial resolution, *Atmos. Chem. Phys.*, 16, 8389–8403,
<https://doi.org/10.5194/acp-16-8389-2016>, 2016.
- Vogel, B., Günther, G., Müller, R., Grooß, J.-U., Hoor, P., Krämer, M., Müller, S., Zahn, A.,
 and Riese, M.: Fast transport from Southeast Asia boundary layer sources to northern
 Europe: rapid uplift in typhoons and eastward eddy shedding of the Asian monsoon
 875 anticyclone, *Atmos. Chem. Phys.*, 14, 12745–12762, <https://doi.org/10.5194/acp-14-12745-2014>, 2014.
- Vogel, B., Müller, R., Günther, G., Spang, R., Hanumanthu, S., Li, D., Riese, M., and Stiller,
 G. P.: Lagrangian simulations of the transport of young air masses to the top of the Asian
 monsoon anticyclone and into the tropical pipe, *Atmos. Chem. Phys.*, 19, 6007–6034,
 880 <https://doi.org/10.5194/acp-19-6007-2019>, 2019.
- Vogel, B., Günther, G., Müller, R., Grooß, J.-U., Afchine, A., Bozem, H., Hoor, P., Krämer,
 M., Müller, S., Riese, M., Rolf, C., Spelten, N., Stiller, G. P., Ungermann, J., and Zahn,
 A.: Long-range transport pathways of tropospheric source gases originating in Asia into
 the northern lower stratosphere during the Asian monsoon season 2012, *Atmos. Chem.*
 885 *Phys.*, 16, 15301–15325, <https://doi.org/10.5194/acp-16-15301-2016>, 2016.
- Wang, Y., Ma, Y.-F., Eskes, H., Inness, A., Flemming, J., and Brasseur, G. P.: Evaluation of
 the CAMS global atmospheric trace gas reanalysis 2003–2016 using aircraft campaign
 observations, *Atmos. Chem. Phys.*, 20, 4493–4521, <https://doi.org/10.5194/acp-20-4493-2020>, 2020.
- 890 Wiegele, A., Glatthor, N., Höpfner, M., Grabowski, U., Kellmann, S., Linden, A., Stiller, G.,
 and Clarmann, T. von: Global distributions of C₂H₆, C₂H₂, HCN, and PAN retrieved from
 MIPAS reduced spectral resolution measurements, *Atmos. Meas. Tech.*, 5, 723–734,
<https://doi.org/10.5194/amt-5-723-2012>, 2012.
- Witze, A.: The Arctic is burning like never before - and that's bad news for climate change,
 895 *Nature*, 585, 336–337, <https://doi.org/10.1038/d41586-020-02568-y>, 2020.
- Xiao, Y., Logan, J. A., Jacob, D. J., Hudman, R. C., Yantosca, R., and Blake, D. R.: Global
 budget of ethane and regional constraints on U.S. sources, *J. Geophys. Res.*, 113, 955,
<https://doi.org/10.1029/2007JD009415>, 2008.

- 900 Xiao, Y., Jacob, D. J., and Turquety, S.: Atmospheric acetylene and its relationship with CO
as an indicator of air mass age, *J. Geophys. Res.*, 112, 955,
<https://doi.org/10.1029/2006JD008268>, 2007.
- Xie, F., Tian, W., and Chipperfield, M. P.: Radiative effect of ozone change on stratosphere-
troposphere exchange, *J. Geophys. Res.*, 113, <https://doi.org/10.1029/2008JD009829>,
2008.
- 905 Yuan, B., Veres, P. R., Warneke, C., Roberts, J. M., Gilman, J. B., Koss, A., Edwards, P. M.,
Graus, M., Kuster, W. C., Li, S.-M., Wild, R. J., Brown, S. S., Dubé, W. P., Lerner, B. M.,
Williams, E. J., Johnson, J. E., Quinn, P. K., Bates, T. S., Lefer, B., Hayes, P. L., Jimenez,
J. L., Weber, R. J., Zamora, R., Ervens, B., Millet, D. B., Rappenglück, B., and Gouw, J.
A. de: Investigation of secondary formation of formic acid: urban environment vs. oil and
910 gas producing region, *Atmos. Chem. Phys.*, 15, 1975–1993, [https://doi.org/10.5194/acp-
15-1975-2015](https://doi.org/10.5194/acp-15-1975-2015), 2015.




Figure 1. Path of the HALO flight on 13 September 2017 during the WISE campaign (large dark brown points) together with GLORIA tangent points (small points with changing colour according to altitude). Measurement times are given in UTC. Coloured framed zones mark areas of special interest as discussed in Sect. 3. Note that only observations performed in the chemistry mode are shown.




Figure 2. Simulated limb emission spectra (with spectral resolution of GLORIA) for a mid-latitude summer standard atmosphere (Remedios et al., 2007) in four microwindows in the spectral region of the C_2H_6 ν_9 band centred at 822 cm^{-1} for a tangent altitude of 8 km . Emissions of individual species contributing to the combined spectrum (all molecules, black line) are shown **(a)**. Retrieved C_2H_6 VMR vertical profile (and a priori profile) of the limb sequence measured at 13 September 2017 at 16:55 UTC combined with absolute and relative errors and the altitude resolution (alt. reso.), determined from the full width at half maximum of the columns of the averaging kernel matrix, together with the observer altitude (z_{obs}). The following individual 1σ errors are shown: spectral noise (red solid line), temperature (green dashed line), line of sight (LOS; blue dotted line), field of view (FOV; dark yellow short dash dotted line) spectroscopic data of disturbing gases (cyan dash dotted line), spectroscopic data of target molecule C_2H_6 (dash dotted magenta line), and total error (black solid line) **(b)**.




Figure 3. Simulated limb emission spectra for four microwindows within the C_2H_2 ν_5 band centred at 730 cm^{-1} for a tangent altitude of 8 km **(a)** and the error budget for a C_2H_2 vertical profile obtained on 13 September 2017 at 16:55 UTC **(b)**. Annotation as per Fig. 2.




Figure 4. Simulated limb emission spectra for three microwindows within the HCOOH ν_6 band centred near 1105 cm^{-1} for a tangent altitude of 8 km **(a)** and the error budget for a HCOOH vertical profile obtained on 13 September 2017 at 16:55 UTC **(b)**. Annotation as per Fig. 2.




Figure 5. Simulated limb emission spectra for two microwindows within the PAN ν_{16} band centred near 792 cm^{-1} for a tangent altitude of 8 km **(a)** and the error budget for a PAN vertical profile obtained on 13 September 2017 at 16:55 UTC **(b)**. Annotation as per Fig. 2.




Figure 6. Horizontal and vertical VMR distributions of (a) O_3 , (b) C_2H_6 , (c) C_2H_2 , (d) $HCOOH$, (e) (PAN) as seen by GLORIA above the North Atlantic region on 13 September 2017. Flight altitude is shown as grey line; dynamical tropopause (2 and 4 potential vorticity units from ECMWF) is plotted as dashed magenta lines. Cyan, blue and green (only PAN) coloured boxes mark regions with enhanced VMR levels (for pollutant species, not for O_3), black boxes comprise a region with low mixing ratios. For all these boxes backward trajectories are calculated (see discussion in Sect. 3).




Figure 7. Horizontal and vertical VMR distributions of GLORIA (repeated from Fig. 6, but temporally smoothed, left column), EMAC_2 (middle column), and CAMS (right column) of (a-c) O₃, (d-f) C₂H₆, (g-h) C₂H₂, (i-k) HCOOH, and (l-n) PAN, as seen on 13 September 2017. The EMAC_2 simulation includes NMVOC emissions enhanced by a factor of 2 as recommended by Monks et al. (2018). No CAMS data for C₂H₂ is available. Annotation as per Fig. 6.




Figure 8. Horizontal and vertical VMR distributions of GLORIA (temporally smoothed, left column), EMAC_2 (enhanced NMVOC emissions) minus GLORIA (middle column), and EMAC (standard NMVOC emissions) minus GLORIA (right column) of (a-c) O_3 , (d-f) C_2H_6 , (g-i) C_2H_2 , (j-l) HCOOH , and (m-o) PAN , as seen on 13 September 2017. Annotation as per Fig. 6.




Figure 9. Backward trajectory calculations performed by CLaMS using ERA-interim wind data. Trajectories start at the GLORIA tangent points and are shown for 20 days (left column), 40 days (middle column), and 60 days (right column) within defined regions: black (a-c), blue (d-f), cyan (g-i), and green (j-l) as displayed in Fig. 6. Trajectory colours denote the potential temperature (which is also a measure of altitude) along the trajectory as indicated in the colour bar. Coloured encircled areas (black, blue, cyan, and green) mark regions where the backward trajectory penetrates the upper edge of the planetary boundary layer (800 hPa). For better clarity, only every tenth trajectory is displayed. Hence, encircled areas may occur where no trajectory is drawn.




Figure 10. Geographical locations of the artificial tracers of air mass origin used in the CLaMS model for the HALO WISE campaign 2017. In some regions, the artificial tracers are defined to separate between continental and maritime areas as well as by different geopotential heights (e.g. Tibetan Plateau). The geographical locations of the artificial tracers of air mass origin used in previous CLaMS simulations can be found in Vogel et al. (2019).




Figure 11. Artificial tracers of air mass origin calculations performed with CLaMS for the GLORIA observation grid showing the horizontal and vertical distribution of fraction of air originating from the boundary layer of different geographical regions as defined in Fig. 10. Results are shown for zones from the North and Central American and Pacific region (**a-f**) and zones for Central Asia (**g**) and South Asia (Sasia) which comprises the subregions INO, IND, NIN, TIB, ECH, BoB, SEA, and Wpool (**g**). The residual part (Res) displayed in (**i**) includes all global regions except the zones shown in (**a-h**) and is of little importance for the GLORIA observations. Annotations as per Fig. 6.