

Atmospheric gaseous hydrochloric and hydrobromic acid in urban Beijing, China: detection, source identification and potential atmospheric impacts

5 Xiaolong Fan^{1#}, Jing Cai^{2#}, Chao Yan², Jian Zhao², Yishuo Guo¹, Chang Li¹, Kaspar R. Dällenbach², Feixue Zheng¹, Zhuohui Lin¹, Biwu Chu^{3,4}, Yonghong Wang², Lubna Dada², Qiaozhi Zha², Wei Du², Jenni Kontkanen², Theo Kurtén⁵, Siddhart Iyer⁶, Joni T Kujansuu^{1,2}, Tuukka Petäjä², Douglas R. Worsnop⁷, Veli-Matti Kerminen², Yongchun Liu¹, Federico Bianchi², Yee Jun Tham^{8,2*}, Lei Yao^{2*}, Markku Kulmala^{1,2,9}

Affiliations:

10 ¹Aerosol and Haze Laboratory, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing 100089, China

² Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki 00560, Finland

15 ³ State Key Joint Laboratory of Environment Simulation and Pollution Control, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

⁴ Center for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China

⁵ Department of Chemistry, University of Helsinki, FI-00014 Helsinki, Finland

⁶ Aerosol Physics Laboratory, Physics Unit, Tampere University, Tampere 33100, Finland

20 ⁷ Aerodyne Research Inc., Billerica, Massachusetts 01821, USA

⁸ School of Marine Sciences, Sun Yat-Sen University, Zhuhai 519082, China

⁹ Joint International Research Laboratory of Atmospheric and Earth System Sciences (JirLATEST), Nanjing University, Nanjing 210023, China.

25 [#] These authors contributed equally.

Correspondence: Lei Yao (lei.yao@helsinki.fi) and Yee Jun Tham (thamyj@mail.sysu.edu.cn)

30

35

40

45 **Abstract**

Gaseous hydrochloric (HCl) and hydrobromic acid (HBr) are vital halogen species that play essential roles in tropospheric physicochemical processes. Yet, the majority of the current studies on these halogen species were conducted in marine or coastal areas. Detection and source identification of HCl and HBr in inland urban areas remain scarce, thus, limiting the full understanding of halogen chemistry and potential atmospheric impacts in the environments with limited influence from the marine sources. Here, both gaseous HCl and HBr were concurrently measured in urban Beijing, China during winter and early spring of 2019. We observed significant HCl and HBr concentrations ranged from a minimum value at 1×10^8 molecules cm^{-3} (4 ppt) and 4×10^7 molecules cm^{-3} (1 ppt) up to 6×10^9 molecules cm^{-3} (222 ppt) and 1×10^9 molecules cm^{-3} (37 ppt), respectively. The HCl and HBr concentrations are enhanced along with the increase of atmospheric temperature, UVB, and levels of gaseous HNO_3 . Based on the air mass analysis and high correlations of HCl and HBr with the burning indicators (HCN and HCNO), the gaseous HCl and HBr are found to be related to anthropogenic burning aerosols. The gas-particle partitioning may also play a dominant role in the elevated daytime HCl and HBr. During the daytime, the reactions of HCl and HBr with OH radicals lead to significant production of atomic Cl and Br, up to 2×10^4 molecules $\text{cm}^{-3} \text{ s}^{-1}$ and 8×10^4 molecules $\text{cm}^{-3} \text{ s}^{-1}$, respectively. The production rate of atomic Br (via $\text{HBr} + \text{OH}$) is 2-3 times higher than that of atomic Cl (via $\text{HCl} + \text{OH}$), highlighting the potential importance of bromine chemistry in the urban area. In polluted days, the production rates of atomic Cl and Br are faster than those on clean days. Furthermore, our observations of elevated HCl and HBr may suggest an important recycling pathway of halogen species in inland megacities, and may provide a plausible explanation for the widespread of halogen chemistry, which could affect the atmospheric oxidation in China.

70

75

80

85

90


95 **1. Introduction**

Tropospheric halogen chemistry plays variety of roles in perturbing the fate of chemical compositions, including ozone (O₃) and volatile organic compounds (VOCs) in the troposphere (Saiz-Lopez and von Glasow, 2012; Simpson et al., 2015; Artiglia et al., 2017). Halogen radicals, in particular the atomic chlorine (Cl·) and bromine (Br·), can deplete the O₃, react rapidly with VOCs with reaction rates up to two orders of magnitude faster than that of the hydroxyl radical (OH) reaction with VOCs and accelerate the depletion of gaseous elemental mercury (Atkinson et al., 2007; Calvert and Lindberg, 2004). Significant halogen-induced O₃ reduction of about 10% of the annually averaged tropospheric ozone column was reported over the tropical marine boundary layer (Saiz-Lopez et al., 2012). However, in polluted coastal regions with high NO_x, the coupling between halogen chemistry and NO_x chemistry contributes to significant enhancement of ozone production of up to 7 ppb (parts per billion by volume) (Li et al., 2020; Sherwen et al., 2017; Sarwar et al., 2014). Besides affecting the ozone chemistry, the oxidation processes of VOCs by halogen radicals can potentially lead to secondary aerosol production. Wang and Hildebrandt Ruiz demonstrated that the chlorine-initiated oxidation of isoprene contributed to the formation of particulate organochloride and the yield of secondary organic aerosol (SOA) ranged from 7 to 36% (Wang and Ruiz, 2017). A recent study also found that the oxidation of alpha-pinene by chlorine atoms yields low-volatility organic compounds, which are essential precursors for secondary particle formation and growth (Wang et al., 2020).

It is known that sea-salt particle is a major source of atomic halogens in the marine environment. The chloride (Cl⁻) and bromide (Br⁻) in the sea-salt particles can be displaced by strong acids (i.e., nitric acid (HNO₃) and sulfuric acid (H₂SO₄)) to release gas-phase hydrogen halides HX (reaction (R1); X = Cl or Br) into the atmosphere (Gard et al., 1998; Thornton et al., 2010). The HX then can react with an OH radical to form a X· via reaction (R2).



On the other hand, the heterogeneous uptake of dinitrogen pentoxide (N₂O₅) onto sea-salt particles can form nitryl halides XNO₂ via reaction (R3) (Finlayson-Pitts et al., 1989; Osthoff et al., 2008; Tham et al., 2014), which is a reservoir of halogen during the nighttime. At sunrise, the XNO₂ undergoes rapid photolysis to liberate highly reactive halogen atom (X·), which subsequently reacts with VOCs to produce HX and peroxy radicals (RO₂; reaction (R4) and (R5)). Besides, the heterogeneous oxidation of Br⁻ by O₃ at the aqueous phase-vapour interface can lead to the formation of a pre-complex intermediate (Br·OOO⁻) which contributes the formation of atmospheric HOBr (Artiglia et al., 2017).



The atmospheric lifetimes of HCl and HBr due to reaction (R2) are approximately 35.6 h and 2.5 h (when OH = 1×10⁷ molecules cm⁻³), respectively, making them a significant daytime recycling source of atomic halogen in the marine atmosphere. Riedel et al. (2012) showed that the reaction of HCl with OH accounts for about 45% of the integrated Cl atom production over the entire day along the Santa Monica Bay of Los Angeles (Riedel et al., 2012). Another ship-borne study reported that the Cl atom production rate peaks at 3×10⁵ molecules cm⁻³ s⁻¹ during the noontime in Southern Coastal California (Crisp et al., 2014). The produced HCl and HBr can also end up in particle phase during the nighttime (Chen et al., 2016; Roberts et al., 2019; Crisp et al., 2014), and further promoting the heterogeneous reaction of N₂O₅ (R3).

The discovery of Thornton et al. (2010) has changed the paradigm of halogen chemistry, where it was

thought to be restricted to the marine environment (Thornton et al., 2010). A significant source of atomic chlorine from the heterogeneous reaction of N_2O_5 onto chloride aerosol (R3) was observed in Boulder, U.S., which is 1400 km from the nearest coastline, indicating that active chlorine chemistry also occurs in the region far from the ocean (Thornton et al., 2010). During the wintertime, the use of road salt could also be a dominant source of atmospheric Cl in the city areas (McNamara et al., 2020). Follow-up studies have confirmed the presence of halogen activation spreading over the continental regions of North America, Canada, Europe and Asia (Mielke et al., 2011; Phillips et al., 2012; Riedel et al., 2013; Tham et al., 2016; Wang et al., 2017; Tham et al., 2018; Liu et al., 2017; Xia et al., 2020; Zhou et al., 2018; McNamara et al., 2020). These findings suggest a crucial role for HCl gas-particle partitioning in sustaining the aerosol chloride concentrations in continental regions for reaction (R3) to take place (Brown and Stutz, 2012).

On the global scale, sea salt sprays were estimated to be the dominant source of halogens such as Cl and Br (Wang et al., 2019a; Keene et al., 1999). Through acid displacement and other heterogeneous processes, 64 Tg a^{-1} and 6.2 Tg a^{-1} gas-phase inorganic Cl and Br from sea salt were emitted to the troposphere, while anthropogenic emissions such as biomass burning, fossil combustion and incineration were supposed to be minor on a global scale (Wang et al., 2019a; Keene et al., 1999). For the emissions of Cl, anthropogenic emissions were quite crucial for both gaseous and particulate Cl in the urban environment and heavily polluted areas. For example, the anthropogenic emissions for gaseous HCl and particulate Cl were 458 and 486 Gg in 2014 in China, of which biomass burning is the largest contributor (Fu et al., 2018). Many recent field studies reported elevated $ClNO_2$ and particulate chloride concentrations in the plumes influenced by biomass burning and coal-fired power plants, suggesting they could be the driving force for the Cl activation process in continental areas (Riedel et al., 2013; Tham et al., 2016; Wang et al., 2017; Liu et al., 2017; Yang et al., 2018). Furthermore, Bannan et al. (2019) showed that the $ClNO_2$ is consistently formed at a landfill site in London, highlighting the potential contribution from landfill emissions of Cl in promoting the reactions (R3) and (R4) (Bannan et al., 2019). Other possible anthropogenic Cl sources include the emissions from industrial, and water and sewage treatment plants (Hara et al., 1989; Graedel and Keene, 1995; Thornton et al., 2010). During the wintertime, the use of road salt could also be a dominant source of atmospheric Cl in the city areas (McNamara et al., 2020).

The atmospheric bromine is much less abundant than chlorine in the stratosphere with the concentrations of around 25 ppt (parts per trillion by volume) compared to 3.7 ppb of chlorine (Bedjanian and Poulet, 2003; Rotermund et al., 2021). HBr is known as a principal bromine sink species for the ozone loss chemistry in the stratosphere showing the average concentration of 1.3 ± 0.39 ppt between 20.0 to 36.5 km altitude (Bedjanian and Poulet, 2003; Nolt et al., 1997; Yang et al., 2005), and also one of the dominant inorganic bromine species in the marine boundary layer, free troposphere and tropical tropopause layer as well (Fernandez et al., 2014; Glasow and Crutzen, 2014; Nolt et al., 1997; Bedjanian and Poulet, 2003). In the urban environment, atmospheric Br was previously known to be strongly affected by traffic emissions since ethylene dibromide ($C_2H_4Br_2$) was used to be as anti-knock compounds to leaded gasoline (Glasow and Crutzen, 2014). Yet, since the phasing out of leaded gasoline, the long-term atmospheric Br exhibited a continuous decreasing trend for 2 to 3 decades in Germany (Lammel et al., 2002), and a similar situation is expected in Beijing as the usage of leaded gasoline was banned from the years around the 2000s in China (Cai et al., 2017).

Despite the advances in the understanding of concentrations and sources of global halogen species, the atmospheric gaseous HCl and HBr in the continental, especially urban environments, are much less studied. Some limited studies focused on the atmospheric HCl, for example, Crisp et al. (2014) summarized that the concentration of HCl is typically less than 1 ppb over the continental regions and McNamara et al. (2020)

measured the concentration of HCl is around 100 ppt from inland sources, while an airborne measurement showed HCl concentrations of around 100 ppt was typically observed over the land area of northeast United States, except near power plant plumes with concentrations over 1 ppb (Crisp et al., 2014;McNamara et al. 2020;Haskins et al., 2018). Furthermore, much less information is available on the presence of HBr in the continental environment. Until very recently, an airborne measurement detected significant levels of gas-phase reactive bromine species in the exhaust of coal-fired power plants (Lee et al., 2018). Therefore, the measurement of gas-phase HCl and HBr in inland urban environments is necessary to fully assess their effects on the tropospheric chemistry, such as gas-particle partitioning effects on the particulate halide concentrations that can undergo rapid activation via reaction (R3). Those would be more important in polluted regions such as the North China Plain, where Beijing is located in and a large amount of chloride were emitted to the atmosphere (Tham et al., 2016;Zhou et al., 2018;Fu et al., 2018).

In this study, we deployed a Chemical Ionization-Atmospheric Pressure interface-long-Time-Of-Flight mass spectrometer (CI-APi-LTOF) to measure the atmospheric gas-phase HCl and HBr from 1 February to 31 March 2019, in urban Beijing, China. To our best knowledge, it is the first time presenting a simultaneous measurement of HCl and HBr with high time-resolution in urban Beijing. Besides, we identify the potential source that contributed to the high levels of gaseous HCl and HBr during wintertime and early springtime. In addition, we estimate the contribution of gaseous HCl and HBr on the production rates of atomic Cl and Br in urban Beijing.

2. Methodology

2.1 Sampling site.

The field measurements were conducted at Beijing University of Chemical Technology (BUCT) monitoring station (39.94° N, 116.30° E), located in an urban area of Beijing, China (Figure 1) where the nearest coastline locates about 150 km away in the southeast. The sampling site is about 130 m north to the Zizhuyuan Road and 550 m west to the West Third Ring Road, which is one of the main roads in Beijing. Besides the effect of traffic, this site is also surrounded by local commercial properties and residential dwellings. Thus, the BUCT sampling site can be regarded as a typical urban site. More information about this sampling site can be found in previous studies (Cai et al., 2020;Kontkanen et al., 2020;Zhou et al., 2020;Chu et al., 2021). The instruments were deployed at the roof of a teaching building, which is approximately 15 m above the ground level.

2.2 CI-APi-LTOF mass spectrometer.


The working principle of CI-APi-LTOF (Aerodyne Research Inc. and ToFwerk AG) has been described elsewhere (Yao et al., 2020;Eisele and Tanner, 1993;Yao et al., 2018), therefore only details relevant to this present work were discussed. A typical mass spectrum during our field measurement was depicted in Figure S1. The dominant reagent ions were nitrate ions (NO_3^- , and $\text{HNO}_3 \cdot \text{NO}_3^-$) and nitrite ions (NO_2^-). Among them, nitrate ions were generated by exposure of sheath flow (pure air with RH ~5%) which carried gaseous HNO_3 . Besides the nitrate ions that acted as dominate reagent ions, nitrite ions were formed from the reaction of a small amount of NO_2 (~1 ppb) in the sheath flow with O_2^- and OH^- which were generated from the exposure of sheath flow (pure air with RH ~5%) to an X-ray source (Hamamatsu L9491) (Figure S5) (Arnold et al., 1995;Skalny et al., 2004). Considering nitrate ions were still the dominant reagent ions (Figure S1), the CI-APi-LTOF was actually operated as a typical nitrate-CI-APi-LTOF.

Ambient air was drawn into the CI-inlet through a 3 quarter-inch stainless steel tube with a flow of ~8 L min^{-1} . A small mixed flow (~0.8 L min^{-1} controlled by a critical orifice with 300 μm diameter) entered the APi-LTOF and be analyzed. The CI-APi-LTOF was operated in the negative V-mode with the mass resolving


225 power of ~ 10000 Th/Th and the mass accuracy better than 5 ppm. Data of CI-APi-LTOF were acquired with
 5 s time resolution, and the recorded data were further analyzed with a MATLAB tofTools package (Junninen
 et al., 2010).

2.3 Detection and quantification of HCl and HBr

230 From Table 1, the gas-phase acidity ($-\Delta G$) of HCl is 1354 kJ mol^{-1} which is larger than that of HNO_3 (1329
 kJ mol^{-1}). Besides, the enthalpy (ΔH) of HNO_3 and Cl^- is $32.8 \text{ kcal mol}^{-1}$, which is higher than that of HCl
 and NO_3^- ($22.9 \text{ kcal mol}^{-1}$) hinting that the reaction of HCl and NO_3^- was unlikely to occur (Figure S4a).
 Additionally, from a previous study, the reaction rate ($< 10^{-12} \text{ molecules cm}^{-3} \text{ s}^{-1}$) between NO_3^- and HCl was
 significantly less than that ($1.4 \times 10^{-9} \text{ molecules cm}^{-3} \text{ s}^{-1}$) of NO_2^- with HCl (Ferguson et al., 1972). Therefore,
 235 the HCl is likely mainly charged by NO_2^- instead of NO_3^- to result in Cl^- formation. The ion-molecule reaction
 between nitrite ions and HCl can be written as follows (Ferguson et al., 1972):



In addition to NO_2^- , the HCl can also react with O_2^- , leading to Cl^- formation via reaction (R7).




Therefore, HCl can be quantified according to:

$$240 \quad [\text{HCl}] = C_{\text{HCl}} \times \frac{(\text{Cl}^-)}{(\text{NO}_2^-) + (\text{O}_2^-)} \quad \text{E (1)}$$

where C_{HCl} (in units of molecules cm^{-3}) is a calibration coefficient of HCl. (Cl^-), (NO_2^-), (O_2^-) represent the
 signals of Cl^- , NO_2^- and O_2^- from CI-APi-LTOF, respectively. Based on ambient data, a very small fraction
 (less than 5%) of Cl^- (or HCl) would react with HNO_3 (or NO_3^-) in the sheath flow to form $\text{Cl}^- \cdot \text{HNO}_3$ (or
 $\text{HCl} \cdot \text{NO}_3^-$). Thus, the signals of $\text{Cl}^- \cdot \text{HNO}_3$ (or $\text{HCl} \cdot \text{NO}_3^-$) were not taken into account for HCl quantification.
 245 The background measurement was carried out by sampling zero air. From Figure S7, the background signals
 were significantly lower than that of ambient air and injected HCl and HBr. The limits of detection (LODs,
 3σ) were 1×10^8 and $1 \times 10^7 \text{ molecules cm}^{-3}$ (i.e., 4 and 0.5 ppt) for HCl and HBr, respectively. Using 4-days
 synchronous gaseous HCl concentrations measured by a Monitor for AeRosols and Gases in Ambient air
 (MARGA, Metrohm Inc., Switzerland), an indirect calibration was adopted to quantify the HCl measured by
 250 the CI-APi-LTOF (Section S5 in Supporting Information). The obtained calibration factor C_{HCl} for HCl is 3
 $\pm 0.1 \times 10^{12} \text{ molecules cm}^{-3}$ (Figure S8b) and the uncertainty of $\pm 30\%$ (Section S5) was applied to the reported
 HCl concentrations. Similar to HCl, the same uncertainty was also adopted for HBr mixing ratios. It should
 be noted that our assumptions lead towards a semi-quantitative estimation of HBr concentrations, due to other
 potential uncertainties (e.g., different sensitivities of HCl and HBr) were not taken into account.

255 On the basis of $-\Delta G$ of HBr, HNO_3 , HNO_2 and HO_2 and the enthalpy (ΔH) calculations (Table 1, Figure
 2 and S4), besides the reaction with NO_2^- and O_2^- , similar with HCl, some of HBr could also react with NO_3^-
 to form Br^- via the reaction (R8) (Ferguson et al., 1972).



Hence, the HBr should be quantified according to:

$$260 \quad [\text{HBr}] = C_{\text{HBr}} \times \frac{(\text{Br}^-)}{(\text{NO}_2^-) + (\text{O}_2^-) + (\text{NO}_3^-)} \quad \text{E (2)}$$

where C_{HBr} (in units of molecules cm^{-3}) is a calibration coefficient of HBr. (Br^-), (NO_2^-), (O_2^-), (NO_3^-)
 represent the signals of Br^- , NO_2^- , O_2^- and NO_3^- from CI-APi-LTOF, respectively. However, as direct
 calibration for HBr was not available, the calibration coefficient of HCl (C_{HCl}) was utilized to semi-quantify
 HBr based on the following equation:

265

$$[HBr] = C_{HCl} \times \frac{(Br^-)}{(NO_2^-) + (O_2^-)} \quad E (3)$$

270

Since the enthalpies (ΔH) of $HBr \cdot NO_3^-$ formed by HBr with NO_3^- (27.3 kcal mol⁻¹) and Br^- with HNO_3 (27.9 kcal mol⁻¹) were very close to each other (Figure S4b), it was difficult to quantify the specific contribution to Br^- from the reaction of HBr with NO_3^- . Also, the ratios of $Br^- \cdot HNO_3$ (or $HBr \cdot NO_3^-$) to Br^- were less than 4%. Therefore, in the equation 3, the reaction pathway of HBr with NO_3^- was not considered. The presented HBr concentrations should be treated as semi-quantitative ones.


275

To confirm these ion-molecule reactions, high concentrations (undetermined) of gaseous HCl and HBr were mixed with zero air generated from a zero-air generator (Aadco 737), and then measured by CI-API-LTOF (Section S4). After the injection of HCl and HBr, the signals of Cl^- , Br^- , $Cl^- \cdot HNO_3$ (or $HCl \cdot NO_3^-$) and $Br^- \cdot HNO_3$ (or $HBr \cdot NO_3^-$) started to increase (Figure S7), confirming that the HCl and HBr can be detected as Cl^- , Br^- , $Cl^- \cdot HNO_3$ and $Br^- \cdot HNO_3$ by CI-API-LTOF.

2.4 Other auxiliary measurements.

Gaseous HCN and HCNO also can be detected by O_2^- through the ion-molecule reactions as follows:

280



285

The $-\Delta G$ of HCN and HCNO are 1433 kJ mol⁻¹ and 1415 kJ mol⁻¹, respectively, which are higher than that of NO_2^- (1393 kJ mol⁻¹) (Table 1), and lower than that of O_2^- (1450 kJ mol⁻¹). Therefore, HCN and HCNO are able to be charged by O_2^- (but not NO_2^-) via deprotonation reaction to lead to CN^- and CNO^- formation. In this study, direct calibrations for HCN and HCNO were not available. Instead, the normalized signals of CN^- and CNO^- by O_2^- were tentatively utilized to indicate the abundance and trend of HCN and HCNO.

290

The meteorological parameters, including temperature and UVB intensities, were recorded by a weather station (Vaisala Inc., Finland). NO_2 was measured with a THERMO 42i NO- NO_2 - NO_x Analyzer (Thermal Environment Instruments Inc. USA). The mass concentrations of particulate chlorine and black carbon (BC) in $PM_{2.5}$ were measured by a time-of-flight aerosol chemical speciation monitor (ToF-ACSM, Aerodyne Research Inc., USA) and an aethalometer (AE33, Magee Inc., USA), respectively (Section S1 in Supporting Information).

295

Meanwhile, we applied 24h air mass back trajectory and Potential Source Contribution Function (PSCF) analyses to help to elucidate the potential source regions (i.e., air masses) of high levels of HCl and HBr. The detailed descriptions of PSCF and air mass trajectory analysis were described in the SI (Section S6) and previous literature (Wang et al., (2014, 2019b)). It is noted that the lifetime of gaseous HCl and HBr could be shorter than the length of the air mass trajectories. These analyses mainly aimed to point out the source regions of pollutant air masses that brought high levels of Cl and Br rather than the real-time origins of air parcels.

3. Results and discussions

300

3.1 HCl and HBr measurement.

305

Figure 3 shows the time series of gaseous HCl and HBr, temperature (T), and ultraviolet radiation b (UVB, 280-315 nm) intensities for the entire measurement period in winter and early spring of 2019 (February to April). High concentrations of HCl and HBr were observed for the whole measurement period with a clear diurnal variation (Figure 3g). The mean concentrations of HCl and HBr are 1×10^9 molecules cm⁻³ (37 ppt) and 2×10^8 molecules cm⁻³ (7 ppt), respectively. The maximum concentrations reach up to 6×10^9 molecules cm⁻³ (222 ppt) for HCl, and 1×10^9 molecules cm⁻³ (37 ppt) for HBr during the daytime. The concentrations

of HCl and HBr showed a similar change in atmospheric temperature and UVB. For the first period of measurement (from 1 to 15 February), HCl and HBr concentrations are lower when the atmospheric temperature is close to 0°C and the UVB intensities are relatively low. Yet, for the later period of March, the HCl and HBr concentrations begin to increase along with the rising of temperature and UVB. In late March, even with higher temperature, due to the less abundant of HNO₃ and particulate chloride, the HCl and HBr concentrations remain at a relatively low level (Figure 3).

The diurnal cycles of HCl and HBr are depicted in Figure 4a and 4b, respectively. The HCl concentrations are typically higher than HBr by approximately an order of magnitude; nevertheless, the diel patterns showed by these two species are quite similar to each other. It is noticed that both HCl and HBr began to increase after sunrise, and relatively high concentration was observed during the daytime (8:00 to 17:00). From Figure 4d, it also can be found that elevated HCl is associated with high temperature and [NO₂]*[OH] value during the daytime. Considering the reaction of NO₂ with OH radical is one of the dominant formation pathways of gaseous HNO₃ during the daytime (Stavrakou et al., 2013), it implies that strong photochemical reactions and the following potential elevated HNO₃ could intensify the HCl releases from particulate chloride in the daytime from 08:00 to 17:00. The OH radical concentrations were calculated using J_{O1D} (Section S8). This phenomenon is consistent with our observation results above where the increase of temperature and UVB could reinforce the formation of chemicals (e.g., HNO₃) that promote the gas-particle partitioning or directly increase gas-phase formation rate of HCl and HBr (Crisp et al., 2014; Riedel et al., 2012), thus further enhancing the HCl and HBr (Figure 3). Although there is no direct measurement of particulate bromide (Br), considering the similarity in diurnal patterns and good correlation ($r = 0.70$) between HBr and HCl (Figure 4c), and HBr tracking well with the temperature and [NO₂]*[OH] (see Figure 3), it is rational to suppose HBr also predominantly derived from gas-particle partitioning process. The contribution by the reaction of bromine atoms with hydrocarbons to form HBr is likely not the dominant pathway as bromine atom is less reactive to hydrocarbons compared to the chlorine atom, and most often reacts with ozone (Simpson et al., 2015).

Our observation of daily averaged mass concentrations of particulate chloride (Cl(p)) in PM_{2.5} showed a similar trend with daily averaged mixing ratios of gaseous HCl (Figure 5a). The difference from the ratios of HCl(g) to Cl(p) in February and March is likely due to the higher temperature in March (Figure 3 and 5a). In contrast, the diurnal variations of HCl and particulate Cl showed the opposite trend at daytime from 08:00 to 17:00 (Figure 5b). The mole ratios of HCl(g) to Cl(p) ranged from <0.1 at nighttime and early morning to >0.3 in the afternoon (Figure 5b). The enhancement of HCl(g)/Cl(p) during the noontime is owing to the large increase of gaseous HCl. It also suggested that the higher temperature and stronger photochemical reactions during the daytime would strongly influence HCl releases from particulate chloride in Beijing, which will be further discussed in the following discussions. During the period between the late afternoon and midnight, the increase of Cl(p) and HCl(g) could be explained by the higher nighttime emissions of residential combustions such as wood and coal burnings in Beijing (Hu et al., 2017; Sun et al., 2016) and high abundance of gaseous HNO₃ are attributed to efficient nocturnal N₂O₅ chemistry (Tham et al., 2018).

These observations showed that there is an abundance of gaseous HCl and HBr in the polluted urban environment. To our best of knowledge, this is the first concurrent observation of gaseous HCl and HBr in a polluted inland urban atmosphere. Although it is well known that the HCl is abundant in the polluted coastal and inland regions, previous studies show that the typical HCl mixing ratios over the continental urban areas are less than 1 ppb (Crisp et al., 2014; Faxon and Allen, 2013; Le Breton et al., 2018; McNamara et al. 2020), which are similar to our observations at Beijing. In contrast, the presence of gaseous HBr in the urban regions is unknown prior to our observation. The significant concentration of HBr in the urban atmosphere of Beijing

is even comparable to the simulated concentrations in the marine environment, where concentration up to 2 ppt was reported (Fernandez et al., 2014). These elevated HCl and HBr in the urban of Beijing may point to the existence of Cl and Br sources in this region.

3.2 Source identification.

355 The natural sources of atmospheric Cl and Br include sea salt spray, wildfires and volcano emissions, while the anthropogenic emissions include coal combustions, traffic emissions as well as other industries such as pesticides, battery industry and waste incinerations (Simpson et al., 2015). Comparing with the sources of particulate Cl and Br that are widely studied and identified in previous literature, the origins of gaseous HCl and HBr are much less studied, due to their much shorter lifetime in the troposphere (Simpson et al., 2015).
360

According to air mass analysis (24h back-trajectory) for HCl and HBr during February and March (Figure 6a and b), the potential source regions of the selected periods with high-level concentrations of HCl (above 75% percentile) were located in the south of North China Plain, such as the south of Hebei province where heavy residential coal, biomass burning and industries emissions occurred (Fu et al., 2018). Those figures further suggested that the high concentrations of HCl seemed not to be strongly affected by marine regions during our sampling period. Instead, the good correlation ($r = 0.67$) between hourly particulate Cl and BC together with the similar trend between particulate Cl and HCl suggested that HCl is likely to have the same original sources with particulate Cl and black carbon (BC) in PM_{2.5} rather than marine sources (Figure 5a and Figure S10a). Hydrocyanic acid (HCN) and isocyanic acid (HCNO), which were typically regarded as tracers for burning emissions, especially biomass burning process (Vigouroux et al., 2012; Adachi et al., 2019; Leslie et al., 2019; Wren et al., 2018; Priestley et al., 2018). Although a recent study showed that HCNO came from both primary emissions and secondary formation in the scale of North China Plain (NCP) during the daytime (Wang et al., 2020), the high correlations between HCN and HCNO (daytime, 08:00-17:00, $r = 0.94$ and nighttime, 18:00-07:00, $r = 0.96$) indicated that in urban Beijing, HCN and HCNO are mainly from primary emission (Figure 7c and 7d) and can be regarded as the tracers of combustion emissions. Thus, high correlations of measured gaseous HCl with HCN ($r = 0.83$) and HCNO ($r = 0.90$) further suggested that the HCl during our sampling period was more likely coming from combustion origins rather than marine source in the urban Beijing (Figure 7a and b). Since gaseous HCl could be affected by both emissions and gas/particle partitioning (shown in Figure 4d), we compared the daily concentrations of gaseous HCl and particulate Cl to minimize the influence of temperature and partitioning. The daily averaged HCl concentration had a high correlation with daily averaged particulate Cl ($r = 0.84$ and 0.70 for winter and early spring periods, respectively) and BC concentration ($r = 0.82$), which is consistent with previous studies that particulate Cl, coal combustion organic aerosol (CCOA) and BC were highly correlated and likely to be from the same source in winter and early spring of Beijing (Hu et al., 2017; Hu et al., 2016).
375
380

385 Similar to HCl, the potential source regions for high Br concentrations were also located in the inland, demonstrating marine sources might not be the dominant source for gaseous HBr in winter of Beijing (Figure 6b). The ratio of particulate Br/Na from previous literature in Beijing was 0.04 (He et al., 2001), which was much higher than those from seawater (0.018) and crustal dust (0.0006 to 0.0008) but much closer to those of biomass burning aerosols (0.01 to 0.06) (Sander et al., 2003). As discussed before, the good correlation ($r = 0.70$) between gaseous HCl and HBr also implied that their similar origins. In our study, moderate correlation coefficients were also observed between gaseous HBr and combustion tracers such as HCN, HCNO (0.63 and 0.62, respectively) and daily BC ($r = 0.60$) (Figure 7a, 7b and S10b). Multiple gaseous organic and inorganic Br compounds such as CH₃Br, Br₂, BrNO₂, BrCl, CH₃Br and CH₂Br₂ were also observed in different combustion processes such as biomass burning, coal combustions and waste
390

395 incineration in previous studies, further supporting the possibilities of combustion origins of the gaseous HBr
in this study (Lee et al., 2018; Keene et al., 1999; Manö and Andreae, 1994). A recent airborne observation
conducted in the U.S. found that high levels of reactive inorganic Br species in the plume from a coal power
plant, likely due to the application of calcium bromide as additives in coal fuel (Lee et al., 2018). Together
all these, in urban Beijing, the measured HBr was more likely coming from combustion sources such as
400 biomass burning and coal combustion in the south of Beijing rather than marine sources. It is also interesting
to note that in a previous marine study conducted at Oahu, Hawaii, gaseous Br was found to be 4 to 10 times
higher than particulate Br (Moyers and Duce, 1972). On the other hand, from a previous observation
conducted in urban Beijing, high levels of both gaseous (7 ng m^{-3}) and particulate (in total suspended particles
(TSP), 18 ng m^{-3}) bromine were measured by offline sampling-organic solvent extraction and Instrumental
405 Neutron Activation Analysis (INAA) method (Tian et al., 2005). Considering the high concentration and
reactivity of Br, gaseous Br from anthropogenic sources may play a more critical role in the urban atmosphere.

3.3 Halogen-atom productions.

To investigate the potential atmospheric implications of HCl and HBr on atmospheric oxidation capacity,
we calculated the production rate of atomic Cl ($P_{Cl\cdot}$) and Br ($P_{Br\cdot}$) via the reactions of HCl and HBr with
410 OH radicals. Figure 8 shows the time series of $P_{Cl\cdot}$, $P_{Br\cdot}$, and the estimated diel concentration of OH calculated
from photolysis rate (J_{OH}) (Section S8). Note that the estimated peak concentrations of OH radicals varied
between $\sim 3 \times 10^5$ to $\sim 4 \times 10^6$ molecules cm^{-3} during noontime. The reaction of HCl with OH radicals lead to a
mean Cl atom production rate of 3×10^3 molecules $\text{cm}^{-3} \text{ s}^{-1}$ during daytime from 08:00 to 17:00 (Figure 8c).
These rates fall within the range of Cl atom production rates ($\sim 10^3$ to 10^6 molecules $\text{cm}^{-3} \text{ s}^{-1}$) reported in
415 polluted environments (Crisp et al., 2014; Hoffmann et al., 2018; McNamara et al 2020). For the reaction of
HBr with OH, it is estimated to produce a daytime mean of 8×10^3 molecules $\text{cm}^{-3} \text{ s}^{-1}$ of Br atom (Figure 8c).
This result shows that in addition to the Cl atom, Br atom could also be present in urban Beijing and may act
as important as the Cl atom in term of reaction with OH, since the $P_{Br\cdot}$ is about 2-3 times faster than the
 $P_{Cl\cdot}$ (Figure 8c). The average HCl and HBr concentrations were observed to be higher during the polluted
420 days (daily mean $\text{PM}_{2.5} \geq 75 \text{ } \mu\text{g m}^{-3}$), which is about 2-3 times higher than the clean days (daily mean $\text{PM}_{2.5}$
 $< 75 \text{ } \mu\text{g m}^{-3}$), as shown in Figure 8b. Consequently, the radical production rate also showed a difference
between clean and polluted days (Figure 8d). From mean diurnal pattern, the values of $P_{Cl\cdot}$ (up to 8×10^3
molecules $\text{cm}^{-3} \text{ s}^{-1}$) and $P_{Br\cdot}$ (up to 2×10^4 molecules $\text{cm}^{-3} \text{ s}^{-1}$) in polluted days were both higher than those of
clean days by up to 2 times (Figure 8d). This hints that the roles of HCl and HBr may be more significant in
425 polluted environments. Recent studies in several polluted sites of China suggested that the photolysis of
 ClNO_2 and Cl_2 are the dominant daytime Cl atom sources leading to Cl atom production rate up to 8×10^6
molecules $\text{cm}^{-3} \text{ s}^{-1}$ (Tham et al., 2016; Liu et al., 2017; Xia et al., 2020), while our observation of Cl atom
production rate from $\text{HCl} + \text{OH}$ could reach up to 2×10^4 molecules $\text{cm}^{-3} \text{ s}^{-1}$ in the daytime. Despite the lower
production rate, the reaction of HCl with OH may also act as important recycling of Cl atom, which ultimately
430 enhanced the atmospheric oxidation capacity (Riedel et al., 2012). In analogous to the chlorine chemistry, the
reaction of HBr with OH could be a significant source of Br atom in the daytime although rapid photolysis
of Br_2 and BrNO_2 is believed to be the major Br atom source in a polluted urban environment as ubiquitous
bromine species (e.g., Br_2 , BrCl and BrNO_2) have been previously observed in residential coal burning and
coal-fired power plant plumes (Lee et al., 2018; Peng et al., 2021).

435 4. Conclusions

In conclusion, we present the first concurrent measurement of both gaseous HCl and HBr in urban Beijing,
a megacity with strong anthropogenic emissions in the North China Plain. Our observation surprisingly shows

440 significant concentrations of HBr in urban Beijing, together with the elevated levels of HCl, throughout the
winter and spring during our sampling period. Gaseous HCl and HBr are most likely originated from
anthropogenic emissions such as burning activities (e.g., biomass burning and fossil fuel combustion) in the
inland region rather than marine sources. Besides, the gas-particle partitioning may play a crucial role in
contributing to elevated levels of HCl and HBr in urban Beijing. In polluted days, the concentrations of HCl
and HBr are higher than those on clean days. These abundant HCl and HBr in the polluted urban troposphere
may further influence the photochemistry of the atmosphere through the following two aspects: (1) direct
445 contributions to the production of highly reactive halogen atom (e.g., Cl· and Br·), which can rapidly oxidize
VOCs (reaction (R5)); (2) replenishing the halide ion (Cl⁻ and Br⁻) in the aerosols for supporting the nocturnal
heterogeneous production of ClNO₂ and BrNO₂, major sources of highly reactive halogen atom at sunrise
(reaction (R3) and (R4)). Our observation of elevated HCl and HBr may indicate an important recycling
pathway of Cl and Br species, and may provide a plausible explanation to the recent observations of
450 widespread halogen activation in polluted areas of China (e.g. Tham et al., 2016; Zhou et al., 2018; Xia et al.,
2020; Peng et al., 2020), which could have a significant influence on the atmospheric oxidation capacity and
secondary aerosol formation. The atomic Cl and Br in polluted days might contribute higher to oxidation
capacity than those on clean days. Furthermore, the additional insight on the HBr levels at Beijing shows that
the bromine chemistry, a previously neglected chemistry, may be important in inland megacities of China.
455 Our results also suggest that understanding of gaseous HCl and HBr would be of much importance to the
photochemistry studies as well as air quality improvement in urban areas of China.

Author Contributions

LY and YJT designed the research. XF, LY, YJT, JC, CY, YG, CL, KRD, FZ, ZL, BC, YM, LD, WD, JK, JTK,
460 JZ, QZ, TK, SI, TP, DRW, VMK, YL, FB and MK carried out the observation, analyzed the data and
interpreted the results. SI and TK provided quantum calculation results. XF, LY, YJT, and JC prepared the
manuscript with contributions from all co-authors.

Declaration of competing interest

465 The authors declare that they have no known competing financial interests.

Data and materials availability:

All data related to this study can be obtained from the corresponding authors (Lei Yao and Yee Jun Tham)
via email.
470

Acknowledgement

The work is supported by ACCC Flagship funded by the Academy of Finland grant number 337549, Academy
of Finland (Center of Excellence in Atmospheric Sciences, project no. 307331, and PROF13 funding, 311932),
the European Research Council via ATM-GTP (742206), CHAPAs (grant no. 850614) and the EMME-CARE
475 project which has received funding from the European Union's Horizon 2020 Research and Innovation.

480

References:

- 485 Adachi, K., Sedlacek, A. J., Kleinman, L., Springston, S. R., and Buseck, P. R.: Spherical tarball particles form through rapid chemical and physical changes of organic matter in biomass-burning smoke, *Proceedings of the National Academy of Sciences*, 116, 201900129, 2019.
- Arnold, S., Morris, R., Viggiano, A., and Jayne, J.: Ion chemistry relevant for chemical ionization detection of SO₃, *J. Geophys. Res-Atmos.*, 100, 14141-14146, 10.1029/95JD01004, 1995.
- 490 Artiglia, L., Edebeli, J., Orlando, F., Chen, S., Lee, M.-T., Corral Arroyo, P., Gilgen, A., Bartels-Rausch, T., Kleibert, A., Vazdar, M., Carignano, M. A., Francisco, J. S., Shepson, P. B., Gladich, I., and Ammann, M.: A surface-stabilized ozonide triggers bromide oxidation at the aqueous solution-vapour interface, *Nat. Commun.*, 8, 700, 10.1038/s41467-017-00823-x, 2017
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume III ‐ gas phase reactions of inorganic halogens, *Atmos. Chem. Phys.*, 7, 981-1191, 10.5194/acp-7-981-2007, 2007.
- 495 Bannan, T. J., Khan, M. A. H., Le Breton, M., Priestley, M., Worrall, S. D., Bacak, A., Marsden, N. A., Lowe, D., Pitt, J., Allen, G., Topping, D., Coe, H., McFiggans, G., Shallcross, D. E., and Percival, C. J.: A Large Source of Atomic Chlorine From ClNO₂ Photolysis at a U.K. Landfill Site, *Geophysical Research Letters*, 46, 8508-8516, 10.1029/2019GL083764, 2019.
- 500 Bedjanian, Y., and Poulet, G.: Kinetics of halogen oxide radicals in the stratosphere, *Chem. Rev.*, 103, 4639, 2003.
- Brown, S. S., and Stutz, J.: Nighttime radical observations and chemistry, *Chemical Society Reviews*, 41, 6405-6447, 10.1039/C2CS35181A, 2012.
- 505 Cai, J., Wang, J., Zhang, Y., Tian, H., Zhu, C., Gross, D. S., Hu, M., Hao, J., He, K., and Wang, S.: Source apportionment of Pb-containing particles in Beijing during January 2013, *Environmental Pollution*, 2017.
- Cai, J., Chu, B., Yao, L., Yan, C., Heikkinen, L. M., Zheng, F., Li, C., Fan, X., Zhang, S., Yang, D., Wang, Y., 510 Kokkonen, T. V., Chan, T., Zhou, Y., Dada, L., Liu, Y., He, H., Paasonen, P., Kujansuu, J. T., Petäjä, T., Mohr, C., Kangasluoma, J., Bianchi, F., Sun, Y., Croteau, P. L., Worsnop, D. R., Kerminen, V.-M., Du, W., Kulmala, M., and Daellenbach, K. R.: Size-segregated particle number and mass concentrations from different emission sources in urban Beijing, *Atmos. Chem. Phys.*, 20, 12721-12740, 10.5194/acp-20-12721-2020, 2020.
- 515 Calvert, J and Lindberg, S.: The potential influence of iodine-containing compounds on the chemistry of the troposphere in the polar spring. II. Mercury depletion. *Atmospheric Environment* 38(30), 5105-5116, 2004.
- Chen, D., Huey, L. G., Tanner, D. J., Salawitch, R. J., Anderson, D. C., Wales, P. A., Pan, L. L., Atlas, E. L., Hornbrook, R. S., Apel, E. C., Blake, N. J., Campos, T. L., Donets, V., Flocke, F. M., Hall, S. R., Hanisco, T. F., Hills, A. J., Honomichl, S. B., Jensen, J. B., Kaser, L., Montzka, D. D., Nicely, J. M., Reeves, J. M., Riemer, D. D., Schauffler, S. M., Ullmann, K., Weinheimer, A. J., and Wolfe, G. M.: Airborne measurements of BrO and the sum of HOBr and Br₂ over the Tropical West Pacific from 1 to 15 km during the CONvective TRansport of Active Species in the Tropics (CONTRAST) experiment, *J. Geophys. Res-Atmos.*, 121, 5605-5612, 578, <https://doi.org/10.1002/2016JD025561>, 2016.
- 520 525 Chu, B. W., Dada, L., Liu, Y. C., Yao, L., Wang, Y. H., Du, W., Cai, J., Dallenbach, K. R., Chen, X. M., Simonen, P., Zhou, Y., Deng, C. J., Fu, Y. Y., Yin, R. J., Li, H. Y., He, X. C., Feng, Z. M., Yan, C., Kangasluoma, J., Bianchi, F., Jiang, J. K., Kujansuu, J., Kerminen, V. M., Petaja, T., He, H., and Kulmala,

- M.: Particle growth with photochemical age from new particle formation to haze in the winter of Beijing, China, *Sci. Total Environ.*, 753, 2021.
- 530 Crisp, T., Lerner, B., Williams, E., Quinn, P., Bates, T., and Bertram, T.: Observations of gas-phase hydrochloric acid in the polluted marine boundary layer, *J. Geophys. Res-Atmos.*, 119, 6897-6915, 10.1002/2013JD020992, 2014.
- Eisele, F. L., and Tanner, D. J.: Measurement of the Gas-Phase Concentration of H₂SO₄ and Methane Sulfonic-Acid and Estimates of H₂SO₄ Production and Loss in the Atmosphere, *J. Geophys. Res-Atmos.*, 535 98, 9001-9010, 1993.
- Faxon, C. B., and Allen, D. T.: Chlorine chemistry in urban atmospheres: a review, *Environmental Chemistry*, 10, 221, 2013.
- Ferguson, E. E., Dunkin, D. B., and Fehsenfeld, F. C.: Reactions of NO₂⁻ and NO₃⁻ with HCl and HBr, *Journal of Chemical Physics*, 57, 1459-1463, 1972.
- 540 Fernandez, R. P., Salawitch, R. J., Kinnison, D. E., Lamarque, J. F., and Saiz-Lopez, A.: Bromine partitioning in the tropical tropopause layer: implications for stratospheric injection, *Atmos. Chem. Phys.*, 14, 17857-17905, 2014.
- Finlayson-Pitts, B. J., Ezell, M. J., and Pitts, J. N.: Formation of chemically active chlorine compounds by reactions of atmospheric NaCl particles with gaseous N₂O₅ and ClONO₂, *Nature*, 337, 241-244, 545 10.1038/337241a0, 1989.
- Fu, X., Wang, T., Wang, S., Zhang, L., Cai, S., Xing, J., and Hao, J.: Anthropogenic Emissions of Hydrogen Chloride and Fine Particulate Chloride in China, *Environ. Sci. Technol.*, 52, 1644-1654, 2018.
- Gard, E. E., Kleeman, M. J., Gross, D. S., Hughes, L. S., Allen, J. O., Morrical, B. D., Ferguson, D. P., Dienes, T., E. Gälli, M., Johnson, R. J., Cass, G. R., and Prather, K. A.: Direct Observation of 550 Heterogeneous Chemistry in the Atmosphere, *Science*, 279, 1184, 10.1126/science.279.5354.1184, 1998.
- Glasow, R. V., and Crutzen, P. J.: Tropospheric Halogen Chemistry, *Treatise on Geochemistry*, 5, 19-69, 2014.
- Graedel, T. E., and Keene, W. C.: Tropospheric budget of reactive chlorine, *Global Biogeochemical Cycles*, 9, 47-77, 1995.
- Hara, H., Kato, T., and Matsushita, H.: The Mechanism of Seasonal Variation in the Size Distributions of 555 Atmospheric Chloride and Nitrate Aerosol in Tokyo, *Bulletin of the Chemical Society of Japan*, 62, 2643-2649, 10.1246/bcsj.62.2643, 1989.
- Haskins, J. D., Jaeglé, L., Shah, V., Lee, B. H., Lopez-Hilfiker, F. D., Campuzano-Jost, P., Schroder, J. C., Day, D. A., Guo, H., Sullivan, A. P., Weber, R., Dibb, J., Campos, T., Jimenez, J. L., Brown, S. S., and Thornton, J. A.: Wintertime Gas-Particle Partitioning and Speciation of Inorganic Chlorine in the Lower 560 Troposphere Over the Northeast United States and Coastal Ocean, *Journal of Geophysical Research: Atmospheres*, 123, 12,897-812,916, 10.1029/2018JD028786, 2018.
- He, K. B., Yang, F. M., Ma, Y. L., Zhang, Q., Yao, X. H., Chan, C. K., Cadle, S., Chan, T., and Mulawa, P.: The characteristics of PM_{2.5} in Beijing, China, *Atmospheric Environment*, 35, 4959-4970, 10.1016/s1352-2310(01)00301-6, 2001.
- 565 Hoffmann, E., Tilgner, A., Wolke, R., and Herrmann, H.: Enhanced Chlorine and Bromine Atom Activation by Hydrolysis of Halogen Nitrates from Marine Aerosols at Polluted Coastal Areas, *Environ. Sci. Technol.*, 53, 10.1021/acs.est.8b05165, 2018.
- Hu, W., Hu, M., Hu, W., Jimenez, J., Yuan, B., Chen, W., Wang, M., Wu, Y., Chen, C., Wang, Z., Peng, J., Zeng, L., and Shao, M.: Chemical composition, sources, and aging process of submicron aerosols in 570 Beijing: Contrast between summer and winter, *Journal of Geophysical Research: Atmospheres*, 121, 1955-1977, 10.1002/2015JD024020, 2016.
- Hu, W., Hu, M., Hu, W., Chen, C., Wu, Y., and Guo, S.: Seasonal variations in high time-resolved chemical

- compositions, sources and evolution of atmospheric submicron aerosols in the megacity Beijing, *Atmospheric Chemistry and Physics*, 9979-10000, 10.5194/acp-17-9979-2017, 2017.
- 575 Junninen, H., Ehn, M., Petaja, T., Luosujarvi, L., Kotiaho, T., Kostianen, R., Rohner, U., Gonin, M., Fuhrer, K., Kulmala, M., and Worsnop, D. R.: A high-resolution mass spectrometer to measure atmospheric ion composition, *Atmos. Meas. Tech.*, 3, 1039-1053, 2010.
- Keene, W. C., Khalil, M. A. K., Erickson, D. J., McCulloch, A., Graedel, T. E., Lobert, J. M., Aucott, M. L., Gong, S. L., Harper, D. B., Kleiman, G., Midgley, P., Moore, R. M., Seuzaret, C., Sturges, W. T.,
580 Benkovitz, C. M., Koropalov, V., Barrie, L. A., and Li, Y. F.: Composite global emissions of reactive chlorine from anthropogenic and natural sources: Reactive Chlorine Emissions Inventory, *Journal of Geophysical Research: Atmospheres*, 104, 8429-8440, 10.1029/1998jd100084, 1999.
- Kontkanen, J., Deng, C., Fu, Y., Dada, L., Zhou, Y., Cai, J., Daellenbach, K. R., Hakala, S., Kokkonen, T. V., Lin, Z., Liu, Y., Wang, Y., Yan, C., Petäjä, T., Jiang, J., Kulmala, M., and Paasonen, P.: Size-resolved
585 particle number emissions in Beijing determined from measured particle size distributions, *Atmos. Chem. Phys.*, 20, 11329-11348, 10.5194/acp-20-11329-2020, 2020.
- Lammel, G., Röhr, A., and Schreiber, H.: Atmospheric lead and bromine in Germany, *Environmental Science and Pollution Research*, 9, 397, 10.1007/BF02987589, 2002.
- Le Breton, M., Hallquist, Å. M., Pathak, R. K., Simpson, D., Wang, Y., Johansson, J., Zheng, J., Yang, Y.,
590 Shang, D., and Wang, H.: Chlorine oxidation of VOCs at a semi-rural site in Beijing: Significant chlorine liberation from ClNO₂ and subsequent gas and particle phase Cl-VOC production, *Atmos. Chem. Phys.*, 18, (17), 13013-13030, 2018.
- Lee, B. H., Lopez-Hilfiker, F. D., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., McDuffie, E. E., Fibiger, D. L., Veres, P. R., Brown, S. S., Campos, T. L., Weinheimer, A. J., Flocke, F. F., Norris, G., O'Mara, K.,
595 Green, J. R., Fiddler, M. N., Bililign, S., Shah, V., Jaegle, L., and Thornton, J. A.: Airborne Observations of Reactive Inorganic Chlorine and Bromine Species in the Exhaust of Coal-Fired Power Plants, *J Geophys Res Atmos*, 123, 11225-11237, 10.1029/2018JD029284, 2018.
- Leslie, M. D., Ridoli, M., Murphy, J. G., and Borduas-Dedekind, N.: Isocyanic acid (HNCO) and its fate in the atmosphere: a review, *Environmental Science: Processes & Impacts*, 21, 2019.
- 600 Li, Q., Badia, A., Wang, T., Sarwar, G., Fu, X., Zhang, L., Zhang, Q., Fung, J., Cuevas, C. A., Wang, S., Zhou, B., and Saiz-Lopez, A.: Potential Effect of Halogens on Atmospheric Oxidation and Air Quality in China, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032058, 10.1029/2019JD032058, 2020.
- Liu, X., Qu, H., Huey, L. G., Wang, Y., Sjostedt, S., Zeng, L., Lu, K., Wu, Y., Hu, M., Shao, M., Zhu, T., and Zhang, Y.: High Levels of Daytime Molecular Chlorine and Nitryl Chloride at a Rural Site on the North
605 China Plain, *Environ. Sci. Technol.*, 51, 9588-9595, 10.1021/acs.est.7b03039, 2017.
- Manö, S., and Andreae, M. O.: Emission of Methyl Bromide from Biomass Burning, *Science*, 263, 1255-1257, 1994.
- McNamara, S. M., Kolesar, K. R., Wang, S., Kirpes, R. M., May, N. W., Gunsch, M. J., Cook, R. D., Fuentes, J. D., Hornbrook, R. S., Apel, E. C., China, S., Laskin, A., and Pratt, K. A.: Observation of Road Salt
610 Aerosol Driving Inland Wintertime Atmospheric Chlorine Chemistry, *ACS Central Science*, 6, 684-694, 10.1021/acscentsci.9b00994, 2020.
- Mielke, L. H., Furgeson, A., and Osthoff, H. D.: Observation of ClNO₂ in a Mid-Continental Urban Environment, *Environ. Sci. Technol.*, 45, 8889-8896, 10.1021/es201955u, 2011.
- Moyers, J. L., and Duce, R. A.: Gaseous and particulate bromine in the marine atmosphere, *Journal of Geophysical Research*, 77, 5330-5338, 10.1029/JC077i027p05330, 1972.
- 615 Nolt, I. G., Ade, P., Alboni, F., Carli, B., Carlotti, M., Cortesi, U., Epifani, M., Griffin, M. J., Hamilton, P. A., Lee, C., Lepri, G., Mencaraglia, F., Murray, A. G., Park, J. H., Park, K., Raspollini, P., Ridolfi, M., and

- Vanek, M. D.: Stratospheric HBr concentration profile obtained from far-infrared emission spectroscopy, *Geophysical Research Letters*, 24, 281-284, 10.1029/97GL00034, 1997.
- 620 Osthoff, H. D., Roberts, J. M., Ravishankara, A. R., Williams, E. J., Lerner, B. M., Sommariva, R., Bates, T. S., Coffman, D., Quinn, P. K., Dibb, J. E., Stark, H., Burkholder, J. B., Talukdar, R. K., Meagher, J., Fehsenfeld, F. C., and Brown, S. S.: High levels of nitryl chloride in the polluted subtropical marine boundary layer, *Nature Geoscience*, 1, 324-328, 10.1038/ngeo177, 2008.
- Peng, X., Wang, W., Xia, M., Chen, H., Ravishankara, A., Li, Q., Alfonso, S.-L., Liu, P., Zhang, F., Zhang, C., Xue, L., Wang, X., George, C., Wang, J., Mu, Y., Chen, J.-M., and Tao, W.: An unexpected large continental source of reactive bromine and chlorine with significant impact on wintertime air quality, 2020.
- 625 Phillips, G. J., Tang, M. J., Thieser, J., Brickwedde, B., Schuster, G., Bohn, B., Lelieveld, J., and Crowley, J. N.: Significant concentrations of nitryl chloride observed in rural continental Europe associated with the influence of sea salt chloride and anthropogenic emissions, *Geophysical Research Letters*, 39, 10.1029/2012GL051912, 2012.
- 630 Priestley, M., Breton, M., Bannan, T., Leather, K., Bacak, A., Reyes Villegas, E., Vocht, F., Shallcross, B., Brazier, T., Khan, A., Allan, J., Shallcross, D., Coe, H., and Percival, C.: Observations of Isocyanate, Amide, Nitrate, and Nitro Compounds From an Anthropogenic Biomass Burning Event Using a ToF-CIMS, *Journal of Geophysical Research: Atmospheres*, 10.1002/2017JD027316, 2018.
- Riedel, T. P., Bertram, T. H., Crisp, T. A., Williams, E. J., Lerner, B. M., Vlasenko, A., Li, S.-M., Gilman, J., de Gouw, J., Bon, D. M., Wagner, N. L., Brown, S. S., and Thornton, J. A.: Nitryl Chloride and Molecular Chlorine in the Coastal Marine Boundary Layer, *Environ. Sci. Technol.*, 46, 10463-10470, 10.1021/es204632r, 2012.
- 640 Riedel, T. P., Wagner, N. L., Dubé, W. P., Middlebrook, A. M., Young, C. J., Öztürk, F., Bahreini, R., VandenBoer, T. C., Wolfe, D. E., Williams, E. J., Roberts, J. M., Brown, S. S., and Thornton, J. A.: Chlorine activation within urban or power plant plumes: Vertically resolved ClNO₂ and Cl₂ measurements from a tall tower in a polluted continental setting, *Journal of Geophysical Research: Atmospheres*, 118, 8702-8715, 10.1002/jgrd.50637, 2013.
- 645 Roberts, T., Dayma, G., and Oppenheimer, C.: Reaction Rates Control High-Temperature Chemistry of Volcanic Gases in Air, *Frontiers in Earth Science*, 7, 154, 10.3389/feart.2019.00154, 2019.
- Rotermund, M. K., Bense, V., Chipperfield, M. P., Engel, A., Grooß, J.-U., Hoor, P., Hüneke, T., Keber, T., Kluge, F., Schreiner, B., Schuck, T., Vogel, B., Zahn, A., and Pfeilsticker, K.: Organic and inorganic bromine measurements around the extratropical tropopause and lowermost stratosphere: Insights into the transport pathways and total bromine, *Atmos. Chem. Phys. Discuss.* [preprint], <https://doi.org/10.5194/acp-2021-202>, in review, 2021.
- 650 Saiz-Lopez, A., Lamarque, J. F., Kinnison, D. E., Tilmes, S., Ordóñez, C., Orlando, J. J., Conley, A. J., Plane, J. M. C., Mahajan, A. S., Sousa Santos, G., Atlas, E. L., Blake, D. R., Sander, S. P., Schauffler, S., Thompson, A. M., and Brasseur, G.: Estimating the climate significance of halogen-driven ozone loss in the tropical marine troposphere, *Atmos. Chem. Phys.*, 12, 3939-3949, 10.5194/acp-12-3939-2012, 2012.
- 655 Saiz-Lopez, A., and von Glasow, R.: Reactive halogen chemistry in the troposphere, *Chem. Soc. Rev.*, 41, 6448-6472, 10.1039/c2cs35208g, 2012.
- Sander, R., Keene, W. C., Pszenny, A. A. P., Arimoto, R., Ayers, G. P., Baboukas, E., Cainey, J. M., Crutzen, P. J., Duce, R. A., Hönninger, G., Huebert, B. J., Maenhaut, W., Mihalopoulos, N., Turekian, V. C., and Van Dingenen, R.: Inorganic bromine in the marine boundary layer: a critical review, *Atmos. Chem. Phys.*, 3, 1301-1336, 10.5194/acp-3-1301-2003, 2003.
- 660

- Sarwar, G., Simon, H., Xing, J., and Mathur, R.: Importance of tropospheric ClNO₂ chemistry across the Northern Hemisphere, *Geophysical Research Letters*, 41, 4050-4058, 10.1002/2014GL059962, 2014.
- 665 Sherwen, T., Evans, M. J., Sommariva, R., Hollis, L. D. J., Ball, S. M., Monks, P. S., Reed, C., Carpenter, L. J., Lee, J. D., Forster, G., Bandy, B., Reeves, C. E., and Bloss, W. J.: Effects of halogens on European air-quality, *Faraday Discussions*, 200, 75-100, 10.1039/C7FD00026J, 2017.
- Simpson, W., Brown, S., Alfonso, S. L., Thornton, J., and Glasow, R.: Tropospheric Halogen Chemistry: Sources, Cycling, and Impacts, *Chemical reviews*, 115, 10.1021/cr5006638, 2015.
- 670 Skalny, J., Mikoviny, T., Matejcik, S., and Mason, N.: An analysis of mass spectrometric study of negative ions extracted from negative corona discharge in air, *International Journal of Mass Spectrometry*, 233, 317-324, 10.1016/j.ijms.2004.01.012, 2004.
- Stavrakou, T., Müller, J. F., Boersma, K. F., van der A, R. J., Kurokawa, J., Ohara, T., and Zhang, Q.: Key chemical NO_x sink uncertainties and how they influence top-down emissions of nitrogen oxides, *Atmospheric Chemistry and Physics*, 13, 9057-9082, 10.5194/acp-13-9057-2013, 2013.
- 675 Sun, Y., Du, W., Fu, P., Wang, Q., Li, J., Ge, X., Zhang, Q., Zhu, C., Ren, L., Xu, W., Zhao, J., Han, T., Worsnop, D. R., and Wang, Z.: Primary and secondary aerosols in Beijing in winter: sources, variations and processes, *Atmospheric Chemistry and Physics*, 16, 8309-8329, 10.5194/acp-16-8309-2016, 2016.
- Tham, Y. J., Yan, C., Xue, L., Zha, Q., Wang, X., and Wang, T.: Presence of high nitryl chloride in Asian coastal environment and its impact on atmospheric photochemistry, *Chinese Science Bulletin*, 59, 356-359, 10.1007/s11434-013-0063-y, 2014.
- 680 Tham, Y. J., Wang, Z., Li, Q., Yun, H., Wang, W., Wang, X., Xue, L., Lu, K., Ma, N., Bohn, B., Li, X., Kecorius, S., Größ, J., Shao, M., Wiedensohler, A., Zhang, Y., and Wang, T.: Significant concentrations of nitryl chloride sustained in the morning: investigations of the causes and impacts on ozone production in a polluted region of northern China, *Atmos. Chem. Phys.*, 16, 14959-14977, 10.5194/acp-16-14959-2016, 2016.
- 685 Tham, Y. J., Wang, Z., Li, Q., Wang, W., Wang, X., Lu, K., Ma, N., Yan, C., Kecorius, S., Wiedensohler, A., Zhang, Y., and Wang, T.: Heterogeneous N₂O₅ uptake coefficient and production yield of ClNO₂ in polluted northern China: roles of aerosol water content and chemical composition, *Atmos. Chem. Phys.*, 18, 13155-13171, 10.5194/acp-18-13155-2018, 2018.
- 690 Thornton, J. A., Kercher, J. P., Riedel, T. P., Wagner, N. L., Cozic, J., Holloway, J. S., Dubé, W. P., Wolfe, G. M., Quinn, P. K., and Middlebrook, A. M.: A large atomic chlorine source inferred from mid-continental reactive nitrogen chemistry, *Nature*, 464, 271-274, 2010.
- Tian, Q., Xu, D., Chai, Z., Lu, Y., and Xiong, Y.: Analysis on the organic bromine in the atmosphere in Beijing, *Journal of Nuclear and Radiochemistry*, 27, 236-238, 252, 2005.
- 695 Vigouroux, C., Stavrakou, T., Whaley, C., Dils, B., Duflot, V., Hermans, C., Kumps, N., Metzger, J.-M., Scolas, F., Vanhaelewyn, G., Müller, J. F., Jones, D., Li, Q., and De Maziere, M.: FTIR time-series of biomass burning products (HCN, C₂H₆, C₂H₂, CH₃OH, and HCOOH) at Reunion Island (21°S, 55°E) and comparisons with model data, *Atmos. Chem. Phys.*, 12, 13733-13786, 10.5194/acpd-12-13733-2012, 2012.
- 700 Wang, D. S., and Ruiz, L. H.: Secondary organic aerosol from chlorine-initiated oxidation of isoprene, *Atmos. Chem. Phys.*, 17, 13491-13508, 10.5194/acp-17-13491-2017, 2017.
- Wang, X., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Zhu, L., Chen, Q., Alexander, B., Sherwen, T., Evans, M. J., Lee, B. H., Haskins, J. D., Lopez-Hilfiker, F. D., Thornton, J. A., Huey, G. L., and Liao, H.: The role of chlorine in global tropospheric chemistry, *Atmos. Chem. Phys.*, 19, 3981-4003, 10.5194/acp-19-3981-2019, 2019a.
- 705 Wang, Y. Q.: MeteoInfo: GIS software for meteorological data visualization and analysis, *Meteorological*

Applications, 21, 2014.

- 710 Wang, Y. Q.: An Open Source Software Suite for Multi-Dimensional Meteorological Data Computation and Visualisation, *Journal of Open Research Software*, 7, 10.5334/jors.267, 2019b.
- Wang, Z., Wang, W., Tham, Y. J., Li, Q., Wang, H., Wen, L., Wang, X., and Wang, T.: Fast heterogeneous N_2O_5 uptake and ClNO_2 production in power plant and industrial plumes observed in the nocturnal residual layer over the North China Plain, *Atmos. Chem. Phys.*, 17, 12361-12378, 10.5194/acp-17-12361-2017, 2017.
- 715 Wang, Z., Yuan, B., Ye, C., Roberts, J., Wisthaler, A., Lin, Y., Li, T., Wu, C., Peng, Y., Wang, C., Wang, S., Yang, S., Wang, B., Qi, J., Wang, C., Song, W., Hu, W., Wang, X., Xu, W., Ma, N., Kuang, Y., Tao, J., Zhang, Z., Su, H., Cheng, Y., Wang, X., and Shao, M.: High Concentrations of Atmospheric Isocyanic Acid (HNCO) Produced from Secondary Sources in China, *Environmental Science & Technology*, 10.1021/acs.est.0c02843, 2020.
- 720 Wren, S. N., John, L., Yuemei, H., Katherine, H., Gang, L., Mihele, C. M., Mittermeier, R. L., Craig, S., Wentzell, J. J. B., and Brook, J. R.: Elucidating real-world vehicle emission factors from mobile measurements over a large metropolitan region: a focus on isocyanic acid, hydrogen cyanide, and black carbon, *Atmos. Chem. Phys.*, 18 (23), 16979-17001, 2018.
- Xia, M., Peng, X., Wang, W., Yu, C., Sun, P., Li, Y., Liu, Y., Xu, Z., Wang, Z., Xu, Z., Nie, W., Ding, A., and 725 Wang, T.: Significant production of ClNO_2 and possible source of Cl_2 from N_2O_5 uptake at a suburban site in eastern China, *Atmos. Chem. Phys.*, 20, 6147-6158, 10.5194/acp-20-6147-2020, 2020.
- Yang, X., Cox, R., Warwick, N., Pyle, J., Carver, G., O'Connor, F., and Savage, N.: Tropospheric bromine chemistry and its impacts on ozone: A model study, *Journal of Geophysical Research*, 110, 10.1029/2005JD006244, 2005.
- 730 Yang, X., Wang, T., Xia, M., Gao, X., Li, Q., Zhang, N., Gao, Y., Lee, S., Wang, X., Xue, L., Yang, L., and Wang, W.: Abundance and origin of fine particulate chloride in continental China, *Science of The Total Environment*, 624, 1041-1051, <https://doi.org/10.1016/j.scitotenv.2017.12.205>, 2018.
- Yao, L., Garmash, O., Bianchi, F., Zheng, J., Yan, C., Kontkanen, J., Junninen, H., Mazon, S. B., Ehn, M., Paasonen, P., Sipila, M., Wang, M., Wang, X., Xiao, S., Chen, H., Lu, Y., Zhang, B., Wang, D., Fu, Q., 735 Geng, F., Li, L., Wang, H., Qiao, L., Yang, X., Chen, J., Kerminen, V. M., Petaja, T., Worsnop, D. R., Kulmala, M., and Wang, L.: Atmospheric new particle formation from sulfuric acid and amines in a Chinese megacity, *Science*, 361, 278-281, 10.1126/science.aao4839, 2018.
- Yao, L., Fan, X., Yan, C., Kurten, T., Daellenbach, K. R., Li, C., Wang, Y., Guo, Y., Dada, L., Rissanen, M. P., Cai, J., Tham, Y. J., Zha, Q., Zhang, S., Du, W., Yu, M., Zheng, F., Zhou, Y., Kontkanen, J., Chan, T., 740 Shen, J., Kujansuu, J. T., Kangasluoma, J., Jiang, J., Wang, L., Worsnop, D. R., Petaja, T., Kerminen, V. M., Liu, Y., Chu, B., He, H., Kulmala, M., and Bianchi, F.: Unprecedented Ambient Sulfur Trioxide (SO_3) Detection: Possible Formation Mechanism and Atmospheric Implications, *Environmental science & technology letters*, 7, 809-818, 10.1021/acs.estlett.0c00615, 2020.
- Zhou, W., Zhao, J., Ouyang, B., Mehra, A., Xu, W., Wang, Y., Bannan, T. J., Worrall, S. D., Priestley, M., Bacak, A., Chen, Q., Xie, C., Wang, Q., Wang, J., Du, W., Zhang, Y., Ge, X., Ye, P., Lee, J. D., Fu, P., 745 Wang, Z., Worsnop, D., Jones, R., Percival, C. J., Coe, H., and Sun, Y.: Production of N_2O_5 and ClNO_2 in summer in urban Beijing, China, *Atmos. Chem. Phys.*, 18, 11581-11597, 10.5194/acp-18-11581-2018, 2018.
- Zhou, Y., Dada, L., Liu, Y., Fu, Y., Kangasluoma, J., Chan, T., Yan, C., Chu, B., Daellenbach, K. R., Bianchi, 750 F., Kokkonen, T. V., Liu, Y., Kujansuu, J., Kerminen, V.-M., Petäjä, T., Wang, L., Jiang, J., and Kulmala, M.: Variation of size-segregated particle number concentrations in wintertime Beijing, *Atmos. Chem. Phys.*, 20, 1201-1216, 10.5194/acp-20-1201-2020, 2020.

Table 1. Gas-phase acidities and deprotonated anion of a few compounds of interest.

Compounds	Formula	$-\Delta G^a$ (kJ mol ⁻¹)	Deprotonated Anion
Hydrobromic acid	HBr	1319	Br ⁻
Nitric acid	HNO ₃	1329	NO ₃ ⁻
hydrochloric acid	HCl	1354	Cl ⁻
Nitrous Acid	HONO	1396	NO ₂ ⁻
Isocyanic Acid	HCNO	1415	CNO ⁻
Hydrocyanic Acid	HCN	1433	CN ⁻
Hydroperoxy radical	HO ₂	1450	O ₂ ⁻
Hypobromous Acid	HOBr	1460	BrO ⁻
Hypochlorous Acid	HOCl	1461	ClO ⁻

^a Gas-phase acidity is defined as $-\Delta G$ for the protonation reaction ($H^+ + A^- \rightarrow HA$). Data are obtained from NIST


755 Chemistry WebBook.

760

765

770

775




780 **Figure 1.** The location of BUCT measurement station. The satellite map was revised from © Google map.

785

790

795




800 **Figure 2.** The calculated enthalpies of HCl·NO₂⁻ formed by HCl with NO₂⁻ and Cl⁻ with HNO₂ and enthalpies
of HBr·NO₂⁻ formed by HBr with NO₂⁻ and Br⁻ with HNO₂ at the DLPNO-CCSD(T)/def2-QZVPP// ωB97X-
D/aug-cc-pVTZ-PP level of theory.

805

810


815

820



825 **Figure 3.** Time profiles of temperature (a), UVB intensities (b), NO₂ concentration (c), OH concentration from calculation (d), [NO₂]*[OH] ($\mu\text{g m}^{-3} * \text{molecules cm}^{-3}$) (e), particulate chloride concentration (Cl(p)) (f) and the mixing ratios of HCl and HBr (g). The data points are in hourly-average interval.

830




835

Figure 4. Diurnal variations of UVB intensities, HCl and HBr concentrations (averaged values \pm one standard deviation) (a and b) and the correlation between HCl and HBr (c). Temperature dependence of gas to particle partitioning ratios of mass concentration of chloride, color-coded by $[\text{NO}_2] \cdot [\text{OH}]$ which was indicated as the abundance of HNO_3 in the daytime (d). In panel c and d, the data points are hourly averaged ones during daytime (8:00-17:00). All snowy and rainy days during the sampling period were excluded.

840

845




850

Figure 5. Time variation of daily averaged concentration of particulate chloride (Cl(p)) measured by ACSM, gaseous HCl (HCl(g)) measured by CI-API-LTOF and mole ratios of HCl(g)/Cl(p) (a) and diurnal variation of HCl(g), Cl(p) and mole ratios of HCl(g)/Cl(p) (b).

855

860

865



870

Figure 6. The results of PSCF analysis for HCl (a) and HBr (b). Darker colors indicate greater potential source distribution. Black stars mark the location of the sampling site.

875

880




Figure 7. The relationship of HCl and HBr concentrations with HCN and HCNO during the daytime (08:00-17:00) (a and b) and the correlations between HCN and HCNO during both daytime (08:00-17:00) (c) and nighttime (18:00-07:00 the next day) (d).

885

890




Figure 8. Time series of calculated production rates of Cl and Br radicals during the observation period (a); diurnal variations of HCl and HBr concentrations in clean and polluted days (b); diurnal variations of production rates of Cl and Br radicals from 08:00 to 17:00, together with calculated OH radical concentrations (c) and production rates of Cl and Br radicals in clean and polluted days (d). The clean and polluted days were classified as daily $\text{PM}_{2.5} < 75 \mu\text{g m}^{-3}$ and $\text{PM}_{2.5} \geq 75 \mu\text{g m}^{-3}$, respectively. The data points are in the hourly-average interval and measured during observation periods from 1 February to 31 March 2019.

895

900