

Measurement report: The effect of aerosol chemical composition on light scattering due to the hygroscopic swelling effect

Rongmin Ren¹, Zhanqing Li², Peng Yan³, Yuying Wang⁴, Hao Wu⁵, Maureen Cribb², Wei Wang¹, Xiao'ai Jin¹, Yanan Li³, and Dongmei Zhang¹

¹State Key Laboratory of Remote Sensing Science, College of Global Change and Earth System Science, Beijing Normal University, Beijing 100875, China

²Department of Atmospheric and Oceanic Science, Earth System Science Interdisciplinary Center,

University of Maryland, College Park, College Park, MD, USA

³CMA Meteorological Observation Center, Centre for Atmosphere Watch and Services, ¹²⁹ Beijing 100081, China ⁴Key Laboratory for Aerosol–Cloud–Precipitation¹²⁸ of China Meteorological Administration, School of Atmospheric Physics, Nanjing University of Information Science and Technology, Nanjing 210044, China ⁵School of Electrical Engineering, Chengdu University of Information Technology, Chengdu 610225, China

Correspondence: Zhanqing Li (zli@atmos.umd.edu)

Received: 4 December 2020 – Discussion started: 18 January 2021 Revised: 18 May 2021 – Accepted: 19 May 2021 – Published:

Abstract. ISSIL iquid ISSIL iquid Issue water in aerosol particles has a significant effect on their optical properties, especially on light scattering, whose dependence on chemical composition is investigated here using measurements made in southern

- ⁵ Beijing in 2019. The effect is measured by the particle light scattering enhancement f(RH), where RH denotes the relative humidity, which is found to be positively and negatively impacted by the proportions of inorganic and organic matter, respectively. Black carbon is also negatively correlated. ¹⁰ The positive impact is more robust when the inorganic mat-
- ter mass fraction was smaller than 40 % (R = 0.93, R: the Pearson's correlation coefficient), becoming weaker as the inorganic matter mass fraction gets larger (R = 0.48). A similar pattern was also found for the negative impact of the s organic matter mass fraction. Nitrate played a more signif-
- icant role in aerosol hygroscopicity than sulfate in Beijing. However, the deliquescence point of ambient aerosols was at about RH = 80% when the ratio of the sulfate mass concentration to the nitrate mass concentration of the aerosol was ²⁰ high (mostly higher than ~ 4). Two schemes to parameterize f(RH) were developed to account for the deliquescent and
- non-deliquescent effects. Using only one f(RH) parameterization scheme to fit all f(RH) processes incurs large errors. A piecewise parameterization scheme is proposed, which can

better describe deliquescence and reduces uncertainties in $_{\rm 25}$ simulating aerosol hygroscopicity.

1 Introduction

Atmospheric aerosols have impacts on visibility, the earth– atmosphere radiation budget, clouds, and precipitation via direct and indirect effects (IPCC, 2013). Both effects are associated with the hygroscopic properties of aerosols and relative humidity (RH) of the atmosphere. The particle light scattering enhancement factor, $f(RH, \lambda)$, is the ratio of the scattering coefficient at an elevated RH level to that under a fixed low RH level (usually RH < 40%) at a certain light wavelength (λ). It has been characterized during international field experiments (Fierz-Schmidhauser et al., 2010a, b; Zieger et al., 2010, 2014) and in particular China (Yan et al., 2009; Zhang et al., 2015; Kuang et al., 2016; L. Liu et al., 2018; C. Zhao et al., 2019; Zhao, 2019; Wu et al., 2020).

Aerosol chemical composition has a strong impact on aerosol hygroscopicity (Fierz-Schmidhauser et al., 2010a, b; Wang et al., 2017, 2018, 2019). Zhang et al. (2015) studied the relationship between the scattering enhancement factor and chemical composition in Lin'an, China, finding that ⁴⁵ nitrate has a stronger effect on aerosol hygroscopicity than sulfate has, which is partially due to the rigid control of sulfur dioxide (SO₂) that reduces the amount of sulfate and increases the content of nitrite (Morgan et al., 2010). Apart from sea salt emissions and gypsum dust emissions during 5 construction containing sulfate, sulfate is mainly formed by

- the oxidation of its gaseous precursor, SO_2 , in the atmosphere. In recent years, SO_2 emissions have been reduced substantially through a series of effective measures taken in China, like controlling the burning of loose coal and desul-
- ¹⁰ furizing industrial equipment (Zhang et al., 2019). Reducing SO₂ in the atmosphere thus directly affects the reduction in the sulfate content of aerosols. The saturated vapor pressure of nitric acid (HNO₃) is higher than that of sulfuric acid (H₂SO₄), so the availability of ammonia (NH₃) is key to the
- ¹⁵ partitioning of HNO₃. HNO₃ is often neutralized by NH₃ after H₂SO₄. Therefore, a reduction in SO₂ means that more NH₃ can be used to neutralize HNO₃, leading to higher nitrate concentrations, such as ammonium nitrate (HN₄NO₃), in aerosols (Monks et al., 2009). Zieger et al. (2014) ana-
- ²⁰ lyzed the correlation between the chemical composition of aerosols and f(RH = 85%, 550 nm) in Melpitz, Germany, and noted negative and positive impacts by organic and black carbon and by inorganic substances, such as ammonium, respectively. However, the correlation between the mass frac-
- ²⁵ tions of individual nitrate (NO₃⁻) and sulfate (SO₄²⁻) ions and f(RH = 85%, 550 nm) were weak. Zieger et al. (2015) also found that the mass fraction of sulfates was strongly correlated with scattering enhancement, while the mass fraction of nitrates had a low correlation in Hyytiälä, Finland.
- ³⁰ Jin et al. (2020) reported that apart from inorganic matter, e.g., SO₄²⁻ and NO₃⁻, organic species also significantly contributed to the aerosol liquid water content. They proposed that in the initial phase of a pollution event, the aerosol liquid water content contributed by organic matter accelerated ³⁵ aqueous-phase reactions, converting gaseous precursors into

secondary aerosols which then absorb more liquid water. Air pollution has been and continues to be a serious prob-

lem in China, especially in megacities like Beijing. A high scattering enhancement factor is one of the most important
factors causing degradation in visibility. Some observational studies of the light hygroscopicity enhancement factor in Beijing have been conducted (Liu et al., 2013; Yang et al., 2015; Zhao et al., 2018; P. Zhao et al., 2019). However, studies exploring the relationship between aerosol chemical com-

- ⁴⁵ position and the particle light scattering enhancement factor are lacking. Moreover, although several optimal expressions of $f(\text{RH}, \lambda)$ for different seasons have been developed (Pan et al., 2009; Yan et al., 2009; Kuang et al., 2016; Wu et al., 2017; Yu et al., 2018; P. Zhao et al., 2019), param-
- et al., 2017; Yu et al., 2018; P. Zhao et al., 2019), param-⁵⁰ eterization of the deliquescent curve of SO₄²⁻ in ambient aerosols has not yet been done for the Beijing–Tianjin–Hebei (BTH) metropolitan region, where aerosol deliquescent phenomena frequently occur (Kuang et al., 2016). It is thus important to develop an optimal parameterization to describe

70

this deliquescent phenomenon to improve model simulations 55 of aerosol hygroscopicity.

In this study, $f(RH, \lambda)$ at three wavelengths for RH ranging from 40% to about 90% was measured by a highresolution humidified nephelometer system deployed in the southern suburban area of Beijing, China. Other aerosol ⁶⁰ chemical and physical properties were also simultaneously measured. Humidograms were classified into two categories, i.e., deliquescent and non-deliquescent, each parameterized separately. The parameterization results of deliquescent processes agreed well with observations. This result is useful for ⁶⁵ improving simulations of $f(RH, \lambda)$ of ambient aerosols during deliquescence in the BTH metropolitan region.

The paper is organized as follows. Section 2 describes the instruments and methods. Section 3 presents and discusses the results of this study, and Sect. 4 provides a summary.

2 Instruments and methods

2.1 Observation site

A comprehensive field experiment was conducted at the climate observatory of the China Meteorological Administration, located in Yizhuang, Beijing, near the southern Fifth 75 Ring Beltway (39.81° N, 116.48° E) surrounded primarily by residential communities and industrial parks (Fig. S1 in the Supplement). Measurements made here can characterize the aerosol chemical and physical properties of a typical suburban area of this megacity in the North China 80 Plain. Equipped with a multitude of instruments measuring, for example, optical, hygroscopic, and chemical properties of aerosols (Z. Li et al., 2019), this study employs only those measurements acquired from 19 September to 4 October 2019. The instruments used in this field experiment in-85 clude a dual-nephelometer system (Aurora 3000, Ecotech), an aerosol chemical speciation monitor (ACSM; Aerodyne Research Inc.), and a seven-wavelength aethalometer (AE33, Magee Scientific). They were all located in a mobility container on the ground. There are two air conditioners inside the 90 container whose temperature was maintained at about 23°C. Sample air $(16.7 \,\mathrm{L\,min^{-1}})$ went through a PM_{2.5} cyclone inlet at about 4 m above the ground, which only allowed particles with an aerodynamic diameter smaller than 2.5 µm to enter, and was then dried by a Nafion dryer (MD-700-36F-3, 95 Perma Pure LLC). The average RH within the sampling line was about 30 %. The sample air was not heated.

2.2 Instruments

A dual-nephelometer system with a high time resolution was used to measure the particle light scattering enhancement ¹⁰⁰ factor of aerosol. After an aerosol sample passed through the Nafion dryer, the dry sample flow (RH < 40 %) was divided into two routes. One sample flow (5 L min⁻¹) went directly into the dry nephelometer. The other sample flow (5 L min⁻¹) passed through an annular concentric humidifying tube (MD-700-6F-3, Perma Pure LLC). The water vapor controlled by the temperature of the liquid water in the outer annulus of the tube passes through a Nafion membrane, humidifying

- $_5$ aerosols in the inner tube (Carrico et al., 1998). The temperature of the liquid water was controlled by adjusting the power of the water baths. The sample flow was then humidified to a given RH and channeled into the wet nephelometer. The scattering coefficients under dry (the mean \pm standard
- ¹⁰ deviation value of RH was $28.75 \pm 5.50 \%$) and wet ambient conditions were measured synchronously by the two nephelometers (Yan et al., 2009). To improve the performance of this system and to decrease the amount of time needed to undergo one aerosol humidifying process, two water baths were
- ¹⁵ used in turn to heat the water circulating in the interior layer of the humidifying tube (Liu and Zhao, 2016).

Since the RH of aerosols inside the nephelometers was constantly changing and real measured data at every moment were needed, the nephelometers operated without Kalman

- ²⁰ filters. Manual "full calibration" and zero check and span check of the two nephelometers were performed at 10:30 (all times in Beijing time, UTC + 8 h)[125] on 19 September 2019. The calibration tolerance of the zero check was $\pm 2 \text{ Mm}^{-1}$, and that of the span check was $\pm 2 \%$ of the span point. Cal-
- ²⁵ ibrations of the two nephelometers in the dry state were consistent (Fig. S2). The truncation and illumination correction of the scattering coefficients has been done following Müller et al. (2011), which was developed specifically for Ecotech nephelometers originating from Anderson and Ogren (1998)
- ³⁰ for TSI nephelometers. For the f(RH) calculations, there is no truncation and illumination correction applied to the scattering coefficients of both dry and humidified types of nephelometer. The comparison of the deviation between corrected and uncorrected f(RH = 85%, 525 nm) is shown in
- ³⁵ Fig. S3. The linear least square regression slop \pm standard deviation is 1.064 \pm 0.002, the intercept \pm standard deviation is -0.082 ± 0.004 , and *R* is 0.999. The fitted line is very close to the line of 1 : 1. The uncorrected *f*(RH = 85 %, 525 nm) is a little lower than the corrected *f*(RH = 85 %, 40 525 nm).

Because the RH levels measured by the probe built into the optical chamber of the wet nephelometer (RH_{chamber}) were imprecise, the RH_{chamber} was corrected in this paper. First, a set of calibrated RH and temperature probes was placed at ⁴⁵ the inlet of the wet nephelometer, and another set was placed at the outlet of the wet nephelometer, obtaining 1 min averages of RH and temperature. We used Vaisala HMP110 probes with accuracies of $\pm 0.2^{\circ}$ C for the 0–40°C temperature range and $\pm 1.5 \%$ RH and $\pm 2.5 \%$ RH for the 0 % RH-⁵⁰ 90 % RH and 90 % RH-100 % RH^{IIS2} ranges, respectively. The temperatures measured by these three probes were different. However, in principle, the dew point temperatures ($T_{dew-point}$) at these three probes at the outlet of the wet neph-⁵⁵ elometer (RH_{outlet} and T_{outlet}) were less affected by the humidifier, RH_{outlet} and T_{outlet} were used to calculate $T_{dew-point}$ at this position using Eq. (1) (Wanielista et al., 1997; James et al., 2015):

$$T_{\text{dew-point}} = \mathrm{RH}_{\mathrm{outlet}}^{\frac{1}{8}} (112 + 0.9T_{\mathrm{outlet}}) + 0.1T_{\mathrm{outlet}} - 112.$$
 (1)

We assume that $T_{\text{dew-point}}$ was approximately the same as that ⁶⁰ in the optical chamber of the wet nephelometer. Based on the temperature in the optical chamber (T_{chamber}) and $T_{\text{dew-point}}$, the actual RH in the optical chamber (RH_{chamber}) can be calculated by rearranging Eq. (1), i.e.,

$$\mathrm{RH}_{\mathrm{chamber}} = \left(\frac{112 - 0.1T_{\mathrm{chamber}} + T_{\mathrm{dew-point}}}{112 + 0.9T_{\mathrm{chamber}}}\right). \tag{2}$$

The dual-nephelometer system with a high time resolution in this study was calibrated with ammonium sulfate, $(NH_4)_2SO_4$, whose deliquescence RH (DRH) was 80% at 298 K (Cheung et al., 2015). Figure S4 shows that the measured phase transition occurs at RH = 80.07%. It illustrates that the RH inside the nephelometer chamber is correct and that the system is functioning properly.

An ACSM measured the mass concentrations of nonrefractory aerosol chemical species, including SO_4^{2-} , NO_3^{-} , ammonium (NH_4^+), chlorine (Chl), and organics (Orgs) in particulate matter with diameters less than 2.5 µm ($PM_{2.5}$). The mass concentration of the equivalent black carbon (eBC) could be retrieved from the measurements of AE33. The Chinese Ministry of Ecology and Environment network and the Beijing Municipal Environmental Monitoring Center ⁸⁰ (http://106.37.208.233:20035/**ISS** and http://www.bjmemc. com.cn/**ISS**) provided mass concentrations of $PM_{2.5}$ measured at the Yizhuang station, about 3 km southeast of the observatory. The LI-COR eddy covariance system (this system includes WindMaster Pro, LI-7500A, and Smart2-00171, LI-COR) measured various meteorological parameters.

2.3 Methods

The particle light scattering enhancement factor, $f(RH, \lambda)$, is defined as

$$f(\mathbf{RH},\lambda) = \frac{\sigma_{\rm sp}(\mathbf{RH},\lambda)}{\sigma_{\rm sp}(\mathbf{RH}_{\rm dry},\lambda)},\tag{3}$$

where $\sigma_{\rm sp}(\rm RH, \lambda)$ represents the scattering coefficient at an elevated RH (usually RH > 40 %), and $\sigma_{\rm sp}(\rm RH_{dry}, \lambda)$ is the scattering coefficient in the dry state (usually RH < 40 %) at wavelength λ . Values of $f(\rm RH, \lambda)$ are generally greater than 1 and increase with increasing RH. In this study, we assume that the aerosol is in the dry state when RH < 40 %. This means that in theory, $f(\rm RH)$ should equal 1 when RH is lower than 40%. However, due to systematic errors and differences in RH measured synchronously by the dry nephelometer and the wet nephelometer, the measured 100 $f(\rm RH < 40\%)$ has small fluctuations and does not equal 1.

Therefore, f(RH > 40%) was normalized as

$$f(\mathrm{RH} > 40\%)_{\mathrm{normalized}} = \left(\frac{f(\mathrm{RH} > 40\%)}{f(\mathrm{RH} < 40\%)_{\mathrm{averaged}}}\right).$$
 (4)

Here, $f(\text{RH} < 40\%)_{\text{averaged}}$ is the corrected coefficient averaged over the whole dataset of RH < 40%.

- ⁵ The absorption coefficient of $PM_{2.5}$ at 520 nm was calculated by the eBC monitor (Han et al., 2015; Zou et al., 2019). To facilitate comparisons, absorption coefficients at 520 nm were transformed into those at 525 nm by assuming that the absorption coefficient is inversely proportional to the wave-
- ¹⁰ length (Bond and Bergstrom, 2006; C. Liu et al., 2018). The quantity $\omega_{0(525 \text{ nm})}$ is the aerosol single-scattering albedo at 525 nm. The dependence of light scattering on wavelength is described by the Ångström exponent ($\alpha_{(\lambda_1 - \lambda_2)}$), an index describing the particle size:

$${}_{45} \alpha_{(\lambda_1 - \lambda_2)} = \frac{\log \sigma_{\rm sp} (\lambda_1) - \log \sigma_{\rm sp} (\lambda_2)}{\log \lambda_2 - \log \lambda_1}.$$
 (5)

Here, $\alpha_{(450-635 \text{ nm})}$ between 450 and 525 nm was calculated using Eq. (5).

The following parameter (F_{org}) denotes the relative amount of organic and inorganic matter:

$$_{20} F_{\rm org} = \frac{C_{\rm c}}{C_{\rm c} + C_{\rm i}},\tag{6}$$

where C_c is the organic matter mass concentration measured by the ACSM, and C_i represents the mass concentration of inorganic salts like (NH₄)₂SO₄, ammonium bisulfate (NH₄HSO₄), and ammonium nitrate (NH₄NO₃).

²⁵ Deliquescence of ambient aerosols was present throughout the study period. To identify this process and describe its magnitude in the 78 % RH–82 % RH range, the hysteresis index η is defined as (Zieger et al., 2010)

$$\eta = 1 - \frac{\gamma_{<78\,\%}}{\gamma_{>82\,\%}}.\tag{7}$$

³⁰ The terms $\gamma_{<78\%}$ and $\gamma_{>82\%}$ are the fit parameters of the f(RH) parametrization scheme,

$$f(\mathbf{RH}) = (1 - \mathbf{RH})^{-\gamma}, \tag{8}$$

at RH < 78 % and RH > 82 %, respectively. The parameter γ is retrieved from Eq. (8) using the whole RH range. It ³⁵ can replace f(RH) in a wider RH range (Doherty et al., 2005; Quinn et al., 2005; Zhang et al., 2015). The theoretical range of η is 0 to 1. The $\gamma_{<78}$ % and $\gamma_{>82}$ % terms, respectively, represent the magnitudes of the scattering enhancement when RH < 78 % and RH > 82 %. If the values ⁴⁰ of $\gamma_{<78}$ % and $\gamma_{>82}$ % are about the same, then η will be close to 0. This suggests that f(RH) increases slowly and continuously when 78 % < RH < 82 %, and no deliquescence is found. However, when the value of $\gamma_{>82}$ % is much higher than $\gamma_{<78}$ %, η approaches 1. This explains why the f(RH)⁴⁵ cycle has a jump at 78 % < RH < 82 %, i.e., very distinct deliquescence occurring in the RH range of 78 % to 82 %. Here, when η is higher than 0.4, deliquescence occurs.

3 Results and discussion

3.1 Overview

Figure 1 depicts the hourly averaged time series of the 50 light-scattering coefficient ($\sigma_{sp,525 \text{ nm}}$), the absorption coefficient ($\sigma_{ap,525 \text{ nm}}$), the single-scattering albedo ($\omega_{0(525 \text{ nm})}$), the scattering Ångström exponent ($\alpha_{(450-635 \text{ nm})}$), and the particle light scattering enhancement factor at RH = 85 %, f(RH = 85%, 525 nm), at the main observatory, and the 55 mass concentration of PM2.5 measured at the Yizhuang station from 19 September to 4 October 2019. During this period, the hourly averaged $\sigma_{sp,525 nm}$ ranged from 3 to 799 Mm^{-1} (Fig. 1a), with a mean \pm standard deviation value of 245 ± 168 Mm⁻¹ (Table 1). The hourly av- 60 eraged $\sigma_{ap,525 \text{ nm}}$ varied from 7 to 135 Mm⁻¹ (Fig. 1b), with a mean \pm standard deviation value of 50 ± 24 Mm⁻¹ (Table 1). Figure 1c and f show that $\omega_{0(525 \text{ nm})}$ increased as the PM_{2.5} concentration increased. The hourly averaged $\omega_{0(525 \text{ nm})}$ during the observation period ranged from 65 0.24 to 0.98, with an overall mean \pm standard deviation value of 0.77 ± 0.15 . The mean \pm standard deviation values of $\omega_{0(525 \text{ nm})}$ during clean (PM_{2.5} \leq 35 µg m⁻³), moderately polluted (35 $\mu g\,m^{-3} < PM_{2.5} \leq 75\,\mu g\,m^{-3}),$ and heavily polluted (PM_{2.5} > 75 μ g m⁻³) periods were 0.65 \pm 0.16, ⁷⁰ 0.83 ± 0.08 , and 0.93 ± 0.04 , respectively (Table 1).

The $\omega_{0(525 \text{ nm})}$ increased as PM_{2.5} pollution increased, indicating that during the contamination process, the proportion of aerosol components with strong scattering properties increased, and the proportion of aerosol components 75 with strong absorbing properties decreased. Higher values of $\omega_{0(525\,\mathrm{nm})}$ usually occurred when the wind was from the northeast at a speed of $1-2 \text{ m s}^{-1}$ and when relatively stronger winds were from the southeast (Fig. 2a), which was frequently accompanied by a high proportion of inorganic 80 matter (Fig. 2e) and low proportions of eBC (Fig. 2d) and organic matter (Fig. 2f). Figure 1d shows that the range of $\alpha_{(450-635\,\text{nm})}$ was narrow at most times during the observation period. Relatively larger particles with lower values of $\alpha_{(450-635 \text{ nm})}$ generally occurred when weak winds were ⁸⁵ from the east and southeast (Fig. 2b).

During the observation period, f(RH = 85%, 525 nm)ranged from 1.15 to 1.86, meaning a 1.15- to 1.86-fold **ISS** increase in the scattering coefficient at RH = 85% compared to dry conditions (Fig. 1e). Daily average values of ⁹⁰ f(RH = 85%, 525 nm) varied between 1.32 and 1.74, with low values (< 1.40) on 19 and 24 September and relatively high values (> 1.70) on 22 and 28 September and 2 October. Organic matter mass fractions were larger than 52%, and inorganic matter mass fractions were generally ⁹⁵ smaller when f(RH = 85%, 525 nm) was less than 1.40. However, high values of f(RH = 85%, 525 nm) in this study were often closely correlated with large fractions of water-soluble ions, such as NO₃⁻ and SO₄²⁻, in PM_{2.5}. Inorganic matter mass fractions were larger than 53%, and or- ¹⁰⁰



Figure 1. Hourly averaged time series (18 September to 4 October 2019) of the (**a**) scattering coefficient ($\sigma_{sp,525 \text{ nm}}$) measured by the dry nephelometer (unit: Mm⁻¹), (**b**) absorption coefficient ($\sigma_{ap,525 \text{ nm}}$; unit: Mm⁻¹), (**c**) single-scattering albedo ($\omega_{0(525 \text{ nm})}$), (**d**) scattering Ångström exponent ($\alpha_{(450-635 \text{ nm})}$), (**e**) particle light scattering enhancement factor at RH = 85 % (f(RH = 85%, 525 nm)), and (**f**) mass concentration of PM_{2.5} (unit: µg m⁻³) measured at the Yizhuang station. The segments of the time series with a blue background represent the occurrence of deliquescence. The timescale is Beijing time (UTC + 8 h). The date in this figure is in the month/day format.

ganic matter mass fractions were relatively small when the f(RH = 85%, 525 nm) was greater than 1.70. The campaign mean \pm standard deviation values of f(RH = 85%, 525 nm) were 1.64 \pm 0.13 (Table 1). Figure 2c reveals that strongly $_{5}$ hygroscopic aerosols with high values of f(RH = 85%, 525 nm) primarily came from the southeast sector. The proportion of secondary inorganics with strong hygroscopic

abilities in aerosols from this direction was high, while the proportion of organic matter with weak hygroscopic abilities was low (Fig. 2e–f). Figure 2d indicates that the mass fraction of eBC with weak hygroscopicity was slightly low in the southeast sector when wind speeds were lower than 4 m s^{-1} . However, when wind speeds were higher than 4 m s^{-1} , the mass fraction of eBC was relatively high in this direction.

Table 1. Average $\sigma_{\text{sp},525 \text{ nm}}$, $\sigma_{\text{ap},525 \text{ nm}}$, $\omega_{0(525 \text{ nm})}$, $\alpha_{(450-635 \text{ nm})}$, f(RH = 85%, 525 nm), and PM_{2.5} mass concentration values at different pollution levels.

	Entire	$PM_{2.5}$ pollution levels (µg m ⁻³)					
	observation period	Very clean $(PM_{2.5} \le 35)$	$\begin{array}{l} \mbox{Moderately polluted} \\ (35 < PM_{2.5} \leq 75) \end{array}$	Heavily polluted $(PM_{2.5} > 75)$			
$\sigma_{\rm sp, 525 nm} ({\rm Mm}^{-1})$	245 ± 168	89 ± 45	279 ± 93	530 ± 75			
$\sigma_{ap,525\rm{nm}}(\rm{Mm}^{-1})$	50 ± 24	43 ± 21	51 ± 20	50 ± 27			
$\omega_{0(525\mathrm{nm})}(-)$	0.77 ± 0.15	0.65 ± 0.16	0.83 ± 0.08	0.93 ± 0.04			
$\alpha_{(450-635\text{nm})}$ (-)	1.55 ± 0.16	1.58 ± 0.15	1.54 ± 0.15	1.33 ± 0.13			
f(RH = 85%, 525 nm) (-)	1.64 ± 0.13	1.49 ± 0.16	1.70 ± 0.06	$1.71{\pm}0.05$			
$PM_{2.5} (\mu g m^{-3})$	44 ± 25	22 ± 9	51 ± 12	90 ± 9			



Figure 2. Wind dependence of (**a**) the single-scattering albedo ($\omega_{0(525 \text{ nm})}$), (**a**) the scattering Ångström exponent ($\alpha_{(450-635 \text{ mm})}$), (**c**) the particle light scattering enhancement factor at RH = 85 % (f(RH = 85%, 525 nm)), (**d**) the mass fraction of eBC, (**e**) the mass fraction of inorganic matter, and (**f**) the mass fraction of organic matter. The circular contours show the average change in wind speed and direction.

Of all data associated with southeast winds, only three cases were identified with wind speeds higher than 4 m s^{-1} , likely winds of short duration so not representative. In the northeast direction, high values of f(RH = 85%, 525 nm) occurred

- $_{5}$ when the wind speed was lower than 2 m s⁻¹. The hygroscopic capacity of aerosols also weakened as the wind speed increased (Fig. 2c). The proportion of secondary inorganics with strong hygroscopicity decreased with increasing wind speed (Fig. 2e), while the proportion of substances with weak
- ¹⁰ hygroscopicity, such as organic matter and eBC, increased with increasing northeasterly wind speeds (Fig. 2d, f). Furthermore, aerosols from the southwest and southern sectors within the wind-speed range of 2 to 4 m s^{-1} had higher scattering enhancement factors (Fig. 2c) mainly because of the
- ¹⁵ deliquescence of sulfates in the ambient aerosols. The specific reasons are explained in detail in Sect. 3.3. Figure S5

shows that apart from the lower values (10th percentile values in Table S1), a small wavelength dependence in scattering enhancement factor is found in all other percentiles, with a stronger wavelength dependence for high values of $_{20}$ f(RH = 85%). Zieger et al. (2014) and Zhang et al. (2015) obtained similar results for Melpitz, Germany, and Linan, China, respectively.

Figure S6a to b show the time series of mass concentrations and mass fractions, respectively, of submicron aerosols, ²⁵ i.e., organic matter, nitrate, sulfate, ammonium, chloride, and eBC in PM_{2.5}. The rightmost pie chart in Fig. S6c shows that during the entire observation period, organic matter was the major component of PM_{2.5}, accounting for 39 %. Nitrate and sulfate comprised similar fractions of PM_{2.5}, i.e., ³⁰ 21 % and 19 %, respectively. The mass fraction of nitrate was slightly larger than that of sulfate. Note that eBC ac-

counted for 11 % of PM2.5 during the entire measurement period. Two special periods were noted. One started in the afternoon of 21 September and ended late morning on 22 September (Sect. I in Fig. S6a). First, the concentrations of 5 all chemical components during this period were high. As shown by the leftmost pie chart in Fig. S6c, the mass fraction of nitrate was the largest, accounting for 33 % of the total mass fraction on average, lasting a long time. However, compared with the proportion of inorganic matter (66%), 10 the mass fraction of organic matter was much smaller, accounting for 27 % of the total mass fraction. Accordingly, the f(RH = 85%, 525 nm) remained at a high level during this period (Fig. 1e). The other special period was on 24 September (Sect. II in Fig. S6a). The mass concentrations 15 of all aerosol species remained low. The middle pie chart in Fig. S6c clearly demonstrates that organic matter comprised the main fraction of PM_{2.5}, accounting for 55 % on average, followed by eBC. The fraction of nitrate was especially small during this clean period. The fraction of sulfate, accounting 20 for 16%, was 4 times that of nitrate, which was an advantage for deliquescence in this period (discussed in more detail in Sect. 3.3). Although the proportions of hydrophobic organic matter and eBC in aerosols during this period was very high, f(RH = 85%, 525 nm) was not the lowest during the whole ₂₅ observation period because the f(RH = 85%, 525 nm) of deliquescence was higher than the normal value. Note that

 $PM_{2.5}$ aerosols at the observatory in suburban Beijing were faintly acidic during the observation period (Fig. S7), benefitting the hygroscopic enhancement of ambient aerosols.

³⁰ 3.2 The relationship between the particle light scattering enhancement factor and aerosol chemical composition

Figure 3 displays f(RH = 85%, 525 nm) as a function of the main chemical component mass fractions. The total aerosol ³⁵ mass concentration is the sum of mass concentrations of all chemical constituents, including nitrate, sulfate, ammonium, chloride, and organic matter measured by the ACSM and eBC retrieved by the AE33. The mass fractions of individual chemical components were calculated by respectively ⁴⁰ dividing the mass concentrations of sulfate, nitrate, ammonium, and eBC by the sum of all chemical constituents. The f(RH = 85%, 525 nm) and eBC were negatively correlated, with a correlation coefficient *R* equal to -0.62 (Fig. 3d). A positive correlation is seen between f(RH = 85%, 525 nm)⁴⁵ and the three other inorganic substances' mass fractions because of their hygroscopic characteristics. The ammonium

- mass fraction had the strongest positive correlation (R = 0.78) with f(RH = 85%, 525 nm) (Fig. 3c). The reason is that ammonium is the common positive ion of $(NH_4)_2SO_4$ ⁵⁰ and NH₄NO₃, two major salts of inorganic substances in aerosols. The relationship between f(RH = 85%, 525 nm)
- and ammonium is thus similar to that between f(RH = 85%, 525 nm)and ammonium is thus similar to that between f(RH = 85%, 525 nm) and inorganic content. The hygroscopic properties

were different for NH₄NO₃ and (NH₄)₂SO₄. As expected, f(RH = 85%, 525 nm) was positively correlated with the ⁵⁵ sum of the nitrate and sulfate mass fractions (slope = 1.03 and R = 0.79; Fig. S8), similar to the correlation between f(RH = 85%, 525 nm) and the inorganic mass fraction. Wu et al. (2017) and Zieger et al. (2014) reported similar results.

Figure 4 shows f(RH = 85%, 525 nm) as a function of 60 the mass fractions of organic and inorganic matter. The mass fractions of inorganic and organic matter were respectively calculated by dividing inorganic matter (the sum of nitrate, sulfate, ammonium, and chloride) and organic matter mass concentrations by the total mass concentration. The inorganic 65 mass fraction was positively correlated with f(RH = 85%), 525 nm) because of the high hygroscopicity of the inorganic compounds, while organic substances were negatively correlated with f(RH = 85%, 525 nm) because of their lower hygroscopicity (P. Zhao et al., 2019). Both correlation coef-70 ficients were similar to those from previous studies (Zieger et al., 2014; Zhang et al., 2015; Wu et al., 2017). Also, the absolute values of both the slopes and corresponding standard deviations found here (0.80 ± 0.04 and 1.00 ± 0.06 for f(RH = 85%, 525 nm) as a function of inorganic and or- 75 ganic matter mass fractions, respectively) were similar to those reported in Lin'an, China $(0.96 \pm 0.02 \text{ and } 1.20 \pm 0.04,$ respectively; Zhang et al., 2015), but much lower than those observed in Melpitz, Germany $(2.2 \pm 0.078 \text{ and } 3.1 \pm 0.1,$ respectively; Zieger et al., 2014). This might be because 80 the f(RH = 85%, 525 nm) measured in Melpitz, Germany, was much higher than that in Lin'an and Beijing. Ambient aerosols in Melpitz, Germany, were partially affected by sea salt, like sodium chloride, transported from the North Sea and being highly hygroscopic. Marine aerosols have a higher 85 hygroscopicity than aerosols influenced more by human activity.

Distinguishing between data points below and above the 40 % organic mass fraction level in Fig. 4, the absolute values of the linear regression slope and R for data below 90 40 % were lower than those for data above 40 %. However, for the inorganic mass fraction (left panels of Fig. 4), the absolute values of the linear regression slope and Rfor data below 40 % were higher than those for data above 40 %. This indicates that the positive correlation between 95 f(RH = 85%, 525 nm) and the inorganic matter mass fraction was very strong when inorganic matter mass fractions were smaller than 40 %. However, when inorganic matter mass fractions were larger than 40%, the increasing tendency in f(RH = 85%, 525 nm) as the proportion of inor- 100 ganic matter increased slowly decreased. Inversely, the negative correlation between f(RH = 85%, 525 nm) and the organic matter mass fraction was slightly weak when organic matter mass fractions were smaller than 40 %. When organic matter mass fractions were larger than 40 %, the decreasing 105 tendency in f(RH = 85%, 525 nm) as the proportion of organic matter decreased became more robust. There may be several reasons for this phenomenon. First, particle size may

2.0 2.0 (b) (a) J(RH=85%,525 nm) .1 8 (RH=85%,525nm) .1 .1 .8 1.2 $y=(0.54\pm0.06)x+(1.55\pm0.01)$ R=0.46 $v=(0.83\pm0.11) x+(1.47\pm0.02) R=0.42$ 1.0 1.0 10 20 30 40 0 10 20 30 40 50 SO₄²⁻ mass fraction (%) NO3⁻ mass fraction (%) 2.0 2.0 (c) (d) J(RH=85%,525 nm) f(RH=85%,525nm) 1.6 14 1.2 1.2 y=(3.47±0.16)x+(1.29±0.02) R=0.78 $.39\pm0.10$ x+(1.81±0.01) R= 1.0 1.05 10 15 10 20 30 40 NH4⁺ mass fraction (%) eBC mass fraction (%)

Figure 3. The particle light scattering enhancement factor f(RH = 85%, 525 nm) as a function of different aerosol chemical component mass fractions measured by the ACSM and the AE33: (a) sulfate (SO_4^{2-}) mass fraction, (b) nitrate (NO_3^{-}) mass fraction, (c) ammonium (NH_4^{+}) mass fraction, and (d) black carbon (eBC) mass fraction. Solid green lines represent bivariate linear regressions. The dark-green shaded areas denote 95% confidence levels, and the light-green shaded areas show the 95% prediction bands for the fits. The linear regression function and the Pearson's correlation coefficient (*R*) are given in each panel.

be one of the most important factors to explain this. For particles with strong hygroscopicity, if their particle number size distribution tends towards large particle sizes, their hygroscopic growth ability may be similar to that of smaller par-5 ticles with weak hygroscopicity (Zieger et al., 2010, 2013;

- Wang et al., 2017, 2018). On the one hand, the f(RH) of particles usually decreases with increasing particle size (Zieger et al., 2013), resulting in a lower f(RH) for larger particles. On the other hand, the larger particles' amplification effect
- ¹⁰ of scattering cross section because of hygroscopic growth is weaker than that of smaller particles (Wu et al., 2017). It may be that when there is a high proportion of inorganic matter and a low proportion of organic matter, the inorganic matter is mainly composed of relatively large particles. It is possi-
- ¹⁵ ble that the f(RH = 85%, 525 nm) of aerosols with a high level of inorganic matter and a low level of organic matter is not as high as expected due to the compensating effect between the size and the chemical composition of aerosol. Second, the higher mass concentration of ambient aerosols is ²⁰ maybe another reason. Overall, when the mass concentration
- of inorganic substances was less than 40%, the total mass concentration of aerosols was relatively low in this experi-

ment. When the proportion of inorganic matter was higher than 40 %, the total mass concentration of aerosols was high, with a clear inhibiting effect of high aerosol mass concentration to the f(RH = 85 %, 525 nm) (Fig. 4e). Finally, it is also possible that when the proportion of inorganic matter is very high, aerosols absorb too much water vapor, leading to insufficient ambient water vapor.

The green dots in Fig. S9a and b represent deliquescence. ³⁰ The range of inorganic mass fraction in these deliquescent processes was from 30 % to 50 %, and the range of organic mass fraction was from 40 % to 60 %. A comparison between Fig. 4a and c and b and d shows that the proportion of sulfate in the total aerosol was much higher than that of nitrate for these deliquescent processes. Meanwhile, the f(RH = 85 %,525 nm) of these deliquescent processes were all above the best-fit regression lines. This demonstrates that the scattering enhancement factor at 85 % RH of deliquescence was generally higher than that of the non-deliquescent process. Kuang et al. (2016) also drew the same conclusion.

Comparing Fig. 4a and c, as the proportion of inorganic salts in the total aerosol increased, the proportion of nitrate in the total aerosol also increased, unlike the proportion of



Figure 4. The particle light scattering enhancement factor f (RH = 85 %, 525 nm) as a function of organic matter mass fraction and inorganic matter mass fraction measured by the ACSM and the AE33: f (RH = 85 %, 525 nm) as a function of (**a**) inorganic matter mass fraction and (**b**) organic matter mass fraction colored by the nitrate mass fraction; f (RH = 85 %, 525 nm) as a function of (**c**) inorganic matter mass fraction and (**d**) organic matter mass fraction colored by the sulfate mass fraction; and f (RH = 85 %, 525 nm) as a function of (**c**) inorganic matter mass fraction of (**c**) inorganic matter mass fraction colored by the sulfate mass fraction; and f (RH = 85 %, 525 nm) as a function of (**e**) inorganic matter mass fraction colored by the sulfate mass concentration of PM_{2.5}. Dotted black lines denote bivariate linear regressions. The red and magenta lines are the best-fit linear regression lines through data points associated with mass fractions smaller than 40 % and larger than 40 %, respectively. The linear regression functions and the Pearson's correlation coefficients (*R*) are given in each panel.



Figure 5. Scatter plots of γ as a function of the relative amounts of organic and inorganic matter (F_{org}): (a) $F_{org} = Org/(Org + (NH_4)_2SO_4 + NH_4HSO_4)$, (b) $F_{org} = Org/(Org + NH_4NO_3)$, and (c) $F_{org} = Org/(Org + (NH_4)_2SO_4 + NH_4HSO_4 + NH_4NO_3)$. Solid red lines show the linear fits, dotted red lines represent the 95% confidence levels, and solid blue lines represent the 95% prediction bands for the fit. The linear regression function and the squared Pearson's correlation coefficient (R^2) are given in each panel.

sulfate in the total aerosol. This demonstrates that nitrate played a primary role in affecting aerosol hygroscopic enhancement during the study period in Beijing. To further compare the correlation between sulfate and the hygroscop-5 icity of aerosols with that between nitrate and the hygroscopicity of aerosols, an ion-pairing scheme was used to calculate the mass concentrations of (NH₄)₂SO₄, NH₄HSO₄, and NH₄NO₃ in aerosols on the basis of the molar numbers of all ions (Gysel et al., 2007). The following is the ion-pairing 10 scheme:

$$n_{\rm NH_4NO_3} = n_{\rm NO_3^-},$$

$$n_{\rm H_2SO_4} = \max\left(0, n_{\rm SO_4^{2-}} - n_{\rm NH_4^+} + n_{\rm NO_3^-}\right),$$

$$n_{\rm NH_4HSO_4} = \min\left(2n_{\rm SO_4^{2-}} - n_{\rm NH_4^+} + n_{\rm NO_3^-}n_{\rm NH_4^+} - n_{\rm NO_3^-}\right),$$

$$n_{\rm (NH_4)_2SO_4} = \max\left(n_{\rm NH_4^+} - n_{\rm NO_3^-} - n_{\rm SO_4^{2-}}, 0\right),$$

$$n_{\rm HNO_3} = 0,$$
(9)

where *n* denotes the number of moles. Figure 5a to c show γ as a function of F_{org} (Eq. 6), where C_i represents the sum of (NH₄)₂SO₄ and NH₄HSO₄, NH₄NO₃ and the sum ¹⁵ of (NH₄)₂SO₄, NH₄HSO₄, and NH₄NO₃ mass concentrations, respectively. Overall, γ and F_{org} are negatively correlated. The coefficient of determination between γ and F_{org} (Org / (Org + NH₄NO₃)) (Fig. 5b) was higher than that between γ and F_{org} (Org / (Org + (Org + (NH₄)₂SO₄ + NH₄HSO₄))

²⁰ (Fig. 5a). The coefficient of determination between γ and F_{org} (Org / (Org + (NH₄)₂SO₄ + NH₄HSO₄ + NH₄NO₃) was the highest (Fig. 5c). This suggests that nitrate played a more significant role than sulfate in affecting aerosol hygroscopic growth during the study period in Beijing. In

- ²⁵ recent years, the Chinese government has made more efforts to control SO₂ emissions (Zhang et al., 2019), e.g., adjusting and optimizing industrial capacities. Clean fuels have also been promoted in the residential sector, with trials for using clean energy in heating in northern China carried out in all ³⁰ "2+26" cities and in the Fenhe River and Weihe River plains.
- In addition, compliance with industrial emission standards

has been strengthened. Desulfurization technology has been also applied to many heavy industrial facilities. However, China has many small-scale manufacturing enterprises, so it is much more difficult to regulate NO_x emissions than SO_2 35 emissions. H. Li et al. (2109) have reported that emissions of SO₂ and NO_x in 2017 dropped by 79.9% and 38.1%, respectively, from 2014 levels in Beijing, China. In 2020, SO₂ and primary PM_{2.5} emissions dropped to 1 million tons, while NO_x and volatile organic compound emissions 40 were still 10 million tons. As a result, the decrease in SO₂ resulted in an increase in NH₄NO₃ (Morgan et al., 2010; Xu et al., 2019; Zhang et al., 2019; H. Li et al., 2019). Several previous studies focusing on megacities like Shanghai and Beijing have all suggested that the increase in nitrate mass concentration played an important role in enhancing the water content of submicron aerosols and reducing visibility under high RH conditions (Sun et al., 2012; Shi et al., 2014; Zhang et al., 2015).

Figure S10a shows the scatterplot of γ as a function 50 of $F_{\text{org}}(\text{Org} + (\text{NH}_4)_2\text{SO}_4 + \text{NH}_4\text{HSO}_4 + \text{NH}_4\text{NO}_3)$, where the color of the data points represents the $SO_4^{2-}/(SO_4^{2-}+SO_2)$ molar ratio. This molar ratio indicates the relative age of aerosols (Quinn et al., 2005) because by gas- and aqueous-phase oxidation processes, 55 SO_2 will convert to SO_4^{2-} . The molar ratio of more aged aerosols is high due to the sufficient time for the conversion. The $SO_4^{2-}/(SO_4^{2-}+SO_2)$ molar ratio is low for younger aerosols. The figure shows that high γ corresponded to high $SO_4^{2-}/(SO_4^{2-}+SO_2)$ molar ratios with a low or- 60 ganic matter content, while low γ corresponded to low $SO_4^{2-}/(SO_4^{2-}+SO_2)$ molar ratios with a high organic matter content, consistent with results reported by Quinn et al. (2005) and Zhang et al. (2015). This demonstrates that the hygroscopicity of aged aerosols is higher than that of 65 younger aerosols. Figure S10b shows that the aerosol scattering coefficients $\log_{10} (\sigma_{sp})$ were relatively low when γ was low and the organic matter mass fraction was large. On the contrary, $\log_{10}(\sigma_{sp})$ was high when γ was high, and the organic mass fraction was small, with a relatively large variation.

3.3 Deliquescence of ambient aerosols

Figure 6a shows the time series of η , with the color of the data ⁵ points representing the ratio of SO_4^{2-} mass concentration to NO_3^- mass concentration. Figure 6b and c show the time series of wind direction and wind speed and ambient temperature and RH, respectively. Overall, deliquescence often occurred under high ambient temperature and low RH condi-10 tions (Fig. 6c). It also more easily occurred when winds with low speeds came from the south or southwest (Fig. 6b). High values of η usually occurred when the ratio of SO₄²⁻ mass concentration to NO₃⁻ mass concentration was high (mostly higher than \sim 4). Figure 7a shows a relatively strong corre-¹⁵ lation between η and the ratio of SO_4^{2-} mass concentration to NO₃⁻ mass concentration (R = 0.62). The blue dots represent the hysteresis index ($\eta > 0.4$) of deliquescence, showing that the corresponding ratios of SO_4^{2-} mass concentration to NO_3^- mass concentration were high (mostly higher

- ²⁰ than \sim 4). The red dots represent the hysteresis index of nondeliquescent processes, showing that the corresponding ratios of SO₄²⁻ mass concentration to NO₃⁻ mass concentration were generally less than 4. Because PM_{2.5} mass concentrations were extremely low on 24 September 2019, data were
- ²⁵ noisy then, and the system error was relatively large (green dots). When these cases of large systematic errors were eliminated, the correlation between η and the ratio of SO₄²⁻ to NO₃⁻ mass concentration increased (R = 0.69). In the North China Plain, Kuang et al. (2016) also observed a similar del-
- ³⁰ iquescent phenomenon associated with (NH₄)₂SO₄, while in Ny-Ålesund, Norway, this deliquescent phenomenon was related to sea salt (Zieger et al., 2010).

According to observational results, there are two environmental conditions of consequence for ambient aerosol deli-³⁵ quescence:

- 1. high ambient temperature and low ambient RH
- 2. relatively good air quality and solar illumination.

Concerning the first condition, Cheung et al. (2015) proposed an indicator describing the neutralization extent of aerosols, $_{40}$ i.e., the molar ratio, MR = (NH_4^+ - NO_3^-) / SO_4^{2^-}. The MR value of the non-deliquescent process was always the lowest in their study. Their results indicate that when the nitrate content of ambient aerosols was slightly high, and the sulfate content was low, aerosol particles did not exhibit del-

- ⁴⁵ iquescence resulting from acidic sulfate. Here, only when the mass concentration ratio of sulfate to nitrate was high (mostly higher than \sim 4) did ambient aerosol particles exhibit deliquescence. Figure 8a and b illustrate the diurnal variations in nitrate, along with ammonium and sulfate. Both the
- ⁵⁰ mass concentration and mass fraction of nitrate reached their largest and smallest values in the early morning and in the

afternoon, respectively. It was closely correlated with the diurnal variation in ambient temperature and RH, where RH gradually rose as the temperature gradually decreased from the late afternoon to the early morning of the next day, fol-55 lowed by a reversal of the trend into the late afternoon of the next day (Fig. 6c). Morino et al. (2006) and Wang et al. (2009) found that the formation of nitrate requires low temperatures and high RH, conditions favorable for the conversion of gaseous nitric acid to solid-phase nitrate. Ambi-60 ent aerosol deliquescence is thus closely related to the environmental conditions of high temperature and low humidity. Note that secondary sulfates are usually produced by photochemical reactions so that the sulfate content of aerosols gradually accumulates from late morning to late afternoon 65 (Huang et al., 2010; Sun et al., 2012). Generally speaking, the ambient temperature was higher, and the humidity was lower in the afternoon, so high values of SO_4^{2-} / NO_3^{-} mostly occurred in the afternoon. This may also explain why the deliquescent cycle occurred most frequently in the afternoon, as 70 shown in Fig. 7b.

The blue-shaded zones in Fig. 1a and f show that deliquescence usually occurred when the air quality was good. To better explain this phenomenon, the observation period was divided into a very clean period ($PM_{2.5} \le 35 \,\mu g \,m^{-3}$) and a 75 moderately polluted period ($PM_{2.5} > 75 \,\mu g \,m^{-3}$). The difference between f(RH = 85%, 525 nm) and f(RH = 80%, 525 nm)525 nm), i.e., DF, or f(RH = 85%, 525 nm) minus f(RH = 80%, 525 nm), was calculated. Figure 8c shows the values of DF during a very clean period and a mod-80 erately polluted period. The DF of deliquescence is larger than that of non-deliquescence because the f(RH) of deliquescence jumps suddenly as the RH increases from 80 % to 85%, while the f(RH) of non-deliquescence increases smoothly as the RH increases. The values of the particle 85 light scattering enhancement factor at high RH for deliquescent processes were usually higher than those for nondeliquescent processes, consistent with the strong hygroscopicity of (NH₄)₂SO₄. Figure 8c shows that the diurnal range of DF during the clean period was larger than that dur-90 ing the moderately polluted period. In particular, from late morning to late afternoon, the DF during the clean period was much higher than that during the moderately polluted period. Figure 8d shows the diurnal variation in SO_4^{2-} / NO_3^{-} . The ratio SO_4^{2-} / NO_3^{-} during the clean period was higher 95 than that during the moderately polluted period, especially from late morning to late afternoon. Overall, good air quality and solar illumination were conducive to photochemical reactions so that more secondary sulfate could be generated during the day, facilitating the deliquescence of ambient 100 aerosols.



Figure 6. Time series of (a) hysteresis index η colored by the SO₄²⁻/NO₃⁻ mass concentration ratio, (b) wind speed (WS; unit: m s⁻¹) colored by wind direction (WD; unit: °), and (c) ambient temperature (*T*; unit: °C) and relative humidity (RH; unit: %). The segments of the time series with a grey background represent the processes of deliquescence. The date in this figure is in the month/day format.



Figure 7. (a) Scatter plot of hysteresis index η as a function of the SO₄²⁻/NO₃⁻ mass concentration ratio. Blue dots represent deliquescence and red dots non-deliquescent processes. Green dots represent those data points with high systematic errors. The inset figure shows the scatter plot excluding the green dots. (b) Scatter plots of the observed f(RH = 85%, 525 nm) values for non-deliquescent (black) and deliquescent (blue) cycles. The linear regression functions and the Pearson's correlation coefficients (*R*) are given in the bottom right corner of (a) for each fitted line.



Figure 8. Average diurnal cycles of (**a**) mass concentration (unit: μ g m⁻³) and (**b**) mass fraction (unit: %) of ammonium (black curves), nitrate (blue curves), and sulfate (red curves) measured by the ACSM. (**c**) The difference between f(RH = 85%, 525 nm) and f(RH = 80%, 525 nm) and (**d**) the mass concentration ratio of sulfate to nitrate (SO₄²⁻/NO₃⁻) during the clean period (blue curve) and the moderately polluted period (red curve). Vertical lines are the standard deviations.

3.4 Parameterizations of f(RH)

3.4.1 Parameterization with the equation $f(\mathbf{RH}) = 1 + m \times \mathbf{RH}^n$

Many empirical expressions have been presented to param-⁵ eterize f(RH) (Kotchenruther and Hobbs, 1998; Carrico et al., 2003; Pan et al., 2009; Fierz-Schmidhauser et al., 2010a; Chen et al., 2014; Brock et al., 2016; Titos et al., 2016; Kuang et al., 2017). The following is the two-parameter scheme introduced by Kotchenruther and Hobbs (1998):

$$f(\mathbf{RH}) = 1 + m \times \mathbf{RH}^n.$$
(10)

The parameter *m* determines the largest value of f(RH = 100%), and the parameter *n* dominates the magnitude of the scattering enhancement, reflecting the curvature of the humidogram.

¹⁵ Deliquescence was frequently observed during the entire measurement campaign. In total, 294 cycles of f(RH)were measured, and 47 cycles (16% of all cycles) showed clear deliquescence (Fig. 9d, e). All f(RH) curves were thus first classified into deliquescent curves and non-deliquescent curves. After averaging PM2.5 concentrations of the corre- 20 sponding cycles, all non-deliquescent curves were further divided into clean ($PM_{2.5} \le 35 \,\mu g \, m^{-3}$), moderately polluted $(35 \,\mu g \,m^{-3} < PM_{2.5} \le 75 \,\mu g \,m^{-3})$, and polluted $(PM_{2.5} > 100 \,m^{-3})$ $75 \,\mu g \,m^{-3}$) categories. The deliquescent curves were divided into clean and moderately clean categories only be- 25 cause deliquescence mainly occurred under good air quality conditions. For cycles without deliquescence (Fig. 9a-c), the measured values were fitted using Eq. (10). For cycles with deliquescence (Fig. 9d, e), f(RH) increased smoothly under low RH conditions then increased sharply. Under 30 low and high RH conditions, the fitted f(RH) values were usually lower than observed values, but the slopes of the two curves were similar. However, when RH approached ~80% where f(RH) sharply increased, the fitted f(RH)values were usually higher than observed values, with different slopes of the two curves. Therefore, segment fitting (Eq. 12) was applied in the parameterization of deliquescent f(RH). The deliquescence observed in our study was primarily caused by $(NH_4)_2SO_4$ in ambient aerosols. The DRH of the pure (NH₄)₂SO₄ aerosols generated in the lab- ⁴⁰ oratory was 80.07 %, measured by our high-resolution hu-



Figure 9. Fitted f(RH) of non-deliquescent curves (**a**–**c**) and deliquescent curves (**d**–**e**) for different pollution levels. Solid red lines represent fitted curves parameterized by Eq. (10). Dotted black lines represent the curves fit by segment function Eq. (12), and the solid blue lines represent the average curves.

midified nephelometer system (Fig. S4). Because the DRH of all deliquescence in this study, according to statistics, was mainly distributed between 78 % and 80 %, Eq. (10) was applied to fit f(RH) values when $RH \ge 82$ % or $RH \le 78$ %. ⁵ For 78 % < RH < 82 %, f(RH) values were parameterized by Eq. (11):

$$f(\mathbf{RH}) = a \times \mathbf{RH} + b. \tag{11}$$

Parameter *a* determines the degree of f(RH) jumps during deliquescence, i.e., the slope of the linear fitting line. Param-¹⁰ eter *b* determines the f(RH, 525 nm) value before deliquescence (RH = 78 %) and after deliquescence (RH = 82 %).

$$\begin{cases} f(\text{RH}) = 1 + m \times \text{RH}^n & \text{RH} \le 78\% \\ f(\text{RH}) = a \times \text{RH} + b & 78\% < \text{RH} < 82\% \\ f(\text{RH}) = 1 + m \times \text{RH}^n & \text{RH} \ge 82\% \end{cases}$$
(12)

Table 2 summarizes the parameters *m* and *n* used in this study and in similar studies. For the non-deliquescence case, the ¹⁵ parameter *n* for the clean period was the largest ($\sim 27 \%$ larger than that for the moderately polluted period and $\sim 16 \%$ larger than that for the polluted period). The difference between the clean and moderately polluted parameter *m* was small, while the parameter *m* in the polluted period

²⁰ was about 0.1 larger than that in the other two periods. As a result, f(RH = 85%, 525 nm) was the largest in the polluted period and the smallest in the clean period. For the deliquescence case, the fitted parameters *m* and *n* for the whole RH range (40% < RH < 90%) were much larger than the *m* and *n* for RH < 78 % and slightly larger than the *m* and $_{25}$ *n* for RH > 82 %. If Eq. (10) was used to parameterize the whole RH range of the deliquescence curves instead of segment fitting, bigger differences between the parameterized fitting results and measured values would occur. This would cause greater uncertainties in the model simulation of aerosol 30 hygroscopicity. Also, there was no significant difference between parameter a under clean ($a = 0.07 \pm 0.006$) and moderately polluted ($a = 0.07 \pm 0.003$) conditions, indicating that the trend and amplitude of the jump growth for aerosol deliquescence are consistent under these two environmental con- 35 ditions. However, the absolute value of parameter b under clean conditions ($b = -4.29 \pm 0.47$) was higher than that under moderately polluted conditions ($b = -3.88 \pm 0.24$), indicating that the DRH for clean periods was slightly higher than that for moderately polluted periods. 40

3.4.2 Steepness of humidograms

The steepness index proposed by Zhang et al. (2015), τ , is defined as

$$\tau = \frac{f'(80\%)}{f'(60\%)} - 1 = \left(\frac{4}{3}\right)^{n-1} - 1 \tag{13}$$

to quantitatively describe the change in the curvature of the $_{45}$ humidogram curves; f'(80%) and f'(60%) are the deriva-

Classification			m	n	а	b	Reference
Non-deliquescence	Very clean Moderately polluted Polluted		$ \begin{array}{ } 1.32 \pm 0.07 \\ 1.29 \pm 0.02 \\ 1.42 \pm 0.04 \end{array} $	$5.05 \pm 0.18 \\ 3.99 \pm 0.06 \\ 4.35 \pm 0.08$			This study
Deliquescence	Very clean	RH < 78~%	0.55 ± 0.02	3.51 ± 0.09			
		$RH > 82 \ \% \ 78 \ \% < RH < 82 \ \%$	1.63 ± 0.10	5.84 ± 0.41	0.07 ± 0.006	-4.29 ± 0.47	
		40~% < RH < 90~%	1.92 ± 0.41	6.96 ± 1.63			
	Moderately polluted	RH < 78~%	0.62 ± 0.01	3.08 ± 0.04			
		$RH > 82 \ \% \ 78 \ \% < RH < 82 \ \%$	1.50 ± 0.04	4.88 ± 0.19	0.07 ± 0.003	-3.88 ± 0.24	
		40~% < RH < 90~%	1.63	5.61			
Clean Polluted Dust				$\begin{array}{c} 6.70 \pm 0.27 \\ 6.27 \pm 0.10 \\ 5.17 \pm 0.4 \end{array}$			Pan et al. (2009)
Locally polluted Northerly polluted Dust-influenced			$ \begin{array}{ } 1.24 \pm 0.29 \\ 1.20 \pm 0.21 \\ 1.02 \pm 0.19 \end{array} $	$5.46 \pm 1.90 \\ 3.90 \pm 1.27 \\ 4.51 \pm 0.80$			Zhang et al. (2015)

Table 2. Curve-fitting parameters of f(RH) at 525 nm for different aerosol types using Eqs. (10) and (12).

tives of the fitted curve of f(RH) given in Eq. (10) at the two different RH values.

A low τ means that the curvature of the humidogram is small, and a high τ means that the slopes of the curve from ⁵ low to high RH sharply change. Figure S11 shows the scatter plot of τ as a function of the nitrate mass fraction, colored by the sulfate mass fraction. In the 0%–15% nitrate mass fraction range, τ decreased sharply as the nitrate mass fraction increased, demonstrating that the curvature of the humido-¹⁰ gram became smaller. For nitrate mass fractions larger than

15 %, τ stabilized to a constant value of ~ 1.6, and the curvature of the humidogram was much smaller.

4 Conclusions

Direct measurements of aerosol hygroscopicity, as expressed ¹⁵ by $f(RH,\lambda)$, were carried out at a site in the southern urban edge of Beijing, aimed at investigating the effect of aerosol water uptake on particle light-scattering properties. The mass concentrations of aerosol chemical components were measured simultaneously by the ACSM and AE-33. Also mea-²⁰ sured were other aerosol parameters, such as the light absorp-

tion coefficient and the mass concentration of $PM_{2.5}$. In total, 294 cycles of f(RH) were measured, and 47 cycles (16% of all cycles) showed clear deliquescence.

The proportion of components making up the chemical ²⁵ composition of aerosols is key to influencing f(RH). In general, f(RH) had a strong positive correlation with the proportion of inorganic matter and a negative correlation with the proportion of organic matter and eBC. High values of f(RH) usually occurred for aged aerosols whose ³⁰ mass fraction of organic matter was small. Low values of f(RH = 85%, 525 nm) often occurred for primary aerosols

with more organic matter. Furthermore, when the mass frac-

tion of inorganic matter was smaller than 40%, the positive correlation between f(RH = 85%, 525 nm) and inorganic matter was much stronger. A similar phenomenon was also found for the negative correlation between organic matter and f(RH = 85%, 525 nm). The compensating effect between the size and chemical composition of aerosol may be one of the main reasons for this phenomenon. High mass concentrations of ambient aerosols and insufficient water vapor in the sample flow may be the other two reasons. Nitrate also played a more significant role in affecting aerosol hygroscopic growth than sulfate in Beijing.

Favorable meteorological conditions for aerosol deliquescence were high ambient temperature and low RH, as well ⁴⁵ as relatively good air quality and strong solar illumination. High temperatures and low RH levels were not conducive to the formation of nitrate, while good air quality and light conditions were conducive to photochemical reactions so that more secondary sulfates could be generated. Only when the ⁵⁰ ratio of the sulfate mass fraction to the nitrate mass fraction was greater than ~ 4 did the deliquescence phenomenon of ambient aerosols easily occur.

All humidograms were first classified as either deliquescent or non-deliquescent. The two kinds of humidograms ⁵⁵ were further classified according to the mass concentration of PM_{2.5}. The two-parameter scheme, $f(RH) = 1 + m \times RH^n$, introduced by Kotchenruther and Hobbs (1998), was applied to fit the non-deliquescent f(RH). The deliquescent f(RH)was parameterized by segment functions (Eq. 12). For the ⁶⁰ deliquescence case, the fitted parameters *m* and *n* for the whole RH range (40 % < RH < 90 %) were much larger than the parameters for RH < 78 % and higher than the parameters for RH > 82 %. This demonstrates that large errors would be incurred if only Eq. (10) was used to parameterize the ⁶⁵ whole RH range of deliquescence curves. The piecewise parameterization scheme (Eq. 12) is a better fit for humidograms representing deliquescence to reduce uncertainties in the model simulation of aerosol hygroscopicity. The curvature of the f(RH) humidogram, described by the steepness index, decreased sharply as the nitrate mass fraction in-

 $_{5}$ creased within the range of 0 %–15 %. When the nitrate mass fraction was larger than 15 %, the steepness index remained constant (~1.6).

Data availability. The data used in this paper can be downloaded online (https://doi.org/10.11922/sciencedb.00785)

¹⁰ *Supplement*. The supplement related to this article is available online at: https://doi.org/10.5194/acp-21-1-2021-supplement.

Author contributions. ZL and PY designed the field experiment. ZL, PY, and RR determined the main goal of this study. RR processed the measurement data and prepared this paper with contri-

¹⁵ butions from all co-authors. PY, HW, and YW provided technical guidance for instrumentation. MC copy-edited the article. WW, XJ, YL, and DZ⁽²¹⁶⁾ participated in the implementation of this experiment.

Competing interests. The authors declare that they have no conflict ²⁰ of interest.

Acknowledgements. This work was supported by the National Key R&D Program of China (no. 2017YFC1501702), the National Science Foundation of China (nos. 91544217, 42005067), the Open Fund of State Key Laboratory of Remote Sensing Science (no. 25 202015), and the Guangdong Basic and Applied Basic Research Fund Committee (2020B1515130003).

Financial support. This research has been supported by the NAME OF FUNDER (grant no. GRANT AGREEMENT NO).

Review statement. This paper was edited by Paul Zieger and re-³⁰ viewed by two anonymous referees.

Anderson, T. L. and Ogren, J. A.: Determining aerosol

Bond, T. C. and Bergstrom, R. W.: Light absorption by carbona-

Brock, C. A., Wagner, N. L., Anderson, B. E., Attwood, A. R.,

27-67, https://doi.org/10.1080/02786820500421521, 2006.

ceous particles: an investigative review, Aerosol Sci. Tech., 40,

Beyersdorf, A., Campuzano-Jost, P., Carlton, A. G., Day, D. A.,

the TSI

Sci.

3563

29.

Tech..

integrat-

57-69.

using

Aerosol

https://doi.org/10.1080/02786829808965551, 1998.

References

radiative properties

ing nephelometer,

ertainties in
y. The cur-
/ the steep-
fraction in-
nitrate massDiskin, G. S., Gordon, T. D., Jimenez, J. L., Lack, D. A., Liao,
J., Markovic, M. Z., Middlebrook, A. M., Ng, N. L., Perring,
A. E., Richardson, M. S., Schwarz, J. P., Washenfelder, R. A.,
Welti, A., Xu, L., Ziemba, L. D., and Murphy, D. M.: Aerosol
optical properties in the southeastern United States in summer
- Part 1: Hygroscopic growth, Atmos. Chem. Phys., 16, 4987–
5007, https://doi.org/10.5194/acp-16-4987-2016, 2016.

- Carrico, C. M., Rood, M. J., and Ogren, J. A.: Aerosol light scattering properties at Cape Grim, Tasmania, during the First Aerosol Characterization Experiment (ACE 1), J. Geophys. Res.-Atmos., 50 103, 16565–16574, https://doi.org/10.1029/98JD00685, 1998.
- Carrico, C. M., Kus, P., Rood, M. J., Quinn, P. K., and Bates, T. S.: Mixtures of pollution, dust, sea salt, and volcanic aerosol during ACE: radiative properties as a function of relative humidity, J. Geophys. Res.-Atmos., 108, 8650, 55 https://doi.org/10.1029/2003JD003405, 2003.
- Chen, J., Zhao, C. S., Ma, N., and Yan, P.: Aerosol hygroscopicity parameter derived from the light scattering enhancement factor measurements in the North China Plain, Atmos. Chem. Phys., 14, 8105–8118, https://doi.org/10.5194/acp-14-8105-2014, 2014.
- Cheung, H. H. Y., Yeung, M. C., Li, Y. J., Lee, B. P., and Chan, C. K.: Relative humidity-dependent TDMA measurements of ambient aerosols at the HKUST supersite in Hong Kong, China, Aerosol Sci. Tech., 49, 643–654, https://doi.org/10.1080/02786826.2015.1058482, 2015.
- Doherty, S. J., Quinn, P. K., Jefferson, A., Carrico, C. M., Anderson, T. L., and Hegg, D.: A comparison and summary of aerosol optical properties as observed in situ from aircraft, ship, and land during ACE-Asia, J. Geophys. Res. Atmos., 110, D04201, https://doi.org/10.1029/2004jd004964, 2005.
- Fierz-Schmidhauser, R., Zieger, P., Gysel, M., Kammermann, L., DeCarlo, P. F., Baltensperger, U., and Weingartner, E.: Measured and predicted aerosol light scattering enhancement factors at the high alpine site Jungfraujoch, Atmos. Chem. Phys., 10, 2319– 2333, https://doi.org/10.5194/acp-10-2319-2010, 2010a.
- Fierz-Schmidhauser, R., Zieger, P., Wehrle, G., Jefferson, A., Ogren, J. A., Baltensperger, U., and Weingartner, E.: Measurement of relative humidity dependent light scattering of aerosols, Atmos. Meas. Tech., 3, 39–50, https://doi.org/10.5194/amt-3-39-2010, 2010b.
- Gysel, M., Crosier, J., Topping, D. O., Whitehead, J. D., Bower, K. N., Cubison, M. J., Williams, P. I., Flynn, M. J., McFiggans, G. B., and Coe, H.: Closure study between chemical composition and hygroscopic growth of aerosol particles during TORCH2, Atmos. Chem. Phys., 7, 6131–6144, https://doi.org/10.5194/acp-7-6131-2007, 2007.
- Han, T., Xu, W., Chen, C., Liu, X., Wang, Q., Li, J., Zhao, X., Du,
 W., Wang, Z., and Sun, Y.: Chemical apportionment of aerosol optical properties during the Asia Economic Cooperation summit in Beijing, China, J. Geophys. Res. Atmos., 120, 12281–12295, 90 https://doi.org/10.1002/2015JD023918, 2015.
- Huang, X.-F., He, L.-Y., Hu, M., Canagaratna, M. R., Sun, Y., Zhang, Q., Zhu, T., Xue, L., Zeng, L.-W., Liu, X.-G., Zhang, Y.-H., Jayne, J. T., Ng, N. L., and Worsnop, D. R.: Highly time-resolved chemical characterization of atmospheric submicron particles during 2008 Beijing Olympic Games using an Aerodyne High-Resolution Aerosol Mass Spectrometer, Atmos. Chem. Phys., 10, 8933–8945, https://doi.org/10.5194/acp-10-8933-2010, 2010.

65

70

75

80

R. Ren et al.: Measurement report

- IPCC: Climate Change 2013: The Physical Science Basis, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013. **INSP**
- James, D. J., Brianna, M. M., Dennis, E., Scott, C. C., and
- Scott, A. B.: Comparison of Single-Point and Continuous Sampling Methods for Estimating Residential Indoor Temperature and Humidity, J. Occup. Environ. Hyg., 12, 785–794, https://doi.org/10.1080/15459624.2015.1047024, 2015.
- Jin, X., Wang, Y., Li, Z., Zhang, F., Xu, W., Sun, Y., Fan, X.,
- Chen, G., Wu, H., Ren, J., Wang, Q., and Cribb, M.: Significant contribution of organics to aerosol liquid water content in winter in Beijing, China, Atmos. Chem. Phys., 20, 901–914, https://doi.org/10.5194/acp-20-901-2020, 2020.
- Kotchenruther, R. A. and Hobbs, P. V.: Humidification factors
- of aerosols from biomass burning in Brazil, J. Geophys. Res.-Atmos., 103, 32081–32089, https://doi.org/10.1029/98JD00340, 1998.
- Kuang, Y., Zhao, C. S., Ma, N., Liu, H. J., Bian, Y. X., Tao, J. C., and Hu, M.: Deliquescent phenomena of ambient aerosols
 on the North China Plain, Geophys. Res. Lett., 43, 8744–8750,

https://doi.org/10.1002/2016g1070273, 2016.

- Kuang, Y., Zhao, C., Tao, J., Bian, Y., Ma, N., and Zhao, G.: A novel method for deriving the aerosol hygroscopicity parameter based only on measurements from a humidified nephelometer system, Atmos. Chem. Phys., 17, 6651–6662,
- 25 fied nephelometer system, Atmos. Chem. Phys., 17, 6651–6662 https://doi.org/10.5194/acp-17-6651-2017, 2017.
- Li, H., Cheng, J., Zhang, Q., Zheng, B., Zhang, Y., Zheng, G., and He, K.: Rapid transition in winter aerosol composition in Beijing from 2014 to 2017: response to clean air actions, Atmos. Chem.
- ³⁰ Phys., 19, 11485–11499, https://doi.org/10.5194/acp-19-11485-2019, 2019.
- Li, Z., Wang, Y., Guo, J., Zhao, C., Cribb, M. C., Dong, X., Fan, J., Gong, D., Huang, J., Jiang, M., Jiang, Y., Lee, S.-S., Li, H., Li, J., Liu, J., Qian, Y., Rosenfeld, D., Shan, S., Sun,
- Y., Wang, H., Xin, J., Yan, X., Yang, X., Yang, X.-Q., Zhang, F., and Zheng, Y.: East Asian Study of Tropospheric Aerosols and their Impact on Regional Clouds, Precipitation, and Climate (EAST-AIRCPC), J. Geophys. Res.-Atmos., 124, 13026–13054, https://doi.org/10.1029/2019jd030758, 2019.
- ⁴⁰ Liu, H. and Zhao, C.: Design of a humidified nephelometer system with high time resolution, Acta Scientiarum Naturalium Universitatis Pekinensis, 52, 999–1004, https://doi.org/10.13209/j.0479-8023.2016.053, 2016.
- Liu, C., Chung, C. E., Yin, Y., and Schnaiter, M.: The absorption Ångström exponent of black carbon: from numerical aspects, At-
- mos. Chem. Phys., 18, 6259–6273, https://doi.org/10.5194/acp-18-6259-2018, 2018.
- Liu, L., Tan, H., Fan, S., Cai, M., Xu, H., Li, F., and Chan, P.: Influence of aerosol hygroscopicity and mix-
- ⁵⁰ ing state on aerosol optical properties in the Pearl River Delta region, China, Sci. Total Environ., 627, 1560–1571, https://doi.org/10.1016/j.scitotenv.2018.01.199, 2018.
 - Liu, X. G., Li, J., Qu, Y., Han, T., Hou, L., Gu, J., Chen, C., Yang, Y., Liu, X., Yang, T., Zhang, Y., Tian, H., and Hu, M.: Formation
- and evolution mechanism of regional haze: a case study in the megacity Beijing, China, Atmos. Chem. Phys., 13, 4501–4514, https://doi.org/10.5194/acp-13-4501-2013, 2013.
- Müller, T., Laborde, M., Kassell, G., and Wiedensohler, A.: Design and performance of a three-wavelength LED-based total scatter

and backscatter integrating nephelometer, Atmos. Meas. Tech., 60 4, 1291–1303, https://doi.org/10.5194/amt-4-1291-2011, 2011.

- Monks, P. S., Granier, C., Fuzzi, S., Stohl, A., Williams, M. L., Akimoto, H., Amann, M., Baklanov, A., Baltensperger, U., Bey, I., Blake, N., Blake, R. S., Carslaw, K. S., Cooper, O. R., Dentener, F. J., Fowler, D., Fragkou, E., Frost, G. J., Generoso, S., Ginoux, P., Grewe, V., Guenther, A., Hansson, H. C., Henne, S., Hjorth, J., Hofzumahaus, A., Huntrieser, H., Isaksen, I. S. A., Jenkin, M. E., Kaiser, J., Kanakidou, M., Klimont, Z., Kulmala, M., Laj, P., Lawrence, M. G., Lee, J. D., Liousse, C., Maione, M., McFiggans, G. B., Metzger, A., Mieville, A., Mous- 70 siopoulos, N., Orlando, J. J., O'Dowd, C. D., Palmer, P. I., Parrish, D. D., Petzold, A., Platt, U., Poschl, U., A.S.H. Prévôt, A. S. H., Reeves, C. E., Reimann, S., Rudich, Y., Sellegri, K., Steinbrecher, R., Simpson, D., ten Brink, H., Theloke, J., van Der Werf, G. R., Vautard, R., Vestreng, V., Vlachokostas, 75 C., and von Glasow, R.: Atmospheric composition change global and regional air quality, Atmos. Environ., 43, 5268-5350, https://doi.org/10.1016/j.atmosenv.2009.08.021, 2009.
- Morgan, W. T., Allan, J. D., Bower, K. N., Esselborn, M., Harris, B., Henzing, J. S., Highwood, E. J., Kiendler-Scharr, A., 80 McMeeking, G. R., Mensah, A. A., Northway, M. J., Osborne, S., Williams, P. I., Krejci, R., and Coe, H.: Enhancement of the aerosol direct radiative effect by semi-volatile aerosol components: airborne measurements in North-Western Europe, Atmos. Chem. Phys., 10, 8151–8171, https://doi.org/10.5194/acp-85 10-8151-2010, 2010.
- Morino, Y., Kondo, Y., Takegawa, N., Miyazaki, Y., Kita, K., Komazaki, Y., Fukuda, M., Miyakawa, T., Moteki, N., and Worsnop, D. R.: Partitioning of HNO₃ and particulate nitrate over Tokyo: effect of vertical mixing, J. Geophys. Res. Atmos., 111, D15215, 90 https://doi.org/10.1029/2005jd006887, 2006.
- Pan, X. L., Yan, P., Tang, J., Ma, J. Z., Wang, Z. F., Gbaguidi, A., and Sun, Y. L.: Observational study of influence of aerosol hygroscopic growth on scattering coefficient over rural area near Beijing mega-city, Atmos. Chem. Phys., 9, 7519–7530, 95 https://doi.org/10.5194/acp-9-7519-2009, 2009.
- Quinn, P. K., Bates, T. S., Baynard, T., Clarke, A. D., Onasch, T. B., Wang, W., Rood, M. J., Andrews, E., Allan, J., Carrico, C. M., Coffman, D., and Worsnop, D.: Impact of particulate organic matter on the relative humidity dependence of light scattering: 100 a simplified parameterization, Geophys. Res. Lett., 32, L22809, https://doi.org/10.1029/2005gl024322, 2005.
- Shi, Y., Chen, J., Hu, D., Wang, L., Yang, X., and Wang, X.: Airborne submicron particulate (PM₁) pollution in Shanghai, China: chemical variability, formation/dissociation ¹⁰⁵ of associated semi-volatile components and the impacts on visibility, Sci. Total Environ., 473, 199–206, https://doi.org/10.1016/j.scitotenv.2013.12.024, 2014.
- Sun, Y., Wang, Z., Dong, H., Yang, T., Li, J., Pan, X., Chen, P., and Jayne, J. T.: Characterization of summer organic ¹¹⁰ and inorganic aerosols in Beijing, China with an aerosol chemical speciation monitor, Atmos. Environ., 51, 250–259, https://doi.org/10.1016/j.atmosenv.2012.01.013, 2012.
- Titos, G., Cazorla, A., Zieger, P., Andrews, E., Lyamani, H., Granadosmunoz, M. J., Olmo, F. J., and Aladosar- 115 boledas, L.: Effect of hygroscopic growth on the aerosol light-scattering coefficient: a review of measurements, tech-

niques and error sources, Atmos. Environ., 141, 494–507, https://doi.org/10.1016/j.atmosenv.2016.07.021, 2016.

- Wang, X., Zhang, Y., Chen, H., Yang, X., Chen, J., and Geng, F.: Particulate nitrate formation in a highly polluted
- urban area: a case study by single-particle mass spectrometry in Shanghai, Environ. Sci. Technol., 43, 3061–3066, https://doi.org/10.1021/es8020155, 2009.
- Wang, Y., Zhang, F., Li, Z., Tan, H., Xu, H., Ren, J., Zhao, J., Du, W., and Sun, Y.: Enhanced hydrophobicity and volatil-
- ity of submicron aerosols under severe emission control conditions in Beijing, Atmos. Chem. Phys., 17, 5239–5251, https://doi.org/10.5194/acp-17-5239-2017, 2017.
- Wang, Y., Li, Z., Zhang, Y., Du, W., Zhang, F., Tan, H., Xu, H., Fan, T., Jin, X., Fan, X., Dong, Z., Wang, Q., and Sun, Y.: Characteri-
- zation of aerosol hygroscopicity, mixing state, and CCN activity at a suburban site in the central North China Plain, Atmos. Chem. Phys., 18, 11739–11752, https://doi.org/10.5194/acp-18-11739-2018, 2018.

Wang, Y., Li, Z., Zhang, R., Jin, X., Xu, W., Fan, X., Wu, H.,

Zhang, F., Sun, Y., Wang, Q., Cribb, M., and Hu, D.: Distinct ultrafine-and accumulation-mode particle properties in clean and polluted urban environments, Geophys. Res. Lett., 46, 10918– 10925, https://doi.org/10.1029/2019GL084047, 2019.

- Wanielista, M., Kersten, R., and Eaglin, R.: Hydrology: Water
- Quantity and Quality Control, 2nd edition, John Wiley & Sons, New York, the United States, 1997. TS10
- Wu, T., Li, Z., Chen, J., Wang, Y., Wu, H., Jin, X., Liang, C., Li, S., Wang, W., and Cribb, M.: Hygroscopicity of different types of aerosol particles: case studies using multi instrument data in megacity Beijing, China, Remote Sens., 12,
- 785, https://doi.org/10.3390/rs12050785, 2020.
- Wu, Y., Wang, X., Yan, P., Zhang, L., Tao, J., Liu, X., Tian, P., Han, Z., and Zhang, R.: Investigation of hygroscopic growth effect on aerosol scattering coefficient at a rural site in the southern North China Plain, Sci. Total Environ., 599, 76–84,
- https://doi.org/10.1016/j.scitotenv.2017.04.194, 2017.
- Xu, W., Sun, Y., Wang, Q., Zhao, J., Wang, J., Ge, X., Xie, C., Zhou, W., Du, W., Li, J., Fu, P., Wang, Z., Worsnop, D. R., and Coe, H.: Changes in aerosol chemistry from 2014 to
- 2016 in winter in Beijing: insights from high-resolution aerosol mass spectrometry, J. Geophys. Res.-Atmos., 124, 1132–1147, https://doi.org/10.1029/2018JD029245, 2019.
- Yan, P., Pan, X., Tang, J., Zhou, X., Zhang, R., and Zeng,
 L.: Hygroscopic growth of aerosol scattering coefficient: a comparative analysis between urban and sub-
- urban sites at winter in Beijing, Particuology, 7, 52–60, https://doi.org/10.1016/j.partic.2008.11.009, 2009.
- Yang, Y. R., Liu, X. G., Qu, Y., An, J. L., Jiang, R., Zhang, Y. H., Sun, Y. L., Wu, Z. J., Zhang, F., Xu, W. Q., and Ma,
- Q. X.: Characteristics and formation mechanism of continuous hazes in China: a case study during the autumn of 2014 in the North China Plain, Atmos. Chem. Phys., 15, 8165–8178, https://doi.org/10.5194/acp-15-8165-2015, 2015.
- Yu, Y., Zhao, C., Kuang, Y., Tao, J., Zhao, G., Shen, C., and Xu, W.:
 A parameterization for the light scattering enhancement factor with aerosol chemical compositions, Atmos. Environ., 191, 370–
 - 377, https://doi.org/10.1016/j.atmosenv.2018.08.016, 2018.
 - Zhang, L., Sun, J. Y., Shen, X. J., Zhang, Y. M., Che, H., Ma, Q. L., Zhang, Y. W., Zhang, X. Y., and Ogren, J. A.: Observa-

tions of relative humidity effects on aerosol light scattering in the ⁶⁰ Yangtze River Delta of China, Atmos. Chem. Phys., 15, 8439– 8454, https://doi.org/10.5194/acp-15-8439-2015, 2015.

- Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu,
 X., Wang, J., He, H., Liu, W., Ding, Y., Lei, Y., Li, J., Wang,
 Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L.,
 Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding,
 A., Gao, J., Fu, Q., Huo, J., Liu, B., Liu, Z., Yang, F., He, K.,
 and Hao, J.: Drivers of improved PM_{2.5} air quality in China
 from 2013 to 2017, P. Natl. Acad. Sci. USA, 116, 24463–24469,
 https://doi.org/10.1073/pnas.1907956116, 2019.
- Zhao, C., Yu, Y., Kuang, Y., Tao, J., and Zhao, G.: Recent progress of aerosol light-scattering enhancement factor studies in China, Adv. Atmos. Sci., 36, 1015–1026, https://doi.org/10.1007/s00376-019-8248-1, 2019.
- Zhao, G., Zhao, C., Kuang, Y., Bian, Y., Tao, J., Shen, C., and ⁷⁵ Yu, Y.: Calculating the aerosol asymmetry factor based on measurements from the humidified nephelometer system, Atmos. Chem. Phys., 18, 9049–9060, https://doi.org/10.5194/acp-18-9049-2018, 2018.
- Zhao, P., Ding, J., Du, X., and Su, J.: High time-resolution ⁸⁰ measurement of light scattering hygroscopic growth factor in Beijing: a novel method for high relative humidity conditions, Atmos. Environ., 215, 116912, https://doi.org/10.1016/j.atmosenv.2019.116912, 2019.
- Zieger, P., Fierz-Schmidhauser, R., Gysel, M., Ström, J., Henne, S., ⁸⁵ Yttri, K. E., Baltensperger, U., and Weingartner, E.: Effects of relative humidity on aerosol light scattering in the Arctic, Atmos. Chem. Phys., 10, 3875–3890, https://doi.org/10.5194/acp-10-3875-2010, 2010.
- Zieger, P., Fierz-Schmidhauser, R., Weingartner, E., and Baltensperger, U.: Effects of relative humidity on aerosol light scattering: results from different European sites, Atmos. Chem. Phys., 13, 10609–10631, https://doi.org/10.5194/acp-13-10609-2013, 2013.
- Zieger, P., Fierz-Schmidhauser, R., Poulain, L., Müller, T. J., Birmili, W., Spindler, G., Wiedensohler, A., Baltensperger, U., and Weingartner, E.: Influence of water uptake on the aerosol particle light-scattering coefficients of the Central European aerosol, Tellus B, 66, 22716, https://doi.org/10.3402/tellusb.v66.22716, 2014.
- Zieger, P., Aalto, P. P., Aaltonen, V., Äijälä, M., Backman, J., Hong, J., Komppula, M., Krejci, R., Laborde, M., Lampilahti, J., de Leeuw, G., Pfüller, A., Rosati, B., Tesche, M., Tunved, P., Väänänen, R., and Petäjä, T.: Low hygroscopic scattering enhancement of boreal aerosol and the implications for a columnar optical closure study, Atmos. Chem. Phys., 15, 7247–7267, https://doi.org/10.5194/acp-15-7247-2015, 2015.
- Zou, J., Yang, S., Hu, B., Liu, Z., Gao, W., Xu, H., Du, C., Wei, J., Ma, Y., Ji, D., and Wang, Y.: A closure study of aerosol optical properties as a function of RH using a *κ*-AMS- ¹¹⁰ BC-Mie model in Beijing, China, Atmos. Environ., 197, 1–13, https://doi.org/10.1016/j.atmosenv.2018.10.015, 2019.

Remarks from the language copy-editor

- Please verify. It is our house standard to list affiliations from the smallest to largest unit, e.g., department before faculty. Is the ordering of Affiliation 2 correct?
- Like CE1, please verify the order from smallest to largest unit.
- Please verify the changes to this affiliation.
- In keeping with the dominant style of the manuscript, no serial (Oxford) commas were removed.
- Please verify the time zone.
- Please verify. When possible, all authors should be included by initials in this section. Thank you.

Remarks from the typesetter

- **TSI** The composition of Figs. 1–4, and 7–9 has been adjusted to our standards.
- TS2 Please confirm.
- TS3 Please add last access date.
- TS4 Please add last access date.
- TS5 Please confirm.
- **TSG** Please provide a reference list entry including creators, title, and date of last access.

Please send a new supplement as a *.pdf without the title, authors, correspondence author, etc. as we will generate a supplement title page during publication (with a citation including the DOI), which will contain this information.

Please note that there is funding information given in the acknowledgements but you have not indicated any funding upon manuscript registration. Therefore, we were not able to complete the financial support statement. Please provide the missing information and double-check your acknowledgements to see whether repeated information can be removed from the acknowledgement. Thanks.

TS9 Please add total pages.

TSIO Please add total pages.