

## Reply to Reviewer #1

This paper analyses CO<sub>2</sub> (and CO) flux anomalies from the Southeast Asia region, using a data assimilation method for estimation of CO<sub>2</sub> fluxes at transport model grid resolution. They have used unique observational datasets and a model framework. The region has already been widely studied by various groups and this paper would be a nice addition to the growing list of literature. My judgement is that the authors demonstrate a capacity that would be utilised in near-real-time in the future or on regular basis. The paper is generally well written and can be accepted for publication after a major revision, in my opinion. Please see my specific comments below for revisions, which I hope are useful for you to consider.

We really appreciate the reviewer for taking valuable time to review our paper and giving us useful comments and suggestions. Described below are our replies to the reviewer's comments with page and line numbers of the revised manuscript. The supplementary manuscript shows changes from the previous one.

line 33: should you say 2nd strongest - please check MEI or something like that?

We appreciate the reviewer for pointing it out. We rather intended it to be "one of the biggest", because the El Niño intensity differs with different indices. We modified it to "This was one of biggest El Niño events". **[Line 33]**

Section 2.1: Just to make sure, you haven't use any other observations apart from the CONTRAIL and VOS in this study. Could you be explicit please.

We appreciate the reviewer's suggestion for clarification. Accordingly, we added the following sentence in the beginning of Section 2.1. "In this inverse analysis, we only used the CONTRAIL and NIES VOS data, because they are predominant in the area we focused on. Here, we briefly describe those observations and further information can be found in the literatures cited therein." **[Lines 99-100]**

line 187: a bit of details or a reference is needed for the BVOC model here, which ones are considered etc. and how CO production is modelled.

First of all, we apologize for incorrect descriptions of the BVOC data of VISIT caused by misunderstanding among co-authors. We now realized that we did not consider the oxidation of BVOC, but only considered the direct emissions of CO from vegetation. Although we recognize that the oxidation of BVOC is important for simulating atmospheric CO, we think it would not significantly affect our results and conclusions in this study, because we performed the CO<sub>2</sub>-only inversion and focused only on biomass burnings. In this regard, we modified the descriptions of BVOC in the manuscript and replaced " $f_{CO,BVOC}$ " with " $f_{CO,vege}$ " in Fig. 2. In the VISIT simulation, CO emissions are estimated with the scheme of Guenther (1997) but using an emission factor by Tao and Jain (2005). The emission rates are estimated by light, temperature, vegetation leaf amount and seasonality, and the scheme has been calibrated with observational data.

We modified the last part of Section 2.2.2 as:

"In the model, the contribution of oxidation from biogenic volatile organic compounds (BVOCs) to CO is not considered yet, but direct CO emissions from vegetation are given at the earth's surface. Although the oxidations of CH<sub>4</sub> and BVOCs are significant sources of atmospheric CO, we treated

the former very simply and did not consider the latter. Therefore, we did not input CO observations to the inverse analysis. In the inversion, the biomass burning emissions of CO, which were predominant in Equatorial Asia, were modified along with those of CO<sub>2</sub>, as described in the next section.” **[Lines 167-171]**

Table 1: GPP, RE errors are quite large, which is nice. why is the error for fire emission is set to greater over the rest, compared to eq. asia?

In fact, the effective error of the prior fire is larger than 80%. This is because the posterior error correlations are considered in time (3 day), which introduces off-diagonal elements in **B** and inflates the errors in Equatorial Asia; that is not the case for the rest of the world. Error numbers summarized in Table 1 were determined after tens of preliminary experiments.

In order to elaborate the flux error settings, we modified the text as follows:

“Similarly, the errors for fossil fuel and fire emissions were introduced, but without error correlations.”

=>

“Similarly, 80% and 100% errors for fire emissions were introduced for Equatorial Asia and the rest of the world, respectively, but without spatial error correlations. Note that the fire errors for Equatorial Asia are practically larger than 80%, because the 3-day temporal correlation inflates the errors. We put a 10% error on the fossil fuel emissions in Equatorial Asia.” **[Lines 218-221]**

line 221: CDIAC data are used for what time period ?

We added the sentence describing how we made the fossil fuel data for 2015.

“Here, the fossil fuel emission data for 2015 were produced from the latest gridded CDIAC data for 2013 by scaling with the global total value for 2015 that is preliminarily reported by Le Quéré et al. (2015).” **[Lines 226-228]**

line 241: how were they derived ? did you model BVOC oxidation ?

Please see our reply to your query to line 187. For the CO emissions of VISIT, we added sentences as follows.

“In the VISIT simulation, CO emissions are estimated with the scheme of Guenther (1997), but using an emission factor by Tao and Jain (2005). The emission rates are estimated by light, temperature, vegetation leaf amount and seasonality, and the scheme has been calibrated with observational data.” **[Lines 250-253]**

line 258: should work in principle because you are starting with an inversion flux, but not sure 2-month of spin-up is sufficient

We consider that the two-month is enough for the spin-up, because the initial mole fraction fields were prepared also by the inversion flux of Niwa et al. (2012). To make it clearer, we added the sentences below.

“Nevertheless, the initial mole fraction fields are consistent with observations to some extent, as they were prepared by the inversion flux of Niwa et al. (2012). This makes the two-month inversion spin-up reasonable.” **[Lines 270-272]**

Figure 3: Are these plots better shown by "Model - Observation" for the lower 2 panels?

We agree with the suggestion and modified Fig. 3 and its caption accordingly.

Figure 4: I wonder if you really need Figure 4

We think Figure 4 is useful for demonstrating probable contribution of the fire emissions to the CO<sub>2</sub> mole fractions observed by CONTRAIL. As shown by Figs. 3 and 4, the observed CO<sub>2</sub> do not show fire-related variations so clearly, Figs 4b and 4c help us to imagine how large the fire contributions are.

Figure 6: Why not show the CO<sub>2</sub> time series as well, as that is the main focus of the paper?

As pointed out, we did not show the CO<sub>2</sub> time series in Fig. 6. That is because we have already shown both the CO<sub>2</sub> and CO observations of NIES VOS in Fig. 5 and from those we can easily recognize the two species are correlated with each other. The aim of Fig. 6 is to investigate if those correlated mole fraction peaks are contributed by fires or not, and, if so, which fire (Sumatra or Borneo) contributed to them. However, we realized that the data ranges of Figs. 5 and 6 were not exactly the same; that of Fig. 5 showed a slightly smaller range (up to 12°N and 120°E) than that of Fig. 6 (up to 15°N and 125°E). Therefore, we expanded the area of Fig. 5 slightly so that its data range is consistent with that of Fig. 6. Furthermore, we added the text “These time series depict the data limited within the range of 95°-125°E and 10°S-15°N consistently with Figs. 5b and d.” in the caption of Fig. 6.

line 343: Delete "We do not consider CH<sub>4</sub>...". I think, none will ask for CH<sub>4</sub> at this point.

Accordingly, we deleted it.

line 352: since you are putting less prior flux uncertainties outside the domain of this analysis, this is bound to happen. do you have another simulation with uniform treatment of prior flux uncertainty ?

As pointed out, this result has some dependency on the prior uncertainties, and actually we put less uncertainties, i.e. strong constraint, on prior fluxes in the outside region. However, their flux contributions to atmospheric CO<sub>2</sub> mole fractions could be big because the area is significantly larger than the target region. Therefore, we do not consider that the small flux uncertainty of the outside region does not necessarily induce the large fire emission estimate in the target region. Furthermore, we tested no fire-emission-based uncertainty, which is almost similar to the case of “uniform treatment of prior flux uncertainty”, and we obtained the similar number for the fire emissions in Equatorial Asia (this is written in the manuscript **[Lines 403-405]**).

line 365: One of the problems here is the over interpretations of the flux differences, without giving a range of uncertainty e.g., how significant are 31 Tg vs 2.7 Tg mentioned

We consider that these numbers between the prior and posterior fluxes are not considerably large. In order to clarify this, we modified the text as follows.

“This small change in October indicates that the simulated mole fractions from the prior flux were well consistent with the CONTRAIL observations.”

=>

“In fact, these flux changes are small compared with the differences between GFED and GFAS (GFED minus GFAS is 92 Tg C and -87 Tg for September and October, respectively), indicating that the simulated mole fractions from the prior flux of C\_GG were overall consistent with the CONTRAIL observations.” **[Lines 390-392]**

Figure 8 & 9: please mention in the caption if the prior flux map is same for both the inversions (C\_GG & C\_NO). and also show the posterior - prior for the C\_NO case. hopefully you will find a place to show the colour bar (on the right?)

According to the suggestion, we added the text

“Note that the prior estimate of (a) was used both for C\_GG and C\_NO, while the prior fire estimate of (d) was used only for C\_GG.”

in the caption of Fig. 8. Furthermore, we modified Figs. 8 and 9 to include the posterior – prior for the C\_NO case.

line 378: Is this statement true? see over the Papua New Guinea or the southern part of Borneo, where are there are large differences in patterns

As pointed out, the pattern is different for Papua New Guinea, but we consider that the pattern is similar to each other for southeast Sumatra and south of Borneo, where the fire was most intense. To limit the statement valid only for the Sumatra and Borne regions, we added text as follows.

“... are well coincident with those of prior fire emissions in southeast Sumatra and south of Borneo.” **[Line 403]**

Figure 11: Nicer to have a common x-axis for these plots for easier reading. line 410: not clear what you mean by "its degree"?

We revised Fig. 11 accordingly. Besides, we changed “its degree” to “the sensitivity” for clarification. **[Line 435]** Please note that footprint quantifies a sensitivity of an observation against surface fluxes.

Figure 12: A bit surprised to see posterior RMSDs are quite often exceeding the prior values for October

Because the NIES VOS is independent of the inversions except for CV\_GG, the RMSD should not be necessarily reduced. This is probably because some representative errors in surface fluxes or atmospheric transport were larger in October than in September, which can be inferred from the larger RMSDs and the smaller correlations. We added this discussion in the revised manuscript as follows.

“Meanwhile, the RMSDs of both CO<sub>2</sub> and CO were not necessarily reduced for October. This is attributable to insufficient representativeness of surface fluxes or atmospheric transport, which can be inferred from the larger RMSDs and the smaller correlations in October than those of September. Nevertheless, the experiment without the prior fire emission (C\_NO) exhibited smaller RMSDs with

the posterior CO<sub>2</sub> and CO fluxes for both September and October. Furthermore, the improvement in the correlation coefficients is remarkable...” **[Lines 461-466]**

Figure 13 and associated text: I am not very sure of usefulness of this discussion. You need to take in to account the ageing of the air mass, i.e., the influence of near and far fields in constructing the emission ratios. It is possible to implement such an analysis using trajectory model for instance.

We did not intend to infer the absolute value of the emission ratio by Fig. 13. As the reviewer says, estimating the absolute value of the emission ratio is difficult due to the aging of the airmass. However, we believe that we could infer if the emission ratio prescribed in the flux data is larger than real or not, by comparing the simulated ratios with those of the observation, in which the aging of the airmass is fairly considered by the atmospheric transport model.

Section 4: not sure if you need these detailed discussions, a table of fire emission estimation can be given in the results section or here for improving readability. Then some of the text, say on the modelling limitations, could be integrated with the conclusions or new subsection in Results

Following the reviewer’s suggestion, we modified the Discussion section. Particularly, we moved a part of the first paragraph to Section 3.2.1 that discusses the obtained fire emissions as shown by Table 3 comparing with other study results. **[Lines 369-375]** Furthermore, we also moved the sentence “with the help of NIES VOS observations, especially its CO data, we demonstrated the validity of our inverse analysis” to the Conclusion section. **[Lines 568-569]**

Meanwhile, we keep the discussion of the modeling limitations, because it is not only the output of the inversion but also includes some perspectives, which we consider suitable for Discussion.

Section 5: Conclusions can focus on the outcome of this paper, with minimal or no referencing to old papers. Some can be kept for the significance of the paper, e.g., the ENSO related flux variability, which has been discussed in the results section.

We have tried to reduce the volume of this section. First, we deleted the sentence referring Niwa et al. (2012). Furthermore, we reduced the last paragraph by moving the sentence “Fields et al. (2016) pointed out ... for 2018 (GIO and MOE, 2020)” to Section 3.2.1 **[Lines 375-378]**.