

Evaluation of the CMIP6 marine subtropical stratocumulus cloud albedo and its controlling factors

Bida Jian¹, Jiming Li^{1*}, Guoyin Wang², Yuxin Zhao¹, Yarong Li¹, Jing Wang¹, Min Zhang³, and Jianping Huang¹

5 ¹ Key Laboratory for Semi-Arid Climate Change of the Ministry of Education, College of Atmospheric Sciences, Lanzhou University, Lanzhou, Gansu, China

² Department of Atmospheric and Oceanic Sciences & Institute of Atmospheric Sciences, Fudan University, Shanghai, China

³ Inner Mongolia Institute of Meteorological Sciences, Hohhot, Inner Mongolia, China

10 *Correspondence to:* Jiming Li (lijiming@lzu.edu.cn)

Abstract. The cloud albedo at the ~~subtropical~~ marine subtropical stratocumulus regions ~~has-plays~~ a key role in regulating the regional energy budget. Based on 12 years of monthly data from multiple satellite datasets, the long-term, monthly and seasonal cycle ~~of~~ averaged cloud albedo at five stratocumulus regions were investigated to inter-compare the atmosphere-only simulations ~~of-between~~ Phase 5 and 6 of the Coupled Model Inter-comparison Project (AMIP5 and AMIP6). Statistical results showed that the long-term regressed cloud albedos were underestimated in most AMIP6 models compared with the satellite-driven cloud albedos, and the AMIP6 models produced a similar spread of AMIP5 ~~at-over~~ all regions. The monthly ~~averaged values~~ ~~mean~~ and seasonal cycle of cloud albedo of AMIP6 ensemble mean showed ~~a~~ better correlation with the satellite-driven observation than that of AMIP5 ensemble mean, ~~-, however~~ ~~However, the AMIP6 model still~~ failed to reproduce the values and amplitude in some regions. By employing the Modern-Era Retrospective Analysis for Research and Applications Version 2 data, this study estimated the relative contributions of different aerosols and meteorological factors on the ~~long-term variation of~~ marine stratocumulus cloud albedo under different cloud liquid water path (LWP) conditions. The multiple regression models can explain ~~~60-65~~ % of the changes in the cloud albedo. Under the monthly mean LWP ≤ 6065 g m⁻², dust and black carbon dominantly contributed to the changes in the cloud albedo, while ~~dust and sulfatesulfur dioxide~~ aerosol contributed the most under the condition of 6065 g m⁻² < LWP ≤ 120 g m⁻². These results suggest that the parameterization of cloud-aerosol interactions is ~~critical-crucial~~ for accurately simulating the cloud albedo in ~~climate~~ models.

15
20
25

1 Introduction

One of the critical parameters in regulating the distribution of solar radiation in the atmosphere and surface is cloud albedo, ~~that-which~~ is the proportion of incoming solar radiation reflected by clouds (Mueller et al., 2011; Wall et al., 2018). A change in the cloud albedo over low-level clouds can cause a significant alteration in the planetary albedo (Engström et al., 2014) and even could offset the warming caused by doubled carbon dioxide (Latham et al., 2008). Recent studies employing the cloud-system-resolving and plume models have shown that changes in the cloud albedo are largely dependent on aerosol

30

and meteorological conditions (Wang et al., 2011; Stuart et al., 2013; Kravitz et al., 2014). However, there are still non-neglectable uncertainties in simulations (Bender et al., 2016).

35 This study specifically focused on the cloud albedo in the subtropical marine stratocumulus regions as it is particularly difficult to reproduce the cloud properties by numerical models (Eyring et al., 2016), which results in a larger uncertainty in energy budget simulations and climate predictions (Wood, 2012). The subtropical marine stratocumulus regions are mainly covered by low-level clouds that usually reflect most of the solar radiation and significantly contribute to the planetary albedo (Seethala et al., 2015). In addition, the contribution of the cloud albedo to planetary albedo over these dark oceans could be tremendous compared with those ~~of-over~~ snow/ice-covered regions with a high surface albedo (Mueller et al., 2011).
40 However, it is a challenge to accurately estimate the cloud albedo in regions where there are different types of clouds for evaluating the cloud albedo resulted from the relationship between the planetary albedo and cloud fractions at a monthly scale (Bender et al., 2011; Bender et al., 2019).

To date, climate models have continuously advanced in main physical processes, model structures and initial conditions to improve the capability to reproduce numerous observed climate events (Van Weverberg et al., 2017; [Huang et al., 2018](#)).
45 Many studies have paid attention to understand~~ing~~ the cloud albedo and its controlling factors over the subtropical marine stratocumulus regions for reducing the uncertainty in models' outputs (Latham et al., 2008; Wood, 2012; Engström et al., 2014; Bender et al., 2019). The cloud albedos obtained from regressing satellite observations at five typical subtropical marine stratocumulus regions exhibited distinct characteristics, ranging from 0.32 to 0.39, and noticeable diurnal variations (Bender et al., 2011; Engström et al., 2014), which may be induced by respective aerosols and meteorological conditions at
50 each region. For example, the southeast Atlantic stratocumulus region (Namibian) is a typical region with massive biomass burning aerosols loading (Wilcox, 2010) while a dominant aerosol loading type in the Canarian region is dust (Waquet et al., 2013). As the value of cloud albedo is usually determined by the cloud optical thickness (COT) and the solar zenith angle (Wood, 2012), the main factors (i.e., the cloud droplets number and sizes) controlling the COT may affect changes in the cloud albedo (George and Wood, 2010; [Xie et al., 2013](#); Bender et al., 2016). ~~Since-Further,~~ the cloud droplets amount and
55 the droplets sizes are affected by cloud condensation nuclei (CCN) [and cloud liquid water content \(Zhao et al., 2012\)](#), it is crucial to understand the interactions in key dynamical and microphysical processes controlling CCN with regard to improving the model capacity to simulate the cloud albedo (Bender et al., 2016; Rosenfeld et al., 2019).

Regarding the microphysical processes, the aerosol-cloud-radiation interactions over the subtropical marine stratocumulus regions have been actively examined in previous studies ([Twomey, 1974, 1977](#); Wang et al., 2011; Bender et al., 2016, 2019; [Zhao et al., 2018](#)).
60 Among them, some studies have demonstrated the effect of aerosols on the marine stratocumulus cloud albedo (Twomey effect). In other words, an increase in aerosols can result in smaller droplet sizes and more droplets, leading to a higher cloud albedo (Twomey, 1974, 1977). However, the cloud-aerosol interactions are complex and varying with aerosol types due to their different effects on clouds. Unfortunately, the Intergovernmental Panel on Climate Change currently lacks confidence in estimating the global aerosol indirect effects (Boucher et al., 2013). Furthermore, the semi-
65 direct effects of absorbing aerosols (e.g., black carbon) are also difficult to be quantified by numerical models (Herbert et al.,

2020). Given different model experiments from the Coupled Model Intercomparison Project phase 5 (CMIP5), Frey et al. (2017) estimated the impact of anthropogenic sulfate and non-sulfate aerosol forcing on changing the cloud albedo and concluded that absorbing aerosols play a key role in offsetting the cloud brightening at a certain degree. Regarding the dynamical processes, previous studies found that the dynamical factors (e.g., vertical velocity or instability) can influence not only the vapor supersaturation, leading to the activation of CCN (Twomey, 1959; Lu et al., 2012; Rosenfeld et al., 2019), but also the cloud droplet number and sizes, by the entrainment and detrainment of air above the clouds (Fuchs et al., 2018; Yang et al., 2019; Scott et al., 2020). Based on satellite observations, Chen et al. (2014) investigated the effects of aerosols on marine warm clouds, they found that the response of LWP to aerosol loading strongly depends on lower tropospheric stability and free-tropospheric moisture. Besides, the free-troposphere relative humidity is also considered as a critical factor in regulating the cloud albedo, because it is closely related to the cloud-top entrainment/drying process that influences the cloud albedo effect (Betts and Ridgway, 1989).

However, most of these studies are based on rapid cloud adjustment to study the effects of specific meteorological factors, or aerosol-cloud interactions over the marine subtropical stratocumulus regions. Few systematic studies focus on the effects of meteorological factors and various aerosol types on the cloud albedo and changes at the monthly scale. Furthermore, it is also crucial to evaluate the performance of current climate models to accurately project the cloud albedo responses to climate change. By the intercomparison of outputs between CMIP3 and CMIP5, Engström et al. (2014) found that the regressed regional averaged cloud albedo and intermodal spread of CMIP5 in the subtropical marine stratocumulus regions are more comparable with the satellite observations compared with those of CMIP3. Given the release of up-to-date CMIP6, as in the previous study, it is necessary to systematically evaluate the performance of CMIP5 and CMIP6 in reproducing the cloud albedo for understanding advances in the skill of climate models to resolve long-standing problems in the marine stratocumulus regions. Based on multiple satellite datasets, this study evaluated the performance of ten CMIP5/AMIP and twenty-eight CMIP6/AMIP outputs. As an essential part of CMIP experiments, the AMIP outputs forced by observed sea surface temperatures (SST) and sea ice concentrations (Eyring et al., 2016) were used in the study. By employing the reanalysis data, this study quantitatively estimated the contributions of each factor to the marine stratocumulus cloud albedo to identify main factors dominating the long-term variations of the marine stratocumulus cloud albedo. This study will provide useful information for comprehensively understanding the impacts of different aerosol types and meteorological factors on cloud albedo changes.

The article is organized as follows. The datasets and methods are given in section 2. The comparison of performances between CMIP5 and CMIP6 is presented in Section 3.2.1. The impacts of different aerosol types and meteorological factors on the cloud albedo are described in Section 3.2. Lastly, section 4 addresses the conclusions and discussion.

2 Datasets and Method

This study compiled multiple satellite datasets, 10 CMIP5/AMIP outputs, 28 CMIP6/AMIP outputs and reanalysis data not only to evaluate the performance of CMIP5 and CMIP6 outputs but also to investigate the variations of the cloud albedo over the typical subtropical marine stratocumulus regions. Since spatial resolutions vary with climate models, all data were interpolated to a $1.0^\circ \times 1.0^\circ$ spatial resolution and monthly temporal resolution for fairly evaluating and intercomparing the performance. The following sections ~~address~~ provide more details on the satellite datasets, CMIP5, CMIP6, and reanalysis data.

2.1 CERES and MODIS

Estimating the cloud albedo ~~is required~~ requires multiple atmospheric variables such as the top-of-atmosphere (TOA) downward, upward (all-sky) shortwave fluxes, cloud liquid water path (LWP) and cloud fractions. In this study, the TOA downward and upward shortwave fluxes were obtained from the Clouds and the Earth's Radiant Energy System (CERES, Wielicki et al., 1996) Single Scanner Footprint (SSF) monthly Ed4A datasets. The LWP and cloud fractions were ~~also~~ obtained from the Moderate Resolution Imaging Spectro-Radiometer (MODIS; Platnick et al., 2003) collection 6.1 level 3 monthly products during the period from 2003 to 2014, i.e., MODIS MYD08_M3 (Aqua) and MOD08_M3 (Terra) products, respectively. The spatial resolutions of these products are $1.0^\circ \times 1.0^\circ$. The CERES TOA shortwave fluxes were converted from broadband (0.2-5.0 μ m) radiances by applying empirical angular distribution models to correct the instrument's incomplete spectral response (Loeb et al., 2001). Then, the real-time fluxes were aggregated to produce 24-hour mean fluxes from empirical diurnal albedo models that create meteorology conditions at the over-flight time (Loeb et al., 2018). It is worth noting that the aforementioned data processing may introduce some potential uncertainties, e.g., diurnal correction error, radiance-to-flux conversion error (one standard deviation, 1σ) and instrument calibration error (1σ). For example, the uncertainty in the monthly combined regional all-sky shortwave flux was 6.2 W m^{-2} (CERES SSF1deg-Hour/Day/Month_Ed4A Data Quality Summary, 2021), where the calculation of the diurnal correction uncertainty was driven by comparisons with Geostationary Earth Radiation Budget data (Doelling et al., 2013). In addition, the cloud fraction, a fraction of MODIS cloudy pixels to the total pixels at each grid box, is determined based on daytime scenes entirely and represents all cloud phases (Platnick et al., 2003). As the CERES and MODIS instruments are both carried onboard Aqua (cross the equator local time: 1130) and Terra (pass the equator local time: 1030) satellites in polar orbits, we averaged the Aqua and Terra products to obtain the observed combined all-sky albedo, cloud fraction, LWP and cloud albedo as in the works of the Engström et al. (2015) and Bender et al. (2017). Time representation errors can be caused by the averaged cloud observations at two time points to represent the daily average. However, recent studies found that the time representation error was significant for short-term studies (up to 14%) while negligible for long-term statistical analysis (Wang and Zhao, 2017; Zhao et al. 2019a).

2.2 CMIP5/AMIP and CMIP6/AMIP

The outputs of 10 CMIP5/AMIP and 28 CMIP6/AMIP include all variables necessary to estimate the cloud albedo, i.e., monthly mean TOA downward, upward (all-sky) fluxes and total cloud fractions (Taylor et al., 2012; Eyring et al., 2016).

130 This study ~~identified–used~~ ten climate models that provide both CMIP5 and CMIP6 outputs and implemented the intercomparison of performance for the regressed cloud albedo during the period from 2003 to 2008. Furthermore, this study evaluated the cloud albedo for twenty-eight CMIP6/AMIP outputs during the period from 2003 to 2014. Tables 1-2 show the characteristics of CMIP5 and CMIP6 models. Note that there is a considerable discrepancy in the total cloud fractions between the CMIP models and MODIS observations, which is caused by different definitions, cloud overlap algorithms and
135 different threshold assumptions for cloud formation (Engström et al., 2015). Moreover, the total cloud fractions in the climate models are usually calculated based on daytime and nighttime cloud fractions, while the observed cloud fractions are only for the daytime. ~~while only daytime cloud fractions are considered to evaluate the observed cloud fractions.~~ As used in Engström et al. (2015), this study also employed the total cloud fractions as there are no available MODIS simulator outputs for CMIP6. Although uncertainty in cloud fraction remains, a previous study also demonstrated that the time representation
140 error was negligible for long-term statistical analysis (Wang and Zhao, 2017).

2.3 MERRA-2

The study employed the Modern-Era Retrospective Analysis for Research and Applications Version 2 (MERRA-2) which provides a long-term aerosol and atmospheric reanalysis record (1980-present) at $0.625^{\circ} \times 0.5^{\circ}$ resolution based on the Goddard Earth Observing System Model, version 5.12.4 (Gelaro et al., 2017). The aerosol reanalysis has been produced by a
145 global data assimilation system that combines satellite- and ground-based observed aerosols with meteorological conditions. Here, the mass mixing ratios of different aerosol types and air density at different levels from the 3-hourly aerosol product (inst3_3d_aer_Nv) and meteorological data from monthly atmosphere product (instM_3d_asm_Np and instM_2d_asm_Nx) ~~are–were~~ collected to represent the monthly regional aerosol and meteorological conditions. The outputs of MERRA-2 reanalysis were used during the common-consistent period from 2003 to 2014 with satellite observations record. As selected
150 in McCoy et al. (2017) and Li et al. (2018), the impacts of different aerosol types on marine stratocumulus cloud albedo were evaluated based on the mass concentrations of hydrophilic black carbon (BC), hydrophilic organic carbon (OC), sulfate aerosol (SO₄), sulfur dioxide (SO₂), the smallest particles dust (DU, i.e., 0.1-1 μm size) and sea salt (SS, 0.03-0.1 μm size) at the 910hPa level. The meteorological variables include the monthly vertical velocity at 900hPa and 700hPa (omega900 and omega700), surface pressure, relative humidity at 700hPa (RH700), air temperature, the eastward wind, the northward
155 wind and surface skin temperature data. In addition, the estimated inversion strength (EIS) and horizontal temperature advection at the surface (SSTadv) factors were also calculated. Finally, all of these meteorological parameters were used to investigate the meteorological effects on the cloud albedo. The units for aerosol mass concentrations, relative humidity, vertical velocity, EIS and SSTadv are kg m⁻³, %, Pa s⁻¹, K and K s⁻¹, respectively. ~~Furthermore, this study employed the~~

relative humidity at 850hPa (RH850) and vertical velocity at 900hPa (ω_{900}) factors to investigate the meteorological effects on the cloud albedo.

2.4 Methods

The planetary albedo (α) can be calculated mainly from the cloud fraction f (Bender et al., 2011) as expressed in Eq. (1):

$$\alpha = \alpha_{cloud}f + \alpha_{clear}(1-f) \quad (1)$$

where, α_{cloud} and α_{clear} denote the albedo under cloudy-sky and clear-sky conditions, respectively. For a given region where the cloud and surface type are homogeneous (i.e., constant α_{cloud} and α_{clear}), namely, a change in α should be driven by a change in the cloud fraction f . The cloud albedo can be estimated by the derivative of Eq. (1) as in Eq. (2):

$$\alpha_{cloud} = d\alpha / df + \alpha_{clear} \quad (2)$$

The invariable α_{cloud} and α_{clear} should be applied for the same cloud type and ocean region. In this light, as the works in Klein and Hartmann (1993), this study also analyzed ~~for~~ only five marine stratocumulus regions: Peruvian (10°S–20°S, 80°W–90°W; A1), Namibian (10°S–20°S, 0°E–10°E; A2), Californian (20°N–30°N, 120°W–130°W; A3), Australian (25°S–35°S, 95°E–105°E; A4) and Canarian (15°N–25°N, 25°W–35°W; A5). Previous study (Engström et al., 2014) has also demonstrated that there is a near-linear relationship between cloud cover and planetary albedo in these regions. Fig. 1 illustrates the locations of the above stratocumulus regions and the near-global distribution of combined planetary albedo averaged from Aqua and Terra during the common period from 2003 to 2014. Here, EIS is defined in Wood and Bretherton (2006):

$$EIS = LTS - \Gamma_m^{850} (Z_{700} - Z_{LCL}) \quad (3)$$

where the lower-tropospheric stability (LTS) is defined as the difference in potential temperature between 700 hPa and the surface, Γ_m^{850} is the moist-adiabatic lapse rate at 850 hPa, Z_{700} and Z_{LCL} is the height of the 700 hPa level and the lifting condensation level relative to the surface, respectively. As in Wood and Bretherton (2006), we assumed the surface relative humidity of 80 % to simplify the calculation of surface dew point temperature. Z_{LCL} was calculated based on the method of Georgakakos and Bras (1984). In addition, SSTadv was obtained by Eq. (4) as in Qu et al., (2015):

$$SSTadv = - \frac{u}{R_E \cos \phi} \frac{\partial SST}{\partial \lambda} - \frac{v}{R_E} \frac{\partial SST}{\partial \phi} \quad (4)$$

where u and v represent the eastward and northward horizontal wind components at 1000hPa, respectively. ϕ and λ are latitude and longitude, respectively. R_E is the mean Earth radius and SST is the surface skin temperature. A positive/negative SSTadv indicates warm/cold advection. The SSTadv can affect the moisture transport within the cloud layer by influencing the surface sensible and latent heat fluxes, and consequently, influence the thickness of marine stratocumulus clouds (George and Wood, 2010).

In the study, to avoid influence from a seasonal cycle, the annual-long-term mean analyses are implemented with deseasonalized monthly mean data processed by removing a mean seasonal cycle, and then adding the monthly mean value to the interannual anomalies data.

The selection of variables is a crucial step to build a multiple linear regression model ~~for~~of the monthly cloud albedo as a function of meteorological factors and aerosol types under two different LWP scenarios (LWP < ~~6065~~ g m⁻² and ~~6065~~ g m⁻² < LWP < 120 g m⁻²). This study selected suitable variables based on correlation analysis. If the correlation between the cloud albedo and a candidate is significant at a 90% confidence level, the variable was considered as a predictor factor.

195 Furthermore, the partial least squares were used to reduce the collinearity between the selected variables (McCoy et al., 2017). The regression model of cloud albedo α_{cloud} is as follows:

$$\alpha_{cloud} = \sum_{i=1}^I a_i M_i + \sum_{j=1}^J b_j \log_{10} A_j + c \quad (35)$$

where a and b are regression coefficients, c is a constant term, M_i represents the i th meteorological predictor, I is the number of meteorological predictor variables, A_j is the j th aerosols predictor, and J is the number of aerosol predictor variables.

The relative contributions of each predictor to the change in the cloud albedo (Huang and Yi, 1991) were evaluated using Eq. (46):

$$R_j = \frac{1}{m} \sum_{i=1}^m [T_{ij}^2 / (\sum_{j=1}^a T_{ij}^2)] \quad (46)$$

where m is the number of the monthly samples, a is the number of predictors, and T_{ij} is the product of the regression coefficients of each term (b_j) and predictor variables (x_{ij}).

After removing the effect of meteorological factors, we further investigated the pure relationship between aerosols and the cloud albedo using the partial correlations between α_{cloud} and $\log_{10}A$, as expressed in Eq. (57):

$$r_{\alpha_{cloud} \log_{10} A \cdot M} = \frac{r_{\alpha_{cloud} \cdot \log_{10} A} - r_{\alpha_{cloud} \cdot M} r_{\log_{10} A \cdot M}}{\sqrt{1 - r_{\alpha_{cloud} \cdot M}^2} \sqrt{1 - r_{\log_{10} A \cdot M}^2}} \quad (57)$$

210 where $r_{\alpha_{cloud} \cdot \log_{10} A}$, $r_{\alpha_{cloud} \cdot M}$ and $r_{\log_{10} A \cdot M}$ is the total correlation between each variable pair and $r_{\alpha_{cloud} \log_{10} A \cdot M}$ is the correlation between α_{cloud} and $\log_{10}A$ which eliminates the effects of meteorological factors M . More details on the partial correlation are described in Jiang et al. (2018) and Engström and Ekman (2010).

3 Results

3.1 Satellite observations and CMIP5/6 simulations

215 The first two columns in Fig. 2 (from a to e) show the estimated long-term mean cloud albedo corresponding to the correlation between planetary albedo and cloud fraction over the five regions from the observation and 22 AMIP5/6 models, including 10 individual models and an ensemble mean for AMIP5 and AMIP6 (represented by AMIP5-MEM and AMIP6-MEM), during the period from 2003 to 2008. For the combined satellite observations, the correlation coefficient values are above 0.85 in all regions. The correlation over the Peruvian region was the largest (~0.95), while a relatively weak correlation (~0.88) appeared in the Canarian region. Such a high correlation between planetary albedo and cloud fraction

further indicates the homogeneity of cloud and surface types over these regions. The regressed cloud albedo from the satellite ranged from 0.30 to 0.42 for the five stratocumulus regions, which is consistent with previous studies (Bender et al., 2011; Engström et al., 2014). As the values averaged over Aqua and Terra albedos and cloud fractions were used as the observation in this study, the regressed cloud albedo values need to be within the range of the Aqua and Terra (Engström et al., 2014). Regarding the AMIP5 and AMIP6 models, a higher correlation (> 0.8) appeared for most models at the five regions, especially higher at the Australian and Canarian regions. At the Peruvian, Namibian and Californian regions, the correlations of the observation were relatively higher than those of most climate models while the observed correlation was approximately close to the median value of model simulations at the Australian and Canarian regions.

Although previous studies indicated that some CMIP6 models updated the cloud physical parameterization in the new version (Seland et al., 2020; Kawai et al., 2020). The correlation coefficients of the AMIP6 models between planetary albedo and cloud fraction showed a lower value than those of the AMIP5, indicating that the linear relationship between cloud fraction and planetary albedo in the AMIP6 models' simulations is not superior to that of AMIP5. While the AMIP6 simulations displayed a similar spread in the estimated cloud albedo for all regions, some AMIP6 models produced a lower correlation coefficient than those of the AMIP5 models (e.g., AMIP6/INM-CM4-8). Notably, the AMIP5-MEM and AMIP6-MEM always produced a worse correlation relationship and more irrational cloud albedo values, indicating that the AMIP5/AMIP6 models have a lack of skill in simulating cloud properties over the marine stratocumulus regions. ~~Note that the performance of NorESM2-LM is superior to that of the previous version of the model (NorESM1-M) by improving the cloud physical parameterization, e.g., the stratiform cloud parameterization that causes underestimation of the cloud fraction and overestimation of the cloud liquid water in NorESM1-M (Seland et al., 2020). This advance in NorESM2-LM may induce the more reasonable representation of the cloud albedo in the stratocumulus regions. Kawai et al. (2020) also concluded that the MRI-ESM2 employed a new stratocumulus scheme to improve the representation of low clouds from the previous version of the model (MRI-CGCM3). However, the cloud albedo simulated from MRI-ESM2 showed a significant improvement only in the Canarian region compared with that of the MRI-CGCM3 simulation.~~

The third and fourth columns in Fig. 2 (from f to j) also shows the estimated long-term mean cloud albedo and the correlation between planetary albedo and cloud fraction over the five regions for the observation and 29 AMIP6 models during 2003 to 2014. The simulated correlation exhibited a larger spread at the Peruvian and Namibian regions than those at other regions, indicating that the AMIP6 models have a lack of capacity to capture the linear relationship between planetary albedo and cloud fraction. The cloud albedos were underestimated in most CMIP6/AMIP models compared with the satellite-based cloud albedos. The Australian (0.30~0.43) and Canarian (0.24~0.42) regions displayed a larger inter-model variability in the cloud albedo than other regions due to a poor skill in simulating the cloud properties (e.g., LWP and COT). Over the Canarian regions, the correlation and cloud albedo of AMIP6-MEM showed good agreement with those of the satellite observation compared with those of the individual AMIP6 models, resulting from the offsetting effect between models.

255 ~~The ACCESS-ESM1-5 and CAMS-CSM1-0 models produce a higher correlation coefficient and more accurate cloud albedo at the Peruvian region. The MPI-ESM1-2-HR, NorESM2-LM, CESM2-FV2 and IPSL-CM6A-LR models simulated the correlation and the cloud albedo comparable to the observations at the Namibian, Californian, Australian and Canarian regions, respectively. In contrast, the EC-Earth3, FGOALS-f3-L, FGOALS-f3-L, INM-CM5-0 and CESM2 models simulated the worst results at the Peruvian, Namibian, Californian, Australian and Canarian regions, respectively.~~ Overall, the AMIP6 models reproduced the cloud albedo and correlation well at the Australian region while a higher uncertainty in

260 model's simulations, i.e., a larger intermodal spread, at the Peruvian region (Engström et al., 2014). Engström et al. (2014) also found that CMIP5 models simulating a higher cloud cover have a tendency to produce a smaller cloud albedo value. Darker clouds can offset the contribution of the higher cloud cover to the planetary albedo, resulting in relatively a consistent model-driven planetary albedo. This is a presentation of the "too few, too bright" problem that persists in GCMs (Nam et al., 2012). To validate whether or not this problem has been improved in the AMIP6 models, we

265 compared the relationship between regressed cloud albedo and cloud fraction (See Fig. S1). The correlations driven by the 28 AMIP6 models were -0.28, 0.19, -0.11, -0.71 and 0.43 for the Peruvian, Namibian, Californian, Australian and Canarian regions, respectively. Compared with the results from the CMIP5 models (Engström et al., 2014), noticeable progress was found at the Namibian and Californian regions while a high negative correlation was simulated at the Australian region, indicating that the new generation models need to be further improved to resolve the longstanding problem.

270 The monthly cloud albedo time series regressed from the satellite, MEM and AMIP6-MEM for the six-year period from 2003 to 2008 over the five regions are shown in Fig. 3(a-e). The temporal correlations (R5/R6) and corresponding confidence value (P5/P6) between simulated (AMIP5-MEM/AMIP6-MEM) and satellite regressed monthly cloud albedo time series are also given in Fig. 3(a-e). Note that the smoothed time series were produced by 12-month smoothing. The statistical results showed that the R5/R6 values were 0.62/0.78, 0.44/0.55, 0.38/0.45, 0.75/0.74 and 0.00/0.05 for the

275 Peruvian, Namibian, Californian, Australian and Canarian regions, respectively. Among them, the correlations only at the Canarian region were insignificant (i.e., P5/P6=1.00/0.70). A high positive correlation appeared at the Australian region (>0.70), indicating that the changes in the cloud albedo are well captured by the models.

Compared with AMIP5-MEM, the regressed monthly cloud albedo of AMIP6-MEM showed a better correlation with the satellite regressed values, ~~while poor performance in reproducing the monthly cloud albedo values and amplitude (the~~

280 ~~difference between the maximum and minimum values of cloud albedo) at the Peruvian and Namibian regions. However, the performance of AMIP5-MEM in reproducing monthly cloud albedo and its amplitude (the difference between the maximum and minimum values of cloud albedo) was better than that of AMIP6-MEM.~~ Furthermore, the monthly cloud albedos obtained from the satellite and models displayed obvious seasonal cycle at all regions except for the Canarian region. This may be related to the fact that a weaker linear relationship between monthly cloud cover and planetary albedo may exist at

285 the Canarian region, resulting in a significant change in the estimated cloud albedo (see Fig. S2). ~~Overall, the performance of AMIP5-MEM in reproducing monthly cloud albedo and amplitude was better than that of AMIP6-MEM.~~

In addition, ~~The the~~ monthly cloud albedo time series for the satellite and AMIP6-MEM for the period from 2003 to 2014 in the five regions are also shown in Fig. 3(f-j), ~~which are consistent with Fig. 3(a-e), indicating that the simulation capability of the AMIP6-MEM in different regions is not improve significantly with the expansion of the simulation time and the increase of the model numbers.~~ The cloud albedos driven by the satellite and model at the Peruvian region ranged from 0.35 to 0.45 and from 0.25 to 0.38, respectively (Fig. 3f). ~~The monthly cloud albedo values of AMIP6-MEM were significantly underestimated compared with those at other regions. In contrast, the monthly cloud albedos of the models showed good agreement with the satellite-based observation at the Namibian and Australian regions, which means that AMIP6-MEM captures the observed seasonal cycle of the cloud albedo well.~~ However, ~~t~~ The amplitudes of the cloud albedo simulated from the model were larger than that of the satellite at the Peruvian, Namibian and Californian regions while smaller at the Australian and Canarian regions. Note that at the Australian region, the monthly cloud albedo exhibited a large variation than that at other regions based on the satellite-based observation, which means that the cloud optical properties (e.g., COT and cloud effective radius) have been considerably changed within the Australian region.

This study further assessed the performance of the AMIP6 models in reproducing the cloud albedo time series. Figs. 4a-e provide the Taylor diagrams (Taylor, 2001) for the five regions, which include the correlation coefficients, the centered root mean square error (RMSE, the green circle), and the standard deviation value between individual AMIP6 models and the satellite-based observations. The centered RMSE and the standard deviation values represent the model's ability to reproduce the phase and amplitude of the variable, respectively. Correlation coefficients greatly varied with regions, ranging from negative (Peruvian, Namibian and Canarian) and positive values. Compared with other regions, most of the models showed a high positive correlation (>0.6) at the Peruvian region ~~while the monthly cloud albedos simulated from the six GCMs (FGOALS-g3, FGOALS-f3-L, INM-CM4-8, INM-CM5-0, MIROC6 and GISS-E2-1-G) showed a negative correlation with the observation.~~ The model-driven cloud albedo was most poorly correlated with the observation at the Canarian region, e.g., < 0.4 or negative values. On the contrary, at the Australia region, all models showed a significant positive correlation (> 0.4). The standard deviation values of the models at the Peruvian, Namibian and California ranged 0.02-0.09, 0.02-0.11 and 0.03-0.10, respectively while 0.03 for the satellite-based observation. This result indicates that most of the models overestimate the amplitude of the cloud albedo time series at the regions. Some models produced the standard deviation values of cloud albedos three times larger than the observation. It ~~is is~~ evident that the standard deviation values of the simulated cloud albedo at the Australian regions ~~are a narrower spread were closer to the observed value~~ than that of other regions, indicating that the AMIP6 models also perform well in simulating the amplitude of the monthly cloud albedo time series at this region. Overall, the intermodal variability in the correlation coefficient, RMSE and standard deviation values was the smallest at the Australian region while the largest at the Peruvian region. ~~Among all the models, the ACCESS-ESM1-5 model outperformed other models at all regions. In contrast, the FGOALS-g3-L and INM-CM5-0 models performed poorly in reproducing the monthly cloud albedo in most regions. These results may be induced by models' capacity in reproducing the linear relationship between cloud cover and planetary albedo for these regions (See Fig. S3).~~

320 Further, Figure 5 shows the annual cycles of the cloud albedo estimated by the satellite and AMIP5/AMIP6 models for the five regions. The seasonal variation in the cloud albedo at each region takes a shape of single peak distribution. In terms of similarity among regions, the cloud albedo at all regions reached the maximum value during the boreal winter season, i.e., December to January in the Northern Hemisphere while June to July in the Southern Hemisphere. Many previous studies (Lin et al., 2009; Wood, 2012; Dong et al., 2014) have demonstrated that the seasonal variations of marine cloud properties (e.g., cloud fraction, LWP and cloud thickness) are strongly affected by meteorological conditions (Lin et al., 2009; Wood, 2012; Dong et al., 2014). Employing a 19-month record of ground-based lidar/radar observations from the Atmospheric Radiation Measurement Program Azores site, for example, Dong et al. (2014) found that the seasonal variations of cloud thickness and LWP are closely related to the seasonal synoptic patterns (e.g., transport of water vapor, relative humidity, high/low pressure system). Furthermore, the influence of aerosols loading is non-neglectable. While the aerosols act as CCN, the concentration of CCN can significantly influence the cloud albedo of low clouds (Twomey, 1974). On the other hand, absorbing aerosols near stratocumulus may enhance absorbing solar energy, resulting in an influence on the dynamical evolution of stratocumulus causing a change in the cloud albedo (Wilcox, 2010). The seasonal cycle of the cloud albedo at the Australia region showed the largest amplitude among the five regions (ranging from 0.37 to 0.52) while the amplitudes at other regions were less than 0.10. Such a result means that the meteorological conditions and aerosol loadings of the cloud system at the Australian region have a relatively larger seasonal variation compared with those at other regions. The COT usually increases with an increase in cloud LWP, resulting in an increase in the cloud albedo (Wood, 2012). Gryspeerdt et al. (2019) also concluded that LWP is the main factor controlling liquid cloud albedo. Thus, this study investigated the seasonal variation of LWP and found that the change in LWP is strongly correlated with the change in cloud albedo at the Peruvian, Australian and Canarian regions (see Fig. S4S3). For the Namibian region, however, many studies have shown that the continuous transportation of absorbing biomass burning aerosols from Africa to the region during the African biomass burning season from August to October (Das et al., 2017) can reside above the clouds, resulting in an increase in the cloud albedo by thickening the stratocumulus (Wilcox, 2010, 2012). Zuidema et al. (2018) also found that the biomass burning aerosols generally exist in the boundary layer at the earlier time of the biomass burning seasons and are mainly located at above the clouds in September to October, which is caused by the northwestward transportation of the biomass burning aerosols from the African continent. However, Fig. 5b shows that the peak of the cloud albedo occurred in July and then continuously decreased from August to October at the Namibian, indicating that the changes in the cloud albedo are difficult to be explained by the negative semi-direct effect of the biomass burning aerosols. This result is consistent with the work of Bender et al. (2016) which concluded that the direct effect and positive semi-direct effect are the main aerosol effects (Wilcox, 2012). That is, clouds become darker under a polluted environment. Regarding the seasonal cycles of cloud droplet number concentration N_d (Li et al., 2018), we found that the seasonal cycles of the cloud albedo at the Namibian region were highly correlated with those of N_d obtained from The Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) (Li et al., 2018), whereas the seasonal cycles of N_d and the cloud albedo showed opposite seasonal changes to each other at the California region. The relationship between the N_d and the cloud

albedo varies with different regions, which may be caused by ~~excluding~~ the effect of meteorological conditions. These
355 results indicate that it is a challenge to study the variability in the cloud albedo over the marine stratocumulus regions under
various meteorological and aerosol conditions.

Fig. 5(a-e) shows the seasonal cycles of cloud albedo at the five regions during a period from 2003 to 2008 for the
AMIP5/AMIP6 and the satellite-based observation. Shading areas in Fig. 5 represent the range of the cloud albedo simulated
by the 22 models. The R5/R6 and P5/P6 values for the seasonal cycles of the cloud albedo obtained from the models and the
360 satellite-based observation are also given in Fig. 5. For the AMIP5-MEM and AMIP6-MEM, the correlations of the cloud
albedo seasonal cycles between the models and the observation are highly positive at all regions ($R5/R6 > 0.6$), except for the
Canarian region ($R5/R6 = 0.22/0.53$). The R values were the largest at the Namibian ($R5/R6 = 0.82/0.92$) and Australian
regions ($R5/R6 = 0.93/0.92$). Overall, the results of AMIP6 were slightly superior to those of AMIP5, especially at the
Canarian region. However, the seasonal cycles of cloud albedo estimated from the AMIP6-MEM at the Canarian region for
365 12 years from 2003 to 2014 (Fig. 5j) exhibited a significant negative correlation with that of the satellite-based observation,
indicating AMIP6-MEM still has a lack of skill to capture the seasonal cycle of the cloud albedo at this region even if the
numbers of AMIP6 models increases. ~~On the contrary, AMIP6-MEM showed a good agreement with the satellite-based
observation at the Australian region (Fig. 5i, $R = 0.93$), indicating a good skill to reproduce the seasonal cloud albedo at this
region. Although the correlation coefficient value at the Peruvian region was the highest ($R = 0.95$), AMIP6-MEM
370 systematically underestimated the monthly cloud albedo values.~~

3.2 The impacts of different aerosol types and meteorological factors on cloud albedo changes

Cloud liquid water may affect the COT, which is subsequently influencing the cloud albedo (Wood, 2012). ~~Bender et al.
(2016) found that for constant cloud fraction, the LWP dominates the change in the cloud albedo.~~ Furthermore, the change of
375 LWP also may influence the relationship between aerosols and cloud properties (Robert et al., 2008; Gryspeerdt et al., 2019;
Douglas and L'Ecuyer, 2019). For example, the effect of aerosols on the cloud albedo may be weakened by a change in the
LWP (Han et al., 2002; Twohy, 2005). Based on in situ observations, recent studies found that the relationship between
aerosol concentration and cloud droplet effective radius changes from negative to positive when liquid water content
increases (Qiu et al., 2017; Zhao et al., 2019b). Considering the effect of LWP, this study evaluated the impact of
380 meteorological parameters and aerosol types on the cloud albedo at different LWP ranges in order to evaluate the influence
of LWP on cloud albedo. Given various LWP ranges, this study evaluated the impact of meteorological parameters and
aerosol types on the cloud albedo to evaluate the influence of LWP on the cloud albedo. Firstly, the 720 monthly sample data
obtained from the five regions were divided into two groups based on the range of monthly mean LWP values: $LWP \leq 6065$
 $g\ m^{-2}$ and $6065\ g\ m^{-2} < LWP \leq 120\ g\ m^{-2}$. Here, the threshold of $65\ g\ m^{-2}$ for LWP was chosen to evenly split the samples.

385 Figure 6a-b shows the regression coefficients in the partial correlation calculation and the relative contributions for
individual variables related to cloud albedo changes under different LWP conditions. ~~The units of vertical velocity and
aerosol mass concentrations are Pa s^{-1} and kg m^{-3} , respectively.~~ Normalized variables were incorporated into the regression
models. There is a considerable discrepancy in the results between the two groups. For the lower LWP bin (i.e., $\text{LWP} \leq 6065$
390 g m^{-2}), the results showed that the regression coefficient related BC/SO2/SS to the cloud albedo was positive while DU and
OC-related coefficients were negative, which indicates that the cloud albedo increases with increasing BC/SO2/SS and
decreases with increasing DU/OC. Fig. 6b also clearly shows that DU, and BC and OC have a larger contribution to the
change in the cloud albedo compared with other predictors, e.g., omega900, EIS and RH850RH700. Under $\text{LWP} > 6065$ g
 m^{-2} , ~~the contribution of DU to the cloud albedo was the largest. In addition, SO2 and SO4 also considerably contributed to
the cloud albedo. the contribution of SO4 to the cloud albedo was the largest. In addition, DU, SO2, BC and RH850 also
395 considerably contributed to the cloud albedo.~~

In addition to the effects of LWP, the difference in the relative contribution may be induced by the regional variability in
aerosol types. A smaller LWP mainly appeared at the Namibian and Canarian where the main aerosol types are DU and BC,
while lower BC loadings were found at the regions with a larger LWP (Fig. S5S4). While the positive coefficient for BC
reflects the indirect effect of aerosols on the cloud albedo, the negative dependency of BC may represent the direct and semi-
400 direct effects of absorbing aerosols (Johnson et al., 2004; Bender et al., 2016). For example, Johnson et al. (2004) found that
absorbing aerosols in clouds can make the clouds warmer and thinner, resulting in a decrease in cloud albedo. Besides,
McCoy et al. (2018) found a negative dependence of N_d on BC at regions with low BC loadings. This means that a decrease
in the cloud albedo may be associated with a decrease in N_d . The dependence of N_d on OC has been also investigated in
previous studies (McCoy et al., 2018; Li et al., 2018) and a negative dependence of N_d on OC has been found in some marine
405 regions. The negative sensitivities of OC to the cloud albedo may attribute to a decrease of N_d with an increase of OC.

Dust is a crucial predictor of the cloud albedo and the coefficient of DU was negative for the two datasets divided in this
study, which may be induced by the semi-direct effects of absorbing aerosols. In literature, many studies have examined the
impacts of dust aerosols on stratocumulus (Doherty and Evan, 2014; Amiri-Farahani et al., 2017). For example, Karydis et al.
(2011) showed that aged dust reduces N_d by consuming the supersaturation of clouds. ~~However, McCoy et al. (2017)~~
410 ~~estimated the indirect effect of aerosol from satellite observations and reanalysis data and found that the dust has a limited
impact on N_d in different stratocumulus regions.~~ Pradelle et al. (2002) employing satellite observations also investigated the
effect of Saharan dust on marine stratocumulus clouds and found that minimum cloud albedo values appeared in regions
with the most dust particles. They also found that the dust in a stratiform cloud may decrease the initial CCN and increase
the effective droplet radius, which causes reducing the cloud albedo (Pradelle and Cautenet 2002). ~~In addition, a recent study~~
415 ~~also showed that the dust aerosol can even further influence the meteorological environment that the clouds form by both
suppressing the SST and affecting the temperature and humidity profile (Sun et al., 2020). Meeoy et al. (2017) estimated the
indirect effect of aerosol from satellite observations and reanalysis data and found that the dust has a limited impact on N_d in~~

~~different stratocumulus regions.~~ A significant influence of dust on the cloud albedo in this study may be driven by the collected samples at the five regions where the cloud albedo and dust highly vary with the regions.

420 Under $LWP \leq 6065 \text{ g m}^{-2}$, the coefficient of SS was a small positive value while the correlation coefficient of sea salt was insignificant under $LWP > 6065 \text{ g m}^{-2}$, which means that these variables are not suitable as a predictor for estimating the cloud albedo. This is consistent with the results of McCoy et al. (2017; 2018) ~~showing that~~ which indicate that the N_d is weakly dependent on the SS although sea salt is an effective CCN. McCoy et al. (2018) have also validated the influence of SS on N_d with up-to-date observations. As submicron SS in the MERRA-2 reanalysis data can be simply predicted from

425 wind speed and SST by a parameterization (Jaeglé et al., 2011; McCoy et al., 2018; Li et al., 2018), the effect of SS on the cloud albedo may be dependent on the relationship between the cloud albedo and near-surface wind speed, which may explain the limited effect of SS on the cloud albedo.

The coefficients of SO_2 were positive for both datasets. In addition, the Twomey effect for SO_2 was further pronounced under the condition with higher LWP. The previous studies (e.g., McCoy et al., 2017; Li et al., 2018) showed that SO_4 plays

430 a key role in modulating N_d . Although their results showed significant positive coefficients of SO_4 with N_d , this study found an unexpected negative correlation of SO_4 with the cloud albedo. Such a result may be driven by the fact that the sulfate aerosol particles and dust are externally mixed. The previous studies showed that sulfate-covered dust can act as CCN, which may induce a decrease in the cloud albedo by enhancing the collision-coalescence progress of droplets (Levin et al., 1996; Rosenfeld et al., 2001).

435 The results of this study showed a weak dependency of the cloud albedo with omega-900, RH700 and EIS. Under $LWP \leq 6065 \text{ g m}^{-2}$, the upward vertical velocity ~~and RH700 has havea- an unexpected negative positive~~ but weak effect on the cloud albedo ~~by enhanced vapor supersaturations, which activates aerosol particles more and subsequently increasing N_d (Twomey, 1959; Reutter et al., 2009) and the relative contributions of omega900 and RH700 are negligible~~. Under $LWP > 6065 \text{ g m}^{-2}$,

440 no significant correlation between the cloud albedo and omega900 was found. Note that the analysis of this study employed the average data at the monthly scale rather than raw satellite measurements at an instantaneous pixel scale, which may make the cloud albedo less sensitive to omega900. ~~In the study, the unexpected negative dependency of the cloud albedo with RH850 was found for the two datasets divided in this study. The coefficient of RH700 was positive and the relative contribution was about 4%. Generally, Drier-drier~~ free-troposphere humidity usually drives stronger entrainment of dry air, which induces evaporating and raising lifted condensation level, resulting in a reduced cloud thickness (Wood, 2012;

445 Eastman and Wood, 2018). The positive dependency of the cloud albedo with EIS was identified for the two datasets divided in this study, which may be caused by stronger inversions linked to increased stability and reduced vertical exchange at cloud top, resulting in thicker low clouds by keeping moisture trapped in the marine boundary layer (MBL) (Scott et al., 2020). Compared to other meteorological factors, the contribution of SSTadv to cloud albedo was larger, non-negligible in both datasets (7%~9%). Under $LWP \leq 65 \text{ g m}^{-2}$, the SSTadv showed a negative coefficient. The cold advection usually

450 thickens clouds by reducing low level stability and transporting more moisture into the MBL (George and Wood, 2010; Scott et al., 2020). Under $LWP > 65 \text{ g m}^{-2}$, the coefficient of SSTadv was positive, which is hard to be explained by the

455 ~~forementioned mechanism. By analysing the correlation between LWP and SSTadv, we found that there was no significant correlation between them. This indicates that the surface temperature advection may affect cloud albedo in other ways than by affecting the moisture in cloud layers. Furthermore, dust can affect the meteorological environment through radiative effects, consequently, the positive coefficient found in this study may be a reflection of their effects (Sun et al., 2020; Huang et al., 2021). The coefficients of the omega700 were negative for both datasets. The downdraft allowed dry air above the cloud to enter the clouds, causing evaporation and making cloud droplets smaller and less, resulting in reducing the cloud albedo (Yang et al., 2019). Note that the role of omega700 was very weak under the condition with higher LWP, and its contribution was negligible. However, Ackerman et al. (2004) showed that the sedimentation of cloud droplets reduces the~~
460 ~~entrainment rate simulated by a dynamics model. If overlying air is in drier and warmer environments, the reduced entrainment will result in a thicker cloud (Bretherton et al., 2007).~~

The analysis of the relative contribution of each predictor variable was similar to the results of the coefficients. Under LWP ≤ 6065 g m⁻², DU and BC contributed approximately 70-63 % variations of the cloud albedo in the regression model. ~~Note that the contribution of omega700 and SSTadv was non-negligible, accounting for 18 %.~~ Under LWP > 6065 g m⁻², ~~SO4 has the largest relative contribution to the cloud albedo changes (~30 %) while~~ the contribution of SO4-DU and DU-SO2 to the change of cloud albedo was about 50-61 %. ~~DU has the largest relative contribution to the cloud albedo changes (~35 %) in both datasets. Note that the contribution of RH850 was non-negligible, accounting for 20 %.~~

The normalized satellite-based and model-driven cloud albedos under different cloud water conditions are shown in Figure 6c-d, where the correlation (R) between the two cloud albedos is given in parentheses. A larger R value indicates a better
470 model. Both of the correlation coefficients are greater than 0.65, indicating the regression model properly captures the changes in the cloud albedo for the two datasets. A considerable part of the variation in cloud albedo can be explained by the change in meteorological parameters and mass concentrations of different aerosol types.

It is also found from Figure 6 that changes in LWP can also cause an alteration of the relationship between aerosol and the cloud albedo. To further investigate the influence of meteorological factors on the relationship, the partial correlations were
475 calculated to eliminate the influence of meteorological parameters individually or simultaneously. If the partial correlation is similar to the total correlation, it means that the influence of meteorological factors on the relationship is limited. In contrast, the influence of meteorological factors on the relationship may be significant if the partial correlation and the total correlation are the opposite sign. Given ~~three-six~~ meteorological parameters (~~omega700, omega900, RH850-RH700, EIS, SSTadv~~ and LWP) considered in this study, the total correlation and partial correlation between the cloud albedo and
480 different aerosols for two sample groups are given in Table 3. ~~The total and partial correlations are similar for BC, OC, SO4~~
~~Under LWP ≤ 6065 g m⁻², indicating that meteorological factors have little influence on the interactions between the aerosols and the cloud albedo. The correlations of SS-all aerosol types was/were weakened~~ when eliminating the effects of meteorological factors. When the influence of ~~EIS and LWP was/were~~ eliminated, the correlation of DU becomes much weaker, indicating that the correlation of DU is sensitive to ~~EIS and LWP. On the contrary, the correlations of BC, OC and~~
485 ~~SO4 were stronger when the influence of LWP was eliminated. In addition, most aerosol types were sensitive to SSTadv~~

except for the SS. The correlation of BC/OC ranged from 0.21/0.20 to -0.03/-0.05 by eliminating the influence of SSTadv, indicating that the relationship between BC/OC and the cloud albedo is extremely sensitive to the influences of SSTadv. On the contrary, the correlation of SO₂ was stronger when the influence of LWP was eliminated. Under LWP > 6065 g m⁻², the correlations of SO₂-all aerosol types ranged from -0.11 to -0.25 varied significantly by eliminating the influence of all meteorological parameters, indicating the relationship between SO₂ and the cloud albedo is extremely sensitive to the influences of meteorological factors. However, the impacts of meteorological factors on the aerosol-cloud interactions were insignificant for the other aerosol types. E. g., the correlation of BC/DU/OC ranged from -0.47/-0.49/-0.45 to -0.01/-0.02/-0.03. This indicates that the cloud-aerosol interaction is more sensitive to the response of meteorological conditions at higher LWP conditions. Although the contribution of meteorological parameters to the change in the cloud albedo is only a small part based on relative contribution calculation, its influence on cloud-aerosol interactions is not negligible.

4 Conclusions and discussion

The cloud albedo at the ~~subtropical~~ marine subtropical stratocumulus regions has a key role in regulating the regional energy budget. However, climate models have a lack of skill to properly capture the cloud properties ~~for over~~ the regions. Therefore, the CMIP6 has more attention to ~~improving-improve~~ some long-standing model biases, e.g., the low cloud simulation over tropical oceans and surface processes (Stouffer et al., 2016). Accordingly, considerable improvements in reproducing the observed seasonal planetary albedo over the subtropical stratocumulus have been found in CMIP6 (Jian et al., 2020). To enhance the confidence in climate predictions, it is necessary to systematically evaluate and compare the performance of CMIP5 and CMIP6 models and to further study the processes that contribute to the cloud albedo using the satellite-driven and reanalysis data. This study investigated the performances of CMIP6 models in reproducing the cloud albedo at the five marine subtropical stratocumulus regions ~~during from~~ 2003 to 2014.

For the long-term regressed values, the cloud albedos were underestimated in most AMIP6 models compared with the satellite-driven cloud albedos. The AMIP6 models produced a similar spread of AMIP5 at all regions, even some AMIP6 models performed worse than AMIP5. ~~The correlation between cloud fraction and planetary albedo for AMIP6 exhibited a larger spread at the Peruvian and Namibian regions while a larger intermodel variability in the cloud albedo at the Australian and Canarian regions compared with other regions.~~ The monthly cloud albedo of AMIP6-MEM showed better correlation with the satellite-driven observation than that of AMIP5-MEM. However, this study found a lack of skill in reproducing the values and amplitude at ~~some regions (e.g., the~~ Peruvian and Namibian), indicating that the cloud parameterization between two generations of AMIP models needs to be further improved to produce more accurate predictions. This study also found that most AMIP6 models overestimated the amplitude of the cloud albedo at all regions except for the Australian region, i.e., simulating higher seasonal variations. Overall, the AMIP6 models performed the best at the Australian region and the worst at the Canarian region. ~~In addition, both the AMIP5-MEM and AMIP6-MEM reproduced the seasonal cycle of the cloud albedo at all regions except for the Canarian region.~~ The seasonal cycle of cloud albedo of AMIP6-MEM ~~was~~ correlated

520 better with satellite-driven observations than that of AMIP5-MEM. For the Australian region, the model-driven seasonal cycle of the cloud albedo was almost consistent with that of the satellite-driven observation, which indicates the superiority of model performance at this region.

Employing the satellite and reanalysis data, we further evaluated the impacts of different aerosol types and meteorological factors on the cloud albedo. Changes in aerosol types and meteorological factors explained ~~~60-65~~ % of the changes in the cloud albedo. However, the controlling factors and their contribution rates varied with LWP conditions. Under the monthly mean LWP ≤ 6065 g m⁻², DU and BC dominantly contributed to the changes in the cloud albedo, while DU and SO₄-SO₂ contributed the most under the condition of 6065 g m⁻² < LWP ≤ 120 g m⁻². Although the contributions of aerosols were significant, the influence of meteorological factors on the cloud-aerosol interactions cannot be ignored.

530 Due to the limitations of polar-orbiting satellite observations, this study did not obtain a complete diurnal cycle of cloud properties and radiation flux, which may induce a bias in the results of this study. The diurnal cycle of marine subtropical stratocumulus cloud albedo is usually significant due to the diurnal cycle of solar energy (Wood, 2012). The maximum cloud thickness usually occurs in the morning and gradually decreasing over the afternoon due to absorbing solar radiation in the cloud layer (Wood et al., 2002; Christensen et al., 2013). It is a challenge to evaluate how much of the cloud albedo bias contributes to the diurnal cycle of cloud albedo. Therefore, it is necessary to evaluate the diurnal cycle of cloud albedo in the marine subtropical stratocumulus regions for reducing the uncertainties in cloud radiation interactions in GCMs. Note that the “too bright, too few” problem was improved at the Namibian and Californian regions in AMIP6. However, even if some models can simulate the cloud albedo more reasonably, it is questionable if other cloud properties can be captured (e.g., total cloud fraction), consequently resulting in significant biases in radiation (see Fig. S3S5). Therefore, we need to pay more attention to improving-improve the calculation of total cloud fraction in the GCMs. Recently, some studies are devoted to improving-improve cloud overlap parameterization for accurately simulate-simulating the cloud fractions in GCMs (Li et al., 2018, 2019). Accordingly, it is also necessary to evaluate the improvement of cloud overlap scheme on cloud radiation interaction using long-term satellite-driven observations and reanalysis data.

Data availability

545 The CERES datasets are available from the CERES website: <https://ceres.larc.nasa.gov/data/#single-scanner-footprint-ssf>. The MODIS datasets are available from the Level-1 and Atmosphere Archive & Distribution System (LAADS) Distributed Active Archive Center (DAAC) website: <https://ladsweb.modaps.eosdis.nasa.gov/archive/allData/61>. The MERRA-2 reanalysis products are downloaded from the MERRA-2 website: <https://disc.gsfc.nasa.gov/datasets?keywords=MERRA-2>. The CMIP5 and CMIP6 products are downloaded from the Earth System Grid Federation (ESGF) website: <https://esgf-node.llnl.gov/projects/esgf-llnl/>.

Competing interests

550 The authors declare that they have no conflict of interest.

Author contribution

BJ and JL organized the paper and carried them out. BJ prepared the manuscript with contributions from all co-authors. JL conceptualized the paper and revised the whole manuscript. GW provided the computing resources. YL downloaded the data and maintain research data. YZ, JW and MZ processed the raw model output data into consistent gridded format for
555 comparison with the satellite dataset. JH provided consultations and acquired the financial support for this study. All authors contributed to the discussion of the results and reviewed the manuscript.

Acknowledgments

This research was jointly supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDA2006010301), the National Science Fund for Excellent Young Scholars (42022037). The authors declare that they
560 have no conflict of interest. We would like to thank the CERES, MODIS, CMIP5, CMIP6 and MERRA-2 science teams for providing excellent and accessible data products that made this study possible.

References

- ~~Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing, *Nature*, 432, 1014-1017, <https://doi.org/10.1038/nature03174>, 2004.~~
- 565 Amiri-Farahani, A., Allen, R. J., Neubauer, D., and Lohmann, U.: Impact of Saharan dust on North Atlantic marine stratocumulus clouds: importance of the semidirect effect, *Atmos. Chem. Phys.*, 17, 6305-6322, <https://doi.org/10.5194/acp-17-6305-2017>, 2017.
- Betts, A. K., and Ridgway, W.: Climatic equilibrium of the atmospheric convective boundary layer over a tropical ocean. *J. Atmos. Sci.*, 46, 2621-2641, [https://doi.org/10.1175/1520-0469\(1989\)046<2621:CEOTAC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1989)046<2621:CEOTAC>2.0.CO;2), 1989.
- 570 Bender, F. A. M., Charlson, R. J., Ekman, A. M. L., and Leahy, L. V.: Quantification of Monthly Mean Regional-Scale Albedo of Marine Stratiform Clouds in Satellite Observations and GCMs, *J. Appl. Meteorol. Clim.*, 50, 2139-2148, <https://doi.org/10.1175/jamc-d-11-049.1>, 2011.

- Bender, F. A. M., Engström, A., and Karlsson, J.: Factors Controlling Cloud Albedo in Marine Subtropical Stratocumulus Regions in Climate Models and Satellite Observations, *J. Clim.*, 29, 3559-3587, <https://doi.org/10.1175/Jcli-D-15-0095.1>, 575 2016.
- Bender, F. A. M., Engström, A., Wood, R., and Charlson, R. J.: Evaluation of Hemispheric Asymmetries in Marine Cloud Radiative Properties, *J. Clim.*, 30, 4131-4147, <https://doi.org/10.1175/Jcli-D-16-0263.1>, 2017.
- Bender, F. A. M., Frey, L., McCoy, D. T., Grosvenor, D. P., and Mohrmann, J. K.: Assessment of aerosol–cloud–radiation correlations in satellite observations, climate models and reanalysis, *Clim. Dynam.*, 52, 4371-4392, 580 <https://doi.org/10.1007/s00382-018-4384-z>, 2018.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y.: Clouds and aerosols, in: Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, 585 USA, 2013.
- ~~[Chen, Y. C., Christensen, M. W., Stephens, G. L., and Seinfeld, J. H.: Satellite-based estimate of global aerosol-cloud radiative forcing by marine warm clouds, *Nat. Geosci.*, 7, 643-646, <https://doi.org/10.1038/NNGEO2214>, 2014.](#)~~
- ~~[Bretherton, C. S., Blossey, P. N., and Uehida, J.: Cloud droplet sedimentation, entrainment efficiency, and subtropical stratocumulus albedo, *Geophys. Res. Lett.*, 34, 5, <https://doi.org/10.1029/2006gl027648>, 2007.](#)~~
- 590 Christensen, M. W., Carrio, G. G., Stephens, G. L., and Cotton, W. R.: Radiative Impacts of Free-Tropospheric Clouds on the Properties of Marine Stratocumulus, *J. Atmos. Sci.*, 70, 3102-3118, <https://doi.org/10.1175/jas-d-12-0287.1>, 2013.
- Das, S., Harshvardhan, H., Bian, H., Chin, M., Curci, G., Protonotariou, A. P., Mielonen, T., Zhang, K., Wang, H., and Liu, X.: Biomass burning aerosol transport and vertical distribution over the South African-Atlantic region, *J. Geophys. Res.*, 122, 6391-6415, <https://doi.org/10.1002/2016jd026421>, 2017.
- 595 ~~[Doelling, D. R., Loeb, N. G., Keyes, D. F., Nordeen, M. L., Morstad, D., Nguyen, C., Wielicki, B. A., Young, D. F., and Sun, M.: Geostationary Enhanced Temporal Interpolation for CERES Flux Products, *J. Atmos. Ocean Tech.*, 30, 1072-1090, <https://doi.org/10.1175/jtech-d-12-00136.1>, 2013.](#)~~
- Doherty, O. M., and Evan, A. T.: Identification of a new dust-stratocumulus indirect effect over the tropical North Atlantic, *Geophys. Res. Lett.*, 41, 6935-6942, <https://doi.org/10.1002/2014gl060897>, 2014.
- 600 Dong, X., Xi, B., Kennedy, A., Minnis, P., and Wood, R.: A 19-Month Record of Marine Aerosol- Cloud-Radiation Properties Derived from DOE ARM Mobile Facility Deployment at the Azores. Part I: Cloud Fraction and Single-Layered MBL Cloud Properties, *J. Clim.*, 27, 3665-3682, <https://doi.org/10.1175/jcli-d-13-00553.1>, 2014.
- Douglas, A., amp, apos, and Ecuyer, T.: Quantifying variations in shortwave aerosol–cloud–radiation interactions using local meteorology and cloud state constraints, *Atmos. Chem. Phys.*, 19, 6251-6268, <https://doi.org/10.5194/acp-19-6251-2019>, 605 2019.

- Eastman, R., and Wood, R.: The Competing Effects of Stability and Humidity on Subtropical Stratocumulus Entrainment and Cloud Evolution from a Lagrangian Perspective, *J. Atmos. Sci.*, 75, 2563-2578, <https://doi.org/10.1175/jas-d-18-0030.1>, 2018.
- 610 Engström, A., Bender, F. A. M., Charlson, R. J., and Wood, R.: The nonlinear relationship between albedo and cloud fraction on near-global, monthly mean scale in observations and in the CMIP5 model ensemble, *Geophys. Res. Lett.*, 42, 9571-9578, <https://doi.org/10.1002/2015GL066275>, 2015.
- Engström, A., Bender, F. A. M., and Karlsson, J.: Improved Representation of Marine Stratocumulus Cloud Shortwave Radiative Properties in the CMIP5 Climate Models, *J. Clim.*, 27, 6175-6188, <https://doi.org/10.1175/jcli-d-13-00755.1>, 2014.
- 615 Engström, A., and Ekman, A. M. L.: Impact of meteorological factors on the correlation between aerosol optical depth and cloud fraction, *Geophys. Res. Lett.*, 37, L18814, <https://doi.org/10.1029/2010gl044361>, 2010.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937-1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Frey, L., Bender, F. A. M., and Svensson, G.: Cloud albedo changes in response to anthropogenic sulfate and non-sulfate 620 aerosol forcings in CMIP5 models, *Atmos. Chem. Phys.*, 17, 9145-9162, <https://doi.org/10.5194/acp-17-9145-2017>, 2017.
- [Fuchs, J., Cermak, J., and Andersen, H.: Building a cloud in the southeast Atlantic: understanding low-cloud controls based on satellite observations with machine learning. *Atmos. Chem. Phys.*, 18, 16537-16552, <https://doi.org/10.5194/acp-18-16537-2018>, 2018.](https://doi.org/10.5194/acp-18-16537-2018)
- 625 Gelaro, R., McCarty, W., Suarez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A., Gu, W., Kim, G. K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Clim.*, 30, 5419-5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 630 [Georgakakos, K. P., and Bras, R. L.: A hydrologically useful station precipitation model: I. Formulation, *Water Resour. Res.*, 20, 1585-1596, <https://doi.org/10.1029/WR020i011p01585>, 1984.](https://doi.org/10.1029/WR020i011p01585)
- George, R. C., and Wood, R.: Subseasonal variability of low cloud radiative properties over the southeast Pacific Ocean, *Atmos. Chem. Phys.*, 10, 4047-4063, <https://doi.org/10.5194/acp-10-4047-2010>, 2010.
- Gryspeerd, E., Goren, T., Sourdeval, O., Quaas, J., Mülmenstädt, J., Dipu, S., Unglaub, C., Gettelman, A., and Christensen, M.: Constraining the aerosol influence on cloud liquid water path, *Atmos. Chem. Phys.*, 19, 5331-5347, 635 <https://doi.org/10.5194/acp-19-5331-2019>, 2019.
- Han, Q. Y., Rossow, W. B., Zeng, J., and Welch, R.: Three different behaviors of liquid water path of water clouds in aerosol-cloud interactions, *J. Atmos. Sci.*, 59, 726-735, [https://doi.org/10.1175/1520-0469\(2002\)059<0726:tdbolw>2.0.co;2](https://doi.org/10.1175/1520-0469(2002)059<0726:tdbolw>2.0.co;2), 2002.

- Herbert, R. J., Bellouin, N., Highwood, E. J., and Hill, A. A.: Diurnal cycle of the semi-direct effect from a persistent absorbing aerosol layer over marine stratocumulus in large-eddy simulations, *Atmos. Chem. Phys.*, 20, 1317-1340, <https://doi.org/10.5194/acp-20-1317-2020>, 2020.
- Huang, J., and Yi, Y.: Inversion of a nonlinear dynamical model from the observation. *Science in China*, 34B, 1246–1251, 1991.
- [Huang, J. P., Huang, J. P., Liu, X. Y., Li, C. Y., Ding, L., and Yu, H. P.: The global oxygen budget and its future projection, *Sci. Bull.*, 63, 1180-1186, <https://doi.org/10.1016/j.scib.2018.07.023>, 2018.](#)
- [Huang, J. P., Liu, X. Y., He, Y. S., Shen, S. Z., Hou, Z. Q., Li, S. G., Li, C. Y., Yao, L. J., and Huang, J. P.: The oxygen cycle and a habitable Earth, *Sci. China Earth Sci.*, 64, 511-528, <https://doi.org/10.1007/s11430-020-9747-1>, 2021.](#)
- Jaegle, L., Quinn, P. K., Bates, T. S., Alexander, B., and Lin, J. T.: Global distribution of sea salt aerosols: new constraints from in situ and remote sensing observations, *Atmos. Chem. Phys.*, 11, 3137-3157, <https://doi.org/10.5194/acp-11-3137-2011>, 2011.
- Jian, B., Li, J., Wang, G., He, Y., Han, Y., Zhang, M., and Huang, J.: The Impacts of Atmospheric and Surface Parameters on Long-Term Variations in the Planetary Albedo, *J. Clim.*, 31, 8705-8718, <https://doi.org/10.1175/Jcli-D-17-0848.1>, 2018.
- Jian, B., Li, J., Zhao, Y., He, Y., Wang, J., and Huang, J.: Evaluation of the CMIP6 planetary albedo climatology using satellite observations, *Clim. Dynam.*, 54, 5145-5161, <https://doi.org/10.1007/s00382-020-05277-4>, 2020.
- Jiang, J. H., Su, H., Huang, L., Wang, Y., Massie, S., Zhao, B., Omar, A., and Wang, Z.: Contrasting effects on deep convective clouds by different types of aerosols, *Nat. Commun.*, 9, 3874, <https://doi.org/10.1038/s41467-018-06280-4>, 2018.
- Johnson, B. T., Shine, K. P., and Forster, P. M.: The semi-direct aerosol effect: Impact of absorbing aerosols on marine stratocumulus, *Q. J. Roy. Meteor. Soc.*, 130, 1407-1422, <https://doi.org/10.1256/qj.03.61>, 2004.
- Karydis, V. A., Kumar, P., Barahona, D., Sokolik, I. N., and Nenes, A.: On the effect of dust particles on global cloud condensation nuclei and cloud droplet number, *J. Geophys. Res.*, 116, D23204, <https://doi.org/10.1029/2011jd016283>, 2011.
- Kawai, H., Yukimoto, S., Koshiro, T., Oshima, N., Tanaka, T., Yoshimura, H., and Nagasawa, R.: Significant improvement of cloud representation in the global climate model MRI-ESM2, *Geosci. Model Dev.*, 12, 2875-2897, <https://doi.org/10.5194/gmd-12-2875-2019>, 2019.
- Klein, S. A., and Hartmann, D. L.: THE SEASONAL CYCLE OF LOW STRATIFORM CLOUDS, *J. Clim.*, 6, 1587-1606, [https://doi.org/10.1175/1520-0442\(1993\)006<1587:tscols>2.0.co;2](https://doi.org/10.1175/1520-0442(1993)006<1587:tscols>2.0.co;2), 1993.
- Kravitz, B., Wang, H., Rasch, P. J., Morrison, H., and Solomon, A. B.: Process-model simulations of cloud albedo enhancement by aerosols in the Arctic, *Philos. T. R. Soc. A.*, 372, <https://doi.org/10.1098/rsta.2014.0052>, 2014.
- Latham, J., Rasch, P., Chen, C.-C., Kettles, L., Gadian, A., Gettelman, A., Morrison, H., Bower, K., and Choulaton, T.: Global temperature stabilization via controlled albedo enhancement of low-level maritime clouds, *Philos. T. R. Soc. A.*, 366, 3969-3987, <https://doi.org/10.1098/rsta.2008.0137>, 2008.

- Levin, Z., Ganor, E., and Gladstein, V.: The effects of desert particles coated with sulfate on rain formation in the eastern Mediterranean, *J. Appl. Meteorol.*, 35, 1511-1523, [https://doi.org/10.1175/1520-0450\(1996\)035<1511:teodpc>2.0.co;2](https://doi.org/10.1175/1520-0450(1996)035<1511:teodpc>2.0.co;2), 1996.
- Li, J., Jian, B., Huang, J., Hu, Y., Zhao, C., Kawamoto, K., Liao, S., and Wu, M.: Long-term variation of cloud droplet number concentrations from space-based Lidar, *Remote Sens. Environ.*, 213, 144-161, <https://doi.org/10.1016/j.rse.2018.05.011>, 2018.
- Li, J., Jian, B., Zhao, C., Zhao, Y., Wang, J., and Huang, J.: Atmospheric Instability Dominates the Long-Term Variation of Cloud Vertical Overlap Over the Southern Great Plains Site, *J. Geophys. Res.*, 124, 9691-9701, <https://doi.org/10.1029/2019jd030954>, 2019.
- Lin, W. Y., Zhang, M. H., and Loeb, N. G.: Seasonal Variation of the Physical Properties of Marine Boundary Layer Clouds off the California Coast, *J. Clim.*, 22, 2624-2638, <https://doi.org/10.1175/2008jcli2478.1>, 2009.
- Loeb, N. G., Doelling, D. R., Wang, H. L., Su, W. Y., Nguyen, C., Corbett, J. G., Liang, L. S., Mitrescu, C., Rose, F. G., and Kato, S.: Clouds and the Earth's Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) Top-of-Atmosphere (TOA) Edition-4.0 Data Product, *J. Clim.*, 31, 895-918, <https://doi.org/10.1175/Jcli-D-17-0208.1>, 2018.
- Loeb, N. G., Priestley, K. J., Kratz, D. P., Geier, E. B., Green, R. N., Wielicki, V. A., Hinton, R. O., and Nolan, S. K.: Determination of unfiltered radiances from the clouds and the Earth's Radiant Energy System instrument, *J. Appl. Meteorol.*, 40, 822-835, [https://doi.org/10.1175/1520-0450\(2001\)040<0822:dourft>2.0.co;2](https://doi.org/10.1175/1520-0450(2001)040<0822:dourft>2.0.co;2), 2001.
- Lu, C., Liu, Y., Niu, S., and Vogelmann, A. M.: Observed impacts of vertical velocity on cloud microphysics and implications for aerosol indirect effects, *Geophys. Res. Lett.*, 39, L21808, <https://doi.org/10.1029/2012gl053599>, 2012.
- McCoy, D. T., Bender, F. A. M., Grosvenor, D. P., Mohrmann, J. K., Hartmann, D. L., Wood, R., and Field, P. R.: Predicting decadal trends in cloud droplet number concentration using reanalysis and satellite data, *Atmos. Chem. Phys.*, 18, 2035-2047, <https://doi.org/10.5194/acp-18-2035-2018>, 2018.
- McCoy, D. T., Bender, F. A. M., Mohrmann, J. K. C., Hartmann, D. L., Wood, R., and Grosvenor, D. P.: The global aerosol - cloud first indirect effect estimated using MODIS, MERRA, and AeroCom, *J. Geophys. Res.*, 122, 1779-1796, <https://doi.org/10.1002/2016jd026141>, 2017.
- Mueller, R., Trentmann, J., Traeger-Chatterjee, C., Posselt, R., and Stoeckli, R.: The Role of the Effective Cloud Albedo for Climate Monitoring and Analysis, *Remote Sens.*, 3, 2305-2320, <https://doi.org/10.3390/rs3112305>, 2011.
- Nam, C., Bony, S., Dufresne, J. L., and Chepfer, H.: The 'too few, too bright' tropical low-cloud problem in CMIP5 models, *Geophys. Res. Lett.*, 39, L21801, <https://doi.org/10.1029/2012gl053421>, 2012.
- Platnick, S., King, M. D., Ackerman, S. A., Menzel, W. P., Baum, B. A., Riedi, J. C., and Frey, R. A.: The MODIS cloud products: Algorithms and examples from Terra, *Ieee T. Geosci. Remote*, 41, 459-473, <https://doi.org/10.1109/tgrs.2002.808301>, 2003.
- Pradelle, F., and Cautenet, G.: Radiative and microphysical interactions between marine stratocumulus clouds and Saharan dust - 2. Modeling, *J. Geophys. Res.*, 107, 4413, <https://doi.org/10.1029/2000jd000156>, 2002.

- 705 Pradelle, F., Cautenet, G., and Jankowiak, I.: Radiative and microphysical interactions between marine stratocumulus clouds and Saharan dust - I. Remote sensing observations, *J. Geophys. Res.*, 107, 4412, <https://doi.org/10.1029/2000jd000155>, 2002.
- Reutter, P., Su, H., Trentmann, J., Simmel, M., Rose, D., Gunthe, S. S., Wernli, H., Andreae, M. O., and Poeschl, U.: Aerosol- and updraft-limited regimes of cloud droplet formation: influence of particle number, size and hygroscopicity on the activation of cloud condensation nuclei (CCN), *Atmos. Chem. Phys.*, 9, 7067-7080, <https://doi.org/10.5194/acp-9-7067-2009>, 2009.
- 710 Roberts, G. C., Ramana, M. V., Corrigan, C., Kim, D., and Ramanathan, V.: Simultaneous observations of aerosol-cloud-albedo interactions with three stacked unmanned aerial vehicles, *Proc. Natl. Acad. Sci. U. S. A.*, 105, 7370-7375, <https://doi.org/10.1073/pnas.0710308105>, 2008.
- 715 Rosenfeld, D., Rudich, Y., and Lahav, R.: Desert dust suppressing precipitation: A possible desertification feedback loop, *Proc. Natl. Acad. Sci. U. S. A.*, 98, 5975-5980, <https://doi.org/10.1073/pnas.101122798>, 2001.
- Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T., and Yu, S.: Aerosol-driven droplet concentrations dominate coverage and water of oceanic low-level clouds, *Science*, 363, eaav0566, <https://doi.org/10.1126/science.aav0566>, 2019.
- Seethala, C., Norris, J. R., and Myers, T. A.: How Has Subtropical Stratocumulus and Associated Meteorology Changed since the 1980s?*, *J. Clim.*, 28, 8396-8410, <https://doi.org/10.1175/jcli-d-15-0120.1>, 2015.
- 720 [Qiu, Y. M., Zhao, C. F., Guo, J. P., and Li, J. M.: 8-Year ground-based observational analysis about the seasonal variation of the aerosol-cloud droplet effective radius relationship at SGP site, *Atmos. Environ.*, 164, 139-146, <https://doi.org/10.1016/j.atmosenv.2017.06.002>, 2017.](https://doi.org/10.1016/j.atmosenv.2017.06.002)
- [Qu, X., Hall, A., Klein, S. A., and DeAngelis, A. M.: Positive tropical marine low-cloud cover feedback inferred from cloud-controlling factors, *Geophys. Res. Lett.*, 42, 7767-7775, <https://doi.org/10.1002/2015GL065627>, 2015.](https://doi.org/10.1002/2015GL065627)
- 725 [Scott, R. C., Myers, T. A., Norris, J. R., Zelinka, M. D., Klein, S. A., Sun, M. G., and Doelling, D. R.: Observed Sensitivity of Low-Cloud Radiative Effects to Meteorological Perturbations over the Global Oceans, *J. Clim.*, 33, 7717-7734, <https://doi.org/10.1175/Jcli-D-19-1028.1>, 2020.](https://doi.org/10.1175/Jcli-D-19-1028.1)
- 730 Seland, Ø., Bentsen, M., Seland Graff, L., Olivíe, D., Toniazzo, T., Gjermundsen, A., Debernard, J. B., Gupta, A. K., He, Y., Kirkevåg, A., Schwinger, J., Tjiputra, J., Schancke Aas, K., Bethke, I., Fan, Y., Griesfeller, J., Grini, A., Guo, C., Ilicak, M., ... Schulz, M.: The Norwegian Earth System Model, NorESM2 - Evaluation of the CMIP6 DECK and historical simulations. *Geosci. Model Dev. Discuss.*, <https://doi.org/10.5194/gmd-2019-378>, 2020. (in review)
- Stouffer, R. J., Eyring, V., Meehl, G. A., Bony, S., Senior, C., Stevens, B., and Taylor, K. E.: CMIP5 Scientific Gaps and Recommendations for CMIP6, *Bull. Am. Meteorol. Soc.*, 98, 95-105, <https://doi.org/10.1175/bams-d-15-00013.1>, 2017.
- 735 Stuart, G. S., Stevens, R. G., Partanen, A. I., Jenkins, A. K. L., Korhonen, H., Forster, P. M., Spracklen, D. V., and Pierce, J. R.: Reduced efficacy of marine cloud brightening geoengineering due to in-plume aerosol coagulation: parameterization and global implications, *Atmos. Chem. Phys.*, 13, 10385-10396, <https://doi.org/10.5194/acp-13-10385-2013>, 2013.

- [Sun, Y., and Zhao, C. F.: Influence of Saharan Dust on the Large-Scale Meteorological Environment for Development of Tropical Cyclone Over North Atlantic Ocean Basin, *J. Geophys. Res.*, 125, e2020JD033454, <https://doi.org/10.1029/2020JD033454>, 2020.](https://doi.org/10.1029/2020JD033454)
- 740 Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *J. Geophys. Res.*, 106, 7183-7192, <https://doi.org/10.1029/2000jd900719>, 2001.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bull. Am. Meteorol.*
- 745 *Soc.*, 93, 485-498, 2012.
- Twohy, C. H.: Evaluation of the aerosol indirect effect in marine stratocumulus clouds: Droplet number, size, liquid water path, and radiative impact, *J. Geophys. Res.*, 110, D08203, <https://doi.org/10.1029/2004jd005116>, 2005.
- Twomey, S.: The nuclei of natural cloud formation. Part II: The supersaturation in natural clouds and the variation of cloud droplet concentration, *Geofis. Pura Appl.*, 243-249, <https://doi.org/10.1007/BF01993560>, 1959.
- 750 Twomey, S.: Pollution and the planetary albedo, *Atmos. Environ.*, 8, 1251-1256. [https://doi.org/10.1016/0004-6981\(74\)90004-3](https://doi.org/10.1016/0004-6981(74)90004-3), 1974.
- Twomey, S.: Influence of pollution on shortwave albedo of clouds, *J. Atmos. Sci.*, 34, 1149-1152. [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.
- Van Weverberg, K., Morcrette, C. J., Petch, J., Klein, S. A., Ma, H. Y., Zhang, C., Xie, S., Tang, Q., Gustafson, W. I., Qian,
- 755 Y., Berg, L. K., Liu, Y., Huang, M., Ahlgrimm, M., Forbes, R., Bazile, E., Roehrig, R., Cole, J., Merryfield, W., Lee, W. S., Cheruy, F., Mellul, L., Wang, Y. C., Johnson, K., and Thieman, M. M.: CAUSES: Attribution of Surface Radiation Biases in NWP and Climate Models near the U.S. Southern Great Plains, *J. Geophys. Res.*, 123, 3612-3644, <https://doi.org/10.1002/2017jd027188>, 2018.
- Wall, C. J., Hartmann, D. L., Thieman, M. M., Smith, W. L., and Minnis, P.: The Life Cycle of Anvil Clouds and the Top-
- 760 of-Atmosphere Radiation Balance over the Tropical West Pacific, *J. Clim.*, 31, 10059-10080, <https://doi.org/10.1175/jcli-d-18-0154.1>, 2018.
- Wang, H., Rasch, P. J., and Feingold, G.: Manipulating marine stratocumulus cloud amount and albedo: a process-modelling study of aerosol-cloud-precipitation interactions in response to injection of cloud condensation nuclei, *Atmos. Chem. Phys.*, 11, 4237-4249, <https://doi.org/10.5194/acp-11-4237-2011>, 2011.
- 765 [Wang, Y., and Zhao, C.: Can MODIS cloud fraction fully represent the diurnal and seasonal variations at DOE ARM SGP and Manus sites?, *J. Geophys. Res.*, 122, 329-343, <https://doi.org/10.1002/2016jd025954>, 2017.](https://doi.org/10.1002/2016jd025954)
- Waquet, F., Peers, F., Ducos, F., Goloub, P., Platnick, S., Riedi, J., Tanré, D., and Thieuleux, F.: Global analysis of aerosol properties above clouds, *Geophys. Res. Lett.*, 40, 5809-5814, <https://doi.org/10.1002/2013gl057482>, 2013.
- Wielicki, B. A., Barkstrom, B. R., Harrison, E. F., Lee III, R. B., Louis Smith, G., and Cooper, J. E.: Clouds and the Earth's
- 770 Radiant Energy System (CERES): An earth observing system experiment, *Bull. Am. Meteorol. Soc.*, 77, 853-868, 1996.
- Wilcox, E. M.: Stratocumulus cloud thickening beneath layers of absorbing smoke aerosol, *Atmos. Chem. Phys.*, 10, 11769-11777, <https://doi.org/10.5194/acp-10-11769-2010>, 2010.

- Wilcox, E. M.: Direct and semi-direct radiative forcing of smoke aerosols over clouds, *Atmos. Chem. Phys.*, 12, 139-149, <https://doi.org/10.5194/acp-12-139-2012>, 2012.
- 775 Wood, R.: Stratocumulus Clouds, *Mon. Weather Rev.*, 140, 2373-2423, <https://doi.org/10.1175/mwr-d-11-00121.1>, 2012.
- [Wood, R., and Bretherton, C. S.: On the relationship between stratiform low cloud cover and lower-tropospheric stability, *J. Clim.*, 19, 6425-6432, <https://doi.org/10.1175/Jcli3988.1>, 2006.](#)
- [Xie, S. C., Liu, X. H., Zhao, C. F., and Zhang, Y. Y.: Sensitivity of CAM5-Simulated Arctic Clouds and Radiation to Ice Nucleation Parameterization, *J. Clim.*, 26, 5981-5999, <https://doi.org/10.1175/Jcli-D-12-00517.1>, 2013.](#)
- 780 [Yang, Y., Zhao, C. F., Dong, X. B., Fan, G. C., Zhou, Y. Q., Wang, Y., Zhao, L. J., Lv, F., and Yan, F.: Toward understanding the process-level impacts of aerosols on microphysical properties of shallow cumulus cloud using aircraft observations, *Atmos. Res.*, 221, 27-33, <https://doi.org/10.1016/j.atmosres.2019.01.027>, 2019.](#)
- [Zhao, C. F., Klein, S. A., Xie, S. C., Liu, X. H., Boyle, J. S., and Zhang, Y. Y.: Aerosol first indirect effects on non-precipitating low-level liquid cloud properties as simulated by CAM5 at ARM sites, *Geophys. Res. Lett.*, 39, L08806, <https://doi.org/10.1029/2012gl051213>, 2012.](#)
- 785 [Zhao, C. F., Lin, Y. L., Wu, F., Wang, Y., Li, Z. Q., Rosenfeld, D., and Wang, Y.: Enlarging Rainfall Area of Tropical Cyclones by Atmospheric Aerosols, *Geophys. Res. Lett.*, 45, 8604-8611, <https://doi.org/10.1029/2018GL079427>, 2018.](#)
- [Zhao, C. F., Chen, Y. Y., Li, J. M., Letu, H. S., Su, Y. F., Chen, T. M., and Wu, X. L.: Fifteen-year statistical analysis of cloud characteristics over China using Terra and Aqua Moderate Resolution Imaging Spectroradiometer observations, *Int. J. Climatol.*, 39, 2612-2629, <https://doi.org/10.1002/joc.5975>, 2019a.](#)
- 790 [Zhao, C. F., Zhao, L. J., and Dong, X. B.: A Case Study of Stratus Cloud Properties Using In Situ Aircraft Observations over Huanghua, China, *Atmos.*, 10, 19, <https://doi.org/10.3390/atmos10010019>, 2019b.](#)
- Zuidema, P., Sedlacek, A. J., III, Flynn, C., Springston, S., Delgadillo, R., Zhang, J., Aiken, A. C., Koontz, A., and Muradyan, P.: The Ascension Island Boundary Layer in the Remote Southeast Atlantic is Often Smoky, *Geophys. Res. Lett.*, 45, 4456-4465, <https://doi.org/10.1002/2017gl076926>, 2018.
- 795

Table 1: The list of CMIP5 models used in the study and their atmospheric horizontal resolutions.

800

	Model name	Origin	Resolution (lon×lat)
1	ACCESS1-0	Commonwealth Scientific and Industrial Research Organization and Bureau of	192×145
2	ACCESS1-3	Meteorology, Australia	192×145

3	FGOALS-g2	Institute of Atmospheric Physics, Chinese Academy of Sciences and Tsinghua University, China	128×60
4	GISS-E2-R	NASA/Goddard Institute for Space Studies, USA	144×90
5	INMCM4	Institute for Numerical Mathematics, Russia	180×120
6	IPSL-CM5A-LR	Institut Pierre Simon Laplace, France	96×96
7	MIROC5	AORI, NIES and JAMSTEC, Japan	256×128
8	MPI-ESM-LR	Max Planck Institute for Meteorology, Germany	192×96
9	MRI-CGCM3	Meteorological Research Institute, Japan	320×160
10	NorESM1-M	Norwegian Climate Centre, Norway	144×96


Table 2: The list of CMIP6 models used in the study and their atmospheric horizontal resolutions.

	Model name	Origin	Resolution (lon×lat)
1	ACCESS-CM2	Commonwealth Scientific and Industrial Research Organization and Bureau of Meteorology, Australia	192×145
2	ACCESS-ESM1-5		192×145
3	FGOALS-g3	Institute of Atmospheric Physics, Chinese Academy of Sciences and Tsinghua University, China	180×80
4	GISS-E2-1-G	NASA/Goddard Institute for Space Studies, USA	144×90
5	INM-CM4-8	Institute for Numerical Mathematics, Russia	180×120
6	IPSL-CM6A-LR	Institut Pierre Simon Laplace, France	144×143
7	MIROC6	AORI, NIES and JAMSTEC, Japan	256×128
8	MPI-ESM1-2-HR	Max Planck Institute for Meteorology, Germany	384×192
9	MRI-ESM2-0	Meteorological Research Institute, Japan	320×160
10	NorESM2-LM	Norwegian Climate Centre, Norway	144×96

11	BCC-CSM2-MR	Beijing Climate Center, China	320×160
12	BCC-ESM1		128×64
13	CAMS-CSM1-0	Chinese Academy of Meteorological Sciences , China	320×160
14	CESM-FV2		144×96
15	CESM2-WACCM	National Center for Atmospheric Research, Climate and Global Dynamics	288×192
16	CESM2	Laboratory, USA	288×192
17	CESM2- WACCM-FV2		144×96
18	CanESM5	Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Canada	128×64
19	E3SM-1-0	LLNL, ANL, BNL, LANL, LBNL, ORNL, PNNL and SNL, USA	360×180
20	EC-Earth3-Veg		512×256
21	EC-Earth3	EC-Earth consortium (27 institutions in Europe)	512×256
22	FGOALS-f3-L	Chinese Academy of Sciences, China	288×180
23	INM-CM5-0	Institute for Numerical Mathematics, Russian Academy of Science, Russia	180×120
24	KACE-1-0-G	National Institute of Meteorological Sciences/Korea Meteorological Administration, Republic of Korea	192×144
25	NESM3	Nanjing University of Information Science and Technology, China	192×96
26	NorCPM1	NorESM Climate modeling Consortium consisting of CICERO, MET-Norway, NERSC, NILU), UIB, UIO and UNI, Norway	144×96
27	SAM0-UNICON	Seoul National University, Republic of Korea	288×192
28	TaiESM1	Research Center for Environmental Changes, Academia Sinica, Taiwan	288×192

Table 3: Total correlations between the cloud albedo and different aerosol types, and the partial correlations to eliminate the influence of three meteorological parameters individually or simultaneously under different LWP conditions. The value above is under the condition of $LWP \leq 650 \text{ g m}^{-2}$. The value in parentheses is under the condition of $650 \text{ g m}^{-2} < LWP \leq 120 \text{ g m}^{-2}$.

	BC	DU	OC	SO ₂	SO ₄	SS
Total correlation	0.25 (-0.42)	-0.49 (-0.51)	0.23 (-0.40)	0.34 (-0.11)	0.34 (-0.51)	-0.26 (-0.01)
Omega900	0.23 (-0.43)	-0.46 (-0.52)	0.22 (-0.41)	0.32 (-0.11)	0.35 (-0.51)	-0.22 (-0.01)
RH850	0.18 (-0.28)	-0.50 (-0.43)	0.17 (-0.26)	0.29 (-0.05)	0.29 (-0.42)	-0.24 (0.03)
LWP	0.35 (-0.31)	-0.34 (-0.31)	0.33 (-0.31)	0.43 (0.13)	0.33 (-0.46)	-0.15 (-0.10)
All parameters	0.24 (-0.07)	-0.31 (-0.12)	0.23 (-0.11)	0.36 (0.25)	0.25 (-0.22)	-0.07 (-0.05)
	BC	DU	OC	SO ₂	SO ₄	SS
Total correlation	0.21 (-0.47)	-0.51 (-0.49)	0.20 (-0.45)	0.32 (-0.10)	0.36 (-0.55)	-0.29 (0.03)
Omega700	0.18 (-0.45)	-0.46 (-0.51)	0.18 (-0.43)	0.27 (-0.11)	0.32 (-0.54)	-0.25 (0.09)
Omega900	0.20 (-0.50)	-0.48 (-0.54)	0.20 (-0.48)	0.30 (-0.12)	0.37 (-0.55)	-0.25 (0.03)
RH700	0.21 (-0.46)	-0.45 (-0.50)	0.22 (-0.44)	0.25 (-0.06)	0.36 (-0.55)	-0.23 (0.01)
EIS	0.14 (-0.30)	-0.38 (-0.43)	0.14 (-0.26)	0.29 (-0.20)	0.18 (-0.33)	-0.14 (0.03)
SSTadv	-0.03 (-0.29)	-0.43 (-0.36)	-0.05 (-0.27)	0.13 (0.05)	0.13 (-0.37)	-0.29 (0.17)
LWP	0.33 (-0.40)	-0.31 (-0.35)	0.30 (-0.39)	0.42 (0.09)	0.34 (-0.53)	-0.15 (-0.08)
All parameters	0.14 (-0.01)	-0.23 (-0.02)	0.11 (-0.03)	0.24 (0.12)	0.20 (0.05)	-0.15 (-0.07)



815 **Figure 1: Near-global distribution of combined planetary albedo averaged from Aqua and Terra during 2003-2014. Red rectangular boxes indicate the five regions used chosen for the analysis: (A1) Peruvian, (A2) Namibian (A3) Californian, (A4) Australian and (A5) Canarian.**














Figure 2: The estimated long-term mean cloud albedo and corresponding correlation coefficient from the relation between planetary albedo and cloud fraction (a-e) from satellite observations (black symbol), 11 AMIP5 (red symbols) and 11 AMIP6 (blue symbols) models during 2003-2008, and (f-j) from satellite observations and 29 AMIP6 models during 2003-2014, over the (a, f) Peruvian, (b, g) Namibian (c, h) Californian, (d, i) Australian and (e, j) Canarian regions.

820



— OBS-OR — OBS-Smooth
 — AMIP5-MEM-OR — AMIP5-MEM-Smooth
 — AMIP6-MEM-OR — AMIP6-MEM-Smooth

— OBS-OR — OBS-Smooth
 — MEM-OR — MEM-Smooth

825 **Figure 3: Monthly mean time series of estimated cloud albedo (a-e) from AMIP5 and AMIP6 multimodel ensemble mean during 2003-2008, and (f-j) from AMIP6 multimodel ensemble mean during 2003-2014 compared with satellite observations, over the (a, f) Peruvian, (b, g) Namibian (c, h) Californian, (d, i) Australian and (e, j) Canarian regions.**






Figure 4: Taylor diagram for monthly estimated cloud albedo between individual AMIP6 model and satellite observations during 2003-2014 over the (a) Peruvian, (b) Namibian (c) Californian, (d) Australian and (e) Canarian regions. The green circles indicate the centered root mean square error.


(a) Peruvian($R_5=0.81, P_5=0.00, R_6=0.92, P_6=0.00$)




(b) Namibian($R_5=0.86, P_5=0.00, R_6=0.89, P_6=0.00$)




(c) Californian($R_5=0.82, P_5=0.00, R_6=0.92, P_6=0.00$)




(d) Australian($R_5=0.93, P_5=0.00, R_6=0.92, P_6=0.00$)




(e) Canarian($R_5=0.22, P_5=0.49, R_6=0.53, P_6=0.08$)




(f) Peruvian($R=0.95, P=0.00$)




(g) Namibian($R=0.92, P=0.00$)




(h) Californian($R=0.76, P=0.00$)




(i) Australian($R=0.93, P=0.00$)



(j) Canarian($R=-0.59, P=0.04$)



835 **Figure 5: Annual cycles of the cloud albedo estimated by (a-e) AMIP5 and AMIP6 multimodel ensemble mean during 2003-2008, and (f-j) AMIP6 multimodel ensemble mean during 2003-2014 compared with satellite observations, over the (a, f) Peruvian, (b, g) Namibian (c, h) Californian, (d, i) Australian and (e, j) Canarian regions. The green and red shading areas indicate the range of the cloud albedo simulated by AMIP5 and AMIP6 models, respectively. The temporal correlations (R5/R6/R value) and P5/P6/P value (if P5/P6/P < 0.10, indicating the correlation R5/R6/R is significant) for the seasonal cycles of the cloud albedo obtained from satellite-based observations and models are given in parentheses.**



840 **Figure 6:** The (a) regression coefficients and corresponding (b) relative contribution of each predictor variables relating to cloud albedo from the multilinear regression models under two LWP conditions: $LWP \leq 60\text{--}65 \text{ g m}^{-2}$ (blue) and $60\text{--}65 \text{ g m}^{-2} < LWP \leq 120 \text{ g m}^{-2}$ (yellow). Note that for ease of comparison, ~~eight~~11 variables are given in the figure, variables without values are not predictive variables of the sample group. And the satellite- and model-driven normalized cloud albedo trained in two sample

groups: (c) $LWP \leq 6065 \text{ g m}^{-2}$ and (d) $6065 \text{ g m}^{-2} < LWP \leq 120 \text{ g m}^{-2}$. The correlations (R value) between satellite- and model-driven normalized cloud albedo are given in parentheses.

845