

This manuscript investigated chemical environment for surface O<sub>3</sub> for six major industrial regions across China in summer 2016. Detailed chemistry-climate model simulations were employed to diagnose ozone sensitivity to precursors and contrast the effectiveness of different measures to reduce surface O<sub>3</sub> concentrations. This manuscript is helpful to understand ozone pollution mechanism in Chinese cities, and within the scope of ACP. I think it is publishable in ACP after my following concerns are addressed.

Line 215: The gross rate of production  $P(O_3)$  actually represents the production rate of  $O_x$  ( $O_3 + NO_2$ ) through the reaction  $HO_2$  ( $RO_2$ ) +  $NO$ . Therefore, the net ozone production rate should include the loss term  $NO_2 + OH$  (Wang et al., 2019. doi.org/10.5194/acp-19-9413-2019). In addition to  $OH + NO_2$  and  $RO_2 + NO_2$ , the loss of  $NO_x$  should also include  $RO_2 + NO$  and  $OH + HONO$ . When calculating OPE, please give specific quantification even though these reactions play a minor role in the loss of  $NO_x$ .

Figure 4 shows significant underestimation for  $NO_2$  in daytime, but overestimation for  $NO_2$  at nighttime. The overestimation of  $NO_2$  at night maybe related to underestimated nighttime chemistry such as the removal of  $NO_3$  and  $N_2O_5$  through heterogeneous uptake (Li et al., 2018; Li et al., 2019). A short discuss should be performed. Additionally, how do these underestimation and overestimation for  $NO_2$  influence your diagnosis of ozone sensitivity? For example, the underestimation of  $NO_2$  in Chongqing will lead to more  $NO_x$ -limited, which likely misleads the actual situation.

Figure 8. shows ozone increased from 70 ppb to over 80 ppb during 2013-2019. However, observed ozone concentrations in Beijing didn't increased significantly during the period or decreased after 2015 in spite that ozone increased over North China Plain (Lu et al., 2018. DOI: 10.1021/acs.estlett.8b00366; Tang et al., 2020. doi.org/10.1016/j.atmosres.2020.105333). This needs further explanations.

Line 270: How do you obtain VOC and  $NO_x$  emissions in 2018 and 2019 given that Cheng et al (2019) just estimated emissions during 2013-2017. Please give specific description.

Line 145: There are only 450 measurement stations in 2013, growing to 1,500 stations in 2017 and 1670 stations in 2019.

Line 300: "summer-mean ozone" should be "daily mean ozone".

#### references

- Li, J., Chen, X., Wang, Z., Du, H., Yang, W., Sun, Y., Hu, B., Li, J., Wang, W., and Wang, T.: Radiative and heterogeneous chemical effects of aerosols on ozone and inorganic aerosols over East Asia, *Science of the Total Environment*, 622, 1327-1342, 2018.
- Li, K., Jacob, D. J., Liao, H., Zhu, J., Shah, V., Shen, L., Bates, K. H., Zhang, Q., and Zhai, S.: A two-pollutant strategy for improving ozone and particulate air quality in China, *Nature Geoscience*, 12,

906-910, 10.1038/s41561-019-0464-x, 2019.