



Revealing the sulfur dioxide emission reductions in China by assimilating surface observations in WRF-Chem

Tie Dai^{1,2*}, Yueming Cheng^{1,2}, Daisuke Goto³, Yingruo Li⁴, Xiao Tang⁵, Guangyu Shi^{1,2}, and Teruyuki Nakajima³

5 ¹State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China.

²Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing University of Information Science and Technology, Nanjing, China.

³National Institute for Environmental Studies, Tsukuba, Japan.

10 ⁴Environmental Meteorology Forecast Center of Beijing-Tianjin-Hebei, China Meteorological Administration, Beijing, China

⁵State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

15 *Correspondence to:* Tie Dai (daitie@mail.iap.ac.cn)

Abstract. The anthropogenic emission of the sulfur dioxide (SO₂) over China has significantly declined as the consequence of clean air actions. In this study, we have developed a new emission inversion system based on a Four-Dimensional Local Ensemble Transform Kalman Filter (4D-LETKF) and the Weather Research and Forecasting model coupled with Chemistry (WRF-Chem) to dynamically update the SO₂ emission grid by grid over China by assimilating the ground-based hourly SO₂ observations. Sensitivity tests for the assimilation system have been conducted firstly to tune four system parameters: ensemble size, horizontal and temporal localization lengths, and perturbation size. Our results reveal that the same random perturbation factors used throughout the whole model grids with assimilating observations within about 180 km can efficiently optimize the SO₂ emission, whereas the ensemble size has only little effect. The temporal localization by assimilating only the subsequent hourly observations can reveal the diurnal variation of the SO₂ emission, which is better than that to update the the magnitude of SO₂ emission every 12 hours by assimilating all the observations within the 12-hour window. The inverted SO₂ emission over China in November 2016 has declined by an average of 49.4% since 2010, which is well in agreement with the “bottom-up” estimation of 48.0%. Larger reductions of SO₂ emission are found over the priori higher source regions such as the Yangtze River Delta (YRD). The simulated SO₂ surface mass concentrations using two distinguished chemical reaction mechanisms are both much more comparable to the observations with the newly inverted SO₂ emission than those with the priori emission. These indicate that the newly developed emission inversion system can



efficiently update the SO₂ emissions based on the routine surface SO₂ observations. The reduced SO₂ emission induces the sulfate and PM_{2.5} surface concentrations to decrease up to 10 $\mu\text{g m}^{-3}$ over the center China.

1 Introduction

35 China and India are the top two emitters of the anthropogenic sulfur dioxide (SO₂) in the world (Li et al., 2017a). SO₂ is a toxic air pollutant and the precursor of sulfate aerosol, leading to the acidification of the atmosphere and the current heavy haze problem in China (Wang et al., 2016a;Huang et al., 2014;Yao et al., 2018). Sulfate aerosol can further perturb the radiative energy budget on Earth through directly scattering solar radiation (Goto et al., 2011) and hydrological cycle by aerosol-cloud interactions (Ramanathan et al., 2001;Sato et al., 2018;Rosenfeld et al., 2019). Hydrophilic polluted
40 continental aerosols such as sulfate serve as cloud condensation nuclei (CCN) and thus have a substantial effect on cloud properties and the initiation of precipitation (Rosenfeld et al., 2008). This, in turn, plays a key role in the climate system. To mitigate climate change and control air quality, the emission control policies especially for SO₂ implemented by China since 2006 cover all the major source sectors and have become increasingly stringent over time (Zhang et al., 2012). Consequently, the decreasing trends of SO₂ loading over China have been revealed by satellite observations, demonstrating SO₂ emissions
45 in China have declined by 75% during 2007-2016 (Wang et al., 2018;Li et al., 2017a). The relative change rate of SO₂ emission in China during 2010-2017 is also estimated as -62% by using the bottom-up emission inventory (Zheng et al., 2018).

The timely precise emission inventories such as SO₂ are the primary inputs to models for air quality prediction and mitigation. All the atmospheric chemistry and aerosol models rely on their descriptions of the emissions virtually, which are
50 mostly from the “bottom-up” emission inventories. The “bottom-up” emission inventories are compiled based on indirect information such as activity data and emission factors (Zhang et al., 2009;Kurokawa et al., 2013;Zheng et al., 2018). Due to the uncertainties of the activity rates and emission factors, large discrepancies of global and regional emissions are identified among different emission inventories (Li et al., 2018;Granier et al., 2011). This demonstrates that there is still no consensus on the best estimates for the emissions of atmospheric compounds. Moreover, the “bottom-up” anthropogenic emission
55 inventories often lag several years behind the present and may quickly become outdated (Zheng et al., 2018), leaving the model without up-to-date emission inventories.

The emission inversion approach can feed historical and near-real-time observations into the models, providing a top-down approach to estimate and timely update the primary emissions of air pollutants (Streets et al., 2013). Generally speaking, variational and ensemble data assimilation approaches are the two widely used methodologies to estimate the emission
60 fluxes of gases (such as NO_x, CO, VOCs) (Tang et al., 2011;Qu et al., 2017;Wu et al., 2020;Miyazaki et al., 2012b;Cheng et al., 2010) and/or aerosols (Dai et al., 2019a;Cohen and Wang, 2014;Peng et al., 2017;Yumimoto et al., 2008). The SO₂ emission inventories over China were updated on monthly or seasonal time scales assuming a linear relationship between SO₂ emissions and satellite observed SO₂ column amounts (Koukouli et al., 2018;Lee et al., 2011), known as the mass balance approach (Martin, 2003), although the sulfur chemistry especially in polluted areas as well as by the interactions of



65 clouds should be nonlinear (Goto et al., 2011;Liao et al., 2003). Fioletov et al. (2015) described a new mass balance
approach to simultaneously estimate the SO₂ lifetimes and emissions from large SO₂ point sources using satellite
measurements. Based on the variational data assimilation approach in the framework of the GEOS-Chem adjoint model,
Wang et al. (2016b) developed a new sophisticated inverse modeling (IM) method to timely update monthly anthropogenic
SO₂ emissions by assimilating the Ozone Monitoring Instrument (OMI) SO₂ satellite measurements. The nonlinear full
70 sulfur chemistry and lifecycle in the atmosphere were accounted for the first time to conduct the top-down estimation of the
anthropogenic SO₂ emissions from the GEOS-Chem adjoint model (Wang et al., 2016b). However, the great limitation to the
application of variational data assimilation approach is the requirement of developing the complicated adjoint model (Henze
et al., 2007). The ensemble data assimilation approach requires neither linearization of the observation operator and nor an
adjoint model, therefore it is much more easily implemented and flexible (Evensen, 2003). Additionally, the ensemble data
75 assimilation and the variational data assimilation use the flow-dependent and pre-calculated model error covariances
respectively (Descombes et al., 2015). Based on the Ensemble Square Root Filter (EnSRF) approach (Chen et al., 2019a), the
recent SO₂ emission changes from the year 2010 in China were successfully updated to improve the model forecast skill.
Retrievals of SO₂ from satellite-based spectrometers are often contaminated by factors such as interference between ozone
and SO₂, and there are significant regional differences between different satellite instruments (Fioletov et al., 2013). This
80 subsequently induces the inconsistency of the inversed regional emissions by assimilating different satellite observations
(Lee et al., 2011). Meanwhile, satellite observations are usually assimilated on the monthly time scale due to data availability.
Compared with satellite observations, the surface SO₂ observations have higher accuracy and temporal frequency. Therefore,
the assimilation of intensive direct surface SO₂ observations can provide more spatial-temporal characteristics of emission
(Chen et al., 2019a). The China National Environmental Monitoring Center (CNEMC) started to monitor hourly
85 concentrations of PM_{2.5} (particulate matter with diameter ≤ 2.5 micrometers), PM₁₀, SO₂, NO₂, CO and O₃ since 2012, and it
had included 1436 monitoring sites from 369 cities by March 2017 (Wu et al., 2018). Those important direct intensive
surface SO₂ observations provide a new chance to estimate the more spatial-temporal characteristics of the SO₂ emission in
China (Chen et al., 2019a).

Due to the limited ensemble members, the Ensemble Kalman filter (EnKF) generally has a spurious long distance correlation
90 problem (Houtekamer and Mitchell, 2001;Miyazaki et al., 2012a). Compared with the EnKF, the Local Ensemble Transform
Kalman Filter (LETKF) can assimilate measurements simultaneously over different model grids in the parallel architecture
(Miyoshi et al., 2007;Hunt et al., 2007). Generally speaking, the LETKF computational time is robust with increasing
observations, while that of most other ensemble Kalman filter is essentially proportional to the number of observations
(Miyoshi et al., 2007). Moreover, the global analysis is linear combinations of the ensemble members in different regions,
95 which is not confined to the limited ensemble members and provides better results in many cases (Ott et al., 2004). A Four-
Dimensional LETKF (4D-LETKF) was recently developed to assimilate hourly aerosol optical properties observed by
satellite, which can avoid frequent switching between the assimilation and the ensemble aerosol forecasting to significantly
reduce computational load (Dai et al., 2019b). In current study, we implement a 4D-LETKF in the Weather Research and



Forecasting model coupled with Chemistry (WRF-Chem). Our major objectives are to investigate whether 4D-LETKF
100 together with the intensive CNEMC SO₂ observations can be applied to quantitatively estimate the spatially resolved
changes of SO₂ emissions in China and how sensitive are the estimated SO₂ emissions to the system parameters of the 4D-
LETKF.

The reminder of the paper is organized as follows. In section 2, the methodology of our emission inversion system is
described in detail. Sect. 3 presents our experimental designs and purposes. The emission inversion results and validations
105 are provided in Sect. 4 before concluding in Sect. 5.

2 Methodology

In order to optimize the SO₂ emissions in this study, we need to formally minimize a scalar cost function J in a Bayesian
framework (Hunt et al., 2007; Huneus et al., 2012). J can be formulated as the sum of the departures of a potential gridded
110 SO₂ emissions x and the corresponding simulated SO₂ surface mass concentrations to the a priori SO₂ emissions x^f and the
CNEMC observed surface SO₂ concentrations y^o :

$$J(x) = 1/2 (x - x^f)^T B^{-1} (x - x^f) + 1/2 (H(x) - y^o)^T R^{-1} (H(x) - y^o) \quad (1)$$

where H is the observation operator that forward the SO₂ emissions to the simulated CNEMC measurements; B and R are
the covariance matrix of the error statistics of the a priori SO₂ emissions and CNEMC observations.

115 2.1 Forward model and observation operator

The relationship between the emission and the surface concentration of short-lived reactive gas SO₂ is mainly determined by
the atmospheric chemical reactions, transport and deposition. The fully coupled “online” Weather Research and Forecasting
model coupled with Chemistry (WRF-Chem) version 4.1.2 (Grell et al., 2005) is served as the forward model to relate the
SO₂ emissions to the simulated observations of surface mass concentration in current study, which can reflect the complex
120 nonlinear relationship between atmospheric chemical concentrations and emissions. Our primary aim is to understand how
sensitive are the estimated SO₂ emissions to the parameters of the assimilation system, which requires huge computing
resources for sensitivity experiments as described later. Therefore, the model is configured with a domain covering most of
China as shown in Fig. 1 with a relatively low horizontal resolution of 50 km and 32 vertical levels (Snyder et al., 2015). A
state-of-the-art and highly nonlinear gas phase chemical mechanism named the second generation Regional Acid Deposition
125 Model (RADM2) (Stockwell et al., 1990) coupled with the Goddard Global Ozone Chemistry Aerosol Radiation and
Transport aerosol model (Chin et al., 2000; Chin et al., 2002) (i.e., chem_opt = 303) is adopted to simulate the atmospheric
sulfur cycle. The RRTMG radiation scheme with prognostic aerosols is selected to consider the aerosol direct effect on
atmospheric radiation and photolysis calculations (Iacono et al., 2008). The other main selected physics are identical to those
of Dai et al. (2019a). The initial and lateral boundary meteorological conditions are from the NCEP Final (FNL) Analysis.
130 To reduce the uncertainties associated with the meteorological fields and facilitate a more straightforward comparison of
simulations and observations, the predicted wind (u , v), temperature (t), and specific humidity (q) by WRF dynamical core
are also nudged to the NCEP FNL analysis every 6 hours (Dai et al., 2018). The WRF-Chem simulated surface gridded SO₂



volume mixing ratios in the unit of parts per million (*ppmv*) are firstly converted to micrograms per cubic meter ($\mu\text{g}/\text{m}^3$) for comparing to the observations (Chen et al., 2019a) and then linearly interpolated to the CNEMC site locations.

135 2.2 SO₂ observations and uncertainties of CNEMC

The quality-assured and controlled measurements of hourly SO₂ surface mass concentration from the CNEMC, which is partly purposefully built for assimilation (Wu et al., 2018), are used to minimize the cost function J . There are a total of 1424 sites in November 2016, and those sites span most of central and eastern China and primarily locate in urban and suburban areas (Peng et al., 2017). Due to unresolved emission variations between urban and suburban, model may have large
140 representativeness errors. To overcome the spatial scale gaps and to produce more representative observations, the super-observation is adopted to average all observations located within a model grid cell (Miyazaki et al., 2012a). Altogether 463 of 7221 model grid cells are covered by the super-observations (Fig. 1). To independently verify the assimilation results, we further randomly eliminate the super-observations located in 155 of the 463 grid cells to be assimilated. The observation error covariance matrix R is assumed diagonal, in other word, the observational error covariance is assumed uncorrelated.
145 The observation error of CNEMC is calculated as same as Chen et al. (2019a), which contains both the measurement and representativeness errors. In the assimilation data quality control process, SO₂ observation leading to absolute innovation exceeding three times of the prior total spread is considered as an outlier and discarded. The innovation is calculated as observation minus the model simulated ensemble mean observation determined from the first guess filed, and the prior total spread is the square root of the sum of the background ensemble variance and the observational error variance (Chen et al.,
150 2019a; Rubin et al., 2016).

2.3 4D-LETKF

The 4D-LETKF assimilation approach generalizes a flow-dependent B from ensemble simulation and finds the minimum of the cost function J as following five formulas (Cheng et al., 2019):

$$\bar{x}^a = \bar{x}^f + X^f \bar{w}^a \quad (2)$$

$$\bar{w}^a = \tilde{P}^a (Y^f)^T R^{-1} (y^o - \bar{y}^f) \quad (3)$$

$$\tilde{P}^a = [(k - 1)I + (Y^f)^T R^{-1} Y^f]^{-1} \quad (4)$$

$$X^a = X^f W^a \quad (5)$$

$$W^a = [(k - 1)\tilde{P}^a]^{1/2} \quad (6)$$

where \bar{x}^f and \bar{x}^a represent the ensemble mean of the first guess (*a priori*) and analysis (*a posteriori*) SO₂ emissions in this
160 study; the ensemble perturbation matrix X is calculated as $x(i) - \bar{x}$, $\{i = 1, 2, \dots, k\}$, which k represents the ensemble size; the matrix \bar{w}^a is the Kalman gain, which specifies the increment between the first guess and the analysis; the vector \bar{y}^f represents the first guess SO₂ surface concentrations averaged over the ensemble members; the matrix Y^f is calculated as $y^f(i) - \bar{y}^f$, $\{i = 1, 2, \dots, k\}$. The ensemble analyses are calculated as the sum of the \bar{x}^a and each of the columns of X^a , which is serving as part of a priori emission information for the next analysis as described later.

165 As shown in Fig. 2, each assimilation cycle with 4D-LETKF includes two steps: a first guess and a state analysis. In our



implementation, the first guess is the WRF-Chem ensemble forecasting for 12 hours with hourly model output. The state analysis optimizes the SO₂ emissions in the past 12 hours. The advantages of 4D-LETKF used here are threefold: (1) each member of the ensemble WRF-Chem simulations is continuously integrated for 12 hours, therefore, this avoids frequent switching between the ensemble WRF-Chem forecasts and the assimilation (Peng et al., 2017; Chen et al., 2019a); (2) the asynchronous observations can be assimilated to optimize the current state (Hunt et al., 2007; Dai et al., 2019b); (3) the assimilation window time of 12 hours could avoid filter convergence and divergence by finite ensemble samples, since more frequent assimilation forces the experiments more closer, inducing the underestimation of the model spread and overconfidence in the first guess state estimate (Schutgens et al., 2010; Miyazaki et al., 2012a; Hunt et al., 2007).

2.4 State variable and forecast model for emission

In this study, the state variable to be optimized is the SO₂ emission. A forecast model for emission is required to propagate observation information and determine the first guess for the next assimilation cycle (Miyazaki et al., 2012a). We adopt the same forecast model for SO₂ emission proposed by Chen et al. (2019a). The forecast model for SO₂ emission weights 75% and 25% toward the SO₂ emission ensemble $E_{t_n}^a$ from the previous analysis and the static initial prior ensemble E_{t_0} as following formula:

$$E_{t_{n+1}}^f = 0.75 \times M E_{t_n}^a M^T + 0.25 \times E_{t_0} \quad (7)$$

where M is the identity matrix. The initial prior ensemble of SO₂ emission is generated by perturbing the freely public available MIX Asian inventory for November 2010 (Li et al., 2017b), and the perturbations follow the lognormal distribution. The mean and the variance of the perturbations are equal to 1 and the MIX SO₂ uncertainty (i.e., 35%) (Li et al., 2017b). In MIX inventory, anthropogenic emissions are aggregated into five sectors: power, industry, residential, transportation, and agriculture. However, only the combined total emission is used in the model and updated in the analysis. It aims to decrease the degree of freedom in the analysis (Miyazaki et al., 2012a). Ten chemical species including both gaseous and aerosol species are included in MIX inventory (Li et al., 2017b). The original monthly MIX anthropogenic emissions with a horizontal resolution of 0.25°×0.25° are remapped to the model resolution of 50 km. The residential, transportation, and agriculture emissions are allocated in the lowest model layer, whereas the power and industry emissions are allocated in the lowest seven model layers with the vertical profiles of the emission factors from the Model Inter-Comparison Study for Asia (MICS-Asia) phase III (Chen et al., 2019b). An improved speciation framework for mapping Asian anthropogenic emissions of non-methane volatile organic compounds (NMVOC) to multiple chemical mechanisms (Li et al., 2014), is adopted to prepare the initial hourly anthropogenic emissions every 12 hours with two separated emission files (i.e., `io_style_emissions` = 2). We do not apply any diurnal variation for the MIX emissions. Therefore, the initial priori emissions are identical throughout the 24 hours. The emissions of aerosol species for WRF-Chem are prepared according to the study of Peng et al. (2017). Notably, only the SO₂ emission is perturbed and optimized by CNEMC SO₂ observations in this study.

The chemical initial conditions (i.e., atmospheric SO₂ concentrations) for the next forward simulation of the WRF-Chem ensemble are also needed to be updated with the optimal emission ensemble from the previous analysis (Peng et al.,



2015;Peters et al., 2005), and this is achieved by recalculation of the WRF-Chem ensemble with the optimized emissions
200 (Fig. 2). In other word, the WRF-Chem ensemble is performed twice in one assimilation cycle.

3 Experimental design

The effectiveness of 4D-LETKF is highly dependent on having sufficient spread in the ensemble members in order for the
observations to impact the first guess (Rubin et al., 2016;Dai et al., 2019b;Hunt et al., 2007). The ensembles represent the
205 uncertainty in the model first guess, therefore, the method for generating the ensemble is an important consideration for an
optimal “top-down” emission inversion. Meanwhile, 4D-LETKF allows a flexible choice of observations to be assimilated
for a specific grid point through horizontal, vertical, and temporal observation localizations (Miyoshi et al., 2007;Dai et al.,
2019b;Cheng et al., 2019). The observation localization gradually reduces the effect of an observation as the increasing
departure from the analysis grid. In this study, the horizontal localization factor is calculated as the Gaussian function
210 (Miyoshi et al., 2007):

$$f(r) = \exp(-r^2/2\sigma^2) \quad (8)$$

where σ is the localization length and r is defined as the physical distance between the observation and the analysis grid,
and we force the localization factor to zero at 3.65 times the localization length (Zhao et al., 2015). In other word, we ignore
observations beyond the cutoff distance. The tuneable horizontal and temporal localization lengths are defined in the
physical distance (km) and time (hour), respectively. The vertical localization is not applied for the SO₂ emission inversion
215 in this study, in other word, we trust the vertical profiles of the emission factors from the Model Inter-Comparison Study for
Asia (MICS-Asia) phase III (Chen et al., 2019b).

A correct choice of the assimilation system parameters such as the ensemble size and correlation length is important to
improve the data assimilation performance (Miyazaki et al., 2012b). A series of sensitivity experiments are performed to
tune the assimilation system as listed in Table 1. A control experiment assuming the same emissions in November 2016 as in
220 November 2010 (i.e., the standard MIX emissions) is conducted as the deterministic simulation to assess the influence of
data assimilation. Considering the GOCART aerosol scheme uses a simple representation of the aerosol chemistry for
reducing the computational load, we also conduct another deterministic simulation using a more sophisticated aerosol
chemical scheme named Model for Simulating Aerosol Interactions and Chemistry (MOSAIC) coupled with the “lumped-
structure” Carbon Bond Mechanism (CBMZ) (Zaveri et al., 2008) (i.e., chem_opt = 9) to investigate the effects of different
225 chemistry and aerosol schemes on SO₂ oxidation. The data assimilation experiments are divided into three groups. In the first
group, same random perturbation factor throughout the whole domain emission grids including vertical and temporal spaces
per member is applied to the MIX SO₂ emission to generate 10 ensemble members for the WRF-Chem ensemble forward
simulation. The same random perturbation factor generates a perfect correlation of emission in both the spatial and temporal
spaces, however, this should not be seen as overly restrictive (Schutgens et al., 2010). Firstly, the “bottom-up” SO₂ emission



230 inventories are to a large extent based on the used activity rates and emission factors (Li et al., 2018). Therefore, with the
same random perturbation factors we effectively create an ensemble of inventories derived with different activity rates and
emission factors. Secondly, the same emission standards for SO₂ emission mitigation are implemented in China (Zheng et al.,
2018), and this induces the SO₂ reductions should be correlated at a certain extent in both spatial and temporal spaces.
Thirdly, the analysis is conducted locally in 4D-LETKF, and the analysis at two grids separated by a distance over about 7.3
235 times the localization length is mostly independent (Schutgens et al., 2010). In this group, the strongest temporal localization
is applied to assimilate only the observations within 1 hour of the local patch center. In other word, the hourly SO₂ emission
is optimized using only the CNEMC SO₂ observation within the subsequent one hour, making the inverted SO₂ emission
variable hour by hour. The difference of the experiments in this group is only the horizontal localization length, which is
assumed as 10km, 30km, 50km, and 100km respectively. The purpose of the experiments in this group is to investigate the
240 effects of horizontal localization length on SO₂ emission inversion. Based on the results in the first group as described latter
in Section 4, the second group of experiments by fixing the horizontal localization length of 50km are subsequently
performed with 10, 20, 40 ensemble members to investigate the effects of ensemble size on SO₂ assimilation. In this group,
we remove the temporal localization to investigate the effects of temporal localization on the SO₂ emission inversion. In
other word, the hourly SO₂ emission is optimized using all the CNEMC SO₂ observations within the 12 hours assimilation
245 window, making the inverted SO₂ emission constant within every 12 hours. In the third group, the experiments are
performed as same as those of the second group except that the ensembles are generated by independently perturbing the
emission in horizontal space but dependently in vertical and temporal spaces. Those last two groups of experiments are used
to investigate the effects of the ensemble size and perturbation factor on SO₂ emission inversion. The sensitivity data
assimilation experiments are all performed for 10 days over the period of 00:00 UTC 8 November to 00:00 UTC 18
250 November 2016. The global model MOZART-4/GEOS-5 provides the initial and lateral boundary conditions used in this
study (<https://www.acom.ucar.edu/wrf-chem/mozart.shtml>, last access: 10 August 2020). The first three days are used as the
spin-up of the data assimilation system, and the subsequent simulation results for one week are analysed in the next section.
Based on the sensitivity tests of the SO₂ emission inversion system, the experiment H50kmT1hE10Ps, which generally
performing better than other experiments, is extended to 00:00 UTC on 1 December 2016. This provides a longer period of
255 20 days to further validate the assimilation system. We also perform a recalculation experiment with the sophisticated
CBMZ/MOSAIC scheme and the updated SO₂ emissions to verify the new SO₂ emission and the associated effects of SO₂
emission reduction.

4 Results

260 4.1 Sensitivity of the inverted SO₂ emission to the assimilation parameters

The spatial distribution of the MIX SO₂ emission in November 2010 at the model lowest layer is shown in Figure 3a, which
is serving as the base of the initial priori SO₂ emission for our experiments in November 2016. The hotspots of the
anthropogenic SO₂ emission are apparently found over the economically developed areas such as the North China Plain



(NCP), the Yangtze River Delta (YRD), and the Peral River Delta (PRD). The Multi-resolution Emission Inventory for
265 China (MEIC, <http://www.meicmodel.org>) developed by Tsinghua University can provide the updated SO₂ emission in
November 2016 (Fig. 3b), which is used as the independent “bottom-up” SO₂ emission to validate our inverted SO₂ emission.
It is apparent that significant negative changes of SO₂ emission are found over the priori higher source regions such as the
NCP, YRD, and PRD between the year of 2010 and 2016, which are in agreement with the changes of the column SO₂
concentrations observed by satellite (Wang et al., 2018). As the consequence of clean air actions (Zheng et al., 2018), the
270 SO₂ emissions over most areas of China show systematic decline from the year 2010 to 2016 (Figure 4a). Can we reveal the
reductions of the SO₂ emission by assimilating the CNEMC observed surface SO₂ concentration?

As shown in Figs. 3c-f and Figs. 4b-e, both spatial distribution and magnitude of the inverted SO₂ emission in November
2016 firstly become closer to the independent MEIC ones but get worse subsequently as increasing the horizontal
localization length of the assimilation system. The experiment with the smallest horizontal localization length (i.e., 10 km)
275 only optimizes the SO₂ emission over the specific grids where there are observations to be assimilated. In such a case, the
significant reductions of the SO₂ emission over the grids with no observation sites are unable to be revealed, such as
Shandong province in the NCP. With a larger localization length, an observation can constrain the emissions in more grids
surrounding the observation and the observation error more gently increases as the distance from the observation location
increased (Hunt et al., 2007). It is obvious that the systemic SO₂ emission reductions especially over the Shandong province
280 are detected by enlarging the horizontal localization length. However, the perfect correlations of the emission perturbations
over the domains with too large horizontal localization length cause spurious error covariance, inducing the more local
emission changes undetectable. This is demonstrated as the inverted SO₂ emissions with a localization length of 100 km tend
to lower than the independent MEIC ones with a mean bias of $-0.44 \text{ mol km}^{-2} \text{ hr}^{-1}$. Generally speaking, the inverted SO₂
emissions with horizontal localization length of 50 km are best in agreement with the MEIC ones with a mean bias of -0.15
285 $\text{mol km}^{-2} \text{ h}^{-1}$ and Root Mean Square Error (RMSE) of $5.34 \text{ mol km}^{-2} \text{ hr}^{-1}$.

With horizontal localization length of 50 km, the spatial distribution of the inverted SO₂ emission by removing the temporal
localization is shown in Fig. 3g. It is clearly found that the inverted SO₂ emissions over the Shandong Province, YRD and
PRD without temporal localization are lower than those with temporal localization, inducing larger negative bias and RMSE
(Fig. 4f). This demonstrates that it is important to reveal the diurnal variations of the SO₂ emission (Wang et al., 2010). The
290 experiment with temporal localization can reveal the hourly variation of the SO₂ emission by assimilating only the
subsequent hourly observations, whereas the experiment without temporal localization only adjust the magnitude of SO₂
emission every 12 hours by assimilating all the observations within the 12-hour window.

As shown in Figs. 3g-i and Figs. 4f-h, there are no significant differences of the horizontal distribution and magnitude of the
inverted SO₂ emission between 10, 20 and 40 ensemble members. This indicates that the ensemble size has little effect on
295 the SO₂ emission inversion when randomly correlated perturbing the emissions. The ensemble forecast with 10 members
seems feasible to reveal the SO₂ reductions in China, although the inverted emissions have not converged properly. This in



turn significantly reduces the required computational resources and time for the forward calculation of the ensemble model, making the dynamical update of air pollutant emissions affordable when assimilating near-real-time observations.

The inverted SO₂ emission with horizontal random uncorrelated perturbations gets closer to the independent MEIC one as increasing the size of the ensemble member (Figs. 3j-l and Figs. 4i-k). However, the performances of the horizontal distribution and magnitude of the inverted SO₂ emission using 40 ensemble members with horizontal random uncorrelated perturbations are even obviously worse than those using 10 ensemble members with horizontal correlated perturbations. This demonstrates that the independent emission perturbations over each model grid tend to underestimate the model spread due to the current limited ensemble members and the cancellation of neighbouring cells (Pagowski and Grell, 2012; Schutgens et al., 2010).

The mean bias and RMSE of SO₂ emission over China by using the MIX SO₂ emission in November 2010 for that in November 2016 are 2.70 and 9.78 mol km⁻²h⁻¹, respectively (Fig. 4a). For the inverted SO₂ emission by data assimilation, the bias and RMSE reduction rates (Miyazaki et al., 2012b) are estimated as follows,

$$\frac{2.70 - |B_{DA}|}{2.70} \times 100. \quad (9)$$

$$\frac{9.78 - |RMSE_{DA}|}{9.78} \times 100. \quad (10)$$

where B_{DA} and $RMSE_{DA}$ are the mean bias and RMSE between the inverted SO₂ emission and the MEIC SO₂ emission in November 2016. As shown in Fig. 5, it is obviously found that (1) the inverted SO₂ emission in every assimilation experiment can both reduce the bias and RMSE; (2) the randomly correlated perturbation factor is superior to the randomly uncorrelated perturbation factor in reducing the bias and RMSE, and it is generally unaffected by the ensemble size; (3) the experiment H50kmT1hE10Ps shows the best performance in both reducing the bias and RMSE, decreasing the bias and RMSE by 94.5% and 45.4% respectively.

4.2 Sensitivity of the surface SO₂ concentration to the emission

Figures 6 and 7 show the horizontal distributions of the biases and RMSEs between the surface SO₂ concentrations simulated in various experiments and the CNEMC observed ones over both the assimilated and independent sites. It is apparent that significant RMSEs and positive biases are found over the priori SO₂ emission hotspot regions such as the NCP, YRD, and PRD in both the two free run experiments, whereas slight RMSEs and negative biases are both found over northwestern China. Furthermore, the horizontal distributions of both the biases and RMSEs of the two free run experiments are generally similar. As given in Table 2, the relative differences of the RMSEs in the FR and FR_CM experiments are both less than 1% over the assimilated and independent sites, although the mean biases in the FR_CM experiment tend to both slightly smaller than those in the FR experiment. Those demonstrate that the biases and RMSEs between the simulated and observed surface SO₂ concentrations are not induced by the uncertainties of the different chemical reaction mechanisms but due to the uncertainties of the used SO₂ emissions. The simulated ensemble mean surface SO₂ concentrations by recalculating the



WRF-Chem with the inverted SO₂ emissions in all assimilation experiments show more comparable to the observations, and the performances of the simulated SO₂ surface concentrations are clearly affected by the inputs of the different inverted SO₂ emissions due to assimilation system parameters. This indicates that the uncertainties of the different chemical reaction mechanisms in simulating SO₂ concentrations are much smaller than those of the SO₂ emissions. In the first group of data assimilation experiments, the largest biases and RMSEs of the simulated and observed SO₂ surface concentrations over both the assimilated and independent sites are found in the H10kmT1hE10Ps experiment. This indicates that the SO₂ emission changes exist grid correlations and the SO₂ emission inversions over only the grids with available assimilated sites are not sufficient to reveal the real SO₂ emission changes in the grids without observation sites. In addition, largest biases and RMSEs over both the assimilated and independent sites are still found in the third group of data assimilation experiments, although the biases and RMSEs are decreasing as the increasement of the ensemble members. This further illustrates there are correlations of the grided SO₂ emission changes and the random perfect correlated emission perturbation factors over the model grids are superior to the random uncorrelated emission perturbations for current emission inversions. The latter is probably due to the current limited ensemble members for reducing the computational resources. However, the sophisticatedly random uncorrelated emission perturbations should have better performances with large or unlimited ensemble members. Similar to the inverted emissions, the experiments in the second group show the ensemble size has little effects on the biases and RMSEs of the SO₂ surface concentrations over both the assimilated and independent sites when the ensemble members are generated by perturbing the emissions perfect correlated over the domain grids. The reductions of the biases of the SO₂ surface concentrations in both the assimilated and independent sites are benefitted from the temporal localization, although the RMSEs are slightly increased. It is interesting that the smallest RMSE of the SO₂ surface concentrations over the independent sites is also found in the H50kmT1hE10Ps experiment with value of 36.20, which the inverted SO₂ emissions are also best in agreement to the independent MEIC ones. This further indicates the assimilation system parameters used in this experiment are suitable for the SO₂ emission inversion, decreasing the biases of SO₂ surface concentrations over assimilated and independent sites by 87.2% and 88.9% respectively.

Figure 8 illustrates the frequency distributions of the deviations of the simulated SO₂ surface concentrations in various experiments minus the observed ones. It is expected that the distributions of the SO₂ surface concentrations deviations for the two free run experiments in China and the three subregions are all positively biased due to the known overestimation of the SO₂ emissions. The distributions of the SO₂ surface concentration deviations with the updated SO₂ emissions in all the data assimilation experiments show reduced biases over both the assimilated and independent sites. However, the distributions of the deviations with the updated SO₂ emissions in the third group of experiments and the H10kmT1hE10Ps experiment are still positively biased, whereas slightly negative biased are found in the second group of experiments and the H100kmT1hE10Ps experiment. The distributions of the SO₂ concentration deviations with the updated SO₂ emissions in the H50kmT1hE10Ps experiment, as expected, shows the best performance with a peak closer to 0 in both the assimilated and independent sites.



4.3 SO₂ reduction in China and associated effects

Based on the aforementioned sensitivity tests of the SO₂ emission inversion system, the experiment H50kmT1hE10Ps is extended to 00:00 UTC on 1 December 2016. This provides a longer period for 20 days to estimate the reduction of the SO₂ emission over China in November over the period 2010-2016. The Bottom-up and Top-down estimations of the SO₂ emission reduction from 2010 to 2016 are calculated by comparing the MEIC and inverted SO₂ emissions by data assimilation in November 2016 to the MIX SO₂ emission in November 2010. As shown in Figure 9, the Top-down estimation of the SO₂ emission reduction over China is 49.4%, which is well agreement with the Bottom-up estimation of 48.0%. In addition, larger SO₂ emission reductions over the three subregions estimated by the Bottom-up approach are correctly revealed by the emission inversion system. The Top-down and Bottom-up estimations of the SO₂ emission reduction over NCP are generally comparable with values of 56.0% and 52.4% respectively. The largest SO₂ emission reductions both with the Top-down and Bottom-up approaches are found over the YRD region with values of 73.1% and 61.8% respectively. The SO₂ emission reduction by Top-down approach are 10% higher than that by the Bottom-up approach over the PRD region. To validate the inverted SO₂ emissions and explore the possible reasons of the overestimation of SO₂ emission reduction over the YRD and PRD by Top-down approach, the time series of the simulated SO₂ surface concentrations in various experiments and the observed ones are shown in Figure 10. The simulated SO₂ surface concentrations especially over the YRD subregion in the two free run experiments show significant positive biases over all the period, revealing the drawback of the prescribed SO₂ emissions in November 2016 as same as that in November 2010. The simulated SO₂ surface concentrations with the inverted SO₂ emissions using both the RADM2/GOCART and CBMZ/MOSAIC chemical reaction mechanisms are much closer to the observations in both the assimilated and independent sites over all the period. This demonstrates the WRF-Chem/4D-LETKF emission inversion system can continuously and dynamically update the SO₂ emissions by assimilating the newly available observations. The SO₂ surface concentrations simulated by the FR_CM experiment are sometime lower than those in the FR experiment especially over the YRD and PRD subregions, indicating the overestimations of the SO₂ emission reduction by the Top-down approach over the YRD and PRD are probability due to the simple aerosol chemistry schemes used in RADM2/GOCART (Chin et al., 2000). This is proved as the simulated SO₂ surface concentrations in YRD with the RADM2/GOCART scheme and the inverted SO₂ emissions over the period 18-22 November 2016 are generally comparable to the observed ones, whereas the simulated SO₂ surface concentrations with the sophisticated CBMZ/MOSAIC scheme and the inverted SO₂ emissions are lower than the observed ones.

Based on the inverted SO₂ emissions from 11 November to 1 December 2016, the daily and diurnal variations of the SO₂ emission reductions over China and the NCP subregion are estimated as shown in Figs. 11a and b respectively. Generally speaking, the daily variation of the SO₂ emission reduction over China is not so significant. Larger SO₂ emission reductions over the period 17 to 19 November in the NCP induced by the first orange alert for heavy winter air pollution in 2016 are clearly detected by the inverted emissions (Shi et al., 2019). Lower SO₂ emission reductions over China and NCP from 21 to 22 November are probably contaminated by the strong cold wind from the northwestern direction, inducing the lowest SO₂



concentrations and underestimating the associated ensemble spread. The latter induces the inverted emission to be overconfident in the background emission (Hunt et al., 2007). The diurnal variations of the SO₂ emission reduction over China and NCP both reveal higher emission reductions in the nighttime, indicating the SO₂ emissions in the nighttime are lower than those in the daytime since the emissions are assumed constant over time in the priori MIX inventory. This is generally reasonable as less human and economic activities happen in the nighttime (Chen et al., 2019b).

Figure 12 shows the spatial distributions of the averaged surface concentrations of the sulfate, ammonium, nitrate, and PM_{2.5} over 11 November to 1 December 2016 simulated with the CBMZ/MOSAIC mechanism and the original MIX emissions, and the absolute and relative changes of the associated aerosol surface concentrations with the newly inverted emissions by data assimilation. It is found that the SO₂ emission reductions induce the sulfate surface concentrations reduced up to 10 μg/m³ (50%) over the center China, and this is due to the sulfate aerosols are dominated by the productions in-cloud oxidations (Chin et al., 2000;Goto et al., 2015) and more cloud are found over the center China (Li et al., 2015;Ma et al., 2014). The nitrate surface concentrations are found slightly increased in the center China as the reductions of sulfate aerosols, and this is due to the emissions of the nitrate precursors (i.e., NO and NO₂) are not updated in this study and NH₄NO₃ is formed only in sulfate-poor aerosols (Zaveri et al., 2008;Chen et al., 2016). The synergy effects of sulfate-nitrate-ammonium induce slightly reductions of ammonium surface concentrations, decreasing the PM_{2.5} surface concentrations about 10 μg m⁻³ (10%) over the center China.

5 Conclusions

The timely precise emission inventories are crucial to air quality prediction and mitigation. To dynamically update the emissions of air pollutants, we have developed a new emission inversion system based on the 4D-LETKF and the fully coupled model named WRF-Chem. Our emission inversion system considers the complex nonlinear relationship between atmospheric chemical concentrations and emissions by ensemble forecasting with perturbed emissions. The emission inversion system is examined to update the outdated MIX SO₂ emissions in November 2010 by assimilating the quality-assured and controlled observations of SO₂ surface concentration from the CNEMC in November 2016. The inverted SO₂ emissions over China by data assimilation for November 2016 are validated with the independent MEIC emissions in November 2016.

Sensitivity tests for the emission inversion system demonstrate that the assumption of the covariance error matrix of the a priori SO₂ emissions has the largest effect on the inverted emissions. The random perfectly correlated emission perturbations throughout the whole model grids with horizontal localization length of 50 km can best reproduce the independent MEIC SO₂ emissions, decreasing the MIX emission bias and RMSE by 94.5% and 45.4% respectively. The independent emission perturbations over each model grid tend to underestimate the model spread due to the current limited ensemble members and the cancellation of neighbouring cells. With the random perfectly correlated emission perturbations, the ensemble size has only little effect on the inverted SO₂ emissions and the ensemble forecast with 10 members seems feasible to reveal the SO₂ reductions in China. The temporal localization by assimilating only the subsequent hourly observations can reveal the



diurnal variation of the SO₂ emission, which is better than that to update the magnitude of SO₂ emission every 12 hours by assimilating all the observations within the 12-hour window.

The known overestimates of the prescribed SO₂ emissions in November 2016 as same as that in November 2010 are successfully detected as the simulated SO₂ surface concentrations especially over the SO₂ emission hotspot subregions with two distinguished chemical reaction mechanisms are both significantly positive biased. The simulated SO₂ surface concentrations with the inverted SO₂ emissions in all assimilation experiments show more comparable to the observations, and the performances of the simulated SO₂ surface concentrations are clearly affected by the inputs of the different inverted SO₂ emissions due to assimilation system parameters. This indicates that the uncertainties of the different chemical reaction mechanisms in simulating SO₂ concentrations are much smaller than those of the SO₂ emissions. The smallest RMSE of the simulated and observed SO₂ surface concentrations over the independent verification sites is also found in the experiment that the inverted SO₂ emissions are best in agreement to the independent MEIC ones, decreasing the biases of SO₂ surface concentrations by 88.9%.

The SO₂ emission reduction over China in November over the period 2010 to 2016 is estimated as 49.4% by assimilating the observations of surface SO₂ concentrations, which is well agreement with the Bottom-up estimation of 48.0%. In addition, larger SO₂ emission reductions over the NCP, YRD and PRD estimated by the Bottom-up approach are correctly revealed by the emission inversion system. Largest SO₂ emission reductions both with the Top-down and Bottom-up approaches are found over the YRD region with values of 73.1% and 61.8% respectively, and the simple parameterizations of the aerosol chemistry in the GOCART scheme may induce the overestimates of the SO₂ emission reductions about 10 percent. The SO₂ emission reductions induce the sulfate and PM_{2.5} surface concentrations to decrease up 10 μg m⁻³ over the center China.

450

Code/Data availability. The WRF-Chem and LETKF source codes are available with the terms and conditions in https://www2.mmm.ucar.edu/wrf/users/download/get_source.html and <https://github.com/takemasa-miyoshi/letkf>. The MIX Asian emission inventory can be found in <http://meicmodel.org/dataset-mix.html>.

455 **Author contributions.** TD designed and performed the experiments used in the study. TD and YC conducted the data analysis. TD prepared the manuscript with help from YC, DG, GS, and TN. DG and TN provided the super-computer resources.

Competing interests. The authors declare that they have no conflict of interest.

460 **Acknowledgments.** This research has been supported by the National Key Research and Development Program of China (grant nos. 2017YFC0209803 and 2016YFC0202001), the Strategic Priority Research Program of the Chinese Academy of



Sciences (grant no. XDA2006010302), the National Natural Science Funds of China (grant nos. 41875133, 41605083, 41905119, and 41590875), and the Youth Innovation Promotion Association CAS (2020078). We are grateful to the relevant researchers who provided NCEP FNL reanalysis data (<https://rda.ucar.edu/datasets/ds083.2/>). The model simulations were performed using the supercomputer resources NIES/HPE Apollo 2000.

465

References

- Chen, D., Liu, Z., Fast, J., and Ban, J.: Simulations of sulfate–nitrate–ammonium (SNA) aerosols during the extreme haze events over northern China in October 2014, *Atmos. Chem. Phys.*, 16, 10707–10724, 10.5194/acp-16-10707-2016, 2016.
- Chen, D., Liu, Z., Ban, J., and Chen, M.: The 2015 and 2016 wintertime air pollution in China: SO₂ emission changes derived from a WRF-Chem/EnKF coupled data assimilation system, *Atmos. Chem. Phys.*, 19, 8619–8650, 10.5194/acp-19-8619-2019, 2019a.
- Chen, L., Gao, Y., Zhang, M., Fu, J. S., Zhu, J., Liao, H., Li, J., Huang, K., Ge, B., Wang, X., Lam, Y. F., Lin, C.-Y., Itahashi, S., Nagashima, T., Kajino, M., Yamaji, K., Wang, Z., and Kurokawa, J.-i.: MICS-Asia III: multi-model comparison and evaluation of aerosol over East Asia, *Atmos. Chem. Phys.*, 19, 11911–11937, 10.5194/acp-19-11911-2019, 2019b.
- Cheng, X., Xu, X., and Ding, G.: An emission source inversion model based on satellite data and its application in air quality forecasts, *Sci. China Earth Sci.*, 53, 752–762, 10.1007/s11430-010-0044-9, 2010.
- Cheng, Y., Dai, T., Goto, D., Schutgens, N. A. J., Shi, G., and Nakajima, T.: Investigating the assimilation of CALIPSO global aerosol vertical observations using a four-dimensional ensemble Kalman filter, *Atmos. Chem. Phys.*, 19, 13445–13467, 10.5194/acp-19-13445-2019, 2019.
- Chin, M., Savoie, D. L., Huebert, B. J., Bandy, A. R., Thornton, D. C., Bates, T. S., Quinn, P. K., Saltzman, E. S., and De Bruyn, W. J.: Atmospheric sulfur cycle simulated in the global model GOCART: Comparison with field observations and regional budgets, *Journal of Geophysical Research: Atmospheres*, 105, 24689–24712, 10.1029/2000jd900385, 2000.
- Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: Tropospheric Aerosol Optical Thickness from the GOCART Model and Comparisons with Satellite and Sun Photometer Measurements, *Journal of the Atmospheric Sciences*, 59, 461–483, 10.1175/1520-0469(2002)059<0461:taotft>2.0.co;2, 2002.
- Cohen, J. B., and Wang, C.: Estimating global black carbon emissions using a top-down Kalman Filter approach, *Journal of Geophysical Research: Atmospheres*, 119, 307–323, 10.1002/2013jd019912, 2014.
- Dai, T., Cheng, Y., Zhang, P., Shi, G., Sekiguchi, M., Suzuki, K., Goto, D., and Nakajima, T.: Impacts of meteorological nudging on the global dust cycle simulated by NICAM coupled with an aerosol model, *Atmospheric Environment*, 190, 99–115, 10.1016/j.atmosenv.2018.07.016, 2018.
- Dai, T., Cheng, Y., Goto, D., Schutgens, N. A. J., Kikuchi, M., Yoshida, M., Shi, G., and Nakajima, T.: Inverting the East Asian Dust Emission Fluxes Using the Ensemble Kalman Smoother and Himawari-8 AODs: A Case Study with WRF-Chem v3.5.1, *Atmosphere*, 10, 10.3390/atmos10090543, 2019a.
- Dai, T., Cheng, Y., Suzuki, K., Goto, D., Kikuchi, M., Schutgens, N. A. J., Yoshida, M., Zhang, P., Husi, L., Shi, G., and Nakajima, T.: Hourly Aerosol Assimilation of Himawari-8 AOT Using the Four-Dimensional Local Ensemble Transform Kalman Filter, *Journal of Advances in Modeling Earth Systems*, 11, 680–711, 10.1029/2018ms001475, 2019b.
- Descombes, G., Auligné, T., Vandenberghe, F., Barker, D. M., and Barré, J.: Generalized background error covariance matrix model (GEN_BE v2.0), *Geoscientific Model Development*, 8, 669–696, 10.5194/gmd-8-669-2015, 2015.



- Evensen, G.: The Ensemble Kalman Filter: theoretical formulation and practical implementation, *Ocean Dynamics*, 53, 343-367, 10.1007/s10236-003-0036-9, 2003.
- 500 Fioletov, V. E., McLinden, C. A., Krotkov, N., Yang, K., Loyola, D. G., Valks, P., Theys, N., Van Roozendaal, M., Nowlan, C. R., Chance, K., Liu, X., Lee, C., and Martin, R. V.: Application of OMI, SCIAMACHY, and GOME-2 satellite SO₂ retrievals for detection of large emission sources, *Journal of Geophysical Research: Atmospheres*, 118, 11,399-311,418, 10.1002/jgrd.50826, 2013.
- Fioletov, V. E., McLinden, C. A., Krotkov, N., and Li, C.: Lifetimes and emissions of SO₂ from point sources estimated from OMI, *Geophys. Res. Lett.*, 42, 1969-1976, 10.1002/2015gl063148, 2015.
- 505 Goto, D., Nakajima, T., Takemura, T., and Sudo, K.: A study of uncertainties in the sulfate distribution and its radiative forcing associated with sulfur chemistry in a global aerosol model, *Atmos. Chem. Phys.*, 11, 10889-10910, 10.5194/acp-11-10889-2011, 2011.
- Goto, D., Nakajima, T., Dai, T., Takemura, T., Kajino, M., Matsui, H., Takami, A., Hatakeyama, S., Sugimoto, N., Shimizu, A., and Ohara, T.: An evaluation of simulated particulate sulfate over East Asia through global model intercomparison, *Journal of Geophysical Research: Atmospheres*, 120, 6247-6270, 10.1002/2014jd021693, 2015.
- 510 Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., Denier van der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J.-F., Liousse, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J.-C., Riahi, K., Schultz, M. G., Smith, S. J., Thompson, A., van Aardenne, J., van der Werf, G. R., and van Vuuren, D. P.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic Change*, 109, 163-190, 10.1007/s10584-011-0154-1, 2011.
- 515 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmospheric Environment*, 39, 6957-6975, 10.1016/j.atmosenv.2005.04.027, 2005.
- Henze, D. K., Hakami, A., and Seinfeld, J. H.: Development of the adjoint of GEOS-Chem, *Atmos. Chem. Phys.*, 7, 2413-2433, 10.5194/acp-7-2413-2007, 2007.
- Houtekamer, P. L., and Mitchell, H. L.: A Sequential Ensemble Kalman Filter for Atmospheric Data Assimilation, *Monthly Weather*
- 520 *Review*, 129, 123-137, 10.1175/1520-0493(2001)129<0123:Asekff>2.0.Co;2, 2001.
- Huang, R. J., Zhang, Y., Bozzetti, C., Ho, K. F., Cao, J. J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., El Haddad, I., and Prevot, A. S.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218-222, 10.1038/nature13774, 2014.
- 525 Huneus, N., Chevallier, F., and Boucher, O.: Estimating aerosol emissions by assimilating observed aerosol optical depth in a global aerosol model, *Atmos. Chem. Phys.*, 12, 4585-4606, 10.5194/acp-12-4585-2012, 2012.
- Hunt, B. R., Kostelich, E. J., and Szunyogh, I.: Efficient data assimilation for spatiotemporal chaos: A local ensemble transform Kalman filter, *Physica D: Nonlinear Phenomena*, 230, 112-126, 10.1016/j.physd.2006.11.008, 2007.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived
- 530 *greenhouse gases: Calculations with the AER radiative transfer models*, *J. Geophys. Res.*, 113, 10.1029/2008jd009944, 2008.
- Koukouli, M. E., Theys, N., Ding, J., Zyrichidou, I., Mijling, B., Balis, D., and van der A, R. J.: Updated SO₂ emission estimates over China using OMI/Aura observations, *Atmospheric Measurement Techniques*, 11, 1817-1832, 10.5194/amt-11-1817-2018, 2018.
- Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K., and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000–2008: Regional Emission inventory in ASia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019-11058, 10.5194/acp-13-11019-2013, 2013.



- Lee, C., Martin, R. V., van Donkelaar, A., Lee, H., Dickerson, R. R., Hains, J. C., Krotkov, N., Richter, A., Vinnikov, K., and Schwab, J. J.: SO₂ emissions and lifetimes: Estimates from inverse modeling using in situ and global, space-based (SCIAMACHY and OMI) observations, *J. Geophys. Res.*, 116, 10.1029/2010jd014758, 2011.
- Li, C., McLinden, C., Fioletov, V., Krotkov, N., Carn, S., Joiner, J., Streets, D., He, H., Ren, X., Li, Z., and Dickerson, R. R.: India Is
540 Overtaking China as the World's Largest Emitter of Anthropogenic Sulfur Dioxide, *Sci Rep*, 7, 14304, 10.1038/s41598-017-14639-8, 2017a.
- Li, J., Huang, J., Stamnes, K., Wang, T., Lv, Q., and Jin, H.: A global survey of cloud overlap based on CALIPSO and CloudSat measurements, *Atmos. Chem. Phys.*, 15, 519-536, 10.5194/acp-15-519-2015, 2015.
- Li, M., Zhang, Q., Streets, D. G., He, K. B., Cheng, Y. F., Emmons, L. K., Huo, H., Kang, S. C., Lu, Z., Shao, M., Su, H., Yu, X., and
545 Zhang, Y.: Mapping Asian anthropogenic emissions of non-methane volatile organic compounds to multiple chemical mechanisms, *Atmos. Chem. Phys.*, 14, 5617-5638, 10.5194/acp-14-5617-2014, 2014.
- Li, M., Zhang, Q., Kurokawa, J.-i., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935-963, 10.5194/acp-17-935-2017, 2017b.
- 550 Li, M., Klimont, Z., Zhang, Q., Martin, R. V., Zheng, B., Heyes, C., Cofala, J., Zhang, Y., and He, K.: Comparison and evaluation of anthropogenic emissions of SO₂ and NO_x over China, *Atmos. Chem. Phys.*, 18, 3433-3456, 10.5194/acp-18-3433-2018, 2018.
- Liao, H., Adams, P. J., Chung, S. H., Seinfeld, J. H., Mickley, L. J., and Jacob, D. J.: Interactions between tropospheric chemistry and aerosols in a unified general circulation model, *Journal of Geophysical Research: Atmospheres*, 108, 4001, 10.1029/2001JD001260, 2003.
- Ma, J., Wu, H., Wang, C., Zhang, X., Li, Z., and Wang, X.: Multiyear satellite and surface observations of cloud fraction over China,
555 *Journal of Geophysical Research: Atmospheres*, 119, 7655-7666, 10.1002/2013jd021413, 2014.
- Martin, R. V.: Global inventory of nitrogen oxide emissions constrained by space-based observations of NO₂ columns, *J. Geophys. Res.*, 108, 10.1029/2003jd003453, 2003.
- Miyazaki, K., Eskes, H. J., and Sudo, K.: Global NO_x emission estimates derived from an assimilation of OMI tropospheric NO₂ columns, *Atmos. Chem. Phys.*, 12, 2263-2288, 10.5194/acp-12-2263-2012, 2012a.
- 560 Miyazaki, K., Eskes, H. J., Sudo, K., Takigawa, M., van Weele, M., and Boersma, K. F.: Simultaneous assimilation of satellite NO₂, O₃, CO, and HNO₃ data for the analysis of tropospheric chemical composition and emissions, *Atmos. Chem. Phys.*, 12, 9545-9579, 10.5194/acp-12-9545-2012, 2012b.
- Miyoshi, T., Yamane, S., and Enomoto, T.: Localizing the Error Covariance by Physical Distances within a Local Ensemble Transform Kalman Filter (LETKF), *SOLA*, 3, 89-92, 10.2151/sola.2007-023, 2007.
- 565 Ott, E., Hunt, B. R., Szunyogh, I., Zimin, A. V., Kostelich, E. J., Corazza, M., Kalnay, E., Patil, D. J., and Yorke, J. A.: A local ensemble Kalman filter for atmospheric data assimilation, *Tellus A*, 56, 415-428, 10.1111/j.1600-0870.2004.00076.x, 2004.
- Pagowski, M., and Grell, G. A.: Experiments with the assimilation of fine aerosols using an ensemble Kalman filter, *Journal of Geophysical Research: Atmospheres*, 117, n/a-n/a, 10.1029/2012jd018333, 2012.
- Peng, Z., Zhang, M., Kou, X., Tian, X., and Ma, X.: A regional carbon data assimilation system and its preliminary evaluation in East Asia,
570 *Atmos. Chem. Phys.*, 15, 1087-1104, 10.5194/acp-15-1087-2015, 2015.
- Peng, Z., Liu, Z., Chen, D., and Ban, J.: Improving PM_{2.5} forecast over China by the joint adjustment of initial conditions and source emissions with an ensemble Kalman filter, *Atmos. Chem. Phys.*, 17, 4837-4855, 10.5194/acp-17-4837-2017, 2017.



- Peters, W., Miller, J. B., Whitaker, J., Denning, A. S., Hirsch, A., Krol, M. C., Zupanski, D., Bruhwiler, L., and Tans, P. P.: An ensemble data assimilation system to estimate CO₂ surface fluxes from atmospheric trace gas observations, *J. Geophys. Res.*, 110, 10.1029/2005jd006157, 2005.
- 575 Qu, Z., Henze, D. K., Capps, S. L., Wang, Y., Xu, X., Wang, J., and Keller, M.: Monthly top-down NO_x emissions for China (2005-2012): A hybrid inversion method and trend analysis, *Journal of Geophysical Research: Atmospheres*, 122, 4600-4625, 10.1002/2016jd025852, 2017.
- Ramanathan, V., Crutzen, P. J., Kiehl, J. T., and Rosenfeld, D.: Aerosols, Climate, and the Hydrological Cycle, *Science*, 294, 2119-2124, 10.1126/science.1064034, 2001.
- 580 Rosenfeld, D., Lohmann, U., Raga, G. B., Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A., and Andreae, M. O.: Flood or Drought: How Do Aerosols Affect Precipitation?, *Science*, 321, 1309, 10.1126/science.1160606, 2008.
- Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T., and Yu, S.: Aerosol-driven droplet concentrations dominate coverage and water of oceanic low-level clouds, *Science*, 363, 10.1126/science.aav0566, 2019.
- 585 Rubin, J. I., Reid, J. S., Hansen, J. A., Anderson, J. L., Collins, N., Hoar, T. J., Hogan, T., Lynch, P., McLay, J., Reynolds, C. A., Sessions, W. R., Westphal, D. L., and Zhang, J.: Development of the Ensemble Navy Aerosol Analysis Prediction System (ENAAAPS) and its application of the Data Assimilation Research Testbed (DART) in support of aerosol forecasting, *Atmos. Chem. Phys.*, 16, 3927-3951, 10.5194/acp-16-3927-2016, 2016.
- Sato, Y., Goto, D., Michibata, T., Suzuki, K., Takemura, T., Tomita, H., and Nakajima, T.: Aerosol effects on cloud water amounts were successfully simulated by a global cloud-system resolving model, *Nature Communications*, 9, 10.1038/s41467-018-03379-6, 2018.
- 590 Schutgens, N. A. J., Miyoshi, T., Takemura, T., and Nakajima, T.: Sensitivity tests for an ensemble Kalman filter for aerosol assimilation, *Atmos. Chem. Phys.*, 10, 6583-6600, 10.5194/acp-10-6583-2010, 2010.
- Shi, Z., Vu, T., Kotthaus, S., Harrison, R. M., Grimmond, S., Yue, S., Zhu, T., Lee, J., Han, Y., Demuzere, M., Dunmore, R. E., Ren, L., Liu, D., Wang, Y., Wild, O., Allan, J., Acton, W. J., Barlow, J., Barratt, B., Beddows, D., Bloss, W. J., Calzolari, G., Carruthers, D., Carslaw, D. C., Chan, Q., Chatzidiakou, L., Chen, Y., Crilley, L., Coe, H., Dai, T., Doherty, R., Duan, F., Fu, P., Ge, B., Ge, M., Guan, D., Hamilton, J. F., He, K., Heal, M., Heard, D., Hewitt, C. N., Hollaway, M., Hu, M., Ji, D., Jiang, X., Jones, R., Kalberer, M., Kelly, F. J., Kramer, L., Langford, B., Lin, C., Lewis, A. C., Li, J., Li, W., Liu, H., Liu, J., Loh, M., Lu, K., Lucarelli, F., Mann, G., McFiggans, G., Miller, M. R., Mills, G., Monk, P., Nemitz, E., Connor, F., Ouyang, B., Palmer, P. I., Percival, C., Popoola, O., Reeves, C., Rickard, A. R., Shao, L., Shi, G., Spracklen, D., Stevenson, D., Sun, Y., Sun, Z., Tao, S., Tong, S., Wang, Q., Wang, W., Wang, X., Wang, X., Wang, Z., Wei, L., Whalley, L., Wu, X., Wu, Z., Xie, P., Yang, F., Zhang, Q., Zhang, Y., Zhang, Y., and Zheng, M.: Introduction to the special issue "In-depth study of air pollution sources and processes within Beijing and its surrounding region (APHH-Beijing)", *Atmos. Chem. Phys.*, 19, 7519-7546, 10.5194/acp-19-7519-2019, 2019.
- 600 Snyder, C., Hacker, J. P., Ha, S. Y., Fossell, K. R., and Berner, J.: Increasing the Skill of Probabilistic Forecasts: Understanding Performance Improvements from Model-Error Representations, *Monthly Weather Review*, 143, 1295-1320, 10.1175/mwr-d-14-00091.1, 2015.
- Stockwell, W. R., Middleton, P., Chang, J. S., and Tang, X.: The second generation regional acid deposition model chemical mechanism for regional air quality modeling, *Journal of Geophysical Research: Atmospheres*, 95, 16343-16367, 10.1029/JD095iD10p16343, 1990.
- Streets, D. G., Canty, T., Carmichael, G. R., de Foy, B., Dickerson, R. R., Duncan, B. N., Edwards, D. P., Haynes, J. A., Henze, D. K., Houyoux, M. R., Jacob, D. J., Krotkov, N. A., Lamsal, L. N., Liu, Y., Lu, Z., Martin, R. V., Pfister, G. G., Pinder, R. W., Salawitch, R. J.,



- 610 and Wecht, K. J.: Emissions estimation from satellite retrievals: A review of current capability, *Atmospheric Environment*, 77, 1011-1042, 10.1016/j.atmosenv.2013.05.051, 2013.
- Tang, X., Zhu, J., Wang, Z. F., and Gbaguidi, A.: Improvement of ozone forecast over Beijing based on ensemble Kalman filter with simultaneous adjustment of initial conditions and emissions, *Atmos. Chem. Phys.*, 11, 12901-12916, 10.5194/acp-11-12901-2011, 2011.
- Wang, G., Zhang, R., Gomez, M. E., Yang, L., Levy Zamora, M., Hu, M., Lin, Y., Peng, J., Guo, S., Meng, J., Li, J., Cheng, C., Hu, T.,
615 Ren, Y., Wang, Y., Gao, J., Cao, J., An, Z., Zhou, W., Li, G., Wang, J., Tian, P., Marrero-Ortiz, W., Secretst, J., Du, Z., Zheng, J., Shang, D., Zeng, L., Shao, M., Wang, W., Huang, Y., Wang, Y., Zhu, Y., Li, Y., Hu, J., Pan, B., Cai, L., Cheng, Y., Ji, Y., Zhang, F., Rosenfeld, D., Liss, P. S., Duce, R. A., Kolb, C. E., and Molina, M. J.: Persistent sulfate formation from London Fog to Chinese haze, *Proc Natl Acad Sci U S A*, 113, 13630-13635, 10.1073/pnas.1616540113, 2016a.
- Wang, T., Wang, P., Theys, N., Tong, D., Hendrick, F., Zhang, Q., and Van Roozendael, M.: Spatial and temporal changes in SO₂ regimes
620 over China in the recent decade and the driving mechanism, *Atmos. Chem. Phys.*, 18, 18063-18078, 10.5194/acp-18-18063-2018, 2018.
- Wang, X., Liang, X.-Z., Jiang, W., Tao, Z., Wang, J. X. L., Liu, H., Han, Z., Liu, S., Zhang, Y., and Grell, G. A.: WRF-Chem simulation of East Asian air quality: Sensitivity to temporal and vertical emissions distributions, *Atmospheric Environment*, 44, 660-669, 10.1016/j.atmosenv.2009.11.011, 2010.
- Wang, Y., Wang, J., Xu, X., Henze, D. K., Wang, Y., and Qu, Z.: A new approach for monthly updates of anthropogenic sulfur dioxide
625 emissions from space: Application to China and implications for air quality forecasts, *Geophys. Res. Lett.*, 43, 9931-9938, 10.1002/2016gl070204, 2016b.
- Wu, H., Tang, X., Wang, Z., Wu, L., Lu, M., Wei, L., and Zhu, J.: Probabilistic Automatic Outlier Detection for Surface Air Quality Measurements from the China National Environmental Monitoring Network, *Advances in Atmospheric Sciences*, 35, 1522-1532, 10.1007/s00376-018-8067-9, 2018.
- 630 Wu, H., Tang, X., Wang, Z., Wu, L., Li, J., Wang, W., Yang, W., and Zhu, J.: High-spatiotemporal-resolution inverse estimation of CO and NO_x emission reductions during emission control periods with a modified ensemble Kalman filter, *Atmospheric Environment*, 236, 10.1016/j.atmosenv.2020.117631, 2020.
- Yao, L., Garmash, O., Bianchi, F., Zheng, J., Yan, C., Kontkanen, J., Junninen, H., Mazon, S. B., Ehn, M., Paasonen, P., Sipilä, M., Wang, M., Wang, X., Xiao, S., Chen, H., Lu, Y., Zhang, B., Wang, D., Fu, Q., Geng, F., Li, L., Wang, H., Qiao, L., Yang, X., Chen, J., Kerminen,
635 V.-M., Petäjä, T., Worsnop, D. R., Kulmala, M., and Wang, L.: Atmospheric new particle formation from sulfuric acid and amines in a Chinese megacity, *Science*, 361, 278, 10.1126/science.aao4839, 2018.
- Yumimoto, K., Uno, I., Sugimoto, N., Shimizu, A., Liu, Z., and Winker, D. M.: Adjoint inversion modeling of Asian dust emission using lidar observations, *Atmos. Chem. Phys.*, 8, 2869-2884, 10.5194/acp-8-2869-2008, 2008.
- Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *J. Geophys. Res.*, 113, 10.1029/2007jd008782, 2008.
640
- Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131-5153, 10.5194/acp-9-5131-2009, 2009.
- Zhang, Q., He, K., and Huo, H.: Cleaning China's air, *Nature*, 484, 161-162, 2012.
- 645 Zhao, Y., Greybush, S. J., Wilson, R. J., Hoffman, R. N., and Kalnay, E.: Impact of assimilation window length on diurnal features in a Mars atmospheric analysis, *Tellus A: Dynamic Meteorology and Oceanography*, 67, 10.3402/tellusa.v67.26042, 2015.



Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095-14111, 10.5194/acp-18-14095-2018, 2018.

650

655

660

665


670

675

680



Figures



685 Figure 1. WRF-Chem model computational domain with the topography. The locations of the assimilated and independent
verification observation sites of the China National Environmental Monitoring Center (CNEMC) are shown as the black
circles and red squares, respectively. The three magenta boxes mark the North China Plain (NCP), the Yangtze River delta
(YRD) and the Peral River delta (PRD) subregions where relatively dense observation sites are available.





Figure 2. Flowchart of the WRF-Chem/4D-LETKF SO₂ emission inversion system by assimilating the SO₂ observations.



690 Figure 3. Spatial distributions of the MIX SO₂ emission in November 2010 (a) and the MEIC SO₂ emission in November 2016 (b) at the model lowest layer. Spatial distributions of the inverted SO₂ emissions in November 2016 in various data assimilation experiments (c-l).




Figure 4. Comparisons of the MIX SO₂ emissions in November 2010 (a) and the inverted SO₂ emission in November 2016 in
695 various data assimilation experiments (b-k) to the MEIC SO₂ emissions in November 2016.





Figure 5. Reductions of the bias and Root Mean Square Error (RMSE) between the inverted SO₂ emissions in various data assimilation experiments and the MEIC ones referring to those between the MIX and MEIC SO₂ emissions.



700 Figure 6. Spatial distributions of the mean biases between the simulated surface SO_2 concentrations in various experiments and the CNEMC observed ones over both the assimilated and independent sites. The locations of the assimilated and independent verification observation sites of the CNEMC are shown as the circles and squares, respectively.






Figure 7. Same as Figure 6 but for the RMSEs.



705

Figure 8. Frequency distributions of the deviations of the simulated SO₂ surface concentrations in various experiments minus the observed ones.



710 Figure 9. SO₂ emission reductions in November over the period 2010 to 2016 in China and the three subregions estimated by
the Bottom-up and Top-down approaches.





Figure 10. Time series of the simulated SO₂ surface concentrations in various experiments and the CNEMC observations.



715 Figure 11. Daily and diurnal variations of the SO₂ emission reductions over China and the NCP subregion based on the inverted emissions.




Figure 12. Spatial distributions of the averaged surface concentrations of the sulfate, ammonium, nitrate and PM_{2.5} over 11 November to 1 December 2016 simulated with the CBMZ/MOSAIC mechanism and the MIX emissions in November 2010, and the absolute and relative changes of the associated aerosol surface concentrations with the updated emissions by data assimilation.

720

725



Tables

Table 1. Experimental design in this study.

Experiments		Design and purpose of the simulation	
Control experiments	FR	Free run using RADM2/GOCART (FR) and CBMZ/MOSAIC (FR_CM) mechanisms with the MIX emission inventory in November 2010 to investigate the effects of different chemistry and aerosol schemes on SO ₂ simulations and provide a reference to evaluate the effects of data assimilation	
	FR_CM		
Data assimilation experiments	H10kmT1hE10Ps H30kmT1hE10Ps H50kmT1hE10Ps H100kmT1hE10Ps	Same random perturbation factors throughout the whole domain emission grids and 12 hours to generate 10 ensembles with assimilated observations within 1h. Experiments with horizontal localization length of 10km, 30km, 50km, and 100km respectively are performed to investigate the effects of horizontal localization length on SO ₂ emission inversion.	
	H50kmT12hE10Ps H50kmT12hE20Ps H50kmT12hE40Ps	Same random perturbation factors throughout the whole domain emission grids and 12 hours to generate 10, 20, 40 ensembles with assimilated observations within 12 h and horizontal localization length of 50km based on above tests. Experiments in this group are performed to investigate the effects of ensemble size on SO ₂ emission inversion.	
	H50kmT12hE10Pi H50kmT12hE20Pi H50kmT12hE40Pi	Horizontal independent random perturbation factors in each emission grid but same throughout 12 hours to generate 10, 20, 40 ensembles with assimilated observations within 12 h and horizontal localization length of 50km. Experiments in this group together with above group are performed to investigate the effects of ensemble size and perturbation factors on SO ₂ emission inversion.	
	Recalculation experiment	CBMZ/MOSAIC with posterior emission	Deterministic simulation with sophisticated CBMZ/MOSAIC scheme is recalculated with the updated SO ₂ emission to verify the updated SO ₂ emissions with an independent mechanism and the associated effects of SO ₂ emission reduction.



Table 2. The mean biases and Root Mean Square Errors (RMSEs) of the simulated SO₂ surface concentrations in various experiments and the CNEMC observed ones over all assimilated and independent sites.

Experiments	Sanity Check		Independent Validation	
	B	RMSE	B	RMSE
FR	44.03	106.04	34.72	78.03
FR_CM	36.49	106.69	27.01	77.45
H10kmT1hE10Ps	12.01	50.89	21.83	59.84
H30kmT1hE10Ps	-4.06	38.20	-0.34	38.57
H50kmT1hE10Ps	-5.65	38.63	-3.84	36.18
H100kmT1hE10Ps	-7.36	39.66	-5.76	36.20
H50kmT12hE10Ps	-9.75	37.36	-7.45	34.59
H50kmT12hE20Ps	-8.80	37.54	-6.20	35.27
H50kmT12hE40Ps	-8.75	37.55	-6.15	35.28
H50kmT12hE10Pi	20.42	75.96	21.62	65.75
H50kmT12hE20Pi	17.38	68.60	20.03	60.90
H50kmT12hE40Pi	13.93	60.12	18.05	57.06