

## Opinion: Papers that shaped Tropospheric Chemistry

Paul S. Monks,<sup>1</sup> A.R. Ravishankara,<sup>2</sup> Erika von Schneidemesser<sup>3</sup> and Roberto Sommariva<sup>1</sup>

1. School of Chemistry, University of Leicester, University Rd., Leicester, LE1 7RH, UK.
2. Departments of Chemistry and Atmospheric Science, Colorado State University, Fort Collins, Colorado, USA.
3. Institute for Advanced Sustainability Studies, Berlinerstrasse 130, 14467 Potsdam, Germany.

### Abstract


Which published papers have transformed our understanding of the chemical processes in the troposphere, and shaped the field of atmospheric chemistry? By way of expert solicitation and interactive peer-review, this paper explores the influence of the ideas in peer-reviewed articles based on the input from our community of atmospheric scientists. We explore how these papers have shaped the development of the field of atmospheric chemistry, and identify the major landmarks in the field of atmospheric chemistry through the lens of those papers' impact on science, legislation and environmental events. We also explore the ways in which one can identify the papers that have most impacted the field and discuss the advantages and disadvantages of the various approaches. Our work highlights the difficulty of creating a simple list and we explore the reasons for this difficulty. The paper also provides a history of the development of our understanding of tropospheric chemistry and points some ways for the future.

## 1. Introduction

Air quality and anthropogenic climate change are two environmental issues of current importance to society. Atmospheric composition is central to both these issues. The atmosphere, and its components, supports life on Earth. In turn, the atmosphere is affected by human population growth and industrialization, as well as all the consequences of those changes. The changes in atmospheric composition also influence the ecosystem on which humans rely.

Air pollution (née composition) and its impacts have a history stretching back to antiquity – see for example the expositions in (Brimblecombe, 1987; Fuller, 2018; Jacobson, 2002; Stern, 1968; Sportisse, 2010; Preining and Davis, 1999; Fowler et al., 2020) and others. Changes in atmospheric composition, with negative impacts particularly on human health (Lelieveld et al., 2015; Landrigan et al., 2018), ecosystems (Fowler et al., 2009) and latterly climate (see for example, (Fiore et al., 2012; von Schneidemesser et al., 2015)), have become primary global concerns during the latter part of the 20<sup>th</sup> and the 21<sup>st</sup> centuries. As an academic subject, air pollution has mostly been systematically studied only since the mid-late 20<sup>th</sup> Century. There have been several recent reviews, (e.g., (Brasseur et al., 2003; Monks et al., 2009; Ravishankara et al., 2015; Ravishankara, 2003)), which have mapped the growth of atmospheric chemistry, but it is not only peer-reviewed papers that provide relevant overviews. It is important to note that when dealing with the development of this subject (or any scientific subject for that matter), much of the baseline knowledge is embodied in textbooks, which for many are the entry point to and the primary reference for the topic (e.g. (Jacob, 1999; Wayne, 2000; Finlayson-Pitts and Pitts, 2000; Seinfeld and Pandis, 2006; Brasseur et al., 1999)).

Figure 1 shows the number of peer-reviewed papers by year that mentioned the phrase “atmospheric chemistry” in the text, as catalogued by the Scopus bibliographic database (<https://www.scopus.com/>). It shows a growth in the later 1970s from around a hundred papers a year to approximately 4,000 a year currently, with a large increase especially over the past two decades. Of course, many more papers discuss atmospheric chemistry, or are relevant to it, without explicitly mentioning these words!



53

54 This paper aims to recognize and highlight some of the most influential peer-reviewed articles that  
 55 have shaped this field. There were many pivotal scientific discoveries and there were many papers  
 56 that spurred action and further research. What were the significant discoveries that shaped the  
 57 atmospheric chemistry of today? And how do we narrow down the list contributions to highlight the  
 58 most impactful ones?

59 There are many ways to choose the papers that described discoveries and influenced atmospheric  
 60 composition and chemistry. Here we have assembled a compilation of influential articles. Our goal is  
 61 not to show what makes a ‘great’ paper, which depends not only on the science, but also the quality  
 62 of the writing, readability, structure of the written work, and the reader – all criteria that are highly  
 63 subjective. Nor are we aiming only for those papers that led to policy and management actions.  
 64 Instead, we try to reflect on *the papers’ science and content* and the influence of the ideas in those  
 65 papers on our community of scientists and on the field’s development. Our approach is to present our  
 66 thoughts – informed by the solicitation for input from colleagues in the field – and share what we  
 67 think are the fundamental discoveries and developments, start a discussion, and allow others to build  
 68 on, reinforce or critique our work.

69 In addition to peer-reviewed papers, and the above mentioned textbooks, we have of course other  
 70 mediums through which we communicate (have communicated) with our peers. These include  
 71 scientific reports, conferences, and meetings. In addition, we have scientific assessments and  
 72 evaluations, which often get more scrutiny and review than the peer-reviewed papers they include.  
 73 These days, other communication media, such as social media, have also become prevalent as formats  
 74 for exchange both within the scientific community and with a broader audience. However, the entire  
 75 community cannot attend all conferences and meetings, the scientific reports are not always  
 76 accessible and often not peer-reviewed, and the assessments are often driven more by policy needs  
 77 rather than by scientific discoveries. Publishing peer-reviewed papers is the closest we come to  
 78 reaching the whole community. We do note that, despite its known issues, the peer-reviewed

literature is still considered the gold standard for quality and reliability. For these reasons, we discuss only peer-reviewed papers here, although we aim to communicate the overarching scientific advances that shaped the field.

### *1.1. How were the papers selected?*

Easily measurable criteria, like the number of citations, are one metric. However, this approach favours papers of a particular vintage and not necessarily the earlier or later papers. Besides, there are several drawbacks to these simple and objective criteria. Citations tend to go down when something is assumed to be common knowledge and makes it into textbooks or compilations. For example, nobody cites Priestley for discovering oxygen or Schönbein for discovering ozone, whenever atmospheric composition is mentioned. Indeed, some of the central concepts of atmospheric chemistry and physics are considered to be common knowledge, and their origins are taken for granted. The number of citations will also be influenced by the journal in which a paper is published, and quite often (we hate to say this) also depends on who else cited them and in which journal they were cited. Citations also depend on how many people are otherwise researching a particular subject. Furthermore, critical assessments and expert data evaluations suppress the citation of the original papers. This is particularly the case, for example, for papers on chemical kinetics and photochemistry, whereby people tend to simply cite the data evaluations such as National Aeronautics and Space Administration Jet Propulsion Laboratory (NASA/JPL, <https://jpldataeval.jpl.nasa.gov/>) or International Union of Pure and Applied Chemistry (IUPAC, <http://iupac.pole-ether.fr/>) panel reports. Similarly, people often cite the quadrennial ozone layer depletion and Intergovernmental Panel on Climate Change (IPCC) assessments, thereby obfuscating the underlying original papers. Other types of papers, such as reviews, tend to get an excessive number of citations (for understandable reasons). Lastly, we cannot overlook the influence of journal availability in different parts of the world. This availability is exacerbated when journal costs go up, and not everybody can access new papers. Nevertheless, there is still a relevance to the number of citations of a paper. We show, for example, the 10 most-cited papers when we were to search on the combination of words “atmospheric and chemistry” in Table 1.

For all the above reasons, we decided to use a different approach here. We solicited the scientific community to obtain input from the experts in the field. To accomplish this, we put out a call through the International Global Atmospheric Chemistry (IGAC) (Melamed et al., 2015) project to its contacts and thereby engaged a broad audience. Despite the broad audience of IGAC, the vast majority of responses came from scientists in North America and Europe. An initial list of influential papers was established by combining the replies received from the expert solicitation to evaluate the most nominated papers. In addition, a variety of perspectives were assembled for the writing team, including different career stages, nationalities, and genders. Despite all these efforts, the selection methods will still inevitably create bias that cannot be escaped. Therefore, in many respects, the chosen papers are not supposed to be a definitive list, but rather a compilation that allows researchers to discuss and reflect on what makes impactful science, and maybe ponder on the landmarks in our subject. Furthermore, we hope that the end product can provide an interesting history and context to those who are joining the community and document the current “perception” of what are the most important papers.

We have noted the drawbacks in our methodology simply to present upfront some of the limitations of what we did in this paper. However, we hope that others will find this work relevant and engaging. Through an open and active peer-review process, we were able to obtain the perspective of a broader

community more reflective of the global composition of the field. To facilitate this, the paper was published first as an open-access, discussion paper that included a public comment period. We hope that this approach overcame some of the limitations and reservations we expressed earlier. We thank all reviewers for their contributions and help in determining the final shape of this overview.

## *1.2. Scope of work*

As with the selection method, one can debate the scope and the methodology for a work such as this. Still, the boundaries we have drawn encompass studies that have shaped our understanding of the atmosphere and the underlying chemical and physical processes, focusing mostly on the troposphere. This includes modelling, field measurements, remote sensing, and laboratory studies (Abbatt et al., 2014). We have also included atmospheric interactions with the biosphere, cryosphere, and hydrosphere.

We selected 2010 as the cut-off year. Our rationale is that for a paper to have been influential in the whole field it must be at least ten years old and thus had time to accrue recognition. We recognize that important papers in newer areas of endeavour are disadvantaged by this criterion. Examples include the subjects of SOA formation (Ehn et al., 2014; Crounse et al., 2013), the chemistry of Criegee intermediates (Welz et al., 2012; Mauldin et al., 2012), galactic rays induced aerosol particle nucleation (Kirkby et al., 2011), and air pollution-climate connections (Shindell et al., 2012). Influential assessments such as bounding black-carbon (Bond et al., 2013) are also missed. However, these areas will undoubtedly be recognized in the coming decades. The ten-year window also allows the scientific community to have had extensive input on a paper's validity, i.e., meeting the criterion of "standing the test of time."

The papers have been grouped into the following general categories and are presented as such in Section 2.

1. *Foundations*
2. *Aerosols and Clouds*
3. *Secondary Organic Aerosols*
4. *Chemical Kinetics, Laboratory Data and Chemical Mechanisms*
5. *Heterogeneous and Multiphase Chemistry*
6. *Chemical Models*
7. *Tropospheric Ozone*
8. *Nitrogen Chemistry*
9. *HOx Chemistry*
10. *Nighttime Chemistry*
11. *Halogen Chemistry*
12. *Volatile Organic Compounds*
13. *Biogenic Emissions and Chemistry*
14. *Biomass Burning*
15. *Emissions and Deposition*
16. *Chemical Transport*
17. *Satellites and the Troposphere*
18. *Stratospheric Chemistry*
19. *Other issues that influenced tropospheric chemistry*

168

169 The groups were chosen to reflect the main areas of research or endeavour, recognizing that this  
170 division could be done in several different ways. There is no assumed equivalence in these groups  
171 regarding their perceived or real importance or impacts. In the following, we discuss the papers in  
172 each group to show why they have been nominated and to put them in the historical context of the  
173 development of atmospheric chemistry as a discipline.

174

175

176

## 2. Survey of Areas

### 2.1. Foundations

Atmospheric chemistry has some long-standing and deep roots. However, it blossomed in the second half of the 20<sup>th</sup> century following concerns about ozone layer depletion and various forms of tropospheric pollution, such as the Los Angeles smog, London smog, and acid precipitation (Table 2). Many note John Dalton's early contributions on the proportion of gases in the atmosphere (Dalton, 1805) and John Tyndall's Bakerian lecture on radiation and gases (Tyndall, 1861) as among the first studies in this field. The work of Arrhenius "On the Influence of Carbonic Acid in the Air upon the Temperature of the Ground" (Arrhenius, 1896) and the subsequent paper of Callendar, "The artificial production of carbon dioxide and its influence on temperature" (Callendar, 1938), laid the groundwork for the linkage between atmospheric chemistry and climate. Concerning aerosols, the seminal work of John Aitken (Aitken, 1888) "On the number of dust particles in the atmosphere" details early work to count the number per cubic centimeter in various indoor and outdoor environments. It is interesting to note that physiologists looking at the number of live germs in the air stimulated Aitken's work. The later work of Köhler (Köhler, 1936) which explored cloud droplet nucleation remains the basis for later work (see the *Aerosols and Clouds* section). The start of atmospheric chemistry as a distinct discipline probably arrived with Chapman's chemical theory of the stratospheric ozone layer in 1930 (Chapman, 1930), which will be further discussed in the *Stratospheric Chemistry* section. This study heralded the importance of atmospheric chemistry on a global scale.

In analyzing the influential papers on atmospheric composition, one cannot help but note the relationship between these papers and the most significant contemporary environmental issues (Table 2). The first of these was the Los Angeles smog, which had its European counterpart, the London "Pea-Soup" (Brimblecombe, 1987). The two events, which in chemical terms have no equivalence, had comparable impacts on public health and opinion. The oft recognized work of Haagen-Smit (Haagen-Smit and Fox, 1954; Haagen-Smit, 1952; Haagen-Smit et al., 1953) in the early 1950s on the Los Angeles smog was the first to coin the term "air pollution" in the modern era. Haagen-Smit showed that automobile exhaust gases can form ozone in the air and should, therefore, be considered a definite source of smog. Figure 2, redrawn from Haagen-Smit (1952), shows a schematic presentation of the reactions in polluted air leading to smog. Notably, the basic features of tropospheric chemical processes, as we understand them today, were already recognized in these early papers, and they showed how ozone could be chemically produced in the troposphere. Brasseur has documented these findings in a very thorough review (Brasseur et al., 2003).

It is widely recognized that both Crutzen (Crutzen, 1973a; Crutzen, 1973b) and Chameides and Walker (1973) found that similar "smog reactions" oxidize methane (CH<sub>4</sub>) and carbon monoxide (CO) to produce substantial amounts of ozone in remote regions of the atmosphere. They estimated chemically produced ozone to be much greater than that transported from the stratosphere, which was believed to be the primary source of this chemical in the troposphere at that point. A few years earlier, in 1970, Hiram Levy II suggested that the hydroxyl radical, which provides the dominant oxidation mechanism in the troposphere, was formed in unpolluted air by the same mechanism that had been described as occurring in polluted air (Levy, 1971). This paper by Levy (1971) is recognized by many as the first description of the chemistry of the lower atmosphere involving hydroxyl radical

reactions of methane and carbon monoxide, hydroperoxyl radicals, and the photolysis of ozone and formaldehyde as radical sources. In particular, he recognized that the very short-lived electronically excited oxygen atom ( $O^1D$ ) is a possible source of the hydroxyl radical (OH), an idea now well established.

Around the same time, Weinstock (1969) explained how cosmic rays lead to the production of radiocarbon dioxide ( $^{14}CO_2$ ), which is incorporated into living plants. This process requires a rapid turnover of radiocarbon monoxide ( $^{14}CO$ ), which was unexpected because the lower atmosphere was thought to be a “chemical desert”. Instead, carbon monoxide appeared to have a turnover time of about one-tenth of a year, primarily driven by hydroxyl radical oxidation. To some, this paper kicked off the research which led to our present understanding of the atmospheric chemistry of the lower atmosphere.




Figure 2 – Schematic representation of the reactions in polluted air leading to smog symptoms (adapted from Haagen-Smit et al. (1953)).

It has been claimed that “acid rain was one of the most important environmental issues during the last decades of the twentieth century” (Grennfelt et al., 2019) (see Table 2). One of the reasons is that acid rain first demonstrated that air quality was not merely a local issue but a regional issue and showed that the atmosphere has no definite boundaries (Fowler et al., 2020). Although the case of acid rain and its effects had been noted and reported by some earlier papers e.g., Odén (1968), for some, it is the paper by Likens and Bormann (1974) that made this issue known to the science community at large. Other early papers (for example, from Urone and Schroeder (1969) and Penkett et al. (1979)) also recognized the vital role of liquid-phase oxidation of sulphur dioxide ( $SO_2$ ) by oxidants such as hydrogen peroxide ( $H_2O_2$ ) and ozone ( $O_3$ ). Current estimates suggest that roughly 50% of the  $SO_2$  oxidation in the lower troposphere occurs through liquid-phase reactions.

The story of lead in the atmosphere is a complex interplay between science, policy and economics (Monks and Williams, 2020), where observations in snow (Murozumi et al., 1969) underpinned the alarming growth and spread of lead pollution and latterly its demise (Boutron et al., 1991).

There is little doubt that one of the most impactful series of papers is that of the eponymous curve by Keeling (Keeling, 1960; Keeling et al., 1979; Pales and Keeling, 1965), showing the steady rise in carbon dioxide ( $CO_2$ ) measured at Mauna Loa observatory (this work has continued uninterrupted by



NOAA/ESRL/CML over the past few decades). Keeling's work was built on the previously mentioned work of Callendar (1938) who compared measurements of CO<sub>2</sub> at Kew, UK (1891-1901) with those in the Eastern USA (1936-1938) and noted an increase in concentration. Although the gas in question is CO<sub>2</sub>, which is often seen only as climate gas, changes in its levels reflect the changing composition of the atmosphere and the effects that it can have, and shows that the two subjects cannot be easily separated. Furthermore, the increase in CO<sub>2</sub> is of central to ocean acidification, a topic not touched upon here but nevertheless very important. The seminal paper by Ramanathan et al. (1985) that highlighted the role of CH<sub>4</sub>, chlorofluorocarbons (CFCs), and nitrous oxide (N<sub>2</sub>O) for climate strengthened the case for the inclusion of chemistry in the climate issue. In many respects, this close-coupling between atmospheric chemistry and climate change was brought to the forefront with the 1995 Nobel Prize being awarded to Paul Crutzen, Mario Molina and F. Sherwood Rowland "for their work in atmospheric chemistry, particularly concerning the formation and decomposition on ozone"<sup>1</sup> (see *Stratospheric Chemistry* section) and, later, with the Nobel Prize to the IPCC.

The role of field campaigns, observations and the attendant models in shaping our understanding of atmospheric chemistry should be recognised as foundational. In general, the adage that the atmosphere is under-observed is still true. Every time a new instrument has been developed to detect a new chemical in the atmosphere, there have been significant advances (Heard, 2006). One could posit that the entire field of atmospheric science started because of detection and quantification of oxygen and ozone in the Earth's atmosphere. Some recent major advances in our field has been through field measurement. For example, observation of the ozone depletion (including the ozone hole), aerosol particles, free radicals, and stable molecules (including ozone layer depleting substances, CO and methane) fundamentally changed the course of the field. Organized systematic probing of the atmosphere has been critical over the past four, or so, decades. Here again, introduction of new instruments (optical, mass spectrometric, etc.) have been game-changing. It is also important to note and highlight the enormous contributions of satellite observations to provide global coverage. Often, as field campaigns and their impact are spread across many papers it is difficult to pull out their specific contributions. Many of the early experiments, encompassing long-range transport, biomass burning and aerosols, particularly using aircraft, have been detailed in (Melamed et al., 2015). Assembling a large number of instruments on a large aircraft to simultaneously measure an array of chemicals was pioneered by Davis (Davis, 1980) and has been a paradigm for field studies ever since.

In addition to organized episodic field measurements, continual measurements of chemicals (often called monitoring) has produced some of the most significant findings about the atmosphere. For example, continual monitoring of surface ozone from Paris or similar stations going back over a hundred years or more has shown the trends in tropospheric pollution due to human activities (Volz and Kley, 1988). The continual monitoring of the Antarctic ozone led to the discovery of the ozone hole. The continual monitoring of CO<sub>2</sub> is the poster-child for climate change! Much of this continual monitoring has been carried out by national agencies and international partnerships. Examples include the US National Oceanic and Atmospheric Administration (NOAA) Earth Systems Research Laboratory (ESRL) Global Monitoring Laboratory's (GML) contributions (Montzka et al., 2007) and international efforts, such as The Advanced Global Atmospheric Gases Experiment (AGAGE) network (Prinn et al., 2001; Prinn et al., 1995), World Meteorological Organisation – Global Atmospheric Watch (WMO-

---

<sup>1</sup> <https://www.nobelprize.org/prizes/chemistry/1995/press-release/>

GAW) (WMO, 2017) and the Network for the Detection of Atmospheric Composition Change (NDACC) (De Mazière et al., 2018).

## 2.2. Aerosols and Clouds

Aerosols in the atmosphere greatly influence both air quality and climate change; they are also a significant media for composition change in the atmosphere. In this section we discuss three main areas of research related to aerosols: (1) Understanding the mechanisms and atmospheric chemistry processes that influence aerosol particle formation, nucleation, and growth, and how aerosols affect composition; (2) The role of aerosols as cloud condensation nuclei and the influence that this process has on climate; and (3) The impact of particulate matter on human health. These areas are, however, related and there is not always a clear division. *Secondary Organic Aerosols* (SOA) and *Heterogeneous and Multiphase Chemistry* are discussed in the corresponding sections (2.3 and 2.5).

The roots of modern aerosol science lie, as previously discussed (see the 2.1. *Foundations* section), in the works of Aitken (1888) and Köhler (1936) on the cloud droplet. Twenty years after Köhler's research, Junge (1955) provided the power-law describing aerosol particle number and identified the stratospheric aerosol layer, now dubbed the 'Junge layer'. Junge concluded: "A real step forward in the understanding of the basic processes in air chemistry can be gained only if aerosol particles and gases are measured simultaneously but separately, and if the aerosol particles, in turn, are separated according to size." This suggestion has been a clarion call for atmospheric scientists ever since.

Junge and Ryan (1958) attempted to elucidate the formation of particles from gas-phase reactants, particularly SO<sub>2</sub> and ammonia (NH<sub>3</sub>), while Fitzgerald (1974) investigated the variation in aerosol particle composition with particle size. They showed that cloud droplet size distribution was insensitive to the specific soluble constituents. Twomey (Twomey, 1977, 1974) suggested that air pollution gives rise to the whitening of clouds and influences the planet's radiative balance. He also indicated that there is a connection between pollution aerosols and cloud reflectance (albedo). This concept is now often referred to as the "Twomey effect." Twomey (1977) expanded on the 1974 work, exploring the balance between the scattering *versus* absorption effect on incoming solar radiation. It is on this basis that much of the current research on the role of aerosols *via* their direct and indirect effects on climate has been built. Bolin and Charlson (1976) estimated that anthropogenic sulphate aerosol from the US and Europe would lead to a global temperature decrease of 0.03-0.06 °C. They recognized early on that "we are already approaching the time when the magnitude of the indirect effects of increasing use of fossil fuel may be comparable to the natural changes of the climate over decades and centuries."

In the early 1970s, Whitby and Knutson developed an instrument to measure particle size distribution in the nanometer to micrometer range (Knutson and Whitby, 1975) – the well-known aerosol particle mobility analyser. They used the measurements from this instrument to introduce a new formulation of the formation and growth of atmospheric aerosol particle size modes – the "Whitby diagram" which is now a common text book figure and is shown in Figure 3. The outcomes of this work show the importance and influence of the development of new instruments that probe the atmosphere.




Figure 3 - Schematic of atmospheric aerosol particle size distribution showing the three modes, the main source of mass for each mode, and the principal processes involved in inserting mass to each mode along with the principal removal mechanisms (Whitby, 1978). (taken from <https://serc.carleton.edu/NAGTWorkshops/metacognition/largeclasses.html>, under CCC)

330

331 The CLAW hypothesis (the acronym taken from the surnames of the proposers Charlson, Lovelock,  
 332 Andreae, and Warren) (Charlson et al., 1987) further connected aerosol science to gas-phase  
 333 chemistry, specifically focused on the feedback loop between ocean ecosystems and Earth's climate.  
 334 This hypothesis built on earlier work by Lovelock et al. (1972) on the oxidation of marine  
 335 dimethylsulphide. Although the conclusions of Charlson et al. (1987) have been questioned (Quinn  
 336 and Bates, 2011), this paper highlighted the interconnections within atmospheric sciences, and  
 337 environmental sciences in general.

338 The work of Friedlander and co-workers (Stelson et al., 1979) further highlighted the role of liquid-  
 339 phase chemistry leading to aerosol particles. A key milestone in our understanding of sulfate  
 340 formation was the recognition that the reaction of the hydroxysulfonyl radical ( $\text{HOSO}_2$ ) with oxygen  
 341 ( $\text{O}_2$ ) is exothermic (Calvert et al., 1978) and leads to gas-phase sulphur trioxide ( $\text{SO}_3$ ), contrary to what  
 342 was accepted at the time. Prior to this finding, there were major difficulties in understanding the  
 343 formation of gas-phase sulphuric acid ( $\text{H}_2\text{SO}_4$ ) formation (Davis et al., 1979) from gas-phase  $\text{SO}_2$   
 344 oxidation, an essential step for the nucleation of new particles from the gas-phase in the atmosphere.

345 This area of research was further developed by Robbin and Damschen (1981) who investigated the  
 346 role of peroxide in the liquid-phase in oxidising  $\text{SO}_2$  which was key to understanding the phenomenon  
 347 of acid rain. Graedel and Weschler (1981) reviewed the chemical transformations in atmospheric  
 348 aerosol particles and raindrops and extended the idea of Martin and Damschen. Stelson and Seinfeld

(1982) evaluated the thermodynamics of ammonium, nitrate, and sulphate aerosols, which was a significant step in understanding particle formation and growth. Pankow's 1994 work (Pankow, 1994b, a) on the absorption model of gas/particle partitioning of organic compounds in the atmosphere is of fundamental importance for models to calculate the amounts of particulate matter (PM) formed and their growth in the urban and regional air, and in the global atmosphere.

Charlson et al. (Charlson et al., 1990; Charlson et al., 1991) produced the first global estimate of the direct aerosol effect that subsequently had a large impact on climate modelling. The role that aerosols on cloud condensation nuclei (CCN) and cloud albedo was also acknowledged, concluding that it may be substantial. How substantial, however, was not quantified at that point because of a lack of knowledge on the relationships involved. A few years later, Boucher and Lohmann (1995) provided an estimate of the indirect effect of anthropogenic aerosols on climate. After many additional years of study based on these foundations and analyses of radiative balance, the total radiative forcing by anthropogenic aerosol is now estimated to be roughly  $-1.1 \text{ W/m}^2$  (IPCC, 2013), thereby solidifying the importance of aerosols in climate change.

Building on the work of Whitby, Mäkelä et al. (1997) conducted continuous monitoring of particles at a forest site in Finland. Beyond confirming the existence of three submicron particle size modes (the nucleation, Aitken, and accumulation modes (see also Covert et al. (1996)), they also observed new particle formation events. These events have been subsequently observed by others and are often depicted in the literature using the famous "banana plots" (Figure 4).




Figure 4 - Contour plots of particle formation event occurring in the morning, followed by a subsequent growing process of the nucleation mode during the afternoon. a) February 21st 1996, b) March 13th 1996 and c) April 13th 1996 (Mäkelä et al., 1997). Thought to be the origin of the “banana plot.”

There has been ample recognition for the research on process representations, such as the mole-fraction-based thermodynamic models (Clegg et al., 1998a, b) and one-parameter model for hygroscopic growth (Petters and Kreidenweis, 2007) and CCN; Facchini et al. (1999) presented experimental work aimed at understanding the role of surface tension in droplet growth and the subsequent effect on cloud albedo and radiative forcing (RF), while Knipping et al. (2000) used a simplified experiment to investigate the role of reactions of gases with ions at the air-water interface.

More specifically, the role of organics in the formation and growth of aerosol particles has been a significant area of research (Kulmala et al., 2000). In addition to the natural hydrocarbons noted above, it has become clear that anthropogenic hydrocarbons such as aromatic compounds are also involved in new particle formation and their growth (Odum et al. (1997).

Jaenicke (2005) was the first to suggest, to the best of our knowledge, that biological particles are an important fraction of atmospheric aerosol particles. This paper prompted the development of a new and exciting field within atmospheric sciences. Fröhlich-Nowoisky et al. (2016) reviewed the role of bioaerosols in health, climate and ecosystems.

### 2.3. Secondary Organic Aerosols

Since the mid-2000s, secondary organic aerosols (SOA) have been the focus of much research, addressing their abundance, sources, and production pathways. One of the foundational works in this area is the recognition of the role of natural and anthropogenic hydrocarbons, and in particular isoprene chemistry in the formation of SOA (Claeys et al., 2004).

The chemical composition of SOA across the globe is still poorly understood (Zhang et al., 2007), although ways to describe the growth of SOA have advanced significantly (Kalberer et al., 2004). Donahue et al. (2006) developed an approach based on the volatility of organics, a concept termed “volatility basis set.” This concept has been extended to a host of volatilities and their classifications. For example, as shown in Figure 5, Robinson et al. (2007) postulated that a large amount of SOA mass is unexplained by current models, and methods used to estimate SOA production do not capture what is measured in the field.




Figure 5 - Aerosol particle formation from the photochemical oxidation of diesel exhaust in an environmental chamber. The grey area indicates the primary aerosol particle (POA + other species). The red area shows the upper-bound estimate of the contribution of known SOA precursors to the suspended aerosol particle mass leaving behind a large fraction that is not accounted for (blue area) (Robinson et al., 2007).

The introduction of the aerosol mass spectrometer (AMS) by Worsnop and colleagues (Canagaratna et al., 2007) along with the pioneering instruments of Prather (Gard et al., 1997) and Murphy et al. (2006) that built on the early work of Sinha (1984), have helped determine aerosol composition. Studies using these instruments have established that organic compounds are ubiquitous in aerosol particles. Zhang et al. (2007) and later Jimenez et al. (2009) explored the chemical composition of PM at different sites across a part of the globe (Figure 6), and their work has now been extended by a large number of groups.

Aimed at addressing some of the ‘missing urban SOA’ in models, Surratt et al. (2010) investigated SOA production from isoprene and Virtanen et al. (2010) showed the amorphous solid state of biogenic secondary organic aerosol particles, challenging the traditional views of the kinetics and thermodynamics of SOA formation and transformation, that assumed low viscosity, liquid-like, particles exchanged chemicals rapidly with the gas-phase.






Figure 6 - Total mass concentration (in micrograms per cubic meter) and mass fractions of nonrefractory inorganic species and organic components in submicrometer aerosol particles measured with the AMS at multiple surface locations in the Northern Hemisphere at mid-latitudes. The organic components were obtained with FA-AMS methods (Zhang et al., 2007). In some studies, the FA-AMS methods identified one OOA factor, whereas in other locations, two types, SV-OOA and LV-OOA, were identified. HOA is a surrogate for urban primary OA, and Other OA includes primary OAs other than HOA that have been identified in several studies, including BBOA. Inset: distributions of O:C for the OA components identified at the different sites (Jimenez et al., 2009).

#### 2.4. Chemical Kinetics, Laboratory Data and Chemical Mechanisms

Chemical kinetics is one of the foundations of atmospheric chemistry (Abbatt et al. (2014)). This includes a number of different areas: investigation of individual chemical reactions; theoretical understanding of elementary reactions; evaluations and compilation of kinetics data; development and compilation of chemical mechanisms for use in models (see the 2.6. *Chemical Models* section); testing and simplification of the models for use in numerical models. Demerjian et al. (1974) is considered by many in the community to be one of the cornerstones of chemical mechanism development and it has been influential in a number of other research areas as well. This paper provided an explicit chemical mechanism for the troposphere in which all the chemical reactions were written as numerically integrated stoichiometric equations to predict photochemical ozone production rates. Previously, all chemical mechanisms had been highly “reduced” (into simple mechanisms) and/or parameterised, with non-stoichiometric equations. Using Demerjian’s approach, many explicit atmospheric chemical mechanisms have been derived, including one of the most widely used, the Master Chemical Mechanism (MCM, Figure 7) (Jenkin et al., 1997; Jenkin et al.,

2003;Saunders et al., 2003). Currently, there are a variety of tropospheric chemistry mechanisms that capture the scope of chemical reactions that are used in a number of models including the 1990 Carter mechanism (Carter, 1990), the regional acid deposition model / regional atmospheric chemistry mechanism (RADM/RACM) (Stockwell et al., 1997;Stockwell et al., 1990), SAPRC-07 (Carter, 2010) and the Chemical Aqueous Phase Radical Mechanism (CAPRAM) (Ervens et al., 2003). One of the key foundational techniques for estimating rate constants is that of structure-activity relationships (Kwok and Atkinson, 1995). In the near future, calculations of rate coefficients based on ab-initio quantum calculations will likely be common.

There is no doubt that the chemical kinetic data compilations have been the backbone of providing much needed experimental data to all chemical mechanisms and models (see the 2.6. *Chemical Models* section). The comprehensive reviews of Atkinson starting in the mid 80s (Atkinson (1986)) and followed by many others, provided a consistent description of the reaction pathways of the alkyl, peroxy and alkoxy radicals produced by the reactions of hydroxyl radicals with a wide range of organic compounds. These papers led the way for the compilation of the IUPAC and NASA/JPL chemical kinetic data evaluation of tropospheric reactions (Atkinson et al., 1989;Crowley et al., 2010;Atkinson et al., 1992;Atkinson et al., 2004;Atkinson et al., 2006;DeMore et al., 1997;Burkholder et al., 2020). (Note that compilation of kinetics data for stratospheric reactions dates back to mid-1970s (Hudson and Reed, 1979)). These works have been the foundation for the development of all chemical mechanisms and have led to the standardisation and improvement of condensed chemical mechanisms used in all chemical models.




Figure 7. Oxidation mechanism of butane in the Master Chemical Mechanism (Jenkin et al., 1997).



The recognition of reactions with negative activation energies and the role of weakly bound adducts were other key steps in improving our understanding of chemical kinetics. When the adduct is strong, we term it an association reaction, which exhibits negative activation energies and pressure dependence. Understanding and representing these type of reactions in atmospheric chemistry has been a major step. In particular, the pioneering work of Troe and coworkers has enabled realistic and simpler representation of these reactions based on the Rice-Ramsperger-Kassel-Marcus (RRKM) theory (Troe, 1979, 1994).

Atmospheric chemistry is often termed atmospheric photochemistry since the initiator for many of the reactions is the production of free radicals, which are directly or indirectly the result of solar radiation. Over the decades, the representation of photochemical processes has been well established. A key element is the calculation of the “j-value” (i.e. the photolysis rate) for a photochemical process, which depends on radiative transfer to obtain the solar flux and laboratory measurements of absorption cross sections and quantum yields. The pioneering works on methods for quickly and accurately calculating j-values are those of Madronich and Flocke (1999) and of Prather and colleagues (Wild et al., 2000).

Moving towards individual reactions, the work of Howard and Evenson (1977) on the reaction between hydroperoxyl radical and nitric oxide,  $\text{HO}_2 + \text{NO}$ , has been recognised as a scrupulously careful study that overturned conventional wisdom on this key reaction in photochemical smog/ozone formation (and in stratospheric chemistry). The work of Vaghjiani and Ravishankara (1991) demonstrated the importance of operating at low  $[\text{OH}]$  to reduce secondary reactions and extended measurements down to low and atmospherically relevant temperatures.

Three papers nearly a decade apart address the fundamental importance of robust laboratory measurements to underpin model-led interpretation of experimental data. The seminal work demonstrating the long-wavelength tail on the ozone photodissociation quantum yield (Ball et al., 1993; Ravishankara et al., 1998) and the related work on the  $\text{O}^1\text{D} + \text{N}_2/\text{O}_2$  reactions (Ravishankara et al., 2002), identified key processes in the formation of OH radicals in the troposphere. Prompted by the findings of Lelieveld et al. (2008) (see the 2.13. *Biogenic Emissions and Chemistry* section) and of Hofzumahaus et al. (2009) (see the 2.9. *HOx Chemistry* section), a pair of laboratory papers published in 2009, about  $\text{HO}_x$  radical regeneration in the oxidation of isoprene (Peeters et al., 2009) and unexpected epoxide formation in the gas-phase photooxidation of isoprene (Paulot et al., 2009) have changed the way we understand the gas and aerosol products and impacts of isoprene chemistry (Kleindienst, 2009). It is worth noting that Peeters et al. (2009), using theoretical electronic-structure calculations, showed the major role of autoxidation chemistry (peroxy – hydroperoxy isomerization). This work changed the traditional view of peroxy radical chemistry and introduced the ideas of isomerization and more complex pathways to atmospheric chemistry.

## 2.5. Heterogeneous and Multiphase Chemistry

Earth’s atmosphere contains various amounts of condensed matter suspended in air. The most visible condensed matter is, of course, clouds. One can also see aerosols when the particle numbers and sizes are large; examples include smog, wildfires, and volcanic eruptions. In addition to clouds, snow and ice provide different mediums that can alter gas-phase chemistry.

Many chemical reactions occur on the surfaces of particles suspended in air, ice/snow on the ground, and within liquid drops. In general, these processes catalyse reactions that would be very slow in the gas-phase, such as those between closed-shell molecules, and/or can produce products that do not form in the gas-phase. For these reasons, heterogeneous and multiphase reactions are of immense interest, although the distinction between heterogeneous and multiphase chemistry is not always clear cut (Ravishankara, 1997). Often "heterogeneous" is taken to mean reactions at surfaces and "multiphase" to mean reactions involving the uptake of gases into (and reaction in) the liquid phase.

The unique contribution of reactions in/on condensed matter burst into the limelight due to their role in the formation of the ozone (Solomon et al., 1986). However, such reactions had been recognized to be important before the ozone hole research, for example in the oxidation of SO<sub>2</sub> (Urone and Schroeder (1969) and Penkett et al. (1979)). Since the 1990s, the roles of heterogeneous and multiphase reactions have been highlighted in many tropospheric processes, as noted here in various sections (see for instance sections 2.1. *Foundations* and 2.4. *Chemical Kinetics, Laboratory Data and Chemical Mechanisms*).

Chameides and Davis (1982) studied the free radical chemistry of cloud droplets and its impact upon the composition of rain showing that the radical chemistry in water droplets could drive production of peroxides, which have the ability to rapidly oxidise sulphur species – a strong link to acid rain. Work of Akimoto et al. (1987) on the photoenhancement of nitrous acid formation in the surface reaction of nitrogen dioxide and water vapour demonstrated the existence of an additional radical source in smog chamber experiments. This built on the earlier work of Pitts et al. (1984) and challenged our understanding of the role of such heterogeneous reactions in the atmosphere. The work of Mozurkewich et al. (1987), Hanson et al. (1992) and subsequently Thornton and Abbatt (2005), on the measurements of HO<sub>2</sub> uptake to aqueous aerosol particles was highly influential in the debate on the aerosol loss of HO<sub>x</sub>, a question that had vexed many modelling studies.

A pioneering paper in tropospheric cloud chemistry is the study by Jacob et al. (1986) in the San Joaquin valley that used a multiphase measurements and modelling approach to study the formation of acid fog. Two further papers have brought heterogeneous chemistry to the fore: Dentener et al. (1996) in their original paper on the role of mineral aerosol as a reactive surface in the global troposphere showed the potential role of mineral dust on sulphur oxides (SO<sub>x</sub>), NO<sub>y</sub> (NO<sub>y</sub> includes nitrogen oxide (NO) and nitrogen dioxide (NO<sub>2</sub>), as well as the compounds produced from the oxidation of those) and O<sub>3</sub> chemistry, and Jacob (2000) who reviewed in more detail the chemistry and ozone *via* HO<sub>x</sub> and NO<sub>y</sub> at the interface of gas phase and suspended particles (including clouds) and led to a highly-cited series of recommendations for future studies.

Another similar area is that of chemical reactions on/in snow and ice. Such reactions were highlighted by Barrie et al. (1988), specifically with regard to the production of halogens on the ice surface, and hinting at the role of the cryosphere as a source of chemical species to the troposphere (see the 2.11 *Halogen Chemistry* section). Given the extent of the cryosphere and in particular of snow (Grannas et al., 2007), findings in the late 1990s demonstrated its role in promoting heterogeneous and multiphase reactions as a significant source of unusual and unexpected chemical species to the atmosphere. One of the most nominated works in this area was that by Honrath et al. (1999) investigating NO<sub>x</sub> production from the illuminated snowpack. Pioneering work of Davis et al. (2001) on the unexpected production of NO<sub>x</sub> in pristine Antarctica is also worthy of note.

## 2.6. Chemical Models

Up front, we want to acknowledge that we are not doing full justice to the important role played by chemical models in the understanding of and developing tropospheric chemistry, informing policymaking, and deciphering field observations. We do, however, note some of the key developments in modelling, which is the way we couple atmospheric motion with chemical processes.

Chemical models are the conduit to represent our knowledge of the chemical and physical processes in the atmosphere within a mathematical (numerical) framework that allows prediction and testing against observations (in the laboratory and the atmosphere). Therefore, models are the tools upon which atmospheric environmental policies are developed. Indeed, the efforts in modelling are vast and they are pivotal tool of tropospheric chemistry. Policies pertaining to climate, air quality, acid precipitation, etc. are based on such model predictions and projections. Further, models have been the tool that have enabled quantification of emissions (the quantity of most interest to policymakers), identification of the sources, and evaluation of impacts. One could argue that our knowledge would be incomplete without models.

Early, simple chemical models (with no chemical transport) were useful tools to elucidate and test the basic theory of photochemical ozone formation (Levy, 1971). The recognition that one cannot treat the chemical transformation without considering atmospheric transport and mixing came early. The original simple 1-D models, often designed with a parameterized vertical transport in terms of an “Eddy Diffusion” concept, were superseded by 2-dimensional models and have now been largely replaced by complex 3-D models. 2-D models of the stratosphere (which are zonal averages with latitude and height being the variable dimensions) have been extremely useful and are still used in assessment activities. (See for example, Garcia and Solomon (1983) and Fleming et al. (1999)). The NASA conference publication 3042 (Jackman et al., 1989) provides an excellent review of sixteen 2-D and a few 3-D stratospheric models that were used in the 1980s and 1990s. Also, chemical transport models (CTMs), which use analysed winds, are often used to separate transport from chemistry. Such models are extremely powerful in accounting for observations, atmospheric budget calculations, and deciphering the roles of various chemical processes taking place in the atmosphere. However, projections and predictions in a changing climate requires coupling of chemistry to meteorological prediction models. Now, free-running, on-line 3-D models, which include chemistry, have been implemented, and the continued enhancements in computing capabilities have greatly improved our modelling capabilities. Logan and co-authors are recognised by many contributors as providing the basic model description of global tropospheric chemistry (Logan et al., 1981). Bey et al. (2001) first described GEOS-Chem, a global, three-dimensional, tropospheric, chemical transport model. Though not the only global tropospheric model, as an open-source model with a large user community and flexibility, it has become a very influential global model. In recent years the Weather Research and Forecasting (WRF)-Chem model has also been used extensively (Grell et al., 2005). There are now numerous 3-D models developed by various organizations. We refer readers to two excellent articles that describe the role of models and their development in detail (Brasseur (2020) and Zhang (2008)). Inverse modelling especially for source and sink attribution has been shown to be a powerful tool (Hein et al., 1997).

In addition to the global models, regional and air-shed models were critical for air quality predictions and are still employed for regulatory use. A series of three papers from the Seinfeld group (Reynolds et al., 1973; Reynolds et al., 1974; Roth et al., 1974) provided the earliest complete descriptions of an air quality policy model. They linked together emission inventories, meteorological data, chemical mechanisms, and air quality network data to evaluate model performance. All subsequent air quality policy models have followed the same general approach and their basic formulation.

Another major use of models is to interpret large scale field measurements. One of the earliest detailed tropospheric chemical modelling studies that integrated highly instrumented intensive field campaign data was that of Harriss (1988) for the ABLE 2A (Amazon Boundary Layer Experiment) campaign in the Amazon Boundary Layer. Now use of multiple models to interpret field data is a common feature of modern tropospheric chemistry research.

Multi-model ensembles of the troposphere as epitomised by Stevenson et al. (2006) and Fiore et al. (2009) (see the 2.16. *Chemical Transport* section) are a powerful tool for generalising the model “understanding” of the atmosphere. This modelling approach makes use of many different models to achieve a more accurate representation of the observations than would be possible by using only one model, thus producing more reliable outcomes for assessments and policies on a global scale. In addition, multiple runs of the same models with slightly different initial conditions are used to examine the range of outcomes. This approach is akin to the use of multiple models and model runs in weather predictions.

## 2.7. Tropospheric Ozone

Ozone is one of the central molecules of atmospheric chemistry and runs through much of the foundations of the discipline, from its role in the stratosphere as a UV shield, to its role as a major greenhouse gas, to its pivotal part in the troposphere as the start and end-product of oxidation chemistry, and its detrimental influence as an air pollutant harmful to human health and ecosystems. Much of the early thinking on ozone was focused on the question of whether tropospheric ozone was a small subset of stratospheric ozone, see for example Galbally (1968) and Fabian and Pruchniewicz (1977). The latter paper showed the value of observational networks based on standardised instrumentation and calibration techniques, together with consistent siting criteria, and raised the issue of seasonal variations in tropospheric ozone and the nature of the processes that drive them. The vertical structure of a layer of high O<sub>3</sub> concentrations in the stratosphere, where O<sub>2</sub> could be directly photolyzed to make oxygen atoms and hence ozone, and declining concentration in the troposphere was indicative of a stratospheric source and a tropospheric sink and this was the prevalent theory prior to the late 1970s (see also the 2.18. *Stratospheric Chemistry* section). A major breakthrough were the two papers by Chameides and Walker, and Crutzen (Chameides and Walker, 1973; Crutzen, 1973b) that showed that ozone can be photochemically generated in the troposphere, just like it is made in smog *via* the reactions involving hydrocarbons and nitrogen oxides.

The importance of ozone as a radiative gas has been known for a long time, with a significant fraction of heating in the stratosphere coming from ozone photolysis followed by its reformation and thus converting sunlight to heat. In 1979, Fishman et al. (Fishman et al., 1979) identified that tropospheric ozone is also a greenhouse gas. Hence, a change in tropospheric ozone will perturb the radiative

energy budget of the Earth-atmosphere system which will in turn perturb the climate system. Ozone thus became the second trace gas after carbon dioxide to be implicated in global warming and climate change.

Large scale mapping of global tropospheric ozone was first undertaken by Logan (Logan, 1985) who looked at seasonal behaviour and trends with a view to understanding anthropogenic influence. This was later complemented by a paper exploring the photochemical origins of tropospheric, rather than stratospheric, ozone in the rural United States (Logan, 1989). As the understanding of the photochemistry of ozone developed, measurements at Niwot Ridge, Colorado (Liu, 1987) aimed to quantify the elements of the ozone budget by season, bringing forward the concept of ozone production efficiency. Lin et al. (1988) explored the non-linearity of tropospheric ozone production with respect to NMHCs and NO<sub>x</sub>. Though this chemistry had been outlined much earlier – e.g. Demerjian et al. (1974) – this work explored it in the background atmosphere with models and measurements. A powerful demonstration of the low-NO<sub>x</sub> ozone destruction chemistry came from measurements made at Cape Grim, Australia, a background station, where Ayers and Penkett (Ayers et al., 1992) and their team(s) used measurements of ozone and peroxides (Figure 8) to show further experimental proof for the photochemical control of ozone in remote locations.




Figure 8 – Average diurnal cycles of peroxide (orange line) and ozone (blue line) in baseline (low NO<sub>x</sub>) air at Cape Grim (Tasmania, Australia) in January 1992. Adapted from Ayers et al. (1992).

Measurements have always been a critical driver in tropospheric chemistry, and the idea to use in-service commercial aircraft as a platform for programs such as MOZAIC/IAGOS (Thouret et al., 1998) has been recognised for the enormous amount of high quality data, which would otherwise be difficult to regularly obtain from the upper troposphere and lower stratosphere. Using such measurements Newell et al. (1999) combined dynamical and chemical tracers to further delineate ozone origin and

budgets. In the same year, Logan published a synthesis of ozone sonde data (Logan, 1999) which gave an unprecedented look at the seasonal and vertical distribution of ozone and became a reference point for the subject. A year later, Thompson et al. (2000) used a combination of shipboard and satellite views of a tropospheric ozone maximum to suggest the occurrence of a tropical Atlantic ozone “paradox”. The “Atlantic paradox” refers to a greater tropospheric ozone column amount over the South Atlantic than the North Atlantic during the West African biomass burning season. This phenomena was further explored using an expanded network of ozonesondes in the southern hemisphere (SHADOZ) (Thompson et al., 2003). In combination with the earlier work of Logan, these became the basis for the measurement-based description of ozone in the troposphere.

A decades worth of knowledge on the relationship between ozone and its precursors was pulled together by Sillman (Sillman, 1999), cementing the concepts of NO<sub>x</sub>- and VOC-sensitive (or NO<sub>x</sub> saturated) chemical regimes. The paper introduced a generation of researchers to isopleth diagrams (the famous Sillman plot Figure 9) and ozone production efficiencies (OPEs).




Figure 9 - . Ozone isopleths (ppb) as a function of the average emission rate for NO<sub>x</sub> and VOC (in 10<sup>12</sup> molec. cm<sup>-2</sup>s<sup>-1</sup>) in 0-D calculations. The isopleths (solid green lines) represent conditions during the afternoon following 3-day calculations with a constant emission rate, at the hour corresponding to maximum O<sub>3</sub>. The short blue dashed line represents the transition from VOC-sensitive to NO<sub>x</sub>-sensitive conditions. Adapted from Sillman and He (2002).

The power of models to explore global tropospheric ozone distributions, budgets and radiative forcing was fully demonstrated in the study by Stevenson et al. (2006), that brought together 26 atmospheric chemistry models to explore both the air quality and the climate roles of ozone (see also the 2.6. *Chemical Models* section). As discussed in the 2.16. *Chemical Transport* section, a similar approach was used by Fiore et al. (2009) to explore the relationship between inter-continental transport and ozone.

## 2.8. Nitrogen Chemistry

Nitrogen oxides are an integral part of tropospheric processes. Nitrogen oxides are released into the troposphere from a variety of biogenic and anthropogenic sources including fossil-fuel combustion, biomass burning, microbial activity in soils and lightning. The concept of the Leighton photostationary state (Leighton, 1961) between NO, NO<sub>2</sub>, and O<sub>3</sub> was well established by the mid-1990s, and early work from Singh and Hanst (1981) highlighted the potential role of peroxyacetyl nitrate (PAN) to be a reservoir for NO<sub>x</sub> in the unpolluted atmosphere. The measurement of NO/NO<sub>2</sub> by chemiluminescence was critical to the widespread measurement of NO<sub>x</sub> (Kley and McFarland, 1980). A landmark paper in the area of nitrogen chemistry was that of Logan (1983) that brought together global and regional budgets for the nitrogen oxides (Table 3). Later, a paper that focused more narrowly on a specific source of NO<sub>x</sub> was that of Yienger and Levy II (1995) who produced an empirical model of global soil-biogenic NO<sub>x</sub> emissions. Higher up in the atmosphere, the work on sources and chemistry of NO<sub>x</sub> by Jaeglé et al. (1998) is recognised for its contribution to the understanding of the NO<sub>x</sub> cycle in the upper troposphere.

HONO, somewhat a Cinderella molecule, whose photolysis can be a major OH source, especially during the early morning was first identified by Perner and Platt (1979), the heterogeneous nature of which has always driven much interest (Kurtenbach et al., 2001).

These works were complemented by a more holistic view of the nitrogen cycle and in particular the concept of reactive nitrogen (Nr) by Galloway et al. (2004) that clearly showed the linkages between the terrestrial ecosystem and the atmosphere and how the nitrogen budget had and would change leading to the important concept of nitrogen cascade (Sutton et al., 2011). In more recent times, extensive work on vehicle NO<sub>x</sub> sources from exhaust remote sensing data (Bishop and Stedman, 1996), as epitomised in Carslaw (2005) should be highlighted. This paper pointed out the trends that can be said to have led to the denouement of the Volkswagen emissions scandal.

## 2.9. HO<sub>x</sub> Chemistry

There is no doubt that the chemistry of OH and HO<sub>2</sub> (known together as HO<sub>x</sub>) has a central role in the atmosphere as well as holding a certain fascination to atmospheric scientists owing to the significant challenges involved in measurement and understanding its impact locally to globally. Much of the history of the measurements of OH and HO<sub>2</sub> is covered in the review of Heard and Pilling (2003). As they wrote “clearly, OH plays a central role in tropospheric chemistry. The in situ measurement of its concentration has long been a goal, but its short lifetime and consequently low concentration provide a serious challenge.”


In order to assess the global impact of OH chemistry in the absence of direct measurements, reactive proxies have been used. Singh (1977) used methyl chloroform to estimate OH abundance since methyl chloroform is exclusively anthropogenic and were emissions are known. This type of work provided a comprehensive picture of the global distribution of OH and, hence, a first overall look at the oxidative capacity of the atmosphere. It was followed, using halocarbon measurements by the AGAGE network, by a global OH determination, while also introducing the atmospheric chemistry community to formal inverse modelling (Prinn et al., 1995). Spivakovsky and co-workers expanded on this work to derive 3-D distributions of OH and used this information to assess the wider impact on the lifetimes of

halocarbons, which have implications for stratospheric ozone (Spivakovsky et al., 2000). Thanks to the availability of long term observations of halocarbons from the AGAGE and NOAA networks, later work using a similar approach found evidence for substantial variations of atmospheric hydroxyl radicals in the previous two decades (Prinn et al., 2001), thus providing a broad overview not only of the global distribution but also of the temporal variability of this crucial species. Such estimates allowed for the quantification of the lifetime of important chemicals such as methane and CFC-substitutes such as the hydrochlorofluorocarbons (HCFCs) and hydrofluorocarbons (HFCs).

The in-situ OH detection in the troposphere has proven elusive for a long time. The use of laser-induced fluorescence provided some of the first clues to its atmosphere concentrations in the 1970s and early 1980s (Davis et al., 1976; Wang et al., 1975), but many of these early measurements were found to have significant artefacts. Long-path UV absorption in Germany showed the OH abundances in German boundary layer to be around  $1\text{--}4 \times 10^6 \text{ cm}^{-3}$  (Perner et al., 1976b). The study by Eisele (1994) at the Fritz Peak Observatory in Colorado, was the first intercomparison experiment of different measurement techniques and provided much needed confidence in the observations of this key molecule. Stevens et al. (1994) developed the low-pressure laser-induced-fluorescence (LIF) instrument, which quickly became one of the most successful and widely used techniques for ambient measurements of OH and HO<sub>2</sub>. As ambient observations of HO<sub>x</sub> became available, they were found useful to test our understanding of tropospheric chemical processes, by comparing them with the results of chemical models (see the 2.6. *Chemical Models* section). Recognised as a foundational paper in this area, Ehhalt (1999) explained with clarity the role of radicals in tropospheric oxidation and what controls their concentrations, using both ambient measurements and calculated concentrations of OH. The OH radical is particularly suited to test our understanding of chemical processes and this was clearly demonstrated in 2009, when the discrepancies between observed and calculated OH and HO<sub>2</sub> in the polluted region of Southern China led Hofzumahaus and co-workers to propose a regeneration pathway for OH, which does not involve NO<sub>x</sub> and thus does not produce O<sub>3</sub> (Hofzumahaus et al., 2009). This, together with the work by Lelieveld and co-workers (Lelieveld et al. 2008), prompted a major reassessment of the isoprene oxidation mechanism by Peeters et al. (2009) who suggested that isomerisation of hydroxyperoxy radicals from isoprene oxidation could be fast enough to regenerate HO<sub>x</sub> in highly forested, low NO<sub>x</sub> environments (see the 2.4. *Chemical Kinetics, Laboratory Data and Chemical Mechanisms* section) and led to a major revision of isoprene chemistry and its role in the troposphere (see the 2.13. *Biogenic Emissions and Chemistry* section).

The sources and sinks of HO<sub>x</sub> radicals have always been a major research focus (Finlayson and Pitts, 1976) and the work of Paulson and Orlando (1996) on the reactions of ozone with alkenes as a source of HO<sub>x</sub> in the boundary layer is widely recognised. Radical chemistry is highly sensitive to the levels of NO and NO<sub>2</sub> and Kleinman's modelling work on hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) concentrations in the boundary layer is recognised for its simple elegance in describing how the HO<sub>x</sub> cycle chemistry is influenced by NO<sub>x</sub> and in giving insight into the differing fates of OH and HO<sub>2</sub> radicals under different NO<sub>x</sub> regimes (Kleinman, 1991) (Figure 10).






The first direct measurements of OH lifetime (Di Carlo et al., 2004) provided evidence of missing reactivity, i.e., that not all sinks of the OH radical are known, which relates to earlier work by Lewis et al. (2000) on unmeasured volatile organic compounds (see the 2.12. *Volatile Organic Compounds* section).

## 2.10. Nighttime Chemistry

There is widespread recognition that the atmosphere's oxidative chemistry is active during the night as well as during the day. Evidence of nighttime chemistry driven by the nitrate radical (NO<sub>3</sub>) and ozone was first observed in the (polluted) troposphere in 1980 by Platt and co-workers (Platt et al., 1980). Much of the early NO<sub>3</sub> work, including laboratory and field studies, is summarised in Wayne's seminal review (Wayne et al., 1991). Platt and colleagues and Plane and colleagues' ground-breaking work based on long-path absorption have shown the importance of NO<sub>3</sub> in the troposphere (Allan et al., 1999; Platt et al., 1979).

Two papers that have been highly influential in shaping our view of nocturnal chemistry are "Nitrogen oxides in the nocturnal boundary layer: Simultaneous in situ measurements of NO<sub>3</sub>" (Brown et al., 2003) and "Variability in Nocturnal Nitrogen Oxide Processing and Its Role in Regional Air Quality" (Brown et al., 2006). Both these papers showed the power of state of the art measurements coupled with models to assess the impact of nocturnal and heterogeneous chemistry on regional air quality. In particular, the paper by Brown et al. (2006) was a powerful demonstration of the role of heterogeneous chemistry and aerosol particle composition in controlling dinitrogen pentoxide (N<sub>2</sub>O<sub>5</sub>) and, therefore NO<sub>3</sub>, concentrations (Figure 11).




The area of  $\text{NO}_3$  chemistry is very active and there have been significant further studies since Brown et al. work. Another area of particular note for nighttime processes are those due to the Criegee intermediate. The role of Criegee intermediates have been known for a while. However, recent ability to isolate and measure the reactivity of this intermediate is showing the importance of this radical.

### 2.11. Halogen Chemistry

In comparison to the atmospheric chemistry in the stratosphere, where halogen chemistry has been well known and characterized for a long time (see the 2.18. *Stratospheric Chemistry* section), the recognition of the role of halogen species in the oxidative chemistry of the troposphere occurred much later. Reviews of the earlier work can be found in Cicerone (1981), Platt and Hönninger (2003), Monks (2005) and the extensive review by von Glasow and Crutzen (von Glasow and Crutzen, 2007).

The role of halogens in the troposphere has been discussed going back to the 1970s (e.g. (Graedel, 1979)). The potential importance of iodine in the troposphere was highlighted by a seminal paper by Chameides and Davis in 1980 (Chameides and Davis, 1980). An important early paper is that from

Barrie in 1988 (Barrie et al., 1988) that demonstrated the dramatic impact of bromine chemistry on Arctic boundary layer ozone (Figure 12). The occurrence of ozone depletion events in the polar boundary layer suggested that halogens could have a significant impact on atmospheric chemistry at low altitudes and not just in the stratosphere. This work brought together halogen and heterogeneous chemistry and led to the discovery of bromine catalyzed ozone depletion on ice-covered surfaces (see 2.5. *Heterogeneous and Multiphase Chemistry*).



One of the foundational papers in the area of halogen chemistry is the modelling study by Vogt et al. (1996) which set the theoretical framework for the sea-salt activation mechanism for halogen release and linked halogen chemistry with the sulphur cycle. While the initial research focus was on chlorine and bromine, Alicke et al. (1999) reported the first iodine oxide observations in the marine boundary layer at Mace Head, Ireland, and proved that iodine can also be an important player in the chemistry of the troposphere. Further investigation found evidence that biogenic iodine species can be responsible for the formation of marine aerosol and cloud condensation nuclei (O'Dowd et al., 2002) recognizing the potential for wide scale impact of iodine chemistry in particle formation. Finlayson-Pitts and her colleagues had suggested the importance of chlorine in tropospheric chemistry based on laboratory data (Finlayson-Pitts et al., 1989) as has been seen earlier by Schroeder and Urone (1974), but it wasn't until 2008 that Osthoff and co-workers (Osthoff et al., 2008) – and the related comment “When air pollution meets sea salt” by von Glasow (2008) – brought attention to the potential for nitryl chloride (ClNO<sub>2</sub>) chemistry to impact ozone formation, nitrogen recycling and VOC

oxidation, with the first ambient observations of this molecule. Also in 2008, the work from Read et al. (2008) clearly showed the global importance of halogens for tropospheric ozone using long-term observations of iodine and bromine oxides (IO, BrO) made at the Cape Verde Atmospheric Observatory. High concentrations of tropospheric Cl<sub>2</sub> first reported by Spicer et al. (1998) have been found in other places.

## 2.12. Volatile Organic Compounds

Volatile organic compounds (VOCs) embraces a wide variety of species emitted from man-made and natural sources. In many respects VOCs are the fuel of the oxidative chemistry in the atmosphere, involved in many gas- and particle-phase processes.

Ehhalt (1974) brought together the details of the methane sources and sinks and put them into a consistent framework that described the life cycle of methane. This conceptual framework has subsequently been expanded to a wide range of trace organic gases. The original understanding of the life cycle of methane has remained largely unchanged over the subsequent 40 years and has formed the basis of the IPCC science assessments on the role of methane in global warming and climate change. Methane itself has been long recognised as important for tropospheric chemistry, but also for climate change as a greenhouse gas and as a source of water vapour to the stratosphere. Specifically, the work of Blake and Rowland (1986) documented the global increase in methane and its implications for climate change.

The large differences in reactivity among the individual VOCs have always been a feature of their chemistry. Darnall et al. (1976) produced a reactivity scale for atmospheric hydrocarbons<sup>2</sup> based on their reaction with the hydroxyl radical, an idea that is still influential to the present day. The concept was further advanced by Carter and Atkinson (1989) who looked at incremental hydrocarbon reactivity, where knowledge of the reactivities of organics with respect to ozone formation in the atmosphere can provide a useful basis for developing appropriate control strategies to reduce ambient ozone levels. It was the beginning of an approach that is often now used in regulation to determine which organic compounds would have the greatest effect in reducing ozone.

VOC transformation can be important in a number of different atmospheric processes. One highly cited early example is the work of Pitts et al. (1978) on the atmospheric reactions of polycyclic aromatic hydrocarbons and their ability to form mutagenic nitro derivatives under typical atmospheric conditions.

While measurement techniques rarely seem to get a mention as being influential, the discipline relies on observations as a critical part of the oeuvre. Already mentioned was the huge impact that accurate techniques to measure the OH radicals had on the development of the field (see 2.9. *HOx chemistry*). Another example is the development of Proton-Transfer Reaction Mass-Spectrometry, which has revolutionised the measurement, in particular, of VOCs (Lindinger et al., 1998) and the earlier work of Lovelock and Lipsky (1960) on the development and application of electron capture detectors that allowed the measurement of VOCs such as dimethyl sulphide and the halocarbons in the troposphere.

---

<sup>2</sup> Hydrocarbons, although often used interchangeably with VOC, do not describe the same group of compounds. Hydrocarbons are a subset of VOC that exclusively contain hydrogen and carbon, and thereby include none of the e.g., oxygenated VOC species.

Research is ongoing as to how many VOCs there are in the atmosphere and what the consequences are of not being able to measure/quantify them all. The work of Lewis et al. (2000) used novel VOC measurements (GC x GC) to find that there was a larger pool of ozone-forming carbon compounds in urban atmospheres than previously posited (Figure 13). The later paper by Goldstein and Galbally (2007) expanded on this work hypothesizing that thousands of VOCs are still unmeasured and unknown, with potentially huge consequences for the carbon budget of the atmosphere.

Continuing work in this area, de Gouw et al. (2005) produced a landmark study that combined analysis of organic carbon in both the gas- and particle-phase in the polluted atmosphere as part of the New England Air Quality study by looking at the evolution of VOCs from their emission sources. The study showed that most of the organic carbon in the particle-phase was formed by secondary anthropogenic processes and that an increasing fraction of the total organic mass was constituted of oxygenated VOCs as a result of the air masses being processed/aged.




Figure 13 - Comprehensive (right) and one-dimensional (left) separations of volatile organic compounds in urban air demonstrating a greater variety and number of VOCs (Lewis et al., 2000)


### 2.13. Biogenic Emissions and Chemistry

Although it has been known for a long time that plants emit organic compounds, the relevance of biogenic VOCs for atmospheric chemical processes was not immediately recognized. The first report

that plants emit volatile organic compounds into the atmosphere was made in 1957 by the Georgian scientist Guivi Sanadze (Sanadze, 1957). Unaware of Sanadze's work in the USSR, Rasmussen and Went independently discovered isoprene emissions in 1964 (Rasmussen and Went, 1964). Sanadze was also the first to show that isoprene emission rates are temperature dependent (Sanadze and Kursanov, 1966). However, the relevance of biogenic VOC for atmospheric chemical processes was not immediately recognized. Went (1960) hypothesised that "volatilisation of terpenes and other plant products results in the production of, first, blue haze, then veil clouds ... ". Although Tingey (Tingey et al., 1979) at the US Environmental Protection Agency did note the potential for isoprene to play a role in regional air quality in 1978, this was not formalised until the ground-breaking work of Chameides and colleagues in 1988 (Chameides et al., 1988) and MacKenzie and colleagues in 1991 (MacKenzie et al., 1991).

In 1992 the seminal review of Fehsenfeld et al. (1992) brought the importance of isoprene and a wide range of other VOCs of biological origin to the attention of the atmospheric chemistry community, opening up an entirely new branch of atmospheric chemistry. Other influential reviews of biogenic VOC emissions, include e.g., the physiology of plants (Kesselmeier and Staudt, 1999) and more recently Sharkey and Monson (2017) reviewed the enigmatic nature of isoprene emissions.

Over time, biogenic chemistry became pivotal for major policy formulations to abate ozone pollution. Underpinning the atmospheric chemistry research that Fehsenfeld et al. (1992) promoted, plant physiologists began working on understanding the biological and environmental controls on biogenic VOC emission rates. This work allowed the development of relatively simple functions to predict the emissions of biogenic VOCs which resulted in the first spatially and temporally resolved global model of biogenic emissions (Guenther et al., 1993). These soon evolved into more sophisticated high resolution global models (Guenther et al., 2000; Guenther et al., 1995), allowing for the emissions of biogenic compounds to be included in atmospheric chemistry models across all scales. Eventually, this work took the form of the widely used MEGAN (Model of Emissions of Gases and Aerosols from Nature) model (Guenther et al., 2006) which is still used in modern Earth system models today (Table 1 and Figure 14).



887

888 Terrestrial vegetation is not the only source of biogenic emissions. Aneja et al. (1979) discussed the  
 889 importance of biogenic sulphur compounds and their role in stratospheric chemistry, while Charlson  
 890 et al. (1987) connected marine biology, atmospheric chemistry and climate into the already  
 891 mentioned CLAW hypothesis (see the 2.2. *Aerosols and Clouds* section).

892 Marine aerosol formation was thought for a long time to be dominated by inorganic components,  
 893 mainly sea-salt and non-sea salt sulphate, but O'Dowd and co-workers (O'Dowd et al., 2004) showed  
 894 that biological activity of plankton blooms can enhance the concentration of cloud condensation  
 895 nuclei, a key aspect of the chemistry-climate feedback mechanism. A similar mechanism is also active  
 896 in relation to biogenic halogen compounds (O'Dowd et al., 2002) which also affects aerosol formation  
 897 as well as the ozone, nitrogen, and sulphur cycles, as discussed in section 2.2. *Aerosols and Clouds*.

898 A paper that has been defined as “controversial but set off a huge amount of activity” is “Atmospheric  
 899 oxidation capacity sustained by a tropical forest” by Lelieveld et al. (2008), which proposed a new



chemical mechanism for low NO<sub>x</sub>, high VOC regions (such as tropical forests), based on modelling studies of a field dataset. Although further studies contradicted this hypothesis, Lelieveld et al. (2008) was instrumental in prompting a large amount of laboratory, theoretical and field studies in the past 10 years. These studies resulted in a major revision of our understanding of biogenic VOC chemistry (see the sections 2.4. *Chemical Kinetics, Laboratory Data and Chemical Mechanisms* and 2.9. *HO<sub>x</sub> chemistry*).

#### 2.14. *Biomass Burning*

Biomass burning, particularly in the tropics, affects terrestrial vegetation dynamics, soil erosion, movement of organic carbon, hemispheric atmospheric composition, air quality and more broadly radiative forcing *via* emissions of trace gases and aerosols (Monks et al., 2009). Crutzen (Crutzen et al., 1979) was the first to highlight biomass burning in the tropics as an important source of atmospheric gases, such as molecular hydrogen (H<sub>2</sub>), CO, N<sub>2</sub>O, NO, chloromethane (CH<sub>3</sub>Cl) and carbonyl sulphide (COS). The importance of biomass burning, based on the observations of a small set of fires, and the appreciation of its potential role was a major step in our understanding of the role of biomass burning in air quality, climate change, and the composition of the troposphere. It is, however, the later paper “Biomass Burning in the Tropics: Impact on Atmospheric Chemistry and Biogeochemical Cycles” by Crutzen and Andreae (Crutzen and Andreae, 1990), one of the top 10 most cited Atmospheric Chemistry papers (Table 1), that has had the greatest impact on this research area, providing quantitative estimates of the amounts of biomass burning taking place around the world and the resulting emissions, recognizing the critical role of biomass burning emissions in the Tropics and from activities in developing countries that were not well documented. Hao and Liu (1994) made a further advance, looking at where and when biomass burning and thereby the related emissions occur. They developed an improved database of the amount of biomass burned owing to deforestation, shifting cultivation, savannah fires, fuel wood use, and clearing of agricultural residues, focused on tropical America, Africa and Asia during the late 1970s.

Simoneit et al. (1999) introduced the important concept that “the monosaccharide derivatives (e.g. levoglucosan) are proposed as specific indicators for cellulose in biomass burning emissions.” They showed that levoglucosan is emitted at such high concentrations that it can be detected in air pollution filter samples at considerable distances from the original combustion source, allowing for source apportionment.

The 2001 paper “Emission of trace gases and aerosols from biomass burning” by Andreae and Merlet (2001), which is also one of the top 10 most cited papers (Table 1 and Figure 15), pulled together emission factors for a large variety of species emitted from biomass fires and is considered a key reference for biomass burning emission factors. Further work in the biomass burning area was later presented by Reid et al. (2005) in a review paper where they looked at measurements of smoke particle size, chemistry, thermodynamic properties, and emission factors from a variety of sources, including laboratory burns, in-situ experiments, remote sensing and modelling. They brought together information from the ‘milieu of small pieces of the biomass-burning puzzle’ and showed that there are large differences in measured particle properties and particle carbon budgets across the literature. van der Werf et al. (2006) investigated interannual variability and the underlying mechanisms regulating variability at continental to global scales using a time series of eight years of satellite and



model data. Total carbon emissions was driven by burning in forested areas, while the amount of burned area was driven by savannah fires, which are influenced by different environmental and human factors than forest fires.




Figure 15 - Vertical profiles of  $O_3$  in the tropical troposphere. The profile over the equatorial Pacific shows no influence from biomass burning, whereas the profile over the Pacific off South America suggests  $O_3$  enhancement due to long-range transport from the tropical continents. The  $O_3$  profiles over Brazil and the Congo show high  $O_3$  concentrations at altitudes between 1 and 4 km due to photochemical production in biomass burning plumes. At higher altitudes,  $O_3$  concentrations are also substantially enhanced, possibly also because of  $O_3$  production by reactions in the emissions of biomass burning (Crutzen and Andreae, 1990).

## 2.15. Emissions and Deposition

Non-chemical sources and sinks of various species are critical components of atmospheric processes and therefore are particularly essential for global and regional models. An early advocate for such was the work of Olivier and the team that created the EDGAR (Emission Database for Global Atmospheric Research) database (Olivier et al., 1994).

Emissions from vehicles and power plants have always been an essential aspect of air quality related policies, and therefore an area where more focused inventory work was needed and done. This approach was pioneered in California in the early 1990s, and the studies by Calvert et al. (1993), Lawson (1993) and Singer and Harley (1996) helped define and verify the California Smog Control Program, providing a solid scientific basis with reliable emissions data. Techniques such as the remote monitoring of traffic generated carbon monoxide (Chaney, 1983; Bishop and Stedman, 1996) are also essential to recognise for the role they played in understanding vehicle emissions.

Agricultural emissions (from both crop and animal production) play an important role in several atmospherically mediated processes of environmental and public health concerns (Chameides et al.,

1999). These atmospheric reactions and processes affect local and regional environmental quality, including odour, particulate matter (PM) exposure, eutrophication, acidification, exposure to toxics, climate, and pathogens (Erisman et al., 2008; Aneja et al., 2009). Agricultural emissions also contribute to the global problems caused by greenhouse gas emissions, specifically nitrous oxide and methane.

The deposition of gases and aerosol particles to the surface is another critical process in the atmosphere. Chamberlain (1966) is credited with the first exposition of the resistance network approach to describe the uptake of gases on surfaces and the identification of transport through the atmospheric boundary layer, through the surface layer, and through the stomata on plants, as important elements of surface uptake. Building upon this work, a comprehensive and widely adopted, parametrization of the dry deposition process for regional and global models was presented in the late 1980s by Wesely (1989). Currently, there has been renewed interest in quantifying and understanding deposition processes. Yet, a systematic description based on fundamental independently measureable physico-chemical parameters is lacking. A complication can arise from a range of oxygenated VOC that can exhibit bi-directional exchange above vegetation (Karl et al., 2010).

#### 2.16. *Chemical Transport*

Transport is an integral part of atmospheric processes and influences atmospheric composition across a range of spatial scales. As early as 1975, Junge (1975) pointed out the importance of the atmospheric residence time of a constituent with respect to global transport and dispersion. Prather's work (Prather, 1994, 1996) provided new insights into timescales for atmospheric oxidation chemistry.

Stratospheric-tropospheric exchange (STE) has always been recognised as a key mechanism in determining tropospheric composition. Early chemical dynamics were demonstrated by Danielsen (1968) that laid the foundations for 3-dimensional modelling of chemical transport looking at stratosphere-troposphere exchange based on radioactivity, ozone, and potential vorticity. Later, Holton and co-authors (Holton et al., 1995) proposed an approach that placed stratosphere-troposphere exchange in the framework of general circulation and helped clarify the roles of the different mechanisms involved and the interplay between large and small scales, by the use of dynamical tracers and potential vorticity. This work is recognised as a big step forward for the understanding of the tropospheric ozone budget. Stohl and colleagues (Stohl et al., 2003) brought together what has been viewed by many as the authoritative work on stratosphere-troposphere exchange.

While the regional nature of air pollution has always been recognised, that is less the case for the impact of trans-continental emissions on air quality. Jacob et al. (1999) showed that there was a need for a global outlook for understanding regional air quality and meeting pollution reduction objectives. This perspective spawned a decade of intense work on intercontinental air pollution and transport. Well recognised in this area is the work by Stohl and colleagues (Stohl et al., 2002) who mapped out the pathways and timescale of intercontinental air pollution transport and brought life to the subjects of atmospheric dynamics and transport of air pollution (Figure 16). Observational studies such as Merrill et al. (1985), Moody et al. (1995), Stohl and Trickl (1999) and Forster et al. (2001) showed the range of mechanisms and impact of long-range transport of air pollution.




Figure 16 - Total columns (a, c) and zonally averaged mixing ratios (b, d), both divided by the respective time interval, of a Asia tracer for ages of 6 – 8 days (a, b) and 25– 30 days (c, d) during DJF. The plots shows the horizontal and vertical impact of a pollution tracer (Stohl et al., 2002).

Moody et al. (1998) explored atmospheric transport history using back trajectories for the Harvard Forest experiment demonstrating the power of trajectory methods at the regional scale. Key tools in the development of this area were, the early simplistic isentropic trajectory methods (Merrill et al., 1985), the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model (Draxler and Hess, 1998), a forerunner of various particle trajectory and dispersion models, which developed into a widely used particle dispersion model, and the FLEXible PARTicle dispersion (FLEXPART) model, a Lagrangian particle dispersion model designed for calculating the long-range and mesoscale dispersion of air pollutants (Stohl et al., 2005).

The ideas of intercontinental source-receptor relationships were embodied in the aforementioned works by Jacob et al. (1999) and Stohl et al. (2002). The long-range transport concept was developed in an effort to explore the source-transport relationships that drive observed ozone concentration in regions farther away from the emission regions (Fiore et al., 2009) (see the 2.6. *Chemical Models* section). This large community collaboration (Fiore et al., 2009) provided valuable insights into the sensitivities of the hemispheric regional background of ozone and how this is controlled by emissions from continental source regions.


Another critical area for atmospheric chemistry is boundary layer dynamics and meteorology. This is particularly important since most emissions are emitted in the boundary layer. Atmospheric dynamics in this important region have been mostly expressed as parameterizations in numerical models (Stull, 1988). The spatial and temporal scales involved in the processes in this region cover a wide range.

The understanding of this region has been mostly based on meteorological and energy/water vapour balance points of view. However, the chemical transformation and dispersion in this region are crucial for how much chemicals actually get out of this region to influence the regional and global atmosphere. Furthermore, the process of dry deposition, a critical loss processes for chemicals, is mostly limited to the boundary layer.

## 2.17. Satellites and the Troposphere

The importance of satellites for the discipline of atmospheric chemistry centres on the ability to give a self-consistent global view of a selected set of tropospheric trace species (Burrows et al., 2011; Martin, 2008; Prospero et al., 2002). The beginning and first demonstration of the effective application of these attributes for the troposphere were the data and the retrievals from the GOME instrument (Burrows et al., 1999) on ERS-2 and SCIAMACHY on Envisat (Bovensmann et al., 1999). Historically, the roots of these early instruments are in stratospheric chemistry, with GOME being deployed to be able to track stratospheric ozone and its key controlling chemical species. Much effort has flowed with measurements from instruments such as OMI, MOPITT, TES, MODIS and ACE as well as shuttle borne instrumentation, e.g. CRISTA (Burrows et al., 2011; Martin, 2008).

The most nominated paper in this area, and one that demonstrated the power of such observations for tropospheric composition research, was "Increase in tropospheric nitrogen dioxide over China observed from space" by Richter et al. (2005), which showed the capability of the satellites to track the build-up of air pollution over vast regions from space (Figure 17). The importance of the work lead by Palmer and co-workers in establishing a method to convert satellite observations to vertical columns for comparison with e.g. models was also widely recognised (Palmer et al., 2001).



Satellite-based instrumentation can measure not only gas-phase trace species but also dust and aerosol particles; mapping of the global distribution of dust (Prospero et al., 2002), the combination of SEAWiFs and TOMS to track Asian dust events (Husar et al., 2001) and the development of the MODIS aerosol algorithm (Remer et al., 2005) provided convincing demonstrations of this capability. A step-change in this area was made with the paper Wang and Christopher (2003) “Intercomparison between satellite-derived aerosol optical thickness and PM<sub>2.5</sub> mass: Implications for air quality studies” which was the first description of the derivation of surface PM<sub>2.5</sub> from satellite aerosol optical depth (AOD), built on by Liu et al. (2004) for the USA. Later, a global picture was developed by van Donkelaar et al. (2006). These type of observations have been extensively used to estimate the global impact of particulate matter (both PM<sub>2.5</sub> and PM<sub>10</sub>) on health.

## 2.18. Stratospheric Chemistry

Tropospheric chemistry has always been influenced by the study of stratospheric chemistry. At the same time, tropospheric chemistry has been pivotal in determining what surface emissions get to the stratosphere. As mentioned earlier (see the 2.1. *Foundations* section), the basis of stratospheric ozone chemistry was laid in the 1930s (Chapman, 1930) (for a full history see Brasseur (2019)), whereas tropospheric chemistry followed a couple of decades later. Interest in stratospheric ozone chemistry increased substantially following the works of Johnston and Crutzen on the role of nitrogen oxides in the stratosphere (Crutzen, 1970; Johnston, 1971). The impact of supersonic transport (Johnston, 1971) and of chlorofluorocarbons (CFCs) (Molina and Rowland, 1974) were important events for stratospheric ozone chemistry. While Crutzen (1970) showed that the nitrogen oxides in the stratosphere come mostly from the nitrous oxides from the ground, Johnson suggested that a fleet of supersonic aircraft could release large amounts of nitrogen oxides into the lower stratosphere causing substantial ozone loss (Johnston, 1971). The potential threat of supersonic transport highlighted the importance of gas-phase catalysis in the atmosphere and, in particular, of the catalytic ozone destruction by HO<sub>x</sub> and NO<sub>x</sub>. These works opened the world’s eyes to the potential for global environmental change from human activities. Soon after, Lovelock (1974) identified CFCs in the troposphere and showed that practically all the CFCs emitted to date were still in the atmosphere. The significant contributions of Hampson (1964), Crutzen, and Johnston, and the recognition of chlorine-catalysed ozone destruction by Stolarski and Cicerone (1974), paved the way for the seminal work of Molina and Rowland (Molina and Rowland, 1974) linking chlorofluorocarbons to ozone layer depletion. The recognition that bromine compounds can also destroy stratospheric ozone (McElroy et al., 1986) further refined the story. The potential role of iodine in stratospheric ozone depletion has also been raised (Solomon et al. 1994), but it is still somewhat unsettled.

The ozone hole (Farman et al., 1985) was an unanticipated shock that awoke the world to the global nature of ozone layer depletion. The origin of the ozone hole was understood in an historic set of studies over a relatively short five-year period. First was the insightful and seminal work of Solomon et al. (1986) that showed that chlorofluorocarbons and other ozone-depleting gases were the key anthropogenic ingredient for the ozone hole. The confluence of cold temperatures that lead to the formation of polar stratospheric clouds (PSCs) and the winter vortex formation over Antarctica provided the opportunity for the massive ozone depletion that resulted in the ozone hole. This work confirmed the suggestion of Farman (Farman et al., 1985) that the ozone hole was due to the increasing abundances of CFCs. In particular, Solomon and co-workers (Solomon et al., 1986)

recognized that stable molecules such as ClONO<sub>2</sub> and HCl could react on solids (and indeed liquids). Along the way, during this intense investigative period, the detection and quantification of the role of ClO as a catalyst by Anderson et al. (1991), as well as De Zafra et al. (1988), was the “smoking gun” that linked the CFCs with the ozone hole. The entire set of field measurements, from the ground, aircraft, and balloons, solidified this linkage.

Less heralded, but equally important, were the laboratory studies that showed that chlorine nitrate (ClONO<sub>2</sub>) and HCl did indeed react on PSCs (Hanson and Ravishankara, 1994; Tolbert et al., 1987; Leu, 1988; Molina, 1991; Molina et al., 1987) and determined the critical rate coefficients for the self-reaction of ClO, the rate-limiting step in the unique catalytic cycles in Antarctica (Cox and Hayman, 1988; Sander et al., 1989; Trolier et al., 1990). Much was learned in later years by studying the Arctic and from the continued observations over the Antarctic. It should be noted that the termolecular reaction of ClO was suggested to be important for high chlorine chemistry by Molina and Molina (Molina and Molina, 1987), and the history of this reaction goes back to Norrish’s work at Cambridge (Norrish and Neville, 1934). One of the lessons from this episode is that natural factors, in this case the formation of a vortex and the occurrence of polar stratospheric clouds, can lead to unexpected consequences when an anthropogenic ingredient (ozone-depleting chemicals) is added to the mix.

The numerical modelling of the stratosphere was an important ingredient for the success of mitigating polar ozone loss, along with the theories of the ozone layer depletion and the ozone hole, the laboratory studies of key processes, and the measurements in the atmosphere. Over the years, these models have enabled a great deal of understanding of the coupling between chemistry and climate. The development of stratospheric chemical transport models (Chipperfield and Pyle, 1988) was a pivotal advancement that enabled quantitative understanding of the ozone layer depletion, including the ozone hole (see the 2.6. *Chemical Models* section).

The weight of science led to the Vienna convention, the Montreal Protocol, and the Protocol’s many amendments and adjustments that are leading to the phasing out of the ozone-depleting gases. The Montreal Protocol is the first international treaty on an environmental issue to be universally ratified and is regarded as one of the most successful. That said, the ozone layer depletion story is not complete. For example, the recognition that nitrous oxide is the remaining major ozone-depleting gas emission (Ravishankara et al. 2009) has connected food production (tropospheric nitrogen cycles) to ozone layer depletion and highlighted the importance of a holistic approach to environmental issues.

#### *2.19 Other issues that influence tropospheric chemistry*

Atmospheric chemistry advances have been influenced by growth in other areas. In particular, the importance of anthropogenic climate change has been instrumental in invigorating atmospheric chemistry studies because many of the major climate forcing agents are chemically active and climate change, in turn, influences tropospheric chemistry (von Schneidemesser et al., 2015). In addition to climate change, other adjacent discoveries and findings have influenced tropospheric chemistry studies.

The global atmospheric and climatic consequences of nuclear war were investigated by both Crutzen and Birks (1982) and Turco et al. (1983). Using models developed for looking at the impact of volcanic eruptions, Turco et al. (1983) concluded that “enhancement of solar ultraviolet radiation due to ozone

depletion, long-term exposure to cold, dark, and radioactivity could pose a serious threat to human survivors and to other species.” Similarly (Crutzen and Birks, 1982) concluded that “the screening of sunlight by the fire-produced aerosol over extended periods during the growing season would eliminate much of the food production in the Northern Hemisphere”.

Air quality has an obvious direct impact on people, and this connection was recognized very early (it was in fact the primary motivation behind the fundamental work of Haagen-Smit). In 1993, Dockery et al. (1993) presented a study of six US cities showing a direct association between air pollution and mortality rates. This paper is a great example of how an adjacent field influences another, in this case atmospheric chemistry and public health. Though association between air pollution and health stretch back to the Los Angeles and London smog, the ‘Six Cities Study’ was a landmark as it demonstrated that the association between air pollution and mortality extended to much lower concentrations than those observed in the smog days.

### 3. Discussion and Summary

A mixture of the history of the discipline and of its the landmark ideas emerges from this exercise of asking the community what they consider to have shaped their research field. Table 2 seeks to bring these elements together to look at the evolution of the leading scientific concepts, their relevance to the environmental legislation (in this sense, we acknowledge an Euro-/US-centric bias), and the most notable environmental events that have shaped the discipline. Atmospheric science often sits at an interesting intersection between the societal interests (e.g. acid precipitation, air quality, ozone layer depletion, and climate) and its scientific venture. Monks and Williams (2020) have recently explored how environmental events in air quality drive policy and how a scientific and societal paradigm shift occurs once the emergency phase has passed.

From an overview of all the nominated papers, several general features are apparent. Ambient measurements are one of the cornerstones of atmospheric science (Abbatt et al., 2014). It is clear that the atmosphere is under-sampled, but over time we have found many ingenious ways to build different measurement strategies from the ground, ships, aircraft, balloon, sonde and satellites. With a focus on chemistry, it is clear that one needs to be able to measure with surety, sensitivity, specificity and speed in the troposphere. Many of the nominated papers reflect the importance of instrument development. Examples include the electrostatic sizers in aerosol science (Knutson and Whitby, 1975), various techniques to measure the hydroxyl radical (Eisele, 1994;Stevens et al., 1994;Perner et al., 1976a), chemiluminescence for NO/NO<sub>2</sub> (Kley and McFarland, 1980), the development of chemical ionization mass spectrometry (Lindinger et al., 1998), the application of the GCxGC-MS technique (Lewis et al., 2000), and aerosol mass-spectrometry measurements, e.g., Zhang et al. (2007). Often, the science underlying the development of these instruments, such as ion-molecule chemistry, is not necessarily acknowledged in the community. The paradigm for field instruments has been the development of analytical methods in the laboratory that are then adopted and/or adapted for field studies. The advances in associated fields such lasers, optics, optical detectors (e.g., camera and diode arrays), mass spectrometry (such as ion-traps and high-resolution time-of-flight mass spectrometry), separation methods (such a various chromatography methods), and meteorological instruments have fundamentally altered atmospheric chemistry in general and tropospheric chemistry in particular. For example, the recent developments in chemical ionization mass spectrometry have led to their

pervasive use in our science. It is fair to say that the ability to separate and measure constituents in the part-per-quadrillion and part-per-trillion mixing ratio range have led to the detections of miniscule amounts of chemicals and their variations. Similarly, the recent revolution in low-cost sensors coupled with the evolution in telecommunications already is and will continue to change our field in the near future. Yet another important area is the development of the details of the chemistry through observations of intermediates and products, etc. These studies have provided some of critical information regarding the details of the chemistry (Cohen and Murphy, 2003).

Another common theme is the critical importance and impact of long-term observations, often termed monitoring, of key atmospheric components, from CO<sub>2</sub> (the “Keeling” curve) to chemically active molecules such as halocarbons (the Antarctic ozone hole, ozone layer depletion and climate change), methane (the changes in the global OH field, background ozone production), and NO<sub>x</sub> (catalyst for tropospheric ozone production, vehicle emissions and acid precipitation). On the other hand, many breakthroughs in understanding the observations emerged because of basic laboratory information on kinetics and photochemistry (e.g., the reaction of HO<sub>2</sub> + NO, the determination of O(<sup>1</sup>D) quantum yields, and the reactions of ClONO<sub>2</sub> and HCl on PSCs.) It is noteworthy that both laboratory studies and long-term observations are currently under funding stress, a situation that is already worrying the community (see, for example, the discussion in Burkholder et al. (2017)).

There is no doubt that atmospheric chemistry is an integrative science: one of the recurring themes in the papers discussed here is the tight relationship between ambient observations, laboratory experiments, and modelling. The integrative power of models has been recognised from the early studies by Levy (1971) to the development of highly sophisticated global transport models by Chipperfield and Pyle (1988) and Bey et al. (2001) up to the more recent demonstrations of the power of model ensembles (Stevenson et al., 2006; Fiore et al., 2009). Much of this progress has parallels in stratospheric chemistry. It is evident in the community that models are a powerful tool to map, test, and predict the atmosphere's past, present, and future. The predictions and projections from these models play essential roles in policy, planning, and management of environmental issues.

Another form of integrative or meta-analysis brings together a range of individual studies to produce a more significant outcome, such as new insights or models. There are some notable examples of this approach in the works on biomass burning by Crutzen and Andreae (1990) and more recently, by Van Der Werf et al. (2010). Similarly, the work to produce isoprene emissions models brought together several global isoprene flux measurements (Guenther et al., 1995; Guenther et al., 2006). Other examples include the work of Zhang et al. (2007) and Jimenez et al. (2009) who integrated various sets of AMS observations to give insight into land distributions of SOA.

There have been developments in fields adjacent to atmospheric chemistry that have shaped atmospheric chemistry progress. Examples include developments in epidemiology (e.g., the Six Cities Study by Dockery et al., 1993), atmospheric dynamics (the role of transport in determining chemical composition, the role of the Antarctic vortex), ocean science (pertaining to deposition to the ocean surface and emissions from the oceans), and in biological/plant science (e.g. (Kesselmeier and Staudt, 1999)). Integrating atmospheric chemistry with these adjacent fields is not only essential but also fruitful and was for many years embodied in the IGBP (Seitzinger et al., 2015), the World Climate Research Program (WCRP) and Earth-system science (now Future Earth) programs. Wider contexts, such as paleoclimate, have allowed an understanding of climate and atmospheric history over the 100,000 year timescales (e.g. (Petit et al., 1999)). That work set the framework for understanding that



the present-day atmospheric burdens of carbon dioxide and methane as important greenhouse gases are unprecedented during the past 420,000 years. They also allow us to estimate the composition of the troposphere in those ancient times.

Similarly, there are concepts that have their roots in tropospheric chemistry and have gone on to have wider impact. The concept of the Anthropocene, most recently highlighted by Crutzen and Stoermer in 2000, indicates that we are in a new geological epoch driven by human activities. The idea was more fully expounded in Crutzen (2002) “the geology of mankind”. There is little doubt that this has been a key idea that has influenced much thinking as well as work far beyond atmospheric science (Table 2). As discussed in detail in the introductory text, we opted here to assemble this compilation of papers by reaching out to the community for nominations, rather than using the number of citations as a primary measure. We will not go into the details again here, suffice it to say that there are inherent advantages and drawbacks to any method one might consider for such a work. In that sense, we would like to acknowledge and thank all those who provided feedback during the open peer-review phase of the manuscript.

We would be remiss in not noting that there remain many areas of tropospheric chemistry that are still in their infancy. The chemistry in the boundary layer and the dynamics of this region is one such. This is particularly noteworthy since humans live and emit in the boundary layer. Another such noteworthy sub-area is deposition of chemicals which are still parameterizations. Developing an understanding of the fundamental steps that are independently measured and understood would be a laudable goal for atmospheric chemists. The state and development of emission inventories so key to models remains an area in need of work. New measurements will provide insights into the ever changing nature of our atmosphere across differing spatial and temporal scales.

The papers highlighted capture a substantial scope of the atmospheric research endeavour over the last 60 years. The challenge now for you, the reader, is to continue to reflect on the papers included here and continue this discussion for not only tropospheric chemistry but the related areas.

#### **4. Author Contributions**

PSM developed the concept and led the writing, PSM, ARR, RS, and EvS solicited input, ARR, RS, EvS contributed to writing and editing.

Acknowledgments:

Work of ARR was supported by Colorado State University. The work of EvS was supported by the IASS Potsdam, with financial support provided by the Federal Ministry of Education and Research of Germany (BMBF) and the Ministry for Science, Research and Culture of the State of Brandenburg (MWFK).

1253 **Table 1 – Top 10 Cited Atmospheric Chemistry Papers (Atmospheric+Chemistry)**

1254 (Scopus, 27/3/20)

#	Paper	Title	Citations
1	Guenther et al. (1995)	A global model of natural volatile organic compound emissions	2760
2	Andreae and Merlet (2001)	Emission of trace gases and aerosols from biomass burning	2350
3	Guenther et al. (2006)	Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature)	2175
4	Jimenez et al. (2009)	Evolution of organic aerosols in the atmosphere	1909
5	Atkinson (2000)	Atmospheric chemistry of VOCs and NOx	1773
6	Crutzen and Andreae (1990)	Biomass burning in the tropics: Impact on atmospheric chemistry and biogeochemical cycles	1686
7	Van Der Werf et al. (2010)	Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997-2009)	1578
8	Atkinson and Arey (2003)	Atmospheric Degradation of Volatile Organic Compounds	1502
9	Grell et al. (2005)	Fully coupled "online" chemistry within the WRF model	1436
10	Lelieveld et al. (2015)	The contribution of outdoor air pollution sources to premature mortality on a global scale	1425

1255

1256 (Web of Science, 27/3/20)

#	Paper	Title	Citations
1	Ramanathan et al. (2001)	Atmosphere - Aerosols, climate, and the hydrological cycle	2278
2	Andreae and Merlet (2001)	Emission of trace gases and aerosols from biomass burning	2168
3	Hallquist et al. (2009)	The formation, properties and impact of secondary organic aerosol: current and emerging issues	1988
4	Jimenez et al. (2009)	Evolution of organic aerosols in the atmosphere	1844
5	Crutzen and Andreae (1990)	Biomass burning in the tropics: Impact on atmospheric chemistry and biogeochemical cycles	1603
6	Atkinson (2000)	Atmospheric chemistry of VOCs and NOx	1596
7	Atkinson et al. (1992)	Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry: Supplement IV. IUPAC Subcommittee on Gas Kinetic Data Evaluation for Atmospheric Chemistry	1488
8	Grell et al. (2005)	Fully coupled "online" chemistry within the WRF model	1332
9	Lelieveld et al. (2015)	The contribution of outdoor air pollution sources to premature mortality on a global scale	1292

10	Bey et al. (2001)	Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation	1212
----	-------------------	--	------

1257

1258

1259 **Table 2 – Science, Regulatory and Environmental Landmarks of the 20<sup>th</sup> and early 21<sup>st</sup> Centuries**

Decade	Science Landmark	Regulatory Landmarks <sup>1</sup>	Environmental Events
1930	Chapman Cycles and Stratospheric Chemistry		
1940			1943 – LA Smog
1950	Air Pollution	1956 - UK Clean Air Act	1952 – Great Smog of London
1960		1963 - US Clean Air Act	
1970	Supersonic Transport Stratospheric Chemistry Tropospheric Chemistry Air Pollution and Clouds	1978 – Ban of Lead in Petrol (USA) 1979 - CLRTAP (UNECE)	1974 – Observations of acid rain
1980	Ozone Hole Chemistry Halogen Chemistry Biogenic Chemistry	1987 - Montreal Protocol 1980 - SO <sub>2</sub> directive (EU)	1985 – Observations of the ozone hole
1990	Air Pollution and Health Satellite Observations of the Troposphere Long-Range Transport of Air Pollutants	1992 – Euro 1 Emission standard 1992 - Ozone Directive (EU) 1997 - Kyoto Protocol 1999 - Goteborg Protocol 1999 – Ban of lead in petrol (EU)	1991 - Mt Pinatubo eruption
2000	SOA Concept of Anthropocene	2001 - NEC Directive (EU)	
2010	Air Pollution and Climate		2015 - Dieselgate
2020			2020 - COVID 19

1260

1261 <sup>1</sup> For more details on the UK/EU perspective see (Williams, 2004;Maynard and Williams, 2018) and  
 1262 for the USA perspective see (Jacobson, 2002); see also (Monks and Williams, 2020).

1263 **Table 3** - A global budget for NO<sub>x</sub> (Logan, 1983)

		10 <sup>12</sup> gm N yr <sup>-1</sup>
Sources		
	Fossil fuel combustion	21 (14-28)
	Biomass Burning	12 (4-24)
	Lightning	8 (8-20)
	Microbial activity in soils	8 (4-16)
	Oxidation of ammonia	1-10
	Photolytic or biological processes in the ocean	<1
	Input from the Stratosphere	≈ 0.5
Total		25-99
Sinks		
	Precipitation	12-42
	Dry Deposition	12-22
Total		24-64

1264


1265    **Appendix 1 (Data runs to 2020 as illustrative)**

1266

### Documents per year by source

Compare the document counts for up to 10 sources. Compare sources and view CiteScore, SJR, and SNIP data

Scopus




1267

1268

### Documents by country or territory

Compare the document counts for up to 15 countries/territories.

Scopus



1269

ABLE	Amazon Boundary-Layer Experiment
ACE	Advanced Composition Explorer
AGAGE	Advanced Global Atmospheric Gases Experiment
AMS	Aerosol Mass Spectrometer
AOD	Aerosol Optical Depth
BBOA	Biomass-Burning Organic Aerosol
CAPRAM	Chemical Aqueous Phase Radical Mechanism
CCN	Cloud Condensation Nuclei
CFC	Chlorofluorocarbon
CLAW	Charlson-Lovelock-Andreae-Warren
CLRTAP	Convention on Long-range Transboundary Air Pollution
CRISTA	Cryogenic Infrared Spectrometers & Telescopes for the Atmosphere
CTM	Chemical Transport Model
EDGAR	Emission Database for Global Atmospheric Research
EPA	United States Environmental Protection Agency
ESRL	Earth System Research Laboratory
FA-AMS	Factor Analysis of Aerosol Mass Spectrometry
FLEXPART	FLEXible PARTicle dispersion model
GC	Gas Chromatography
GC-MS	Gas Chromatography-Mass Spectrometry
GEOS-Chem	Goddard Earth Observing System - Chemical Transport Model
GML	Global Monitoring Laboratory
GOME	Global Ozone Monitoring Experiment
HCFC	Hydrochlorofluorocarbon
HOA	Hydrocarbon-like Organic Aerosol
HYSPLIT	Hybrid Single-Particle Lagrangian Integrated Trajectory model
IAGOS	In-service Aircraft for a Global Observing System
IGAC	International Global Atmospheric Chemistry
IGBP	International Geosphere-Biosphere Programme
IPCC	Intergovernmental Panel on Climate Change
IUPAC	International Union of Pure and Applied Chemistry
JPL	NASA Jet Propulsion Laboratory
LIF	Laser Induced Fluorescence
LV-OOA	Low-volatility Oxygenated Organic Aerosol
MCM	Master Chemical Mechanism
MEGAN	Model of Emissions of Gases and Aerosols from Nature
MODIS	Moderate Resolution Imaging Spectroradiometer
MOPITT	Measurements Of Pollution In The Troposphere

MOZAIC	Measurements of OZone and water vapour by in-service Airbus airCraft
NASA	National Aeronautics and Space Administration
NEC	National Emissions reduction Commitments
NMHC	Non-Methane Hydrocarbon
NOAA	National Oceanic and Atmospheric Administration
OA	Organic Aerosol
OMI	Ozone Monitoring Instrument
OOA	Oxygenated Organic Aerosol
OPE	Ozone Production Efficiency
PAN	Peroxyacetyl Nitrate
PM	Particulate Matter
POA	Primary Organic Aerosol
PSC	Polar Stratospheric Clouds
RACM	Regional Atmospheric Chemistry Mechanism
RADM	Regional Acid Deposition Model
RF	Radiative Forcing
RRKM	Rice-Ramsperger-Kassel-Marcus
SAPRC	Statewide Air Pollution Research Center
SCIAMACHY	SCanning Imaging Absorption spectroMeter for Atmospheric CartographY
SeaWIFS	Sea-Viewing Wide Field-of-View Sensor
SHADOZ	Southern Hemisphere ADDitional OZonesondes
SOA	Secondary Organic Aerosol
STE	Stratospheric-Tropospheric Exchange
SV-OOA	Semi-volatile Oxygenated Organic Aerosol
TES	Technology Experiment Satellite
TOMS	Total Ozone Mapping Spectrometer
UNECE	United Nations Economic Commission for Europe
UV	Ultraviolet
VOC	Volatile Organic Compound
WCRP	World Climate Research Program
WRF	Weather Research and Forecasting model
WRF-Chem	Weather Research and Forecasting model coupled to Chemistry

1271



## 1272    **References**

- 1273    Abbatt, J., George, C., Melamed, M., Monks, P., Pandis, S., and Rudich, Y.: New Directions:  
1274    Fundamentals of atmospheric chemistry: Keeping a three-legged stool balanced, *Atmospheric*  
1275    *Environment*, 84, 390-391, 10.1016/j.atmosenv.2013.10.025, 2014.
- 1276    Aitken, J.: On the Number of Dust Particles in the Atmosphere, *Nature*, 37, 428-430,  
1277    10.1038/037428a0, 1888.
- 1278    Akimoto, H., Takagi, H., and Sakamaki, F.: Photoenhancement of the nitrous acid formation in the  
1279    surface reaction of nitrogen dioxide and water vapor: Extra radical source in smog chamber  
1280    experiments, *International Journal of Chemical Kinetics*, 19, 539-551, 10.1002/kin.550190606, 1987.
- 1281    Alicke, B., Hebestreit, K., Stutz, J., and Platt, U.: Iodine oxide in the marine boundary layer, *Nature*,  
1282    397, 572 - 573, 1999.
- 1283    Allan, B. J., Carslaw, N., Coe, H., Burgess, R. A., and Plane, J. M. C.: Observations of the nitrate radical  
1284    in the marine boundary layer, *Journal of Atmospheric Chemistry*, 33, 129-154,  
1285    10.1023/A:1005917203307, 1999.
- 1286    Anderson, J. G., Toohey, D. W., and Brune, W. H.: Free Radicals Within the Antarctic Vortex: The Role  
1287    of CFCs in Antarctic Ozone Loss, *Science*, 251, 39-46, 10.1126/science.251.4989.39, 1991.
- 1288    Andreae, M. O., and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global*  
1289    *Biogeochemical Cycles*, 15, 955-966, 10.1029/2000gb001382, 2001.
- 1290    Aneja, V. P., Overton, J. H., Cupitt, L. T., Durham, J. L., and Wilson, W. E.: Carbon disulphide and  
1291    carbonyl sulphide from biogenic sources and their contributions to the global sulphur cycle, *Nature*,  
1292    282, 493-496, 10.1038/282493a0, 1979.
- 1293    Aneja, V. P., Schlesinger, W. H., and Erisman, J. W.: Effects of agriculture upon the air quality and  
1294    climate: Research, policy, and regulations, *Environmental Science and Technology*, 43, 4234-4240,  
1295    10.1021/es8024403, 2009.
- 1296    Arrhenius, S.: On the Influence of Carbonic Acid in the Air upon the Temperature of the Ground  
1297    *Philosophical Magazine and Journal of Science*, 41, 237-276, 1896.
- 1298    Atkinson, R.: Kinetics and Mechanisms of the Gas-Phase Reactions of the Hydroxyl Radical with  
1299    Organic Compounds under Atmospheric Conditions, *Chemical Reviews*, 86, 69-201,  
1300    10.1021/cr00071a004, 1986.
- 1301    Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, R. F., Jr., Kerr Chairman, J. A., and Troe, J.: Evaluated  
1302    Kinetic and Photochemical Data for Atmospheric Chemistry: Supplement III. IUPAC Subcommittee on  
1303    Gas Kinetic Data Evaluation for Atmospheric Chemistry, *Journal of Physical and Chemical Reference*  
1304    *Data*, 18, 881-1097, 10.1063/1.555832, 1989.
- 1305    Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, R. F., Jr., Kerr, J. A., and Troe, J.: Evaluated Kinetic  
1306    and Photochemical Data for Atmospheric Chemistry: Supplement IV. IUPAC Subcommittee on Gas  
1307    Kinetic Data Evaluation for Atmospheric Chemistry, *Journal of Physical and Chemical Reference Data*,  
1308    21, 1125-1568, 10.1063/1.555918, 1992.
- 1309    Atkinson, R.: Atmospheric chemistry of VOCs and NO(x), *Atmospheric Environment*, 34, 2063-2101,  
1310    10.1016/S1352-2310(99)00460-4, 2000.
- 1311    Atkinson, R., and Arey, J.: Atmospheric Degradation of Volatile Organic Compounds, *Chemical*  
1312    *Reviews*, 103, 4605-4638, 10.1021/cr0206420, 2003.
- 1313    Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Haynes, R. G., Jenkin, M. E.,  
1314    Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry:  
1315    Volume I - Gas phase reactions of Ox, HOx, NOx and SOx species, *Atmospheric Chemistry and*  
1316    *Physics*, 4, 1461-1738, 2004.
- 1317    Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi,  
1318    M. J., Troe, J., and Subcommittee, I.: Evaluated kinetic and photochemical data for atmospheric  
1319    chemistry: Volume II &ndash; gas phase reactions of organic species, *Atmos. Chem. Phys.*, 6, 3625-  
1320    4055, 10.5194/acp-6-3625-2006, 2006.

1321 Ayers, G. P., Penkett, S. A., Gillett, R. W., Bandy, B., Galbally, I. E., Meyer, C. P., Elsworth, C. M.,  
 1322 Bentley, S. T., and Forgan, B. W.: Evidence for photochemical control of ozone concentrations in  
 1323 unpolluted marine air, *Nature*, 360, 446-449, 10.1038/360446a0, 1992.  
 1324 Ball, S. M., Hancock, G., Murphy, I. J., and Rayner, S. P.: The relative quantum yields of  $O_2(a^1\Delta_g)$  from  
 1325 the photolysis of ozone in the wavelength range  $270\text{nm} \leq \lambda \leq 329\text{nm}$ , *Geophysical Research Letters*,  
 1326 20, 2063-2066, 10.1029/93gl02494, 1993.  
 1327 Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J., and Rasmussen, R. A.: Ozone destruction  
 1328 and photochemical reactions at polar sunrise in the lower Arctic atmosphere, *Nature*, 334, 138-140,  
 1329 1988.  
 1330 Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L.  
 1331 J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology:  
 1332 Model description and evaluation, *Journal of Geophysical Research Atmospheres*, 106, 23073-23095,  
 1333 10.1029/2001JD000807, 2001.  
 1334 Bishop, G. A., and Stedman, D. H.: Measuring the Emissions of Passing Cars, *Accounts of Chemical*  
 1335 *Research*, 29, 489-495, 10.1021/ar950240x, 1996.  
 1336 Blake, D. R., and Rowland, F. S.: World-wide increase in tropospheric methane, 1978–1983, *Journal*  
 1337 *of Atmospheric Chemistry*, 4, 43-62, 10.1007/BF00053772, 1986.  
 1338 Bolin, B., and Charlson, R. J.: On the role of the tropospheric sulfur cycle in the shortwave radiative  
 1339 climate of the earth, *Ambio*, 5, 47-54, 1976.  
 1340 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Bernsten, T., DeAngelo, B. J., Flanner, M. G.,  
 1341 Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G.,  
 1342 Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K.,  
 1343 Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T.,  
 1344 Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific  
 1345 assessment, *Journal of Geophysical Research: Atmospheres*, 118, 5380–5552, 10.1002/jgrd.50171,  
 1346 2013.  
 1347 Boucher, O., and Lohmann, U.: The sulfate-CCN-cloud albedo effect, *Tellus B: Chemical and Physical*  
 1348 *Meteorology*, 47, 281-300, 10.3402/tellusb.v47i3.16048, 1995.  
 1349 Boutron, C. F., Görlach, U., Candelone, J.-P., Bolshov, M. A., and Delmas, R. J.: Decrease in  
 1350 anthropogenic lead, cadmium and zinc in Greenland snows since the late 1960s, *Nature*, 353, 153-  
 1351 156, 10.1038/353153a0, 1991.  
 1352 Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noel, S., Rozanov, V. V., Chance, K. V., and  
 1353 Goede, A. P. H.: SCIAMACHY: Mission objectives and measurement modes, *Journal of the*  
 1354 *Atmospheric Sciences*, 56, 127-150, 1999.  
 1355 Brasseur, G.: *The Ozone Layer - from Discovery to Recovery*, American Meteorological Society,  
 1356 Boston, USA, 2019.  
 1357 Brasseur, G. P., Orlando, J. J., and Tyndall, G. S.: *Atmospheric Chemistry and Global Change*, OUP,  
 1358 1999.  
 1359 Brasseur, G. P., Prinn, R. G., and Pszenny, A. A. P.: *Atmospheric Chemistry in a Changing World*,  
 1360 *Global Change - IGBP Series*, Springer, 2003.  
 1361 Brasseur, G. P.: The Importance of Fundamental Science for Society: The Success Story of Ozone  
 1362 Research, *Perspectives of Earth and Space Scientists*, 1, e2020CN000136,  
 1363 <https://doi.org/10.1029/2020CN000136>, 2020.  
 1364 Brimblecombe, P.: *The Big Smoke: The History of Air Pollution in London since Medieval Times*  
 1365 *Methuen & Co., New York*, 1987.  
 1366 Brown, S. S., Stark, H., Ryerson, T. B., Williams, E. J., Nicks Jr, D. K., Trainer, M., Fehsenfeld, F. C., and  
 1367 Ravishankara, A. R.: Nitrogen oxides in the nocturnal boundary layer: Simultaneous in situ  
 1368 measurements of  $NO_3$ ,  $N_2O_5$ ,  $NO_2$ ,  $NO$ , and  $O_3$ , *J. Geophys. Res.*, 108, 4299,  
 1369 doi:10.1029/2002JD002917, 2003.

1370 Brown, S. S., Ryerson, T. B., Wollny, A. G. C., Brock, A., Peltier, R., Sullivan, A. P., Weber, R. J., Dube,  
 1371 W. P., Trainer, M., Meagher, J. F., Fehsenfeld, F. C., and Ravishankara, A. R.: Variability in Nocturnal  
 1372 Nitrogen Oxide Processing and Its Role in Regional Air Quality, *Science*, 311, 67-70, 2006.  
 1373 Burkholder, J. B., Abbatt, J. P. D., Barnes, I., Roberts, J. M., Melamed, M. L., Ammann, M., Bertram, A.  
 1374 K., Cappa, C. D., Carlton, A. G., Carpenter, L. J., Crowley, J. N., Dubowski, Y., George, C., Heard, D. E.,  
 1375 Herrmann, H., Keutsch, F. N., Kroll, J. H., McNeill, V. F., Ng, N. L., Nizkorodov, S. A., Orlando, J. J.,  
 1376 Percival, C. J., Picquet-Varrault, B., Rudich, Y., Seakins, P. W., Surratt, J. D., Tanimoto, H., Thornton, J.  
 1377 A., Tong, Z., Tyndall, G. S., Wahner, A., Weschler, C. J., Wilson, K. R., and Ziemann, P. J.: The Essential  
 1378 Role for Laboratory Studies in Atmospheric Chemistry, *Environmental Science & Technology*, 51,  
 1379 2519-2528, 10.1021/acs.est.6b04947, 2017.  
 1380 Burkholder, J. B., Abbatt, J. P. D., Cappa, C., Dibble, T. S., Colb, C. E., Orkin, C. L., Wilmouth, D. M.,  
 1381 Sander, S. P., Barker, J. R., Crounse, J. D., Huie, R. E., Kurylo, M. J., Percival, C. J., and Wine, P. H.:  
 1382 Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, JPL, JPL, 2020.  
 1383 Burrows, J. P., Weber, M., Buchwitz, M., Rozanov, V., Ladstatter-Weissenmayer, A., Richter, A.,  
 1384 DeBeek, R., Hoogen, R., Bramstedt, K., Eichmann, K. U., and Eisinger, M.: The global ozone  
 1385 monitoring experiment (GOME): Mission concept and first scientific results, *Journal of the*  
 1386 *Atmospheric Sciences*, 56, 151-175, 1999.  
 1387 Burrows, J. P., Platt, U., and Borrell, P.: The Remote Sensing of Tropospheric Composition from  
 1388 Space, *Physics of Earth and Space Environments*, Springer-Verlag Berlin, Heidelberg, 551 pp., 2011.  
 1389 Callendar, G. S.: The artificial production of carbon dioxide and its influence on temperature,  
 1390 *Quarterly Journal of the Royal Meteorological Society*, 64, 223-240, 10.1002/qj.49706427503, 1938.  
 1391 Calvert, J. G., Su, F., Bottenheim, J. W., and Strausz, O. P.: Mechanism of the homogeneous oxidation  
 1392 of sulfur dioxide in the troposphere, *Atmospheric Environment* (1967), 12, 197-226, 10.1016/0004-  
 1393 6981(78)90201-9, 1978.  
 1394 Calvert, J. G., Heywood, J. B., Sawyer, R. F., and Seinfeld, J. H.: Achieving acceptable air quality: Some  
 1395 reflections on controlling vehicle emissions, *Science*, 261, 37-45, 10.1126/science.261.5117.37, 1993.  
 1396 Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B.,  
 1397 Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J.,  
 1398 DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and microphysical  
 1399 characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, *Mass*  
 1400 *Spectrometry Reviews*, 26, 185-222, 10.1002/mas.20115, 2007.  
 1401 Carslaw, D. C.: Evidence of an increasing NO<sub>2</sub>/NO<sub>x</sub> emissions ratio from road traffic emissions,  
 1402 *Atmospheric Environment*, 39, 4793-4802, 10.1016/j.atmosenv.2005.06.023, 2005.  
 1403 Carter, W. P. L., and Atkinson, R.: A Computer Modeling Study of Incremental Hydrocarbon  
 1404 Reactivity *Environmental Science & Technology*, 23, 864-880, 1989.  
 1405 Carter, W. P. L.: A detailed mechanism for the gas-phase atmospheric reactions of organic  
 1406 compounds, *Atmospheric Environment. Part A. General Topics*, 24, 481-518,  
 1407 [https://doi.org/10.1016/0960-1686\(90\)90005-8](https://doi.org/10.1016/0960-1686(90)90005-8), 1990.  
 1408 Carter, W. P. L.: Development of the SAPRC-07 chemical mechanism, *Atmospheric Environment*, 44,  
 1409 5324-5335, <https://doi.org/10.1016/j.atmosenv.2010.01.026>, 2010.  
 1410 Chamberlain, A. C.: Transport of gases to and from grass and grass-like surfaces, *Proc. R. Soc. Lond.*  
 1411 *A*, 290, 236-265, 1966.  
 1412 Chameides, W. L., and Walker, J. C. G.: A photochemical theory for tropospheric ozone, *J. Geophys.*  
 1413 *Res.*, 78, 8751-8760, 1973.  
 1414 Chameides, W. L., and Davis, D. D.: Iodine: Its possible role in tropospheric photochemistry, *Journal*  
 1415 *of Geophysical Research: Oceans*, 85, 7383-7398, <https://doi.org/10.1029/JC085iC12p07383>, 1980.  
 1416 Chameides, W. L., and Davis, D. D.: The free radical chemistry of cloud droplets and its impact upon  
 1417 the composition of rain, *Journal of Geophysical Research*, 87, 4863-4877, 10.1029/JC087iC07p04863,  
 1418 1982.

1419 Chameides, W. L., Lindsay, R. W., Richardson, J., and Kiang, C. S.: The role of biogenic hydrocarbons  
 1420 in urban photochemical smog: Atlanta as a case study, *Science*, 241, 1473-1475,  
 1421 10.1126/science.3420404, 1988.

1422 Chameides, W. L., Yu, H., Liu, S. C., Bergin, M., Zhou, X., Mearns, L., Wang, G., Kiang, C. S., Saylor, R.  
 1423 D., Luo, C., Huang, Y., Steiner, A., and Giorgi, F.: Case study of the effects of atmospheric aerosols  
 1424 and regional haze on agriculture: An opportunity to enhance crop yields in China through emission  
 1425 controls?, *Proceedings of the National Academy of Sciences of the United States of America*, 96,  
 1426 13626-13633, 10.1073/pnas.96.24.13626, 1999.

1427 Chaney, L. W.: The remote measurement of traffic generated carbon monoxide, *Journal of the Air*  
 1428 *Pollution Control Association*, 33, 220-222, 10.1080/00022470.1983.10465568, 1983.

1429 Chapman, S.: On ozone and atomic oxygen in the upper atmosphere, *The London, Edinburgh, and*  
 1430 *Dublin Philosophical Magazine and Journal of Science*, 10, 369-383, 10.1080/14786443009461588,  
 1431 1930.

1432 Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton,  
 1433 atmospheric sulphur, cloud albedo and climate, *Nature*, 326, 655-661, 10.1038/326655a0, 1987.

1434 Charlson, R. J., Langner, J., and Rodhe, H.: Sulphate aerosol and climate, *Nature*, 348, 22,  
 1435 10.1038/348022a0, 1990.

1436 Charlson, R. J., Langner, J., Rodhe, H., Leovy, C. B., and Warren, S. G.: Perturbation of the Northern  
 1437 Hemisphere radiative balance by backscattering from anthropogenic sulfate aerosols, *Tellus, Series*  
 1438 *A-B*, 43 A-B, 152-163, 1991.

1439 Chipperfield, M. P., and Pyle, J. A.: Two-dimensional modelling of the Antarctic lower stratosphere,  
 1440 *Geophysical Research Letters*, 15, 875-878, 10.1029/GL015i008p00875, 1988.

1441 Cicerone, R. J.: Halogens in the atmosphere, *Reviews of Geophysics*, 19, 123-139,  
 1442 <https://doi.org/10.1029/RG019i001p00123>, 1981.

1443 Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P.,  
 1444 Andreae, M. O., Artaxo, P., and Maenhaut, W.: Formation of Secondary Organic Aerosols Through  
 1445 Photooxidation of Isoprene, *Science*, 303, 1173-1176, 10.1126/science.1092805, 2004.

1446 Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system  $\text{H}^+\text{-NH}_4^+\text{-SO}_4^{2-}\text{-}$   
 1447  $\text{NO}_3^-\text{-H}_2\text{O}$  at tropospheric temperatures, *Journal of Physical Chemistry A*, 102, 2137-2154, 1998a.

1448 Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system  $\text{H}^+\text{-NH}_4^+\text{-Na}^+\text{-}$   
 1449  $\text{SO}_4^{2-}\text{-NO}_3^-\text{-Cl}^-\text{-H}_2\text{O}$  at 298.15 K, *Journal of Physical Chemistry A*, 102, 2155-2171, 1998b.

1450 Cohen, R. C., and Murphy, J. G.: Photochemistry of  $\text{NO}_2$  in Earth's Stratosphere: Constraints from  
 1451 Observations, *Chemical Reviews*, 103, 4985-4998, 10.1021/cr020647x, 2003.

1452 Covert, D. S., Wiedensohler, A., Aalto, P., Heintzenberg, J., McMurry, P. H., and Leck, C.: Aerosol  
 1453 number size distributions from 3 to 500 nm diameter in the arctic marine boundary layer during  
 1454 summer and autumn, *Tellus, Series B: Chemical and Physical Meteorology*, 48, 197-212,  
 1455 10.3402/tellusb.v48i2.15886, 1996.

1456 Cox, R. A., and Hayman, G. D.: The stability and photochemistry of dimers of the ClO radical and  
 1457 implications for Antarctic ozone depletion, *Nature*, 332, 796-800, 10.1038/332796a0, 1988.

1458 Crounse, J. D., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., and Wennberg, P. O.: Autoxidation of  
 1459 organic compounds in the atmosphere, *Journal of Physical Chemistry Letters*, 4, 3513-3520,  
 1460 10.1021/jz4019207, 2013.

1461 Crowley, J. N., Ammann, M., Cox, R. A., Hynes, R. G., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J.,  
 1462 and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume v  
 1463 -heterogeneous reactions on solid substrates, *Atmospheric Chemistry and Physics*, 10, 9059-9223,  
 1464 10.5194/acp-10-9059-2010, 2010.

1465 Crutzen, P.: A discussion of the chemistry of some minor constituents in the stratosphere and  
 1466 troposphere, *Pure and Applied Geophysics*, 106, 1385-1399, 10.1007/BF00881092, 1973a.

1467 Crutzen, P. J.: The influence of nitrogen oxides on the atmospheric ozone content, *Quarterly Journal*  
 1468 *of the Royal Meteorological Society*, 96, 320-325, 10.1002/qj.49709640815, 1970.

1469 Crutzen, P. J.: Photochemical reactions initiated by and influencing ozone in the unpolluted  
1470 troposphere, *Tellus*, 26, 47-57, 1973b.

1471 Crutzen, P. J., Heidt, L. E., Krasnec, J. P., Pollock, W. H., and Seiler, W.: Biomass burning as a source of  
1472 atmospheric gases CO, H<sub>2</sub>, N<sub>2</sub>O, NO, CH<sub>3</sub>Cl and COS, *Nature*, 282, 253-256, 10.1038/282253a0, 1979.

1473 Crutzen, P. J., and Birks, J. W.: The atmosphere after a nuclear war: twilight at noon, *Ambio*, 11, 114-  
1474 125, 1982.

1475 Crutzen, P. J., and Andreae, M. O.: Biomass burning in the tropics: Impact on atmospheric chemistry  
1476 and biogeochemical cycles, *Science*, 250, 1669-1678, 1990.

1477 Crutzen, P. J.: Geology of mankind, *Nature*, 415, 23, 10.1038/415023a, 2002.

1478 Dalton, J.: Experimental enquiry into the proportion of the several gases or elastic fluids, constituting  
1479 the atmosphere., *Memoirs of the Literary and Philosophical Society of Manchester*, 1, 244-258,  
1480 1805.

1481 Danielsen, E. F.: Stratospheric-Tropospheric Exchange Based on Radioactivity, Ozone and Potential  
1482 Vorticity, *Journal of the Atmospheric Sciences*, 25, 502-518, 10.1175/1520-  
1483 0469(1968)025<0502:STEBOR>2.0.CO;2, 1968.

1484 Darnall, K. R., Lloyd, A. C., Winer, A. M., and Pitts, J. N.: Reactivity scale for atmospheric  
1485 hydrocarbons based on reaction with hydroxyl radical, *Environmental Science & Technology*, 10,  
1486 692-696, 10.1021/es60118a008, 1976.

1487 Davis, D., Nowak, J. B., Chen, G., Buhr, M., Arimoto, R., Hogan, A., Eisele, F., Mauldin, L., Tanner, D.,  
1488 Shetter, R., Lefer, B., and McMurry, P.: Unexpected high levels of NO observed at South Pole,  
1489 *Geophysical Research Letters*, 28, 3625-3628, 10.1029/2000GL012584, 2001.

1490 Davis, D. D., Heaps, W., and McGee, T.: Direct measurements of natural tropospheric levels of OH via  
1491 an aircraft borne tunable dye laser, *Geophysical Research Letters*, 3, 331-333,  
1492 <https://doi.org/10.1029/GL003i006p00331>, 1976.

1493 Davis, D. D., Ravishankara, A. R., and Fischer, S.: SO<sub>2</sub> oxidation via the hydroxyl radical: Atmospheric  
1494 fate of HSO<sub>x</sub> radicals, *Geophysical Research Letters*, 6, 113-116, 10.1029/GL006i002p00113, 1979.

1495 Davis, D. D.: Project Gametag: An overview, *Journal of Geophysical Research: Oceans*, 85, 7285-  
1496 7292, <https://doi.org/10.1029/JC085iC12p07285>, 1980.

1497 de Gouw, J. A., Middlebrook, A. M., Warneke, C., Goldan, P. D., Kuster, W. C., Roberts, J. M.,  
1498 Fehsenfeld, F. C., Worsnop, D. R., Canagaratna, M. R., Pszenny, A. A. P., Keene, W. C., Marchewka,  
1499 M., Bertman, S. B., and Bates, T. S.: Budget of organic carbon in a polluted atmosphere: Results from  
1500 the New England Air Quality Study in 2002, *Journal of Geophysical Research D: Atmospheres*, 110, 1-  
1501 22, 10.1029/2004JD005623, 2005.

1502 De Mazière, M., Thompson, A. M., Kurylo, M. J., Wild, J. D., Bernhard, G., Blumenstock, T., Braathen,  
1503 G. O., Hannigan, J. W., Lambert, J. C., Leblanc, T., McGee, T. J., Nedoluha, G., Petropavlovskikh, I.,  
1504 Seckmeyer, G., Simon, P. C., Steinbrecht, W., and Strahan, S. E.: The Network for the Detection of  
1505 Atmospheric Composition Change (NDACC): history, status and perspectives, *Atmos. Chem. Phys.*,  
1506 18, 4935-4964, 10.5194/acp-18-4935-2018, 2018.

1507 De Zafra, R. L., Jaramillo, M., Parrish, A., Solomon, P., Connor, B., and Barrett, J.: High concentrations  
1508 of chlorine monoxide at low altitudes in the Antarctic spring stratosphere: Diurnal variation, *Nature*,  
1509 328, 408-411, 1988.

1510 Demerjian, K. L., Kerr, J. A., and Calvert, J. G.: MECHANISM OF PHOTOCHEMICAL SMOG FORMATION,  
1511 *Adv in Environ Sci and Technol*, 4, 1-262, 1974.

1512 DeMore, W. B., Sander, S. P., Golden, D. M., Hampson, R. F., Kurylo, M. J., Howard, C. J.,  
1513 Ravishankara, A. R., Kolb, C. E., and Molina, M. J.: Chemical Kinetics and Photochemical Data for Use  
1514 in Stratospheric Modeling, NASA, JPL, 1997.

1515 Dentener, F. J., Carmichael, G. R., Zhang, Y., Lelieveld, J., and Crutzen, P. J.: Role of mineral aerosol as  
1516 a reactive surface in the global troposphere, *Journal of Geophysical Research: Atmospheres*, 101,  
1517 22869-22889, 10.1029/96JD01818, 1996.

1518 Di Carlo, P., Brune, W. H., Martinez, M., Harder, H., Leshner, R., Ren, X. R., Thornberry, T., Carroll, M.  
 1519 A., Young, V., Shepson, P. B., Riemer, D., Apel, E., and Campbell, C.: Missing OH reactivity in a forest:  
 1520 Evidence for unknown reactive biogenic VOCs, *Science*, 304, 722-725, 2004.  
 1521 Dockery, D. W., Pope, C. A., Xu, X., Spengler, J. D., Ware, J. H., Fay, M. E., Ferris, B. G., and Speizer, F.  
 1522 E.: An association between air pollution and mortality in six US cities. *New England J. Med.*, 29, 1753-  
 1523 1759, 1993.  
 1524 Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled partitioning, dilution, and  
 1525 chemical aging of semivolatile organics, *Environmental Science and Technology*, 40, 2635-2643,  
 1526 10.1021/es052297c, 2006.  
 1527 Draxler, R. R., and Hess, G. D.: An overview of the HYSPLIT\_4 modelling system for trajectories,  
 1528 dispersion and deposition, *Australian Meteorological Magazine*, 47, 295-308, 1998.  
 1529 Ehhalt, D. H.: The atmospheric cycle of methane, *Tellus*, 26, 58-70, 10.1111/j.2153-  
 1530 3490.1974.tb01952.x, 1974.  
 1531 Ehhalt, D. H.: Photooxidation of trace gases in the troposphere, *Physical Chemistry Chemical Physics*,  
 1532 1, 5401-5408, 1999.  
 1533 Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F.,  
 1534 Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I. H., Rissanen, M., Jokinen, T.,  
 1535 Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, L. B., Jørgensen,  
 1536 S., Kjaergaard, H. G., Canagaratna, M., Maso, M. D., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.  
 1537 M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility  
 1538 secondary organic aerosol, *Nature*, 506, 476-479, 10.1038/nature13032, 2014.  
 1539 Eisele, F. L.: Intercomparison of tropospheric OH and ancillary trace gas measurements at Fritz Peak  
 1540 Observatory, Colorado, *Journal of Geophysical Research*, 99, 18,605-618,626, 1994.  
 1541 Erisman, J. W., Sutton, M. A., Galloway, J., Klimont, Z., and Winiwarter, W.: How a century of  
 1542 ammonia synthesis changed the world, *Nature Geoscience*, 1, 636-639, 10.1038/ngeo325, 2008.  
 1543 Ervens, B., George, C., Williams, J. E., Buxton, G. V., Salmon, G. A., Bydder, M., Wilkinson, F.,  
 1544 Dentener, F., Mirabel, P., Wolke, R., and Herrmann, H.: CAPRAM 2.4 (MODAC mechanism): An  
 1545 extended and condensed tropospheric aqueous phase mechanism and its application, *Journal of*  
 1546 *Geophysical Research: Atmospheres*, 108, 10.1029/2002JD002202, 2003.  
 1547 Fabian, P., and Pruchniewicz, P. G.: Meridional distribution of ozone in the troposphere and its  
 1548 seasonal variations, *J. Geophys. Res.*, 82, 2063-2073, 1977.  
 1549 Facchini, M. C., Mircea, M., Fuzzi, S., and Charlson, R. J.: Cloud albedo enhancement by surface-  
 1550 active organic solutes in growing droplets, *Nature*, 401, 257-259, 10.1038/45758, 1999.  
 1551 Farman, J. C., Gardiner, B. G., and Shanklin, J. D.: Large losses of total ozone in Antarctica reveal  
 1552 seasonal ClO x/NO<sub>x</sub> interaction, *Nature*, 315, 207-210, 10.1038/315207a0, 1985.  
 1553 Fehsenfeld, F., Calvert, J., Fall, R., Goldan, P., Guenther, A. B., Hewitt, C. N., Lamb, B., Liu, S., Trainer,  
 1554 M., Westberg, H., and Zimmerman, P.: Emissions of volatile organic compounds from vegetation and  
 1555 the implications for atmospheric chemistry, *Global Biogeochemical Cycles*, 6, 389-430,  
 1556 10.1029/92GB02125, 1992.  
 1557 Finlayson-Pitts, B. J., Ezell, M. J., and Pitts, J. N.: Formation of chemically active chlorine compounds  
 1558 by reactions of atmospheric NaCl particles with gaseous N<sub>2</sub>O<sub>5</sub> and ClONO<sub>2</sub>, *Nature*, 337, 241-244,  
 1559 10.1038/337241a0, 1989.  
 1560 Finlayson-Pitts, B. J., and Pitts, J. N.: Chemistry of the upper and lower atmosphere - theory,  
 1561 experiments, and applications, 969 pp, Academic Press, San Diego, 2000.  
 1562 Finlayson, B. J., and Pitts, J. N.: Photochemistry of the Polluted Troposphere, *Science*, 192, 111-119,  
 1563 10.1126/science.192.4235.111, 1976.  
 1564 Fiore, A. M., Dentener, F. J., Wild, O., Cuvelier, C., Schultz, M. G., Hess, P., Textor, C., Schulz, M.,  
 1565 Doherty, R. M., Horowitz, L. W., MacKenzie, I. A., Sanderson, M. G., Shindell, D. T., Stevenson, D. S.,  
 1566 Szopa, S., Van Dingenen, R., Zeng, G., Atherton, C., Bergmann, D., Bey, I., Carmichael, G., Collins, W.  
 1567 J., Duncan, B. N., Faluvegi, G., Folberth, G., Gauss, M., Gong, S., Hauglustaine, D., Holloway, T.,  
 1568 Isaksen, I. S. A., Jacob, D. J., Jonson, J. E., Kaminski, J. W., Keating, T. J., Lupu, A., Marmer, E.,



1569 Montanaro, V., Park, R. J., Pitari, G., Pringle, K. J., Pyle, J. A., Schroeder, S., Vivanco, M. G., Wind, P.,  
 1570 Wojcik, G., Wu, S., and Zuber, A.: Multimodel estimates of intercontinental source-receptor  
 1571 relationships for ozone pollution, *J. Geophys. Res.-Atmos.*, 114, D04301, 10.1029/2008jd010816,  
 1572 2009.  
 1573 Fiore, A. M., Naik, V., Spracklen, D. V., Steiner, A., Unger, N., Prather, M., Bergmann, D., Cameron-  
 1574 Smith, P. J., Cionni, I., Collins, W. J., Dalsoren, S., Eyring, V., Folberth, G. A., Ginoux, P., Horowitz, L.  
 1575 W., Josse, B., Lamarque, J. F., MacKenzie, I. A., Nagashima, T., O'Connor, F. M., Righi, M., Rumbold, S.  
 1576 T., Shindell, D. T., Skeie, R. B., Sudo, K., Szopa, S., Takemura, T., and Zeng, G.: Global air quality and  
 1577 climate, *Chem Soc Rev*, 41, 6663-6683, 10.1039/c2cs35095e, 2012.  
 1578 Fishman, J., Ramanathan, V., Crutzen, P. J., and Liu, S. C.: Tropospheric ozone and climate, *Nature*,  
 1579 282, 818-820, 10.1038/282818a0, 1979.  
 1580 Fitzgerald, J. W.: Effect of Aerosol Composition on Cloud Droplet Size Distribution: A Numerical  
 1581 Study, *Journal of the Atmospheric Sciences*, 31, 1358-1367, 10.1175/1520-  
 1582 0469(1974)031<1358:Eoacoc>2.0.Co;2, 1974.  
 1583 Fleming, E. L., Jackman, C. H., Stolarski, R. S., and Considine, D. B.: Simulation of stratospheric tracers  
 1584 using an improved empirically based two-dimensional model transport formulation, *Journal of*  
 1585 *Geophysical Research: Atmospheres*, 104, 23911-23934, <https://doi.org/10.1029/1999JD900332>,  
 1586 1999.  
 1587 Forster, C., Wandering, U., Wotawa, G., James, P., Mattis, I., Althausen, D., Simmonds, P., O'Doherty,  
 1588 S., Jennings, S. G., Kleefeld, C., Schneider, J., Trickl, T., Kreipl, S., Jäger, H., and Stohl, A.: Transport of  
 1589 boreal forest fire emissions from Canada to Europe, *J. Geophys. Res.*, 106, 22887-22906., 2001.  
 1590 Fowler, D., Pilegaard, K., Sutton, M. A., Ambus, P., Raivonen, M., Duyzer, J., Simpson, D., Fagerli, H.,  
 1591 Fuzzi, S., Schjoerring, J. K., Granier, C., Neftel, A., Isaksen, I. S. A., Laj, P., Maione, M., Monks, P. S.,  
 1592 Burkhardt, J., Daemmgen, U., Neirynck, J., Personne, E., Wichink-Kruit, R., Butterbach-Bahl, K.,  
 1593 Flechard, C., Tuovinen, J. P., Coyle, M., Gerosa, G., Loubet, B., Altimir, N., Gruenhage, L., Ammann,  
 1594 C., Cieslik, S., Paoletti, E., Mikkelsen, T. N., Ro-Poulsen, H., Cellier, P., Cape, J. N., Horvath, L., Loreto,  
 1595 F., Niinemets, U., Palmer, P. I., Rinne, J., Misztal, P., Nemitz, E., Nilsson, D., Pryor, S., Gallagher, M.  
 1596 W., Vesala, T., Skiba, U., Brüeggemann, N., Zechmeister-Boltenstern, S., Williams, J., O'Dowd, C.,  
 1597 Facchini, M. C., de Leeuw, G., Flossman, A., Chaumerliac, N., and Erisman, J. W.: Atmospheric  
 1598 composition change: Ecosystems-Atmosphere interactions, *Atmospheric Environment*, 43, 5193-  
 1599 5267, 10.1016/j.atmosenv.2009.07.068, 2009.  
 1600 Fowler, D., Brimblecombe, P., Burrows, J., Heal, M. R., Grennfelt, P., Stevenson, D. S., Jowett, A.,  
 1601 Nemitz, E., Coyle, M., Lui, X., Chang, Y., Fuller, G. W., Sutton, M. A., Klimont, Z., Unsworth, M. H., and  
 1602 Vieno, M.: A chronology of global air quality, *Philosophical Transactions of the Royal Society A:*  
 1603 *Mathematical, Physical and Engineering Sciences*, 378, 20190314, doi:10.1098/rsta.2019.0314, 2020.  
 1604 Fröhlich-Nowoisky, J., Kampf, C. J., Weber, B., Huffman, J. A., Pöhlker, C., Andreae, M. O., Lang-Yona,  
 1605 N., Burrows, S. M., Gunthe, S. S., Elbert, W., Su, H., Hoor, P., Thines, E., Hoffmann, T., Després, V. R.,  
 1606 and Pöschl, U.: Bioaerosols in the Earth system: Climate, health, and ecosystem interactions,  
 1607 *Atmospheric Research*, 182, 346-376, <https://doi.org/10.1016/j.atmosres.2016.07.018>, 2016.  
 1608 Fuller, G. W.: *The Invisible Killer: The Rising Threat of Global Air Pollution - and how we can fight*  
 1609 *back*, Melville House, 2018.  
 1610 Galbally, I.: Some measurements of ozone variation and destruction in the atmospheric surface  
 1611 layer, *Nature*, 218, 456-457, 10.1038/218456a0, 1968.  
 1612 Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., Asner,  
 1613 G. P., Cleveland, C. C., Green, P. A., Holland, E. A., Karl, D. M., Michaels, A. F., Porter, J. H., Townsend,  
 1614 A. R., and Vorosmarty, C. J.: Nitrogen cycles: past, present, and future, *Biogeochemistry*, 70, 153-  
 1615 226, 10.1007/s10533-004-0370-0, 2004.  
 1616 Garcia, R. R., and Solomon, S.: A numerical model of the zonally averaged dynamical and chemical  
 1617 structure of the middle atmosphere, *Journal of Geophysical Research: Oceans*, 88, 1379-1400,  
 1618 <https://doi.org/10.1029/JC088iC02p01379>, 1983.

1619 Gard, E., Mayer, J. E., Morrical, B. D., Dienes, T., Fergenson, D. P., and Prather, K. A.: Real-Time  
 1620 Analysis of Individual Atmospheric Aerosol Particles: Design and Performance of a Portable ATOFMS,  
 1621 Analytical Chemistry, 69, 4083-4091, 10.1021/ac970540n, 1997.  
 1622 Goldstein, A. H., and Galbally, I. E.: Known and Unexplored Organic Constituents in the Earth's  
 1623 Atmosphere, Environ. Sci. Technol., 1 March, 1515-1521, 2007.  
 1624 Graedel, T. E.: KINETIC PHOTOCHEMISTRY OF THE MARINE ATMOSPHERE., J. Geophys. Res., 84, 273-  
 1625 286, 1979.  
 1626 Graedel, T. E., and Weschler, C. J.: Chemistry within aqueous atmospheric aerosols and raindrops,  
 1627 Reviews of Geophysics, 19, 505-539, 10.1029/RG019i004p00505, 1981.  
 1628 Grannas, A. M., Jones, A. E., Dibb, J. E., Ammann, M., Anastasio, C., Beine, H. J., Bergin, M.,  
 1629 Bottenheim, J., Boxe, C. S., Carver, G., Chen, G., Crawford, J., Dominé, F., Frey, M. M., Guzmán, M. I.,  
 1630 Heard, D. E., Helmig, D., Hoffmann, M. R., Honrath, R. E., Huey, L. G., Hutterli, M. A., Jacobi, H.-W.,  
 1631 Klán, P., Lefer, B., McConnell, J., Plane, J., Sander, R., Savarino, J., Shepson, P. B., Simpson, W. R.,  
 1632 Sodeau, J. R., von Glasow, R., Weller, R., Wolff, E. W., and Zhu, T.: An overview of snow  
 1633 photochemistry: evidence, mechanisms and impacts, Atmos. Chem. Phys., 7, 4329-4373, 2007.  
 1634 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully  
 1635 coupled "online" chemistry within the WRF model, Atmospheric Environment, 39, 6957-6975,  
 1636 10.1016/j.atmosenv.2005.04.027, 2005.  
 1637 Grennfelt, P., Engleryd, A., Forsius, M., Hov, Ø., Rodhe, H., and Cowling, E.: Acid rain and air  
 1638 pollution: 50 years of progress in environmental science and policy, Ambio, 10.1007/s13280-019-  
 1639 01244-4, 2019.  
 1640 Guenther, A., Hewitt, C., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau,  
 1641 M., McKay, W., Pierce, T., Scholes, R., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.:  
 1642 A global model of natural volatile organic compound emissions, J. Geophys. Res., 100, 8873-8892,  
 1643 1995.  
 1644 Guenther, A., Geron, C., Pierce, T., Lamb, B., Harley, P., and Fall, R.: Natural emissions of non-  
 1645 methane volatile organic compounds, carbon monoxide, and oxides of nitrogen from North America,  
 1646 Atmospheric Environment, 34, 2205-2230, [https://doi.org/10.1016/S1352-2310\(99\)00465-3](https://doi.org/10.1016/S1352-2310(99)00465-3), 2000.  
 1647 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global  
 1648 terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from  
 1649 Nature), Atmospheric Chemistry and Physics, 6, 3181-3210, 10.5194/acp-6-3181-2006, 2006.  
 1650 Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and  
 1651 monoterpene emission rate variability: Model evaluations and sensitivity analyses, Journal of  
 1652 Geophysical Research: Atmospheres, 98, 12609-12617, 10.1029/93jd00527, 1993.  
 1653 Haagen-Smit, A. J.: Chemistry and Physiology of Los Angeles Smog, Industrial & Engineering  
 1654 Chemistry, 44, 1342-1346, 10.1021/ie50510a045, 1952.  
 1655 Haagen-Smit, A. J., Bradley, C. E., and Fox, M. M.: Ozone Formation in Photochemical Oxidation of  
 1656 Organic Substances, Industrial & Engineering Chemistry, 45, 2086-2089, 10.1021/ie50525a044, 1953.  
 1657 Haagen-Smit, A. J., and Fox, M. M.: Photochemical ozone formation with hydrocarbons and  
 1658 automobile exhaust, Air Repair, 4, 105-136, 10.1080/00966665.1954.10467649, 1954.  
 1659 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J.,  
 1660 Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y.,  
 1661 Jang, M., Jenkin, M., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T.,  
 1662 Monod, A., Prevot, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The Formation,  
 1663 Properties and Impact of Secondary Organic Aerosol: Current and Emerging Issues, Atmos. Chem.  
 1664 Phys., 9, 5155-5236, 10.5194/acp-9-5155-2009 2009.  
 1665 Hampson, J.: Photochemical behavior of the ozone layer, Can. Armament Res. and Dev. Estab.,  
 1666 Valcartier, Quebec, Canada, 1964.  
 1667 Hanson, D. R., Burkholder, J. B., Howard, C. J., and Ravishankara, A. R.: Measurement of OH and HO<sub>2</sub>  
 1668 radical uptake coefficients on water and sulfuric acid surfaces, Journal of Physical Chemistry, 96,  
 1669 4979-4985, 10.1021/j100191a046, 1992.



1670 Hanson, D. R., and Ravishankara, A. R.: Reactive uptake of ClONO<sub>2</sub> onto sulfuric acid due to reaction  
 1671 with HCl and H<sub>2</sub>O, *Journal of Physical Chemistry*, 98, 5728-5735, 10.1021/j100073a026, 1994.  
 1672 Hao, W. M., and Liu, M. H.: Spatial and temporal distribution of tropical biomass burning, *Global*  
 1673 *Biogeochemical Cycles*, 8, 495-504, 1994.  
 1674 Harriss, R. C.: The Amazon Boundary Layer Experiment ( ABLE 2A): dry season 1985, *Journal of*  
 1675 *Geophysical Research*, 93, 1351-1360, 10.1029/JD093iD02p01351, 1988.  
 1676 Heard, D. E., and Pilling, M. J.: Measurement of OH and HO<sub>2</sub> in the Troposphere, *Chemical Reviews*,  
 1677 103, 5163-5198, 2003.  
 1678 Heard, D. E.: *Analytical Techniques for Atmospheric Measurement*, Wiley-Blackwell, 2006.  
 1679 Hein, R., Crutzen, P. J., and Heimann, M.: An inverse modeling approach to investigate the global  
 1680 atmospheric methane cycle, *Global Biogeochemical Cycles*, 11, 43-76, 10.1029/96GB03043, 1997.  
 1681 Hofzumahaus, A., Rohrer, F., Lu, K. D., Bohn, B., Brauers, T., Chang, C. C., Fuchs, H., Holland, F., Kita,  
 1682 K., Kondo, Y., Li, X., Lou, S. R., Shao, M., Zeng, L. M., Wahner, A., and Zhang, Y. H.: Amplified Trace  
 1683 Gas Removal in the Troposphere, *Science*, 324, 1702-1704, 10.1126/science.1164566, 2009.  
 1684 Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B., and Pfister, L.: Stratosphere-  
 1685 troposphere exchange, *Reviews of Geophysics*, 33, 403-439, 10.1029/95rg02097, 1995.  
 1686 Honrath, R. E., Peterson, M. C., Guo, S., Dibb, J. E., Shepson, P. B., and Campbell, B.: Evidence of NO<sub>x</sub>  
 1687 production within or upon ice particles in the Greenland snow-pack, *Geophys. Res. Lett.*, 26, 695-  
 1688 698, 1999.  
 1689 Howard, C. J., and Evenson, K. M.: Kinetics of the reaction of HO<sub>2</sub> with NO, *Geophysical Research*  
 1690 *Letters*, 4, 437-440, 10.1029/GL004i010p00437, 1977.  
 1691 Hudson, R. D., and Reed, E. I.: STRATOSPHERE: PRESENT AND FUTURE, NASA Reference Publication,  
 1692 1979.  
 1693 Husar, R. B., Tratt, D. M., Schichtel, B. A., Falke, S. R., Li, F., Jaffe, D., Gassó, S., Gill, T., Laulainen, N.  
 1694 S., Lu, F., Reheis, M. C., Chun, Y., Westphal, D., Holben, B. N., Gueymard, C., McKendry, I., Kuring, N.,  
 1695 Feldman, G. C., McClain, C., Frouin, R. J., Merrill, J., DuBois, D., Vignola, F., Murayama, T., Nickovic,  
 1696 S., Wilson, W. E., Sassen, K., Sugimoto, N., and Malm, W. C.: Asian dust events of April 1998, *Journal*  
 1697 *of Geophysical Research Atmospheres*, 106, 18317-18330, 10.1029/2000JD900788, 2001.  
 1698 IPCC: *Climate Change 2013 – The Physical Science Basis*, Cambridge University Press, Cambridge,  
 1699 1552 pp., 2013.  
 1700 Jackman, C., Seals, R., and Prather, M.: Two-dimensional intercomparison of stratospheric models,  
 1701 1989.  
 1702 Jacob, D. J., Munger, J. W., Waldman, J. M., and Hoffmann, M. R.: The H<sub>2</sub>SO<sub>4</sub> - HNO<sub>3</sub> - NH<sub>3</sub> system at  
 1703 high humidities and in fogs. 1. Spatial and temporal patterns in the San Joaquin Valley of California,  
 1704 *Journal of Geophysical Research*, 91, 1073-1088, 10.1029/JD091iD01p01073, 1986.  
 1705 Jacob, D. J.: *Introduction to Atmospheric Chemistry*, Princeton University Press, Princeton, USA,  
 1706 1999.  
 1707 Jacob, D. J., Logan, J. A., and Murti, P. P.: Effect of rising Asian emissions on surface ozone in the  
 1708 United States, *Geophysical Research Letters*, 26, 2175-2178, 10.1029/1999GL900450, 1999.  
 1709 Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmospheric Environment*, 34, 2131-  
 1710 2159, 10.1016/S1352-2310(99)00462-8, 2000.  
 1711 Jacobson, M. Z.: *Atmospheric Pollution: History, Science and Regulation*, Cambridge University Press,  
 1712 2002.  
 1713 Jaeglé, L., Jacob, D. J., Wang, Y., Weinheimer, A. J., Ridley, B. A., Campos, T. L., Sachse, G. W., and  
 1714 Hagen, D. E.: Sources and chemistry of NO<sub>x</sub> in the upper troposphere over the United States,  
 1715 *Geophysical Research Letters*, 25, 1705-1708, 10.1029/97gl03591, 1998.  
 1716 Jaenicke, R.: Abundance of cellular material and proteins in the atmosphere, *Science*, 308, 73,  
 1717 10.1126/science.1106335, 2005.  
 1718 Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic  
 1719 compounds: A protocol for mechanism development, *Atmos. Environ.*, 31, 81-104, 1997.

1720 Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the  
 1721 Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile  
 1722 organic compounds, *Atmos. Chem. Phys.*, 3, 181-193, 10.5194/acp-3-181-2003, 2003.  
 1723 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P.  
 1724 F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson,  
 1725 A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A.,  
 1726 Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R.,  
 1727 Cubison, M. J., Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K.,  
 1728 Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell,  
 1729 L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina,  
 1730 K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C.,  
 1731 Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols  
 1732 in the Atmosphere, *Science*, 326, 1525, 10.1126/science.1180353, 2009.  
 1733 Johnston, H.: Reduction of stratospheric ozone by nitrogen oxide catalysts from supersonic transport  
 1734 exhaust, *Science*, 173, 517-522, 10.1126/science.173.3996.517, 1971.  
 1735 Junge, C.: THE SIZE DISTRIBUTION AND AGING OF NATURAL AEROSOLS AS DETERMINED FROM  
 1736 ELECTRICAL AND OPTICAL DATA ON THE ATMOSPHERE, *Journal of Meteorology*, 12, 13-25,  
 1737 10.1175/1520-0469(1955)012<0013:Tsdiao>2.0.Co;2, 1955.  
 1738 Junge, C. E., and Ryan, T. G.: Study of the SO<sub>2</sub> oxidation in solution and its role in atmospheric  
 1739 chemistry, *Quarterly Journal of the Royal Meteorological Society*, 84, 46-55,  
 1740 10.1002/qj.49708435906, 1958.  
 1741 Junge, C. E.: BASIC CONSIDERATIONS ABOUT TRACE CONSTITUENTS IN THE ATMOSPHERE AS  
 1742 RELATED TO THE FATE OF GLOBAL POLLUTANTS, *Adv Environ Sci Technol*, 8, 7-25, 1975.  
 1743 Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S. H., Fisseha, R.,  
 1744 Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of Polymers as  
 1745 Major Components of Atmospheric Organic Aerosols, *Science*, 303, 1659-1662,  
 1746 10.1126/science.1092185, 2004.  
 1747 Karl, T., Harley, P., Emmons, L., Thornton, B., Guenther, A., Basu, C., Turnipseed, A., and Jardine, K.:  
 1748 Efficient Atmospheric Cleansing of Oxidized Organic Trace Gases by Vegetation, *Science*, 1192534,  
 1749 10.1126/science.1192534, 2010.  
 1750 Keeling, C. D.: The Concentration and Isotopic Abundances of Carbon Dioxide in the Atmosphere,  
 1751 *Tellus*, 12, 200-203, 10.1111/j.2153-3490.1960.tb01300.x, 1960.  
 1752 Keeling, C. D., Mook, W. G., and Tans, P. P.: Recent trends in the <sup>13</sup>C/<sup>12</sup>C ratio of atmospheric carbon  
 1753 dioxide, *Nature*, 277, 121-123, 10.1038/277121a0, 1979.  
 1754 Kesselmeier, J., and Staudt, M.: Biogenic Volatile Organic Compounds (VOC): An Overview on  
 1755 Emission, Physiology and Ecology, *Journal of Atmospheric Chemistry*, 33, 23-88,  
 1756 10.1023/A:1006127516791, 1999.  
 1757 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L.,  
 1758 Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G.,  
 1759 Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn,  
 1760 M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A.,  
 1761 Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J.,  
 1762 Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J.  
 1763 H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P.  
 1764 E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R.,  
 1765 Baltensperger, U., and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in  
 1766 atmospheric aerosol nucleation, *Nature*, 476, 429-435, 10.1038/nature10343, 2011.  
 1767 Kleindienst, T. E.: Epoxying Isoprene Chemistry, *Science*, 325, 687-688, 10.1126/science.1178324,  
 1768 2009.  
 1769 Kleinman, L. I.: Seasonal dependence of boundary layer peroxide concentration: the low and high  
 1770 NO<sub>x</sub> regimes, *Journal of Geophysical Research*, 96, 20,721-720,733, 1991.

1771 Kley, D., and McFarland, M.: Chemiluminescence detector for NO and NO<sub>2</sub>, *Atmospheric Technology*,  
1772 12, 63-69, 1980.

1773 Knipping, E. M., Lakin, M. J., Foster, K. L., Jungwirth, P., Tobias, D. J., Gerber, R. B., Dabdub, D., and  
1774 Finlayson-Pitts, B. J.: Experiments and simulations of ion-enhanced interfacial chemistry on aqueous  
1775 NaCl aerosols, *Science*, 288, 301-306, 10.1126/science.288.5464.301, 2000.

1776 Knutson, E. O., and Whitby, K. T.: Aerosol classification by electric mobility: apparatus, theory, and  
1777 applications, *Journal of Aerosol Science*, 6, 443-451, 10.1016/0021-8502(75)90060-9, 1975.

1778 Köhler, H.: The nucleus in and the growth of hygroscopic droplets, *Transactions of the Faraday  
1779 Society*, 32, 1152-1161, 10.1039/TF9363201152, 1936.

1780 Kulmala, M., Pirjola, L., and Mäkelä, J. M.: Stable sulphate clusters as a source of new atmospheric  
1781 particles, *Nature*, 404, 66-69, 10.1038/35003550, 2000.

1782 Kurtenbach, R., Becker, K. H., Gomes, J. A. G., Kleffmann, J., Lörzer, J. C., Spittler, M., Wiesen, P.,  
1783 Ackermann, R., Geyer, A., and Platt, U.: Investigations of emissions and heterogeneous formation of  
1784 HONO in a road traffic tunnel, *Atmospheric Environment*, 35, 3385-3394, 10.1016/S1352-  
1785 2310(01)00138-8, 2001.

1786 Kwok, E. S. C., and Atkinson, R.: Estimation of hydroxyl radical reaction rate constants for gas-phase  
1787 organic compounds using a structure-reactivity relationship: An update, *Atmospheric Environment*,  
1788 29, 1685-1695, 10.1016/1352-2310(95)00069-B, 1995.

1789 Landrigan, P. J., Fuller, R., Acosta, N. J. R., Adeyi, O., Arnold, R., Basu, N. N., Baldé, A. B., Bertollini, R.,  
1790 Bose-O'Reilly, S., Boufford, J. I., Breyse, P. N., Chiles, T., Mahidol, C., Coll-Seck, A. M., Cropper, M. L.,  
1791 Fobil, J., Fuster, V., Greenstone, M., Haines, A., Hanrahan, D., Hunter, D., Khare, M., Krupnick, A.,  
1792 Lanphear, B., Lohani, B., Martin, K., Mathiasen, K. V., McTeer, M. A., Murray, C. J. L.,  
1793 Ndahimananjara, J. D., Perera, F., Potočník, J., Preker, A. S., Ramesh, J., Rockström, J., Salinas, C.,  
1794 Samson, L. D., Sandilya, K., Sly, P. D., Smith, K. R., Steiner, A., Stewart, R. B., Suk, W. A., van Schayck,  
1795 O. C. P., Yadama, G. N., Yumkella, K., and Zhong, M.: The Lancet Commission on pollution and health,  
1796 *The Lancet*, 391, 462-512, 10.1016/S0140-6736(17)32345-0, 2018.

1797 Lawson, D. R.: "Passing the test" – human behavior and California's smog check program, *Air and  
1798 Waste*, 43, 1567-1575, 10.1080/1073161X.1993.10467226, 1993.

1799 Leighton, P. A.: *Photochemistry of Air Pollution*, Academic Press, New York, 1961.

1800 Lelieveld, J., Butler, T. M., Crowley, J. N., Dillon, T. J., Fischer, H., Ganzeveld, L., Lawrence, M. G.,  
1801 Martinez, M., Taraborrelli, D., and Williams, J.: Atmospheric oxidation capacity sustained by a  
1802 tropical forest, *Nature*, 452, 737-740 2008.

1803 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air  
1804 pollution sources to premature mortality on a global scale, *Nature*, 525, 367-371,  
1805 10.1038/nature15371, 2015.

1806 Leu, M. T.: Laboratory studies of sticking coefficients and heterogeneous reactions important in the  
1807 Antarctic stratosphere, *Geophysical Research Letters*, 15, 17-20, 10.1029/GL015i001p00017, 1988.

1808 Levy, H.: NORMAL ATMOSPHERE - LARGE RADICAL AND FORMALDEHYDE CONCENTRATIONS  
1809 PREDICTED, *Science*, 173, 141-&, 10.1126/science.173.3992.141, 1971.

1810 Lewis, A. C., Carslaw, N., Marriott, P. J., Kinghorn, R. M., Morrison, P., Lee, A. L., Bartle, K. D., and  
1811 Pilling, M. J.: A larger pool of ozone-forming carbon compounds in urban atmospheres, *Nature*, 405,  
1812 778-781, 10.1038/35015540, 2000.

1813 Likens, G. E., and Bormann, F. H.: Acid rain: A serious regional environmental problem, *Science*, 184,  
1814 1176-1179, 10.1126/science.184.4142.1176, 1974.

1815 Lin, X., Trainer, M., and Liu, S. C.: On the nonlinearity of the tropospheric ozone production, *Journal  
1816 of Geophysical Research: Atmospheres*, 93, 15879-15888, 10.1029/JD093iD12p15879, 1988.

1817 Lindinger, W., Hansel, A., and Jordan, A.: On-line monitoring of volatile organic compounds at pptv  
1818 levels by means of proton-transfer-reaction mass spectrometry (PTR-MS) - Medical applications,  
1819 food control and environmental research, *Int. J. Mass Spectrom.*, 173, 191-241, 1998.

1820 Liu, S. C.: Ozone production in the rural troposphere and the implications for regional and global  
 1821 ozone distributions, *Journal of Geophysical Research*, 92, 4191-4207, 10.1029/JD092iD04p04191,  
 1822 1987.

1823 Liu, Y., Park, R. J., Jacob, D. J., Li, Q., Kilaru, V., and Sarnat, J. A.: Mapping annual mean ground-level  
 1824 PM<sub>2.5</sub> concentrations using Multiangle Imaging Spectroradiometer aerosol optical thickness over  
 1825 the contiguous United States, *Journal of Geophysical Research: Atmospheres*, 109,  
 1826 <https://doi.org/10.1029/2004JD005025>, 2004.

1827 Logan, J. A., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Tropospheric chemistry: a global  
 1828 perspective, *Journal of Geophysical Research*, 86, 7210-7254, 10.1029/JC086iC08p07210, 1981.

1829 Logan, J. A.: Nitrogen oxides in the troposphere: global and regional budgets, *Journal of Geophysical*  
 1830 *Research*, 88, 10785-10807, 10.1029/JC088iC15p10785, 1983.

1831 Logan, J. A.: Tropospheric ozone - seasonal behaviour, trends, and anthropogenic influence, *J.*  
 1832 *Geophys. Res.-Atmos.*, 90, 10463-10482, 1985.

1833 Logan, J. A.: Ozone in rural areas of the United States, *Journal of Geophysical Research*, 94, 8511-  
 1834 8532, 10.1029/JD094iD06p08511, 1989.

1835 Logan, J. A.: An analysis of ozonesonde data for the troposphere: Recommendations for testing 3-D  
 1836 models and development of a gridded climatology for tropospheric ozone, *Journal of Geophysical*  
 1837 *Research Atmospheres*, 104, 16115-16149, 10.1029/1998JD100096, 1999.

1838 Lovelock, J. E., and Lipsky, S. R.: Electron Affinity Spectroscopy—A New Method for the Identification  
 1839 of Functional Groups in Chemical Compounds Separated by Gas Chromatography, *Journal of the*  
 1840 *American Chemical Society*, 82, 431-433, 10.1021/ja01487a045, 1960.

1841 Lovelock, J. E., Maggs, R. J., and Rasmussen, R. A.: Atmospheric Dimethyl Sulphide and the Natural  
 1842 Sulphur Cycle, *Nature*, 237, 452-453, 10.1038/237452a0, 1972.

1843 Lovelock, J. E.: Atmospheric halocarbons and stratospheric ozone, *Nature*, 252, 292-294,  
 1844 10.1038/252292a0, 1974.

1845 MacKenzie, R. A., Harrison, R. M., Colbeck, I., and Nicholas Hewitt, C.: The role of biogenic  
 1846 hydrocarbons in the production of ozone in urban plumes in southeast England, *Atmospheric*  
 1847 *Environment. Part A. General Topics*, 25, 351-359, [https://doi.org/10.1016/0960-1686\(91\)90306-R](https://doi.org/10.1016/0960-1686(91)90306-R),  
 1848 1991.

1849 Madronich, S., and Flocke, S.: The Role of Solar Radiation in Atmospheric Chemistry, *The Handbook*  
 1850 *of Environmental Chemistry (Reactions and Processes)*. edited by: Boule, P., Springer-Verlag, Berlin,  
 1851 Heidelberg, 1999.

1852 Mäkelä, J. M., Aalto, P., Jokinen, V., Pohja, T., Nissinen, A., Palmroth, S., Markkanen, T., Seitsonen, K.,  
 1853 Lihavainen, H., and Kulmala, M.: Observations of ultrafine aerosol particle formation and growth in  
 1854 boreal forest, *Geophysical Research Letters*, 24, 1219-1222, 10.1029/97GL00920, 1997.

1855 Martin, R. V.: Satellite remote sensing of surface air quality, *Atmospheric Environment*, 42, 7823-  
 1856 7843, 10.1016/j.atmosenv.2008.07.018, 2008.

1857 Mauldin Iii, R. L., Berndt, T., Sipilä, M., Paasonen, P., Petäjä, T., Kim, S., Kurtén, T., Stratmann, F.,  
 1858 Kerminen, V. M., and Kulmala, M.: A new atmospherically relevant oxidant of sulphur dioxide,  
 1859 *Nature*, 488, 193-196, 10.1038/nature11278, 2012.

1860 Maynard, R. L., and Williams, M. L.: Regulation of Air Quality in the European Union, in: *Regulatory*  
 1861 *Toxicology in the European Union*, The Royal Society of Chemistry, 539-556, 2018.

1862 McElroy, M. B., Salawitch, R. J., Wofsy, S. C., and Logan, J. A.: Reductions of Antarctic ozone due to  
 1863 synergistic interactions of chlorine and bromine, *Nature*, 321, 759-762, 10.1038/321759a0, 1986.

1864 Melamed, M. L., Monks, P. S., Goldstein, A. H., Lawrence, M. G., and Jennings, J.: The international  
 1865 global atmospheric chemistry (IGAC) project: Facilitating atmospheric chemistry research for 25  
 1866 years, *Anthropocene*, 12, 17-28, <http://dx.doi.org/10.1016/j.ancene.2015.10.001>, 2015.

1867 Merrill, J. T., Bleck, R., and Avila, L.: Modeling atmospheric transport to the Marshall Islands, *Journal*  
 1868 *of Geophysical Research: Atmospheres*, 90, 12927-12936,  
 1869 <https://doi.org/10.1029/JD090iD07p12927>, 1985.

1870 Molina, L. T., and Molina, M. J.: Production of chlorine oxide ( $\text{Cl}_2\text{O}_2$ ) from the self-reaction of the  
1871 chlorine oxide (ClO) radical, *The Journal of Physical Chemistry*, 91, 433-436, 10.1021/j100286a035,  
1872 1987.

1873 Molina, M. J., and Rowland, F. S.: Stratospheric sink for chlorofluoromethanes: Chlorine atomc-  
1874 atalysed destruction of ozone, *Nature*, 249, 810-812, 10.1038/249810a0, 1974.

1875 Molina, M. J., Tso, T. L., Molina, L. T., and Wang, F. C. Y.: Antarctic stratospheric chemistry of chlorine  
1876 nitrate, hydrogen chloride, and ice: Release of active chlorine, *Science*, 238, 1253-1257,  
1877 10.1126/science.238.4831.1253, 1987.

1878 Molina, M. J.: Heterogeneous chemistry on polar stratospheric clouds, *Atmospheric Environment*  
1879 Part A, General Topics, 25, 2535-2537, 10.1016/0960-1686(91)90170-C, 1991.

1880 Monks, P. S.: Gas-phase radical chemistry in the troposphere, *Chem Soc Rev*, 34, 376-395, Doi  
1881 10.1039/B307982c, 2005.

1882 Monks, P. S., Granier, C., Fuzzi, S., Stohl, A., Williams, M. L., Akimoto, H., Amann, M., Baklanov, A.,  
1883 Baltensperger, U., Bey, I., Blake, N., Blake, R. S., Carslaw, K., Cooper, O. R., Dentener, F., Fowler, D.,  
1884 Fragkou, E., Frost, G. J., Generoso, S., Ginoux, P., Grewe, V., Guenther, A., Hansson, H. C., Henne, S.,  
1885 Hjorth, J., Hofzumahaus, A., Huntrieser, H., Isaksen, I. S. A., Jenkin, M. E., Kaiser, J., Kanakidou, M.,  
1886 Klimont, Z., Kulmala, M., Laj, P., Lawrence, M. G., Lee, J. D., Liousse, C., Maione, M., McFiggans, G.,  
1887 Metzger, A., Mieville, A., Moussiopoulos, N., Orlando, J. J., O'Dowd, C. D., Palmer, P. I., Parrish, D. D.,  
1888 Petzold, A., Platt, U., Poschl, U., Prevot, A. S. H., Reeves, C. E., Reimann, S., Rudich, Y., Sellegri, K.,  
1889 Steinbrecher, R., Simpson, D., ten Brink, H., Theloke, J., van der Werf, G. R., Vautard, R., Vestreng, V.,  
1890 Vlachokostas, C., and von Glasow, R.: Atmospheric composition change - global and regional air  
1891 quality, *Atmospheric Environment*, 43, 5268-5350, DOI 10.1016/j.atmosenv.2009.08.021, 2009.

1892 Monks, P. S., and Williams, M. L.: What does success look like for air quality policy? A perspective,  
1893 *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*,  
1894 378, 20190326, doi:10.1098/rsta.2019.0326, 2020.

1895 Montzka, S. A., Calvert, P., Hall, B. D., Elkins, J. W., Conway, T. J., Tans, P. P., and Sweeney, C.: On the  
1896 global distribution, seasonality, and budget of atmospheric carbonyl sulfide (COS) and some  
1897 similarities to CO<sub>2</sub>, *Journal of Geophysical Research: Atmospheres*, 112,  
1898 <https://doi.org/10.1029/2006JD007665>, 2007.

1899 Moody, J. L., Oltmans, S. J., Levy, H., and Merrill, J. T.: Transport, climatology of tropospheric ozone -  
1900 Bermuda, 1988-1991, *J. Geophys. Res.-Atmos.*, 100, 7179-7194, 10.1029/94jd02830, 1995.

1901 Moody, J. L., Munger, J. W., Goldstein, A. H., Jacob, D. J., and Wofsy, S. C.: Harvard Forest regional-  
1902 scale air mass composition by Patterns in Atmospheric Transport History (PATH), *Journal of*  
1903 *Geophysical Research: Atmospheres*, 103, 13181-13194, 10.1029/98jd00526, 1998.

1904 Mozurkewich, M., McMurry, P. H., Gupta, A., and Calvert, J. G.: Mass Accomodation Coefficient for  
1905 HO<sub>2</sub> Radicals on Aqueous Particles, *J. Geophys. Res.*, 92, 4163-4170, 1987.

1906 Murozumi, M., Chow, T. J., and Patterson, C.: Chemical concentrations of pollutant lead aerosols,  
1907 terrestrial dusts and sea salts in Greenland and Antarctic snow strata, *Geochimica et Cosmochimica*  
1908 *Acta*, 33, 1247-1294, [https://doi.org/10.1016/0016-7037\(69\)90045-3](https://doi.org/10.1016/0016-7037(69)90045-3), 1969.

1909 Murphy, D. M., Cziczo, D. J., Froyd, K. D., Hudson, P. K., Matthew, B. M., Middlebrook, A. M., Peltier,  
1910 R. E., Sullivan, A., Thomson, D. S., and Weber, R. J.: Single-peptide mass spectrometry of  
1911 tropospheric aerosol particles, *Journal of Geophysical Research Atmospheres*, 111,  
1912 10.1029/2006JD007340, 2006.

1913 Newell, R. E., Thouret, V., Cho, J. Y. N., Stoller, P., Marengo, A., and Smit, H. G.: Ubiquity of quasi-  
1914 horizontal layers in the troposphere, *Nature*, 398, 316-319, 1999.

1915 Norrish, R. G. W., and Neville, G. H. J.: The decomposition of ozone photosensitised by chlorine,  
1916 *Journal of the Chemical Society (Resumed)*, 1864-1872, 1934.

1917 O'Dowd, C. D., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Young, J. Y.,  
1918 and Putaud, J. P.: Biogenically driven organic contribution to marine aerosol, *Nature*, 431, 676-680,  
1919 10.1038/nature02959, 2004.

1920 O'Dowd, C. D., Bahreini, R., Flagan, R. C., Seinfeld, J. H., Hämerl, K., Pirjola, L., Kulmala, M., and  
 1921 Hoffmann, T.: Marine aerosol formation from biogenic iodine emissions, *Nature*, 417, 632-636,  
 1922 10.1038/nature00775, 2002.

1923 Odén, S.: The acidification of air precipitation and its consequences in the natural environment, NFR,  
 1924 1968.

1925 Odum, J. R., Jungkamp, T. P. W., Griffin, R. J., Flagan, R. C., and Seinfeld, J. H.: The atmospheric  
 1926 aerosol forming potential of whole gasoline vapor, *Science*, 276, 96-99, 1997.

1927 Olivier, J. G. J., Bouwman, A. F., van der Maas, C. W. M., and Berdowski, J. J. M.: Emission database  
 1928 for global atmospheric research (Edgar), *Environmental Monitoring and Assessment*, 31, 93-106,  
 1929 10.1007/BF00547184, 1994.

1930 Osthoff, H. D., Roberts, J. M., Ravishankara, A. R., Williams, E. J., Lerner, B. M., Sommariva, R., Bates,  
 1931 T. S., Coffman, D., Quinn, P. K., Dibb, J. E., Stark, H., Burkholder, J. B., Talukdar, R. K., Meagher, J.,  
 1932 Fehsenfeld, F. C., and Brown, S. S.: High levels of nitryl chloride in the polluted subtropical marine  
 1933 boundary layer, *Nature Geoscience*, 1, 324-328, 10.1038/ngeo177, 2008.

1934 Pales, J. C., and Keeling, C. D.: The concentration of atmospheric carbon dioxide in Hawaii, *Journal of*  
 1935 *Geophysical Research* (1896-1977), 70, 6053-6076, 10.1029/JZ070i024p06053, 1965.

1936 Palmer, P. I., Jacob, D. J., Chance, K., Martin, R. V., Spurr, R. J. D., Kurosu, T. P., Bey, I., Yantosca, R.,  
 1937 Fiore, A., and Li, Q.: Air mass factor formulation for spectroscopic measurements from satellites:  
 1938 Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment, *Journal of*  
 1939 *Geophysical Research: Atmospheres*, 106, 14539-14550, 10.1029/2000jd900772, 2001.

1940 Pankow, J. F.: An absorption model of gas/particle partitioning of organic compounds in the  
 1941 atmosphere, *Atmospheric Environment*, 28, 185-188, [https://doi.org/10.1016/1352-2310\(94\)90093-](https://doi.org/10.1016/1352-2310(94)90093-0)  
 1942 [0](https://doi.org/10.1016/1352-2310(94)90093-0), 1994a.

1943 Pankow, J. F.: An absorption model of the gas/aerosol partitioning involved in the formation of  
 1944 secondary organic aerosol, *Atmospheric Environment*, 28, 189-193, [https://doi.org/10.1016/1352-](https://doi.org/10.1016/1352-2310(94)90094-9)  
 1945 [2310\(94\)90094-9](https://doi.org/10.1016/1352-2310(94)90094-9), 1994b.

1946 Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kürten, A., Clair, J. M. S., Seinfeld, J. H., and Wennberg, P.  
 1947 O.: Unexpected epoxide formation in the gas-phase photooxidation of isoprene, *Science*, 325, 730-  
 1948 733, 10.1126/science.1172910, 2009.

1949 Paulson, S. E., and Orlando, J. J.: The reactions of ozone with alkenes: An important source of HOx in  
 1950 the boundary layer, *Geophysical Research Letters*, 23, 3727-3730, 10.1029/96gl03477, 1996.

1951 Peeters, J., Nguyen, T. L., and Vereecken, L.: HOx radical regeneration in the oxidation of isoprene,  
 1952 *Phys. Chem. Chem Phys.*, 11, 5935 - 5939, 2009.

1953 Penkett, S. A., Jones, B. M. R., Brich, K. A., and Eggleton, A. E. J.: The importance of atmospheric  
 1954 ozone and hydrogen peroxide in oxidising sulphur dioxide in cloud and rainwater, *Atmospheric*  
 1955 *Environment* (1967), 13, 123-137, 10.1016/0004-6981(79)90251-8, 1979.

1956 Perner, D., Ehhalt, D. H., Pätz, H. W., Platt, U., Röth, E. P., and Volz, A.: OH - Radicals in the lower  
 1957 troposphere, *Geophysical Research Letters*, 3, 466-468, 10.1029/GL003i008p00466, 1976a.

1958 Perner, D., Ehhalt, D. H., Pätz, H. W., Platt, U., Röth, E. P., and Volz, A.: OH - Radicals in the lower  
 1959 troposphere, *Geophysical Research Letters*, 3, 466-468, <https://doi.org/10.1029/GL003i008p00466>,  
 1960 1976b.

1961 Perner, D., and Platt, U.: Detection of nitrous acid in the atmosphere by differential optical  
 1962 absorption, *Geophysical Research Letters*, 6, 917-920, 10.1029/GL006i012p00917, 1979.

1963 Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J. M., Basile, I., Bender, M., Chappellaz, J.,  
 1964 Davis, M., Delaygue, G., Delmotte, M., Kotiyakov, V. M., Legrand, M., Lipenkov, V. Y., Lorius, C.,  
 1965 Pépin, L., Ritz, C., Saltzman, E., and Stievenard, M.: Climate and atmospheric history of the past  
 1966 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429-436, 10.1038/20859, 1999.

1967 Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and  
 1968 cloud condensation nucleus activity, *Atmospheric Chemistry and Physics*, 7, 1961-1971,  
 1969 10.5194/acp-7-1961-2007, 2007.



1970 Pitts, J., Van Cauwenberghe, K., Grosjean, D., Schmid, J., Fitz, D., Belser, W., Knudson, G., and Hynds,  
1971 P.: Atmospheric reactions of polycyclic aromatic hydrocarbons: facile formation of mutagenic nitro  
1972 derivatives, *Science*, 202, 515-519, 10.1126/science.705341, 1978.  
1973 Pitts, J. N., Sanhueza, E., Atkinson, R., Carter, W. P. L., Winer, A. M., Harris, G. W., and Plum, C. N.:  
1974 An investigation of the dark formation of nitrous acid in environmental chambers, *International*  
1975 *Journal of Chemical Kinetics*, 16, 919-939, <https://doi.org/10.1002/kin.550160712>, 1984.  
1976 Platt, U., Perner, D., and Paetz, H. W.: SIMULTANEOUS MEASUREMENT OF ATMOSPHERIC CH<sub>2</sub>O, O<sub>3</sub>,  
1977 AND NO<sub>2</sub> BY DIFFERENTIAL OPTICAL ABSORPTION, *Journal of Geophysical Research*, 84, 6329-6335,  
1978 10.1029/JC084iC10p06329, 1979.  
1979 Platt, U., Perner, D., Winer, A. M., Harris, G. W., and Pitts Jr, J. N.: Detection of NO<sub>3</sub> in the polluted  
1980 troposphere by differential optical absorption, *Geophysical Research Letters*, 7, 89-92,  
1981 10.1029/GL007i001p00089, 1980.  
1982 Platt, U., and Hönninger, G.: The role of halogen species in the troposphere, *Chemosphere*, 52, 325-  
1983 338, 2003.  
1984 Prather, M. J.: Lifetimes and eigenstates in atmospheric chemistry, *Geophysical Research Letters*, 21,  
1985 801-804, 10.1029/94GL00840, 1994.  
1986 Prather, M. J.: Time scales in atmospheric chemistry: Theory, GWPs for CH<sub>4</sub> and CO, and runaway  
1987 growth, *Geophysical Research Letters*, 23, 2597-2600, 10.1029/96GL02371, 1996.  
1988 Preining, O., and Davis, E. J.: History of Aerosol Science, Symposium on the History of Aerosol  
1989 Science, Vienna, Austria, 1999.  
1990 Prinn, R. G., Weiss, R. F., Miller, B. R., Huang, J., Alyea, F. N., Cunnold, D. M., Fraser, P. J., Hartley, D.  
1991 E., and Simmonds, P. G.: Atmospheric trends and lifetime of CH<sub>3</sub>CCl<sub>3</sub> and global OH concentrations,  
1992 *Science*, 269, 187-192, 10.1126/science.269.5221.187, 1995.  
1993 Prinn, R. G., Huang, J., Weiss, R. F., Cunnold, D. M., Fraser, P. J., Simmonds, P. G., McCulloch, A.,  
1994 Harth, C., Salameh, P., O'Doherty, S., Wang, R. H. J., Porter, L., and Miller, B. R.: Evidence for  
1995 substantial variations of atmospheric hydroxyl radicals in the past two decades, *Science*, 292, 1882-  
1996 1888, 10.1126/science.1058673, 2001.  
1997 Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., and Gill, T. E.: Environmental characterization  
1998 of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping  
1999 Spectrometer (TOMS) absorbing aerosol product, *Reviews of Geophysics*, 40, 2-1-2-31,  
2000 10.1029/2000RG000095, 2002.  
2001 Quinn, P. K., and Bates, T. S.: The case against climate regulation via oceanic phytoplankton sulphur  
2002 emissions, *Nature*, 480, 51-56, 10.1038/nature10580, 2011.  
2003 Ramanathan, V., Cicerone, R. J., Singh, H. B., and Kiehl, J. T.: Trace gas trends and their potential role  
2004 in climate change, *Journal of Geophysical Research*, 90, 5547-5566, 10.1029/JD090iD03p05547,  
2005 1985.  
2006 Ramanathan, V., Crutzen, P. J., Kiehl, J. T., and Rosenfeld, D.: Atmosphere - Aerosols, climate, and  
2007 the hydrological cycle, *Science*, 294, 2119-2124, 10.1126/science.1064034, 2001.  
2008 Rasmussen, R., and Went, F. W.: Volatile organic matter of plant origin in the atmosphere, *Science*,  
2009 144, 566, 10.1126/science.144.3618.566-a, 1964.  
2010 Ravishankara, A. R.: Heterogeneous and multiphase chemistry in the troposphere, *Science*, 276,  
2011 1058-1065, 10.1126/science.276.5315.1058, 1997.  
2012 Ravishankara, A. R., Hancock, G., Kawasaki, M., and Matsumi, Y.: Photochemistry of ozone: Surprises  
2013 and recent lessons, *Science*, 280, 60-61, 10.1126/science.280.5360.60, 1998.  
2014 Ravishankara, A. R., Dunlea, E. J., Blitz, M. A., Dillon, T. J., Heard, D. E., Pilling, M. J., Strekowski, R. S.,  
2015 Nicovich, J. M., and Wine, P. H.: Redetermination of the rate coefficient for the reaction of O(<sup>1</sup>D)  
2016 with N<sub>2</sub>, *Geophysical Research Letters*, 29, 35-31-35-34, 10.1029/2002GL014850, 2002.  
2017 Ravishankara, A. R.: Introduction: Atmospheric Chemistry Long-Term Issues, *Chemical Reviews*, 103,  
2018 4505-4508, 10.1021/cr020463i, 2003.  
2019 Ravishankara, A. R., Rudich, Y., and Pyle, J. A.: Role of Chemistry in Earth's Climate, *Chemical*  
2020 *Reviews*, 115, 3679-3681, 10.1021/acs.chemrev.5b00226, 2015.

2021 Read, K. A., Mahajan, A. S., Carpenter, L. J., Evans, M. J., Faria, B. V. E., Heard, D. E., Hopkins, J. R.,  
 2022 Lee, J. D., Moller, S. J., Lewis, A. C., Mendes, L., McQuaid, J. B., Oetjen, H., Saiz-Lopez, A., Pilling, M.  
 2023 J., and Plane, J. M. C.: Extensive halogen-mediated ozone destruction over the tropical Atlantic  
 2024 Ocean, *Nature*, 453, 1232-1235, 2008.  
 2025 Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part  
 2026 II: Intensive physical properties of biomass burning particles, *Atmospheric Chemistry and Physics*, 5,  
 2027 799-825, 10.5194/acp-5-799-2005, 2005.  
 2028 Remer, L. A., Kaufman, Y. J., Tanré, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R. R., Ichoku, C., Levy,  
 2029 R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N.: The MODIS aerosol algorithm,  
 2030 products, and validation, *Journal of the Atmospheric Sciences*, 62, 947-973, 10.1175/JAS3385.1,  
 2031 2005.  
 2032 Reynolds, S. D., Roth, P. M., and Seinfeld, J. H.: Mathematical modeling of photochemical air  
 2033 pollution—I: Formulation of the model, *Atmospheric Environment* (1967), 7, 1033-1061,  
 2034 [https://doi.org/10.1016/0004-6981\(73\)90214-X](https://doi.org/10.1016/0004-6981(73)90214-X), 1973.  
 2035 Reynolds, S. D., Liu, M.-K., Hecht, T. A., Roth, P. M., and Seinfeld, J. H.: Mathematical modeling of  
 2036 photochemical air pollution—III. Evaluation of the model, *Atmospheric Environment* (1967), 8, 563-  
 2037 596, [https://doi.org/10.1016/0004-6981\(74\)90143-7](https://doi.org/10.1016/0004-6981(74)90143-7), 1974.  
 2038 Richter, A., Burrows, J. P., Nuss, H., Granier, C., and Niemeier, U.: Increase in tropospheric nitrogen  
 2039 dioxide over China observed from space, *Nature*, 437, 129-132, 10.1038/nature04092, 2005.  
 2040 Robbin, M. L., and Damschen, D. E.: Aqueous oxidation of sulfur dioxide by hydrogen peroxide at low  
 2041 pH, *Atmospheric Environment* (1967), 15, 1615-1621, 10.1016/0004-6981(81)90146-3, 1981.  
 2042 Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P.,  
 2043 Lane, T. E., Pierce, J. R., and Pandis, S. N.: Rethinking Organic Aerosols: Semivolatile Emissions and  
 2044 Photochemical Aging, *Science*, 315, 1259-1262, doi:10.1126/science.1133061, 2007.  
 2045 Roth, P. M., Roberts, P. J. W., Mei-Kao, L., Reynolds, S. D., and Seinfeld, J. H.: Mathematical modeling  
 2046 of photochemical air pollution—II. A model and inventory of pollutant emissions, *Atmospheric*  
 2047 *Environment* (1967), 8, 97-130, [https://doi.org/10.1016/0004-6981\(74\)90023-7](https://doi.org/10.1016/0004-6981(74)90023-7), 1974.  
 2048 Sanadze, G. A.: The nature of gaseous substances emitted by leaves of Robinia pseudoacacia,  
 2049 *Soobshch Akad Nauk Gruz SSR*, 19, 83–86, 1957.  
 2050 Sanadze, G. A., and Kursanov, A. N.: Light and temperature curves of the evolution of C<sub>5</sub>H<sub>8</sub>,  
 2051 *Fiziol.Rast.*, 13, 458–461, 1966.  
 2052 Sander, S. P., Friedl, R. R., and Yung, Y. L.: Rate of formation of the ClO dimer in the polar  
 2053 stratosphere: Implications for ozone loss, *Science*, 245, 1095-1098, 10.1126/science.245.4922.1095,  
 2054 1989.  
 2055 Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the  
 2056 Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile  
 2057 organic compounds, *Atmos. Chem. Phys.*, 3, 161-180, 10.5194/acp-3-161-2003, 2003.  
 2058 Schroeder, W. H., and Urone, P.: Formation of nitrosyl chloride from salt particles in air,  
 2059 *Environmental Science & Technology*, 8, 756-758, 10.1021/es60093a015, 1974.  
 2060 Seinfeld, J. H., and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate*  
 2061 *Change*, 2nd ed., Wiley, 2006.  
 2062 Seitzinger, S. P., Gaffney, O., Brasseur, G., Broadgate, W., Ciais, P., Claussen, M., Erisman, J. W.,  
 2063 Kiefer, T., Lancelot, C., Monks, P. S., Smyth, K., Syvitski, J., and Uematsu, M.: International  
 2064 Geosphere–Biosphere Programme and Earth system science: Three decades of co-evolution,  
 2065 *Anthropocene*, 12, 3-16, <http://dx.doi.org/10.1016/j.ancene.2016.01.001>, 2015.  
 2066 Sharkey, T. D., and Monson, R. K.: Isoprene research – 60 years later, the biology is still enigmatic,  
 2067 *Plant, Cell & Environment*, 40, 1671-1678, <https://doi.org/10.1111/pce.12930>, 2017.  
 2068 Shindell, D., Kuylenstierna, J. C. I., Vignati, E., van Dingenen, R., Amann, M., Klimont, Z., Anenberg, S.  
 2069 C., Muller, N., Janssens-Maenhout, G., Raes, F., Schwartz, J., Faluvegi, G., Pozzoli, L., Kupiainen, K.,  
 2070 Höglund-Isaksson, L., Emberson, L., Streets, D., Ramanathan, V., Hicks, K., Oanh, N. T. K., Milly, G.,



2071 Williams, M., Demkine, V., and Fowler, D.: Simultaneously Mitigating Near-Term Climate Change and  
 2072 Improving Human Health and Food Security, *Science*, 335, 183-189, 10.1126/science.1210026, 2012.  
 2073 Sillman, S.: The relation between ozone, NO<sub>x</sub> and hydrocarbons in urban and polluted rural  
 2074 environments, *Atmospheric Environment*, 33, 1821-1845, 10.1016/s1352-2310(98)00345-8, 1999.  
 2075 Sillman, S., and He, D.: Some theoretical results concerning O<sub>3</sub>-NO<sub>x</sub>-VOC chemistry and NO<sub>x</sub>-VOC  
 2076 indicators, *Journal of Geophysical Research Atmospheres*, 107, 10.1029/2001JD001123, 2002.  
 2077 Simoneit, B. R. T., Schauer, J. J., Nolte, C. G., Oros, D. R., Elias, V. O., Fraser, M. P., Rogge, W. F., and  
 2078 Cass, G. R.: Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles,  
 2079 *Atmospheric Environment*, 33, 173-182, 10.1016/S1352-2310(98)00145-9, 1999.  
 2080 Singer, B. C., and Harley, R. A.: A Fuel-Based Motor Vehicle Emission Inventory, *Journal of the Air and*  
 2081 *Waste Management Association*, 46, 581-593, 10.1080/10473289.1996.10467492, 1996.  
 2082 Singh, H. B.: Preliminary estimation of average tropospheric HO concentrations in the northern and  
 2083 southern hemispheres, *Geophysical Research Letters*, 4, 453-456, 10.1029/GL004i010p00453, 1977.  
 2084 Singh, H. B., and Hanst, P. L.: Peroxyacetyl nitrate (PAN) in the unpolluted atmosphere: An important  
 2085 reservoir for nitrogen oxides, *Geophysical Research Letters*, 8, 941-944, 10.1029/GL008i008p00941,  
 2086 1981.  
 2087 Sinha, M. P.: Laser-induced volatilization and ionization of microparticles, *Review of Scientific*  
 2088 *Instruments*, 55, 886-891, 10.1063/1.1137851, 1984.  
 2089 Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of Antarctic ozone,  
 2090 *Nature*, 321, 755-758, 10.1038/321755a0, 1986.  
 2091 Spicer, C. W., Chapman, E. G., Finlayson-Pitts, B. J., Plastridge, R. A., Hubbe, J. M., Fast, J. D., and  
 2092 Berkowitz, C. M.: Unexpectedly high concentrations of molecular chlorine in coastal air, *Nature*, 394,  
 2093 353-356, 10.1038/28584, 1998.  
 2094 Spivakovsky, C. M., Logan, J. A., Montzka, S. A., Balkanski, Y. J., Foreman-Fowler, M., Jones, D. B. A.,  
 2095 Horowitz, L. W., Fusco, A. C., Brenninkmeijer, C. A. M., Prather, M. J., Wofsy, S. C., and McElroy, M.  
 2096 B.: Three-dimensional climatological distribution of tropospheric OH: Update and evaluation, *Journal*  
 2097 *of Geophysical Research Atmospheres*, 105, 8931-8980, 10.1029/1999JD901006, 2000.  
 2098 Sportisse, B.: *Fundamentals in Air Pollution*, Springer, 293 pp., 2010.  
 2099 Stelson, A. W., Friedlander, S. K., and Seinfeld, J. H.: A note on the equilibrium relationship between  
 2100 ammonia and nitric acid and particulate ammonium nitrate, *Atmospheric Environment* (1967), 13,  
 2101 369-371, 10.1016/0004-6981(79)90293-2, 1979.  
 2102 Stelson, A. W., and Seinfeld, J. H.: Thermodynamic prediction of the water activity, NH<sub>4</sub>NO<sub>3</sub>  
 2103 dissociation constant, density and refractive index for the NH<sub>4</sub>NO<sub>3</sub>-(NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O system at 25°C,  
 2104 *Atmospheric Environment* (1967), 16, 2507-2514, 10.1016/0004-6981(82)90142-1, 1982.  
 2105 Stern, A. C.: *Air Pollution*, 2 ed., Academic Press, New York, 1968.  
 2106 Stevens, P. S., Mather, J. H., and Brune, W. H.: Measurement of tropospheric OH and HO<sub>2</sub> by laser-  
 2107 induced fluorescence at low pressure, *Journal of Geophysical Research*, 99, 3543-3557,  
 2108 10.1029/93JD03342, 1994.  
 2109 Stevenson, D. S., Dentener, F. J., Schultz, M. G., Ellingsen, K., van Noije, T. P. C., Wild, O., Zeng, G.,  
 2110 Amann, M., Atherton, C. S., Bell, N., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J.,  
 2111 Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A.,  
 2112 Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J. F., Lawrence, M. G., Montanaro, V., Muller,  
 2113 J. F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H.,  
 2114 Shindell, D. T., Strahan, S. E., Sudo, K., and Szopa, S.: Multimodel ensemble simulations of present-  
 2115 day and near-future tropospheric ozone, *J. Geophys. Res.-Atmos.*, 111, 10.1029/2005jd006338,  
 2116 2006.  
 2117 Stockwell, W. R., Middleton, P., Chang, J. S., and Tang, X.: The second generation regional acid  
 2118 deposition model chemical mechanism for regional air quality modeling, *Journal of Geophysical*  
 2119 *Research: Atmospheres*, 95, 16343-16367, 10.1029/JD095iD10p16343, 1990.

2120 Stockwell, W. R., Kirchner, F., Kuhn, M., and Seefeld, S.: A new mechanism for regional atmospheric  
 2121 chemistry modeling, *Journal of Geophysical Research: Atmospheres*, 102, 25847-25879,  
 2122 10.1029/97jd00849, 1997.  
 2123 Stohl, A., and Trickl, T.: A text book example of long range transport: simultaneous observation of  
 2124 ozone maxima of stratospheric and North American origin in the FT over Europe, *J. Geophys. Res.*,  
 2125 104, 30,445-430,462., 1999.  
 2126 Stohl, A., Eckhardt, S., Forster, C., James, P., and Spichtinger, N.: On the pathways and timescales of  
 2127 intercontinental air pollution transport, *Journal of Geophysical Research: Atmospheres*, 107,  
 2128 10.1029/2001JD001396, 2002.  
 2129 Stohl, A., Bonasoni, P., Cristofanelli, P., Collins, W., Feichter, J., Frank, A., Forster, C., Gerasopoulos,  
 2130 E., Gäggeler, H., James, P., Kentarchos, T., Kromp-Kolb, H., Krüger, B., Land, C., Meloen, J.,  
 2131 Papayannis, A., Priller, A., Seibert, P., Sprenger, M., Roelofs, G. J., Scheel, H. E., Schnabel, C.,  
 2132 Siegmund, P., Tobler, L., Trickl, T., Wernli, H., Wirth, V., Zanis, P., and Zerefos, C.: Stratosphere-  
 2133 troposphere exchange: A review, and what we have learned from STACCATO, *Journal of Geophysical*  
 2134 *Research: Atmospheres*, 108, 10.1029/2002jd002490, 2003.  
 2135 Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle  
 2136 dispersion model FLEXPART version 6.2, *Atmospheric Chemistry and Physics*, 5, 2461-2474, 2005.  
 2137 Stolarski, R. S., and Cicerone, R. J.: Stratospheric Chlorine: a Possible Sink for Ozone, *Canadian*  
 2138 *Journal of Chemistry*, 52, 1610-1615, 10.1139/v74-233, 1974.  
 2139 Stull, R. B.: *An Introduction to Boundary Layer Meteorology*, Kluwer Academic, Dordrecht, 1988.  
 2140 Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan,  
 2141 R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic  
 2142 aerosol formation from isoprene, *Proceedings of the National Academy of Sciences*, 107, 6640-6645,  
 2143 10.1073/pnas.0911114107, 2010.  
 2144 Sutton, M. A., Howard, C. M., Erisman, J. W., Billen, G., Bleeker, A., Grenfelt, P., Grinsven, H. v., and  
 2145 Grizzetti, B.: *The European Nitrogen Assessment: Sources, Effects and Policy Perspectives*,  
 2146 Cambridge University Press, Cambridge, 664 pp., 2011.  
 2147 Thompson, A. M., Doddridge, B. G., Witte, J. C., Hudson, R. D., Luke, W. T., Johnson, J. E., Johnson, B.  
 2148 J., Oltmans, S. J., and Weller, R.: A tropical atlantic paradox: Shipboard and satellite views of a  
 2149 tropospheric ozone maximum and wave-one in January-February 1999, *Geophysical Research*  
 2150 *Letters*, 27, 3317-3320, 10.1029/1999GL011273, 2000.  
 2151 Thompson, A. M., Witte, J. C., Oltmans, S. J., Schmidlin, F. J., Logan, J. A., Fujiwara, M., Kirchhoff, V.  
 2152 W. J. H., Posny, F., Coetzee, G. J. R., Hoegger, B., Kawakami, S., Ogawa, T., Fortuin, J. P. F., and  
 2153 Kelder, H. M.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 tropical ozone  
 2154 climatology 2. Tropospheric variability and the zonal wave-one, *Journal of Geophysical Research:*  
 2155 *Atmospheres*, 108, 10.1029/2002JD002241, 2003.  
 2156 Thornton, J., and Abbatt, J. P. D.: Measurements of HO<sub>2</sub> uptake to aqueous aerosol: Mass  
 2157 accommodation coefficients and net reactive loss, *Journal of Geophysical Research D: Atmospheres*,  
 2158 110, 1-12, 10.1029/2004JD005402, 2005.  
 2159 Thouret, V., Marenco, A., Logan, J. A., Nédélec, P., and Grouhel, C.: Comparisons of ozone  
 2160 measurements from the MOZAIC airborne program and the ozone sounding network at eight  
 2161 locations, *Journal of Geophysical Research Atmospheres*, 103, 25695-25720, 10.1029/98JD02243,  
 2162 1998.  
 2163 Tingey, D. T., Manning, M., Grothaus, L. C., and Burns, W. F.: The Influence of Light and Temperature  
 2164 on Isoprene Emission Rates from Live Oak, *Physiologia Plantarum*, 47, 112-118,  
 2165 <https://doi.org/10.1111/j.1399-3054.1979.tb03200.x>, 1979.  
 2166 Tolbert, M. A., Rossi, M. J., Malhotra, R., and Golden, D. M.: Reaction of chlorine nitrate with  
 2167 hydrogen chloride and water at antarctic stratospheric temperatures, *Science*, 238, 1258-1260,  
 2168 10.1126/science.238.4831.1258, 1987.  
 2169 Troe, J.: Predictive possibilities of unimolecular rate theory, *Journal of Physical Chemistry*, 83, 114-  
 2170 126, 10.1021/j100464a019, 1979.

2171 Troe, J.: The Polanyi lecture. The colourful world of complex-forming bimolecular reactions, *Journal*  
 2172 *of the Chemical Society, Faraday Transactions*, 90, 2303-2317, 10.1039/FT9949002303, 1994.  
 2173 Trolier, M., Mauldin Iii, R. L., and Ravishankara, A. R.: Rate coefficient for the termolecular channel of  
 2174 the self-reaction of ClO, *Journal of Physical Chemistry*, 94, 4896-4907, 10.1021/j100375a027, 1990.  
 2175 Turco, R. P., Toon, O. B., Ackerman, T. P., Pollack, J. B., and Sagan, C.: Nuclear winter: Global  
 2176 consequences of multiple nuclear explosions, *Science*, 222, 1283-1292,  
 2177 10.1126/science.222.4630.1283, 1983.  
 2178 Twomey, S.: Pollution and the planetary albedo, *Atmospheric Environment* (1967), 8, 1251-1256,  
 2179 10.1016/0004-6981(74)90004-3, 1974.  
 2180 Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of the*  
 2181 *Atmospheric Sciences*, 34, 1149-1152, 10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2, 1977.  
 2182 Tyndall, J.: The Bakerian Lecture: On the Absorption and Radiation of Heat by Gases and Vapours,  
 2183 and on the Physical Connexion of Radiation, Absorption, and Conduction, *Philosophical Transactions*  
 2184 *of the Royal Society of London*, 151, 1-36, 1861.  
 2185 Urone, P., and Schroeder, W. H.: SO<sub>2</sub> in the atmosphere: A wealth of monitoring data, but few  
 2186 reaction rate studies, *Environmental Science and Technology*, 3, 436-445, 10.1021/es60028a006,  
 2187 1969.  
 2188 Vaghjiani, G. L., and Ravishankara, A. R.: New measurement of the rate coefficient for the reaction of  
 2189 OH with methane, *Nature*, 350, 406-409, 10.1038/350406a0, 1991.  
 2190 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr, A. F.:  
 2191 Interannual variability of global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*,  
 2192 6, 3423-3441, 2006.  
 2193 Van Der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C.,  
 2194 Defries, R. S., Jin, Y., and Van Leeuwen, T. T.: Global fire emissions and the contribution of  
 2195 deforestation, savanna, forest, agricultural, and peat fires (1997-2009), *Atmospheric Chemistry and*  
 2196 *Physics*, 10, 11707-11735, 10.5194/acp-10-11707-2010, 2010.  
 2197 van Donkelaar, A., Martin, R. V., and Park, R. J.: Estimating ground-level PM<sub>2.5</sub> using aerosol optical  
 2198 depth determined from satellite remote sensing, *Journal of Geophysical Research: Atmospheres*,  
 2199 111, 10.1029/2005jd006996, 2006.  
 2200 Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., Yli-Pirilä, P., Leskinen, J., Mäkelä, J. M.,  
 2201 Holopainen, J. K., Pöschl, U., Kulmala, M., Worsnop, D. R., and Laaksonen, A.: An amorphous solid  
 2202 state of biogenic secondary organic aerosol particles, *Nature*, 467, 824-827, 10.1038/nature09455,  
 2203 2010.  
 2204 Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the  
 2205 remote marine boundary layer, *Nature*, 383, 327-330, 1996.  
 2206 Volz, A., and Kley, D.: Evaluation of the Montsouris Series of Ozone Measurements Made in the  
 2207 19th-Century, *Nature*, 332, 240-242, 1988.  
 2208 von Glasow, R., and Crutzen, P. J.: Tropospheric halogen chemistry in: *Treatise on Geochemistry*,  
 2209 edited by: Holland, H. D., and Turekian, K. K., Elsevier-Pergamon, Oxford, 1 - 67, 2007.  
 2210 von Glasow, R.: Pollution meets sea salt, *Nature Geoscience*, 1, 292, 10.1038/ngeo192, 2008.  
 2211 von Schneidemesser, E., Monks, P. S., Allan, J. D., Bruhwiler, L., Forster, P., Fowler, D., Lauer, A.,  
 2212 Morgan, W. T., Paasonen, P., Righi, M., Sindelarova, K., and Sutton, M. A.: Chemistry and the  
 2213 Linkages between Air Quality and Climate Change, *Chem Rev*, 115, 3856-3897,  
 2214 10.1021/acs.chemrev.5b00089, 2015.  
 2215 Wang, C. C., Davis, L. I., Wu, C. H., Japar, S., Niki, H., and Weinstock, B.: Hydroxyl Radical  
 2216 Concentrations Measured in Ambient Air, *Science*, 189, 797-800, 10.1126/science.189.4205.797,  
 2217 1975.  
 2218 Wang, J., and Christopher, S. A.: Intercomparison between satellite-derived aerosol optical thickness  
 2219 and PM<sub>2.5</sub> mass: Implications for air quality studies, *Geophysical Research Letters*, 30,  
 2220 <https://doi.org/10.1029/2003GL018174>, 2003.

2221 Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosa-mas, C. E., Hjorth, J., LeBras, G., Moortgat,  
 2222 G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The Nitrate Radical - Physics, Chemistry,  
 2223 and the Atmosphere, *Atmospheric Environment Part a-General Topics*, 25, 1-203, 1991.  
 2224 Wayne, R. P.: *Chemistry of Atmospheres*, 3rd ed., OUP, Oxford, 2000.  
 2225 Weinstock, B.: Carbon monoxide: Residence time in the atmosphere, *Science*, 166, 224-225,  
 2226 10.1126/science.166.3902.224, 1969.  
 2227 Welz, O., Savee, J. D., Osborn, D. L., Vasu, S. S., Percival, C. J., Shallcross, D. E., and Taatjes, C. A.:  
 2228 Direct kinetic measurements of Criegee intermediate ( $\text{CH}_2\text{OO}$ ) formed by reaction of  $\text{CH}_2\text{I}$  with  $\text{O}_2$ ,  
 2229 *Science*, 335, 204-207, 2012.  
 2230 Went, F. W.: Blue Hazes in the Atmosphere, *Nature*, 187, 641-643, 10.1038/187641a0, 1960.  
 2231 Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale  
 2232 numerical models, *Atmospheric Environment* (1967), 23, 1293-1304, 10.1016/0004-6981(89)90153-  
 2233 4, 1989.  
 2234 Whitby, K. T.: The physical characteristics of sulfur aerosols, *Atmospheric Environment* (1967), 12,  
 2235 135-159, 10.1016/0004-6981(78)90196-8, 1978.  
 2236 Wild, O., Zhu, X., and Prather, M. J.: Fast-J: Accurate simulation of in- and below-cloud photolysis in  
 2237 tropospheric chemical models, *Journal of Atmospheric Chemistry*, 37, 245-282,  
 2238 10.1023/A:1006415919030, 2000.  
 2239 Williams, M.: Air pollution and policy—1952–2002, *Science of The Total Environment*, 334-335, 15-  
 2240 20, <https://doi.org/10.1016/j.scitotenv.2004.04.026>, 2004.  
 2241 WMO: WMO Global Atmosphere Watch (GAW) Implementation Plan: 2016-2023, WMO, Geneva,  
 2242 2017.  
 2243 Yienger, J. J., and Levy II, H.: Empirical model of global soil-biogenic  $\text{NO}_x$  emissions, *Journal of*  
 2244 *Geophysical Research: Atmospheres*, 100, 11447-11464, 10.1029/95jd00370, 1995.  
 2245 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A.,  
 2246 Middlebrook, A. M., Suni, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D.,  
 2247 Onasch, T., Jayne, J., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider,  
 2248 J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R.,  
 2249 Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and  
 2250 dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern  
 2251 Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, L13801, doi:10.1029/2007GL029979, 2007.  
 2252 Zhang, Y.: Online-coupled meteorology and chemistry models: history, current status, and outlook,  
 2253 *Atmos. Chem. Phys.*, 8, 2895-2932, 10.5194/acp-8-2895-2008, 2008.  
 2254