

Study of Different Carbon Bond 6 (CB6) Mechanisms by Using a Concentration Sensitivity Analysis

Le Cao, Simeng Li, and Luhang Sun

Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Nanjing University of Information Science and Technology, Nanjing, China

Correspondence: L. Cao
(le.cao@nuist.edu.cn)

Abstract. Since the year 2010, different versions of the Carbon Bond 6 (CB6) mechanism have been developed, to accurately estimate the contribution to the air pollution by the chemistry. **In order to better understand the differences in simulation results brought about by the modifications between different versions of the CB6 mechanism,** in the present study, we investigated the behavior of three different CB6 mechanisms (CB6r1, CB6r2 and CB6r3) in simulating ozone (O_3), nitrogen oxides (NO_x) and formaldehyde (HCHO) **under two different emission conditions,** by applying a concentration sensitivity analysis in a box model. The results show that when the surface emission is **weak,** the O_3 level predicted by CB6r1 is approximately 7 ppb higher than that predicted by CB6r2 and CB6r3, specifically due to the change in the sink of **acyl peroxy radicals with high order carbons (i.e. species CXO3) in the mechanism and the difference in the ozone dependence on the isoprene emission.** In contrast, **although CB6r1 estimates slightly higher values of NO_x and HCHO than the other two mechanisms at an early stage of the simulation,** the levels of NO_x and HCHO estimated by these three CB6 mechanisms **at the end of the 7-day simulation** are mostly similar, when the surface emission is **weak.** After **the increase of** the surface emission, the simulated profiles of O_3 , NO_x and HCHO obtained by CB6r2 and CB6r3 were found nearly the same **during the simulation period, but CB6r1 tends to estimate substantially higher values than CB6r2 and CB6r3.** The deviation between the O_3 levels provided by CB6r1 and the other two CB6 mechanisms (i.e. CB6r2 and CB6r3) was found enlarged **compared with the weak-emission scenario,** because of the **weaker** dependence of ozone on the emission of isoprene in CB6r1 **than those in CB6r2 and CB6r3 in this scenario.** Moreover, HCHO predicted by CB6r1 was found larger than that predicted by CB6r2 and CB6r3, which is caused by an enhanced dependence of HCHO on the emission of isoprene in CB6r1. Regarding to NO_x , **it was found that CB6r1 gives a higher value than the other two mechanisms, which is caused by the relatively stronger connection between the NO_x prediction and the release of NO and NO_2 in CB6r1, due to the change in the product of the reaction between isoprene and NO_3 in CB6r1.** Consequently, more emitted NO_x is involved in the reaction system denoted by CB6r1, which enables a following NO_x formation and thus a higher NO_x prediction of CB6r1.

1 Introduction

Air pollution occurs when the concentration of particles or gases in the atmosphere **is high, which brings** a harmful effect to the human beings and the environment of the earth. It was estimated that in 2007, approximately 3.45 million people were killed

25 worldwide due to the air pollution (Zhang et al., 2017a). Thus, it is needed to investigate the physicochemical properties of the
air pollution so that the formulation of the control strategy by the government can be guided.

Atmospheric transport model (ATM) is an efficient tool for revealing factors dominating the air pollution. Usually the ATM
includes a variety of processes that are responsible for the concentration change of pollutants in the atmosphere, such as the
production/consumption by the local chemistry, horizontal advection and vertical convection. By using ATMs, the contribution
30 to the concentration change of the pollutants by each process can be numerically estimated.

Gas-phase chemical reaction mechanism is an essential part of the ATM. It can transform the emissions and the chemical
reactions occurring in the atmosphere into the corresponding change of the species, which enables following computations
of the ATM. To the present, several atmospheric gas-phase chemical reaction mechanisms have been proposed, such as the
detailed Master Chemical Mechanism (MCM) (Jenkin et al., 1997, 2003, 2012, 2015; Saunders et al., 2003; Bloss et al., 2005)
35 and the global chemical transport model MOZART (Model for Ozone and Related chemical Tracers) mechanism (Emmons
et al., 2010). Among these chemical mechanisms, condensed mechanisms such as Carbon Bond Mechanisms (Gery et al., 1989;
Zaveri and Peters, 1999; Yarwood et al., 2005, 2010) and SAPRC mechanisms (Carter, 2000a,b, 2010) are widely applied in
ATMs due to their relatively small size and adequate accuracy. In these condensed mechanisms, different techniques are used
to lump volatile organic compounds (VOCs) into functional groups while the treatment of the inorganic chemistry is mostly
40 similar.

Carbon Bond Mechanism (CBM) is a kind of condensed mechanism, which lumps VOCs by chemical moiety (Gery et al.,
1989; Stockwell et al., 2019). In CBM mechanisms, the carbon bond is treated as a reaction unit, and the carbon bonds with
the same bonding state are treated as a group, while the exact location of the carbon bonds in the molecule is neglected.
CBM is conveniently implemented in the ATMs because of its small size and high accuracy in predicting the concentration
45 change of the pollutants. However, due to the lumping technique, biases are inevitably brought into computations. Thus, many
updates were made to the CBM mechanism to reduce biases, such as adding an explicit representation of species with the same
molecular type (e.g. species ALDX for higher-order aldehydes).

As mentioned above, CBM mechanism has been updated for several generations. In 1989, CB-IV was proposed by Gery
et al. (1989), and it was then widely used in many air quality models such as WRF-Chem (Grell et al., 2005) and CMAQ (Byun
50 and Schere, 2006). In 2005, based on the CB-IV mechanism, Yarwood et al. (2005) released CB05 by explicitly adding higher
order aldehydes (ALDX) and internal olefins (IOLE) into the mechanism. A large amount of smog chamber experiments were
also used to validate CB05, and it was reported that CB05 behaves better than CB-IV against the chamber data (Yarwood et al.,
2005). Later on, an update to CB05 was made by Whitten et al. (2010) by combining a new toluene mechanism with CB05,
namely CB05TU mechanism. It was proved that the CB05TU mechanism improves upon the CB05 mechanism in simulating
55 toluene related reactions (Whitten et al., 2010).

The latest version of the CBM mechanism is CB6 (Yarwood et al., 2010), because it is the 6-th generation of this mechanism
family, and it was released to deal with the tightening of the ozone standard in the US. Long-lived and relatively abundant
organic compounds formed by peroxy radical reactions (RO_2 - RO_2) are taken into account in CB6. Moreover, the isoprene
chemistry and the aromatic chemistry are extensively revised, to improve the modeling of the formation of secondary organic

60 aerosols (SOAs). It was shown that CB6 performs better in simulating the maximum value of ozone as well as the ozone formation rate compared with the CB05 mechanism (Yarwood et al., 2010). From then on, several updates were made to CB6 so that currently there are four versions of CB6 available, i.e. CB6r1, CB6r2, CB6r3 and CB6r4. In CB6r1, the mechanism previously proposed by Yarwood et al. (2010) was revised again (Yarwood et al., 2012), and several reactions and products were corrected. New experimental data (EUPHORE experiments) were also adopted to validate the CB6r1 mechanism (Yarwood et al., 2012). After that, in 2013, Ruiz Hildebrandt and Yarwood (2013) included the interactions between organic aerosols and total reactive nitrogen (NO_y) in the mechanism, and then gave the CB6r2 mechanism. In CB6r2, organic nitrates were divided into two groups, simple alkyl nitrates (NTR1) that remain in the gas phase and multi-functional nitrates (NTR2) that can partition into organic aerosols. Because of the inclusion of the multi-functional aerosol nitrates (i.e. NTR2), lower recycling efficiency of nitrogen oxides (NO_x) from nitrates is acquired using CB6r2, leading to a lower ozone production relative to CB6r1 (Ruiz Hildebrandt and Yarwood, 2013). The third version of CB6 is CB6r3 (Emery et al., 2015), which was developed to account for the influence of the low temperature on the formation of organic nitrates. It aims at modeling the winter ozone formation event in Uinta Basin in the US under cold conditions, and it was found that the inclusion of the temperature dependence in CB6r3 would cause an ozone reduction in winter environments, due to an enhanced formation of organic nitrates (Emery et al., 2015). The latest version of the CB6 mechanism is CB6r4 (Emery et al., 2016), which was designed by combing CB6r3 with a 16-reaction skeletal iodine mechanism, to consider the ozone depletion by the iodine chemistry. It was found that CB6r2 and CB6r4 perform similarly in simulating ozone across the continental US, but CB6r4 tends to predict a lower ozone than CB6r2, possibly due to the depletion of ozone by the iodine chemistry in the marine boundary layer (Emery et al., 2016). Currently, the CB6r3 mechanism is available in the latest version of the CMAQ model (Community Multiscale Air Quality model, available at: www.epa.gov/cmaq) (Byun and Schere, 2006), while CB6r2 and CB6r4 are both included in the CAMx model (Comprehensive Air Quality Model with Extensions, available at: www.camx.com) (ENVIRON, 2015).

80 Many investigations have been made using the CB6 mechanisms. To name a few, Luecken et al. (2019) used CB6r3 to simulate ozone, oxidized nitrogen (NO_y) and hazardous air pollutants (HAPs) across the continental US. In their study, a comparison between CB6r3, CB05TU, and CB05 as well as the observational data was performed. It was shown that these chemical mechanisms behave similarly for the ozone prediction, and CB6r3 performs the best in simulating the vertical distribution of peroxyacyl nitrates. Marvin et al. (2017) used five chemical mechanisms including CB6r2 to evaluate the impact of the isoprene chemistry on the simulation of formaldehyde (HCHO) in the summertime southeast US. They also suggested a set of modifications to CB6r2 that can improve the comparison of the modeled HCHO to observations. Zhang et al. (2017b) used the CAMx model (ENVIRON, 2015) with the implementation of the CB6r2 mechanism to estimate the biogenic isoprene emissions in US by using two different emission models, BEIS (Pierce et al., 1998; Bash et al., 2016) and MEGAN (Guenther et al., 2006, 2012), and they found that the MEGAN model predicts more isoprene emissions than the BEIS model. Recently, by implementing seven different chemical mechanisms including CB6r3 into a box model constrained by the observational data, Derwent (2017) investigated the responses of the ozone production rate and the mixing ratio of hydroxyl radicals (i.e. OH) to a reduction of NO_x and VOCs in these chemical mechanisms. It was found that when the constrained values of NO_x and VOCs in the box model are reduced, different mechanisms behave differently, especially in the prediction of OH. Later,

95 Derwent (2017) used the same model to study the response of the OH mixing ratio to the representation of oxidation and degradation of VOCs in thirteen different mechanisms including CB6r3, and Derwent (2017) found that the influence brought about by aromatics such as toluene and o-xylene on the change of OH differs a lot between different chemical mechanisms.

Despite the studies mentioned above, the internal properties of these CB6 mechanisms such as the relationship between the ozone formation and the surface emissions are still not thoroughly investigated and compared. Moreover, the corresponding change brought about by the modifications between different versions of the CB6 mechanism also needs further investigation. 100 Therefore, in this study, we performed a concentration sensitivity analysis on different versions of the CB6 mechanism (CB6r1, CB6r2, and CB6r3) to see the dependence of the formation of ozone (O_3), nitrogen oxides (NO_x , $NO+NO_2$) and formaldehyde (HCHO) on each reaction of the mechanism as well as the surface emission and dry deposition. By doing that, we were able to figure out reasons causing the deviations between the results obtained by using different CB6 mechanisms. The factors 105 dominating the formation and consumption of the focused species (O_3 , NO_x and HCHO) in these mechanisms can also be revealed.

The structure of this paper is as follows. In Sect. 2, CB6 mechanisms used in this study are introduced, and the method used to analyze the mechanism as well as the governing equations are also described. Sect. 3 gives the results of the concentration sensitivity analysis and the related discussions. In Sect. 4, major conclusions achieved in this study are summarized. Future 110 work is also prospected in this section.

2 Description of the Mechanisms and the Numerical Method

In the present study, we first implemented different versions of the CB6 mechanism (i.e. CB6r1, CB6r2, and CB6r3) into a box model, KINAL (Turányi, 1990a), to simulate the temporal evolution of O_3 , NO_x and HCHO under two different emission conditions. The surface emission intensity implemented in the model was assumed weak at first, which represents an emission condition in rural regions (Saylor and Ford, 1995; Sandu et al., 1997). By doing that, chemical reactions playing an important 115 role in the change of the focused species can be indicated. Then, sensitivities of the focused species (O_3 , NO_x and HCHO) to each reaction of the mechanisms were computed, to reveal the influence brought about by the modifications between these CB6 mechanisms. Later, the surface emission was increased in the model, and a same procedure was performed on these mechanisms again, so that the behavior of these CB6 mechanisms under a typical heavily polluted condition in urban regions 120 can be investigated.

The CB6 mechanisms studied in this paper contain approximately 80 chemical species and 220 reactions. The CB6r1 version contains 80 species and 222 reactions, and the CB6r2 version contains 81 species and 215 reactions. The CB6r3 version has 82 species and 221 reactions, including reactions accounting for the temperature dependence of the alkyl nitrate formation. Complete listings of all the reactions of these mechanisms are given in Tab. A1 of the appendix. The updates in CB6r2 and 125 CB6r3 compared with their previous version are also marked in Tab. A1. Compared with CB6r1, CB6r2 divides the organic nitrates generated from alkanes, olefins, aromatics and oxygenated VOCs (i.e. the species named NTR in CB6r1) into two groups, NTR1 that exists exclusively in the gas phase and NTR2 that can partition into organic aerosols. As a result of this

speciation, in CB6r2, the organic nitrates, NTR1 and NTR2, undergo the following reactions:



130




135 Reaction (R1) denotes the photolysis of NTR1, which enables a recycling of NO_x and a following ozone formation enhancement. Reaction (R2) represents an addition reaction leading to the conversion from NTR1 to NTR2. Reaction (R3) means that the organic nitrate partitioning within the aerosols undergoes hydrolysis and forms HNO_3 . Ruiz Hildebrandt and Yarwood (2013) reported that because of this speciation, CB6r2 has a lower recycling efficiency of NO_x from organic nitrates than CB6r1. The levels of O_3 and NO_x predicted by CB6r2 are thus lower than those predicted by CB6r1.

Regarding to CB6r3, it decomposes the formation process of alkyl nitrates from alkanes in CB6r2:



140



145 into seven reactions:



150





155



By making this modification, the dependence of the alkyl nitrate yield on the pressure and the temperature can be considered in CB6r3, especially under cold conditions. For this purpose, two new operators, XPRP and XPAP, were also added. Under a standard condition (pressure: 1 atm, temperature: 298 K), the formation of the alkyl nitrates (NTR1 and NTR2) in CB6r3 through Reactions (R7)-(R13) is equal to that in CB6r2 through Reactions (R4)-(R6) (Emery et al., 2015).

CB6r4 improves upon CB6r3 by adding a condensed iodine mechanism to consider the iodine-induced ozone destruction (Emery et al., 2016). However, CB6r4 was not investigated in this study, because the halogen chemistry is not the focus of the present study. A comparison between CB6r4 and other CB6 mechanisms in a halogen-rich environment is attributed to a future work.

We implemented the CB6 mechanisms mentioned above into a box model, KINAL (Turányi, 1990a), to capture the time variations of O_3 , NO_x and HCHO by solving Eq. (1):

$$\frac{dc}{dt} = f(c, k) + E - D. \quad (1)$$

In Eq. (1), c is a column vector of species concentrations. k is a vector of reaction rate constants and t denotes time. E represents a source term of the local surface emission, and in the present model the surface emission is parameterized as a group of reactions having products and a constant reaction rate but without reactants. D in Eq. (1) is a loss term representing the dry deposition process of atmospheric constituents, and this process is parameterized in the model as a series of reactions having reactants but without forming any product. KINAL is a box model provided for the analysis of complex reaction systems. Stiff kinetic differential equations can be solved in KINAL, and it was proved that KINAL performs robustly and efficiently (Turányi, 1990a,b; Cao et al., 2014, 2016, 2019). A background air composition (see Tab. 1), adapted from Saylor and Ford (1995) and Sandu et al. (1997), was used as the initial condition of the model. This air composition represents a heavily polluted atmosphere, in which the background level of NO_x is in the order of 1-100 ppb. Two different scenarios, “weak emission” and “strong emission”, were simulated, and the emission intensities belonging to these two scenarios are

180 listed in Tab. 1. They denote typical emission conditions in rural regions and urban regions, respectively (Saylor and Ford, 1995; Sandu et al., 1997). A 7-day simulation was performed and the simulation starts at noon (12:00) of the first day. The time variations of O₃, NO_x and HCHO were recorded every hour during the simulated period.

After obtaining the temporal evolution of O₃, NO_x and HCHO, relative concentration sensitivities of these species to different CB6 mechanisms were computed to reveal the dependence of these species on each reaction of the mechanism, surface emissions, and the rate of dry deposition for each atmospheric constituent. The relative concentration sensitivity \tilde{S}_{ij} can be expressed as

$$\tilde{S}_{ij} = \frac{\partial \ln c_i}{\partial \ln k_j} = \frac{k_j}{c_i} \frac{\partial c_i}{\partial k_j} = \frac{k_j}{c_i} S_{ij}, \quad (2)$$

which shows the importance of the j -th reaction for the concentration change of the i -th chemical species. In Eq. (2), c_i is the concentration of the i -th chemical species, and k_j denotes the rate constant of the j -th reaction. $S_{ij} = \partial c_i / \partial k_j$ is the absolute concentration sensitivity, and the unit of S_{ij} depends on the order of the j -th reaction. In order to compare the sensitivity coefficients belonging to different reactions, S_{ij} is normalized by being multiplied with k_j/c_i so that a dimensionless sensitivity coefficient, \tilde{S}_{ij} , is obtained. The relative concentration sensitivity \tilde{S}_{ij} thus represents the percentage change in the i -th species concentration due to a small perturbation in the rate of the j -th reaction. The evaluation of the concentration sensitivity is helpful for discovering the interdependence between the solution of Eq. (1) and input parameters of the model such as the reaction rate constants and the intensity of the surface emission.

The reaction rate constants of the mechanisms were taken from IUPAC database (Atkinson et al., 2004, 2006, 2007, 2008; Crowley et al., 2010; Ammann et al., 2013) and NASA/JPL database (Sander et al., 2006), and a constant temperature 298 K was assumed for the calculation of the reaction rates. Photolytic reaction rates were estimated by using TUV (Tropospheric Ultraviolet and Visible) radiation model (Madronich and Flocke, 1997, 1999), assuming a 300 Dobson overhead ozone column and a 1 km measuring height. Data of cross section and quantum yield for each photolyzed species were taken from CMAQ model version 5.3 (Byun and Schere, 2006). When the local time resides between 4:30 (sunrise) and 19:30 (sunset), the photolytic reaction rates vary with the solar zenith angle (SZA), while the photolytic reactions are switched off if the local time is out of this range. With respect to the dry deposition process, a first-order rate coefficient (k_d) indicating the loss caused by dry deposition is calculated using the following equation:

$$205 \quad k_d = v_d/L, \quad (3)$$

in which v_d denotes the dry deposition velocity, and the values of v_d used in the present study for different atmospheric constituents are given in Tab. 2. L in Eq. (3) is the boundary layer height, and is assumed as 1 km in the model.

In the following section, computational results are presented and discussed.

3 Results and Discussions

210 We first show the temporal evolution of O₃, NO_x and HCHO obtained by using CB6r1, CB6r2, and CB6r3 under the given initial condition (see Tab. 1), applying a weak surface emission. The differences between the results using different mechanisms are

also analyzed. Then the concentration sensitivities of the focused species (O_3 , NO_x and HCHO) to different CB6 mechanisms are displayed, to indicate the internal difference between these mechanisms. Later on, results with the implementation of a strong surface emission are shown. By doing that, the dependence of different CB6 mechanisms on the surface emission under a typical heavily polluted condition in urban regions can be compared and investigated.

3.1 Temporal evolution of O_3 , NO_x and HCHO (weak emission)

Figure 1 shows the temporal profiles of O_3 , NO_x and HCHO predicted by CB6r1, CB6r2 and CB6r3 in the weak emission scenario. Due to the small intensity of the surface emission in this scenario, the differences between these predictions are able to reflect different capabilities of these mechanisms in converting the initial concentrations into the change of the species. It is seen that under this condition, ozone profiles simulated by these three mechanisms show a notable deviation (see Fig. 1a). In most of the simulated period, the ozone mixing ratio predicted by CB6r1 is higher than those predicted by CB6r2 and CB6r3. Figure 1(a) shows that at the beginning of the simulation (before day 1.5), CB6r1 and CB6r3 behave similarly, while CB6r2 predicts a higher ozone. However, as the reaction proceeds (after day 1.5), CB6r2 starts to predict a lower ozone than CB6r1, and the simulated profile of CB6r2 approaches to that obtained by CB6r3. In contrast, the ozone predicted by CB6r1 becomes higher than those predicted by CB6r2 and CB6r3. During the end of the 7-day simulation, the daily averaged ozone predicted by CB6r1 over the 7-th day is approximately 7 ppb higher than those predicted by CB6r2 and CB6r3, and the ozone levels given by CB6r2 and CB6r3 are almost identical.

For NO_x simulations, Fig. 1(b) shows that ~~these three mechanisms give almost identical results during the end of the computation.~~ at the beginning of the simulation, NO_x declines rapidly from the initial value (7 ppb) to less than 1 ppb, due to the conversion to PAN (peroxyacetyl nitrate) and HNO_3 . At the end of the simulation, the mixing ratio of NO_x becomes lower than 0.5 ppb. The major nitrogen containing compound during this time period is HNO_3 , as PAN is thermally decomposed and photolyzed during the daytime. In the comparison of temporal profiles belonging to different mechanisms, CB6r1 consistently estimates a slightly higher NO_x than the other two mechanisms, but the deviation between the estimations of these mechanisms becomes smaller approaching the end of the 7-day simulation. However, it should be noted that the difference in the predicted NO_x using different mechanisms may become larger when the surface emission intensity increases, due to different capability in transforming emissions into the change of the species for each mechanism.

With respect to HCHO predictions, it is seen in Fig. 1(c) that the deviation between the results of these three mechanisms is more pronounced at the start stage of the simulation. During this time period, CB6r1 predicts a much higher HCHO than the other two mechanisms, especially at noon of every day. However, at the end of the simulation, although CB6r1 still estimates a higher HCHO than CB6r2 and CB6r3, the difference becomes smaller, and CB6r2 and CB6r3 give a similar HCHO. ~~In model simulations, usually a spin-up process is used to minimize the uncertainty induced by the implementation of an inaccurate initial condition. Based on our results, we found that if HCHO is focused, the spin-up process is also capable of reducing the deviation between the HCHO predictions by different CB6 mechanisms, when the surface emission can be neglected.~~

In summary, we found that when the surface emission is weak, the ozone concentration predicted by CB6r1 is mostly higher than that obtained by using CB6r2 or CB6r3. When the end of the simulation approaches, ozone simulated by CB6r1

is approximately 7 ppb larger than those simulated by CB6r2 and CB6r3. In contrast to that, after a 7-day computation, the NO_x and HCHO levels obtained by using these three CB6 mechanisms are more similar. At the beginning of the simulation, CB6r1 gives significantly higher values of NO_x and HCHO than the other two CB6 mechanisms. However, when the end of the simulation comes, the difference tends to disappear.

250 3.2 Concentration sensitivity analysis of different CB6 mechanisms (weak emission)

We then conducted a concentration sensitivity analysis on different CB6 mechanisms under the weak emission condition, and from these results we were able to identify the relative importance of each reaction in these mechanisms for the change of the focused species, and discover the reasons causing the deviations between the simulation results of different CB6 mechanisms.

The ozone sensitivity to the CB6r3 mechanism averaged over the last day of the computation is shown in Fig. 2. A positive sensitivity means that an increase of the reaction rate would accelerate the formation of O_3 , while a negative value denotes a decline of O_3 when the reaction rate increases. Note that Reactions (R222)-(R234) in Fig. 2(d) represent surface emissions belonging to different chemical species, and Reactions (R235)-(R243) denote dry depositions for different atmospheric constituents. It is seen from Fig. 2 that apart from the reactions standing for surface emissions and dry depositions, chemical reactions with large sensitivities mostly possess a reaction number less than 52 (i.e. before Reaction (R52) in the mechanism). Because Reactions (R1)-(R52) in the mechanism represent the inorganic chemistry while reactions after (R52) are mostly VOC-involved reactions (see Tab. A1 in the appendix), it demonstrates an important role of the inorganic chemistry in this simulation, possibly due to the high initial value of NO_x and the weak VOC emissions in this scenario. From the sensitivity analysis of CB6r3 shown in Fig. 2, we were also able to figure out the most important reactions for the change of ozone, which can be divided into two groups. The first reaction group includes Reactions (R1) $\text{NO}_2 + h\nu \rightarrow \text{NO} + \text{O}$, (R3) $\text{O}_3 + \text{NO} \rightarrow \text{NO}_2$, (R25) $\text{HO}_2 + \text{NO} \rightarrow \text{OH} + \text{NO}_2$, (R26) $\text{NO}_2 + \text{O}_3 \rightarrow \text{NO}_3$ and (R45) $\text{NO}_2 + \text{OH} \rightarrow \text{HNO}_3$, which are reactions denoting the inter-conversion of NO_x and the loss of NO_x through the formation of NO_3 and HNO_3 . It demonstrates the significance of reactive nitrogen oxides in determining the final ozone level. The other important reaction group includes (R9) $\text{O}_3 + h\nu \rightarrow \text{O}(^1\text{D})$, (R10) $\text{O}(^1\text{D}) + \text{M} \rightarrow \text{O} + \text{M}$, (R11) $\text{O}(^1\text{D}) + \text{H}_2\text{O} \rightarrow 2\text{OH}$, and (R13) $\text{O}_3 + \text{HO}_2 \rightarrow \text{OH}$. These reactions represent the ozone loss due to the formation of hydroxyl radicals (i.e. OH). With respect to the other chemical reactions in the mechanism, their sensitivities are all smaller than 0.1 (see Figs. 2b, c and d), denoting a minor influence on the change of ozone by these chemical reactions.

Regarding to the ozone sensitivities to surface emissions (i.e. Reactions (R222)-(R234) in Fig. 2d) and dry depositions (i.e. Reactions (R235)-(R243) in Fig. 2d), it is seen that although the implemented surface emission in this scenario is weak, the emission still exerts a strong influence on the change of ozone, indicated by relatively large sensitivity coefficients belonging to NO and NO_2 emissions (~ 0.3 , see Fig. 2d). The values of the ozone sensitivities to the NO and NO_2 emissions are comparable to those corresponding to chemical reactions denoting the inter-conversion of NO_x (i.e. (R1) and (R3) in Fig. 2a). In contrast to that, although the emission intensity of isoprene in this scenario is relatively large, the dependence of ozone on the isoprene emission is minor, reflected by the small ozone sensitivity to the isoprene emission (i.e. Reaction (R232) in Fig. 2d). In addition,

it was also found in Fig. 2(d) that dry deposition is the largest sink of ozone in this weak emission scenario, when CB6r3 is implemented.

We then computed the ozone sensitivities to the other two mechanisms (i.e. CB6r2 and CB6r1). Because these figures are similar to Fig. 2, we show these results in the supplementary material of the manuscript (Figs. S1 and S2). The similarity between these figures also denotes a consistent treatment of the inorganic chemistry and a similar lumping technique of VOCs in these CB6 mechanisms. By comparing Fig. S1 with Fig. 2, we found that the averaged ozone sensitivity to CB6r2 over the last day is almost identical to the sensitivity to CB6r3, thus leading to a similar ozone prediction by these two mechanisms, which has been shown before (see Fig. 1a). The only major difference between the ozone sensitivities to CB6r2 and CB6r3 is that CB6r3 improves upon CB6r2 by adding Reactions (R217)-(R220) to include the temperature dependence of the alkyl nitrate formation. Therefore, the sensitivities of these reactions are absent for CB6r2, shown in Fig. S1 of the supplements. However, as mentioned above, under a condition of a 298 K temperature and 1 atm pressure, the formation of the alkyl nitrate in CB6r2 and CB6r3 are equivalent. Thus, the addition of these reactions in the CB6r3 mechanism would not significantly affect the predicted ozone in this scenario. However, at a different temperature, this update in CB6r3 may bring a large change in ozone, indicated by the relatively large sensitivities of ozone to Reactions (R219) $\text{XPAR} \rightarrow \text{XO}_2\text{N} + \text{RO}_2$ and (R220) $\text{XPAR} \rightarrow 0.13\text{ALDX} + 0.87\text{ROR} + 0.13\text{XO}_2\text{H} + 0.87\text{XO}_2 + \text{RO}_2 - 0.13\text{PAR}$, shown in Fig. 2(d).

We then tried to figure out reactions causing the higher ozone prediction of CB6r1. From a comparison between Fig. S1 and Fig. S2 in the supplementary material, we found two factors heavily responsible for the higher ozone prediction of CB6r1. One is the modification of Reaction (R66) about the sink of CXO3 in CB6r1. CXO3 represents acylperoxy radicals with three and higher carbons, and is able to oxidize NO and thus form ozone. In CB6r1, the form of Reaction (R66) is as follows, $\text{CXO}_3 + \text{RO}_2 \rightarrow \text{CXO}_3$, while in CB6r2, the form is $\text{CXO}_3 + \text{RO}_2 \rightarrow 0.8\text{ALD}_2 + 0.8\text{XO}_2\text{H} + 0.8\text{RO}_2$. In CB6r1, the total amount of CXO3 is unaltered through Reaction (R66), thus leading to a negligible ozone sensitivity to (R66). However, the update of Reaction (R66) in CB6r2 causes this reaction to be a major sink of CXO3 in the mechanism. As a result, the ozone significance of Reaction (R66) increases in CB6r2. Moreover, due to this enhanced importance of Reaction (R66), it was found in CB6r2 that the significance of many other CXO3 related reactions, e.g. (R62) $\text{CXO}_3 + \text{NO}_2 \rightarrow \text{PANX}$, (R63) $\text{PANX} \rightarrow \text{NO}_2 + \text{CXO}_3$, (R65) $\text{CXO}_3 + \text{HO}_2 \rightarrow 0.41\text{PACD} + 0.15\text{AACD} + 0.15\text{O}_3 + 0.44\text{ALD}_2 + 0.44\text{XO}_2\text{H} + 0.44\text{RO}_2 + 0.44\text{OH}$, (R67) $\text{CXO}_3 + \text{CXO}_3 \rightarrow 2\text{ALD}_2 + 2\text{XO}_2\text{H} + 2\text{RO}_2$, and (R110) $\text{ALDX} + \text{OH} \rightarrow \text{CXO}_3$ drops, from a moderate value in CB6r1 to a small value in CB6r2. The formation of ozone in CB6r2 is thus getting suppressed due to the additional consumption of CXO3 through Reaction (R66). This finding also denotes an important role of CXO3 in determining ozone in the CB6 mechanisms. Thus, more attention should be paid to CXO3 related reactions in future mechanism developments. The significance of CXO3 in the mechanism for the conversion of NO to NO₂ and the formation of ozone has also been identified by Luecken et al. (2008) in a model study on the behavior of three chemical mechanisms including CB-IV, CB05 and SAPRC99. Aside from the change of Reaction (R66), we also found the ozone sensitivity to the isoprene emission shifts from a small value in CB6r2 and CB6r3 to a moderate positive value in CB6r1 (see Fig. S2(d)). Due to this sensitivity shift, when CB6r1 is used in the model, an enhanced emission of isoprene in this scenario would result in a significant elevation of the predicted ozone. The reason for this sensitivity shift is attributed to the modification of Reaction (R157) between CB6r1

and the other two mechanisms. In CB6r1, Reaction (R157) is in the form of $\text{ISOP} + \text{NO}_3 \rightarrow 0.65 \text{ INTR} + \text{other products}$,
315 so that the emitted isoprene is partly converted to INTR (i.e. organic nitrates from isoprene reactions), while in CB6r2 and
CB6r3 the product INTR is updated as NTR2. The product INTR in CB6r1 is able to react with OH, forming many organic
compounds such as NTR, ALD2 and ALDX that can promote the ozone formation. In contrast, NTR2, the multi-functional
nitrate formed through Reaction (R157) in CB6r2 and CB6r3, is then converted to HNO_3 , which is relatively inactive for the
ozone formation. As a result, the influence caused by the isoprene emission on the change of ozone in CB6r2 and CB6r3 is
320 weaker than that in CB6r1, which is also confirmed by the smaller ozone sensitivity to Reaction (R157) in CB6r2 and CB6r3.
Moreover, because of the enhanced ozone dependence on the isoprene emission in CB6r1, the ozone sensitivities to many
reactions that are associated with isoprene and the products of isoprene reactions, e.g. (R156) $\text{ISOP} + \text{O}_3 \rightarrow \text{products}$, (R92)
 $\text{NTR} + h\nu \rightarrow \text{products}$, (R106) $\text{ALD2} + \text{OH} \rightarrow \text{C2O3}$, and (R110) $\text{ALDX} + \text{OH} \rightarrow \text{CXO3}$, in CB6r1 were also found larger
than those in CB6r2 and CB6r3 (see Fig. S2). Thus, it can be concluded that the dependence of ozone in CB6r1 on the isoprene
325 emission is different from those in CB6r2 and CB6r3, due to the change in the products of Reaction (R157), which will be
discussed further in a later context.

The NO_x sensitivity to the CB6r3 mechanism is displayed in Fig. 3. It is seen that in CB6r3, except the reactions representing
emissions and depositions, chemical reactions (R1) $\text{NO}_2 + h\nu \rightarrow \text{NO} + \text{O}$, (R3) $\text{O}_3 + \text{NO} \rightarrow \text{NO}_2$ and (R26) $\text{NO}_2 + \text{O}_3 \rightarrow$
 NO_3 , are the most determining reactions for the change of NO_x . It is because that the formation of NO_3 in the presence
330 of O_3 is a major chemical pathway for the loss of NO_x , especially in the nighttime. Other important chemical reactions for
the change of NO_x include (R9) $\text{O}_3 + h\nu \rightarrow \text{O}(^1\text{D})$, (R10) $\text{O}(^1\text{D}) + \text{M} \rightarrow \text{O} + \text{M}$, (R11) $\text{O}(^1\text{D}) + \text{H}_2\text{O} \rightarrow 2\text{OH}$, and (R25)
 $\text{HO}_2 + \text{NO} \rightarrow \text{OH} + \text{NO}_2$, which are related to the formation of hydroxyl radicals (i.e. OH). It is not surprising as the reaction
between OH and NO_2 that forms HNO_3 acts as a large sink of reactive nitrogen oxides. Regarding to the other reactions in the
mechanism, their sensitivities are much smaller, thus bringing a negligible influence on the change of NO_x . With respect to the
335 surface emissions, it can be found in Fig. 3 that the emissions of NO and NO_2 would elevate the predicted level of NO_x , which
is natural. In contrast, the dependence of NO_x on the isoprene emission is minor in this scenario using CB6r3, indicated by
the corresponding small sensitivity (see Reaction (R232) in Fig. 3d). Regarding to the reactions signifying dry depositions, it
was found in Fig. 3(d) that the dry deposition of ozone is able to strongly elevate the predicted level of NO_x . The reason is that
ozone is critical for the conversion of NO_2 to NO_3 , which is a major loss of NO_x in this scenario as discussed above. Thus, the
340 decline of ozone due to dry deposition would substantially inhibit the formation of NO_3 , thus elevating the concentration of
 NO_x .

The sensitivities of NO_x to the other two CB6 mechanisms, CB6r2 and CB6r1, are shown in Figs. S3 and S4 of the supple-
mentary material, respectively. It was found that the sensitivity of NO_x to CB6r2 shown in Fig. S3 is strongly similar to the
sensitivity to CB6r3 displayed in Fig. 3. The largest change in the NO_x sensitivity between CB6r3 and CB6r2 is the addition
345 of reactions representing the temperature dependence of alkyl nitrate formation in CB6r3, i.e. Reactions (R217)-(R220) in
Fig. 3(d). However, as mentioned above, the scheme for the temperature dependence in CB6r3 is equivalent to that in CB6r2
under the situation used in this study. Thus, adding these reactions into CB6r3 would not exert a significant influence on the
change of NO_x . But the moderate sensitivity coefficients belonging to Reactions (R219) and (R220) shown in Fig. 3(d) denote

that under a different temperature condition, the change of NO_x brought about by this update might be larger. Apart from this
350 change, other reactions that largely modified between CB6r3 and CB6r2 possess a small sensitivity coefficient. Thus, choosing
CB6r3 or CB6r2 in this weak-emission scenario would not significantly influence the predicted NO_x . However, for the CB6r1
mechanism, Fig. S4(d) shows that the NO_x sensitivity to the isoprene emission in CB6r1 is substantially larger than that in
CB6r2 or CB6r3. It indicates that the dependence of NO_x on the isoprene emission in CB6r1 is heavier than that in CB6r2 and
CB6r3. The reason is also attributed to the change of Reaction (R157), which is similar to the conclusion achieved in the ozone
355 sensitivity analysis discussed above. In CB6r1, the product INTR is generated through Reaction (R157) rather than NTR2 in
CB6r2 and CB6r3. As a result, in CB6r1, the emitted isoprene can be more converted to organic compounds such as NTR.
Then, the photolysis of NTR, i.e. Reaction (R92) $\text{NTR} + h\nu \rightarrow \text{NO}_2 + \text{XO}_2\text{H} + \text{RO}_2$, would increase the NO_x concentration,
leading to a higher NO_x prediction of CB6r1 than those of CB6r2 and CB6r3. This mechanism is also indicated by the increased
 NO_x sensitivity possessed by Reaction (R92) in CB6r1 compared to that in CB6r2 and CB6r3 (see Fig. S4).

360 At last, for the HCHO sensitivity, it is seen in Fig. 4 that in CB6r3, the largest HCHO decay pathway is the dry deposition,
denoted by its most negative sensitivity coefficient (see Fig. 4d). And another major chemical pathway for the destruction of
HCHO is the photolysis of HCHO, i.e. Reactions (R97) $\text{HCHO} + h\nu \rightarrow 2\text{HO}_2 + \text{CO}$ and (R98) $\text{HCHO} + h\nu \rightarrow \text{CO}$, both
of which possess relatively large absolute values of the HCHO sensitivity. In contrast, major HCHO formation pathways are
found including Reactions (R72) $\text{MEO}_2 + \text{HO}_2 \rightarrow 0.9\text{MEPX} + 0.1\text{HCHO}$ and (R124) $\text{CH}_4 + \text{OH} \rightarrow \text{MEO}_2 + \text{RO}_2$. It is
365 due to the large amount of CH_4 in the initial condition, which is a major source of HCHO through its oxidation. Moreover, it
was also found in Fig. 4(c) that Reaction (R156) $\text{ISOP} + \text{O}_3 \rightarrow 0.6\text{HCHO} + \text{other products}$ plays an important role in the
formation of HCHO, due to the emission of isoprene in this scenario. This strongly enhanced HCHO formation by the release
of isoprene is also demonstrated by the large positive sensitivity coefficient possessed by the isoprene emission, i.e. (R232) in
Fig. 4(d). Thus, it can be concluded that except the initial amount of CH_4 , the emission intensity of isoprene is also a critical
370 factor determining the predicted value of HCHO in this weak emission scenario.

By comparing the HCHO sensitivity to CB6r3 shown in Fig. 4 with the HCHO sensitivities to CB6r2 (Fig. S5 of the
supplements) and CB6r1 (Fig. S6 of the supplements), it was found that the largest change in the HCHO sensitivity between
CB6r3 and CB6r2 is again the addition of reactions representing the temperature dependence in CB6r3. However, similar to
the findings discussed above, the prediction of HCHO by CB6r3 is not heavily affected by the addition of these reactions under
375 a standard condition and thus is similar to that by CB6r2. Most interestingly, different from the situations in simulating ozone
and NO_x , it was found in Fig. 4(d) that the reactions representing the temperature dependence (i.e. Reactions (R217)-(R220))
possess relatively small HCHO sensitivities (<0.05). Thus, it can be expected that even under a different temperature condition,
the influence on the HCHO prediction caused by the change of the temperature is also possibly small.

With respect to the HCHO sensitivity to CB6r1 (see Fig. S6 of the supplements), some changes were found. First, different
380 from the small negative sensitivity to Reaction (R157) in CB6r2 and CB6r3, the HCHO sensitivity to Reaction (R157) in CB6r1
has a moderate positive value. We figured out that it is also caused by the difference in the product of this reaction between
CB6r1 and the other two mechanisms. Through Reaction (R157), the emitted isoprene in CB6r1 can be more conveniently
converted to active organic compounds such as NTR that can be oxidized and generate HCHO. Therefore, the predicted

HCHO in CB6r1 is higher than that in CB6r2 and CB6r3 in this scenario (shown in Fig. 1c). This conclusion is also confirmed
385 by the increased importance of Reaction (R92) $\text{NTR} + h\nu \rightarrow \text{NO}_2 + \text{XO}_2\text{H} + \text{RO}_2$ in CB6r1 for the change of HCHO (see
Fig. S6), and the elevated HCHO sensitivity to the isoprene emission in CB6r1 (~ 0.53) compared to that in CB6r2 and CB6r3
(~ 0.48). Another special finding from the analysis of the HCHO sensitivity to CB6r1 is that in CB6r1, a reverse in the signs
of HCHO sensitivities to (R1) $\text{NO}_2 + h\nu \rightarrow \text{NO} + \text{O}$ and (R3) $\text{O}_3 + \text{NO} \rightarrow \text{NO}_2$ occurs (see Fig. S6), compared to those in
CB6r2 and CB6r3. The reason is also attributed to the difference in the HCHO dependence on the isoprene emission between
390 CB6r1 and the other two mechanisms. The occurrence of Reaction (R3) is able to increase the NO_2/NO ratio, which further
promotes the formation of NO_3 . Because the reaction between NO_3 and isoprene, i.e. Reaction (R157), plays a more important
role in the prediction of HCHO in CB6r1 than in CB6r2/CB6r3 as discussed above, the occurrence of Reaction (R3) in CB6r1
can thus remarkably promote the formation of HCHO by accelerating the isoprene+ NO_3 reaction. In contrast, in CB6r2 or
CB6r3, the reaction between isoprene and NO_3 plays a relatively minor role in determining HCHO due to the product update
395 from INTR to NTR2. Therefore, instead of accelerating the HCHO formation, NO_2 formed through Reaction (R3) in CB6r2
and CB6r3 consumes OH and thus suppresses the formation of HCHO by retarding the oxidation of CH_4 . As a consequence,
the sensitivity of HCHO to Reaction (R3) becomes negative in CB6r2 and CB6r3, shown in Fig. S5 of the supplements and
Fig. 4 of the manuscript, respectively.

In general, the sensitivity analysis shows that when the surface emission is weak, One of the updates in CB6r2 and CB6r3
400 compared with CB6r1 that can strongly affect the ozone prediction is the change in the sink of CXO3, i.e. Reaction (R66).
Because of this modification, the significance of many CXO3 related reactions also changes, causing a lower ozone prediction
of CB6r2 and CB6r3 than that of CB6r1. Apart from that, the lower ozone predicted by CB6r2 and CB6r3 was also found
contributed by the weaker dependence of ozone on the isoprene emission in CB6r2 and CB6r3 than that in CB6r1, due to the
change in the product of the reaction between isoprene and NO_3 . On the contrary, the ozone sensitivities to CB6r2 and CB6r3
405 are approximately the same, thus leading to a similar O_3 prediction. With respect to NO_x and HCHO, it was found that the
difference in the product of the reaction between isoprene and NO_3 causes a stronger dependence of NO_x and HCHO on the
emission of isoprene in CB6r1 than that in CB6r2 or CB6r3. As the isoprene emission in this scenario promotes the formation
of NO_x and HCHO, the levels of NO_x and HCHO given by CB6r1 are thus slightly higher than those estimated by CB6r2 and
CB6r3. In contrast, reactions that largely modified between CB6r2 and CB6r3 mostly have small sensitivities so that these
410 updates exert a negligible impact on the predictions of NO_x and HCHO, and the predictions given by CB6r2 and CB6r3 are thus
similar. However, under a different temperature condition, the predictions of O_3 and NO_x by CB6r3 might be largely different
from those predicted by CB6r2 and CB6r1, indicated by the moderate values of the sensitivities belonging to the temperature
dependent reactions in CB6r3. In contrast, the estimation of HCHO might not be significantly affected, according to the small
HCHO sensitivities to these temperature-dependent reactions in CB6r3.

415 3.3 Temporal evolution of ozone, NO_x and HCHO (strong emission)

According to the discussions above, it is known that the difference between the estimations of atmospheric constituents using
different CB6 mechanisms is not only caused by the change in the forms of reactions between these mechanisms, but also

determined by the different dependence of the mechanism on surface emissions and dry depositions. Thus, we continued to increase the intensity of the surface emission to investigate the behavior of these CB6 mechanisms under a strong emission condition. The emission intensity for each chemical species in this strong emission scenario has been given in Tab. 1.

The temporal profiles of O_3 , NO_x and HCHO in the strong emission scenario is shown in Fig. 5. It is seen from Fig. 5(a) that after the increase of the surface emission, the ozone concentration keeps steady during the whole simulated period, instead of dropping to a low level in the weak emission scenario. The ozone level at the end of the 7-day simulation is within a range of 60-130 ppb, much higher than that in the weak emission scenario (~ 20 -40 ppb). By comparing ozone profiles obtained by using different CB6 mechanisms in Fig. 5(a), we found the ozone predictions by CB6r2 and CB6r3 approximately the same, while CB6r1 predicts a much higher value. It was calculated that the averaged ozone over the 7-th day predicted by CB6r1 is approximately 24 ppb higher than that predicted by CB6r2 or CB6r3. Thus, after the increase of the surface emission, the deviation in the predicted ozone between CB6r1 and the other two CB6 mechanisms is enlarged, compared with that in the weak-emission scenario. It demonstrates that the CB6r1 mechanism has a stronger transformation ability in converting the surface emission into the change of ozone than the other two mechanisms. As a result, simulations using CB6r1 would yield a much higher ozone than that using CB6r2 or CB6r3, even though a same intensity of the surface emission is applied, and the deviation in the predicted ozone would become larger when the applied surface emission increases in the model. In a previous regional modeling of the air quality across the continental US (Ruiz Hildebrandt and Yarwood, 2013), it was reported that CB6r1 predicts a higher ozone than CB6r2. Thus, the results of the present study are consistent with the conclusions of Ruiz Hildebrandt and Yarwood (2013).

The change of NO_x with time is displayed in Fig. 5(b). It shows that NO_x declines rapidly from the relatively high initial value (7 ppb) to a stable level, 1-2 ppb. This final value range is also much higher than that in the weak-emission scenario (< 0.5 ppb). An obvious diurnal variation of NO_x is exhibited, and a peak value was found in the early morning of each day. Figure 5(b) also shows that CB6r2 and CB6r3 give similar NO_x predictions, while CB6r1 behaves differently. CB6r1 predicts a higher NO_x than CB6r2 and CB6r3 in most of the simulated period, and the difference grows when the end of the simulation approaches. We figured out the reason for this difference is that in CB6r1, Reaction (R66) has a form of $CXO_3 + RO_2 \rightarrow CXO_3$, which has been discussed above. As a result, the amount of CXO_3 is unaltered through Reaction (R66) in CB6r1. Then CXO_3 in CB6r1 turns to be more involved in reactions forming PANX (peroxyacyl nitrate with three and higher carbons). This trend is also confirmed by the relatively higher amount of PANX predicted by CB6r1 than those predicted by CB6r2 and CB6r3 (not shown here). As a result, during the nighttime, an additional amount of NO_x in CB6r1 is converted to PANX, causing a lower prediction of NO_x in CB6r1. In contrast, when the sun rises, the photolysis of PANX leads to a more rapid increase of NO_x in CB6r1, compared with that in CB6r2 and CB6r3.

With respect to HCHO, we found that due to the inclusion of the strong surface emission in this scenario, the level of HCHO keeps increasing when the simulation proceeds. The emitted species that are responsible for the enhancement of HCHO will be investigated in a later context. The temporal change of HCHO shows a strong diurnal variation, in which it peaks in the afternoon and reaches the trough in the early morning of every day. The predicted HCHO profiles using CB6r2 and CB6r3 are found almost identical. In a box model study of Marvin et al. (2017), they also found that using CB6r3 causes a negligible

impact (<1%) on the simulated HCHO compared to using CB6r2, which is consistent with the findings of the present study. In contrast to that, CB6r1 consistently yields a higher value of HCHO than CB6r2 and CB6r3, and the deviation is more pronounced during the daytime. At the 7-th day of the simulation, the peak value of HCHO at noon obtained by CB6r1 is around 14-15 ppb, while the lowest value is approximately 7 ppb. The deviation between the peak values of HCHO predicted by CB6r1 and the other two mechanisms is approximately 2.5 ppb. These values are all much higher than those in the weak-emission scenario (see Fig. 1c), due to the inclusion of the stronger surface emission.

In summary, due to the inclusion of a stronger surface emission, an enhancement of the predicted O₃, NO_x and HCHO was found, compared with the weak-emission scenario. Moreover, simulated results of CB6r2 and CB6r3 are almost identical, while CB6r1 consistently gives higher values of O₃, NO_x and HCHO than the other two mechanisms. Moreover, the deviations between the estimations by CB6r1 and the other two CB6 mechanisms in this scenario were found enlarged compared to those in the weak-emission scenario, reflecting a stronger transformation ability of CB6r1 converting the surface emissions into the change of atmospheric constituents. The most special finding in this scenario is the prediction of NO_x, as the daytime level of NO_x predicted by CB6r1 is higher than that predicted by CB6r2 and CB6r3, but the nighttime NO_x given by CB6r1 is lower than that given by the other two mechanisms. It is because that NO_x in CB6r1 participate more frequently in the formation of PANX, leading to a larger NO_x consumption in the nighttime and a more rapid NO_x formation in the daytime.

3.4 Concentration sensitivity analysis of different CB6 mechanisms (strong emission)

The concentration sensitivity analysis was applied on these CB6 mechanisms again, after implementing the strong surface emission. Figure 6 shows the ozone sensitivity to the CB6r3 mechanism. Note that Reactions (R222)-(R234) in Fig. 6(d) represent the sensitivities to the surface emissions. From a global view, it can be found that after increasing the surface emission, the importance of many reactions in the mechanism increases, compared with the weak-emission case. Reactions that the significance changes the most are (1) NO_x related reactions: (R1) NO₂ + hν → NO + O and (R3) O₃ + NO → NO₂; (2) terminal olefins (OLE) related reactions: (R142) OLE + OH → 0.78HCHO + 0.49ALD2 + 0.49ALDX + 0.98XO2H + 0.2XO2 + 0.02XO2N + 1.2RO2 - 0.73PAR and (R143) OLE + O₃ → 0.29ALD2 + 0.56HCHO + 0.27ALDX + 0.15XO2H + 0.15RO2 + 0.33OH + 0.08HO₂ + 0.38CO + 0.07GLY + 0.07MGLY + 0.09FACD + 0.13AACD + 0.04H₂O₂ - 0.79PAR; (3) isoprene related reactions: (R149) ISOP + OH → ISO2 + RO2 and (R157) ISOP + NO₃ → 0.35NO₂ + 0.65NTR2 + 0.64XO2H + 0.33XO2 + 0.03XO2N + RO2 + 0.35HCHO + 0.35ISPD. Moreover, Fig. 6(d) shows that the surface emissions of many chemical species exert a strong influence on the change of ozone, reflected by the relatively large ozone sensitivities to Reactions (R222)-(R234). It can be seen in Fig. 6 that the ozone sensitivities to the surface emissions are comparable to those belonging to the NO_x related reactions (i.e. (R1) and (R3)). In addition, similar to the weak-emission scenario, dry deposition, denoted by Reaction (R233) in Fig. 6(d), acts as a major loss pathway of ozone, indicated by the large absolute value of its ozone sensitivity.

Among the surface emissions, the most influential emitted species for the change of O₃ is NO (see Fig. 6d). It is not surprising as the oxidation of the emitted NO by the hydroperoxy radical and methylperoxy radicals would form O₃. Aside from this, the release of NO₂ would also increase the O₃ level, through its photolytic decomposition. In contrast to the NO_x emissions, the increase of VOC emissions in this scenario would decrease the formation of O₃, which is indicated in Fig. 6(d) by the negative

ozone sensitivities to the VOC emissions. Moreover, it was found that the ozone sensitivity to the emissions of NO_x is larger than that to the VOC emissions. This trend has also been revealed by Luecken et al. (2018), showing that the dependence of ozone on NO_x is approximately three times as heavy as that on hydrocarbons in their model studies. In the present study, the most influential VOC for the change of O_3 is found as ISOP (isoprene). Isoprene can react rapidly with OH and NO_3 , which substantially contributes to the formation and consumption of ozone. Thus, among the emissions of VOCs, more attention should be paid to the isoprene emission applied in air quality models in order to achieve a more accurate ozone prediction. In previous studies, it was shown that different biogenic emission models (e.g. MEGAN and BEIS) may yield different isoprene emission estimates (Bash et al., 2016; Zhang et al., 2017b). Thus, the choice of the biogenic emission model in the settings of the air quality model would strongly influence the modeled ozone, according to the findings of the present study. Moreover, an enhancement of the emissions of terminal olefins (OLE), toluene (TOL), xylene and other polyalkyl aromatics (XYL) would also reduce the ozone level. From the response of the ozone concentration to the surface emission, we concluded that in this scenario, an emission control of NO_x especially NO is effective in reducing O_3 , while an emission control of VOCs leads to an increase of ozone, when CB6r3 is implemented.

The ozone sensitivities to the reactions of CB6r2 and CB6r1 as well as the surface emissions and dry depositions are shown in Figs. S7 and S8 of the supplementary material. First, by comparing Fig. 6 with Fig. S7 in the supplements, we found that after the inclusion of the strong surface emission, the ozone sensitivities to CB6r3 and CB6r2 are still approximately the same. ~~Although the update of Reaction (R158) mentioned above between CB6r2 and CB6r3 still causes a sensitivity shift from a positive value to a negative value, the impact caused by this sensitivity shift is negligible compared with the influence brought about by the inclusion of the surface emission.~~ As a result, under this condition, O_3 predicted by CB6r3 is almost equal to that predicted by CB6r2, which has been displayed in Fig 5(a). For the CB6r1 mechanism, the ozone sensitivity displayed in Fig. S8 of the supplements shows a remarkable difference, especially in the dependence of O_3 on the surface emission. It was found that in this strong-emission scenario, the dependence of O_3 on the emission of isoprene is weaker in CB6r1 than that in CB6r2 and CB6r3 (see the sensitivity to Reaction (R233) in Fig. S8 of the supplements). On the contrary, the ozone sensitivities to the emissions of other hydrocarbons in CB6r1 mostly keep unchanged. Thus, in CB6r1, the emitted VOC that O_3 depends on the most becomes XYL (xylene and other polyalkyl aromatics) instead of ISOP (isoprene). As a result, the O_3 destruction caused by the isoprene emission is suppressed in CB6r1. The importance of many isoprene related reactions, such as Reaction (R149) $\text{ISOP} + \text{OH} \rightarrow \text{ISO}_2 + \text{RO}_2$ also becomes weaker in CB6r1 than that in CB6r2 and CB6r3. The reason for the less dependence of ozone on the isoprene emission in CB6r1 under this strong emission condition is again attributed to the change in the product of Reaction (R157), i.e. $\text{ISOP} + \text{NO}_3$. As discussed above in the weak-emission scenario, Reaction (R157) in CB6r1 is able to promote the formation of ozone by converting the emitted isoprene into organic nitrates such as INTR and NTR, which can be more conveniently recycled than the product NTR2 in CB6r2 and CB6r3. Because in this strong-emission scenario, the release of VOCs tends to suppress the ozone formation, the positive influence of the isoprene emission on the change of ozone brought by Reaction (R157) in CB6r1 can thus offset the negative dependence of ozone on the emission of isoprene, leading to a weaker dependence of ozone in CB6r1. This positive influence on the formation of ozone caused by Reaction (R157) in CB6r1 is also reflected by the shift of the ozone sensitivity to Reaction (R157) from negative in CB6r2 and CB6r3 to positive in CB6r1 (see

Fig. S8), as well as the relatively large sensitivities possessed by INTR and NTR reactions, i.e. (R170) $\text{INTR} + \text{OH}$ and (R92) $\text{NTR} + h\nu$ in CB6r1. Therefore, under the condition of a strong surface emission, due to the less dependence of O_3 on the emitted isoprene, the O_3 level predicted by CB6r1 is higher than that predicted by CB6r2 and CB6r3, which has been shown in Fig 5(a). In the weak-emission scenario shown before, we found the modification in Reaction (R66) about the sink of CXO_3 to be another major factor causing the difference between the simulation results of CB6r1 and CB6r2/CB6r3. However, it can be seen in Fig. S8 that after increasing the surface emission, the ozone sensitivity to Reaction (R66) is negligible, compared to the sensitivity to the surface emission. Therefore, the difference in the predicted ozone between different CB6 mechanisms is mostly caused by the change of the O_3 dependence on the surface emission, especially the release of isoprene.

The sensitivities of NO_x to the reactions of the CB6r3 mechanism as well as the surface emissions and dry depositions are displayed in Fig. 7. It can be seen that after the increase of the surface emission, the most dominant factor for the change of NO_x is still the direct emission of NO_x (i.e. Reactions (R222) and (R223) in Fig. 7d). Moreover, due to the enhanced NO_x emission in this scenario, the significance of Reactions (R1) $\text{NO}_2 + h\nu \rightarrow \text{NO} + \text{O}$, (R3) $\text{O}_3 + \text{NO} \rightarrow \text{NO}_2$ and (R26) $\text{NO}_2 + \text{O}_3 \rightarrow \text{NO}_3$, which represent the inter-conversion of reactive nitrogen oxides and the loss of NO_x due to the formation of NO_3 , increases, compared with that in the weak-emission scenario. In contrast, for the reactions associated with the formation of OH: (R9) $\text{O}_3 + h\nu \rightarrow \text{O}(^1\text{D})$, (R10) $\text{O}(^1\text{D}) + \text{M} \rightarrow \text{O} + \text{M}$, (R11) $\text{O}(^1\text{D}) + \text{H}_2\text{O} \rightarrow 2\text{OH}$, and (R25) $\text{HO}_2 + \text{NO} \rightarrow \text{OH} + \text{NO}_2$, which used to be important in the weak-emission scenario, their NO_x sensitivities decrease. It is because that after the enhancement of the surface emission, the ozone level is elevated, so that NO_2 is more involved in Reaction (R26) $\text{NO}_2 + \text{O}_3 \rightarrow \text{NO}_3$. As a result, the importance of the reaction between NO_2 and OH for the change of NO_x drops, leading to a less dependence of NO_x on the formation of OH under this situation.

In this strong-emission scenario using CB6r3, the emitted species that NO_x depends on the most is NO (see Fig. 7d), which is natural as the direct emission of NO would strongly increase NO_x . The release of NO_2 also elevates the NO_x level predicted by CB6r3. In contrast to that, the release of different VOCs exerts a different influence on the concentration change of NO_x . The emissions of ISO (isoprene) and OLE (terminal olefins) would increase the simulated level of NO_x , while the emissions of TOL (toluene) and XYL (xylene and other polyalkyl aromatics) decrease it. Figure. 7(d) also shows that the dependence of NO_x on the emissions of VOCs is remarkably lower than that on the direct emissions of NO_x , which is similar to the conclusion achieved in the ozone sensitivity analysis. tends to suppress the formation of NO_x , denoted by the negative sensitivities to VOC emissions. The most influential VOC for the change of NO_x is also the species ISOP (isoprene), identifying the important role of isoprene in the CB6r3 mechanism. In addition, Fig. 7(d) shows that in this scenario, the dry deposition of ozone still plays the role of enhancing the formation of NO_x as that in the weak-emission scenario, due to the deceleration of NO_3 formation by the decline of ozone.

From a comparison between the NO_x sensitivities to CB6r2 (shown in Fig. S9 of the supplements) and CB6r3 (see Fig. 7), again we found that the NO_x sensitivity to CB6r2 is almost identical to that to CB6r3, thus leading to a similar prediction of NO_x by these two mechanisms. However, for CB6r1 (see Fig. S10 of the supplements), the sensitivities of NO_x to the surface emissions of NO and NO_2 (0.84 and 0.13) are relatively larger than those in CB6r2/CB6r3 (0.76-0.79 and 0.11). The reason for the heavier dependence of NO_x on the surface emissions in CB6r1 is that the reaction between isoprene and NO_3 in

560 CB6r1 facilitates the formation of organic nitrates (i.e. NTR, INTR) as discussed above. The formed organic nitrates are then photolyzed or react with OH, forming NO_x . As a result, the emitted NO_x is more involved in the chemical system represented by CB6r1, leading to a higher dependence of NO_x on the direct release of NO and NO_2 in CB6r1. As a consequence, the NO_x level predicted by CB6r1 is higher than those given by CB6r2 and CB6r3 with the same emission intensity of NO_x . ~~similar to the findings in the sensitivity analysis of O_3 , the NO_x mixing ratio predicted by CB6r1 depends less on the surface emission of isoprene. Aside from that, it was found in CB6r1 that the dependence of NO_x on all the emissions becomes weaker than that in CB6r2 and CB6r3. It means that in the CB6r1 mechanism, the influence caused by the addition of the surface emission is smaller, and the change of NO_x is more associated with the local chemical reactions such as the PANX formation. This enhanced NO_x dependence on the local chemistry in CB6r1 is also indicated by the increasing sensitivities of many reactions in the mechanism, such as the PANX formation and consumption reactions, i.e. (R62) $\text{CXO}_3 + \text{NO}_2 \rightarrow \text{PANX}$ and (R63) $\text{PANX} \rightarrow \text{NO}_2 + \text{CXO}_3$ (see Fig. S10 of the supplements), compared with that in CB6r3 and CB6r2 (Fig. 7 and Fig. S9 of the supplementary material). It thus leads to an enhanced formation of PANX and a stronger diurnal change in the NO_x concentration in CB6r1, which has been displayed in Fig. 5(e).~~

570 At last, we focused on the averaged sensitivities of HCHO to these three different CB6 mechanisms. Figure 8 shows the HCHO sensitivities to the chemical reactions, **surface emissions and dry depositions** for the CB6r3 mechanism. It was found that processes that play an important role in the change of HCHO include (R96) $\text{HCHO} + \text{OH} \rightarrow \text{HO}_2 + \text{CO}$, (R97) $\text{HCHO} + h\nu \rightarrow 2\text{HO}_2 + \text{CO}$, and (R98) $\text{HCHO} + h\nu \rightarrow \text{CO}$, which are reactions consuming HCHO, **and the dry deposition process denoted by Reaction (R243) in Fig. 8(d)**. These important HCHO decay pathways are the same to those in the weak-emission case. **In addition, a strong negative role of Reaction (R26), $\text{NO}_2 + \text{O}_3 \rightarrow \text{NO}_3$, in determining HCHO is also identified under this strong emission condition (see Fig. 8a).** The reason is that in CB6r3, the emitted isoprene reacts with NO_3 generated from Reaction (R26), forming NTR2. NTR2 is then converted to the inactive HNO_3 and thus exerts a minor influence on the change of HCHO. Therefore, the occurrence of Reaction (R26) in CB6r3 can substantially suppress the formation of HCHO caused by the release of isoprene, especially under this strong NO_x emission condition. It is also interesting to note that Reaction (R124) $\text{CH}_4 + \text{OH} \rightarrow \text{MEO}_2 + \text{RO}_2$ that used to strongly promote the formation of HCHO in the weak-emission scenario no longer possesses a large sensitivity. It denotes a decreased importance of the initial CH_4 in the formation of HCHO after increasing the surface emission in the model. Instead, the release of VOCs would significantly promote the formation of HCHO.

585 From the dependence of HCHO on the surface emissions displayed in Fig. 8(d), we found that an increase in the emission intensity of VOCs especially isoprene and terminal olefins would significantly enhance the HCHO formation, and the influence caused by the emissions of isoprene and terminal olefins is approximately equal, indicated by the similar sensitivities to these two emissions. This strong influence of the emissions of isoprene and other olefins on the change of HCHO has also been identified in many previous studies (Luecken et al., 2006, 2008; Wolfe et al., 2016; Marvin et al., 2017). Moreover, in the present study, we found that the increase of the NO_x emission leads to an elevation of HCHO. **It is because that the release of NO_x would significantly increase the ozone level in this scenario. As Reaction (R156) $\text{ISOP} + \text{O}_3 \rightarrow 0.6\text{HCHO} + 0.65\text{ISPD} + 0.15\text{ALDX} + 0.2\text{CXO}_3 + 0.35\text{PAR} + 0.27\text{OH} + 0.2\text{XO}_2 + 0.2\text{RO}_2 + 0.07\text{HO}_2 + 0.07\text{CO}$ is a major pathway for the formation of HCHO, reflected by the strong HCHO dependence on Reaction (R156) (see Fig. 8c), the increase of ozone due to the enhanced**

590

NO_x emission would thus promote the formation of HCHO, leading to a positive dependence of HCHO on the release of NO_x. It was also found in this study that HCHO is more sensitive to VOCs than NO_x, which is in accordance with the conclusions achieved in the previous sensitivity study of HCHO to precursor species (Luecken et al., 2018). The reason is that HCHO can be formed under both NO_x-rich and NO_x-poor conditions, resulting in a weaker dependence of HCHO on the NO_x emissions than that on the VOC emissions.

In a previous modeling study conducted by Luecken et al. (2019) using CB6r3, they found an underestimation of HCHO in a comparison with observations across the US. Luecken et al. (2019) suggested that the underestimation of HCHO might be caused by the uncertainties in biogenic emissions including direct HCHO emissions and other VOC emissions. Based on our findings, we suggested that the underestimation of HCHO might be caused by the underestimation of isoprene and other alkene emissions. In contrast, the direct emission of HCHO may possibly only exert a minor impact on the change of HCHO, according to the sensitivity analysis of CB6r3 in the present study. In the study of Luecken et al. (2019), they also performed a sensitivity test by doubling the isoprene emission, and it was found that the simulated HCHO is elevated due to the enhancement of the isoprene emission. This is also in accordance with our findings in the present study.

By comparing the HCHO sensitivities to CB6r3 (Fig 8) with those to CB6r2 (Fig. S11 in the supplements), we noticed that the most dominant reactions for the change of HCHO approximately the same in these two mechanisms, thus leading to a similar HCHO prediction by these two mechanisms. However, the sensitivity of HCHO to CB6r1 displayed in Fig. S12 of the supplements shows that in CB6r1, the significance of the isoprene emission on the change of HCHO (~0.39) is approximately 10% higher than that in CB6r2 and CB6r3 (~0.36). As a result, a same increment in the isoprene emission would lead to a relatively larger increase in HCHO predicted by CB6r1 relative to that predicted by CB6r2 or CB6r3. This is also the reason for the relatively higher HCHO prediction by CB6r1 shown in Fig. 5. Again, it is attributed to the higher ozone prediction of CB6r1 caused by the change of Reaction (R157). Because the major HCHO formation pathway in this simulation scenario is the reaction between isoprene and ozone, under the condition with higher ozone estimated by CB6r1, larger amount of the emitted isoprene can be converted to HCHO, thus causing the higher HCHO prediction and the heavier dependence of HCHO on the isoprene emission in CB6r1. This finding again indicates the importance of revising the isoprene chemistry in future updates to the CB6 mechanism.

In summary, in the situation with the inclusion of the strong surface emission in the model, we found that the ozone level predicted by CB6r2 and CB6r3 depends heavily on the surface emission especially the release of NO and isoprene. In contrast, the dependence of ozone on the isoprene is weaker in CB6r1. Aside from that, the importance of many isoprene related reactions in CB6r1 for the change of ozone decreases, which is shown in the sensitivity analysis. These changes in the ozone sensitivity lead to a higher ozone prediction of CB6r1 relative to that of CB6r2 and CB6r3, even though a same surface emission condition is applied. With respect to the change of NO_x, in CB6r2 and CB6r3, the most influential emissions are also the release of NO and isoprene. However, in CB6r1, the dependence of the predicted NO_x on the surface emissions of NO and NO₂ is heavier than that in CB6r2 and CB6r3, due to the change in the product of the reaction between isoprene and NO₃, resulting in a higher NO_x prediction of CB6r1. ~~but more on the local chemistry. Chemical reactions especially the formation of PANX play a more important role in determining NO_x, leading to a higher NO_x prediction during the daytime and a lower NO_x prediction during~~

~~the nighttime in CB6r1, respectively.~~ At last, for HCHO, the sensitivity analysis shows that the change of HCHO relies more on the emissions of VOCs instead of the emissions of NO_x , and the enhancement of the VOC emissions particularly isoprene and terminal olefins would significantly promote the formation of HCHO. However, the dependence of HCHO on the release of isoprene in CB6r1 is stronger than that in the other two mechanisms, thus leading to a higher prediction of HCHO in CB6r1 under the same emission condition.

4 Conclusions and Future Work

In the present study, we found that different versions of the CB6 mechanism perform differently in simulating O_3 , NO_x and HCHO, although the same initial condition and the same intensity of the surface emission is set up. When the surface emission is **weak**, CB6r1 predicts a higher ozone value than the other two mechanisms, and the deviation is approximately 7 ppb. The sensitivity analysis suggests that the higher ozone prediction by CB6r1 is partly caused by the modification of the chemical loss pathways of acylperoxy radicals with three and higher carbons (i.e. species CXO3) in the mechanism. Due to this modification, less CXO3 is consumed in CB6r1 than that in CB6r2 and CB6r3, resulting in a higher ozone prediction by CB6r1. **Moreover, the ozone sensitivity to the isoprene emission in CB6r1 was found larger than that in CB6r2 and CB6r3, which also contributes to the higher ozone prediction of CB6r1 under the same isoprene emission condition.** Regarding to NO_x and HCHO, ~~the predictions by these three CB6 mechanisms are similar during the end of the simulation;~~ the estimations given by CB6r1 are higher than those given by CB6r2 and CB6r3, but the deviations between the simulation results become smaller during the end of the 7-day computation. The sensitivity analysis also shows that the update in CB6r3 about the temperature dependence of organic nitrate formation might exert a strong influence on the prediction of ozone and NO_x under a different temperature condition, while the impact of the temperature change on HCHO might be minor.

After implementing a **strong** surface emission into the model, we found the simulated levels of O_3 , NO_x and HCHO elevated, compared with those in the **weak**-emission scenario. It was also found that the ozone concentration predicted by CB6r2 and CB6r3 depends on the emissions of NO and isoprene the most, while in CB6r1 the dependence of ozone on the isoprene emission is weaker. Because in this simulation, the isoprene emission tends to suppress the ozone formation, ozone predicted by CB6r1 is thus higher than those predicted by CB6r2 and CB6r3 with the same emission intensity. With respect to the NO_x prediction, in CB6r1, **the association between the mixing ratio of NO_x and the release of NO and NO_2 was found stronger than that in CB6r2 and CB6r3.** It is because that the released NO_x is more involved in the reaction system represented by CB6r1, thus leading to a higher NO_x prediction of CB6r1, compared with that given by CB6r2 or CB6r3. ~~NO_x depends more on the local chemistry such as the formation of PANX than that in CB6r2 and CB6r3. As a result, NO_x in CB6r1 is more involved in the formation of reservoirs such as PANX, thus leading to a lower NO_x prediction at nighttime and a higher NO_x prediction at daytime compared with that in CB6r2 and CB6r3.~~ At last, we found that the HCHO predictions of these three CB6 mechanisms rely mostly on the emissions of NO, isoprene and terminal olefins. However, in CB6r1, the association between HCHO and the isoprene emission is stronger, resulting in a higher HCHO prediction relative to that in CB6r2 and CB6r3 with the same isoprene emission.

660 The present study has its limitations. The conclusions achieved in this study are mostly valid under conditions that have
 been presented and analyzed in this box model study, and these conditions may possibly differ from those present in 3-D
 model simulations of the atmosphere. Therefore, in the future, we plan to test the behavior of these CB6 mechanisms under
 different environmental conditions with different surface emission intensities, especially the conditions that are implemented
 in 3-D model studies of the atmosphere. The influence caused by the varying of the temperature on the concentration change
 665 of the focused species, especially for CB6r3 should also be investigated. Moreover, the latest version of the CB6 mechanism,
 CB6r4 (Emery et al., 2016), should be studied and compared with the three CB6 mechanisms investigated in the present study,
 particularly in a halogen-rich environment. In addition, the conclusions achieved in this box-model study need to be confirmed
 in simulations using multi-dimensional air quality models. At present, we are conducting three-dimensional simulations using
 CMAQ (Byun and Schere, 2006) and CAMx (ENVIRON, 2015) to discover the difference in modeling O₃, NO_x and HCHO
 670 by using these different versions of the CB6 mechanism, which is attributed to a future publication.

Code and data availability. The source code of the model and the data of the computational results shown in this article can be acquired
 from the link: https://pan.baidu.com/s/1Gi_Tb-SlrMi0IvD-4FBYJQ, using the password: bpc1.

Appendix A

Table A1. Complete listings of chemical reactions belonging to different CB6 mechanisms used in the present study. The updates between
 different versions of the CB6 mechanism are also marked. The abbreviation “-” denotes that there is no change in the form of this reaction
 between different CB6 mechanisms.

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R1)	NO ₂ + hν → NO + O	(R1)	-	(R1)	-
(R2)	O + O ₂ + M → O ₃ + M	(R2)	-	(R2)	-
(R3)	O ₃ + NO → NO ₂	(R3)	-	(R3)	-
(R4)	O + NO + M → NO ₂ + M	(R4)	-	(R4)	-
(R5)	O + NO ₂ → NO	(R5)	-	(R5)	-
(R6)	O + NO ₂ → NO ₃	(R6)	-	(R6)	-
(R7)	O + O ₃ →	(R7)	-	(R7)	-
(R8)	O ₃ + hν → O	(R8)	-	(R8)	-
(R9)	O ₃ + hν → O(¹ D)	(R9)	-	(R9)	-
(R10)	O(¹ D) + M → O + M	(R10)	-	(R10)	-

Table A1. (continued).

Reaction Number	CB6r1^a	Reaction Number	CB6r2^b	Reaction Number	CB6r3^c
(R11)	$O(^1D) + H_2O \rightarrow 2OH$	(R11)	-	(R11)	-
(R12)	$O_3 + OH \rightarrow HO_2$	(R12)	-	(R12)	-
(R13)	$O_3 + HO_2 \rightarrow OH$	(R13)	-	(R13)	-
(R14)	$OH + O \rightarrow HO_2$	(R14)	-	(R14)	-
(R15)	$HO_2 + O \rightarrow OH$	(R15)	-	(R15)	-
(R16)	$OH + OH \rightarrow O$	(R16)	-	(R16)	-
(R17)	$OH + OH \rightarrow H_2O_2$	(R17)	-	(R17)	-
(R18)	$OH + HO_2 \rightarrow$	(R18)	-	(R18)	-
(R19)	$HO_2 + HO_2 \rightarrow H_2O_2$	(R19)	-	(R19)	-
(R20)	$HO_2 + HO_2 + H_2O \rightarrow H_2O_2$	(R20)	-	(R20)	-
(R21)	$H_2O_2 + h\nu \rightarrow 2OH$	(R21)	-	(R21)	-
(R22)	$H_2O_2 + OH \rightarrow HO_2$	(R22)	-	(R22)	-
(R23)	$H_2O_2 + O \rightarrow OH + HO_2$	(R23)	-	(R23)	-
(R24)	$NO + NO + O_2 \rightarrow 2NO_2$	(R24)	-	(R24)	-
(R25)	$HO_2 + NO \rightarrow OH + NO_2$	(R25)	-	(R25)	-
(R26)	$NO_2 + O_3 \rightarrow NO_3$	(R26)	-	(R26)	-
(R27)	$NO_3 + h\nu \rightarrow NO_2 + O$	(R27)	-	(R27)	-
(R28)	$NO_3 + h\nu \rightarrow NO$	(R28)	-	(R28)	-
(R29)	$NO_3 + NO \rightarrow 2NO_2$	(R29)	-	(R29)	-
(R30)	$NO_3 + NO_2 \rightarrow NO + NO_2$	(R30)	-	(R30)	-
(R31)	$NO_3 + O \rightarrow NO_2$	(R31)	-	(R31)	-
(R32)	$NO_3 + OH \rightarrow HO_2 + NO_2$	(R32)	-	(R32)	-
(R33)	$NO_3 + HO_2 \rightarrow OH + NO_2$	(R33)	-	(R33)	-
(R34)	$NO_3 + O_3 \rightarrow NO_2$	(R34)	-	(R34)	-
(R35)	$NO_3 + NO_3 \rightarrow 2NO_2$	(R35)	-	(R35)	-
(R36)	$NO_3 + NO_2 \rightarrow N_2O_5$	(R36)	-	(R36)	-
(R37)	$N_2O_5 \rightarrow NO_3 + NO_2$	(R37)	-	(R37)	-
(R38)	$N_2O_5 + h\nu \rightarrow NO_3 + NO_2$	(R38)	-	(R38)	-
(R39)	$N_2O_5 + H_2O \rightarrow 2HNO_3$	(R39)	-	(R39)	-
(R40)	$NO + OH \rightarrow HONO$	(R40)	-	(R40)	-

Table A1. (continued).

Reaction Number	CB6r1^a	Reaction Number	CB6r2^b	Reaction Number	CB6r3^c
(R41)	$\text{NO} + \text{NO}_2 + \text{H}_2\text{O} \rightarrow 2\text{HONO}$	(R41)	-	(R41)	-
(R42)	$\text{HONO} + \text{HONO} \rightarrow \text{NO} + \text{NO}_2$	(R42)	-	(R42)	-
(R43)	$\text{HONO} + h\nu \rightarrow \text{NO} + \text{OH}$	(R43)	-	(R43)	-
(R44)	$\text{HONO} + \text{OH} \rightarrow \text{NO}_2$	(R44)	-	(R44)	-
(R45)	$\text{NO}_2 + \text{OH} \rightarrow \text{HNO}_3$	(R45)	-	(R45)	-
(R46)	$\text{HNO}_3 + \text{OH} \rightarrow \text{NO}_3$	(R46)	-	(R46)	-
(R47)	$\text{HNO}_3 + h\nu \rightarrow \text{OH} + \text{NO}_2$	(R47)	-	(R47)	-
(R48)	$\text{HO}_2 + \text{NO}_2 \rightarrow \text{PNA}$	(R48)	-	(R48)	-
(R49)	$\text{PNA} \rightarrow \text{HO}_2 + \text{NO}_2$	(R49)	-	(R49)	-
(R50)	$\text{PNA} + h\nu \rightarrow 0.59\text{HO}_2 + 0.59\text{NO}_2 + 0.41\text{OH} + 0.41\text{NO}_3$	(R50)	-	(R50)	-
(R51)	$\text{PNA} + \text{OH} \rightarrow \text{NO}_2$	(R51)	-	(R51)	-
(R52)	$\text{SO}_2 + \text{OH} \rightarrow \text{SULF} + \text{HO}_2$	(R52)	-	(R52)	-
(R53)	$\text{C}_2\text{O}_3 + \text{NO} \rightarrow \text{NO}_2 + \text{MEO}_2 + \text{RO}_2$	(R53)	-	(R53)	-
(R54)	$\text{C}_2\text{O}_3 + \text{NO}_2 \rightarrow \text{PAN}$	(R54)	-	(R54)	-
(R55)	$\text{PAN} \rightarrow \text{C}_2\text{O}_3 + \text{NO}_2$	(R55)	-	(R55)	-
(R56)	$\text{PAN} + h\nu \rightarrow 0.6\text{NO}_2 + 0.6\text{C}_2\text{O}_3 + 0.4\text{NO}_3 + 0.4\text{MEO}_2 + 0.4\text{RO}_2$	(R56)	-	(R56)	-
(R57)	$\text{C}_2\text{O}_3 + \text{HO}_2 \rightarrow 0.41\text{PACD} + 0.15\text{AACD} + 0.15\text{O}_3 + 0.44\text{MEO}_2 + 0.44\text{RO}_2 + 0.44\text{OH}$	(R57)	-	(R57)	-
(R58)	$\text{C}_2\text{O}_3 + \text{RO}_2 \rightarrow \text{C}_2\text{O}_3$	(R58)	-	(R58)	-
(R59)	$\text{C}_2\text{O}_3 + \text{C}_2\text{O}_3 \rightarrow 2\text{MEO}_2 + 2\text{RO}_2$	(R59)	-	(R59)	-
(R60)	$\text{C}_2\text{O}_3 + \text{CXO}_3 \rightarrow \text{MEO}_2 + \text{ALD}_2 + \text{XO}_2\text{H} + 2\text{RO}_2$	(R60)	-	(R60)	-
(R61)	$\text{CXO}_3 + \text{NO} \rightarrow \text{NO}_2 + \text{ALD}_2 + \text{XO}_2\text{H} + \text{RO}_2$	(R61)	-	(R61)	-
(R62)	$\text{CXO}_3 + \text{NO}_2 \rightarrow \text{PANX}$	(R62)	-	(R62)	-
(R63)	$\text{PANX} \rightarrow \text{NO}_2 + \text{CXO}_3$	(R63)	-	(R63)	-
(R64)	$\text{PANX} + h\nu \rightarrow 0.6\text{NO}_2 + 0.6\text{CXO}_3 + 0.4\text{NO}_3 + 0.4\text{ALD}_2 + 0.4\text{XO}_2\text{H} + 0.4\text{RO}_2$	(R64)	-	(R64)	-

Table A1. (continued).

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R65)	$CXO_3 + HO_2 \rightarrow 0.41PACD + 0.15AACD + 0.15O_3 + 0.44ALD_2 + 0.44XO_2H + 0.44RO_2 + 0.44OH$	(R65)	-	(R65)	-
(R66)	$CXO_3 + RO_2 \rightarrow CXO_3$	(R66)	$CXO_3 + RO_2 \rightarrow 0.8ALD_2 + 0.8XO_2H + 0.8RO_2$	(R66)	-
(R67)	$CXO_3 + CXO_3 \rightarrow 2ALD_2 + 2XO_2H + 2RO_2$	(R67)	-	(R67)	-
(R68)	$RO_2 + NO \rightarrow NO$	(R68)	-	(R68)	-
(R69)	$RO_2 + HO_2 \rightarrow HO_2$	(R69)	-	(R69)	-
(R70)	$RO_2 + RO_2 \rightarrow$	(R70)	-	(R70)	-
(R71)	$MEO_2 + NO \rightarrow HCHO + HO_2 + NO_2$	(R71)	-	(R71)	-
(R72)	$MEO_2 + HO_2 \rightarrow 0.9MEPX + 0.1HCHO$	(R72)	-	(R72)	-
(R73)	$MEO_2 + C_2O_3 \rightarrow HCHO + 0.9HO_2 + 0.9MEO_2 + 0.1AACD + 0.9RO_2$	(R73)	-	(R73)	-
(R74)	$MEO_2 + RO_2 \rightarrow 0.685HCHO + 0.315MEOH + 0.37HO_2 + RO_2$	(R74)	-	(R74)	-
(R75)	$XO_2H + NO \rightarrow NO_2 + HO_2$	(R75)	-	(R75)	-
(R76)	$XO_2H + HO_2 \rightarrow ROOH$	(R76)	-	(R76)	-
(R77)	$XO_2H + C_2O_3 \rightarrow 0.8HO_2 + 0.8MEO_2 + 0.2AACD + 0.8RO_2$	(R77)	-	(R77)	-
(R78)	$XO_2H + RO_2 \rightarrow 0.6HO_2 + RO_2$	(R78)	-	(R78)	-
(R79)	$XO_2 + NO \rightarrow NO_2$	(R79)	-	(R79)	-
(R80)	$XO_2 + HO_2 \rightarrow ROOH$	(R80)	-	(R80)	-
(R81)	$XO_2 + C_2O_3 \rightarrow 0.8MEO_2 + 0.2AACD + 0.8RO_2$	(R81)	-	(R81)	-
(R82)	$XO_2 + RO_2 \rightarrow 0.6HO_2 + RO_2$	(R82)	$XO_2 + RO_2 \rightarrow RO_2$	(R82)	-
(R83)	$XO_2N + NO \rightarrow NTR$	(R83)	$XO_2N + NO \rightarrow 0.5NTR_1 + 0.5NTR_2$	(R83)	-
(R84)	$XO_2N + HO_2 \rightarrow ROOH$	(R84)	-	(R84)	-
(R85)	$XO_2N + C_2O_3 \rightarrow 0.8HO_2 + 0.8MEO_2 + 0.2AACD + 0.8RO_2$	(R85)	-	(R85)	-
(R86)	$XO_2N + RO_2 \rightarrow 0.6HO_2 + RO_2$	(R86)	$XO_2N + RO_2 \rightarrow RO_2$	(R86)	-
(R87)	$MEPX + OH \rightarrow 0.6MEO_2 + 0.6RO_2 + 0.4HCHO + 0.4OH$	(R87)	-	(R87)	-
(R88)	$MEPX + h\nu \rightarrow MEO_2 + RO_2 + OH$	(R88)	-	(R88)	-
(R89)	$ROOH + OH \rightarrow 0.54XO_2H + 0.06XO_2N + 0.6RO_2 + 0.4OH$	(R89)	-	(R89)	-
(R90)	$ROOH + h\nu \rightarrow HO_2 + OH$	(R90)	-	(R90)	-

Table A1. (continued).

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R91)	$\text{NTR} + \text{OH} \rightarrow \text{HNO}_3 + \text{XO}_2\text{H} + \text{RO}_2$	(R91)	$\text{NTR1} + \text{OH} \rightarrow \text{NTR2}$	(R91)	-
(R92)	$\text{NTR} + h\nu \rightarrow \text{NO}_2 + \text{XO}_2\text{H} + \text{RO}_2$	(R92)	$\text{NTR1} + h\nu \rightarrow \text{NO}_2$	(R92)	-
(R93)	$\text{FACD} + \text{OH} \rightarrow \text{HO}_2$	(R93)	-	(R93)	-
(R94)	$\text{AACD} + \text{OH} \rightarrow \text{MEO}_2 + \text{RO}_2$	(R94)	-	(R94)	-
(R95)	$\text{PACD} + \text{OH} \rightarrow \text{C}_2\text{O}_3$	(R95)	-	(R95)	-
(R96)	$\text{HCHO} + \text{OH} \rightarrow \text{HO}_2 + \text{CO}$	(R96)	-	(R96)	-
(R97)	$\text{HCHO} + h\nu \rightarrow 2\text{HO}_2 + \text{CO}$	(R97)	-	(R97)	-
(R98)	$\text{HCHO} + h\nu \rightarrow \text{CO}$	(R98)	-	(R98)	-
(R99)	$\text{HCHO} + \text{O} \rightarrow \text{OH} + \text{HO}_2 + \text{CO}$	(R99)	-	(R99)	-
(R100)	$\text{HCHO} + \text{NO}_3 \rightarrow \text{HNO}_3 + \text{HO}_2 + \text{CO}$	(R100)	-	(R100)	-
(R101)	$\text{HCHO} + \text{HO}_2 \rightarrow \text{HCO}_3$	(R101)	-	(R101)	-
(R102)	$\text{HCO}_3 \rightarrow \text{HCHO} + \text{HO}_2$	(R102)	-	(R102)	-
(R103)	$\text{HCO}_3 + \text{NO} \rightarrow \text{FACD} + \text{NO}_2 + \text{HO}_2$	(R103)	-	(R103)	-
(R104)	$\text{HCO}_3 + \text{HO}_2 \rightarrow 0.5\text{MEPX} + 0.5\text{FACD} + 0.2\text{OH} + 0.2\text{HO}_2$	(R104)	-	(R104)	-
(R105)	$\text{ALD}_2 + \text{O} \rightarrow \text{C}_2\text{O}_3 + \text{OH}$	(R105)	-	(R105)	-
(R106)	$\text{ALD}_2 + \text{OH} \rightarrow \text{C}_2\text{O}_3$	(R106)	-	(R106)	-
(R107)	$\text{ALD}_2 + \text{NO}_3 \rightarrow \text{C}_2\text{O}_3 + \text{HNO}_3$	(R107)	-	(R107)	-
(R108)	$\text{ALD}_2 + h\nu \rightarrow \text{MEO}_2 + \text{RO}_2 + \text{CO} + \text{HO}_2$	(R108)	-	(R108)	-
(R109)	$\text{ALDX} + \text{O} \rightarrow \text{CXO}_3 + \text{OH}$	(R109)	-	(R109)	-
(R110)	$\text{ALDX} + \text{OH} \rightarrow \text{CXO}_3$	(R110)	-	(R110)	-
(R111)	$\text{ALDX} + \text{NO}_3 \rightarrow \text{CXO}_3 + \text{HNO}_3$	(R111)	-	(R111)	-
(R112)	$\text{ALDX} + h\nu \rightarrow \text{MEO}_2 + \text{RO}_2 + \text{CO} + \text{HO}_2$	(R112)	$\text{ALDX} + h\nu \rightarrow \text{ALD}_2 + \text{XO}_2\text{H} + \text{RO}_2 + \text{CO} + \text{HO}_2$	(R112)	-
(R113)	$\text{GLYD} + \text{OH} \rightarrow 0.2\text{GLY} + 0.2\text{HO}_2 + 0.8\text{C}_2\text{O}_3$	(R113)	-	(R113)	-
(R114)	$\text{GLYD} + h\nu \rightarrow 0.74\text{HCHO} + 0.89\text{CO} + 1.4\text{HO}_2 + 0.15\text{MEOH} + 0.19\text{OH} + 0.11\text{GLY} + 0.11\text{XO}_2\text{H} + 0.11\text{RO}_2$	(R114)	-	(R114)	-
(R115)	$\text{GLYD} + \text{NO}_3 \rightarrow \text{HNO}_3 + \text{C}_2\text{O}_3$	(R115)	-	(R115)	-
(R116)	$\text{GLY} + \text{OH} \rightarrow 1.7\text{CO} + 0.3\text{XO}_2 + 0.3\text{RO}_2 + \text{HO}_2$	(R116)	$\text{GLY} + \text{OH} \rightarrow 1.8\text{CO} + 0.2\text{XO}_2 + 0.2\text{RO}_2 + \text{HO}_2$	(R116)	-
(R117)	$\text{GLY} + h\nu \rightarrow 2\text{HO}_2 + 2\text{CO}$	(R117)	-	(R117)	-
(R118)	$\text{GLY} + \text{NO}_3 \rightarrow \text{HNO}_3 + \text{CO} + \text{HO}_2 + \text{XO}_2 + \text{RO}_2$	(R118)	$\text{GLY} + \text{NO}_3 \rightarrow \text{HNO}_3 + 1.5\text{CO} + 0.5\text{XO}_2 + 0.5\text{RO}_2 + \text{HO}_2$	(R118)	-

Table A1. (continued).

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R119)	MGLY + $h\nu$ → C2O3 + HO ₂ + CO	(R119)	-	(R119)	-
(R120)	MGLY + NO ₃ → HNO ₃ + C2O3 + XO2 + RO2	(R120)	-	(R120)	-
(R121)	MGLY + OH → C2O3 + CO	(R121)	-	(R121)	-
(R122)	H ₂ + OH → HO ₂	(R122)	-	(R122)	-
(R123)	CO + OH → HO ₂	(R123)	-	(R123)	-
(R124)	CH ₄ + OH → MEO2 + RO ₂	(R124)	-	(R124)	-
(R125)	ETHA + OH → 0.991ALD2 + 0.991XO2H+ 0.009XO2N + RO ₂	(R125)	-	(R125)	-
(R126)	MEOH + OH → HCHO + HO ₂	(R126)	-	(R126)	-
(R127)	ETOH + OH → 0.95ALD2 + 0.9HO ₂ + 0.1XO2H + 0.1RO ₂ + 0.078HCHO + 0.011GLYD	(R127)	-	(R127)	-
(R128)	KET + $h\nu$ → 0.5ALD2 + 0.5C2O3+ 0.5XO2H + 0.5CXO3 + 0.5MEO2 + RO2- 2.5PAR	(R128)	-	(R128)	-
(R129)	ACET + $h\nu$ → 0.38CO + 1.38MEO2+ 1.38RO2 + 0.62C2O3	(R129)	-	(R129)	-
(R130)	ACET + OH → HCHO + C2O3 + XO2 + RO2	(R130)	-	(R130)	-
(R131)	PRPA + OH → 0.71ACET + 0.26ALDX+ 0.26PAR + 0.97XO2H + 1.00RO2 + 0.03XO2N	(R131)	-	(R131)	PRPA + OH → XPRP
(R132)	PAR + OH → 0.11ALDX + 0.76ROR+ 0.11XO2H + 0.76XO2 + RO2 - 0.11PAR+ 0.13XO2N	(R132)	-	(R132)	PAR + OH → XPAR -
(R133)	ROR → 0.2KET + 0.42ACET + 0.74ALD2+ 0.37ALDX + 0.04XO2N + 0.94XO2H + 0.98RO2+ 0.02ROR - 2.7PAR	(R133)	-	(R133)	-
(R134)	ROR + O ₂ → KET + HO ₂	(R134)	-	(R134)	-
(R135)	ROR + NO ₂ → NTR	(R135)	ROR + NO ₂ → NTR2	(R135)	ROR + NO ₂ → NTR1
(R136)	ETHY + OH → 0.7GLY + 0.7OH+ 0.3FACD + 0.3CO + 0.3HO ₂	(R136)	-	(R136)	-
(R137)	ETH + O → HCHO + HO ₂ + CO + 0.7XO2H + 0.7RO2 + 0.3OH	(R137)	-	(R137)	-
(R138)	ETH + OH → XO2H + RO2+ 1.56HCHO + 0.22GLYD	(R138)	-	(R138)	-
(R139)	ETH + O ₃ → HCHO + 0.51CO+ 0.16HO ₂ + 0.16OH + 0.37FACD	(R139)	-	(R139)	-
(R140)	ETH + NO ₃ → 0.5NO ₂ + 0.5NTR+ 0.5XO2H + 0.5XO2 + RO2 + 1.12HCHO	(R140)	ETH + NO ₃ → 0.5NO ₂ + 0.5NTR1+ 0.5XO2H + 0.5XO2 + RO2 + 1.12HCHO	(R140)	-
(R141)	OLE + O → 0.2ALD2 + 0.3ALDX+ 0.1HO ₂ + 0.2XO2H + 0.2CO + 0.2HCHO+ 0.01XO2N + 0.21RO2 + 0.2PAR + 0.1OH	(R141)	-	(R141)	-

Table A1. (continued).

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R142)	OLE + OH → 0.78HCHO + 0.49ALD2+ 0.49ALDX + 0.98XO2H + 0.2XO2 + 0.02XO2N+ 1.2RO2 - 0.73PAR	(R142)	-	(R142)	-
(R143)	OLE + O ₃ → 0.29ALD2 + 0.56HCHO+ 0.27ALDX + 0.15XO2H + 0.15RO2 + 0.33OH+ 0.08HO ₂ + 0.38CO + 0.07GLY + 0.07MGLY+ 0.09FACD + 0.13AACD + 0.04H ₂ O ₂ - 0.79PAR	(R143)	-	(R143)	-
(R144)	OLE + NO ₃ → 0.5NO ₂ + 0.5NTR+ 0.48XO2 + 0.48XO2H + 0.04XO2N + RO2+ 0.5HCHO + 0.25ALD2 + 0.38ALDX - PAR	(R144)	OLE + NO ₃ → 0.5NO ₂ + 0.5NTR1+ 0.48XO2 + 0.48XO2H + 0.04XO2N + RO2+ 0.5HCHO + 0.25ALD2 + 0.38ALDX - PAR	(R144)	-
(R145)	IOLE + O → 1.24ALD2 + 0.66ALDX+ 0.1XO2H + 0.1RO2 + 0.1CO + 0.1PAR	(R145)	-	(R145)	-
(R146)	IOLE + OH → 1.30ALD2 + 0.7ALDX+ XO2H + RO2	(R146)	-	(R146)	-
(R147)	IOLE + O ₃ → 0.73ALD2 + 0.44ALDX+ 0.13HCHO + 0.24CO + 0.5OH + 0.3XO2H+ 0.3RO2 + 0.24GLY + 0.06MGLY + 0.29PAR+ 0.08AACD + 0.08H ₂ O ₂	(R147)	-	(R147)	-
(R148)	IOLE + NO ₃ → 0.5NO ₂ + 0.5NTR+ 0.48XO2 + 0.48XO2H + 0.04XO2N + RO2+ 0.5ALD2 + 0.62ALDX + PAR	(R148)	IOLE + NO ₃ → 0.5NO ₂ + 0.5NTR1+ 0.48XO2 + 0.48XO2H + 0.04XO2N + RO2+ 0.5ALD2 + 0.62ALDX + PAR	(R148)	-
(R149)	ISOP + OH → ISO2RO2+	(R149)	-	(R149)	-
(R150)		(R150)	ISOP + O → 0.75ISPD + 0.5HCHO+ 0.25XO2 + 0.25RO2 + 0.25HO ₂ + 0.25CXO3+ 0.25PAR	(R150)	-
(R151)	ISO2 + NO → 0.12INTR + 0.88NO ₂ + 0.8HO ₂ + 0.66HCHO + 0.66ISPD + 0.08XO2H+ 0.08RO2 + 0.05IOLE + 0.04GLYD + 0.12PAR+ 0.04GLY + 0.04MGLY + 0.09OLE + 0.12ALDX	(R151)	ISO2 + NO → 0.1INTR + 0.9NO ₂ + 0.67HCHO + 0.9ISPD + 0.82HO ₂ + 0.08XO2H + 0.08RO2	(R151)	-
(R152)	ISO2 + HO ₂ → 0.88ISPX + 0.12OH+ 0.12HO ₂ + 0.12HCHO + 0.12ISPD	(R152)	-	(R152)	-
(R153)	ISO2 + C2O3 → 0.71HO ₂ + 0.58HCHO+ 0.58ISPD + 0.07XO2H + 0.04IOLE + 0.04GLYD+ 0.1PAR + 0.03GLY + 0.04MGLY + 0.08OLE+ 0.1ALDX + 0.8MEO2 + 0.2AACD + 0.87RO2	(R153)	ISO2 + C2O3 → 0.6HCHO + ISPD+ 0.73HO ₂ + 0.07XO2H + 0.8MEO2 + 0.2AACD+ 0.87RO2	(R153)	-
(R154)	ISO2 + RO2 → 0.8HO ₂ + 0.66HCHO+ 0.66ISPD + 0.08XO2H + 0.05IOLE + 0.04GLYD+ 0.12PAR + 0.04GLY + 0.04MGLY + 0.09OLE+ 0.12ALDX + 1.08RO2	(R154)	ISO2 + RO2 → 0.6HCHO + ISPD+ 0.73HO ₂ + 0.07XO2H + 1.07RO2	(R154)	-

Table A1. (continued).

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R155)	ISO ₂ → 0.8HO ₂ + 0.04OH + 0.04HCHO + 0.8ISPD	(R155)	ISO ₂ → HO ₂ + HPLD	(R155)	-
(R156)	ISOP + O ₃ → 0.6HCHO + 0.65ISPD + 0.15ALDX + 0.2CXO ₃ + 0.35PAR + 0.27OH + 0.2XO ₂ + 0.2RO ₂ + 0.07HO ₂ + 0.07CO	(R156)	-	(R156)	-
(R157)	ISOP + NO ₃ → 0.35NO ₂ + 0.65INTR + 0.64XO ₂ H + 0.33XO ₂ + 0.03XO ₂ N + RO ₂ + 0.35HCHO + 0.35ISPD	(R157)	ISOP + NO ₃ → 0.35NO ₂ + 0.65NTR ₂ + 0.64XO ₂ H + 0.33XO ₂ + 0.03XO ₂ N + RO ₂ + 0.35HCHO + 0.35ISPD	(R157)	-
(R158)	ISPD + OH → 0.1XO ₂ N + 0.38XO ₂ + 0.32XO ₂ H + 0.79RO ₂ + 0.84PAR + 0.38C ₂ O ₃ + 0.21CXO ₃ + 0.38GLYD + 0.24MGLY + 0.24HCHO + 0.07OLE + 0.08CO + 0.03ALDX	(R158)	ISPD + OH → 0.06XO ₂ N + 0.52XO ₂ + 0.24XO ₂ H + 0.15MGLY + 0.27MEO ₂ + 0.12GLY + 0.35GLYD + 0.23C ₂ O ₃ + 0.12CXO ₃ + 0.24PAR + 0.26ACET + 0.2CO + 0.14HO ₂ + 1.09RO ₂	(R158)	ISPD + OH → 0.02XO ₂ N + 0.52XO ₂ + 0.12MGLY + 0.12MEO ₂ + 0.27GLYD + 0.27C ₂ O ₃ + 0.46OPO ₃ + 0.12PAR + 0.14ACET + 0.14CO + 0.14HO ₂ + 0.66RO ₂
(R159)	ISPD + O ₃ → 0.02ALD ₂ + 0.15HCHO + 0.23CO + 0.85MGLY + 0.36PAR + 0.11C ₂ O ₃ + 0.06XO ₂ H + 0.06RO ₂ + 0.27OH + 0.09HO ₂	(R159)	ISPD + O ₃ → 0.04ALD ₂ + 0.23HCHO + 0.53MGLY + 0.17GLY + 0.17ACET + 0.54CO + 0.46OH + 0.15FACD + 0.4HO ₂ + 0.14C ₂ O ₃	(R159)	-
(R160)	ISPD + NO ₃ → 0.64CO + 0.28HCHO + 0.36ALDX + 1.28PAR + 0.85HO ₂ + 0.07CXO ₃ + 0.07XO ₂ H + 0.07RO ₂ + 0.85NTR + 0.15HNO ₃	(R160)	ISPD + NO ₃ → 0.72HNO ₃ + 0.14NTR ₂ + 0.14NO ₂ + 0.14XO ₂ + 0.14XO ₂ H + 0.11GLYD + 0.11MGLY + 0.72PAR + 0.72CXO ₃ + 0.28RO ₂	(R160)	-
(R161)	ISPD + hν → 0.33CO + 0.07ALD ₂ + 0.9HCHO + 0.83PAR + 0.33HO ₂ + 0.7XO ₂ H + 0.7RO ₂ + 0.97C ₂ O ₃	(R161)	ISPD + hν → 0.76HO ₂ + 0.34XO ₂ H + 0.16XO ₂ + 0.34MEO ₂ + 0.21C ₂ O ₃ + 0.26HCHO + 0.24OLE + 0.24PAR + 0.17ACET + 0.13GLYD + 0.84RO ₂	(R161)	-
(R162)	ISPX + OH → 0.9EPOX + 0.93OH + 0.07ISO ₂ + 0.07RO ₂ + 0.03IOLE + 0.03ALDX	(R162)	-	(R162)	-
(R163)		(R163)	HPLD → OH + ISPD + HO ₂	(R163)	HPLD → OH + ISPD
(R164)		(R164)	HPLD + NO ₃ → HNO ₃ + ISPD	(R164)	-
(R165)	EPOX + OH → EPX ₂ + RO ₂	(R165)	-	(R165)	-
(R166)	EPX ₂ + HO ₂ → 0.28GLYD + 0.28GLY + 0.28MGLY + 1.12OH + 0.82HO ₂ + 0.38HCHO + 0.07FACD + 0.25CO + 2.17PAR	(R166)	-	(R166)	-
(R167)	EPX ₂ + NO → 0.28GLYD + 0.28GLY + 0.28MGLY + 0.12OH + 0.82HO ₂ + 0.38HCHO + NO ₂ + 0.25CO + 2.17PAR	(R167)	-	(R167)	-
(R168)	EPX ₂ + C ₂ O ₃ → 0.22GLYD + 0.22GLY + 0.22MGLY + 0.1OH + 0.66HO ₂ + 0.3HCHO + 0.2CO + 1.74PAR + 0.8MEO ₂ + 0.2AACD + 0.8RO ₂	(R168)	-	(R168)	-
(R169)	EPX ₂ + RO ₂ → 0.28GLYD + 0.28GLY + 0.28MGLY + 0.12OH + 0.82HO ₂ + 0.38HCHO + 0.25CO + 2.17PAR + RO ₂	(R169)	-	(R169)	-
(R170)	INTR + OH → 0.63XO ₂ + 0.37XO ₂ H + RO ₂ + 0.44NO ₂ + 0.18NO ₃ + 0.1INTR + 0.59HCHO + 0.33GLYD + 0.18FACD + 2.70PAR + 0.1OLE + 0.08ALDX + 0.27NTR	(R170)	INTR + OH → 0.63XO ₂ + 0.37XO ₂ H + RO ₂ + 0.44NO ₂ + 0.18NO ₃ + 0.1INTR + 0.59HCHO + 0.33GLYD + 0.18FACD + 2.70PAR + 0.1OLE + 0.08ALDX + 0.27NTR	(R170)	-

Table A1. (continued).

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R171)	TERP + O → 0.15ALDX + 5.12PAR	(R171)	-	(R171)	-
(R172)	TERP + OH → 0.75XO2H + 0.5XO2+ 0.25XO2N + 1.5RO2 + 0.28HCHO + 1.66PAR+ 0.47ALDX	(R172)	-	(R172)	-
(R173)	TERP + O ₃ → 0.57OH + 0.07XO2H+ 0.69XO2 + 0.18XO2N + 0.94RO2 + 0.24HCHO+ 0.001CO + 7PAR + 0.21ALDX + 0.39CXO3	(R173)	-	(R173)	-
(R174)	TERP + NO ₃ → 0.47NO ₂ + 0.28XO2H+ 0.75XO2 + 0.25XO2N + 1.28RO2 + 0.47ALDX+ 0.53NTR	(R174)	TERP + NO ₃ → 0.47NO ₂ + 0.28XO2H+ 0.75XO2 + 0.25XO2N + 1.28RO2 + 0.47ALDX+ 0.53NTR2	(R174)	-
(R175)	BENZ + OH → 0.53CRES + 0.35BZO2+ 0.35RO2 + 0.12OPEN + 0.12OH + 0.53HO ₂	(R175)	-	(R175)	-
(R176)	BZO2 + NO → 0.92NO ₂ + 0.08NTR+ 0.92GLY + 0.92OPEN + 0.92HO ₂	(R176)	BZO2 + NO → 0.92NO ₂ + 0.08NTR2+ 0.92GLY + 0.92OPEN + 0.92HO ₂	(R176)	-
(R177)	BZO2 + C2O3 → GLY + OPEN+ HO ₂ + MEO2 + RO2	(R177)	-	(R177)	-
(R178)	BZO2 + HO ₂ →	(R178)	-	(R178)	-
(R179)	BZO2 + RO2 → GLY + OPEN+ HO ₂ + RO2	(R179)	-	(R179)	-
(R180)	TOL + OH → 0.18CRES + 0.65TO2+ 0.72RO2 + 0.1OPEN + 0.1OH + 0.07XO2H+ 0.18HO ₂	(R180)	-	(R180)	-
(R181)	TO2 + NO → 0.86NO ₂ + 0.14NTR+ 0.42GLY + 0.44MGLY + 0.66OPEN + 0.2XOPN+ 0.86HO ₂	(R181)	TO2 + NO → 0.86NO ₂ + 0.14NTR2+ 0.42GLY + 0.44MGLY + 0.66OPEN + 0.2XOPN+ 0.86HO ₂	(R181)	-
(R182)	TO2 + C2O3 → 0.48GLY + 0.52MGLY+ 0.77OPEN + 0.23XOPN + HO ₂ + MEO2+ RO2	(R182)	-	(R182)	-
(R183)	TO2 + HO ₂ →	(R183)	-	(R183)	-
(R184)	TO2 + RO2 → 0.48GLY + 0.52MGLY+ 0.77OPEN + 0.23XOPN + HO ₂ + RO2	(R184)	-	(R184)	-
(R185)	XYL + OH → 0.15CRES + 0.54XLO2+ 0.6RO2 + 0.24XOPN + 0.24OH + 0.06XO2H+ 0.15HO ₂	(R185)	-	(R185)	-

Table A1. (continued).

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R186)	XLO2 + NO → 0.86NO ₂ + 0.14NTR+ 0.22GLY + 0.68MGLY + 0.3OPEN + 0.56XOPN+ 0.86HO ₂	(R186)	XLO2 + NO → 0.86NO ₂ + 0.14NTR2+ 0.22GLY + 0.68MGLY + 0.3OPEN + 0.56XOPN+ 0.86HO ₂	(R186)	-
(R187)	XLO2 + HO ₂ →	(R187)	-	(R187)	-
(R188)	XLO2 + C2O3 → 0.26GLY + 0.77MGLY + 0.35OPEN + 0.65XOPN + HO ₂ + MEO2+ RO2	(R188)	-	(R188)	-
(R189)	XLO2 + RO2 → 0.26GLY + 0.77MGLY+ 0.35OPEN + 0.65XOPN + HO ₂ + RO ₂	(R189)	-	(R189)	-
(R190)	CRES + OH → 0.06CRO + 0.12XO2H+ HO ₂ + 0.13OPEN + 0.73CAT1 + 0.06CO+ 0.06XO2N + 0.18RO2 + 0.06HCHO	(R190)	CRES + OH → 0.03GLY + 0.03OPEN+ HO ₂ + 0.2CRO + 0.73CAT1 + 0.02XO2N+ 0.02RO2	(R190)	-
(R191)	CRES + NO ₃ → 0.30CRO + HNO ₃ + 0.24XO2 + 0.36XO2H + 0.48ALDX + 0.24HCHO+ 0.24MGLY + 0.12OPEN + 0.1XO2N + 0.7RO2+ 0.24CO	(R191)	CRES + NO ₃ → 0.3CRO + HNO ₃ + 0.48XO2 + 0.12XO2H + 0.24GLY + 0.24MGLY+ 0.48OPO3 + 0.1XO2N + 0.7RO2	(R191)	-
(R192)	CRO + NO ₂ → CRON	(R192)	-	(R192)	-
(R193)	CRO + HO ₂ → CRES	(R193)	-	(R193)	-
(R194)	CRON + OH → CRNO	(R194)	CRON + OH → NTR2 + 0.5CRO	(R194)	-
(R195)	CRON + NO ₃ → CRNO + HNO ₃	(R195)	CRON + NO ₃ → NTR2 + 0.5CRO+ HNO ₃	(R195)	-
(R196)		(R196)	CRON + hν → HONO + HO ₂ + HCHO + OPEN	(R196)	-
(R197)	XOPN + hν → CAO2 + 0.7HO ₂ + 0.7CO + 0.3C2O3 + RO2	(R197)	XOPN + hν → 0.4GLY + XO2H+ 0.7HO ₂ + 0.7CO + 0.3C2O3	(R197)	-
(R198)	XOPN + OH → MGLY + CAO2+ XO2H + RO2	(R198)	XOPN + OH → MGLY + 0.4GLY+ 2XO2H + 2RO2	(R198)	-
(R199)	XOPN + O ₃ → 1.2MGLY + 0.5OH+ 0.6C2O3 + 0.1ALD2 + 0.5CO + 0.3XO2H+ 0.3RO2	(R199)	-	(R199)	-
(R200)	XOPN + NO ₃ → 0.5NO ₂ + 0.5NTR+ 0.45XO2H + 0.45XO2 + 0.1XO2N + RO2+ 0.25OPEN + 0.25MGLY	(R200)	XOPN + NO ₃ → 0.5NO ₂ + 0.5NTR2+ 0.45XO2H + 0.45XO2 + 0.1XO2N + RO2+ 0.25OPEN + 0.25MGLY	(R200)	-
(R201)	OPEN + hν → OPO3 + HO ₂ + CO	(R201)	-	(R201)	-
(R202)	OPEN + OH → 0.6OPO3 + 0.4RO2+ 0.4CAO2	(R202)	OPEN + OH → 0.6OPO3+ 0.4XO2H + 0.4RO2 + 0.4GLY	(R202)	-
(R203)	OPEN + O ₃ → 1.4GLY + 0.24MGLY+ 0.5OH + 0.12C2O3 + 0.08HCHO + 0.02ALD2+ 1.98CO + 0.56HO ₂	(R203)	-	(R203)	-
(R204)	OPEN + NO ₃ → OPO3 + HNO ₃	(R204)	-	(R204)	-

Table A1. (continued).

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R205)	CAT1 + OH → CAO2 + RO2	(R205)	CAT1 + OH → 0.14HCHO + 0.2HO ₂ + 0.5CRO	(R205)	-
(R206)	CAT1 + NO ₃ → CRO + HNO ₃	(R206)	-	(R206)	-
(R207)	OPO3 + NO → NO ₂ + XO2H + RO2 + ALDX	(R207)	OPO3 + NO → NO ₂ + 0.5GLY + 0.5CO + 0.8HO ₂ + 0.2CXO3	(R207)	-
(R208)	OPO3 + NO ₂ → OPAN	(R208)	-	(R208)	-
(R209)	OPAN → OPO3 + NO ₂	(R209)	-	(R209)	-
(R210)	OPO3 + HO ₂ → 0.41PACD + 0.15AACD + 0.15O ₃ + 0.44ALDX + 0.44XO2H + 0.44RO2 + 0.44OH	(R210)	-	(R210)	-
(R211)	OPO3 + C2O3 → MEO2 + XO2 + ALDX + 2RO2	(R211)	-	(R211)	-
(R212)	OPO3 + RO2 → 0.8XO2H + 0.8ALDX + 1.8RO2 + 0.2AACD	(R212)	-	(R212)	-
		(R213)	OPAN + OH → 0.5NO ₂ + 0.5GLY + CO + 0.5NTR2	(R213)	-
		(R214)	PANX + OH → ALD2 + NO ₂	(R214)	-
				(R215)	NAPH + OH → 0.15CRES + 0.54XLO2 + 0.6RO2 + 0.24XOPN + 0.24OH + 0.06XO2H + 0.15HO ₂
				(R216)	ECH4 + OH → MEO2 + RO2
				(R217)	XPRP → XO2N + RO2
				(R218)	XPRP → 0.73ACET + 0.27ALDX + 0.27PAR + XO2H + RO2
				(R219)	XPAR → XO2N + RO2
				(R220)	XPAR → 0.13ALDX + 0.87ROR + 0.13XO2H + 0.87XO2 + RO2 - 0.13PAR
		(R215)	NTR2 → HNO ₃	(R221)	-
(R213)	CRNO + NO ₂ → 2NTR				
(R214)	CRNO + O ₃ → CRN2				
(R215)	CRN2 + NO → CRNO + NO ₂				
(R216)	CRN2 + HO ₂ → CRPX				
(R217)	CRPX + hν → CRNO + OH				
(R218)	CRPX + OH → CRN2				
(R219)	CAO2 + NO → 0.86NO ₂ + 0.14NTR + 1.2HO ₂ + 0.34HCHO + 0.34CO				
(R220)	CAO2 + HO ₂ →				
(R221)	CAO2 + C2O3 → HO ₂ + 0.4GLY + MEO2 + RO2				
(R222)	CAO2 + RO2 → HO ₂ + 0.4GLY + RO2				

Table A1. (continued).

Reaction Number	CB6r1 ^a	Reaction Number	CB6r2 ^b	Reaction Number	CB6r3 ^c
(R223) ^d	→ NO	(R216)	-	(R222)	-
(R224) ^d	→ NO ₂	(R217)	-	(R223)	-
(R225) ^d	→ CO	(R218)	-	(R224)	-
(R226) ^d	→ HCHO	(R219)	-	(R225)	-
(R227) ^d	→ ALD2	(R220)	-	(R226)	-
(R228) ^d	→ IOLE	(R221)	-	(R227)	-
(R229) ^d	→ ALDX	(R222)	-	(R228)	-
(R230) ^d	→ ETH	(R223)	-	(R229)	-
(R231) ^d	→ TOL	(R224)	-	(R230)	-
(R232) ^d	→ XYL	(R225)	-	(R231)	-
(R233) ^d	→ ISOP	(R226)	-	(R232)	-
(R234) ^d	→ PAR	(R227)	-	(R233)	-
(R235) ^d	→ OLE	(R228)	-	(R234)	-
(R236) ^e	O ₃ →	(R229)	-	(R235)	-
(R237) ^e	NO →	(R230)	-	(R236)	-
(R238) ^e	NO ₂ →	(R231)	-	(R237)	-
(R239) ^e	HNO ₃ →	(R232)	-	(R238)	-
(R240) ^e	H ₂ O ₂ →	(R233)	-	(R239)	-
(R241) ^e	N ₂ O ₅ →	(R234)	-	(R240)	-
(R242) ^e	CO →	(R235)	-	(R241)	-
(R243) ^e	HONO →	(R236)	-	(R242)	-
(R244) ^e	HCHO →	(R237)	-	(R243)	-

Notes: ^a Yarwood et al. (2012).

^b Ruiz Hildebrandt and Yarwood (2013).

^c Emery et al. (2015).

^d Added to the mechanism to represent surface emissions, not included in the original mechanism.

^e Added to the mechanism to represent dry depositions, not included in the original mechanism.

Acknowledgements. This work was supported by the National Key R&D Program of China (Grant No. 2017YFC0209801) and the National Natural Science Foundation of China (Grant No. 41375044). The numerical calculations in this paper have been done on the high performance

675

computing system in the High Performance Computing Center, Nanjing University of Information Science & Technology. The authors also like to thank Rolf Sander from Max-Planck Institute for Chemistry for his valuable suggestions on the improvements of the model.

Author contributions. Le Cao conceived the idea of the article and wrote the python script for converting the data format. Le Cao also configured and performed the computations. Simeng Li revised the chemical mechanisms and wrote the paper with Le Cao together. Luhang
680 Sun revised the paper and gave valuable suggestions. All the authors listed have read and approved the final manuscript.

Competing interests. The authors declare no conflict of interest.

References

- 685 Ammann, M., Cox, R. A., Crowley, J. N., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume VI – heterogeneous reactions with liquid substrates, *Atmospheric Chemistry and Physics*, 13, 8045–8228, <https://doi.org/10.5194/acp-13-8045-2013>, 2013.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O_x, HO_x, NO_x and SO_x species, *Atmospheric Chemistry and Physics*, 4, 1461–1738, <https://doi.org/10.5194/acp-4-1461-2004>, 2004.
- 690 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and Subcommittee, I.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II - gas phase reactions of organic species, *Atmospheric Chemistry and Physics*, 6, 3625–4055, <https://doi.org/10.5194/acp-6-3625-2006>, 2006.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume III - gas phase reactions of inorganic halogens, *Atmospheric Chemistry and Physics*, 7, 981–1191, <https://doi.org/10.5194/acp-7-981-2007>, 2007.
- 695 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume IV – gas phase reactions of organic halogen species, *Atmospheric Chemistry and Physics*, 8, 4141–4496, <https://doi.org/10.5194/acp-8-4141-2008>, 2008.
- Bash, J. O., Baker, K. R., and Beaver, M. R.: Evaluation of improved land use and canopy representation in BEIS v3.61 with biogenic VOC measurements in California, *Geoscientific Model Development*, 9, 2191–2207, <https://doi.org/10.5194/gmd-9-2191-2016>, 2016.
- 700 Bloss, C., Wagner, V., Jenkin, M. E., Volkamer, R., Bloss, W. J., Lee, J. D., Heard, D. E., Wirtz, K., Martin-Reviejo, M., Rea, G., Wenger, J. C., and Pilling, M. J.: Development of a detailed chemical mechanism (MCMv3.1) for the atmospheric oxidation of aromatic hydrocarbons, *Atmos. Chem. Phys.*, 5, 641–664, <https://doi.org/10.5194/acp-5-641-2005>, 2005.
- Byun, D. and Schere, K. L.: Review of the Governing Equations, Computational Algorithms, and Other Components of the Models-3 Community Multiscale Air Quality (CMAQ) Modeling System, *Applied Mechanics Reviews*, 59, 51–77, <https://doi.org/10.1115/1.2128636>, 705 2006.
- Cao, L., Sihler, H., Platt, U., and Gutheil, E.: Numerical analysis of the chemical kinetic mechanisms of ozone depletion and halogen release in the polar troposphere, *Atmos. Chem. Phys.*, 14, 3771–3787, <https://doi.org/10.5194/acp-14-3771-2014>, 2014.
- Cao, L., Wang, C., Mao, M., Grosshans, H., and Cao, N.: Derivation of the reduced reaction mechanisms of ozone depletion events in the Arctic spring by using concentration sensitivity analysis and principal component analysis, *Atmos. Chem. Phys.*, 16, 14853–14873, 710 <https://doi.org/10.5194/acp-16-14853-2016>, 2016.
- Cao, L., Gao, M., Li, S., Yi, Z., and Meng, X.: Sensitivity analysis of the dependence of the Carbon Bond Mechanism IV (CBM-IV) on the initial air composition under an urban condition, *Atmospheric Environment*, 215, 116860, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2019.116860>, 2019.
- Carter, W. P.: Documentation of the SAPRC-99 chemical mechanism for VOC reactivity assessment, Tech. Rep. 329, California Air Resources Board, 2000a.
- 715 Carter, W. P.: Implementation of the SAPRC-99 chemical mechanism into the Models-3 framework, Tech. rep., United States Environmental Protection Agency, 2000b.

- Carter, W. P.: Development of the SAPRC-07 chemical mechanism, *Atmos. Environ.*, 44, 5324 – 5335, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2010.01.026>, 2010.
- 720 Crowley, J. N., Ammann, M., Cox, R. A., Hynes, R. G., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume V – heterogeneous reactions on solid substrates, *Atmospheric Chemistry and Physics*, 10, 9059–9223, <https://doi.org/10.5194/acp-10-9059-2010>, 2010.
- Derwent, R.: Intercomparison of chemical mechanisms for air quality policy formulation and assessment under North American conditions, *Journal of the Air & Waste Management Association*, 67, 789–796, <https://doi.org/10.1080/10962247.2017.1292969>, <https://doi.org/10.1080/10962247.2017.1292969>, PMID: 28278034, 2017.
- 725 Emery, C., Jung, J., Koo, B., and Yarwood, G.: Final Report, Improvements to CAMx Snow Cover Treatments and Carbon Bond Chemical Mechanism for Winter Ozone, Tech. rep., Ramboll Environ, 2015.
- Emery, C., Koo, B., Hsieh, W., Wentland, A., Wilson, G., and Yarwood, G.: Technical Memorandum to Chris Misenis at U.S. EPA, Tech. rep., 2016.
- 730 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geoscientific Model Development*, 3, 43–67, <https://doi.org/10.5194/gmd-3-43-2010>, <https://gmd.copernicus.org/articles/3/43/2010/>, 2010.
- ENVIRON: User's Guide: Comprehensive Air Quality Model with Extensions (CAMx), Version 6.50, Tech. rep., 2015.
- 735 Gery, M. W., Whitten, G. Z., Killus, J. P., and Dodge, M. C.: A photochemical kinetics mechanism for urban and regional scale computer modeling, *J. Geophys. Res. Atmos.*, 94, 12 925–12 956, 1989.
- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmos. Environ.*, 39, 6957 – 6975, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, <https://doi.org/10.5194/acp-6-3181-2006>, 2006.
- 740 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- 745 Hauglustaine, D. A., Granier, C., Brasseur, G. P., and Mégie, G.: The importance of atmospheric chemistry in the calculation of radiative forcing on the climate system, *Journal of Geophysical Research: Atmospheres*, 99, 1173–1186, <https://doi.org/https://doi.org/10.1029/93JD02987>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/93JD02987>, 1994.
- Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31, 81–104, [https://doi.org/https://doi.org/10.1016/S1352-2310\(96\)00105-7](https://doi.org/https://doi.org/10.1016/S1352-2310(96)00105-7), 1997.
- 750 Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 181–193, <https://doi.org/10.5194/acp-3-181-2003>, 2003.
- Jenkin, M. E., Wyche, K. P., Evans, C. J., Carr, T., Monks, P. S., Alfarra, M. R., Barley, M. H., McFiggans, G. B., Young, J. C., and Rickard, A. R.: Development and chamber evaluation of the MCM v3.2 degradation scheme for β -caryophyllene, *Atmos. Chem. Phys.*, 12, 5275–5308, <https://doi.org/10.5194/acp-12-5275-2012>, 2012.
- 755

- Jenkin, M. E., Young, J. C., and Rickard, A. R.: The MCM v3.3.1 degradation scheme for isoprene, *Atmos. Chem. Phys.*, 15, 11 433–11 459, <https://doi.org/10.5194/acp-15-11433-2015>, 2015.
- Luecken, D., Hutzell, W., and Gipson, G.: Development and analysis of air quality modeling simulations for hazardous air pollutants, *Atmospheric Environment*, 40, 5087 – 5096, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2005.12.044>, 2006.
- 760 Luecken, D., Phillips, S., Sarwar, G., and Jang, C.: Effects of using the CB05 vs. SAPRC99 vs. CB4 chemical mechanism on model predictions: Ozone and gas-phase photochemical precursor concentrations, *Atmospheric Environment*, 42, 5805 – 5820, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2007.08.056>, 2008.
- Luecken, D., Yarwood, G., and Hutzell, W.: Multipollutant modeling of ozone, reactive nitrogen and HAPs across the continental US with CMAQ-CB6, *Atmospheric Environment*, 201, 62 – 72, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2018.11.060>, 2019.
- 765 Luecken, D. J., Napelenok, S. L., Strum, M., Scheffe, R., and Phillips, S.: Sensitivity of Ambient Atmospheric Formaldehyde and Ozone to Precursor Species and Source Types Across the United States, *Environmental Science & Technology*, 52, 4668–4675, <https://doi.org/10.1021/acs.est.7b05509>, 2018.
- Madronich, S. and Flocke, S.: Theoretical Estimation of Biologically Effective UV Radiation at the Earth's Surface, in: *Solar Ultraviolet Radiation*, edited by Zerefos, C. S. and Bais, A. F., pp. 23–48, Springer Berlin Heidelberg, Berlin, Heidelberg, 1997.
- 770 Madronich, S. and Flocke, S.: *The Role of Solar Radiation in Atmospheric Chemistry*, pp. 1–26, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/978-3-540-69044-3_1, 1999.
- Marvin, M. R., Wolfe, G. M., Salawitch, R. J., Canty, T. P., Roberts, S. J., Travis, K. R., Aikin, K. C., de Gouw, J. A., Graus, M., Hanisco, T. F., Holloway, J. S., Hübler, G., Kaiser, J., Keutsch, F. N., Peischl, J., Pollack, I. B., Roberts, J. M., Ryerson, T. B., Veres, P. R., and Warneke, C.: Impact of evolving isoprene mechanisms on simulated formaldehyde: An inter-comparison supported by in situ observations
- 775 from SENEX, *Atmospheric Environment*, 164, 325 – 336, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2017.05.049>, 2017.
- Pierce, T., Geron, C., Bender, L., Dennis, R., Tonnesen, G., and Guenther, A.: Influence of increased isoprene emissions on regional ozone modeling, *Journal of Geophysical Research: Atmospheres*, 103, 25 611–25 629, <https://doi.org/https://doi.org/10.1029/98JD01804>, 1998.
- Ruiz Hildebrandt, L. and Yarwood, G.: Interactions between Organic Aerosol and NO_y: Influence on Oxidant Production, Tech. rep., 2013.
- Sander, S. P., Finlayson-Pitts, B. J., Friedl, R. R., Golden, D. M., Huie, R. E., Keller-Rudek, H., Kolb, C. E., Kurylo, M. J., Molina, M. J.,
- 780 Moortgat, G. K., Orkin, V. L., Ravishankara, A. R., and Wine, P. W.: *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies*. Evaluation No. 15 (JPL Publication 06-2), Jet Propulsion Laboratory, Pasadena, CA, 2006.
- Sandu, A., Verwer, J., Loon, M. V., Carmichael, G., and Seinfeld, J.: Benchmarking stiff ODE solvers for atmospheric chemistry problems I: implicit versus explicit, *Atmos. Environ.*, 31, 3151–3166, 1997.
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, <https://doi.org/10.5194/acp-3-161-2003>, 2003.
- Saylor, R. D. and Ford, G. D.: On the comparison of numerical methods for the integration of kinetic equations in atmospheric chemistry and transport models, *Atmospheric Environment*, 29, 2585 – 2593, [https://doi.org/https://doi.org/10.1016/1352-2310\(95\)00187-4](https://doi.org/https://doi.org/10.1016/1352-2310(95)00187-4), 1995.
- Seinfeld, J. and Pandis, S.: *Atmos. Chem. Phys.: from air pollution to climate change*, A Wiley-Interscience publications, Wiley, 2006.
- 790 Seyfioglu, R., Odabasi, M., and Cetin, E.: Wet and dry deposition of formaldehyde in Izmir, Turkey, *Science of The Total Environment*, 366, 809–818, <https://doi.org/https://doi.org/10.1016/j.scitotenv.2005.08.005>, 2006.
- Stockwell, W. R., Saunders, E., Goliff, W. S., and Fitzgerald, R. M.: A Perspective on the Development of Gas-phase Chemical Mechanisms for Eulerian Air Quality Models, *Journal of the Air & Waste Management Association* (1995), 70, 2019.

- Turányi, T.: KINAL - a program package for kinetic analysis of reaction mechanisms., *Comput. Chem.*, 14, 253–254, 1990a.
- 795 Turányi, T.: Sensitivity analysis of complex kinetic systems. Tools and applications, *J. Math. Chem.*, 5, 203–248, <https://doi.org/10.1007/BF01166355>, 1990b.
- Whitten, G. Z., Heo, G., Kimura, Y., McDonald-Buller, E., Allen, D. T., Carter, W. P., and Yarwood, G.: A new condensed toluene mechanism for Carbon Bond: CB05-TU, *Atmospheric Environment*, 44, 5346 – 5355, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2009.12.029>, 2010.
- 800 Wolfe, G. M., Kaiser, J., Hanisco, T. F., Keutsch, F. N., de Gouw, J. A., Gilman, J. B., Graus, M., Hatch, C. D., Holloway, J., Horowitz, L. W., Lee, B. H., Lerner, B. M., Lopez-Hilifiker, F., Mao, J., Marvin, M. R., Peischl, J., Pollack, I. B., Roberts, J. M., Ryerson, T. B., Thornton, J. A., Veres, P. R., and Warneke, C.: Formaldehyde production from isoprene oxidation across NO_x regimes, *Atmospheric Chemistry and Physics*, 16, 2597–2610, <https://doi.org/10.5194/acp-16-2597-2016>, 2016.
- Yarwood, G., Rao, S., Yocke, M., and Whitten, G.: Updates to the carbon bond chemical mechanism: CB05, Tech. rep., U.S. Environmental
805 Protection Agency, 2005.
- Yarwood, G., Jung, J., Whitten, G., Heo, G., J, M., and M, E.: Updates to the Carbon Bond Mechanism for Version 6 (CB6), in: 9th Annual CMAS Conference, pp. 1–4, Chapel Hill, NC, 2010.
- Yarwood, G., Heo, G., Carter, W., and GZ, W.: Final Report, Environmental Chamber Experiments to Evaluate NO_x Sinks and Recycling in Atmospheric Chemical Mechanisms, Tech. rep., Texas Air Quality Research Program, The University of Texas at Austin, 2012.
- 810 Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, *J. Geophys. Res. Atmos.*, 104, 30 387–30 415, <https://doi.org/10.1029/1999JD900876>, 1999.
- Zhang, Q., Jiang, X., Tong, D., Davis, S. J., Zhao, H., Geng, G., Feng, T., Zheng, B., Lu, Z., Streets, D. G., Ni, R., Brauer, M., van Donkelaar, A., Martin, R. V., Huo, H., Liu, Z., Pan, D., Kan, H., Yan, Y., Lin, J., He, K., and Guan, D.: Transboundary health impacts of transported global air pollution and international trade, *Nature*, 543, <https://doi.org/10.1038/nature21712>, 2017a.
- 815 Zhang, R., Cohan, A., Biazar, A. P., and Cohan, D. S.: Source apportionment of biogenic contributions to ozone formation over the United States, *Atmospheric Environment*, 164, 8–19, 2017b.

Table 1. The initial air composition and the surface emission intensities used in **two different simulation scenarios**. This initial condition was adapted from Saylor and Ford (1995) and Sandu et al. (1997), and represents a heavily polluted environment with a 70% relative humidity.

Species	Initial Concentration (ppb)	Weak Emission (ppb h ⁻¹)	Strong Emission (ppb h ⁻¹)
NO	5	0.01	0.25
NO ₂	2	0.01	0.05
HONO	1	-	-
O ₃	100	-	-
CO	300	-	2.00
HCHO	10	-	0.20
ALD2	2.2	-	0.04
IOLE	6.7	-	0.13
ALDX	1.1	-	0.02
PAN	1	-	-
ETH	10	-	0.20
TOL	10	-	0.20
XYL	10	-	0.20
ISOP	10	0.10	1.00
PAR	50	-	2.00
OLE	10	-	1.00
H ₂	560	-	-
CH ₄	1850	-	-
H ₂ O	2.17 × 10 ⁷	-	-

Table 2. Dry deposition velocities used in model simulations for different atmospheric constituents.

Species	Deposition Velocity (cm s^{-1})	Reference
O ₃	0.4	Seinfeld and Pandis (2006)
NO	0.016	Seinfeld and Pandis (2006)
NO ₂	0.1	Seinfeld and Pandis (2006)
HNO ₃	4.0	Seinfeld and Pandis (2006)
H ₂ O ₂	0.5	Seinfeld and Pandis (2006)
N ₂ O ₅	4.0	Hauglustaine et al. (1994)
CO	0.03	Seinfeld and Pandis (2006)
HONO	4.0	Hauglustaine et al. (1994)
HCHO	6.0	Seyfioglu et al. (2006)






Figure 1. Simulated temporal evolutions of O_3 , NO_x and HCHO by using different versions of the CB6 mechanism, when the surface emission is weak.




Figure 2. Averaged ozone sensitivity to the CB6r3 mechanism over the 7-th day, when the surface emission is **weak**. Note that the horizontal scales of the sub-figures are different. All the values of the sensitivities shown in this figure can be found in Tab. S1 of the supplementary material.




Figure 3. Averaged NO_x sensitivity to the CB6r3 mechanism over the 7-th day, when the surface emission is **weak**. Note that the horizontal scales of the sub-figures are different. All the values of the sensitivities shown in this figure can be found in Tab. S1 of the supplementary material.




Figure 4. Averaged HCHO sensitivity to the CB6r3 mechanism over the 7-th day, when the surface emission is **weak**. Note that the horizontal scales of the sub-figures are different. All the values of the sensitivities shown in this figure can be found in Tab. S1 of the supplementary material.






Figure 5. Simulated temporal evolution of O_3 , NO_x and HCHO by using different versions of the CB6 mechanism, when the surface emission is **strong**.




Figure 6. Averaged ozone sensitivity to the CB6r3 mechanism over the 7-th day, when the surface emission is **strong**. All the values of the sensitivities shown in this figure can be found in Tab. S1 of the supplementary material.




Figure 7. Averaged NO_x sensitivity to the CB6r3 mechanism over the 7-th day, when the surface emission is **strong**. All the values of the sensitivities shown in this figure can be found in Tab. S1 of the supplementary material.




Figure 8. Averaged HCHO sensitivity to the CB6r3 mechanism over the 7-th day, when the surface emission is **strong**. All the values of the sensitivities shown in this figure can be found in Tab. S1 of the supplementary material.