

A New Conceptual Model for Adiabatic Fog

Felipe Toledo¹, Martial Haeffelin², Eivind Wærsted³, and Jean-Charles Dupont⁴

¹Laboratoire de Météorologie Dynamique, École Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau, France

²Institut Pierre Simon Laplace, École Polytechnique, CNRS, Institut Polytechnique de Paris, 91128 Palaiseau, France

³Laboratoire de Météorologie Dynamique, École Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau, France.

Current affiliation: The Norwegian Meteorological Institute, Henrik Mohns Plass 1, 0313, Oslo, Norway

⁴Institut Pierre-Simon Laplace, École Polytechnique, UVSQ, Université Paris-Saclay, 91128 Palaiseau, France

Correspondence: Felipe Toledo (ftoledo@lmd.polytechnique.fr)

Abstract.

We Visibility reduction caused by fog can be hazardous for human activities, especially for the transport sector. Previous studies show that this problem could be mitigated by improving nowcasting of fog dissipation. To address this issue, we propose a new paradigm to describe the temporal evolution of continental fog layers. This paradigm defines fog which could potentially 5 improve our understanding of the life cycle of adiabatic continental fogs, and of the conditions that must take place for fog dissipation.

For this purpose, adiabatic fog is defined as a layer saturated from filled with suspended liquid water droplets, extending from an upper boundary all the way down to the surface, with a saturated adiabatic temperature profile. In this layer, the surface to a known upper boundary, and whose liquid water path (LWP) exceeds must exceed a critical value, the critical liquid water 10 path (CLWP). When the LWP is less than the CLWP the fog water cannot, the amount of fog liquid water is not sufficient to extend all the way down to the surface, leading to a surface horizontal visibility greater than 1 km. On the opposite Conversely, when the LWP is larger than exceeds the CLWP, the fog water extends all the way to amount of fog water is enough to reach the surface, inducing a horizontal visibility less than 1 km. The excess water with respect to the critical value is then defined as the reservoir liquid water path (RLWP).

15 The new fog paradigm is formulated as a conceptual model that relates the liquid water path of adiabatic fog with its thickness and surface liquid water content, and allows the critical and reservoir liquid water paths to be computed. Both variables can be tracked in real time using vertical profiling measurements, enabling a real time diagnostic of fog status.

The conceptual model is tested using data from seven years of measurements performed at the SIRTA observatory, combining 20 cloud radar, microwave radiometer, ceilometer, scatterometer and weather station measurements. In this time period we found 80 fog events with reliable measurements, with 56 of these lasting more than three hours.

The paper presents the conceptual model and its capability to derive the LWP from the fog CTH and surface horizontal visibility with an RMS uncertainty of 10.5 g m^{-2} . The impact of fog liquid water path and fog top height variations on fog life cycle (formation to dissipation) is presented based on four case studies, and statistics derived from 56 fog events. Our results show in particular that the, based on measurements and an empirical parametrization for the adiabaticity, validate the 25 applicability of the model. The calculated reservoir liquid water path is consistently positive during the mature phase of the

~~fog and that it fog, and~~ starts to decrease quasi monotonously about one hour before dissipation, reaching a near-zero value at the time of dissipation. ~~The Hence, the~~ reservoir liquid water path and its time derivative could ~~hence~~ be used as ~~an indicator for indicators of the~~ life cycle stage~~and support short range forecasting~~, ~~to support nowcasting~~ of fog dissipation.

1 Introduction

30 Fog occurs due to multiple processes that lead to ~~saturation of the air near the surface, through cooling water vapor saturation in the air close to the surface. Water vapor saturation can be caused by a reduction of air temperature, such as due to radiative cooling, turbulent heat exchange, diffusion, adiabatic cooling through lifting, advection, and through moistening of the air, such as. It can also occur by air moistening, due to water~~ evaporation from the surface, evaporation of drizzle, advection of moist air, and vertical mixing (Brown and Roach, 1976; Gultepe et al., 2007; Dupont et al., 2012). ~~Similarly~~ ~~On the contrary,~~ 35 fog dissipates as a result of warming and drying of the air near the surface, and also through the removal of droplets by precipitation (Brown and Roach, 1976; Haeflalin et al., 2010; Wærsted et al., 2017, 2019).

Stable fog and adiabatic fog should be distinguished because radiative, thermodynamic, dynamic and microphysical processes are significantly contrasted in the two types of fog. In a stable fog layer, the equivalent potential temperature increases with height, which inhibits vertical mixing. The surface is therefore weakly coupled with the fog top. Stable fog remains shallow and contains small amounts of liquid water, limiting the radiative cooling of the fog layer. In contrast, in an adiabatic fog the stability is close to neutral, enabling rapid vertical mixing, so that the surface and fog top are strongly coupled (Price, 2011; Porson et al., 2011). An adiabatic fog behaves similarly to stratocumulus clouds on top of convective boundary layers (Cermak and Bendix, 2011). The processes of ~~stratocumulus clouds adiabatic fogs~~ have been studied extensively in the past with large-eddy simulation (LES) and numerical weather prediction (NWP) models (Nakanishi, 2000; Porson et al., 2011; Bergot, 2013, 45 Wærsted et al., 2019).

An adiabatic fog or stratiform cloud cools at its top from emission of long wave radiation, which destabilises the cloud and leads to convective mixing. When the cloud is coupled with the land surface, the destabilising process can be further strengthened by heat fluxes from below due to soil heat (Price, 2011). A thermal inversion develops right above the cooling cloud fog top and limits the coupling between the cloud and free atmosphere above. The thermal inversion defines the upper 50 boundary of the adiabatic fog. The lower boundary of the stratiform cloud layer varies in time and space depending the amount of liquid water present in the cloud. For the adiabatic fog, the lower boundary is defined by the surface and is therefore fixed. Hence a fog layer may not grow geometrically deeper when the amount of liquid water increases.

Cermak and Bendix (2011) define fog and stratiform clouds based on cloud layer top altitude and liquid water content that follows a sub-adiabatic profile. A fog layer is thus defined as a stratiform cloud that contains sufficient liquid water to reach 55 down to the surface.

Wærsted et al. (2019) showed using a large eddy-simulation model and remote sensing measurements that dissipation of fog can occur due to both reduction of liquid water content of the fog layer and increase of fog top height. Dissipation is defined here as removal of fog droplets leading to visibility increasing above 1 km at screen-level height. The simulations reveal a

similar behavior as proposed by Cermak and Bendix (2011). For a given fog top height, if the liquid water path contained in the
60 fog layer becomes insufficient, the fog base lifts from the ground, which can be interpreted as fog dissipation through lifting
into a stratiform cloud.

In adiabatic clouds, the thickness can be approximated from liquid water path. Brenguier et al. (2000) state that liquid water path is proportional to the square of cloud thickness. A precise quantification of the relationship between fog thickness and fog liquid water path is lacking in the literature.

65 In this article we present a conceptual model that relates the liquid water path of adiabatic fog to its geometrical thickness and surface liquid water content. The conceptual model enables an estimation of the minimum amount of column liquid water that is necessary to reach a visibility less than 1000 meters at the surface, defined as the critical liquid water path, and a calculation of the excess water that enhances fog persistence, defined as the reservoir liquid water path. The model also enables a quantification of the impact of liquid water path and geometrical thickness variations on the reservoir, a characteristic that
70 could be later used to improve fog forecasting tools.

The conceptual model theory is explained in Section 2. In Section 3, we present all measurements used to construct and evaluate the conceptual model. In Section 4 we derive a parametrization for fog adiabaticity using historical data, and we compare the conceptual model predictions with fog thickness, liquid water path and surface liquid water content observations. In Section 5 we present case studies to exemplify how conceptual model variables enable us to understand fog evolution, and
75 statistical results of fog behavior during its formation, middle life and dissipation phases.

2 Fog Conceptual Model

2.1 Fog LWP Conceptual Model

The hypothesis of this work is that there is a general rule, dependent on fog macroscopic properties, that must be fulfilled when the when a fog layer persists at the surface level. To deduce this rule layer is well-mixed, the persistence or not of fog at surface level will be determined by vertically integrated quantities of the whole fog layer, and in particular the integrated liquid water content. To test this hypothesis we develop a unidimensional model for a fog column, based on previous models for stratus clouds.

For stratus clouds, cloud Liquid Water Content (LWC) increases with height can be modelled using Eq. (1) (Albrecht et al., 1990) (Betts, 1982; Albrecht et al., 1990; Cermak and Bendix, 2011). In this equation, z is the vertical distance above the Cloud Base
85 Height (CBH), which increases until reaching the Cloud Top Height (CTH). $\Gamma_{ad}(T, P)$ is the negative of the change in saturation mixing ratio with height for an ideal adiabatic cloud, and $\alpha(z)$ is the local adiabaticity, defined as the ratio between the real and the ideal adiabatic liquid water content change with height. $\Gamma_{ad}(T, P)$ is a quantity that depends on the local temperature T and pressure P . The equation used for its calculation can be found in appendix A.

$$\frac{dLWC(z)}{dz} = \alpha(z) \Gamma_{ad}(T, P) \quad (1)$$

90 This model can also be applied for well mixed fog layers, where the adiabatic profile assumption is valid. ~~This happens when fog is opaque, and thus radiative cooling happens~~ Fog layers that are radiatively opaque will cool almost exclusively at the fog top. ~~This cooling introduces instability, enhancing vertical mixing due to~~ and therefore tend towards static instability, which causes mixing through convective turbulence. During day time, convection is reinforced by sensible heat release from the surface. This mixing induces the formation of a saturated adiabatic temperature profile in fog layers (Roach et al., 1976; 95 Boutle et al., 2018; Wærsted et al., 2019).

However, there is one key difference in fog layers that must be considered when integrating (1). In stratus clouds, it is assumed that the LWC at the cloud base is zero, because condensation is starting gradually from unsaturated air, and therefore there is a smooth transition between dry and moist air.

This smooth transition does not ~~hold in the case of~~ occur in fog layers. ~~Here, CBH is not limited by the dry-moist air transition but rather by a solid boundary, the surface. The surface limits vertical fog development, and causes an excess LWC~~ In this case, the cloud base is fixed by the surface height, and has a positive LWC. These characteristics are the reason for the visibility reduction at the surface. It is worth noting that for adiabatic fog, the surface presence could produce a larger accumulation of LWC with respect to other clouds of the same thickness. This could happen because in this fog type, water vapor condensation can occur rapidly at the fog base when compared to a cloud. ~~This larger LWC is what drives the visibility reduction at the surface~~ top, due to radiative cooling (e.g. Wærsted et al. (2017)), and this LWC would be redistributed in a layer of a fixed vertical extent. Vertical redistribution would happen because in adiabatic fog, the stability is close to neutral and therefore vertical circulation caused by surface heating, or cloud top radiative cooling, are possible (Smith et al., 2018).

105 Thus, when integrating Eq. (1) it is necessary to account for a non-zero Surface Liquid Water Content (LWC_0). Since fog (and stratus clouds) are shallow, ~~the~~ ~~their~~ LWC increases with height, and $\Gamma_{ad}(T, P)$ can be assumed constant for the whole 110 layer (Albrecht et al., 1990; Braun et al., 2018). This leads to the LWC formulation of Eq. (2).

$$LWC(z) = \int_{z'=0}^{z'=z} \alpha(z') \Gamma_{ad}(T, P) \underline{z} \underline{dz'} + LWC_0 \quad (2)$$

The blue curve of Fig. 1 (a) illustrates how LWC behaves in well mixed fog. For most of the fog layer thickness, LWC increases with height due to upward motions of moisture from the surface and within the cloud (Oliver et al., 1978; Manton, 1983; Walker, 2003; Cermak and Bendix, 2011). Then, when approaching fog top from below, the LWC change with height 115 decreases until becoming a net reduction of LWC near the top. This decrease is due to entrainment of dry-air at the top, which leads to a quick decline in droplet size and LWC (Brown and Roach, 1976; Roach et al., 1982; Driedonks and Duynkerke, 1989; Hoffmann and Roth, 1989; Boers and Mitchell, 1994; Cermak and Bendix, 2011).

120 Fog LWP is defined as the integral of LWC(z) in the fog column. Its formulation as a function of adiabaticity is presented in Eq. (3b), where z is the height above the surface. Since in fog the CBH is always at the surface, fog thickness is completely defined by its CTH.

$$LWP = \int_{z=0}^{z=CTH} \underline{\underline{LWC(z)}} dz \quad (3a)$$

$$= \int_{z=0}^{z=CTH} \left(\int_{z'=0}^{z'=z} \alpha(z') \Gamma_{ad}(T, P) \underline{\underline{dz'}} + LWC_0 \right) dz \quad (3b)$$

$$LWP = \frac{1}{2} \alpha_{eq} \Gamma_{ad}(T, P) CTH^2 + LWC_0 CTH \quad (3c)$$

125 To simplify the calculation of the integral in Eq. (3b), which requires the knowledge of the adiabaticity profile $\alpha(z)$, we introduce the Equivalent Adiabaticity α_{eq} term. The Equivalent Adiabaticity is defined as ~~a~~^{the} constant adiabaticity value that would ~~provide~~^{give} the same LWP as that derived from ~~value~~, when replacing $\alpha(z')$ in Eq. (3b). ~~In our study, α_{eq} is estimated using a parametrization derived from 7 years of fog observations at the SIRTA observatory (see Sect. 4.2). It is worth mentioning that this parameter is also defined in literature as the in-cloud mixing parameter β (e.g. Cermak and Bendix (2011)), which is equivalent to α_{eq} and can be easily transformed using the rule $\alpha_{eq} = (1 - \beta)$.~~


130 ~~The Equivalent Adiabaticity~~ The equivalent adiabaticity enables the definition of the Fog LWP Conceptual Model, indicated Conceptual Model LWP, in Eq. (3c).

The Conceptual Model LWP has the same value as Fog LWP, but its LWC(z) profile is different because it uses a constant adiabaticity value.

135 ~~The relationship between Fog and Conceptual Model LWP~~ This difference is illustrated in Fig. 1 (a). Fog LWP is the light blue surface, bound by the fog LWC curve with varying adiabaticity with height. Whereas, the Conceptual Model LWP corresponds to the dashed area. Its LWC increases linearly with height because of the constant adiabaticity value. This figure shows that both Fog and the Conceptual Model have the same Surface LWC for a given LWP value. Considering that surface LWC can be linked to visibility, this implies that for a given fog LWP value, the Conceptual Model should predict realistic visibility values at the surface.

140 ~~In our study, α_{eq} is estimated using a parametrization derived from 7 years of fog observations at the SIRTA observatory (see Sect. 4.2). It is worth mentioning that this parameter is also defined in literature as the in-cloud mixing parameter β (e.g. Betts (1982); Cermak and Bendix (2011)), which is equivalent to α_{eq} and can be easily transformed using the rule $\alpha_{eq} = (1 - \beta)$.~~

(a) Relationship between Fog and Conceptual Model LWC and LWP



(b) Conceptual Model Critical and Reservoir LWP




Figure 1. (a) Illustration of the relationship between Fog, Conceptual Model and adiabatic LWC with vs height. In all cases LWC changes with height from its surface value until reaching fog top (CTH). Fog and Conceptual Model LWP have the same value. (b) Representation of the Critical LWP (CLWP) and Reservoir LWP (RLWP) with respect to fog LWP. CLWP is predicted LWP value that fog should have when visibility equals 1000 meters at the surface (with an associated surface LWC defined as LWC_c). RLWP is the difference between fog and the CLWP, and represents the excess water that enables fog persistence.

2.2 Critical and Reservoir LWP

145 Wærsted (2018) found that fog dissipation by lifting of its base is explained by a deficit in LWP considering a given fog thickness. This motivated the definition of a Critical Liquid Water Path (CLWP), which is the minimum amount of LWP needed for a cloud to reach the surface, and reduce horizontal visibility below 1000 meters.

150 CLWP is formulated from Eq. (3c), assuming a Critical Liquid Water Content LWC_c at the surface. LWC_c is the LWC that would cause a 1000 meters visibility, calculated using the parametrization derived by Gultepe et al. (2006) (appendix B). This parametrization indicates that the LWC_c has a value of $\approx 0.02 \text{ gm}^{-3}$.

$$CLWP = \frac{1}{2} \alpha_{eq} \Gamma_{ad}(T, P) CTH^2 + LWC_c CTH \quad (4)$$

When fog is present, its LWP value must be always larger than the CLWP. This property motivates the definition of an additional parameter, the Reservoir Liquid Water Path (RLWP). RLWP is a quantitative metric on how far fog is from dissipation, and is calculated using Eq. (5).

$$155 \quad RLWP = LWP - CLWP = LWP - \frac{1}{2} \alpha_{eq} \Gamma_{ad}(T, P) CTH^2 - LWC_c CTH \quad (5)$$

The relationship between CLWP and RLWP is illustrated in Fig. 1 (b). In this case, we have a fog with a given cloud top height CTH and a liquid water content LWP, that are associated ~~to with~~ a liquid water content LWC_0 at the surface. This LWC is greater than the critical value LWC_c , because visibility is less than 1000 m. The CLWP of this fog, indicated by the red surface to the left, is calculated using Eq. (4). Its value indicates the minimum LWP that fog can have before reducing surface 160 LWC below its critical value, which could cause an increase of visibility above 1000 meters. All excess liquid water above the CLWP value creates the RLWP, indicated by the green surface to the right, and corresponds to all the excess LWP that must be removed before fog can dissipate at the surface.

3 Dataset and Data Treatment Methodology

The dataset used to study the Conceptual Model formulation consists on seven years of fog observations made at the SIRTA 165 atmospheric observatory, from July of 2013 to March of 2020 (Haeffelin et al., 2005). This observatory is located 156 m above sea level, approximately 20 km south of Paris ($48^{\circ}43'N$, $2^{\circ}12'E$) in a location with a relatively high fog incidence (about 30 fog events per year).

The observatory data must be treated to transform raw measurements into Conceptual Model variables. Section 3.1 indicates which instruments are used in this study, Sec. 3.2 describes how fog events are detected, and how their formation and dissipation 170 time is identified, and Sec. 3.3 explains the processing of raw observations into Conceptual Model variables.

After data treatment, an additional data quality control stage is performed to remove from the data pool the fog cases with measurements taken under non optimal conditions. The criteria used is explained in Sec. 3.4. A summary of the complete data processing is shown in Fig. 2.

3.1 Observations

175 The SIRTA observatory is equipped with a large array of instruments, tailored for observing fog and fog processes (Haeffelin et al., 2010; Wærsted, 2018). A subset of these instruments is selected for studying the proposed conceptual model, based on the required inputs. These instruments are listed in Table 1.

180 Data from three remote sensing instruments is used: a CL31 Ceilometer, a BASTA Cloud Radar and a HATPRO Microwave Radiometer. The CL31 is a widely used instrument for Cloud Base Height (CBH) detection, with a vertical resolution of 15 meters (Kotthaus et al., 2016). In this study it is used to retrieve the CBH of low stratus clouds preceding fog events, and to track CBH lifting during temporary or definitive dissipation of the fog layer.

185 The Cloud Radar BASTA is a 95 GHz FMCW radar used to retrieve vertical profiles of cloud reflectivity, up to 12 km of height (Delanoë et al., 2016). It operates continuously alternating between 12.5, 25 and 100 m resolution modes every 12 seconds. The 12.5 m mode has the highest vertical resolution and therefore it is used to retrieve fog CTH. Meanwhile, the 100 m mode is the most sensitive and reaches the highest altitude of 12 km, and therefore is used to detect the presence of clouds above the fog layer.

190 The multi-wavelength microwave radiometer (MWR) HATPRO measures the integrated LWP of the atmospheric column. The manufacturer specified uncertainty of the LWP product is of $\pm 20 \text{ g m}^{-2}$, but for relatively small LWP ($< 40 \text{ g m}^{-2}$), investigations indicate that the uncertainty is within $\pm 5\text{--}10 \text{ g m}^{-2}$, at least when the fog forms in clear sky so that a possible time-independent bias can be corrected for (Marke et al., 2016; Wærsted et al., 2017). When no other cloud is present above the fog layer, LWP measured by the MWR will correspond to fog LWP. Thus, MWR and Cloud Radar data can be combined to perform reliable fog LWP retrievals.

195 These remote sensing instruments are complemented by a weather station 2 meters above the surface, and two Scatterometers, at 4 and 20 meters above the surface. The weather station provides the thermodynamic data necessary to calculate the saturated adiabatic lapse rate $\Gamma_{ad}(T, P)$, and the 4-m scatterometer provides the visibility data used to detect fog events and to calculate fog LWC at the surface. Visibility data is also used to complement the CL31 CBH estimation for very low cloud layers.

3.2 Fog event detection

200 Fog periods are identified using a scheme based on previous work done by Tardif and Rasmussen (2007); Wærsted et al. (2019). This method requires the re-sampling of the surface visibility time series to 5 minute blocks. Each 5 min block is assigned a "fog" or "clear" value, depending on the distribution of visibility in its time period. A block is assigned the "fog" value when more than half of the visibility measurements are less than 1000 m, and is assigned "clear" otherwise.




Figure 2. Summary of the data treatment and calculation methodology. The procedure can be separated in three main stages: First, data preparation consists in identifying fog periods from historical visibility measurements, and in gathering raw instrumental information for these periods. Second, data is re-sampled and homogenized into 5 minute time blocks. First order products such as fog CTH, LWP, among others are calculated. Third, the data treated in the second stage is used to calculate conceptual model variables. An additional data quality control stage is included, to check if fog the variables of each identified period were retrieved under reliable instrument operation operating conditions of the instruments.

Table 1. List of instruments and measurements used in this study.

Instrument	Measured Quantity	Vertical Range and Resolution (RA) and Resolution (RE)	Time Res.
905 nm Ceilometer <i>Vaisala CL31</i>	Attenuated backscatter ($\text{m}^{-1} \text{ sr}^{-1}$)	RA 0-7600 m, RE 15 m	60 s
14-Ch. Microwave Radiometer <i>RPG HATPRO</i>	Liquid Water Path (g m^{-2})	Integrated column	60 s
95 GHz FMCW Cloud Radar <i>BASTA</i>	Radar Equivalent Reflectivity (dBZ)	RA 85-6000 m, RE 12.5 m RA 100-12000 m, RE 100 m	12 s 12 s
550 nm Scatterometer <i>Degreane DF320/DF20+</i>	Visibility (m)	At 4 m above ground At 20 m above ground	60 s 60 s
Thermometer <i>Guilcor PT100</i>	Air Temperature (K)	At 2 m above ground	60 s
Barometer <i>Druck RPT410F</i>	Surface Pressure (Pa)	At 2 m above ground	60 s

After assigning values to each block of the complete visibility time series, we analyze groups of five consecutive blocks in a sliding manner. These five contiguous blocks are defined as a construct, and its value is positive when the central and at least
205 two other are fog blocks, and negative otherwise.

A fog event forms when a positive construct is encountered, with a formation time defined as the central time of the first fog block in the construct. Conversely, a fog event dissipates when the last positive construct is followed by either a negative construct or three consecutive clear blocks. Fog dissipation time is set as the central time of the block immediately after the last fog block in the last positive construct. Fog events separated by less than 1 hr are merged, and all fog events lasting less
210 than 1 hr are discarded. This algorithm provides the formation and dissipation time of 217 fog events between July 2013 and March 2020. **It's worth noting that this method, based on visibility measurements only, does not classify the fog type. Hence, all fog types are considered in this study.**

3.3 Data processing

After identifying the fog events, it is necessary to process raw measurements from the instruments into information that can
215 be used by the conceptual model. To **enable the study of study** the conceptual model variables during fog **itself events**, and the time period surrounding **itthem**, observational data is automatically processed **and re-sampled to 5 min time blocks, covering the period** from 3 hours before fog formation to 3 hours after fog dissipation.

CBH is retrieved using a threshold value of $2 \cdot 10^{-4} \text{ m}^{-1} \text{ sr}^{-1}$ on the CL31 attenuated backscatter measurements, following the method of Haeffelin et al. (2016). When the liquid layer is closer than 15 m to the ground, the CL31 cannot identify the
220 CBH anymore and therefore the Scatterometer measurements are checked, setting the CBH as 0 m when visibility drops below

1000 m. Both CBH and visibility measurements are averaged to five minute time blocks, matching the blocks used by the fog detection algorithm.

The Cloud Radar is used to retrieve fog CTH and to detect the presence of higher clouds above the fog layer, based on its vertical reflectivity profile (Wærsted et al., 2019). To retrieve CTH, reflectivity signals in each radar gate are analyzed, starting 225 from the gate closest to the CBH and checking one gate at a time, going upwards. CTH is estimated as the height of the gate under the first gate where no cloud signal is detected. A gate is considered to have a valid cloud signal if more than half of the reflectivity samples in a five minute time block are not removed by the automatic noise filtering algorithm of the radar (Delanoë et al., 2016). As with CBH, time blocks used in CTH retrievals match those defined for fog detection.

A limitation of this method is that the minimum detectable CTH is of 85 meters. Under this height, radar interference 230 becomes very significant, making the differentiation between a valid cloud signal and noise very difficult. ~~Therefore, we decide to not use data associated with CTH retrievals below 85 meters~~In this situation the CTH retrieval is not possible, and therefore the associated time block would not have a valid CTH value.

~~Additionally, radar data treatment creates~~ Radar data is also used to create a flag indicating the possible presence of liquid clouds above the fog layer ~~if when~~ another valid signal is observed above fog CTH, within the first kilometer for the 12.5 m 235 resolution mode, or within the first 6000 m for the 100 m resolution mode. This flag is used in LWP retrievals, as explained below.

The HATPRO Microwave Radiometer ~~is used to perform~~ performs LWP retrievals of fog. ~~LWP measured by HATPRO is averaged every 60 s, which are then averaged and re-sampled~~ to the 5 min time blocks and then is filtered using the radar higher cloud flag block grid. Additionally, when a given time block has an associated flag indicating the possible presence of 240 higher liquid clouds, the LWP sample is declared not valid. This is done to ~~avoid~~ ensure that the LWP samples are reliable, by avoiding a possible fog LWP overestimation when liquid clouds are present~~above the fog layer~~.

Time series of surface temperature and pressure are all averaged to match the 5 minute time blocks. The saturated adiabatic lapse rate $\Gamma_{ad}(T, P)$ is calculated for each of these time blocks using these measurements and the equations in appendix A.

In this scheme, it is important to note that to have a valid sample of conceptual model variables in a given 5 min time block, the block must have valid measurements of fog CTH, LWP, surface visibility, and surface temperature and pressure. Therefore, it is possible to have fog cases without valid samples of conceptual model variables for some time periods. We decided to use these cases (if they comply with the data quality control of Sect. 3.4), and to consider all the samples with valid conceptual model calculations for the statistical analyses.

3.4 Data quality control

250 After data treatment is complete for all automatically detected fog events, a manual check is done to remove cases where data is unreliable. This happens when instruments operate under non optimal conditions, or when the upper liquid cloud flagging algorithm did not work correctly.

This control consist on accepting or removing complete fog cases and their associated dataset. A fog case is removed from the data pool if measurements taken when the fog takes place comply with at least one of the following criteria:

255 1. Data is taken during or after strong precipitation: Strong precipitation wet the Microwave Radiometer radome, leading
to unreliable LWP retrievals for an unpredictable period of time that can last up to hours, even when following all
maintenance instructions (Görsdorf et al., 2020). Additionally, strong rain leads to difficulties in identifying the fog CTH
because the strong reflectivity from rain hides the weaker returns from suspended fog droplets.

260 2. There are no valid data blocks: No CTH or LWP retrievals could be made for the given fog event. This can happen when
fog is thinner than 85 meters, or when liquid clouds are present above fog for the complete event duration.

265 3. Fog and Cloud borders are not well identified: In some cases the automatic cloud border detection algorithm fails,
leading to unfiltered LWP retrievals with liquid clouds above, or to a bad estimation of fog CTH when upper clouds
are too close to the fog layer. The latter can be seen in the radar data as multilayer fog formed by the union of two
previously independent cloud layers. This situation departs from the single well mixed layer assumption, and therefore
the conceptual model is not applicable.

The quicklooks for the accepted and rejected fog cases are available in the article supplementary material. After this stage we end with 80 valid fog cases and 137 rejected cases, where 50 were removed because of criterion 1, 69 because of criterion 2 and 18 because of criterion 3. These 80 valid fog cases ~~pass to the next stage~~ have at least one valid sample of conceptual model variables (see Sect. 3.3), which are then used in the next stages of data analysis and results.

270 **4 Data Analysis and Results**

4.1 Fog Adiabaticity

A key parameter in the calculation of the CLWP is the Equivalent Fog adiabaticity α_{eq} (Eq. (4)). This parameter has been previously studied in literature for boundary layer stratocumulus and stratus clouds, where typically observed values of α_{eq} range between 0.6 and 0.9 (Slingo et al., 1982; Boers et al., 1990; Boers and Mitchell, 1994; Braun et al., 2018). In this situation, clouds have an adiabatic profile and are buoyant (Betts, 1982). Buoyancy is important because it is necessary to have dissipation by lifting of the fog base.

~~It~~ Hence, it is interesting to study whether these ~~adiabaticity~~ values also apply to fog, ~~since fog which~~ is a special ~~ease of~~ cloud whose vertical development is limited by ~~cloud case with~~ a solid lower boundary at the surface. Therefore, we use ~~our~~ the complete database to calculate α_{eq} by closure, with Eq. (6). This equation is an inversion of the conceptual model formulation of Eq. (3c), and enables an estimation of the adiabaticity ~~while~~ correcting the impact of ~~excessive LWC accumulation caused by the solid lower boundary~~ the LWC accumulation at the fog base. We only perform α_{eq} retrievals when visibility is below 2000 m, in order to remain close to fog conditions, ~~where the conceptual model is valid~~.

$$\alpha_{eq}^{closure} = \frac{2(LWP - LWC_0 CTH)}{\Gamma_{ad}(T, P) CTH^2} \quad (6)$$

Figure 3 (a) shows the relationship between resulting equivalent adiabaticity $\alpha_{eq}^{closure}$, versus CTH and LWP. The results
285 indicate that fog layers with small LWP are characterized by large ranges $\alpha_{eq}^{closure}$ increases for greater values of LWP and
CTH. In addition, negative adiabaticity values are found for lower LWP values, specially below 30 g m^{-2} .

To study this behavior in more detail, Figure 3 (b) shows a boxplot with the statistics of $\alpha_{eq}^{closure}$, and sometimes it can even
reach for different LWP ranges. Here we observe that negative adiabaticity values become frequent when the LWP is below the
30-40 g m^{-2} range, until occurring for more than half of the samples when the LWP is below 20 g m^{-2} .

290 This can be explained by considering that fog with LWP less than 30 g m^{-2} is not optically thick (Wærsted et al., 2017).
Under this condition, the liquid water condensation happens everywhere in the liquid layer, but it is mostly driven by surface
cooling (Wærsted et al., 2017). This process is associated with stable atmospheric conditions, where vertical mixing is almost
negligible (Zhou and Ferrier, 2008). Under this regime, the LWC will be distributed according to the cooling and condensation
rate at each height, and therefore it is likely possible to have situations where surface LWC is greater than LWC values above,
295 especially during radiation fog formation. This situation would lead to the observed negative α_{eq} values.

When fog LWP begins to surpass 30 surpasses the 30-40 g m^{-2} , its adiabaticity converges to 0.7, which, as stated
in the previous lines, is a value consistent with typical observations of boundary layer stratocumulus
(Slingo et al., 1982; Boers et al., 1990; Boers and Mitchell, 1994; Cermak and Bendix, 2011; Braun et al., 2018). This can be
explained because fog gradually becomes opaque to infrared radiation when its LWP surpasses 30 g m^{-2} (Wærsted et al., 2017).
300 In this scenario, LWC generation is mostly driven by radiative cooling at the fog top. This radiative cooling induces a temperature
gradient between the fog top and the surface, leading to convective motions. An increase in the intensity of convection will
be correlated with an increase in fog CTH, because the additional energy would enhance boundary layer development. Then,
as fog becomes deeper, it is expected that the relatively stronger convective motions associated would drive the vertical liquid
water mixing closer to what is observed in boundary layer clouds. Our results are consistent with this description. $\alpha_{eq}^{closure}$
305 increases in average with fog CTH until reaching a plateau at $\alpha \approx 0.7$, a value consistent with typical observations of boundary
layer stratocumulus (Slingo et al., 1982; Boers et al., 1990; Boers and Mitchell, 1994; Cermak and Bendix, 2011; Braun et al., 2018)
- This result and theory also indicate that dissipation by base lifting should happen when the LWP is at or above the 30-40 g m^{-2} range, when the layer is adiabatic and buoyant.

Finally, we can also observe that adiabaticity sometimes reaches values slightly greater than 1, which can be associated with
310 periods when fog is superadiabatic. This is possibly caused by an excess of liquid water with respect to the extent of the fog
column, which may be caused by the surface presence, as introduced in Sect. 2.

The data presented

4.2 Adiabaticity parametrization as a function of CTH

The strong correlation between adiabaticity and CTH observed in Fig. 3 (ba) suggests that α_{eq} can be parametrized as a function
315 of CTH. The parametrization curve is calculated by minimizing the error of the model presented in Eq. (7) with respect to the




Figure 3. (a) ~~Closure Equivalent~~ adiabaticity ~~results~~ versus fog CTH and LWP. The equivalent adiabaticity is calculated ~~by closure~~, using Eq. (6) and samples with visibility inferior to 2000 m. (b) Boxplot with the distribution of ~~closure~~ the equivalent adiabaticity for each radar bin, calculated by closure, for different LWP ranges. In both figures only samples with the derived parametrization superimposed (Eq. visibility below 2000 m are considered. (7)).

median α_{eq} value at each radar range bin (see Fig. 4). To reduce uncertainty due to lack of data, we only use range only bins with more than 20 valid samples are used.

$$\alpha_{eq}(CTH) = \alpha_0 \left(1 - e^{-\frac{CTH - H_0}{L}} \right) \quad (7)$$

The retrieved value for each coefficient are $\alpha_0 = 0.66$, $H_0 = 107.3$ m and $L = 50.2$ m. These parameters come from fog statistical behavior, and can be interpreted as follows: α_0 is the equivalent adiabaticity value that fog reaches when it has completely transitioned into an adiabatic regime. H_0 is the usual height at which LWC starts to increase with height. L indicates, based on adiabaticity, that the transition from stable to adiabatic fog is possible when CTH reaches 150 meters, and very likely when CTH is above 250 meters ($H_0 + L$ and $H_0 + 3L$ respectively).

In principle, the adiabaticity parametrization is valid for CTH values below 462.5 m, where the parametrization is derived. Beyond this height there is not enough data to guarantee its reliability; however it is likely that adiabaticity should remain close




Figure 4. Boxplot with the distribution of equivalent adiabaticity for each radar CTH bin, with the derived parametrization superimposed (Eq. (7)). Equivalent adiabaticity is calculated by closure using Eq. (6). Only samples with visibility below 2000 m are considered.

to the convergence value of 0.66 based on our observations and on what has been previously published in literature (Slingo et al., 1982; Boers et al., 1990; Boers and Mitchell, 1994; Cermak and Bendix, 2011; Braun et al., 2018).

4.3 Conceptual model validation

In this section we study fog statistical data to study how it behaves with respect to the conceptual model. Figure 5 (a) shows 330 all CTH, LWP and surface LWC measurements taken when fog is present (visibility less than 1000 m). Data is separated in different temperature ranges. Modeled LWP and CLWP curves are shown. LWP and CLWP theoretical curves are calculated using Eqs. (3c) and (4) respectively, with the $\alpha_{eq}(CTH)$ parametrization derived in Sec. 4.2. Each hexagon color is given by the mean LWC₀, calculated using all the data in their respective CTH+LWP space. Hexagons with less than 5 samples within their surface are removed, since they are likely to be associated with non replicable, noisy data.

335 This figure shows a good agreement between the theoretical curves and observed results. Most LWP samples are higher than the critical value, as the model predicts when visibility is less than 1000 meters. Additionally, it can be seen that for a fixed CTH, LWP increases with LWC₀. This behavior seems to be well captured in the current Conceptual Model formulation, as the difference between the three lines shows (each theoretical LWP line has a different LWC₀ value, indicated in the legend).

Figure 5 (b) shows data samples taken when visibility is between 1000 and 2000 meters, as an scatterplot. As in Sec. 4.2.1, 340 the 2000 m superior limit to visibility is selected, to remain close to fog conditions where the conceptual model is valid. LWP of these data samples should be less than the CLWP line for these visibility values, however we observe that sometimes they can also be larger. This can be explained by two main reasons: CLWP is calculated for a single temperature while data temperature varies within a range, and because of instrumental uncertainties. HATPRO LWP uncertainty is around 10 g m⁻², while radar CTH retrieval has a resolution of 12.5 m. This uncertainty is present in this retrieved data, and is also likely to be propagated 345 inside the $\alpha_{eq}(CTH)$ parametrization, introducing some variability in the results. However this is not deemed critical, since

variability around the CLWP line is smaller than 10 g m^{-2} , and because the fog life cycle studies of Sec. 5) verify that LWP is lower than the critical value before fog formation and after fog dissipation.

Finally, we perform an evaluation on how well the Conceptual Model predicts fog LWP, based on CTH, Temperature, Pressure and surface LWC inputs. These variables are used to calculate the Conceptual Model LWP with Eq. (3c), with the $\alpha_{eq}(CTH)$ parametrization of Sec. 4.2, and compared against HATPRO LWP retrievals. Results are shown in Fig. 6. Here we can see that most samples are close to the 1-1 line for LWP values less than approximately 190 g m^{-2} . Beyond this LWP value some deviation appears, however there is not enough data available to verify if this is a systematic error of the model or on how data was taken. Despite this deviation, the good agreement between modeled and observed LWP can be seen in the linear fit, with a slope equal to 1, and in the RMSE of just 10.5 g m^{-2} , which is very close to the LWP retrieval uncertainty.

355 4.4 Drivers of RLWP temporal variations

Equation (5), indicates that changes in both LWP and CTH can contribute to RLWP depletion, and therefore to fog dissipation. To quantify the relative impact of LWP and CTH changes in RLWP, we calculate the time derivative of Eq. (5). By assuming constant temperature and pressure, and using the $\alpha(CTH)$ parametrization of Sec. 4.2, we obtain Eq. (8).

This equation shows that RLWP changes are proportional to LWP variations, and to CTH variations weighted by the function $F(CTH, \Gamma_{ad}, \alpha_{eq})$. This function, written explicitly in Eqs. (9a) and (9b), converts CTH variations into g m^{-2} units, and thus enables a comparison between both effects.

$$\frac{dRLWP}{dt} = \frac{dLWP}{dt} - F(CTH, \Gamma_{ad}, \alpha_{eq}) \frac{dCTH}{dt} \quad (8)$$

$$F(CTH, \Gamma_{ad}, \alpha_{eq}) = \frac{1}{2} \frac{\partial \alpha_{eq}(CTH)}{\partial CTH} \Gamma_{ad}(T, P) CTH^2 + \alpha_{eq}(CTH) \Gamma_{ad}(T, P) CTH + LWC_c \quad (9a)$$

$$\frac{\partial \alpha_{eq}(CTH)}{\partial CTH} = \frac{\alpha_0}{L} e^{-\frac{CTH - H_0}{L}} \quad (9b)$$

365 Equation (8) implies that RLWP depletion, and thus fog dissipation, can occur by LWP reduction and/or by CTH growth. It also indicates that it is possible to have compensating effects enhancing fog persistence, for example fog that is reducing its LWP could persist if its CTH is also decreasing (which can happen under strong subsidence). Another implication is that it is possible to have fog dissipation even if LWP is increasing quickly, through a fast increase in CTH. The case studies of Sec. 5.1 show how useful this separation between LWP and CTH effects can be, by analyzing some examples of the 370 previously mentioned scenarios. Section 5.2.3 shows statistical results of fog RLWP, LWP and CTH time derivatives just before dissipation.




Figure 5. Observations of CTH, LWP and LWC at the surface for different temperature and visibility ranges. Data associated [with](#) visibility values below 1000 m is to the left (title (a)), while data measured with visibility values between 1000 and 2000 m is to the right (title (b)). Conceptual model theoretical LWP and CLWP lines for different conditions, indicated in the legend, are superimposed. The adiabaticity values used in the conceptual model calculation [come from](#) are calculated using the adiabaticity parametrization of Sec. 4.2.




Figure 6. 2D histogram comparing HATPRO and Conceptual Model LWP values, for data retrieved when visibility is less than 2000 m. Conceptual model LWP is calculated using fog CTH, fog LWC at the surface derived from visibility, surface temperature, surface pressure and the adiabaticity parametrization of Eq. (7). Under these conditions, the conceptual model predicts LWP with an RMSE of 10.5 g m^{-2} and an almost perfect linear relationship.

5 Fog life cycle

5.1 Case studies

We present ~~4~~³ case studies to illustrate the behavior and role of changes in LWP and CTH on presence of fog at the surface 375 during the fog life cycle (Figs. 7, 8, ?? and 9). For each case we provide a ~~6-panel~~^{5-panel} figure that illustrates the time series of fog/stratus layer boundaries, reflectivity profile, 4-m and 20-m horizontal visibilities, the fog/stratus layer measured LWP and computed RLWP, temperature and closure adiabaticity; and the change rate of RLWP, with the individual contributions from LWP and CTH variations.

In all ~~four~~^{three} cases, we observe that fog is present at the ground (4-m height visibility $< 1 \text{ km}$) when the RLWP is greater 380 than 0 g m^{-2} . RLWP changes at a rate of $+/- 10 \text{ g m}^{-2} \text{ Hr}^{-1}$, with values reaching $+/- 30 \text{ g m}^{-2} \text{ Hr}^{-1}$ at times. The LWP estimation of all case studies is done directly ~~with using~~ the HATPRO, ~~since as radar images show, there are no cloud signals verifying that the radar does not detect signals from liquid clouds below 6 km of height (panel (b) of each case study figure)~~.

Case study 1 (Fig. 7): Radiative fog occurring during fall season (31 October 2015) that forms six hours before sunrise and dissipates about three hours after sunrise at 10:25 UTC. The fog layer is about 200 m thick during the entire fog life cycle with 385 a water content of $30\text{-}60 \text{ g m}^{-2}$. This LWP range and the adiabaticity values close to 0.6 indicates that fog is optically thick and can be considered as a well-mixed layer for most of its duration. The RLWP is not large, mostly near $+ 10 \text{ g m}^{-2}$, with a maximum value of 30 g m^{-2} observed 2-3 hours before sunrise. CTH changes are relatively slow during the entire fog life

cycle, with values less than 50 m Hr^{-1} . From 03 to 05 UTC, the CTH increases which acts as RLWP depletion of nearly $-20 \text{ g m}^{-2} \text{ Hr}^{-1}$, while at the same time the LWP increases with a rate reaching $+50 \text{ g m}^{-2} \text{ Hr}^{-1}$ resulting in a net increase of RLWP. After 05 UTC, the trends in CTH and LWP reverse. The CTH subsides slowly (about -20 m Hr^{-1}) contributing positively on the RLWP at a rate of nearly $+5-10 \text{ g m}^{-2} \text{ Hr}^{-1}$, while the LWP initiates a progressive and nearly monotonous decrease of $-10 \text{ g m}^{-2} \text{ Hr}^{-1}$ that brings the RLWP to 0 g m^{-2} at 09 UTC. The progressive drying of the fog layer is also identifiable in the closure adiabaticity value, which starts to decrease just after sunrise. After 09 UTC, the near-surface visibility initiates a rapid increase, exceeding 1 km at 10:25 UTC, time at which the entire fog layer is dissipated. The complete layer dissipation and the increasing temperature makes it highly unlikely that fog will re-form in the coming hours. Note on Fig. 7 (f) that LWP and CTH contributions to RLWP are nearly always of opposite signs, but not equal in magnitude.

Case study 2 (Fig. 8): Another radiative fog that occurs in the fall season, just a few days apart from case study 1 (26 October 2015). It forms just three hours before sunrise and dissipates about 3.5 hours after sunrise at 10:55 UTC. The fog layer is about 200 m thick during the mature phase of the fog life cycle and nearly doubles between sunrise and time of dissipation, while the water content remains above 50 g m^{-2} . After fog formation, RLWP reaches 30 g m^{-2} in about one hour and remains at this level for about 2 hours. Fog adiabaticity indicates that after the first hour from formation fog remains in a well mixed state. Around sunrise, RLWP initiates a nearly monotonous decreasing trend of $-10 \text{ g m}^{-2} \text{ Hr}^{-1}$ that will last until fog dissipation. The negative RLWP rate is driven by the rise of CTH that contribute negatively on RLWP with a rate that exceeds $-20 \text{ g m}^{-2} \text{ Hr}^{-1}$ only partially compensated by $+20 \text{ g m}^{-2} \text{ Hr}^{-1}$ LWP increase rates. Oscillations in LWP and CTH contributions to RLWP are clearly visible in Fig. 8 (f). When there is strong cooling at the fog layer top, LWP ~~increases~~ and vertical circulation ~~is intensified. This increases~~ ~~increase. This in turn increases the~~ mixing with the layer above fog, resulting in a CTH increase. On the contrary, processes associated with CTH subsidence tend to decrease LWP rates (Wærsted, 2018). In this case study, the depletion of RLWP is clearly driven by the CTH increase and the fog LWP still exceeds 75 g m^{-2} at the time of dissipation.

Case study 3 (Fig. ??): ~~This third case occurred also in the fall season a few days apart from the two previous cases. It is characterized by a very late dissipation time 14:40 UTC, which is eight hours after sunrise. We will focus here on differences with the two previous cases that can explain the persisting character of this fog layer. During the fog life cycle of case 3, the CTH behaves similarly to that of case 1, reaching about 200 m agl and revealing both positive and negative evolutions with a rate of about $\pm 25 \text{ m Hr}^{-1}$. During the fog life cycle of case 3, the LWP behaves similarly to that of case 2, ranging between 50 and 100 g m $^{-2}$, which leads to a fog layer with a significant RLWP reaching $+50 \text{ g m}^{-2}$ during the first five hours of its life cycle. Adiabaticity remains always close to 0.6, indicating that the layer is well mixed for the complete fog duration. At 12:00 UTC, nearly six hours after sunrise, the RLWP is still greater than $+30 \text{ g m}^{-2}$, a clear sign that the fog is not about to dissipate, as confirmed by the 200-m near surface visibility. Between sunrise and noon, the rate of change of RLWP switches from positive to negative values (mostly within $\pm 10 \text{ g m}^{-2} \text{ Hr}^{-1}$). Contributions to RLWP changes from LWP and CTH changes are of opposite signs, with LWP being the dominant contributor, and their values cross the zero line nearly simultaneously eight times during the fog life cycle (Fig. ?? (f)). This is clear evidence of the feedback mechanisms that occur in fog between CTH changes and LWP, that tend to dampen the evolution of RLWP and lead to fog layers that persist for many hours even during daytime. The fog dissipation at 14:40 results primarily from a significant decrease of LWP after 12:00 UTC with a~~




Figure 7. Case study 1. (a) Cloud Base Height (CBH), Cloud Top Height (CTH), and the cloud radar 12.5 m resolution reflectivity profile for the first 1000 m of height. (b) **Cloud-radar 100-m resolution reflectivity profile up to 10 km of height.** (c) 4-m and 20-m horizontal visibilities. (d) Fog/Stratus layer measured LWP and computed RLWP. (e) Temperature and closure adiabaticity (calculated only when visibility is less than 2000 m). (f) Change rate of RLWP, with the individual contributions from LWP and CTH variations. In each panel, the time of fog formation and fog dissipation are clearly marked as well as the time of sunrise.




Figure 8. Case study 2. (a) Cloud Base Height (CBH), Cloud Top Height (CTH), and the cloud radar 12.5 m resolution reflectivity profile for the first 1000 m of height. (b) **Cloud-radar 100-m resolution reflectivity profile up to 10 km of height.** (c) Fog/Stratus layer measured LWP and computed RLWP. (d) Temperature and closure adiabaticity (calculated only when visibility is less than 2000 m). (e) Change rate of RLWP, with the individual contributions from LWP and CTH variations. In each panel, the time of fog formation and fog dissipation are clearly marked as well as the time of sunrise.

comparatively stable CTH. At the time of dissipation, when RLWP reaches 0 g m^{-2} , there is still about 25 g m^{-2} of LWP in the 200 m thick fog layer, which becomes insufficient to maintain the fog at the ground.

425 Case study 3. (a) Cloud Base Height (CBH), Cloud Top Height (CTH), and the cloud radar 12.5 m resolution reflectivity profile for the first 1000 m of height. (b) Cloud radar 100 m resolution reflectivity profile up to 10 km of height. (c) 4 m and 20-m horizontal visibilities. (d) Fog/Stratus layer measured LWP and computed RLWP. (e) Temperature and closure adiabaticity (calculated only when visibility is less than 2000 m). (f) Change rate of RLWP, with the individual contributions from LWP and CTH variations. In each panel, the time of fog formation and fog dissipation are clearly marked as well as the 430 time of sunrise.

Case study 4 (Fig. 9): Here we have a typical case of a very low stratus cloud layer with CTH near 250 m agl and an LWP that ranges $25\text{--}50 \text{ g m}^{-2}$. This combination leads to a negative RLWP that is insufficient for the stratus to deepen all the way to the surface. As expected for low stratus clouds, the value of closure adiabaticity is close to 0.6 for all valid samples (when visibility is less than 2000 m, to have valid conceptual model conditions with positive LWC at the surface). The stratus is 435 present from 18:00 UTC onwards during twelve hours with a near-surface visibility of about 2-3 km. From 18 until 23 UTC, RLWP is clearly negative changing frequently from negative to positive rates of change (about $+\text{--} 5 \text{ g m}^{-2} \text{ h}^{-1}$) as the contributions of LWP and CTH changes oscillate from positive to negative values (as also seen in Case 3). At 01 UTC, the stratus reaches a new equilibrium with an LWP hovering around 50 g m^{-2} , which brings the RLWP very close to 0 g m^{-2} . The fog CBH is then below 20 agl, as evidenced by the visibility values measured at 20 m agl (Fig. 9 (c)). Between 04:30 and 06:30 440 UTC, the RLWP becomes again negative and the stratus base lifts. A strong increase in LWP ($+40 \text{ g m}^{-2} \text{ h}^{-1}$) starting after 06:00 UTC leads to a positive RLWP after 06:30 UTC and the stratus layers deepens all the way to the surface. The trend in LWP reverses around 08 UTC ($-20 \text{ g m}^{-2} \text{ h}^{-1}$) while the CTH remains mostly constant hence reducing the RLWP towards 0 g m^{-2} before 10 UTC. This case study shows that the RLWP is also a good indicator of the possibility for a very low stratus layer to deepen into fog and then reversely for the fog to lift into a low stratus.

445 5.2 Fog life cycle statistics

Taking advantage of our large database, we study the behavior of fog RLWP and its time derivative $d\text{RLWP}/dt$ statistically, for three different periods: fog formation, mature stage and dissipation. The objective is to identify patterns that these fog variables follow at each stage. This could lead to the development of new indicators to enhance the capabilities of fog forecasting models.

Fog formation statistics are taken between 90 minutes before and 90 minutes after the time block were fog formation is 450 identified from visibility measurements (Sec. 3.2). Likewise, for the dissipation period the analyzed data is taken from 90 minutes before to 90 minutes after the dissipation time block. All remaining blocks between 90 minutes after fog formation, and 90 minutes before fog dissipation, are considered to be fog middle life data. Because of how the fog stages are defined, the cases included in this statistical analysis must have a duration of at least 3 hours. This is valid for 56 cases, which are used for statistical analysis in the following sections.




Figure 9. Case study 4-3. (a) Cloud Base Height (CBH), Cloud Top Height (CTH), and the cloud radar 12.5 m resolution reflectivity profile for the first 1000 m of height. (b) **Cloud-radar 100-m resolution reflectivity profile up to 10 km of height.** (c) 4-m and 20-m horizontal visibilities. (d) Fog/Stratus layer measured LWP and computed RLWP. (e) Temperature and closure adiabaticity (calculated only when visibility is less than 2000 m). (f) Change rate of RLWP, with the individual contributions from LWP and CTH variations. In each panel, the time of fog formation and fog dissipation are clearly marked as well as the time of sunrise.

455 The time derivative of the RLWP (and the sliding mean used in Fig. 10 (b.2)) is estimated by calculating the slope of a linear fit on RLWP data within ± 30 minutes of a given time block. The retrieved slope value is declared valid only if at least 75% of the RLWP samples used in its calculation are valid.

5.2.1 Fog formation

460 Figure 10 (a.1) shows the statistical behavior of RLWP between 90 minutes before and 90 minutes after for formation. It can be seen that at fog formation there is a transition from negative to positive RLWP values. The relatively lower amount of samples before -35 minutes from fog formation happen because there are less fog cases where the cloud has formed that early, or that have an identifiable CTH above 85 meters. Yet, we can see that RLWP cannot be significantly lower than -10 g m^{-2} if fog will form within 30 minutes.

465 Additionally, in Fig. 10 (a.2) we can see that $d\text{RLWP}/dt$ becomes positive about one hour before formation, and remains consistently positive for another hour after formation. This first hour after fog formation is when fog reservoir grows the most, reaching a change rate of 10 to $25 \text{ g m}^{-2} \text{ hr}^{-1}$, and it may be critical in establishing fog persistence for the coming hours. After this first hour, fog RLWP stabilizes around 10 to 20 g m^{-2} and the increase per hour is reduced until entering the mature stage.

470 All 56 fog cases lasting more than 3 hours are considered for the statistics. However, since radiation fog is formed from a shallow layer close to the surface, these cases usually do not provide valid data points because their CTH cannot be retrieved with the radar (it can only observe CTH values above 85 m). Therefore, most of the data points before and around formation time are contributed by stratus lowering fog events.

5.2.2 Fog mature stage

475 A histogram with RLWP values in the period defined as fog mature stage is shown in Fig. 10 (b.1). We can see that approximately 90% of the time fog has a positive RLWP value, with a median value of 20.1 g m^{-2} over all cases and reaching up to 60 g m^{-2} . Negative RLWP values in fog mature stage are explained by short-term temporary lifting of fog from the surface, most likely caused by RLWP oscillations.

480 Figure 10 (b.2) shows the statistics of $d\text{RLWP}/dt$ versus the sliding mean value of RLWP using the same data points involved in the slope calculation. This figure shows that RLWP and its time derivative are not correlated, and that most of the time $d\text{RLWP}/dt$ remains within $\pm 20 \text{ g m}^{-2} \text{ hr}^{-1}$. The very low median value of $d\text{RLWP}/dt = -0.2 \text{ g m}^{-2} \text{ hr}^{-1}$ shows that fog does not have a clear tendency of RLWP increase or decrease in the long term. Thus, during this stage of fog life cycle, RLWP remains positive most of the time, with variations driven by oscillations in the value of $d\text{RLWP}/dt$.

485 The statistics for this period defined as fog mature stage are derived using the 56 fog events lasting more than 3 hours. In the fog mature stage several radiation fog cases will be developed beyond 85 m of CTH, and therefore both stratus lowering and radiation fog cases contribute to the statistics.




Figure 10. The boxplots of panels (a.1) and (a.2) represent RLWP and $d(RLWP)/dt$ statistics for each time block 90 minutes before and after fog formation. Boxplot shows the 25th, 50th and 75th percentiles, and the maximum and minimum values. [The number of samples per bin is shown in Fig. S2 of the supplementary material](#). Panels (b.1) and (b.2) show RLWP and $d(RLWP)/dt$ statistics during fog middle life, between 90 minutes after fog formation and 90 minutes before fog dissipation, calculated using 4064 and 3952 samples respectively. [Only fog events longer than 3 hours are considered](#) [The ordinate axis of panel \(b.1\) is associated with the cumulative and normalized distributions](#).

5.2.3 Fog dissipation

In the latter stage of fog life cycle, shown in Fig. 11 (a.1), RLWP decreases consistently from positive values associated to with the middle of the life cycle until reaching negative values after fog dissipation. Additionally, there are almost no RLWP samples above 30 g m^{-2} observed in the last 30 minutes before dissipation. Hence, an RLWP value above 30 g m^{-2} may be interpreted as an indicator of fog persistence.

Figure 11 (a.2) shows that the monotonous decrease in RLWP begins about 60 minutes before fog dissipation, and can commonly reach values of about -10 to -30 g m^{-2} h^{-1} . These negative values in the time derivative continue after fog dissipation, and can be explained by further lifting or drying of the remaining low stratus cloud (Wærsted et al., 2019).

To observe To study what is the main driver of fog dissipation, we calculate Fig. 11 (b) shows the calculated $d\text{RLWP}/dt$, $d\text{LWP}/dt$ and $-F(\text{CTH}, \Gamma_{ad}, \alpha_{eq}) \cdot d\text{CTH}/dt$ trends, defined in Sec. 4.4, using the last 60 minutes of data before dissipation. Theoretically, dissipation can only happen when the RLWP decreases, which only happens when the sum of the LWP and CTH time derivative terms is negative (Eq. (8)). This matches the results of Fig. 11, which has most points in the quadrants leading to the aforementioned condition. The few points that show a RLWP increase before dissipation, to the right of the dashed line, are associated with uncertain retrievals due to low absolute RLWP values, or fast RLWP depletion in the few minutes just before dissipation (time trends are calculated using a one hour linear fit).

Overall, data shows that fog dissipation happens in fits. Additionally, observations confirm that fog dissipates under the same scenarios predicted by the Conceptual Model in Sect. 4.4. Fog dissipates Here the conceptual model predicts that fog could dissipate, even when the LWP increases when the layer thickening effect is increasing, if the RLWP reduction from layer thickening is larger (large strong CTH increase), and leads to a net reduction of RLWP. Alternatively, Conversely, fog can also dissipate when LWP decreases, even if the LWP decreases, even when the CTH subsides. Finally, some cases dissipate with the contribution of both effects, LWP decrease and layer thickening. Data also shows that some regimes are forbidden in the last 60 minutes of fog life cycle, specifically any case in which the addition of $d\text{LWP}/dt$ and $-F(\text{CTH}, \Gamma_{ad}, \alpha_{eq}) \cdot d\text{CTH}/dt$ is positive. This condition can be helpful for identifying conditions where fog is likely to persist or dissipate.

6 Conclusions

This work presents a Conceptual Model for adiabatic fog that relates fog liquid water path with its thickness, surface liquid water content and adiabaticity. The model predicts that LWP can be split into two contributions: the first is proportional to the adiabaticity and the square of CTH, and the second is the product of surface LWC and CTH. The later dependency is due to an excessive accumulation of water with respect to an equally thick cloud, which happens only appears in fog because the surface presence limits vertical development.

This excess accumulation of water motivates the definition of two diagnostic parameters, which later will prove to be key in understanding fog evolution: the Critical LWP and the Reservoir LWP. The Critical LWP (CLWP) is the minimum amount of column water that would fill the fog layer and cause a visibility reduction down to 1000 m at the surface. The Critical LWP can be calculated using the conceptual model, by imposing a surface LWC equivalent to a 1000 m visibility. Meanwhile, the




Figure 11. The boxplots of panels (a.1) and (a.2) show RLWP and dRLWP/dt statistics for each time block, 90 minutes before and after fog dissipation. These statistics are derived using 56 fog events, however there may be less than this amount of valid samples for each bin. The number of valid samples per bin is shown in Fig. S3 of the supplementary material. Panel (b) uses shows the last 60 minutes of data before dissipation to calculate the impact of LWP and CTH variations in RLWP depletion, using data from the last 60 minutes before dissipation. The dashed line indicates the theoretical limit where fog dissipation is possible (only to the left of this line). In quadrants II and III cloud base lifting contributes to RLWP decrease, while in Quadrants III and IV the LWP decrease contributes to RLWP depletion. This panel contains 40 valid samples from 56 fog cases, calculated using the method explained at the beginning of Sect. 5.2

Reservoir LWP (RLWP) is the difference between fog LWP and the Critical value, and represents the excess of water that
520 enables fog persistence. Case studies and statistical results show that the Reservoir LWP is positive when fog is present, and reaches 0 g m^{-2} at about the same time as fog dissipation.

The model is used to statistically study fog adiabaticity. Important conclusions are that thinner fog ~~has relatively lower adiabaticity values, less than the~~, ~~with a LWP less than 20 g m^{-2} , have adiabaticity values below 0.6~~~~value commonly found in literature for boundary layer clouds, even reaching~~, and ~~can even reach~~ negative values. This happens when the fog layer
525 is not yet opaque during the fog formation stage, when LWC distribution is not even and may be larger closer to the surface. ~~Meanwhile~~~~In this situation fog is not buoyant and therefore it may not lift when the RLWP reaches 0 g m^{-2} . Conversely,~~ when fog is developed, its adiabaticity value gets closer to previously observed values for boundary layer fog, converging at approximately 0.66 for fog ~~thicker than 250 meters. An adiabaticity with a LWP greater than $30-40 \text{ g m}^{-2}$. Here the fog layer is adiabatic, and therefore the fog base should lift when the RLWP depletes down to 0 g m^{-2} . Adiabaticity results are highly~~
530 ~~variable for LWP values between $20-30 \text{ g m}^{-2}$, and therefore it may be necessary to include additional observations to discern the adiabaticity of the fog layer in this LWP range.~~

~~Another result from the study of adiabaticity is an adiabaticity parametrization as a function of fog thickness~~~~is derived to~~
which can be used to estimate fog LWP and to perform conceptual model calculations.

~~Using this parametrization, the conceptual model enables an~~~~The~~ estimation of fog LWP ~~with has~~ an RMSE of 10.5 g m^{-2} , which is close to the uncertainty in LWP measurement of 10 g m^{-2} . ~~Additionally, data shows that the modeled LWP dependency, validating the modeled dependency of the LWP on surface LWC, temperature and CTH is well captured by the model formulation, pressure and CTH.~~

The temporal derivative of the ~~Reservoir LWP~~RLWP is studied, obtaining an analytic formulation that enables the ~~separation of quantification of the contribution of LWP and CTH contributions to reservoir depletion, and thus in causing variations to the depletion of the reservoir, and therefore leading to~~ fog dissipation. This formulation ~~predicts~~, which is validated by observations, indicates that fog dissipation will depend on the ~~relationship ratio~~ between LWP and CTH ~~changes~~ variations, and that ~~it is possible for fog to dissipate even if fog can dissipate by lifting as long as the net RLWP trend is negative, even if 1. LWP and CTH are both increasing, 2. LWP is decreasing and CTH increasing and 3. LWP is increasing or CTH is decreasing if the other term impact on reducing the reservoir is larger. This prediction is verified by statistically studying the LWP and CTH dissipation trends, where in some cases it was found that dissipation happened under antagonist LWP and CTH effects are both decreasing.~~

~~In addition, the database is used to statistically observe the Reservoir LWP time derivative, finding a significant increase of Reservoir LWP. Statistical observations of the fog life cycle indicate that the RLWP increases, in general, about 60 minutes before and after fog formation. This is followed by ~~oseillating Reservoir LWP variations of lower magnitude, sometimes created by larger but compensating LWP and CTH variations, sustaining a stable positive Reservoir LWP value~~ positive RLWP values, during fog middle life. ~~Fog life cycle ends when the Reservoir, that may oscilate or vary depending on the LWP and CTH evolution. Then, about 60 minutes before dissipation, the RLWP starts to decrease consistently, a trend that starts roughly 60 minutes before fog dissipation until reaching 0 g m^{-2} at dissipation time.~~~~

555 The aforementioned conclusions and the paper results indicate that the Reservoir LWP RLWP and its time derivative can be used as indicators of the fog life cycle stage and as a diagnostic tool to predict, at the local scale. This enables its potential use as an additional diagnostic variable, to quantify how close fog is from dissipation at the local scale. Reservoir LWP forecasting is also conceivable by eventually including fog processes linked to. This may complement visibility measurements at key sites affected by fog, such as airports and land roads, and help improving their logistics to reduce costs and the probability of accidents (Tardif and Rasmussen, 2007).

560 At present, the RLWP provides an estimation, in real time, of the excess of water of fog that enables the fog layer to remain at the surface. This can already be used as a diagnostic to estimate how likely fog persistence is for the coming minutes, based on the instant RLWP value and its trend (fog dissipation nowcasting). For example, results indicate that fog will not dissipate in the next 30 min if its RLWP is greater than 30 g m^{-2} . Additionally, the RLWP must have a decreasing trend before dissipation, and therefore a positive trend would indicate fog persistence. This result could be improved by introducing 565 forecasting tools to the conceptual model scheme. Forecasting when the RLWP will become 0 g m^{-2} would provide a proxy to predict fog dissipation by base lifting. This forecasting could be done, for example, by considering physical processes. They provide information on fog evolution, and could be used to estimate how the LWP and CTH variations in the calculations, and thus the RLWP, will evolve in the near future (e.g. Waersted et al. (2019)).

570 Regarding future work, it could be interesting to implement this framework on LES simulations, to improve our understanding of adiabaticity and of the surface effect in the excess accumulation of water that happens in fog with respect to equally thick clouds. Another interesting perspective would be to test conceptual model calculations using the output of fog large-eddy simulations (LES). If the conceptual model variables behave as theoretically expected in these simulations, they could be used to further study the impact of microphysics or surface properties on fog adiabaticity.

575 Other area of future work is to improve the exploitation of the measurements. For example, the conceptual model formulation could be tested with data obtained in interest would be to study the conceptual model at other sites with frequent fog events. When fog is adiabatic ($\text{LWP} > 30-40 \text{ g m}^{-2}$), the observed equivalent adiabaticity results is consistent with values observed at other sites. This hints that the conceptual model could be applicable at other sites with frequent fog events, to test how general is similar fog types (continental mid-latitude fogs), with possible variations in the adiabaticity parametrization - due to local conditions. This remains to be verified using real observations.

580 It would also be of interest to study how the direct retrieval of adiabaticity profiles from cloud radar reflectivity profiles could improve the quality of fog RLWP estimation. For example by enabling the retrieval of adiabaticity profiles instead of relying on a single average value. Other observations of interest include the quantification of local processes and larger area effects, such as advection, to enable the forecasting of RLWP. RLWP forecasting would greatly improve our real time assessment of fog dissipation tendency. However, these improvements would require a significant improvement on cloud radar 585 calibration standards. As is explained in the introduction, cloud microphysic retrievals are very sensitive to the radar calibration. Additionally, to enable the comparison of data between different sites, standarized calibration methodologies must be put in place. Thus, significant work is done to rise solution proposals for these issues. These proposals are presented in the following

~~two chapters of the thesis be used to improve the accuracy of the RLWP estimation, compared to the use of a single equivalent value.~~

590 *Data availability.* All data used in this study is hosted by the SIRTA observatory. Data access can be requested for free following the conditions indicated in the SIRTA data policy (https://sirta.ipsl.fr/data_policy.html).

SIRTA observatory website: <https://sirta.ipsl.fr/>

Data request form: https://sirta.ipsl.fr/data_form.html

Appendix A: Calculation of $\Gamma_{ad}(T, P)$

595 The inverse of the saturation mixing ratio change with height $\Gamma_{ad}(T, P)$ is calculated using the formulation published by Albrecht et al. (1990) and Braun et al. (2018), shown in Eq. (A1).

$$\Gamma_{ad}(T, P) = \left[\frac{(\epsilon + w_s) w_s l_v}{R_d T^2} \Gamma_w - \frac{g w_s P}{(P - e_s) R_d T} \right] \rho_d \quad (A1)$$

A description and the equations necessary to calculate each term used in the calculation of $\Gamma_{ad}(T, P)$ are given in Tab. A1.

Table A1. List of all the terms needed for the calculation of $\Gamma_{ad}(T, P)$.

Term	Definition	Calculation	Units
T	Surface temperature		K
P	Surface pressure		Pa
l_v	Latent heat of vaporization	$2.5 \cdot 10^6$	$\text{J Kg}^{-1} \text{ K}^{-1}$
c_p	Specific heat of dry air at constant pressure	1005	$\text{J Kg}^{-1} \text{ K}^{-1}$
g	Acceleration of gravity	9.81	m s^{-2}
R_d	Dry air ideal gas constant	287.0	$\text{J Kg}^{-1} \text{ K}^{-1}$
R_v	Water vapor ideal gas constant	461.5	$\text{J Kg}^{-1} \text{ K}^{-1}$
ϵ	Ratio of R_d to R_v	$\frac{R_d}{R_v}$	
e_s	Vapor saturation pressure	$611.2 \cdot \exp\left(\frac{17.67(T-273.15)}{T-29.65}\right)$	Pa
w_s	Saturation mixing ratio	$\epsilon \frac{e_s}{P-e_s}$	
ρ_d	Dry air density	$\frac{P-e_s}{R_d T}$	Kg m^{-3}
Γ_w	Moist adiabatic lapse rate	$\frac{g}{c_p} \left(1 + \frac{l_v w_s}{R_d T}\right) / \left(1 + \frac{\epsilon l_v^2 w_s}{R_d c_p T^2}\right)$	K m^{-1}
$\Gamma_{ad}(T, P)$		Eq. (A1)	Kg m^{-4}

Appendix B: Visibility-LWC parametrization

600 Surface LWC estimation from visibility measurements is done by inverting Gultepe et al. (2006) Eq. (6). This results in Eq. (B1), where LWC is Liquid Water Content in Kg m^{-3} and VIS is the visibility in meters.

$$LWC = 0.0187 \cdot 10^{-3} \cdot \left(\frac{VIS}{1000} \right)^{-1.041} \quad (B1)$$

Author contributions. FT and MH developed the conceptual model and its formulation, based on initial work by EW and MH. FT and EW developed the code used for data analysis. FT and MH defined the paper structure and content. MH and JCD manage the SIRTA observatory, 605 which provided the used dataset. All authors reviewed the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We acknowledge all the SIRTA observatory technical team for their extraordinary work on retrieving long term and high quality datasets. SIRTA measurements were performed in the framework of the ACTRIS, supported by the European Commission under the Horizon 2020 – Research and Innovation Framework Programme, H2020-INFRADEV-2019-2. We also acknowledge Marc-Antoine Drouin and Cristophe Boitel of the SIRTA observatory for their help on data access. We acknowledge the french Association Nationale Recherche Technologie (ANRT) and the company Meteomodem for their contribution in the funding of this work. [Finally, this publication is based upon work from COST Action PROBE, supported by COST \(European Cooperation in Science and Technology\).](#)

610

References

Albrecht, B. A., Fairall, C. W., Thomson, D. W., White, A. B., Snider, J. B., and Schubert, W. H.: Surface-based remote sensing of the observed and the Adiabatic liquid water content of stratocumulus clouds, *Geophysical Research Letters*, 17, 89–92, 615 <https://doi.org/10.1029/GL017i001p00089>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/GL017i001p00089>, 1990.

Bergot, T.: Small-scale structure of radiation fog: a large-eddy simulation study, *Quarterly Journal of the Royal Meteorological Society*, 139, 1099–1112, 616 2013.

Bergot, T.: Large-eddy simulation study of the dissipation of radiation fog, *Quarterly Journal of the Royal Meteorological Society*, 142, 1029–1040, 620 2016.

Betts, A. K.: Cloud Thermodynamic Models in Saturation Point Coordinates, *Journal of Atmospheric Sciences*, 39, 2182 – 2191, 621 [https://doi.org/10.1175/1520-0469\(1982\)039<2182:CTMISP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1982)039<2182:CTMISP>2.0.CO;2), https://journals.ametsoc.org/view/journals/atsc/39/10/1520-0469_1982_039_2182_ctmisp_2_0_co_2.xml, 1982.

Boers, R. and Mitchell, R. M.: Absorption feedback in stratocumulus clouds influence on cloud top albedo, *Tellus A*, 46, 229–241, 622 1994.

Boers, R., Melfi, S. H., and Palm, S. P.: Cold-Air Outbreak during GALE: Lidar Observations and Modeling of Boundary Layer Dynamics, 623 *Monthly Weather Review*, 119, 1132–1150, 1990.

Boutle, I., Price, J., Kudzotsa, I., Kokkola, H., and Romakkaniemi, S.: Aerosol–fog interaction and the transition to well-mixed radiation fog, *Atmospheric Chemistry and Physics*, 18, 7827–7840, 624 <https://doi.org/10.5194/acp-18-7827-2018>, <https://acp.copernicus.org/articles/18/7827/2018/>, 2018.

Braun, R. A., Dadashazar, H., MacDonald, A. B., Crosbie, E., Jonsson, H. H., Woods, R. K., Flagan, R. C., Seinfeld, J. H., and Sorooshian, A.: Cloud Adiabaticity and Its Relationship to Marine Stratocumulus Characteristics Over the Northeast Pacific Ocean, *Journal of Geophysical Research: Atmospheres*, 123, 13,790–13,806, 625 <https://doi.org/10.1029/2018JD029287>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD029287>, 2018.

Brenguier, J.-L., Pawlowska, H., Schüller, L., Preusker, R., Fischer, J., and Fouquart, Y.: Radiative properties of boundary layer clouds: 630 Droplet effective radius versus number concentration, *Journal of the atmospheric sciences*, 57, 803–821, 2000.

Brown, R. and Roach, W.: The physics of radiation fog: II—a numerical study, *Quarterly Journal of the Royal Meteorological Society*, 102, 335–354, 635 1976.

Cermak, J. and Bendix, J.: Detecting ground fog from space – a microphysics-based approach, *International Journal of Remote Sensing*, 32, 3345–3371, <https://doi.org/10.1080/01431161003747505>, <https://doi.org/10.1080/01431161003747505>, 2011.

Delanoë, J., Protat, A., Vinson, J.-P., Brett, W., Caudoux, C., Bertrand, F., Parent du Chatelet, J., Hallali, R., Barthes, L., Haeffelin, M., et al.: 640 Basta: a 95-GHz fmcw doppler radar for cloud and fog studies, *Journal of Atmospheric and Oceanic Technology*, 33, 1023–1038, 2016.

Driedonks, A. and Duynkerke, P.: Current problems in the stratocumulus-topped atmospheric boundary layer, *Boundary-Layer Meteorology*, 46, 275–303, 645 1989.

Dupont, J.-C., Haeffelin, M., Protat, A., Bouniol, D., Boyouk, N., and Morille, Y.: Stratus–fog formation and dissipation: a 6-day case study, *Boundary-layer meteorology*, 143, 207–225, 2012.

Gultepe, I., Müller, M. D., and Boybeyi, Z.: A New Visibility Parameterization for Warm-Fog Applications in Numerical Weather Prediction Models, *Journal of Applied Meteorology and Climatology*, 45, 1469–1480, <https://doi.org/10.1175/JAM2423.1>, <https://doi.org/10.1175/JAM2423.1>, 2006.

Gultepe, I., Tardif, R., Michaelides, S., Cermak, J., Bott, A., Bendix, J., Müller, M. D., Pagowski, M., Hansen, B., Ellrod, G., et al.: Fog research: A review of past achievements and future perspectives, *Pure and applied geophysics*, 164, 1121–1159, 2007.

Görsdorf, U., Knist, C., and Lochmann, M.: First results of the cloud radar and microwave radiometer comparison campaign at Lindenberg, aCTRIS Week 2020, 2020.

Haeffelin, M., Barthès, L., Bock, O., Boitel, C., Bony, S., Bouniol, D., Chepfer, H., Chiriaco, M., Cuesta, J., Delanoë, J., et al.: SIRTA, a ground-based atmospheric observatory for cloud and aerosol research, in: *Annales Geophysicae*, vol. 23, pp. 253–275, 2005.

Haeffelin, M., Bergot, T., Elias, T., Tardif, R., Carrer, D., Chazette, P., Colomb, M., Drobinski, P., Dupont, E., Dupont, J.-C., et al.: PARIS-FOG: Shedding new light on fog physical processes, *Bulletin of the American Meteorological Society*, 91, 767–783, 2010.

Haeffelin, M., Laffineur, Q., Bravo-Aranda, J.-A., Drouin, M.-A., Casquero-Vera, J.-A., Dupont, J.-C., and De Backer, H.: Radiation fog formation alerts using attenuated backscatter power from automatic lidars and ceilometers, *Atmospheric Measurement Techniques*, 9, 5347, 2016.

Hoffmann, H.-E. and Roth, R.: Cloudphysical parameters in dependence on height above cloud base in different clouds, *Meteorology and Atmospheric Physics*, 41, 247–254, 1989.

Kotthaus, S., O'Connor, E., Münkel, C., Charlton-Perez, C., Haeffelin, M., Gabey, A. M., and Grimmond, C. S. B.: Recommendations for processing atmospheric attenuated backscatter profiles from Vaisala CL31 ceilometers, *Atmospheric Measurement Techniques*, 9, 3769–3791, <https://doi.org/10.5194/amt-9-3769-2016>, <https://amt.copernicus.org/articles/9/3769/2016/>, 2016.

Manton, M.: The physics of clouds in the atmosphere, *Reports on Progress in Physics*, 46, 1393, 1983.

Marke, T., Ebelt, K., Löhnert, U., and Turner, D. D.: Statistical retrieval of thin liquid cloud microphysical properties using ground-based infrared and microwave observations, *Journal of Geophysical Research: Atmospheres*, 121, 14,558–14,573, <https://doi.org/10.1002/2016JD025667>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016JD025667>, 2016.

Nakanishi, M.: Large-eddy simulation of radiation fog, *Boundary-layer meteorology*, 94, 461–493, 2000.

Oliver, D., Lewellen, W., and Williamson, G.: The interaction between turbulent and radiative transport in the development of fog and low-level stratus, *Journal of the Atmospheric Sciences*, 35, 301–316, 1978.

Porson, A., Price, J., Lock, A., and Clark, P.: Radiation fog. Part II: Large-eddy simulations in very stable conditions, *Boundary-layer meteorology*, 139, 193–224, 2011.

Price, J.: Radiation fog. Part I: observations of stability and drop size distributions, *Boundary-layer meteorology*, 139, 167–191, 2011.

Roach, W., Brown, R., Caughey, S., Crease, B., and Slingo, A.: A field study of nocturnal stratocumulus: I. Mean structure and budgets, *Quarterly Journal of the Royal Meteorological Society*, 108, 103–123, 1982.

Roach, W. T., Brown, R., Caughey, S. J., Garland, J. A., and Readings, C. J.: The physics of radiation fog: I – a field study, *Quarterly Journal of the Royal Meteorological Society*, 102, 313–333, <https://doi.org/10.1002/qj.49710243204>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49710243204>, 1976.

Slingo, A., Brown, R., and Wrench, C.: A field study of nocturnal stratocumulus; III. High resolution radiative and microphysical observations, *Quarterly Journal of the Royal Meteorological Society*, 108, 145–165, 1982.

Smith, D. K. E., Renfrew, I. A., Price, J. D., and Dorling, S. R.: Numerical modelling of the evolution of the boundary layer during a radiation fog event, *Weather*, 73, 310–316, [https://doi.org/https://doi.org/10.1002/wea.3305](https://doi.org/10.1002/wea.3305), <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/wea.3305>, 2018.

Tardif, R. and Rasmussen, R. M.: Event-based climatology and typology of fog in the New York City region, *Journal of applied meteorology and climatology*, 46, 1141–1168, 2007.

Wærsted, E.: Description of physical processes driving the life cycle of radiation fog and fog–stratus transitions based on conceptual models, Ph.D. thesis, Paris Saclay, 2018.

Wærsted, E. G., Haeffelin, M., Dupont, J.-C., Delanoë, J., and Dubuisson, P.: Radiation in fog: quantification of the impact on fog liquid water based on ground-based remote sensing, *Atmospheric Chemistry and Physics*, 17, 10 811–10 835, <https://doi.org/10.5194/acp-17-10811-2017>, <https://www.atmos-chem-phys.net/17/10811/2017/>, 2017.

Wærsted, E. G., Haeffelin, M., Steeneveld, G.-J., and Dupont, J.-C.: Understanding the dissipation of continental fog by analysing the LWP budget using idealized LES and in situ observations, *Quarterly Journal of the Royal Meteorological Society*, 145, 784–804, <https://doi.org/10.1002/qj.3465>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3465>, 2019.

695 Walker, M.: The science of weather: Radiation fog and steam fog, *Weather*, 58, 196–197, 2003.

Zhou, B. and Ferrier, B. S.: Asymptotic analysis of equilibrium in radiation fog, *Journal of Applied Meteorology and Climatology*, 47, 1704–1722, 2008.