

Authors' Responses to Reviewers' Comments

Journal: Atmospheric Chemistry and Physics

Manuscript ID: acp-2020-142

5 **Manuscript Title:** Polycyclic aromatic hydrocarbons (PAHs), oxy- and nitro-PAHs in ambient air of Arctic town Longyearbyen, Svalbard

Authors: Tatiana Drotikova, Aasim M. Ali, Anne Karine Halse, Helena C. Reinardy, Roland Kallenborn

We thank the two referees for their constructive comments, which have led to considerable improvement of the manuscript.

10 After discussion within the author team, we have comprehensively addressed all recommendations provided. As it was suggested, multivariate statistical analysis was repeated with the PAH diagnostic ratios as variables. To focus on the initial goal of the section 4, discussions explaining the samples grouping and weather influences were omitted from the PCA outcomes. Therefore, we now introduce PCA loading plots instead of the biplots. General indications of the weather influence are now included in the "Gas/particle partitioning" section 3.2.2. We recently obtained new official information on
15 the settlement's statistics of vehicles, fuel type, coal transportation, and marine port traffic, and this information is now included in the manuscript. Consequently, the "Source identification" section 4 was reorganized and rewritten in response to the comments from Reviewer 2, and we are convinced that now it provides better insights on the potential local sources of PAHs, nitro- and oxy-PAHs in Longyearbyen.

Below we provide our point-by-point responses (in blue color font) to each referee comment (in black color font; note "old" line numbers). Revised text is in red color font. Please, note new numeration of the revised text lines, as well as tables and figures in the supplement information to the manuscript. We hope that these details will clarify the current content and structure of the manuscript.

Anonymous Referee #1

25 Reviewer's general comment:

I enjoyed reading this paper; it is well written and scientifically sound. Furthermore, it makes an important contribution to the Arctic literature, providing results and interpretation of possible sources PAHs, oxy- and nitro-PAHs using ambient samples. I recommend the paper be accepted for publication with minor revisions.

Author's response:

30 We appreciate the reviewer's careful consideration of our manuscript and constructive comments. We have addressed the reviewer's specific comments below.

Reviewer's comment:

35 Line 33 insinuates that the highest PAH levels observed in the Arctic are in the winter, and many of these would settle in the particulate phase because of the lower temperatures in lines 55 it is suggested that the oxygenated and nitro PAHs would be higher in the summers due to the increased presence of oxidants and photodegradation. Wondering if there are actual environmental measurement studies that support this, or if the authors have considered measuring ambient levels in this region in the summer.

Author's response:

40 To the best of our knowledge, the data on oxy- and nitro-PAHs atmospheric formation in the high Arctic does not exist. Unfortunately, we can not provide any references on this topic. In currently planned follow-up studies we will provide the year around obtained Svalbard data for PAHs, nitro- and oxy-PAHs, and we consider to include the atmospheric transformation of PAHs topic too.

45 Reviewer's comment:

Check phenanthrene spelling in introduction and wherever else relevant.

Author's response:

It was corrected twice in the abstract. Thank you.

50 Reviewer's comment on Figure 4:

I think it's interesting that the PAHs that seem to exist mostly in the gaseous phase (e.g. PHE, FLU, Ant, 2NNAp) tend to associate more with the lower temperature dry samples (A4 to A7). Curious as to whether the authors can clarify why.


Author's response:

55 For LMW PAHs, water solubility (for Ace, Acy, Flu, Phe, 1- and 2-NNAp, 9-Flu, and 9,10-AntQ) and polarity (for LMW nitro- and oxy-PAHs) play an additional role in wet scavenging processes (Shahpoury et al., 2018). The gas phase removal from the atmosphere is due to substance dissolution in water droplets (Shahpoury et al., 2018), which enhances the scavenging effect at higher humidity. The higher humidity conditions on the warmer days resulted in higher levels of LMW PAHs on the dry colder days. Although, as confirmed by earlier studies, snow scavenging may also be an important, and sometimes a dominating scavenging process for LMW PAHs (Wania et al., 1999). As can be seen on the plot provided
60 below, concentrations of the LMW PAHs were reduced due to snowing event on the day 5. These observations show that this topic is quite complicated and requires long-term study investigation of LMW PAHs behavior in the high Arctic atmosphere. Wider consideration of the issue is outwith the scope of the study.

Another possible reason could be higher emissions from sources (cars, power plant) in colder period. Temporal variation in wind speed and wind direction might be of importance too but we did not reveal it in this study.

- 65
- Shahpoury, P., Kitanovski, Z., Lammel, G. J. A. C., and Physics: Snow scavenging and phase partitioning of nitrated and oxygenated aromatic hydrocarbons in polluted and remote environments in central Europe and the European Arctic, *Atmos. Chem. Phys.*, 18, 13495-13510, <https://doi.org/10.5194/acp-18-13495-2018>, 2018.
 - Wania, F., Mackay, D., and Hoff, J. T.: The importance of snow scavenging of polychlorinated biphenyl and polycyclic aromatic hydrocarbon vapors, *Environ. Sci. Technol.*, 33, 195-197, <https://doi.org/10.1021/es980806n>,
70 1999.

Individual PAH values for Adventdalen air samples



Reviewer's comment on Adventdalen PCA:

I can't seem to understand what the co-authors are trying to convey here. Why would A1 and A2 being negatively correlated indicate specific humidity is an essential parameter for the removal from the atmosphere? I thought it was interesting that A2 and A5 (the highest precipitation events) were quite separated on the PCA plot. Initially, I thought the differences could allude to the differences in scavenging efficiencies of snow and rain for PACs (i.e. where PAC concentrations in A5 lower than in A2?) But, then at the UNIS station you don't see the same separation for corresponding samples.

Author's response:

Intensive precipitation occurred on day 2 (rain, 4.2 mm) and day 5 (snow, 2.8 mm) compared to the other days with little (A1 and A7, 0.1-0.2 mm) or no precipitation (A3 and A4). Hence, we expected that the samples A2 and A5, and perhaps A1 and A7 too, would be clustered all together due to scavenging effect and expected lower concentrations of PAHs and the derivatives on those days. Surprisingly, samples A2 and A5 were found to be separated and two groups of samples were observed. Within each group, A2 and A5 were clustered with other samples collected on the days with little or no precipitation. Therefore we concluded that the amount of precipitation influencing the PC1 to a lesser extent.

If considering the difference in scavenging efficiencies of snow and rain for PAHs as defining criteria, then the samples taken on the days without precipitation (A3 and A4) would expected to be separated from the others, and two (days with precipitation and days without precipitation) or three (rain, snow, no precipitation) groups would be observed. This PCA outcome was not seen.




Figure 1. PCA for Adventdalen samples

It was easy to notice that the samples taken on the cold days, later in September, were clustered together and separated from the warmer days. Arctic air is dry and cold, and ambient temperatures have strong influence on the actual amount of water vapor in the air (specific humidity). As consequence of similar ambient temperatures, specific humidity was found similar within each group (4.1 ± 0.1 for the warm group and 2.8 ± 0.7 for the cold group). Furthermore, all the studied compounds (except 2+3-NFlt, 2-NFlu and 9-NFlt) showed statistically significant ($p < 0.05$) negative correlation with specific humidity (Table S17). In addition, the detected PAH concentrations were significantly lower on the rainy day 2 ($\sum 46$ PAHs 430.9 pg m^{-3}) compared to the day 5 with snowing event ($\sum 46$ PAHs 635.3 pg m^{-3}). The gas phase removal from the atmosphere is due to substance dissolution in water droplets, which enhances the scavenging effect at higher humidity. Hence, we suggest here that the actual amount of water vapor in the air might be an essential parameter for removal from the atmosphere, and attributed it to PC1.

However, this conclusion is only to be considered indicative based on the here preformed pilot study (with few observations only). Hence, further investigations are needed to confirm this assumption and add more statistical power to our hypothesis. Please note, ours study did not aim to focus on weather influence. We therefore have now removed all discussions explaining the samples grouping to focus on the potential sources identification. Thought brief indications of the weather influence are now included in the gas/particle partitioning section 3.2.2:

Page 7, line 281: The influence of wet deposition was indicated by a significant negative correlation between amount of precipitation and concentrations of several particle-bound HMW PAHs (Chry, BbkFlt, IPyr, BPer, BaFlu-11, and BaAnt-7,12) as well as semi-volatile Phe, Flt, and Pyr, which are more predominant in gaseous phase (Spearman correlation, $p < 0.05$, Table S16. Effective wet scavenging of Phe, Flt, and Pyr has been suggested (Škrdlíková et al., 2011). Furthermore, a strong negative correlation with mass of water vapor in the air (specific humidity) was determined for most of the compounds (Spearman correlation, $p < 0.05$, Table S17). Particle associated HMW compounds are readily scavenged by precipitation, while water solubility and polarity (for nitro- and oxy-PAHs) play an additional role in wet scavenging processes (Shahpoury et al., 2018). The gas phase removal from the atmosphere is due to substance dissolution in water droplets, which enhances the scavenging effect at higher humidity. Higher sensitivity of gas scavenging compared with particle scavenging towards liquid water content was also indicated by Škrdlíková et al. (2011).

Reviewer's comment on A2 and A5 samples separation:

But, then at the UNIS station you don't see the same separation for corresponding samples.

Author's response:

We did not expect to see similar PCA outcome for these two locations. The UNIS sampling station is located closer to the town and thus more influenced by emissions from passenger cars traffic, PP, marine port etc. Changes in the rate of emission

from the sources due to human-influenced diurnal cycle may cause large variability in the PAH levels. Such disturbances are eliminated in Adventdalen, which is rural (an approximately background) sampling location with significantly less number of potential sources, and thus well suitable for such weather influences observations.

Reviewer's comment on the Figures 4-6:

Could the captions of Figures 4-6 include what A1-A7 and U1-U7 represent? i.e. A1- A3 = humid, etc. It is stated in the text, but its easier for the readers to follow if these are added under the figures.

Author's response:

As we explained, the discussion about weather influence was withdrawn from PCA. Now we introduced loading plots instead of the biplots. Please, see the updated version of the manuscript attached as separated file and the revised text is marked in red color font.

Reviewer's comment on the Table S12:

It would be interesting to include a table like Table S12 but a more comprehensive summary of PAHs levels in other Arctic regions. Because I found myself wondering how air concentrations were comparable to Arctic regions without a power-plant, for example.

Author's response:

Based on the comments from both the referees, we considered available studies on the Arctic as well as rural locations worldwide. The dedicated table with appropriate references is added for comparison. Please, see Table S14 in the updated version of the supplementary information to the manuscript. We have added the following text in the Section 3.2.1:

Page 6, line 241: The measured PAH concentrations in the present study were in agreement with the 2 decades average data reported for the Arctic monitoring stations in Svalbard (Zeppelin) and Finland (Pallas), and were about an order of magnitude higher compared to the Canadian Arctic (Alert) concentrations (Yu et al., 2019). The PAH levels observed in Longyearbyen were significantly (up to 2 orders of magnitude) lower compared to rural sites in Europe and China (Table S14).

Reviewer's comment:

This manuscript provides a study of PAHs and PAH derivatives (nitro- and oxy-PAHs) in the ambient air of two locations in Svalbard. The study includes results about their atmospheric concentrations, gas/particle partitioning and a tentative of source identification based on measurements in ambient air and at the emission of a coal power plant identified as a probable major contributor in the area of the study. Overall, the manuscript is well written, and the results are scientifically relevant especially in the context of the Arctic region which is still poorly documented about such toxic compounds. However, my major concerns are about the source identification methodology and results together with some statements about the gas/particle partitioning processes of these compounds. Thus, I would recommend the publication of this paper in ACP after some major revisions detailed directly into the pdf file of the text. Please also note the supplement to this comment: <https://www.atmos-chem-phys-discuss.net/acp-2020-142/acp-2020-142-RC2-supplement.pdf>.

Author's response:

We would like to thank the reviewer for his insightful comments towards improving our manuscript. We appreciate all the constructive suggestions and we have carefully addressed it, as explained below.

Lines 41-42: PAHs are regulated in many countries, eg. US, Canada, Holland, Sweden, Switzerland, and Denmark (Bandowe and Meusel, 2017).

Reviewer's comment:

Let's say USA, Europe and Canada. Specify in ambient air. Please cite the European Directive 2004/107/CE instead and for USA, USA EPA.

- *European Official Journal: Directive 2004/107/CE of the European Parliament and of the Council of 15 December 2004 relating to arsenic, cadmium, mercury, nickel and polycyclic aromatic hydrocarbons in ambient air (26/01/2005)., Official Journal, L23, 3–16, 2005.*
- *US EPA, 2011. Polycyclic Aromatic Hydrocarbons on the Gulf Coastline <http://www.epa.gov/bpspill/pahs.html>.*

Author's response:

We appreciate the comment and added information about PAH regulation guidelines. Text updated for ambient air regulation: Page 1, line 40: **Atmospheric PAHs are regulated in USA, Canada, UK, and Europe (EU Directive 2004/107/EC, 2005; US EPA, 2011; UK Air DEFRA, 2007; Ontario Ministry of the Environment and Climate Change, 2016).**

Line 46: PAHs are byproducts of different incomplete combustion processes, mainly fossil fuels and biomass burning.

Reviewer's comment:

- *Ravindra, K., Sokhi, R. and Van Grieken, R.: Atmospheric polycyclic aromatic hydrocarbons: Source attribution, emission factors and regulation, Atmospheric Environment, 42, 2895–2921, doi: DOI:10.1016/j.atmosenv.2007.12.010, 2008.*

Author's response:

The reference was added. Thank you.

Line 56: PAHs react with a number of atmospheric oxidants, most notably the hydroxyl radical, ozone, the nitrate radical, and nitrogen dioxide (Keyte et al., 2013). This leads to their transformation into more toxic oxygenated and nitrated PAH derivatives (oxy-PAHs and nitro-PAHs). Oxy- and nitro-PAHs are also constituents of raw coal and can be emitted with PAHs following the same combustion processes (Huang et al., 2014b).

Reviewer's comment:

More generally from combustion processes including fossil fuel combustion, biomass burning, etc.

Author's response:

195 We mentioned in the text: "PAHs are byproducts of different incomplete combustion processes, mainly fossil fuels and biomass burning" and "Oxy- and nitro-PAHs are also constituents of raw coal and can be emitted with PAHs following the same combustion processes". Thus the text remains without changes. These sentences are sufficiently explaining the potential oxy- and nitro-PAH sources from combustion processes. Thus, the text remained unchanged.

200 Line 61: the level of nitro- and oxy-PAHs in the Arctic atmosphere is unknown (Balmer and Muir, 2017).

Reviewer's comment:

is poorly documented

Author's response:

205 Based upon the reviewer's remarks, we have repeated the literature search regarding studies on atmospheric levels of nitro- and oxy-PAHs. Our updated literature survey confirmed that the statement of Balmer and Muir (2017) remains valid. Thus, the text was not changed.

Reviewer's comment, lines 99-103:

210 It would be useful for the reader to specify what are the typical emissions of the regulatory pollutants from this power plant (PM, SO₂, NO₂, etc.) especially with such treatment systems. It should refer to Table S12 for PM. This show the large efficiency of the flue gas treatments applied in this PP.

Author's response:

Additional information is provided in the main text. Text included in the manuscript line 103-104:

215 Page 3, line 103: **Low emissions are reported: dust 1.5±0.2, SO₂ 0.3±0.1, NO_x 244±19, CO 63±5 mg Nm⁻³ (Lundgjerdingen, 2017), reflecting high efficiency of the flue gas cleaning system.**

Reviewer's comment, line 112:

WFGD scrubber

Author's response:

220 Corrected accordingly

Reviewer's comment, line 115:

Tissuquartz? (no binding)

Author's response:

225 Yes, correct. Pallflex QFF, pure quartz, no binder, product ID 7202

<https://shop.pall.com/us/en/products/zid7202?CategoryName=ID31&CatalogID=Laboratory>. The updated text states:

Page 3, line 116: The particulate phase was collected on quartz fiber filter (QFF, pre-burned at 450°C for 6 h; Ø = 47 mm, **no binder**, Pallflex, USA), and the gaseous phase on polyurethane foam (PUF, Soxhlet pre-cleaned in toluene for 24 h followed by 24 h acetone wash; Ø = 50 mm, L = 75 mm, Klaus Ziemer GmbH, Germany).

230

Reviewer's comment, lines 116-117:

Using Soxhlet, PFE? Please provide some details

Author's response:

235 PUFs were Soxhlet pre-cleaned (VDI-cleaned, method BGI 505.47) by the producer, Klaus Ziemer GmbH, Germany. The cleaning method specification is now included.

Reviewer's comment, line 118:

It is maybe not of great importance here. It should maybe discuss here. What is a typical PM size distribution from coal combustion especially after all the cleaning steps of the PP?

240 Author's response:

Mass-size distribution was not measured neither by the accredited company (Lundgjerdingen, 2017) working for PP nor by us. PM emission and PM size distribution depends on temperature of the ESP inlet flue gas (Li et al., 2018), voltage and current of ESP fields, number of electric fields, as well as retention time of ESP (Liu, 2018). Thus, the size distribution may vary largely, as well as PM emission itself shown in the Table S12. Removal efficiency is higher for larger particles, while finer particles are more difficult to eliminate (Li et al., 2018; Liu, 2018). According to the study by Li et al. (2018), coarse particles proportion in flue gas after ESP was reduced from 74 to 24%, while proportion of fine and ultra-fine (<1 µm) particles together increased from 10 to 38%, though medium size particles (2.5-10 µm) were the most abundant (about 40%) in the particulate emission. Besides, removal efficiency is higher by a combination of ESP and WFGD (up to 99.9% as stated in the manuscript lines 181-182) and the PM size distribution may be changed. Thus, we do not see how this literature based information (not measured by us) about PM size distribution can be incorporated into the PP sampling description section 2.2.1, as well as in the results discussion. We believe that the provided data about ultra-low particles emission ($1.5 \pm 0.2 \text{ mg m}^{-3}$; line 184) is sufficient for the recent study scope and, considering the provided PM emissions from other coal-fired power plants in Table S12.

255 Lines 122-123: The prevailing wind direction in Longyearbyen and Adventdalen is from the southeast. In summer, when the soil surface in Adventdalen becomes warmer than water surface in Adventfjörd, the wind direction can temporary change to northwesterly.

Reviewer's comment:

Any reference or data to support this statement?

260 Author's response:

Appropriate references added

- Dekhtyareva, A., Edvardsen, K., Holmén, K., Hermansen, O., and Hansson, H. C.: Influence of local and regional air pollution on atmospheric measurements in Ny-Ålesund, *International Journal of Sustainable Development and Planning*, 11, 578-587, 10.2495/sdp-v11-n4-578-587, 2016.
- Esau, I., Argentini, S., Przybylak, R., Repina, I., and Sjöblom, A.: Svalbard meteorology, *Advances in Meteorology*, 2012, 8-164, 2012.

Reviewer's comment, line 124:

the peak of marine

270 Author's response:

Corrected accordingly

Line 129: For each station, 6 high volume air samples were collected for particulate (QFF, pre-burned at 450°C for 6 h; Ø = 103 mm, no binder, Munktel/Ahlstrom, Finland)

275 Reviewer's comment:

Why not Pallflex like for the PP? No binder?

Author's response:

The providing company informed us that the Ø = 103 mm size Pallflex QFF for large volume air sampling would be out of stock for several month. Therefore, we simply chosen the similar type, no binder QFF available from a different company.

The quality of the filters was similar: "MK 360 conditioned by high temperature pre-heating; Made of extremely pure micro-quartz fibers; Binder free; Particularly suitable for low amount of particles". The additional information "no binder" is now included.

Lines 129-130: For each station, 6 high volume air samples were collected for particulate (QFF, pre-burned at 450°C for 6 h; Ø = 103 mm, Munktell/Ahlstrom, Finland) and gaseous (PUF, pre-cleaned in toluene for 24 h followed by 24 h acetone wash; Ø = 65 mm, L = 100 mm, Klaus Ziemer GmbH, Germany) phases.

Reviewer's comment:

Using Soxhlet, PFE? Please provide some details. Why not the same brand as for the PP?

Author's response:

The PUFs used for PP chimney and ambient air sampling are from the same producer (Klaus Ziemer GmbH, Germany) but different size because different pumps were used (portable low volume Digitel and High volume TISCH). The additional information on the cleaning method, Soxhlet, is now included in the text.

Reviewer's comment, line 135:

analysis

Author's response:

Corrected accordingly

Reviewer's comment, line 135:

(number of PP samples and blanks) to be specified in the previous section

Author's response:

To avoid repetition in Sections 2.2.1 and 2.2.2, extra details were included for clarification. Updated text:

Page 4, line 134: All samples (PP, UNIS, and Adventdalen) were kept intact inside the sampling unit after collection. In order to reduce the risk of post-collection contamination, the unit was sealed in two plastic bags for transportation to the lab, where samples were removed from the unit, sealed with layers of aluminum foil, and stored airtight in two plastic bags. Samples were kept frozen at -20°C until analysis. A total of 18 samples (18 QFFs and 18 PUFs) and 8 field blanks (4 for PP and 4 for UNIS and Adventdalen) were collected.

Reviewer's comment, line 138:

GC-EI-MS/MS and GC-ECNI-MS, give the names

Author's response:

The clarifying information was included. The updated text states:

Page 4, line 140: 16 PAHs, 8 oxy- and 21 nitro-PAHs (Table S3) were quantified using gas chromatography in combination with electron ionization mass spectrometry GC-EI-MS/MS and gas chromatography in combination with electron capture negative ion mass spectrometry GC-ECNI-MS, respectively.

Reviewer's comment, line 143:

QuEChERS-like extraction, full name?

Author's response:

320 The updated text includes this information:

Page 4, line 145: QFF samples were extracted with dichloromethane using a quick easy cheap effective rugged and safe QuEChERS-like procedure developed previously for the analysis of particulate bound PAHs (Albinet et al., 2013; Albinet et al., 2014).

325 Line 144: PUF samples were Soxhlet extracted with dichloromethane for 24 h.

Reviewer's comment:

Soxhlet extraction, manual or automatic Soxhlet?

Author's response:

All samples were manually Soxhlet extracted. The text remains the same.

330

Line 155-156: The method efficiency was tested using QFF (n=4) and PUF (n=4) spiked samples (Table S7). Acceptable recoveries ranged between 63-109% for dPAHs, 56-68% for dOxy-PAHs, and 44-89% for dNitro-PAHs (Table S8).

Reviewer's comment:

This is a first step but it would have better to evaluate the extraction efficiency by analyzing a standard reference material especially for the particulate phase

335

Author's response:

We agree, it might have been beneficial and will be the subject for follow-up studies. However, we do not intend to use our analytical approach for accredited reporting and thus consider the method as completely validated.

340 Reviewer's comment, line 175:

Actually, on Table 1, it would have been better to provide the emission factors for the power plant for all the individual species (PAHs and derivatives) instead of the concentration in the flue gas.

It would be also interesting to compare these results to results from the bibliography event for coal power plants with different flue gas treatment systems as done based on the concentrations below (Table S12)

345

Author's response:

The goal of analyzing the PP stack emission samples was only to obtain PAH chemical profile in order to determine the markers to be used for source identification. An estimation of the PP cleaning system efficiency would be a different study with different requirements (isokinetic sampling) and extra equipment for simultaneous measurements of other emissions like PM, NO_x, SO_x and perhaps others. We therefore decided to omit this interesting aspect for our studies for avoiding overloading the manuscript.

350

For the determination of emission factors, as suggested, accurate data on the PP coal load during sampling is needed. Unfortunately, this internal information is not available to us. We can only operate with annual coal consumption but it varies between summer and winter seasons. Therefore, correct EFs could not be calculated, and also is considered as marginal for our study.

355

Lines 216-217: Nap is not a common indicator due to its ubiquitous presence and often high blank sampling material contamination; thus, Nap was not considered as potential marker.

Reviewer's comment:

Blank contamination by Nap, was observed in this study.

360

Author's response:

The text was corrected as follows:

Page 6, line 219: Due to its ubiquitous presence, Nap was not considered as suitable marker.

Line 218: Phe, Flu, Flt, Pyr, 9-Flu, and 9,10-AntQ were the main PAHs detected in the Longyearbyen PP flue gas

Reviewer's comment:

PACs. 9-Fluo and 9,10-Ant are oxy-PAHs

Author's response:

The corrected text states:

Page 6, line 219: Phe, Flu, Flt, Pyr, 9-Flu, and 9,10-AntQ were the main PAHs and oxy-PAHs detected in the Longyearbyen PP flue gas

Reviewer's comment on Figure 2, line 218:

I would probably modify this figure as follows. 4 pie charts: 1) proportions of PAHs, oxy-PAHs and nitro-PAHs over total PACs; 2) pie for PAHs only; 3) Pie for oxy-PAHs only; 4) Pie for nitro-PAHs only

Author's response:

The Figure 2 was modified as suggested

Line 226: The concentrations of PAHs measured at UNIS was a factor of 2 higher than at Adventdalen

Reviewer's comment:

to change "was" to "were"

Author's response:

Corrected accordingly

Line 226-227: The concentrations of PAHs measured at UNIS was a factor of 2 higher than at Adventdalen, while the PAH profiles were similar.

Reviewer's comment:

They (PAH chemical profiles) are not shown neither in the main text nor in the SM.

Author's response:

The PAH, oxy- and nitro-PAH chemical profiles are now included in the SI (Figure S2). The updated text states:

Page 6, line 232: The UNIS and Adventdalen chemical profiles of PAHs and oxy-PAHs were similar, while the profiles of nitro-PAHs differ (Fig. S2). Proportions of 1- and 2-NNap were higher in UNIS samples (about 60% of $\Sigma 22$ nitro-PAHs), while 9-NAnt and 2+3-NFlt showed higher contributions into the nitro-PAH profile of samples from Adventdalen (about 55% of $\Sigma 22$ nitro-PAHs).

Reviewer's comment, line 227:

the number of compounds quantified should be specified all along the text. Same for PP. Same for nitro-PAHs

Author's response:

The text is amended accordingly

Reviewer's comment, line 229:

the number of compounds quantified should be specified all along the text. Same for PP

Author's response:

The text is amended accordingly

405 Lines 231-233: The PAH concentrations measured at Longyearbyen (UNIS and Adventdalen) were two orders of magnitude higher than those detected at the Zeppelin station and the same order of magnitude as in Birkenes (southern mainland Norway) (NILU, 2019) (Table S13) for the same period (autumn 2018).

Reviewer's comment:

These results could be also compared to the ones reported by Yu *et al.*, 2019.

- 410
- Yu, Y., Katsoyiannis, A., Bohlin-Nizzetto, P., Brorström-Lundén, E., Ma, J., Zhao, Y., Wu, Z., Tych, W., Mindham, D., Sverko, E., Barresi, E., Dryfhout-Clark, H., Fellin, P. and Hung, H.: Polycyclic Aromatic Hydrocarbons Not Declining in Arctic Air Despite Global Emission Reduction, *Environ. Sci. Technol.*, 53(5), 2375–2382, doi:10.1021/acs.est.8b

Author's response:

415 A dedicated Table S14 with appropriate references is added for comparison. The text is adapted accordingly:

Page 6, line 240: The measured PAH concentrations in the present study were in agreement with the 2 decades average data reported for the Arctic monitoring stations in Svalbard (Zeppelin) and Finland (Pallas), and were about an order of magnitude higher compared to the Canadian Arctic (Alert) concentrations (Yu *et al.*, 2019). The PAH levels observed in Longyearbyen were significantly (up to 2 orders of magnitude) lower compared to rural sites in Europe and China (Table S14).

420

Line 223: Table S13, UNIS and Adventdalen atmospheric concentrations (G+P) of $\Sigma 16$ PAHs compared to national and regional background concentrations detected in autumn 2018.

Reviewer's comment:

425 In addition to this Table, it would be easier to show a graph presenting the average (+SD) PAH chemical profiles for each site.

Author's response:

A graphic presentation is provided in the SI Figure S3 and the text added.

Page 6, line 239: The PAH profiles were dominated by Phe and Flu at all sites. A higher proportion of Phe was observed in Longyearbyen samples.

430

Lines 234- 235: Phe and Flu also dominated the PAH profile at Zeppelin station, which may indicate similar sources of contamination.

Reviewer's comment:

435 I won't be so conclusive in terms of sources

Author's response:

We agree, the revised text states:

Page 6, line 239: The PAH profiles were dominated by Phe and Flu at all sites. A higher proportion of Phe was observed in Longyearbyen samples.

440

Reviewer's comment, lines 236:

The data about nitro- and oxy-PAHs are scarce in the literature for the arctic area but it would be interesting to compare these results in terms of concentrations; difference of order of magnitude between PAHs, oxy-PAHs, and nitro-PAHs; chemical profiles; with results from the literature for other locations worldwide (including for rural, altitude or remote sites). For

instance, is the dominance of 9-Fluo and 9,10-AntQ is specific of your sites or it is something commonly observed in ambient air? Same for 1-NN and 2-NFlt? Same for both locations.

Author's response:

A dedicated Table S14 with appropriate references is added for comparison. The text is adopted accordingly. The revised text provides this comparison:

Page 7, line 249: The sum of oxy-PAHs detected at UNIS were similar to a rural site in eastern England (Alam et al., 2014), and a factor of 2 higher than at a rural location in the Czech Republic (Lammel et al., 2020), but were significantly lower than in rural southern China (Huang et al., 2014a) and the French Alps (Albinet et al., 2008).

Page 7, line 256: Overall, nitro-PAH concentrations were similar to those reported for the Pallas and Råö Scandinavian stations (Brorström-Lundén et al., 2010), and the rural site in the Czech Republic which is representative of central European background levels (Lammel et al., 2020).

Line 236-240: Concentrations of 9-Flu and 9,10-AntQ were the highest among measured oxy-PAHs in the present study. The 9-Flu level ($270.3 \pm 146.9 \text{ pg m}^{-3}$ at UNIS and $139.4 \pm 24.9 \text{ pg m}^{-3}$ in Adventdalen) was the same order of magnitude as reported for the background monitoring stations in the north of Finland (Pallas) and in the south of Sweden (Råö) (Brorström-Lundén et al., 2010), while 9,10-AntQ ($163.5 \pm 57.4 \text{ pg m}^{-3}$ at UNIS and $71.7 \pm 39.2 \text{ pg m}^{-3}$ in Adventdalen) was an order of magnitude higher in Longyearbyen. The nitro-PAH levels in our study were overall lower than other reported background sites.

Reviewer's comment on the last sentence:

That would need some references

Author's response:

The nitro-PAHs levels were compared with the same background stations data as for oxy-PAHs reported by Brorström-Lundén et al. (2010). The reference was already included. The revised text states:

Page 7, line 256: Overall, nitro-PAH concentrations were similar to those reported for the Pallas and Råö Scandinavian stations (Brorström-Lundén et al., 2010), and the rural site in the Czech Republic which is representative of central European background levels (Lammel et al., 2020).

Line 244: Gas/particle partitioning

Reviewer's comment:

As mentioned above, these results should be compared to the one available in the literature for other locations worldwide (in winter season to get comparable temperatures).

- Delgado-Saborit, J. M., Alam, M. S., Godri Pollitt, K. J., Stark, C. and Harrison, R. M.: Analysis of atmospheric concentrations of quinones and polycyclic aromatic hydrocarbons in vapour and particulate phases, *Atmospheric Environment*, 77, 974–982, doi:10.1016/j.atmosenv.2013.05.080, 2013.
- Alam, M. S., Delgado-Saborit, J. M., Stark, C. and Harrison, R. M.: Using atmospheric measurements of PAH and quinone compounds at roadside and urban background sites to assess sources and reactivity, *Atmospheric Environment*, 77, 24–35, doi:10.1016/j.atmosenv.2013.04.068, 2013.
- Huang, B., Ming Liu, Bi, X., Chaemfa, C., Ren, Z., Wang, X., Sheng, G. and Fu, J.: Phase distribution, sources and risk assessment of PAHs, NPAHs and OPAHs in a rural site of Pearl River Delta region, China, *Atmospheric Pollution Research*, 5(2), 210–218, doi:10.5094/APR.2014.026, 2014.
- Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E. and Jaffrezo, J.-L.: Nitrated and oxygenated derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleys: Part 1: Concentrations, sources and gas/particle partitioning, *Atmospheric Environment*, 42(1), 43–54, doi:10.1016/j.atmosenv.2007.10.009, 2008.
- Shahpoury, P., Lammel, G., Albinet, A., Sofuoglu, A., Dumanoglu, Y., Sofuoglu, S. C., Wagner, Z. and Zdimal, V.: Evaluation of a conceptual model for gas-particle partitioning of polycyclic aromatic hydrocarbons using poly-parameter linear free energy relationships, *Environ. Sci. Technol.*, doi:10.1021/acs.est.6b02158, 2016.

- Tomaz, S., Shahpoury, P., Jaffrezo, J.-L., Lammel, G., Perraudin, E., Villenave, E. and Albinet, A.: One-year study of polycyclic aromatic compounds at an urban site in Grenoble (France): Seasonal variations, gas/particle partitioning and cancer risk estimation, *Science of The Total Environment*, 565, 1071–1083, doi:10.1016/j.scitotenv.2016.05.137, 2016.

Author's response:

We appreciate the constructive support of the reviewer and implemented the below reports in the discussion of the Gas/particle partitioning section. The additional text provided the comparison:

Page 8, line 291: In general, the obtained %PM were in agreement with the earlier reported (Table S18). Higher %PM of 9,10-AntQ and several nitro-PAHs (1- and 2-NNap, 2-NFlu, 9-NAnt, 9-NPhen) were detected in French alpine sites in winter (Albinet et al., 2008), while higher %PM of Flt, 9-Flu, and cPPhen-4 found in the present study contrasts with those reported for temperate urban and rural sites in China and Europe (Huang et al., 2014a; Tomaz et al., 2016). Sources difference, weather influence such as precipitation and temperature, as well as different atmospheric conditions (e.g., number of suspended particles, mass size particle distribution, and specific humidity), are likely responsible for these variations.

Lines 245-246: Gas/particle partitioning is an important process that controls transport, degradation, and distribution patterns of contaminants in and between environmental compartments (Huang et al., 2014a).

Reviewer's comment:

This not the right reference to support this.

- Franklin, J.; Atkinson, R.; Howard, P. H.; Orlando, J. J.; Seigneur, C.; Wallington, T. J.; Zetzsch, C., *Quantitative determination of persistence in air. In Evaluation of Persistence and Long-Range Transport of Chemicals in the Environment*; Klečka, G., et al., Ed.; SETAC Press: Pensacola, USA, 2000; pp 7-62.
- Bidleman, T. F. *Atmospheric processes. Environ. Sci. Technol.* 1988, 22 (4), 361-367.
- Finlayson-Pitts, B. J., Pitts, J.N. *Chemistry of the Upper and Lower Atmosphere: Theory, Experiments, Application*; Academic Press: San Diego, USA, 2000.
- Lammel, G.; Sehili, A. M.; Bond, T. C.; Feichter, J.; Grassl, H. *Gas/particle partitioning and global distribution of polycyclic aromatic hydrocarbons – a modelling approach. Chemosphere* 2009, 76 (1), 98-106.

Author's response:

The reference list was updated. Thank you.

Lines 248-250: In general, LMW PAHs were found in the gas phase, while HMW PAHs were present in the particulate phase (Table 1), which is in accordance to their physico-chemical parameters, such as octanol-air partition coefficient (Table S3).

Reviewer's comment:

Other parameters are also to consider like the PM chemical composition, the vapor pressure (and so temperature and molecular weight).

- Pankow, J. F. *Review and comparative analysis of the theories on partitioning between the gas and aerosol particulate phases in the atmosphere. Atmos. Environ.* 1987, 21 (11), 2275-2283.
- Lohmann, R.; Lammel, G. *Adsorptive and absorptive contributions to the gas-particle partitioning of polycyclic aromatic hydrocarbons: state of knowledge and recommended parametrization for modeling. Environ. Sci. Technol.* 2004, 38 (14), 3793-3803.
- Shahpoury, P., Lammel, G., Albinet, A., Sofuoglu, A., Dumanoglu, Y., Sofuoglu, S. C., Wagner, Z. and Zdimal, V.: *Evaluation of a conceptual model for gas-particle partitioning of polycyclic aromatic hydrocarbons using poly-parameter linear free energy relationships, Environ. Sci. Technol.*, doi:10.1021/acs.est.6b02158, 2016.
- Tomaz, S., Shahpoury, P., Jaffrezo, J.-L., Lammel, G., Perraudin, E., Villenave, E. and Albinet, A.: One-year study of polycyclic aromatic compounds at an urban site in Grenoble (France): Seasonal variations, gas/particle partitioning and cancer risk estimation, *Science of The Total Environment*, 565, 1071–1083, doi:10.1016/j.scitotenv.2016.05.137, 2016.

Author's response:

The list of the relevant physical-chemical properties was extended accordingly. Weather influence is discussed later in this section. We revised the text as follows:

Page 7, line 264: In general, LMW PAHs were found in the gas phase, while HMW PAHs were present in the particulate phase (Table 1), which is in accordance to their physico-chemical parameters, such as octanol-air partition coefficient, vapor pressure, and molecular (Table S3) (Tomaz et al., 2016; Shahpoury, 2016).

Page 7, line 270: %PM also depends on aerosol surface area, organic matter, and black carbon content (Lohmann and Lammel, 2004).

Line 254: Strong negative

correlations (Spearman coefficient > 0.65) of percentage of PAH determined in particulate phase (%PM) with ambient temperature and specific humidity were determined for Chry, 1-NNap, 2-NNap, cPhe-4, and 2+3-NFlt, although they were not statistically significant (Table S14).

Reviewer's comment:

These compounds, and most of the compounds shown in Table S14, are mainly associated to the gas phase. In my opinion, it is difficult to conclude anything about the link between humidity and gas/particle partitioning based in such conditions.

Author's response:

We agree with this note. The text has now been amended to reflect temperature influence on the partitioning of the earlier listed in this paragraph compounds, for which large migration between the phases was observed. Potential influence of humidity is discussed later in this section (lines 284-290). The text was updated:

Page 7, line 266: Repartitioning between phases (Fig. 3) mainly impacted semi-volatile compounds with three and four aromatic rings (Flt, Pyr, BaAnt, Chry; 2-NFlu, 9-Flu, cPhe-4, 9,10-AntQ, 9-NAnt, and 2+3-NFlt) as a response to changing meteorological conditions (Hu et al., 2019). Strong negative correlations of percentage of PAH determined in particulate phase (%PM) with ambient temperature were confirmed for most of these compounds (Table S15). %PM also depends on aerosol surface area, organic matter, and black carbon content (Lohmann and Lammel, 2004).

Lines 265-268: The influence of wet deposition was indicated by a significant negative correlation between concentrations of several HMW PAHs (Pyr, Chry, BbkFlt, IPyr, BPer, BaFlu-11, and BaAnt-7,12) and precipitation (Spearman correlation, $p < 0.05$, Table S15), resulting in a lower amount of particle-bound PAHs transported from the town, and thus lower %PM in Adventdalen.

Reviewer's comment:

What about particulate nitro-PAHs like 2+3-NFlt?

Author's response:

The negative correlation between concentrations and amount of precipitation was also found for 2+3-NFlt (80% PM) but it was not significant (Spearman correlation = -0.145, $p < 0.05$), and thus not included in the Table S15.

Reviewer's comment:

Why Flt has not the same behavior like Pyr? The authors should consider the following article in their discussion (paper cited later in the article, line 348):

- Škrdlíková, L., Landlová, L., Klánová, J. and Lammel, G.: Wet deposition and scavenging efficiency of gaseous and particulate phase polycyclic aromatic compounds at a central European suburban site, *Atmospheric Environment*, 45(25), 4305–4312, doi:10.1016/j.atmosenv.2011.04.072, 2011.

Author's response:

We thank the reviewer for this insight full comment, and we think that these compounds do have similar behavior. We have now included compound specific aspects and discussions on environmental behavior of these compounds in a dedicated section. The new text states:

Page 7, line 281: The influence of wet deposition was indicated by a significant negative correlation between amount of precipitation and concentrations of several particle-bound HMW PAHs (Chry, BbkFlt, IPyr, BPer, BaFlu-11, and BaAnt-7,12) as well as semi-volatile Phe, Flt, and Pyr, which are more predominant in gaseous phase (Spearman correlation, $p < 0.05$, Table S16. Effective wet scavenging of Phe, Flt, and Pyr has been suggested (Škrdlíková et al., 2011). Furthermore, a strong negative correlation with mass of water vapor in the air (specific humidity) was determined for most of the compounds (Spearman correlation, $p < 0.05$, Table S17). Particle associated HMW compounds are readily scavenged by precipitation, while water solubility and polarity (for nitro- and oxy-PAHs) play an additional role in wet scavenging processes (Shahpoury et al., 2018). The gas phase removal from the atmosphere is due to substance dissolution in water droplets, which enhances the scavenging effect at higher humidity. Higher sensitivity of gas scavenging compared with particle scavenging towards liquid water content was also indicated by Škrdlíková et al. (2011).

Line 269: “Source identification” section

Reviewer’s comment:

Reviewer’s comment on “Source identification” section: For all of this part, I would suggest the authors instead of running the PCA only using the PAC concentrations, to also run it using the diagnostic ratios from Table 4 for both UNIS and Adventdalen. Separately or together with the concentrations (maybe not all the compounds but a selection of the key ones). Please also consider these publications in the choice of the ratios and in the discussion:

- Dvorská, A., Lammel, G. and Klánová, J.: Use of diagnostic ratios for studying source apportionment and reactivity of ambient polycyclic aromatic hydrocarbons over Central Europe, *Atmospheric Environment*, 45(2), 420–427, doi:10.1016/j.atmosenv.2010.09.063, 2011.
- Katsoyiannis, A., Sweetman, A. J. and Jones, K. C.: PAH molecular diagnostic ratios applied to atmospheric sources: a critical evaluation using two decades of source inventory and air concentration data from the UK, *Environ. Sci. Technol.*, 45(20), 8897–8906, doi:10.1021/es202277u, 2011.

In addition, the authors should consider the ratios used in the study by Yu et al., 2019.

- Yu, Y., Katsoyiannis, A., Bohlin-Nizzetto, P., Brorström-Lundén, E., Ma, J., Zhao, Y., Wu, Z., Tych, W., Mindham, D., Sverko, E., Barresi, E., Dryfhout-Clark, H., Fellin, P. and Hung, H.: Polycyclic Aromatic Hydrocarbons Not Declining in Arctic Air Despite Global Emission Reduction, *Environ. Sci. Technol.*, 53(5), 2375–2382, doi:10.1021/acs.est.8b

Author’s response:

All the suggested papers were studied, and the ratios correlated best with the source specific PAHs were preferred. The applicability and scientific value of diagnostic ratios for PAH source elucidation was comprehensively discussed in the revised version. The revised text states:

Page 8, line 314: Diagnostic ratios may be affected by large-scale mixing of PAHs in the atmosphere, differing emission rates of PAH from the same source, influence of changing environmental conditions, and atmospheric processing of individual PAH compounds with different atmospheric lifetimes and reactivities (Alam et al., 2013; Tobiszewski and Namieśnik, 2012; Katsoyiannis and Breivik, 2014). Ratios based on highly reactive compounds such as Ant and BaAnt were not included, while more stable HMW PAHs diagnostic ratios were interpreted with greater confidence (Galarneau, 2008; Alam et al., 2014). Yunker et al. (2002) previously proposed the ratio of IPyr/(IPyr+BPer) to recognize vehicle from coal combustion emissions. BbkFlt/BPer was selected as an additional marker ratio for traffic due to the greater capacity to discriminate diesel and gasoline emissions, as well as its wider value range (Kuo et al., 2013). The Flt/(Flt+Pyr) ratio is widely used for source identification and, in particular, to understand if PAHs are mainly emitted from petroleum sources or

from combustion processes (Yu et al., 2019). The Flu/(Flu+Pyr) ratio was selected as a specific indicator for coal combustion due to its strong correlation with the local PP determined markers, and the ratio value was also in agreement with literature (Yunker et al., 2002; Katsoyiannis and Breivik, 2014).

We are thankful for the suggestion to include the diagnostic ratios into PCA, which were adopted in the revised version. The multivariate statistical part was completely revised and rewritten (revised text is marked in red color font). Similarly to Albinet et al. (2007), we have removed the samples grouping interpretation because this discussion was distracting from the main point of our goal in this section. Our focus here is intended to be on the evaluation of the potential sources but not the weather influence. However, general discussion on the weather influences is now included in the gas/particle partitioning part (page 7, lines 281-292), as mentioned earlier.

- Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., and Jaffrezo, J. L.: Nitrated and oxygenated derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleys Part 1: Concentrations, sources and gas/particle partitioning, *Atmos. Environ.*, 42, 43-54, 10.1016/j.atmosenv.2007.10.009, 2008.

Reviewer's comment, lines 278-279:

If the use of highly reactive compounds like BaA, BaP etc should be avoided, why the authors used the "diagnostic ratios" BaA/(BaA+Chry) and BaP/BPer or BaP/(BaP+Chry) (and in Table 3 Ant/(Ant+Phen)) in their source evaluation?

Author's response:

We were calculating those ratios because they have been applied and shown useful in previous studies. Hence, these calculations allow a better comparability with similar studies previously reported. However, these ratios are removed from the revised version of the manuscript.

Reviewer's comment, lines 281-282:

Just above it is mentioned: "Due to high atmospheric reactivity of Ant and BaAnt, utilization as source apportionment should be avoided". The authors should justify finally the use of this ratio. The authors should also consider the following publications about the use of the PAH diagnostic ratios in terms of source apportionment (and cite them to moderate their conclusions sometimes):

- Dvorská, A., Lammel, G. and Klánová, J.: Use of diagnostic ratios for studying source apportionment and reactivity of ambient polycyclic aromatic hydrocarbons over Central Europe, *Atmospheric Environment*, 45(2), 420–427, doi:10.1016/j.atmosenv.2010.09.063, 2011.
- Katsoyiannis, A., Sweetman, A. J. and Jones, K. C.: PAH molecular diagnostic ratios applied to atmospheric sources: a critical evaluation using two decades of source inventory and air concentration data from the UK, *Environ. Sci. Technol.*, 45(20), 8897–8906, doi:10.1021/es202277u, 2011.

Author's response:

All the suggested papers were studied, and the ratios correlated best with the source specific PAHs were preferred. As example of the decision making: Dvorská et al. (2011) suggested Flt/(Flt+PYR) ratio as coal combustion characteristic but no correlation with the local PP detected marker was found (PCA was applied). Another suggested coal specific IPyr/(IPyr+BghiPer) ratio was not applicable for our study because IPyr and BghiPer were not detected in the local PP plume as a result of an effective exhaust cleaning system. Thus, another ratios were chosen.

Reviewer's comment, lines 290-291:

I have exactly the same comment as Reviewer 1. I would have expected different scavenging due to rainfalls or snowfalls. Were A5 concentrations lower than A2?

Author's response:

Yes, the reviewer assume correctly. The concentrations on the rainy day 2 were significantly lower compared to the detected levels on the day 5 with snow event, $\Sigma 46$ PAHs 430.9 pg m⁻³ on day 2 (rain) and ($\Sigma 46$ PAHs 635.3 pg m⁻³ on day 5 (snow). This was only true for Adventdalen. At UNIS the levels on the days 2 and 5 were nearly equal. The UNIS sampling station is located closer to the town and thus more influenced by emissions from different settlement sources. Higher emissions from any source on the day 2 can be a possible reason of non-observed scavenging effect of rain at UNIS. For instance, more boats were in the marine port of Longyearbyen during the day 2 sampling period (week 34 on the new Figure S5) compared to all the other days. A passenger cruise ship Artania (MMSI: 311000608) was notably the biggest vessel (230 x 32 m, 44656 t gross tonnage) in the harbor on day 2.

<https://www.marinetraffic.com/en/ais/details/ships/shipid:371216/mmsi:311000608/imo:8201480/vessel:ARTANIA>

However, to focus on the potential sources only (the original goal of this section of the manuscript), the in-depth discussion on the difference between 2nd and 5th sampling days is now excluded since this topic is considered outside the scope of the study. More general discussion about weather influence is now included in the section 3.2.2 about the gas/particle partitioning.

Line 292: This indicates that the mass of water vapor in the air (specific humidity), in contrast to relative humidity, is an essential parameter for removal from the atmosphere.

Reviewer's comment:

Does it means like fog processing (by opposition to scavenging due to the rainfalls)? I suppose yes, so the following sentences (lines 299-303) should move here "The gas phase removal from the atmosphere is due to substance dissolution in water droplets (Shahpoury et al., 2018), which enhances the scavenging effect at higher humidity. A strong negative correlation with humidity was determined for all quantified LMW PAHs, significant for Acy, Pyr, 1-NNap, and 9-Flu (Spearman correlation, p<0.05; Table S17). Presence of Ant and Flt (gas-phase PAHs with low polarity and water solubility) in the group is likely due to the same source of origin."

Author's response:

We agree it is probably correct to equate elevated humidity with fog processing. However, as reported earlier, general discussion on this topic is now moved to the Section 3.2.2. The new text states:

Page 7, line 281: The influence of wet deposition was indicated by a significant negative correlation between amount of precipitation and concentrations of several particle-bound HMW PAHs (Chry, BbkFlt, IPyr, BPer, BaFlu-11, and BaAnt-7,12) as well as semi-volatile Phe, Flt, and Pyr, which are more predominant in gaseous phase (Spearman correlation, p<0.05, Table S16. Effective wet scavenging of Phe, Flt, and Pyr has been suggested (Škrdlíková et al., 2011). Furthermore, a strong negative correlation with mass of water vapor in the air (specific humidity) was determined for most of the compounds (Spearman correlation, p<0.05, Table S17). Particle associated HMW compounds are readily scavenged by precipitation, while water solubility and polarity (for nitro- and oxy-PAHs) play an additional role in wet scavenging processes (Shahpoury et al., 2018). The gas phase removal from the atmosphere is due to substance dissolution in water droplets, which enhances the scavenging effect at higher humidity. Higher sensitivity of gas scavenging compared with particle scavenging towards liquid water content was also indicated by Škrdlíková et al. (2011).

Reviewer's comment, line 304:

Let's not call that tracers cause it implies chemical stability, specificity to the emission sources that are not the case for any PAC. "based on some source indicators..."

Author's response:

This term is now removed as suggested.

710

Lines 304-305: Based on the tracers and their loadings (Table S16), PC1 can be assigned to local PP coal burning (Flu, Phe, Flt, Pyr, 9-Flu, 9,10-AntQ).

Reviewer's comment:

Is it based on the local PP chemical profile obtained here? It should be specified

715

Author's response:

Yes, the most abundant compounds detected in the local PP emission were used as markers in this study. It is clarified in the revised text:

Page 8, line 323: **The Flu/(Flu+Pyr) ratio was selected as a specific indicator for coal combustion due to its strong correlation with the local PP determined markers, and the ratio value was also in agreement with literature (Yunker et al., 2002; Katsoyiannis and Breivik, 2014).**

720

Reviewer's comment, line 306:

What is the proportion of diesel vs gasoline vehicle engines in Svalbard?

Author's response:

725

Additional information is added as text and Table S19 in the updated SI.

Page 10, line 376: **1114 private cars were registered in Longyearbyen in 2018 (sbb.no), including old and modern (Euro 3-7 emission standard) technology cars, approximately equally balanced between gasoline and diesel fuel.**

Reviewer's comment, lines 309-310:

730

This would need references to support the secondary origin of these compounds and their use in such case

Author's response:

Appropriate references are added.

Page 9, line 364: These results indicate atmospheric formation as an additional source of 9-NAnt and 2+3-NFlt, in agreement with other studies (Lin et al., 2015b; Hayakawa et al., 2000; Shahpoury et al., 2018).

735

Line 321: Relative contribution of primary and secondary sources of nitro-PAHs could be tested by applying a 2-NFlt/1-NPyr ratio (Zielinska et al., 1989).

Reviewer's comment:

It is not the right reference. Prefer:

740

- Ciccioni, P., Cecinato, A., Brancaleoni, E., Frattoni, M., Zacchei, P., Miguel, A.H., De Castro Vasconcellos, P., 1996. Formation and transport of 2-nitrofluoranthene and 2-nitropyrene of photochemical origin in the troposphere. *J. Geophys. Res.* 101, 19567e19581. <http://dx.doi.org/10.1029/95JD02118>.
- Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., 2007. Polycyclic aromatic hydrocarbons (PAHs), nitrated PAHs and oxygenated PAHs in ambient air of the Marseilles area (South of France): concentrations and sources. *Sci. Total Environ.* 384, 280e292. <http://dx.doi.org/10.1016/j.scitotenv.2007.04.028>.
- Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., Jaffrezo, J.-L., 2008a. Nitrated and oxygenated derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleys: Part 1: concentrations, sources and gas/particle partitioning. *Atmos. Environ.* 42, 43e54. <http://dx.doi.org/10.1016/j.atmosenv.2007.10.009>.
- Bamford, H.A., Baker, J.E., 2003. Nitro-polycyclic aromatic hydrocarbon concentrations and sources in urban and suburban atmospheres of the Mid-Atlantic region. *Atmos. Environ.* 37, 2077e2091. [http://dx.doi.org/10.1016/S1352-2310\(03\)00102-X](http://dx.doi.org/10.1016/S1352-2310(03)00102-X).

745

750

- Ringuet, J., Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., 2012b. Diurnal/nocturnal concentrations and sources of particulate-bound PAHs, Oxy- PAHs and Nitro-PAHs at traffic and suburban sites in the region of Paris (France). *Sci. Total Environ.* 437, 297e305. <http://dx.doi.org/10.1016/j.scitotenv.2012.07.072>.
- Tomaz, S., Jaffrezo, J.-L., Favez, O., Perraudin, E., Villenave, E. and Albinet, A.: Sources and atmospheric chemistry of oxy- and nitro-PAHs in the ambient air of Grenoble (France), *Atmospheric Environment*, 161, 144–154, doi:10.1016/j.atmosenv.2017.04.042, 2017.

Author's response:

We thank the reviewer for the additional references and extended the citations accordingly

Reviewer's comment, lines 322-323:

Again, what is the proportion of diesel vs gasoline vehicle engines in Svalbard? Could you really specify diesel? If diesel is so predominant, 1-NPyr should have been detected. See the following paper:

- Keyte, I. J., Albinet, A. and Harrison, R. M.: On-road traffic emissions of polycyclic aromatic hydrocarbons and their oxy- and nitro- derivative compounds measured in road tunnel environments, *Science of The Total Environment*, 566–567, 1131–1142, doi:10.1016/j.scitotenv.2016.05.152, 2016.

Author's response:

The proportion of diesel and gasoline passenger cars is equal (Table S19) but, based on own experience as a local permanent resident of Longyearbyen, we confirm that passenger cars are rarely driven outside the settlement. Due to proximity to the active coal mine N7 (about 7 km from the Adventdalen sampling station, Figure 1), regular coal transportation (5-10 times per day) by trucks and occasional tourist busses are the main vehicles in Adventdalen, both diesel driven.

The suggested study by Keyte et al. (2016) is mainly on passenger vehicles and not well suitable for modern heavy duty vehicles, considering last five years advances in the exhaust aftertreatment including gas recirculation, diesel particulate filter, diesel oxidation catalyst, selective catalytic reduction, ammonia oxidation catalyst and others.

The mining company provided us with the specific documentation on fuel quality, engine technology type of the trucks used for coal transportation. We confirm that the vehicles are only 1-2 years old and operate on high quality diesel (new Table S23) used for all vehicles in Longyearbyen. Recent studies on a modern technology heavy duty vehicles report substantial reduction of nitro-PAH concentrations, and 1-NPyr is not the dominating compound.

For instance, Liu et al. (2015) reports 8-10 orders of magnitude mitigation of 6-NChry, 7-NBaAnt, 4-NPyr and other nitro-PAHs, and 2 orders of magnitude reduction of 1-NPyr level as result of the similar to Longyearbyen trucks exhaust aftertreatment system.

- Liu, Z. G., Wall, J. C., Ottinger, N. A., and McGuffin, D.: Mitigation of PAH and Nitro-PAH Emissions from Nonroad Diesel Engines, *Environ. Sci. Technol.*, 49, 3662-3671, 10.1021/es505434r, 2015.

Thus, considering diesel motors modern technologies, 1,NPyr is not a strong marker of diesel emission from modern trucks, and due to actually low traffic in Adventdalen, 1-NPyr level was below the method detection limit. The new insights are now included in the text:

Page 9, lines 332-345: Because of the rural position, car traffic is much lower at this location. At the same time, due to the proximity to an active mine (Fig. 1), heavy-duty vehicles (coal trucks, tourist busses, geotechnical drilling machinery) are thus the main candidate source for PAH vehicle emissions. Produced coal is regularly delivered from the coal mine to PP and storage area in the harbor on a road situated in 150 m distance from the Adventdalen sampling station. Coal is transported by Volvo FH540 trucks (built in 2018-2020) driven on diesel CFPP-12 (NS-EN 590) (Nilssen P., Store Norske, personal communication; Table S23). The trucks have Euro 6 standard (HC 0.13, CO 1.5, NO_x 0.4, PM 0.01 g (kWh)⁻¹) (DieselNet, 2020) compliant Volvo D13K engines (540 hp, 405 kW) engines fitted with exhaust gas recirculation, diesel particulate filter, diesel oxidation catalyst, selective catalytic reduction, and ammonia oxidation catalyst (Volvo Trucks, 2020). These allow high operation temperatures and high efficiencies in reducing particle and NO_x emissions. Numerous studies showed substantial reduction in gaseous and particulate emissions of PAH, nitro- and oxy-PAHs as the result of such

mitigation in particle and NO_x emissions (Hu et al., 2013b; Gerald Liu et al., 2010; Khalek et al., 2015; Huang et al., 2015). Up to 10 orders of magnitude reduction in emission from similar to Volvo D13K heavy-duty engine was reported for several nitro-PAHs (6-NChry, 1-NPyr, 2-NPyr, 4-NPyr, 7-NBaAnt) (Liu et al., 2015; Gerald Liu et al., 2010), which were not detected in the present study most likely due to low vehicle number in Adventdalen.

Lines 329-330: However, studies by Wania et al. (1999) report that snow scavenging may be an important, and sometimes a dominating scavenging process for lighter PAHs, mediated via a process of adsorption to the air-ice (Wania et al., 1999).

Reviewer's comment:

That should be specified before, lines 290-293.

Author's response:

The discussion on the difference between snow and rain scavenging efficiency is not included in the text anymore. Thus, this sentence was deleted.

Line 331: Traffic emission (mainly diesel exhaust) and the Longyearbyen coal-burning PP were the main local sources of PAHs and nitro- and oxy-PAHs in Adventdalen, and atmospheric transformation of PAHs is an additional source of nitro-PAHs.

Reviewer's comment:

(mainly diesel exhaust) to be moderated

Author's response:

We are convinced that the new provided above text supports this statement better than before. The sentence was not changed.

Lines 334-335: LMW PAHs were scavenged by snow, while the level of humidity was an essential parameter for total PAH removal from the atmosphere.

Reviewer's comment:

OK. See comment line 290 and the one from reviewer 1.

Author's response:

The discussion on the difference between snow and rain scavenging efficiency is not included in the text anymore. Higher sensitivity of gas scavenging than particle scavenging towards liquid water content was also concluded by Škrdlíková, L., Landlová, L., Klánová, J., and Lammel, G.: *Wet deposition and scavenging efficiency of gaseous and particulate phase polycyclic aromatic compounds at a central European suburban site*, *Atmos. Environ.*, 45, 4305-4312, <https://doi.org/10.1016/j.atmosenv.2011.04.072>, 2011. Our proposal on the importance of humidity level on the total scavenging is in agreement with this long-term study.

Line 342: several of these compounds (BPer, IPyr, and BaFlu-11) were reported to be emitted after gasoline burning as well (Zielinska et al., 2004; Albinet et al., 2007)

Reviewer's comment to include the additional references:

- Keyte, I. J., Albinet, A. and Harrison, R. M.: *On-road traffic emissions of polycyclic aromatic hydrocarbons and their oxy- and nitro- derivative compounds measured in road tunnel environments*, *Science of The Total Environment*, 566–567, 1131–1142, doi:10.1016/j.scitotenv.2016.05.152, 2016.
- Rogge, W.F., Hildemann, L.M., Mazurek, M.A., Cass, G.R., Simoneit, B.R.T., 1993a. *Sources of fine organic aerosol 0.2. Noncatalyst and catalyst-equipped automobiles and heavy duty diesel trucks*. *Environ. Sci. Technol.* 27, 636–651.

- Miguel, A.H., Kirchstetter, T.W., Harley, R.A., Hering, S.V., 1998. On-road emissions of particulate polycyclic aromatic hydrocarbons and black carbon from gasoline and diesel vehicles. *Environ. Sci. Technol.* 32, 450–455.
- Li, C.K., Kamens, R.M., 1993. The use of polycyclic aromatic hydrocarbons as source signatures in receptor modeling. *Atmos. Environ.* 27, 523–532.

Author's response:

References added. Thank you.

Line 342: Absence of 1-NPyr and 2-NFlu, the principal compounds of diesel exhaust (Albinet et al., 2007), supported gasoline combustion as more dominant source as well.

Reviewer's comment:

It is the same at Adventdalen

Author's response:

For the UNIS sample location we now report general traffic emission (diesel and gasoline) as an important potential emission source. Although, explanations on the below detection limit level of 1-NPyr are provided:

Page 10, line 388: The diesel emission predominance was found for two out of the six sampling days, although particulate phase 1-NPyr, a marker of diesel emissions, was not detected. 1-NPyr forms in the combustion chamber of diesel engines by the addition of nitrogen oxide or nitrogen dioxide to free Pyr radicals (IARC, 2014). Its generation is facilitated by the high engine temperatures (IARC, 2014; Karavalakis et al., 2010; Guan et al., 2017; Huang et al., 2015), which likely can not be reached in Longyearbyen due to short driving distances and low speed limit. The use of high quality ultra-low sulfur fuel with substantially reduced emissions of NO_x leads to reduced nitration of PAHs during fuel combustion (Heeb et al., 2008; Zhao et al., 2020b), and together with low total vehicle number, resulting in low nitro-PAH emissions. Occurred atmospheric deposition may be of influence too.

Line 343: Absence of 1-NPyr and 2-NFlu, the principal compounds of diesel exhaust (Albinet et al., 2007), supported gasoline combustion as more dominant source as well.

Reviewer's comment to include additional reference:

- Keyte, I. J., Albinet, A. and Harrison, R. M.: On-road traffic emissions of polycyclic aromatic hydrocarbons and their oxy- and nitro- derivative compounds measured in road tunnel environments, *Science of The Total Environment*, 566–567, 1131–1142, doi:10.1016/j.scitotenv.2016.05.152, 2016.

Author's response:

1-NPyr and 2-NFlu are not used in this context in the revised version of the manuscript.

Lines 349-351: Zhang et al. (2015) reported that scavenging of particulate-phase PAHs is about 20 times more efficient than scavenging of gas phase PAHs.

Reviewer's comment:

This should be cited in the discussion above for Adventdalen (line 290-303)

Author's response:

This sentence does not exist in the updated version

Lines 353-354: The first group was characterized by the presence of 9,10-AntQ, Phe, and Flu, attributable to the PP emission source.

Reviewer's comment:

Is it based on the local PP chemical profile obtained here? It should be specified

Author's response:

Yes, it is based on the local PP emission profile. It was specified in the text.

Page 6, line 219: Further, Phe, Flu, Flt, Pyr, 9-Flu, and 9,10-AntQ were the main PAHs and oxy-PAHs detected in the Longyearbyen PP flue gas (Fig. 2), therefore the presence and diagnostic ratios (Table 3) of these compounds were used as markers of the PP source in the present work.

Line 354: Diagnostic ratios of Flu/(Flu+Pyr) and BaAnt/(BaAnt+Chry) (Table 4) also indicated that the PP is a source of PAHs and nitro- and oxy-PAHs at UNIS.

Reviewer's comment:

OK. Justify the use of such ratio BaAnt/(BaAnt+Chry) cause the sampling site is close to the PP and the time residence in the atmosphere to reach the sampling location is low.

Author's response:

The use of the BaAnt/(BaAnt+Chry) ratio was omitted as explained in the text (page 8, lines 314-326).

Reviewer's comment, line 355:

marker or indicator (not indicator)

Author's response:

The term is now changed all over the text.

Reviewer's comment, line 359:

I would be curious to see where the authors have seen that these compounds (4-NBip and 1,5-DNNap) are characteristics of diesel exhausts

Author's response:

Traffic emission source was concluded by (Alam et al., 2015), and heavy-duty diesel vehicles emission as source of 4-NBip by (Hu et al., 2013a). However, 4-NBip and 1,5-DNNap are not used in this context in the revised version.

- Alam, M. S., Keyte, I. J., Yin, J., Stark, C., Jones, A. M., and Harrison, R. M.: Diurnal variability of polycyclic aromatic compound (PAC) concentrations: Relationship with meteorological conditions and inferred sources, *Atmos. Environ.*, 122, 427-438, <https://doi.org/10.1016/j.atmosenv.2015.09.050>, 2015.
- Hu, S., Herner, J. D., Robertson, W., Kobayashi, R., Chang, M. C. O., Huang, S.-M., Zielinska, B., Kado, N., Collins, J. F., Rieger, P., Huai, T., and Ayala, A.: Emissions of polycyclic aromatic hydrocarbons (PAHs) and nitro-PAHs from heavy-duty diesel vehicles with DPF and SCR, *Journal of the Air & Waste Management Association*, 63, 984-996, [10.1080/10962247.2013.795202](https://doi.org/10.1080/10962247.2013.795202), 2013.

Reviewer's comment, line 359:

This one is secondary (2+3-NFlt) and not at all from diesel exhausts

Author's response:

These two isomers, 2-NFlt and 3-NFlt, were not baseline separated on a chromatogram. Thus we report its sum concentration as 2+3-NFlt. According to Alam et al. (2015) 2-NFlt forms in atmosphere, while 3-NFlt was reported in several studies in diesel exhaust (Heeb et al., 2008; Liu et al., 2015), as well as several references are given in SI Table in (Keyte et al., 2013). That is why the presence of 2+3-NFlt in diesel emission was stated. However, 2+3-NFlt is not used in this context in the revised version.

- Alam, M. S., Keyte, I. J., Yin, J., Stark, C., Jones, A. M., and Harrison, R. M.: Diurnal variability of polycyclic aromatic compound (PAC) concentrations: Relationship with meteorological conditions and inferred sources, *Atmos. Environ.*, 122, 427-438, <https://doi.org/10.1016/j.atmosenv.2015.09.050>, 2015.

- Heeb, N. V., Schmid, P., Kohler, M., Gujer, E., Zennegg, M., Wenger, D., Wichser, A., Ulrich, A., Gfeller, U., Honegger, P., Zeyer, K., Emmenegger, L., Petermann, J.-L., Czerwinski, J., Mosimann, T., Kasper, M., and Mayer, A.: *Secondary Effects of Catalytic Diesel Particulate Filters: Conversion of PAHs versus Formation of Nitro-PAHs*, *Environ. Sci. Technol.*, 42, 3773-3779, 10.1021/es7026949, 2008.
- Liu, Z. G., Wall, J. C., Ottinger, N. A., and McGuffin, D.: *Mitigation of PAH and Nitro-PAH Emissions from Nonroad Diesel Engines*, *Environ. Sci. Technol.*, 49, 3662-3671, 10.1021/es505434r, 2015.
- Keyte, I. J., Harrison, R. M., and Lammel, G.: *Chemical reactivity and long-range transport potential of polycyclic aromatic hydrocarbons – a review*, *Chem. Soc. Rev.*, 42, 9333-9391, <https://doi.org/10.1039/C3CS60147A>, 2013.

Reviewer's comment, lines 360-362:

This would be more convincing based on the diagnostic ratio. I would suggest the authors instead of running the PCA only using the PAC concentrations, to also run it using the diagnostic ratios from Table 4 for both UNIS and Adventdalen.

Author's response:

The followed the suggestion and revised the PCAs of UNIS and Adventdalen data accordingly. It allowed better interpretation of the potential sources. We thank the reviewer for this valuable suggestion.

Lines 379-380: In contrast to the study by Yu et al. (2019), no strong indication for LRAT biomass burning emissions was found for this set of air samples.

Reviewer's comment:

For that, it would have been necessary to use other markers like retene for PAHs and even better, levoglucosan. Biomass burning from forest fires? It should be specified. Were any major forest fires during the sampling period? The authors should provide the back trajectories for the studied sampling period. The samplings were in summer so the impact of wood combustion for residential heating purposes from Northern Europe or Russia would be negligible.

Author's response:

This sentence was deleted. To note, we sampled on the days with predicted unusual NW wind (in order to catch the PP emission), thus air mainly arrived from the north and Greenland, the areas without biomass burning. We provided the back trajectories in the updated SI (Figure S4), and the updated text states:

Page 8, line 288: **Due to changes in the Arctic front, more frequent precipitation, and very low levels of wood and coal burning for residential heating in the northern hemisphere in the summer, the LRAT of PAHs to the Arctic is low in summer. Sampling was performed on days with predicted northwesterly wind, and according to the 5-day back trajectory analysis, the air arriving to Svalbard in the sampling period mainly came from the north and from Greenland (Fig. S4). As discussed in Section 3.2.1, two orders of magnitude lower PAH concentrations were detected at the Zeppelin monitoring station compared to the levels in Longyearbyen on the same time. Thus, local emissions were the main sources of PAHs in Longyearbyen in this study.**

Reviewer's comment on Table 4:

I would not use these ratios BaPyr/(BaPyr +Chry) and BaPyr/BPer cause of non-stability of BaPyr. In addition, they are not mentioned in the text.

Author's response:

This is corrected. We appreciate the constructive comment of the reviewer.

Reviewer's comment on Figure 2 title:

PAH and PAH derivatives.

Author's response:

Corrected accordingly.

Polycyclic aromatic hydrocarbons (PAHs), oxy- and nitro-PAHs in ambient air of the Arctic town Longyearbyen, Svalbard

Tatiana Drotikova^{1,2*}, Aasim M. Ali², Anne Karine Halse³, Helena C. Reinardy^{4,1}, and Roland Kallenborn^{2,1}

¹University Centre in Svalbard (UNIS), Department of Arctic Technology, Longyearbyen, Svalbard, Norway

²Norwegian University of Life Sciences (NMBU), Faculty of Chemistry, Biotechnology and Food Sciences, Ås, Norway

³Norwegian Institute for Air Research (NILU), Department of Environmental Chemistry, Kjeller, Norway

⁴Scottish Marine Institute (SAMS), Department of Aquaculture and Environment, Oban, Argyll, PA37 1QA, UK

Correspondence to: Tatiana M. Drotikova (tatiana.drotikova@unis.no)

Abstract. Polycyclic aromatic hydrocarbons (PAHs) are not declining in Arctic air despite reductions in their global emissions. In Svalbard, the Longyearbyen coal-fired power plant is considered to be one of the major local source of PAHs. Power plant stack emissions and ambient air samples, collected simultaneously 1 km (UNIS) and 6 km (Adventdalen) transect distance, were analyzed (gaseous and particulate phases separately) for 22 nitro-PAHs, 9 oxy-PAHs and 16 parent PAHs by GC/ECNI/MS and GC-MS/MS. Results confirm low level of PAH emissions ($\sum 16$ PAHs = $1.5 \mu\text{g kg}^{-1}$ coal) from the power plant. ~~Phenathrene~~Phenanthrene, 9,10-anthraquinone, 9-fluorenone, fluorene, fluoranthene, and pyrene accounted for 85% of the plant emission (not including naphthalene). A dilution effect was observed for the transect ambient air samples, 1.26 ± 0.16 and $0.63 \pm 0.14 \text{ ng m}^{-3}$ sum all 47 PAH derivatives for UNIS and Adventdalen, respectively. The PAH profile was homogeneous for these recipient stations with ~~phenathrene~~phenanthrene and 9-fluorenone being most abundant. ~~Principal component~~Multivariate statistical analysis, ~~in combination with PAH diagnostic ratios and literature data on different source specific markers,~~ confirmed coal combustion, ~~gasoline,~~marine and ~~diesel~~vehicle traffic as the predominant sources of PAHs. Secondary atmospheric formation of 9-nitroanthracene and 2+3-nitrofluoranthene was evaluated and concluded. ~~Results also indicate that ambient PAH concentrations were affected by precipitation events, and specific humidity is an essential parameter influencing PAH scavenging from the air.~~PAHs partitioning between gaseous and particulate phases showed a strong dependence on ambient temperatures and humidity. The present study contributes important data which can be utilized to eliminate uncertainties in model predictions that aim to assess the extent and impacts of Arctic atmospheric contaminants.

1 Introduction

Traditionally, Arctic regions are considered to be pristine and remote from the majority of potential large-scale emission sources in industrialized middle latitude countries (Armitage et al., 2011; Macdonal et al., 2000; Barrie et al., 1992). Atmospheric transport is the most efficient way for polycyclic aromatic hydrocarbons (PAHs), released in the lower latitudes, to reach the Arctic (Friedman et al., 2014). Long range atmospheric transport (LRAT) to Arctic regions has strong seasonality with an increased tendency during winter and spring (Willis et al., 2018). This is driven by a different mean circulation direction across the Arctic in winter compared to summer, the extension and significantly increased permeability of the Arctic front in winter, and the absence of wet removal of particles during transport (Willis et al., 2018). These factors explain observed maximum near-surface pollutant concentrations during winter and minimum levels during summer (Klonecki, 2003). Fossil fuel sources dominate total aerosol organic carbon in Arctic winter air, with a predominance of alkanes, PAHs, and phthalates (Fu et al., 2009). During the past decades, the background monitoring of atmospheric pollutants in Ny-Ålesund, Svalbard, and Alert, Canada, have been an important data repository for information on occurrence and LRAT of anthropogenic contaminants including persistent organic pollutants and PAHs in the Arctic regions. The data demonstrates ubiquitous distribution of PAHs on a global scale, including the Arctic. “Confirmed occurrence of a

pollutant in a polar environment” is an important criterion considered by conventions, including the United Nations Economic Commission for Europe (UNECE), Stockholm, Basel, and Rotterdam (Fiedler et al., 2019). UNECE has incorporated PAHs in the Convention on Long-range Transboundary Air Pollution (UNECE, 1998). ~~PAHs are regulated in many countries, eg. US, Canada, Holland, Sweden, Switzerland, and Denmark (Bandowe and Meusel, 2017).~~ Atmospheric PAHs are regulated in USA, Canada, UK, and Europe (EU Directive 2004/107/EC, 2005; US EPA, 2011; UK Air DEFRA, 2007; Ontario Ministry of the Environment and Climate Change, 2016). PAHs are also included in the list of target chemicals of the Convention for the Protection of the Marine Environment of the North-East Atlantic (OSPAR). PAH concentrations are not declining in the Arctic despite global emission reductions (Yu et al., 2019), and PAHs are listed as “chemicals of emerging concern in the Arctic” (Balmer and Muir, 2017).

PAHs are byproducts of different incomplete combustion processes, mainly fossil fuels and biomass burning. (Ravindra et al., 2008a). Their toxic and carcinogenic effects on both human health and ecosystems are well documented (Kim et al., 2013; Reynaud and Deschaux, 2006; Macdonald et al., 2010). Under unique Arctic weather conditions, with extreme temperatures, wind, and light seasonality, atmospheric PAHs may behave differently compared to in temperate climatic conditions. Low temperatures favor partitioning of semi-volatile PAHs from gas phase to particulate phase, which makes them more persistent in the Arctic environment (Lammel, 2015). Due to extended winter darkness in the Arctic, photodegradation of PAHs is limited for several months. The transition from dark polar winter to the light spring and summer brings large increase in the amount of available solar radiation and oxidants in the Arctic troposphere (Willis et al., 2018). PAHs react with a number of atmospheric oxidants, most notably the hydroxyl radical, ozone, the nitrate radical, and nitrogen dioxide (Keyte et al., 2013). This leads to their transformation into more toxic oxygenated and nitrated PAH derivatives (oxy-PAHs and nitro-PAHs). Oxy- and nitro-PAHs are also constituents of raw coal and can be emitted with PAHs following the same combustion processes (Huang et al., 2014b). Oxy- and nitro-PAHs have high toxicity (Onduka et al., 2012); they can act as direct mutagens, carcinogens, and oxidative stressors in biota (Durant et al., 1996). The biological effects of nitro- and oxy-PAHs can be greater than those of the parent PAHs (Kielhorn et al., 2003). In remote locations they are found at concentrations near detection limits and thus are mostly not included in monitoring programs, and the level of nitro- and oxy-PAHs in the Arctic atmosphere is unknown (Balmer and Muir, 2017).

The Arctic is warming at a higher rate than the global average and visible changes happen rapidly here. Thus, it is a key area for modeling studies on climate effects on contaminants with a main focus on LRAT from lower latitudes. As a consequence, local Arctic sources are usually disregarded, and lack of information on local emission sources is a source of uncertainty in model predictions that often deviate significantly from observations (Schmale et al., 2018). Local emission sources may be of high importance in winter, when strong temperature atmospheric inversions can be frequent in Arctic region (Bradley et al., 1992). These episodes inhibit the mass and heat fluxes from the surface to the atmosphere, and consequently the dilution of surface emissions (Janhall et al., 2006; Li et al., 2019). This trapping of emissions results in poor air quality and can be potentially harmful to local people. Climate change introduces additional sources of PAHs to the Arctic region. In the past decade, human activities such as resource exploration, research, tourism, fisheries, and maritime traffic have increased substantially due to warming and corresponding reduction of sea ice, opening up new shipping routes (Jörundsdóttir et al., 2014). Warming may also enhance volatilization of low molecular weight (LMW) PAHs from ground surfaces (Friedman et al., 2014) and melting sea ice (Yu et al., 2019). Reactivity of PAHs in the gas phase is significantly greater than when associated with particles (Keyte et al., 2013), therefore increasing air temperatures can be expected to lead to increased levels of toxic nitro- and oxy-PAHs.

The need for a comprehensive assessment of local contaminant sources in Svalbard was acknowledged and initiated in the international Arctic Monitoring and Assessment Programme (AMAP); with the major focus on persistent organic pollutants (Pedersen et al., 2011), there is a scarcity of data on local sources of PAHs in Svalbard. A back-trajectory analysis of twenty

years data for three representative PAHs (Phe, Pyr, and BaPyr; see Table 1 for full names) suggested that Svalbard is impacted by air masses coming from eastern Russia, northern Europe, and northwest Russia during winter (Yu et al., 2019). Overall, combined European and Russian emissions accounted for more than 80% of episodic high-concentration events in Svalbard in 2007 (Balmer and Muir, 2017; Friedman and Selin, 2012). However, observed concentrations of Phe and Pyr from the Zeppelin station, Svalbard, were higher than model simulations, indicating important contributions of local sources of PAHs to the Arctic atmosphere, too (Yu et al. (2019). This study focused on the main settlement in Svalbard, Longyearbyen, with a population of approximately 2300 inhabitants and a high level of (partially seasonal) human activities (transport, coal mining, industry, tourism, and research). The local coal-fired power plant (PP) was hypothesized to be the major local source of PAHs, and the overall objectives of this study were to: (1) evaluate PAH emissions from the local power plant, (2) examine concentrations and profile changes with distance from the PP, (3) quantify concentrations of PAHs and nitro- and oxy-PAHs, in both gaseous and particulate phases, and (4) determine other potential local sources of PAHs, and nitro- and oxy-PAHs.

2 Material and methods

2.1 Sampling site

Svalbard is an archipelago located between latitudes 77° and 81°N in the Western Barents Sea. Longyearbyen, being the largest populated settlement, was chosen as the study area. The local PP was installed in Longyearbyen in the 1980s and provides the community with sufficient electricity (45 000 MW) and central heating supply (70 000 MW) throughout the year (Bøckman, 2019). The PP is fueled by coal produced in a nearby mine at Breinosa (mine No. 7). This coal has a distinct quality (brown, high volatile bituminous coal with vitrinite reflectance $R_o=0.78\%$ (Marshall et al., 2015)) and is well suited for energy production. Coal consumption is about 25-30 thousand ton per year. The PP has two boilers, 32 MW each. The coal burning temperature is about 1000 °C (Bøckman, 2019). Since December 2015, the flue gas purification system consists of a selective non-catalytic reduction (SNCR) system, an electrostatic precipitator (ESP), and a wet flue-gas desulfurization (WFGD) scrubber. After SNCR the NO_x content in the flue gas is reduced by 50% by spraying urea solution as a reduction agent into the boiler. Further, in the ESP step, dust is electrically charged and deflected toward the collection electrodes. In the WFGD scrubber, the flue gas is cooled and desulfurized by sea water. Low emissions are reported: dust 1.5 ± 0.2 , SO_2 0.3 ± 0.1 , NO_x 244 ± 19 , CO 63 ± 5 mg Nm⁻³ (Lundgjerdinen, 2017), reflecting high efficiency of the flue gas cleaning system. For PP emission analysis, stack emission air samples were collected at source (PP), and two locations at transect distance: the roof of the University Centre in Svalbard (UNIS, urban location, 1 km from PP) and the former northern lights observatory in Adventdalen (Adventdalen, rural location, 6 km from PP, 7 km to the active coal mine No. 7) (Fig. 1). Sampling at UNIS and Adventdalen was performed simultaneously.

2.2 Sample collection

2.2.1 Power plant

A total of 6 low volume (1.3-3.0 m³) samples of the PP stack emission were collected (Table S1) under normal operating conditions, collected on 27th September (PP1-PP3) and 2nd October (PP4-PP6) 2018. Sampling was performed downstream the WFGD scrubber, after all flue gas cleaning steps. The sampling probe (inner Ø = 11 mm) was situated to face the direction of the flue gas. A custom-made low volume, battery powered, air sampler (Digitel, Switzerland) was used to pump the flue gas through the sampling material placed in a stainless steel cartridge (16249, Sartorius Stedim Biotech GmbH, Germany). The particulate phase was collected on quartz fiber filter (QFF, pre-burned at 450°C for 6 h; Ø = 47 mm, no

binder, Pallflex, USA), and the gaseous phase on polyurethane foam (PUF, Soxhlet pre-cleaned in toluene for 24 h followed by 24 h acetone wash; Ø = 50 mm, L = 75 mm, Klaus Ziemer GmbH, Germany). Although, the pump was operated at the maximum speed (35 L min⁻¹, which corresponds to 6.1 m s⁻¹ probe intake flow speed), an isokinetic sampling regime was not achieved. The flue gas parameters (temperature 8.9±0.5°C, moisture 28±2%, flow speed 18.1±0.8 m s⁻¹, and density 1.24±0.2 kg m⁻³) were measured during the sampling by FKT3DP1A multi meter equipped with S-type Pitote probe (FlowKinetics LLC, USA).

2.2.2 UNIS and Adventdalen

The prevailing wind direction in Longyearbyen and Adventdalen is from the southeast. In summer, when the soil surface in Adventdalen becomes warmer than water surface in Adventfjorden, the wind direction can temporary change to northwesterly. To focus on PP emission and avoid the peak (Dekhtyareva et al., 2016; Esau et al., 2012). To focus on PP emission and avoid the peak of marine traffic in the summer, simultaneous sampling at UNIS and Adventdalen was carried out from 28th August to 28th of September 2018 (Table S1), on days with predicted northwesterly wind direction, (Fig. S1), using high volume air samplers (TISCH-1000-BLXZ, TISCH Environmental Inc., USA) equipped with dual chamber sampling module (particle filter, stainless screen and vapor filter, glass cartridge). About 370 m³ of ambient air was collected over 24 h per sample (Table S1). For each station, 6 high volume air samples were collected for particulate (QFF, pre-burned at 450°C for 6 h; Ø = 103 mm, no binder, Munktell/Ahlstrom, Finland) and gaseous (PUF, Soxhlet pre-cleaned in toluene for 24 h followed by 24 h acetone wash; Ø = 65 mm, L = 100 mm, Klaus Ziemer GmbH, Germany) phases. Weather parameters including ambient temperature, relative humidity, UV radiation, wind direction, and precipitation were recorded (Table S2). All samples (PP, UNIS, and Adventdalen) were kept intact inside the sampling unit after collection. In order to reduce the risk of post-collection contamination, the unit was sealed in two plastic bags for transportation to the lab, where samples were removed from the unit, sealed with layers of aluminum foil, and stored airtight in two plastic bags. Samples were kept frozen at -20°C until analyzed. A total of 18 samples (18 QFFs and 18 PUFs) and 8 field blanks (4 for PP and 4 for UNIS and Adventdalen) were collected.

2.3 Analytical procedure

16 PAHs, 8 oxy- and 21 nitro-PAHs (Table S3) were quantified using gas chromatography in combination with electron ionization mass spectrometry GC-EI-MS/MS and gas chromatography in combination with electron capture negative ion mass spectrometry GC-ECNI-MS, respectively. Full details on analytical methods, including equipment and procedures, are outlined in the SI (Text S1, Tables S4 and S5). In brief, all QFFs (particulate phase) and PUFs (gaseous phase) samples were extracted separately by two different methods, followed by the same clean-up procedure. Several ²H-labelled PAH (dPAH) surrogates (16 dPAHs, 3 dOxy-PAHs, and 6 dNitro-PAHs), were added to samples prior to extraction. QFF samples were extracted with dichloromethane using a quick easy cheap effective rugged and safe QuEChERS-like procedure developed previously for the analysis of particulate bound PAHs (Albinet et al., 2013; Albinet et al., 2014). PUF samples were Soxhlet extracted with dichloromethane for 24 h. The extracts were concentrated and cleaned up first with neutral alumina Al₂O₃, and then with neutral silica SiO₂. Elutes were dried under gentle nitrogen stream and redissolved in approximately 100 µL *n*-hexane. The purified samples were spiked with three labelled standards to evaluate the surrogate recoveries.

2.4 Quality assurance

Detailed information on method validation and quality control is provided in SI (Text S2). Field (n=4 for PP, n=4 for UNIS and Adventdalen combined) and laboratory (n=3 for PP, n=3 for UNIS and Adventdalen combined) blanks were analysed in order to evaluate possible contamination during sample transport and analysis. The method detection limit (MDL) was

determined based on blank values for each sampling material type (Table S6). High contamination of PUF blank samples by 9,10-PheQ (for UNIS and Adventdalen), and 2-NFlu (for PP) was found; these compounds were excluded from the final results. No blank correction was performed for the concentration calculations. Samples with PAH concentrations below instrumental limit of quantification (LOQ) were replaced by LOQ/2 for statistical analysis. The method efficiency was tested using QFF (n=4) and PUF (n=4) spiked samples (Table S7). Acceptable recoveries ranged between 63-109% for dPAHs, 56-68% for dOxy-PAHs, and 44-89% for dNitro-PAHs (Table S8).

2.5 Statistical analysis

Statistical analyses of compound concentrations were performed with Minitab 18 Statistical Software (Minitab LLC, Pennsylvania, USA). Normality and homogeneity of variances were tested with Shapiro-Wilk and Levene's tests, respectively. Mann-Whitney U test was performed to test significant differences between sampling locations ~~(UNIS and Adventdalen)~~. Spearman's correlation was used to investigate relationships between different variables. The statistical significance was set at $p < 0.05$, unless stated.

Principle component analysis (PCA) was performed for PAH source apportionment. A 6×29 matrix (sample number \times 29 detected compounds, including 14 PAHs and 15 nitro- and oxy-PAHs) dataset was used to assess the source contribution to PAHs for each location, and 12×29 matrix dataset to investigate PAH profile differences between UNIS and Adventdalen. Total PAH concentrations (gaseous and particulate, G+P), were used to minimize the influence of partitioning, ageing, and photochemical degradation (Kim et al., 2009). PCA was based on a correlation matrix to standardize scales and weight all variables equally (Holmes et al., 2017). PCA was first applied on the concentrations matrix only, and then weather parameters were carefully included in order to explain the observed sample groupings.

3 Results and discussion

3.1 Longyearbyen power plant PAH emission profile

Individual concentrations and phase distribution (percentage on particulate matter, %PM) of target PAH are summarized in Table 1. The sum of total (G+P) concentration of the 16 priority PAHs ($\sum 16$ PAHs; U.S. Environmental Protection Agency) in the purified flue gases emitted from the PP is $0.106 \mu\text{g m}^{-3}$, which corresponds to $1.5 \mu\text{g kg}^{-1}$ coal. Currently, there is no PAH emissions standard for coal-fired power plants in Norway. However, compared to the Canadian emission limits of PAHs for municipal solid waste incinerators of $5 \mu\text{g m}^{-3}$ (Li et al., 2016), the Longyearbyen PP emissions is a factor of 3 lower. About 94% of 16 PAHs were emitted in a gas phase, in agreement with earlier studies (Li et al., 2016; Wang et al., 2015; Yang et al., 1998). The emission profile of the Longyearbyen PP is dominated by LMW PAHs (2 and 3 rings), which represents 89% of $\sum 16$ PAHs emission; high molecular weight (HMW) PAHs (5-7 rings) were not detected, likely due to their low vapor pressure and thus association to particles. A combination of ESP and WFGD has a removal efficiency of PM up to 99.9% (Wang et al., 2019). Fine cooling of the PP flue gas (8.9 ± 0.5 °C) by cold sea water facilitates high PM collection efficiency as well (Noda and Makino, 2010; Wang et al., 2019). As a result, PP dust emissions are below the ultra-low standard of 5 mg m^{-3} (Zhao et al., 2017) at $1.5 \pm 0.2 \text{ mg m}^{-3}$ (Lundgjerdingen, 2017). The PAH emissions profile was dominated by Nap and Phe, accounting for 53 % and 27 % of $\sum 16$ PAHs, followed by Flu, Flt and Pyr. Nap and Phe are often reported as major emitted compounds from power plants equipped with analogous exhaust cleaning systems and/or burning the same type of coal (Hsu et al., 2016; Li et al., 2016; Wang et al., 2015). A similar PAH emissions profile was reported by Hsu et al. (2016) for the power plant in central Taiwan (Table S12). A higher flue gas dust concentration and different coal sources resulted in 40% emissions of four ringed PAHs compared to 11% for Longyearbyen PP. Operation

conditions and boiler type can have significant effects on emitted PAH profiles and concentrations (Wang et al., 2015), as well as combustion temperature (Peng et al., 2016), and geological maturity (Huang et al., 2014b).

Nitro- and oxy-PAHs are constituents of raw coal and can also be produced from parent PAH compounds during high temperature coal combustion (Huang et al., 2014b). The yields of individual nitro-PAHs from the PP was 1 - 2 orders of magnitude lower than those of their corresponding parent PAHs, and individual concentrations were at or below 1.7 ng m^{-3} ; 1-NNap was the most abundant nitro-PAH. Huang et al. (2014b) investigated the same type of coal (bituminous, $R_0 = 0.77 \%$), burned at lower temperatures in a honeycomb briquette stove; nitro-PAHs were absent in the raw coal and calculated nitro-PAH/PAH ratios were >1 confirming formation of nitro-PAH compounds during coal combustion. In contrast, in the present work, the same daughter to parent PAH ratios were < 1 (Table 2), indicating an absence of nitro-PAH formation during coal combustion or possible thermal degradation of nitro-PAH at 1000°C .

The yields of oxy-PAHs were orders of magnitude higher than nitro-PAHs because oxy-PAHs can be produced by reaction of PAH with O^\bullet or OH^\bullet radicals generated continuously by radical chain reactions during combustion (Huang et al., 2014b). 9-Flu and 9,10-AntQ were the most abundant among the oxy-PAHs (12.4 and 15.6 ng m^{-3} , respectively), and concentration of 9,10-PheQ was a factor of six lower. The calculated ratios of oxy-PAH to corresponding parent PAH were lowest for 9,10-PheQ/Phe and highest for 9,10-AntQ/Ant (Table 2). This can be due to a higher content of Phe in coal, as well as different reaction rates of Phe, Ant, and Flu with O^\bullet or OH^\bullet radicals. Difference between the reaction rates of Flu and Ant can possibly be explained by different reaction pathways; Flu undergoes H atom abstraction at the 9-position to form 9-Flu, while Ant requires OH^\bullet attack on the aromatic ring (Brubaker and Hites, 1998). Ant and Phe have essentially the same 3-ring structure, only differing by the relative position of their aromatic rings. However, Ant appears to be significantly more reactive, due to the sterically unhindered molecular structure of Ant (Keyte et al., 2013). Formation of specific PAHs is also a temperature dependent process (Peng et al., 2016).

Ant, BaAnt, and Chry are often used as ~~tracers~~indicators of coal combustions (Zheng et al., 2019; Wu et al., 2014; Wang et al., 2009), however, their concentrations in the flue gas of the Longyearbyen PP were negligible. This demonstrates the strong importance of determining indicatory PAH profiles for individual combustion sources for correct source identification. PAH emissions from different coal plants are hard to compare because they are affected by many factors including coal type, boiler load, combustion mode (Wang et al., 2015), and flue gas cleaning systems. Nap was the most abundant PAH emitted from the Longyearbyen PP. ~~Nap is not a common tracer due~~Due to its ubiquitous presence ~~and often high blank sampling material contamination; thus~~, Nap was not considered as ~~potentialsuitable~~ marker. Further, Phe, Flu, Flt, Pyr, 9-Flu, and 9,10-AntQ were the main PAHs and oxy-PAHs detected in the Longyearbyen PP flue gas (Fig. 2), therefore the presence and diagnostic ratios (Table 3) of these compounds were used as markers of the PP source in the present work. In Yu et al. (2019), coal combustion was identified as the main source (68% contribution) of PAHs, at the Zeppelin monitoring station at Ny-Ålesund, Svalbard, and Phe, Flu, Flt, and Pyr were the main contributors, most likely attributable to the Longyearbyen PP located 115 km southeast of Ny-Ålesund. Overall, total flue gas emissions were $960\,000 \text{ Nm}^3 \text{ day}^{-1}$ (Lundgjerdingen, 2017), and a daily emissions of $\Sigma 16$ PAHs and sum of the nitro- and oxy-PAHs are approximately 98.7 g of and 35.6 g , respectively.

3.2 UNIS and Adventdalen

3.2.1 Ambient concentrations and PAH profiles

The concentrations of PAHs and oxy-PAHs measured at UNIS ~~was were~~ a factor of 2 higher than at Adventdalen, while ~~the nitro-PAH profiles were similar. $\Sigma 16$ levels differed less (Table 1). $\Sigma 15$ PAHs was were~~ 749.2 ± 72.6 (UNIS) and $369.1 \pm 66.7 \text{ pg m}^{-3}$ at UNIS and (Adventdalen, respectively (Table 1). ΣOxy PAHs were approximately a factor of 2 lower, with average values of-); $\Sigma 8$ Oxy-PAHs were 471.0 ± 150.8 (UNIS) and $233.1 \pm 68.3 \text{ pg m}^{-3}$ (Adventdalen); Σnitro $\Sigma 22$ nitro-PAHs were an

order of magnitude lower than both parent PAHs and oxy-PAHs, with average values of 36.8 ± 6.2 (UNIS) and 27.2 ± 11.1 pg m^{-3} (Adventdalen). The UNIS and Adventdalen chemical profiles of PAHs and oxy-PAHs were similar, while the profiles of nitro-PAHs differ (Fig. S2). Proportions of 1- and 2-NNap were higher in UNIS samples (about 60% of $\Sigma 22$ nitro-PAHs), while 9-NAnt and 2+3-NFlt showed higher contributions into the nitro-PAH profile of samples from Adventdalen (about 55% of $\Sigma 22$ nitro-PAHs).

Among the parent PAHs, Phe (~~ranged ranging~~ from 191.7 to 470.0 pg m^{-3}) and Flu (~~ranged ranging~~ from 38.5 to 236.0 pg m^{-3}) were the most abundant ~~at both sites in the present study~~. The PAH concentrations measured ~~at in~~ Longyearbyen (UNIS and Adventdalen) were ~~two~~2 orders of magnitude higher than those detected at the Zeppelin station and the same order of magnitude as in Birkenes (southern mainland Norway) (NILU, 2019) (Table S13) for the same period (~~autumn 2018~~). (Table S13). The PAH profiles were dominated by Phe and Flu at all sites (Fig. S3). ~~also dominated the PAH profile at Zeppelin station, which may indicate similar sources of contamination. A higher proportion of Phe was observed in Longyearbyen samples. The measured PAH concentrations in the present study were in agreement with the 2 decades average data reported for the Arctic monitoring stations in Svalbard (Zeppelin) and Finland (Pallas), and were about an order of magnitude higher compared to the Canadian Arctic (Alert) concentrations (Yu et al., 2019). The PAH levels observed in Longyearbyen were significantly (up to 2 orders of magnitude) lower compared to rural sites in Europe and China (Table S14).~~

~~Concentrations~~Among measured oxy-PAHs, concentrations of 9-Flu and 9,10-AntQ were the highest ~~among measured oxy-PAHs~~ in the present study. The 9-Flu level (270.3 ± 146.9 pg m^{-3} at UNIS and 139.4 ± 24.9 pg m^{-3} in Adventdalen) was the same order of magnitude as reported for ~~the Pallas and Råö (southern Sweden) background monitoring stations in the north of Finland (Pallas) and in the south of Sweden (Råö)~~ (Brorström-Lundén et al., 2010), while 9,10-AntQ (163.5 ± 57.4 pg m^{-3} at UNIS and 71.7 ± 39.2 pg m^{-3} in Adventdalen) was an order of magnitude higher in Longyearbyen. ~~The nitro-PAH levels in our study were overall lower than other reported background sites. 1-NNap and 2+3-NFlt were the most abundant nitro-PAHs detected at UNIS and Adventdalen air samples; the, and was equal to the winter levels in central European background air (Lammel et al., 2020). The sum of oxy-PAHs detected at UNIS were similar to a rural site in eastern England (Alam et al., 2014), and a factor of 2 higher than at a rural location in the Czech Republic (Lammel et al., 2020), but were significantly lower than in rural southern China (Huang et al., 2014a) and the French Alps (Albinet et al., 2008).~~

~~1-NNap and 2+3-NFlt were the most abundant nitro-PAHs detected at UNIS and Adventdalen. The level of 2+3-NFlt (9.5 ± 1.6 pg m^{-3} at UNIS and 12.3 ± 7.7 pg m^{-3} in Adventdalen) was an order of magnitude higher than that at Råö and Pallas stations; 1-NNap average detected concentrations were 17.0 ± 3.0 pg m^{-3} at UNIS and 5.0 ± 3.2 pg m^{-3} in Adventdalen. (Brorström-Lundén et al., 2010), and 1-NNap average concentrations were 17.0 ± 3.0 pg m^{-3} at UNIS and 5.0 ± 3.2 pg m^{-3} in Adventdalen. Overall, nitro-PAH concentrations were similar to those reported for the Pallas and Råö Scandinavian stations (Brorström-Lundén et al., 2010), and the rural site in the Czech Republic which is representative of central European background levels (Lammel et al., 2020).~~

3.2.2 Gas/particle partitioning

Gas/particle partitioning is an important process that controls transport, degradation, and distribution patterns of contaminants in and between environmental compartments (Huang et al., 2014a). (Finlayson-Pitts and Pitts Jr, 1999; Lammel et al., 2009; Franklin, 2000). The sampling campaign in the present study was conducted from late Arctic summer until early autumn and during this period the air temperature varied from ± 6.8 °C in August to -4.4 °C in September and several precipitation events (snow and rain) occurred. In general, LMW PAHs were found in the gas phase, while HMW PAHs were present in the particulate phase (Table 1), which is in accordance to their physico-chemical parameters, such as octanol-air partition coefficient (Table S3), vapor pressure, and molecular weight (Table S3; Tomaz et al., 2016; Shahpoury, 2016). Repartitioning between phases (Fig. 3) mainly impacted semi-volatile compounds with three and four aromatic rings (Flt,

Pyr, BaAnt, Chry; 2-NFlu, 9-Flu, cPPhe-4, 9,10-AntQ, 9-NAnt, and 2+3-NFlt) as a response to changing meteorological conditions (Hu et al., 2019). ~~Strong negative correlations (Spearman coefficient > 0.65) of percentage of PAH determined in particulate phase (%PM) with ambient temperature and specific humidity were determined for Chry, 1-NNap, 2-NNap, cPPhe-4, and 2+3-NFlt, although they were not statistically significant (Table S14). Likely, it was confounded by diurnal variations in the rate of PAH emissions from different local sources. Specific humidity played an essential role in PAH wet scavenging as it is further discussed in Section 4.1.~~ Strong negative correlations of percentage of PAH determined in particulate phase (%PM) with ambient temperature were confirmed for most of these compounds (Table S15). %PM also depends on aerosol surface area, organic matter, and black carbon content (Lohmann and Lammel, 2004).

Compared to Adventdalen, the urban UNIS location ensure a higher level of PAHs emitted from different nearby anthropogenic sources, including the PP. ~~Low~~ Furthermore, low ambient temperature reinforces partitioning of freshly emitted gaseous PAHs to the particulate phase. As a result, %PM at UNIS was higher than in Adventdalen. Deposition (wet and dry) and chemical reactions with atmospheric oxidants are important removal processes of PAH from air (Keyte et al., 2013). On the local scale, within an hour of travel time from PP to Adventdalen, it is not expected that photolytically-initiated transformation of the freshly emitted PAHs has a strong influence on gas phase concentrations and consequently on %PM. Dry deposition rates vary depending on the type of adsorbing particle (mass, size, aerodynamic properties, shape, and chemical composition) and the atmospheric conditions (Weinbruch et al., 2018). ~~The influence of wet deposition was indicated by a significant negative correlation between concentrations of several HMW PAHs (Pyr, Chry, BbkFlt, IPyr, BPer, BaFlu-11, and BaAnt-7,12) and precipitation (Spearman correlation, $p < 0.05$, Table S15), resulting in a lower amount of particle-bound PAHs transported from the town, and thus lower %PM in Adventdalen, and may be a dominant PAH removal process in source areas (Sharma and McBean, 2002).~~

The influence of wet deposition was indicated by a significant negative correlation between amount of precipitation and concentrations of several particle-bound HMW PAHs (Chry, BbkFlt, IPyr, BPer, BaFlu-11, and BaAnt-7,12) as well as semi-volatile Phe, Flt, and Pyr, which are more predominant in gaseous phase (Spearman correlation, $p < 0.05$, Table S16. Effective wet scavenging of Phe, Flt, and Pyr has been suggested (Škrdliková et al., 2011). Furthermore, a strong negative correlation with mass of water vapor in the air (specific humidity) was determined for most of the compounds (Spearman correlation, $p < 0.05$, Table S17). Particle associated HMW compounds are readily scavenged by precipitation, while water solubility and polarity (for nitro- and oxy-PAHs) play an additional role in wet scavenging processes (Shahpoury et al., 2018). The gas phase removal from the atmosphere is due to substance dissolution in water droplets, which enhances the scavenging effect at higher humidity. Higher sensitivity of gas scavenging compared with particle scavenging towards liquid water content was also indicated by Škrdliková et al. (2011).

In general, the obtained %PM were in agreement with the earlier reported (Table S18). Higher %PM of 9,10-AntQ and several nitro-PAHs (1- and 2-NNap, 2-NFlu, 9-NAnt, 9-NPhen) were detected in French alpine sites in winter (Albinet et al., 2008), while higher %PM of Flt, 9-Flu, and cPPhen-4 found in the present study contrasts with those reported for temperate urban and rural sites in China and Europe (Huang et al., 2014a; Tomaz et al., 2016). Sources difference, weather influence such as precipitation and temperature, as well as different atmospheric conditions (e.g., number of suspended particles, mass size particle distribution, and specific humidity), are likely responsible for these variations.

4 Source identification

~~PCA was the main tool applied on Adventdalen (n=6) and UNIS (n=6) samples to determine possible PAH sources at each location, as well as on all the analyzed ambient samples together (n=12) to identify the difference between the two transect locations. Total PAH (G+P) concentrations were used to minimize the influence of partitioning, aging, and photochemical~~

degradation. PAH diagnostic ratios (Table 4) were utilized as an additional supportive tool taking into account their possible shift due to large scale mixing of PAHs in the atmosphere, different emission rates of PAH from the same source, influence of changing environmental conditions (Katsoyiannis and Breivik, 2014), and atmospheric processing (Alam et al., 2013). Transport of PAHs over short distances from a source can be enough to cause a change in interpretation and identification of the source (Katsoyiannis and Breivik, 2014). Different PAHs have diverse reactivity with other atmospheric species and half lives in the atmosphere

Due to changes in the Arctic front, more frequent precipitation, and very low levels of wood and coal burning for residential heating in the northern hemisphere in the summer, the LRAT of PAHs to the Arctic is low in summer. Sampling was performed on days with predicted northwesterly wind, and according to the 5-day back trajectory analysis, the air arriving to Svalbard in the sampling period mainly came from the north and from Greenland (Fig. S4). As discussed in Section 3.2.1, two orders of magnitude lower PAH concentrations were detected at the Zeppelin monitoring station compared to the levels in Longyearbyen on the same time. Thus, local emissions were the main sources of PAHs in Longyearbyen in this study.

Besides the PP emission, vehicles are another obvious local source of PAHs. In 2018, 1558 vehicles, including cars, lorries, and busses were registered in Longyearbyen (Statistics Norway, 2018). Table S19. Longyearbyen maintains about 50 km of paved and unpaved roads dedicated for traffic (Bore R.R., 2012). Sampling was conducted at the end of summer in order to avoid peak emissions from marine traffic and to focus on PP emissions; however, it is likely that some of the 718 registered private boats were active in Adventfjorden and several larger ocean-going vessels were in the port around the sampling period (weeks 34-38 in Fig. S5). Thus, shipping emissions could not be eliminated as a potential source of PAHs. To note, there is no local waste incineration and wood burning.

PCA was applied to samples from Adventdalen (n=6) and UNIS (n=6) to determine potential PAH sources at each location. Total PAH (G+P) concentrations were used to minimize the influence of partitioning, aging, and photochemical degradation. Selected PAH diagnostic ratios (Table 3) and weather parameters were utilized as additional supportive tools for sources interpretation, and their values were used as variables. Diagnostic ratios may be affected by large-scale mixing of PAHs in the atmosphere, differing emission rates of PAH from the same source, influence of changing environmental conditions, and atmospheric processing of individual PAH compounds with different atmospheric lifetimes and reactivities (Alam et al., 2013; Tobiszewski and Namieśnik, 2012; Katsoyiannis and Breivik, 2014). Ratios based on highly reactive compounds such as Ant and BaAnt were not included, while more stable HMW PAHs diagnostic ratios were interpreted with greater confidence (Galarneau, 2008; Alam et al., 2014). Due to high atmospheric reactivity of Ant and BaAnt, utilization as source apportionment should be avoided, while HMW PAH diagnostic ratios may be exploited with greater confidence owing to their increased stability (Alam et al., 2013). Although, based on similarities in characteristic travel distances among pairs of PAHs, Katsoyiannis and Breivik (2014) named BaAnt/(BaAnt+Chry) ratio as the most robust for air concentrations. Conclusions on the chosen diagnostic ratios are incorporated in the PCA findings discussion. Two principal components (PCs) for Adventdalen (79%) and three PCs for UNIS (80%) were studied in details. Yunker et al. (2002) previously proposed the ratio of IPyr/(IPyr+BPer) to recognize vehicle from coal combustion emissions. BbkFlt/BPer was selected as an additional marker ratio for traffic due to the greater capacity to discriminate diesel and gasoline emissions, as well as its wider value range (Kuo et al., 2013). The Flt/(Flt+Pyr) ratio is widely used for source identification and, in particular, to understand if PAHs are mainly emitted from petroleum sources or from combustion processes (Yu et al., 2019). The Flu/(Flu+Pyr) ratio was selected as a specific indicator for coal combustion due to its strong correlation with the local PP determined markers, and the ratio value was also in agreement with literature (Yunker et al., 2002; Katsoyiannis and Breivik, 2014). Two principal components (PCs) for Adventdalen (74%) and two PCs for UNIS (74%) were focused on.

4.1 Adventdalen

Two PCs explain 79% of the total variance (Table S16) of Adventdalen data. PC1 (55%) revealed a distinct profile dependence on ambient conditions (temperature, specific humidity, and UV radiation). Two groups, “humid samples” (A1, A2, A3) and “dry samples” (A4, A5, A7), were identified (Fig. 4). The first group corresponds to the sampling days with higher values of temperature, specific humidity, and UV radiation (Table S2). Most of the target compounds in the “humid samples” were at lower concentrations compared to the “dry samples”. The majority of PAHs were negatively correlated with specific humidity, temperature, UV radiation, and precipitation (Fig. 4). Samples A2 and A5, corresponding to two days with heavy precipitation events (raining at +5 °C and snowing at -3 °C, respectively), were not grouped together but sharply separated by the PC1. This indicates that the mass of water vapor in the air (specific humidity), in contrast to relative humidity, is an essential parameter for removal from the atmosphere. In the present study, many compounds were negatively correlated with humid conditions, particularly the intense rainfall episode (sample A2, Fig. 4). Heavier molecular weight compounds of this group (Chry, 9,10-AntQ, cPPhe-4, 9-NPhe, BaFlu-11, and BaAnt-7,12), which have lower vapor pressure and thus volatility, were full or partly bound to particles, 38–100% PM (Table 1). Therefore, they were readily scavenged by precipitation (rainfall). For those which are lighter and more volatile, water solubility (for Ace, Aey, Flu, Phe, 1- and 2-NNap, 9-Flu, and 9,10-AntQ) and polarity (for nitro- and oxy-PAHs) play an additional role in wet scavenging processes (Shahpoury et al., 2018). The gas-phase removal from the atmosphere is due to substance dissolution in water droplets (Shahpoury et al., 2018), which enhances the scavenging effect at higher humidity. A strong negative correlation with humidity was determined for all quantified LMW PAHs, significant for Aey, Pyr, 1-NNap, and 9-Flu (Spearman correlation, $p < 0.05$; Table S17). Presence of Ant and Flt (gas-phase PAHs with low polarity and water solubility) in the group is likely due to the same source of origin.

Based on the tracers and their loadings (Table S16), PC1 can be assigned to local PP coal burning (Flu, Phe, Flt, Pyr, 9-Flu, 9,10-AntQ) and vehicular emissions (Pyr, Chry, 1-NNap, cPPhe-4, BaFlu-11, BaAnt-7,12). The diagnostic ratios $IPyr/(IPyr+BPer)$ and $IPyr/BPer$ indicated mainly contribution from diesel emissions (Table 4) and several of these compounds (cPPhe-4, 1-NNap, and BaAnt-7,12) were reported to be emitted after diesel burning as well (Røgge et al., 1993; Albinet et al., 2007; Zhao et al., 2018).

Besides, The first and the second PCs described 51% and 23% of the total variance, respectively (Table S20). Three groups of compounds suggest three different potential sources. The first group include Flt, Pyr, cPPhe-4, BaFlu-11, and BaAnt-7,12. Strong correlations between their concentrations and the $IPyr/(IPyr+BPer)$ ratio suggest a traffic origin for these compounds (Yunker et al., 2002), and specifically diesel emissions (Table 3; Table S21; Ravindra et al., 2006; Ravindra et al., 2008a). Because of the rural position, car traffic is much lower at this location. At the same time, due to the proximity to an active mine (Fig. 1), heavy-duty vehicles (coal trucks, tourist busses, geotechnical drilling machinery) are thus the main candidate source for PAH vehicle emissions. Produced coal is regularly delivered from the coal mine to PP and storage area in the harbor on a road situated in 150 m distance from the Adventdalen sampling station. Coal is transported by Volvo FH540 trucks (built in 2018–2020) driven on diesel CFPP-12 (NS-EN 590) (Nilssen P., Store Norske, personal communication; Table S23). The trucks have Euro 6 standard (HC 0.13, CO 1.5, NO_x 0.4, PM 0.01 g (kWh)⁻¹) (DieselNet, 2020) compliant Volvo D13K engines (540 hp, 405 kW) engines fitted with exhaust gas recirculation, diesel particulate filter, diesel oxidation catalyst, selective catalytic reduction, and ammonia oxidation catalyst (Volvo Trucks, 2020). These allow high operation temperatures and high efficiencies in reducing particle and NO_x emissions. Numerous studies showed substantial reduction in gaseous and particulate emissions of PAH, nitro- and oxy-PAHs as the result of such mitigation in particle and NO_x emissions (Hu et al., 2013b; Gerald Liu et al., 2010; Khalek et al., 2015; Huang et al., 2015). Up to 10 orders of magnitude reduction in emission from similar to Volvo D13K heavy-duty engine was reported for several nitro-PAHs (6-NChry, 1-NPyr, 2-NPyr, 4-NPyr, 7-NBaAnt) (Liu et al., 2015; Gerald Liu et al., 2010), which were not detected in the present study

most likely due to low vehicle number in Adventdalen. However, Flt, Phe, and Pyr have been widely reported to be emitted after diesel emissions (Albinet et al., 2007; Ravindra et al., 2008a; Wingfors, 2001) and BaFlu-11, BaAnt-7,12, cPPhen as well (Nyström et al., 2016; Ahmed et al., 2018; Rogge et al., 1993). 9-Flu, 9,10-AntQ, and 1-NNap were the main oxy- and nitro-PAHs emitted from modern technology heavy-duty diesel engine (Liu et al., 2015; Gerald Liu et al., 2010; Guan et al., 2017), supporting the traffic origin of the group 1 compounds too.

9-Flu, 9,10-AntQ, and 1-NNap, together with Phe and Flu (group 2), seems to have double origin. On a PCA loading plot these compounds have similar proximity to the traffic emission ratio (PC1), as well as to the coal combustion ratio (PC2). As earlier reported, Phe, Flu, 9,10-AntQ, 9-Flu, and 1-NNap together accounted for 74% of the total PAH emission from the local PP (Fig. 2) and Mann-Whitney U statistical test (n=6, p<0.05) showed no significant difference between values of the Flu/(Flu+Pyr) coal combustion diagnostic ratio based on the measured Flu and Pyr concentrations in Adventdalen and in the local PP stack emission.

The PC1 emphasized a positive correlation of 2+3-NFlt and 9-NAnt with temperature, humidity, and UV radiation, as well as negative correlation with a (group of 3) with temperature, humidity, and UV radiation, as well as a negative correlation with primary PAHs (Fig. 4, Table S17), suggestive of suggesting a secondary source of origin. The daughter-to-parent PAH ratios, 9-NAnt/Ant and 2+3-NFlt/Flt (Table S18; S24, showed statistically significant correlations with temperature, humidity, and UV radiation (Spearman correlation, p<0.10; Table S19; S25). Moreover, 2+3-NFlt and 9-NAnt had a strong positive correlation with each other and negatively correlated with their parent compounds (Spearman correlation, Table S20; S26, by reason of assumed chemical transformation. It should be noted that 9-NAnt and 2+3-NFlt were detected in the PP flue gas at low levels (0.08 ng/m³ and 0.5 ng/m³, respectively), and further statistical analysis (Spearman correlation, Table S20; S26, Fig. 4) showed no correlation with established PP tracers (Phe, 9,10-AntQ, and 9-Flu), suggesting a different source of origin. These results indicate atmospheric formation as an additional source of 9-NAnt and 2+3-NFlt, in agreement with other studies (Lin et al., 2015b; Hayakawa et al., 2000; Shahpoury et al., 2018). Sampling close to a major source of NO_x emission, such as the local power plant, can result in concentrations of NO₃ and NO₂ at high enough levels for atmospheric transformation of PAHs to occur (Keyte et al., 2013). Relative contribution of primary and secondary sources of nitro-PAHs could be tested/evaluated by applying a 2-NFlt/1-NPyr ratio (Zielinska et al., 1989; Ringuet et al., 2012; Tomaz et al., 2017; Ciccioli et al., 1996), but 1-NPyr was not detected in our study. We conclude that PC1 (55%) is associated with local PP coal burning, atmospheric transformation, and traffic emission (mainly diesel exhaust) sources.

PC2 did not reveal additional PAH sources and confirmed inputs from the PP coal burning (Phe, Flu, and 9-Flu) and diesel exhaust (4-NBip, 1- and 2-NNap, 2+3-NFlt, Hu et al., 2013b; Keyte et al., 2013; Alam et al., 2015). Moreover, PC2 revealed a negative correlation of LMW PAHs, namely 4-NBip, 2-NNap, Phe, Flu, 1-NNap, 9-Flu, and Aey (in loading order) with precipitation, mainly snow on day five (sample A5, Fig. 4). These PAHs were detected in the gas phase. Commonly, wet scavenging of PAHs is considered to be ineffective, unless the substance is particle associated (Škrdlíková et al., 2011). However, studies by Wania et al. (1999) report that snow scavenging may be an important, and sometimes a dominating scavenging process for lighter PAHs, mediated via a process of adsorption to the air-ice (Wania et al., 1999).

The two PCs explain 79.74% of the total variance. Traffic emission (mainly diesel exhaust) and the Longyearbyen coal-burning PP were concluded as the main local sources of PAHs and nitro- and oxy-PAHs in Adventdalen, and atmospheric transformation of PAHs is an additional source of nitro-PAHs. The statistical analysis has also emphasized the importance of weather conditions on the spatial distribution and concentrations of PAHs. LMW PAHs were scavenged by snow, while the level of humidity was an essential parameter for total PAH removal from the atmosphere.

The UNIS sampling location is only 1 km away from the local coal fired PP and the PAH profile is therefore influenced by local activities in and around the town. Three PCs explain 80.3% of the total variance (Table S21). PC1 (41%) was equally loaded with Flt, Pyr, BaAnt, Chry, BbkFlt, BaPyr, IPyr, BPer, 2 NNap, BaAnt 7,12, BaFlu 11, and BZT, which are commonly found in traffic emissions. The diagnostic ratios IPyr/(IPyr+BPer) and IPyr/BPer mainly indicated contribution from gasoline emissions (Table 4) and several of these compounds (BPer, IPyr, and BaFlu 11) were reported to be emitted after gasoline burning as well. Absence of 1 NPyr and 2 NFlu, the principal compounds of diesel exhaust (Albinet et al., 2007), supported gasoline combustion as more dominant source as well. Precipitation data was included in PCA as variables to explain sample groupings. On the PC1 biplot (Fig. 5) samples U1, U2, and U5, collected on days with substantial precipitation (snow and rain; Table S2), are grouped together. All PC1 contributors showed negative correlations with precipitation. For Pyr, Chry, BbkFlt, BPer, and BaAnt 7,12 the negative correlation with precipitation was statistically significant (Spearman correlation, $p < 0.10$; Table S15). The majority of these PAH derivatives were found mainly in the particulate fraction (Table 1), and lighter compounds were less affected by precipitation. This is in agreement with particle scavenging dominant during wet deposition (Škrdliková et al., 2011). Zhang et al. (2015) reported that scavenging of particulate phase PAHs is about 20 times more efficient than scavenging of gas phase PAHs.

PC2 (25%) was highly loaded by wind direction (Fig. 5, Table S21), and samples were split between two groups depending on prevailing wind direction during sampling. The first group was characterized by the presence of 9,10-AntQ, Phe, and Flu, attributable to the PP emission source. Diagnostic ratios of Flu/(Flu+Pyr) and BaAnt/(BaAnt+Chry) (Table 4) also indicated that the PP is a source of PAHs and nitro- and oxy-PAHs at UNIS. 9-Flu, a tracer for the PP, may have other possible sources, including diesel and gasoline vehicle exhaust, coal powder, road dust particles (Keyte et al., 2013), and may be locally produced, transported from longer range, or secondarily formed in the atmosphere (Kojima et al., 2010). 9-Flu is associated with the second sample grouping (U2, U6, U7), collected during conditions of wind blowing from the west and west-southwest (Fig. S1), and they are characterized by the presence of 4-NBip, 1,5-DNNap, 2+3-NFlt, and 2-NFlu, known markers for diesel emissions (Hu et al., 2013b; Keyte et al., 2013; Helmig et al., 1992). This can be explained by coal transportation by trucks, mining work, geotechnical drilling, and boat traffic in the fjord. Therefore, PC2 was attributed to coal burning emissions from the PP and diesel emissions.

The proximity of the UNIS sampling location to central Longyearbyen, as well as to the PP and the port, makes the UNIS location more complex for source identification. UNIS is mainly influenced by passenger car traffic, although heavy-duty vehicles also pass the station. 1114 private cars were registered in Longyearbyen in 2018 (sbb.no), including old and modern (Euro 3-7 emission standard) technology cars, approximately equally balanced between gasoline and diesel fuel. Gasoline 95 (with up to 5% bioethanol) and diesel CFPP-12 (with up to 7% biodiesel NS-EN 14214) are the exclusive fuels used in summer time, and comply with the Norwegian standard NS-EN 228 and NS-EN 590, respectively, with ultra-low sulfur content (< 10 ppm or 0.001%) (Storø J., LNS Spitsbergen, personal communication). Details of the fuel parameters can be found in Table S22-S23.

Muñoz et al. (2018) undertook a study under similar vehicle and fuel conditions to Longyearbyen and reported a predominance of LMW PAHs for both fuels. Flt, Pyr, Phe and BPer, BaPyr, Chry, BbkFlt were found to be the most abundant compounds in gaseous and particulate phases respectively, in agreement with earlier studies (Nyström et al., 2016). Similar PAHs pattern was found for UNIS samples. On the PCA plot (Fig. 5) Flt, Pyr, BPer, BaPyr, Chry, and BbkFlt are grouped together (group 1) and have equally high loadings on the PC1 (Table S27). The concentrations of the compounds are significantly correlated with the traffic ratios BbkFlt/BPer ($p < 0.05$, Table S28) suggesting the same origin of the compounds. The BbkFlt/BPer ratio varied from 0.43 to 0.72 indicating either diesel or gasoline emissions (Kuo et al., 2013). The diesel emission predominance was found for two out of the six sampling days, although particulate phase 1-NPyr, a marker of

diesel emissions, was not detected. 1-NPyr forms in the combustion chamber of diesel engines by the addition of nitrogen oxide or nitrogen dioxide to free Pyr radicals (IARC, 2014). Its generation is facilitated by the high engine temperatures (IARC, 2014; Karavalakis et al., 2010; Guan et al., 2017; Huang et al., 2015), which likely can not be reached in Longyearbyen due to short driving distances and low speed limit. The use of high quality ultra-low sulfur fuel with substantially reduced emissions of NO_x leads to reduced nitration of PAHs during fuel combustion (Heeb et al., 2008; Zhao et al., 2020b), and together with low total vehicle number, resulting in low nitro-PAH emissions. Occurred atmospheric deposition may be of influence too.

Gaseous phase 1-NNap and 2-NNap have large loadings on the PC1. They are often reported in traffic emissions (Alam et al., 2015; Albinet et al., 2007; Keyte et al., 2016), as well as oxy-PAHs such as BaFlu-11, BaAnt-7,12 and BZT (Nyström et al., 2016; Albinet et al., 2007; Ahmed et al., 2018; Karavalakis et al., 2010). All these nitro- and oxy-PAHs have a strong positive correlation with the traffic ratio (Table S28). Thus, we conclude traffic (diesel and gasoline) is the source for Flt, Pyr, Chry, BPer, BaPyr, 1-NNap, 2-NNap, BaFlu-11, BaAnt-7,12, and BZT at the UNIS location.

A second group of compounds (Phe, Flu and 9,10-AntQ) was strongly correlated with the coal combustion ratio Flu/(Flu+Pyr) (Fig. 5, Table S28), supported by their predominance (along with 9-Flu) in the PP emissions (Fig. 2). 9-Flu may have other possible sources, including diesel and gasoline vehicle exhaust, coal powder, road dust particles (Keyte et al., 2013), and may be locally produced, transported from longer range, or secondarily formed in the atmosphere (Kojima et al., 2010). Interestingly, despite heavy rain during sampling, 9-Flu was found in its maximum concentration (about 2-fold higher the average detected level) on the second sampling day, which may suggest a strong local emission on that day in addition to the daily PP emissions. 9-Flu showed a strong positive correlation with Flt/(Flt+Pyr) ratio, indicating petrol or marine fuel sources (Zhang et al., 2019). The ratio did not correlate with the traffic emitted compounds and the traffic ratio (group 1), thus marine fuel emission was considered as a potential source. Despite the intention to collect air samples at the end of summer to avoid the peak marine traffic, 4 large boats (fishing, 2 cruise vessels, and oil tanker; Fig. S5) and some private boats were registered in Longyearbyen harbor during sampling day 2 (Kystdatahuset, 2018). Our assumption is supported by reports of 9-Flu, cPPhe-4, and 9,10-AntQ as major oxy-PAHs in ship emissions (Czech et al., 2017; Zhao et al., 2020a; Zhao et al., 2019). The cruise ship Artania (MMSI: 311000608) was notably the biggest (230 x 32 m, 44656 t gross tonnage) among boats registered in Longyearbyen port on the day 2 sampling period (Marine Traffic, 2020). According to Svalbard environmental law, vessels entering Svalbard coastal waters are required to use distillate marine fuel (DMA ISO 8217:2017) instead of heavy marine oil to satisfy regulations on the fuel sulfur content below 1.0% (Governor of Svalbard, 2020). Ultra-low sulfur diesel CFPP-12 (NS-EN 590, with sulfur content below 0.001%, SI) is also used for private boats. Such predominance of distillate marine diesel explains the strong correlation of 9-Flu with the marine fuel ratio and no correlation with 9,10-AntQ, which is mainly emitted from heavy fuel oil (Huang et al., 2018). Use of high quality fuels decrease the emissions of particles (Anderson et al., 2015). A reduction of up to 94% particulate PAH emission was reported when burning low-sulfur fuel compared to heavy fuel oil (Huang et al., 2018; Gregoris et al., 2016; Kotchenruther, 2017; Czech et al., 2017). This explains the absence of particle-bound PAHs correlating with the marine ratio Flt/(Flt+Pyr).

PC3 (15%) was loaded mainly with 9 NAnt, cPPhe 4, 9 Flu, and 2+3 NFlt (Table S21). These compounds may originate from combustion processes, often from vehicle emissions. However, on the PC3 score and loading plots (Fig. S2) it can be interpreted that these compounds were negatively correlated with other traffic emission tracers such as 1-NNap, 2-NNap, BZT, 2- and 4-NBip, and BaFlu, suggesting different sources of origin. Several studies have shown that 9 NAnt, cPPhe 4, 9-Flu, and 2+3 NFlt can be secondarily formed via chemical reactions in the atmosphere (Singh et al., 2017; Lin et al., 2015a; Zhao et al., 2018). The compounds were also associated with the sample U2, collected on the warmest day with high UV

radiation and humidity levels. Thus, a secondary origin of 9-NAnt and 2+3-NFlt at UNIS was concluded, in agreement with source apportionment in Adventdalen, and PC3 was attributed to atmospheric transformation of PAHs.

The three PCs explain 79.7% of the total variance of the UNIS samples. PP coal burning, gasoline traffic, and diesel marine shipping emissions, and atmospheric transformations were determined as the main sources of PAHs and nitro- and oxy-PAHs at UNIS.

4.3 Transect ambient samples



PCA was performed to compare samples collected at UNIS and Adventdalen locations (n=12; Fig. 6). The same potential sources of PAHs and nitro- and oxy-PAHs for the Longyearbyen-Adventdalen vicinity were confirmed, which explains their similar profiles. Samples from UNIS and Adventdalen were clearly separated due to the higher concentrations of almost all the detected compounds at UNIS, originating from Longyearbyen activities: coal combustion in the PP, vehicle traffic, and marine traffic. LRAT input, including secondary formation of the derivatives, was more evident in Adventdalen where town emissions influence is reduced. In contrast to the study by Yu et al. (2019), no strong indication for LRAT biomass burning emissions was found for this set of air samples.

5 Conclusion

Results provide insights into local sources of atmospheric PAHs and nitro- and oxy-PAHs in Svalbard. Source markers for the coal-burning PP in Longyearbyen were determined, and generally low emissions of PAHs confirmed an efficient exhaust cleaning system. However, PAHs are emitted daily from coal-burning, and due to a large volume of flue gas emissions, the PP remains an important local anthropogenic source of atmospheric contaminants. Ambient air concentrations of PAHs were substantially affected by rain and snow, and specific humidity plays an important role in PAH removal from the atmosphere. Overall, PAH concentrations were the same order of magnitude as detected at other background Scandinavian air sampling stations, and two orders of magnitude higher than in Ny-Ålesund, Svalbard. The gas/particle partitioning of PAHs and nitro- and oxy-PAHs was dependent on air temperature and specific humidity, and mainly impacted semi-volatile compounds with three and four aromatic rings. Traffic emissions were another contributor. Vehicle and marine traffic were other contributors to PAH emissions, with larger input from gasoline-driven cars at UNIS and diesel vehicles at the remote site due to mining and geotechnical work in Adventdalen. The results also revealed secondary atmospheric formation as an additional source of some nitro-PAHs, 2+3-NFlt and 9-NAnt. The present study contributes to understanding fate and distribution of PAHs in the Arctic, and it provides important information on the phase-separated concentrations of PAHs, and nitro- and oxy-PAHs in Arctic air, as well as markers of the Longyearbyen PP emissions. This data can eliminate uncertainties in model predictions that aim to assess the extent and impacts of Arctic atmospheric contaminants. Furthermore, the knowledge on the local emissions level can be important in case of temperature inversion in the lower atmosphere when vertical dilution is limited and contaminants are trapped near the ground, which may be adverse to public health.

Data availability. The dataset used in this paper is included in the Supplement, and further information is available from the corresponding author tatiana.drotikova@unis.no.

Supplement. The supplement related to this article is available online at: ~~xxx~~ <https://www.atmos-chem-phys-discuss.net/acp-2020-142/acp-2020-142-supplement.pdf>.

Author contribution. RK, AKH, and HR designed the campaign. TD conducted the field and lab works. TD with support from AA and RK optimized, validated and performed GC analysis and further quantification. TD processed and interpreted PCA outcome. TD prepared the manuscript with contributions from all co-authors. TD, AA, and AKH prepared the Supplementary materials section.

Competing interests. The authors declare that there is no conflict of interest.

Acknowledgements. We gratefully acknowledge Longyearbyen Lokaltstyre (Longyearbyen Community Council), personally Kim Rune Røkenes (a former leader of Energyverket), for the support on performing the PP exhaust sampling. We also thank Rasmus Bøckman (Lokaltstyre, Energyverket) for providing information on the PP system operating parameters; Morten Hogsnes and Kristin Lundgjerdingen (Applica Test & Certification AS) for sharing their knowledge on PP flue gas sampling; Siiri Wickström (UNIS) for helping with weather prediction; Marcos Porcires (UNIS) for on-site installation of weather station; Øyvind Mikkelsen (Norwegian University of Science and Technology) for teaching on PCA topic; Malte Jochmann (UNIS/Store Norske) for fruitful discussions on Svalbard coal quality. This research was financially supported by UNIS, NMBU, and the Svalbard Environmental Protection Fund (AtmoPart project).

References

- Ahmed, T. M., Bergvall, C., and Westerholm, R.: Emissions of particulate associated oxygenated and native polycyclic aromatic hydrocarbons from vehicles powered by ethanol/gasoline fuel blends, *Fuel*, 214, 381-385, 10.1016/j.fuel.2017.11.059, 2018.
- Alam, M. S., Delgado-Saborit, J. M., Stark, C., and Harrison, R. M.: Using atmospheric measurements of PAH and quinone compounds at roadside and urban background sites to assess sources and reactivity, *Atmos. Environ.*, 77, 24-35, <https://doi.org/10.1016/j.atmosenv.2013.04.068>, 2013.
- Alam, M. S., Delgado-Saborit, J. M., Stark, C., and Harrison, R. M.: Investigating PAH relative reactivity using congener profiles, quinone measurements and back trajectories, *Atmos. Chem. Phys.*, 14, 2467-2477, <https://doi.org/10.5194/acp-14-2467-2014>, 2014.
- Alam, M. S., Keyte, I. J., Yin, J., Stark, C., Jones, A. M., and Harrison, R. M.: Diurnal variability of polycyclic aromatic compound (PAC) concentrations: Relationship with meteorological conditions and inferred sources, *Atmos. Environ.*, 122, 427-438, <https://doi.org/10.1016/j.atmosenv.2015.09.050>, 2015.
- Albinet, A., Leoz-Garziandia, E., Budzinski, H., and Villenave, E.: Polycyclic aromatic hydrocarbons (PAHs), nitrated PAHs and oxygenated PAHs in ambient air of the Marseilles area (South of France): Concentrations and sources, *Sci. Total Environ.*, 384, 280-292, <https://doi.org/10.1016/j.scitotenv.2007.04.028>, 2007.
- Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., and Jaffrezo, J. L.: Nitrated and oxygenated derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleysPart 1: Concentrations, sources and gas/particle partitioning, *Atmos. Environ.*, 42, 43-54, 10.1016/j.atmosenv.2007.10.009, 2008.
- Albinet, A., Tomaz, S., and Lestremau, F.: A really quick easy cheap effective rugged and safe (QuEChERS) extraction procedure for the analysis of particle-bound PAHs in ambient air and emission samples, *Sci. Total Environ.*, 450-451, 31-38, <https://doi.org/10.1016/j.scitotenv.2013.01.068>, 2013.
- Albinet, A., Nalin, F., Tomaz, S., Beaumont, J., and Lestremau, F.: A simple QuEChERS-like extraction approach for molecular chemical characterization of organic aerosols: application to nitrated and oxygenated PAH derivatives (NPAH and OPAH) quantified by GC-NICIMS, *Anal. Bioanal.Chem.*, 406, 3131-3148, <https://doi.org/10.1007/s00216-014-7760-5>, 2014.
- Anderson, M., Salo, K., Hallquist, Å. M., and Fridell, E.: Characterization of particles from a marine engine operating at low loads, *Atmos. Environ.*, 101, 65-71, <https://doi.org/10.1016/j.atmosenv.2014.11.009>, 2015.

- Armitage, J. M., Quinn, C. L., and Wania, F.: Global climate change and contaminants-an overview of opportunities and priorities for modelling the potential implications for long-term human exposure to organic compounds in the Arctic, *J Environ Monit*, 13, 1532-1546, <https://doi.org/10.1039/c1em10131e>, 2011.
- 1585 Balmer, J., and Muir, D.: Polycyclic aromatic hydrocarbons (PAHs), in: AMAP Assessment 2016: Chemicals of emerging Arctic concern, edited by: Hung, H., Letcher, R., and Yu, Y., Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway, 219-238, 2017.
- Bandowe, B. A. M., and Meusel, H.: Nitrated polycyclic aromatic hydrocarbons (nitro-PAHs) in the environment – A review, *Sci. Total Environ.*, 581-582, 237-257, <https://doi.org/10.1016/j.scitotenv.2016.12.115>, 2017.
- 1590 Barrie, L. A., Gregor, D., Hargrave, B., Lake, R., Muir, D., Shearer, R., Tracey, B., and Bidleman, T.: Arctic contaminants: Sources, occurrence and pathways, *Sci Total Environ*, 122, 1-74, [https://doi.org/10.1016/0048-9697\(92\)90245-n](https://doi.org/10.1016/0048-9697(92)90245-n), 1992.
- Bore R.R., A. I., Kristiansen J.E., Modig I.: Dette er Svalbard 2012. Hva tallene forteller (in Norwegian), Oslo, Norway, 24, 2012.
- 1595 Bradley, R. S., Keimig, F. T., and Diaz, H. F.: Climatology of surface-based inversions in the North American Arctic, *J. Geophys. Res.*, 97, 15699, <https://doi.org/10.1029/92JD01451>, 1992.
- Brorström-Lundén, E., Remberger, M., Kaj, L., Hansson, K., Palm Cousins, A., and Andersson, H.: Results from the Swedish national screening programme 2008, IVL Swedish Environmental Research Institute, Göteborg, Sweden, 69, 2010.
- 1600 Brubaker, W. W., and Hites, R. A.: OH reaction kinetics of polycyclic aromatic hydrocarbons and polychlorinated dibenzo-p-dioxins and dibenzofurans, *J. Phys. Chem. A*, 102, 915-921, <https://doi.org/10.1021/jp9721199>, 1998.
- Bøckman, R.: Fremtidens energiutfordringer på Svalbard (in Norwegian), Longyearbyen Lokalstyre, Norway, Open file at www.uit.no, last access: 28 January 2020, 10, 2019.
- 1605 Ciccioli, P., Cecinato, A., Brancaleoni, E., Frattoni, M., Zacchei, P., Miguel, A. H., and De Castro Vasconcellos, P.: Formation and transport of 2-nitrofluoranthene and 2-nitropyrene of photochemical origin in the troposphere, 101, 19567-19581, 10.1029/95jd02118, 1996.
- Czech, H., Stengel, B., Adam, T., Sklorz, M., Streibel, T., and Zimmermann, R.: A chemometric investigation of aromatic emission profiles from a marine engine in comparison with residential wood combustion and road traffic: Implications for source apportionment inside and outside sulphur emission control areas, *Atmos. Environ.*, 167, 212-222, <https://doi.org/10.1016/j.atmosenv.2017.08.022>, 2017.
- 1610 Dekhtyareva, A., Edvardsen, K., Holmén, K., Hermansen, O., and Hansson, H. C.: Influence of local and regional air pollution on atmospheric measurements in Ny-Ålesund, *International Journal of Sustainable Development and Planning*, 11, 578-587, 10.2495/sdp-v11-n4-578-587, 2016.
- EU emission standards for heavy-duty truck and bus engines: www.dieselnet.com/standards/, access: 05 June, 2020.
- 1615 Durant, J. L., Busby Jr, W. F., Lafleur, A. L., Penman, B. W., and Crespi, C. L.: Human cell mutagenicity of oxygenated, nitrated and unsubstituted polycyclic aromatic hydrocarbons associated with urban aerosols, *Mutation Research/Genetic Toxicology*, 371, 123-157, [http://dx.doi.org/10.1016/S0165-1218\(96\)90103-2](http://dx.doi.org/10.1016/S0165-1218(96)90103-2), 1996.
- 1620 Dvorská, A., Lammel, G., and Klánová, J.: Use of diagnostic ratios for studying source apportionment and reactivity of ambient polycyclic aromatic hydrocarbons over Central Europe, *Atmos. Environ.*, 45, 420-427, <https://doi.org/10.1016/j.atmosenv.2010.09.063>, 2011.
- Esau, I., Argentini, S., Przybylak, R., Repina, I., and Sjöblom, A.: Svalbard meteorology, *Advances in Meteorology*, 2012, 8-164, 2012.
- 1625 EU Directive 2004/107/EC: Directive 2004/107/EC of the European Parliament and the Council of 15 December 2004 relating to arsenic, cadmium, mercury, nickel and polycyclic aromatic hydrocarbons in ambient air. In: Official Journal of the European Union 2005.
- Fiedler, H., Kallenborn, R., Boer, J. d., and Sydnese, L. K.: The Stockholm Convention: A tool for the global regulation of persistent organic pollutants, *Chem. Int.*, 41, 4-11, <https://doi.org/10.1515/ci-2019-0202>, 2019.
- 1630 Finlayson-Pitts, B. J., and Pitts Jr, J. N.: Chemistry of the upper and lower atmosphere: theory, experiments, and applications, Elsevier, 1999.
- Franklin, J. A., R.; Howard, P. H.; Orlando, J. J.; Seigneur, C.; Wallington, T. J.; Zetzsch, C.: Quantitative determination of persistence in air, in: Evaluation of Persistence and Long-Range Transport of Chemicals in the Environment, edited by: Klečka, G., et al., SETAC Press, Pensacola, USA, 7-62, 2000.

1635 Friedman, C. L., and Selin, N. E.: Long-range atmospheric transport of polycyclic aromatic hydrocarbons: a global 3-D model analysis including evaluation of Arctic sources, *Environ. Sci. Technol.*, 46, 9501-9510, <https://doi.org/10.1021/es301904d>, 2012.

Friedman, C. L., Zhang, Y., and Selin, N. E.: Climate change and emissions impacts on atmospheric PAH transport to the Arctic, *Environ. Sci. Technol.*, 48, 429-437, <https://doi.org/10.1021/es403098w>, 2014.

1640 Fu, P., Kawamura, K., Chen, J., and Barrie, L. A.: Isoprene, monoterpene, and sesquiterpene oxidation products in the high Arctic aerosols during late winter to early summer, *Environ. Sci. Technol.*, 43, 4022-4028, <https://doi.org/10.1021/es803669a>, 2009.

Galarneau, E.: Source specificity and atmospheric processing of airborne PAHs: Implications for source apportionment, *Atmos. Environ.*, 42, 8139-8149, 10.1016/j.atmosenv.2008.07.025, 2008.

1645 Gerald Liu, Z., Berg, D. R., Vasys, V. N., Dettmann, M. E., Zielinska, B., and Schauer, J. J.: Analysis of C1, C2, and C10 through C33 particle-phase and semi-volatile organic compound emissions from heavy-duty diesel engines, *Atmos. Environ.*, 44, 1108-1115, <https://doi.org/10.1016/j.atmosenv.2009.11.036>, 2010.

Heavy fuel oil ban in the protected areas of Svalbard: <https://www.sysselmannen.no/en/heavy-fuel-oil-ban-in-the-protected-areas/>, access: 05 June 2020, 2020.

1650 Gregoris, E., Barbaro, E., Morabito, E., Toscano, G., Donateo, A., Cesari, D., Contini, D., and Gambaro, A.: Impact of maritime traffic on polycyclic aromatic hydrocarbons, metals and particulate matter in Venice air, *Environmental Science and Pollution Research*, 23, 6951-6959, 10.1007/s11356-015-5811-x, 2016.

Guan, C., Cheung, C. S., Li, X., and Huang, Z.: Effects of oxygenated fuels on the particle-phase compounds emitted from a diesel engine, *Atmospheric Pollution Research*, 8, 209-220, <https://doi.org/10.1016/j.apr.2016.08.005>, 2017.

1655 Hayakawa, K., Murahashi, T., Akutsu, K., Kanda, T., Tang, N., Kakimoto, H., Toriba, A., and Kizu, R.: Comparison of polycyclic aromatic hydrocarbons and nitropolycyclic aromatic hydrocarbons in airborne and automobile exhaust particulates, *Polycyclic Aromat. Compd.*, 20, 179-190, <https://doi.org/10.1080/10406630008034784>, 2000.

1660 Heeb, N. V., Schmid, P., Kohler, M., Gujer, E., Zennegg, M., Wenger, D., Wichser, A., Ulrich, A., Gfeller, U., Honegger, P., Zeyer, K., Emmenegger, L., Petermann, J.-L., Czerwinski, J., Mosimann, T., Kasper, M., and Mayer, A.: Secondary Effects of Catalytic Diesel Particulate Filters: Conversion of PAHs versus Formation of Nitro-PAHs, *Environ. Sci. Technol.*, 42, 3773-3779, 10.1021/es7026949, 2008.

Helmig, D., Arey, J., Atkinson, R., Harger, W. P., and McElroy, P. A.: Products of the OH radical-initiated gas-phase reaction of fluorene in the presence of NO_x, *Atmos. Environ. A-Gen*, 26, 1735-1745, [https://doi.org/10.1016/0960-1686\(92\)90071-r](https://doi.org/10.1016/0960-1686(92)90071-r), 1992.

1665 Holmes, D., Moody, P., Dine, D., and Trueman, L.: Research methods for the biosciences, Oxford University press, Oxford, United Kingdom, 2017.

Hsu, W. T., Liu, M. C., Hung, P. C., Chang, S. H., and Chang, M. B.: PAH emissions from coal combustion and waste incineration, *J. Hazard. Mater.*, 318, 32-40, <http://doi.org/10.1016/j.jhazmat.2016.06.038>, 2016.

1670 Hu, H., Tian, M., Zhang, L., Yang, F., Peng, C., Chen, Y., Shi, G., Yao, X., Jiang, C., and Wang, J.: Sources and gas-particle partitioning of atmospheric parent, oxygenated, and nitrated polycyclic aromatic hydrocarbons in a humid city in southwest China, *Atmos. Environ.*, 206, 1-10, <https://doi.org/10.1016/j.atmosenv.2019.02.041>, 2019.

1675 Hu, S., Herner, J. D., Robertson, W., Kobayashi, R., Chang, M. C. O., Huang, S.-M., Zielinska, B., Kado, N., Collins, J. F., Rieger, P., Huai, T., and Ayala, A.: Emissions of polycyclic aromatic hydrocarbons (PAHs) and nitro-PAHs from heavy-duty diesel vehicles with DPF and SCR, *Journal of the Air & Waste Management Association*, 63, 984-996, 10.1080/10962247.2013.795202, 2013a.

1680 Hu, S., Herner, J. D., Robertson, W., Kobayashi, R., Chang, M. C. O., Huang, S.-M., Zielinska, B., Kado, N., Collins, J. F., Rieger, P., Huai, T., and Ayala, A.: Emissions of polycyclic aromatic hydrocarbons (PAHs) and nitro-PAHs from heavy-duty diesel vehicles with DPF and SCR, *J. Air Waste MA.*, 63, 984-996, <https://doi.org/10.1080/10962247.2013.795202>, 2013b.

Huang, B., Liu, M., Bi, X., Chaemfa, C., Ren, Z., Wang, X., Sheng, G., and Fu, J.: Phase distribution, sources and risk assessment of PAHs, NPAHs and OPAHs in a rural site of Pearl River Delta region, China, *Atmos. Pollut. Res.*, 5, 210-218, <https://doi.org/10.5094/APR.2014.026>, 2014a.

1685 Huang, C., Hu, Q., Wang, H., Qiao, L., Jing, S. a., Wang, H., Zhou, M., Zhu, S., Ma, Y., Lou, S., Li, L., Tao, S., Li, Y., and Lou, D.: Emission factors of particulate and gaseous compounds from a large cargo vessel operated under real-world conditions, *Environ. Pollut.*, 242, 667-674, <https://doi.org/10.1016/j.envpol.2018.07.036>, 2018.

Huang, L., Bohac, S. V., Chernyak, S. M., and Batterman, S. A.: Effects of fuels, engine load and exhaust after-treatment on diesel engine SVOC emissions and development of SVOC profiles for receptor modeling, *Atmos. Environ.*, 102, 228-238, [10.1016/j.atmosenv.2014.11.046](https://doi.org/10.1016/j.atmosenv.2014.11.046), 2015.

Huang, W., Huang, B., Bi, X., Lin, Q., Liu, M., Ren, Z., Zhang, G., Wang, X., Sheng, G., and Fu, J.: Emission of PAHs, NPAHs and OPAHs from residential honeycomb coal briquette combustion, *Energy & Fuels*, 28, 636-642, <https://doi.org/10.1021/ef401901d>, 2014b.

IARC: Diesel and Gasoline Engine Exhausts and Some Nitroarenes. 1-Nitropyrene. , IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, International Agency for Research on Cancer, Lyon, France, 2014.

Janhall, S., Olofson, K., Andersson, P., Pettersson, J., and Hallquist, M.: Evolution of the urban aerosol during winter temperature inversion episodes, *Atmos. Environ.*, 40, 5355-5366, <https://doi.org/10.1016/j.atmosenv.2006.04.051>, 2006.

Jörundsdóttir, H. Ó., Jensen, S., Hylland, K., Holth, T. F., Gunnlaugsdóttir, H., Svavarsson, J., Ólafsdóttir, Á., El-Taliawy, H., Rigét, F., Strand, J., Nyberg, E., Bignert, A., Hoydal, K. S., and Halldórsson, H. P.: Pristine Arctic: Background mapping of PAHs, PAH metabolites and inorganic trace elements in the North-Atlantic Arctic and sub-Arctic coastal environment, *Sci. Total Environ.*, 493, 719-728, <http://dx.doi.org/10.1016/j.scitotenv.2014.06.030>, 2014.

Karavalakis, G., Stournas, S., Ampatzoglou, D., Bakeas, E., and Spanos, A.: Regulated and Unregulated Emissions of a Euro 4 SUV Operated with Diesel and Soy-based Biodiesel Blends, *SAE Int. J. Fuels Lubr.*, 2, 115-131, 2010.

Katsoyiannis, A., and Breivik, K.: Model-based evaluation of the use of polycyclic aromatic hydrocarbons molecular diagnostic ratios as a source identification tool, *Environ. Pollut.*, 184, 488-494, <https://doi.org/10.1016/j.envpol.2013.09.028>, 2014.

Kavouras, I. G., Koutrakis, P., Tsapakis, M., Lagoudaki, E., Stephanou, E. G., Von Baer, D., and Oyola, P.: Source apportionment of urban particulate aliphatic and polynuclear aromatic hydrocarbons (PAHs) using multivariate methods, *Environ. Sci. Technol.*, 35, 2288-2294, <https://doi.org/10.1021/es001540z>, 2001.

Keyte, I. J., Harrison, R. M., and Lammel, G.: Chemical reactivity and long-range transport potential of polycyclic aromatic hydrocarbons – a review, *Chem. Soc. Rev.*, 42, 9333-9391, <https://doi.org/10.1039/C3CS60147A>, 2013.

Keyte, I. J., Albinet, A., and Harrison, R. M.: On-road traffic emissions of polycyclic aromatic hydrocarbons and their oxy- and nitro- derivative compounds measured in road tunnel environments, 566-567, 1131-1142, [10.1016/j.scitotenv.2016.05.152](https://doi.org/10.1016/j.scitotenv.2016.05.152), 2016.

Khalek, I. A., Blanks, M. G., Merritt, P. M., and Zielinska, B.: Regulated and unregulated emissions from modern 2010 emissions-compliant heavy-duty on-highway diesel engines, *Journal of the Air & Waste Management Association*, 65, 987-1001, [10.1080/10962247.2015.1051606](https://doi.org/10.1080/10962247.2015.1051606), 2015.

Kielhorn, J., Wahnschaffe, U., and Mangelsdorf, I.: Environmental health criteria 229. Selected nitro- and nitro-oxy-polycyclic aromatic hydrocarbons, World Health Organization, 511 pp., 2003.

Kim, D., Kumfer, B. M., Anastasio, C., Kennedy, I. M., and Young, T. M.: Environmental aging of polycyclic aromatic hydrocarbons on soot and its effect on source identification, *Chemosphere*, 76, 1075-1081, <https://doi.org/10.1016/j.chemosphere.2009.04.031>, 2009.

Kim, K.-H., Jahan, S. A., Kabir, E., and Brown, R. J. C.: A review of airborne polycyclic aromatic hydrocarbons (PAHs) and their human health effects, *Environ. Int.*, 60, 71-80, <https://doi.org/10.1016/j.envint.2013.07.019>, 2013.

Klonecki, A.: Seasonal changes in the transport of pollutants into the Arctic troposphere-model study, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2002jd002199>, 2003.

Kojima, Y., Inazu, K., Hisamatsu, Y., Okochi, H., Baba, T., and Nagoya, T.: Influence of secondary formation on atmospheric occurrences of oxygenated polycyclic aromatic hydrocarbons in airborne particles, *Atmos. Environ.*, 44, 2873-2880, <https://doi.org/10.1016/j.atmosenv.2010.04.048>, 2010.

Kotchenruther, R. A.: The effects of marine vessel fuel sulfur regulations on ambient PM2.5 at coastal and near coastal monitoring sites in the U.S., *Atmos. Environ.*, 151, 52-61, <https://doi.org/10.1016/j.atmosenv.2016.12.012>, 2017.

Kuo, C.-Y., Chien, P.-S., Kuo, W.-C., Wei, C.-T., and Rau, J.-Y.: Comparison of polycyclic aromatic hydrocarbon emissions on gasoline- and diesel-dominated routes, 185, 5749-5761, [10.1007/s10661-012-2981-6](https://doi.org/10.1007/s10661-012-2981-6), 2013.

Longyearbyen port traffic <https://kystdatahuset.no/>, access: 05 June 2020, 2018.

Lammel, G., Sehili, A. M., Bond, T. C., Feichter, J., and Grassl, H.: Gas/particle partitioning and global distribution of polycyclic aromatic hydrocarbons – A modelling approach, 76, 98-106, [10.1016/j.chemosphere.2009.02.017](https://doi.org/10.1016/j.chemosphere.2009.02.017), 2009.

- Lammel, G.: Long-range atmospheric transport of polycyclic aromatic hydrocarbons is worldwide problem - results from measurements at remote sites and modelling, *Acta Chimica Slovenica*, 729-735, <https://doi.org/10.17344/acsi.2015.1387>, 2015.
- 1745 Lammel, G., Kitanovski, Z., Kukucka, P., Novak, J., Arangio, A. M., Codling, G. P., Filippi, A., Hovorka, J., Kuta, J., Leoni, C., Pribylova, P., Prokes, R., Sanka, O., Shahpoury, P., Tong, H. J., and Wietzoreck, M.: Oxygenated and Nitrated Polycyclic Aromatic Hydrocarbons in Ambient Air-Levels, Phase Partitioning, Mass Size Distributions, and Inhalation Bioaccessibility, *Environmental Science & Technology*, 54, 2615-2625, 10.1021/acs.est.9b06820, 2020.
- 1750 Li, J., Li, X., Li, M., Lu, S., Yan, J., Xie, W., Liu, C., and Qi, Z.: Influence of air pollution control devices on the polycyclic aromatic hydrocarbon distribution in flue gas from an ultralow-emission coal-fired power plant, *Energy & Fuels*, 30, 9572-9579, <http://doi.org/10.1021/acs.energyfuels.6b01381>, 2016.
- Li, J., Chen, H., Li, Z., Wang, P., Fan, X., He, W., and Zhang, J.: Analysis of low-level temperature inversions and their effects on aerosols in the lower atmosphere, *Adv. Atmos. Sci.*, 36, 1235-1250, <https://doi.org/10.1007/s00376-019-9018-9>, 2019.
- 1755 Li, X., Li, J., Wu, D., Lu, S., Zhou, C., Qi, Z., Li, M., and Yan, J.: Removal effect of the low-low temperature electrostatic precipitator on polycyclic aromatic hydrocarbons, *Chemosphere*, 211, 44-49, <https://doi.org/10.1016/j.chemosphere.2018.07.080>, 2018.
- Lin, Y., Ma, Y., Qiu, X., Li, R., Fang, Y., Wang, J., Zhu, Y., and Hu, D. J. J. o. G. R. A.: Sources, transformation, and health implications of PAHs and their nitrated, hydroxylated, and oxygenated derivatives in PM_{2.5} in Beijing, *J. Geophys. Res.*, 120, 7219-7228, <https://doi.org/10.1002/2015JD023628>, 2015a.
- 1760 Lin, Y., Qiu, X., Ma, Y., Ma, J., Zheng, M., and Shao, M.: Concentrations and spatial distribution of polycyclic aromatic hydrocarbons (PAHs) and nitrated PAHs (NPAHs) in the atmosphere of North China, and the transformation from PAHs to NPAHs, *Environ. Pollut.*, 196, 164-170, <https://doi.org/10.1016/j.envpol.2014.10.005>, 2015b.
- 1765 Liu, S.: Distribution of PAHs and trace elements in coal fly ash collected from a 5-stage electrostatic precipitator, *J. Electrostat.*, Volume 96, December 2018, Pages 144-150, 10.1016/j.elstat.2018.10.012, 2018.
- Liu, Z. G., Wall, J. C., Ottinger, N. A., and McGuffin, D.: Mitigation of PAH and Nitro-PAH Emissions from Nonroad Diesel Engines, *Environ. Sci. Technol.*, 49, 3662-3671, 10.1021/es505434r, 2015.
- 1770 Lohmann, R., and Lammel, G.: Adsorptive and Absorptive Contributions to the Gas-Particle Partitioning of Polycyclic Aromatic Hydrocarbons: State of Knowledge and Recommended Parametrization for Modeling, *Environ. Sci. Technol.*, 38, 3793-3803, 10.1021/es035337q, 2004.
- Lundgjerdingen, K. S.: Teknisk rapport Longyearbyen Energiverket (in Norwegian), Applica Test & Certification AS, Longyearbyen, Norway, Available by request from Longyearbyen Lokaltstyre, 23, 2017.
- 1775 Macdonal, R. W., Barrie, L. A., Bidleman, T. F., Diamond, M. L., Gregor, D. J., Semkin, R. G., Strachan, W. M., Li, Y. F., Wania, F., Alae, M., Alexeeva, L. B., Backus, S. M., Bailey, R., Bewers, J. M., Gobeil, C., Halsall, C. J., Harner, T., Hoff, J. T., Jantunen, L. M., Lockhart, W. L., Mackay, D., Muir, D. C., Pudykiewicz, J., Reimer, K. J., Smith, J. N., and Stern, G. A.: Contaminants in the Canadian Arctic: 5 years of progress in understanding sources, occurrence and pathways, *Sci. Total Environ.*, 254, 93-234, [https://doi.org/10.1016/S0048-9697\(00\)00434-4](https://doi.org/10.1016/S0048-9697(00)00434-4), 2000.
- 1780 Macdonald, C., Lockhart, L., and Gilman, A.: Effects of oil and gas activity on the environment and human health, in: Assessment 2007: Oil and gas activities in the Arctic - effect and potential effects, Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway, 5_1-5_164, 2010.
- Vessel information:
<https://www.marinetraffic.com/en/ais/details/ships/shipid:371216/mmsi:311000608/imo:8201480/vessel:ARTANIA>, access: 05 June 2020, 2020.
- 1785 Marshall, C., Uguna, J., Large, D. J., Meredith, W., Jochmann, M., Friis, B., Vane, C., Spiro, B. F., Snape, C. E., and Orheim, A.: Geochemistry and petrology of palaeocene coals from Spitzbergen — Part 2: Maturity variations and implications for local and regional burial models, *Int. J. Coal Geol.*, 143, 1-10, <https://doi.org/10.1016/j.coal.2015.03.013>, 2015.
- 1790 Muñoz, M., Haag, R., Honegger, P., Zeyer, K., Mohn, J., Comte, P., Czerwinski, J., and Heeb, N. V.: Co-formation and co-release of genotoxic PAHs, alkyl-PAHs and soot nanoparticles from gasoline direct injection vehicles, *Atmos. Environ.*, 178, 242-254, <https://doi.org/10.1016/j.atmosenv.2018.01.050>, 2018.
- NILU: Observation data of atmospheric PAHs at Zeppelin and Birkenes stations. Norwegian Institute for Air Research (Ed.), 2019.

1795 Noda, N., and Makino, H.: Influence of operating temperature on performance of electrostatic precipitator for pulverized coal combustion boiler, *Adv. Powder Technol.*, 21, 495-499, <https://doi.org/10.1016/j.appt.2010.04.012>, 2010.

Nyström, R., Sadiktsis, I., Ahmed, T. M., Westerholm, R., Koegler, J. H., Blomberg, A., Sandström, T., and Boman, C.: Physical and chemical properties of RME biodiesel exhaust particles without engine modifications, *Fuel*, 186, 261-269, [10.1016/j.fuel.2016.08.062](https://doi.org/10.1016/j.fuel.2016.08.062), 2016.

1800 Onduka, T., Ojima, D., Kakuno, A., Ito, K., Koyama, J., and Fujii, K.: Nitrated polycyclic aromatic hydrocarbons in the marine environment: acute toxicities for organisms at three trophic levels, *Jpn. J. Environ. Toxicol.*, 15, 1-10, <https://doi.org/10.11403/jset.15.1>, 2012.

Ontario Ministry of the Environment and Climate Change: Ontario Ambient Air Quality Criteria. <https://www.ontario.ca/page/ontarios-ambient-air-quality-criteria-sorted-contaminant-name>, 2016.

1805 Pedersen, H., Kallenborn, R., Ottesen, R., Gabrielsen, G., Schrum, C., Evenset, A., Ruus, A., Benjaminsen, H., Sagerup, K., and Christensen, G.: PCBs on Svalbard, Norwegian Environment Agency and Governor of Svalbard, Longyearbyen, Norway, 99 pp., 2011.

Peng, N., Li, Y., Liu, Z., Liu, T., and Gai, C.: Emission, distribution and toxicity of polycyclic aromatic hydrocarbons (PAHs) during municipal solid waste (MSW) and coal co-combustion, *Sci. Total Environ.*, 565, 1201-1207, <https://doi.org/10.1016/j.scitotenv.2016.05.188>, 2016.

1810 Ravindra, K., Bencs, L., Wauters, E., de Hoog, J., Deutsch, F., Roekens, E., Bleux, N., Berghmans, P., and Van Grieken, R.: Seasonal and site-specific variation in vapour and aerosol phase PAHs over Flanders (Belgium) and their relation with anthropogenic activities, *Atmos. Environ.*, 40, 771-785, <https://doi.org/10.1016/j.atmosenv.2005.10.011>, 2006.

1815 Ravindra, K., Sokhi, R., and Vangrieken, R.: Atmospheric polycyclic aromatic hydrocarbons: Source attribution, emission factors and regulation, *Atmos. Environ.*, 42, 2895-2921, [10.1016/j.atmosenv.2007.12.010](https://doi.org/10.1016/j.atmosenv.2007.12.010), 2008a.

Ravindra, K., Wauters, E., and Van Grieken, R.: Variation in particulate PAHs levels and their relation with the transboundary movement of the air masses, *Sci. Total Environ.*, 396, 100-110, <https://doi.org/10.1016/j.scitotenv.2008.02.018>, 2008b.

1820 Reynaud, S., and Deschaux, P.: The effects of polycyclic aromatic hydrocarbons on the immune system of fish: A review, *Aquat. Toxicol.*, 77, 229-238, [http://10.1016/j.aquatox.2005.10.018](https://doi.org/10.1016/j.aquatox.2005.10.018), 2006.

Ringuet, J., Albinet, A., Leoz-Garziandia, E., Budzinski, H., and Villenave, E.: Diurnal/nocturnal concentrations and sources of particulate-bound PAHs, OPAHs and NPAHs at traffic and suburban sites in the region of Paris (France), *Sci. Total Environ.*, 437, 297-305, [10.1016/j.scitotenv.2012.07.072](https://doi.org/10.1016/j.scitotenv.2012.07.072), 2012.

1825 Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R.: Sources of fine organic aerosol. 2. Nuncatalyst and catalyst-equipped automobiles and heavy-duty diesel trucks, *Environ. Sci. Technol.*, 27, 636-651, <https://doi.org/10.1021/es00041a007>, 1993.

Schmale, J., Arnold, S. R., Law, K. S., Thorp, T., Anenberg, S., Simpson, W. R., Mao, J., and Pratt, K. A.: Local Arctic air pollution: A neglected but serious problem, *Earth's Future*, 6, 1385-1412, <https://doi.org/10.1029/2018ef000952>, 2018.

1830 Shahpoury, P., Kitanovski, Z., Lammel, G. J. A. C., and Physics: Snow scavenging and phase partitioning of nitrated and oxygenated aromatic hydrocarbons in polluted and remote environments in central Europe and the European Arctic, *Atmos. Chem. Phys.*, 18, 13495-13510, <https://doi.org/10.5194/acp-18-13495-2018>, 2018.

1835 Shahpoury, P., Lammel, G., Albinet, A., Sofuoglu, A., Dumanoglu, Y., Sofuoglu, S. C., Wagner, Z. and Zdimas, V.: Evaluation of a conceptual model for gas-particle partitioning of polycyclic aromatic hydrocarbons using poly-parameter linear free energy relationships, *Environ. Sci. Technol.*, doi:10.1021/acs.est.6b02158, 2016.

Sharma, M., and McBean, E. A.: Atmospheric PAH Deposition: Deposition Velocities and Washout Ratios, *J. Environ. Eng.*, 128, 186-195, doi:10.1061/(ASCE)0733-9372(2002)128:2(186), 2002.

1840 Singh, D. K., Kawamura, K., Yanase, A., and Barrie, L. A.: Distributions of polycyclic aromatic hydrocarbons, aromatic ketones, carboxylic acids, and trace metals in Arctic aerosols: Long-range atmospheric transport, photochemical degradation/production at polar sunrise, *Environ. Sci. Technol.*, 51, 8992-9004, <https://doi.org/10.1021/acs.est.7b01644>, 2017.

Škrdlíková, L., Landlová, L., Klánová, J., and Lammel, G.: Wet deposition and scavenging efficiency of gaseous and particulate phase polycyclic aromatic compounds at a central European suburban site, *Atmos. Environ.*, 45, 4305-4312, <https://doi.org/10.1016/j.atmosenv.2011.04.072>, 2011.

1845 Registered vehicles, by region, statistical variable and year: <https://www.ssb.no/statbank/table/11823/>, access: 05 June 2020, 2018.

- Tobiszewski, M., and Namieśnik, J.: PAH diagnostic ratios for the identification of pollution emission sources, *Environ. Pollut.*, 162, 110-119, <https://doi.org/10.1016/j.envpol.2011.10.025>, 2012.
- Tomaz, S., Shahpoury, P., Jaffrezo, J.-L., Lammel, G., Perraudin, E., Villenave, E., and Albinet, A.: One-year study of polycyclic aromatic compounds at an urban site in Grenoble (France): Seasonal variations, gas/particle partitioning and cancer risk estimation, *Sci. Total Environ.*, 565, 1071-1083, 10.1016/j.scitotenv.2016.05.137, 2016.
- Tomaz, S., Jaffrezo, J.-L., Favez, O., Perraudin, E., Villenave, E., and Albinet, A.: Sources and atmospheric chemistry of oxy- and nitro-PAHs in the ambient air of Grenoble (France), *Atmos. Environ.*, 161, 144-154, <https://doi.org/10.1016/j.atmosenv.2017.04.042>, 2017.
- UK Air DEFRA: The Air Quality Strategy for England, Scotland, Wales and Northern Ireland. Department for Environment, Food and Rural A. (Ed.), 2007.
- UNECE: The 1998 Aarhus protocol on persistent organic pollutants (POPs), https://www.unece.org/env/lrtap/pops_h1.html, 49, 1998.
- Polycyclic Aromatic Hydrocarbons on the Gulf Coastline: [https://archive.epa.gov/emergency/bpspill/web/html/pahs.html#:~:text=Polycyclic%20Aromatic%20Hydrocarbons%20\(PAHs\)%20are,commonly%20called%20'weathered%20oil'](https://archive.epa.gov/emergency/bpspill/web/html/pahs.html#:~:text=Polycyclic%20Aromatic%20Hydrocarbons%20(PAHs)%20are,commonly%20called%20'weathered%20oil'), 2011.
- Volvo D13 Engine family: https://www.volvotrucks.us/-/media/vtna/files/shared/powertrain/revised4147_101-volvo_d13_engine-brochure_low-res.pdf, access: 05 June, 2020.
- Wang, G., Ma, Z., Deng, J., Li, Z., Duan, L., Zhang, Q., Hao, J., and Jiang, J.: Characteristics of particulate matter from four coal-fired power plants with low-low temperature electrostatic precipitator in China, *Sci. Total Environ.*, 662, 455-461, <https://doi.org/10.1016/j.scitotenv.2019.01.080>, 2019.
- Wang, K., Shen, Y., Zhang, S., Ye, Y., Shen, Q., Hu, J., and Wang, X.: Application of spatial analysis and multivariate analysis techniques in distribution and source study of polycyclic aromatic hydrocarbons in the topsoil of Beijing, China, *Environ. Geol.*, 56, 1041-1050, <https://doi.org/10.1007/s00254-008-1204-5>, 2009.
- Wang, R., Liu, G., and Zhang, J.: Variations of emission characterization of PAHs emitted from different utility boilers of coal-fired power plants and risk assessment related to atmospheric PAHs, *Science of The Total Environment*, 538, 180-190, 10.1016/j.scitotenv.2015.08.043, 2015.
- Wania, F., Mackay, D., and Hoff, J. T.: The importance of snow scavenging of polychlorinated biphenyl and polycyclic aromatic hydrocarbon vapors, *Environ. Sci. Technol.*, 33, 195-197, <https://doi.org/10.1021/es980806n>, 1999.
- Weinbruch, S., Benker, N., Kandler, K., Schutze, K., Kling, K., Berlinger, B., Thomassen, Y., Drotikova, T., and Kallenborn, R.: Source identification of individual soot agglomerates in Arctic air by transmission electron microscopy, *Atmos. Environ.*, 172, 47-54, <https://doi.org/10.1016/j.atmosenv.2017.10.033>, 2018.
- Willis, M. D., Leaitch, W. R., and Abbatt, J. P. D.: Processes controlling the composition and abundance of Arctic aerosol, *Rev. Geophys.*, <https://doi.org/10.1029/2018rg000602>, 2018.
- Wingfors, H.: Characterisation and determination of profiles of polycyclic aromatic hydrocarbons in a traffic tunnel in Gothenburg, Sweden, 35, 6361-6369, 10.1016/s1352-2310(01)00389-2, 2001.
- Wu, D., Wang, Z., Chen, J., Kong, S., Fu, X., Deng, H., Shao, G., and Wu, G.: Polycyclic aromatic hydrocarbons (PAHs) in atmospheric PM_{2.5} and PM₁₀ at a coal-based industrial city: Implication for PAH control at industrial agglomeration regions, China, *Atmos. Res.*, 149, 217-229, <https://doi.org/10.1016/j.atmosres.2014.06.012>, 2014.
- Yang, H.-H., Lee, W.-J., Chen, S.-J., and Lai, S.-O.: PAH emission from various industrial stacks, *J. Hazard. Mater.*, 60, 159-174, [https://doi.org/10.1016/s0304-3894\(98\)00089-2](https://doi.org/10.1016/s0304-3894(98)00089-2), 1998.
- Yu, Y., Katsoyiannis, A., Bohlin-Nizzetto, P., Brorström-Lundén, E., Ma, J., Zhao, Y., Wu, Z., Tych, W., Mindham, D., Sverko, E., Barresi, E., Dryfhout-Clark, H., Fellin, P., and Hung, H.: Polycyclic aromatic hydrocarbons not declining in Arctic air despite global emission reduction, *Environ. Sci. Technol.*, 53, 2375-2382, <https://doi.org/10.1021/acs.est.8b05353>, 2019.
- Yunker, M. B., Macdonald, R. W., Vingarzan, R., Mitchell, R. H., Goyette, D., and Sylvestre, S.: PAHs in the Fraser River basin: a critical appraisal of PAH ratios as indicators of PAH source and composition, *Org. Geochem.*, 33, 489-515, [https://doi.org/10.1016/s0146-6380\(02\)00002-5](https://doi.org/10.1016/s0146-6380(02)00002-5), 2002.
- Zhang, F., Chen, Y., Cui, M., Feng, Y., Yang, X., Chen, J., Zhang, Y., Gao, H., Tian, C., Matthias, V., and Liu, H.: Emission factors and environmental implication of organic pollutants in PM emitted from various vessels in China, *Atmos. Environ.*, 200, 302-311, 10.1016/j.atmosenv.2018.12.006, 2019.

- Zhang, L., Cheng, I., Muir, D., and Charland, J. P.: Scavenging ratios of polycyclic aromatic compounds in rain and snow in the Athabasca oil sands region, *Atmos. Chem. Phys.*, 15, 1421-1434, <https://doi.org/10.5194/acp-15-1421-2015>, 2015.
- 1905 Zhao, J., Zhang, J., Sun, L., Liu, Y., Lin, Y., Li, Y., Wang, T., and Mao, H.: Characterization of PM 2.5 -bound nitrated and oxygenated polycyclic aromatic hydrocarbons in ambient air of Langfang during periods with and without traffic restriction, *Atmos. Res.*, 213, 302-308, <https://doi.org/10.1016/j.atmosres.2018.06.015>, 2018.
- 1910 Zhao, J., Zhang, Y., Wang, T., Sun, L., Yang, Z., Lin, Y., Chen, Y., and Mao, H.: Characterization of PM2.5-bound polycyclic aromatic hydrocarbons and their derivatives (nitro-and oxy-PAHs) emissions from two ship engines under different operating conditions, *Chemosphere*, 225, 43-52, [10.1016/j.chemosphere.2019.03.022](https://doi.org/10.1016/j.chemosphere.2019.03.022), 2019.
- Zhao, J., Zhang, Y., Chang, J., Peng, S., Hong, N., Hu, J., Lv, J., Wang, T., and Mao, H.: Emission characteristics and temporal variation of PAHs and their derivatives from an ocean-going cargo vessel, *Chemosphere*, 249, 126194, [10.1016/j.chemosphere.2020.126194](https://doi.org/10.1016/j.chemosphere.2020.126194), 2020a.
- 1915 Zhao, S., Duan, Y., Yao, T., Liu, M., Lu, J., Tan, H., Wang, X., and Wu, L.: Study on the mercury emission and transformation in an ultra-low emission coal-fired power plant, *Fuel*, 199, 653-661, <https://doi.org/10.1016/j.fuel.2017.03.038>, 2017.
- Zhao, T., Yang, L., Huang, Q., Zhang, W., Duan, S., Gao, H., and Wang, W.: PM2.5-bound polycyclic aromatic hydrocarbons (PAHs) and nitrated-PAHs (NPAHs) emitted by gasoline vehicles: Characterization and health risk assessment, *Sci. Total Environ.*, 727, 138631, <https://doi.org/10.1016/j.scitotenv.2020.138631>, 2020b.
- 1920 Zheng, L., Ou, J., Liu, M., Chen, Y., Tang, Q., and Hu, Y.: Seasonal and spatial variations of PM10-bounded PAHs in a coal mining city, China: Distributions, sources, and health risks, *Ecotox. Environ. Safe.*, 169, 470-478, <https://doi.org/10.1016/j.ecoenv.2018.11.063>, 2019.
- 1925 Zielinska, B., Arey, J., Atkinson, R., and McElroy, P. A.: Formation of methylnitronaphthalenes from the gas-phase reactions of 1-and 2-methylnaphthalene with hydroxyl radicals and nitrogen oxide (N2O5) and their occurrence in ambient air, *Environ. Sci. Technol.*, 23, 723-729, <https://doi.org/10.1021/es00064a011>, 1989.

Table 1. Concentrations of PAHs (G+P) and percentage in the particulate phase (%PM) in Longyearbyen power plant, UNIS, and Adventdalen. Data are average, minimum and maximum, n=6 for each location*.

Compound name	Abbreviated name	Power plant			UNIS			Adventdalen		
		Mean ng m ⁻³	Min-Max ng m ⁻³	Mean %PM	Mean pg m ⁻³	Min-Max pg m ⁻³	Mean %PM	Mean pg m ⁻³	Min-Max pg m ⁻³	Mean %PM
Naphthalene	Nap	51.82	32.74-59.82	7.4	<MDL	<MDL	-	<MDL	<MDL	-
Acenaphthylene	Acy	2.30	1.22-3.80	0.6	16.89	7.14-29.15	0.0	2.40	1.10-5.13	0.0
Acenaphthene	Ace	0.87	0.30-2.18	8.4	48.48	24.29-72.99	0.0	3.84	1.25-6.62	0.0
Fluorene	Flu	7.61	3.68-12.16	4.6	170.50	136.5-236.0	1.1	59.96	38.49-95.82	1.8
Phenanthrene	Phe	27.32	12.01-44.87	5.6	409.20	368.5-470.0	6.5	236.30	191.7-270.8	3.7
Anthracene	Ant	1.06	0.23-2.13	0.0	18.04	12.29-25.52	0.0	14.25	10.46-19.33	3.5
Fluoranthene	Flt	6.99	1.43-12.49	3.9	28.45	24.14-36.06	40.5	19.14	10.76-38.54	23.2
Pyrene	Pyr	4.40	1.08-7.35	8.3	39.47	30.72-47.84	26.8	27.17	20.91-35.89	15.8
Benzo(a)anthracene	BaAnt	0.13	0.04-0.20	0.0	2.17	0.01-5.83	68.2	n.d.	n.d.	-
Chrysene	Chry	0.28	0.06-0.42	0.0	7.32	2.60-13.47	81.7	3.12	0.11-7.11	64.1
Benzo(b+k)fluoranthene	BbkFlt	n.d.	n.d.	-	2.23	0.01-5.87	100.0	0.75	0.01-2.79	100.0
Benzo(a)pyrene	BaPyr	n.d.	n.d.	-	0.89	0.01-2.46	100.0	0.34	0.01-1.16	100.0
Indeno(1.2.3-cd)pyrene	IPyr	n.d.	n.d.	-	1.63	0.07-3.79	100.0	0.71	0.07-2.67	100.0
Dibenzo(a,h)anthracene	DBAnt	n.d.	n.d.	-	n.d.	n.d.	-	n.d.	n.d.	-
Benzo(g,h,i)perylene	BPer	n.d.	n.d.	-	3.92	1.44-8.12	100.0	1.21	0.08-3.83	100.0
Σ16 PAHs		102.8	61.9-139.1	-	749.2	687.4-866.9	-	369.1	279.0-454.5	-
9-Fluorenone	9-Flu	12.35	5.57-19.54	19.2	270.30	128.2-543.8	41.7	139.40	110.2-177.2	25.5
9,10-Anthraquinone	9,10-AntQ	15.76	4.60-47.00	21.3	163.50	105.2-269.1	37.5	71.70	11.4-118.4	43.9
4H-Cyclopenta(def)-phenanthrene-4-one	cPPhe-4	1.30	0.51-2.55	15.8	27.23	20.16-35.80	65.5	18.77	11.97-39.10	38.1
9,10-Phenanthrenequinone	9,10-PheQ	2.13	0.96-4.40	0.0	<MDL	<MDL	-	<MDL	<MDL	-
Benzo(a)fluoren-11-one	BaFlu-11	0.16	0.08-0.23	27.6	6.07	1.79-11.08	100.0	2.23	0.71-4.36	100.0
Benzanthrone	BZT	0.87	0.14-1.31	0.0	1.76	0.02-4.32	96.7	0.10	0.02-0.58	100.0
Benzo(a)anthracene-7,12-dione	BaAnt-7,12	n.d.	n.d.	-	2.20	0.01-4.86	100.0	0.93	0.01-2.21	100.0
6H-Benzo(cd)pyren-6-one	BPyr-6	n.d.	n.d.	-	n.d.	n.d.	-	n.d.	n.d.	-
Σ8 OPAHs		32.6	15.8-73.1	-	471.0	325.9-741.4	-	233.1	124.7-337.1	-
1-Nitronaphthalene	1-NNap	2.19	0.99-4.69	61.7	16.97	13.36-21.53	0.1	5.02	1.91-9.84	1.5
2-Nitronaphthalene	2-NNap	0.26	0.11-0.40	31.7	5.08	2.44-7.33	3.1	1.88	1.29-2.83	5.4
2-Nitrobiphenyl	2-NBip	0.16	0.07-0.29	39.9	0.99	0.82-1.20	10.1	0.98	0.81-1.29	5.9
4-Nitrobiphenyl	4-NBip	n.d.	n.d.	-	2.23	1.51-2.68	0.0	2.45	0.29-4.10	0.0
1,5-Dinitronaphthalene	1,5-DNNap	n.d.	n.d.	-	0.80	0.05-2.17	80.0	0.93	0.05-3.72	53.9
5-Nitroacenaphthene	5-NAce	n.d.	n.d.	-	0.15	0.05-0.38	0.0	0.30	0.05-1.62	0.0
2-Nitrofluorene	2-NFlu	0.04	0.02-0.14	0.0	0.21	0.07-0.78	15.1	0.59	0.07-1.05	4.2
9-Nitroanthracene	9-NAnt	0.08	0.02-0.23	0.0	0.62	0.19-0.91	n.d.	2.26	0.12-4.70	57.8
9-Nitrophenanthrene	9-NPhe	n.d.	n.d.	-	0.20	0.09-0.37	n.d.	0.44	0.09-1.17	25.0
3-Nitrophenanthrene	3-NPhe	0.76	0.0003-1.93	96.1	n.d.	n.d.	-	n.d.	n.d.	-
2-Nitroanthracene	2-NAnt	0.31	0.07-0.62	0.0	n.d.	n.d.	-	n.d.	n.d.	-
2+3-Nitrofluoranthene	2+3-NFlt	0.52	0.06-1.14	0.0	9.50	7.32-11.37	94.5	12.30	4.68-26.66	79.8
4-Nitropyrene	4-NPyr	0.11	0.03-0.17	0.0	n.d.	n.d.	-	n.d.	n.d.	-
1-Nitropyrene	1-NPyr	n.d.	n.d.	-	n.d.	n.d.	-	n.d.	n.d.	-
2,7-Dinitrofluorene	2,7-DNFlu	0.06	0.001-0.14	0.0	n.d.	n.d.	-	n.d.	n.d.	-
7-Nitrobenzo(a)anthracene	7-NBaAnt	0.58	0.11-0.93	0.0	n.d.	n.d.	-	n.d.	n.d.	-
6-Nitrochrysene	6-NChry	n.d.	n.d.	-	n.d.	n.d.	-	n.d.	n.d.	-
1,3-Dinitropyrene	1,3-DNPyr	n.d.	n.d.	-	n.d.	n.d.	-	n.d.	n.d.	-

1,6-Dinitropyrene	1,6-DNPyr	n.d.	n.d.	-	n.d.	n.d.	-	n.d.	n.d.	-
1,8-Dinitropyrene	1,8-DNPyr	n.d.	n.d.	-	n.d.	n.d.	-	n.d.	n.d.	-
6-Nitrobenzo(a)pyrene	6-NBaPyr	n.d.	n.d.	-	n.d.	n.d.	-	n.d.	n.d.	-
Σ22 NPAHs		4.5	2.0-7.8	-	36.8	30.3-46.1	-	27.2	13.5-44.4	-

1930 *Full results are given in SI (Table S9-S11)

<MDL below method detection limit

n.d. not detected

1935 **Table 2.** Ratios of individual oxy- and nitro-PAHs to their corresponding parent PAHs in Longyearbyen power plant (G+P; n=6).

Ratio	Mean	STD
Nitro-PAH/PAH		
2-NFlu/Flu	0.004	0.005
3-NPhe/Phe	0.028	0.028
2-NAnt/Ant	0.150	0.107
9-NAnt/Ant	0.040	0.028
2+3-NFlt/Flt	0.030	0.023
7-NBaAnt/BaAnt	5.37	3.87
Oxy-PAH/PAH		
9,10-PheQ/Phe	0.08	0.01
cPPhen-4/Pyr	0.31	0.10
BaFlu-11/Chry	0.65	0.34
9-Flu/Phe	0.47	0.13
9-Flu/Flu	1.67	0.29
9,10-AntQ/Ant	12.17	7.30

1940

Table 3. PAHs diagnostic ratios for the Longyearbyen power plant (G+P; n=6).



Ratio	Mean	STD
Ant/(Ant+Phe)	0.044	0.002
Flt/(Flt+Pyr)	0.606	0.025
Flu/(Flu+Pyr)	0.638	0.110
BaAnt/(BaAnt+Chry)	0.332	0.027

Table 4.3. Source identification based on diagnostic ratios derived from total (G+P) concentrations; average of individual ratio values (n=6) with standard deviation are presented

	Ratio	Potential sources	Reference
IPyr/(IPyr+BPer)			
Power plant	n.d.	-	
UNIS	0.32±0.01	<0.35 Gasoline	(Ravindra et al., 2008b)
Adventdalen	0.45±0.05	0.35-0.70 Diesel	(Kavouras et al., 2001), (Ravindra et al., 2008b)
Flu/(Flu+Pyr)			
Power plant	0.64±0.11	-	
UNIS	0.81±0.04	>0.5 Coal combustion >0.5 Diesel	(Yunker et al., 2002), (Katsoyiannis and Breivik, 2014) (Ravindra et al., 2008b)
Adventdalen	0.68±0.05	0.64 Local power plant >0.5 Coal combustion	This study (Yunker et al., 2002), (Katsoyiannis and Breivik, 2014)
Flt/(Flt+Pyr)			
Power plant	n.d.	-	
UNIS	0.42±0.02	0.31–0.42 marine fuel	(Zhang et al., 2019)
Adventdalen	0.40±0.08	<0.5 Petrol emission	(Yunker et al., 2002)
BbkFlt/BPer			
Power plant	n.d.	-	
UNIS	0.48±0.03	<0.4 Gasoline	(Kuo et al., 2013)
Adventdalen	0.87±0.25	0.78 Diesel	(Kuo et al., 2013)







Figure 2. Air sampling transect locations in the vicinity of Longyearbyen.




PAHs, nitro- and oxy-PAHs



PAHs



Oxy-PAHs



Nitro-PAHs





Figure 3. Proportion of PAHs and PAH derivatives (G+P; excluding Nap) in the Longyearbyen power plant emission (n=6).



1955 **Figure 4.** Percentage of (a) PAHs and (b) nitro- and oxy-PAHs determined in particulate phase (% PM) at UNIS (n=6) and Adventdalen (n=6); individual standard deviations are used to calculate the intervals.





Figure 5. Principal component analysis [biplot](#) of PC1 and PC2 for Adventdalen samples (G+P; n=6).




Figure 6. Principal component analysis biplo+loading plot of PC1 and PC2 for UNIS samples (G+P; n=6).




Figure 6. Score (a) and loading (b) plots of PC1 and PC2 for Adventdalen (A) and UNIS (U) samples together (G+P; n=12).