

1 **Sensitivity Analysis of the Surface Ozone and Fine**
2 **Particulate Matter to Meteorological Parameters in China**

3 Zhihao Shi¹, Lin Huang¹, Jingyi Li¹, Qi Ying², Hongliang Zhang^{3,4}, Jianlin Hu^{1*}

4
5 ¹Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution
6 Control, Collaborative Innovation Center of Atmospheric Environment and Equipment
7 Technology, Nanjing University of Information Science & Technology, Nanjing 210044,
8 China

9 ²Zachry Department of Civil and Environmental Engineering, Texas A&M University,
10 College Station, TX 77843, USA

11 ³Department of Environmental Science and Engineering, Fudan University, Shanghai
12 200438, China

13 ⁴Institute of Eco-Chongming (SIEC), Shanghai 200062, China

14
15 *Corresponding authors:

16 Jianlin Hu, Email: jianlinhu@nuist.edu.cn. Phone: +86-25-58731504.

17 **Abstract**

18 Meteorological conditions play important roles in the formation of ozone (O₃) and
19 fine particulate matter (PM_{2.5}). China has been suffering from serious regional air
20 pollution problems, characterized by high concentrations of surface O₃ and PM_{2.5}. In
21 this study, the Community Multiscale Air Quality (CMAQ) model was used to quantify
22 the sensitivity of surface O₃ and PM_{2.5} to key meteorological parameters in different
23 regions of China. Six meteorological parameters were perturbed to create different
24 meteorological conditions, including temperature (T), wind speed (WS), absolute
25 humidity (AH), planetary boundary layer height (PBLH), cloud liquid water content
26 (CLW) and precipitation (PCP). Air quality simulations under the perturbed
27 meteorological conditions were conducted in China in January and July of 2013. The
28 changes in O₃ and PM_{2.5} concentrations due to individual meteorological parameters
29 were then quantified. T has a great influence on the daily maximum 8-h average O₃ (O₃-
30 8h) concentrations, which leads to O₃-8h increases by 1.7 ppb K⁻¹ in January in
31 Chongqing and 1.1 ppb K⁻¹ in July in Beijing. WS, AH, and PBLH have a smaller but
32 notable influence on O₃-8h with maximum change rates of 0.3, -0.15, and 0.14 ppb %⁻¹
33 ¹, respectively. T, WS, AH, and PBLH have important effects on PM_{2.5} formation of in
34 both January and July. In general, PM_{2.5} sensitivities are negative to T, WS, and PBLH
35 and positive to AH in most regions of China. The sensitivities in January are much
36 larger than in July. PM_{2.5} sensitivity to T, WS, PBLH, and AH in January can be up to -
37 5 μg m⁻³ K⁻¹, -3 μg m⁻³ %⁻¹, -1 μg m⁻³ %⁻¹, and +0.6 μg m⁻³ %⁻¹, respectively, and in
38 July can be up to -2 μg m⁻³ K⁻¹, -0.4 μg m⁻³ %⁻¹, -0.14 μg m⁻³ %⁻¹, and +0.3 μg m⁻³ %⁻¹,
39 respectively. Other meteorological factors (CLW and PCP) have negligible effects on
40 O₃-8h (less than 0.01 ppb %⁻¹) and PM_{2.5} (less than 0.01 μg m⁻³ %⁻¹). The results suggest
41 that surface O₃ and PM_{2.5} concentrations can change significantly due to changes in

42 meteorological parameters and it is necessary to consider these effects when developing
43 emission control strategies in different regions of China.

44

45 Keywords: sensitivity, meteorological conditions, fine particulate matter, ozone,

46 CMAQ model

1. Introduction

China has serious air pollution problems and fine particulate matter (PM_{2.5}) and ozone (O₃) are the two major air pollutants (Lin et al., 2010; Hu et al., 2016; Lu et al., 2019; Wu et al., 2019). The annual average PM_{2.5} concentrations were higher than 50 µg m⁻³ in 26 out of the total 31 provincial capital cities in mainland China during 2013-2014 (Wang et al., 2014a), and the national 4th highest daily maximum 8-hour average O₃ (O₃-8h) is 86.0 ppb during the warm-seasons (April–September) in 2013-2017, which is 6.3–30% higher than that in other industrialized regions of the world (Lu et al., 2018). PM_{2.5} alone caused 0.87-1.36 million deaths every year in China, and long-term exposure to O₃ was responsible for an extra 254 000 deaths (Apte et al., 2015; Cohen et al., 2017; Hu et al., 2017b; Silver et al., 2018). China has made remarkable improvement in air quality during recent years (Zhang et al., 2017; Zhao et al., 2017; China, 2018; Zheng et al., 2018), however, air pollution is still severe, making it the fourth-ranked healthy risk factor (Stanaway et al., 2018).

Surface PM_{2.5} and O₃ concentrations are determined by atmospheric processes of emissions, transport and dispersion, chemical transformation (due to gas-phase, aqueous-phase and aerosol chemistry), and dry and wet deposition. These processes are affected by meteorological conditions. Studies have shown that the surface O₃ and PM_{2.5} concentrations are sensitive to different meteorological parameters. For example, Dawson et al. (2007b) have investigated the sensitivity of surface O₃ to different meteorological parameters in the eastern United States (US) using the comprehensive air quality model with extensions (CAM_X). The results showed that temperature (T) had

69 the greatest influence on daily O₃-8h of 0.34 ppb K⁻¹, followed by absolute humidity
70 (AH) of 0.025 ppb %⁻¹. Bernard et al. (2001) also confirmed that T presented a notable
71 positive correlation with the surface O₃ concentration. The effects of meteorological
72 parameters on PM_{2.5} are even more complicated. Tran and Mölders (2011) showed that
73 elevated PM_{2.5} concentrations tended to occur under the condition of calm wind, low T
74 and relative humidity in Fairbanks, Alaska. Olvera Alvarez et al. (2018) used a land use
75 regression model to analyze the effects of different meteorological parameters on PM_{2.5}
76 in El Paso, Texas and obtained the same conclusion in winter, but in spring, the high
77 PM_{2.5} level was associated with high wind speed (WS) and low humidity. Dawson et al.
78 (2007a) studied the effects of individual meteorological parameters in the Eastern US
79 and found that PM_{2.5} concentration decreased markedly as the increased precipitation
80 (PCP) in winter, but in summer, the main meteorological factors affecting the PM_{2.5}
81 concentration were T, WS and planetary boundary layer height (PBLH). Dawson et al.
82 (2009) simulated the effects of climate change on regional and urban air quality in the
83 Eastern US, and found PM_{2.5} concentration decreased by 0.3 µg m⁻³ in January mostly
84 due to increasing in PCP and increased by 2.5 µg m⁻³ in July largely due to decreasing
85 in PBLH and WS. Horne and Dabdub (2017) altered various meteorological parameters
86 to investigate their effects on O₃, PM_{2.5} and secondary organic aerosols (SOA), and
87 found that the T predominated the effects of meteorology in California.

88 Many studies have proved that meteorological conditions play very important
89 roles in air pollution events in China. Studies found that the pollutant concentrations
90 could vary up to several times, due to meteorological changes with the same emission

91 sources (Zhang et al., 2010; Xing et al., 2011; Zheng et al., 2015; Cai et al., 2017; Liu
92 et al., 2017; Ning et al., 2018; Yang et al., 2018; Li et al., 2019b). For example, Xing et
93 al. (2011) studied that the difference between the effects of 2007 and 2008
94 meteorological conditions on air quality during the 2008 Beijing Olympics. They found
95 higher humidity in August 2008 was beneficial to the formation of SO_4^{2-} by up to ~60%,
96 and lower T prevented the evaporation of NO_3^- by up to ~60%. Liu et al. (2017) reported
97 that the monthly mean $\text{PM}_{2.5}$ concentrations in the Jing-Jin-Ji (JJJ) area in December
98 2015 increased by 5%~137% due to the unfavorable weather conditions such as low
99 WS and high humidity.

100 A few studies investigated the relationships between air quality and meteorological
101 conditions in China. Zhang et al. (2015) conducted a correlation analysis between air
102 quality and meteorology in three megacities Beijing, Shanghai and Guangzhou in China.
103 The result showed that air pollutants were significantly negatively correlated with WS,
104 and O_3 had a positive correlation with T. Yin et al. (2016) found that the relationship
105 between WS and $\text{PM}_{2.5}$ has complicated influence, with higher PM at low and high WS
106 than in light to moderate winds in Beijing from 2008 to 2014. Xu et al. (2018) examined
107 the variations of $\text{PM}_{2.5}$ concentration in January 2017 in China compared to that in
108 January 2016 and found meteorological conditions of low WS, high humidity, low
109 PBLH and low PCP contributed to $\text{PM}_{2.5}$ concentration worsening by 29.7%, 42.6%
110 and 7.9% in the JJJ region, the Pearl River Delta (PRD) region and the Cheng-Yu Basin
111 (CYB) region, respectively. Ma et al. (2019) analyzed the effects of meteorology on air
112 pollution in the Yangtze River Delta (YRD) region during 2014-2016 and found $\text{PM}_{2.5}$

113 was highly negatively correlated to WS, while O₃ concentration was positively
114 correlated to T but negatively related to relative humidity. Zhu et al. (2017) reported
115 that the surface concentrations of O₃ increased by 2-6 ppb in January and 8-12 ppb in
116 July 2014 in PRD, mainly due to the increase in T and the decrease in NO_x emissions.

117 These studies have investigated the impacts of meteorological conditions on PM_{2.5}
118 and O₃ in certain regions of China, however, quantitative sensitivity of PM_{2.5} and O₃ to
119 meteorological parameters has not been examined. The objective of this study is to
120 quantify the sensitivity of O₃ and PM_{2.5} to different meteorological parameters in winter
121 and summer in different regions of China. The paper is constructed as following,
122 Section 2 describes the method used to estimate the sensitivity, and Section 3 presents
123 the effects of each meteorological variable on O₃ and PM_{2.5} in China and in five
124 representative cities. Conclusions are then summarized in Section 4.

125

126 **2. Methods**

127 The sensitivity of O₃ and PM_{2.5} associated with changes in meteorological
128 parameters was quantified using the Community Multiscale Air Quality (CMAQ)
129 model version 5.0.2. The meteorological parameters include T, WS, AH, PBLH, PCP,
130 and cloud liquid water content (CLW). A base case was firstly simulated with
131 meteorological fields predicted by the Weather Research and Forecasting (WRF) model
132 v3.7.1 (<http://www.wrf-model.org/>) using the NCEP FNL Operational Model Global
133 Tropospheric Analyses dataset as the initial and boundary conditions. The base case has
134 been described in a previous study and the model configurations of the base case were
135 reported there (Hu et al., 2015). The WRF predicted meteorological parameters and the

136 CMAQ predicted surface O₃ and PM_{2.5} have been evaluated against observations at 422
137 sites in 60 major cities in China, and the accuracy of the model performance has been
138 validated (Hu et al., 2016).

139 A suit of perturbation scenarios was created, and in each scenario, a certain
140 meteorological parameter was perturbed to a certain extent. The details of the
141 perturbation scenarios are listed in Table 1. Among those changes, the T was absolute
142 changes, and other parameters are relative variations. The magnitude ranges of
143 perturbations are based on IPCC AR5 report and the study of Dawson et al. (2007) and
144 the references therein. For each parameter, three positive and three negative
145 perturbations were then designed within its range to have a more comprehensive
146 examination on the sensitivity of PM_{2.5} and O₃ to this parameter. All perturbations were
147 implemented uniformly in space on the modeling domain and in time through the
148 modeling periods. The perturbations on temperature, wind speed, and absolute humidity
149 were made in all layers. To separate the effects of individual meteorological parameters,
150 only one parameter was changed in each case while all other parameters were kept
151 unchanged. Therefore, cloud dissipating or forming in response to changing
152 temperature was not considered in the simulations. Please note that this type of
153 perturbations are not what happens in the real world where meteorological parameters
154 are inter-linked. When perturbing horizontal wind speed, to avoid unphysical situations
155 that mass would not be conserved, the vertical wind speed was adjusted in the vertical
156 transport calculation based on the air density changes to conserve mass.

157 Then the CMAQ model was re-run to predict the air quality under the perturbed

158 meteorological condition. The emissions and other inputs were kept unchanged in each
159 perturbed meteorological scenarios, therefore the difference of O₃ and PM_{2.5}
160 concentrations between each of the perturbation case and the base case was due to the
161 change in the specific meteorological parameter, and the sensitivity of O₃ and PM_{2.5} to
162 individual meteorological parameters could be quantitatively determined. It is
163 worthwhile to note that some meteorological parameters could have significant impacts
164 on emissions, such as the effect of T on biogenic VOC and soil NO_x emissions, the
165 cloud cover/convection on lightning NO_x emissions, the effect of T on power plant
166 NO_x emissions (high T leads to higher electricity demand in summer), which would
167 affect air quality. Therefore, the sensitivities in this study only include the ‘direct’
168 effects of individual meteorological parameters on air quality. A full evaluation of the
169 impacts of climate/weather changes on air quality should consider effects of the
170 emissions changes.

171 The modeling domain covers East Asia, including entire China, with a horizontal
172 resolution of 36×36 km². The base case and perturbation cases were conducted in
173 January and July in 2013, representing the winter and the summer conditions,
174 respectively. In addition to the regional analysis, five representative megacities were
175 selected, i.e., Beijing, Shanghai, Guangzhou, Chongqing, Xi’an (Fig.1). These cities are
176 located in the North China Plain (NCP), YRD, PRD, CYB, and Guanzhong Plain,
177 respectively, where serious air pollution problems often occur. In this study, O₃-8h was
178 used in the O₃ analyses, and 24h average PM_{2.5} was used in the PM_{2.5} analyses, if not
179 specifically stated. The O₃-T relationship is examined using the method in Rasmussen

180 et al. (2012) in the five cities. Observed and predicted O₃-T relationships were estimated
181 using the daily observed and predicted O₃-8h concentrations and daily maximum T in
182 July (O₃ observations became available from March 2013 in China, so no O₃
183 observations in January). The results are shown in Fig. S1. CMAQ predicts positive O₃-
184 T relationship in most cities except in Beijing, and the model tends to underestimated
185 the daily O₃-T relationship except in Shanghai. The underestimation of O₃-T by the
186 CMAQ model in this study is consistent with the findings in Rasmussen et al. (2012).
187 Please note that we only have 1 month data and we use daily O₃-8h and daily maximum
188 temperature in the evaluation, while a much more meaningful evaluation should be
189 performed to use monthly averaged O₃ and monthly average temperature over a long-
190 term period Rasmussen et al. (2012).

191

192 **3. Results and Discussion**

193 **3.1 Impacts of meteorological parameters on surface O₃**

194 Figs. 2(a) and 2(b) show the spatial-distribution of the predicted monthly average
195 O₃-8h concentrations in January and July, respectively. In January, the highest average
196 concentrations are about 70 ppb in the Sichuan Basin, and the concentrations in
197 southern and eastern China are generally higher than those in northern China. In July,
198 the highest average concentrations are over 80ppb in the large areas of NCP and YRD,
199 CYB, and Guangzhou areas in the PRD.

200 Fig. 3 shows the spatial distribution of the concentration changes of O₃-8h in
201 January and July due to change of T + 1.0 K, WS - 10%, AH +10% PBLH - 20%, CLW
202 + 10% case, and PCP + 10%, respectively. Fig. S1-S3 shows the results due to other

203 extent changes in these parameters. When T increases 1.0 K (Fig. 3(a)), O₃-8h increases
204 1-2 ppb in most area of eastern and central China in January and in NCP and YRD in
205 July, which is consistent with the high O₃ spatial distribution in the base case (shown in
206 Fig. 2). O₃-8h decreases up to 4 ppb in January in Northeast China, and up to 2 ppb in
207 the Southwest border of China and the East China Sea, which are the areas of low O₃
208 concentrations (generally less than the background O₃ concentration of 35 ppb).
209 Therefore, the effect of T on O₃ is dependent on the O₃ formation regime. An increase
210 in T promotes O₃ formation chemistry in net O₃ formation areas (O₃ concentrations
211 greater than 35 ppb), but accelerates O₃ consumption chemistry in the net O₃ loss areas
212 (O₃ concentrations less than 35 ppb).

213 Fig. 3(b) shows that the differences of O₃-8h in January and July when WS is 10%
214 less than the base case in 2013. The influence of wind on O₃ concentration is complex,
215 but generally, slower WS decreases O₃ in January in most parts of China, particularly
216 in Sichuan by up to 3 ppb, but increases O₃ in July by a few ppb over the most areas in
217 eastern and central China. Therefore, the impact of WS on O₃ appears opposite in winter
218 and summer. Weaker winds slow down the dispersion of NO_x and VOCs, which is
219 conducive to O₃ formation in summer when the vertical mixing is strong, but increases
220 O₃ titration in the surface in winter due to weaker vertical mixing.

221 Fig. 3(c) displays that the surface O₃ is expected to decrease generally less than 1
222 ppb when AH increases by 10% (relative change) both in January and July in most land
223 areas of China except in the northeast area. Fig. 3(d) shows that a 20% decrease of
224 PBLH leads to O₃-8h decreases by a few ppb in most area in January, while in July O₃-

225 8h increases in eastern and central regions, especially in YRD, CYB and areas in Hubei-
226 Hunan-Jiangxi in the central China. Sensitivity of O₃ to CLW and PCP is relatively
227 small. Fig. 3(e) demonstrates that O₃-8h changes -0.03 ppb to 0.03 ppb in January and
228 July for a 10% increase in CLW. Fig. 3(f) demonstrates that a 10% increase in PCP
229 results in -0.1 to 0.2 ppb changes in O₃-8h. O₃ changes due to the six meteorological
230 factors with different extents of perturbation (Figs. S1-S3) shows the similar trends and
231 spatial patterns.

232 **3.2 Impacts of meteorological parameters on surface PM_{2.5}**

233 Figs. 2(c) and 2(d) show the spatial distribution of the monthly average surface
234 PM_{2.5} concentrations in January and July. PM_{2.5} in January reaches over 200 μg m⁻³ in
235 JJJ, SYB, central China, and urban areas in the Northeast China. PM_{2.5} is much lower
236 in July, generally lower than 50 μg m⁻³, but is high (up to 70 μg m⁻³) in areas in the JJJ,
237 YRD, and central China regions.

238 Fig. 4 shows the spatial distribution of PM_{2.5} changes due to the same changes of
239 meteorological factors, as in Fig. 3. The PM_{2.5} results of other cases of the sensitivity
240 study are shown in Figs. S4-S6 of the Supplementary Materials. The results indicate
241 that in January, a 1.0 K increase in T leads to up to 5-6 μg m⁻³ decrease of PM_{2.5} in JJJ
242 and central China; in July, a 1.0 K increase in T causes PM_{2.5} increase by about 1 μg m⁻³
243 in southern China but decrease by 1-3 μg m⁻³ in the JJJ and east coast region. A 10%
244 decrease in WS causes PM_{2.5} increase up to over 40 μg m⁻³ in January and up to 5 μg
245 m⁻³ in July. A 10% relative increase in AH leads to PM_{2.5} increase of up to 6 μg m⁻³ in
246 January and up to 2 μg m⁻³ in JJJ and northeast regions but slightly decrease of less than

247 1 $\mu\text{g m}^{-3}$ in southern China in July. A 20% decrease of PBLH causes $\text{PM}_{2.5}$ increases by
248 up to 20 $\mu\text{g m}^{-3}$ in January and up to 4 $\mu\text{g m}^{-3}$ in July. The impact of CLW and PCP on
249 $\text{PM}_{2.5}$ is small, and generally increase in CLW increases surface $\text{PM}_{2.5}$ and increase in
250 PCP decreases $\text{PM}_{2.5}$.

251 The changes in the total $\text{PM}_{2.5}$ mass concentrations are determined by the changes
252 in the chemical components of $\text{PM}_{2.5}$. Fig. S7 displays the fraction of $\text{PM}_{2.5}$ species
253 (elemental carbon (EC), primary organic carbon (POC), secondary organic aerosol
254 (SOA), sulfate (SO_4^{2-}), nitrate (NO_3^-), and ammonium (NH_4^+)) in five representative
255 cities. Secondary inorganic aerosols (SO_4^{2-} , NO_3^- , NH_4^+) are the major PM components,
256 accounting for over 50% of $\text{PM}_{2.5}$ in January and about 40% in July. Fig. 5 and Fig. 6
257 show the changes of the major $\text{PM}_{2.5}$ components due to the same changes of
258 meteorological factors as in Fig. 4 in January and in July, respectively. The results show
259 that the effects of the meteorological parameters on the total $\text{PM}_{2.5}$ (shown in Fig. 4)
260 are mainly due to their effects on SO_4^{2-} , NO_3^- , and NH_4^+ in January, and due to the
261 changes in SO_4^{2-} , NO_3^- , NH_4^+ , and SOA in July. In general, PBLH, WS, and PCP are
262 negatively correlated to SO_4^{2-} , NO_3^- , and NH_4^+ formation, but AH and CLW are
263 positively correlated to these components. SOA concentrations are much higher in July
264 than in January due to the contribution from biogenic emissions (Hu et al., 2017a). SOA
265 formation is affected by reaction rates (positively affected by T), availability of oxidants
266 (such as changes in O_3), and hydrogen ion strength (affected by changes in SO_4^{2-} , NO_3^- ,
267 NH_4^+). SOA concentrations mainly increase in south China.

268 It is worthwhile noting that the effects of T on SO_4^{2-} and NO_3^- (changes of NH_4^+

269 is determined by changes of SO_4^{2-} and NO_3^-). Both in January and July, increase in T
270 decreases SO_4^{2-} and NO_3^- in the major areas of eastern China. The NO_3^- decreases is
271 expected because volatile NH_4NO_3 favors more in gas phase in higher temperature, and
272 this result is consistent with studies in other regions (Dawson et al., 2007a; Horne and
273 Dabdub, 2017). SO_4^{2-} is found to increase with T increase in those studies because faster
274 gas- and aqueous- phase reactions of SO_4^{2-} . However, our finding of SO_4^{2-} in China is
275 opposite. The CMAQ-Sulfur Tracking Model (CMAQ-STM) was further used to track
276 the SO_4^{2-} formation from different processes. The results confirm that the SO_4^{2-}
277 production from gas- and aqueous- phase increases with T increase. But meanwhile
278 SO_4^{2-} production from heterogeneous reactions is reduced more when T is increased.
279 Heterogeneous SO_4^{2-} formation has been proposed as a major SO_4^{2-} formation pathway
280 during China haze events (Wang et al., 2014b; Gen et al., 2019; Huang et al., 2019; Li
281 et al., 2019a) and in this study it accounts for up to ~75% of total SO_4^{2-} production. The
282 treatment of heterogeneous SO_4^{2-} formation currently is modeled as a surface-
283 controlled uptake process, in which the formation rate is determined by the aerosol
284 surface area and the uptake coefficient of SO_2 on particle surface (Ying et al., 2014).
285 When T is increased, the particle surface area decreases (as particle mass concentration
286 decreases due to a combined effect of other components), resulting in decrease in the
287 heterogeneous SO_4^{2-} formation.

288 An additional simulation was run to illustrate the combined effects of perturbations
289 in all meteorological parameters (T+1.0K, WS-10%, AH+10%, PBLH-20%,
290 CLW+10%, and PCP+10%) on O_3 and $\text{PM}_{2.5}$ in January and July. The results are shown

291 in Fig. S8. The average O₃-8h concentration in this combined-change simulation drop
292 by ~2 ppb in January, except in the northeast. In July, O₃ in eastern China and the
293 Sichuan basin rose by 2 ppb. The changes in PM_{2.5} resulting from this combined-change
294 simulation were significantly higher, compared to the basecase concentrations, even
295 increased by up to 50 µg m⁻³ in January.

296

297 **3.3. Quantitative sensitivity of O₃ and PM_{2.5} to individual** 298 **meteorology parameters**

299 The quantitative sensitivity of O₃ and PM_{2.5} concentrations to individual
300 meteorological parameter is calculated by linear fitting of the changes in monthly-
301 average concentrations under all of the six perturbed cases of the meteorological
302 parameter. Figs. S8-S10 in Supplementary Materials show the calculation examples of
303 T, WS, and AH on O₃ at the five major cities of Beijing, Chongqing, Guangzhou,
304 Shanghai and Xi'an, and Figs. S11-S13 shows the examples for the PM_{2.5} cases. Fig.7
305 demonstrates the sensitivities of O₃-8h and PM_{2.5} and its components to each
306 meteorological parameter in the five cities. In January, T has a positive impact on O₃ in
307 all cities, and the largest impact is in Chongqing with a rate of +1.69 ppb K⁻¹. In July,
308 O₃ also shows a strong positive sensitivity to T in Beijing with +1.06 ppb K⁻¹ and in
309 Shanghai with +0.98 ppb K⁻¹, but has a small negative sensitivity (-0.15 ppb K⁻¹) in
310 Xi'an and a moderate negative sensitivity (-0.74 ppb K⁻¹) in Guangzhou. The O₃
311 sensitivity to T in Guangzhou in July shows a highly nonlinear trend and is very
312 different from other cities (Fig. S8(c)). More studies are needed to investigate the effects

313 of T on O₃ pollution in the YRD region during summertime. WS and PBLH both have
314 positive effects on O₃-8h in January, the effects vary significantly among cities, with
315 0.004-0.3 ppb %⁻¹ for WS and 0.04-0.14 ppb %⁻¹ for PBLH. AH has a negative effect
316 on O₃-8h in January, ranging from -0.01 to -0.15 ppb %⁻¹. But in July, the impacts of
317 WS, AH, and PBLH are negative in most cities, with a range of -0.05 to -0.18, -0.05 to
318 -0.13, and -0.02 to -0.07 ppb %⁻¹, respectively. Generally speaking, T, WS, AH and
319 PBLH led to rather larger O₃ changes. The sensitivity of O₃ to CLW and PCP is even
320 minimal (less than 0.1 ppb %⁻¹) and mostly negative.

321 Negative sensitivities are found for surface PM_{2.5} concentrations to T, WS, PBLH,
322 and PCP, and positive sensitivities for PM_{2.5} to AH and CLW. The sensitivity of T in the
323 five cities ranges from -1.5 to -3.6 μg m⁻³ K⁻¹ in January and -0.3 to -1.65 μg m⁻³ K⁻¹ in
324 July. PM_{2.5} is also very sensitive to WS in January, with a range of -0.8 to -2.97 μg m⁻³
325 %⁻¹, while the sensitivity (-0.03 to -0.19 μg m⁻³ %⁻¹) becomes much smaller in July.
326 The sensitivity to PBLH is -0.12 to -0.58 μg m⁻³ %⁻¹ in January and -0.003 to -0.23 μg
327 m⁻³ %⁻¹ in July. The sensitivity to AH is 0.16 to 0.30 μg m⁻³ %⁻¹ in January and 0.05 to
328 0.27 μg m⁻³ %⁻¹ in July. Sensitivity to CLW and PCP is small in January and July, mostly
329 less than 0.01 μg m⁻³ %⁻¹. The PM_{2.5} sensitivities can be explained by the major
330 components of SO₄²⁻, NO₃⁻, and NH₄⁺ in January and by SO₄²⁻, NO₃⁻, NH₄⁺, and SOA
331 in July.

332 Fig. 8 shows the spatial variations of the sensitivity of O₃-8h and PM_{2.5} to the
333 meteorological parameters. The sensitivity of O₃-8h to temperature is more significant

334 in Sichuan and southern provinces of China in January, and in NCP and YRD in July,
335 up to +2 ppb K⁻¹ in both January and July. O₃-8h sensitivity to WS is diverse in space,
336 and is generally positive in Sichuan and southern provinces in January; and it is negative
337 in east China but positive in west China. O₃-8h sensitivity to AH is generally negative
338 in both months in most regions of China, except the northeast in January and southwest
339 in July. O₃-8h sensitivity to PBLH is mostly positive in January but becomes negative
340 in YRD, CYB, NCP, and central China in July. O₃-8h sensitivity to CLW and PCP is
341 negligible.

342 Fig. 9 displays the spatial variations of the sensitivity of surface PM_{2.5} to the
343 meteorological parameters. PM_{2.5} sensitivities to the meteorological parameters are
344 more consistent in January and July than the cases of O₃, i.e., negative sensitivity to T,
345 WS, PBLH, PCP, and positive to AH and CLW in most regions of China in both months.
346 On the other hand, PM_{2.5} sensitivities are more profound in January than in July. PM_{2.5}
347 sensitivity to T is up to -5 μg m⁻³ K⁻¹ in January and up to -2 μg m⁻³ K⁻¹ in July. PM_{2.5}
348 sensitivity to WS is up to -3 μg m⁻³ %⁻¹ in January, and up to -0.4 μg m⁻³ %⁻¹ in July.
349 PM_{2.5} sensitivity to PBLH is up to -1 μg m⁻³ %⁻¹ in January, and up to -0.14 μg m⁻³ %⁻¹
350 in July. PM_{2.5} sensitivity to AH is up to +0.6 μg m⁻³ %⁻¹ in January, and up to 0.3 μg m⁻³
351 %⁻¹ in July. The sensitivities to CLW and PCP is small, compared to the other four
352 meteorological parameters. PM_{2.5} sensitivity to T is negative in most land areas of China
353 in January and in NCP and YRD in July because of the negative effects of T on SO₄²⁻,
354 NO₃⁻, and NH₄⁺, as discussed in previous sections. PM_{2.5} sensitivity to T is positive in
355 south China in July due to more SOA with higher T.

4. Conclusions

Meteorological conditions can have a great influence on surface O_3 and $PM_{2.5}$ concentrations. In this study, the sensitivities of O_3 -8h and $PM_{2.5}$ to T, WS, AH, PBLH, PCP, and CLW are quantitatively estimated in January and July, respectively in China. The response of O_3 -8h to T is important and the sensitivity can be up to +2 ppb K^{-1} in both January and July, and the sensitivity is dependent on the O_3 chemistry formation or loss regime, i.e., positive in the net O_3 formation areas, and negative in the O_3 consumption areas. In general, $PM_{2.5}$ sensitivities are negative to T, WS, PBLH, and PCP and positive to AH and CLW in most regions of China in both January and July. The sensitivities in January are much larger than in July. $PM_{2.5}$ sensitivities to T, WS, AH, and PBLH are important. The $PM_{2.5}$ sensitivities to these meteorological parameters are through major effects on SO_4^{2-} , NO_3^- , NH_4^+ , and SOA. The sensitivities of O_3 and $PM_{2.5}$ to CLW and PCP are negligible. The results show that O_3 and $PM_{2.5}$ concentrations in China are greatly affected by meteorological conditions, therefore changes in these meteorological parameters due to climate change or inter-annual meteorological variations could potentially alter O_3 and $PM_{2.5}$ concentrations significantly, and it should consider these effects when developing emission control strategies. The results also show that the O_3 and $PM_{2.5}$ sensitivities to meteorological parameters have substantial spatial variations. Future studies can further investigate how the changes in meteorological conditions affect the effectiveness of emission control plans in reaching the designed air quality objectives in the different regions of China.

378

379 **Data availability.** All of the modeling results will be available online after we
380 publish the paper.

381 **Author contributions.** ZS, and JH designed research. ZS, LH, JL, QY, HZ
382 and JH contributed to model development and configuration. ZS, LH, JL, and JH
383 analyzed the data. ZS prepared the manuscript and all coauthors helped improve the
384 manuscript.

385 **Competing interests.** The authors declare that they have no conflict of
386 interest.

387 **Acknowledgment**

388 This work was supported by the National Key R&D Program of China
389 (2018YFC0213800), the National Natural Science Foundation of China (41975162,
390 41675125 and 41705102), and Jiangsu Environmental Protection Research Project
391 (2016015).

392

393 **References**

- 394 Apte, J.S., Marshall, J.D., Cohen, A.J., Brauer, M., 2015. Addressing global mortality from ambient PM_{2.5}.
395 *Environmental science & technology* 49, 8057-8066.
- 396 Bernard, S.M., Samet, J.M., Grambsch, A., Ebi, K.L., Romieu, I., 2001. The potential impacts of climate
397 variability and change on air pollution-related health effects in the United States. *Environmental health*
398 *perspectives* 109, 199-209.
- 399 Cai, W., Li, K., Liao, H., Wang, H., Wu, L., 2017. Weather conditions conducive to Beijing severe haze
400 more frequent under climate change. *Nature Climate Change* 7, 257.
- 401 China, 2018. Air Quality Targets Set by the Action Plan Have Been Fully Realized(2018).
- 402 Cohen, A.J., Brauer, M., Burnett, R., Anderson, H.R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef,
403 B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y.,
404 Martin, R., Morawska, L., Pope, C.A., Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R., van
405 Donkelaar, A., Vos, T., Murray, C.J.L., Forouzanfar, M.H., 2017. Estimates and 25-year trends of the global
406 burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of
407 Diseases Study 2015. *The Lancet* 389, 1907-1918.
- 408 Dawson, J., Adams, P., Pandis, S., 2007a. Sensitivity of PM_{2.5} to climate in the Eastern US: a modeling
409 case study. *Atmospheric chemistry and physics* 7, 4295-4309.
- 410 Dawson, J.P., Adams, P.J., Pandis, S.N., 2007b. Sensitivity of ozone to summertime climate in the eastern
411 USA: A modeling case study. *Atmospheric environment* 41, 1494-1511.
- 412 Dawson, J.P., Racherla, P.N., Lynn, B.H., Adams, P.J., Pandis, S.N., 2009. Impacts of climate change on
413 regional and urban air quality in the eastern United States: Role of meteorology. *Journal of Geophysical*
414 *Research: Atmospheres* 114.
- 415 Gen, M.S., Zhang, R.F., Huang, D.D., Li, Y.J., Chan, C.K., 2019. Heterogeneous SO₂ Oxidation in Sulfate
416 Formation by Photolysis of Particulate Nitrate. *Environmental Science & Technology Letters* 6, 86-91.
- 417 Horne, J.R., Dabdub, D., 2017. Impact of global climate change on ozone, particulate matter, and
418 secondary organic aerosol concentrations in California: A model perturbation analysis. *Atmospheric*
419 *Environment* 153, 1-17.
- 420 Hu, J., Wang, P., Ying, Q., Zhang, H., Chen, J., Ge, X., Li, X., Jiang, J., Wang, S., Zhang, J., Zhao, Y., Zhang,
421 Y., 2017a. Modeling biogenic and anthropogenic secondary organic aerosol in China. *Atmos. Chem. Phys.*
422 *17*, 77-92.
- 423 Hu, J., Wu, L., Zheng, B., Zhang, Q., He, K., Chang, Q., Li, X., Yang, F., Ying, Q., Zhang, H., 2015. Source
424 contributions and regional transport of primary particulate matter in China. *Environ Pollut* 207, 31-42.
- 425 Hu, J.L., Chen, J.J., Ying, Q., Zhang, H.L., 2016. One-year simulation of ozone and particulate matter in
426 China using WRF/CMAQ modeling system. *Atmospheric Chemistry and Physics* 16, 10333-10350.
- 427 Hu, J.L., Huang, L., Chen, M.D., Liao, H., Zhang, H.L., Wang, S.X., Zhang, Q., Ying, Q., 2017b. Premature
428 Mortality Attributable to Particulate Matter in China: Source Contributions and Responses to Reductions.
429 *Environmental Science & Technology* 51, 9950-9959.
- 430 Huang, L., An, J.Y., Koo, B., Yarwood, G., Yan, R.S., Wang, Y.J., Huang, C., Li, L., 2019. Sulfate formation
431 during heavy winter haze events and the potential contribution from heterogeneous SO₂ + NO₂
432 reactions in the Yangtze River Delta region, China. *Atmospheric Chemistry and Physics* 19, 14311-14328.
- 433 Li, M.M., Wang, T.J., Xie, M., Li, S., Zhuang, B.L., Huang, X., Chen, P.L., Zhao, M., Liu, J.E., 2019a.
434 Formation and Evolution Mechanisms for Two Extreme Haze Episodes in the Yangtze River Delta Region
435 of China During Winter 2016. *J Geophys Res-Atmos* 124, 3607-3623.

436 Li, X., Gao, Z., Li, Y., Gao, C.Y., Ren, J., Zhang, X., 2019b. Meteorological conditions for severe foggy haze
437 episodes over north China in 2016–2017 winter. *Atmospheric Environment* 199, 284-298.

438 Lin, J., Nielsen, C.P., Zhao, Y., Lei, Y., Liu, Y., McElroy, M.B., 2010. Recent Changes in Particulate Air
439 Pollution over China Observed from Space and the Ground: Effectiveness of Emission Control.
440 *Environmental Science & Technology* 44, 7771-7776.

441 Liu, T., Gong, S., He, J., Yu, M., Wang, Q., Li, H., Liu, W., Zhang, J., Li, L., Wang, X., Li, S., Lu, Y., Du, H.,
442 Wang, Y., Zhou, C., Liu, H., Zhao, Q., 2017. Attributions of meteorological and emission factors to the
443 2015 winter severe haze pollution episodes in China's Jing-Jin-Ji area. *Atmospheric Chemistry and*
444 *Physics* 17, 2971-2980.

445 Lu, H., Lyu, X., Cheng, H., Ling, Z., Guo, H., 2019. Overview on the spatial–temporal characteristics of the
446 ozone formation regime in China. *Environmental Science: Processes & Impacts*.

447 Lu, X., Hong, J., Zhang, L., Cooper, O.R., Schultz, M.G., Xu, X., Wang, T., Gao, M., Zhao, Y., Zhang, Y., 2018.
448 Severe Surface Ozone Pollution in China: A Global Perspective. *Environmental Science & Technology*
449 *Letters* 5, 487-494.

450 Ma, T., Duan, F., He, K., Qin, Y., Tong, D., Geng, G., Liu, X., Li, H., Yang, S., Ye, S., Xu, B., Zhang, Q., Ma, Y.,
451 2019. Air pollution characteristics and their relationship with emissions and meteorology in the Yangtze
452 River Delta region during 2014–2016. *Journal of Environmental Sciences* 83, 8-20.

453 Ning, G., Wang, S., Yim, S.H.L., Li, J., Hu, Y., Shang, Z., Wang, J., Wang, J., 2018. Impact of low-pressure
454 systems on winter heavy air pollution in the northwest Sichuan Basin, China. *Atmospheric Chemistry*
455 *and Physics* 18, 13601-13615.

456 Olvera Alvarez, H.A., Myers, O.B., Weigel, M., Armijos, R.X., 2018. The value of using seasonality and
457 meteorological variables to model intra-urban PM_{2.5} variation. *Atmospheric Environment* 182, 1-8.

458 Silver, B., Reddington, C.L., Arnold, S.R., Spracklen, D.V., 2018. Substantial changes in air pollution across
459 China during 2015–2017. *Environmental Research Letters* 13, 114012.

460 Stanaway, J.D., Afshin, A., Gakidou, E., Lim, S.S., Abate, D., Abate, K.H., Abbafati, C., Abbasi, N.,
461 Abastabar, H., Abd-Allah, F.J.T.L., 2018. Global, regional, and national comparative risk assessment of
462 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195
463 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017.
464 *Lancet* 392, 1923-1994.

465 Tran, H.N.Q., Mölders, N., 2011. Investigations on meteorological conditions for elevated PM_{2.5} in
466 Fairbanks, Alaska. *Atmospheric Research* 99, 39-49.

467 Wang, Y., Ying, Q., Hu, J., Zhang, H., 2014a. Spatial and temporal variations of six criteria air pollutants
468 in 31 provincial capital cities in China during 2013–2014. *Environ Int* 73, 413-422.

469 Wang, Y.X., Zhang, Q.Q., Jiang, J.K., Zhou, W., Wang, B.Y., He, K.B., Duan, F.K., Zhang, Q., Philip, S., Xie,
470 Y.Y., 2014b. Enhanced sulfate formation during China's severe winter haze episode in January 2013
471 missing from current models. *J Geophys Res-Atmos* 119.

472 Wu, C., Hu, W., Zhou, M., Li, S., Jia, Y., 2019. Data-driven regionalization for analyzing the spatiotemporal
473 characteristics of air quality in China. *Atmospheric Environment* 203, 172-182.

474 Xing, J., Zhang, Y., Wang, S., Liu, X., Cheng, S., Zhang, Q., Chen, Y., Streets, D.G., Jang, C., Hao, J., Wang,
475 W., 2011. Modeling study on the air quality impacts from emission reductions and atypical
476 meteorological conditions during the 2008 Beijing Olympics. *Atmospheric Environment* 45, 1786-1798.

477 Xu, Y., Xue, W., Lei, Y., Zhao, Y., Cheng, S., Ren, Z., Huang, Q., 2018. Impact of Meteorological Conditions
478 on PM_{2.5} Pollution in China during Winter. *Atmosphere* 9, 429.

479 Yang, Y., Zheng, X., Gao, Z., Wang, H., Wang, T., Li, Y., Lau, G.N., Yim, S.H., 2018. Long - term trends of

480 persistent synoptic circulation events in planetary boundary layer and their relationships with haze
481 pollution in winter half year over eastern China. *Journal of Geophysical Research: Atmospheres* 123,
482 10,991-911,007.

483 Yin, Q., Wang, J., Hu, M., Wong, H., 2016. Estimation of daily PM2.5 concentration and its relationship
484 with meteorological conditions in Beijing. *Journal of Environmental Sciences* 48, 161-168.

485 Ying, Q., Cureño, I.V., Chen, G., Ali, S., Zhang, H., Malloy, M., Bravo, H.A., Sosa, R., 2014. Impacts of
486 Stabilized Criegee Intermediates, surface uptake processes and higher aromatic secondary organic
487 aerosol yields on predicted PM2.5 concentrations in the Mexico City Metropolitan Zone. *Atmospheric*
488 *Environment* 94, 438-447.

489 Zhang, H., Wang, Y., Hu, J., Ying, Q., Hu, X.-M., 2015. Relationships between meteorological parameters
490 and criteria air pollutants in three megacities in China. *Environmental Research* 140, 242-254.

491 Zhang, J., Reid, J.S., Alfaro-Contreras, R., Xian, P., 2017. Has China been exporting less particulate air
492 pollution over the past decade? *Geophysical Research Letters* 44, 2941-2948.

493 Zhang, L., Liao, H., Li, J., 2010. Impacts of Asian summer monsoon on seasonal and interannual variations
494 of aerosols over eastern China. *Journal of Geophysical Research: Atmospheres* 115.

495 Zhao, B., Jiang, J.H., Gu, Y., Diner, D., Worden, J., Liou, K.-N., Su, H., Xing, J., Garay, M., Huang, L., 2017.
496 Decadal-scale trends in regional aerosol particle properties and their linkage to emission changes.
497 *Environmental Research Letters* 12.

498 Zheng, B., Chevallier, F., Ciais, P., Yin, Y., Deeter, M.N., Worden, H.M., Wang, Y., Zhang, Q., He, K., 2018.
499 Rapid decline in carbon monoxide emissions and export from East Asia between years 2005 and 2016.
500 *Environmental Research Letters* 13.

501 Zheng, X., Fu, Y., Yang, Y., Liu, G., 2015. Impact of atmospheric circulations on aerosol distributions in
502 autumn over eastern China: observational evidence. *Atmospheric Chemistry and Physics* 15, 12115.

503 Zhu, K., Xie, M., Wang, T., Cai, J., Li, S., Feng, W., 2017. A modeling study on the effect of urban land
504 surface forcing to regional meteorology and air quality over South China. *Atmospheric Environment* 152,
505 389-404.

506

507

508

509 Tables and Figures


510

511 Table 1. Meteorological perturbations imposed in this study

| Meteorological Parameter | Changes in Values Examined |
|------------------------------|---|
| Temperature (T) | $\pm 0.5\text{K}$, $\pm 1.0\text{K}$, $\pm 1.5\text{K}$ |
| Wind speed (WS) | $\pm 5\%$, $\pm 10\%$, $\pm 20\%$ |
| Absolute Humidity (AH) | $\pm 5\%$, $\pm 10\%$, $\pm 20\%$ |
| Boundary layer height (PBLH) | $\pm 10\%$, $\pm 20\%$, $\pm 30\%$ |
| Cloud liquid content (CLW) | $\pm 5\%$, $\pm 10\%$, $\pm 20\%$ |
| Precipitation (PCP) | $\pm 5\%$, $\pm 10\%$, $\pm 20\%$ |


512

513



514

515 **Fig.1** Location map of China and the five cities.




516

517 **Fig.2** Spatial distributions of monthly average O_3 -8 h (ppb) in (a) January and (b) July, and
 518 monthly average $PM_{2.5}$ ($\mu g m^{-3}$) in (c) January and (d) July 2013.


519

520




521

522 **Fig.3** Changes in monthly average O₃-8h (ppb) in January and July, 2013 due to (a) T+1.0K, (b)
 523 WS-10%, (c) AH+10%, (d) PBLH-20%, (e) CLW+10%, and (f) PCP+10%.



524

525 **Fig.4** Changes in monthly average $\text{PM}_{2.5}$ concentration ($\mu\text{g m}^{-3}$) in January and July, 2013 due to
 526 (a) T+1.0K, (b) WS-10%, (c) AH+10%, (d) PBLH-20%, (e) CLW+10%, and (f) PCP+10%.



527


528 **Fig.5** Changes in monthly average $PM_{2.5}$ component concentration ($\mu g m^{-3}$) in January due to (a)

529 T+1.0K, (b) WS-10%, (c) AH+10%, (d) PBLH-20%, (e) CLW+10%, and (f) PCP+10%.

530

531

532




533

534 **Fig.6** Changes in monthly average PM_{2.5} component concentration (µg m⁻³) in July due to (a)

535 T+1.0K, (b) WS-10%, (c) AH+10%, (d) PBLH-20%, (e) CLW+10%, and (f) PCP+10%.


536



537

538

539 **Fig.7** Sensitivity of O₃-8h, PM_{2.5} and its components to meteorological parameter of (a) T, (b) WS,
 540 (c) AH, (d) PBLH, (e) CLW, and (f) PCP in five cities in China. The unit of sensitivity is ppb K⁻¹
 541 for O₃-8h to T, and is ppb %⁻¹ for O₃-8h to other meteorological parameters; and the unit is μg m⁻³
 542 K⁻¹ for PM_{2.5} and its components to T, and is μg m⁻³ %⁻¹ for PM_{2.5} and its components to other
 543 meteorological parameters.




545

546

547

Fig.8 Sensitivity of O_3 -8h mean to meteorological perturbations (a) T, (b) WS, (c) AH, (d) PBLH, (e) CLW, (f) PCP in China. The value in T is measured in $ppb K^{-1}$, and others is $ppb \%^{-1}$.



548

549 **Fig.9** Sensitivity of $PM_{2.5}$ mean to meteorological perturbations (a) T, (b) WS, (c) AH, (d) PBLH,

550 (e) CLW, (f) PCP in China. The value in T is measured in $\mu g m^{-3} K^{-1}$, and others is $\mu g m^{-3} \%^{-1}$.

551

552