



1 **Sensitivity Analysis of the Surface Ozone and Fine**
2 **Particulate Matter to Meteorological Parameters in China**

3 Zhihao Shi¹, Lin Huang¹, Jingyi Li¹, Qi Ying², Hongliang Zhang^{3,4}, Jianlin Hu^{1*}

4
5 ¹Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution
6 Control, Collaborative Innovation Center of Atmospheric Environment and
7 Equipment Technology, Nanjing University of Information Science & Technology,
8 Nanjing 210044, China

9 ²Zachry Department of Civil and Environmental Engineering, Texas A&M University,
10 College Station, TX 77843, USA

11 ³Department of Environmental Science and Engineering, Fudan University, Shanghai
12 200438, China

13 ⁴Institute of Eco-Chongming (SIEC), Shanghai 200062, China

14
15 *Corresponding authors:

16 Jianlin Hu, Email: jianlinhu@nuist.edu.cn. Phone: +86-25-58731504.



17 Abstract

18 Meteorological conditions play important roles in the formation of ozone (O₃)
19 and fine particulate matter (PM_{2.5}). China has been suffering from serious regional air
20 pollution problems, characterized by high concentrations of surface O₃ and PM_{2.5}. In
21 this study, the Community Multiscale Air Quality (CMAQ) model was used to
22 quantify the sensitivity of surface O₃ and PM_{2.5} to key meteorological parameters in
23 different regions of China. Six meteorological parameters were perturbed to create
24 different meteorological conditions, including temperature (T), wind speed (WS),
25 absolute humidity (AH), planetary boundary layer height (PBLH), cloud liquid water
26 content (CLW) and precipitation (PCP). Air quality simulations under the perturbed
27 meteorological conditions were conducted in China in January and July of 2013. The
28 changes in O₃ and PM_{2.5} concentrations due to individual meteorological parameters
29 were then quantified. T has the greatest impact on the daily maximum 8-h average O₃
30 (O₃-8h) concentrations, which leads to O₃-8h increases by 1.7 ppb K⁻¹ in January in
31 Chongqing and 1.1 ppb K⁻¹ in July in Beijing. WS, AH, and PBLH have a smaller but
32 notable influence on O₃-8h with maximum change rates of 0.3, -0.15, and 0.14
33 ppb %⁻¹, respectively. T, WS, AH, and PBLH have important effects on
34 PM_{2.5} formation of in both January and July. In general, PM_{2.5} sensitivities are
35 negative to T, WS, and PBLH and positive to AH in most regions of China. The
36 sensitivities in January are much larger than in July. PM_{2.5} sensitivity to T, WS, PBLH,
37 and AH in January can be up to -5 μg m⁻³ K⁻¹, -3 μg m⁻³ %⁻¹, -1 μg m⁻³, and +0.6 μg
38 m⁻³ %⁻¹, respectively, and in July can be up to -2 μg m⁻³ K⁻¹, -0.4 μg m⁻³ %⁻¹, -0.14 μg
39 m⁻³ %⁻¹, and +0.3 μg m⁻³ %⁻¹, respectively. Other meteorological factors (CLW and
40 PCP) have negligible effects on O₃-8h (less than 0.01 ppb %⁻¹) and PM_{2.5} (less than
41 0.01 μg m⁻³ %⁻¹). The results suggest that surface O₃ and PM_{2.5} concentrations can



42 change significantly due to changes in meteorological parameters and it is necessary
43 to consider these effects when developing emission control strategies in different
44 regions of China.

45

46 Keywords: sensitivity, meteorological conditions, fine particulate matter, ozone,

47 CMAQ model



48 1. Introduction

49 China has serious air pollution problems and fine particulate matter (PM_{2.5}) and
50 ozone (O₃) are the two major air pollutants (Lin et al., 2010; Hu et al., 2016; Lu et al.,
51 2019; Wu et al., 2019). The annual average PM_{2.5} concentrations were higher than 50
52 µg m⁻³ in 26 out of the total 31 provincial capital cities in mainland China during
53 2013-2014 (Wang et al., 2014a), and the national 4th highest daily maximum 8-hour
54 average O₃ (O₃-8h) is 86.0 ppb during the warm-seasons (April–September) in
55 2013-2017, which is 6.3–30% higher than that in other industrialized regions of the
56 world (Lu et al., 2018). PM_{2.5} alone caused 0.87-1.36 million deaths every year in
57 China, and long-term exposure to O₃ was responsible for an extra 254 000 deaths
58 (Apte et al., 2015; Cohen et al., 2017; Hu et al., 2017b; Silver et al., 2018). China has
59 made remarkable improvement in air quality during recent years (Zhang et al., 2017;
60 Zhao et al., 2017; China, 2018; Zheng et al., 2018), however, air pollution is still
61 severe, making it the fourth-ranked healthy risk factor (Stanaway et al., 2018).

62 Surface PM_{2.5} and O₃ concentrations are determined by atmospheric processes of
63 emissions, transport and dispersion, chemical transformation (due to gas-phase,
64 aqueous-phase and aerosol chemistry), and dry and wet deposition. These processes
65 are affected by meteorological conditions. Studies have shown that the surface O₃ and
66 PM_{2.5} concentrations are sensitive to different meteorological parameters. For
67 example, Dawson et al. (2007b) have investigated the sensitivity of surface O₃ to
68 different meteorological parameters in the eastern United States (US) using the
69 comprehensive air quality model with extensions (CAM_x). The results showed that



70 temperature (T) had the greatest influence on daily O_3 -8h of 0.34 ppb K^{-1} , followed by
71 absolute humidity (AH) of $0.025 \text{ ppb \%}^{-1}$. Bernard et al. (2001) also confirmed that T
72 presented a notable positive correlation with the surface O_3 concentration. The effects
73 of meteorological parameters on $PM_{2.5}$ are even more complicated. Tran and Mölders
74 (2011) showed that elevated $PM_{2.5}$ concentrations tended to occur under the condition
75 of calm wind, low T and relative humidity in Fairbanks, Alaska. Olvera Alvarez et al.
76 (2018) used a land use regression model to analyze the effects of different
77 meteorological parameters on $PM_{2.5}$ in El Paso, Texas and obtained the same
78 conclusion in winter, but in spring, the high $PM_{2.5}$ level was associated with high wind
79 speed (WS) and low humidity. Dawson et al. (2007a) studied the effects of individual
80 meteorological parameters in the Eastern US and found that $PM_{2.5}$ concentration
81 decreased markedly as the increased precipitation (PCP) in winter, but in summer, the
82 main meteorological factors affecting the $PM_{2.5}$ concentration were T, WS and
83 planetary boundary layer height (PBLH). Dawson et al. (2009) simulated the effects
84 of climate change on regional and urban air quality in the Eastern US, and found
85 $PM_{2.5}$ concentration decreased by $0.3 \mu\text{g m}^{-3}$ in January mostly due to increasing in
86 PCP and increased by $2.5 \mu\text{g m}^{-3}$ in July largely due to decreasing in PBLH and WS.
87 Horne and Dabdub (2017) altered various meteorological parameters to investigate
88 their effects on O_3 , $PM_{2.5}$ and secondary organic aerosols (SOA), and found that the T
89 predominated the effects of meteorology in California.

90 Many studies have proved that meteorological conditions play very important
91 roles in air pollution events in China. Studies found that the pollutant concentrations



92 could vary up to several times, due to meteorological changes with the same emission
93 sources (Zhang et al., 2010; Xing et al., 2011; Zheng et al., 2015; Cai et al., 2017; Liu
94 et al., 2017; Ning et al., 2018; Yang et al., 2018; Li et al., 2019b). For example, Xing
95 et al. (2011) studied that the difference between the effects of 2007 and 2008
96 meteorological conditions on air quality during the 2008 Beijing Olympics. They
97 found higher humidity in August 2008 was beneficial to the formation of SO_4^{2-} by up
98 to ~60%, and lower T prevented the evaporation of NO_3^- by up to ~60%. Liu et al.
99 (2017) reported that the monthly mean $\text{PM}_{2.5}$ concentrations in the Jing-Jin-Ji (JJJ)
100 area in December 2015 increased by 5%~137% due to the unfavorable weather
101 conditions such as low WS and high humidity.

102 A few studies investigated the relationships between air quality and
103 meteorological conditions in China. Zhang et al. (2015) conducted a correlation
104 analysis between air quality and meteorology in three megacities Beijing, Shanghai
105 and Guangzhou in China. The result showed that air pollutants were significantly
106 negatively correlated with WS, and O_3 had a positive correlation with T. Yin et al.
107 (2016) found that the relationship between WS and $\text{PM}_{2.5}$ has complicated influence,
108 with higher PM at low and high WS than in light to moderate winds in Beijing from
109 2008 to 2014. Xu et al. (2018) examined the variations of $\text{PM}_{2.5}$ concentration in
110 January 2017 in China compared to that in January 2016 and found meteorological
111 conditions of low WS, high humidity, low PBLH and low PCP contributed to $\text{PM}_{2.5}$
112 concentration worsening by 29.7%, 42.6% and 7.9% in the JJJ region, the Pearl River
113 Delta (PRD) region and the Cheng-Yu Basin (CYB) region, respectively. Ma et al.



114 (2019) analyzed the effects of meteorology on air pollution in the Yangtze River Delta
115 (YRD) region during 2014-2016 and found $PM_{2.5}$ was highly negatively correlated to
116 WS, while O_3 concentration was positively correlated to T but negatively related to
117 relative humidity. Zhu et al. (2017) reported that the surface concentrations of O_3
118 increased by 2-6 ppb in January and 8-12 ppb in July 2014 in PRD, mainly due to the
119 increase in T and the decrease in NO_x emissions.

120 These studies have investigated the impacts of meteorological conditions on
121 $PM_{2.5}$ and O_3 in certain regions of China, however, quantitative sensitivity of $PM_{2.5}$
122 and O_3 to meteorological parameters has not been examined. The objective of this
123 study is to quantify the sensitivity of O_3 and $PM_{2.5}$ to different meteorological
124 parameters in winter and summer in different regions of China. The paper is
125 constructed as following, Section 2 describes the method used to estimate the
126 sensitivity, and Section 3 presents the effects of each meteorological variable on O_3
127 and $PM_{2.5}$ in China and in five representative cities. Conclusions are then summarized
128 in Section 4.

129

130 2. Methods

131 The sensitivity of O_3 and $PM_{2.5}$ associated with changes in meteorological
132 parameters was quantified using the Community Multiscale Air Quality (CMAQ)
133 model version 5.0.2. The meteorological parameters include T, WS, AH, PBLH, PCP,
134 and cloud liquid water content (CLW). A base case was firstly simulated with
135 meteorological fields predicted by the Weather Research and Forecasting (WRF)
136 model v3.7.1 (<http://www.wrf-model.org/>) using the NCEP FNL Operational Model



137 Global Tropospheric Analyses dataset as the initial and boundary conditions. The base
138 case has been described in a previous study and the model configurations of the base
139 case were reported there (Hu et al., 2015). The WRF predicted meteorological
140 parameters and the CMAQ predicted surface O₃ and PM_{2.5} have been evaluated
141 against observations at 422 sites in 60 major cities in China, and the accuracy of the
142 model performance has been validated (Hu et al., 2016).

143 A suit of perturbation scenarios was created, and in each scenario, a certain
144 meteorological parameter was perturbed to a certain extent. The details of the
145 perturbation scenarios are listed in Table 1. Among those changes, the T was absolute
146 changes, and other parameters are relative variations. Then the CMAQ model was
147 re-run to predict the air quality under the perturbed meteorological condition. The
148 emissions and other inputs were kept unchanged in each perturbed meteorological
149 scenarios, therefore the difference of O₃ and PM_{2.5} concentrations between each of the
150 perturbation case and the base case was due to the change in the specific
151 meteorological parameter, and the sensitivity of O₃ and PM_{2.5} to individual
152 meteorological parameters could be quantitatively determined.

153 The modeling domain covers East Asia, including entire China, with a horizontal
154 resolution of 36 × 36 km². The base case and perturbation cases were conducted in
155 January and July in 2013, representing the winter and the summer conditions,
156 respectively. In addition to the regional analysis, five representative megacities were
157 selected, i.e., Beijing, Shanghai, Guangzhou, Chongqing, Xi'an (Fig.1). These cities
158 are located in the North China Plain (NCP), YRD, PRD, CYB, and Guanzhong Plain,



159 respectively, where serious air pollution problems often occur. In this study, O₃-8h
160 was used in the O₃ analyses, and 24h average PM_{2.5} was used in the PM_{2.5} analyses, if
161 not specifically stated.

162

163 **3. Results and Discussion**

164 **3.1 Impacts of meteorological parameters on surface O₃**

165 Figs. 2(a) and 2(b) show the spatial-distribution of the predicted monthly average
166 O₃-8h concentrations in January and July, respectively. In January, the highest average
167 concentrations are about 70 ppb in the Sichuan Basin, and the concentrations in
168 southern and eastern China are generally higher than those in northern China. In July,
169 the highest average concentrations are over 80ppb in the large areas of NCP and YRD,
170 CYB, and Guangzhou areas in the PRD.

171 Fig. 3 shows the spatial distribution of the concentration changes of O₃-8h in
172 January and July due to change of T + 1.0 K, WS - 10%, AH +10% PBLH -
173 20%, CLW + 10% case, and PCP + 10%, respectively. Fig. S1-S3 shows the results
174 due to other extent changes in these parameters. When T increases 1.0 K (Fig. 3(a)),
175 O₃-8h increases 1-2 ppb in most area of eastern and central China in January and in
176 NCP and YRD in July, which is consistent with the high O₃ spatial distribution in the
177 base case (shown in Fig. 2). O₃-8h decreases up to 4 ppb in January in Northeast
178 China, and up to 2 ppb in the Southwest border of China and the East China Sea,
179 which are the areas of low O₃ concentrations (generally less than 35 ppb). Therefore,
180 the effect of T on O₃ is dependent on the O₃ formation regime. An increase in T
181 promotes O₃ formation chemistry in net O₃ formation areas, but accelerates O₃



182 consumption chemistry in the net O₃ loss areas.

183 Fig. 3(b) shows that the differences of O₃-8h in January and July when WS is 10%
184 less than the base case in 2013. The influence of wind on O₃ concentration is complex,
185 but generally, slower WS decreases O₃ in January in most parts of China, particularly
186 in Sichuan by up to 3 ppb, but increases O₃ in July by a few ppb over the most areas
187 in eastern and central China. Therefore, the impact of WS on O₃ appears opposite in
188 winter and summer. Weaker winds slow down the dispersion of NO_x and VOCs,
189 which is conducive to O₃ formation in summer when the vertical mixing is strong, but
190 increases O₃ titration in the surface in winter due to weaker vertical mixing.

191 Fig. 3(c) displays that the surface O₃ is expected to decrease generally less than 1
192 ppb when AH increases by 10% (relative change) both in January and July in most
193 land areas of China except in the northeast area. Fig. 3(d) shows that a 20% decrease
194 of PBLH leads to O₃-8h decreases by a few ppb in most area in January, while in July
195 O₃-8h increases in eastern and central regions, especially in YRD, CYB and areas in
196 Hubei-Hunan-Jiangxi in the central China. Sensitivity of O₃ to CLW and PCP is
197 relatively small. Fig. 3(e) demonstrates that O₃-8h changes -0.03 ppb to 0.03 ppb in
198 January and July for a 10% increase in CLW. Fig. 3(f) demonstrates that a 10%
199 increase in PCP results in -0.1 to 0.2 ppb changes in O₃-8h. O₃ changes due to the six
200 meteorological factors with different extents of perturbation (Figs. S1-S3) shows the
201 similar trends and spatial patterns.

202 **3.2 Impacts of meteorological parameters on surface PM_{2.5}**

203 Figs. 2(c) and 2(d) show the spatial distribution of the monthly average surface



204 $PM_{2.5}$ concentrations in January and July. $PM_{2.5}$ in January reaches over $200 \mu\text{g m}^{-3}$ in
205 JJJ, SYB, central China, and urban areas in the Northeast China. $PM_{2.5}$ is much lower
206 in July, generally lower than $50 \mu\text{g m}^{-3}$, but is high (up to $70 \mu\text{g m}^{-3}$) in areas in the JJJ,
207 YRD, and central China regions.

208 Fig. 4 shows the spatial distribution of $PM_{2.5}$ changes due to the same changes of
209 meteorological factors, as in Fig. 3. The $PM_{2.5}$ results of other cases of the sensitivity
210 study are shown in Figs. S4-S6 of the Supplementary Materials. The results indicate
211 that in January, a 1.0 K increase in T leads to up to $5\text{-}6 \mu\text{g m}^{-3}$ decrease of $PM_{2.5}$ in JJJ
212 and central China; in July, a 1.0 K increase in T causes $PM_{2.5}$ increase by about $1 \mu\text{g}$
213 m^{-3} in southern China but decrease by $1\text{-}3 \mu\text{g m}^{-3}$ in the JJJ and east coast region. A 10%
214 decrease in WS causes $PM_{2.5}$ increase up to over $40 \mu\text{g m}^{-3}$ in January and up to $5 \mu\text{g}$
215 m^{-3} in July. A 10% relative increase in AH leads to $PM_{2.5}$ increase of up to $6 \mu\text{g m}^{-3}$ in
216 January and up to $2 \mu\text{g m}^{-3}$ in JJJ and northeast regions but slightly decrease of less
217 than $1 \mu\text{g m}^{-3}$ in southern China in July. A 20% decrease of PBLH causes $PM_{2.5}$
218 increases by up to $20 \mu\text{g m}^{-3}$ in January and up to $4 \mu\text{g m}^{-3}$ in July. The impact of
219 CLW and PCP on $PM_{2.5}$ is small, and generally increase in CLW increases surface
220 $PM_{2.5}$ and increase in PCP decreases $PM_{2.5}$.

221 The changes in the total $PM_{2.5}$ mass concentrations are determined by the
222 changes in the chemical components of $PM_{2.5}$. Fig. S7 displays the fraction of $PM_{2.5}$
223 species (elemental carbon (EC), primary organic carbon (POC), secondary organic
224 aerosol (SOA), sulfate (SO_4^{2-}), nitrate (NO_3^-), and ammonium (NH_4^+)) in five
225 representative cities. Secondary inorganic aerosols (SO_4^{2-} , NO_3^- , NH_4^+) are the major



226 PM components, accounting for over 50% of $PM_{2.5}$ in January and about 40% in July.
227 Fig. 5 and Fig. 6 show the changes of the major $PM_{2.5}$ components due to the same
228 changes of meteorological factors as in Fig. 4 in January and in July, respectively. The
229 results show that the effects of the meteorological parameters on the total $PM_{2.5}$
230 (shown in Fig. 4) are mainly due to their effects on SO_4^{2-} , NO_3^- , and NH_4^+ in January,
231 and due to the changes in SO_4^{2-} , NO_3^- , NH_4^+ , and SOA in July. In general, PBLH, WS,
232 and PCP are negatively correlated to SO_4^{2-} , NO_3^- , and NH_4^+ formation, but AH and
233 CLW are positively correlated to these components. SOA concentrations are much
234 higher in July than in January due to the contribution from biogenic emissions (Hu et
235 al., 2017a). SOA formation is affected by reaction rates (positively affected by T),
236 availability of oxidants (such as changes in O_3), and hydrogen ion strength (affected
237 by changes in SO_4^{2-} , NO_3^- , NH_4^+). SOA concentrations mainly increase in south
238 China.

239 It is worthwhile noting that the effects of T on SO_4^{2-} and NO_3^- (changes of NH_4^+
240 is determined by changes of SO_4^{2-} and NO_3^-). Both in January and July, increase in T
241 decreases SO_4^{2-} and NO_3^- in the major areas of eastern China. The NO_3^- decrease is
242 expected because volatile NH_4NO_3 favors more in gas phase in higher temperature,
243 and this result is consistent with studies in other regions (Dawson et al., 2007a; Horne
244 and Dabdub, 2017). SO_4^{2-} is found to increase with T increase in those studies because
245 faster gas- and aqueous- phase reactions of SO_4^{2-} . However, our finding of SO_4^{2-} in
246 China is opposite. The CMAQ-Sulfur Tracking Model (CMAQ-STM) was further
247 used to track the SO_4^{2-} formation from different processes. The results confirm that



248 the SO_4^{2-} production from gas- and aqueous- phase increases with T increase. But
249 meanwhile SO_4^{2-} production from heterogeneous reactions is reduced more when T is
250 increased. Heterogeneous SO_4^{2-} formation has been proposed as a major SO_4^{2-}
251 formation pathway during China haze events (Wang et al., 2014b; Gen et al., 2019;
252 Huang et al., 2019; Li et al., 2019a) and in this study it accounts for up to ~75% of
253 total SO_4^{2-} production. The treatment of heterogeneous SO_4^{2-} formation currently is
254 modeled as a surface-controlled uptake process, in which the formation rate is
255 determined by the aerosol surface area and the uptake coefficient of SO_2 on particle
256 surface (Ying et al., 2014). When T is increased, the particle surface area decreases (as
257 particle mass concentration decreases due to a combined effect of other components),
258 resulting in decrease in the heterogeneous SO_4^{2-} formation.

259

260 **3.3. Quantitative sensitivity of O_3 and $\text{PM}_{2.5}$ to individual** 261 **meteorology parameters**

262 The quantitative sensitivity of O_3 and $\text{PM}_{2.5}$ concentrations to individual
263 meteorological parameter is calculated by linear fitting of the changes in
264 monthly-average concentrations under all of the six perturbed cases of the
265 meteorological parameter. Figs. S8-S10 in Supplementary Materials show the
266 calculation examples of T, WS, and AH on O_3 at the five major cities of Beijing,
267 Chongqing, Guangzhou, Shanghai and Xi'an, and Figs. S11-S13 shows the examples
268 for the $\text{PM}_{2.5}$ cases. Fig.7 demonstrates the sensitivities of O_3 -8h and $\text{PM}_{2.5}$ and its
269 components to each meteorological parameter in the five cities. In January, T has a



270 positive impact on O_3 in all cities, and the largest impact is in Chongqing with a rate
271 of $+1.69 \text{ ppb K}^{-1}$. In July, O_3 also shows a strong positive sensitivity to T in Beijing
272 with $+1.06 \text{ ppb K}^{-1}$ and in Shanghai with $+0.98 \text{ ppb K}^{-1}$, but has a small negative
273 sensitivity (-0.15 ppb K^{-1}) in Xi'an and a moderate negative sensitivity (-0.74 ppb K^{-1})
274 in Guangzhou. The O_3 sensitivity to T in Guangzhou in July shows a highly nonlinear
275 trend and is very different from other cities (Fig. S8(c)). More studies are needed to
276 investigate the effects of T on O_3 pollution in the YRD region during summertime.
277 WS and PBLH both have positive effects on O_3 -8h in January, the effects vary
278 significantly among cities, with 0.004 - 0.3 ppb \%^{-1} for WS and 0.04 - 0.14 ppb \%^{-1} for
279 PBLH. AH has a negative effect on O_3 -8h in January, ranging from -0.01 to -0.15
280 ppb \%^{-1} . But in July, the impacts of WS, AH, and PBLH are negative in most cities,
281 with a range of -0.05 to -0.18 , -0.05 to -0.13 , and -0.02 to $-0.07 \text{ ppb \%}^{-1}$, respectively.
282 Generally speaking, the sensitivity of O_3 to T is obviously higher than that of WS, AH,
283 and PBLH. The sensitivity of O_3 to CLW and PCP is even minimal (less than 0.01
284 ppb \%^{-1}) and mostly negative.

285 Negative sensitivities are found for surface $PM_{2.5}$ concentrations to T, WS, PBLH,
286 and PCP, and positive sensitivities for $PM_{2.5}$ to AH and CLW. The sensitivity of T in
287 the five cities ranges from -1.5 to $-3.6 \mu\text{g m}^{-3} \text{ K}^{-1}$ in January and -0.3 to $-1.65 \mu\text{g m}^{-3}$
288 K^{-1} in July. $PM_{2.5}$ is also very sensitive to WS in January, with a range of -0.8 to -2.97
289 $\mu\text{g m}^{-3} \%^{-1}$, while the sensitivity (-0.03 to $-0.19 \mu\text{g m}^{-3} \%^{-1}$) becomes much smaller in
290 July. The sensitivity to PBLH is -0.12 to $-0.58 \mu\text{g m}^{-3} \%^{-1}$ in January and -0.003 to
291 $-0.23 \mu\text{g m}^{-3} \%^{-1}$ in July. The sensitivity to AH is 0.16 to $0.30 \mu\text{g m}^{-3} \%^{-1}$ in January



292 and 0.05 to $0.27 \mu\text{g m}^{-3} \%^{-1}$ in July. Sensitivity to CLW and PCP is small in January
293 and July, mostly less than $0.01 \mu\text{g m}^{-3} \%^{-1}$. The $\text{PM}_{2.5}$ sensitivities can be explained by
294 the major components of SO_4^{2-} , NO_3^- , and NH_4^+ in January and by SO_4^{2-} , NO_3^- , NH_4^+ ,
295 and SOA in July.

296 Fig. 8 shows the spatial variations of the sensitivity of O_3 -8h and $\text{PM}_{2.5}$ to the
297 meteorological parameters. The sensitivity of O_3 -8h to temperature is more significant
298 in Sichuan and southern provinces of China in January, and in NCP and YRD in July,
299 up to $+2 \text{ ppb K}^{-1}$ in both January and July. O_3 -8h sensitivity to WS is diverse in space,
300 and is generally positive in Sichuan and southern provinces in January; and it is
301 negative in east China but positive in west China. O_3 -8h sensitivity to AH is generally
302 negative in both months in most regions of China, except the northeast in January and
303 southwest in July. O_3 -8h sensitivity to PBLH is mostly positive in January but
304 becomes negative in YRD, CYB, NCP, and central China in July. O_3 -8h sensitivity to
305 CLW and PCP is negligible.

306 Fig. 9 displays the spatial variations of the sensitivity of surface $\text{PM}_{2.5}$ to the
307 meteorological parameters. $\text{PM}_{2.5}$ sensitivities to the meteorological parameters are
308 more consistent in January and July than the cases of O_3 , i.e., negative sensitivity to T,
309 WS, PBLH, PCP, and positive to AH and CLW in most regions of China in both
310 months. On the other hand, $\text{PM}_{2.5}$ sensitivities are more profound in January than in
311 July. $\text{PM}_{2.5}$ sensitivity to T is up to $-5 \mu\text{g m}^{-3} \text{K}^{-1}$ in January and up to $-2 \mu\text{g m}^{-3} \text{K}^{-1}$ in
312 July. $\text{PM}_{2.5}$ sensitivity to WS is up to $-3 \mu\text{g m}^{-3} \%^{-1}$ in January, and up to $-0.4 \mu\text{g}$



313 $\text{m}^{-3} \%^{-1}$ in July. $\text{PM}_{2.5}$ sensitivity to PBLH is up to $-1 \mu\text{g m}^{-3} \%^{-1}$ in January, and up to
314 $-0.14 \mu\text{g m}^{-3} \%^{-1}$ in July. $\text{PM}_{2.5}$ sensitivity to AH is up to $+0.6 \mu\text{g m}^{-3} \%^{-1}$ in January,
315 and up to $0.3 \mu\text{g m}^{-3} \%^{-1}$ in July. The sensitivities to CLW and PCP is small, compared
316 to the other four meteorological parameters. $\text{PM}_{2.5}$ sensitivity to T is negative in most
317 land areas of China in January and in NCP and YRD in July because of the negative
318 effects of T on SO_4^{2-} , NO_3^- , and NH_4^+ , as discussed in previous sections. $\text{PM}_{2.5}$
319 sensitivity to T is positive in south China in July due to more SOA with higher T.

320 **4. Conclusions**

321 Meteorological conditions can have a great influence on surface O_3 and $\text{PM}_{2.5}$
322 concentrations. In this study, the sensitivities of O_3 -8h and $\text{PM}_{2.5}$ to T, WS, AH, PBLH,
323 PCP, and CLW are quantitatively estimated in January and July, respectively in China.
324 O_3 -8h is most sensitive to T and the sensitivity can be up to $+2 \text{ppb K}^{-1}$ in both
325 January and July, and the sensitivity is dependent on the O_3 chemistry formation or
326 loss regime, i.e., positive in the net O_3 formation areas, and negative in the O_3
327 consumption areas. In general, $\text{PM}_{2.5}$ sensitivities are negative to T, WS, PBLH, and
328 PCP and positive to AH and CLW in most regions of China in both January and July.
329 The sensitivities in January are much larger than in July. $\text{PM}_{2.5}$ sensitivities to T, WS,
330 AH, and PBLH are important. The $\text{PM}_{2.5}$ sensitivities to these meteorological
331 parameters are through major effects on SO_4^{2-} , NO_3^- , NH_4^+ , and SOA. The
332 sensitivities of O_3 and $\text{PM}_{2.5}$ to CLW and PCP are negligible. The results show that O_3
333 and $\text{PM}_{2.5}$ concentrations in China are greatly affected by meteorological conditions,
334 therefore changes in these meteorological parameters due to climate change or



335 inter-annual meteorological variations could potentially alter O₃ and PM_{2.5}
336 concentrations significantly, and it should consider these effects when developing
337 emission control strategies. The results also show that the O₃ and PM_{2.5} sensitivities to
338 meteorological parameters have substantial spatial variations. Future studies can
339 further investigate how the changes in meteorological conditions affect the
340 effectiveness of emission control plans in reaching the designed air quality objectives
341 in the different regions of China.

342

343 **Data availability.** All of the modeling results will be available online after we
344 publish the paper.

345 **Author contributions.** ZS, and JH designed research. ZS, LH, JL, QY, HZ
346 and JH contributed to model development and configuration. ZS, LH, JL, and JH
347 analyzed the data. ZS prepared the manuscript and all coauthors helped improve the
348 manuscript.

349 **Competing interests.** The authors declare that they have no conflict of
350 interest.

351 **Acknowledgment**

352 This work was supported by the National Key R&D Program of China
353 (2018YFC0213800), the National Natural Science Foundation of China (41975162,
354 41675125 and 41705102), and Jiangsu Environmental Protection Research Project
355 (2016015).

356



357 **References**

- 358 Apte, J.S., Marshall, J.D., Cohen, A.J., Brauer, M., 2015. Addressing global mortality from ambient PM2.
359 5. *Environmental science & technology* 49, 8057-8066.
- 360 Bernard, S.M., Samet, J.M., Grambsch, A., Ebi, K.L., Romieu, I., 2001. The potential impacts of climate
361 variability and change on air pollution-related health effects in the United States. *Environmental*
362 *health perspectives* 109, 199-209.
- 363 Cai, W., Li, K., Liao, H., Wang, H., Wu, L., 2017. Weather conditions conducive to Beijing severe haze
364 more frequent under climate change. *Nature Climate Change* 7, 257.
- 365 China, 2018. Air Quality Targets Set by the Action Plan Have Been Fully Realized(2018).
- 366 Cohen, A.J., Brauer, M., Burnett, R., Anderson, H.R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef,
367 B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y.,
368 Martin, R., Morawska, L., Pope, C.A., Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R.,
369 van Donkelaar, A., Vos, T., Murray, C.J.L., Forouzanfar, M.H., 2017. Estimates and 25-year trends of the
370 global burden of disease attributable to ambient air pollution: an analysis of data from the Global
371 Burden of Diseases Study 2015. *The Lancet* 389, 1907-1918.
- 372 Dawson, J., Adams, P., Pandis, S., 2007a. Sensitivity of PM 2.5 to climate in the Eastern US: a modeling
373 case study. *Atmospheric chemistry and physics* 7, 4295-4309.
- 374 Dawson, J.P., Adams, P.J., Pandis, S.N., 2007b. Sensitivity of ozone to summertime climate in the
375 eastern USA: A modeling case study. *Atmospheric environment* 41, 1494-1511.
- 376 Dawson, J.P., Racherla, P.N., Lynn, B.H., Adams, P.J., Pandis, S.N., 2009. Impacts of climate change on
377 regional and urban air quality in the eastern United States: Role of meteorology. *Journal of*
378 *Geophysical Research: Atmospheres* 114.
- 379 Gen, M.S., Zhang, R.F., Huang, D.D., Li, Y.J., Chan, C.K., 2019. Heterogeneous SO2 Oxidation in Sulfate
380 Formation by Photolysis of Particulate Nitrate. *Environmental Science & Technology Letters* 6, 86-91.
- 381 Horne, J.R., Dabdub, D., 2017. Impact of global climate change on ozone, particulate matter, and
382 secondary organic aerosol concentrations in California: A model perturbation analysis. *Atmospheric*
383 *Environment* 153, 1-17.
- 384 Hu, J., Wang, P., Ying, Q., Zhang, H., Chen, J., Ge, X., Li, X., Jiang, J., Wang, S., Zhang, J., Zhao, Y., Zhang,
385 Y., 2017a. Modeling biogenic and anthropogenic secondary organic aerosol in China. *Atmos. Chem.*
386 *Phys.* 17, 77-92.
- 387 Hu, J., Wu, L., Zheng, B., Zhang, Q., He, K., Chang, Q., Li, X., Yang, F., Ying, Q., Zhang, H., 2015. Source
388 contributions and regional transport of primary particulate matter in China. *Environ Pollut* 207, 31-42.
- 389 Hu, J.L., Chen, J.J., Ying, Q., Zhang, H.L., 2016. One-year simulation of ozone and particulate matter in
390 China using WRF/CMAQ modeling system. *Atmospheric Chemistry and Physics* 16, 10333-10350.
- 391 Hu, J.L., Huang, L., Chen, M.D., Liao, H., Zhang, H.L., Wang, S.X., Zhang, Q., Ying, Q., 2017b. Premature
392 Mortality Attributable to Particulate Matter in China: Source Contributions and Responses to
393 Reductions. *Environmental Science & Technology* 51, 9950-9959.
- 394 Huang, L., An, J.Y., Koo, B., Yarwood, G., Yan, R.S., Wang, Y.J., Huang, C., Li, L., 2019. Sulfate formation
395 during heavy winter haze events and the potential contribution from heterogeneous SO2 + NO2
396 reactions in the Yangtze River Delta region, China. *Atmospheric Chemistry and Physics* 19,
397 14311-14328.
- 398 Li, M.M., Wang, T.J., Xie, M., Li, S., Zhuang, B.L., Huang, X., Chen, P.L., Zhao, M., Liu, J.E., 2019a.
399 Formation and Evolution Mechanisms for Two Extreme Haze Episodes in the Yangtze River Delta



- 400 Region of China During Winter 2016. *J Geophys Res-Atmos* 124, 3607-3623.
- 401 Li, X., Gao, Z., Li, Y., Gao, C.Y., Ren, J., Zhang, X., 2019b. Meteorological conditions for severe foggy haze
402 episodes over north China in 2016–2017 winter. *Atmospheric Environment* 199, 284-298.
- 403 Lin, J., Nielsen, C.P., Zhao, Y., Lei, Y., Liu, Y., McElroy, M.B., 2010. Recent Changes in Particulate Air
404 Pollution over China Observed from Space and the Ground: Effectiveness of Emission Control.
405 *Environmental Science & Technology* 44, 7771-7776.
- 406 Liu, T., Gong, S., He, J., Yu, M., Wang, Q., Li, H., Liu, W., Zhang, J., Li, L., Wang, X., Li, S., Lu, Y., Du, H.,
407 Wang, Y., Zhou, C., Liu, H., Zhao, Q., 2017. Attributions of meteorological and emission factors to the
408 2015 winter severe haze pollution episodes in China's Jing-Jin-Ji area. *Atmospheric Chemistry and*
409 *Physics* 17, 2971-2980.
- 410 Lu, H., Lyu, X., Cheng, H., Ling, Z., Guo, H., 2019. Overview on the spatial–temporal characteristics of
411 the ozone formation regime in China. *Environmental Science: Processes & Impacts*.
- 412 Lu, X., Hong, J., Zhang, L., Cooper, O.R., Schultz, M.G., Xu, X., Wang, T., Gao, M., Zhao, Y., Zhang, Y.,
413 2018. Severe Surface Ozone Pollution in China: A Global Perspective. *Environmental Science &*
414 *Technology Letters* 5, 487-494.
- 415 Ma, T., Duan, F., He, K., Qin, Y., Tong, D., Geng, G., Liu, X., Li, H., Yang, S., Ye, S., Xu, B., Zhang, Q., Ma, Y.,
416 2019. Air pollution characteristics and their relationship with emissions and meteorology in the
417 Yangtze River Delta region during 2014–2016. *Journal of Environmental Sciences* 83, 8-20.
- 418 Ning, G., Wang, S., Yim, S.H.L., Li, J., Hu, Y., Shang, Z., Wang, J., Wang, J., 2018. Impact of low-pressure
419 systems on winter heavy air pollution in the northwest Sichuan Basin, China. *Atmospheric Chemistry*
420 *and Physics* 18, 13601-13615.
- 421 Olvera Alvarez, H.A., Myers, O.B., Weigel, M., Armijos, R.X., 2018. The value of using seasonality and
422 meteorological variables to model intra-urban PM_{2.5} variation. *Atmospheric Environment* 182, 1-8.
- 423 Silver, B., Reddington, C.L., Arnold, S.R., Spracklen, D.V., 2018. Substantial changes in air pollution
424 across China during 2015–2017. *Environmental Research Letters* 13, 114012.
- 425 Stanaway, J.D., Afshin, A., Gakidou, E., Lim, S.S., Abate, D., Abate, K.H., Abbafati, C., Abbasi, N.,
426 Abastabar, H., Abd-Allah, F.J.T.L., 2018. Global, regional, and national comparative risk assessment of
427 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195
428 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study
429 2017. 392, 1923-1994.
- 430 Tran, H.N.Q., Mölders, N., 2011. Investigations on meteorological conditions for elevated PM_{2.5} in
431 Fairbanks, Alaska. *Atmospheric Research* 99, 39-49.
- 432 Wang, Y., Ying, Q., Hu, J., Zhang, H., 2014a. Spatial and temporal variations of six criteria air pollutants
433 in 31 provincial capital cities in China during 2013–2014. *Environ Int* 73, 413-422.
- 434 Wang, Y.X., Zhang, Q.Q., Jiang, J.K., Zhou, W., Wang, B.Y., He, K.B., Duan, F.K., Zhang, Q., Philip, S., Xie,
435 Y.Y., 2014b. Enhanced sulfate formation during China's severe winter haze episode in January 2013
436 missing from current models. *J Geophys Res-Atmos* 119.
- 437 Wu, C., Hu, W., Zhou, M., Li, S., Jia, Y., 2019. Data-driven regionalization for analyzing the
438 spatiotemporal characteristics of air quality in China. *Atmospheric Environment* 203, 172-182.
- 439 Xing, J., Zhang, Y., Wang, S., Liu, X., Cheng, S., Zhang, Q., Chen, Y., Streets, D.G., Jang, C., Hao, J., Wang,
440 W., 2011. Modeling study on the air quality impacts from emission reductions and atypical
441 meteorological conditions during the 2008 Beijing Olympics. *Atmospheric Environment* 45, 1786-1798.
- 442 Xu, Y., Xue, W., Lei, Y., Zhao, Y., Cheng, S., Ren, Z., Huang, Q., 2018. Impact of Meteorological
443 Conditions on PM_{2.5} Pollution in China during Winter. *Atmosphere* 9, 429.



444 Yang, Y., Zheng, X., Gao, Z., Wang, H., Wang, T., Li, Y., Lau, G.N., Yim, S.H., 2018. Long - term trends of
445 persistent synoptic circulation events in planetary boundary layer and their relationships with haze
446 pollution in winter half year over eastern China. *Journal of Geophysical Research: Atmospheres* 123,
447 10,991-911,007.

448 Yin, Q., Wang, J., Hu, M., Wong, H., 2016. Estimation of daily PM2.5 concentration and its relationship
449 with meteorological conditions in Beijing. *Journal of Environmental Sciences* 48, 161-168.

450 Ying, Q., Cureño, I.V., Chen, G., Ali, S., Zhang, H., Malloy, M., Bravo, H.A., Sosa, R., 2014. Impacts of
451 Stabilized Criegee Intermediates, surface uptake processes and higher aromatic secondary organic
452 aerosol yields on predicted PM2.5 concentrations in the Mexico City Metropolitan Zone. *Atmospheric
453 Environment* 94, 438-447.

454 Zhang, H., Wang, Y., Hu, J., Ying, Q., Hu, X.-M., 2015. Relationships between meteorological
455 parameters and criteria air pollutants in three megacities in China. *Environmental Research* 140,
456 242-254.

457 Zhang, J., Reid, J.S., Alfaro-Contreras, R., Xian, P., 2017. Has China been exporting less particulate air
458 pollution over the past decade? *Geophysical Research Letters* 44, 2941-2948.

459 Zhang, L., Liao, H., Li, J., 2010. Impacts of Asian summer monsoon on seasonal and interannual
460 variations of aerosols over eastern China. *Journal of Geophysical Research: Atmospheres* 115.

461 Zhao, B., Jiang, J.H., Gu, Y., Diner, D., Worden, J., Liou, K.-N., Su, H., Xing, J., Garay, M., Huang, L., 2017.
462 Decadal-scale trends in regional aerosol particle properties and their linkage to emission changes.
463 *Environmental Research Letters* 12.

464 Zheng, B., Chevallier, F., Ciais, P., Yin, Y., Deeter, M.N., Worden, H.M., Wang, Y., Zhang, Q., He, K., 2018.
465 Rapid decline in carbon monoxide emissions and export from East Asia between years 2005 and 2016.
466 *Environmental Research Letters* 13.

467 Zheng, X., Fu, Y., Yang, Y., Liu, G., 2015. Impact of atmospheric circulations on aerosol distributions in
468 autumn over eastern China: observational evidence. *Atmospheric Chemistry and Physics* 15, 12115.

469 Zhu, K., Xie, M., Wang, T., Cai, J., Li, S., Feng, W., 2017. A modeling study on the effect of urban land
470 surface forcing to regional meteorology and air quality over South China. *Atmospheric Environment*
471 152, 389-404.

472
473
474



475 Tables and Figures


476

477 Table 1. Meteorological perturbations imposed in this study

Meteorological Parameter	Changes in Values Examined
Temperature (T)	$\pm 0.5\text{K}$, $\pm 1.0\text{K}$, $\pm 1.5\text{K}$
Wind speed (WS)	$\pm 5\%$, $\pm 10\%$, $\pm 20\%$
Absolute Humidity (AH)	$\pm 5\%$, $\pm 10\%$, $\pm 20\%$
Boundary layer height (PBLH)	$\pm 10\%$, $\pm 20\%$, $\pm 30\%$
Cloud liquid content (CLW)	$\pm 5\%$, $\pm 10\%$, $\pm 20\%$
Precipitation (PCP)	$\pm 5\%$, $\pm 10\%$, $\pm 20\%$


478

479



480

481 **Fig.1** Location map of China and the five cities.




482

483 **Fig.2** Spatial distributions of monthly average O_3 -8 h (ppb) in (a) January and (b) July, and
484 monthly average $PM_{2.5}$ ($\mu g m^{-3}$) in (c) January and (d) July 2013.

485


486



487

488 **Fig.3** Changes in monthly average O₃-8h (ppb) in January and July, 2013 due to (a) T+1.0K, (b)


489 WS-10%, (c) AH+10%, (d) PBLH-20%, (e) CLW+10%, and (f) PCP+10%.



490


491 **Fig.4** Changes in monthly average $\text{PM}_{2.5}$ concentration ($\mu\text{g m}^{-3}$) in January and July, 2013 due to

492 (a) T+1.0K, (b) WS-10%, (c) AH+10%, (d) PBLH-20%, (e) CLW+10%, and (f) PCP+10%.



493
494
495
496
497
498


Fig.5 Changes in monthly average $\text{PM}_{2.5}$ component concentration ($\mu\text{g m}^{-3}$) in January due to (a) T+1.0K, (b) WS-10%, (c) AH+10%, (d) PBLH-20%, (e) CLW+10%, and (f) PCP+10%.



499

500 **Fig.6** Changes in monthly average $PM_{2.5}$ component concentration ($\mu\text{g m}^{-3}$) in July due to (a)
501 T+1.0K, (b) WS-10%, (c) AH+10%, (d) PBLH-20%, (e) CLW+10%, and (f) PCP+10%.

502




503
 504

505 **Fig.7** Sensitivity of O_3 -8h, $PM_{2.5}$ and its components to meteorological parameter of (a) T, (b) WS,
 506 (c) AH, (d) PBLH, (e) CLW, and (f) PCP in five cities in China. The unit of sensitivity is $ppb\ K^{-1}$
 507 for O_3 -8h to T, and is $ppb\ \%^{-1}$ for O_3 -8h to other meteorological parameters; and the unit is $\mu g\ m^{-3}$
 508 K^{-1} for $PM_{2.5}$ and its components to T, and is $\mu g\ m^{-3}\ \%^{-1}$ for $PM_{2.5}$ and its components to other
 509 meteorological parameters.




510



511

512 **Fig.8** Sensitivity of O_3 -8h mean to meteorological perturbations (a) T, (b) WS, (c) AH, (d) PBLH,

513 (e) CLW, (f) PCP in China. The value in T is measured in $ppb K^{-1}$, and others is $ppb \%^{-1}$.



514
 515 **Fig.9** Sensitivity of $PM_{2.5}$ mean to meteorological perturbations (a) T, (b) WS, (c) AH, (d) PBLH,
 516 (e) CLW, (f) PCP in China. The value in T is measured in $\mu g m^{-3} K^{-1}$, and others is $\mu g m^{-3} \%^{-1}$.