

Measurement Report: Aircraft Observations of Ozone, Nitrogen Oxides, and Volatile Organic Compounds over Hebei Province, China

Sarah E. Benish¹, Hao He¹, Xinrong Ren^{1,2}, Sandra J. Roberts³, Ross J. Salawitch^{1,3}, Zhanqing Li^{1,4}, Fei Wang^{4,5}, Yuying Wang⁶, Fang Zhang⁴, Min Shao⁷, Sihua Lu⁷, Russell R. Dickerson¹

¹ Department of Atmospheric and Oceanic Science, University of Maryland, College Park, MD 20742, USA.

² Air Resources Laboratory, National Oceanic and Atmospheric Administration, College Park, MD 20740, USA.

³ Department of Chemistry and Biochemistry, University of Maryland, College Park, MD 20742, USA.

⁴ State Key Laboratory of Earth Surface Processes and Resource Ecology, College of Global Change and Earth System Science, Beijing Normal University, Beijing, 100875, China.

⁵ Key Laboratory for Cloud Physics, Chinese Academy of Meteorological Sciences, Beijing, 100081, China.

⁶ Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, School of Atmospheric Physics, Nanjing University of Information Science and Technology, Nanjing, 21004, China.

⁷ College of Environmental Science and Engineering, Peking University, Beijing, 100871, China.

Correspondence to: Sarah E. Benish (sebenish@umd.edu)

Abstract. To provide insight into the planetary boundary layer (PBL) production of ozone (O₃) over the North China Plain, the Air chemistry Research in Asia (ARIAs) campaign conducted aircraft measurements of air pollutants over Hebei Province, China between May and June 2016. We evaluate vertical profiles of trace gas species including O₃, nitrogen oxides (NO_x), carbon monoxide (CO), and volatile organic compounds (VOCs) and relate to rates of O₃ production. This analysis shows measured O₃ levels ranged from 45 to 146 ppbv, with the peak median concentration (~92 ppbv) occurring between 1000 and 1500 m. The NO_x concentrations exhibited strong spatial and altitudinal variations, with a maximum of 53 ppbv. Ratios of CO/CO₂ indicate the prevalence of low efficiency combustion from biomass burning and residential coal burning, but indicate some success of regional pollution controls compared to earlier studies in China. Concentrations of total measured VOCs reveals alkanes dominate the total measured volume mixing ratio of VOCs (68%) and sources include vehicular emissions, fuel and solvent evaporation, and biomass burning. Alkanes and alkenes/alkynes are responsible for 74% of the total VOC reactivity assessed by calculating the OH loss rates, while aromatics contribute the most to the total Ozone Formation Potential (OFP) (43%) with toluene, m/p-xylene, ethylene, propylene, and i-pentane playing significant roles in the aloft production of O₃ in this region. In the PBL below 500 m, box model calculations constrained by measured precursors indicate the peak rate of mean O₃ production was ~7 ppbv/hour. Pollution frequently extended above the PBL into the lower free troposphere around 3000 m, where NO₂ mixing ratios (~400 pptv) led to net production rates of O₃ up to ~3 ppbv/hour; this pollution can travel substantial distances downwind. The O₃ sensitivity regime is determined to be NO_x-limited throughout the PBL, while more VOC-limited at low altitudes near urban centers, demonstrating that control of both VOCs and NO_x is needed to reduce aloft O₃ pollution over Hebei.

1 Introduction

Explosive urbanization and rapid industrialization contributed to high ground-level ozone (O₃) and particulate matter (PM) over the past several decades in the North China Plain (NCP) (Johnson et al., 2006; Ran et al., 2011; Shao et al., 2009; Zhang et al., 2014). Household burning of coal used for cooking and heating, emissions from gasoline, diesel, and liquified petroleum gas (LPG) vehicles, as well as large-scale burning of winter wheat residues in the NCP are some of the many sources responsible for O₃ precursors, such as nitrogen oxides (NO_x=NO+NO₂) and volatile organic compounds (VOCs) (Chen et al.,

2017; Long et al., 2016; Stavrakou et al., 2016). Ozone is harmful to both the human respiratory system (Bell et al., 2006; 40 Jerrett et al., 2009) and to photosynthetic processes by vegetation (Avnery et al., 2011; Reich and Amundson, 1985), while some VOCs, such as benzene and chloroform, are known to be hemotoxic and carcinogenic (Environmental Protection Agency - Integrated Risk Information System, 2003; Lan et al., 2004). Several studies using the NASA Ozone Monitoring Instrument (OMI) have found reductions of some pollutants like sulfur dioxide (SO₂) over the NCP (He et al., 2012; Krotkov et al., 2016; Li et al., 2010, 2017), but NO₂ pollution still remains severe in China (Figure 1a).

45 Ozone is created through the oxidation of NO by hydroperoxyl radicals (HO₂) and organic peroxy radicals (RO₂), products of carbon monoxide (CO) and VOC oxidation. When one of these precursors is the limiting reactant, the rate of O₃ production is considered VOC- or NO_x-sensitive (Finlayson-Pitts and Pitts, 1999; Sillman et al., 1990). The role of VOCs on the formation of O₃ depends on the characteristics of the environment, including the main emission sources of primary pollutants and ambient temperature (Pusede et al., 2014), and the interaction of aerosols within the PBL to reduce photolysis (An et al., 2019). High 50 aerosol concentrations have been shown to decrease photolysis and hinder summer surface O₃ formation by 25 ppbv on average in Xi'an, China (Feng et al., 2016), which pose a challenge for pollution control strategies.

Natural emissions are the largest source of VOCs globally and react more efficiently with OH than most anthropogenic compounds (Di Carlo et al., 2004), but exhibit a strong seasonal, diurnal, and spatial dependence (Li et al., 2013). Biogenic VOCs have been found to play a significant role in the formation of O₃ at the surface (Ma et al., 2019; Zong et al., 2018) and 55 throughout the boundary layer in the NCP (Wang et al., 2008a), as well as influence production of PM_{2.5} (Guo et al., 2014) and secondary organic aerosols (SOA) (Wu et al., 2020b). In particular, isoprene has been estimated to account for 27% of the total O₃ production in June 2010 in Beijing (Mo et al., 2018), suggesting the need to consider biogenic isoprene emissions in formulating O₃ control strategies. Quantifying the abundance of NO_x and the suite of VOC chemicals throughout the lower troposphere is urgently needed to better understand the photochemistry of O₃ production in the NCP, which in turn will lead 60 to the development of successful mitigation strategies.

In-situ airborne measurements provide valuable information regarding the horizontal and vertical distributions of air pollutants over a large spatial area. Airborne measurements are necessary to characterize air pollution over large cities, as well as surrounding areas. Ozone and PM are produced throughout the planetary boundary layer (PBL), so aircraft observations can lead to a more complete picture of pollution formation and transport than is available only from surface observations. While 65 several airborne campaigns have deployed to investigate the regionally transported pollution problem in East Asia, including the NASA Korea-United States Air Quality Study (KORUS-AQ) (Al-Saadi et al., (2015), see https://www-air.larc.nasa.gov/missions/korus-aq/docs/White_paper_NASA_KORUS-AQ.pdf), that occurred at the same time as our measurements, few airborne studies characterize the source region of severe smog within the Hebei Province region of China.

Through Chinese/American partnerships with Peking University, Beijing Normal University, and the University of Maryland, 70 we conducted a field campaign in Hebei Province, China called Air chemistry Research In Asia (ARIAs). The ARIAs

campaign was designed to characterize and quantify the composition of trace gases and aerosol optical properties over Hebei to improve tools used to evaluate the effectiveness of air pollution reduction policies. Since air pollution transport from Asia typically peaks in early to mid-spring (Liu et al., 2003), we hoped to provide detailed altitude profiles over the Asian source region to enable Lagrangian experiments with KORUS-AQ, but only two sustained transport events occurred (Peterson et al., 2019). Despite the infrequent transboundary pollution events, ARIAs observations generated valuable characteristic pollution signatures that helped describe combustion efficiency and its impact downwind (Halliday et al., 2019) to correct model biases of CO in global chemistry-climate models (Gaubert et al., 2020), and to show that MOPITT bias increases at high CO concentrations (Tang et al., 2020). Furthermore, ARIAs measurements characterized aerosol optical properties in the planetary boundary layer and free troposphere during clean and polluted conditions (Wang et al., 2018a), as well as used in the validation of MAX-DOAS profiles of NO₂, SO₂, HONO, HCHO, CHOCHO, and aerosols (Wang et al., 2019b).

The Ministry of Environmental Protection of the People's Republic of China reported six of the top ten cities with the worst air quality in 2016 were located in Hebei (including the capital city of Shijiazhuang). The North China Plain is one of the most polluted regions in the world, but implementation of pollution reduction measures through the Five-Year Plans has allowed for decreasing trends of many pollutants. In particular, Zhang et al. (2020) found an increased number of days of clean/light haze and a decreased number of days with heavy haze, along with a significant decline of SO₂ concentrations. Similarly, using observations from MODIS and OMI, Si et al. (2019) found AOD and SO₂ to decrease from 2006 to 2015, while NO₂ rose by 4.79% in the NCP during this period. While surface NO₂ decreased 20% from May 2014 to December 2018 throughout China, there are still a large number of measurement stations with increasing trends of NO₂ due to changes in meteorological conditions and aerosol emissions (Fan et al., 2020), illustrating the need for more research characterizing air pollution in this region. In this study, we analyze concentrations of O₃, NO_x, NO_y, CO, and VOCs obtained during 11 research flights between May and June 2016. The VOC chemical reactivity and impact on O₃ production is assessed and we utilize an observation-constrained box model to evaluate photochemical properties of the production of O₃ that occurs throughout the lower free troposphere.

2 Materials and Methods

2.1 Air sampling and analysis

The ARIAs campaign included 11 research flights from May-June 2016 in Hebei Province (Fig. 1b). Flight days were chosen based on meteorological conditions associated with smog events, such as higher temperatures, little cloud cover, low relative humidity, weak winds, and shallow PBL height (Tang et al., 2012). The Y-12 aircraft was based at Luancheng Airport (114.59°E, 37.91°N, 58 m above sea level (ASL)), located in southeast Shijiazhuang (population around 10 million), a major economic center in Hebei, including pharmaceutical and textile industries, machinery and chemical manufacturing, construction, and electronics production. Flight sampling occurred east of the Taihang Mountains and the Y-12 flew vertical

spirals from ~300 m to ~3500 m over Shijiazhuang as well as three other locations: Julu (115.02°E, 37.22°N, 30 m ASL), Quzhou (114.96°E, 36.76°N, 40 m ASL), and Xingtai (114.36°E, 37.18°N, 182 m ASL) (see Table S1 for a description of flight paths, weather conditions, and statistics of measured trace gases).

05 Various instruments aboard the Y-12 aircraft collected trace gas, aerosol, and meteorological data. The aircraft instrumentation (Table 1) included different gas and particle sample inlets on the top of the fuselage and pressure/temperature/humidity sensors (Cloud Water Inertial Probe (CWIP), Rain Dynamics) installed under one wing of the aircraft (Fig. S1). Flight position data were recorded using a portable global positioning system (GPS) and the CWIP. The aircraft was equipped with the following trace gas analyzers: (1) a Picarro cavity ring down spectrometer (CRDS) for measurements of CH₄, CO₂, CO, and H₂O; (2) a
10 Thermal Electron Corporation (TECO) Model 49C UV absorption O₃ analyzer; (3) a TECO Model 43C pulsed fluorescence SO₂ analyzer; (4) a Los Gatos Research Model RMT-200 CRDS NO₂ analyzer; and (5) a TECO Model 42C NO-NO_y analyzer. Power constraints and a converter issue led to limited NO_y, NO_x, and CO measurements during the campaign, particularly in the lower free troposphere (LFT). We remove observations of NO/NO_y over three spiral locations due to limited measurements. Negative values indicate readings around the detection limit, usually at high altitudes. The aircraft was also equipped with an
15 inlet to measure aerosols up to ~5.0 μm diameter and aerosol optical properties, including a nephelometer (TSI Model 3563) to measure aerosol scattering, a particle soot absorption photometer (PSAP) to measure aerosol absorption, and an aethalometer (Magee Model AE31) and a Single-Particle Soot Photometer (SP2, Droplet Measurement Technologies) to measure black carbon. Observed aerosol optical properties have been summarized by F. Wang et al. (2018); further details on aircraft instrumentation are given by Ren et al. (2018).

20 Twenty-six whole air samples (WAS) were collected directly into 3.2 L fused silica lined electropolished stainless steel canisters (Entech Instrument Inc., Simi Valley, CA) at a variety of pressure altitudes from 400 m to 3500 m between 1:30 and 9:00 UTC (9:30 and 17:00 local time). The sampling period for the WAS canisters was approximately 1-2 minutes during the spirals. Samples were analyzed for 54 VOCs and 16 halocarbons. Since the halogenated species have negligible effects on O₃ production, we exclude these species from the analysis presented here. We also exclude 2 WAS canisters from this analysis
25 due to evidence of contamination after collection (Text S1). Limited samples collected over one province in one season may not be able to represent O₃ chemistry for all of China, but the scarcity of airborne VOC measurements in this region makes these data valuable for characterizing the composition of air throughout and above the PBL, demonstrating how the production of O₃ aloft differs from that at the surface.

The VOC analytical techniques used by the College of Environmental Sciences and Engineering at Peking University (PKU)
30 in Beijing have been summarized in the past (Mo et al., 2015; Wang et al., 2010a), and we briefly describe the method here. The WAS canisters were cleaned following a standard sampling procedure, pressured with nitrogen and vacuumed three times to 2.6 Pa. The hydrocarbons were quantified using a gas chromatograph equipped with a mass selective detector (GC-MSD, Hewlett Packard 5975/7890, USA) and a flame ionization detector (FID) coupled with a cryofocusing pre-concentration system (Entech Instrument 7100A, Simi Valley, CA). This system used a Dean Switch™ (Agilent Technologies, Santa Clara, CA,

35 USA) to introduce the effluent into a DB-624 column (60 m × 0.25 mm × 1.8 μm; J&W Scientific, Folsom, CA, USA) with
an MSD to separate and analyze C4–C12 hydrocarbons and halocarbons. A PLOT (Al/KCl) column (30 m × 0.25 mm × 3.0
μm; J&W Scientific) with an FID was used to measure the C2–C4 hydrocarbons.

The Photochemical Assessment Monitoring Stations (PAMS) (55 NMHCs) and Toxic Organic-15 (TO-15) standard mixtures
were used to calibrate the GC-MSD/FID system that measured the C2–C12 VOCs. Samples with known concentrations of
40 four VOCs (bromochloromethane, 1,4-difluorobenzene, chlorobenzene-d5 and 1-bromo-3-fluorobenzene) were used as
internal standards for each sample to calibrate the system. The GC-MSD/FID system was calibrated at five concentrations,
ranging from 0.5 to 8 ppbv, for each of these four compounds before sample analysis. Correlation coefficients, ranging from
0.987 to 0.999 showed that the integral areas of peaks were proportional to the concentrations of the target compounds. A gas
standard (diluting from 1 ppmv to 2 ppbv) was measured each day to check the stability of the system. Summary statistics of
45 the VOCs along with the method detection limit (MDL) (ranging from 0.002 to 0.027 ppbv) are reported in Table S2. Total
uncertainty for VOC measurements reflects instrument noise, plus uncertainty in calibration standards, contamination, and
pressurization. Best estimate of the total uncertainty is ±20% with 95% confidence due to uncertainties associated with airborne
sampling platforms. Intercomparison experiments of VOC measurements between PKU and other laboratories showed good
agreement (Liu et al., 2008b).

50 The Atmosphere-Aerosol-Boundary Layer-Cloud (A²BC) Interaction Joint Experiment campaign collected meteorological,
aerosol, and trace gas information from a ground-based site in Xingtai (114.36°E, 37.18°N, 182 m ASL) from May to
December 2016 (Wang et al., 2018b, 2019b). An intensive observation period in May and June 2016 was conducted to coincide
with ARIAs. Data from A²BC instruments used in our analysis include: (1) a NO_x analyzer with a molybdenum converter
(Ecotech model 9841A); (2) an infrared absorption CO analyzer (Ecotech model 9830A); and (3) a UV absorption O₃ analyzer
55 (Ecotech model 9810A). Results of NO₂, SO₂, HONO, HCHO, CHOCHO, and aerosols derived from the Differential Optical
Absorption Spectrometer (DOAS) are summarized by Yang Wang et al., (2019). The A²BC site is located in northwest Xingtai,
nestled in the east foothill of the Taihang Mountains. Agricultural crops surround the site, consisting heavily of winter wheat,
that is harvested, with the stubble burned in June (Liu and Si, 2011). Xingtai is a city with approximately 7 million people and
is surrounded by industry including coal mining and coal-burning power plants, cement and steel industries, chemical
60 processing, iron-smelting, and glass manufacturing.

2.2 Box Model Simulations

A box model called Framework in 0-Dimensional Atmospheric Modelling (F0AMv3.1) (Wolfe et al., 2016) is used to evaluate
oxidation processes to understand O₃ photochemical production both at the surface and aloft. The box model simulations cover
the Y-12 flight tracks during seven flights and daytime hours at the A²BC supersite in Xingtai (where the Y-12 conducted
65 spirals) using the Carbon Bond Mechanism, version 6, revision 2 (CB6r2). Both the Y-12 flights and surface simulations define
a physical loss lifetime of 24 hours to mitigate build-up of long-lived oxidation products over multiple days of integration.

For the ARIAs flight data, the model is constrained by 1-minute average observed concentrations of VOCs, NO₂, CO, and O₃. Due to the limited number of grab canisters per flight, VOCs are constrained based on the altitude of the sampling relative to the height of the PBL, which is determined using potential temperature and water vapor vertical profiles for each flight. All WAS canister data collected below the top of the PBL during a flight are averaged. Data from all of the WAS canisters for the entire campaign collected above the research flight's PBL are averaged for that flight. Periodic missing Y-12 NO₂ data due to internal auto-zeroing are linearly interpolated since gaps were short (~2 minutes). The chemical system defined by each set of observations is integrated 5 days forward in time, in 1-hour time steps with diurnal variation of solar zenith angle (SZA), in order for calculated reactive intermediates to achieve diel steady state. Reaction rate constants are calculated using aircraft measurements of pressure, temperature, and relative humidity. The SZA is determined based on the time and location of the aircraft, and used to calculate photolysis rates as described below.

For the A²BC surface data, the model is constrained by 5-minute average concentrations of VOCs, NO₂, CO, and O₃ on days that a flight occurred. For May 17, surface data for NO₂ is filled with 1-hour average data collected for other days of the month, due to missing surface measurements on this day. The average concentrations from the WAS canisters below 500 m are used as ground concentrations since A²BC did not measure VOCs at the surface. Similar to the flight data, the chemical system for the surface observations is integrated for 3 days forward in time, in 1-hour time steps with time-varying SZA, to reach diel steady state. Reaction rate constants are calculated from ground measurements of pressure, temperature, and relative humidity. Time and ground elevation are used to calculate the SZA, which controls photolysis frequencies as described below.

Photolysis frequencies, not measured during ARIAs or at the A²BC supersite, evolve over the course of a model step and are calculated by combining cross sections and quantum yields with solar spectra derived from the NCAR Tropospheric Ultraviolet and Visible (TUV) version 5.2 radiation model. At the start of the model run, input solar zenith angle, altitude or elevation, O₃ column, and surface albedo are used for linear interpolation across TUV lookup tables (F0AM's "hybrid" method). We use SZA and altitude/elevation from ARIAs/A²BC measurements and constant values for ozone column (325 DU) and surface albedo (0.17), which we estimate based on concurrent data from the OMI level-3 OMDOAO3e data product (https://disc.gsfc.nasa.gov/datasets/OMDOAO3e_003/summary?keywords=OMDOAO3e_003). A correction factor of 0.8, determined by trial and error, is used to scale j-values to better agree with the observed NO/NO₂ ratio.

The impact of aerosols on O₃ production depends on the optical properties as well as the vertical distribution (Dickerson et al., 1997; Kelley et al., 1995). In the presence of scattering and absorbing aerosols, photolysis frequencies will be altered, thus changing the O₃ formation and atmospheric oxidizing capability (Wu et al., 2020a). Previous research over China has shown that as AOD increases, the extinction effect of aerosols on photolysis frequencies decreases due to a higher proportion of scattering aerosols under high AOD conditions (Wang et al., 2019a). Optical depth, single scattering albedo, and angstrom exponent during ARIAs (see Wang et al., 2018a) are used in the TUV online calculator (https://cprm.acom.ucar.edu/Models/TUV/Interactive_TUV/) to assess the impact of aerosols on photolysis frequencies. Most of the aerosol particles during ARIAs were concentrated in the lowest 2 km of the atmosphere with a single scattering albedo

00 at 550 nm of 0.85 and an average AOD ~0.2. The impact of aerosol optical properties measured during ARIAs on photolysis frequencies is small compared to the default setting, so no additional adjustments are made to the model values.

The method described here to constrain VOCs introduces large uncertainty due to the sparsity of measurements obtained over a large area that potentially consists of a wide variety of chemical compositions. However, the production of O₃ aloft is not well characterized over Hebei, so our observations may help improve the understanding of air pollution for this region, despite
05 this limitation. Additionally, unlike a 3-dimensional chemical transport model, the box model simulations do not include advection or emissions. These processes, while important, are not included in the box model since O₃ precursors were measured and used to constrain the box model calculations. Box modelling is used to gain an understanding of O₃ production and its sensitivity to ambient levels of NO_x and VOCs based upon measured meteorological parameters and the concentration of a wide variety of chemical species.

10 2.2.1 Ozone Production and Sensitivity Calculations

The photochemical production of O₃ during the daytime is determined by the production rate of NO₂ molecules from the HO₂+NO and RO₂+NO reactions minus the loss mechanisms (Finlayson-Pitts and Pitts, 1999). Thus, the net O₃ production rate, net(P(O₃)) can be estimated following Equation 1:

$$\begin{aligned} net(P(O_3)) = & k_{HO_2+NO}[HO_2][NO] + \sum_{i=1}^n k_{RO_{2i}+NO}[RO_{2i}][NO] - P(RONO_2) \\ & - k_{OH+NO_2+M}[OH][NO_2][M] - k_{HO_2+O_3}[HO_2][O_3] - k_{OH+O_3}[OH][O_3] \\ & - k_{O(^1D)+H_2O}[O(^1D)][H_2O] - L(O_3 + alkenes) \end{aligned} \quad (1)$$

15

where k denotes the different reaction rate constants and RO_{2i} is the concentration of individual organic peroxy radicals. The terms subtracted from the production of O₃ are the loss mechanisms: the formation of nitrates, P(RONO₂), the reaction of OH and NO₂ to form nitric acid, the reactions of OH and HO₂ with O₃, the reaction of O(¹D) with H₂O, and the reactions of O₃ with alkenes. Additional terms not included here are the rate of O₃ loss by dry deposition and direct loss on aerosol surfaces
20 (dilution is the only physical loss in the current F0AM setup).

We evaluate the sensitivity of O₃ production to NO_x and VOCs using the ratio of L_N/Q, where L_N is the radical loss through reactions with NO and Q is the primary radical production (Kleinman, 2005a). When L_N/Q is much less than 0.5, the O₃ production regime is NO_x-limited; when L_N/Q ratio is much higher than 0.5, the regime is VOC-limited. Different environments can have varying amounts of organic nitrates that impact the cut-off value of L_N/Q, so this value could vary
25 around 0.5 (Kleinman, 2005b).

3 Results and discussion

3.1 Observations of nitrogen oxides, carbon monoxide, and ozone

Our observations confirm heavy loadings of air pollution over Hebei. Vertical profiles show peak median concentrations of NO (1.6 ppbv), NO₂ (4.4 ppbv) and NO_y (25.7 ppbv) below 500 m with large variability (Fig. 2). Median concentrations of NO and NO₂ drop off gradually with altitude, while median NO_y remains close to ~15 ppbv throughout most of the profile. Between 500 and 1000 m, sufficient levels of NO_x are observed (median=3.8 ppbv), indicating continued production of O₃ in the PBL. Above 3000 m, median concentrations of NO and NO₂ fall to 350 pptv and 106 pptv, respectively (not measured simultaneously), still sufficient to produce O₃ as air parcels travels downwind. Median mixing ratios of O₃ and CO remain high (~80 ppbv and ~120 ppbv, respectively) throughout the altitudes sampled by the Y-12.

Unlike previous airborne studies over Beijing from 1994-2005 that found increased O₃ concentrations below 1 km with constant levels (~52 ppbv) between 1 and 2 km (Ding et al., 2008), our O₃ concentrations peaked between 1000 and 1500 m (median = 91.6 ppbv). Low ratios of NO_x/NO_y (<0.30) indicate significant O₃ production had already occurred, but the strong correlation (R=0.71, Fig. S2) between 1-minute NO_z (NO_y-NO_x) and O_x (O₃+NO₂), an empirical estimate of the O₃ production efficiency (OPE), below 1500 m demonstrates moderate production of O₃ continued during sampling. The OPE of ~3.5 during ARIAs is smaller than the average OPE value of ~8 obtained during 2013 DISCOVER-AQ flights in Houston (Mazduca et al., 2016), likely due to the higher NO_x concentrations observed in Hebei than Texas.

Maps of O₃ and NO₂ on the Y-12 flight tracks (Fig. 3) show the largest concentrations around the spiral locations as well as between the three most northern cities, Shijiazhuang, Julu, and Xingtai. Regions of elevated NO₂ do not always correspond with high O₃ concentrations. The flight with the maximum observed NO₂ mixing ratio (35.3 ppbv) during ARIAs occurred on May 17 around 8:30 am LST. The aircraft was flying a flat transect at 500 m from Shijiazhuang to Julu when a large peak of NO₂, CO₂ (500 ppmv), and NO (15 ppbv) was sampled. Concentrations of O₃ were low during the time of the peak (~60 ppbv), indicating NO-O₃ titration, but O₃ levels were quite high (>90 ppbv) throughout the remainder of the flight. The maximum O₃ concentration (142.5 ppbv) was measured on May 21 during descent into Luancheng Airport in Shijiazhuang. Observations of NO_x were not available for this flight, but elevated CO concentrations (565 ppbv) were observed. High concentrations of O₃ were also observed away from the large megacities. For instance, an O₃ plume (~125 ppbv) was sampled on June 6 at 1500 m over a more suburban area between Shijiazhuang and Julu with NO₂ levels ~500 pptv.

Vertical profiles of trace gases over the four spiral locations (Fig. 4) generally show the highest concentrations over the two largest cities, Shijiazhuang and Xingtai. These two megacities exhibit the greatest variability, below 500 m altitude, of all trace gases discussed here below 500 m. At 3000 m, Xingtai demonstrates the most NO₂ (~800 pptv), while the other spiral locations show ~300 pptv. Median profiles of NO_y below 500 m are highest over Julu (27.6 ppbv). Median vertical profiles of CO are relatively consistent (~300 ppbv) below 2000 m over the spiral locations, while Julu shows the highest median concentration between 2500 and 3000 m (209.1 ppbv). Measurements of CO above Xingtai indicate a large spread in observations at all

altitudes from the lowest 500 m (10th percentile= 258 ppbv, 90th percentile=1049 ppbv) up to 2000 m (10th percentile=97.7 ppbv, 90th percentile = 135 ppbv). This variability may be partially explained by the possible burning of wheat straw during early summer 2016. Strong correlations between ethane and acetylene, two biomass burning markers (see Sect. 3.2), further suggest wheat residue burning over Xingtai. Median vertical profiles of O₃ below 500 m were 10-25 ppbv higher in Shijiazhuang (median=96.2 ppbv) than the other spiral locations. Concentrations of O₃ are generally stable or slightly increasing in the lowest 2000 m, and median O₃ is 75-80 ppbv even as high as 2500-3000 m. Xingtai shows the smallest variability of aloft O₃ levels above 2000 m, likely due to the position of this city on the leeward side of the Taihang Mountains.

The vertical profiles of O₃ compared to concurrent surface measurements in Xingtai indicates the A²BC site usually observed larger average concentrations than observed aloft, but this difference was highly dependent upon time of day (Fig. S3). The early afternoon profiles on May 8 showed average surface concentrations only slightly higher than the Y-12 measurements at ~400 m, while the mid-afternoon profiles on May 21 showed ~25 ppbv higher surface O₃ concentrations than Y-12 observations. At low altitudes (~700 m), the late morning flight (around 11:00 LST) on May 28 observed levels of O₃ ranging from 72-80 ppbv, comparable to average surface concentrations of 78 ppbv at the same time. By contrast, the afternoon flight (approximately 17:00 LST) at the same altitude later that day observed ~25 ppbv lower levels of O₃ compared to the surface (average=121 ppbv). All profiles on June 11 showed 10-30 ppbv lower average surface concentrations than measured during the Y-12 spirals.

The overall measured concentrations (1-second data, standard deviation, minimum, and maximum values) of NO_x, CO, and O₃ in this study are compared with other airborne studies in China including KORUS-AQ flights when outflow was directly from China (Table 2). Comparable to our range of NO_x levels from concentrations near the detection limit to 53.2 ppbv, autumn flights in the Yangtze River Delta in 2007 documented large variability in NO_x concentrations, ranging from 3 to 40 ppbv (Geng et al., 2009), while April 2006 observations in northern China similarly find a mean concentration ~5 ppbv (Wang et al., 2008b). The minimum CO concentration during ARIAs (80.5 ppbv) was measured in the lower free troposphere, which is a much smaller minimum concentration than reported by earlier studies. The warm-sector PBL air ahead of a cold front in April 2007 in Shenyang Province in northeast China found ~300 ppbv CO between 1000 and 4000 m (Dickerson et al., 2007), generally larger than most ARIAs profiles (except for July). The maximum value of CO during ARIAs (over 6 ppmv) agrees better with the literature, although there are few reported aircraft measurements of CO in Northeast Asia. Average and maximum O₃ concentrations during ARIAs were much higher than in other studies, but comparable to KORUS-AQ measurements from May 24-29 when the flow of air was direct from China. Since the majority of past airborne studies occurred over the sea areas during other seasons, it is not surprising that an urbanized environment like Hebei experienced much larger amounts of O₃ than previously reported.

The ratios between combustion tracers can be used to understand the source and efficiency. During high-efficiency combustion in modern power plants, fuel carbon is converted to CO₂ with near unit efficiency, resulting in low CO/CO₂ (<0.10%), while low-efficiency combustion (cold or smoldering processes or low-technology combustion) yields larger ratios. The regression

of 1-second CO against CO₂ (Fig. 5a) shows high linear correlation ($R=0.90$) and high ratios of CO/CO₂ (3.1%) together with large amounts of SO₂. These measurements are illustrative of low-efficiency fossil fuel combustion, likely from residential coal burning as these observations were all collected at ~500 m, and are compared to other studies in Table 3. Our results indicating the prevalence of low-efficiency combustion agree with KORUS-AQ airborne data over the West Sea with 2.8% CO/CO₂ (Tang et al., 2018), as well as with December 2017 surface measurements at Jingdezhen station in central China of 2.6% when air mass transport was from northern China (Xia et al., 2020). Compared to earlier studies in rural and urban areas of Beijing in the mid-2000s (Han et al., 2009; Wang et al., 2010b) and to 2011 measurements in Nanjing (Huang et al., 2015), the ARIAs CO/CO₂ ratio is 0.1-2.7% lower, evident of some success of regional pollution control strategies. By contrast, our CO/CO₂ ratio is higher than satellite-derived ratios over megacities that have implemented extensive pollution control measures (Silva et al., 2013). Similarly, compared to airborne measurements from the 2015 Wintertime INvestigation of Transport, Emissions, and Reactivity (WINTER) campaign in the Baltimore/Washington, D.C. region (Ren et al., 2018), our CO/CO₂ ratio is about a factor of 6 larger. Higher CO/CO₂ ratios (~6%) with less than 0.1 ppm SO₂, as seen briefly during three ARIAs flights, are more in line with emissions from burning of wheat straw in Hebei of ~6% (Cao et al., 2008), and other inefficient, biofuel combustion.

The $\Delta\text{CO}/\Delta\text{NO}_y$ ratio (equivalent to the slope in a CO vs. NO_y plot) (Fig. 5b) is an indicator for distinguishing plumes with efficient O₃ formation. Typical values of this ratio are ~40 in background air and between ~4-7 in fresh emissions plumes in Houston (Neuman et al., 2009). The $\Delta\text{CO}/\Delta\text{NO}_y$ ratio of 23.5 measured during ARIAs indicates some photochemical aging and contributions from fossil fuel or biomass burning, but high values of CO, NO_y, and SO₂ suggests sampling of air parcels heavily influenced by power plants. The CO/NO_x emission ratio (Fig. 5c) from ARIAs agrees with higher emission ratios of gasoline vehicles, while higher amounts of CO, NO_x, and SO₂ indicate coal burning from the residential sector or inefficient electric generating units. While most of these observations are reflective of the prevalence of low efficiency fossil fuel combustion, the aircraft sampled a plume on June 6 while flying spirals over Julu containing 0.9% CO/CO₂ and 0.4% SO₂/CO₂ (Fig. S4), likely due to a coal-burning power plant operating at high combustion efficiency, either using a sulfur scrubber or burning low sulfur fuel.

15 3.2 Observations and sources of VOCs

The total measured VOC mixing ratios ranged from 4 to 23 ppbv, largely dependent upon the altitude of collection, and was mostly dominated by alkanes (Fig. 6). The samples associated with the largest concentrations of O₃ were all collected at altitudes ~500 m during a period with stagnant high pressure. Generally, the samples collected below 500 m showed larger amounts of alkenes/alkynes and aromatics than canisters collected elsewhere in the PBL. The top VOCs ranked by mean volume mixing ratio (Table 4) shows that alkanes dominate the total measured VOC mixing ratio during ARIAs (68%), followed by alkenes/alkynes (17%), and aromatics (15%). The top 10 VOC species are C2-C5 alkanes, C2-C3 alkenes/alkynes,

benzene, and toluene. The observed mixing ratios of ethane and propane are 2.65 ppbv and 1.39 ppbv, respectively, which together accounts for ~52% of the total alkane mixing ratio.

25 The levels of ambient VOCs during ARIAs are generally lower than prior surface observations since measurements were taken in the PBL away from primary sources. Prior ground-based studies have similarly found alkanes to contribute the majority (>50%) of the total VOC concentration in late spring in the Beijing-Tianjin-Hebei region (Li et al., 2015; Tang et al., 2009; Yuan et al., 2013). The most abundant species during ARIAs are comparable to previous studies finding ethane, propane, and acetylene among the most prevalent, but likely have different sources based on the study location (Jia et al., 2016; Li et al., 2015; Mo et al., 2015; Tang et al., 2009). In the Beijing-Tianjin-Hebei region, ambient acetylene, ethylene, and other light alkanes have been attributed to emissions from gasoline vehicles (Li et al., 2015), while in Guangzhou, the widespread use of
30 LPG has resulted in high levels of propane (Guo et al., 2011). Additionally, our observations have higher amounts of branched alkanes, such as 2,2,4-trimethylpentane and 2-methylheptane (both components of gasoline), but lower amounts of isoprene due to collection over mostly urban regions with lower ambient temperatures than the summer months. Since isoprene with a lifetime of hours (Seinfeld and Pandis, 2006) in the summer typically exhibits a strong vertical gradient in the PBL (Huang et al., 2017), we find the mean amount of isoprene measured during ARIAs is about 7 times lower than average May 2014 surface
35 measurements in Beijing (Li et al., 2015), as well as ~200 pptv lower than June-July 2007 airborne measurements in the PBL in NE China (Xue et al., 2011). Next, we examine the potential sources contributing to observations of VOCs by comparing with ratios and correlations from known sources.

Since CO can be a marker for anthropogenically emitted hydrocarbons, particularly combustion products, we first use the
40 ratios of various VOCs to CO to reveal insight into changes in emissions in the region. Ratios of VOCs to CO can vary substantially among cities (Baker et al., 2008; Warneke et al., 2007), but in general can provide details about fuel types and combustion efficiency between metropolitan regions. Despite ARIAs measurements sampling in close proximity to local VOCs sources, most VOCs do not correlate strongly with CO, reflective of the lack of common source signatures and some photochemical aging of the sampled air masses. We report slopes of VOCs/CO in Table S2 when $R > 0.50$. Ethane has the
45 strongest correlation with CO ($R=0.72$) and the slope (2.5 pptv/ppbv) agrees well with ratios from urban areas of the United States in 1999-2005 (2.4 pptv/ppbv) (Baker et al., 2008) as well as with charcoal burning emission ratios (Andreae and Merlet, 2001). The ARIAs emission ratio of benzene/CO (1.8 pptv/ppbv) is slightly higher than found in urban regions of the United States (0.7, Baker et al., 2008) and Mexico City (0.93-1.20, Apel et al., 2010), likely due to higher emissions by widespread combustion of coal and agricultural residues (Zhang et al., 2015). By contrast, the ARIAs emission ratios of ethylene and
50 acetylene to CO (2.9 and 1.4 pptv/ppbv, respectively) are lower than observed in urban areas in the United States (4.1 and 3.4 pptv/ppbv, respectively) and Mexico City (7.90-8.40 and 8.20-9.60 pptv/ppbv, respectively), where the dominant source was reported to be transportation-related (Baker et al., 2008). The lower ratio of ethylene/CO is comparable to emission ratios reported from charcoal burning (2.3 pptv/ppbv) (Andreae and Merlet, 2001).

55 Ethane is the most abundant VOC in this study and correlates well with indicators for biomass and coal burning ($R>0.81$), such as acetylene, ethylene, benzene, and SO_2 . The ratio of acetylene to ethane (Fig. 7a) during ARIAs is 0.59, comparable to the ratio found in a plume of fresh biomass burning in Canada (Blake et al., 1994) and within the range of crop residue burning (~0.2-0.6) found in other studies in China (Chen et al., 2017). High ratios of benzene/propane (1.12) are comparable to dry grass combustion samples collected in the central Pearl River Delta (PRD) (1.6) (Wang et al., 2005) and further confirm the presence of VOCs due to biomass burning.

60 The C3 and C4 alkanes, including propane and the butanes, are the three main components of LPG and their correlation acts as an indicator for LPG leakage. In this study, a moderate correlation ($R\sim 0.50$) is found between n-butane and propane and i-butane with n-butane. The ratio of n-butane/propane during ARIAs is 0.60, which agrees well with ratios from vehicle emissions (Liu et al., 2008a), but is lower than slopes measured in the PRD (2.1) (Lai et al., 2009), where VOCs originated from LPG leakage. Additionally, propane correlates well with acetylene and ethylene (Figure 7a), two well-known vehicular emission tracers.

65 Since acetylene and propane have comparable photochemical lifetimes with respect to OH attack, the ratio can be used to assess the relative importance of fossil fuel combustion and LPG leakage (Goldan et al., 2000). LPG contains propane but not acetylene (acetylene/propane <1) while combustion of fossil fuels commonly produces small amounts of propane relative to acetylene (acetylene/propane >1) (Conner et al., 1995; Gilman et al., 2013; Russo et al., 2010; Watson et al., 2001). In this study, the acetylene/propane ratio (Fig. 7a) is greater than 1, indicating emissions from vehicles (Fraser et al., 1998). These results suggest vehicles are largely responsible for the C3 and C4 alkanes as well as the C2 alkenes/alkynes observations.

70 The C5 alkanes and some C6 alkanes like 2,3-dimethylbutane and 2-methylpentane are found in vehicular exhaust and in gasoline vapor (Tsai et al., 2006). The i-pentane to n-pentane ratio is commonly used to identify the contributions of natural gas, vehicular emissions, and fuel evaporation since these alkanes have similar boiling points, vapor pressures, and reaction rate coefficients with OH. In areas heavily dominated by natural gas drilling, ratios lie between 0.82-0.89 (Gilman et al., 2013), while higher ratios are associated with vehicle emissions (2.2-3.8) and fuel evaporation (1.8-4.6) (Jobson et al., 2004; McGaughey et al., 2004; Russo et al., 2010; Wang et al., 2013). In this study, i-pentane and n-pentane are highly correlated ($R=0.93$), indicating a common source of these compounds. The slope is 10.3, higher than reported in previous studies in China (Li et al., 2019), and the large i-pentane concentrations are likely reflective of gasoline evaporation due to the extremely volatile nature of i-pentane. The influence of fuel evaporative emissions is further identified by strong correlations between C4-C7 alkanes and alkenes typical of fuel evaporative emissions. Strong correlations of many long-chain alkanes (C6-C7 and octane) with i-pentane ($R>0.73$ except for cyclohexane) but absence of correlations with acetylene indicates solvent evaporation may be another source of long-chain alkanes.

80 Typically, the ratio of cis-2-butene/trans-2-butene is used to determine the source of C4 alkenes (Li et al., 2015; Velasco et al., 2007). However, in this study, all measurements of cis-2-butene and trans-2-butene are below the detection limit, so assessing

the ratio and correlation is not possible. Previous studies in this region in China have attributed C4 alkenes to vehicular emissions (Li et al., 2015).

90 The correlation between the C7-C8 aromatics is strong ($R > 0.76$) and revealing of typical signatures from incomplete combustion. The toluene/ethylbenzene ratio (10.7) is higher than traffic and urban emission ratios (~ 5 -8), but closer to ratios associated with biomass burning (9.41) (Monod et al., 2001; Parrish et al., 1998). Toluene also correlates with all C7-C9 alkanes ($R > 0.64$) and with *i*-pentane ($R = 0.85$), compounds from diesel and gasoline evaporation. High levels of toluene reported in Hong Kong by Ho et al., (2004) were suggested to be emitted from gasoline evaporation, while Chan et al., (2006b) attributed the high toluene levels in different PRD cities to industrial solvent usage.

95 There is an excellent correlation ($R > 0.99$) between *o*-xylene and *m/p*-xylene (Fig. 7b) and the slope (0.33) is comparable to the emission ratio found in a tunnel study (0.35) (Liu et al., 2008a). The *o*-xylene/ethylbenzene (0.60, Fig. 7b) slope is lower than vehicle exhaust emission ratios (1.2-1.8) (Conner et al., 1995; Jobson et al., 2004; Kirchstetter et al., 1996; Rogak et al., 1998; Sagebiel et al., 1996), but the correlation is extremely strong, suggesting the preferential loss of xylenes during transport due to their higher reactivity. These correlations and ratios suggest incomplete combustion from vehicular emissions and biomass burning are an important source of C7 and C8 aromatics.

00 The ratio between benzene/toluene (B/T) is a useful indicator to distinguish between vehicular emissions and other combustion sources. A ratio ~ 0.5 is often attributed to vehicular sources (Brocco et al., 1997; Perry and Gee, 1995), while ratios larger than 1 have been reported for coal or charcoal burning (Andreae and Merlet, 2001; Moreira Dos Santos et al., 2004). Benzene was observed at high mean ratios over Hebei (0.51 ppbv) and the average B/T ratio is 1.8 ± 1.6 ppbv/ppbv. The correlation of some hydrocarbons can highlight the differences between $B/T > 1$ ($N = 17$) and $B/T < 1$ ($N = 9$). The correlation found between 05 benzene and acetylene when all samples are grouped together (Fig. 7a) substantially improves just considering “traffic-related” samples ($B/T < 1$) ($R = 0.93$), suggesting a contribution of vehicular sources to benzene and acetylene measurements.

3.3 The effect of VOCs on ozone formation

10 In order to effectively reduce O_3 concentrations, it is crucial to understand the relative importance of individual VOCs in terms of the production of O_3 because each VOC exhibits different chemical reactivities. In this section, we present results using the loss rate of each VOC species with OH and ozone formation potential (OFP) assuming no influence of aerosols. Since the aerosol effect on O_3 formation is dependent upon time of day (solar zenith angle), meteorology, levels of local and neighboring aerosols, and the VOC/ NO_x ratio, the calculations presented here are simplified compared to the more complicated chemical composition of the atmosphere, but are still useful to help inform control strategies.

3.3.1 OH loss rate of VOC species

15 The calculation of the first-order loss rate of OH with different VOCs, termed OH reactivity, provides a measure of the potential to produce HO₂ and RO₂, key intermediate species in the production of O₃ (Stroud et al., 2008). Since the reaction with OH accounts for the majority of loss of most VOCs, the rate constant (obtained from the Master Chemical Mechanism version 3.3.1 (MCM3.3.1) and the National Institute of Standards and Technology (NIST) Chemical Kinetics database (www.kinetics.nist.gov/) for the reaction between OH and various hydrocarbons reflects the overall reactivity of that hydrocarbon (Finlayson-Pitts and Pitts, 1999). OH reactivity for each VOC species (VOC_i) is defined by Equation 2:

$$OHR(VOC_i) = k_{OH+VOC_i} * [VOC_i] \quad (2)$$

Where k_{OH+VOC_i} is the reaction rate constant between OH and VOC_i. Among the VOC groups, alkanes and alkenes/alkynes both contribute the most to the total VOC reactivity, accounting for 37% each. Aromatics accounted for 26% of the total VOC reactivity. The relative contribution of the top 10 VOCs ranked by mean OH reactivity (Table 5) shows ethylene, propylene, and isoprene among the top measured alkene species, together contributing ~33% to total OH reactivity. Among the alkanes, 2-methylpentane and i-pentane contribute the most (13%) to total OH reactivity, followed by the branched pentanes and propane. Aromatic compounds such as toluene and m/p-xylene constitute 13% to total OH reactivity. Previous ground-based summer studies in China have found larger contributions of isoprene to OH reactivity, ranging from ~10-30% (Li et al., 2015; Xue et al., 2017), than ARIAs (7.2%).

30 3.3.2 Ozone formation potential of NMHCs

Since OH reactivity only provides a qualitative identification of the most reactive species and does not reflect products and their production of further free radicals, we next consider the contribution to the formation of O₃ using ozone formation potential (OFP). The OFP of a VOC relies on the quantity maximum incremental reactivity (MIR), which represents the amount of O₃ formed from the addition of a small amount of the VOC species in interest under high NO_x conditions. Values of MIR (unit: g O₃ formed/ g VOC) have been calculated based on model simulations evaluated with smog chamber measurements (Carter, 2010, 1994). The OFP is calculated according to Equation 3:

$$OFP(VOC_i) = MIR_{VOC_i} * [VOC_i] \quad (3)$$

This method gives an estimate of only the first 24 hours after initial release. The median measured VOC/NO_x ratio for all WAS canisters was 4.9 ppbv/ppbv. In comparison, the ratio of reactive organic gas to NO_x (ROG/NO_x) in Los Angeles is 7.6 ppbv/ppbv (Carter, 1994). VOCs experience photochemical loss from emission sources near the surface to measured aloft concentrations. Estimation of OFP from aircraft observations throughout the PBL indicates how formation of O₃ may be different from previous surface studies.

To identify the major contributors to O₃ formation in this region, the 10 species with the highest mean OFP are listed in Table 4. Aromatic compounds are the largest contributor to total OFP (43%), followed by alkanes (30%) and alkenes/alkynes (27%).
45 Toluene and ethylene make the largest contributions (19.6% and 15.7%, respectively) to total OFP. The high MIR of these compounds (MIR=4.0 g O₃/g VOC and 9.00 g O₃/g VOC, respectively) and large mixing ratios (4.9% and 5.7% of the total measured VOC volume mixing ratio) drives their important contribution to O₃ formation. The relatively short lifetime of ethylene (~1.4 days) combined with the large range of measured mixing ratios (0.18 to 3.54 ppbv) suggests sampling of air masses with little to moderate photochemical processing, indicating the large range of influence on OFP. The most reactive
50 compound in terms of OFP is trans-2-butene (MIR=15.16 g O₃/g VOC), but its low concentration results in only 0.2% to total OFP. At the other extreme, ethane accounts for a relatively high percentage of total measured VOC volume mixing ratio (17.0%) yet only contributes 2.1% to OFP due to its low reactivity (MIR=0.49 g O₃/g VOC).

Previous studies in China report aromatics and alkenes account for the most OFP (Cai et al., 2010; Cheng et al., 2010; Jia et al., 2016; Liang et al., 2017; Wang et al., 2010a, 2016; Xie et al., 2008; Zheng et al., 2009). At a surface site in Beijing (May
55 2014), Li et al. (2015) found m/p-xylene, ethylene, toluene, propylene, and o-xylene are most influential to OFP, while at a ground station in Tianjin (August 2018), Han et al. (2020) found that ethylene, isoprene, toluene, m/p-xylene, and propylene were important contributors to OFP. Our study supports a larger contribution of anthropogenic VOCs than biogenic VOCs in spring, although summer studies indicate a major role for isoprene to the formation of O₃ in the NCP (Han et al., 2020; Zong et al., 2018). Since isoprene is mostly emitted by biogenic sources during the warmer summer months with strong solar
60 radiation and when soil moisture is sufficient for plant growth, we expect isoprene to have a larger impact on O₃ production in the summer than during spring, the time of our study. J. H. Tang et al. (2007) concluded ethylene, toluene, and m/p-xylene are the main contributors to OFP during spring 2005 at the surface in the PRD, citing emissions from industry and vehicular exhaust. Our study agrees with past research in urban areas in China identifying the most reactive VOCs in terms of OFP; O₃ appears to be formed more slowly above the surface and in nonurban areas, but production is still substantial.

National measures for Chinese VOCs abatement were released in 2015, mainly focused on the reduction of anthropogenic
65 VOCs from sources in the petrochemical industry, organic chemical industry, packaging printing, and industrial coating, not considering reactivity or chemical speciation (Li et al., 2018). A 2010 VOC emission inventory study concluded the top 15 OFP species (including m/p-xylene, toluene, propylene, o-xylene, and ethylbenzene) contributed 69% of total OFP, but only accounted for 30% of the total emission of VOCs by mass (Liang et al., 2017). Our analysis of the top 10 species ranked by
70 mean OFP shows these compounds contribute 68% to total OFP but only represent 37% of the total volume mixing ratio. Li et al., (2018) classifies industrial coal burning, biomass burning, and motorcycles to the top three VOC emission sources in Shijiazhuang, but OFP is highest for furniture coating, automobile coating, diesel vehicles, fuel evaporation, and gasoline vehicles. These results confirm that reactivity scales and emissions rates should be considered together when formulating control strategies for O₃.

75 3.4 Photochemical ozone production rate and sensitivity

In this section, we describe calculated net photochemical production rates of O₃ using the box model constrained by aircraft observations. Ozone production rates calculated from the box model are high in major urban center, particularly Shijiazhuang and Xingtai, but also between these cities (Fig. 8a). The highest rates (>10 ppbv/hour) are generally found closer to the surface, but in some instances upwards of 2000 m. The largest net production rate of O₃ (over 16 ppbv/hour) was located along the
80 Taihang Mountains between Shijiazhuang and Xingtai. This large net production rate occurred ~2000 m on June 11, 2016 when NO, NO_y, NO₂, and O₃ were ~2 ppbv, ~18 ppbv, ~3 ppbv, and ~75 ppbv, respectively.

Vertical profiles of production, loss, and net rates of O₃ (Fig. 9) show that HO₂+NO made more O₃ than RO₂+NO during the campaign. The major loss of O₃ was due to the termination of NO₂ through its reaction with OH below 2500 m. Reaction with O(¹D) is the main loss of O₃ above 2500 m. A maximum of net O₃ production for the mean profile was observed in the lowest
85 500 m of ~7 ppbv/hour. In the PBL between 1500-2000 m, where median NO and NO₂ were 534 and 625 pptv, respectively, O₃ production rates were ~4 ppbv/hour. In the lower FT from 2500 to 3000 m, peak net O₃ production rates still reached ~3ppbv/hour and were conducive to long-range transport.

Values of L_N/Q (Fig. 8b) indicate production rates of O₃ are mostly NO_x-sensitive (i.e., L_N/Q < 0.5) in the PBL over Hebei and some of the largest net production rates of O₃ are associated with NO_x-sensitivity. In order to control aloft O₃ production that
90 has the potential to be transported downwind, NO_x is the most important precursor to control. However, at low altitudes near urban centers, the production rate of O₃ tends to be more VOC-sensitive (i.e., L_N/Q > 0.5), particularly during morning flights. In urban regions of China, an O₃ formation transition from VOC-limited at the surface to NO_x-limited at ~1 km has been documented (Chen et al., 2013; Han et al., 2020). Additionally, many studies conclude O₃ production in urban areas of China
95 is VOC-sensitive in spring, while likely more NO_x-sensitive in more rural areas (Ran et al., 2011; Xue et al., 2013). Using updated emissions from a nonlinear joint analytical inversion of VOCs and NO_x from the Ozone Mapping and Profile Suite Nadir Mapper (OMPS-NM) formaldehyde and OMI NO₂ columns during KORUS-AQ in WRF-CMAQ, Souri et al. (2020) found the maximum daily 8 hour average surface O₃ over the NCP to increase by 4.56 ppbv, suggesting that emission control strategies on VOCs should be prioritized. Pusede et al., (2014) assessed the temperature dependence of emission control scenarios to lower O₃ in San Joaquin Valley, California and concluded reducing organic emissions at moderate and high
00 temperatures with co-occurring NO_x decreases will further diminish the number of O₃ violations. Thus, the control of NO_x as well as VOCs may be necessary to control both aloft and near-ground O₃ production in the NCP.

4 Summary

High concentrations of O₃ and its precursors were pervasive over Hebei Province, China in Spring 2016. In this study, we quantify the composition and photochemical nature of the lower troposphere associated with smog events. Measurements of

05 trace gases including O₃, CO, NO_x, NO_y, and of aerosol optical properties were acquired in May and June 2016. Twenty-six
samples analyzed for 54 VOCs were taken aboard a Y-12 research aircraft mostly in the PBL. Our observations confirm heavy
loadings of pollution over Hebei.

The major conclusions of our study are:

1. We observed high amounts O₃, ranging from 45 ppbv to 145 ppbv, with the highest values found over Shijiazhuang.
10 The highest NO_x concentrations were observed over Xingtai below 500 m. The highest NO_x and CO concentrations
were 53.2 ppbv and 6054 ppbv, respectively. Ratios of CO/CO₂ indicate inefficient combustion from residential coal
and biomass burning throughout the region but have decreased in China since the early 2000s suggesting the
implementation of successful pollution control strategies.
 2. Concentrations of total measured VOCs reveals alkanes contribute the most by volume mixing ratio (68%), while
15 alkenes/alkynes and aromatics together supply the most (74%) to the calculated OH loss rate. Aromatics constitute
most (43%) to the total calculated OFP and toluene, ethylene, m/p-xylene, propylene, and i-pentane play significant
roles in the aloft formation of O₃ in this region. In contrast to other surface studies in summer, we find a lower
contribution of biogenic sources (e.g. isoprene) to the formation of O₃ in the PBL. Sources of VOCs include vehicular
emissions, biomass burning, and fuel and solvent evaporation.
 - 20 3. High amounts of NO_x and VOCs throughout the PBL over nonurban parts of Hebei Province were found to generate
O₃ at a peak mean rate of ~7 ppbv/hour below 500 m. The lower free troposphere (from ~2500 to ~3000 m) was also
frequently polluted with CO and NO₂ averaging ~125 ppbv and ~140 pptv with peak net production rates of O₃ ~3
ppbv/hour, allowing for continued formation of O₃ as the air mass travels downwind. The O₃ production regime is
found to be NO_x-limited throughout the PBL over Hebei, while more VOC-limited at low altitudes near urban centers.
- 25 Our measurements in spring 2016 over Hebei cannot represent all of China or the seasonal variation of O₃ photochemistry, but
measurements from an airborne platform make a valuable addition to the understanding of one of the most polluted regions in
China, and indeed the world. The photochemistry of O₃ production is highly dependent upon the interaction of radiation and
aerosols within the PBL and future work is needed to assess optical properties of aerosols at wavelengths relevant to photolysis
of O₃ to O(¹D) and thus OH. We show that to improve air quality in Hebei Province, both NO_x and VOCs from vehicles and
30 fuel evaporation should be targeted. While VOCs are already targeted for emission reduction in China, the egregious
concentrations of O₃ observed in this study further confirm the formation of a reactivity-oriented control strategy is urgent.

Author contributions

The ARIAs campaign was supervised by RD, ZQ, and XR. XR, HH, and FW conducted the measurements on board the
research aircraft and VOCs were analyzed by MS and SL. A²BC observations were collected by ZL, FW, YW, and FZ. SR
35 helped set up the box model. SB carried out the scientific analysis of the aircraft data and drafted the manuscript with
contributions from all co-authors.

Acknowledgments, Samples, and Data

This work was supported by National Science Foundation (NSF 9188-1524) and the National Institute of Standards and Technology (NIST). The authors are grateful for the flight crew and the many scientists who helped to collect ARIAs and A²BC observations. The ARIAs flight data are available at: <https://www-air.larc.nasa.gov/cgi-bin/ArcView/korusaq?OTHER=1#top>. Background global map is from Esri, available at: <https://www.arcgis.com/home/item.html?id=21b4ba14d9e5472d97afcbb819f7368e#overview>. Additionally, we thank Glenn Wolfe of NASA for his support of the F0AM box modelling effort and Gabriele Pfister and Frank Flocke at NCAR for providing helpful discussion during manuscript preparation. The scientific results and conclusions, as well as any views or opinions expressed herein, are those of the authors and do not necessarily reflect the views of NIST, NSF, or NOAA.

References

- Al-Saadi, J., Carmichael, G., Crawford, J., Emmons, L., Kim, S., Song, C. K., Chang, L. S., Lee, G., Kim, J. and Park, R.: NASA Contributions to KORUS-AQ: An International Cooperative Air Quality Field Study in Korea., 2015.
- An, J., Wang, J., Zhang, Y. and Zhu, B.: Source Apportionment of Volatile Organic Compounds in an Urban Environment at the Yangtze River Delta, China, *Arch. Environ. Contam. Toxicol.*, 72(3), 335–348, doi:10.1007/s00244-017-0371-3, 2017.
- An, Z., Huang, R. J., Zhang, R., Tie, X., Li, G., Cao, J., Zhou, W., Shi, Z., Han, Y., Gu, Z. and Ji, Y.: Severe haze in northern China: A synergy of anthropogenic emissions and atmospheric processes, *Proc. Natl. Acad. Sci. U. S. A.*, 116(18), 8657–8666, doi:10.1073/pnas.1900125116, 2019.
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cycles*, 15(4), 955–966, doi:10.1029/2000GB001382, 2001.
- Apel, E. C., Emmons, L. K., Karl, T., Flocke, F., Hills, A. J., Madronich, S., Lee-Taylor, J., Fried, A., Weibring, P., Walega, J., Richter, D., Tie, X., Mauldin, L., Campos, T., Weinheimer, A., Knapp, D., Sive, B., Kleinman, L., Springston, S., Zaveri, R., Ortega, J., Voss, P., Blake, D., Baker, A., Warneke, C., Welsh-Bon, D., de Gouw, J., Zheng, J., Zhang, R., Rudolph, J., Junkermann, W. and Riemer, D. D.: Chemical evolution of volatile organic compounds in the outflow of the Mexico City Metropolitan area, *Atmos. Chem. Phys.*, 10(5), 2353–2375, doi:10.5194/acp-10-2353-2010, 2010.
- Avnery, S., Mauzerall, D. L., Liu, J. and Horowitz, L. W.: Global crop yield reductions due to surface ozone exposure: 2. Year 2030 potential crop production losses and economic damage under two scenarios of O₃ pollution, *Atmos. Environ.*, 45(13), 2297–2309, doi:10.1016/j.atmosenv.2011.01.002, 2011.
- Baker, A. K., Beyersdorf, A. J., Doezema, L. A., Katzenstein, A., Meinardi, S., Simpson, I. J., Blake, D. R. and Sherwood Rowland, F.: Measurements of nonmethane hydrocarbons in 28 United States cities, *Atmos. Environ.*, 42(1), 170–182, doi:10.1016/j.atmosenv.2007.09.007, 2008.
- Barletta, B., Meinardi, S., Rowland, F. S., Chan, C. Y., Wang, X., Zou, S., Lo, Y. C. and Blake, D. R.: Volatile organic compounds in 43 Chinese cities, *Atmos. Environ.*, 39(32), 5979–5990, doi:10.1016/j.atmosenv.2005.06.029, 2005.
- Bell, M. L., Peng, R. D. and Dominici, F.: The Exposure – Response Curve for Ozone and Risk of Mortality and the Adequacy of Current Ozone Regulations, *Environ. Health Perspect.*, 114(4), 532–536, doi:10.1289/ehp.8816, 2006.

- Blake, D. R., Smith, T. W., Chen, T.-Y., Whipple, W. J. and Rowland, F. S.: Effects of biomass burning on summertime nonmethane hydrocarbon concentrations in the Canadian wetlands, *J. Geophys. Res.*, 99(D1), 1699, doi:10.1029/93jd02598, 1994.
- Brocco, D., Fratarcangeli, R., Lepore, L., Petricca, M. and Ventrone, I.: Determination of aromatic hydrocarbons in urban air of Rome, *Atmos. Environ.*, 31(4), 557–566, doi:10.1016/S1352-2310(96)00226-9, 1997.
- 75 Cai, C. J., Geng, F. H., Tie, X. X., Yu, Q., Peng, L. and Zhou, G. Q.: Characteristics of ambient volatile organic compounds (VOCs) measured in Shanghai, China, *Sensors*, 10(8), 7843–7862, doi:10.3390/s100807843, 2010.
- Cao, G., Zhang, X., Gong, S. and Zheng, F.: Investigation on emission factors of particulate matter and gaseous pollutants from crop residue burning, *J. Environ. Sci.*, 20(1), 50–55, doi:10.1016/S1001-0742(08)60007-8, 2008.
- Di Carlo, P., Brune, W. H., Martinez, M., Harder, H., Leshner, R., Ren, X., Thornberry, T., Carroll, M. A., Young, V., Shepson, P. B., Riemer, D., Apel, E. and Campbell, C.: Missing OH Reactivity in a Forest: Evidence for Unknown Reactive Biogenic VOCs, *Science* (80-), 304(5671), 722–725, doi:10.1126/science.1094392, 2004.
- 80 Carter, W. P. L.: Updated Maximum Incremental Reactivity Scale and Hydrocarbon Bin Reactivities for Regulatory Applications., 2010.
- Carter, W. P. L.: Development of Ozone Reactivity Scales for Volatile Organic Compounds, *Air Waste*, 44(7), 881–899, doi:10.1080/1073161X.1994.10467290, 1994.
- 85 Chan, L. Y., Chu, K. W., Zou, S. C., Chan, C. Y., Wang, X. M., Barletta, B., Blake, D. R., Guo, H. and Tsai, W. Y.: Characteristics of nonmethane hydrocarbons (NMHCs) in industrial, industrial-urban, and industrial-suburban atmospheres of the Pearl River Delta (PRD) region of south China, *J. Geophys. Res. Atmos.*, 111(11), 1–9, doi:10.1029/2005JD006481, 2006.
- Chen, J., Li, C., Ristovski, Z., Milic, A., Gu, Y., Islam, M. S., Wang, S., Hao, J., Zhang, H., He, C., Guo, H., Fu, H., Miljevic, B., Morawska, L., Thai, P., LAM, Y. F., Pereira, G., Ding, A., Huang, X. and Dumka, U. C.: A review of biomass burning: Emissions and impacts on air quality, health and climate in China, *Sci. Total Environ.*, 579, 1000–1034, doi:10.1016/j.scitotenv.2016.11.025, 2017.
- 90 Chen, P., Quan, J., Zhang, Q., Tie, X., Gao, Y., Li, X. and Huang, M.: Measurements of vertical and horizontal distributions of ozone over Beijing from 2007 to 2010, *Atmos. Environ.*, 74, 37–44, doi:10.1016/j.atmosenv.2013.03.026, 2013.
- Cheng, H. R., Guo, H., Saunders, S. M., Lam, S. H. M., Jiang, F., Wang, X. M., Simpson, I. J., Blake, D. R., Louie, P. K. K. and Wang, T. J.: Assessing photochemical ozone formation in the Pearl River Delta with a photochemical trajectory model, *Atmos. Environ.*, 44(34), 4199–4208, doi:10.1016/j.atmosenv.2010.07.019, 2010.
- 95 Choi, J., Park, R. J., Lee, H., Lee, S., Jo, D. S., Jeong, J. I., Henze, D. K., Woo, J., Ban, S., Lee, M., Lim, C., Park, M., Shin, H. J., Cho, S., Peterson, D. and Song, C.: Impacts of local vs . trans-boundary emissions from different sectors on PM 2.5 exposure in South Korea during the KORUS-AQ campaign, *Atmos. Environ.*, 203, 196–205, doi:10.1016/j.atmosenv.2019.02.008, 2019.
- Conner, T. L., Lonneman, W. A. and Seila, R. L.: Transportation-related volatile hydrocarbon source profiles measured in Atlanta, *J. Air Waste Manag. Assoc.*, 45(5), 383–394, doi:10.1080/10473289.1995.10467370, 1995.
- 00 Dickerson, R. R., Kondragunta, S., Stenchikov, G., Civerolo, K. , Doddridge, B. G. and Holben, B. N.: The Impact of Aerosols on Solar Ultraviolet Radiation and Photochemical Smog, *Science* (80-), 278(5339), 827–830, doi:10.1126/science.278.5339.827, 1997.
- Dickerson, R. R., Li, C., Li, Z., Marufu, L. T., Stehr, J. W., McClure, B., Krotkov, N., Chen, H., Wang, P., Xia, X., Ban, X., Gong, F., Yuan, J. and Yang, J.: Aircraft observations of dust and pollutants over northeast China : Insight into the meteorological mechanisms of transport, *J. Geophys. Res.*, 112, 1–13, doi:10.1029/2007JD008999, 2007.
- 05 Ding, A. J., Wang, T., Thouret, V., Cammas, J.-P. and Nédélec, P.: Tropospheric ozone climatology over Beijing: analysis of aircraft data from the MOZAIC program, *Atmos. Chem. Phys.*, 8(1), 1–13, doi:10.5194/acp-8-1-2008, 2008.
- Environmental Protection Agency - Integrated Risk Information System: Benzene ; CASRN 71-43-2., 2003.

- 10 Fan, H., Zhao, C. and Yang, Y.: A comprehensive analysis of the spatio-temporal variation of urban air pollution in China during 2014–2018, *Atmos. Environ.*, 220, 117066, doi:10.1016/j.atmosenv.2019.117066, 2020.
- Finlayson-Pitts, B. J. and Pitts, J. N.: *Chemistry of the Upper and Lower Atmosphere*, Academic Press, San Diego, CA., 1999.
- Fraser, M. P., Cass, G. R. and Simoneit, B. R. T.: Gas-phase and particle-phase organic compounds emitted from motor vehicle traffic in a Los Angeles roadway tunnel, *Environ. Sci. Technol.*, 32(14), 2051–2060, doi:10.1021/es970916e, 1998.
- 15 Gaubert, B., Emmons, L. K., Raeder, K., Tilmes, S., Miyazaki, K., Jr, A. F. A., Elguindi, N., Granier, C., Tang, W., Barré, J., Worden, M., Buchholz, R. R., Edwards, D. P., Franke, P. and Anderson, J. L.: Correcting model biases of CO in East Asia: impact on oxidant distributions during KORUS-AQ, *Atmos. Chem. Phys. Discuss.*, doi:https://doi.org/10.5194/acp-2020-599, 2020.
- Geng, F., Zhang, Q., Tie, X., Huang, M., Ma, X. and Deng, Z.: Aircraft measurements of O₃, NO_x, CO, VOCs, and SO₂ in the Yangtze River Delta region, *Atmos. Environ.*, 43(3), 584–593, doi:10.1016/j.atmosenv.2008.10.021, 2009.
- 20 Gilman, J. B., Lerner, B. M., Kuster, W. C. and De Gouw, J. A.: Source signature of volatile organic compounds from oil and natural gas operations in northeastern Colorado, *Environ. Sci. Technol.*, 47(3), 1297–1305, doi:10.1021/es304119a, 2013.
- Goldan, P. D., Parrish, D. D., Kuster, W. C., Trainer, M., McKeen, S. A., Holloway, J., Jobson, B. T., Sueper, D. T. and Fehsenfeld, F. C.: Airborne measurements of isoprene, CO, and anthropogenic hydrocarbons and their implications, *J. Geophys. Res. Atmos.*, 105(D7), 9091–9105, doi:10.1029/1999JD900429, 2000.
- 25 Guo, S., Tan, J., Ma, Y., Yang, F., Yu, Y. and Wang, J.: Characteristics of atmospheric non-methane hydrocarbons during high PM10 episodes and normal days in Foshan, China, *Atmos. Res.*, 101(3), 701–710, doi:10.1016/j.atmosres.2011.04.022, 2011.
- Guo, S., Hu, M., Zamora, M. L., Peng, J., Shang, D., Zheng, J., Du, Z., Wu, Z., Shao, M., Zeng, L., Molina, M. J. and Zhang, R.: Elucidating severe urban haze formation in China, *Proc. Natl. Acad. Sci. U. S. A.*, 111(49), 17373–17378, doi:10.1073/pnas.1419604111, 2014.
- Halliday, H. S., Digangi, J. P., Choi, Y., Diskin, G. S., Pusede, S. E. and Rana, M.: Using Short-Term CO/CO₂ Ratios to Assess Air Mass Differences over the Korean Peninsula during KORUS-AQ, *J. Geophys. Res. Atmos.*, 124, 10,951–10,972, doi:10.1029/2018JD029697, 30 2019.
- Han, S., Kondo, Y., Oshima, N., Takegawa, N., Miyazaki, Y., Hu, M., Lin, P., Deng, Z., Zhao, Y., Sugimoto, N. and Wu, Y.: Temporal variations of elemental carbon in Beijing, *J. Geophys. Res. Atmos.*, 114(23), 1–16, doi:10.1029/2009JD012027, 2009.
- Han, S., Yao, Q., Tie, X., Zhang, Y., Zhang, M., Li, P. and Cai, Z.: Analysis of surface and vertical measurements of O₃ and its chemical production in the NCP region, China, *Atmos. Environ.*, 241, 117759, doi:10.1016/j.atmosenv.2020.117759, 2020.
- 35 Hatakeyama, S., Takami, A., Wang, W. and Tang, D.: Aerial observation of air pollutants and aerosols over Bo Hai, China, *Atmos. Environ.*, 39(32), 5893–5898, doi:10.1016/j.atmosenv.2005.06.025, 2005.
- He, H., Li, C., Loughner, C. P., Li, Z., Krotkov, N. A., Yang, K., Wang, L., Zheng, Y., Bao, X., Zhao, G. and Dickerson, R. R.: SO₂ over central China: Measurements, numerical simulations and the tropospheric sulfur budget, *J. Geophys. Res.*, 117, 1–15, doi:10.1029/2011JD016473, 2012.
- 40 Ho, K. F., Lee, S. C., Guo, H. and Tsai, W. Y.: Seasonal and diurnal variations of volatile organic compounds (VOCs) in the atmosphere of Hong Kong, *Sci. Total Environ.*, 322(1–3), 155–166, doi:10.1016/j.scitotenv.2003.10.004, 2004.
- Huang, M., Carmichael, G. R., Crawford, J. H., Wisthaler, A., Zhan, X., Hain, C. R., Lee, P. and Guenther, A. B.: Biogenic isoprene emissions driven by regional weather predictions using different initialization methods: Case studies during the SEAC4RS and DISCOVER-AQ airborne campaigns, *Geosci. Model Dev.*, 10(8), 3085–3104, doi:10.5194/gmd-10-3085-2017, 2017.
- 45 Huang, X., Wang, T., Talbot, R., Xie, M., Mao, H., Li, S., Zhuang, B., Yang, X., Fu, C., Zhu, J., Huang, X. and Xu, R.: Temporal characteristics of atmospheric CO₂ in urban Nanjing, China, *Atmos. Res.*, 153, 437–450, doi:10.1016/j.atmosres.2014.09.007, 2015.

- Inomata, Y., Iwasaka, Y., Osada, K., Hayashi, M., Mori, I., Kido, M., Hara, K. and Sakai, T.: Vertical distributions of particles and sulfur gases (volatile sulfur compounds and SO₂) over East Asia: Comparison with two aircraft-borne measurements under the Asian continental outflow in spring and winter, *Atmos. Environ.*, 40(3), 430–444, doi:10.1016/j.atmosenv.2005.09.055, 2006.
- 50 Jerrett, M., Burnett, R. T., Pope, C. A., Ito, K., Thurston, G., Krewski, D., Shi, Y., Calle, E. and Thun, M.: Long-Term Ozone Exposure and Mortality, *N. Engl. J. Med.*, 360(11), 1085–1095, doi:10.1056/NEJMoa0803894, 2009.
- Jia, C., Mao, X., Huang, T., Liang, X., Wang, Y., Shen, Y., Jiang, W., Wang, H., Bai, Z., Ma, M., Yu, Z., Ma, J. and Gao, H.: Non-methane hydrocarbons (NMHCs) and their contribution to ozone formation potential in a petrochemical industrialized city, Northwest China, *Atmos. Res.*, 169, 225–236, doi:10.1016/j.atmosres.2015.10.006, 2016.
- 55 Jobson, B. T., Berkowitz, C. M., Kuster, W. C., Goldan, P. D., Williams, E. J., Fesenfeld, F. C., Apel, E. C., Karl, T., Lonneman, W. A. and Riemer, D.: Hydrocarbon source signatures in Houston, Texas: Influence of the petrochemical industry, *J. Geophys. Res. D Atmos.*, 109(24), 1–26, doi:10.1029/2004JD004887, 2004.
- Johnson, D., Utembe, S. R. and Jenkin, M. E.: Simulating the detailed chemical composition of secondary organic aerosol formed on a regional scale during the TORCH 2003 campaign in the southern UK, *Atmos. Chem. Phys.*, 6(2), 419–431, doi:10.5194/acp-6-419-2006, 2006.
- 60 Kelley, P., Dickerson, R. R., Luke, T. and Kok, G. L.: Rate of NO₂ photolysis from the surface to 7.6 km altitude in clear-sky and clouds, *Geophys. Res. Lett.*, 22(19), 2621–2624, 1995.
- Kirchstetter, T. W., Singer, B. C., Harley, R. A., Kendall, G. R. and Ghan, W.: Impact of oxygenated gasoline use on California light-duty vehicle emissions, *Environ. Sci. Technol.*, 30(2), 661–670, doi:10.1021/es950406p, 1996.
- 65 Kleinman, L.: A comparative study of ozone production in five U.S. metropolitan areas, *J. Geophys. Res.*, 110(D2), D02301, doi:10.1029/2004JD005096, 2005a.
- Kleinman, L.: The dependence of tropospheric ozone production rate on ozone precursors, *Atmos. Environ.*, 39(3), 575–586, doi:10.1016/j.atmosenv.2004.08.047, 2005b.
- Krotkov, N. A., McLinden, C. A., Li, C., Lamsal, L. N., Celarier, E. A., Marchenko, S. V., Swartz, W. H., Bucsela, E. J., Joiner, J., Duncan, B. N., Folkert Boersma, K., Pepijn Veeffkind, J., Levelt, P. F., Fioletov, V. E., Dickerson, R. R., He, H., Lu, Z. and Streets, D. G.: Aura OMI observations of regional SO₂ and NO₂ pollution changes from 2005 to 2015, *Atmos. Chem. Phys.*, 16(7), 4605–4629, doi:10.5194/acp-16-4605-2016, 2016.
- 70 Lai, C. H., Chang, C. C., Wang, C. H., Shao, M., Zhang, Y. and Wang, J. L.: Emissions of liquefied petroleum gas (LPG) from motor vehicles, *Atmos. Environ.*, 43(7), 1456–1463, doi:10.1016/j.atmosenv.2008.11.045, 2009.
- 75 Lan, Q., Zhang, L., Li, G., Vermeulen, R., Weinberg, R. S., Dosemeci, M., Rappaport, S. M., Shen, M., Alter, B. P., Wu, Y., Kopp, W., Waidyanatha, S., Rabkin, C., Guo, W., Chanock, S., Hayes, R. B., Linet, M., Kim, S., Yin, S., Rothman, N. and Smith, M. T.: Hematotoxicity in Workers Exposed to Low Levels of Benzene, *Science (80-)*, 306(5702), 1774–1776, doi:10.1126/science.1102443, 2004.
- Li, B., Sai, S., Ho, H., Gong, S., Ni, J., Li, H., Han, L., Yang, Y. and Qi, Y.: Characterization of VOCs and their related atmospheric processes in a central Chinese city during severe ozone pollution periods, *Atmos. Chem. Phys.*, 617–638, 2019.
- 80 Li, C., Krotkov, N. A., Dickerson, R. R., Li, Z., Yang, K. and Chin, M.: Transport and evolution of a pollution plume from northern China: A satellite-based case study, *J. Geophys. Res.*, 115, D00K03, doi:10.1029/2009JD012245, 2010.
- Li, C., McLinden, C., Fioletov, V., Krotkov, N., Carn, S., Joiner, J., Streets, D., He, H., Ren, X., Li, Z. and Dickerson, R. R.: India Is Overtaking China as the World's Largest Emitter of Anthropogenic Sulfur Dioxide, *Sci. Rep.*, 7(1), 14304, doi:10.1038/s41598-017-14639-8, 2017.

- 85 Li, G., Wei, W., Shao, X., Nie, L., Wang, H., Yan, X. and Zhang, R.: A comprehensive classification method for VOC emission sources to tackle air pollution based on VOC species reactivity and emission amounts, *J. Environ. Sci.*, 67, 78–88, doi:10.1016/j.jes.2017.08.003, 2018.
- Li, L., Xie, S., Zeng, L., Wu, R. and Li, J.: Characteristics of volatile organic compounds and their role in ground-level ozone formation in the Beijing-Tianjin-Hebei region, China, *Atmos. Environ.*, 113, 247–254, doi:10.1016/j.atmosenv.2015.05.021, 2015.
- Li, L. Y., Chen, Y. and Xie, S. D.: Spatio-temporal variation of biogenic volatile organic compounds emissions in China, *Environ. Pollut.*, 90, 182, 157–168, doi:10.1016/j.envpol.2013.06.042, 2013.
- Liang, X., Chen, X., Zhang, J., Shi, T., Sun, X., Fan, L., Wang, L. and Ye, D.: Reactivity-based industrial volatile organic compounds emission inventory and its implications for ozone control strategies in China, *Atmos. Environ.*, 162(2), 115–126, doi:10.1016/j.atmosenv.2017.04.036, 2017.
- Liu, H., Jacob, D. J., Bey, I., Yantosca, R. M., Duncan, B. N. and Sachse, G. W.: Transport pathways for Asian pollution outflow over the Pacific: Interannual and seasonal variations, *J. Geophys. Res. D Atmos.*, 108(20), doi:10.1029/2002jd003102, 2003.
- 95 Liu, J. and Si, W.: Using NDVI and air temperature to monitoring winter-wheat phenology in Xingtai, Hebei, China, 2011 Int. Conf. Control. Autom. Syst. Eng. CASE 2011, doi:10.1109/ICCASE.2011.5997798, 2011.
- Liu, Y., Shao, M., Fu, L., Lu, S., Zeng, L. and Tang, D.: Source profiles of volatile organic compounds (VOCs) measured in China: Part I, *Atmos. Environ.*, 42(25), 6247–6260, doi:10.1016/j.atmosenv.2008.01.070, 2008a.
- 00 Liu, Y., Shao, M., Lu, S., Chang, C., Wang, J. and Chen, G.: Volatile Organic Compound (VOC) measurements in the Pearl River Delta (PRD) region, China, *Atmos. Chem. Phys.*, 8, 1531–1545, 2008b.
- Long, X., Tie, X., Cao, J., Huang, R., Feng, T., Li, N., Zhao, S., Tian, J., Li, G. and Zhang, Q.: Impact of crop field burning and mountains on heavy haze in the North China Plain: A case study, *Atmos. Chem. Phys.*, 16(15), 9675–9691, doi:10.5194/acp-16-9675-2016, 2016.
- Ma, M., Gao, Y., Wang, Y., Zhang, S., Ruby Leung, L., Liu, C., Wang, S., Zhao, B., Chang, X., Su, H., Zhang, T., Sheng, L., Yao, X. and Gao, H.: Substantial ozone enhancement over the North China Plain from increased biogenic emissions due to heat waves and land cover in summer 2017, *Atmos. Chem. Phys.*, 19(19), 12195–12207, doi:10.5194/acp-19-12195-2019, 2019.
- Mazzuca, G. M., Ren, X., Loughner, C. P., Estes, M., Crawford, J. H., Pickering, K. E., Weinheimer, A. J. and Dickerson, R. R.: Ozone production and its sensitivity to NO_x and VOCs: Results from the DISCOVER-AQ field experiment, Houston 2013, *Atmos. Chem. Phys.*, 16(22), 14463–14474, doi:10.5194/acp-16-14463-2016, 2016.
- 10 McGaughey, G. R., Desai, N. R., Allen, D. T., Seila, R. L., Lonneman, W. A., Fraser, M. P., Harley, R. A., Pollack, A. K., Ivy, J. M. and Price, J. H.: Analysis of motor vehicle emissions in a Houston tunnel during the Texas Air Quality Study 2000, *Atmos. Environ.*, 38(20), 3363–3372, doi:10.1016/j.atmosenv.2004.03.006, 2004.
- Mo, Z., Shao, M., Lu, S., Qu, H., Zhou, M., Sun, J. and Gou, B.: Process-specific emission characteristics of volatile organic compounds (VOCs) from petrochemical facilities in the Yangtze River Delta, China, *Sci. Total Environ.*, 533, 422–431, doi:10.1016/j.scitotenv.2015.06.089, 2015.
- 15 Mo, Z., Shao, M., Wang, W., Liu, Y., Wang, M. and Lu, S.: Evaluation of biogenic isoprene emissions and their contribution to ozone formation by ground-based measurements in Beijing, China, *Sci. Total Environ.*, 627, 1485–1494, doi:10.1016/j.scitotenv.2018.01.336, 2018.
- Monod, A., Sive, B. C., Avino, P., Chen, T., Blake, D. R. and Sherwood Rowland, F.: Monoaromatic compounds in ambient air of various cities: A focus on correlations between the xylenes and ethylbenzene, *Atmos. Environ.*, 35(1), 135–149, doi:10.1016/S1352-2310(00)00274-0, 2001.
- 20 Moreira Dos Santos, C. Y., De Almeida Azevedo, D. and De Aquino Neto, F. R.: Atmospheric distribution of organic compounds from

- urban areas near a coal-fired power station, *Atmos. Environ.*, 38(9), 1247–1257, doi:10.1016/j.atmosenv.2003.11.026, 2004.
- Neuman, J. A., Nowak, J. B., Zheng, W., Flocke, F., Ryerson, T. B., Trainer, M., Holloway, J. S., Parrish, D. D., Frost, G. J., Peischl, J.,
25 Atlas, E. L., Bahreini, R., Wollny, A. G. and Fehsenfeld, F. C.: Relationship between photochemical ozone production and NO_x oxidation in Houston, Texas, *J. Geophys. Res.*, 114, D00F08, doi:10.1029/2008JD011688, 2009.
- Parrish, D. D., Trainer, M., Young, V., Goldan, P. D., Kuster, W. C., Jobson, B. T., Fehsenfeld, F. C., Lonneman, W. A., Zika, R. D., Farmer, C. T., Riemer, D. D. and Rodgers, M. O.: Internal consistency tests for evaluation of measurements of anthropogenic hydrocarbons in the troposphere, *J. Geophys. Res. Atmos.*, 103(D17), 22339–22359, doi:10.1029/98JD01364, 1998.
- 30 Perry, R. and Gee, I. L.: Vehicle emissions in relation to fuel composition, *Sci. Total Environ.*, 169(1–3), 149–156, doi:10.1016/0048-9697(95)04643-F, 1995.
- Peterson, D. A., Hyer, E. J., Han, S. O., Crawford, J. H., Park, R. J., Holz, R., Kuehn, R. E., Eloranta, E., Knote, C., Jordan, C. E. and Lefer, B. L.: Meteorology influencing springtime air quality, pollution transport, and visibility in Korea, *Elementa*, 7(1), doi:10.1525/elementa.395, 2019.
- 35 Pusede, S. E., Gentner, D. R., Wooldridge, P. J., Browne, E. C., Rollins, A. W., Min, K. E., Russell, A. R., Thomas, J., Zhang, L., Brune, W. H., Henry, S. B., Digangi, J. P., Keutsch, F. N., Harrold, S. A., Thornton, J. A., Beaver, M. R., St. Clair, J. M., Wennberg, P. O., Sanders, J., Ren, X., Vandenberg, T. C., Markovic, M. Z., Guha, A., Weber, R., Goldstein, A. H. and Cohen, R. C.: On the temperature dependence of organic reactivity, nitrogen oxides, ozone production, and the impact of emission controls in San Joaquin Valley, California, *Atmos. Chem. Phys.*, 14(7), 3373–3395, doi:10.5194/acp-14-3373-2014, 2014.
- 40 Ran, L., Zhao, C. S., Xu, W. Y., Lu, X. Q., Han, M., Lin, W. L., Yan, P., Xu, X. B., Deng, Z. Z., Ma, N., Liu, P. F., Yu, J., Liang, W. D. and Chen, L. L.: VOC reactivity and its effect on ozone production during the HaChi summer campaign, *Atmos. Chem. Phys.*, 11(10), 4657–4667, doi:10.5194/acp-11-4657-2011, 2011.
- Reich, P. B. and Amundson, R. G.: Ambient levels of ozone reduce net photosynthesis in tree and crop species, *Science* (80-.), 230(4725), 566–570, doi:10.1126/science.230.4725.566, 1985.
- 45 Ren, X., Salmon, O. E., Hansford, J. R., Ahn, D., Hall, D., Benish, S. E., Stratton, P. R., He, H., Sahu, S., Grimes, C., Heimburger, A. M. F., Martin, C. R., Cohen, M. D., Stunder, B., Salawitch, R. J., Ehrman, S. H., Shepson, P. B. and Dickerson, R. R.: Methane Emissions from the Baltimore-Washington Area Based on Airborne Observations: Comparison to Emissions Inventories, *J. Geophys. Res. Atmos.*, 1–14, doi:10.1029/2018JD028851, 2018.
- Rogak, S. N., Pott, U., Dann, T. and Wang, D.: Gaseous Emissions from Vehicles in a Traffic Tunnel in Vancouver, British Columbia, *J. Air Waste Manag. Assoc.*, 48(7), 604–615, doi:10.1080/10473289.1998.10463713, 1998.
- 50 Russo, R. S., Zhou, Y., White, M. L., Mao, H., Talbot, R. and Sive, B. C.: Multi-year (2004–2008) record of nonmethane hydrocarbons and halocarbons in New England: Seasonal variations and regional sources, *Atmos. Chem. Phys.*, 10(10), 4909–4929, doi:10.5194/acp-10-4909-2010, 2010.
- Sagebiel, J. C., Zielinska, B., Pierson, W. R. and Gertler, A. W.: Real-world emissions and calculated reactivities of organic species from motor vehicles, *Atmos. Environ.*, 30(12), 2287–2296, doi:10.1016/1352-2310(95)00117-4, 1996.
- 55 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics*, 2nd ed., John Wiley & Sons, Inc., New Jersey., 2006.
- Shao, M., Zhang, Y., Zeng, L., Tang, X., Zhang, J., Zhong, L. and Wang, B.: Ground-level ozone in the Pearl River Delta and the roles of VOC and NO_x in its production, *J. Environ. Manage.*, 90(1), 512–518, doi:10.1016/j.jenvman.2007.12.008, 2009.
- Si, Y., Wang, H., Cai, K., Chen, L., Zhou, Z. and Li, S.: Long-term (2006–2015) variations and relations of multiple atmospheric pollutants based on multi-remote sensing data over the North China Plain, *Environ. Pollut.*, 255, 113323, doi:10.1016/j.envpol.2019.113323, 2019.
- 60

- Sillman, S., Logan, A. and Wofsy, C.: The Sensitivity of Ozone to Nitrogen Oxides and Hydrocarbons in Regional Ozone Episodes, *J. Geophys. Res.*, 95, 1837–1851, 1990.
- Silva, S. J., Arellano, A. F. and Worden, H. M.: Toward anthropogenic combustion emission constraints from space-based analysis of urban CO₂/CO sensitivity, *Geophys. Res. Lett.*, 40(18), 4971–4976, doi:10.1002/grl.50954, 2013.
- 65 Sourì, A., Nowlan, C., González Abad, G., Zhu, L., Blake, D., Fried, A., Weinheimer, A., Woo, J.-H., Zhang, Q., Chan Miller, C., Liu, X. and Chance, K.: An Inversion of NO_x and NMVOC Emissions using Satellite Observations during the KORUS-AQ Campaign and Implications for Surface Ozone over East Asia, *Atmos. Chem. Phys.*, 1, 1–39, doi:10.5194/acp-2020-220, 2020.
- Stavrakou, T., Müller, J., Bauwens, M., Smedt, I. De, Lerot, C. and Roozendaal, M. Van: Substantial Underestimation of Post-Harvest Burning Emissions in the North China Plain Revealed by Multi-Species Space Observations, *Nat. Publ. Gr.*, 6, doi:10.1038/srep32307, 2016.
- 70 Stroud, C. A., Morneau, G., Makar, P. A., Moran, M. D., Gong, W., Pabla, B., Zhang, J., Bouchet, V. S., Fox, D., Venkatesh, S., Wang, D. and Dann, T.: OH-reactivity of volatile organic compounds at urban and rural sites across Canada: Evaluation of air quality model predictions using speciated VOC measurements, *Atmos. Environ.*, 42(33), 7746–7756, doi:10.1016/j.atmosenv.2008.05.054, 2008.
- Tang, G., Wang, Y., Li, X., Ji, D., Hsu, S. and Gao, X.: Spatial-temporal variations in surface ozone in Northern China as observed during 2009–2010 and possible implications for future air quality control strategies, *Atmos. Chem. Phys.*, 12(5), 2757–2776, doi:10.5194/acp-12-2757-2012, 2012.
- 75 Tang, J. H., Chan, L. Y., Chan, C. Y., Li, Y. S., Chang, C. C., Liu, S. C., Wu, D. and Li, Y. D.: Characteristics and diurnal variations of NMHCs at urban, suburban, and rural sites in the Pearl River Delta and a remote site in South China, *Atmos. Environ.*, 41(38), 8620–8632, doi:10.1016/j.atmosenv.2007.07.029, 2007.
- Tang, J. H., Chan, L. Y., Chang, C. C., Liu, S. and Li, Y. S.: Characteristics and sources of non-methane hydrocarbons in background atmospheres of eastern, southwestern, and southern China, *J. Geophys. Res. Atmos.*, 114(3), doi:10.1029/2008JD010333, 2009.
- 80 Tang, W., Arellano, A. F., DiGangi, J. P., Choi, Y., Diskin, G. S., Agustí-Panareda, A., Parrington, M., Massart, S., Gaubert, B., Lee, Y., Kim, D., Jung, J., Hong, J., Hong, J. W., Kanaya, Y., Lee, M., Stauffer, R. M., Thompson, A. M., Flynn, J. H. and Woo, J. H.: Evaluating high-resolution forecasts of atmospheric CO and CO₂ from a global prediction system during KORUS-AQ field campaign, *Atmos. Chem. Phys.*, 18(15), 11007–11030, doi:10.5194/acp-18-11007-2018, 2018.
- 85 Tang, W., Worden, H. M., Deeter, M. N., Edwards, D. P., Emmons, L. K., Martínez-Alonso, S., Gaubert, B., Buchholz, R. R., Diskin, G. S., Dickerson, R. R., Ren, X., He, H. and Kondo, Y.: Assessing Measurements of Pollution in the Troposphere (MOPITT) carbon monoxide retrievals over urban versus non-urban regions, *Atmos. Meas. Tech.*, 13(3), 1337–1356, doi:10.5194/amt-13-1337-2020, 2020.
- Tsai, W. Y., Chan, L. Y., Blake, D. R. and Chu, K. W.: Vehicular fuel composition and atmospheric emissions in South China: Hong Kong, Macau, Guangzhou, and Zhuhai, *Atmos. Chem. Phys.*, 3281–3288, 2006.
- 90 Velasco, E., Lamb, B., Westberg, H., Allwine, E., Sosa, G. and Jobson, B. T.: Distribution, magnitudes, reactivities, ratios and diurnal patterns of volatile organic compounds in the Valley of Mexico during the MCMA 2002 & 2003 field campaigns, *Atmos. Chem. Phys.*, 329–353, 2007.
- Wang, B., Shao, M., Lu, S. H., Yuan, B., Zhao, Y., Wang, M., Zhang, S. Q. and Wu, D.: Variation of ambient non-methane hydrocarbons in Beijing city in summer 2008, *Atmos. Chem. Phys.*, 5911–5923, doi:10.5194/acp-10-5911-2010, 2010a.
- 95 Wang, F., Li, Z., Ren, X., Jiang, Q., He, H., Dickerson, R. R., Dong, X. and Lv, F.: Vertical distributions of aerosol optical properties during the spring 2016 ARIAs airborne campaign in the North China Plain, *Atmos. Chem. Phys. Discuss.*, 18, 1–22, doi:10.5194/acp-2017-1021, 2018a.
- Wang, G., Cheng, S., Wei, W., Zhou, Y., Yao, S. and Zhang, H.: Characteristics and source apportionment of VOCs in the suburban area of

- Beijing, China, *Atmos. Pollut. Res.*, 7(4), 711–724, doi:10.1016/j.apr.2016.03.006, 2016.
- 00 Wang, M., Shao, M., Lu, S. H., Yang, Y. D. and Chen, W. T.: Evidence of coal combustion contribution to ambient VOCs during winter in Beijing, *Chinese Chem. Lett.*, 24(9), 829–832, doi:10.1016/j.ccl.2013.05.029, 2013.
- Wang, Q., Han, Z., Wang, T. and Zhang, R.: Impacts of biogenic emissions of VOC and NO_x on tropospheric ozone during summertime in eastern China, *Sci. Total Environ.*, 395(1), 41–49, doi:10.1016/j.scitotenv.2008.01.059, 2008a.
- 05 Wang, T., Guo, H., Blake, D. R., Kwok, Y. H., Simpson, I. J. and Li, Y. S.: Measurements of trace gases in the inflow of South China Sea background air and outflow of regional pollution at Tai O, Southern China, *J. Atmos. Chem.*, 52(3), 295–317, doi:10.1007/s10874-005-2219-x, 2005.
- Wang, W., Ma, J., Hatakeyama, S., Liu, X., Chen, Y., Takami, A., Ren, L. and Geng, C.: Aircraft measurements of vertical ultrafine particles profiles over Northern China coastal areas during dust storms in 2006, *Atmos. Environ.*, 42(22), 5715–5720, doi:10.1016/j.atmosenv.2008.03.042, 2008b.
- 10 Wang, W., Li, X., Shao, M., Hu, M., Zeng, L., Wu, Y. and Tan, T.: The impact of aerosols on photolysis frequencies and ozone production in Beijing during the 4-year period 2012 – 2015, *Atmos. Chem. Phys.*, (2), 9413–9429, 2019a.
- Wang, Y., Munger, J. W., Xu, S., McElroy, M. B., Hao, J., Nielsen, C. P. and Ma, H.: CO₂ and its correlation with CO at a rural site near Beijing: Implications for combustion efficiency in China, *Atmos. Chem. Phys.*, 10(18), 8881–8897, doi:10.5194/acp-10-8881-2010, 2010b.
- 15 Wang, Y., Li, Z., Zhang, Y., Du, W., Zhang, F., Tan, H., Xu, H., Jin, X., Fan, X., Dong, Z., Wang, Q. and Sun, Y.: Characterization of aerosol hygroscopicity, mixing state, and CCN activity at a suburban site in the central North China Plain, *Atmos. Chem. Phys. Discuss.*, 1–34, doi:10.5194/acp-2017-1100, 2018b.
- Wang, Y., Dörner, S., Donner, S., Böhnke, S., Smedt, I., Dickerson, R. R., Dong, Z., He, H., Li, Z., Li, Z., Li, D., Liu, D., Ren, X., Theys, N., Wang, Y., Wang, Y., Wang, Z., Xu, H., Xu, J. and Wagner, T.: Vertical profiles of NO₂, SO₂, HONO, HCHO, CHOCHO and aerosols derived from MAX-DOAS measurements at a rural site in the central western North China Plain and their relation to emission sources and effects of regional transport, *Atmos. Chem. Phys.*, (2), 5417–5449, 2019b.
- 20 Warneke, C., McKeen, S. A., de Gouw, J. A., Goldan, P. D., Kuster, W. C., Holloway, J. S., Williams, E. J., Lerner, B. M., Parrish, D. D., Trainer, M., Fehsenfeld, F. C., Kato, S., Atlas, E. L., Baker, A. and Blake, D. R.: Determination of urban volatile organic compound emission ratios and comparison with an emissions database, *J. Geophys. Res. Atmos.*, 112(10), doi:10.1029/2006JD007930, 2007.
- Watson, J. G., Chow, J. C. and Fujita, E. M.: Review of volatile organic compound source apportionment by chemical mass balance, *Atmos. Environ.*, 35(9), 1567–1584, doi:10.1016/S1352-2310(00)00461-1, 2001.
- 25 Wolfe, G. M., Marvin, M. R., Roberts, S. J., Travis, K. R. and Liao, J.: The framework for 0-D atmospheric modeling (F0AM) v3.1, *Geosci. Model Dev.*, 9(9), 3309–3319, doi:10.5194/gmd-9-3309-2016, 2016.
- Wu, J., Bei, N., Hu, B., Liu, S., Wang, Y., Shen, Z., Li, X., Liu, L., Wang, R., Liu, Z., Cao, J., Tie, X., Molina, L. T. and Li, G.: Aerosol–photolysis interaction reduces particulate matter during wintertime haze events, *Proc. Natl. Acad. Sci. U. S. A.*, 117(18), 9755–9761, doi:10.1073/pnas.1916775117, 2020a.
- 30 Wu, K., Yang, X., Chen, D., Gu, S., Lu, Y., Jiang, Q., Wang, K., Ou, Y., Qian, Y., Shao, P. and Lu, S.: Estimation of biogenic VOC emissions and their corresponding impact on ozone and secondary organic aerosol formation in China, *Atmos. Res.*, 231(April 2019), 104656, doi:10.1016/j.atmosres.2019.104656, 2020b.
- Xia, L., Zhang, G., Liu, L., Li, B., Zhan, M., Kong, P. and Wang, H.: Atmospheric CO₂ and CO at Jingdezhen station in central China: Understanding the regional transport and combustion efficiency, *Atmos. Environ.*, 222, 117104, doi:10.1016/j.atmosenv.2019.117104, 2020.
- 35 Xie, X., Shao, M., Liu, Y., Lu, S., Chang, C. C. and Chen, Z. M.: Estimate of initial isoprene contribution to ozone formation potential in

- Beijing, China, *Atmos. Environ.*, 42(24), 6000–6010, doi:10.1016/j.atmosenv.2008.03.035, 2008.
- Xue, L., Wang, T., Simpson, I. J., Ding, A., Gao, J., Blake, D. R., Wang, X., Wang, W., Lei, H. and Jin, D.: Vertical distributions of non-methane hydrocarbons and halocarbons in the lower troposphere over northeast China, *Atmos. Environ.*, 45(36), 6501–6509, doi:10.1016/j.atmosenv.2011.08.072, 2011.
- 40 Xue, L. K., Wang, T., Gao, J., Ding, a. J., Zhou, X. H., Blake, D. R., Wang, X. F., Saunders, S. M., Fan, S. J., Zuo, H. C., Zhang, Q. Z. and Wang, W. X.: Ozone production in four major cities of China: sensitivity to ozone precursors and heterogeneous processes, *Atmos. Chem. Phys. Discuss.*, 13(10), 27243–27285, doi:10.5194/acpd-13-27243-2013, 2013.
- Xue, Y., Ho, S. S. H., Huang, Y., Li, B., Wang, L., Dai, W., Cao, J. and Lee, S.: Source apportionment of VOCs and their impacts on surface ozone in an industry city of Baoji, Northwestern China, *Sci. Rep.*, doi:10.1038/s41598-017-10631-4, 2017.
- 45 Yang, X., Wang, X., Yang, W., Xu, J., Ren, L., He, Y., Liu, B., Bai, Z., Meng, F. and Hu, M.: Aircraft measurements of SO₂, NO_x, CO, and O₃ over the coastal and offshore area of Yellow Sea of China, *Environ. Monit. Assess.*, 188(9), doi:10.1007/s10661-016-5533-7, 2016.
- Yuan, B., Hu, W. W., Shao, M., Wang, M., Chen, W. T., Lu, S. H., Zeng, L. M. and Hu, M.: VOC emissions, evolutions and contributions to SOA formation at a receptor site in eastern China, *Atmos. Chem. Phys.*, 13(17), 8815–8832, doi:10.5194/acp-13-8815-2013, 2013.
- 50 Zhang, Q., Yuan, B., Shao, M., Wang, X., Lu, S., Lu, K., Wang, M., Chen, L., Chang, C. C. and Liu, S. C.: Variations of ground-level O₃ and its precursors in Beijing in summertime between 2005 and 2011, *Atmos. Chem. Phys.*, 14(12), 6089–6101, doi:10.5194/acp-14-6089-2014, 2014.
- Zhang, Z., Wang, X., Zhang, Y., Lü, S. and Huang, Z.: Ambient air benzene at background sites in China’s most developed coastal regions : Exposure levels , source implications and health risks, *Sci. Total Environ.*, 511, 792–800, doi:10.1016/j.scitotenv.2015.01.003, 2015.
- 55 Zheng, J., Shao, M., Che, W., Zhang, L., Zhong, L., Zhang, Y. and Streets, D.: Speciated VOC Emission Inventory and Spatial Patterns of Ozone Formation Potential in the Pearl River Delta , China, *Enviromental Sci. Technol.*, 43(22), 8580–8586, 2009.
- Zong, R., Yang, X., Wen, L., Xu, C., Zhu, Y., Chen, T., Yao, L., Wang, L., Zhang, J., Yang, L., Wang, X., Shao, M., Zhu, T., Xue, L. and Wang, W.: Strong ozone production at a rural site in the North China Plain: Mixed effects of urban plumes and biogenic emissions, *J. Environ. Sci.*, 71, 261–270, doi:10.1016/j.jes.2018.05.003, 2018.
- 60 Zou, Y., Deng, X. J., Zhu, D., Gong, D. C., Wang, H., Li, F., Tan, H. B., Deng, T., Mai, B. R., Liu, X. T. and Wang, B. G.: Characteristics of 1 year of observational data of VOCs, NO_x and O₃ at a suburban site in Guangzhou , China, *Atmos. Chem. Phys.*, 15, 6625–6636, doi:10.5194/acp-15-6625-2015, 2015.

65

70

Table 1. Y-12 research aircraft instrumentation during ARIAs.

Variable	Method
Aircraft Position	Global Positioning System (GPS)
Meteorology (Temperature, Relative humidity, Pressure, 2-D Wind)	Cloud water inertial probe (CWIP)
Greenhouse Gases (CO ₂ /CH ₄ /CO/H ₂ O)	Cavity Ring Down Spectroscopy Picarro Model G24201-m
Ozone (O ₃)	UV-absorption, TECO 49C
Sulfur dioxide (SO ₂)	Pulsed fluorescence, TECO 43C
Nitrogen dioxide (NO ₂)	Cavity enhanced absorption spectroscopy, Los Gatos RMT-200 CRDS
NO/NO _y	Chemiluminescence, modified TECO 42C with an external Molybdenum converter at 375°C
Aerosol Scattering, b _{scat} (450, 500, 700, nm)	Nephelometer, TSI Model 3563
Aerosol Absorption, b _{abs} (565 nm)	Particle Soot Absorption Photometer (PSAP)
Black Carbon (370, 470, 520, 590, 660, 880, 950 nm)	Aethalometer, Magee Model AE31
Black Carbon	Single-Particle Soot Photometer (SP2)
VOCs	Grab Canisters, GC-MSD/FID

95 **Table 2. Aircraft monitoring results (all altitudes mostly in the PBL, 1-second data) in comparison with other airborne studies in the region. All units ppbv.**

	This Study, ARIAs		KORUS-AQ*	Yellow Sea, coastal and offshore ^a		YRD ^b	Northeast China ^c	Northern China ^d	Bohai Sea ^e	Japan Sea ^f
	May-June 2016		May 24-29	April 2011		October 2007	April 5, 2007	April 2006	March 2002	April 1996
ppbv	Ave (STD)	Min/Max	Ave. (STD)	Ave	Min/Max	Min/Max	Ahead of cold front	Ave	Min/Max	Min/Max
NO _x	5.1 (7.9)	0/53.2	1.3 (4.9)	2.45	0.49/9.58	3/40	-	5.01	-/18	-/-
CO	290.7 (309.6)	80.5/6054	258.2 (144.5)	980	630/1950	3000/7000	~300	-	-/-	-/-
O ₃	85.0 (15.6)	45.0/145.6	89.8 (17.5)	76.3	43.0/126.5	20/60	~70	43.8	35/65	70/90

* Statistics calculated for 1-second data during three flights at all altitudes during the “extreme pollution period” (Choi et al., 2019) where the KORUS-AQ DC-8 flew over the Yellow Sea to measure outflow from China.

^a Yang et al., (2016).

^b Geng et al., (2009).

^c Dickerson et al., (2007)

^d Wang et al., (2008).

^e Hatakeyama et al., (2005).

^f Inomata et al., (2006).

00

05

10

15

20

Table 3. Comparison of CO/CO₂ ratios during ARIAs to other ground-based and aloft measurements in China and developed regions of the world.

Study	Location	Year	CO/CO ₂ (%)
This Study*	North China Plain	May-June 2016	3.1
Wang et al., 2010	Miyuan, rural Beijing	Winter 2004	5.8
		Winter 2008	3.8
Huang et al., 2015	Nanjing, China	2011	3.4-4.2
Silva et al., 2013	Space-based Megacities	June 2009-May 2010	Beijing/Tianjin: 4.3 London: 0.6 Mumbai: 1.4 New York: 1.3
Han et al., 2009	Beijing, China	2005-2006	Fall: 3.0 Winter: 4.4
Tang et al., 2018*	West Sea	May-June 2016	2.8
	Seoul		0.9
Xia et al., 2020	Jingdezhen station, central China, airflow from N China	December 2017	2.6
	Jingdezhen station, airflow from SW China	18-21 January 2017	1.4
Ren et al., 2018	Baltimore/Washington, D.C.	Winter 2016	0.53

*=Aircraft studies

60 **Table 4. Comparison of the top 10 most abundant species measured in this study with other ground observations in China (Units: ppbv).**

	This Study			43 Cities ^a	QZ ^b	GZ ^d	FS ^e	LZ ^f	BJ ^b	NJ ^c
	May-June 2016			January-February 2001	July 2014	June 2011-May 2012	December 2008	June 2013-August 2013	May 2014	June-August 2014
	Ave	%	Range	Range	Ave	Ave	Ave	Ave	Ave	Range
Ethane	2.65	17.0	1.80-4.15	3.7-17.0	3.53	3.66	16.91	-	4.37	0.04-19.35
Propane	1.39	8.9	0.98-1.89	1.5-20.8	1.31	4.34	16.26	3.40	2.44	0.20-6.74
Ethylene	0.88	5.7	0.18-3.54	2.1-34.8	1.92	2.99	28.46	-	2.33	0.27-23.68
Acetylene	0.80	5.1	0.23-1.93	2.9-58.3	1.94	-	32.82	-	2.17	0.02-7.96
Toluene	0.76	4.9	0.03-4.40	0.4-11.2	0.48	4.59	18.87	1.01	1.33	0.01-5.40
i-Pentane	0.67	4.3	0.03-5.44	0.3-18.8	0.60	1.72	1.84	2.43	0.99	0.01-4.32
i-Butane	0.62	4.0	0.06-3.96	0.4-4.6	-	2.67	4.66	2.43	1.03	0.01-3.48
Benzene	0.51	3.3	0.06-2.18	0.7-10.4	0.81	0.62	6.00	1.94	0.82	0.02-11.79
2,2,4-Trimethylpentane	0.43	2.8	0.01-5.42	-	-	0.22	-	0.10	-	0.01-0.63
2-Methylheptane	0.40	2.6	0.01-5.52	-	-	0.08	0	1.49	-	0.01-0.46

^a43 Cities, China (Barletta et al., 2005).

^bQZ, Quzhou, BJ, Beijing (Li et al., 2015).

65 ^cNJ, Nanjing, Yangtze River Delta (An et al., 2017).

^dGZ, Guangzhou, Pearl River Delta (Zou et al., 2015).

^eFS, Foshan, Pearl River Delta, haze days (Guo et al., 2011).

^fLZ, Lanzhou (Jia et al., 2016).

70

75

80


85 **Table 5. Top 10 VOC species (mean and percentage breakdown) which contribute to ozone formation based on OH reactivity and Ozone formation potential during ARIAs.**

OH Reactivity			Ozone Formation Potential		
Species	Mean (s ⁻¹)	%	Species	Mean (ppbv O ₃)	%
Ethylene	0.18	15.7	Toluene	5.81	19.6
Propylene	0.11	9.6	Ethylene	4.65	15.7
Toluene	0.10	8.9	m/p-Xylene	1.87	6.3
2-Methylheptane	0.09	7.9	Propylene	1.72	5.8
Isoprene	0.08	7.2	i-Pentane	1.47	5.0
i-Pentane	0.06	5.1	2,2,4-Trimethylpentane	1.30	4.4
m/p-Xylene	0.05	4.1	2-Methylheptane	1.02	3.4
2,3,4-Trimethylpentane	0.03	3.0	i-Butane	0.93	3.1
2,2,4-Trimethylpentane	0.03	2.8	o-Xylene	0.72	2.4
Propane	0.03	2.7	1,2,4-Trimethylbenzene	0.67	2.3

90

95

00



05 **Figure 1. (a) May and June 2016 OMI tropospheric column NO₂ from NASA Goddard Earth Sciences Data and Information Services Center. The North China Plain is clearly seen in the center with high column NO₂ concentrations; the black rectangle indicates the ARIAs campaign domain and corresponds to the region shown in panel b. (b) Map of 11 ARIAs flight tracks (colored by flight number) and location of VOC samples (black triangles). The background map is provided by Esri, Copyright: ©2009.**





Figure 2. Box and whisker plots of 1-second profiles of NO, NO₂, NO_y, CO, and O₃ for data collected in 500 m bins. The whiskers show the 10th and 90th percentiles, the box denotes the 25th and 75th percentiles, and the central red line indicates the median value within each bin. Average PBL height for all ARIAs flight is ~1500 m. The total number of observations at altitudes above 2500 m of NO and NO_y is small (~2,200 or about ~30 minutes of measurements) since the NO/NO_y instrument cannot measure both species simultaneously.

10



15 **Figure 3. Maps of the ARIAs flight track colored by the 1-second measured mixing ratio (ppbv) of NO_2 (a) and O_3 (b). The background map is provided by Esri, Copyright: ©2009.**





Figure 4. Vertical profiles of 1-second NO (red), NO₂ (blue), NO_y (green), CO (orange), and O₃ (purple) in 500 m bins over the 4 spiral locations: Xingtai (a), Shijiazhuang (b), Quzhou (c), and Julu (d). The dashed lines indicate the 10th and 90th percentiles, the solid line is the median and the dotted line is the mean. We remove observations of NO/NO_y above 2500 m over three spiral locations due to limited measurements.



25 **Figure 5.** Scatter plots of 1-second (a) CO and CO₂, (b) CO and NO_y, and (c) CO and NO_x colored by the SO₂ mixing ratio for all ARIAs flights. The dashed line shows the linear regression for each plot. The dotted line in panel a indicates the higher ratio commonly associated with biomass and biofuel burning.




Figure 6. Total VOC mixing ratio for each WAS canister during ARIAs colored by alkanes (red), alkenes/alkynes (green), and aromatics (blue). The concurrent O₃ mixing ratio, divided by 5, is shown in grey diamonds (using left y-axis) and the pressure altitude of the sample is denoted by orange triangles (using right y-axis).

30






Figure 7. (a) Scatterplot of acetylene with ethane (green upward triangles), propane (orange diamonds), ethylene (purple downward triangles), and benzene (navy x's) for all WAS canisters during ARIAs. (b) Regression plots of o-xylene with m/p-xylene (red circles) and ethylbenzene (yellow squares). The dashed lines show the results of a linear least squares regression line for all data points.



35 **Figure 8. (a) Map of the net production rates of O₃ calculated using F0AM box model results along the Y-12 flight track during ARIAs. (b) Map of L_N/Q , an O₃ sensitivity indicator, along the Y-12 flight path. Ozone production is VOC-sensitive when $L_N/Q > 0.5$ and NO_x-sensitive when $L_N/Q < 0.5$ (Kleinman, 2005a). The size of the dots in both plots is proportional to the production rate of O₃. The background map is provided by Esri, Copyright: ©2009.**



40

Figure 9. Vertical profiles of the rate of production of O₃ (left), O₃ loss rate (middle), and net O₃ production rate (right) during ARIAs.