# Heterogeneous nucleation of water vapor on different types of black carbon particles

Ari Laaksonen<sup>1,2</sup>, Jussi Malila<sup>3</sup>, Athanasios Nenes<sup>4,5</sup>

#### 5

<sup>1</sup>Finnish Meteorological Institute, 00101 Helsinki, Finland.

<sup>2</sup>Department of Applied Physics, University of Eastern Finland, 70211 Kuopio, Finland.

<sup>3</sup>Nano and Molecular Systems Research Unit, 90014 University of Oulu, Finland.

<sup>4</sup>Laboratory of Atmospheric Processes and their Impacts, School of Architecture, Civil & Environmental Engineering, École
 Polytechnique Federale de Lausanne, 1015, Lausanne, Switzerland

<sup>5</sup> Institute of Chemical Engineering Sciences, Foundation for Research and Technology Hellas (FORTH/ICE-HT), 26504, Patras, Greece

Correspondence to: Ari Laaksonen (ari.laaksonen@fmi.fi)

- 15 Abstract: Heterogeneous nucleation of water vapor on insoluble particles affects cloud formation, precipitation, the hydrological cycle and climate. Despite its importance, heterogeneous nucleation remains a poorly understood phenomenon that relies heavily on empirical information for its quantitative description. Here, we examine heterogeneous nucleation of water vapor on and cloud drop activation of different types of soots, both pure black carbon particles, and black carbon particles mixed with secondary organic matter. We show that the recently developed adsorption nucleation theory
- 20 quantitatively predicts the nucleation of water and droplet formation upon particles of the various soot types. A surprising consequence of this new understanding is that, with sufficient adsorption site density, soot particles can activate into cloud droplets even when completely lacking any soluble material.

#### 25 1 Introduction

When water vapor becomes supersaturated – i.e., its relative humidity exceeds 100% – it is in a metastable state, and can form liquid water or ice. External surfaces can facilitate the phase transition, through a process known as heterogeneous nucleation. Industrial processes (e.g. dropwise condensation), biological systems (e.g. infection strategy of plant pathogens), everyday situations (e.g. fogging of glasses and windshields), and fundamentally important atmospheric phenomena (clouds,

30 frost and hoar frost) are all driven by the heterogeneous nucleation of water (Pruppacher and Klett, 1997; Franks, 2003).

Despite its ubiquity and importance, heterogeneous nucleation of water vapor remains poorly understood even after more than a century of research (Pruppacher and Klett, 1997; Möller, 2008). This poor understanding is expressed by the lack of an established heterogeneous nucleation theory that provides quantitative comprehension of the process, and translates to a large uncertainty regarding the role of aerosol–cloud interactions in the climate system (Seinfeld et al., 2016). All traditional

35 formulations are variants of classical nucleation theory (CNT) (Fletcher, 1958), which provide notoriously poor predictions of water drop nucleation (Mahata and Alofs, 1975). Molecular simulations (Lupi et al., 2014) are able to reveal aspects of heterogeneous nucleation phenomena, but on their own cannot provide a theoretical framework for describing heterogeneous nucleation in atmospheric and climate models – and are impractical for implementation in models themselves.

A shortcoming of CNT is that it does not recognize the existence of any water on surfaces prior to the formation of a

- 40 macroscopic droplet (Fig. 1a). Another shortcoming is that all the energetics of interaction between the surface and the nucleating droplet are expressed by one parameter, the contact angle (Fletcher, 1958). In reality, water adsorbs to the surface already at sub-saturated conditions (relative humidity below 100%), which can play a critical role in the nucleation process. Notably, the energetics of adsorbed water differs considerably from what is implied by the single value provided by the contact angle. In fact, the heat of adsorption changes with increasing adsorption layer thickness (Hill, 1949a). Not describing
- 45 the process of water adsorption therefore omits important physics that may hinder a quantitative description of heterogeneous nucleation. Unlike CNT, the adsorption nucleation theory (ANT) (Laaksonen, 2015; Laaksonen and Malila 2016) accounts for water adsorption prior to onset of heterogeneous nucleation (Figs. 1a and 1b).

Soot particles formed in combustion processes often contain sufficient amounts of water-soluble impurities so that their CCN activity can be described using the  $\kappa$ -Köhler theory (Petters and Kreidenweis, 2007), provided that the  $\kappa$ -value is known. An

50 exception is the major fraction of soot particles produced by aircraft engines, which contain only small amounts of impurities, and are rather hydrophobic (Demirdjian et al., 2006). As those particles may participate in contrail or cirrus ice formation (Ikhenaze et al., 2020) either directly after emission or at some later point in time, it is important to be able to characterize their water uptake and nucleation properties theoretically as well as possible.

Water insoluble atmospheric particles often have rough surfaces characterized by cavities that display physical and chemical

55 heterogeneity. This means that nucleation is not equally likely for all locations on the surface, which is another challenge for CNT. Here we show that when both adsorption and appropriate geometric considerations are included in the theoretical description of water nucleation, a quantitative theory of heterogeneous nucleation of water on aerosol is established. We demonstrate the new framework for different types of black carbon (BC) particles, which have long been known to be climate active – but with large uncertainty on their impact direct radiative forcing and on cloud formation and lifetime.



Figure 1. a) A schematic showing the principal differences between ANT and CNT. ANT is based on the premise that vapor molecules are adsorbed on the surface already at subsaturation, and the molecular nature of adsorption is 65 reflected in the values of the FHH parameters A and B that are obtained from experimental adsorption equilibria. The premise of CNT is that liquid droplets appear on the surface once the vapor becomes supersaturated. When experimental contact angles are used, CNT predicts higher critical supersaturations  $S^*$ , at which unlimited condensation starts, than ANT. In practice, the contact angle used for CNT is an adjustable parameter to fit 70 experiments, while for ANT it is the bulk (observed) value, adjusted owing to the interaction of adsorbed water molecules on the surface. b) Schematic Köhler type curves for various heterogeneous nucleation pathways of water on a 60 nm spherical particle. Curve 1 depicts the saturation ratio as a function of the diameter of individual droplets with contact angle of 30°. Nucleation takes place at the maximum of the curve. Curve 2 is the Köhler curve for the FHH activation mechanism, corresponding to a contact angle of 0°. Curve 3 corresponds to a situation where the droplets 75 fill the surface of the solid particle just below S = 1, and coalesce to form a uniform liquid film, which then grows along curve 2 until nucleation occurs at the maximum. Curve 4 is otherwise similar to curve 3, but the droplets fill the surface

at S = 1.02 and nucleation occurs instantaneously because the saturation ratio is higher than the maximum of curve 2. Note that the blue curves have been slightly offset along the *x*-axis for clarity.

#### **2** Theoretical framework

- 80 It has been shown with atomic force microscopy (Cao et al., 2011) that adsorbed water exists as discrete "patches" (droplets) of liquid water, rather than a uniform film, on non-wettable surfaces. These patches then grow with increasing relative humidity (RH). Unless they fill the surface and coalesce to form a film, the growth continues past 100% RH to the supersaturated regime (Rowley and Innes, 1942), owing to the Kelvin effect, which causes the equilibrium vapor pressure to increase over curved surfaces. At a characteristic level of supersaturation (the so called "critical supersaturation"), however,
- 85 the droplets experience heterogeneous nucleation, where unconstrained condensation of water vapor ensues to form a water droplet.

To account for the above considerations, we develop a theory that is based on physisorption and captures the progressive accumulation of water, from monolayer "patches" on a surface (at low RH) up to macroscopic amounts of condensed material throughout the adsorption–nucleation transition (at the critical supersaturation). For this, we adopt a multilayer

formulation that has minimal coefficients and can be informed by standard adsorption experiments. Our formulation (Laaksonen, 2015; Laaksonen and Malila, 2015; Laaksonen et al., 2016) uses a combination of the Frenkel–Halsey–Hill (FHH) multilayer adsorption theory (Frenkel, 1946; Halsey, 1948; Hill, 1949b) and the Kelvin effect (Defay et al., 1966). We treat the surface as covered with individual, adsorbed droplets with an average distance *s* from their nearest neighbours. If the average droplet size and the average distance between them is known, the experimentally observable adsorption layer
thickness can be calculated. Surface heterogeneities imply droplets with a distribution of radii and contact angles; but as a first approximation we assume all droplets to have the same size and contact angle (these can be considered as average values). The equilibrium condition for a single droplet is given by

$$\ln(S) = -\frac{A}{N_a^B} + \frac{2\gamma v_w}{kTR} \tag{1}$$

where *S* denotes saturation ratio,  $\gamma$  is surface tension,  $v_w$  is the volume of a water molecule, *k* is the Boltzmann constant, *T* is 100 temperature, *R* is the droplet radius, and *A* and *B* are FHH parameters. The average number of monolayers,  $\overline{N_d}$ , over the droplet projection area on the adsorbent surface is given by  $\overline{N_d} = \overline{\delta} / \delta_m$ , where  $\overline{\delta}$  denotes the average water layer thickness, and  $\delta_m$  is the thickness of a monolayer given as  $v_w/\sigma$ , where  $\sigma$  is the cross-sectional area of an adsorbed water molecule.  $\overline{N_d}$  and *R* can be related via a relation that depends on the contact angle  $\theta$  and the curvature of the adsorbent surface (see Sec. 2.1. for details of the theory). Thus, unlike in CNT, contact angle is a purely geometric parameter

- 105 describing the behavior of bulk water on the surface. Molecular simulations have shown that the contact angle of a water nanodroplet on graphitic surface equals the macroscopic value for droplets having radii larger than about 4 nm (Sergi et al, 2012). For nanodroplets on hydrophilic clays, the contact angle does not vary for droplets containing 256–1000 molecules (Zheng et al., 2017) – which means that any change in contact angle refers to nanodroplets with radii of the order of 1 nm. The sizes of critical clusters calculated using our theory are typically much larger than that, justifying the use of
- experimental contact angles in the heterogeneous nucleation calculations. In case of a zero contact angle, water forms a uniform film on the adsorbent surface. If uniform adsorption occurs on spherical particles with radius  $R_p$ , the average number of monolayers is  $\overline{N} = (R - R_p) / \delta_m$  (Sorjamaa and Laaksonen, 2007; Laaksonen and Malila, 2016).

The FHH interaction parameters *A* and *B* describe the strength of molecular interaction between the adsorbent and the first adsorption layer, and the exponential decay of the interaction strength as a function of distance, respectively. With the van

- der Waals model fluid, *A* and *B* have well defined values (note, however, that *A* depends on the assumed molecular crosssection of the adsorbing molecule), but in real systems they must be treated as empirical constants that can be determined from measured adsorption isotherms. In case of droplet-wise adsorption, the macroscopically determined adsorption layer thickness is a statistical measure that can be related to the droplet size if the average distance between the adsorbed droplets, *s*, and the contact angle are known. In practice, *s* can be obtained together with *A* and *B* from the adsorption isotherms
- 120 (Laaksonen, 2015; Laaksonen and Malila, 2016).

125

We note that there is considerable uncertainty with the value of the experimental contact angle, due to contact angle hysteresis (Tadmor, 2004). The difference between the advancing and receding angles can be on the order of 10 degrees or even more. In any case, with graphite the critical supersaturation is surprisingly insensitive to an uncertainty of around  $\pm 10^{\circ}$ in the value of  $\theta$ . The reason for this is that the contact angle is included in the equations used when the theory is fitted to adsorption experiments in order to determine the values of the FHH parameters *A* and *B*. Thus, if the contact angle is changed by a few degrees, the model has to be re-fitted to the adsorption data, which can be done simply by changing the value of the *A*-parameter. It turns out that this does not either compromise the quality of the fit to the adsorption experiments or noticeably change the calculated critical supersaturation unless the contact angle is changed by more than about plus or minus 10 degrees (see Fig. A2).

- Figure 1b shows a schematic picture of nucleation of water vapor on a spherical particle in our model framework. A single droplet on a BC particle grows along the red curve 1 until it reaches the critical supersaturation  $S_{C1}^*$  and nucleates. However, if there are several similar droplets growing on the particle, and the average distance between them is small enough so that they coalesce to form a uniform liquid film at sufficiently low saturation ratio (blue curve 3), the new equilibrium as well as the new critical supersaturation  $S_{U}^*$  can be calculated simply by using a zero contact angle (red curve 2). Depending on the
- 135 value of *s*, it is also possible that the droplet coalescence occurs at some supersaturation  $S_{Co}^*$  between  $S_{U}^*$  and  $S_{Cl}^*$ , in which case nucleation is immediate (blue curve 4).

Classical nucleation theory is based on the idea that nucleation is a stochastic phenomenon whereas ice nucleation theories that assume specific temperature dependent ice nucleation sites are deterministic. The adsorption nucleation theory is deterministic as one of its premises is that nucleation occurs on specific adsorption sites, and there is no stochastic element

140 included in the theory. Such an element could be introduced similarly as in the soccer ball model (Niedermeier et al., 2014), by assuming a distribution of contact angles between the adsorption sites. However, without knowledge of e.g. the possible widths of such distributions, we prefer not to include this idea at present. In the future, we aim to use molecular dynamics to explore this issue with chemically and physically heterogeneous surfaces.

#### 145 **2.1 Adsorption nucleation theory**

The starting point of the theory is that adsorbing vapor molecules are assumed to cluster around adsorption sites to form spherical caps of liquid drops having a contact angle  $\theta$ . The equilibrium condition is given by Eq. (1). The geometry for a spherical seed particle is as shown in Fig. 2. The average over  $\delta^B$  is given by

$$\overline{\delta^{B}} = \left[1 - \cos\Phi\right]^{-1} \int_{0}^{\Phi} \delta^{B} \sin\alpha d\alpha$$
<sup>(2)</sup>

150 with

$$\delta = -R_p + R\cos\beta + \sqrt{(R\cos\beta)^2 - R^2 + d^2}$$
(3)

$$d = \sqrt{R_p^2 + R^2 - 2R_p R \cos\theta} \tag{4}$$

$$\cos\Phi = (R_p - R\cos\theta)/d \tag{5}$$

$$\cos\beta = (R_p + \delta - d\cos\alpha)/R \tag{6}$$

*S* can now be computed as a function of *R* using Eq. (1) with  $\overline{N_d^B} = \overline{\delta^B} / \delta_m^B$  evaluated numerically from Eqs. (2) – (6). The resulting curve shows a maximum at the critical supersaturation *S*<sup>\*</sup> (and critical radius *R*<sup>\*</sup>), marking the onset of heterogeneous nucleation.



160 Figure 2. The geometry used in deriving equations for nucleation of a spherical drop on a spherical nanoparticle.

The values of the adsorption parameters *A* and *B*, as well as that of the distance between active sites *s*, can be obtained from measured adsorption isotherms. If a portion of the adsorption data is clearly in the multilayer regime and the data points align linearly when ln(-ln(*S*)) is plotted vs. ln(*N*), parameters *A* and *B* can be determined by fitting the classical FHH equation ln(*S*)  $165 = -AN^B$  to the data (here, *N* denotes the macroscopically observable surface coverage). If the contact angle is known, the distance between active sites can be obtained from the sub-monolayer part of the adsorption isotherm. To do that, a relation is needed between the size of an individual adsorbed droplet, and the macroscopic surface coverage. Making the approximation  $\overline{N_d^B} \approx \overline{N_d}^B$ , Eq. (1) can be translated into

$$\ln(S) = -A \left[ \frac{\pi \zeta^2}{\eta^2 N} \right]^{B/3} + \frac{2\gamma}{3kT} \left[ \frac{\pi \eta \zeta^2}{N} \right]^{1/3}, \qquad \theta \le 90^\circ, \tag{7}$$

170 
$$\ln(S) = -A \left(\frac{\eta}{\beta}\right)^B \left[\frac{\pi g(\theta)}{3s^2 N}\right]^{\frac{B}{3}} + \frac{2\gamma v}{3kT} \left[\frac{\pi g(\theta)}{3s^2 N}\right]^{\frac{1}{3}}, \quad \theta > 90^{\circ}$$
(8)

with  $\zeta = 3v\sin\theta/s$ ,  $\eta = \sigma f(\theta)$ ,  $g(\theta) = 4 - (1 + \cos\theta)^2(2 - \cos\theta)$ , and  $f(\theta) = \frac{(1 - \cos\theta)^2(2 + \cos\theta)}{\sin^2\theta}$ ,  $\theta \le 90^\circ$ ;

$$f(\theta) = 2 - 3\cos\theta, \theta > 90^{\circ}.$$

190

Distance between adsorption sites s can now be determined by fitting Eq. (7) or (8) to the sub-monolayer data (note that it is best to use the portion of the data just below a monolayer, as the low coverage data may be affected by pores or other factors

175 not taken into account in the above equations.

When the adsorbent surface is poorly wettable, the adsorption data may not extend to the multilayer regime at all. In such a case, all of the parameters need to be obtained from a best fit of either Eq. (7) or Eq. (8) to the data set.

As noted in relation to Fig. 1b, nucleation can also occur due to coalescence of adsorbed droplets. The criteria we use for the coalescence nucleation are that (i) the projection areas of adsorbed droplets equal the surface area of the seed particle, and that

180 (ii) the supersaturation exceeds the critical supersaturation calculated for a seed particle that is perfectly wettable.

#### 2.2 Liquid drop nucleation on insoluble particles mixed with soluble material

If  $\theta = 0^\circ$ , i.e. the seed particle is completely wettable, the saturation ratio can be written in terms of dry and wet particle radii as

185 
$$\ln(S) = -A \left(\frac{R-R_p}{D_W}\right)^{-B} + \frac{2\gamma v_W}{kTR}$$
(9)

where we have approximated monolayer thickness with the diameter of a water molecule,  $D_w$  (Laaksonen and Malila, 2016). For a partially soluble seed particle – for example, a black carbon particle coated with low-volatility organic or inorganic substances after atmospheric processing – Eq. (9) needs to be further modified to take the hygroscopicity of solution into account. The following equation (Kumar et al, 2011) can be written in terms of the hygroscopicity parameter  $\kappa$  (Petters and Kreidenweis, 2007):

$$\ln(S) = -A \left(\frac{R-R_p}{D_w}\right)^{-B} - \frac{\varepsilon_s R_p^3 \kappa}{\left(R^3 - \varepsilon_i R_p^3\right)} + \frac{2\gamma v_w}{kTR}$$
(10)

where  $\varepsilon_s$  and  $\varepsilon_i$  are the volume fractions of soluble and insoluble species in the seed particles.

#### 3. Results and discussion

#### 195 **3.1 Constraining water adsorption parameters on carbon**

We now focus on determination of water vapor adsorption parameters on graphite and on black carbon (BC) particles. Figure 2 shows a logarithmic FHH plot of several adsorption isotherms. The logarithm of the number of monolayers is shown as a function of ln(-ln *S*) because according to the FHH adsorption model, the multilayer adsorption data should align linearly in such a plot (note that saturation ratio decreases from left to right). Starting from the bottom, the lowermost measurement data

- 200 set (Young et al, 1954) shows adsorption on graphitized carbon black treated at 400 °C for 12 hours prior to the measurements. Based on the minute amounts of water adsorbed at even the highest relative humidity, and information obtained from heat of immersion measurements, Young et al. (1954) suggested that clusters of water adsorb at hydrophilic surface sites which are far apart and may arise from surface oxides formed at the edge atoms of graphite, which has since then been corroborated by quantum chemical calculations (Oubal et al., 2010). This description is in complete agreement
- with the droplet-wise FHH adsorption model (see the lowermost solid line in Fig. 3 where the model reproduces the data). The model parameters are shown in Fig. 3; the contact angle value was obtained from Fowkes and Harkins (1940), and the parameters *A*, *B*, and *s* were determined by optimizing the model fit to the experimental data.



210 Figure 3. An FHH-plot of experimental (markers) and modelled (lines) adsorption isotherms of water vapor on graphite and different soots. Note that in the abscissa, increasing supersaturation is to the left. See text for details.

When BC is exposed to atmospheric conditions, the number of hydrophilic surface sites is expected to increase with time as different contaminants adsorb on the carbon and may undergo oxidation (Tritscher et al., 2011; Aria et al., 2016; Grimonprez

et al., 2018; Friebel and Mensah, 2019). Contaminants may also be present due to imperfect combustion, in case the carbon was produced by burning. This suggests that adsorption isotherms of BC containing impurities can be accounted for by increasing the density of the adsorption sites in the FHH model (without changing the other adsorption parameters), constrained by observations. The data in the upper part of Fig. 3 (Kuznetsov et al., 2003; Popovicheva, 2008) show water adsorption on spark discharge

- soot, and thermal soot produced by natural gas pyrolysis. The materials were not treated at high temperature prior to the adsorption measurements, and are thus likely to contain more adsorption sites from surface contaminants than the graphitized carbon black used by Young et al. (1954). The blue and red solid curves represent the corresponding adsorption isotherms, obtained by simply reducing the average distance of adsorption sites in the FHH model from 75 nm to 6.8 nm and 5 nm, respectively, and keeping the other parameters at the graphitized carbon black values. The data for RH > 80% (ln(-ln
- (S) (S) (-1.5) agree well with the theoretical curves, whereas at lower relative humidities the data show more adsorption than the model. The latter is caused by the effects of soot porosity, which is indicated by a hysteresis between the measured adsorption and desorption curves.

When the density of hydrophilic adsorption sites is sufficiently high, the growing water clusters will start coalescing as the RH increases above some threshold value, and a liquid film will be formed on the BC surface. If the film formation occurs at

RH below 100%, the conditions for activation of a BC particle into a cloud droplet (where the particle is said to act as a Cloud Condensation Nuclei, or CCN) can be described using a zero contact angle and the total surface area of the particle, while the full adsorption nucleation theory with non-zero contact angle is required when the active site density is low (curve 3 in Fig. 1b). We next examine whether the CCN activation of different types of BC particles can be described with the theoretical framework as constrained above, using model parameters shown in Fig. 3.

235

#### 3.2 Water nucleation on black carbon particles

Figure 4 depicts CCN activation data for hydrophobic and hydrophilic BC particles. The most hydrophobic particles (Kotzick et al., 1997) were produced using a spark discharger. The particles were in contact only with argon and nitrogen gases prior to the activation measurements (black spheres), and thus contained very little adsorption sites. As can be seen, all

240 model curves are below the spark discharger data. However, the model with parameters obtained using the adsorption data for graphitized carbon black (Fig. 3) is in excellent agreement with the CCN activation data for spark discharger particles that were subjected to ozonolysis prior to the activation measurement (Kotzick et al., 1997) (red squares), and therefore have an increased density of adsorption sites.



Figure 4. Experimental (markers) and theoretical (lines) critical supersaturations as a function of particle size for cloud drop activation of hydrophobic and hydrophilic soot particles. See text for details.

Palas soot consists of agglomerates of primary soot spherules with a diameter on the order of 5–10 nm (Gysel et al., 2012).

- 250 There is a possibility that the heterogeneous nucleation measurements would be explained by nucleation of capillary condensed pendular water rings formed around the contact points of primary spherules. Such pendular rings would not be formed much below 100 % relative humidity as the contact angle is close to 90 degrees, and thus the meniscus cannot in practice be concave. Crouzel and Marlow (1995) modelled water ring nucleation assuming equal sized spheres in contact. For a contact angle of 85 degrees, they calculated that the critical supersaturation is about 2% for 400 nm diameter spheres,
- 255 about 4% for 200 nm particles, and about 8% for 100 nm particles. The size dependence is so strong that with 5–10 nm spheres, it is unlikely that the nucleation would take place at significantly lower supersaturation than we have calculated for Fig. 4 (red line).

The blue diamonds in Fig. 4 show activation of soot particles produced by burning camp stove fuel (white gas) (Hagen et al., 1989). Such particles can be expected to contain a large amount of oxidized species on their surfaces, and may therefore

- already have a liquid film of water at activation. The blue curve shows model prediction with the same FHH parameters *A* and *B* as the red curve, but a zero contact angle. It appears that the model has somewhat gentler size dependence than the rather scattered data, but otherwise the agreement is quite satisfactory. The CCN activation of pigment black particles (Dalirian et al., 2018), also match the theoretical line within uncertainty limits (with the exception of the largest particle size, which has the uncertainty range just below the theoretical line).
- Very interestingly, the three datapoints of Dusek et al. (2006) are clearly above those of Dalirian et al. (2018). The principal difference between the soots used in these two studies is that Dusek et al. (2006) applied heat treatment, thus very likely reducing the adsorption site density. Indeed, a theoretical line calculated with average adsorption site distance 4.4 nm and assuming the coagulation nucleation to occur, matches the data perfectly. When a site distance of just 0.2 nm smaller is assumed, the theoretical prediction becomes equal to the CCN activation prediction for particles larger than about 50 nm.
- 270 With smaller particles, the coagulation nucleation appears to match the camp stove gas soot data better than the CCN activation theory. In general, accounting for droplet-wise adsorption and resulting alternative pathways for heterogeneous nucleation (Fig. 1) allows us to obtain a much-improved description on the nucleation of water on BC particles than CNT

(Kotzick et al., 1997) or adsorption nucleation theories not accounting for initial adsorption on distinct sites (Henson, 2007; Sorjamaa and Laaksonen, 2007).

- 275 Further enhancement in BC hydrophilicity, on top of increased adsorption site density due to contaminant oxidation, may occur e.g. as a result of heterogeneous chemistry leading to a substantial amount of water-soluble material on the BC particles. The red diamonds in Fig, 5 (Wittbom et al., 2014) are for particles aged in a smog chamber prior to the activation measurements, and were estimated to contain between 8 and 91 percent by mass of water soluble substance, respectively (the data points were deciphered from Figs. 10 and 11 in Wittbom et al. (2014)). The blue crosses were calculated using the
- theory which describes the combined effect of adsorption and water-soluble material when the volume fraction ε and the hygroscopicity parameter κ of the water-soluble material are known (Kumar et al., 2011; Petters and Kreidenweis, 2007; see previous section for details). The FHH parameters were kept the same as for the pure black carbon particles, and the κ-value was assumed to be 0.13 (Wittbom et al., 2014). The model predictions, shown by the blue crosses, are once again in excellent agreement with the data. Only in three cases the theory provides clear underestimates, the worst occurring at the lowest SOA mass fraction of about 0.08, as can be seen in Fig. 5b, while Fig. 5a shows that the experimental point is very close to the Kelvin line. Comparison to Fig. 4 indicates that this data point is compatible with activation of particles that do

not contain water soluble material.



290 Figure 5. Experimental and theoretical critical supersaturations of water vapor on mixed soot particles containing BC and SOA as a function of a) volume equivalent diameter of dry particles and b) SOA mass fraction. The data are a subset of the data shown in Figures 10 and 11 of Wittbom et al. (2014), and were identified by matching supersaturations (to a precision < 0.01%) in the two figures.

#### 4. Conclusions

295 In this study, we have shown that the adsorption nucleation theory can predict the critical supersaturations for heterogeneous nucleation and CCN activation of water vapor on different types of black carbon particles, using a single set of experimental adsorption and contact angle parameters.

It is quite obvious that surface heterogeneities cause adsorbed droplets to have a range of radii and contact angles, but to make the theory usable, we assume that we can resort to average values. This applies to the FHH parameters A and B as well;

300 it is likely that there would be variability if isolated surface regions were examined. However, the variability of the heterogeneities – whether chemical or physical – is reflected in the values of the "average" adsorption parameters obtained from macroscopic adsorption isotherms. Our hypothesis is that the same parameters can be made use of when extending the theory to the supersaturated regime to predict critical supersaturations. The good match between our theoretical predictions and the experimental critical supersaturations provides clear indication that our hypothesis is well supported. This also

implies that the heterogeneities occur at a spatial scale smaller than the surface area covered by the critical nuclei.

Our findings concerning adsorption nucleation on BC have a direct relevance for climate studies, as even the sign of the radiative forcing due to BC aerosol–cloud interactions remains uncertain (Bond et al., 2013). Notably, atmospheric lifetime of BC particles is regulated by condensation of water and subsequent cloud processes, which contributes further to the uncertainty of the total radiative forcing (Samseth et al., 2018).

310

#### Appendix A: Water properties used in the calculations

T denotes absolute temperature (K), T' is temperature in degrees Celsius, and  $R_g$  is the molar gas constant.

Equilibrium vapor pressure of liquid water (Pa) (Murphy and Koop, 2010):

 $p_w = \exp\{54.842763 - 6763.22/T - 4.21\ln(T) + 0.000367T + \tanh[0.0415(T-218.8)][53.878-1331.22/T-10.000367T] + \hbar[0.000367T] + \hbar[0.00037T] + \hbar[0.000367T] + \hbar[0.00037T] + \hbar[0.000367T] + \hbar[0.00037T] + \hbar[0.000$ 

315  $9.44523\ln(T)+0.014025T$ ]

Density of liquid water (g cm<sup>-3</sup>) (Marcolli, 2017):

 $\rho_{w} = 1.8643535 - 0.0725821489T + 2.5194368 \cdot 10^{-3}T^{2} - 4.9000203 \cdot 10^{-5}T^{3} + 5.860253 \cdot 10^{-7}T^{4} - 4.5055151 \cdot 10^{-9}T^{5} + 2.2616353 \cdot 10^{-11}T^{6} - 7.3484974 \cdot 10^{-14}T^{7} + 1.4862784 \cdot 10^{-16}T^{8} - 1.6984748 \cdot 10^{-19}T^{9} + 8.3699379 \cdot 10^{-23}T^{10}$ 

320 Surface tension of water (N m<sup>-1</sup>) (Hrubý et al., 2014):

 $\gamma_w = 0.2358(1 - T/647.096)^{1.256}[1 - 0.625(1 - T/647.096)]$ 

#### **Appendix B: Sensitivity analysis**

The contact angle value used in the calculations shown above is 86°; here we examine how much the critical supersaturation is affected by changing the contact angle by up to plus/minus 10 degrees. As noted in the main text, when the contact angle

- is given a new value, the model needs to be re-fitted to the adsorption data by changing the value of the FHH parameter *A*. Figure A1 shows the values of *A* as a function of the contact angle in the range 76°–96° when this is done (the values of the parameters *B* and *s* were kept at 1.93 and 75 nm, respectively). Figure A2 shows the experimental adsorption data together with the model for  $(\theta, A) = (76^\circ, 7.9)$ , (86°, 12) and (96°, 18). As can be seen, all curves fit the data well, although there is some deviation at low saturation ratios. Figure A3 shows the calculated critical supersaturations for water nucleation. The
- 330 dry particle diameter (100 nm) was chosen based on the experimental nucleation data of Kotzick et al. (1997). The change of contact angle by 20 degrees (and the simultaneous change of the *A* parameter as shown in Fig. A1) causes a change of less than 0.7% in critical supersaturation. This is much less than the experimental uncertainty of the critical supersaturations of ozonolyzed Palas soot particles (Fig. 4).



Figure A1. Pairs of contact angle and the FHH parameter A that produce good model fits to the adsorption data.



**Figure A2.** Model fits to adsorption data for three pairs of  $\theta$  and *A*.



**Figure A3.** Critical supersaturations in water nucleation as a function of contact angle. The *A*-parameter values used in the calculations are as shown in Fig. A1. Nucleation temperature is +25 °C and dry particle diameter 100 nm.

Acknowledgments. Al acknowledges support by the Academy of Finland, C-Main project (grant no. 309141) and the Center of Excellence programme (grant no. 307331). JM acknowledges support by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 717022). AN acknowledges support by the project PyroTRACH (ERC-2016-COG) funded from H2020-EU.1.1. - Excellent Science - European Research

Council (ERC), project ID 726165 and from the European Union's Horizon 2020 project FORCeS under grant agreement No 821205.

Code and Data availability. Code and model results are available upon request

350

*Competing interests.* The authors declare that they have no conflict of interest.

355 *Author contributions*. AL performed all theoretical calculations and drafted the manuscript. All co-authors discussed the results. JM and AN commented on the manuscript and contributed to writing of the final version.

365

370

- Aria, A. I., Kidambi, P. R., Weatherup, R. S., Xiao, L., Williams, J. A., and Hofmann, S.: Time evolution of the wettability of supported graphene under ambient air exposure, J. Phys. Chem. C 120, 2215–2224,
- 360 https://doi.org/10.1021/acs.jpcc.5b10492, 2016.
  - Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, J. Geophys. Res. Atmos. 118, 5380–5552,
    - https://doi.org/10.1002/jgrd.50171, 2013.
  - Cao, P., Xu, K., Varghese, J. O., and Heath, J. R.: The microscopic structure of adsorbed water on hydrophobic surfaces under ambient conditions, Nano Lett. 11, 5581-5586, https://doi.org/10.1021/nl2036639, 2011.
  - Crouzet, Y., and Marlow, W. H.: Calculations of the equilibrium vapor pressure of water over adhering 50–200-nm spheres, Aerosol Sci. Tech. 22, 43-59, https://doi.org/10.1080/02786829408959727, 1995.
    - Dalirian, M., Ylisirniö, A., Buchholz, A., Schlesinger, D., Ström, J., Virtanen, A., and Riipinen, I.: Cloud droplet activation of black carbon particles coated with organic compounds of varying solubility, Atmos. Chem. Phys., 18, 12477– 12489, https://doi.org/10.5194/acp-18-12477-2018, 2018.

Defay, R, Prigogine, I., and Bellemans, A.: Surface Tension and Adsorption, Longmans, London, 1966.

- 375 Demirdjian, B., Ferry, D., Suzanne, J., Popovicheva, O.B., Persiantseva, N.M., and Shonija, N.K.: Heterogeneities in the microstructure and composition of aircraft engine combustor soot: impact on the water uptake, J. Atmos. Chem. 56, 83–103, https://doi.org/10.1007/s10874-006-9043-9, 2007.
  - Dusek, U., Reischl, G. P., and Hitzenberger, R. CCN activation of pure and coated carbon black particles, Environ. Sci. Tech. 40, 1223–1240, https://doi.org/10.1021/es0503478, 2006.
- Fletcher, N.H.: Size effect in heterogeneous nucleation. J. Chem. Phys. 29, 572–576, https://doi.org/10.1063/1.1744540, 1958.

Franks, F.: Nucleation of ice and its management in ecosystems, Phil. Trans. R. Soc. Lond. A 361, 557–574, https://doi.org/10.1098/rsta.2002.1141, 2003.

- Fowkes, F. M., and Harkins, W. D.: The state of monolayers adsorbed at the interface solid-aqueous solution, J. Am. Chem.
- 385 Soc. 62, 3377–3386, https://doi.org/10.1021/ja01869a029, 1940.

Frenkel, J.: Kinetic Theory of Liquids, Oxford University Press, London, 1946.

Friebel, F., and Mensah, A. A.: Ozone concentration versus temperature: atmospheric aging of soot particles. Langmuir 35, 14437–14450, https://doi.org/10.1021/acs.langmuir.9b02372, 2019.

Grimonprez, S., Faccinetto, A., Batut, S., Wu, J., Desgroux, P., and Petitprez, D.: Cloud condensation nuclei from the

- activation with ozone of soot particles sampled from a kerosene diffusion flame. Aerosol Sci.Tech. 52, 814-827,
  2018. https://doi.org/10.1080/02786826.2018.1472367.
  - Hagen, D. E., Trueblood, M. B., and White, D. R.: Hydration properties of combustion aerosols, Aerosol Sci. Tech. 10, 63–69, https://doi.org/10.1080/02786828908959221, 1989.

Halsey, G., Physical Adsorption on non-uniform surfaces, J. Chem. Phys. 16, 931–937, https://doi.org/10.1063/1.1746689, 1948.

- Henson, B. F.: An adsorption model of insoluble particle activation: Application to black carbon, J. Geophys. Res. 112, D24S16, https://doi.org/10.1029/2007JD008549, 2007.
- Hill, T. L.: Statistical mechanics of adsorption. V. Thermodynamics and heat of adsorption, J. Chem. Phys. 17, 520–535, https://doi.org/10.1063/1.1747314, 1949a.
- Hill, T. L., Physical adsorption and the free volume model for liquids, J. Chem. Phys. 17, 590, https://doi.org/10.1063/1.1747341, 1949b.
  - Hrubý, J., Vinš, V., Mareš, R., and Kalová, J.: Surface tension of supercooled water: no inflection point down to -25 °C, J. Phys. Chem. Lett. 5, 425–428, https://doi.org/10.1021/jz402571a, 2014.

Ikhenazene, R., Pirim, C., Noble, J. A., Irimiea, C., Carpentier, Y., Ortega I. K., Ouf, F.-X., Focsa, C., and Chazallon, B.: Ice

405 nucleation activities of carbon-bearing materials in deposition mode: From graphite to airplane soot surrogates, J.
 Phys. Chem. C 124, 489–503, https://doi.org/10.1021/acs.jpcc.9b08715, 2020.

Kim, S., Kim, D., Kim, J., An, S., and Jhe, W.: Direct evidence for curvature-dependent surface tension in capillary condensation: Kelvin equation at molecular scale, Phys. Rev. X 8, 041046, https://doi.org/10.1103/PhysRevX.8.041046, 2018.

410 Kotzick, R., Panne, U., and Niessner, R.: Changes in condensation properties of ultrafine carbon particles subjected to oxidation by ozone, J. Aerosol Sci. 28, 725–735, https://doi.org/10.1016/S0021-8502(96)00471-5, 1997.

Kumar, P., Sokolik, I. N., and Nenes, A.: Measurements of cloud condensation nuclei activity and droplet activation kinetics of fresh unprocessed regional dust samples and minerals, Atmos. Chem. Phys., 11, 3527–3541, https://doi.org/10.5194/acp-11-3527-2011, 2011.

- Kuznetsov, B. V., Rakhmanova, T. A., Popovicheva, O. B., and Shonija, N. K.: Water adsorption and energetic properties of spark discharge soot: specific features of hydrophobicity, J. Aerosol Sci. 34, 1465–1479, https://doi.org/10.1016/S0021-8502(03)00366-5, 2003.
  - Laaksonen, A.: A unifying model for adsorption and nucleation of vapors on solid surfaces, J. Phys. Chem. A 119, 3736–3745, https://doi.org/10.1021/acs.jpca.5b00325, 2015.
- 420 Laaksonen, A., and Malila, J.: An adsorption theory of heterogeneous nucleation of water vapour on nanoparticles, Atmos. Chem. Phys. 16, 135–143, https://doi.org/10.5194/acp-16-135-2016, 2016.
  - Laaksonen, A., Malila, J., Nenes, A., Hung, H.-M., and Chen, J.–P.: Surface fractal dimension, water adsorption efficiency, and cloud nucleation activity of insoluble aerosol, Sci. Rep. 6, 25504, https://doi.org/10.1038/srep25504, 2016.
  - Lupi, L., Kastelowitz, N., and Molinero, V.: Vapor deposition of water on graphitic surfaces: formation of amorphous ice,
- 425 bilayer ice, ice I, and liquid water, J. Chem. Phys. 141, 18C508, https://doi.org/10.1063/1.4895543, 2014.
  - Mahata, P. C., and Alofs, D. J., Insoluble condensation nuclei: the effect of contact angle, surface roughness and adsorption, J. Atmos. Sci. 32, 116–122, https://doi.org/10.1175/1520-0469(1975)032<0116:ICNTEO>2.0.CO;2, 1975.

Marcolli, C.: Pre-activation of aerosol particles by ice preserved in pores, Atmos. Chem. Phys.,

17, 1595–1622, https://doi.org/10.5194/acp-17-1595-2017, 2017.

430 Möller, D.: On the history of the scientific exploration of fog, dew, rain and other atmospheric water, Erde 139, 11–44, 2008.

- Murphy, D. M., and Koop, T.: Review of the vapour pressures of ice and supercooled water for atmospheric applications, Q. J. R. Meteorol. Soc. 131, 1539–1565, https://doi.org/10.1256/qj.04.94, 2005.
- Niedermeier, D., Ervens, B., Clauss, T., Voigtländer, J., Wex, H., Hartmann, S., and Stratmann, F.: A computationally
   efficient description of heterogeneous freezing: A simplified version of the Soccer ball model, Geophys. Res. Lett.
   41, 736–741, https://doi.org/10.1002/2013GL058684, 2014.
  - Oubal, M., Picaud, S., Rayez, M.-T., and Rayez, J.-C.: Water adsorption on oxidized single atomic vacancies present at the surface of small carbonaceous nanoparticles modeling soot, ChemPhysChem 11, 4088–4096, https://doi.org/10.1002/cphc.201000364, 2010.
- 440 Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, Atmos. Chem. Phys., 7, 1961–1971, https://doi.org/10.5194/acp-7-1961-2007, 2007.
  - Popovicheva, O. B., Persiantseva, N. M., Tishkova, V., Shonija, N. K., and Zubareva, N. A.: Quantification of water uptake by soot particles, Environ. Res. Lett. 3, 025009, https://doi.org/ 10.1088/1748-9326/3/2/025009, 2008.

Pruppacher, H. R., and Klett, J. D.: Microphysics of Clouds and Precipitation, Kluwer Academic,

445 Dordrecht, ed. 2, 1997.

- Rowley, H. H., and Innes, W. B.: Relationships between the spreading pressure, adsorption, and wetting, J. Phys. Chem. 46, 694–705, https://doi.org/10.1021/j150421a002, 1942.
- Samseth, B. H., Stern, C. W., Andrews, E., Kahn, R. A., Myhre, G., Schulz, M., and Schuster, G. L.: Aerosol absorption: Progress towards global and regional constraints, Curr. Clim. Change Rep. 4, 65–83,
- 450 https://doi.org/10.1007/s40641-018-0091-4, 2018.
- Seinfeld, J. H., Bretherton, C. S., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S. J., Guenther, A. B., Kahn, R. A., Kracunas, I. P., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G., and Wood, R.: Improving our fundamental understanding of the role of aerosol-cloud interactions in the climate system, Proc. Natl. Acad. Sci. U.S.A. 113, 5781–5790, https://doi.org/10.1073/pnas.1514043113, 2016.

Sergi, D., Schocci, G., and Ortona, A.: Molecular dynamics simulations of the contact angle

between water droplets and graphite surfaces, Fluid Phase Equilibr. 332, 173–177, https://doi.org/10.1016/j.fluid.2012.07.010, 2012.

Sorjamaa, R., and Laaksonen, A., The effect of H<sub>2</sub>O adsorption on cloud drop activation of insoluble particles: a theoretical framework, Atmos. Chem. Phys. 7, 6175–6180, https://doi.org/10.5194/acp-7-6175-2007, 2007.

- Tadmor, R.: Line energy and the relation between advancing, receding, and Young contact angles, Langmuir 20, 7659–7664, https://doi.org/10.1021/la049410h, 2004.
- Tritscher, T., Jurányi, Z., Martin, M., Chirico, R., Gysel, M., Heringa, M. F., De-Carlo, P. F., Sierau, B., Prévôt, A. S. H., Weingartner, E., and Baltensperger, U.: Changes of hygroscopicity and morphology during ageing of diesel soot, Environ. Res. Lett. 6, 034026, https://doi.org/10.1088/1748-9326/6/3/034026, 2011.
- Wittbom, C., Eriksson, A. C., Rissler, J., Carlsson, J. E., Roldin, P., Nordin, E. Z., Nilsson, P. T., Swietlicki, E., Pagels, J. H., and Svenningsson, B.: Cloud droplet activity changes of soot aerosol upon smog chamber ageing, Atmos. Chem. Phys., 14, 9831–9854, https://doi.org/10.5194/acp-14-9831-2014, 2014.
- Young, G. J., Chessick, J. J., Healey, F. H., and Zettlemoyer, A. C., Thermodynamics of the adsorption of water on graphon
  from heats of immersion and adsorption data, J. Phys. Chem. 58, 313–315, https://doi.org/10.1021/j150514a006,
  1954.
  - Zheng, Y, and Zaoui, A.: Wetting and nanodroplet contact angle of the clay 2:1 surface: The case of Na-montmorillonite (001), Appl. Surf. Sci. 396, 717–722, https://doi.org/10.1016/j.apsusc.2016.11.015, 2017.

475

460

We thank all reviewers for useful comments. We would like to note that we found a minor bug in the program used for ANT calculations. After fixing the bug, the adsorption site distances used for calculating the coagulation nucleation curves in Fig. 4 are 4.2 and 4.4 nm instead of 3.6 and 3.8 nm. This change does not affect the discussion and conclusions in any way.

## Reviewer 2

- <sup>490</sup> The authors describe a new approach to parameterize the CCN activity of insoluble particles. Instead of the solute effect, they use the FHH-adsorption-isotherm to de-scribe the water adsorption on the surface of insoluble carbon black particles. The manuscript is well written and well structured. The scientific content is exciting and fills a long-existing knowledge gap. Therefore, I only have very few minor comments.
- 495 We thank Reviewer 2, our replies to the detailed comments are below

P3 L53 "as well" Corrected.

500 P3 L 58 "climate active" is a bit vague. I would suggest referring to radiative forcing and cloud formation directly

New formulation: ... different types of black carbon (BC) particles, which have long been known to be climate active – but with large uncertainty on their impacts on cloud formation and radiative forcing.

## 505

P7L130 It is a bit confusing when you say "droplet". Does this refer to a single droplet or to the water patches on the material

It refers to a single droplet growing on a BC particle. We have changed the sentences as follows: A single droplet on a BC particle grows along the red curve 1 until it reaches the critical supersaturation S<sub>cl</sub>\* and nucleates. However, if there are several similar droplets growing on the particle, and the average distance between them is small enough so that they coalesce to form a uniform liquid film at sufficiently low saturation ratio...

P11 L204 (Aria et al., 2016) investigated the wettability of graphene after exposure to ambient air. While this can be presentative for certain aspects of atmospheric black carbon particles, it should not be taken as an example for an atmospheric aging process. To point out, that exposure to atmospheric condition increases the interaction of black carbon with water, the following studies should be considered as an additional reference

Tritscher, T.; Jurányi, Z.; Martin, M.; Chirico, R.; Gysel, M.; Heringa, M. F.; De-Carlo, P. F.; Sierau, B.; Prévôt, A. S. H.; Weingartner, E.; et al. Changes of Hygroscopicity and Morphology 520 during Ageing of Diesel Soot. Environ. Res. Lett. 2011, 6 (3), 34026. https://doi.org/10.1088/1748-9326/6/3/034026. Grimon-prez, S.; Faccinetto, A.; Batut, S.; Wu, J.; Desgroux, P.; Petitprez, D. Cloud Condensation Nuclei from the Activation with Ozone of Soot Particles Sampled from a (8), Kerosene Diffusion Flame.Aerosol Sci.Technol.2018. 52 814-827. https://doi.org/10.1080/02786826.2018.1472367. Friebel, F.: Mensah, A. A. Ozone 525 Concentration versus Temperaturea Ă r: Atmospheric Aging of Soot Particles. Langmuir 2019, 35 (45), 14437–14450. https://doi.org/10.1021/acs.langmuir.9b02372

We have added these references.

## 530 **Reviewer 3**

The authors present measurements of heterogeneous nucleation of water on different types of carbon black. They tend to understand these measurements with a recently developed adsorption nucleation theory. The interesting finding is that with sufficient adsorption sites, soot particles can activate even when no soluble material is present, which is in contrast to the kappa-

535 Köhler theory. The paper is well-written and fits into the journal ACP. However, the paper is short and eventually should be published as the new MS type "ACP Letters". Anyway, there are some shortcomings which should be discussed before publication:

We thank reviewer 3 and would like to note that although ms is not very long, it is clearly over the 2500 word limit set for ACP letters. Our detailed replies are below.

## 540

Main comments The contact angle is a rather imprecise value as mentioned by the authors (line 121-130). However, the explanation why in the case of graphite the contact angle should be more trustable (+/-5) is not very convincing.

As a matter of fact, our sensitivity analysis shows that changing the contact angle by +/- 10 degrees and the FHH A-parameter simultaneously so that the FHH model still fits measured adsorption data well, the critical supersaturation is affected by less than 1% (we changed the y-axis of Fig. A3 to match those of Figs. 4 and 5). This is much less than the experimental uncertainty of the critical supersaturations of ozonolyzed Palas soot particles (Fig. 4). We have clarified this in the text, and removed to references to +/- 5 degrees.

More arguments should be presented considering the structure, morphology and chemistry of the respective graphite type and its impact on the contact angle and on the heterogeneous nucleation. Häusler et al. (2018) have measured the impact of graphene and graphene oxides on heterogeneous ice nucleation and found that structure, morphology and chemistry have an

555 important impact. The authors might discuss how they can parameterize such findings for the heterogeneous nucleation of water vapor.

Häusler et al. showed the importance of graphene lattice order as well as degree of surface oxidation and surface defects on immersion freezing. Compared with vapor-liquid nucleation, freezing is impacted by possible ice epitaxy, and therefore understanding the exact surface structure is even more important than in droplet formation. Nevertheless, we now discuss structure of the Palas soot in conjunction with possible impact of capillary menisci formed between primary soot spherules to the heterogeneous nucleation (see also our reply to J.C. Corbin).

565 Niedermeier et al. (2014) have developed a soccer ball model for heterogeneous ice nucleation relying on different contact angles on the surface of a nucleus. The authors might discuss how this model compares to their adsorption nucleation theory.

The soccer ball model is intermediate between stochastic and singular (or deterministic) nucleation theories. Classical nucleation theory is based on the idea that nucleation is stochastic phenomenon whereas ice nucleation theories that assume specific temperature dependent ice nucleation sites are deterministic. The adsorption nucleation theory is deterministic as one of its premises is that nucleation occurs on specific adsorption sites, and there is no stochastic element included in the theory. Such an element could be introduced similarly as in the soccer ball model, by assuming a distribution of contact angles between the adsorption sites. However, without knowledge of e.g. the possible widths of such distributions, we prefer not to include this idea at present. Having said that, molecular dynamics could possibly be used in the future to explore this issue with chemically and physically heterogeneous surfaces. We have added these considerations in the manuscript.

## 580

560

## Minor comments

Fig. 3: The caption of fig. 3 mentions more details in the text. However, the text doesnot explain how to read this figure and for what reason InN is plotted against In(-InS).

The reason is that according to the FHH theory, multilayer adsorption data should align linearly in such a plot. This has been added to the text.

Fig. A1: The x-axis theta should have the unit degree ( $\circ$ ). The label of the y-axis on the righthand site should be deleted. Fig.

590

A3: The supersaturation S\* should have the unit percent (%). The contact angle(theta) should have the unit degree ( $\circ$ )

Fixed

ReferencesHaeusler, T., Gebhardt, P., Iglesias, D., Rameshan, C., Marchesan, S., Eder, D.,Grothe, H. (2018), Ice Nucleation Activity of Graphene and Graphene Oxides, J. Phys.Chem. C 122, 15, 8182–8190, https://doi.org/10.1021/acs.jpcc.7b10675Niedermeier, D., B. Ervens, T. Clauss, J. Voigtländer, H. Wex, S. Hartmann, and F.Stratmann (2014), A computationally efficient description of heterogeneous freezing:A simplified version of the Soccer ball model, Geophys. Res. Lett., 41, 736–741, doi:10.1002/2013GL058684.

## J. C. Corbin

- <sup>605</sup> To this very interesting work, I would like to suggest that the authors briefly address the role of contact region between soot spherules (also known as monomers) in their discussion. Since real soot particles typically contain at least a few spherules, menisci between these spherules may alter the curvature of any condensed phase water. While this well-known phenomenon (Butt and Kappl, 2009) is probably outside of the scope of the present work, it may be worthwhile
- 610 to note at what point and by approximately what degree it might alter the authors' conclusions. For example, since capillary menisci form below saturation, how would the authors conclusions change for a particle in

which capillary condensation is normally imagined? I do realize that the macroscopic phenomenon of capillary condensation is not treated by a nucleation theory. I also realize that

- 615 experimental data have already been presented. Hence, I expect that only a brief comment would be necessary to clarify this question. I apologize if such a comment was already made, and I missed it. One might also hypothesize that the junctions between soot spherules may be important in the sense of being either more or less heterogeneous than the spherules. However, I am not aware of experimental evidence in support of this hypothesis.
- 620 Reference:Butt,Hans Jürgen,and Michael Kappl.2009."Normal CapillaryForces."Advances in Colloid and Interface Science 146 (1–2):48–60.https://doi.org/10.1016/j.cis.2008.10.002.

We would like to thank Dr. Corbin for this useful insight. There are two factors impacting possible capillary condensation. The first is that the contact angle we use is nearly 90 degrees, which means that a meniscus cannot in practice be concave, which in turn implies that capillary condensation will not occur at undersaturation. The second is the primary spherule sizes. Palas soot primary spherules have diameters on the order of 5-10 nm. Crouzet and Marlow (Aerosol Sci. Tech. 22, 43-59, 1995) have modelled the nucleation of

pendular water ring formed around the contact point of two equal sized spheres. It can be read from their Fig. 6 and Table 2 that when contact angle is 85 degrees, the supersaturation
 required for nucleation to occur is about 8%. However, this is when the sphere radius is 50 nm. It can also be seen from Table 2 that the critical supersaturation increases rapidly with sphere radius; for 200 nm spheres it would be just 2%. We can thus conclude that with Palas soot, it is unlikely that this type of capillary nucleation would take place at significantly lower supersaturation than we have calculated for our Fig. 4. We have added discussion in the manuscript.

## List of changes to the manuscript

640

650

Line 59: Added a few words related to climate impacts of BC

Lines 123-129: Replaced plus/minus 5 degrees by plus/minus 10 degrees when discussing contact angle uncertainty.

Lines 137 – 143: Added discussion related to stochastic vs deterministic nucleation theories.

Lines 197-199: Added a sentence to explain the logarithmic FHH plot.

Lines 214-215: Added references provided by Reviewer 2.

Lines 249 – 257: Added a paragraph discussing the possible capillary condensation/nucleation.

Lines 330-333: Added sentences pointing out the magnitude of the contact angle ucertainty in ANT calculations compared with nucleation experiment uncertainty.

Fig. 4: Made a new figure with correct adsorption site distances.

Fig. A1: Made a new version as suggested by Reviewer 3.

Fig. A3: Made a new version as suggested by Reviewer 3.