

~~Arctic marine~~ The ice nucleating aerosol: a laboratory study activity of Arctic sea surface microlayer samples and marine algal cultures

Luisa Ickes^{1,2*}, Grace C. E. Porter³, Robert Wagner², Michael P. Adams³, Sascha Bierbauer², Allan K. Bertram⁴, Merete Bilde⁵, Sigurd Christiansen⁵, Annica M. L. Ekman¹, Elena Gorokhova⁶, Kristina Höhler², Alexei A. Kiselev², Caroline Leck¹, Ottmar Möhler², Benjamin J. Murray³, Thea Schiebel², Romy Ullrich², and Matthew Salter⁶

¹Department of Meteorology & Bolin Centre for Climate Studies, Stockholm University, Stockholm, Sweden

²Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

³School of Earth and Environment, University of Leeds, Leeds, United Kingdom

⁴Department of Chemistry, University of British Columbia, Vancouver, Canada

⁵Department of Chemistry, Aarhus University, Aarhus, Denmark

⁶Department of Environmental Science and Analytical Chemistry & Bolin Centre for Climate Studies, Stockholm University, Stockholm, Sweden

*Now at: Department of Space, Earth and Environment, Chalmers, Gothenburg, Sweden

Correspondence: Luisa Ickes (luisa.ickes@misu.su.se)

Abstract. In recent years, sea spray and the biological material it contains has received increased attention as a source of ice nucleating particles (INPs). Such INPs may play a role in remote marine regions, where other sources of INPs are scarce or absent. [In the Arctic, these INPs can influence water-ice partitioning in low-level clouds and thereby the cloud lifetime, with consequences for the surface energy budget, sea ice formation and melt, and climate.](#) Marine aerosol is of diverse nature, so identifying sources of INPs is challenging. One fraction of marine bioaerosol, phytoplankton and their exudates, has been a particular focus of marine INP research. In our study we attempt to address three main questions. Firstly, we compare the ice nucleating ability of two common phytoplankton species with Arctic seawater microlayer samples using the same instrumentation to see if these phytoplankton species produce ice nucleating material with sufficient activity to account for the ice nucleation observed in Arctic microlayer samples. We present first measurements of the ice nucleating ability of two predominant phytoplankton species, *Melosira arctica*, a common Arctic diatom species and *Skeletonema marinoi*, a ubiquitous diatom species across oceans worldwide. To determine the potential effect of nutrient conditions and characteristics of the algal culture, such as the amount of organic carbon associated with algal cells, on the ice nucleation activity, the *Skeletonema marinoi* was grown under different nutrient regimes. From comparison of the ice nucleation data of the algal cultures to those obtained from a range of sea surface microlayer (SML) samples obtained during three different field expeditions to the Arctic (ACCACIA, NETCARE, ASCOS) we found that although these diatoms do produce ice nucleating material, they were not as ice active as the investigated microlayer samples. Secondly, to improve our understanding of local Arctic marine sources as atmospheric INP we applied [several two](#) aerosolisation techniques to analyse the ice nucleating ability of aerosolised microlayer and algae samples. The aerosols were generated either by direct nebulisation of the undiluted bulk solutions, or by the addition of the samples to a sea spray simulation chamber filled with artificial seawater. The latter method generates aerosol

20 particles using a plunging jet to mimic the process of oceanic wave-breaking. We observed that the aerosols produced using
this approach can be ice active indicating that the ice nucleating material in seawater can indeed transfer to the aerosol phase.
Thirdly, we attempted to measure ice nucleation activity across the entire temperature range relevant for mixed-phase clouds
using a suite of ice nucleation measurement techniques- an expansion cloud chamber, a continuous flow diffusion chamber,
and a cold stage. In order to compare the measurements made using the different instruments, we have normalised the data in
25 relation to the mass of salt present in the nascent sea spray aerosol. At temperatures above 248 K some of the SML samples
were very effective at nucleating ice, but there was substantial variability between the different samples. In contrast, there was
much less variability between samples below 248 K. [We discuss our results in the context of aerosol-cloud interactions in the
Arctic with a focus on furthering our understanding of which INP types may be important in the Arctic atmosphere.](#)

1 Introduction

30 Clouds have a strong impact on the energy balance and therefore play an important role in the Earth's climate system
(Chahine, 1992; Boucher et al., 2013). They are particularly important in the high-latitudes, one of the regions most sensi-
tive to global warming (Stocker et al., 2013), where they not only influence the energy budget (Garrett et al., 2009; Mor-
rison et al., 2012), but also the subsequent melting and freezing of sea ice (Intrieri et al., 2002; Pithan and Mauritsen,
2014). As such, they are involved in several climate feedback processes. The radiative characteristics of clouds depend on
35 their microphysical structure, e.g. if the cloud consists of water droplets or ice crystals. Mixed-phase clouds which are
comprised of both ice crystals and super-cooled water droplets are common in the high Arctic (Shupe et al., 2006). For-
mation of liquid cloud droplets requires the presence of an aerosol particle that facilitates water vapour condensation on
its surface (so-called cloud condensation nuclei – CCN). Aerosol particles are also necessary for the initiation of primary
ice formation within these clouds by a process known as heterogeneous freezing (so-called ice nucleating particles – INP).
40 Typically, only a small fraction of aerosol particles has the ability to nucleate ice. [The-Despite increasing interest in INP
\(Szyrmer and Zawadzki, 1997; Hoose and Möhler, 2012; DeMott et al., 2010\), it is still uncertain which](#) types of aerosol parti-
cles that constitute good INP [are uncertain \(DeMott et al., 2010\) in the atmosphere \(Kanji et al., 2017\)](#). Aerosol particles known
to nucleate ice crystals by heterogeneous freezing in mixed-phase clouds include mineral dust, volcanic ash and primary biolo-
gical particles, such as pollen, fungi and bacteria, and fragments of those (Hoose and Möhler, 2012). Those are aerosol particles
45 with a predominantly terrestrial source. However, there are regions which are relatively isolated from terrestrial sources, such
as the summer high Arctic, remote parts of North Atlantic, North Pacific and Southern Ocean. In such regions, sea spray aerosol
could be an important source of INP ([Burrows et al., 2013; Yun and Penner, 2013; Vergara-Temprado et al., 2017; Huang et al., 2018](#)) ([Burrows et al., 2013](#)).

The potential for marine environments to act as sources of INPs was first investigated during the [1970s and 80s-1960s](#) (see Ta-
ble 1). This area of research has attracted renewed attention in more recent years. Indeed, recent observations indicate that bio-
50 genic material present at both the interface between the ocean and atmosphere, the so-called sea surface microlayer (SML), and
within nascent sea spray aerosol can be ice active, e.g. [Knopf et al. \(2011\); Wilson et al. \(2015\); DeMott et al. \(2016\); Irish et al. \(2017\)](#) ([Knopf et al. \(2011\)](#)).
Previous studies can be separated into three main groups: (i) ambient ice nucleation measurements in marine environments,

(ii) studies investigating the ice nucleating potential of seawater and SML samples, and (iii) studies concerned with the ice nucleating potential of different phytoplankton species and their exudates (Table 1). One of the key recent studies concerned with whether sea spray aerosol contains significant amounts of INPs was conducted by DeMott et al. (2016) who examined the ice nucleation potential of laboratory generated nascent sea spray aerosol particles and compared their findings with measurements of ambient marine aerosol. Critically, they observed that laboratory generated sea spray aerosol has a similar ice nucleation activity to ambient marine aerosols and that the ice nucleating activity of nascent sea spray aerosol strongly increased in association with phytoplankton blooms. Given these observations, the authors conclude that the INP present in sea spray aerosol are likely linked to organic matter associated with phytoplankton blooms. DeMott et al. (2016) also showed that different INP types were active at different temperatures (~~266.15, 258.15, 250.15, 247.15, 243.15 K~~). Despite the finding that significant amounts of ice active material are present in nascent sea spray aerosol, the measured number concentration of INP in ambient marine aerosol was still several orders of magnitude lower than equivalent measurements in ambient terrestrial aerosol. Another relevant study was conducted by Wilson et al. (2015) who analysed SML samples collected in the Atlantic and Arctic oceans. The ice activity of these samples was highly variable with the temperature at which half of the sample droplets froze, the so-called median freezing temperature, ranging from approximately 265 to 248 K. Based on tests with samples that have been filtered and heated, these authors concluded that submicron biogenic material was likely responsible for the ice activity of seawater samples from a range of locations. This suggests that whole cells are not responsible for the observed ice nucleation (Schnell and Vali, 1975; Wilson et al., 2015; Irish et al., 2017). Further, exudates of the marine diatom *Thalassiosira pseudonana*, a widespread phytoplankton species, have been shown to nucleate ice (Knopf et al., 2011; Wilson et al., 2015; Ladino et al., 2016); hence it has been proposed that organic material associated with phytoplankton cell exudates may explain the ice nucleation activity of marine SML samples. However, Knopf et al. (2011) also found that intact cells are effective INP in the mixed-phase temperature regime. Another hypothesis is that bacteria play a role as shown by e.g. Fall and Schnell (1985).

Motivated by these previous studies we have analysed the freezing potential of two common phytoplankton species, *Melosira arctica* and *Skeletonema marinoi*. *Skeletonema marinoi* is a very common diatom species, especially in temperate coastal regions during the spring bloom (Kooistra et al., 2008). *Melosira arctica* on the other hand is the most productive algae in the Arctic Ocean (Booth and Horner, 1997). [Melosira arctica was found along with polymer gels in high Arctic cloud water samples \(Orellana et al., 2011\)](#). Environmental factors, such as light and nutrient supply, have a high potential to affect the biochemical composition of phytoplankton and thus biogenic exudate material. ~~It has been suggested that absolute cell concentrations are not the sole determining factor for aerosol flux and that aerosol size distribution can be affected by the growth conditions of the microorganisms (Alpert et al., 2015)~~[The degree to which both the flux and composition of sea spray aerosol is affected by biological activity in the surface ocean is a longstanding question in the field. However, studies have suggested that the aerosol flux may not only be impacted by the absolute cell concentrations of phytoplankton but also their growing conditions, e.g. Alpert et al. \(2015\)](#). Thus those environmental factors have an effect on the presence of INPs coming from marine sources as well. Therefore, algae grown under different nutrient regimes may differ in their INP ability, which is investigated in this study. *Skeletonema marinoi* was cultivated with different nutrition levels in order to mimic nutrient limitation and growth inhibition in phytoplankton. This leads to a variation in the carbon content of each cell and thus in the

cell suspensions, which enables us to investigate the resulting effects of different growth rates and cell carbon content on ice nucleation. Our aim here was to investigate whether changing these cell properties has any impact on the ice nucleation activity
90 of the phytoplankton.

Another goal of this study was to improve our understanding of whether Arctic marine regions may have local sources of marine INPs. Although it has been found that organic matter with marine origin is prevalent in aerosol particles present in the high Arctic during summer ([Leck et al., 2002](#)) e.g. [Leck et al. \(2002\)](#) and that marine organic matter nucleates ice e.g. Wilson et al. (2015), the ice nucleating potential of the aerosolised organic matter has not been examined in detail [for the](#)
95 [Arctic region](#). Therefore, we have determined the heterogeneous ice nucleating ability of artificial seawater containing two phytoplankton species cultured in the laboratory along with samples of SML collected during a series of field campaigns in the North Atlantic and Arctic Oceans. Within this study, two different aerosolisation techniques were utilised to test the impact of the aerosol generation method on the ice nucleation behaviour of the resulting particles.

Measurements have been made with a variety of ice nucleation measurement techniques and all measurements were conducted under conditions relevant for mixed-phase clouds, i.e. above about 235 K and at water saturation. We have utilised a
100 number of different experimental methods to derive the ice nucleating ability of our samples, with the ultimate goal of merging these different measurements across the full temperature range relevant for mixed-phase clouds. Through comparison of the ice nucleation activity of artificial seawater containing *Melosira arctica* with that of the SML samples we aim to shed light on how representative relevant algal cultures are for Arctic marine INP.

A description of the methods of sample collection and cultivation as well as the experimental setup and ice nucleation
105 measurement techniques are introduced in Sect. 2. The results of the ice nucleation measurements and a comparison with previous marine INP measurements found in the literature are presented in Sect. 3. Since we have made measurements across the full temperature range relevant for mixed-phase clouds (273.15 K until 233.15 K) this section is split into three parts. The first part (Sect. 3.1) focuses on the measurements at temperatures above 248 K referred to as the "high temperature regime"
110 throughout this article, while the second part (Sect. 3.2) focuses on the measurements conducted at temperatures below 248 K referred to as the "low temperature regime" throughout this article. In the final part (Sect. 3.3) we present an integrated spectrum over the full temperature range. Finally, we conclude this study with a summary of the major findings and discussion of potential atmospheric implications of our results (Sect. 4).

2 Methods and experimental setup

To determine the ice nucleating ability of our samples we have used three independent methods (Fig. 1). Firstly, bulk cell
115 suspensions of the algal cultures and field samples were aerosolised using a nebuliser and the generated particles were injected into the Aerosol Interaction and Dynamics in the Atmosphere (AIDA) aerosol and cloud chamber (Möhler et al., 2008). The ice nucleation behaviour of the particles was then either measured in situ in the AIDA chamber by performing an expansion cooling experiment, or by probing the particles *ex-situ* with a continuous flow diffusion chamber (CFDC) called INKA [Ice Nucleation
120 instrument of the Karlsruhe Institute of Technology; Schiebel (2017)]. Secondly, for a subset of the samples, a certain volume

Table 1. An overview of previous laboratory and field studies which have either investigated nascent sea spray aerosol particles as INP or ambient INP in marine regions (including SML/seawater samples) [in the temperature range of mixed-phase clouds](#). The location ("Loc.") of each of the field studies is given. Laboratory studies are indicated as "Lab". The "Data" column indicates how the ice nucleation activity was estimated- usual measures are as amount of INP per m³ or L, the frozen fraction *FF* as a function of temperature or the median freezing temperature *T*₅₀, i.e. the temperature at which 50% of the droplets were frozen. Where relevant, the "Subst." column states specific substances or species that were studied. The column "Instr." provides information about the instrument(s) used in the study. [Here different cloud chambers are simplified by the term "cloud chambers", with the reference of the basic principle given in the index \(1: Warner \(1957\), 2: Bigg \(1957\), 3: Bigg et al. \(1963\), 4: Langer et al. \(1967\), 5: Stevenson \(1968\), 6: Gagin and Arroyo \(1969\), 7: Langer and Rodgers \(1975\)\).](#) [CFDC stands for continuous flow diffusion chamber. Cold stage, freezing assays etc. are all described with the term "drop freez."](#)

(i) Ambient ice nucleation measurements in marine environments

Study	Loc.	Data	Subst.	Instr.
Kline and Brier 1958	Washington DC	INP/L	Airborne	Cloud chamber¹
Isono et al. 1959	Tokyo (Pacific)	INP/L	Airborne	Cloud chamber²
Battan and Riley 1960	Arizona (Gulf of Mexico)	INP/L	Airborne	Cloud chamber¹
Kline 1960	Washington DC	INP/L	Airborne	Cloud chamber ¹
Bigg 1973	Southern Ocean (SO)	INP/m ³	Airborne	Filter & cloud chamber⁶
Radke et al. 1976	Alaska	INP/L	Airborne	Filter & dyn. chamber ⁴
Schnell 1977	Canada (Atlantic)	INP/m ³	Airborne	Filter & therm.-diff.chamber drop freez.
Flyger and Heidam 1978	Northern Greenland	INP/L	Airborne	Filter & cloud chamber⁵
Borys and Grant 1983	Arctic	INP/m³	Airborne	Filter & cloud chamber⁵
Nagamoto et al. 1984	Florida	INP/m ³	Airborne	Filter & dyn.-chamber cloud chamber⁷
Fountain and Ohtake 1985	Alaska	INP/L	Airborne	Filter & cloud chamber³
Rosinski et al. 1986	Pacific	Freez. T, INP/m³	Airborne	Filter & drop freez. & cloud chamber⁷
Rosinski et al. 1987	Pacific	Freez. T, INP/m³	Airborne	Filter & dyn. chamber ⁷
Borys 1989	Arctic	INP/m³	Airborne	Filter & cloud chamber⁷
Bigg 1990	SO, Hawaii	INP/m³	Airborne	Filter & cloud chamber^{3,5,6}
Rosinski et al. 1995	East China Sea	INP/m ³	Airborne	Filter & dyn. chamber⁷
Rosinski 1995	Washington State (North Pacific)	INP/m³	Airborne	Filter & dyn. chamber⁷
Bigg 1996	Arctic	INP/m³	Airborne	Filter & cloud chamber⁵
DeMott et al. 2016	Caribbean, Arctic, Canada , Canada, Pacific, Pacific, Lab (MART)	INP/L	Airborne	CFDC, filter (CSU)& drop freez.
Mason et al. 2015	Lab (MART) Canada (North Pacific)	INP/L	Airborne	Filter & drop freez.
Ladino et al. 2016	Canada (North Pacific)	INP/L	Airborne	CFDC
Wex et al. Creamean et al. 2018	Arctic	INP/L	Airborne	Filter & drop freez.
McCluskey et al. 2018	Mace head	INP/L	Airborne	CFDC, filter & drop freez.
McCluskey et al. 2018	SO	INP/m³	Airborne	CFDC, filter & drop freez.
Si et al. 2018	Canada/Arctic	INP/L	Airborne	Filter & drop freez.
Welti et al. 2018	Atlantic	INP/m³	Airborne	CFDC, filter & drop freez.

Table 1. Continued.

(i) Ambient ice nucleation measurements in marine environments

<u>Study</u>	<u>Loc.</u>	<u>Data</u>	<u>Subst.</u>	<u>Instr.</u>
<u>Creamean et al. 2019</u>	Arctic	INP/L	Airborne	Drop -Filter & drop freez.
<u>Gong et al. 2019</u>	<u>Cyprus</u>	<u>INP/L</u>	<u>Airborne</u>	<u>Filter & drop freez.</u>
Irish et al. 2019a	Arctic	INP/L	Airborne	Filter & drop freez.
Creamean et al. <u>Ladino et al. 2019</u>	<u>Gulf of Mexico</u>	<u>INP/L</u>	<u>Airborne</u>	<u>Filter & drop freez.</u>
<u>Si et al. 2019</u>	Arctic	INP/L	Airborne	Drop -Filter & drop freez.
<u>Wex et al. 2019</u>	<u>Arctic</u>	<u>INP/L</u>	<u>Airborne</u>	<u>Filter & drop freez.</u>
<u>Hartmann et al. 2020</u>	<u>Arctic</u>	<u>INP/L</u>	<u>Airborne</u>	<u>Filter & drop freez.</u>
<u>Gong et al. 2020</u>	<u>Cape Verde (Atlantic)</u>	<u>INP/L</u>	<u>Airborne</u>	<u>Filter & drop freez.</u>
<u>Welti et al. 2020</u>	<u>Arctic, Atlantic, Pacific, SO</u>	<u>INP/m³</u>	<u>Airborne</u>	<u>CFDC, filter & drop freez.</u>

of the bulk solutions was added to 20 L of artificial seawater in the mobile Aarhus University sea spray simulation chamber called AEGOR (Christiansen et al., 2019). Aerosol particles generated by bubble bursting in AEGOR were injected into the AIDA chamber in the same manner as the particles generated using the nebuliser and their ice nucleation activity was measured both in AIDA expansion cooling experiment and with INKA. Thirdly, the INP abundance within the liquid samples used to generate aerosols was determined using the microliter nucleation by immersed particle instrument (μ l-NIPI), where droplets of the bulk solutions were pipetted onto a cold stage (Whale et al., 2015).

Additionally, it was investigated if material from the same algal cultures and SML samples affects the ability of sea spray aerosols to act as CCN. The measurements of the CCN-derived hygroscopicity and the implication on Arctic clouds are presented in a companion study, see Christiansen et al. (2020, submitted to J. Geophys. Res.).

130 2.1 Samples and sample treatment

Two types of samples were investigated in this study: algal cultures (*Skeletonema marinoi* and *Melosira arctica*) and SML samples. One diatom species (*Skeletonema marinoi*) was grown under different conditions. The SML samples were collected during three field expeditions in the Arctic region [ACCACIA (Wilson et al., 2015), NETCARE (Irish et al., 2019b) and ASCOS (Gao et al., 2012)]. Table 2 provides an overview of how the samples were analysed and summarises all the measurements conducted during this campaign.

Culture conditions and nutrient regimes for algae

The two diatoms were cultured axenically in Guillard's f/2+Si medium in two-liter glass bottles on a shaking table (0.5 rpm/min) inside a climate chamber. Algal growth rate and number of cells per colony were monitored using the cell counter TC20 (Bio-Rad). *Skeletonema marinoi* (CCAP 1077/5; Göteborg University Marine Algal Culture Collection, GUMACC) was

Table 1. Continued.

(ii) Studies investigating the ice nucleating potential of seawater and SML samples

Study	Loc.	Data	Subst.	Instr.
Birstein and Anderson 1953	Artificial	Freez. T	Sea salt	Cloud chamber
Brier and Kline 1959	Washington DC	INP/L	Seawater	Cloud chamber¹
Schnell and Vali 1975	Pacific (N/S), Caribbean Atlantic	INP/m ³	Seawater	Drop freez.
Schnell and Vali 1976	Canada, California, Bahamas	INP/m ³	Seawater	Drop freez.
Schnell 1977	Atlantic (Canada)	INP/m ³	Seawater	Drop freez.
Parker et al. 1985	Antarctica	<i>FF</i>	Sea ice	Drop freez.
Rosinski et al. 1988	Gulf of Mexico	INP/m ³	Seawater	Dyn-chamber Cloud chamber⁷
Wilson et al. 2015	Arctic, Canada N. Pacific, Atlantic	<i>FF</i> , n_m	Seawater	Drop freez.
Irish et al. 2017	Arctic	<i>FF</i>	Seawater	Drop freez.
McCluskey et al. 2017	Lab (MART)	INP/L	Seawater	CFDC, filter (CSU) & drop freez.
Irish et al. 2019b	Arctic	INP/L & , <i>FF</i>	Seawater	Drop freez.
Creamean et al. 2019	Arctic	INP/L	Seawater	Drop freez.
Wilbourn et al. 2020	North Atlantic	Freez. T, <i>FF</i>	Aerosolized Sea water	Drop freez.
Gong et al. 2020	Cape Verde (Atlantic)	INP/L	Seawater	Drop freez.
Wolf et al. 2020	Florida Straits, North Pacific	Active site density	Sea water	CFDC

(iii) Studies concerned with the ice nucleating potential of different phytoplankton species and their exudates

Study	Loc.	Data	Subst.	Instr.
Schnell 1975	Lab	INP/m ³	Phytoplankton	Drop freez.
Parker et al. 1985	Lab	<i>FF</i>	Mar. bacteria	Drop freez.
Fall and Schnell 1985	Lab	T_{50}	Mar. bacteria	Drop freez.
Alpert et al. 2011	Lab	Freez. T	Aqu. NaCl, diatoms	Drop freez.
Alpert et al. 2011	Lab	Freez. T	Phytoplankton	Drop freez.
Knopf et al. 2011	Lab	Freez. T	Mar. diatoms	Drop freez.
Ladino et al. 2016	Lab	<i>FF</i>	Phytoplankton, mar. bacteria	CFDC
McCluskey et al. 2017	Lab	INP/L	Phytoplankton	CFDC, filter (CSU) & drop freez.
DeMott et al. 2018	Lab	T_{50}	Fatty acids	Drop freez.
Tesson and Šantl Temkiv 2018	Lab	Freez. T	Micro-algae	Drop freez.
Wilbourn et al. 2020	North Atlantic	Freez. T	Phytoplankton	Drop freez.




Figure 1. Schematic of the various aerosolisation (sea spray chamber AEGOR and nebuliser) and ice nucleation [Aerosol Interaction and Dynamics in the Atmosphere (AIDA) aerosol and cloud chamber, Ice Nucleation Instrument of the Karlsruhe Institute of Technology (INKA) and microliter nucleation by immersed particle instrument (μ L-NIPI)] measurement techniques employed in this study.

140 isolated from the Long Island Sound (Milford Harbour, USA). *Melosira arctica* (MATV-1402; Helsinki University) originated from the Western Gulf of Finland, the Baltic Sea.

Skeletonema marinoi (SM) was grown at 26 PSU, 293.15 K using a 12:12 h light:dark cycle at $90 \mu\text{mol photons m}^{-2} \text{s}^{-1}$. Concentrations of nitrate and phosphate in the media were adjusted to conform to three experimental conditions in order to manipulate growth rates and cell carbon content: (1) nutrient-replete conditions (SM100; high growth, high nutrient content of cells), (2) 60% nutrient-saturation (SM60; high growth but low nutrient content), and (3) low-nutrient treatment (SM10; low growth, low nutrient content). The respective nitrate and phosphate concentrations were 5 and $1 \mu\text{M}$ in SM100, 3 and $0.6 \mu\text{M}$ in SM60, and 0.2 and $0.1 \mu\text{M}$ in SM10 treatments. Nutrients, but also their stoichiometric ratios, determine growth. In treatment (1), we had algae that were both dividing fast and had a large cell size. In treatment (2), the cell division rate was high, but the cell size was small (phosphorus limitation). In treatment (3), both parameters were low (nitrogen and phosphorus limitation).

150 The algae were harvested, i.e. the entire culture volume was transferred to a plastic bag and frozen, when reaching a density of $\sim 3 \times 10^5$ and $\sim 5 \times 10^6$ cells/mL in the nutrient-replete and nutrient-sufficient (SM100 and SM60) conditions, respectively. Due to poor growth in SM10, the culture was harvested simultaneously with the other two treatments before reaching comparable cell densities.

Melosira arctica (MA) was grown at 6 PSU, 278.15 K and a 16:8 h light:dark cycle at $60 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ and harvested 155 when they reached $\sim 2 \times 10^5$ cells/mL. This culture is referred to as MA100.

Immediately after collection, the harvested algae were frozen for storage and transport at 193.15 K. We assume that freezing the samples does not influence the results of the experiments, an assumption supported by the literature (Schnell and Vali, 1976; Irish et al., 2019b). Prior to freezing, a sub-sample of known volume from each species/treatment was collected on a

Table 2. Overview of the measurements conducted in this study. The first column lists all the different samples investigated (see Section 2.1) including information on the campaigns during which the field samples were collected. The type of the sample is given in the second column. The aerosolisation techniques used for the AIDA measurements is denoted in the third column while the fourth column lists all the ice nucleation instruments used to probe the sample. The fourth column shows the date of the experiments. The fifth and six column presents the results of the fitted aerosol size distribution: the median particle diameter D and the width (geometric standard deviation) for all AIDA expansions (rounded to two digits after the comma).

Sample name	Type	Aerosolisation techniques		Date (AIDA expansion)	Fitted	Fitted
		(AIDA)	Instruments		med. D [μm]	std. dev. (AIDA)
Sigma-Aldrich sea salt	Artificial	Nebuliser, AEGOR <u>AEGOR</u>	AIDA, μl -NIPI	27.01.2017,	<u>0.59</u>	<u>1.41</u>
				30.01.2017,	<u>0.31</u>	<u>2.34</u>
				06.02.2017	<u>0.43</u>	<u>2.68</u>
SM100	Cultured	Nebuliser, AEGOR <u>AEGOR</u>	AIDA, μl -NIPI, INKA	06.02.2017, 07.02	<u>0.71</u>	<u>1.38</u>
				<u>21.02</u> .2017,	<u>0.76</u>	<u>1.37</u>
				08.02 <u>07.02</u> .2017, 21.02	<u>0.43</u>	<u>2.46</u>
				<u>08.02</u> .2017	<u>0.46</u>	<u>2.23</u>
SM60	Cultured	Nebuliser	AIDA, μl -NIPI, INKA	08.02.2017	<u>0.73</u>	<u>1.43</u>
SM10	Cultured	Nebuliser, AEGOR <u>AEGOR</u>	AIDA, μl -NIPI, INKA	16.02.2017,	<u>0.72</u>	<u>1.47</u>
				17.02.2017	<u>0.49</u>	<u>2.83</u>
MA100	Cultured	Nebuliser, AEGOR <u>AEGOR</u>	AIDA, μl -NIPI, INKA	22.02.2017,	<u>0.41</u>	<u>1.33</u>
				23.02.2017	<u>0.74</u>	<u>2.8</u>
STN2 (NETCARE)	SML	Nebuliser	AIDA, μl -NIPI	10.02.2017	<u>0.82</u>	<u>1.30</u>
STN3 (NETCARE)	SML	Nebuliser	AIDA, μl -NIPI, INKA	15.02.2017	<u>0.77</u>	<u>1.38</u>
STN7 (NETCARE)	SML	Nebuliser	AIDA, μl -NIPI, INKA	15.02.2017	<u>0.77</u>	<u>1.39</u>
SML5 (ACCACIA)	SML	Nebuliser, AEGOR <u>AEGOR</u>	AIDA, μl -NIPI, INKA	01.02.2017,	<u>0.59</u>	<u>1.47</u>
				02.02.2017	<u>0.7</u>	<u>2.61</u>
SML8 (ACCACIA)	SML	Nebuliser, AEGOR <u>AEGOR</u>	AIDA, μl -NIPI, INKA	31.01.2017	<u>0.88</u>	<u>1.21</u>
					<u>0.40</u>	<u>2.9</u>
SML16 (ACCACIA)	SML	Nebuliser	AIDA, μl -NIPI, INKA	03.02.2017	<u>0.86</u>	<u>1.25</u>
SML17 (ACCACIA)	SML	Nebuliser	AIDA, μl -NIPI, INKA	09.02.2017	<u>0.83</u>	<u>1.36</u>
SML19 (ACCACIA)	SML	Nebuliser	AIDA, μl -NIPI, INKA	03.02.2017	<u>0.90</u>	<u>1.27</u>
ASCOS (< 5 kDa)	SML	Nebuliser	AIDA, μl -NIPI	23.02.2017	<u>0.75</u>	<u>1.43</u>
ASCOS (foam)	SML	Nebuliser	AIDA, μl -NIPI	24.02.2017	<u>0.85</u>	<u>1.3</u>
ASCOS (5 kDa to 0.22 μm)	SML	Nebuliser	AIDA, μl -NIPI	24.02.2017	<u>0.18</u>	<u>1.27</u>

Table 3. Characteristics of the bulk samples used during the study: non-purgeable organic carbon content, the water activity of the artificial seawater, algal cultures and two SML samples, the algae cells per mL of the cultures and the carbon cell content of the cultures. For the diluted samples we give in brackets-parentheses how many mL of sample where added to 20 L of artificial seawater (3.5 wt% solution of the synthetic Sigma-Aldrich sea salt mixture in ultrapure water) in the AEGOR sea spray tank (see Sect. 2.2). For the samples indicated with "pure" the undiluted sample was used. The water activity of the samples was estimated directly using the dewpoint. These measurements were repeated three times, resulting in the standard deviations (STD) given here.

Sample name	Non-purgeable organic carbon	Water activity	Water activity	Algae cells	Carbon cell content
	[mg C L ⁻¹]	Dewpoint (dewpoint)	STD (dewpoint)	[mL ⁻¹]	[μgC mL ⁻¹]
SM100 (pure)	14.3	0.9871	0.0004	5280000	105.6
SM10 (pure)	5.1	0.9916	0.0005	350000	9.8
MA100 (pure)	10.9	0.9861	0.0006	188700	245.31
Sigma-Aldrich sea salt (pure)	1.1	0.9854	0.0004		
SM100 (79 mL in AEGOR)	2.3	0.9861	0.0008	20774	0.42
SM100 (406 mL in AEGOR)	1.7	0.9838	0.0006	105051	2.1
SM10 (approx. 900 mL in AEGOR)	0.9	0.9855	0.0002	15072	0.42
MA100 (893 mL in AEGOR)	3.2	0.9861	0.0006	1	10.49
SML8 (200 mL in AEGOR)	1.6	0.9866	0.0004		
SML5 (100 mL in AEGOR)	1.1	0.9857	0.0002		

0.2 μm filter for dry weight (DW), C and N analysis. The non-purgeable organic carbon content and the water activity of each sample was measured after the experiments. These values are summarised in Table 3.

Field samples

The SML samples were collected from different locations in the Arctic. A subset of the samples were collected during the Aerosol-Cloud Coupling and Climate Interactions in the Arctic (ACCACIA) expedition in July and August, 2013 in the Arctic Atlantic [East of Greenland and North of Spitsbergen, for more details see Wilson et al. (2015)]. Another subset of samples were collected as part of the Network on Climate and Aerosols: Addressing Key Uncertainties in Remote Canadian Environments (NETCARE) project during July and August, 2016 in the Eastern Canadian Arctic [for more details see Irish et al. (2019b)]. During ACCACIA and NETCARE a remote-controlled sampling catamaran was used for collection [ACCACIA: Knulst et al. (2003); Matrai et al. (2008); NETCARE: Shinki et al. (2012)]. Previous analysis of these samples in terms of ice nucleating ability can be found in the respective publications (Wilson et al., 2015; Irish et al., 2019b). The third subset of samples originates from the Arctic Summer Cloud Ocean Study (ASCOS) in August 2008 (Tjernström et al., 2014). The surface microlayer

water was collected from an open lead using the same sampling catamaran used during the ACCACIA campaign. The sample investigated in this study was collected on August 17 in 2008 at ca. 88°N and treated afterwards in three different ways. Two subsamples were subjected to a two-step ultrafiltration procedure. Firstly, the sample was passed through Millipore membrane filters (nominal pore size 0.22 μm) under mild vacuum. Secondly, the filtered samples were ultrafiltered and diafiltered through a tangential flow filtration system (TFF, Millipore) equipped with cartridges with a molecular weight cut off of 5 kDa. The fraction that passed through the 0.22 μm filters but not the TFF system is referred to as high molecular weight dissolved organic matter (5 kDa to 0.22 μm). To obtain even greater separation into low molecular weight dissolved organic matter, sample which passed through the TFF system was further filtered in an Amicon® stirred cell (< 5 kDa). The third subsample is a foam layer sample. Seawater without pre-filtration was fed directly into a pre-cleaned glass tower (15.3 L, 2 m in height). Purified zero air was forced into the system through a sintered glass frit (nominal pore size 15 - 25 μm) from the bottom of the tower at a flow rate of 150 mL min^{-1} . After the bubble experiment, seawater at the uppermost layer (about 3 cm) together with foamy substances were slowly overflowed into a collecting flask by an additional feeding of seawater from the middle of the tower. The collected water from the top layer consisting of both foam and background seawater is referred to as foam layer sample. The foam sample should be similar to an unfiltered SML sample (as obtained during ACCACIA and NETCARE). More details on the methods of filtration applied during ASCOS can be found in Gao et al. (2012).

All samples were immediately frozen at 193.15 K for storage and transport. The field samples are labelled according to the original names in the respective publications: the samples originating from the field expedition ACCAIA are called SML (purple, green and turquoise colours in the figures), the samples from NETCARE STN (blue colours in the figures) and the samples from ASCOS are called ASCOS (red and yellow colours in the figures). The numbers refer to the original sample numbers.

2.2 Aerosolisation techniques

Two different techniques were used to aerosolise samples into the AIDA cloud chamber. Firstly, undiluted samples were aerosolised using an ultrasonic nebuliser (GA2400, SinapTec) and injected directly into the AIDA chamber. An injection period of 20-30 minutes was sufficient to fill the AIDA chamber with an aerosol number concentration of approx. 550 cm^{-3} . Secondly, we used the temperature-controlled sea spray simulation chamber, AEGOR, with the aim of generating bubble-bursting aerosols in a more representative manner (Christiansen et al., 2019). The sea spray tank was filled with 20 L of artificial seawater (3.5 wt% solution of the synthetic Sigma-Aldrich sea salt mixture, product number S9883, in ultrapure water). Sigma-Aldrich sea salt is nominally purely inorganic and should not contain any biological or other ice nucleating components. Thereafter, a certain volume of the investigated sample, as specified in Table 3, was added and the aerosol generation process was started. The cell concentrations of algae in the [experiment-AEGOR tank](#) (see Table 3) ranging from 1 to 10⁶ cells mL^{-1} are representative for a strong phytoplankton bloom (Henderson et al., 2008; Borkman and Smayda, 2009; Saravanan and Godhe, 2010; Suikkanen et al., 2011; Canesi and Rynearson, 2016). In AEGOR sea spray aerosols are generated by a plunging jet that entrains air into the sea spray tank and thus leads to bubble bursting, emitting aerosol particles to the head space (flow

rate of the jet 5 L min^{-1} , nozzle diameter 4 mm). Bubble formation using this technique mimics bubble formation through
205 wave breaking. Bubbles rising through the water column scavenge surface active organic material and transport it to the surface
where it forms a microlayer. Subsequently, bubble-bursting transfers this surface active organic material to the aerosol phase.
Since the efficiency of particle generation by the sea spray simulation chamber was much lower than the nebuliser, injection of
particles generated using this approach into the AIDA chamber was conducted over a period of 14-16 h, resulting in an aerosol
particle concentration of approx. $300\text{-}400 \text{ cm}^{-3}$. Because of this time-consuming procedure, only a subset of the bulk solutions
210 was used for aerosol generation with AEGOR (Table 2). The temperature of the AEGOR tank was set to 293.15 K for the SM
culture samples, 277.15 K for the MA culture sample and 275.15 K for the SML samples.

Aerosolising an SML sample with a nebuliser is very different from aerosolisation due to bubble-bursting for a number of
reasons. Firstly, only a small volume of sample is required for nebulisation so pure SML samples could be aerosolised (we
had limited sample volume) while the sea spray simulation chamber requires a higher volume of sample as they were added to
215 20 L of artificial seawater (we used up to 900 mL sample volume). As such, the SML samples underwent significant dilution
when added to artificial seawater in the sea spray simulation chamber. Secondly, the process of aerosol generation by bubble
bursting is quite different to aerosol generation in a nebuliser. As such, those aerosols generated in the sea spray simulation
chamber are likely more representative of aerosols generated by oceanic bubble-bursting (Collins et al., 2014; King et al.,
2012; Prather et al., 2013). Given these differences, ~~once we have accounted for the relevant dilution factor in the sea spray~~
220 ~~simulation chamber (see Table 3)~~, comparison of the ice activity of aerosol generated by these two techniques should enable
us to determine whether INP material is preferentially aerosolised by bubble-bursting.

We expect that aerosolisation of the samples with the nebulizer results in an upper estimate of INP because the undiluted
SML (or cultured) samples are aerosolised whereas AEGOR is aerosolising a dilution of the samples with artificial seawater,
which could result in a lower estimate of INP. However, it is not only the dilution factor in the sea spray simulation chamber (see
225 Table 3), which has to be accounted for. The aerosolisation process itself is different in AEGOR compared to the nebuliser.
In the nebuliser the suspension is well mixed, while in AEGOR the aerosol particles are formed from an organic enriched
surface microlayer at the top of the tank. That leads to different expectations depending on the sample type. For the SML
samples we would not expect such a huge difference due to this aspect. Here, we aerosolise in one case the pure well mixed
SML (nebuliser), while in the other case we aerosolise the SML that has formed in AEGOR, which should be similar to the
230 original SML sample. For the cultured samples, however, we would expect a larger influence. In AEGOR the phytoplankton
material is floating at the surface of the tank leading to organic enriched aerosol particles during the aerosolisation, while
the nebuliser might produce less enriched aerosol particles due to the mixing of the sample. Note that this might depend on
the algae culture as well. Another crucial aspect of the two different aerosolisation methods is the size distribution and the
resulting chemical composition of the generated aerosol. It was demonstrated in the laboratory and as well measured in the
235 field, that for sea spray aerosol the organic composition of the aerosol particles and the generated size distribution are related
(O'Dowd, C. D. et al., 2004; Prather et al., 2013). One interesting aspect of our study is to see the influence of all the aspects
mentioned above and to check if the diluted samples aerosolised with AEGOR show a similar or a lower freezing signal
compared to the aerosolised pure samples.

2.3 Aerosol size and number measurements

240 The aerosol particle number concentration was measured using a condensation particle counter (CPC3010, TSI). The aerosol
particle number size distributions were measured with a scanning mobility particles sizer (SMPS, TSI; mobility diameter
0.014 - 0.820 μm) and an aerodynamic particle spectrometer (APS, TSI; aerodynamic diameter 0.523 - 19.81 μm). In the
AIDA chamber, typically held at 250 K and a relative humidity of 78% during aerosol injection (see Sect. 2.4 [for more details
and an explanation on the low temperature](#)), the aerosol particles were suspended as supercooled aqueous solution droplets. It
245 is important to consider, however, that the size distribution measurements were done at room temperature (298 K) by sampling
air from the cold interior of the aerosol chamber (Fig. 1). The water vapour content at 250 K corresponds to a relative humidity
of only 2.4% after warming to 298 K (Murphy and Koop, 2005). We thus assume that the measured size distributions represent
the effloresced, dry particle sizes of the algal culture and SML particles (Koop et al., 2000). A dynamic shape factor of 1.08
and a particle density of 2.017 g cm^{-3} (Zieger et al., 2017) for sea salt were used to convert the mobility and aerodynamic
250 diameters of the SMPS and APS measurements into the volume-equivalent spherical diameters. Fig. 2 shows the combined
size spectra of the SMPS and APS measurements, plotted as surface area size distributions, for two exemplary aerosol particle
populations produced by the nebuliser and AEGOR (SM100 and SML8).

The comparison of both aerosolisation techniques for the algae and the field samples shows that the nebuliser produces rather
uniformly sized particles with a median diameter of about 0.8 μm in the surface area size distributions. In contrast, the bubble
255 bursting process simulated in AEGOR leads to a much broader surface area size distribution with a smaller median diameter.

[However, both the nebulizer and AEGOR are not producing very narrow size distributions \(see Fig. 2\).](#) The majority of our
aerosolised samples yielded surface area size distributions very similar to those shown in Fig. 2. For each sample a log-normal
fit was created based on least-squares. The fits are expressed as a function of the median equal-volume sphere diameter, the
geometric standard deviation σ and the aerosol surface area concentration. The median diameter of the particles generated
260 with the nebuliser was typically in the range from 0.71 to 0.90 μm with a distribution width σ between 1.21 and 1.47. Smaller
particles with median diameters of 0.59, 0.41, and 0.18 μm were obtained for the SML5, MA100, and ASCOS (high mol.
weight, 5 kDa - 0.22 μm) samples, respectively, which is probably related to lower salt concentrations in the respective solu-
tions. Aerosol generation with AEGOR yielded median diameters between 0.4 and 0.7 μm and distribution widths σ between
2.2 and 2.9.

265 2.4 Ice nucleation measurement techniques

The combination of instrumental methods used in this study facilitates measurement of the ice nucleating ability of marine
organic aerosols over a wide temperature range. The ice nucleation activity was measured using three different ice nucleation
instruments: AIDA, INKA, and the μl -NIPI, which all have their highest sensitivities in different temperature ranges. While
the μl -NIPI is sensitive in the temperature regime above 248 K, AIDA and INKA are only sensitive in the temperature regime
270 below 248 K for the type of samples analysed in this study. All three measurement techniques are explained in detail in the
following sections.




Figure 2. Measured size distributions and fits to the data for two different samples: an algae sample (SM100) and a field sample (SML8). The samples were aerosolised using a nebuliser (solid line) or the sea spray simulation chamber AEGOR (dashed line). The aerosol size measurements are done with an APS (circles) and a SMPS (triangles). D denotes the equal-volume sphere diameter of the aerosol particles, S the surface area concentration.

AIDA

The AIDA facility comprises two aerosol chambers (Fig. 1) (Möhler et al., 2008). The term AIDA chamber refers to the 84.3 m³ sized aluminium vessel that is enclosed in an isolating containment and can be operated at any temperature between ambient and 183 K. The A smaller 3.7 m³-sized stainless steel vessel is located in the vicinity of AIDA. It is referred to as the APC (aerosol preparation and characterisation) chamber and can only be operated at ambient temperature. As indicated in Sect. 2.2, the aerosol particles were directly injected into the AIDA chamber to probe their ice nucleation activity by expansion cooling experiments. For practical reasons, the same aerosol particles were additionally injected into the APC chamber, acting as a reservoir for long-term measurements of the particles' ice nucleation behaviour with the INKA instrument (see next section) and for the CCN measurements (see Christiansen et al. 2020, submitted to J. Geophys. Res.).

The operation of the AIDA chamber as a cloud simulation chamber for studying ice nucleation has been thoroughly described previously (Möhler et al., 2003; Möhler et al., 2005; Wagner and Möhler, 2013). Briefly, a mechanical pump is used for a controlled reduction of the chamber pressure starting from ambient to about 800 hPa. Expansion cooling generates supersaturations with respect to ice and/or supercooled liquid water, triggering the formation of ice crystals and supercooled water droplets by various nucleation mechanisms (Vali, 1985; Vali et al., 2015). In the present study, the ice nucleation activity of the algal cultures and SML samples was investigated in the immersion freezing mode at mixed-phase cloud temperatures. For

aerosol injection, the AIDA chamber was typically held at a temperature of 250 K and a relative humidity with respect to supercooled water (RH_w) of about 78%, as controlled by an ice layer on the inner walls of the aluminium vessel. RH_w was measured in situ by tuneable diode laser (TDL) absorption spectroscopy with an uncertainty of $\pm 5\%$ (Fahey et al., 2014). With increasing RH_w during expansion cooling, the injected aqueous solution droplets continuously took up water vapour from the gas phase, and were finally activated to $\geq 10 \mu\text{m}$ -sized cloud droplets when RH_w exceeded 100%. The number concentration and size of the cloud droplets were measured with two optical particle counters (OPCs) Welas 1 and 2 (Palas GmbH) with an overall detection range of 0.7 - 240 μm . Cloud formation was typically observed after 3 K of expansion cooling, i.e., at a temperature of about 247 K. Whereas pure supercooled water droplets would only freeze homogeneously when the gas temperature further dropped to about 238 K during expansion cooling (Benz et al., 2005), the activated algal culture and SML aerosol particles exhibited heterogeneous ice nucleation modes due to immersion freezing at temperatures above 238 K. The number concentration of the nucleated ice crystals, N_{ice} , was separately deduced from the OPC records by using an optical threshold size to subtract the scattering signals of the smaller-sized supercooled cloud droplets. By dividing N_{ice} through the seed aerosol particle number concentration, the ice active fraction, FF , of the aerosol particle population was calculated. By further dividing FF through the average dry surface area of a particle, A_{aer} (determined from the size distribution measurements shown in Fig. 2) the ice nucleation active surface site density, n_s , of the polydisperse particle population could be computed, e.g. Hoose and Möhler (2012):

$$n_s(T) = \frac{FF(T)}{A_{\text{aer}}} \quad (1)$$

This equation is an approximation, which is valid for small values of $FF(T)$ (Hoose and Möhler, 2012) and was tested to be applicable for the dataset presented here. It is also assumed that n_s is independent of size.

The uncertainty of the deduced ice nucleation active surface site densities (n_s) was estimated to $\pm 40\%$ (Ullrich et al., 2017). In the following we estimate a lower detection limit of n_s in the AIDA experiments. The minimum detectable ice particle number concentration, as limited by the size of the detection volume of the OPC sensors, is about 0.05 cm^{-3} , equalling to one detected ice crystals in a sampling period of about 10 s. Together with the typical seed aerosol particle number concentration of about 500 cm^{-3} (Sect. 2.2), the lower detection limit for FF can thus be estimated to about 10^{-4} . The average dry surface area of the aerosol particles generated with the nebuliser was around $1 \mu\text{m}^2$, yielding a lower detection limit for n_s of about 10^8 m^{-2} (Eq. 1). In comparison with recent literature n_s values for laboratory and field sea spray aerosol particles (DeMott et al., 2016), n_s only exceeded such values at temperatures below about 248 K. This illustrates why the starting temperature of the expansion cooling runs was chosen as low as 250 K, thus limiting the ice nucleation data to temperatures below about 247 K. During our study we also probed a number of samples (STN2, STN3 and SM100) at a higher starting temperature of 258 K. However, we did not observe any ice formation above the detection limit down to a temperature of 248 K. For this reason, the AIDA data cover the above-defined low temperature regime of the ice nucleation spectra.

In addition to the expansion cooling cycles with the algal and SML samples, we conducted three control runs with the synthetic Sigma-Aldrich sea salt mixture, both using AEGOR and the nebuliser for aerosol generation. Here, the deduced n_s were close to the estimated detection limit of $1 \cdot 10^8 \text{ m}^{-2}$ at temperatures between 247 and 238 K. The small amount of

heterogeneously formed ice crystals could be due to traces of insoluble components in the synthetic salt mixture or due to ice nucleation on background aerosol particles in the cloud chamber. All aerosols exhibited n_s values 2–50 times larger than this background signal (see Sect. 3.2). To account for possible contamination originating in the nebuliser or AEGOR a background subtraction was conducted using these reference experiments with a pure Sigma-Aldrich sea salt solution and subsequent
325 estimation of the average background n_s value. The estimated background from these reference experiments was consistent and independent of temperature. It is higher for AEGOR compared to the nebuliser, probably due to the more complex setup of aerosolisation in the former.

INKA

Most of the samples that were probed in the AIDA chamber were also tested on their ice nucleation activity using the INKA
330 cylindrical continuous flow diffusion chamber (Schiebel, 2017). As explained above, the APC chamber was used as an aerosol particle reservoir for the INKA measurements. The APC chamber was held at 298 K and RH < 5%, meaning that the injected solution droplets generated with the nebuliser or AEGOR readily effloresced to form crystalline particles. Upon injection into the INKA instrument, aerosols are exposed to well controlled temperature and relative humidity conditions by flowing through a chamber with iced walls held at different temperatures. The sample air flow is sheathed by ~~dry~~-particle free synthetic air
335 (initially dry) in order to position the aerosol lamina between the walls and to allow for the calculation of the thermodynamic conditions within the lamina (Rogers, 1988). The residence time of the aerosol is 10 to 15 s, depending on the actual settings. Any droplets that might have formed in this section will shrink in a subsequent chamber section with no temperature difference between the iced walls. The formed ice particles will persist in this so-called evaporation section. The thus increased size difference between droplets and ice particles at the chamber outlet allows for an easy ice particle detection with an optical
340 particle counter (Climet CI-3100). INKA scans the ice nucleation activity by continuously increasing the sample's relative humidity at constant temperature settings. Due to a larger detection volume of the Climet OPC compared to the Welas sensors used in the AIDA experiments, the lower detection limit for n_s with INKA is about 10^7 m^{-2} . In inter-comparison studies using natural soil dust aerosol (DeMott et al., 2018a) or commercially available cellulose particles (Hiranuma et al., 2019) INKA has shown a good agreement with AIDA and other ice nucleation instruments. In the present study, most experiments have been
345 conducted above 241.15 K to enable a clear differentiation from homogeneous freezing events and to allow direct comparison with AIDA results.

μl -NIPI

The μl -NIPI is a cold stage instrument, used with a substrate to probe the ice nucleation in immersion mode of μl volume droplets (Whale et al., 2015). To do so, the droplets of the sample under investigation (if not explicitly otherwise mentioned
350 this is a bulk sample) are pipetted onto a silanised glass slide, which serves as a hydrophobic substrate. It is a “bulk” technique analysing the suspension directly under the assumption that the sample is well mixed, so that particles are distributed ~~uniformly~~randomly, and each droplet is representative of the sample as a whole, meaning each one has an approximately equal probability of containing an INP active at a given temperature. The droplets are then cooled at a rate of 1 K min^{-1} until the

droplets are all frozen. The temperature values of the individual freezing events are optically detected using a camera and
355 offline analysis. The number of droplet freezing events detected throughout the temperature ramp are then converted into a
fraction frozen at each temperature. This fraction frozen, or FF curve, represents the raw freezing events. In order to calculate
a concentration of INP per liquid unit volume of sample, $K(T)$, the FF must be thought of as the probability of freezing, and
so the equation below can be used to deduce the cumulative nucleus concentration per unit volume of sample used (Vali, 1971):

$$FF(T) = \frac{N_{\text{frozen droplets}}(T)}{N_{\text{droplets}}} \quad (2)$$

360
$$K(T) = \frac{-\ln(1 - FF(T))}{V_{\text{droplet}}} \cdot D, \quad (3)$$

where V_{droplet} is the volume of a droplet, N_{droplets} is the total number of droplets on the cold stage at the beginning of
the freezing experiment, $N_{\text{frozen droplets}}$ is the amount of droplets frozen at a certain temperature and D is the dilution factor
relative to the undiluted sample, relevant for the samples coming from AEGOR and a couple of dilution experiments conducted
with the algal cultures (in all other cases D is 1).

365 $K(T)$ can then be weighted to physical aspects of the sample such as the surface area of the particles or the mass of salt in
the sample in order to directly compare to other instruments using the same sample.

In contrast to AIDA and INKA the $\mu\text{l-NIPI}$ is sensitive to INP in a relatively high temperature range. Given the relatively large
size of the pipetted droplets, this technique is better suited to the investigation of freezing by rare INPs i.e. there is a greater
probability of having an INP within the droplet which subsequently freezes the whole droplet.

370 **3 Results**

In this section, we first address the ice nucleation measurements with the $\mu\text{l-NIPI}$ instrument in the temperature regime above
248 K (Sect. 3.1). The AIDA and INKA results for temperatures below 248 K are presented in Sect. 3.2. Finally, Sect. 3.3
outlines an approach to combine the AIDA/INKA and $\mu\text{l-NIPI}$ data into a single dataset to examine the ice nucleation behaviour
of the algal cultures and Arctic SML samples over the full temperature range relevant for freezing in the mixed-phase cloud
375 regime.

3.1 Temperature regime above 248 K (Bulk samples)

The frozen-fraction curves measured with $\mu\text{l-NIPI}$ for the field and algal samples are shown in Fig. 3. Among the field samples
there is a large spread in ice nucleation activity with a median freezing temperature T_{50} ($FF=0.5$, i.e. half of the droplets are
frozen) of approx. 262 to 245 K, i.e. a spread of 17 K. While the ice nucleation is very variable throughout the samples, the
380 dependence on temperature (slope of the curves) is mostly similar. A number of the samples exhibited ice nucleation activity
at relatively high temperatures (>263.15 K), with the ASCOS high molecular weight sample (ASCOS high mol. w., 5 kDa to
0.22 μm) and SML5 being the most ice active. Both algal samples studied were also ice active, although they were clearly less

ice active than the field samples despite their relatively high cell concentration (compared to natural seawater). For example, the T_{50} of the culture samples is approx. 252 to 246 K (range of 6 K), so within the colder part of the variability of the field samples (see Fig. 3). Further to this, no large differences (a difference of T_{50} of approx. 5 K) were observed between the different diatom species or when comparing the different nutrient conditions for SM. However, it should be noted that there was ~~an intra-specific~~ variability within the individual cultures and that storage changed the ice nucleation activity. For example, the same SM100 culture ~~that~~ was delivered to AIDA in two separate bags ~~showed different activity between the two bags~~. We refer to one bag as SM100a, the other one SM100b. A third sample (SM100c, a sub-sample of SM100b) was analysed two months after the campaign after having been stored at or below 253 K. SM100d, also a sub-sample of SM100b, was used for some further tests 10 months after the campaign (as well stored at or below 253 K). Note that the results of SM100d should be used with caution and not directly compared to the other ones, since this sample was unfrozen several times and stored for a quite a long period of time, which might not be ideal.

Comparing SM100a, SM100b and SM100c, it can be seen that the freezing properties of the SM100 sample is variable, ~~as both samples have different gradients~~. SM100a and SM100b look similar, with most of the spectrum at low temperature in the background. They show differences at higher temperatures, where SM100a displayed activity but SM100b did not. SM100c showed different activity from SM100a and SM100b with the freezing shifted to higher temperatures that leads to the whole curve being outside of the background (compared to SM100b). The gradients between all three curves is also different, with SM100a having the shallowest slope. Additional to the variability of the sample itself (different bags - SM100a and SM100b), it seems that the sample changed with time, so age, storage and multiple freezing cycles may all have had effects on the sample.

The STN samples have been analysed previously using a similar droplet freezing technique albeit using a 10 times faster cooling rate (10 K min^{-1}) (Irish et al., 2019b). Comparison of these measurements with our measurements of the same samples highlight the differences. We observed up to an order of magnitude higher $K(T)$ values [and up to a 10 K difference for the same $K(T)$] than those reported in (Irish et al., 2019b), which might have been influenced by the difference in the cooling rate. The temperature at which 50% of the droplets are frozen has been shown to decrease with increased cooling rate in Wright and Petters (2013); Herbert et al. (2014), also this dependence was shown to be rather small. Nevertheless, a shift of 10 K for a factor of 10 change in cooling rate is unlikely. The SML samples from Wilson et al. (2015) were analysed using the same droplet freezing technique as in this study. Samples SML5, SML8 and SML16 exhibited ice activity at similar temperatures to those presented in Wilson et al. (2015), while samples SML17 and SML19 exhibited lower ice activity, with lower temperatures of freezing for the same fraction frozen. Therefore, we conclude that some samples were unaffected by long-term storage (being frozen at 193.15 K), while the activities of other samples changed. This indicates that some ice active components are altered through the freezing, storage and thawing process. Note that this contradicts earlier assumptions based on findings of Schnell and Vali (1976); Irish et al. (2019b). ~~It indicates that microlayer samples~~ However, Polen et al. (2016) has shown that the biological INP Snomax is sensitive to storage. An alteration of INP characteristics of our microlayer samples indicates that they contain different ice active components which have different properties and may be related to different biological processes. In this paper we use the re-measured droplet freezing results to compare the ice nucleation activity between instruments.

The influence of bubbling the samples in the sea spray chamber AEGOR on the ice nucleation activity was investigated by comparing pure samples with three different sub-samples taken out of AEGOR after bubbling: one bulk sub-sample (collected from the bottom of AEGOR), one scoop sub-sample (collected by scooping a falcon tube along the surface liquid) and a microlayer sub-sample [collected by the glass-plate technique as per the methods of (Harvey, 1966)]. Note that all these samples are bulk samples. Upon introduction to AEGOR there was a significant dilution of the sample with artificial seawater (Table 3). The ice nucleation activity of the SML5 sub-samples as described above is shown in Fig. 4. ~~In the FF-curve (left hand side of Fig. 4) there is a clear reduction in the ice nucleation activity of all three of the sub-samples compared to the pure SML5 sample. The AEGOR samples freeze at lower temperatures. This is consistent with the sample being diluted when introduced to the sea spray simulation chamber. When the same data are~~ To take the dilution into account the data is plotted with respect to the volume of sample used, as INP/L, ~~the datapoints for the undiluted pure sample and diluted sub-samples align (right hand side of Fig. 4), as the dilution has been taken into account~~ (see Eq. 3). Interestingly, the bulk and microlayer sub-samples exhibit lower ice activity than the scoop sub-sample. However, it is important to note that most points from the bulk and microlayer samples are in the baseline of the μ l-NIPI experiment, and can therefore be seen as upper limits. It is notable however, that the 'microlayer' sample obtained with a glass plate had a lower activity than scooping the surface water, which might suggest that the ice active components may only have an intermediate affinity for the glass plate. Nevertheless, the fact that the upper layers of water in the AEGOR are enhanced in INP suggests that organic INP material scavenged by bubbles resides at the water surface and is likely surface-active (i.e. material which preferentially resides at an interface). As such, this material may be scavenged by the bubbling in the chamber and be preferentially aerosolised during the bubble bursting process.

3.2 Temperature regime below 248 K (Aerosolised samples)

The ice nucleation results of the AIDA and INKA measurements, expressed as ice nucleation active site densities versus temperature $n_s(T)$, are shown in Fig. 5 (SML samples) and Fig. 6 (algal cultures). With respect to the experiments where AEGOR was used for aerosol generation, some samples did not exhibit a detectable freezing signal above the background (SM100, SM10, and SML8) and are therefore not included. As a comparison to our data, Fig. 5 includes a recently published dataset consisting of field measurements of sea spray aerosols and laboratory data of particles released during an algae bloom generated in a marine aerosol reference tank (DeMott et al., 2016). Furthermore, we show a parameterisation of the temperature-dependent n_s values for desert dust particles (Niemand et al., 2012).

~~In contrast to the large variability of the ice nucleation activity evident in the μ l-NIPI measurements at higher temperatures (Fig. 3), the~~ The various SML samples show ~~much less little~~ variation at temperatures below 248 K when probed in the AIDA chamber, meaning that the SML samples all exhibited similar ice nucleation activity (n_s of 10^9 m^{-2} at temperatures between 240 - 244 K) and the individual $n_s(T)$ -curves of the AIDA measurements form a rather compact block of data (Fig. 5). One notable exception is the ASCOS high-molecular weight sample (ASCOS high mol. w., 5 kDa to $0.22 \mu\text{m}$). Whereas the foam and < 5 kDa ASCOS samples fall into the range of n_s values observed for the other SML and STN microlayer samples, n_s for the high-molecular weight sample is about one order of magnitude higher. This agrees with the μ l-NIPI observations, where




Figure 3. Fraction frozen curve, a measure of the fraction of droplets frozen at discrete temperatures, for:

a) 9 different SML field samples coming from three different Arctic field expeditions (ACCACIA, NETCARE, ASCOS) measured with the μ I-NIPI (droplet freezing technique, bulk, undiluted samples). The field sample from ASCOS was treated in three different ways (see Sect. 2.1).

b) Two cultured diatom species measured with the μ I-NIPI (droplet freezing technique, bulk): *Skeletonema marinoi* (SM) and *Melosira arctica* (MA). The SM sample was investigated for two different nutrient regimes (see Sect. 2.1). Two duplicate samples of SM100 (SM100a and ~~SM100c~~SM100b) are reflecting the variability of the sample. One sample (SM100a and SM100b are from two bags collected from the same culture. SM100c is a sub-sample of SM100b after 2 months storage. SM100d, a sub-sample of SM100b, long was in storage) for 10 months, and was then nebulised and then re-tested to see-determine the effect of the aerosolisation on the sample.

The points with reduced opacity represent upper limits for those data points, as they could have been affected by background signal.

Note that the temperature in both plots was not corrected for freezing point depression caused by salts because the water activity was not available for all samples.

455 this particular sample also proved to be one of the most ice active. The ASCOS high-molecular weight sample consists of the high molecular weight dissolved organic matter of the collected SML sample. More specifically, it was shown in Orellana et al. (2011) and Gao et al. (2012) that this sample mostly contained ~~of~~ marine colloidal gels. This might lead to an enrichment of ice active organic material and explains the high ice nucleation activity of this sample. Note that this sample is highly concentrated. The size range of the filtration of the sample indicates that macromolecules are responsible for the freezing of the sample. Most bacteria, cell debris, etc. are likely to be removed by the ultrafiltration. The size distribution of the nebulised ASCOS high-molecular weight sample resulted in particles with the smallest diameters compared with other samples, which might have an influence on the ice nucleation activity as well since the chemical composition of sea spray aerosol is highly size




Figure 4. ~~Frozen fraction curve (left) and cumulative~~ Cumulative INP concentration per unit volume field sample SML5 ~~(right)~~ for the pure sample in comparison to different dilutions (sub-samples from AEGOR: bulk, microlayer, scoop; see text for details). ~~The points with reduced opacity (frozen fraction curve) represent upper limits for those data points, as they could have been affected by background signal.~~ Where the lower error bar is unchanged from the previous point (cumulative INP concentration), there may have been no additional INP detected above the background signal. Note that the temperature in this plot was not corrected for freezing point depression caused by salts because the water activity was not available for all samples.

460 dependent. This sample might consist of smaller particles with a larger organic mass fraction compared to the other samples. Other field samples that proved to be particularly ice active in the high temperature regime like SML5, however, do not show superior ice nucleation activity at temperatures below 248 K. This is an indication that different types of ice active materials might cause the freezing in the different temperature ranges, an issue that will be further discussed in Sect. 3.3 when combining the AIDA and μ I-NIPI data sets.

465 In order to facilitate the comparison of the AIDA measurements with previous studies of ambient marine aerosols, we chose to represent the DeMott et al. (2016) data in Fig. 5 by a grey shaded area that encompasses the observed range of nucleation site density values n_s . A similar representation was used by McCluskey et al. (2017), who have determined n_s for nascent sea spray aerosol particles during phytoplankton blooms in the laboratory. These data are not separately depicted because they fall into the regime of the DeMott et al. (2016) dataset. A particular subset of the DeMott et al. (2016) data is highlighted in
 470 Fig. 5 by the grey stars. These data points refer to a laboratory experiment in the Marine Aerosol Reference Tank (MART) following the peak of the phytoplankton bloom. The n_s values derived from the AIDA measurements for the field samples fall into the range of former observations, albeit towards the upper, more ice active regime of the data by DeMott et al. (2016). The

MART data for the artificially enhanced phytoplankton bloom is in good agreement with the upper thresholds of n_s for our field samples. Given that most of the AIDA measurements were made by aerosolising the undiluted SML solutions with the nebuliser, it can be expected that this dataset indeed represents an upper limit of the ice nucleation activity of natural sea spray aerosol particles.

The experiments where AEGOR was used for aerosol generation shed some light on how much of the ice active material in the SML bulk solutions may be released during the process of air entrainment, bubble scavenging and bubble bursting. For both sample types investigated, the algal cultures and natural SML samples, we find examples where the ice nucleation activity observed of particles generated using the AEGOR tank remains similar to the ice activity of aerosols generated by nebulising the pure sample despite the strong dilution of the samples with artificial seawater in the AEGOR tank (SML5, Fig. 5; MA100, Fig. 6). This suggests that in some cases, the organic INP material is indeed preferentially scavenged by the bubbling in the seawater tank and aerosolised during the bubble bursting process. For other samples, however, the ice nucleation activity was reduced to below the detection limit (n_s of 10^8 m^{-2}) after the dilution in AEGOR (SML8, SM10, and SM100). This variability in the AEGOR experiments might explain why the previous field measurements of sea spray aerosol particles show a huge spread in the n_s values, whereas the laboratory nebuliser data fall into a narrow range at the upper end of the ice nucleation activity scale. Note that this upper limit of the ice nucleation activity of the field samples, however, is still one order of magnitude lower than the n_s parameterisation for mineral dust [Fig. 5, Niemand et al. (2012)], underlining the relatively poor heterogeneous ice nucleation activity of sea spray aerosol particles compared to other atmospherically relevant types of INPs in the temperature range ~~below~~ above 248 K. In the (High) Arctic both transported dust and sea spray aerosol (transported or locally originated) can be present (see Willis et al. (2018) for a thorough review of literature). However, which source is dominant for ice nucleation might be locally very different. In regions dominated by sea spray aerosol the fraction of organic matter within the aerosol population is another uncertainty.

At low temperatures, the algal cultures had similar ice nucleation activities compared to the field samples, with *Melosira arctica* being slightly more ice active than *Skeletonema marinoi*. For *Skeletonema marinoi* grown under replete and deplete nutrient conditions, the culture with the highest nutrient limitation and inhibited growth (SM10) had somewhat lower n_s values compared to SM100 and SM60, but this trend is only distinct in the AIDA data and not as clearly visible in the INKA measurement. For comparison, we added previously published $n_s(T)$ values for two other algae, the diatom *Thalassiosira pseudonana* (Knopf et al., 2011) and the green algae *Nannochloris atomus* (Alpert et al., 2011a) [the data points were taken from Murray et al. (2012)]. The ice nucleation activities of these two species are in reasonable agreement with the data presented here. They lie towards the lower end of the AIDA data and fully overlap with the range of the n_s from the INKA measurements.

With respect to the comparison between the AIDA and INKA measurements, the INKA results tend to be shifted to lower n_s values, although the INKA data partly overlaps with the AIDA data within the respective error bars. As previous INP measurements for insoluble aerosol particles such as soil dust have shown good agreement between AIDA and INKA (DeMott et al., 2018a), the deviation for the current study with soluble, marine aerosol particles might be related to the particles' phase state. For soluble aerosols, the different time scales and particles' phase state evolution in the AIDA and INKA measurements might

affect the observed INP data. In AIDA, the aerosol particles are initially suspended as aqueous solution droplets, gradually take up water when the expansion cooling run is started, are activated to μm -sized cloud droplets when the relative humidity exceeds 100%, and potentially nucleate ice by immersion freezing upon further reduction of the temperature during expansion cooling. These processes occur on an overall time scale of approx. 5 min. For the INKA measurements, the aerosol particles are suspended as effloresced crystals in the APC chamber. During a very short time period of only 10 to 15 s in the first section of the CFDC chamber, the particles have to undergo the complex trajectory of deliquescence, droplet activation, and freezing. The short residence time in INKA might prevent equilibration of the aerosol to the instrument conditions. Thus, it is possible that at certain locations there is not enough water vapour present to fully activate the aerosol particles to cloud droplets and that this effect may account for the slightly lower n_s values compared to the AIDA measurements. Note that efflorescence might as well change the INP activity of the aerosol particles.

3.3 Combined ~~temperature regime – full ice nucleation spectra~~ bulk and aerosol phase measurements

One of the central aims of this study was to analyse the ice nucleation behaviour of Arctic SML samples and two different algal cultures over the full temperature range relevant for freezing in mixed phase clouds. We also wanted to assess if the ice nucleation material is transferred from the bulk to the aerosol phase. The samples were measured with different instruments sensitive to different temperature regimes: AIDA and INKA below 248 K (aerosol phase) and μl -NIPI above 248 K (bulk). Here we attempt to directly compare the AIDA and μl -NIPI datasets ~~and combine them into a single dataset~~. The INKA dataset is not included in the comparison since the AIDA dataset is more comprehensive and has a finer temperature resolution than the INKA data.

To enable comparison and answer the question if the ice nucleating material is transferred from the bulk to the aerosol phase, both datasets (AIDA and μl -NIPI) require normalisation so that the ice nucleation behaviour can be expressed with the same quantity as a function of temperature. We have chosen to normalise both sets of data to the mass of salt present in the solution droplets since this quantity can be estimated for both approaches. Thus, the ice nucleation behaviour is expressed as ice nucleation active site density per mass of salt (n_m ; $[n_m] = g^{-1}$). It is more obvious how to treat and harmonise ice nucleation data using materials like mineral dust which have a relatively well-defined surface area. The surface area of an aerosol dispersion can be used to derive n_s in much the same way as dust particles in bulk suspension. However, when the ice nucleating material in a sample is soluble or forms colloidal suspensions then it is less clear how to treat it. This is especially complex for the marine system, where the bulk sample can be very different from what is aerosolised into the atmosphere - one question that we want to investigate a bit further by comparing the AIDA and the μl -NIPI datasets. While we can, and have, derived n_s values for the AIDA and INKA data where the surface area is the surface area of the dry aerosol, we cannot do this for the bulk suspension measurements from the μl -NIPI instrument. Similarly, while we have a measure of organic mass for the bulk microlayer samples we do not have a measurement of the organic mass in the aerosol phase, hence we cannot normalise to organic mass. Solution volume cannot be used, since the volume of the solution of the aerosol changes as its concentration alters to come to equilibrium with the chamber conditions. Hence, we have chosen to normalise to the mass of salt, a quantity which can be readily estimated from both the bulk and aerosol experiments. When contrasting the resulting n_m values




Figure 5. Surface active site density n_s as a measure for ice nucleation activity at different temperatures for 11 different SML samples from the AIDA (coloured full circles and triangles) and INKA (open squares) measurements. The field sample from ASCOS was treated in three different ways (see Sect. 2.1). Different symbols show the different aerosolisation techniques for the AIDA measurement (nebuliser in circles, AEGOR in triangles). The AIDA n_s data were corrected for the background ice nucleation mode observed in the reference experiments with purely inorganic Sigma-Aldrich sea salt solution droplets (see Sect. 2.4). The data of DeMott et al. (2016) is shown as a grey shaded area (fit and shifted fits to the upper and lower limit of the data) and grey stars (MART phytoplankton bloom), see text for details.

it should be borne in mind that the spread in activities is likely an indication of the range of concentrations of the ice active components as well as variability in the activity of those components. ~~Also, the~~ The objective of our work was to compare droplet freezing assay results with aerosolised measurements, rather than to derive a quantity which could be used to predict atmospheric INP. Ideally, we would quote active sites per unit mass of the nucleating component, but if the identity and mass of the nucleating component is unknown this is not possible (as in this case). However, this approach enables us to ~~study the ice-nucleating activity of two common phytoplankton species and Arctic microlayer samples over a wide range of mixed-phase~~

545




Figure 6. Surface active site density as a measure for ice nucleation activity at different temperatures for the two different diatom species (SM and MA) from the AIDA and INKA measurements. For SM, three samples grown under different nutrient regimes to generate cultures with different exudate properties (SM10, SM60, SM100) are shown. Literature n_s data for *Thalassiosira pseudonana* and *Nannochloris atomus* are shown as a comparison. The AIDA n_s data were corrected for the background ice nucleation mode observed in the reference experiments with purely inorganic Sigma-Aldrich sea salt solution droplets (see Sect. 2.4).

~~cloud conditions using several instruments and test the consistency of these~~ investigate if the bulk and the aerosolised samples behave similarly and if both ice nucleation techniques complement each other when normalised and brought into one context.

550 For the μ l-NIPI data we derive the salt concentration for each sample in g/L using the measured water activity of the samples and the parameterisation linking the water activity and salt concentration of seawater presented by Tang et al. (1997). To calculate the ice nucleation active site density per mass of salt, the measured INP/L is simply divided by the salt concentration in g/L. For the samples where no water activity was measured as part of this study (see Table 3), the values from Wilson et al. (2015) (for the ACCAIA SML samples) or an average of all SML samples (for the NETCARE STN samples) was used. We added an additional uncertainty of 20% (arbitrary) to the error bars for the n_m values of the samples where the water activity

555 was not directly measured. The ASCOS samples are not included in the unified dataset. Their water activity could not be directly measured because the remaining sample volume was too small. Furthermore, these samples were treated differently to the other microlayer samples so an average water activity might not be a good representation for these samples.

For the AIDA data the measured FF was normalised with the measured mass concentration of dry particles (as obtained from the SMPS and APS measurements, see discussion in Sect. 2.2), instead of using the particles' surface area concentration for normalisation that yielded the n_s data shown in Figs. 5 and 6. The underlying assumption is that the dominating constituents in terms of mass is salt with a density of $2.017 \pm 0.006 \text{ g cm}^{-3}$ [Sigma-Aldrich sea salt; Zieger et al. (2017)]. Considering the composition of marine aerosols as presented in Gantt and Meskhidze (2013) this assumption is fair for the typical sizes of aerosol particles aerosolised into AIDA.

~~The combined temperature spectra for the~~ 565 INP transfer from the bulk to the aerosol phase

The combined ice nucleation activity of the field samples is shown in Fig. 7. ~~As a striking result, there is much more variability in the ice nucleation activity of the samples when analysed with the $\mu\text{l-NIPI}$ than with AIDA (approx. 15 K vs. 5 K). This larger variability in the high temperature range has been observed in other studies, too, e.g. for soil or agricultural dust (O'Sullivan et al., 2014; Schiebel, 2017; Suski et al., 2018). One explanation for this behaviour could be that there are multiple INP types in seawater, just like there are in terrestrial samples, leading to a high diversity of the INP spectra at high temperatures. At low temperature the ice nucleation activity is much less variable and low throughout all samples.~~

570 ~~Most samples feature a rather continuous slope in the temperature-dependent INP spectrum. One notable exception is the STN7 sample, which shows a pronounced, step-wise change in the ice nucleation behaviour at about 263 K.~~

~~There is a gap in the temperature (and n_m) range covered by the AIDA and the $\mu\text{l-NIPI}$ data sets. Therefore, we cannot fully assess the validity of our normalisation approach with respect to the mass of salt. However, for the SML samples, it is reasonable to assume that the composition of the aerosolised solution droplets probed in the AIDA chamber is very similar to that of the corresponding bulk solutions used in the $\mu\text{l-NIPI}$ measurements. As discussed below, this assumption is not necessarily valid for the algal cultures.~~

The combined temperature spectra for the ice nucleation activity of the algal samples is shown in Fig. 8 and Fig. 9; the 580 samples were split in two figures for clarity. ~~Figure 8 shows the $n_m(T)$ spectra for the SM100 culture and the variability including two SM100 samples (a and b for biological variability; c and d for storage effects) as discussed in section 3.1. The latter (Fig. 9) shows the spectra for MA100 and SM10. To bridge the gap in the ice nucleation spectra~~

We first turn to the comparison between the AIDA and the $\mu\text{l-NIPI}$ data, we did additional dilution experiments with $\mu\text{l-NIPI}$ to extend the temperature regime of the $\mu\text{l-NIPI}$ data to lower temperature. Diluting the SM100 and MA100 sample has the effect of reducing the freezing temperature and increasing n_m . Thus the curves from the undiluted samples can be extended to 585 lower temperatures. That works well for SM100 and partly also MA100. For MA100 there is a gap between the n_m curves, with the diluted sample having higher n_m values compared to the undiluted sample in the same temperature regime. The slope

of the n_m curve continues to be steep throughout the dilutions. However, there are some points which may have been affected by the background signal, which are denoted by the larger lower error bar value. It is not clear why there is such a difference in the behaviour after dilution between the SM100 and MA100 samples, and further investigation into the differences in their composition and how this related to their ice nucleating ability is necessary.

We now turn to the comparison measurements for the algal and field samples focusing on the difference between aerosolised and bulk samples. A significant difference between the AIDA and $\mu\text{l-NIPI}$ measurements for the algal and field samples measurement is that one is derived from an aerosolised sample and one is derived directly from the pipetted culture medium. Comparison between $\mu\text{l-NIPI}$, AIDA and other instruments in a recent intercomparison was very good (DeMott et al., 2018b). Inspection of the data in Figure 7 and Fig. 7, Fig. 8 and Fig. 9 suggests that the data from the two techniques might be consistent, but n_m would have to be extremely steep at the intermediate temperatures. The discontinuity of the AIDA and the NIPI data, i.e. the shift of the AIDA data to higher n_m values might be related to a change of physical characteristics upon aerosolisation. A significant difference between the AIDA and $\mu\text{l-NIPI}$ measurement is that one is derived from an aerosolised sample and one is derived directly from the pipetted culture medium. As mentioned above, aerosolisation aerosolisation. Aerosolisation may alter the physical characteristics of the ice nucleating material compared to when it is in the culture medium through breaking up aggregates or disrupting cells. This was shown for *Pseudomonas syringae* cells in the study of Alsved et al. (2018). Hence, it is feasible that the ice nucleation activities of the aerosolised samples in the AIDA experiments are higher than those in the $\mu\text{l-NIPI}$ experiments. However, there is a recognisable difference between both types of samples. The aerosolisation technique might exert more of an influence on the cultured samples compared to the microlayer samples, where the INP are thought to be associated with submicron organic detritus, rather than intact cells. For the SML samples, it is therefore reasonable to assume that the composition of the aerosolised solution droplets probed in the AIDA chamber is very similar to that of the corresponding bulk solutions used in the $\mu\text{l-NIPI}$ measurements. Indeed, the n_m spectrum looks more uniform as compared to the algal cultures. Most samples feature a rather continuous slope in the temperature-dependent INP spectrum. One notable exception is the STN7 sample, which shows a pronounced, step wise change in the ice nucleation behaviour at about 263 K.

For the algal cultures the assumption that the aerosolised and bulk samples are similar is not necessarily valid. In order to investigate if the process of nebulising influences the ice nucleating activity of cell suspensions, we nebulised a SM100 sample, collected the mist-nebulised sample as a bulk liquid and retested its ice nucleating activity using the $\mu\text{l-NIPI}$. Nebulisation increased the activity of the sample (see Fig. 3). We suggest that this might be consistent with the break up or rupture of cells in the vigorous nebulisation process, which might then release macromolecular ice nucleating materials. Alternatively, there might be agglomerated cells or colloidal particles inside the sample. That means that ice active sites can be either inaccessible or simply concentrated in a few particles. These aggregates might remain relatively intact during pipetting, but may be disrupted on nebulisation. It would have the effect of dispersing the ice nucleating entities throughout the aqueous suspension, thus increasing the probability of freezing across the droplet distribution when nebulising the sample. However, nebulising MQ water (not shown) showed that some impurities can likely be introduced by the nebuliser itself. These hypotheses deserve further investigation in the future. Given these factors, the aerosolisation technique might exert more of an influence on the cultured samples compared to the microlayer samples since the INP in the latter are thought to be associated with submicron

~~organic detritus, rather than intact cells.~~

Further to this, we have the hypothesis that the aerosolised material entering AIDA was very different compared to the pure
625 cultures. For example, first analysis of electron microscopic pictures of aerosol particles contained in AIDA (representative
for particles aerosolised with a nebuliser into AIDA) during the experiments with *Skeletonema marinoi* showed no cells or
obvious cell fragments visible (see left picture of Fig. 10). This is consistent with the microlayer being dominantly composed
of organic detritus and might be a result of biochemical processes within the microlayer. In contrast the right picture of Fig. 10,
where SM100 droplets were pipetted directly from the solution, shows clearly cells, which are then also present in the droplets
630 analysed with $\mu\text{l-NIPI}$. However, a more detailed analysis would be needed to give a final answer on the difference of the
aerosol particles in AIDA compared with aerosol particles within pipetted droplets.

Dilution tests bulk measurements

Figure 8 shows the $n_m(T)$ spectra for the SM100 culture and the variability including two SM100 samples (a and b for
biological variability; c and d for storage effects) as discussed in section 3.1. The latter (Fig. 9) shows the spectra for MA100
635 and SM10. To bridge the gap in the ice nucleation spectra between the AIDA and the $\mu\text{l-NIPI}$ data, we did additional dilution
experiments with $\mu\text{l-NIPI}$ to extend the temperature regime of the $\mu\text{l-NIPI}$ data to lower temperature. Diluting the SM100 and
MA100 sample has the effect of reducing the freezing temperature and increasing n_m . Thus the curves from the undiluted
samples can be extended to lower temperatures. That works well for SM100 and partly also MA100. For MA100 the slope of
the n_m curve continues to be steep throughout the dilutions. However, there are some points which may have been affected
640 by the background signal, which are denoted by the larger lower error bar value. It is not clear why there is such a difference
in the behaviour after dilution between the SM100 and MA100 samples, and further investigation into the differences in their
composition and how this is related to their ice nucleating ability is necessary.

Temperature dependent difference in ice nucleation behaviour

As a striking result, there is much more variability in the ice nucleation activity of the samples when analysed with the $\mu\text{l-NIPI}$
645 than with AIDA (approx. 15 K vs. 5 K). This larger variability in the high temperature range has been observed in other
studies, too, e.g. for soil or agricultural dust (O'Sullivan et al., 2014; Schiebel, 2017; Suski et al., 2018). One explanation for
this behaviour could be that there are multiple INP types in seawater, just like there are in terrestrial samples, leading to a high
diversity of the INP spectra at high temperatures. At low temperature the ice nucleation activity is much less variable and low
throughout all samples.

650 4 Conclusions

In this study the ice nucleation activity of several bulk and aerosolised SML samples from the Arctic region was investigated
and compared with pure and aerosolised samples of two diatom species-cultures (*Skeletonema marinoi* and *Melosira arctica*).
The measurements were conducted with a suite of ice nucleation instruments (AIDA, INKA, $\mu\text{l-NIPI}$) which are sensitive in




Figure 7. Normalised AIDA and μ I-NIPI measurements for 7 field samples showing a full ice nucleation spectrum represented as ice nucleation active site density per mass of sea salt n_m . For the AIDA measurements both aerosolisation techniques (nebuliser and AEGOR) are included. The points of the μ I-NIPI measurements which could have been affected by background signal and represent upper limits are indicated by a lower error bar that is unchanged from the previous point, as there may have been no additional INP detected above the background signal. The temperature in this plot was corrected for freezing [point](#) depression caused by salts for the μ I-NIPI measurements.

different temperature regimes across the whole mixed-phase cloud temperature range (below and above 248 K). In order to
 655 ~~make direct comparisons between~~ [compare](#) the different approaches [and the ability of the ice nucleating material to transfer to the aerosol phase](#) we have normalised all of the measurements by the salt mass present in the [bulk and aerosolised](#) samples. Normalisation in this manner results in an ice nucleation active site density per mass of salt n_m .

~~With regard to our~~ [Our](#) three main objectives ~~;-first were:~~ [first](#), the comparison of the ice nucleating ability of two common phytoplankton species with Arctic microlayer samples, second, [the impact of the aerosolisation technique on the results](#), and
 660 third, [the sample variability over the entire mixed-phase cloud temperature range](#); ~~-~~ [Concerning these objectives](#) we can draw the following conclusions:




Figure 8. Ice nucleation active site density per mass of sea salt n_m estimated from the AIDA and μ I-NIPI measurements for the SM100 culture samples. For the μ I-NIPI measurements, the SM100 samples were additionally diluted with ultrapure water. Note that the dilution was conducted up to 8 weeks after the main campaign (Leeds, UK). The points of the μ I-NIPI measurements which could have been affected by the background signal and represent upper limits are indicated by the lower error bar unchanged from the previous point, as there may have been no additional INP detected above the background signal. The temperature in this plot was corrected for freezing point depression caused by salts for the μ I-NIPI measurements.

When comparing the full temperature spectrum of the algal cultures with the field samples it is evident that the culture samples are similar to the field samples in the low temperature regime but are not within-among the most ice active samples of the spectrum in the high temperature regime. The two algae species-As the investigated algae species show less ice activity in the temperature regime above 248 K compared to the natural field samples, we conclude that they, especially *Melosira arctica*, cannot explain the freezing at the high temperatures, they only represent the less ice active share of the natural field samples. A normalisation of the samples to the atmospheric algal content would be needed to quantify this observation. This result indicates that the INPs active at the highest temperatures are not triggered-by-either-one of the two types of phytoplankton cells




Figure 9. Ice nucleation active site density per mass of sea salt n_m estimated from the AIDA and $\mu\text{l-NIPI}$ measurements for the SM10 and MA culture samples. For the AIDA measurements both aerosolisation techniques (nebuliser and AEGOR) are included. For the $\mu\text{l-NIPI}$ measurements, the MA100 sample was additionally diluted with ultrapure water. Note that the dilution was conducted up to 8 weeks after the main campaign (Leeds, UK). The points of the $\mu\text{l-NIPI}$ measurements which could have been affected by background signal and represent upper limits are indicated by the lower error bar unchanged from the previous point, as there may have been no additional INP detected above the background signal. The temperature in this plot was corrected for freezing [point](#) depression caused by salts for the $\mu\text{l-NIPI}$ measurements.

studied or their exudates. However, since we have only tested two mono-species grown axenically and harvested at exponential
670 growth phase, we cannot rule out ice nucleation being triggered by a consortia of microorganisms facilitating break-up of cells
and mass-release of organic matter from a phytoplankton bloom. The freshly produced pure algae cultures are different from
the diluted field samples, which are highly diverse in terms of composition. Aged algal cultures may exhibit a different freezing
behaviour.

For *Skeletonema marinoi*, the culture was grown at different nutrition conditions to test the dependence of the freezing on
675 the algal characteristics, such as total organic carbon (cell organic carbon and all dissolved organic [carbon](#)), cell wall structure,




Figure 10. Electron microscope pictures of SM10 (aerosolised by AEGOR) collected from AIDA (left) and SM100 in droplets pipetted directly from the solution (right).

colony length etc.. No significant difference could be found when comparing the ice nucleation behaviour of the samples grown at different rates and under varying nutrient limitation, so there is no clear evidence for a correlation between the total organic carbon content of the culture sample (see Table 3) and the freezing of the sample.

A key aspect of this study is that we have used both a sea spray simulation chamber and a nebuliser to introduce samples into AIDA (low temperature regime). Using a sea spray simulation chamber (AEGOR) allowed us to test the effect of mimicking the process of bubble bursting on the ice activity of the aerosol generated. A larger spread was observed in general for the SML samples diluted in AEGOR - some retained the activity of the undiluted sample, in some cases the IN ability decreased below the detection limit. Lower ice nucleation active site densities (for the cases where the IN ability decreased below the detection limit) can be explained by the difference in the size distribution of the aerosols generated by the two approaches.

Analysing the ice nucleation spectra over the whole temperature regime, the SML field samples exhibit a high variability in ice nucleation activity in the temperature regime above 248 K compared with lower temperatures. Above 248 K the variation in the median freezing temperature T_{50} is approx. 15 K with some samples showing a strong freezing signal at high temperatures ($T_{50} \approx 262$ K), while below 248 K the spread of T_{50} is only approx. 5 K. The behaviour of the samples in the different temperature regimes might be related to different types of INP active in the different regimes. In the temperature range below 248 K the results of this study are in the upper range of the values measured by DeMott et al. (2016), which show a larger spread compared to the results of the nebulised samples. This larger spread could be explained by the aerosolisation (see paragraph above). However, neither the SML nor the algal samples exhibit a strong freezing signal in the low temperature regime (below 248 K) compared to desert dust. There was no significant freezing above the detection limit in the AIDA chamber (around

695 $2 \times 10^8 \text{ m}^{-2}$) at temperatures higher than 246 K. The ice nucleation active surface site densities were generally at least one order of magnitude lower than those for desert dust.

We also tentatively show that nebulisation enhances the ice nucleating ability of some cell cultures. We suggest that the aerosolisation process [using a nebuliser](#) might rupture individual cells allowing ice nucleating macro-molecules to be dispersed through the aerosol population. Alternatively aggregates of cells or colloidal material may be broken up during aerosolisation. This [might be unlikely to be relevant for environmental conditions. Here, this](#) may lead to the aerosolised samples in the AIDA chamber having a greater ice nucleating activity than they would otherwise have. Pipetting of droplets, as done for the μl -NIPI measurements, might be much less likely to exert sufficient force on the samples to break up cells, aggregates or colloidal material. Our hypothesis is that this process is particularly important for cell cultures and is less important in microlayer samples which consist of organic 'detritus' rather than intact cells (i.e. the organic material is already well dispersed).

In the experiments with microlitre volume droplets (μl -NIPI), which are sensitive to rarer ice nucleating particles, some of the SML samples have values of $T_{50} \approx 262 \text{ K}$. This indicates that there is a low concentration of relatively active ice nucleating entities in these samples. The high variability observed in the high temperature regime suggests that there is a substantial variability in the presence of INP in the samples. What gives rise to this variability and what factors control it is a particularly important outstanding question. Previous work has shown that both the type and concentration of INP varies substantially throughout the development and decay of a phytoplankton bloom ([Wang et al., 2015](#))([Wang et al., 2015](#); [McCluskey et al., 2017](#); [Wilbourn et al., 2020](#)). There are perhaps various types of marine INP from different biological sources present in these natural samples. While our results, and those in the literature e.g. Knopf et al. (2011); Alpert et al. (2011a), show that phytoplankton can nucleate ice, it is also feasible that bacteria exploiting organic detritus from a plume might nucleate ice (Fall and Schnell, 1985). The presence of bacterial proteinacious ice nucleating material would be consistent with the observation that INP in microlayer samples are heat sensitive, e.g. Wilson et al. (2015); Irish et al. (2017). However, a heat treatment test (~~not shown~~[see appendix](#)) on SM100 did not give a [strong-clear](#) indication for this hypothesis: in the low temperature regime no heat sensitivity of freezing~~and in the high temperature regime only a weak heat sensitivity of freezing~~, [but a significant deactivation of freezing on heating](#) could be seen [in the high temperature regime](#) for that sample. Since bacteria tend to be larger than 200 nm and bacterial ice active proteins are cell-membrane bound, one would expect to lose the ice activity associated with bacteria when filtering the sample through a $0.2 \mu\text{m}$ filter (Maki et al., 1974; Murray et al., 2012). This could not be seen in our results of the differently treated ASCOS samples, where the filtered ASCOS sample $< 0.22 \mu\text{m}$ did not show any reduction in ice nucleation activity. The ice nucleation activity of this sample indicated that macromolecules are responsible for the freezing, which were highly concentrated in the sample. ~~The fact~~[It was suggested in literature](#) that marine INP are very small and heat sensitive ([Schnell and Vali, 1975](#); [Vali et al., 1976](#); [Wilson et al., 2015](#); [Irish et al., 2017, 2019b](#)), [which](#) is consistent with an ice nucleating protein [responsible for the INP activity](#), similar to those found in terrestrial fungi (Pouleur et al., 1992; O'Sullivan et al., 2015, 2016). Marine viruses may also fit this size requirement, although we are not aware of any studies on them for ice nucleation. A different candidate could be bacterial vesicles which are 50 - 200 nm particles and can retain the ice nucleating activity of their parent bacterium (Phelps et al., 1986). Another possibility is that the ice nucleating ability of the organic material in seawater is in part due to riverine input. River water is known to harbour large quantities of macromolecular INP (Larsen

et al., 2017; Moffett et al., 2018) and the observed anti-correlation between INP and salinity is consistent with a significant
730 riverine input of INP to some marine environments (Irish et al., 2019b). Given the massive diversity of the high temperature
INP observed in seawater in this and previous studies, e.g. Schnell and Vali (1975); Schnell (1977); Wilson et al. (2015); Irish
et al. (2017, 2019b), it is likely that the sources of these INP are also highly variable and heterogeneous, much as they are in
the terrestrial environment.

From our study it is difficult to answer the question whether Arctic regions may have local marine sources of INPs and how
735 much they influence Arctic mixed-phase clouds. At temperature above 248 K the ice nucleation activity of the investigated
samples was very diverse, with some samples reaching a quite high median freezing temperature of 262 K, thus potentially
being able to trigger freezing in Arctic mixed-phase clouds. The measurements in the temperature regime below 248 K on the
other hand did not show that the samples were particularly ice active, especially when compared to dust, despite the fact that
the results show an upper limit for n_s . Both measurements differentiated in the way the samples were analysed (bulk vs. aerosol
740 phase). This was most relevant for the cultured samples, giving some hint that aerosolisation of cell cultures may change the
ice nucleation activity of these, a process that could be important in the environment as well.

Appendix A: Heat test SM100




Figure A1. Heat test of SM100, measured with a) AIDA (Surface active site density n_s as a measure for ice nucleation activity at different temperatures) and b) μl -NIPI (Fraction frozen curve). The temperature in the μl -NIPI plot was corrected for freezing point depression caused by salts.

Code and data availability. The data will be available at the KITopen data repository (<https://www.bibliothek.kit.edu/cms/kitopen.php>).

Author contributions. **Conceptualisation measurement campaign:** MS, RW; **Measurement campaign:** MS, RW (overall lead); RW, LI (AIDA); GCEP, MPA (NIPI); SB, KH (INKA); SC (CCN); AAK, LI (ESEM); **Samples:** EG (algal cultures), GCEP (SML), CL (ASCOS); **Analysis of the data:** LI, RU, RW (AIDA); GCEP, MPA (NIPI); SB, KH, TS (INKA); SC (CCN); AAK (ESEM); **Visualisation:** LI; Fig. 1: LI, MS; Fig. 10: AAK; **Writing:** LI, MS, RW, KH, BJM, GCEP; **Review & editing:** All authors.

Competing interests. The authors declare that no competing interests are present.

Acknowledgements. We gratefully acknowledge the support of the Engineering and Infrastructure group of IMK-AAF, in particular Olga Dombrowski, Rainer Buschbacher, Tomasz Chudy, Steffen Vogt, and Georg Scheurig. This study was supported by the Helmholtz-Gemeinschaft Deutscher Forschungszentren as part of the program "Atmosphere and Climate". We thank Victoria E. Irish for shipment and help with the SML samples (STN). We thank Nadine Hoffmann (IMK-AAF) for preparing the SM samples for the ESEM measurements. We thank EUROCHAMP-2020 for TNA (Trans-national Access) support and funding and the Bolin Centre for Climate Research for supporting our data workshop held in Stockholm in 2017. LI was supported by the Swiss National Science Foundation (Early Postdoc.Mobility) and the Swedish Science Foundation (Vetenskapsrådet), with grant number 2015-05318. MES was supported by the Swedish Science Foundation (Vetenskapsrådet) with grant number 2016-05100. [CL was supported by the Swedish Science Foundation \(Vetenskapsrådet\) with grant number 2016-03518.](#) BJM acknowledge the European Research Council (MarineIce, 648661), MB and SC Aarhus University and Hakon Lund Foundation and AKB the Natural Sciences and Engineering Research Council of Canada for funding. [We thank the two anonymous reviewers for valuable feedback.](#)

760 **References**

- Alpert, P. A., Aller, J. Y., and Knopf, D. A.: Initiation of the ice phase by marine biogenic surfaces in supersaturated gas and supercooled aqueous phases, *Phys. Chem. Chem. Phys.*, 13, 19 882–19 894, <https://doi.org/10.1039/C1CP21844A>, 2011a.
- Alpert, P. A., Aller, J. Y., and Knopf, D. A.: Ice nucleation from aqueous NaCl droplets with and without marine diatoms, *Atmos. Chem. Phys.*, 11, 5539–5555, <https://doi.org/10.5194/acp-11-5539-2011>, 2011b.
- 765 Alpert, P. A., Kilhau, W. P., Bothe, D. W., Radway, J. C., Aller, J. Y., and Knopf, D. A.: The influence of marine microbial activities on aerosol production: A laboratory mesocosm study, *Journal of Geophysical Research: Atmospheres*, 120, 8841–8860, <https://doi.org/10.1002/2015JD023469>, 2015.
- Alsved, M., Holm, S., Christiansen, S., Smidt, M., Rosati, B., Ling, M., Boesen, T., Finster, K., Bilde, M., Löndahl, J., and Šantl Temkiv, T.: Effect of Aerosolization and Drying on the Viability of *Pseudomonas syringae* Cells, *Frontiers in Microbiology*, 9, 3086, <https://doi.org/10.3389/fmicb.2018.03086>, 2018.
- 770 Battan, L. J. and Riley, J. J.: Ice-crystal nuclei and maritime air, *Journal of Meteorology*, 17, 675–676, 1960.
- Benz, S., Megahed, K., Möhler, O., Saathoff, H., Wagner, R., and Schurath, U.: T-dependent rate measurements of homogeneous ice nucleation in cloud droplets using a large atmospheric simulation chamber, *J. Photochem. Photobiol. A*, 176, 208–217, <https://doi.org/10.1016/j.jphotochem.2005.08.026>, 2005.
- 775 Bigg, E. K.: A new Technique for Counting Ice-Forming Nuclei in Aerosols, *Tellus*, 9, 394–400, <https://doi.org/10.1111/j.2153-3490.1957.tb01895.x>, 1957.
- Bigg, E. K.: Ice nucleus concentrations in remote areas, *J. Atmos. Sci.*, 30, 1153–1157, 1973.
- Bigg, E. K.: Long-term trends in ice nucleus concentrations, *Atmospheric research*, 25, 409–415, 1990.
- Bigg, E. K.: Ice forming nuclei in the high Arctic, *Tellus B: Chemical and Physical Meteorology*, 48, 223–233, 1996.
- 780 Bigg, E. K., Mossop, S. C., Meade, R. T., and Thorndike, N. S. C.: The measurement of ice nucleus concentrations by means of Millipore filters, *Journal of Applied Meteorology*, 2, 266–269, 1963.
- Birstein, S. J. and Anderson, C. E.: Preliminary report on sea salt as an ice nucleus, *Journal of Meteorology*, 10, 166–166, 1953.
- Booth, B. C. and Horner, R. A.: Microalgae on the arctic ocean section, 1994: species abundance and biomass, *Deep Sea Research Part II: Topical Studies in Oceanography*, 44, 1607–1622, [https://doi.org/10.1016/S0967-0645\(97\)00057-X](https://doi.org/10.1016/S0967-0645(97)00057-X), 1997.
- 785 Borkman, D. G. and Smayda, T.: Multidecadal (1959–1997) changes in *Skeletonema* abundance and seasonal bloom patterns in Narragansett Bay, Rhode Island, USA, *Journal of Sea Research*, 61, 84–94, <https://doi.org/10.1016/j.seares.2008.10.004>, 2009.
- Borys, R. D.: Studies of ice nucleation by arctic aerosol on AGASP-II, *Journal of Atmospheric Chemistry*, 9, 169–185, 1989.
- Borys, R. D. and Grant, L. O.: The Effects of long-range transport of air pollutants on Arctic cloud-active aerosol, *Atmospheric science paper*; no. 367, 1983.
- 790 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y.: Clouds and Aerosols, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, 2013.
- 795 Brier, G. W. and Kline, D. B.: Ocean water as a source of ice nuclei, *Science*, 130, 717–718, 1959.

- Burrows, S. M., Hoose, C., Pöschl, U., and Lawrence, M. G.: Ice nuclei in marine air: biogenic particles or dust?, *Atmos. Chem. Phys.*, 13, 245–267, <https://doi.org/10.5194/acp-13-245-2013>, 2013.
- Canesi, K. L. and Rynearson, T. A.: Temporal variation of *Skeletonema* community composition from a long-term time series in Narragansett Bay identified using high-throughput DNA sequencing, *Mar. Ecol. Prog. Ser.*, 556, 1–16, <https://doi.org/10.3354/meps11843>, 2016.
- 800 Chahine, M. T.: The hydrological cycle and its influence on climate, *Nature*, 359, 373–380, 1992.
- Christiansen, S., Salter, M. E., Gorokhova, E., Nguyen, Q. T., and Bilde, M.: Sea Spray Aerosol Formation: Laboratory Results on the Role of Air Entrainment, Water Temperature, and Phytoplankton Biomass, *Environmental science & technology*, 53, 13 107–13 116, 2019.
- Collins, D. B., Zhao, D. F., Ruppel, M. J., Laskina, O., Grandquist, J. R., Modini, R. L., Stokes, M. D., Russell, L. M., Bertram, T. H., Grassian, V. H., Deane, G. B., and Prather, K. A.: Direct aerosol chemical composition measurements to evaluate the physico-chemical differences between controlled sea spray aerosol generation schemes, *Atmospheric Measurement Techniques*, 7, 3667–3683, <https://doi.org/10.5194/amt-7-3667-2014>, 2014.
- 805 Creamean, J. M., Kirpes, R. M., Pratt, K. A., Spada, N. J., Maahn, M., de Boer, G., Schnell, R. C., and China, S.: Marine and terrestrial influences on ice nucleating particles during continuous springtime measurements in an Arctic oilfield location, *Atmospheric Chemistry and Physics*, 18, 18 023–18 042, <https://doi.org/10.5194/acp-18-18023-2018>, 2018.
- 810 Creamean, J. M., Cross, J. N., Pickart, R., McRaven, L., Lin, P., Pacini, A., Hanlon, R., Schmale, D. G., Cenicerros, J., Aydell, T., Colombi, N., Bolger, E., and DeMott, P. J.: Ice Nucleating Particles Carried From Below a Phytoplankton Bloom to the Arctic Atmosphere, *Geophys. Res. Lett.*, 46, 8572–8581, <https://doi.org/10.1029/2019GL083039>, 2019.
- DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T., and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *Proceedings of the National Academy of Science of the United States of America (PNAS)*, 107, 11 217–11 222, 2010.
- 815 DeMott, P. J., Hill, T. C. J., McCluskey, C. S., Prather, K. A., Collins, D. B., Sullivan, R. C., Ruppel, M. J., Mason, R. H., Irish, V. E., Lee, T., Hwang, C. Y., Rhee, T. S., Snider, J. R., McMeeking, G. R., Dhaniyala, S., Lewis, E. R., Wentzell, J. J. B., Abbatt, J., Lee, C., Sultana, C. M., Ault, A. P., Axson, J. L., Diaz Martinez, M., Venero, I., Santos-Figueroa, G., Stokes, M. D., Deane, G. B., Mayol-Bracero, O. L., Grassian, V. H., Bertram, T. H., Bertram, A. K., Moffett, B. F., and Franc, G. D.: Sea spray aerosol as a unique source of ice nucleating particles, *Proceedings of the National Academy of Sciences*, 113, 5797–5803, <https://doi.org/10.1073/pnas.1514034112>, 2016.
- 820 DeMott, P. J., Mason, R. H., McCluskey, C. S., Hill, T. C. J., Perkins, R. J., Desyaterik, Y., Bertram, A. K., Trueblood, J. V., Grassian, V. H., Qiu, Y., Molinero, V., Tobo, Y., Sultana, C. M., Lee, C., and Prather, K. A.: Ice nucleation by particles containing long-chain fatty acids of relevance to freezing by sea spray aerosols, *Environ. Sci.: Processes Impacts*, 20, 1559–1569, <https://doi.org/10.1039/C8EM00386F>, 2018a.
- 825 DeMott, P. J., Möhler, O., Cziczo, D. J., Hiranuma, N., Petters, M. D., Petters, S. S., Belosi, F., Bingemer, H. G., Brooks, S. D., Budke, C., Burkert-Kohn, M., Collier, K. N., Danielczok, A., Eppers, O., Felgitsch, L., Garimella, S., Grothe, H., Herenz, P., Hill, T. C. J., Höhler, K., Kanji, Z. A., Kiselev, A., Koop, T., Kristensen, T. B., Krüger, K., Kulkarni, G., Levin, E. J. T., Murray, B. J., Nicosia, A., O’Sullivan, D., Peckaus, A., Polen, M. J., Price, H. C., Reicher, N., Rothenberg, D. A., Rudich, Y., Santachiara, G., Schiebel, T., Schrod, J., Seifried, T. M., Stratmann, F., Sullivan, R. C., Suski, K. J., Szakáll, M., Taylor, H. P., Ullrich, R., Vergara-Temprado, J., Wagner, R., Whale, T. F., Weber, D., Welti, A., Wilson, T. W., Wolf, M. J., and Zenker, J.: The Fifth International Workshop on Ice Nucleation phase 2 (FIN-02): Laboratory intercomparison of ice nucleation measurements, *Atmospheric Measurement Techniques Discussions*, 2018, 1–44, <https://doi.org/10.5194/amt-2018-191>, 2018b.
- 830

- Fahey, D. W., Gao, R.-S., Möhler, O., Saathoff, H., Schiller, C., Ebert, V., Krämer, M., Peter, T., Amarouche, N., Avallone, L. M., Bauer, R., Bozóki, Z., Christensen, L. E., Davis, S. M., Durrý, G., Dyroff, C., Herman, R. L., Hunsmann, S., Khaykin, S. M., Mackrodt, P., Meyer, J.,
835 Smith, J. B., Spelten, N., Troy, R. F., Vömel, H., Wagner, S., and Wienhold, F. G.: The AquaVIT-1 intercomparison of atmospheric water vapor measurement techniques, *Atmospheric Measurement Techniques*, 7, 3177–3213, <https://doi.org/10.5194/amt-7-3177-2014>, 2014.
- Fall, R. and Schnell, R. C.: Association of an ice-nucleating pseudomonad with cultures of the marine dinoflagellate, *Heterocapsa niei*, *J. Mar. Res.*, 43, 257–265, <https://doi.org/10.1357/002224085788437370>, 1985.
- Flyger, H. and Heidam, N. Z.: Ground level measurements of the summer tropospheric aerosol in Northern Greenland, *Journal of Aerosol Science*, 9, 157–168, [https://doi.org/10.1016/0021-8502\(78\)90075-7](https://doi.org/10.1016/0021-8502(78)90075-7), 1978.
840
- Fountain, A. G. and Ohtake, T.: Concentrations and source areas of ice nuclei in the Alaskan atmosphere, *Journal of climate and applied meteorology*, 24, 377–382, 1985.
- Gagin, A. and Arroyo, M.: A thermal diffusion chamber for the measurement of ice nucleus concentration, *J. Rech. Atmos.*, 55, 115–122, 1969.
- 845 Gantt, B. and Meskhidze, N.: The physical and chemical characteristics of marine primary organic aerosol: a review, *Atmos. Chem. Phys.*, 13, 3979–3996, <https://doi.org/10.5194/acp-13-3979-2013>, 2013.
- Gao, Q., Leck, C., Rauschenberg, C., and Matrai, P. A.: On the chemical dynamics of extracellular polysaccharides in the high Arctic surface microlayer, *Ocean Science*, 8, 401–418, <https://doi.org/10.5194/os-8-401-2012>, 2012.
- Garrett, T. J., Maestas, M. M., Krueger, S. K., and Schmidt, C. T.: Acceleration by aerosol of a radiative-thermodynamic cloud feedback
850 influencing Arctic surface warming, *Geophys. Res. Lett.*, 36, 2009.
- Gong, X., Wex, H., Müller, T., Wiedensohler, A., Höhler, K., Kandler, K., Ma, N., Dietel, B., Schiebel, T., Möhler, O., and Stratmann, F.: Characterization of aerosol properties at Cyprus, focusing on cloud condensation nuclei and ice-nucleating particles, *Atmospheric Chemistry and Physics*, 19, 10 883–10 900, <https://doi.org/10.5194/acp-19-10883-2019>, 2019.
- Gong, X., Wex, H., van Pinxteren, M., Triesch, N., Fomba, K. W., Lubitz, J., Stolle, C., Robinson, T.-B., Müller, T., Herrmann, H., and
855 Stratmann, F.: Characterization of aerosol particles at Cabo Verde close to sea level and at the cloud level – Part 2: Ice-nucleating particles in air, cloud and seawater, *Atmospheric Chemistry and Physics*, 20, 1451–1468, <https://doi.org/10.5194/acp-20-1451-2020>, 2020.
- Hartmann, M., Adachi, K., Eppers, O., Haas, C., Herber, A., Holzinger, R., Hünerbein, A., Jäkel, E., Jentsch, C., van Pinxteren, M., Wex, H., Willmes, S., and Stratmann, F.: Wintertime Airborne Measurements of Ice Nucleating Particles in the High Arctic: A Hint to a Marine, Biogenic Source for Ice Nucleating Particles, *Geophysical Research Letters*, 47, e2020GL087 770, <https://doi.org/10.1029/2020GL087770>,
860 2020.
- Harvey, G. W.: Microlayer collection from the sea surface: a new method and initial results, *Limnol. Oceanogr.*, 11, 608–613, <https://doi.org/10.4319/lo.1966.11.4.0608>, 1966.
- Henderson, R., Chips, M., Cornwell, N., Hitchins, P., Holden, B., Hurley, S., Parsons, S. A., Wetherill, A., and Jefferson, B.: Experiences of algae in UK waters: a treatment perspective, *Water and Environment Journal*, 22, 184–192, <https://doi.org/10.1111/j.1747-6593.2007.00100.x>, 2008.
865
- Herbert, R. J., Murray, B. J., Whale, T. F., Dobbie, S. J., and Atkinson, J. D.: Representing time-dependent freezing behaviour in immersion mode ice nucleation, *Atmos. Chem. Phys.*, 14, 8501–8520, <https://doi.org/10.5194/acp-14-8501-2014>, 2014.
- Hiranuma, N., Adachi, K., Bell, D. M., Belosi, F., Beydoun, H., Bhaduri, B., Bingemer, H., Budke, C., Clemen, H.-C., Conen, F., Cory, K. M., Curtius, J., DeMott, P. J., Eppers, O., Grawe, S., Hartmann, S., Hoffmann, N., Höhler, K., Jantsch, E., Kiselev, A., Koop, T.,
870 Kulkarni, G., Mayer, A., Murakami, M., Murray, B. J., Nicosia, A., Petters, M. D., Piazza, M., Polen, M., Reicher, N., Rudich, Y., Saito,

- A., Santachiara, G., Schiebel, T., Schill, G. P., Schneider, J., Segev, L., Stopelli, E., Sullivan, R. C., Suski, K., Szakáll, M., Tajiri, T., Taylor, H., Tobo, Y., Ullrich, R., Weber, D., Wex, H., Whale, T. F., Whiteside, C. L., Yamashita, K., Zelenyuk, A., and Möhler, O.: A comprehensive characterization of ice nucleation by three different types of cellulose particles immersed in water, *Atmos. Chem. Phys.*, 19, 4823–4849, <https://doi.org/10.5194/acp-19-4823-2019>, 2019.
- 875 Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 9817–9854, <https://doi.org/10.5194/acp-12-9817-2012>, 2012.
- Huang, W. T. K., Ickes, L., Tegen, I., Rinaldi, M., Ceburnis, D., and Lohmann, U.: Global relevance of marine organic aerosol as ice nucleating particles, *Atmos. Chem. Phys.*, 18, 11 423–11 445, <https://doi.org/10.5194/acp-18-11423-2018>, 2018.
- Intrieri, J. M., Fairall, C. W., Shupe, M. D., Persson, P. O. G., Andreas, E. L., Guest, P. S., and Moritz, R. E.: An annual cycle of Arctic surface cloud forcing at SHEBA, *Journal of Geophysical Research: Oceans*, 107, 2002.
- 880 Irish, V. E., Elizondo, P., Chen, J., Chou, C., Charette, J., Lizotte, M., Ladino, L. A., Wilson, T. W., Gosselin, M., Murray, B. J., Polishchuk, E., Abbatt, J. P. D., Miller, L. A., and Bertram, A. K.: Ice-nucleating particles in Canadian Arctic sea-surface microlayer and bulk seawater, *Atmos. Chem. Phys.*, 17, 10 583–10 595, <https://doi.org/10.5194/acp-17-10583-2017>, 2017.
- Irish, V. E., Hanna, S. J., Willis, M. D., China, S., Thomas, J. L., Wentzell, J. J. B., Cirisan, A., Si, M., Leaitch, W. R., Murphy, J. G., Abbatt, J. P. D., Laskin, A., Girard, E., and Bertram, A. K.: Ice nucleating particles in the marine boundary layer in the Canadian Arctic during summer 2014, *Atmos. Chem. Phys.*, 19, 1027–1039, <https://doi.org/10.5194/acp-19-1027-2019>, 2019a.
- 885 Irish, V. E., Hanna, S. J., Xi, Y., Boyer, M., Polishchuk, E., Ahmed, M., Chen, J., Abbatt, J. P. D., Gosselin, M., Chang, R., Miller, L. A., and Bertram, A. K.: Revisiting properties and concentrations of ice-nucleating particles in the sea surface microlayer and bulk seawater in the Canadian Arctic during summer, *Atmos. Chem. Phys.*, 19, 7775–7787, <https://doi.org/10.5194/acp-19-7775-2019>, 2019b.
- 890 Isono, K., Komabayasi, M., and Ono, A.: The Nature and the Origin of Ice Nuclei in the. Atmosphere, *Journal of the Meteorological Society of Japan. Ser. II*, 37, 211–233, 1959.
- Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Krämer, M.: Overview of Ice Nucleating Particles, *Meteorological Monographs*, 58, 1.1–1.33, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0006.1>, 2017.
- King, S. M., Butcher, A. C., Rosenoern, T., Coz, E., Lieke, K. I., de Leeuw, G., Nilsson, E. D., and Bilde, M.: Investigating Primary Marine Aerosol Properties: CCN Activity of Sea Salt and Mixed Inorganic–Organic Particles, *Environmental Science & Technology*, 46, 10 405–10 412, <https://doi.org/10.1021/es300574u>, 2012.
- 895 Kline, D. B.: Recent Observations of Freezing Nuclei Variations at Ground Level, pp. 240–246, American Geophysical Union (AGU), <https://doi.org/10.1029/GM005p0240>, 1960.
- Kline, D. B. and Brier, G. W.: A note on freezing nuclei anomalies, *Monthly Weather Review*, 86, 329–333, [https://doi.org/10.1175/1520-0493\(1958\)086<0329:ANOFNA>2.0.CO;2](https://doi.org/10.1175/1520-0493(1958)086<0329:ANOFNA>2.0.CO;2), 1958.
- 900 Knopf, D. A., Alpert, P. A., Wang, B., and Aller, J. Y.: Stimulation of ice nucleation by marine diatoms, *Nature Geosci*, 4, 88–90, <https://doi.org/10.1038/ngeo1037>, 2011.
- Knulst, J. C., Rosenberger, D., Thompson, B., and Paatero, J.: Intensive Sea Surface Microlayer Investigations of Open Leads in the Pack Ice during Arctic Ocean 2001 Expedition, *Langmuir*, 19, 10 194–10 199, <https://doi.org/10.1021/la035069+>, 2003.
- 905 Kooistra, W. H., Sarno, D., Balzano, S., Gu, H., Andersen, R. A., and Zingone, A.: Global Diversity and Biogeography of Skeletonema Species (Bacillariophyta), *Protist*, 159, 177–193, <https://doi.org/10.1016/j.protis.2007.09.004>, 2008.

- Koop, T., Kapilashrami, A., Molina, L. T., and Molina, M. J.: Phase transitions of sea-salt/water mixtures at low temperatures: Implications for ozone chemistry in the polar marine boundary layer, *Journal of Geophysical Research: Atmospheres*, 105, 26 393–26402, <https://doi.org/10.1029/2000JD900413>, 2000.
- 910 Ladino, L. A., Yakobi-Hancock, J. D., Kilthau, W. P., Mason, R. H., Si, M., Li, J., Miller, L. A., Schiller, C. L., Huffman, J. A., Aller, J. Y., Knopf, D. A., Bertram, A. K., and Abbatt, J. P. D.: Addressing the ice nucleating abilities of marine aerosol: A combination of deposition mode laboratory and field measurements, *Atmos. Environ.*, 132, 1–10, <https://doi.org/10.1016/j.atmosenv.2016.02.028>, 2016.
- Ladino, L. A., Raga, G. B., Alvarez-Ospina, H., Andino-Enrriquez, M. A., Rosas, I., Martínez, L., Salinas, E., Miranda, J., Ramírez-Díaz, Z., Figueroa, B., Chou, C., Bertram, A. K., Quintana, E. T., Maldonado, L. A., García-Reynoso, A., Si, M., and Irish, V. E.: Ice-nucleating
 915 particles in a coastal tropical site, *Atmospheric Chemistry and Physics*, 19, 6147–6165, <https://doi.org/10.5194/acp-19-6147-2019>, 2019.
- Langer, G. and Rodgers, J.: An Experimental Study of the Detection of Ice Nuclei on Membrane Filters and Other Substrata, *Journal of Applied Meteorology*, 14, 560–570, [https://doi.org/10.1175/1520-0450\(1975\)014<0560:AESOTD>2.0.CO;2](https://doi.org/10.1175/1520-0450(1975)014<0560:AESOTD>2.0.CO;2), 1975.
- Langer, G., Rosinski, J., and Edwards, C. P.: A Continuous Ice Nucleus Counter and its Application to Tracking in the Troposphere, *Journal of Applied Meteorology*, 6, 114–125, [https://doi.org/10.1175/1520-0450\(1967\)006<0114:ACINCA>2.0.CO;2](https://doi.org/10.1175/1520-0450(1967)006<0114:ACINCA>2.0.CO;2), 1967.
- 920 Larsen, J. A., Conen, F., and Alewell, C.: Export of ice nucleating particles from a watershed, *Royal Society Open Science*, 4, 170 213, <https://doi.org/10.1098/rsos.170213>, 2017.
- Leck, C., Norman, M., Bigg, E. K., and Hillamo, R.: Chemical composition and sources of the high Arctic aerosol relevant for cloud formation, *Journal of Geophysical Research: Atmospheres*, 107, AAC 1–1–AAC 1–17, <https://doi.org/10.1029/2001JD001463>, 2002.
- Maki, L. R., Galyan, E. L., Chang-Chien, M.-M., and Caldwell, D. R.: Ice nucleation induced by *Pseudomonas syringae*, *Appl. Environ. Microbiol.*, 28, 456–459, 1974.
- 925 Mason, R. H., Si, M., Li, J., Chou, C., Dickie, R., Toom-Sauntry, D., Pöhlker, C., Yakobi-Hancock, J. D., Ladino, L. A., Jones, K., Leaitch, W. R., Schiller, C. L., Abbatt, J. P. D., Huffman, J. A., and Bertram, A. K.: Ice nucleating particles at a coastal marine boundary layer site: correlations with aerosol type and meteorological conditions, *Atmospheric Chemistry and Physics*, 15, 12 547–12 566, <https://doi.org/10.5194/acp-15-12547-2015>, 2015.
- 930 Matrai, P. A., Tranvik, L., Leck, C., and Knulst, J. C.: Are high Arctic surface microlayers a potential source of aerosol organic precursors?, *Mar. Chem.*, 108, 109–122, <https://doi.org/10.1016/j.marchem.2007.11.001>, 2008.
- McCluskey, C. S., Hill, T. C. J., Malfatti, F., Sultana, C. M., Lee, C., Santander, M. V., Beall, C. M., Moore, K. A., Cornwell, G. C., Collins, D. B., Prather, K. A., Jayarathne, T., Stone, E. A., Azam, F., Kreidenweis, S. M., and DeMott, P. J.: A Dynamic Link between Ice Nucleating Particles Released in Nascent Sea Spray Aerosol and Oceanic Biological Activity during Two Mesocosm Experiments, *J. Atmos. Sci.*, 74, 151–166, <https://doi.org/10.1175/JAS-D-16-0087.1>, 2017.
- 935 McCluskey, C. S., Hill, T. C. J., Humphries, R. S., Rauker, A. M., Moreau, S., Stratton, P. G., Chambers, S. D., Williams, A. G., McRobert, I., Ward, J., Keywood, M. D., Harnwell, J., Ponsonby, W., Loh, Z. M., Krummel, P. B., Protat, A., Kreidenweis, S. M., and DeMott, P. J.: Observations of Ice Nucleating Particles Over Southern Ocean Waters, *Geophysical Research Letters*, 45, 11,989–11,997, <https://doi.org/10.1029/2018GL079981>, 2018a.
- 940 McCluskey, C. S., Ovadnevaite, J., Rinaldi, M., Atkinson, J., Belosi, F., Ceburnis, D., Marullo, S., Hill, T. C. J., Lohmann, U., Kanji, Z. A., O’Dowd, C., Kreidenweis, S. M., and DeMott, P. J.: Marine and Terrestrial Organic Ice-Nucleating Particles in Pristine Marine to Continentally Influenced Northeast Atlantic Air Masses, *Journal of Geophysical Research: Atmospheres*, 123, 6196–6212, <https://doi.org/10.1029/2017JD028033>, 2018b.

- 945 Moffett, B., Hill, T., and DeMott, P.: Abundance of biological ice nucleating particles in the Mississippi and its major tributaries, *Atmosphere*, 9, 307, 2018.
- Möhler, O., Stetzer, O., Schaefers, S., Linke, C., Schnaiter, M., Tiede, R., Saathoff, H., Krämer, M., Mangold, A., Budz, P., Zink, P., Schreiner, J., Mauersberger, K., Haag, W., Kärcher, B., and Schurath, U.: Experimental investigation of homogeneous freezing of sulphuric acid particles in the aerosol chamber AIDA, *Atmos. Chem. Phys.*, 3, 211–223, <https://doi.org/10.5194/acp-3-211-2003>, 2003.
- 950 Morrison, H., de Boer, G., Feingold, G., Harrington, J., Shupe, M. D., and Sulia, K.: Resilience of persistent Arctic mixed-phase clouds, *Nature Geoscience*, 5, 11–17, 2012.
- Murphy, D. M. and Koop, T.: Review of the vapour pressures of ice and supercooled water for atmospheric applications, *Quart. J. Roy. Meteor. Soc.*, 131, 1539–1565, <https://doi.org/10.1256/qj.04.94>, 2005.
- Murray, B. J., O'Sullivan, D., Atkinson, J. D., and Webb, M. E.: Ice nucleation by particles immersed in supercooled cloud droplets, *Chem. Soc. Rev.*, 41, 6519–6554, <https://doi.org/10.1039/C2CS35200A>, 2012.
- 955 Möhler, O., Büttner, S., Linke, C., Schnaiter, M., Saathoff, H., Stetzer, O., Wagner, R., Krämer, M., Mangold, A., Ebert, V., and Schurath, U.: Effect of sulfuric acid coating on heterogeneous ice nucleation by soot aerosol particles, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/10.1029/2004JD005169>, 2005.
- Möhler, O., Benz, S., Saathoff, H., Schnaiter, M., Wagner, R., Schneider, J., Walter, S., Ebert, V., and Wagner, S.: The effect of organic coating on the heterogeneous ice nucleation efficiency of mineral dust aerosols, *Environ. Res. Lett.*, 3, 025 007, <https://doi.org/10.1088/1748-9326/3/2/025007>, 2008.
- 960 Nagamoto, C. T., Rosinski, J., Haagenson, P. L., Michalowska-Smak, A., and Parungo, F.: Characteristics of ice-forming nuclei in continental-maritime air, *J. Aerosol Sci.*, 15, 147–166, 1984.
- Niemand, M., Möhler, O., Vogel, B., Vogel, H., Hoose, C., Connolly, P., Klein, H., Bingemer, H., DeMott, P., Skrotzki, J., and Leisner, T.: A Particle-Surface-Area-Based Parameterization of Immersion Freezing on Desert Dust Particles, *J. Atmos. Sci.*, 69, 3077–3092, <https://doi.org/10.1175/JAS-D-11-0249.1>, 2012.
- 965 O'Dowd, C. D., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Yoon, Y. J., and Putaud, J.-P.: Biogenically driven organic contribution to marine aerosol, *Nature*, 431, 676, 2004.
- Orellana, M. V., Matrai, P. A., Leck, C., Rauschenberg, C. D., Lee, A. M., and Coz, E.: Marine microgels as a source of cloud condensation nuclei in the high Arctic, *Proceedings of the National Academy of Sciences*, 108, 13 612–13 617, <https://doi.org/10.1073/pnas.1102457108>, 970 2011.
- O'Sullivan, D., Murray, B. J., Malkin, T. L., Whale, T. F., Umo, N. S., Atkinson, J. D., Price, H. C., Baustian, K. J., Browse, J., and Webb, M. E.: Ice nucleation by fertile soil dusts: relative importance of mineral and biogenic components, *Atmos. Chem. Phys.*, 14, 1853–1867, <https://doi.org/10.5194/acp-14-1853-2014>, 2014.
- O'Sullivan, D., Murray, B. J., Ross, J. F., Whale, T. F., Price, H. C., Atkinson, J. D., Umo, N. S., and Webb, M. E.: The relevance of nanoscale 975 biological fragments for ice nucleation in clouds, *Scientific reports*, 5, 8082, 2015.
- O'Sullivan, D., Murray, B. J., Ross, J. F., and Webb, M. E.: The adsorption of fungal ice-nucleating proteins on mineral dusts: a terrestrial reservoir of atmospheric ice-nucleating particles, *Atmos. Chem. Phys.*, 16, 7879–7887, <https://doi.org/10.5194/acp-16-7879-2016>, 2016.
- Parker, L. V., Sullivan, C. W., Forest, T. W., and Ackley, S. F.: Ice nucleation activity of Antarctic marine microorganisms, *Antarctic J*, 20, 126–127, 1985.
- 980 Phelps, P., Giddings, T. H., Prochoda, M., and Fall, R.: Release of cell-free ice nuclei by *Erwinia herbicola*, *J. Bacteriol.*, 167, 496–502, 1986.

- Pithan, F. and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models, *Nature Geoscience*, 7, 181–184, 2014.
- Polen, M., Lawlis, E., and Sullivan, R. C.: The unstable ice nucleation properties of Snomax® bacterial particles, *Journal of Geophysical Research: Atmospheres*, 121, 11,666–11,678, <https://doi.org/10.1002/2016JD025251>, 2016.
- 985 Pouleur, S., Richard, C., Martin, J.-G., and Antoun, H.: Ice Nucleation Activity in *Fusarium acuminatum* and *Fusarium avenaceum*, *Appl. Environ. Microbiol.*, 58, 2960–2964, 1992.
- Prather, K. A., Bertram, T. H., Grassian, V. H., Deane, G. B., Stokes, M. D., DeMott, P. J., Aluwihare, L. I., Palenik, B. P., Azam, F., Seinfeld, J. H., Moffet, R. C., Molina, M. J., Cappa, C. D., Geiger, F. M., Roberts, G. C., Russell, L. M., Ault, A. P., Baltusaitis, J., Collins, D. B., Corrigan, C. E., Cuadra-Rodriguez, L. A., Ebben, C. J., Forestieri, S. D., Guasco, T. L., Hersey, S. P., Kim, M. J., Lambert, W. F., Modini, R. L., Mui, W., Pedler, B. E., Ruppel, M. J., Ryder, O. S., Schoepp, N. G., Sullivan, R. C., and Zhao, D.: Bringing the ocean into the laboratory to probe the chemical complexity of sea spray aerosol, *Proceedings of the National Academy of Sciences*, 110, 7550–7555, <https://doi.org/10.1073/pnas.1300262110>, 2013.
- 990 Radke, L. F., Hobbs, P. V., and Pinnons, J. E.: Observations of Cloud Condensation Nuclei, Sodium-Containing Particles, Ice Nuclei and the Light-Scattering Coefficient Near Barrow, Alaska, *J. Appl. Meteor.*, 15, 982–995, [https://doi.org/10.1175/1520-0450\(1976\)015<0982:OOCNS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1976)015<0982:OOCNS>2.0.CO;2), 1976.
- 995 Rogers, D. C.: Development of a continuous flow thermal gradient diffusion chamber for ice nucleation studies, *Atmos. Res.*, 22, 149–181, [https://doi.org/10.1016/0169-8095\(88\)90005-1](https://doi.org/10.1016/0169-8095(88)90005-1), 1988.
- Rosinski, J.: Cloud condensation nuclei as a real source of ice forming nuclei in continental and marine air masses, *Atmospheric research*, 38, 351–359, 1995.
- 1000 Rosinski, J., Haagenson, P. L., Nagamoto, C. T., and Parungo, F.: Ice-forming nuclei of maritime origin, *J. Aerosol Sci.*, 17, 23–46, [https://doi.org/10.1016/0021-8502\(86\)90004-2](https://doi.org/10.1016/0021-8502(86)90004-2), 1986.
- Rosinski, J., Haagenson, P. L., Nagamoto, C. T., and Parungo, F.: Nature of ice-forming nuclei in marine air masses, *Journal of aerosol science*, 18, 291–309, 1987.
- Rosinski, J., Haagenson, P., Nagamoto, C., Quintana, B., Parungo, F., and Hoyt, S.: Ice-forming nuclei in air masses over the Gulf of Mexico, *J. Aerosol Sci.*, 19, 539–551, [https://doi.org/10.1016/0021-8502\(88\)90206-6](https://doi.org/10.1016/0021-8502(88)90206-6), 1988.
- 1005 Rosinski, J., Nagamoto, C. T., and Zhou, M. Y.: Ice-forming nuclei over the East China Sea, *Atmospheric research*, 36, 95–105, 1995.
- Saravanan, V. and Godhe, A.: Genetic heterogeneity and physiological variation among seasonally separated clones of *Skeletonema marinoi* (Bacillariophyceae) in the Gullmar Fjord, Sweden, *European Journal of Phycology*, 45, 177–190, <https://doi.org/10.1080/09670260903445146>, 2010.
- 1010 Schiebel, T.: Ice Nucleation Activity of Soil Dust Aerosols, PhD thesis; Karlsruher Institut für Technologie (KIT), <https://doi.org/10.5445/IR/1000076327>, 12.04.02; LK 01, 2017.
- Schnell, R. C.: Ice nuclei produced by laboratory cultured marine phytoplankton, *Geophys. Res. Lett.*, 2, 500–502, 1975.
- Schnell, R. C.: Ice Nuclei in Seawater, Fog Water and Marine Air off the Coast of Nova Scotia: Summer 1975, *J. Atmos. Sci.*, 34, 1299–1305, [https://doi.org/10.1175/1520-0469\(1977\)034<1299:INISFW>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1299:INISFW>2.0.CO;2), 1977.
- 1015 Schnell, R. C. and Vali, G.: Freezing nuclei in marine waters, *Tellus*, 27, 321–323, <https://doi.org/10.1111/j.2153-3490.1975.tb01682.x>, 1975.
- Schnell, R. C. and Vali, G.: Biogenic Ice Nuclei: Part I. Terrestrial and Marine Sources, *J. Atmos. Sci.*, 33, 1554–1564, [https://doi.org/10.1175/1520-0469\(1976\)033<1554:BINPIT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033<1554:BINPIT>2.0.CO;2), 1976.

- Shinki, M., Wendeberg, M., Vagle, S., Cullen, J. T., and Hore, D. K.: Characterization of adsorbed microlayer thickness on an oceanic glass
1020 plate sampler, *Limnology and Oceanography: Methods*, 10, 728–735, <https://doi.org/10.4319/lom.2012.10.728>, 2012.
- Shupe, M. D., Matrosov, S. Y., and Uttal, T.: Arctic Mixed-Phase Cloud Properties Derived from Surface-Based Sensors at SHEBA, *J. Atmos.
Sci.*, 63, 697–711, <https://doi.org/10.1175/JAS3659.1>, 2006.
- Si, M., Irish, V. E., Mason, R. H., Vergara-Temprado, J., Hanna, S. J., Ladino, L. A., Yakobi-Hancock, J. D., Schiller, C. L., Wentzell, J. J. B.,
1025 Abbatt, J. P. D., Carslaw, K. S., Murray, B. J., and Bertram, A. K.: Ice-nucleating ability of aerosol particles and possible sources at three
coastal marine sites, *Atmospheric Chemistry and Physics*, 18, 15 669–15 685, <https://doi.org/10.5194/acp-18-15669-2018>, 2018.
- Si, M., Evoy, E., Yun, J., Xi, Y., Hanna, S. J., Chivulescu, A., Rawlings, K., Veber, D., Platt, A., Kunkel, D., Hoor, P., Sharma, S., Leaitch,
W. R., and Bertram, A. K.: Concentrations, composition, and sources of ice-nucleating particles in the Canadian High Arctic during spring
2016, *Atmospheric Chemistry and Physics*, 19, 3007–3024, <https://doi.org/10.5194/acp-19-3007-2019>, 2019.
- Stevenson, C. M.: An improved Millipore filter technique for measuring the concentrations of freezing nuclei in the atmosphere, *Quarterly
1030 Journal of the Royal Meteorological Society*, 94, 35–43, 1968.
- Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M.: *Climate
Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental
Panel on Climate Change*, Cambridge University Press, 2013.
- Suikkanen, S., Hakanen, P., Spilling, K., and Kremp, A.: Allelopathic effects of Baltic Sea spring bloom dinoflagellates on co-occurring
1035 phytoplankton, *Mar. Ecol. Prog. Ser.*, 439, 45–55, <https://doi.org/10.3354/meps09356>, 2011.
- Suski, K. J., Hill, T. C. J., Levin, E. J. T., Miller, A., DeMott, P. J., and Kreidenweis, S. M.: Agricultural harvesting emissions of ice-nucleating
particles, *Atmos. Chem. Phys.*, 18, 13 755–13 771, <https://doi.org/10.5194/acp-18-13755-2018>, 2018.
- Szyrmer, W. and Zawadzki, I.: Biogenic and anthropogenic sources of ice-forming nuclei: A review, *Bulletin of the American Meteorological
Society*, 78, 209–228, 1997.
- 1040 Tang, I. N., Tridico, A. C., and Fung, K. H.: Thermodynamic and optical properties of sea salt aerosols, *Journal of Geophysical Research:
Atmospheres*, 102, 23 269–23 275, <https://doi.org/10.1029/97JD01806>, 1997.
- Tesson, S. V. M. and Šantl Temkiv, T.: Ice Nucleation Activity and Aeolian Dispersal Success in Airborne and Aquatic Microalgae, *Frontiers
in Microbiology*, 9, 2681, <https://doi.org/10.3389/fmicb.2018.02681>, 2018.
- Tjernström, M., Leck, C., Birch, C. E., Bottenheim, J. W., Brooks, B. J., Brooks, I. M., Bäcklin, L., Chang, R. Y.-W., de Leeuw, G., Di Liberto,
1045 L., de la Rosa, S., Granath, E., Graus, M., Hansel, A., Heintzenberg, J., Held, A., Hind, A., Johnston, P., Knulst, J., Martin, M., Matrai,
P. A., Mauritsen, T., Müller, M., Norris, S. J., Orellana, M. V., Orsini, D. A., Paatero, J., Persson, P. O. G., Gao, Q., Rauschenberg, C.,
Ristovski, Z., Sedlar, J., Shupe, M. D., Sierau, B., Sirevaag, A., Sjogren, S., Stetzer, O., Swietlicki, E., Szczodrak, M., Vaattovaara, P.,
Wahlberg, N., Westberg, M., and Wheeler, C. R.: The Arctic Summer Cloud Ocean Study (ASCOS): overview and experimental design,
Atmos. Chem. Phys., 14, 2823–2869, <https://doi.org/10.5194/acp-14-2823-2014>, 2014.
- 1050 Ullrich, R., Hoose, C., Möhler, O., Niemand, M., Wagner, R., Höhler, K., Hiranuma, N., Saathoff, H., and Leisner, T.: A New Ice Nucleation
Active Site Parameterization for Desert Dust and Soot, *J. Atmos. Sci.*, 74, 699–717, <https://doi.org/10.1175/JAS-D-16-0074.1>, 2017.
- Vali, G.: Quantitative evaluation of experimental results an the heterogeneous freezing nucleation of supercooled liquids, *J. Atmos. Sci.*, 28,
402–409, 1971.
- Vali, G.: Atmospheric ice nucleation—A review, *J. Rech. Atmos.*, 19, 105–115, 1985.
- 1055 Vali, G., Christensen, M., Fresh, R. W., Galyan, E. L., Maki, L. R., and Schnell, R. C.: Biogenic ice nuclei. Part II: Bacterial sources, *Journal
of the Atmospheric Sciences*, 33, 1565–1570, 1976.

- Vali, G., DeMott, P. J., Möhler, O., and Whale, T. F.: Technical Note: A proposal for ice nucleation terminology, *Atmos. Chem. Phys.*, 15, 10263–10270, <https://doi.org/10.5194/acp-15-10263-2015>, 2015.
- 1060 Vergara-Temprado, J., Murray, B. J., Wilson, T. W., O’Sullivan, D., Browse, J., Pringle, K. J., Ardon-Dryer, K., Bertram, A. K., Burrows, S. M., Ceburnis, D., DeMott, P. J., Mason, R. H., O’Dowd, C. D., Rinaldi, M., and Carslaw, K. S.: Contribution of feldspar and marine organic aerosols to global ice nucleating particle concentrations, *Atmos. Chem. Phys.*, 17, 3637–3658, <https://doi.org/10.5194/acp-17-3637-2017>, 2017.
- Wagner, R. and Möhler, O.: Heterogeneous ice nucleation ability of crystalline sodium chloride dihydrate particles, *Journal of Geophysical Research: Atmospheres*, 118, 4610–4622, <https://doi.org/10.1002/jgrd.50325>, 2013.
- 1065 Wang, X., Sultana, C. M., Trueblood, J., Hill, T. C. J., Malfatti, F., Lee, C., Laskina, O., Moore, K. A., Beall, C. M., McCluskey, C. S., Cornwell, G. C., Zhou, Y., Cox, J. L., Pendergraft, M. A., Santander, M. V., Bertram, T. H., Cappa, C. D., Azam, F., DeMott, P. J., Grassian, V. H., and Prather, K. A.: Microbial Control of Sea Spray Aerosol Composition: A Tale of Two Blooms, *ACS Central Science*, 1, 124–131, <https://doi.org/10.1021/acscentsci.5b00148>, 2015.
- Warner, J.: An instrument for the measurement of freezing nucleus concentration, *Bulletin de L’Observatoire du Puy de Dome*, pp. 33–46, 1070 1957.
- Welti, A., Müller, K., Fleming, Z. L., and Stratmann, F.: Concentration and variability of ice nuclei in the subtropical maritime boundary layer, *Atmospheric Chemistry and Physics*, 18, 5307–5320, <https://doi.org/10.5194/acp-18-5307-2018>, 2018.
- Welti, A., Bigg, E. K., DeMott, P. J., Gong, X., Hartmann, M., Harvey, M., Henning, S., Herenz, P., Hill, T. C. J., Hornblow, B., Leck, C., Löffler, M., McCluskey, C. S., Rauker, A. M., Schmale, J., Tatzelt, C., van Pinxteren, M., and Stratmann, F.: Ship-based measurements of 1075 ice nuclei concentrations over the Arctic, Atlantic, Pacific and Southern Ocean, *Atmospheric Chemistry and Physics Discussions*, 2020, 1–22, <https://doi.org/10.5194/acp-2020-466>, 2020.
- Wex, H., Huang, L., Zhang, W., Hung, H., Traversi, R., Becagli, S., Sheesley, R. J., Moffett, C. E., Barrett, T. E., Bossi, R., Skov, H., Hünerbein, A., Lubitz, J., Löffler, M., Linke, O., Hartmann, M., Herenz, P., and Stratmann, F.: Annual variability of ice-nucleating particle concentrations at different Arctic locations, *Atmos. Chem. Phys.*, 19, 5293–5311, <https://doi.org/10.5194/acp-19-5293-2019>, 2019.
- 1080 Whale, T. F., Murray, B. J., O’Sullivan, D., Wilson, T. W., Umo, N. S., Baustian, K. J., Atkinson, J. D., Workneh, D. A., and Morris, G. J.: A technique for quantifying heterogeneous ice nucleation in microlitre supercooled water droplets, *Atmospheric Measurement Techniques*, 8, 2437–2447, <https://doi.org/10.5194/amt-8-2437-2015>, 2015.
- Wilbourn, E. K., Thornton, D. C. O., Ott, C., Graff, J., Quinn, P. K., Bates, T. S., Betha, R., Russell, L. M., Behrenfeld, M. J., and Brooks, S. D.: Ice Nucleation by Marine Aerosols Over the North Atlantic Ocean in Late Spring, *Journal of Geophysical Research: Atmospheres*, 1085 125, e2019JD030913, <https://doi.org/10.1029/2019JD030913>, 2020.
- Willis, M. D., Leaitch, W. R., and Abbatt, J. P.: Processes Controlling the Composition and Abundance of Arctic Aerosol, *Reviews of Geophysics*, 56, 621–671, <https://doi.org/10.1029/2018RG000602>, 2018.
- Wilson, T. W., Ladino, L. A., Alpert, P. A., Breckels, M. N., Brooks, I. M., Browse, J., Burrows, S. M., Carslaw, K. S., Huffman, J. A., Judd, C., Kilhau, W. P., Mason, R. H., McFiggans, G., Miller, L. A., Najera, J. J., Polishchuk, E., Rae, S., Schiller, C. L., Si, M., Temprado, 1090 J. V., Whale, T. F., Wong, J. P. S., Wurl, O., Yakobi-Hancock, J. D., Abbatt, J. P. D., Aller, J. Y., Bertram, A. K., Knopf, D. A., and Murray, B. J.: A marine biogenic source of atmospheric ice-nucleating particles, *Nature*, 525, 234–238, <https://doi.org/10.1038/nature14986>, 2015.
- Wolf, M. J., Goodell, M., Dong, E., Dove, L. A., Zhang, C., Franco, L. J., Shen, C., Rutkowski, E. G., Narducci, D. N., Mullen, S., Babbín, A. R., and Cziczo, D. J.: A Link between the Ice Nucleation Activity of Sea Spray Aerosol and the Biogeochemistry of Seawater, *Atmospheric Chemistry and Physics Discussions*, 2020, 1–28, <https://doi.org/10.5194/acp-2020-416>, 2020.

- 1095 Wright, T. P. and Petters, M. D.: The role of time in heterogeneous freezing nucleation, *Journal of Geophysical Research: Atmospheres*, 118, 3731–3743, <https://doi.org/10.1002/jgrd.50365>, 2013.
- Yun, Y. and Penner, J. E.: An evaluation of the potential radiative forcing and climatic impact of marine organic aerosols as heterogeneous ice nuclei, *Geophys. Res. Lett.*, 40, 4121–4126, <https://doi.org/10.1002/grl.50794>, 2013.
- Zieger, P., Väisänen, O., Corbin, J. C., Partridge, D. G., Bastelberger, S., Mousavi-Fard, M., Rosati, B., Gysel, M., Krieger, U. K., Leck, C.,
1100 Nenes, A., Riipinen, I., Virtanen, A., and Salter, M. E.: Revising the hygroscopicity of inorganic sea salt particles, *Nature Communications*, 8, 15 883, 2017.