

Responses to Reviewer 1

The manuscript “Optical properties and composition of viscous organic particles found in the Southern Great Plains” presents the observation results of HVOP during April and May 2016. The works focused to discriminate between ASOP and TB in the HVOP. The studies provide different methods to characterize the ASOP. However, my primary concern is that the authors did not make it clear the contribution of ASOP to the BrC and how important the ASOP is in the atmosphere.

The authors would like to thank Reviewer #1 for their comments, suggestions, and time as well. We have made changes to the manuscript accordingly and have indicated below where these changes were made.

To address the primary concern about the magnitude of ASOP contribution to BrC and the atmosphere, Line 23 was added to the abstract:

“one out of the three samples investigated had a significant number of ASOP while the other two samples contained TB.”

Along with Line 33:

“In addition, our results show that ASOP may only be atmospherically relevant during times when suitable emission conditions are met”

And Line 429 in the conclusion begins more discussion:

“Of the three sampling days focused on in this study, only one indicated an appreciable number of ASOP present. Because the conditions necessary for ASOP emission depend both on soil properties and precipitation characteristics, the dominant source of HVOP will often be TB due to the frequency of biomass burning instances along with the large number of particles they emit. ASOP are likely to contribute to aerosol properties, optical and otherwise, only during short time periods where the emission conditions are met.”

Major comments

1. The experimental section “a portion of one SOM sample was used to generate particles by bubbling N₂ gas through the liquid using a fritted glass bubbler and then collected on substrates to mimic the hypothesized mechanism of ASOP formation.” The authors should give the detail information about the bubbling simulation processes and parameters.

More detail added on Line 125 which now reads:

“In addition, an aliquot of 30-40 mL of one SOM sample was used to generate particles by bubbling N₂ gas (at 8 L/min) through the liquid using a fritted glass bubbler. The N₂ gas, now carrying particles generated from the bursting of bubbles at the air-water interface was directed into a cascade impactor (Sioutas Personal Cascade Impactor #225-370, SKC) where particles were collected on a pre-loaded microscopy grid.”

2. “where ASOP comprised up to 80% of the particles by number.” The impactor can only provide the mass concentration. But the ASOP fraction was obtained based on particle number. Why? What is used to determine the ASOP number?

Rephrased to emphasize how the ASOP particles were identified on Line 124:

“...to produce 300-500 nm diameter particles, where subsequent tilted SEM imaging revealed that ASOP comprised up to 80% of the particles by number.”

3. Precipitation results in the emissions of ASOP, but the AAE did not present an obvious increase during the rain events. Why?

Added some additional discussion on Line 200 which now reads:

“While ASOP emissions are expected during rain events (Joung and Buie, 2015), precipitation scavenging is also occurring. The net effect of these two competing processes likely depends on many environmental conditions and is not yet clear. During a rain event, ASOP will contribute to an increased AAE but because AAE is a bulk optical property the presence of other absorbing aerosols like black carbon or the washout of mineral dust (large particles with a high AAE) can dampen the effect that rain events have on AAE (Bergstrom et al., 2007).”

4. In Fig. 1, the AAE did not correlate with the particle number. Why?

To expand on this Line 257 now reads:

“Even though elevated HVOP numbers were identified on a number of days, Fig. 1 shows that there is no clear relationship between the particle concentration and AAE. One reason for this is the presence of other absorbing or non-absorbing aerosols which will increase the measured particle concentration while affecting the AAE differently than HVOP are expected to.”

And Line 263 now reads:

“This correlation also uses only aerosols impacted onto MOUDI stages 7 and 8, whose combined aerodynamic size range (0.18 – 0.56 μm) covers much of the ambient aerosol surface area distribution (Seinfeld and Pandis, 2006). Because optical properties like AAE will be most sensitive to changes in this size range, we exclude particles which are counted in the overall particle concentration but contribute less to bulk optical properties.”

5. “ASOP may make up a small fraction of the HVOP seen during this sampling period.”

Indicating that most of the HVOP comes from the anthropogenic sources, such as coal-fired power plant, oil refinery and natural gas power plants. My concern is that why we should focus on the ASOP, as it may have few impacts on aerosol radiative forcing and climate effects.

To address concerns about the importance of ASOP see additions to abstract and conclusion mentioned above.

More clarification was added to better define the sampling period this paragraph was referring to. Line 296 now reads:

“Although it is possible ASOP may make up a fraction of the HVOP seen during the April 28th sample, immediately east of the sampling site are a coal-fired power plant, an oil refinery, and natural gas power plants which could possibly contribute to the HVOP seen on this day.”

6. “Our results support the bubble bursting mechanism of particle generation during rainfall resulting in the ejection of soil organics into the atmosphere.” But the paper did not discuss how the ASOP ejected from soil to atmosphere during rainfall. The authors should include this parts in the paper.

The aerosol generation mechanism was expanded upon in the introduction at Line 60 which now reads:

“Briefly, as falling water droplets make contact with the porous soil surface, air from within the soil is trapped beneath the resulting water layer. Bubbles form as the droplet sinks and mixes with the soil. These bubbles then rise to the air-water interface bringing dissolved soil organics with them where a cavity forms and then ruptures producing aerosols containing compounds entrained from the soil (Joung and Buie, 2015).”

Responses to Reviewer 2

Fraund et al. conducted observations of characteristics of high viscosity organic particles (HVOP) apportioned as either airborne soil organic particles (ASOP) or tar balls (TB) during the HI-SCALE campaign. Formation mechanisms and properties of ASOP or TB have been paid much attention in recent years in the atmospheric chemistry community because of their role in climate forcing. There are few studies observing ASOP and TB in the same campaign and comparing their differences. As the authors pointed out that it is challenging to differentiate between ASOP and TB, this study is timely and I recommend the publication after the following comments can be addressed.

The authors would like to thank Reviewer #2 for their detailed comments, suggestions, and time as well as for recommending this manuscript for publication. We have made changes to the manuscript accordingly and have indicated below where these changes were made.

Major comments:

(1) The analysis of the three particular periods (April 28th, May 5th and 14th) with elevated AAE is reasonable and comprehensive. However, I am curious how these ASOP or TB went through chemical and physical changes after they were emitted? For example, the authors expected that HVOP in the May 14th samples would be present for at least 10 hours after the rain event (Line 295). Why the AAE gradually increased after the rain event and reached the peak ~10 hours later (Fig. 1)? Did ASOP need this long time to be formed or accumulated? Would chemical ageing affect the particle composition during the 10 hours and make ASOP more viscous (Adachi et al., 2019)? What is the reason AAE decreased quickly after it reached the peak? Does it indicate the life time of HVOP is relatively short? Or they were transported to areas downwind? I understand it may not be easy to clearly know the transformation of ASOP and TB, at least some discussions should be added.

Discussion on the aging of TB and ASOP was added to Line 350, which now reads:

“The chemical composition of TB has been reported in literature, with fresh TB being comprised of various biomass tar products with substantial degree of aromaticity. The nonpolar products were the most strongly associated with the wavelength dependence in absorption seen in TB and were found in greater number in fresh TB (Li et al., 2019). Photochemical oxidation in the presence of O₃ or OH radicals was shown to bleach this wavelength dependence after about 3.5 days (Sumlin et al., 2017). Photooxidation was suggested to break the network of the conjugated double bonds of the TB chromophores, resulting in more oxygenated (carbonyl substituted) products which concentrated on the surface of the TB (Hand et al., 2005; Tóth et al., 2014). ASOP are comprised, in part, of high molecular weight humic-like substances. These compounds contain multiple conjugated ring systems which likely serve as chromophores in a similar way to the tar materials in TB (Kumada, 1955). The extensive conjugated systems may be driving the enhanced C=C peak in Fig. 5 for ASOP associated spectra. Also, because humic-like substances in ASOP are substituted by many different functional groups, rather than the nonpolar components of TB, more reactive sites may be available for oxidation. This may lead to faster atmospheric processing and bleaching of ASOP wavelength dependent absorption. This aging could also serve to increase the viscosity of ASOP in the same way that it does to TB (Adachi et al., 2019).”

Extra discussion on the 10-hour delay seen in the April 14th sample was also added to Line 321:

“It is also possible that emission conditions for ASOP (soil and rainfall characteristics) were ideal somewhere along the air mass trajectory. This would have the effect of bringing in a high concentration of ASOP after the aerosol plume traveled. In addition, production of viscous SOA particles was likely taking place at the same time. The peak in AAE seen during the May 14th sampling period occurs as the sun is rising and ozone concentration rapidly increases (this also coincides with an increase in particle concentration). Although viscous SOA particles would contribute to the AAE, they are formed <100 nm in size, limiting their effect on optical properties. The peak in AAE here, and in the other cases, drops down quickly due to the particle-laden air mass moving away from the sampling site.”

(2) Line 322-325: The authors described the solubility of the non-spherical particles, which reminded me that what is the hygroscopicity of ASOP and TB? In recent years there are a lot of experiments measuring the viscosity of SOA formed from various precursors (summarized in Reid et al., 2018) at different relative humidity (RH). I see ASOP and TB are quite different from SOA but is it possible to measure the viscosity values of ASOP and TB varied with RH? Are ASOP and TB too small to measure their viscosity? As highly viscous secondary organic particles were observed frequently in ambient air (Virtanen et al., 2010 and other studies afterwards), I suggest at least add a few sentences describing the HVOP in this study is somehow different from those highly viscous secondary organic aerosol particles.

A comparison between viscous SOA and ASOP/TB is made on Line 371:

“Viscous SOA have been observed previously, under both laboratory and ambient conditions (Virtanen et al., 2010). The HVOP discussed in the current work differ from these viscous SOA by virtue of their mode of formation with both TB and ASOP being comprised of larger organic compounds. Furthermore, the particles themselves are larger. Viscous SOA are formed and observed as much smaller particles (<100 nm) than TB and ASOP (300 – 700 nm).”

A discussion of ASOP/TB hygroscopicity was also added to Line 362, now reading:

“Another factor differentiating their aging processes could be their hygroscopicity. Reports have shown TB change very little in morphology even when cycled from 0 to 100% RH, having a growth factor of ~1.09 (Adachi and Buseck, 2011; Semeniuk et al., 2007). However, because ASOP are formed via dissolved SOM that is ejected during precipitation, they are expected to be more hygroscopic. Indeed, Wang (2016) showed the results of RH cycling in supplemental figures and found a growth factor of ~1.15 at 85% RH along with droplet activation at 98% RH.”

Minor comments:

(1) Line 38: delete “a” in “strong a climate : : :”. Deleted “a”

Line 40: delete “aerosol” in “organic aerosol carbon”. Deleted “aerosol”

Line 78: should be Springston 2016 not 2011. Reference fixed

(2) Line 119: Why the SOM aquatic samples were collected on May 17th? Why not choose 14th, same day that HVOP were sampled? Added to Line 122: “..., prior to offline analysis of microscopy samples”

(3) Line 186: The threshold (10 mm/hr) defining the rain events is different from 5 mm/hr described in Line 100. All threshold values changed to 10 mm/hr

(4) Line 207: Is it certain that an elevated AAE suggests the presence of spherical HVOP? I think better replace “can” to “may”. Changed “can” to “may”

(5) Line 217: There are two Moffet et al. together; please delete one. Also Line 234 for Veghte et al and Line 250 for Wang et al.

Reworded Line 227 to read “Following the procedure described in Moffet (2010) for chemical speciation maps, each... or as a region with high C=C bonding (Moffet et al., 2010).”

Repeated instances of Veghte et al and Wang et al were removed.

(6) Table 1: what does IOP1 indicate?

Table 1 caption now reads: “Ambient sampling information...”

(7) Supplement: there are two “Figure S1”.

The second supplemental figure was renamed “Figure S2”

(8) Line 256: Cite Wang et al. 2016 after “lab-generated SOA particles”.

Inserted Wang et al., 2016 citation

(9) Figure 1: It seems some grey vertical bars indicating BrC appearance do not have elevated AAE > 1.7, for instance, the night of May 1st. Why it is included in the “BrC events”?

Modified Figure 1 caption to read “...Grey vertical bars indicate sampling periods where microscopy samples were collected... Arrows indicate the three sampling periods investigated in the current work where BrC particles were expected.”

References:

Adachi, K., Sedlacek, A. J., Kleinman, L., Springston, S. R., Wang, J., Chand, D., Hubbe, J. M., Shilling, J. E., Onasch, T. B., Kinase, T., Sakata, K., Takahashi, Y. and Buseck, P. R.: Spherical tarball particles form through rapid chemical and physical changes of organic matter in biomass-burning smoke, *Proceedings of the National Academy of Sciences*, 116, 19336-19341, 10.1073/pnas.1900129116, 2019.

Reid, J. P., Bertram, A. K., Topping, D. O., Laskin, A., Martin, S. T., Petters, M. D., Pope, F. D. and Rovelli, G.: The viscosity of atmospherically relevant organic particles, *Nat. Commun.*, 9, 956, 10.1038/s41467-018-03027-z, 2018.

Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., Yli-Pirilä, P., Leskinen, J., Mäkelä, J. M., Holopainen, J. K., Pöschl, U. and Kulmala, M.: An amorphous solid state of biogenic secondary organic aerosol particles, *Nature*, 467, 824-827, <https://doi.org/10.1038/nature09455>, 2010.

Müller, L., Reinnig, M.-C., Warnke, J., and Hoffmann, Th.: Unambiguous identification of esters as oligomers in secondary organic aerosol formed from cyclohexene and cyclohexene/_-pinene ozonolysis, *Atmos. Chem. Phys.*, 8, 1423–1433, <https://doi.org/10.5194/acp-8-1423-2008>, 2008.

Optical properties and composition of viscous organic particles found in the Southern Great Plains

Matthew Fraund¹, Daniel J. Bonanno¹, Swarup China², Don Q. Pham¹, Daniel Veghte³, Johannes Weis^{4,5,a}, Gourihar Kulkarni², Ken Teske⁶, Mary K. Gilles⁴, Alexander Laskin^{2,b}, Ryan C. Moffet^{1,c}

- 5 1. Department of Chemistry, University of the Pacific, Stockton, California, 95204, USA
2. Environmental and Molecular Sciences Laboratory, Pacific Northwest National Laboratory (PNNL), Richland, Washington 99352, USA
3. Center for Electron Microscopy and Analysis, The Ohio State University, Columbus, Ohio, 43212, USA
4. Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California, 94720, USA
10 5. Department of Chemistry, University of California, Berkeley, California, 94720, USA
6. Atmospheric Radiation Monitoring (Southern Great Plains Climate Research Facility), Billings, Oklahoma, 74630 USA
a. Current address: Physikalisches Institut, Universität Würzburg, Am Hubland, 97074 Würzburg, Germany
b. Current address: Chemistry Department, Purdue University, West Lafayette, Indiana, 47907, USA
15 c. Current address: Sonoma Technology, Petaluma, California, 94954, USA

Correspondence to: Matthew Fraund (m_fraud@u.pacific.edu)

Abstract. Atmospheric high viscosity organic particles (HVOP) were observed in samples of ambient aerosols collected on April and May 2016 in the Southern Great Plains of the United States. These particles were apportioned as either airborne soil organic particles (ASOP) or tar balls (TB) from biomass burning based on spectro-microscopic imaging and assessment of meteorological records of smoke and precipitation data. Regardless of their apportionment, the number fractions of HVOP were positively correlated ($R^2 = 0.85$) with increased values of Ångström absorption exponent (AAE) measured *in-situ* for ambient aerosol at the site. Extending this correlation to 100% HVOP yields an AAE of 2.6, similar to previous literature reports of the class of light absorbing organic particles known as brown carbon (BrC). One out of the three samples investigated had a significant number of ASOP while the other two samples contained TB. Although there are chemical similarities between ASOP and TB, they can be distinguished based on composition inferred from near edge absorption X-ray fine structure (NEXAFS) spectroscopy. ASOP were distinguished from TB based on their average $-\text{COOH}/\text{C}=\text{C}$ and $-\text{COOH}/\text{COH}$ peak ratios, with ASOP having lower ratios. NEXAFS spectra of filtered soil organic brine particles nebulized from field samples of standing water deposited after rain were consistent with ASOP when laboratory particles were generated by bubble bursting at the air-organic brine interface. However, particles generated by nebulizing the bulk volume of soil organic brine had particle composition different from ASOP. These observations are consistent with the raindrop generation mechanism responsible for ASOP emissions in the area of study. In contrast, nebulized samples carry with them higher fractions soil inorganics dissolved in the bulk volume of soil brine that are not aerosolized by the raindrop mechanism. Our results support the bubble bursting mechanism of particle generation during rainfall resulting in the ejection of soil organics into the atmosphere. In addition, our results show that ASOP may only be atmospherically relevant during times when suitable emission conditions are met.

20
25
30

35 1 Introduction:

Regional and global atmospheric transport models are commonly used to predict the impact of aerosols on radiative forcing (Feng et al., 2013). The efficacy of these models relies on estimates of the types, number concentrations, spatial distribution, and emission sources of aerosols. One challenge of continued scientific discussion is how (and to what extent) industry and other anthropogenic activities contribute to climate forcing. Emissions of soot, one of the most well-studied anthropogenic aerosols emitted by fossil fuel combustion, have been shown to have a strong ~~a~~-climate warming factor comparable to carbon dioxide (Bond et al., 2013). However, soot is not the only light absorbing carbon containing aerosol of concern. Less absorbing, but often more abundant light absorbing organic ~~aerosol~~-carbon aerosol, known as brown carbon (BrC), has been the subject of increased investigation, with studies showing that BrC may account for a substantial fraction of the total aerosol radiative forcing (Feng et al., 2013;Zhang et al., 2017). These particles exhibit a wavelength dependent light absorption, absorbing stronger at shorter wavelengths, giving them a brown appearance. Ongoing research on the role of BrC in global radiative forcing suggests it is insufficiently represented in models (Feng et al., 2013;Laskin et al., 2015;Bond and Bergstrom, 2006).

BrC particles are defined in very general terms of their chemical composition and optical properties (Bond et al., 2013;Laskin et al., 2015). BrC refers to a broad category of organic particles comprised of many different chromophores originating from a variety of sources (Andreae and Gelencsér, 2006). Sources of BrC include, but are not limited to: biomass burning (Rizzo et al., 2013;Laskin et al., 2015), biogenic fungi, humic like substances (Andreae and Gelencsér, 2006;Wang et al., 2016), and secondary organics (Laskin et al., 2015). One example of BrC particles are tar balls (TB) which are most commonly found in biomass burning emissions downwind of smoldering fires (Chakrabarty et al., 2010). TB of 50-300 nm in size have a characteristic spherical morphology observed in many studies, indicative of their highly viscous or “glassy” phase state. Recently, larger spherical organic particles (~500 nm) of a different BrC type (Veghte et al., 2017) have been also observed in rural Oklahoma, with their composition consistent with soil organic matter (SOM). These newly discovered BrC particles have been termed airborne soil organic particles (ASOP) (Wang et al., 2016). The study reported formation of highly viscous submicron particles following the impaction of rain droplets on the soil surfaces which eject ASOP into the ambient air (Wang et al., 2016). Mechanism of plausible ASOP emissions was further corroborated by laboratory experiments with water droplets falling onto surfaces of soil proxies. Briefly, As falling water droplets make contact with the porous soil surface, air from within the soil is trapped beneath the rapidly expanding droplet resulting water layer. Bubbles form as the droplet sinks and mixes with the soil. These bubbles then rise to the surface of the droplet-air-water interface bringing dissolved soil organics with them where a cavity forms and then ruptures producing film and jet droplets aerosols containing compounds entrained from the soil (Joung and Buie, 2015).

65 A defining characteristic of BrC particles is the spectral dependence of light absorption in the visible wavelengths. One way of quantifying this dependence is to measure the absorption coefficient (σ_{ap}) as a function of wavelength, to calculate the absorption Ångström exponent (AAE) (Backman et al., 2014). The AAE value is essentially the slope of a log-log plot of

absorption coefficient and wavelength, with higher values corresponding to enhanced absorption in shorter wavelengths. Because soot absorbs nearly equally across the entire visible spectrum, it has an AAE value of approximately 1. In contrast, reported AAE values for BrC are substantially higher, typically in a range of 2.5-6 (Rizzo et al., 2013; Lawless et al., 2004; Hoffer et al., 2006) but could be as high as 9-11 (Andreae and Gelencsér, 2006; Lin et al., 2017). An AAE value of 2.5 has been used as a lower limit to attribute absorption, at least in part, to BrC (Lack and Langridge, 2013). In addition to the AAE, absorption due to BrC has also been investigated through calculation of the complex refractive index (Veghte et al., 2017).

To better characterize highly viscous organic particles (HVOP) appearing as solid spheres, samples of ambient particles were collected at the atmospheric radiation measurement (ARM) facility in Lamont, Oklahoma located in the southern great plains (SGP) as part of the Holistic Interactions of Shallow Clouds, Aerosols, and Land-Ecosystems (HI-SCALE) field campaign (Fast et al., 2019). To compare the spectroscopic signatures of atmospheric particles with SOM typical for the area of study, aquatic samples of the surface layer of muddy puddles were also collected around the sampling site. These aquatic samples were then filtered, nebulized, and the resulting particles were impacted onto microscopy substrates. The purpose of collecting these samples was to use them to compare particle morphology and composition when AAE was high, as indicated by online measurements performed at the site (Springston et al., 2016). The particle samples were analyzed with both scanning electron microscopy (SEM) and scanning transmission X-ray microscopy coupled with near-edge X-ray absorption fine structure (STXM-NEXAFS) spectroscopy. SEM images were taken at a 75° tilted angle with respect to the surface normal to identify HVOP suspected to be either ASOP or TB. STXM was then used to obtain chemical images of particles at the C K-edge spectral range to distinguish ASOP and TB based on their carbon NEXAFS spectra (Wang et al., 2016).

The present work evaluates the appearance of ASOP following rain events and determines their particle-specific spectroscopic characteristics that would enable distinguishing them from biomass burning TB, SOA, or other anthropogenic sources (Parworth et al., 2015; Sheridan et al., 2001). Characterizing the properties and emission sources of unaccounted BrC particles like ASOP is vital to ensure the proper representation of BrC in local and regional climate models. Multiple experimental records (both real-time and spectro-microscopic) were used here to assess the underreported ASOP. Using these techniques in tandem with longstanding atmospheric measurements will also help in advancing our knowledge of how ASOPs fit into the broader class of BrC and how ASOP may affect radiative forcing by aerosols.

2 Experimental:

95 2.1 Sample Collection

Samples of atmospheric particles were collected at the ARM SGP field site located in north central Oklahoma [36° 36' 18.0" N, 97° 29' 6.0" W], at an altitude of 320 meters (Ferrare et al., 2006). Sampling was performed before forecasted and after observed rain events during the rainy season from April 26th through May 17th, during day and night periods separately. Within this timeframe, samples were collected on 17 days with each sampling period beginning between 8:00 to

100 21:00 local time. The total sampling time was approximately eight hours (unless interrupted by a major rain event) with
samplers operated using a 30/30 min on/ off duty cycle. Similarly, 4 nighttime samples were collected between the hours of
18:00 and 6:00 local time with a duty cycle of 60/30 min on/off. Rain events interrupted two sampling days (May 8th and May
16th) in which no sampling took place. The primary objective was to observe particle types during sunny days following major
rain events, where a threshold precipitation rate of 5-10 mm/hr was used to define a rain event, when enhanced ASOP
105 concentration would be likely. Note that ASOP may be present during some periods without a recent local rain event due to
transport from elsewhere.

Particles were collected by impaction using a Micro-Orifice Uniform Deposit Impactor (MOUDI, MSP 100) attached
to a rotating motor that rotates the eight impaction surfaces (known as stages) to facilitate uniform particle deposition. The
impactor was connected to a ¾ max horsepower (0.56 kW) vacuum pump (General Electric Motors & Industrial Systems, 10
110 PSI rating). With a 30 L/min sampling flow rate, the micro-orifice nozzles reduce jet velocity, pressure drop, particle bounce
and re-entrainment. This MOUDI was connected to a mesh covered sampling inlet which was tilted downward, avoiding
unwanted collection of descending debris, insects, and other sources of contamination, and was positioned ~6 meters above
the ground.

Samples for analysis were selected from two stages with the following particle size cutoff ranges: stage 7 (0.32 to
115 0.56 µm) and stage 8 (0.18 to 0.32 µm). These stages were chosen to cover part of the size ranges for TB (50 to 300 nm) (Tóth
et al., 2014) and ASOP (300 to 800 nm) (Wang et al., 2016) while also constraining the number of samples to analyze.
Substrates of Si₃N₄ film supported by a silicon wafer (0.5 X 0.5 mm² Si₃N₄ window, 100 nm membrane thickness, 5 X 5 mm²
Si frame; Silson, Inc.) and filmed TEM copper grids (Carbon type B film, Copper 400 mesh grids; Ted Pella, Inc.) were used
as impaction substrates.

120 In addition to impaction samples of ambient particles, aquatic samples containing SOM brine were collected in 50-
200 mL aliquots via syringes from mud puddles surrounding the SGP field site. This was performed to evaluate a connection
between organics from terrestrial aquatic samples, the hypothesized source of ASOP, and ambient ASOP. Four SOM aquatic
samples were collected on May 17th 2016, prior to offline analysis of microscopy samples. The obtained samples were then
nebulized in laboratory experiments using a Collison nebulizer (3 jet MRE, CH Technologies USA) and collected on stage 8
125 of a 10-stage impactor (110-R, MSP, Inc.) to produce 300-500 nm diameter particles, where subsequent tilted SEM imaging
revealed that ASOP comprised up to 80% of the particles by number. In addition, an aliquot of 30-40 mL portion of one SOM
sample was used to generate particles by bubbling N₂ gas (at 8 L/min)-through the liquid using a fritted glass bubbler. The N₂
gas, now carrying particles generated from the bursting of bubbles at the air-water interface, was directed into a cascade
impactor (Sioutas Personal Cascade Impactor #225-370, SKC) where particles were collected on a pre-loaded microscopy grid
130 and then collected on substrates to mimic the hypothesized mechanism of ASOP formation. Stage D of the Sioutas impactor
was used (D₅₀ 0.25 µm) to best correspond with the size ranges of MOUDI samples taken

2.2 STXM Measurements and Image Processing.

The STXM instruments (beamline 5.3.2.2 ALS, Berkeley, CA, USA, and the SM1 beamline CLS, Saskatoon, SK, Canada) used in this work are located in the Advanced Light Source (ALS) at the Lawrence Berkley National Laboratory and at the Canadian Light Source (CLS) at the university of Saskatchewan (Kilcoyne et al., 2003;Regier et al., 2007). Briefly, monochromatic soft X-rays are focused down to a spot size ranging from 20 to 40 nm in diameter. The sample is raster scanned after a region of sufficient particle concentration is found and individual images are captured at selected photon energies. Maps were collected in addition to spectra, which are images consisting of 8 energies around the elemental absorption k-edges, four energies corresponding to C absorption at 278.0 eV, 285.4 eV, 288.6 eV, and 320.0 eV, two for N absorption at 398.0 eV and 430.0 eV, and 2 for O absorption at 525.0 eV and 550.0 eV. These energies are used to identify and characterize the basic chemical composition maps of particles on a pixel-by-pixel basis (Fraund et al., 2017). The two C energies at 285.4 and 288.6 eV are used for the identification of soot or elemental carbon (EC). The absorption peak at 285.4 eV occurs due to the excitation of the $C\ 1s \rightarrow \pi^*_{C=C}$ transition (* indicating the excited state) which is indicative of sp^2 hybridized carbon-carbon bonds ($C=C$). This excitation of sp^2 hybridized carbon is prominent for soot (Bond and Bergstrom, 2006). To identify soot, the intensity of the sp^2 peak (relative to the pre-edge at 278 eV after normalization to 320 eV) must be equal to or exceed 35% of that of highly ordered pyrolytic graphite (Moffet et al., 2010). Absorption at 288.6 eV is representative of the $C\ 1s \rightarrow \pi^*_{R(C^*=O)OH}$ transition characteristic of carboxylic acid groups (COOH) that are very common in atmospheric organic carbon. These photon energies aid in differentiating individual particles based on the molecular speciation of carbon, making this method convenient for analysis of the field collected particles (Moffet et al., 2010). Spatial displacement between images within a stack does occur and, whenever needed, it has been corrected for by utilizing the image registration algorithm developed by Guizar-Sicarios⁷ (Fraund et al., 2017;Guizar-Sicairos et al., 2008).

2.3 SEM Measurements

The SEM analysis of particle samples was performed at the Environmental Molecular Sciences Laboratory (EMSL) located at Pacific Northwest National Laboratory (PNNL). Particles were imaged using a computer-controlled scanning electron microscope (FEI, Quanta 3D FEG, Hillsboro, AL, USA). SEM images were initially taken orthogonally to the substrate until a particle laden region on the substrate was identified. The substrate mount was then tilted by 75° in order to identify spherical particles. Based on the tilted images, particles which had an aspect ratio (height divided by width) greater than 0.8 were identified as spherical HVOP. The SEM images were categorized based on aspect ratio to determine the fraction of HVOP (high aspect ratio) compared to flat, non-spherical particles (low aspect ratio) for ensembles of approximately 400 total particles. This counting was performed manually for both ambient and lab-generated samples.

2.4 PSAP Measurements.

The Particle Soot Absorption Photometer (3- λ PSAP) instrument measures light transmission through aerosol filter samples at three wavelengths: red (660 nm), green (522 nm) and blue (470 nm) (Springston, 2018). PSAP measurement taken at the SGP ARM facility is a standard data product (ARM, 2019). Equation (1) shows how absorption coefficients σ_{ap} are calculated from raw PSAP data using spot size (A), sample volume (V), and average filter transmittances for incident (I_o) and transmitted (e) light through particle laden filter.

$$\sigma_{ap} = \frac{A}{V} \ln \left[\frac{I_o}{I} \right] \quad (1)$$

Implementing corrections specified in Bond *et al.* (1999), as shown in Eq. (2), minimizes noise present in the PSAP data, which is a result of inherent unit-to-unit variability in field instrumentation. Additionally, these corrections mitigate systematic error from filter loading. The absorption reported by the PSAP instrument (σ_{PSAP}) includes an inherent calibration for a given measurement period which monitors the change in transmission by using the previous sample as a blank.

$$\sigma_{PSAP} = \frac{\sigma_{ap}}{2(0.5398\tau + 0.355)} \quad (2)$$

The raw absorption coefficient of a sample (σ_{ap}) at a given time is normalized by filter transmission (τ), which is reset after the installation of a new filter ($\tau = 1$ for an unloaded filter) (Springston, 2018). The corrected absorption coefficients were used to calculate the *AAE* values. Backman *et al.* (2014) introduced the following equation relating Ångström Exponent (*AE*) to the wavelength (λ) dependent extinction (Backman et al., 2014):

$$\ln [\sigma_E(\lambda)] = -AE \ln[\lambda] + C \quad (3)$$

here, σ_E refers to the extinction coefficient at wavelength λ . While the *AE* represents the spectral dependence of combined scattering and absorption, the *AAE* is specific to absorption and is obtained by substituting σ_{ap} for σ_E (Backman et al., 2014). By taking Eq. (3) at a given wavelength and subtracting Eq. (1) at another wavelength, the constant *C* can be removed, resulting in a more practical equation:

$$AAE = \frac{-\ln \left[\frac{\sigma_{ap}(\lambda_1)}{\sigma_{ap}(\lambda_2)} \right]}{\ln \left[\frac{\lambda_1}{\lambda_2} \right]} \quad (4)$$

Using Eq. (4), PSAP data was used to calculate the *AAE* of ambient aerosols. Appearance of BrC was then inferred from PSAP time records based on elevated values of *AAE* > 1.7 (Kirchstetter et al., 2004).

3.1 Optical Properties of ~~Spherical Particles~~High Viscosity Organic Particles (SP)

Spherical HVOP have similar spectral characteristics to BrC regardless of their origin. Both TB and ASOP show characteristic increased absorption for shorter wavelengths of visible light compared to longer wavelengths (Alexander et al., 2008; Veghte et al., 2017). During periods when they are prevalent, overall aerosol optical properties should start to resemble those of BrC. To investigate the presence of BrC with this wavelength dependence, PSAP data was utilized to determine time records of AAE as shown in Fig. 1 alongside the corrected absorption coefficients. Time records of CO mixing ratios and particle number concentrations are also shown in Fig. 1 to provide further information on the composition of air mass. Figure 1 shows how both red and blue absorption coefficients and AAE change over the course of this study. Rain events were defined as periods of time where rainfall exceeded 10 mm/hr. If two rain events were observed less than 30 minutes apart, they were considered one event. The vertical gray bars show sampling periods when aerosol with BrC properties was detected and which particle samples were imaged with STXM and SEM.

Both the absorption coefficient and AAE time series data were collected with minute time resolution; the data has been averaged over 30-minute time windows to emphasize longer term data trends instead of short-term fluctuations. The AAE sometimes shows an increase after rain events, though it is not consistent; because aerosol production from rain is more complex than rainfall amounts (depending on droplet size and impact velocity as well as soil characteristics) it is difficult to see a direct correlation between rain and AAE or particle concentration. While ASOP emissions are expected during rain events, precipitation scavenging is also occurring (Joung and Buie, 2015). The net effect of these two competing processes likely depends on many environmental conditions and is not yet clear. During a rain event, ASOP will contribute to an increased AAE but because AAE is a bulk optical property the presence of other absorbing aerosols like black carbon or the washout of mineral dust (large particles with a high AAE) can dampen the effect that rain events have on AAE (Bergstrom et al., 2007).

From this time record, a few samples stand out: the night of April 28th, May 5th, and May 14th as these samples had elevated AAE >1.7. On May 5th and April 28th measured AAE values were above 2, warranting analysis of particle samples (Lack and Langridge, 2013). While the May 14th sample doesn't show particularly high AAE values, compared to the entire time series, it was collected after a heavy rain storm passed through the area. Also, the particle concentration immediately following the rain event on May 14th shows a significant level of precipitation scavenging, reducing the number of background particles ~~other than ASOP~~ present during sampling, concentrating any ASOP produced (Hegg et al., 2011). Of note, the lower the σ_{ap} drops, due to rain or otherwise, the more pronounced the effect PSAP noise has on the calculated AAE. In addition to having high AAE values, both the April 28th and May 5th periods showed elevated particle and CO concentrations indicating plausible anthropogenic activity or biomass burning plumes. Figure S1 shows the time series of ~~four~~ trace gases (N₂O, CO, O₃ and SO₂) and particle concentration which can be used to identify anthropogenically influenced air plumes. In addition to elevated CO levels, the April 28th and May 5th samples show elevated SO₂ and N₂O concentrations while some of the highest

O₃ concentrations seen over the sampling periods were observed during May 5th. In contrast, the May 14th period exhibits low or fairly average concentrations of these trace gasses.

3.2 Chemical Imaging of High Viscosity Organic Particles

220 Bulk optical properties, like an elevated AAE, ~~can~~may suggest the presence of spherical HVOP, therefore these measurements were used to select samples for detailed chemical imaging of particles. First, tilted SEM images were taken and HVOP fractions were observed in individual samples, ranging from 5% to near 70% by number. Figure 2 shows representative microscopy and spectro-microscopy images for three days where HVOP fractions were high. The top row shows the tilted SEM images used to identify HVOP. Magenta arrows point to a few identified HVOP to highlight how much they stick out
225 above the substrate compared to the others. Of note, the SEM images also show presence of what looks like fractal soot particles in the April 28th and May 5th samples.

Second, the same samples were later imaged by the STXM spectro-microscopy. The middle and bottom row of images shown in Fig. 2 are STXM chemical speciation maps and total carbon absorbance (TCA) maps, respectively. Following the procedure described in Moffet (2010) for ~~In the~~ chemical speciation maps, each pixel is assigned as either inorganic dominant,
230 organic dominant, or as a region with high C=C bonding ~~following procedure describe in Moffet, et al. 2010~~ (Moffet et al., 2010). The TCA maps indicate the thickness of each of the particles (as calculated using previously published thickness equations (Fraund et al., 2019; O'Brien et al., 2015)) normalized by the individual area equivalent diameter. Values close to 0 represent flat particles, while values closer to 1 represent taller, nearly spherical particles. As was suggested in the SEM images, soot is present in both the April 28th and May 5th samples, while the May 14th sample has only organic and inorganic particles.
235 In the April 28th and May 5th samples, circular soot-containing particles are associated with the highest TCA. In contrast, the May 14th sample contained high TCA spherical particles comprised of organic dominant material only.

Sample collection information from the 7 samples where the fractions of HVOP were calculated is presented in Table 1 below. The highest HVOP fraction was observed with the samples taken on May 5th. The prevalence of these particles can be seen in Fig. 2 in the top row. Elevated fractions of HVOP were also found for the April 28th night sample taken at 18:30
240 and for the May 14th sample. While the April 28th and May 5th sample showed elevated HVOP fractions for both stage 7 and 8 samples, the May 14th sample is unique in that a higher HVOP fraction was only found for the smaller stage. In addition, the April 28th and the May 5th sample both have elevated particle concentrations and CO mixing ratios, suggesting more polluted conditions possibly due to anthropogenic activity or biomass burning events. In contrast, the time period corresponding to May 14th sample shows the lowest particle concentration and CO mixing ratio. Also of note is that the April 28th and May 5th samples
245 were both taken long after the last rain, with May 5th being taken multiple days afterwards whereas the May 14th sample was taken 10 hours after the last rain.

Optical properties of individual HVOP from this same data set have been investigated in our previous work, ~~Veghte, et al. in 2017~~ (Veghte et al., 2017). There, the complex refractive index from 200 to 1200 nm was calculated for HVOP found in the April 28th sample using electron energy loss spectroscopy (EELS). The imaginary part (k) of the refractive index is

250 related to light absorption and can be related to the absorption coefficient using $\sigma_{ap} = 4\pi k/\lambda$ (Jennings et al., 1979). Absorption coefficients over the 200 – 1200 nm wavelength range were calculated from ~~the a~~ published imaginary refractive index plot (Veghte et al., 2017). From this plot published in Veghte, ~~et al. (2103)~~, the AAE was calculated using ~~the~~ σ_{ap} values for 660 and 470 nm and a resulted in a value of 1.41. This is close to the value calculated in the present work (1.42, Table 1) for the April 28th sample (a difference of only 0.01) showing that the two methods agree, at least for one sample.

255 **Table 1. Ambient sampling informationes collected during IOPI. Stage 8 values in parentheses where available.**

Start Date (CDT) ¹	MOUDI Stage	Duration (hr)	HVOP (%)	AAE (Red/Blue) ²	Hours Since Last Rain ³	Particle Conc. (cm ⁻³) ⁴	CO (ppb) ⁵	# Particles Imaged SEM STXM		HVOP Source
26 Apr 14:00	7	1.5	<5	0.808	>72	2100	150	450	0	<u>Low HVOP</u>
28 Apr 9:45	7	5	3	1.20	30	4100	140	200	0	<u>Low HVOP</u>
28 Apr 18:30	7 (8)	10	23 (25)	1.42	39	10300	160	900	28	<u>Biomass Burning, TB</u>
1 May 11:30	7 (8)	4	12 (15)	1.23	45	1000	130	500	0	<u>Low HVOP</u>
2 May 20:00	7	10	13	1.15	80	1900	130	900	185	<u>Low HVOP</u>
5 May 8:00	7 (8)	13	70 (60)	2.01	140	6000	170	1000	32 (214)	<u>Biomass Burning, TB</u>
14 May 11:00	7 (8)	5	10 (35)	1.29	10	650	120	300	50 (86)	<u>Biogenic, ASOP</u>

¹Central Daylight Time (UTC -5) ²Particle Soot Absorption Photometer (PSAP) ³Video Disdrometer ⁴Condensation Particle Counter (CPC) ⁵ARM/Aerosol Observatory System (AOS)

260 Even though elevated HVOP numbers were identified on a number of days, Fig. 1 shows that there is no clear relationship between the particle concentration and AAE. One reason for this is the presence of other absorbing or non-absorbing aerosols which will increase the measured particle concentration while affecting the AAE differently than HVOP are expected to. To address this, From Table 1 a correlation plot was made (using values found in Table 1) between the AAE values and the HVOP fractions and a strong correlation was found ($R^2 = 0.85$) as seen in Fig. S2. Extrapolating this linear correlation to 100% HVOP yields an AAE value of 2.6, consistent with previously reported AAE values for BrC (Lack and Langridge, 2013). This not only suggests that the HVOP found here are BrC, but that they warrant consideration by models due to their measureable effect on bulk aerosol-radiation interactions as they can occupy a significant fractions of the aerosol fine mode. This correlation also uses only aerosols impacted onto MOUDI stages 7 and 8, whose combined aerodynamic size range (0.18 – 0.56 μm) covers much of the ambient aerosol surface area distribution (Seinfeld and Pandis, 2006). Because optical properties like AAE will be most sensitive to changes in this size range, we exclude particles which are counted in the overall particle concentration but contribute less to bulk optical properties. This not only suggests that the HVOP found here are BrC, but that they warrant consideration by models due to their measureable effect on bulk aerosol-radiation interactions as they can occupy a significant fractions of the aerosol fine mode.

270 The appearance of viscous HVOP at the SGP site has been reported ~~earlier by Wang et al. in 2016~~ previously (Wang et al., 2016). There, they showed that the viscous HVOP at SGP had an elevated total carbon absorption (TCA, defined by the pre-edge OD subtracted from the post-edge OD) compared to other carbonaceous particles of similar sizes. High TCA values indicate particles that were not deformed upon impaction suggesting a high viscosity. Figure 3 shows TCA values of

individual particles plotted against circular equivalent diameter (CED) for four samples reported in this work. The April 28th, May 5th, and May 14th samples all have elevated TCA whereas the May 2nd sample shows lower carbon absorption, in line with the TCA values characteristic of lab-generated SOA particles (Wang et al., 2016). Note that the May 5th sample, where the highest HVOP fraction was identified, has the smallest particles with TCA values above the ambient organic particle regions. Contrast this with the May 2nd sample, which shows very few particles with high TCA values and a correspondingly low HVOP fraction.

So far, the above analysis applies to a general class of HVOP, which can include both ASOP and TB particles. Additional considerations are necessary, however, before any conclusions are tied to ASOP exclusively. The two samples, with the highest HVOP fractions and AAE values, were those collected more than 39 hours after a rain event. Because emission of ASOP is associated with bursting of bubbles at flooded soil surfaces after rain, the HVOP found in the April 28th and May 5th samples are likely not locally emitted ASOP, while ASOP might be present in the sample of May 14th.

To investigate the nature and source of HVOP and determine which can be confidently classified as ASOP, smoke and fire from biomass burning sources (NOAA:OSPO, 2019) and precipitation data (NWS, 2019), were used along with the calculations of air backward trajectories using a hybrid single particle Lagrangian integrated trajectory model (HYSPLIT) (Stein et al., 2015; Rolph et al., 2017). This data for the events when the three samples had elevated HVOP fractions are shown in Fig. 4 below: Apr 28th, May 5th, and May 14th. Additional information about the HYSPLIT trajectory conditions, as well as trajectories calculated from multiple starting altitudes, is available in Fig. S3.

April 28th had a moderate fraction of HVOP ($25 \pm 0.6\%$) along with the second highest AAE over the sampling periods studied here. This sample was also taken about 39 hours after the last rain, which makes ASOP less likely to be found. Figure 4 shows that the corresponding air mass trajectories passed over a few burning fires ~~and while. Smoke-smoke~~ was present in some of the fires surrounding the sampling site, ~~but the~~ air mass ~~trajectories~~ did not pass through these regions. Precipitation data shows rainfall in some of the surrounding states but none close to the sampling site. The storm surrounding Oklahoma may have produced some ASOP which could have been transported to the sampling site, although elevated particle concentration and slightly enhanced CO, SO₄, and N₂O concentration at the time of sampling suggested biomass burning as more ~~likely-prominent~~ emission source (Koppmann et al., 2005). ~~Although it is possible~~ ASOP may make up a ~~small~~ fraction of the HVOP seen during ~~this sampling period~~ ~~the April 28th sample, however. Immediately-immediately~~ east of the sampling site are a coal-fired power plant, an oil refinery, and natural gas power plants ~~which could possibly contribute to the HVOP seen on this day. Furthermore, Aas~~ seen in Fig. 4 (as well as in more detail in Fig. S3), the April 28th sample has an initial eastward direction.

May 5th has the highest HVOP fraction and the highest average AAE but it also had been days (140 hours) since the last rain event, making ASOP unlikely. There were many fires surrounding the sampling site compared to the other sampling periods and the back trajectories show air masses passing directly over some of these fires suggesting the presence of associated smoke emissions. Precipitation data shows that no rainfall was observed anywhere near the sampling site. Figure 1 also shows that this sampling date coincided with a slight particle concentration enhancement and the highest CO mixing ratios observed

310 over this period of the field campaign. Figure S1 also shows enhancement in O₃ and SO₂ concentrations. Because TB are found within smoke plumes, the high HVOP fraction observed in the April 28th and May 5th samples may be primarily (or exclusively) due to TB (Pósfai et al., 2004). The presence of combustion byproducts like TB and soot is supported by the STXM carbon speciation maps shown in Fig. 2, where particles with elevated C=C bonding constituents are often attributed to soot.

315 The last sample date shown (May 14th) has regions of smoke away from the sampling site with backward trajectories heading from just outside the smoke filled region. However, because a rain event was recorded 10 hours prior to sampling where significant precipitation scavenging was observed (see the particle concentration decrease in Fig. 1), no influence from biomass burning was observed in this sample. The precipitation map shows that precipitation was observed over the sampling site as well as many of the surrounding areas (Radke et al., 1980). Because the microscopy samples were taken shortly after it had rained, the 35 ± 1.3% fraction of HVOP observed at that time may likely be related to ASOP, contributing to the spike
320 in AAE seen during this sampling period.

With the HVOP observed in the microscopy images being around 0.6 µm in diameter, it is expected that they would be aloft and present for the sampling period for at least 10 hours after the rain event (Williams et al., 2002). Any particles of this size travelling from further away, such as ASOP produced elsewhere, through some of the surrounding storm are less likely to be scavenged due to the lowered precipitation scavenging efficiency (Greenfield scavenging gap) at approximately
325 0.1 µm (Radke et al., 1980). This could have effectively concentrated ASOP particles as the air mass travelled through the surrounding storm. It is also possible that emission conditions for ASOP (soil and rainfall characteristics) were ideal somewhere along the airmass trajectory. This would have the effect of bringing in a high concentration of ASOP after the aerosol plume traveled. In addition, production of viscous SOA particles was likely taking place at the same time (Virtanen et al., 2010). The peak in AAE seen during the May 14th sampling period occurs as the sun is rising and ozone concentration rapidly increases (this also coincides with an increase in particle concentration). Although viscous SOA particles would contribute to the AAE, they are formed <100 nm in size, limiting their effect on optical properties. The peak in AAE here, and in the other cases, drops down quickly due to the particle-laden airmass moving away from the sampling site.
330

The influence of smoke shown in Fig. 4 may account for the enhancement of HVOP fractions without rainfall in the April 28th and May 5th samples, likely due to TB. The carbon STXM/NEXAFS spectra of TB have been recorded previously
335 and its characteristic features are shown in Fig. 5 (Tivanski et al., 2007). The same figure compares the STXM/NEXAFS spectra for both ambient particles collected during this study and lab generated ASOP proxies. The right panel of Fig. 5 also includes three characteristic spectra of HVOP particles from the April 28th, May 5th and May 14th samples. Even though May 5th and April 28th had the highest AAE values and the highest HVOP fractions, many hours since the last rain along with the presence of smoke suggest they might be TB, consistent with their NEXAFS spectral features. Three apparent peaks are
340 common for these spectra: the C=C peak at 285.3 eV, the COH peak at 286.7 eV, and the R-(C=O)OH peak at 288.6 eV, all of which are present in the previously reported TB spectra (Tivanski et al., 2007). May 5th spectra also show a small feature around 289.5 eV which is present (and more prominent) in the TB spectra, a peak which is associated with alkyl carbon bonded

to oxygen, often alcohols. This similarity reinforces the assumption that May 5th contains a large amount (70%) of HVOP attributable to TB.

345 Upon comparison with the April 28th and May 5th samples, NEXAFS spectra of HVOP particles from the May 14th
sample (taken 10 hours after raining) show a slightly enhanced C=C peak and an almost absent COH peak. These same features
can be seen in the previously reported NEXAFS spectra of ASOP (Wang et al., 2016; Veghte et al., 2017) and “free light” SOM
isolated in that work for comparison. One reason for the difference in COH peak intensity may be due to the presence of
levoglucosan or other plant derived products such as polysaccharides, tannins, or lignin fragments (Marín-Spiotta et al., 2008)
350 (which contains multiple –OH groups) in the samples affected by smoke plumes, a common product of biomass burning from
the pyrolysis of carbohydrates (Lakshmanan and Hoelscher, 1970). Another differentiating factor is the ratio of intensities
between the –COOH peak at 288.6 eV and the C=C peak at 285.3 eV. In the TB spectrum, the –COOH peak is much higher
than the C=C peak compared to the ASOP spectrum (Wang et al., 2016) and this difference is borne out in the spectra collected
for the current study as well.

355 The chemical composition of TB has been reported in literature, with fresh TB being comprised various biomass tar
products with a substantial degree of aromaticity. The nonpolar products were the most strongly associated with the wavelength
dependence in absorption seen in TB and were found in greater number in fresh TB (Li et al., 2019). Photochemical oxidation
in the presence of O₃ or OH radicals was shown to bleach this wavelength dependence after about 3.5 days (Sumlin et al.,
2017). Photooxidation was suggested to break network of conjugated double bonds in the TB chromophores, resulting in more
360 oxygenated (carbonyl substituted) products which concentrated on the surface of the TB (Hand et al., 2005; Tóth et al., 2014).
ASOP are comprised, in part, of high molecular weight humic-like substances. These compounds contain multiple conjugated
ring systems which likely serve as chromophores in a similar way to the tar materials in TB (Kumada, 1955). The extensive
conjugated systems may be driving the enhanced C=C peak in Fig. 5 for ASOP associated spectra. Also, because humic-like
substances in ASOP are substituted by many different functional groups, rather than the nonpolar components of TB, more
365 reactive sites may be available for oxidation. This may lead to faster atmospheric processing and bleaching of ASOP
wavelength dependent absorption. This aging could also serve to increase the viscosity of ASOP in the same way that it does
to TB (Adachi et al., 2019). Another factor differentiating their aging processes could be their hygroscopicity. Reports have
shown TB change very little in morphology even when cycled from 0 to 100% RH, having a growth factor of ~1.09 (Adachi
and Buseck, 2011; Semeniuk et al., 2007). However, because ASOP are formed via dissolved SOM that is ejected during
370 precipitation, they are expected to be more hygroscopic. Indeed, Wang (2016) showed the results of RH cycling in
supplemental figures and found a growth factor of ~1.15 at 85% RH along with droplet activation at 98% RH.

Also shown is a spectrum of organic particles not associated with HVOP. This spectrum is characterized by small
C=C and COH absorptions with an intense –COOH peak. The large –COOH peak seen in non-SP-HVOP organics is indicative
of much higher contributions of carboxylic functional groups which define the solubility of the organic particles. This increased
375 solubility would lower the viscosity of the non-viscous organic particles due to their substantial water content and thus these
particles would be deformed and flatten upon impact. Viscous SOA have been observed previously, under both laboratory

and ambient conditions (Virtanen et al., 2010). The HVOP discussed in the current work differ from these viscous SOA by virtue of their mode of formation with both TB and ASOP being comprised of larger organic compounds. Furthermore, the particles themselves are larger. Viscous SOA are formed and observed as much smaller particles (<100 nm) than TB and ASOP (300 – 700 nm).

The left-panel of Fig. 5 shows STXM/NEXAFS spectra of ASOP proxies generated from the SOM brine. The top four spectra from the puddle water samples all show a fairly strong carbonate signal at around 290.1 eV along with two broad potassium peaks (L_2 and L_3) at about 298 eV. Also, the corresponding inset carbon speciation map shows a large, inorganic dominant region (teal blue). The presence of large inorganic regions is not consistent with the mainly organic particles seen previously. However, because these samples were nebulized from bulk solutions to begin with, water soluble carbonates from the soil must have been present upon nebulization. To better model the bubble bursting mechanism of generating ASOP, dry N_2 gas was bubbled through puddle water samples and particles resulted from the bubble bursting at the air-solution interface were collected. In these experiments, generated particles showed almost a complete reduction of the carbonate peak and a small reduction in the potassium peaks, plus the carbon speciation map (Fig. 5 inset) showed an entirely organic dominant particle. Comparison between the spectra of ASOP proxies generated in the ‘bubbling experiment’ with the ambient spectra of HVOP from May 14th shown in the right panel indicates substantial similarities between their spectral features. This includes the diminished $-COOH/C=C$ peak ratio and the relative absence of a COH peak.

For more quantitative comparison two sets of peak ratios were calculated. The first between the $-COOH$ and $C=C$ absorptions and another between the $-COOH$ and COH peaks. The peak ratios are plotted in Fig. 6 below for all of the ambient samples, the proxy particles from the “bubbling” experiment, and the two literature spectra. As noted above, the TB spectra have higher $-COOH/C=C$ and $-COOH/COH$ peak ratios. The separation between these two ratios is also the largest for the TB spectrum. While the April 28th and May 5th peak ratios are not quite as high, they all bear a strong resemblance to the TB spectrum. Perhaps alone the similarity would not be enough to define these ambient samples as TB; however, coupled with the other data presented here the peak ratios support the SP-HVOP seen in the May 5th sample as being TB. The April 28th peak ratios are further separated than the literature ASOP peak ratios are, but closer together than the TB literature peak ratios. With the possibility of ASOP travelling from surrounding states (with active rainfall) as well as the proximity of smoke plumes it is likely that the HVOP observed on April 28th are comprised of both TB and ASOP. The May 14th sample peak ratios however, are much different and are more comparable to the ASOP peak ratios. The $-COOH/C=C$ ratio is much lower in both cases and the $-COOH/COH$ ratio is of the same value. From the lab generated aerosols, the bubbling sample peak ratios are also similar to the ASOP peak ratios, suggesting that bubbling reproduces the mechanism of ASOP generation. Another noteworthy observation is the difference between the April 28th or May 5th peak ratios and the May 14th peak ratios. Although each of these samples showed the presence of HVOP and each sample’s AAE suggested that these HVOP were BrC, there is a stark contrast between the smoke influenced samples (Apr 28th and May 5th) and the rainfall influence samples (May 14th), suggesting different sources of SP-HVOP in these two cases.

410 4 Conclusion

BrC particles like TB and ASOP and their place in the global aerosol budget are yet insufficiently understood. Here, it was shown that presence of HVOP is correlated with the BrC properties of overall aerosols, indicated by the elevated values of AAE. On multiple days HVOP were observed to comprise a significant fraction of the fine mode aerosols.

415 Tilted SEM was used to identify HVOP fractions in a number of samples taken during this study, and the fractions of HVOP present for each sample were determined based on the aspect ratios of individual particles. The HVOP fractions showed a strong correlation with the average AAE over the sampling periods, with an R^2 of 0.85. When extrapolated to 100% fraction of HVOP an AAE of 2.6 was calculated, consistent with literature reported values of BrC (Lack and Langridge, 2013). These observed AAE values suggest that BrC relevant particles can be identified by methods of chemical imaging based on three dimensional morphology coupled with chemical composition. These samples were further classified into samples with TB and 420 samples with ASOP based on their NEXAFS characteristics and by comparing smoke and precipitation data during corresponding collection periods.

Chemical imaging showing the differences between ASOP and TB laden samples was performed using STXM/NEXAFS spectromicroscopy. Samples unaffected by recent rain, collected while smoke plumes were present showed a higher $-\text{COOH}/\text{C}=\text{C}$ peak ratio and an elevated $-\text{COH}$ peak intensity. The elevated $-\text{COH}$ peak is likely due to the presence 425 of sugars such as levoglucosan or other less oxidized molecules. The sample from May 14th was collected 10 hours after a rain event and had less influence from smoke plumes. This sample showed a much more subdued $-\text{COH}$ peak and a smaller $-\text{COOH}/\text{C}=\text{C}$ peak ratio. Comparing the ambient spectra collected here with previously collected spectra supported the presence of TB in the smoke-affected samples and the presence of ASOP in the sample taken after a rain event. Peak ratios between $-\text{COOH}$ and $\text{C}=\text{C}$ and between $-\text{COOH}$ and COH were calculated to serve as a quantitative metric that can be potentially used 430 to differentiate between TB and ASOP, and, probably more generally, between the smoke-affected samples and the samples with particles induced by rainfall.

HVOP are a subclass of BrC particles which can define the nature of aerosol-radiation interactions during time periods when they are prevalent. Differentiating between types of HVOP like TB and ASOP has proven to be a challenging task that relies on subtle differences in chemical composition and atmospheric conditions at the time of sampling. Of the three sampling 435 days focused on in this study, only one indicated an appreciable number of ASOP present. Because the conditions necessary for ASOP emission depend both on soil properties and precipitation characteristics, the dominant source of HVOP will often be TB due to the frequency of biomass burning instances along with the large number of particles they emit. ASOP are likely to contribute to aerosol properties, optical and otherwise, only during short time periods where the emission conditions are met. Further questions still exist about ASOP specifically. How do soil characteristics affect the composition of ASOP? How 440 are ASOP transformed as they travel through the atmosphere? What are the emissions factors of ASOP? Answering questions like these may improve the quality of models in regions where large areas with open soils such as agricultural fields and grasslands are exposed to intensive rains, especially during rainy seasons when ASOP might be prevalent.

Data availability. The data set used here is available for download as a .zip file at <https://doi.org/10.17605/OSF.IO/G8FPW>
445 (Fraund, 2020).

Code availability. MatLab code used for the current work is available as a Supplement.

Author contributions. Conceptualization, M.F., D.J.B., S.C., D.V., M.K.G., A.L., and R.C.M.; Software, M.F., D.J.B. and
450 R.C.M.; Formal Analysis, M.F. and D.J.B.; Investigation, M.F., D.J.B., S.C., D.Q.P., D.V., J.W., G.K., and R.C.M.; Resources,
K.T., M.K.G., A.L., and R.C.M.; Data Curation, M.F., D.J.B., S.C., D.V., and R.C.M.; Writing – Original Draft, M.F. and
D.J.B.; Writing – Review & Editing, all authors; Visualization, M.F., D.J.B., and S.C.; Supervision, M.K.G., A.L., and R.C.M.
; Project Administration, M.K.G., A.L., R.C.M.; Funding Acquisition, M.K.G., A.L., and R.C.M.

455 *Competing interests.* The authors declare that they have no conflict of interest

Acknowledgements. We acknowledge support from the US Department of Energy’s Atmospheric System Research program,
Office of Biological and Environmental Research (OBER), award DE-SC0018948. PSAP records obtained from the
Atmospheric Radiation Measurement (ARM) User Facility of OBER was used in this study. Chemical imaging of particles
460 was performed at The Advanced Light Source at Lawrence Berkeley National Laboratory. The beamline staff of 11.0.2 and
5.3.2 helped make the current work possible, specifically: David Shapiro, David Kilcoyne, Matthew Markus, and Hendrik
Ohldag. A portion of the research was performed using EMSL (grid.436923.9), a DOE Office of Science User Facility
sponsored by the Office of Biological and Environmental Research. Part of the research described in this paper was performed
at the Canadian Light Source, a national research facility of the University of Saskatchewan, which is supported by the Canada
465 Foundation for Innovation (CFI), the Natural Sciences and Engineering Research Council (NSERC), the National Research
Council (NRC), the Canadian Institutes of Health Research (CIHR), the Government of Saskatchewan, and the University of
Saskatchewan. The CLS work was done at the SM beamline 10ID-1 with help of its staff: Jian Wang, Yingshen Lu, and Jan
Geilhufe. The NOAA HYSPLIT transport and dispersion model (<http://www.ready.noaa.gov>) was used to calculate backward
trajectories for this publication. We would acknowledge Shih-Ming Huang from Sonoma Technology Inc. for suggesting the
470 use of the archived precipitation data from the National Weather Service’s (NWS) Advanced Hydrologic Prediction Service
reported here for the selected periods of our study.

References

Adachi, K., and Buseck, P. R.: Atmospheric tar balls from biomass burning in Mexico, *Journal of Geophysical Research: Atmospheres*, 116, 2011.

- 475 Adachi, K., Sedlacek, A. J., Kleinman, L., Springston, S. R., Wang, J., Chand, D., Hubbe, J. M., Shilling, J. E., Onasch, T. B., and Kinase, T.: Spherical tarball particles form through rapid chemical and physical changes of organic matter in biomass-burning smoke, *Proceedings of the National Academy of Sciences*, 116, 19336-19341, 2019.
- Alexander, D. T., Crozier, P. A., and Anderson, J. R.: Brown carbon spheres in East Asian outflow and their optical properties, *Science*, 321, 833-836, 2008.
- 480 Andreae, M., and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmospheric Chemistry and Physics*, 6, 3131-3148, 2006.
- Data Discovery: <https://adc.arm.gov/discovery/>, access: May-10, 2019.
- Backman, J. C. G., Virkkula, A. O., Vakkari, V., Beukes, P., van Zyl, P., Josipovic, M., Piketh, S., Tiitta, P., Chiloane, K., and Petaja, T.: Differences in aerosol absorption Ångström exponents between correction algorithms for particle soot absorption photometer measured on South African Highveld, *Atmospheric Measurement Techniques Discussions*, 2014.
- 485 Bergstrom, R. W., Pilewskie, P., Russell, P. B., Redemann, J., Bond, T. C., Quinn, P. K., and Sierau, B.: Spectral absorption properties of atmospheric aerosols, 2007.
- Bond, T. C., and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol science and technology*, 40, 27-67, 2006.
- 490 Bond, T. C., Doherty, S. J., Fahey, D., Forster, P., Berntsen, T., DeAngelo, B., Flanner, M., Ghan, S., Kärcher, B., and Koch, D.: Bounding the role of black carbon in the climate system: A scientific assessment, *Journal of Geophysical Research: Atmospheres*, 118, 5380-5552, 2013.
- Chakrabarty, R., Moosmüller, H., Chen, L.-W., Lewis, K., Arnott, W., Mazzoleni, C., Dubey, M., Wold, C., Hao, W., and Kreidenweis, S.: Brown carbon in tar balls from smoldering biomass combustion, *Atmospheric Chemistry and Physics*, 10, 6363-6370, 2010.
- 495 Fast, J. D., Berg, L. K., Alexander, L., Bell, D., D'Ambro, E., Hubbe, J., Kuang, C., Liu, J., Long, C., and Matthews, A.: Overview of the HI-SCALE Field Campaign: A New Perspective on Shallow Convective Clouds, *Bulletin of the American Meteorological Society*, 100, 821-840, 2019.
- Feng, Y., Ramanathan, V., and Kotamarthi, V.: Brown carbon: a significant atmospheric absorber of solar radiation?, *Atmospheric Chemistry and Physics*, 13, 8607-8621, 2013.
- 500 Ferrare, R., Turner, D., Clayton, M., Schmid, B., Redemann, J., Covert, D., Elleman, R., Ogren, J., Andrews, E., and Goldsmith, J. E.: Evaluation of daytime measurements of aerosols and water vapor made by an operational Raman lidar over the Southern Great Plains, *Journal of Geophysical Research: Atmospheres*, 111, 2006.
- Fraund, M., Pham, D. Q., Bonanno, D., Harder, T. H., Wang, B., Brito, J., de Sá, S. S., Carbone, S., China, S., and Artaxo, P.: 505 Elemental Mixing State of Aerosol Particles Collected in Central Amazonia during GoAmazon2014/15, *Atmosphere*, 8, 173, 2017.
- Fraund, M., Park, T., Bonanno, D., Pham, D. Q., and Moffet, R. C.: Quantitative capabilities of STXM to measure spatially resolved organic volume fractions of mixed organic/inorganic particles, *Atmospheric Measurement Techniques*, 12, 1619, 2019.
- 510 Guizar-Sicairos, M., Thurman, S. T., and Fienup, J. R.: Efficient subpixel image registration algorithms, *Optics Letters*, 33, 156-158, 2008.
- Hand, J. L., Malm, W., Laskin, A., Day, D., Lee, T.-b., Wang, C., Carrico, C., Carrillo, J., Cowin, J. P., and Collett Jr, J.: Optical, physical, and chemical properties of tar balls observed during the Yosemite Aerosol Characterization Study, *Journal of Geophysical Research: Atmospheres*, 110, 2005.
- 515 Hegg, D. A., Clarke, A. D., Doherty, S. J., and Ström, J.: Measurements of black carbon aerosol washout ratio on Svalbard, *Tellus B: Chemical and Physical Meteorology*, 63, 891-900, 2011.
- Hoffer, A., Gelencsér, A., Guyon, P., Kiss, G., Schmid, O., Frank, G., Artaxo, P., and Andreae, M.: Optical properties of humic-like substances (HULIS) in biomass-burning aerosols, *Atmospheric Chemistry and Physics*, 6, 3563-3570, 2006.
- Jennings, S., Pinnick, R., and Gillespie, J.: Relation between absorption coefficient and imaginary index of atmospheric aerosol constituents, *Applied optics*, 18, 1368-1371, 1979.
- 520 Joung, Y. S., and Buie, C. R.: Aerosol generation by raindrop impact on soil, *Nature communications*, 6, 6083, 2015.
- Kilcoyne, A., Tyliczszak, T., Steele, W., Fakra, S., Hitchcock, P., Franck, K., Anderson, E., Harteneck, B., Rightor, E., and Mitchell, G.: Interferometer-controlled scanning transmission X-ray microscopes at the Advanced Light Source, *Journal of Synchrotron Radiation*, 10, 125-136, 2003.

- 525 Kirchstetter, T. W., Novakov, T., and Hobbs, P. V.: Evidence that the spectral dependence of light absorption by aerosols is affected by organic carbon, *Journal of Geophysical Research: Atmospheres*, 109, 2004.
- Koppmann, R., Czapiewski, K. v., and Reid, J.: A review of biomass burning emissions, part I: gaseous emissions of carbon monoxide, methane, volatile organic compounds, and nitrogen containing compounds, *Atmospheric Chemistry and Physics Discussions*, 5, 10455-10516, 2005.
- 530 Kumada, K.: Absorption spectra of humic acids, *Soil Science and Plant Nutrition*, 1, 29-30, 10.1080/00380768.1955.10434360, 1955.
- Lack, D., and Langridge, J.: On the attribution of black and brown carbon light absorption using the Ångström exponent, *Atmospheric Chemistry and Physics*, 13, 10535-10543, 2013.
- Lakshmanan, C. M., and Hoelscher, H. E.: Production of Levoglucosan by Pyrolysis of Carbohydrates Pyrolysis in Hot Inert Gas Stream, *Starch-Stärke*, 22, 261-264, 1970.
- 535 Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of atmospheric brown carbon, *Chemical reviews*, 115, 4335-4382, 2015.
- Lawless, P. A., Rodes, C. E., and Ensor, D. S.: Multiwavelength absorbance of filter deposits for determination of environmental tobacco smoke and black carbon, *Atmospheric Environment*, 38, 3373-3383, 2004.
- 540 Li, C., He, Q., Schade, J., Passig, J., Zimmermann, R., Meidan, D., Laskin, A., and Rudich, Y.: Dynamic changes in optical and chemical properties of tar ball aerosols by atmospheric photochemical aging, *Atmos. Chem. Phys.*, 19, 139-163, 10.5194/acp-19-139-2019, 2019.
- Lin, P., Bluvshstein, N., Rudich, Y., Nizkorodov, S. A., Laskin, J., and Laskin, A.: Molecular chemistry of atmospheric brown carbon inferred from a nationwide biomass burning event, *Environmental science & technology*, 51, 11561-11570, 2017.
- 545 Marín-Spiotta, E., Swanston, C. W., Torn, M. S., Silver, W. L., and Burton, S. D.: Chemical and mineral control of soil carbon turnover in abandoned tropical pastures, *Geoderma*, 143, 49-62, 2008.
- Moffet, R. C., Henn, T., Laskin, A., and Gilles, M. K.: Automated Chemical Analysis of Internally Mixed Aerosol Particles Using X-ray Spectromicroscopy at the Carbon K-Edge†, *Analytical Chemistry*, 82, 7906-7914, 2010.
- Hazard Mapping System Fire and Smoke Product: <https://www.ospo.noaa.gov/Products/land/hms.html>, access: May-10, 2019.
- 550 Quantitative Precipitation Estimates: <https://water.weather.gov/precip/>, access: May-10, 2019.
- O'Brien, R. E., Wang, B., Laskin, A., Riemer, N., West, M., Zhang, Q., Sun, Y., Yu, X. Y., Alpert, P., and Knopf, D. A.: Chemical imaging of ambient aerosol particles: Observational constraints on mixing state parameterization, *Journal of Geophysical Research: Atmospheres*, 120, 9591-9605, 2015.
- 555 Parworth, C., Fast, J., Mei, F., Shippert, T., Sivaraman, C., Tilp, A., Watson, T., and Zhang, Q.: Long-term measurements of submicrometer aerosol chemistry at the Southern Great Plains (SGP) using an Aerosol Chemical Speciation Monitor (ACSM), *Atmospheric Environment*, 106, 43-55, 2015.
- Pósfai, M., Gelencsér, A., Simonics, R., Arató, K., Li, J., Hobbs, P. V., and Buseck, P. R.: Atmospheric tar balls: Particles from biomass and biofuel burning, *Journal of Geophysical Research: Atmospheres*, 109, 2004.
- Radke, L. F., Hobbs, P. V., and Eltgroth, M. W.: Scavenging of aerosol particles by precipitation, *Journal of Applied Meteorology*, 19, 715-722, 1980.
- 560 Regier, T., Krochak, J., Sham, T., Hu, Y., Thompson, J., and Blyth, R.: Performance and capabilities of the Canadian Dragon: The SGM beamline at the Canadian Light Source, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 582, 93-95, 2007.
- Rizzo, L. V., Artaxo, P., Mueller, T., Wiedensohler, A., Paixao, M., Cirino, G. G., Arana, A., Swietlicki, E., Roldin, P., and Fors, E. O.: Long term measurements of aerosol optical properties at a primary forest site in Amazonia, *Atmospheric Chemistry and Physics*, 13, 2391-2413, 2013.
- 565 Rolph, G., Stein, A., and Stunder, B.: Real-time Environmental Applications and Display sYstem: READY, *Environmental Modelling & Software*, 95, 210-228, 2017.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, 2006.
- 570 Semeniuk, T. A., Wise, M. E., Martin, S. T., Russell, L. M., and Buseck, P. R.: Hygroscopic behavior of aerosol particles from biomass fires using environmental transmission electron microscopy, *Journal of Atmospheric Chemistry*, 56, 259-273, 2007.

Sheridan, P., Delene, D., and Ogren, J.: Four years of continuous surface aerosol measurements from the Department of Energy's Atmospheric Radiation measurement Program Southern Great Plains Cloud and Radiation Testbed site, *Journal of Geophysical Research: Atmospheres*, 106, 20735-20747, 2001.

575 AirNow - Tech Navigator: <http://www.airnowtech.org/navigator/>, access: May-10, 2019.

Springston, S., Andrews, E., and Salwen, C.: Particle Soot Absorption Photometer (AOSPSAP3W1M). ARM, 10.5439/1225037, 2016

Springston, S.: Radiance Research Particle Soot/Absorption Photometer Instrument Handbook, DOE ARM Climate Research Facility, Washington, DC (United States), 2018.

580 Stein, A., Draxler, R. R., Rolph, G. D., Stunder, B. J., Cohen, M., and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bulletin of the American Meteorological Society*, 96, 2059-2077, 2015.

Sumlin, B. J., Pandey, A., Walker, M. J., Pattison, R. S., Williams, B. J., and Chakrabarty, R. K.: Atmospheric photooxidation diminishes light absorption by primary brown carbon aerosol from biomass burning, *Environmental Science & Technology Letters*, 4, 540-545, 2017.

585 Tivanski, A. V., Hopkins, R. J., Tyliczszak, T., and Gilles, M. K.: Oxygenated interface on biomass burn tar balls determined by single particle scanning transmission X-ray microscopy, *The Journal of Physical Chemistry A*, 111, 5448-5458, 2007.

Tóth, A., Hoffer, A., Nyiró-Kósa, I., Pósfai, M., and Gelencsér, A.: Atmospheric tar balls: aged primary droplets from biomass burning?, *Atmospheric Chemistry and Physics*, 14, 6669-6675, 2014.

590 Veghte, D. P., China, S., Weis, J., Kovarik, L., Gilles, M. K., and Laskin, A.: Optical Properties of Airborne Soil Organic Particles, *ACS Earth and Space Chemistry*, 1, 511-521, 2017.

Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., Yli-Pirilä, P., Leskinen, J., Mäkelä, J. M., Holopainen, J. K., Pöschl, U., and Kulmala, M.: An amorphous solid state of biogenic secondary organic aerosol particles, *Nature*, 467, 824-827, 2010.

Wang, B., Harder, T. H., Kelly, S. T., Piens, D. S., China, S., Kovarik, L., Keiluweit, M., Arey, B. W., Gilles, M. K., and Laskin, A.: Airborne soil organic particles generated by precipitation, *Nature géoscience*, 9, 433, 2016.

595 Williams, J., De Reus, M., Krejci, R., Fischer, H., and Ström, J.: Application of the variability-size relationship to atmospheric aerosol studies: estimating aerosol lifetimes and ages, 2002.

Zhang, Y., Forrister, H., Liu, J., Dibb, J., Anderson, B., Schwarz, J. P., Perring, A. E., Jimenez, J. L., Campuzano-Jost, P., and Wang, Y.: Top-of-atmosphere radiative forcing affected by brown carbon in the upper troposphere, *Nature Geoscience*, 10, 486-489, 2017.

600





Figure 1: Time series for a) absorption Ångström exponent (AAE) calculated from red and blue absorption coefficients, b) absorption coefficients measured at red (660 nm), green (522 nm), and blue (470 nm) wavelengths, and c) CO mixing ratio and ambient particle concentration. Grey vertical bars indicate sampling periods where microscopy samples were collected. ~~of BrC appearance when particle samples were analyzed.~~ Red vertical bars represent rain events. Arrows indicate the three sampling periods investigated in the current work where BrC particles were expected. PSAP data for May 8th is not available due to instrument error.


605




610 Figure 2: (Top row) Tilted (75°) SEM images show differences in **SPHVOP** fractions between three samples. Each of these samples were collected on a MOUDI stage 8, which selected for particles in the 150-360 nm size range. Magenta arrows point to characteristic HVOP. (Middle row) Carbon speciation maps with red representing regions with enhanced C=C bonding, green representing organics, and teal representing inorganics. (Bottom row) Total carbon absorption (TCA) images calculated from dividing thickness by the area equivalent diameter of each particle. The carbon speciation and TCA maps show the same fields of view. All scale bars are 1 μm in length

625

Figure 4: Smoke, fire, and precipitation data along with HYSPLIT back trajectories for three sample dates. The red circle represents the sampling site while the small red triangles represent fires. The gray overlays seen in the top row represent detected smoke particles (overlapping smoke plumes are shown in darker shades of gray). The bottom row shows the 24-hour average precipitation amount over the sampling date. The top row maps were obtained using the AirNow-Tech navigator using the Hazard Mapping System smoke product from NOAA, Source: U.S. EPA AirNow-Tech (SonomaTechnologiesInc., 2019). HYSPLIT trajectories for April 28th and May 5th are for 24 hours. The May 14th back trajectory was truncated at 10 hours due to a rain event with significant precipitation scavenging. Precipitation maps were made using the NWS AHPS (NWS, 2019).



630 Figure 5: Comparison of NEXAFS spectra between laboratory generated ASOP proxies (left) and four ambient ASOP samples (right). Inset carbon speciation maps are shown for representative particles with green representing organic dominant regions and teal representing inorganic dominant regions. TCA images, like the ones shown in Fig. 2 are shown as well. Only one image is shown for the ambient samples as they all look similar. The ASOP proxy spectra are show for particles generated by nebulizing aquatic samples collected at the site and by bubbling N₂ gas through one of the aquatic samples.



635

Figure 6: Plot of peak ratios for: (blue dots) carboxylic acid peak at 288.6 eV and the carbon double bond peak at 285.3 eV and (red dots) carboxylic acid peak and the COH peak at 286.7 eV. ASOP Proxies refers to particles generated via bubbling through the aquatic sample of SOM.




Figure S1. Time series of a) absorption Ångström exponent (AAE) b) ozone and sulfur dioxide particle concentration and bc carbon monoxide and nitrous oxide sulfur dioxide. Grey vertical bars indicate periods of Brown Carbon (BrC) appearance when particle samples were analyzed. Red vertical bars represent rain events. Arrows indicate the three sampling periods investigated in the current work.




Figure S1S2. Linear correlation between fractions of Spherical Particles (SP) high viscosity organic particles (HVOP) and the red/blue absorption Ångström exponent (AAE). A linear fit is shown, having an R^2 value of 85%. Extrapolated to 100% SP yields an AAE value of 2.5, consistent with typical BrC characteristics (Lack and Langridge, 2013).




Figure S3. Hybrid single particle Lagrangian integrated trajectory (HYSPLIT) back trajectories for three sample dates in 2016 at different starting altitudes. The starting location (represented by the black star) is the Lamont, OK central ARM facility at 36.60 N and 97.49 W. The April 28th and May 5th plots are 24 hour trajectories while the May 14th plots are 10 hour trajectories. Different colored trajectories within a single plot represent new trajectories every 3 hours after the initial starting time. Archived data was taken from the North American Mesoscale (NAM) sigma-pressure hybrid model.