

Dear Prof. Jianzhong Ma,

Thank you for your valuable comments and suggestions. We have made a revision accordingly. The point to point responses are provided below in Italic. The
5 *comparison of our manuscript between this version and the previous version is also provided. Please let us know should you have future questions.*

Thanks again for your time and efforts on our paper.

10 *Best regards,*
Jane Liu on behalf of all the co-authors

Comments to the Author:

15 Dear Prof. Jane Liu and all authors,

I had sent the revised manuscript to two referees for further review, but one of them did not give his/her review report until now. I think your manuscript can be accepted after the following issues have been addressed.

20 Figure 2b shows the 7-day backward trajectories (in meters above the ground) arriving at the TP surface. Since no trajectories are shown for the domain within the TP, it is difficult to see the transport pathways of air masses arriving at a specific region, e.g., the central TP. Air masses from the west (blue dots) are still at high
25 altitudes (above 1000 m) at the western edge of the TP. Are 1-day backward trajectories for these air masses included in the plot? Will these air masses enter into the TP domain from the west or other side(s) of the TP? The authors may consider revising Fig. 2b and/or providing additional discussion for better understanding this

subpanel.

30

While Fig. 2a presents low surface BC concentrations on the north of the TP, Fig. 2c shows high BC transported fluxes there. How about the regional distribution of BC column density? Or, does an elevated BC layer existing on the north of the TP?

35 Best regards, Jianzhong

Response:

Figure 2 is shown to be an example of the estimate of BC transport to the TP surface and to help to understand the calculation of BC transport flux to the TP surface and its spatial distribution.

Following the editor's advice, we replaced the Figure 2a in the last version with a new figure showing BC total column, and replaced Figure 2b with a new figure showing all 7-day backward trajectories arriving at the TP surface in the month. The distribution of surface BC concentrations is moved to Supplement.

In this example, both BC total column and BC concentrations at the surface are high in central China around 110°E (Figures 2a and S3). A large number of trajectories pass through central China and finally arrive at the TP surface (Figure 2b). Therefore, the amount of BC transported from central China to the TP is high (Figure 2c). Trajectories from Xinjiang province, northwestern China (to the north of the TP) are also at low altitudes (Figure 2b). However, BC total column and BC concentrations at the surface are low there (Figures 2a and S3). Therefore, the amount of BC transported from Xinjiang province to the TP surface is lower than that from central China. Air masses from the Middle East and central Asian countries are from high altitudes and transported downward to the surface over different locations in the TP (Figure 2b). Although these trajectories are in large numbers, the amount of BC transported from the Middle East and central Asian countries is relatively small

(Figure 2c) because the backward trajectories appear at high altitudes (Figure 2b) and BC concentrations over there are low (Figure 2a).

Please see the revisions in lines 255-266.

60

Impacts of atmospheric transport and biomass burning on the interannual variation in black carbon aerosols over the Tibetan Plateau

Han Han^{1,#}, Yue Wu^{1,2,#}, Jane Liu^{3,1}, Tianliang Zhao⁴, Bingliang Zhuang¹, Honglei Wang⁴, Yichen Li¹, Huimin Chen¹, Ye Zhu⁵, Hongnian Liu¹, Qin'geng Wang⁶, Shu Li¹, Tijian Wang¹, Min Xie¹, and Mengmeng Li¹

¹School of Atmospheric Sciences, Nanjing University, Nanjing, China

²Suzhou Meteorological Bureau, Suzhou, China

³Department of Geography and Planning, University of Toronto, Toronto, Canada

⁴School of Atmospheric Physics, Nanjing University of Information Science & Technology, Nanjing, China

⁵Shanghai Public Meteorological Service Centre, Shanghai, China

⁶School of the Environment, Nanjing University, Nanjing, China

[#]These authors contributed equally to this work.

Correspondence: Jane Liu (janejj.liu@utoronto.ca)

Abstract

Atmospheric black carbon (BC) in the Tibetan Plateau (TP) can largely impact regional and global climate. Still, studies on the interannual variation in atmospheric BC over the TP, and associated variation in BC sources and controlling factors are rather limited. In this study, we characterize the variations in atmospheric BC over the TP surface layer through analysis of 20-year (1995-2014) simulations from a global chemical transport model, GEOS-Chem. The results show that, surface BC concentrations over the TP vary largely in space and by season, reflecting complicated interplays of BC sources from different origins. Of all areas in the TP, surface BC

90 concentrations are highest over the eastern and southern TP, where surface BC are
susceptible to BC transport from East Asia and South Asia, respectively. Applying a
backward-trajectory method that combines the BC concentrations from GEOS-Chem
simulations and trajectories from the Hybrid Single-Particle Lagrangian Integrated
Trajectory (HYSPLIT) model, we assess the contributions of worldwide source
95 regions to surface BC in the TP. We estimate that on the 20-year average, 77% surface
BC in the TP comes from South Asia (43%) and East Asia (35%). Regarding seasonal
variation in nonlocal influences, South Asia and East Asia are dominant source
regions in winter and summer, respectively, in terms of both magnitude and affected
areas in the TP. Interannually, surface BC over the TP is largely modulated by
100 atmospheric transport of BC from nonlocal regions year-round and by biomass
burning in South Asia, mostly in spring. We find that the extremely strong biomass
burning in South Asia in the spring of 1999 greatly enhanced surface BC
concentrations in the TP (31% relative to the climatology). We find that the strength
of the Asian monsoon correlates significantly with the interannual variation in the
105 amount of BC transported to the TP from nonlocal regions. In summer, a stronger East
Asian summer monsoon and a South Asian summer monsoon tend to, respectively,
lead to more BC transport from central China and northeastern South Asia to the TP.
In winter, BC transport from central China is enhanced in years with a strong East
Asia winter monsoon or a Siberian High. A stronger Siberian High can also bring
110 more BC from northern South Asia to the TP. This study underscores the impacts of
atmospheric transport and biomass burning on the interannual variation in surface BC
over the TP. It reveals a close connection between the Asian monsoon and
atmospheric transport of BC from nonlocal regions to the TP.

115 **1 Introduction**

Black carbon (BC) is a carbonaceous aerosol formed from combustion of carbon-
based fuels and materials. BC in the atmosphere is a major air pollutant and a strong

absorber of solar radiation (Bond et al., 2013). Atmospheric BC can greatly influence regional (Ramanathan and Carmichael, 2008; Zhuang et al., 2018) and global (Allen et al., 2012; Chung et al., 2012) climate through multiple mechanisms. It can cause atmospheric heating (Cappa et al., 2012) and surface dimming (Flanner et al., 2009) and influence cloud formation and development processes (Jacobson et al., 2012). Furthermore, after its deposition on snow or ice, BC reduces the surface albedo and accelerates the melting of glaciers and snow cover (Hansen and Nazarenko, 2004; Flanner et al., 2007).

The Tibetan Plateau (TP) has an average altitude of over 4 km and an area of 2.6 million km², known as the Third Pole. Because of its special geography, the TP can greatly impact regional and global climate through dynamic and thermal processes (Wu et al., 2015; Li et al., 2018). The TP has a large number of glaciers (over 0.1 million km²) and a wide coverage of snow (59% of the TP in winter) (Qin et al., 2006; Yao et al., 2012). Although atmospheric BC in the TP is among the lowest in the world, BC there can alter the climate (Lau et al., 2010; Jiang et al., 2017), ecosystem (Kang et al., 2019), and hydrology (Barnett et al., 2005) in the TP, consequently influencing the living environment of billions of people in the world. Atmospheric BC is an important factor driving the surface warming in the TP due to its strong absorption of solar radiation (He et al., 2014a). After its deposition to the TP ground, BC in the snow reduces the surface albedo (Ming et al., 2009; Qian et al., 2011; Qu et al., 2014). He et al. (2014a) suggested that the annual mean direct radiative forcing and snow albedo forcing of BC in the TP are respectively ~ 2.3 and ~ 2.9 W m⁻² over the snow-covered regions. Because of the effects on radiation, BC deposited on snow can further reduce the duration of snow cover in the TP (Ménégoz et al., 2014). Zhang et al. (2018) estimated that such reduction is around 3.1 ± 0.1 days. Furthermore, BC in both atmosphere and cryosphere over the TP is responsible for retreats of the snow cover (Menon et al., 2010; Xu et al., 2016) and glaciers (Xu et al., 2009; Ming et al.,

2012; Niu et al., 2020) in the past decades.

Atmospheric BC concentrations in the TP vary with location and season, which was revealed by limited observations over different regions in southern (Marinoni et al., 2010; Putero et al., 2014; Chen et al., 2018), northern (Zhao et al., 2012), and southeastern (Cao et al., 2011; Wang et al., 2018, 2019) TP. Over these regions, seasonal variations in atmospheric BC show different patterns. Chen et al. (2018) observed that in the Himalayas over the southern TP, surface BC concentrations reached the highest in spring (over $0.9 \mu\text{g m}^{-3}$ in April) and the lowest in summer (under $0.1 \mu\text{g m}^{-3}$ in July) from May 2015 to May 2017. Zhao et al. (2012) reported that in the Qilian Shan over the northern TP, surface BC concentrations were less than $0.2 \mu\text{g m}^{-3}$ from May 2009 to March 2011 with the highest in summer and lowest in autumn. Wang et al. (2016) suggested that surface BC concentrations show a seasonality of winter high and spring low at a site in the southeastern TP, while a pattern of winter low and spring high was at a site in the central TP from November 2012 to June 2013. Because BC observations in the TP are limited due to the harsh environment and sparse sites, we can take advantages of numerical simulations to investigate atmospheric BC and its complicated variations over the entire TP in space and by season. Furthermore, how these variations fluctuate from year to year over decades is an issue worthy of exploration.

Due to weak anthropogenic activities, the contribution of local emissions to atmospheric BC in the TP is low (Zhang et al., 2015). Concentrations of atmospheric BC in the TP are greatly influenced by the long-range transport of BC from nonlocal regions (Kopacz et al., 2011; Lu et al., 2012; Kang et al., 2019). South Asia and East Asia are suggested to be two main source regions of atmospheric BC in the TP (Lu et al., 2012). Zhang et al. (2015) estimated that in 2001, local contribution to BC column burden in the TP was only around 10%, while the contributions from South Asia and

East Asia are respectively about 50% and 20%. Some studies also investigated the
175 pathways of BC transport to the TP and derived their characteristics (Cao et al., 2011).
The Asian summer monsoon system was identified as an important influencing factor
for transport of atmospheric species from South Asia to the TP (Chen et al., 2013; Han
et al., 2014; Xu et al., 2014; Zhang et al., 2015). In summer, BC from northern India
can be transported to the middle and upper troposphere and then crossing the
180 Himalayas to the TP via southwesterly winds (Yang et al., 2018). BC emitted in East
Asia can be uplifted to upper layers by the summer monsoon circulation and then
transported to the northeastern TP (Zhang et al., 2015). The midlatitude westerlies are
favorable to BC transport from central Asia and northern India to the western TP
(Chen et al., 2018) but unfavorable to BC transport from eastern China to the TP (Cao
185 et al., 2011). Although previous studies explored the mechanisms of BC transport to
the TP, large uncertainties remain in the quantified fractional contributions of BC
transport from different source regions to the TP (Yang et al., 2018). More
importantly, how BC transport to the TP varies interannually and what are underlying
mechanisms for the variation are unclear. Therefore, it is necessary to examine how
190 BC transport to the TP varies from year to year and how the Asian monsoon affect the
variation.

Previous observations and simulations showed that biomass burning is a major
source of atmospheric BC in the TP (Lu et al., 2012; Zhang et al., 2015). Zhang et al.
195 (2015) estimated that biomass burning together with biofuel emissions can contribute
to around half of the annual mean BC column burden over the TP. Engling et al.
(2011) reported that BC emissions from fire events in Southeast Asia in spring could
probably increase the BC concentrations over a mountain site in the southeastern part
of the TP. Putero et al. (2014) suggested that over half of the high BC episodes in the
200 southern Himalayas were likely affected by the fire events in South Asia. However,
these studies demonstrated the influences of biomass burning in a relative short term

or during some fire events, few investigated the influences in a long term over a decade (Mao and Liao, 2016). The influence of biomass burning on the interannual variation in atmospheric BC over the TP warrants an in-depth study.

205

In this study, we aim to assess the impacts of atmospheric transport and biomass burning on surface BC concentrations over the TP, especially on the interannual variation in BC during 1995-2014. To estimate BC transport from different source regions to the TP, we adopt an approach based on a global chemical transport model, GEOS-Chem (Bey et al., 2001), and a trajectory model, the Hybrid Single-Particle Lagrangian Integrated Trajectory model (HYSPLIT) (Draxler and Hess, 1998; Stein et al., 2015). In the following, the method and models are described in section 2. Section 3 discusses the seasonal variations in surface BC over the TP and in BC transport from source regions to the TP based on the mean status of the 20-year simulations. The interannual variation in surface BC over the TP and the impacts of biomass burning and transport on this variation are analyzed in section 4. Discussion and conclusions are provided in sections 5 and 6, respectively. In this paper, BC refers to BC aerosols in the atmosphere. Surface BC refers to atmospheric BC aerosols in the surface layer.

210

215

220

2 Data and methods

2.1 GEOS-Chem simulations

A global chemical transport model, GEOS-Chem (version v9-02, <http://geos-chem.org>) (Bey et al., 2001), is used to simulate global BC concentrations. GEOS-Chem is driven by the NASA Modern-Era Retrospective Analysis for Research and Applications (MERRA) meteorological data (Rienecker et al., 2011). In this study, we focused on how surface BC in the TP responds to interannual variations in natural influences including biomass burning and meteorology. Therefore, anthropogenic emissions in our simulations were allowed to vary seasonally but not interannually,

225

230 i.e. anthropogenic emissions in 2000 including their seasonality were used for each
year of the study period. We conducted three GEOS-Chem simulations: CTRL,
FixBB, and FixMet. The three simulations covered the study period from 1995 to
2014 (using 1994 for spin-up) at 2° latitude by 2.5° longitude horizontal resolution
with 47 vertical layers. In CTRL, both biomass burning emissions and meteorological
235 fields varied interannually. In FixBB, interannual meteorology was allowed and fire
emissions were fixed in 2005, to remove the impact of the interannual variation in
biomass burning on BC concentrations in the TP. In FixMet, emissions from biomass
burning were allowed to vary interannually and meteorology was fixed in 2005, to
remove the impact of interannual meteorology.

240

In the simulations, global anthropogenic BC emissions were based on Bond et al.
(2007), with an annual emission of 4.4 Tg C in 2000 (Leibensperger et al., 2012).
Global biomass burning emissions of BC were from the Global Fire Emissions
Database version 3 (GFED3) inventory (van der Werf et al., 2010), which covers the
245 period of 1997-2011. BC in GEOS-Chem is represented by two tracers: hydrophobic
and hydrophilic (Park et al., 2003). Freshly emitted BC is mostly (80%) hydrophobic
(Cooke et al., 1999). Hydrophobic BC becomes hydrophilic typically in a few days
(McMeeking et al., 2011), which is simply assumed as 1.15 days in the model, called
an e-folding time (Cooke et al., 1999; Park et al., 2005; He et al., 2014b). Simulations
250 of aerosol dry and wet depositions follows Liu et al. (2001). Dry deposition of
aerosols is simulated using a resistance-in-series model (Walcek et al., 1986)
dependent on local surface type and meteorological conditions, while wet deposition
scheme includes scavenging in convective updrafts, as well as in-cloud and below-
cloud scavenging from convective and large-scale precipitation. Dry deposition is
255 generally smaller than wet deposition (He et al., 2014b; K. Li et al., 2016). Tracer
advection is computed every 15 minutes with a flux-form semi-Lagrangian method
(Lin and Rood, 1996). The tracer moist convection scheme follows Allen et al.

(1996a, b), using GEOS convection, entrainment, and detrainment mass fluxes. The deep convection is parameterized using the relaxed Arakawa-Schubert scheme
260 (Arakawa and Schubert, 1974; Moorthi and Suarez, 1992) and for the shallow convection, the scheme in Hack (1994) is used.

GEOS-Chem has been widely used for BC simulations in the world (Park et al., 2003, 2005; K. Li et al., 2016) and over the TP and its surroundings (Kopacz et al.,
265 2011; Lu et al., 2012; He et al., 2014a; He et al., 2014b; Mao and Liao, 2016). The performance of GEOS-Chem in simulating BC over the TP and surroundings was systematically evaluated by He et al. (2014b) using in situ measurements of BC in surface air, BC in snow, and BC absorption aerosol optical depth. He et al. (2014b) found that the simulated BC in surface air are compared statistically well with
270 observations at sites away from urban areas and the model can generally capture the seasonality of the observations, whereas the BC concentrations in the TP are likely to be underestimated by the model. In Supplement, we evaluated the spatial and seasonal variations in GEOS-Chem BC simulations using observational data from literature. We found a significant correlation ($r = 0.99$, $p < 0.05$) between the observations and
275 simulations at the rural and remote sites (Figure S1). We further compared surface BC concentrations from the GEOS-Chem simulations with those from MERRA2 reanalysis (The Modern-Era Retrospective analysis for Research and Applications version 2, M2TMNXAER, https://cmr.earthdata.nasa.gov/search/concepts/C1276812866-GES_DISC.html). The
280 magnitude and spatial distribution of surface BC concentrations in the TP from GEOS-Chem and MERRA2 are highly consistent, with correlation coefficients over 0.98 in the four seasons (Figure S2). Furthermore, surface BC from GEOS-Chem and MERRA2 shows similar seasonality over the TP as well as over East Asia, South Asia, and Southeast Asia (Table S2). Regarding the interannual variations of surface
285 BC over these regions, the similarity between the simulations and reanalysis is high in

spring, summer, and autumn but low in winter (Table S2).

2.2 Fire and meteorological data

The nighttime fire count product retrieved from ATSR (Along Track Scanning Radiometer) using Algorithm 2, available from European Space Agency (290 http://due.esrin.esa.int/page_wfa.php), were used to verify the biomass burning emissions in GEOS-Chem. ATSR is onboard the Second European Remote-Sensing Satellite (ERS-2). The spatial resolution of the data is 1 km, and the sensor achieves a global coverage every three days. The ATSR satellite data with the period of 1997- (295 2011 were gridded to the GFED3 grids with a resolution of $0.5^\circ \times 0.5^\circ$ in longitude and latitude.

The meteorological data used in this study for analysis and for driving the backward trajectories are the NCEP/NCAR (National Centers for Environmental Prediction/National Center for Atmospheric Research) reanalysis, available from the (300 <https://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis.surface.html>). The data include geopotential height and wind. The horizontal resolution is 2° latitude \times 2.5° longitude.

305

2.3 Transport estimation

Combining GEOS-Chem simulations and HYSPLIT (version 4, http://www.arl.noaa.gov/HYSPLIT_info.php, Draxler and Hess, 1998; Stein et al., 2015) trajectories, we estimated the contributions of different source regions in the (310 world to surface BC in the TP during 1995-2014. HYSPLIT is an atmospheric transport and dispersion model (Fleming et al., 2012), developed by the Air Resources Laboratory of the National Oceanic and Atmospheric Administration (NOAA). Meteorological inputs to HYSPLIT are the NCEP/NCAR reanalysis at a resolution of

2.5° latitude × 2.5° longitude. We evenly divided the TP into 70 GEOS-Chem grids.

315 Considering that the average lifetime of atmospheric BC is about a week, we simulated 7-day backward trajectories originated from each of the 70 grids. The trajectories were initialized four times a day (00, 06, 12 and 18 UTC) during 1995-2014. The starting altitude for the trajectories is 100 m above ground which is within the typical planetary boundary layer in the TP (Ram et al., 2010). We divided the world into seven regions (Figure 1b), including the TP, central Asia, East Asia, South Asia, Southeast Asia, the region of other Asia, Europe, and Africa, and the rest of the world. BC concentrations from CTRL simulation were used in the estimation of BC from different source regions.

325 Lu et al. (2012) proposed a novel approach that combined BC emissions with the backward trajectories from the TP to quantify the origins of BC in the TP. Modifying Lu et al. (2012)'s approach, we combine BC concentrations, instead of BC emissions in Lu et al. (2012), with the backward trajectories from the TP for the same purpose. We assume that BC aerosols have a lifetime of D days and the back trajectories using HYSPLIT are simulated for D days ($D=7$ in this study). To make the estimation stable, the amount of BC transported to a TP surface grid on a day is assumed to be a mean of the BC transport along the backward trajectories originated from that grid in the past D days, i.e.,

$$BC_{imported} = \frac{\sum_{d=1}^D BC_d}{D} \quad (1)$$

335 where BC_d is the amount of BC that are transported to that TP surface grid along the backward trajectory on a previous day d ($d=1, 2, \dots, D$).

Equation (1) provides a way to estimate the amount of BC that is transported to the TP from any model grid outside the TP during a period of interest. For a grid $g_{i,j,k}$ outside the TP (i, j, k are indices for the model grid in longitude, latitude, and altitude coordinates, respectively), the total amount of BC transported from $g_{i,j,k}$ to

the TP during a period of interest ($C_{i,j,k}$) can ~~be estimated~~ be estimated by

$$C_{i,j,k} = \frac{\sum_{n=1}^N c \times v}{D \times M} \quad (2)$$

where n is an index for the number of trajectories. N is the total number of trajectories
345 that have passed through the grid $g_{i,j,k}$ during the period of interest, for example, in a
month. c is the daily BC concentrations at $g_{i,j,k}$ when trajectory n passing $g_{i,j,k}$, and v is
the volume of $g_{i,j,k}$. M is the number of trajectories in a day ($M=4$ in this study).

Therefore, the total amount of BC transported to the TP ($T_{i,j}$) from the entire
tropospheric column above a surface grid $g_{i,j,0}$ in a source region outside the TP during
350 the period of interest can be assessed by

$$T_{i,j} = \sum_{k=1}^K C_{i,j,k} \quad (3)$$

where K is the number of model layers in the troposphere.

Finally, the amount of BC transported from a nonlocal source region to the TP
355 surface can be summed up and the fractional contributions of different source regions
to surface BC in the TP can be quantified. This developed method is inspired by Lu et
al. (2012) and is robust and stable because it is not sensitive to the number of
trajectories taken in a day (M) and the number of days taken for the trajectories (D).

360 Using this method, the amount of BC transported from a nonlocal grid to the TP
surface is determined by both BC concentrations in that grid and the number of
trajectories passing through that grid within the tropospheric column. An example of
the estimation of BC transport to the TP surface in April 2005 is shown in Figure 2. In
this example, both BC total column and BC concentrations at the surface are high in
365 central China around 110°E (~~Figure~~ Figures 2a and S3). A large number of trajectories
pass through central China and finally arrive at the TP surface (Figure 2b). Therefore,
the amount of BC transported from central China to the TP is high (Figure 2c). ~~In~~
contrast, although Trajectories from Xinjiang province, northwestern China (to the
north of the TP) are also at low altitudes (Figure 2b). However, BC total column and

370 BC concentrations at the surface are both low there (Figures 2a and S3). Therefore,
the amount of BC transported from Xinjiang province to the TP surface is lower than
that from central China. Air masses from the Middle East and central Asian countries
are from high altitudes and transported downward to the surface over different
locations in the TP (Figure 2b). Although these trajectories pass through central Asia
375 are also are in large numbers, the amount of BC transported from the Middle East and
central Asia to the TP Asian countries is relatively small (Figure 2c) because BC
concentrations over central Asia are low and the the backward trajectories appear at
high altitudes (Figures 2a and 2b). Figure 2b) and BC concentrations over there are
low (Figure 2a).

380

3 Seasonal variations in surface black carbon over the Tibetan Plateau and in black carbon transport to the Tibetan Plateau

In this section, the analysis is based on the mean status of the 20-year simulations. Figure 3 shows the amount of BC transported from each of the GEOS-Chem grids
385 through the tropospheric column to the TP surface in the four seasons, which varies greatly in space and among seasons. Obviously, surface BC in the TP mainly originates from South Asia and East Asia, especially from the regions near the southern and eastern borders of the TP, including central China, northeastern South Asia, and northern South Asia. These spatial distributions of nonlocal contributions to
390 the BC in the TP in four seasons are similar to those in Lu et al. (2012). We also found a good agreement ($r=0.72$, $p<0.05$) in the estimation of imported BC between this study and Lu et al. (2012) at several sites in the TP, although our estimates are higher than those from Lu et al. (2012). The backward-trajectory approach modified in this study shows strong performance in identifying the source regions for BC over the TP.

395

The simulated annual mean surface BC concentrations over the entire TP are shown in Figure 4a. The BC concentrations are high along the eastern and southern

borders and low in the center of the TP. This spatial variation in surface BC concentrations dominates in the four seasons (not shown). BC concentrations over the TP show strong spatial gradient, which is likely due to the blocking of BC transport by the mountains with high elevations (Cao et al., 2011; Zhao et al., 2017). Figures 4b-4f show the dominant nonlocal influences to surface BC over the TP. In terms of affected areas, South Asia and East Asia are the most important source regions, which respectively impact 67% and 25% areas of the TP in the annual mean BC (Figure 4b). In terms of the amount of BC imported, 77% of surface BC in the TP comes from South Asia (43%) and East Asia (35%). Because of the leeward location of East Asia under prevailing westerlies, the influence of East Asia is constrained mainly in northern and eastern TP. Using a backward-trajectory method, Lu et al. (2012) estimated that the contributions of South Asia and East Asia are respectively 67% and 17%. Using a tracer tagging approach, Zhang et al. (2015) suggested that the contributions of the two regions are respectively 50% and 19%. Our estimate of fractional contribution of both South Asia and East Asia is comparable with the two previous studies, while the estimated fractional contribution of East Asia is higher than them.

415

Seasonally, the contribution of South Asia to surface BC in the TP is strongest in winter (55%) and weakest in summer (33%). South Asia is identified as the dominant contributor for 83% and 41% areas of the TP in winter and summer, respectively (Figure 4). In contrast, the contribution of East Asia is highest in summer (45%) and lower in winter (23%). East Asia is identified as the dominant contributor for 34% and 11% areas of the TP respectively in summer and winter (Figure 4). These seasonal variations in the contributions of South Asia and East Asia are generally in agreement with earlier studies (Lu et al., 2012; Zhang et al., 2015; Yang et al., 2018) with various numerical methods. For instance, using the emission perturbation method, Yang et al. (2018) suggested that the contribution of South Asia is 61% in non-monsoon season

425

(October-April) and 19% in monsoon season (May-September).

We further divided the TP into five subregions, namely, eastern TP, southern TP, western TP, northern TP, and central TP (Figure 4a). The 20-year means in the subregions show different BC levels, seasonalities, and dominant BC source regions (Figures 4-6). Over the eastern TP, the mean surface BC concentrations are the highest among the five subregions (Figure 5b). Above 70% of surface BC in the eastern TP is transported from East Asia (Figure 6b).

In the southern TP, the mean surface BC concentrations are the 2nd highest among the five subregions, which are high in spring and low in the other seasons (Figure 5c). Such seasonality is likely resulted from the high fire emissions over South Asia in spring, the favorable atmospheric circulation for BC transport to the southern TP in spring, and the strong wet deposition of BC by the monsoon precipitation in summer (Chen et al., 2018). This seasonality is in consistency with observations in literature (Marinoni et al., 2010; Cong et al., 2015). South Asia is the dominant source region for surface BC in the southern TP year-round, with fractional contributions of over 73% (Figure 6c). The dominant contribution of South Asia to the southern TP was also suggested in previous studies (He et al., 2014a; Zhang et al., 2015; Yang et al., 2018). The second dominant source region for the southern TP is Southeast Asia, which contributes 13% BC to surface BC in the region in the annual mean.

Over the western TP, the mean BC concentrations are the 3rd lowest among the five subregions, with a seasonality of high BC in winter and spring and low BC in summer and autumn (Figure 5d). The higher values in spring and winter agree with the BC measurements at sites in the western Himalayas (Nair et al., 2013). BC transport from South Asia contributes to 82% of surface BC in winter and 70% in

summer (Figure 6d). Such seasonality with winter high and summer low in the
455 fractional contribution of South Asia to surface BC over the western TP were also
suggested by Zhang et al. (2015).

In the northern TP, the mean BC concentrations are the 2nd lowest among the five
subregions, which are at maximum in winter and minimum in spring (Figure 5e). This
460 seasonality is different from an observational study, which reported that, over the
Qilian Shan in the northern TP, surface BC concentrations are highest in summer and
lowest in autumn (Zhao et al., 2012). The dominant source region for surface BC over
the northern TP is central Asia in spring and autumn, East Asia in summer, and South
Asia in winter (Figure 6e). Influenced by the prevailing westerlies, central Asia
465 contribute to surface BC in the western (Figure 6d) and northern TP (Figure 6e) more
than to other TP subregions. In the annual mean, central Asia can contribute to 9%
and 27% surface BC in the western (Figure 6d) and northern (Figure 6e) TP,
respectively.

470 Among the five subregions, the central TP is with the lowest BC concentrations
(Figure 5f). Seasonally, BC concentrations over the central TP is higher in spring and
winter than in summer and autumn. This simulated seasonality is different from the
observed one at a site in the central TP reported in Wang et al. (2016), who showed
that BC concentrations are higher in spring and lower in winter from November 2012
475 to June 2013. South Asia is the dominant source region for surface BC in the central
TP and its influence is strongest in winter, contributing to 77% of surface BC in this
subregion. In contrast, the influence of East Asia is strongest in summer, contributing
to 34% of surface BC there.

480 The local contribution of the TP and the contribution of Southeast Asia are much
lower than those of South Asia and East Asia (Figure 6). In the annual mean, local

contribution to surface BC in the TP is approximately 11% (Figure 6a). Among all the TP subregions, the fraction in the local contribution is highest in the central TP, being ~14% in the annual mean (Figure 6f). Using a tagging tracer method, Zhang et al. (2015) also suggested that local emissions contribute about 10% of BC in the TP. Influenced by the prevailing westerlies, BC from Southeast Asia cannot be efficiently transported to the TP by atmospheric circulation. The annual mean contribution of Southeast Asia to surface BC in the TP is ~7% (Figure 6a). Seasonally, the contribution of Southeast Asia is highest in spring (10%) (Figure 6a) when fire BC emissions are highest there. Geographically, the contribution of Southeast Asia is highest in the southern TP (13% in the annual mean) (Figure 6c) because of its proximity to Southeast Asia.

4 Interannual variation in surface black carbon over the Tibetan Plateau

4.1 Influences of biomass burning on the interannual variation in surface black carbon over the Tibetan Plateau

Figure 7 shows the anomalies of surface BC concentrations averaged over the TP from the three GEOS-Chem simulations. Note that in these simulations, only natural influences of meteorological and biomass burning in the interannual variation of surface BC in the TP are considered. The simulations from CTRL and FixBB are significantly correlated with each other in all the seasons ($r=0.45$, $p<0.05$ in spring, $r>0.8$, $p<0.05$ in the other three seasons), indicating the important role of meteorology in the interannual variation in surface BC concentrations in the TP. Remarkably, in spring (Figure 7a), the correlation coefficient of BC anomalies between CTRL and FixMet simulations reaches 0.8786 ($p<0.05$), indicating the importance of biomass burning to the interannual variation in BC in spring. The largest anomaly of BC concentrations from CTRL simulation is in 1999. The comparison between CTRL and FixMet simulations suggests that this strong anomaly is largely explained by biomass burning. Even if we exclude the extreme year 1999, the correlation ($r=0.77$, $p<0.05$)

510 between CTRL and FixMet simulations remains significant, indicating the strong influence of biomass burning on the variation in springtime surface BC from year to year. In Supplement (Figure [S3S4](#)), we applied the mean absolute deviation and absolute percent departure from the mean to quantitatively represent the strength of the interannual variation in surface BC over the TP, similar to Mao and Liao (2016).
515 Figure [S3S4](#) further confirms that biomass burning is an important driver of the interannual variation in surface BC over the TP in spring, while meteorology is more important than biomass burning in the other seasons.

To further examine the influence of biomass burning in spring, we integrally
520 analyzed data from ATSR satellite fire counts, GFED3 fire emissions, and the GEOS-Chem simulations. Both ATSR and GFED3 data show that fires occur frequently over the Indo-Gangetic Plain, central India, and Southeast Asia (Figures 1 and 8). Fire activities in Asia are well described in the GFED3 inventory that is used in the GEOS-Chem simulations (Figures 8b and 8c). We found the interannual variation in BC
525 anomalies in the TP from CTRL simulation is significantly correlated to the total number of fire counts in the Indo-Gangetic Plain ($r=0.76$, $p<0.05$), and central India ($r=0.67$, $p<0.05$). The correlation is insignificant to the total number of fire counts in Southeast Asia ($r=0.19$, $p>0.05$). In spring of 1999, extreme fire activities occurred in the Indo-Gangetic Plain and central India (Figure 8b). Driven by the favorable
530 atmospheric circulation, the strong BC emissions from the extreme fire activities greatly enhanced surface BC concentrations in the TP (Figure 8d). In CTRL simulation, positive BC anomalies appear over the entire TP, with a regional mean of $0.08 \mu\text{g m}^{-3}$ or 31% relative to the 1995-2014 climatology (Figure 7a). Even when 1999 spring is removed from the time series, biomass burning still shows a large
535 impact on interannual surface BC over the TP in spring (Figure [S3S4](#)). Additionally, in winter, biomass burning was extremely strong in 1998 (Figure 7d), which enhanced the regional mean surface BC concentrations in the TP by $0.01 \mu\text{g m}^{-3}$ or 5% relative

to the climatology.

540 In the following, the important role of meteorology in modulating the interannual variation in surface BC over the TP is to be explored in sections 4.2 and 4.3 from a perspective of the influences of the Asian monsoon on BC transport to the TP.

545 **4.2 Influences of the Asian summer monsoon on the interannual variation in transport of black carbon to the Tibetan Plateau from nonlocal regions in summer**

The TP can largely impact the Asian monsoon system through thermal and dynamic processes (Wu et al., 2015). In the meantime, the Asian monsoon can significantly influence the transport of atmospheric species to the TP (Xu et al., 2014). In this section, we show the influences of two Asian monsoon subsystems on the interannual variation in BC transport to the TP in summer. We employed a unified dynamical monsoon index to represent the strength of East Asian summer monsoon (EASM) and South Asian summer monsoon (SASM). The index was proposed by Li and Zeng (2002) and it has been widely applied to quantify the impact of the Asian monsoon on air pollution in Asia (Mao et al., 2017; Lu et al., 2018). Using this index, Han et al. (2019) found a close correlation between the EASM and ozone transport from nonlocal regions to East Asia. The calculation of the index was introduced in Li and Zeng (2002) and Han et al. (2019). The index is respectively termed as EASM index (EASMI) and SASM index (SASMI), when it is applied to represent the strength of EASM and SASM. A higher EASMI indicates a stronger EASM and a higher SASMI indicates a stronger SASM.

Figure 9a shows the spatial distribution of the correlation between BC transport to the TP and wind at 850 hPa at each of the grids in summer. As known from Figure 3b, central China is a dominant source region for the TP in summer, accounting for

63% of the imported BC from East Asia to the TP surface. In Figure 9a, BC from central China correlates significantly with the zonal wind at 850 hPa in central China (Figure 9a), with a regional mean correlation coefficient of -0.55 ($p < 0.05$). Westward winds (negative in the zonal component of wind vector) over central China favor BC transport from East Asia to the TP. Furthermore, the EASMI also correlates negatively with the zonal wind at 850 hPa over central China (Figure 9b). When the EASM is stronger, the zonal wind in the monsoon circulation weakens over this region (Yang et al., 2014; Han et al., 2019), suggesting that westward winds may occur more frequently or with higher speed. Therefore, BC transport to the TP from central China is enhanced (Figure 9c), as a significantly positive correlation is found between the strength of the EASM and BC transport from central China to the TP surface ($r = 0.49$, $p < 0.05$) and between the strength of the EASM and BC transport from central China to the eastern TP surface ($r = 0.48$, $p < 0.05$). This is further confirmed by the differences in BC transport to the TP surface in summers between strong and weak EASM years (Figure 9d).

Figure 10 shows how the SASM impacts the BC transport from South Asia over the TP surface in summer. Serving as a heat source in the Asian summer monsoon system, the TP promotes strong convection and modulates the meridional circulation (Xu et al., 2014). Driven by the meridional circulation, BC in South Asia can be transported northward and upward to the TP (Figure 3b). BC transport from northeastern South Asia to the TP accounts for 30% of the total BC transport from South Asia. Interannually, BC transport from northeastern South Asia is significantly correlated with the meridional wind at 500 hPa ($r = 0.65$, $p < 0.05$, Figure 10a), which is also closely correlated to the strength of the SASM (Figure 10b). In strong SASM years, an anomalous cyclone locates over the northern South Asia at 500 hPa and correspondingly the meridional wind over the northeastern South Asia is increased (Figure 10d). This well explains why the interannual variation in BC transport from

northeastern South Asia correlates positively with the strength of the SASM ($r=0.55$,
595 $p<0.05$ for the TP, $r=0.56$, $p<0.05$ for the STP, Figure 10c). Among all source regions,
the differences in BC transport from northeastern South Asia to the TP is largest in
summers between strong and weak EASM years (Figure 10d).

4.3 Influences of the Asian winter monsoon on the interannual variation in the 600 transport of black carbon to the Tibetan Plateau from nonlocal region in winter

The Asian winter monsoon is a predominant climate feature in Asia and an important
modulator of the distribution and transport of air pollutants (Mao et al., 2017; Zhu et
al., 2017). However, the impact of the Asian winter monsoon on the interannual
variation in BC transport to the TP is scantily studied. Here, we assess such impact
605 with two climate indices. We measure the intensity of East Asian winter monsoon
(EAWM) by an index defined by Jhun and Lee (2004). The EAWM index (EAWMI)
represents the EAWM intensity by the meridional wind shear associated with the jet
stream in the upper troposphere. It can be calculated by the difference in the regional
averaged zonal wind speed at 300 hPa between the areas 27.5-37.5°N, 110-170°E and
610 50-60°N, 80-140°E. Using the EAWMI, it is found that the EAWM is closely
correlated with the interannual variation in pollution transport over East Asia (Q. Li et
al., 2016; Han et al., 2019). Furthermore, the Siberian High is a key component of the
EAWM system (Wu and Wang, 2002) and its strength can be described using an index
defined by Wu and Wang (2002). This Siberian High index (SHI) can be calculated
615 from the regional mean sea level pressure over the area of the Siberian High (40-
60°N, 80-120°E). The EAWMI and SHI are highly correlated ($r=0.72$, $p<0.05$).

Figure 11 illustrates a connection between the EAWM and BC transport from
East Asia to the TP surface. We mainly focused on BC transport from central China,
620 as this area contributes to 54% of the total BC transport from East Asia to the TP
(Figure 3d). BC transport from central China to the TP surface layer correlates

significantly with the zonal wind at 850 hPa over central China ($r=-0.73$, $p<0.05$, Figure 11a). The zonal wind over this region is also correlated with the strength of the EAWM ($r = -0.5$, $p<0.05$) and the strength of the Siberian High ($r=-0.65$, $p<0.05$) (Figure 11b). A significant correlation ($r=0.59$, $p<0.05$) is found between the strength of the EAWM and BC transport from central China to the TP surface (Figure 11c). When the EAWM is stronger, the more frequent or stronger westward winds can enhance BC transport from central China to the TP (Figure 11d).

630 A connection between BC transport from South Asia to the TP surface and the Siberian High in winter is shown in Figure 12. BC over the northern South Asia can be transported efficiently to the TP by the prevailing subtropical westerlies. Northern South Asia contributes to 70% of BC transported from South Asia to the TP surface (Figure 3d). The contribution of northern South Asia to surface BC in the TP is significantly related ($r=0.72$, $p<0.05$) to the zonal wind at 500 hPa over the TP (Figure 12a). The westerlies over the TP are also correlated with the strength of Siberian High (Figure 12b). The elevated zonal wind in the middle troposphere over the TP in winters with strong Siberian High can enhance the BC transport from northern South Asia to the TP (Figure 12d). Significant correlations are found between the strength of the Siberian High and BC transport from northern South Asia to the TP surface (640 $r=0.66$, $p<0.05$) and between the strength of the Siberian High and BC transport from northern South Asia to the southern TP surface ($r=0.65$, $p<0.05$) (Figure 12c). In addition, the contribution of northern South Asia to surface BC in the western TP increases significantly with the meridional wind at 500 hPa over the western TP (645 $r=0.64$, $p<0.05$). The differences in BC transport from northern South Asia to the TP is largest in winter between strong and weak SHI years (Figure 12d).

5 Discussion

The findings in this study provide an enhanced understanding of the long-range

650 transport of BC to the TP. We comprehensively assessed the BC transport from
worldwide source regions to the TP. Our results reveal the source regions of surface
BC over the entire TP in the four seasons, which was investigated by limited studies
(Lu et al., 2012; Zhang et al., 2015). The influences of South Asia and East Asia on
the TP were noticed by previous studies. Most of them were focused on limited
655 locations (Cao et al., 2011; Engling et al., 2011; Chen et al., 2018) or in one or few
seasons (Zhao et al., 2017; Wang et al., 2018). Here, we further quantified the
influence of South Asia and East Asia over the entire TP in the four seasons, in terms
of both fractional contribution (Figure 6) and affected areas in the TP (Figure 4).
Moreover, we identified three key areas within South Asia and East Asia and found
660 that the contribution of BC from there to surface BC in the TP is highest among South
Asia and East Asia (Figure 3).

The estimate of the fractional contributions from various source regions to
surface BC in the TP from this study is generally comparable with that from literature
665 (Lu et al., 2012; Zhang et al., 2015; Yang et al., 2018). (1) The total contribution of
South Asia and East Asia to surface BC in the TP is estimated to be 77%, between
69% by Zhang et al. (2015) and 84% by Lu et al. (2012). (2) The contribution of BC
from South Asia is strongerlargest in winter and weakersmallest in summer, which
was also suggested by Lu et al. (2012), Zhang et al. (2015), and Yang et al. (2018).
670 Yang et al. (2018) modeled that the contribution of South Asia is 61% in non-
monsoon season (October-April) and 19% in monsoon season (May-September). (3)
The ~~estimate of~~ local contribution is estimated to be ~10%, comparable with that byin
Zhang et al. (2015), indicating the dominantpredominant role of BC transport from
nonlocal regions. Nevertheless, there are some disagreements between this and
675 previous studies in various extents. A noticeable disagreement is that this study
estimates that the annual mean contribution of East Asia is estimated to be
approximately 35% (Figure 6, also see Figure S5S6 for the estimated contribution at a

slightly higher level according to an emission perturbation simulation), while the estimates by Lu et al. (2012) and Zhang et al. (2015) are respectively 17% and 19%.

680 The ~~diserepaney~~discrepancies may be associated with the differences in region definitions and the estimation models.

Biomass burning is an important source for BC aerosols over the TP (Zhang et al., 2015). It was observed that BC emissions from biomass burning in South Asia
685 could be transported to the TP by the atmospheric circulation (Cong et al., 2015), and thus resulted in high BC episodes in the southern TP (Engling et al., 2011; Putero et al., 2014). Limited numerical studies assessed the impact of biomass burning on surface BC in the TP over a long-term period (Mao and Liao, 2016). Here, we demonstrated that biomass burning is an important driver of the interannual variation
690 in surface BC over the TP in spring (Figure 7). In particular, we found that there were extremely strong fire activities over the Indo-Gangetic Plain from 1998 winter to 1999 spring that largely enhanced surface BC concentrations over the entire TP (Figures 7 and 8). This extreme anomaly in fire activities and associated influence on BC over
the TP may have not been ~~fully~~ documented in such detail.

695
The Asian monsoon can influence the atmospheric circulation over the TP and its surroundings (Xu et al., 2014; Han et al., 2019). We found that the Asian monsoon system can significantly modulate the interannual variation in BC transport from South Asia and East Asia to the TP (Figures 9-12). We revealed that the EASM can
700 modulate the westward transport of BC from central China to the TP in summer (Figure 9), while previous studies mostly focused on the transport pathway build by the SASM (Zhao et al., 2017; Kang et al., 2019). In winter, this study shows significant influences of the EAWM on the BC transport from northern South Asia and central China to the TP (Figures 11 and 12). These results can also shed some
705 light on the transport mechanisms of other atmospheric species to the TP, such as

water vapour.

Regarding methodology, numerical simulations, including adjoint (Kopacz et al., 2011), backward-trajectory (Lu et al., 2012), tagging tracer (Zhang et al., 2015), and emission perturbation (Yang et al., 2018) simulations, have been used to identify BC transport from sources to the TP. Each of the methods has its advantages and limitations (Lu et al., 2012; Zhang et al., 2015). Modifying an approach proposed by Lu et al. (2012), we developed an efficient and stable method which shows strong performance in revealing the fractional contribution of BC transport from different source regions to the TP by season and year. We also run a set of emission perturbation simulations using GEOS-Chem to compare with the outcomes of the trajectory method being developed in this study. The emission perturbation method can estimate the contribution of a source region to BC concentrations in a receptor region (i.e. the TP in this study) by turning off BC emissions in that source region. The detailed description on our simulation and its outcomes are provided in Supplement (Figure [S5S6](#)).

Comparing the results between the backward-trajectory and the emission perturbation methods (Figure 6 vs. Figure [S5S6](#)), we can observe the following: (1) Both methods show that South Asia and East Asia are two dominant source regions for surface BC in the TP. The total contributions of South Asia and East Asia estimated by the two methods are comparable, which are respectively 77% (Figure 6a) and 82% (Figure [S5aS6a](#)). (2) The two methods suggest similar seasonality for the contributions of South Asia and East Asia. The contribution of South Asia to surface BC in the TP is stronger in winter and spring, while the contribution of East Asia is stronger in summer and autumn. (3) The influences of BC transport on different TP subregions estimated by the two methods are generally consistent. For example, both methods show that over 70% of surface BC in the eastern TP comes from East Asia,

and over 70% of surface BC in the southern TP comes from South Asia. Therefore,
735 both methods show comparable results, although there are some discrepancies in
various details.

Overall, the backward-trajectory method developed in this study can reasonably
quantify the relative contributions of different source regions to surface BC in the TP.

740 This method has following advantages. (1) The transport pathway of BC can be
visibly expressed (Figures 2 and 3). (2) The spatial ~~distribution of variation in~~ the
contribution of source regions can be showed explicitly (Figures 3 and 4). (3) It is
feasible for users of a chemical transport model to investigate the source-receptor
relationships if adjoint and tagged modes are unavailable to them. Note that this
745 method assumes a fixed lifetime of atmospheric BC (i.e. 7 days in this study), which
might lead to some uncertainties. The emission perturbation method does not require
this assumption. ~~It~~ The emission perturbation method is reliable and straightforward.
However, ~~the emission perturbation method it~~ cannot provide the spatial information
showed in Figures 2-4. It can ~~only~~ provide an overall assessment only for the total
750 contribution from each of the source regions defined, ~~which~~. The number of these
regions are often limited because of constraints of computation expenses.

This study is subject to some limitations. Numerical simulations have advantages
of covering large areas over the entire TP and long periods, such as 20 years in this
755 study. However, according to this and earlier studies, there are discrepancies between
observations and simulations from GEOS-Chem. He et al. (2014b) suggested that BC
aerosols over the TP may be underestimated by GEOS-Chem. Simulations in this
study have a weaker seasonal variation in BC concentrations than observations from
previous studies (Figure S1). All of these imply uncertainty in simulating the absolute
760 BC concentrations over the TP by GEOS-Chem. Furthermore, the simulations are
with a resolution of 2° latitude by 2° longitude. Such a resolution may not fully

capture processes in the sub-grid scale, such as the mountain-valley wind (Cong et al., 2015). Using a regional model, Zhang et al. (2020) demonstrated that compared with simulations with lower resolution, simulations with higher resolution can better
765 resolve the effects of topography and consequently yield stronger transport flux of BC from South Asia to the TP. Therefore, our results may somewhat underestimate the contribution of South Asia because of the model resolution. Using regional models at higher resolutions in the future can better describe the terrain effect in the TP.

770 In addition, anthropogenic BC emission inventories used in GEOS-Chem simulations may add uncertainties in our estimates. Figure [S6S7](#) compares the anthropogenic emissions in 2000 used in this study and those in 2010 from the Task Force on Hemispheric Transport of Air Pollution Phase 2 (TF HTAP2). It shows that anthropogenic BC emissions in most areas of South Asia have substantially increased
775 from 2000 to 2010 (Figure [S6S7](#)). On regional mean, anthropogenic BC emissions in East Asia have also increased from 2000 to 2010 (Figure [S6S7](#)). Inside East Asia, anthropogenic BC emissions in most regions of eastern China have increased, while decreased in Korea and Japan from 2000 to 2010 (Figure [S6S7](#)). Therefore, our results may underestimate the BC transport from South Asia and East Asia to the TP
780 in 2010. After 2010, anthropogenic BC emissions have changed as well (Zheng et al., 2018). Zheng et al. (2018) reported that anthropogenic BC emissions in China have reduced 27% from 2010 to 2017. BC emissions from biomass burning used in this study cover 1997-2011 and fire emissions in 2011 are used for simulation years after 2011. However, fire activities in South Asia have strong interannual variations and
785 have been significantly increased from 2001 to 2016 (Earl et al., 2018). Therefore, fire emissions used here may lead to biases in the BC simulations after 2011. This study is focused on natural drivers (biomass burning and meteorology) that are connective to the interannual variation in BC over the TP. The impact of interannual anthropogenic emissions warrants further studies.

6 Conclusions

Using a global chemical transport model, GEOS-Chem, we characterized the variation in surface BC over the TP in 20 years from 1995 to 2014. Applying an approach that combines the BC ~~simulations~~concentrations from GEOS-Chem and backward trajectories from HYSPLIT, we identified the source regions for surface BC in the TP and demonstrated the influences of atmospheric transport and biomass burning on the interannual variation in surface BC over the TP. The major conclusions are drawn as follows.

800 Based on the 20-year mean, surface BC concentrations are higher in the southern and eastern TP than in the other TP subregions. Surface BC in the TP is mainly influenced by two source regions: East Asia and South Asia. The two regions totally contribute 77% surface BC in the TP. The influence of East Asia is dominant in summer while the influence of South Asia is dominant in winter. By subregion, 805 surface BC in the southern and eastern TP comes mainly from South Asia and East Asia, respectively. Over the western and central TP, surface BC comes mainly from South Asia. Over the northern TP, the ~~most important~~dominate source region is central Asia in spring and autumn, East Asia in summer, and South Asia in winter.

810 Interannually, from 1995 to 2014, biomass burning can explain over ~~75~~74% of the variation in springtime surface BC concentrations over the TP if biomass burning and meteorology are both considered in GEOS-Chem simulations. Springtime surface BC in the TP is significantly correlated to the total number of fire counts from ATSR satellite data at 1 km resolution over the Indo-Gangetic Plain in South Asia ($r=0.76$, 815 $p<0.05$). In the spring of 1999, the extremely strong biomass burning in South Asia largely elevated surface BC concentrations ($0.08 \mu\text{g m}^{-3}$ or 31% relative to the climatology) over the TP. The strong biomass burning in South Asia in the winter of

1998 also enhanced BC concentrations over the TP. The implication of such large disturbances to climate in the TP and the relevant regions is yet to be explored.

820

The interannual variation in surface BC over the TP are greatly influenced by meteorology. Specifically, the Asian monsoon system alters the long-range transport of BC to the TP by modulating the atmospheric circulation. In summer, when the EASM is stronger than the normal, the more frequent or stronger westward wind in the lower troposphere can enhance BC transport from central China to the TP. When the SASM is stronger, the increased meridional wind over the northeastern South Asia in the middle troposphere can enhance BC transport from northeastern South Asia to the TP. In winter, when the EAWM is stronger than the normal, the reduced zonal wind in the lower troposphere tends to increase BC transport from central China to the TP. In winters with a stronger Siberian High, the enhanced zonal wind in the middle troposphere over the TP tends to carry more BC from northern South Asia to the TP.

825

830

Data availability

The GEOE-Chem model is publicly available at <http://geos-chem.org>. The HYSPLIT model can be acquired from http://www.arl.noaa.gov/HYSPLIT_info.php. The meteorological and fire data were download from <https://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis.surface.html> and <https://earth.esa.int/web/guest/home>, respectively.

835

840

Author contributions

JL, YW, ~~and HH~~, and TZ designed the research. HH and YW performed the study under supervision of JL ~~and TZ~~. HH, YW, YL, HC, YZ analyzed the data. HH, YW, and JL wrote the paper with valuable input from TZ, BZ, HW, HL, QW, SL, TW, MX, and ML.

845

Competing interests

The authors declare no conflict of interest.

Acknowledgments

850 We are grateful to the following model and data providers. The GEOE-Chem model is developed and managed by the Atmospheric Chemistry Modeling Group at Harvard University. The HYSPLIT model is developed by NOAA Air Resources Laboratory. The meteorological and fire data were respectively acquired from NOAA Earth System Research Laboratory and European Space Agency. We thank the anonymous
855 reviewers for their constructive comments and suggestions.

Financial support

This research is supported by the Chinese Ministry of Science and Technology National Key R&D Program of China (grant nos. 2019YFA0606803 and
860 2016YFA0600204) and the Natural Science Foundation of China (grant nos. 91744209, 91544230, and 41375140).