

Answer to the reviewers :

We are grateful to the reviewers for very detailed comments that help clarify several aspects of the manuscript. Reviewers' comments are highlighted in red, while our response follows in black.

Reviewer 2

Minor suggestion for revision:

\Psi is defined as rimed fraction in line 92; it would be helpful to repeat this definition when the values of \Psi are discussed in lines 228 ff.

Ψ is now defined as rimed fraction also in line 231 in the revised text.

Reviewer 3.

General Comments

The authors have done a very nice job of pulling material into the main manuscript or otherwise clarifying statements, measurements and modeling parameters in responding to my original comments. I have just a few suggestions in hopes to make crystal clear what advances are made in this paper versus what pieces of the puzzle remain for investigation. In particular, the paper still somewhat dances around the point that primary ice nucleation based on current information appears not capable of providing the requisite starting ice concentrations for the BR process to then provide the critical secondary mechanism needed to explain ultimate ice concentrations achieved in pockets of clouds over the region. A qualitative word, “meaningful” is used to describe ice concentrations of relevance, and yet, what has been measured so far in the region of interest is then not “meaningful.” Just a few word changes will clarify this point.

Specific Comments

1) Abstract, line 36: I suggest “necessary and sufficient”. This seems to me to be true. Sufficient alone implies to one that this is not difficult to explain. I understand that some possible explanations have been provided, but there remain significant efforts to prove any such points. I do not mean to detract from the significant results of the paper, only to emphasize mildly that this is a missing piece.

“necessary and sufficient” has now been added to the abstract

2) Section 4.4, lines 399-404: I believe that it is not an issue of calibration or even potentially missing INP sources, as this presently suggests. I can suggest the emphasized words in “None of the utilized primary ice nucleation parameterizations were **developed or are likely representative for...**”. The following sentence appears

irrelevant and should be removed. The pristine or limited nature of the INP populations in the region is the important point, and no new source is likely to explain any missing populations. If trying to say that new parameterizations will be needed to accurately represent sources, that would be fine, but it is not articulated that way in this revised wording.

It is now clarified that none of the utilized primary ice nucleation parameterizations **'are likely representative for'** the pristine conditions encountered over the high-latitude Southern Ocean. The following sentence has been removed, as suggested by the reviewer, to avoid confusion

3) Paragraph starting on line 440: This is excellent. One minor note on line 437. Terrestrial INPs are “assumed to be” or “expected to be” higher? The noted model study reference discusses contrasts between land and oceanic INP concentrations, but does not demonstrate that Antarctica is a specific elevated source of INPs.

It is now clarified that terrestrial INPs are “**expected to be**” higher

4) Conclusions, lines 455-457: Again the same point as above. Suggest (new words emphasized in bold and others crossed out) “...such conditions could likely be achieved through ice seeding (as likely happens in the examined case) or through INP transport from the Antarctic continent, where INP concentrations are generally **predicted to be** higher (Vergara-Temprado et al., 2018). **These points remain for future confirmation.**”

Rephrased exactly as suggested by the reviewer

5) Appendix A, line 500: Suggest to replace “have been calibrated” with “were not developed to be representative of”. Also, add “parameterizations” after Cooper and DeMott.

The sentence has been rephrased: 'none of these schemes is likely accurate as they have not been developed for such pristine conditions' . The word "parameterizations" after Cooper and DeMott is now added in line 501

**Secondary ice production in summer clouds over the Antarctic coast:
an underappreciated process in atmospheric models**

Georgia Sotiropoulou^{1,2}, Étienne Vignon^{3,4}, Gillian Young⁵, Hugh Morrison^{6,7}, Sebastian J.
5 O'Shea⁸, Thomas Lachlan-Cope⁹, Alexis Berne³, Athanasios Nenes^{1,10}

¹Laboratory of Atmospheric Processes and their Impacts (LAPI), Ecole Polytechnique
Fédérale de Lausanne (EPFL), Lausanne, Switzerland

²Department of Meteorology, Stockholm University & Bolin Center for Climate Research,
10 Sweden

³Environmental Remote Sensing Laboratory (LTE), EPFL, Lausanne, Switzerland

⁴Laboratoire de Météorologie Dynamique (LMD), IPSL, Sorbonne Université/CNRS, UMR
8539, Paris, France

⁵School of Earth and Environment, University of Leeds, UK

15 ⁶National Center for Atmospheric Research, Boulder, CO, USA

⁷ARC Centre for Excellence in Climate System Science, University of New South Wales,
Sydney, Australia

⁸Centre for Atmospheric Science, University of Manchester, UK

⁹British Antarctic Survey, Cambridge, UK

20 ¹⁰ICE-HT, Foundation for Research and Technology Hellas (FORTH), Patras, Greece

Correspondence to: georgia.sotiropoulou@epfl.ch, athanasios.nenes@epfl.ch

Abstract

25 The correct representation of Antarctic clouds in atmospheric models is crucial for accurate
projections of the future Antarctic climate. This is particularly true for summer clouds which
play a critical role in the surface melting of the ice-shelves in the vicinity of Weddell Sea. The
pristine atmosphere over the Antarctic coast is characterized by low concentrations of Ice
Nucleating Particles (INPs), which often result in the formation of supercooled liquid clouds.
30 However, when ice formation occurs, the ice crystal number concentrations (ICNCs) are
substantially higher than those predicted by existing primary ice nucleation parameterizations.
The rime-splintering mechanism, thought to be the dominant secondary ice production (SIP)
mechanism at temperatures between -8 and -3°C, is also weak in the Weather and Research
Forecasting model. Including a parameterization for SIP due to break-up (BR) from collisions

35 between ice particles improves ICNC representation in the modeled mixed-phase clouds,
suggesting that BR could account for the enhanced ICNCs often found in Antarctic clouds.
The model results indicate that a minimum concentration of about 0.1 L^{-1} primary ice crystals
is **necessary and** sufficient to initiate significant break-up, while there is little sensitivity to
40 increasing INPs. The BR mechanism is currently not represented in most weather prediction
and climate models; including this process can have a significant impact on the Antarctic
radiation budget.

1. Introduction

Predictions of Antarctic climate are hampered by the poor representation of mixed-phase
clouds over the Southern Ocean and the Antarctic Seas (Haynes et al., 2011; Flato et al.,
45 2013; Bodas-Salcedo et al., 2014; Hyder et al., 2018). Model simulations reveal significant
discrepancies in the Antarctic surface radiation budget, associated with cloud biases that are
driven by errors in the representation of the cloud microphysical structure (Lawson and
Gettelman, 2014; King et al., 2015; Listowski and Lachlan-Cope, 2017). A correct
representation of the cloud radiative impacts largely depends on the parameterization of cloud
50 microphysical processes (Listowski and Lachlan-Cope, 2017; Hines et al., 2019; Young et al.,
2019) and precipitation (Vignon et al., 2019), which determine the concentration and
characteristics of liquid drops and ice crystals.

Ice crystals form at temperatures above -38°C through heterogeneous nucleation
(Pruppacher and Klett, 1997); this means that the presence of insoluble aerosols that act as ice
55 nucleating particles (INPs) is required. However, Antarctica and Southern Ocean are
relatively clean regions and INPs are sparse (McCluskey et al., 2018; Schmale et al., 2019;
Welti et al., 2020). Thus it is especially surprising that enhanced ice crystal number
concentrations (ICNCs) have been observed in Antarctic clouds (Lachlan-Cope et al., 2016;
O'Shea et al., 2017). Secondary ice processes are believed to magnify ICNCs in polar clouds
60 with important implications for the surface radiative balance (Young et al., 2019), yet the
underlying mechanisms remain highly uncertain (Field et al., 2017).

The only well-established SIP mechanism that has been extensively implemented in
weather forecast and climate models is rime-splintering (Hallett and Mossop, 1974), also
known as the Hallett-Mossop process (H-M), which refers to the production of ice splinters
65 after collisions of supercooled droplets with ice particles (Hallett and Mossop, 1974;
Heymsfield and Mossop, 1984). This process is effective only in a limited temperature range,
between -8 and -3°C , and requires the presence of supercooled liquid droplets both smaller

Georgia 5/11/2020 10:48

Comment [1]: Comment #1, reviewer 3

and larger than 13 μm and 24 μm , respectively (Mossop and Hallett, 1974; Choulaton et al., 1980). However, recent studies have shown that H-M cannot sufficiently explain the
70 enhanced ICNCs observed in both Arctic (Sotiropoulou et al., 2020) and Antarctic (Young et al., 2019) clouds. While some Antarctic studies (Vergara-Temprado et al., 2018; Young et al., 2019) suggest that the underestimation of ice multiplication in models might be related to uncertainties in the description of the H-M process, we argue that this is likely driven by the fact that almost no models include other SIP mechanisms.

75 Another SIP mechanism, identified in recent laboratory studies (Leisner et al., 2014; Lauber et al., 2018), is the generation of ice fragments from shattering of relatively large frozen drops. This process however, while very efficient in convective clouds (Korolev et al., 2019), has been found ineffective in polar regions (Fu et al., 2019; Sotiropoulou et al., 2020). This is in agreement with Lawson et al. (2017) and Sullivan et al. (2018a) who have shown
80 that drop-shattering occurs in clouds with a relatively warm cloud base.

Mechanical break-up (BR) of ice particles that collide with each other is another process that results in ice multiplication (Vardiman, 1978; Takahashi et al., 1995) and it operates over a wide temperature range with maximum efficiency around -15°C . Limited knowledge of the BR mechanism comes from few laboratory experiments (Vardiman, 1978;
85 Takahashi et al., 1995) and small-scale modeling (Fridlind et al., 2007; Yano and Phillips, 2011, 2016; Phillips et al., 2017a,b; Sullivan et al., 2018a; Sotiropoulou et al., 2020). To the authors knowledge only two attempts have been made to incorporate this process in mesoscale models (Sullivan et al., 2018b; Hoarau et al., 2018). Specifically, Hoarau et al. (2018) assumed a constant number of fragments (F_{BR}) generated per snow-graupel collision in
90 Meso-NH model, while Sullivan et al. (2018b) implemented a temperature-dependent relationship for F_{BR} in COSMO-ART for several types of collisions (e.g. crystal-graupel, graupel-hail, etc), based on the results of Takahashi et al. (1995). Phillips et al. (2017a) recently developed a physically-based description of F_{BR} , which is a function of collisional kinetic energy and accounts for the effect of the colliding particles' size and rimed fraction
95 (Ψ). While being more advanced than any other parameterization proposed for BR, this scheme has never been implemented in mesoscale models before; it has only been tested in small-scale models for convective clouds (Phillips et al., 2017b; Qu et al., 2020) and Arctic stratocumulus (Sotiropoulou et al., 2020).

100 Sotiropoulou et al. (2020) recently showed that the observed ICNCs in Arctic clouds within the H-M temperature zone can be explained only by the combination of BR with the H-M process, which results in a 10 to 20-fold enhancement of the primary ice crystals. Based

on their results, we postulate that BR may also play a critical role in Antarctic clouds and can potentially explain the discrepancy between the observed and modeled ICNCs in the region (Young et al., 2019). To test this hypothesis, we implement parameterizations of the BR process in the Morrison microphysics scheme (Morrison et al., 2005) (hereafter M05) in the Weather and Research Forecasting (WRF) model V4.0.1 and examine their influence on the Antarctic clouds observed during the Microphysics of Antarctic Clouds (MAC) field campaign (O’Shea et al., 2017; Young et al., 2019).

110 **2. Observations**

2.1. MAC Instrumentation

The MAC field campaign was conducted in November–December 2015 over coastal Antarctica and the Weddell Sea, with the aim to offer detailed measurements of the microphysical and aerosol properties of the coastal Antarctic atmosphere. MAC included an extensive suite of airborne and ground-based instruments, a detailed description of which can be found in O’Shea et al. (2017). Here we only offer a brief recap of the instrumentation used in this study.

Cloud particle size distributions were derived using the images from a 2D Stereo (2DS, SPEC Inc., USA; Lawson et al., 2006) probe with a nominal size range from 10 to 1280 μm (10 μm pixel resolution). Shattering effects at the probes’ inlet were corrected by applying “antishatter” tips (Korolev et al., 2011) and inter-arrival time (IAT) post analysis (Crosier et al., 2011). The 2DS is a single particle instrument, measuring all particles that pass through its sample volume, which depends on particle size and the data integration period. For example, at 300 μm , 1 count measured using over a 1-sec averaging window equals to a concentration of 0.27 L^{-1} ; the uncertainty due to counting statistics is 100%. Total uncertainty is even higher but cannot be quantified.

Aerosol particle measurements of sizes 0.25 to 32 μm were made using the Grimm optical particle counter (GRIMM model 1.109), while a Cloud Aerosol Spectrometer (CAS, DMT; Baumgardner et al., 2001; Glen and Brooks, 2013) measured particles between 0.6 to 50 μm . Following the methodology of Young et al. (2019) and O’Shea et al. (2017), we only consider Grimm measurements of particles with diameter smaller than 1.6 μm in our analysis to avoid including data subject to inlet losses at larger particle sizes. Finally, the aircraft also included instrumentation to measure temperature, turbulence, humidity, radiation and surface temperature (King et al., 2008).

2.2 Case study

For our investigations we focus on the MAC case examined in Young et al. (2019), for which they showed that the H-M process, as currently parameterized in WRF, cannot explain the observed ICNCs. Young et al. (2019) utilized measurements from two MAC flights, M218 and M219, combined in one case study; both flights were conducted on 27 November 2015 over the Weddell Sea (Fig. 1): M218 between 15.3-16.7 UTC and M219 between 20.45-22.5 UTC. On that day, a low pressure system persisted over the eastern Weddell Sea, resulting in a southeasterly flow reaching the aircraft with air mass back trajectories from around the low pressure system, towards the Antarctic Peninsula and southern Patagonia (O'Shea et al., 2017).

The temperature and microphysical conditions encountered during these flights are representative of the MAC campaign (see Table 1 in O'Shea et al., 2017, and Fig. S6 in Young et al., 2019). Cloud measurements were collected mainly within the lowest 1.1 km above sea-level (a.s.l.) during both flights and within a temperature range of ~ -9 to -3°C . The sampled stratocumulus clouds were dominated by supercooled liquid drops, while ice formation occurred in isolated ice patches characterized by substantially enhanced ICNCs; the mean (max) ICNCs in these cloud regions were 1.16 (9.03) L^{-1} and 3.33 (87.31) L^{-1} for M218 and M219, respectively. The mean concentration of aerosols with sizes between 0.5 - $1.6 \mu\text{m}$ was 0.56 cm^{-3} and 0.41 cm^{-3} (cm^{-3} at standard temperature and pressure) for the two flights. Such low aerosol conditions and concurrent high ICNC concentrations within this temperature range are frequently found in West Antarctic Peninsula (Lachlan-Cope et al., 2016). Moreover, similar cloud droplet concentrations (N_{drop}) were measured during both flights (Young et al., 2019): the mean N_{drop} was 82.7 cm^{-3} for M218 and 100.4 cm^{-3} for M219, which are comparable with previous observations from the Antarctic Peninsula (Lachlan-Cope et al., 2016).

3. Modeling Methods

3.1. Model set-up

This study is conducted with the WRF model (Skamarock et al., 2008), version 4.0.1, by applying the same model set-up as in Young et al. (2019). Two domains with a respective horizontal resolution of 5 km and 1 km are used, where the inner one is two-way nested to the parent domain (Fig. 1). The polar stereographic projection is applied. The outer domain is

170 centered at 74.2°N, 30°E and includes 201×201 grid points, while the second domain
consists of 326×406 grids. Both domains have a high vertical resolution with 70 eta levels,
25 of which correspond to lowest 2 km of the atmosphere. The model top is set to 50 hPa. The
simulation period spans from 26 to 28 November 2014, 00:00 UTC, allowing for a 24-hour
spin up period before the day of interest (27 November). The model timestep is set to 30 (6)
175 sec for the outer (inner) domain, while output data are produced every 30 minutes.

Input data for the initial, lateral and boundary conditions for the simulations are
obtained from the European Centre for Medium-Range Weather Forecasting reanalysis (Dee
et al., 2011), as recommended by Bromwich et al. (2013). For both shortwave and longwave
radiation components, the RRTMG radiation scheme (Rapid Radiative Transfer Model for
180 GCMs) is applied. The Mellor-Yamada-Nakanishi-Niino (MYNN; Nakanishi and Niino,
2006) 2.5-level closure planetary boundary layer (PBL) and surface options are also
implemented, in combination with the Noah Land Surface Model (Noah LSM; Chen and
Dudhia, 2001), which includes a simplified thermodynamic sea-ice model. Given the short
run length, time-varying sea ice concentrations are not utilized. Young et al. (2019) used the
185 Polar WRF V3.6.1 to represent fractional sea-ice, a capability not available in standard WRF
V3.6. However, this option has been made available in the more recent V4.0.1 that we use in
this study. Following Young et al. (2019), the sea-ice albedo is set to 0.82, with a default
thickness of 3 m, and snow accumulation depth on sea ice is allowed to vary between 0.001 m
and 1.0 m.

190 A so-called ‘cumulus parameterization’ for shallow-convection subgrid processes is
not activated in both domains to ensure all cloud processes are represented by the grid-scale
microphysics scheme. Note that 5 km is a general upper limit for a convection-resolving
resolution (Klemp, 2006; Prein et al., 2015). Cloud microphysics are parameterized following
Morrison et al. (2005), hereafter M05. M05 performs well in reproducing Antarctic clouds,
195 resulting in improved representation of the liquid phase and thus the cloud radiative effects
compared to less advanced microphysical schemes (Listowski and Lachlan-Cope, 2017;
Hines et al., 2019). This bulk microphysics scheme predicts mixing ratios and number
concentrations for cloud ice, rain, snow and graupel species. While the mass mixing ratio of
cloud water is a prognostic variable, N_{drop} is constant parameter. The default value of the
200 scheme is 200 cm^{-3} ; here N_{drop} is set to 92 cm^{-3} , which is the mean value of M218 and M219
flight measurements (see Section 2.2).

3.2 Sensitivity Simulations

205 A detailed description of the ice formation processes in M05 and the implemented BR
parameterizations is offered in Appendix A and B, respectively. We assume that collisions
that include at least one large particle (thus ice-snow, ice-graupel and graupel-snow, snow-
snow and graupel-graupel) result in ice multiplication; contribution from collisions between
small ice particles (cloud ice) are neglected. In addition to the control (CNTRL) simulation,
210 which corresponds to the default set-up of M05 and accounts only for H-M, we perform seven
sensitivity simulations with varying description of F_{BR} . We also perform an additional
simulation as in CNTRL except with no H-M, and thus no SIP at all, which is referred as
NOSIP in the text.

In two sensitivity simulations with active break-up we assume, as in Hoarau et al.
(2018), a constant number of fragments generated per collision. This number is constrained
215 by in-situ measurements from the Arctic (Schwarzenboeck et al., 2009) which indicated that
one-branch ice-crystals are more common in polar clouds, resulting in ejection of a single
fragment after collision with another ice particle. However, this analysis (Schwarzenboeck et
al., 2009) included only dendritic crystals with size larger than 300 μm . Based on these results
we perform two simulations: FRAG1 assumes all collision types generate one fragment
220 without any size restrictions, while FRAG1siz allows for ice multiplication only if the particle
that undergoes fragmentation is larger than 300 μm . Note that because cloud ice with
characteristic diameter larger than 250 μm is converted to snow in the M05 scheme, collisions
that include cloud ice are assumed to not result in any multiplication in FRAG1siz.

The standard temperature-dependent formula of Takahashi et al. (1995) for F_{BR} ,
225 applied in Sullivan et al. (2018b), is tested here in the TAKAH simulation. However,
Takahashi et al. (1995) used 2-cm hailballs in their experiments, which is an unrealistic set-
up. For this reason we perform an additional simulation, TAKAHsc, in which this relationship
is further scaled with size (see Appendix B).

Finally, the Phillips parameterization is implemented in three simulations with varying
230 **rimed fraction** (Ψ) for the cloud ice/snow particles that undergo fragmentation; Ψ is not
predicted in most bulk microphysics scheme, including M05, and thus it is prescribed as a
constant. Note that F_{BR} is a function of Ψ only for the ice crystals or snowflakes that undergo
break-up, but not for graupel (Appendix B). Graupel is assumed to have $\Psi \geq 0.5$, while the
other ice types are characterized by lower rimed fraction. For this reason, we will consider
235 values of Ψ for cloud ice and snow between 0.2 (lightly rimed) and 0.4 (heavily rimed)
(Phillips et al., 2017a, b). These simulations are referred as PHIL0.2, PHIL0.3 and PHIL0.4 in
the text, where the number indicates the assumed values of Ψ for cloud ice and snow.

Georgia 5/11/2020 10:47

Comment [2]: Reviewer 2

4. Results

240 4.1 BR effect on microphysical properties

In Fig. 2a the modeled total ice number concentrations (cloud ice + snow + graupel, N_{isg}) derived for the region encompassing the 2 MAC flights (Fig. 1) are compared with measurements derived from the 2D Stereo (2DS) probe (see Section 2.1 for details). ICNCs in Fig. 2 are interpolated to match the time resolution of the temperature measurements. Then
245 cloud ice statistics are calculated for $N_{isg} > 0.005 \text{ L}^{-1}$, an indicator for the presence of an ice patch (O'Shea et al., 2017; Young et al., 2019). Moreover, since 2DS cannot resolve the shape (thus cloud phase) of particles smaller than $80 \mu\text{m}$, only modeled ice particles with sizes larger than this threshold are considered in Fig. 2, like in Young et al. (2019). While mean and maximum statistics are discussed below, additional statistical metrics (e.g. median
250 and interquartile range) are shown in Fig. S1 (Text S1).

The mean observed N_{isg} for the whole MAC campaign generally fluctuates between $0.5\text{--}4.5 \text{ L}^{-1}$. The variation in N_{isg} with temperature is somewhat larger for our case study (November 27), as maximum mean concentration goes up to $\sim 6.4 \text{ L}^{-1}$ at $T = -6.5^\circ\text{C}$. Consistently lower concentrations are observed for temperatures $\leq -7^\circ\text{C}$, but the temperature
255 statistics are poor for this temperature range as very few observations are available (Fig. 2a). The CNTRL simulation consistently underestimates the mean observations, producing mean $N_{isg} \sim 0.1 \text{ L}^{-1}$ over the examined temperature range (Fig. 2a). NOSIP produces similar results to CNTRL, suggesting that the H-M process included in default M05 (CNTRL) is hardly effective at all.

260 PHIL0.2 and PHIL0.3 also produce similar mean ICNCs to CNTRL (Fig. 2a, b), suggesting that lightly to moderately rimed ice particles do not contribute to ice multiplication through collisional break-up. The higher rimed fraction in PHIL0.4 results in very good agreement with mean observations (Fig. 2a), especially over the whole MAC campaign FRAG1siz also agrees well with mean observations, but when the size restrictions are ignored
265 (FRAG1) the model gives substantial ICNC overestimation. TAKAH simulation also produces unrealistically high mean N_{isg} , while TAKAHsc is in closer agreement with observations. The largest deviations from observations for TAKAHsc are observed for temperatures below -7°C , where no good measurement statistics are available (see discussion above).

270 Overall, CNTRL, PHIL0.2 and PHIL0.3 cannot reproduce the observed spectrum (Fig. 2b) and substantially underestimate the frequency of ICNCs larger than 1 L^{-1} . PHIL0.4,

FRAG1siz and TAKAHsc, however, can successfully reproduce the observed range of values (Fig. 2b), but the relative frequency remains somewhat underestimated. FRAG1 is in closest agreement with the observed spectrum, while TAKAH often overestimates the relative frequency (Fig. 2b). Maximum ICNCs in FRAG1 and TAKAH are 6403 and 2600 L⁻¹, respectively, which are about 70 and 30 times larger than the observed maximum value: 88 L⁻¹. This suggests that BR parameterizations that do not account for the impact of size are rather unrealistic. The maximum ICNCs in PHIL0.4, FRAG1siz and TAKAHsc are 174 L⁻¹, 150 L⁻¹ and 173 L⁻¹, which agree to within a factor of two with observations, while they are substantially underestimated in CNTRL (7.8 L⁻¹), PHIL0.2 (4.7 L⁻¹) and PHIL0.3 (5.2 L⁻¹).

Vertical distributions of cloud ice (N_i), graupel (N_g), snow (N_s) and total ICNC (N_{isg}) number concentration are examined in Fig. 3(a-d) for all simulations except those that produce unrealistically large concentrations (FRAG1 and TAKAH). The observed ICNCs are also shown in Fig. 3a and 3c. For consistency with M05, which converts all cloud ice particles with characteristic diameters larger than 250 μm to snow, the same threshold is adopted for splitting the observational dataset in these two ice categories. Graupel concentrations cannot be distinguished in the measurements (hence no ‘Nov 27’ profile in Fig. 3b); however, the model simulations that are in better agreement with observations (Fig. 2) suggest that these are negligible compared to cloud ice/snow concentrations. Graupel concentrations in Fig. 3b are shown for the whole size spectrum. In contrast, cloud ice (Fig. 3a), snow (Fig. 3c) and total ICNCs (Fig. 3d) include only particles with size larger than 80 μm for consistency with the observations shown in the same panel.

PHIL0.2 and PHIL0.3 produce slightly larger N_i (Fig. 3a) than CNTRL, but reduced N_g (Fig. 3b) values and similar or reduced N_s (Fig. 3c); these mean N_i and N_s profiles are orders of magnitude lower than the observed values. PHIL0.4, FRAG1siz and TAKAHsc produce similar N_i to the observations (Fig. 3a), while N_s is slightly underestimated (Fig. 3c). FRAG1siz is in somewhat better agreement with N_i observations than the other two simulations, especially at heights above 750 m a.s.l (Fig. 3a); this is also reflected in total ICNCs (Fig. 3d). Activating BR generally results in reduction of N_g (Fig. 3b). This decrease is larger than one order of magnitude in the three best performing simulations, compared to CNTRL, however we cannot assess which of these graupel profiles better represents reality. Nevertheless, we can overall conclude that PHIL0.4, FRAG1siz and TAKAHsc result in improved agreement of the vertical distribution of total ICNCs with observations compared to the rest of the simulations (Fig. 3d), including the default set-up of M05. Moreover, cloud ice concentrations (Fig. 3a) are comparable to snow concentrations (Fig. 3c) in these three

simulations, in agreement with observations. In contrast, simulations with deactivated or negligible BR result in substantially larger number of snow than cloud ice particles. This indicates that BR shifts the ice particle spectra to smaller sizes, which results in a more realistic representation of the ice microphysical characteristics.

310 The simulated liquid water content (LWC) is compared with CAS observations in Fig. 4. All simulations, except TAKAH, produce similar or slightly overestimated mean LWC at temperatures $\leq -3.5^{\circ}\text{C}$; at -3°C the mean observed values are higher (Fig. 4a). An overestimation of LWC in these runs is more evident in Fig. 4b; the observed spectrum does not include values larger than 0.5 g m^{-2} , while the simulated spectra are wider. An exception to this is TAKAH simulation, which underestimates mean LWCs and glaciates clouds at temperatures below -7°C (Fig. 4a), while it produces a narrower LWC spectrum compared to the observed (Fig. 4b). Apart from TAKAH, the remaining simulations produce similar liquid water properties with minor improvements in the runs with reduced LWC values, e.g. in FRAG1 (Fig. 4a). Nevertheless, while the produced range of LWC values in FRAG1 is somewhat closer to the observed, it still underestimates the relative frequency for most of the observed spectrum (Fig. 4b).

4.2 BR effect on surface radiation

325 To examine how deviations in ICNCs affect climate, mean radiative fluxes at the surface and at the top of the atmosphere (TOA) for all model simulations are presented in Table 1. Note that mass mixing ratio fields for all cloud species are provided from the microphysics to the radiation scheme, but no information on droplet and ice effective radius is exchanged.

330 Increasing BR multiplication has a pronounced impact on shortwave radiation, as it results in decreasing sunlight reflection and thus increasing downward surface radiation (SWD_{SFC}). Upward surface radiation (SWU_{SFC}) is a function of SWD and thus exhibits similar behaviour. This is due to the fact that increased BR efficiency (Fig. 2) results in decreased liquid water path (LWP) and cloud albedo. The difference between CNTRL and the simulations that improve ICNC representation (PHIL0.4, FRAG1siz and TAKAHsc) fluctuates between $11.9\text{-}25.7 \text{ W m}^{-2}$ for SWD_{SFC} and $6.7\text{-}12.4 \text{ W m}^{-2}$ for SWU_{SFC} (Table 1).

335 Cloud longwave radiative effects are mainly determined by cloud liquid properties, since liquid water is more opaque to longwave radiation than ice particles. However, no substantial differences in mean LWP are indicated for CNTRL (40.1 g m^{-2}), PHIL0.2 (33.2 g m^{-2}), PHIL0.3 (40.2 g m^{-2}), PHIL0.4 (29.1 g m^{-2}) and FRAG1siz (30.1 g m^{-2}), since LWP values fall within the black body emission range (Stephens, 1978). Optically thinner clouds

340 are produced in TAKAHsc (23.1 g m⁻²), and especially in FRAG1 (8.2 g m⁻²) and TAKAH
(3.2 g m⁻²) runs. Note that most simulations, including CNTRL, produce wider LWC spectra
than the observed by overestimating cloud liquid (Fig. 4b). Generally, decreasing liquid
content is in better agreement with observations (see section 4.1), suggesting that including
the BR process in M05 likely shifts the simulated LWPs towards more realistic values.
345 However, excessive ice multiplication, as in TAKAH, results in unrealistic liquid properties
(Fig. 4a) and thus errors in surface radiation.

Pronounced reduction in (LWD_{SFC}) is only found for the simulations FRAG1siz,
FRAG1 and TAKAHsc, which have a mean LWP well below 30 g m⁻², the lowest limit of the
black body emission range (Stephens, 1978). In all other simulations, the reduction in cloud
350 liquid due to BR is not large enough to alter the cloud emissivity significantly. The upward
longwave component (LWU_{SFC}) is only slightly affected in all simulations (<~ 1.3 W m⁻²).
Young et al. (2019) showed that underestimation of ICNCs results in significant positive and
negative biases in the surface Cloud Radiative Forcing (CRF) over the coastal areas; our
results agree with their findings, as CRF biases vary between -78 W m⁻² and +86 W m⁻² for
355 the most realistic simulations (Fig. S2, Text S2). Furthermore, the difference between
CNTRL and the realistic simulations in upward radiation flux at TOA (Table 1) is also more
pronounced for the shortwave component (SWU_{TOA}), fluctuating between 4.7-9.2 W m⁻², and
less significant for LWU_{TOA} (1.4-3.6 W m⁻²). Ultimately, both surface and TOA radiation
results indicate that a correct representation of SIP in the atmospheric models is critical for
360 the projection of the energy budget and thus for the future Antarctic climate.

4.3 Sensitivity to uncertainties in H-M description

To investigate the interactions between BR and H-M, we compare simulations in which the
H-M efficiency is either enhanced or turned off. Young et al. (2019) remove all liquid
365 thresholds from the H-M description, allowing for the process to become active over the
whole droplet spectrum. However, this change resulted in very weak ICNC enhancement in
their simulations. Here, we further remove all graupel/snow thresholds from H-M description
(Appendix A), which implies that there no size restrictions for the initiation of the process.
This modification is applied to CNTRL, PHIL0.3 and PHIL0.4 set-up, resulting in three
370 additional sensitivity tests: CNTRL_NOHRES, PHIL0.3_NOHRES and
PHIL0.4_NOHRES, respectively. Furthermore, in addition to NOSIP which corresponds to
CNTRL set-up but without H-M, another two simulations are performed with BR active but
H-M completely deactivated: PHIL0.3_NOHM and PHIL0.4_NOHM.

375 Mean ICNCs in CNTRL_NOTHRES are enhanced by on average a factor of three
compared to CNTRL (Fig. 5a). However this simulation underestimates concentrations at
temperatures larger than -7°C ; the mean observed value at this range is 2 L^{-1} , while the
simulated mean is 0.3 L^{-1} . Good agreement between CNTRL_NOTHRES and observations is
only achieved at temperatures $< -7^{\circ}\text{C}$, where statistical metrics for the two MAC cases are
poor (see Section 4.1). While PHIL0.3 did not result in any substantial multiplication, mean
380 ICNCs in PHIL0.3_NOTHRES are 5 times larger. The difference between PHIL0.4 and
PHIL0.4_NOTHRES is generally small for temperature warmer than -7°C , not exceeding a
factor of two, but it becomes substantially larger at colder temperatures. However, while the
95th percentiles for CNTRL_NOTHRES and PHIL0.3_NOTHRES are similar and more
comparable to observations, PHIL0.4_NOTHRES produces values larger than 10 L^{-1} at all
385 temperatures considered (Fig. 5b).

Excluding the temperature range ($< -7^{\circ}\text{C}$) that does not include substantial number of
measurements to evaluate model results, mean ICNC observations generally lay between
PHIL0.3_NOTHRES and PHIL0.4_NOTHRES in this set of simulations, while
CNTRL_NOTHRES produces somewhat lower values (Fig. 5a). However, this set-up
390 overestimates H-M efficiency, as it doesn't include any size limitations, which is not
consistent with current knowledge on the H-M mechanism derived from laboratory studies
(Hallet and Mossop, 1974; Choulaton et al., 1980). Nevertheless, the adapted thresholds are
ad-hoc, tuned for different conditions; these should be refined to get a more realistic H-M
effect in polar clouds.

395 Deactivating H-M completely does not substantially impact the results. This further
confirms the fact that the prescribed ad-hoc thresholds prevent the initiation of the process in
the studied conditions. Furthermore, it indicates that BR mechanism can explain the observed
ICNCs independently of whether H-M is active or not.

400 4.4 Sensitivity to uncertainties in primary ice formation

None of the utilized primary ice nucleation parameterizations are **likely representative** for the
pristine conditions encountered over the high-latitude Southern Ocean, thus it is likely that
primary ice formation is overestimated in this case. To examine how the uncertainty in
parameterizations for primary ice affects SIP efficiency, we perform two sets of simulations
405 by dividing or multiplying the efficiency of all primary ice production mechanisms
(immersion freezing, contact freezing and deposition/condensation-freezing nucleation) by a
factor of 10. Specifically, the first set with diminished ice nucleation includes

Georgia 5/11/2020 10:48

Comment [3]: Comment #2, reviewer 3

CNTRL_INP0.1, PHIL0.3_INP0.1 and PHIL0.4_INP0.1, while the second set with enhanced nucleation consists of CNTRL_INP10, PHIL0.3_INP10 and PHIL0.4_INP10.

410 Decreasing primary ice production by a factor of 10 inhibits BR multiplication (Fig. 6). Note that while maximum N_{isg} in PHIL0.3_INP0.1 and PHIL0.4_INP0.1 is 8.8 L^{-1} and 11 L^{-1} respectively, these simulations produce lower mean values (Fig. 6a) and N_{isg95} (Fig. 6b) than CNTRL_INP0.1 with maximum N_{isg} 5.7 L^{-1} . This is partly because a larger number of values in CNTRL_INP0.1 fall below the 0.005 L^{-1} threshold and are not included in the
415 presented mean statistics. Nevertheless, it is clear that the fewer primary ice crystals in all these sensitivity tests result in decreased frequency of ice-ice particle collisions, which is not sufficient to initiate significant BR multiplication.

Increasing primary ice production by an order of magnitude in CNTRL_INP10 still results in underestimated ice concentrations than observed, providing additional evidence for
420 the significant role of SIP in these conditions. The increased concentration of primary ice crystals enhances BR efficiency in PHIL0.3_INP10 compared to PHIL0.3, however the produced mean concentrations still are lower than the observed; N_{isg95} in PHIL0.3_INP10 only slightly exceeds unity (Fig. 6b). In contrast, PHIL0.4_INP10 produces similar mean ICNCs with PHIL0.4. N_{isg95} is also similar at warmer temperatures between two simulations,
425 while larger deviations are observed at temperatures $\leq -7^\circ\text{C}$ (Fig. 6b). This suggests that increasing concentrations of available ice particles tend to decrease the efficiency of BR mechanism, hence possible overestimations in primary ice production are likely offset by decreasing production of secondary ice.

In summary, the above results indicate that BR cannot be initiated when the available
430 primary ice concentrations are substantially lower than 0.1 L^{-1} , which is the mean primary ICNCs produced in NOSIP simulation (Fig. 2a). Yet, INPs over Southern Ocean are often substantially lower (McCluskey et al., 2018; Schmale et al., 2019; Welti et al., 2020). Ice seeding from clouds above the boundary layer was suggested by Young et al. (2019) as a key contributor to the primary ICNC levels for the studied case (see their Supporting
435 Information). Another process that can likely result in optimal conditions for BR to be initiated is aerosol transport from the Antarctic continent, where terrestrial INPs are **expected to be higher** (Vergara-Temprado et al., 2018). Moreover, a combination of these processes and the H-M mechanism, whose efficiency is substantially restricted in the current version of M05, might also provide the necessary concentrations to initiate BR; this was also the case for
440 Arctic stratocumulus conditions in Sotiropoulou et al. (2020). Understanding these interactions between different processes in the Antarctic region would likely provide insights

Georgia 5/11/2020 10:50

Comment [4]: Comment #3, reviewer 3

to the conditions that favor the development of isolated ice patches with substantially high ICNCs within predominantly supercooled liquid clouds. In higher INP conditions, which are likely less representative of the coastal Antarctic climate, the sensitivity of BR parameterization is expected to be lower.

5. Conclusions

Our results indicate that collisional break-up of ice crystals can explain observations of enhanced ICNCs in coastal Antarctic clouds, but this process requires the presence of $\sim 0.1 \text{ L}^{-1}$ primary ice crystals (as produced in NOSIP simulation) for initiation. This likely is a key threshold that can lead the development of isolated ice patches with enhanced ICNCs in predominantly supercooled liquid clouds (Grosvenor et al., 2012; O'Shea et al., 2017). Over the Southern Ocean, when INPs are generally sparse (McCluskey et al., 2018; Schmale et al., 2019; Welti et al., 2020), such conditions could likely be achieved through ice seeding (as likely happens in the examined case) or through INP transport from the Antarctic continent, where **INP concentrations are predicted to be higher (Vergara-Temprado et al., 2018). These points remain for future confirmation.**

Although BR has been observed in polar conditions before (Rangno and Hobbs, 2001; Schwarzenboeck et al., 2009), this mechanism is currently not implemented in most weather prediction and climate models. The more advanced Phillips et al. (2017a) parameterization produces realistic ICNCs in Antarctic clouds as long as a high rimed fraction is prescribed for the particles that undergo fracture, in agreement with Sotiropoulou et al. (2020). This indicates that our conclusions may not hold for winter clouds in the region, which contain less supercooled liquid water (Listowski et al., 2019) and are less prone to riming. However, for the studied case, a comparison of vapor deposition rates with riming rates (which include mass changes due to collisions with droplets/raindrops and due to contact/immersion freezing) for CNTRL simulation indicate that these two are on average comparable for cloud ice, while riming rates are substantially larger than vapor deposition rates for snow (not shown). These results suggest that prescribing a high rimed fraction for cloud ice and snow in M05 is not unreasonable; nevertheless Ψ in reality is highly variable for different temperature and microphysical conditions. More simplified parameterizations also produce improved results as long as the impact of the dependence of F_{BR} on the ice particle size is accounted for.

The very few existing BR descriptions in mesoscale models either do not account for size limitations (Sullivan et al., 2018b) or do not account for all collision types (Hoarau et al., 2018), which limits their realism. Increasing ICNCs from BR alters significantly the radiative

Georgia 5/11/2020 10:50

Comment [5]: Comment #4, reviewer 3

effects of summer mixed-phase Antarctic clouds; these clouds play a critical role in the surface melting of ice-shelves in the vicinity of Weddell Sea (Gilbert et al., 2020) and thus their accurate microphysical representation in models is of great importance.

480 **Appendix A: Ice formation processes in M05 scheme**

The standard M05 scheme includes three primary ice production mechanisms through heterogeneous nucleation (immersion freezing, contact freezing and deposition/condensation-freezing nucleation), and one SIP process (H-M).

Immersion freezing of cloud droplets and rain is based on the work of Bigg (1953).

485 This mechanism is active below -4°C and produces a raindrop freezing rate that depends on the degree of supercooling and the number concentration and volume of supercooled drops. The Meyers et al. (1992) description is used for contact freezing, also active below -4°C . The effective diffusivity of the contact nuclei to the drops are estimated from Brownian motion similar to Young (1974): $D_{ap} = R T (6 p r_i N_A m)^{-1} [1 + 0.0737 T (2880P)^{-1} r_i^{-1}]$, where R is the
490 universal gas constant, N_A is Avogadro's number, m is the dynamic viscosity of air, T is the temperature, P is the air pressure, and the radius of ice nuclei r_i is assumed to be 1×10^{-7} m. The factor in the brackets $[\]$ is a correction factor accounting for the mean free path of air molecules relative to the size of the ice nuclei (all units are MKS).

The default parameterization for deposition/condensation-freezing ice nucleation in
495 M05 is from Cooper (1986), which depends only on temperature and is active below -8°C in liquid saturated conditions or when ice supersaturation exceeds 8%. However, the aerosol-aware DeMott et al. (2010) parameterization for heterogeneous nucleation has been shown to compare better with in-cloud ice measurements over the Antarctic Peninsula than Cooper (Listowski and Lachlan-Cope, 2017), although none of these schemes is likely accurate as
500 they have not been developed for such pristine conditions. Nevertheless, both Cooper and DeMott parameterizations produce similar primary ice concentrations over the temperature range covered by the observations, but Cooper predicts more primary ice at lower temperatures ($<13^{\circ}\text{C}$), which might affect the representation of higher-altitude clouds (see Supporting Information in Young et al. 2019). For this reason, we apply the DeMott
505 description in our simulations, where the mean aerosol concentration of particles with sizes between $0.5\text{-}1.6 \mu\text{m}$ for the two flights (0.49scm^{-3}) is used as input (Young et al., 2019). Uncertainty to this formulation is investigated through a number of sensitivity tests (section 4.4).

The H-M parameterization, adapted from Reisner et al. (1998), is based on the

Georgia 5/11/2020 10:51

Comment [6]: Comment #5, reviewer 3

510 laboratory experiments conducted by Hallett and Mossop (1974), who found a maximum of
 ~350 splinters per milligram of rime generated around -5°C:

$$\frac{dN_{iHM}}{dt} = \rho \cdot 3.5 \cdot 10^8 f(T) \frac{dm_{rime}}{dt} \quad (1)$$

where dN_{iHM}/dt is the number of new fragments produced at a given timestep, $f(T)$ is the
 temperature-dependent efficiency of the process, ρ is the air density, and dm_{rime}/dt is the mass
 515 production rate of rime on snow or graupel due to accretion of cloud and rain drops. $f(T)$ is 0
 for $T < -8^\circ\text{C}$ and $T > -3^\circ\text{C}$, 1 for $T = -5^\circ\text{C}$, and increases linearly between these two extremes
 for $T \geq -8^\circ\text{C}$ and $T \leq -3^\circ\text{C}$.

Furthermore, for H-M to become activated in M05, two conditions must be met: (a)
 snow (or graupel) mass mixing ratios must be greater than 0.1 g kg^{-1} and (b) cloud liquid (or
 520 rain) water mass mixing ratios should be greater than 0.5 (or 0.1 g kg^{-1}). To achieve a good
 agreement between modeled and observed ICNCs for the simulated case, Young et al. (2019)
 had to remove condition (b) and multiply the H-M efficiency by a factor of 10.

Appendix B: Parameterizing collisional break-up in M05

525 There are three types of ice particles considered in the M05 scheme: small (cloud) ice, snow,
 and graupel. Ice multiplication is allowed after cloud ice-snow, cloud ice-graupel, graupel-
 snow, snow-snow and graupel-graupel collisions. The standard M05 scheme includes a
 description for collisions between cloud ice and snow to represent the accretion process,
 following the “continuous collection” approach:

$$530 \quad \frac{dN_{iAC}}{dt} = \frac{\pi}{4} \rho E_{col} \Gamma(b_s + 3) a_s \frac{N_i N_{0s}}{\lambda_s^{(b_s+3)}} \quad (2)$$

dN_{iAC}/dt is the rate of ice crystal number concentration collected by snow. N_{0s} and λ_s are the
 intercept and slope parameters of the snow size distribution, represented by an inverse
 exponential function, and Γ is the Euler gamma function. a_s and b_s are the characteristic
 parameters for snow in the fallspeed-diameter relationship (Morrison et al., 2005); a_s includes
 535 a density correction factor (Heymsfield et al., 2007). Note that the diameter (d_i) and terminal
 velocity (u_i) of cloud ice particles are considered much smaller than those of snow: $d_i \ll d_s$ and
 $u_i \ll u_s$, so that they are neglected in Eq. (2). E_{col} is the collection (sticking) efficiency between
 ice particles, set to 0.1; hence, it is assumed that only 10% of cloud ice particles that collide
 540 with snow are actually collected. We assume the remaining 90% of collisions result in ice
 particle break-up, hence the following relationship gives the rate of cloud ice-snow collisions
 that contribute to ice multiplication:

$$\frac{dN_{iis}}{dt} = \frac{\pi}{4} \rho (1 - E_{col}) \Gamma(b_s + 3) a_s \frac{N_i N_{0s}}{\lambda_s^{(b_s+3)}} \quad (3)$$

In the default M05, collisions between cloud ice and graupel particles are neglected as it is assumed that the collection efficiency of such collisions is negligible. To represent cloud ice-graupel collisions for ice multiplication, we use Eq. (3), but the size distribution and fallspeed parameters of snow are replaced by those for graupel. Moreover, since cloud ice is not collected by graupel particles, we assume that 100% of these collisions result in cloud ice break-up:

$$\frac{dN_{iig}}{dt} = \frac{\pi}{4} \rho \Gamma(b_g + 3) a_g \frac{N_i N_{0g}}{\lambda_g^{(b_g+3)}} \quad (4)$$

In the default M05 scheme, collisions between snow and graupel are also neglected because it is assumed that the collection efficiency for such collisions is negligible. For this study, graupel-snow collisions are treated using expressions similar to those for raindrop-snow collisions in M05. These are adapted from Ikawa and Saito (1991) and represent collisions between relatively large precipitation-sized particles:

$$\frac{dQ_{isg}}{dt} = \pi^2 \rho_s \rho \left| \Delta u_{m_{sg}} \right| \frac{N_{0s} N_{0g}}{\lambda_s^3} \left(\frac{5}{\lambda_s^3 \lambda_g} + \frac{2}{\lambda_s^2 \lambda_g^2} + \frac{0.5}{\lambda_s \lambda_g^3} \right) \quad (5)$$

$$\frac{dN_{isg}}{dt} = \frac{\pi}{2} \rho \left| \Delta u_{n_{sg}} \right| N_{0s} N_{0g} \left(\frac{1}{\lambda_s^3 \lambda_g} + \frac{1}{\lambda_s^2 \lambda_g^2} + \frac{1}{\lambda_s \lambda_g^3} \right) \quad (6)$$

$$\text{where } \left| \Delta u_{m_{sg}} \right| = \left((1.2u_{ms} - 0.95u_{mg})^2 + 0.08u_{mg}u_{ms} \right)^{1/2} \quad (7)$$

$$\text{and } \left| \Delta u_{n_{sg}} \right| = \left((1.7u_{ns} - u_{ng})^2 + 0.3u_{ng}u_{ns} \right)^{1/2} \quad (8)$$

dQ_{isg}/dt and dN_{isg}/dt represent the bulk rates that snow mass and number concentration collide with graupel and contribute to ice multiplication through fragmentation. Corrections in the mass (or number) -weighted difference in terminal velocity $\Delta u_{m_{sg}}$ (or $\Delta u_{n_{sg}}$) of the colliding particles (Eq. 7,8) are adapted from Mizuno (1990) and Reisner et al. (1998), to account for underestimates when $u_{ns} \approx u_{ng}$.

M05 also includes a description for collisions between snowflakes to represent snow aggregation, following Passarelli (1978):

$$\frac{dN_{sAG}}{dt} = \frac{1108a_s E_{col}}{4 \times 720} \pi^{\frac{1-b_s}{3}} \rho^{\frac{2+b_s}{3}} \rho_s^{\frac{-2-b_s}{3}} Q_s^{\frac{2+b_s}{3}} N_s^{\frac{4-b_s}{3}} \quad (9)$$

Based on this expression we parameterize the number of snow-snow collisions that contribute to ice multiplication as:

$$\frac{dN_{iss}}{dt} = \frac{1108a_s(1-E_{col})}{4 \times 720} \pi^{\frac{1-b_s}{3}} \rho^{\frac{2+b_s}{3}} \rho_s^{\frac{-2-b_s}{3}} Q_s^{\frac{2+b_s}{3}} N_s^{\frac{4-b_s}{3}} \quad (10)$$

Because snow aggregation does not result in any mass transfer, the snow mass involved in

these collisions is not calculated by the default M05 scheme. We obtain a description of dQ_{iss}/dt by applying the size distribution and fallspeed parameters of snow in the analytical solution for self-collection derived by Verlinde et al. (1990):

$$\frac{dQ_{iss}}{dt} = \frac{914\pi^2}{48\rho\rho_s} (1 - E_{col}) a_s d_s^{b_s+5} N_s^2 \quad (11)$$

575 To test the consistency of Eq. (10) and (11), which were derived using different methods, we repeated the CNTRL and PHIL0.4 simulations but with the Eq. (9) and (10) replaced by the analytical solution for the change in number concentration from self-collection derived by Verlinde and Cotton (1993). The sensitivity of the results to this modification was found to be insignificant.

580 Graupel-graupel collisions are also parameterized in a similar manner. Since there is no graupel aggregation (collection efficiency of such collisions is assumed to be negligible), 100% of the collisions are assumed to contribute to break-up:

$$\frac{dN_{igg}}{dt} = \frac{1108a_g}{4 \times 720} \pi^{\frac{1-b_g}{3}} \rho^{\frac{2+b_g}{3}} \rho_g^{\frac{-2-b_g}{3}} Q_g^{\frac{2+b_g}{3}} N_g^{\frac{4-b_g}{3}} \quad (12)$$

$$\frac{dQ_{igg}}{dt} = \frac{836\pi^2}{48\rho\rho_g} a_g d_g^{b_g+5} N_g^2 \quad (13)$$

585 The value 1108 in Eq. (10) is valid for $b_s=0.4$ (Passarelli, 1978); in M05 $b_s=0.41$ and $b_g=0.37$, thus adapting this value for both snow-snow (10) and graupel-graupel (12) collisions is a reasonable approximation.

Following the methodology of Sullivan et al. (2018b) in TAKAH simulation, the number of fragments generated due to ice-ice particle collisions (F_{BR}) is:

590
$$F_{BR} = 280 (T - 252)^{1.2} e^{-(T-252)/5} \quad (14)$$

However, Takahashi et al. (1995) used 2-cm hailballs in their experiments, thus to further include the influence of size in this formulation, we implement a size-scaled expression in TAKAHsc simulation, assuming that F_{BR} depends linearly on D , decreasing to 0 at $D = 0$:

$$F_{BR} = 280 (T - 252)^{1.2} e^{-(T-252)/5} \frac{D}{D_o} \quad (15)$$

595 where D (in meters) is the size of the ice particle that undergoes fracturing and $D_o=0.02$ m, the size of hailballs used by Takahashi et al. (1995).

The Phillips et al. (2017a) parameterization allows for varying treatment of F_{BR} depending on the ice crystal type and habit.

$$F_{BR} = \alpha A \left(1 - \exp \left\{ - \left[\frac{CK_o}{\alpha A} \right]^\psi \right\} \right) \quad (16)$$

600 where : $K_o = \frac{m_1 m_2}{m_1 + m_2} (\Delta u_{n_{12}})^2$,

$$\psi = 3.5 \times 10^{-3}$$

$$a = \pi D^2$$

605 m_1, m_2 are the masses of the colliding particles and Δu_{n12} is the difference in their terminal velocities. The correction applied in Eq. (8) is also adapted here to account for underestimates when $u_{n1} \approx u_{n2}$. D (in meters) is the size of the smaller ice particle which undergoes fracturing and a is its surface area. The parameterization was developed based on particles with diameters $500 \mu\text{m} < D < 5 \text{ mm}$, however Phillips et al. (2017a) suggest that it can be used for particle sizes outside the recommended range as long as the input variables to the scheme are set to the nearest limit of the range. C is the asperity-fragility coefficient and ψ is a correction term for the effects of sublimation based on the field observations by Vardiman (1978). For cloud ice-snow, cloud ice-graupel, snow-graupel and snow-snow collisions:

$$610 \quad A = 1.58 \cdot 10^7 (1 + 100\Psi^2) \left(1 + \frac{1.33 \cdot 10^{-4}}{D^{1.5}}\right),$$

$$\gamma = 0.5 - 0.25\Psi,$$

$$C = 7.08 \times 10^6 \psi$$

The above parameters adapted from Phillips et al. (2017a) concern planar crystals or snow with rimed fraction $\Psi < 0.5$ that undergo fracturing: $\Psi \leq 0.2$ corresponds to lightly rimed particles, while $\Psi \approx 0.4$ represents highly rimed crystals/snow. The choice of the ice habit is based on particle images collected during the MAC flights, which indicate the presence of needles and planar particles (O'Shea et al., 2017); needles are often considered secondary ice (Field et al., 2017). However, the M05 scheme does not explicitly consider habit and assumes spherical particles for all processes except sedimentation, for which the fallspeed-diameter relationships are for non-spherical ice.

620 For graupel-graupel collisions the parameters implemented in Eq. (16) are somewhat different (Phillips et al., 2017a):

$$A = \frac{a_0}{3} + \max\left(\frac{2a_0}{3} - \frac{a_0}{9} |T - 258|, 0\right)$$

$$\gamma = 0.3,$$

$$C = 6.3 \times 10^6 \psi$$

625 Finally, an upper limit for the number of fragments produced per collision is imposed, set to $F_{BRmax} = 100$; this is the same for all collision types (Phillips et al., 2017a).

We estimate the production rate of fragments for cloud ice-snow collisions and cloud ice -graupel collisions using Eq. (3) or (4) and one of the proposed formulations for F_{BR} above: $\frac{dN_{is}}{dt} F_{BR}$ and $\frac{dN_{ig}}{dt} F_{BR}$. For both of these collision types we assume that the cloud ice particles undergo break-up and the new smaller ice fragments remain within the same ice

630 particle category. For snow-graupel collisions, where the snow particle is assumed to undergo
fracturing, the production term $\frac{dN_{isg}}{dt} F_{BR}$ is added to the cloud ice category. In this case mass
transfer from the snow to the cloud ice category also occurs, but according to Phillips et al.
(2017a) this is only 0.1% of the snow mass that collides with graupel (5). Snow-snow and
graupel-graupel collisions are handled in the same way: $\frac{dN_{iss}}{dt} F_{BR}$ and $\frac{dN_{igg}}{dt} F_{BR}$ are added to
635 the cloud ice number equation, while 0.1% of $\frac{dQ_{iss}}{dt}$ (11) and $\frac{dQ_{igg}}{dt}$ (13) is added to the
corresponding mass equation.

Code and data availability: MAC data are available at
<https://catalogue.ceda.ac.uk/uuid/da17dab196f74d64af3ccbc35624027b>. The modified
640 Morrison scheme is available upon request

Competing interests: The authors declare that they have no conflict of interest.

Author contribution: GS and AN conceived and led this study. EV helped with the model
645 configuration and set-up, and provided Fig. 1. GY provided the observations and the model
set-up for the MAC case. SJO post-processed MAC data. GS implemented the BR
parameterizations, performed the WRF simulations, analyzed the results and, together with
AN, led the manuscript writing. All authors contributed to the scientific interpretation,
discussion and writing of the manuscript.

650 **Acknowledgements:** GS and AN acknowledge support from Laboratory of Atmospheric
Processes and Their Impacts (LAPI) at the École Polytechnique Fédérale de Lausanne (EPFL)
the project IC-IRIM (project ID 2018-01760) funded by the Swedish Research Council for
Sustainable Development (FORMAS), the project PyroTRACH (ERC-2016-COG) funded
655 from H2020-EU.1.1. - Excellent Science - European Research Council (project ID 726165)
and the project FORCeS funded from Horizon H2020-EU.3.5.1. (project ID 821205). EV and
AB acknowledge the financial support from EPFL-ENAC through the LOSUMEA project.
The National Center for Atmospheric Research is sponsored by the U.S. National Science
Foundation. GY acknowledges support from the UK Natural Environment Research Council
660 (grant no.: NE/R009686/1). We are also grateful to MAC scientific crew for the observational
datasets used in this study.

References:

- 665 Baumgardner, D., Jonsson, H., Dawson, W., O'Connor, D., and R. Newton: The cloud, aerosol
and precipitation spectrometer: A new instrument for cloud investigations. *Atmos. Res.*, 59,
251–264. [https://doi.org/10.1016/S0169-8095\(01\)00119-3](https://doi.org/10.1016/S0169-8095(01)00119-3), 2001
- 670 Bodas-Salcedo, A., Williams, K. D., Ringer, M. A., Beau, I., Cole, J. N. S., Dufresne, J.-L., et
al.: Origins of the solar radiation biases over the Southern Ocean in CFMIP2 Models, *J.*
Clim., 27, 41–56. <https://doi.org/10.1175/JCLI-D-13-00169.1>, 2014
- Bigg, E. K. : The formation of atmospheric ice crystals by the freezing of droplets. *Q. J. Roy.*
Meteorol. Soc., 79, 510–519. <https://doi.org/10.1002/qj.49707934207>, 1953
- 675 Bromwich, D. H., Otieno, F. O., Hines, K. M., Manning, K. W., and Shilo, E.: Comprehensive
evaluation of polar weather research and forecasting model performance in the Antarctic,
Journal of Geophysical Research: Atmospheres, 118, 274–292, doi:10.1029/2012jd018139,
<http://dx.doi.org/10.1029/2012JD018139>, 2013.
- 680 Brown, P. and Francis, P.: Improved measurements of the ice water content in cirrus using a
total-water probe, *J. Atmos. Ocean. Tech.*, 12, 410–414, 1995.
- Chen, F., and Dudhia, J.: Coupling an Advanced Land Surface–Hydrology Model with the
Penn State– NCAR MM5 Modeling System. Part I: Model Implementation and Sensitivity.
685 *Monthly Weather Rev.*, 129, 569–585, 2001.
- Choulaton, T. W., D. J. Griggs, B. Y. Humood, and Latham, J. : Laboratory studies of
riming, and its relation to ice splinter production. *Quart. J. Roy. Meteor. Soc.*, 106, 367–374,
doi:<https://doi.org/10.1002/qj.49710644809>, 1980.
690
- Crosier, J., Choulaton, T. W., Westbrook, C. D., Blyth, A. M., Bower, K. N., Connolly, P. J.,
Dearden, C., Gallagher, M. W., Cui, Z., and Nicol, J. C.: Microphysical properties of cold
frontal rainbands, *Q. J. Roy. Meteorol. Soc.*, 140, 1257–1268, doi:10.1002/qj.2206, 2013.
- 695 Cooper, W.A.: Ice initiation in natural clouds. *Meteorological Monographs*, 21, 29–32.

<https://doi.org/10.1175/0065-9401-21.43.29>, 1986

700 Dee, D.P., Uppala, S.M., Simmons, A.J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,
Balmaseda, M.A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A.C.M., van de Berg, L.,
Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L.,
Healy, S.B., Hersbach, H., Holm, E.V., Isaksen, I., Kalberg, P., Kohler, M., Matricardi, M.,
705 McNally, A.P., Monge-Sanz, B.M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P.,
Tavolato, C., Thepaut, J.-N., Vitart, F.: The ERA-Interim reanalysis: Configuration and
performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597,
<https://doi.org/10.1002/qj.828>, 2011.

DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H.,
Richardson, M. S., Eidhammer, T., and Rogers, D. C.: Predicting global atmospheric ice
nuclei distributions and their impacts on climate, *Proc. Nat. Acad. Sci.*,
710 [doi:10.1073/pnas.0910818107](https://doi.org/10.1073/pnas.0910818107), 2010.

Field, P., Lawson, P., Brown, G., Lloyd, C., Westbrook, D., Moisseev, A., Miltenberger, A.,
Nenes, A., Blyth, A., Choularton, T., Connolly, P., Bühl, J., Crosier, J., Cui, Z., Dearden, C.,
DeMott, P., Flossmann, A., Heymsfield, A., Huang, Y., Kalesse, H., Kanji, Z., Korolev, A.,
715 Kirchgaessner, A., Lasher-Trapp, S., Leisner, T., McFarquhar, G., Phillips, V., Stith, J., and
Sullivan, S.: Chapter 7: Secondary ice production - current state of the science and
recommendations for the future, *Meteor. Monogr.*, [doi:10.1175/AMSMONOGRAPHS-D-16-0014.1](https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0014.1), 2017.

720 Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S., Collins, W., Cox, P., Driouech,
F., Emori, S., Eyring, V., Forest, C., Gleckler, P., Guilyardi, E., Jakob, C., Kattsov, V.,
Reason, C., and Rummukainen, M.: Evaluation of Climate Models. In: *Climate Change 2013:
The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report
of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M.
725 Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)], 2013.

Fridlind, A. M., Ackerman, A. S., McFarquhar, G., Zhang, G., Poellot, M. R., DeMott, P. J.,
Prenni, A. J., and Heymsfield, A. J.: Ice properties of single-layer stratocumulus during the
Mixed-Phase Arctic Cloud Experiment: 2. Model results., *J. Geophys. Res.*, 112, D24202,

730 <https://doi.org/10.1029/2007JD008646>, 2007.

Fu, S., Deng, X., Shupe, M.D., and Huiwen X.: A modelling study of the continuous ice formation in an autumnal Arctic mixed-phase cloud case, *Atmos. Res.*, 228, 77-85, <https://doi.org/10.1016/j.atmosres.2019.05.021>, 2019

735

Gilbert, E, Orr, A, King, JC, et al. Summertime cloud phase strongly influences surface melting on the Larsen C ice shelf, *Antarctica. Q J R Meteorol Soc.* 2020; 1-16. <https://doi.org/10.1002/qj.3753>

740 Glen, A., and Brooks, S. D.: A new method for measuring optical scattering properties of atmospherically relevant dusts using the Cloud and Aerosol Spectrometer with Polarization (CASPOL). *Atmospheric Chemistry & Physics*, 13, 1345–1356, <https://doi.org/10.5194/acp-13-1345-2013>, 2013

745 Grosvenor, D. P., Choularton, T. W., Lachlan-Cope, T., Gallagher, M. W., Crosier, J., Bower, K. N., Ladkin, R. S., and Dorsey, J. R.: In-situ aircraft observations of ice concentrations within clouds over the Antarctic Peninsula and Larsen Ice Shelf, *Atmos. Chem. Phys.*, 12, 11275–11294, <https://doi.org/10.5194/acp-12-11275-2012>, 2012.

750 Hallett, J. and Mossop, S. C.: Production of secondary ice particles during the riming process, *Nature*, 249, 26–28, doi:10.1038/249026a0, 1974.

Haynes, J.M., C. Jakob, W.B. Rossow, G. Tselioudis, and J. Brown: Major Characteristics of Southern Ocean Cloud Regimes and Their Effects on the Energy Budget. *J.*

755 *Climate*, 24, 5061-5080, <https://doi.org/10.1175/2011JCLI4052.1>, 2011

Heymsfield, A.J., A. Bansemer, and C.H. Twohy: Refinements to Ice Particle Mass Dimensional and Terminal Velocity Relationships for Ice Clouds. Part I: Temperature Dependence. *J. Atmos. Sci.*, 64, 1047–1067, <https://doi.org/10.1175/JAS3890.1>, 2007

760

Hines, K. M., Bromwich, D. H., Wang, S.-H., Silber, I., Verlinde, J., and Lubin, D.: Microphysics of summer clouds in central West Antarctica simulated by the Polar Weather Research and Forecasting Model (WRF) and the Antarctic Mesoscale Prediction System

- (AMPS), *Atmos. Chem. Phys.*, 19, 12431–12454, <https://doi.org/10.5194/acp-19-12431-2019>,
765 2019.
- Hoarau, T., Pinty, J.-P., and Barthe, C.: A representation of the collisional ice break-up process in the two-moment microphysics LIMA v1.0 scheme of Meso-NH, *Geosci. Model Dev.*, 11, 4269–4289, <https://doi.org/10.5194/gmd-11-4269-2018>, 2018.
- 770 Hyder, P., Edwards, J. M., Allan, R. P., Hewitt, H. T., Bracegirdle, T. J., Gregory, J. M., et al.: Critical Southern Ocean climate model biases traced to atmospheric model cloud errors. *Nature Communications*, 9, 3625. <https://doi.org/10.1038/s41467-018-05634-2>, 2018
- Ikawa, M., and Saito, K.: Description of a Non-hydrostatic Model Developed at the Forecast
775 Research Department of the MR, MRI Tech. Rep. 28, 238 pp, 1991
- King J. C., Gadian, A., Kirchgaessner, A., Kuipers, Munneke, P., Lachlan-Cope, T. A., Orr, A., Reijmer, C., van den Broeke, M. R., van Wessem, J. M., and Weeks, M.: Validation of the summertime surface energy budget of Larsen C Ice Shelf (Antarctica) as represented in three
780 high-resolution atmospheric models, *J. Geophys. Res.*, 120, 1335–1347, <https://doi.org/10.1002/2014JD022604>, 2015.
- King, J. C., Lachlan-Cope, T. A., Ladkin, R. S., and Weiss, A.: Airborne Measurements in the Stable Boundary Layer over the Larsen Ice Shelf, Antarctica, *Boundary-Layer Meteorology*,
785 127, 413–428, doi:10.1007/s10546-008-9271-4, 2008.
- Klemp, J. B.: Advances in the WRF model for convection-resolving forecasting, *Adv. Geosci.*, 7, 25–29, <https://doi.org/10.5194/adgeo-7-25-2006>, 2006.
- 790 Korolev, A. V., Emery, E. F., Strapp, J.W., Cober, S. G., Isaac, G. A., Wasey, M., and Marcotte, D.: Small ice particles in tropospheric clouds: fact or artifact?, *B. Am. Meteorol. Soc.*, 92, 967–973, doi:10.1175/2010BAMS3141.1, 2011.
- Korolev, A., Heckman, I., Wolde, M., Ackerman, A. S., Fridlind, A. M., Ladino, L., Lawson, P., Milbrandt, J., and Williams, E.: A new look at the environmental conditions favorable to secondary ice production, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2019-611>, in review, 2019.
- 795

- 800 Lachlan-Cope, T., Listowski, C., and O'Shea, S: The microphysics of clouds over the
Antarctic peninsula—Part 1: Observations. *Atmos. Chem. Phys.*, 16(24), 15,605–15,617.
<https://doi.org/10.5194/acp-16-15605-2016>, 2016
- 805 Lauber, A., Kiselev, A., Pander, T., Handmann, P., and Leisner, T.: Secondary ice formation
during freezing of levitated droplets, *J. Atmos. Sci.*, 75, 2815–
2826, <https://doi.org/10.1175/JAS-D-18-0052.1>, 2018.
- 810 Lawson, R. P., O'Connor, D., Zmarzly, P., Weaver, K., Baker, B., Mo, Q., and Jonsson, H. :
The 2D-S (Stereo) probe: Design and preliminary tests of a new airborne, high-speed, high-
resolution particle imaging probe. *Journal of Atmospheric and Oceanic Technology*, 23,
1462–1477. <https://doi.org/10.1175/JTECH1927.1>, 2006
- 815 Lawson, P., Gurganus, C., Woods, S., and Brientjes, R.: Aircraft observations of cumulus
microphysics ranging from the tropics to midlatitudes: implications for a “new” secondary ice
process, *J. Atmos. Sci.*, 74, 2899–2920, <https://doi.org/10.1175/JAS-D-17-0033.1>, 2017.
- 820 Leisner, T., Pander, T., Handmann, P., and Kiselev, A.: Secondary ice processes upon
heterogeneous freezing of cloud droplets, 14th Conf. on Cloud Physics and Atmospheric
Radiation, Amer. Meteor. Soc, Boston, MA, 2014.
- 825 Listowski, C., and Lachlan-Cope, T.: The microphysics of clouds over the antarctic
peninsula—Part 2:Modelling aspects within Polar WRF. *Atmos. Chem. Phys.*,17(17),
10,195–10,221. <https://doi.org/10.5194/acp-17-10195-2017>, 2017
- 830 Listowski, C., Delanoë, J., Kirchgassner, A., Lachlan-Cope, T., and King, J.: Antarctic
clouds, supercooled liquid water and mixed phase, investigated with DARDAR: geographical
and seasonal variations, *Atmos. Chem. Phys.*, 19, 6771–6808, <https://doi.org/10.5194/acp-19-6771-2019>, 2019. Meyers, M. P., DeMott, P. J., and Cotton, W. R.: New primary ice-
nucleation parameterizations in an explicit cloud model. *Journal of Applied Meteorology*, 31,
708–721. [https://doi.org/10.1175/1520-0450\(1992\)031<0708:NPINPI>2.0.CO;2](https://doi.org/10.1175/1520-0450(1992)031<0708:NPINPI>2.0.CO;2), 1992
- 830 McCluskey, C. S., Hill, T. C. J. , Humphries, R. S., Rauker, A. M., Moreau, S., Stratton, P.

- G., Chambers, S. D., Williams, A. G., McRobert, I., Ward, J., Keywood, M.D., Harnwell, J., Ponsonby, W., Loh, Z.M., Krummel, P. B., Protat, A., Kreidenweis, S.M., and DeMott, P.J.: Observations of ice nucleating particles over Southern Ocean waters. *Geophysical Research Letters*, 45, 11,989–11,997. <https://doi.org/10.1029/2018GL079981>, 2018
- 835
- Mizuno, H.: Parameterization of the accretion process between different precipitation elements. *J. Meteor. Soc. Japan*, 57, 273-281, 1990
- Mossop, S. C., and Hallett, J.: Ice crystal concentration in cumulus clouds: Influence of the drop spectrum. *Science*, 186, 632–634. <https://doi.org/10.1126/science.186.4164.632>, 1974.
- 840
- Mossop, S. C. (1985). Secondary ice particle production during rime growth: The effect of drop size distribution and rimer velocity. *Quart. J. Roy. Meteor. Soc.*, 111(470), 1113–1124. <https://doi.org/10.1002/qj.49711147012>
- 845
- Morrison, H., Curry, J.A., and Khvorostyanov, V.I.: A New Double-Moment Microphysics Parameterization for Application in Cloud and Climate Models. Part I: Description, *Atmos. Sci.*, 62, 3683-3704 62, 2005
- 850
- Nakanishi, N., and Niino, H.: An improved Mellor–Yamada level-3 model: Its numerical stability and application to a regional prediction of advection fog. *Boundary-Layer Meteorology*, 119, 397–407, 2006
- O’Shea, S. J., Choulaton, T. W., Flynn, M., Bower, K. N., Gallagher, M., Crosier, J., 855 Williams, P., Crawford, I., Fleming, Z. L., Listowski, C., Kirchgaessner, A., Ladkin, R. S., and Lachlan-Cope, T.: In situ measurements of cloud microphysics and aerosol over coastal Antarctica during the MAC campaign, *Atmos. Chem. Phys.*, 17, 13049–13070, <https://doi.org/10.5194/acp-17-13049-2017>, 2017.
- 860
- Passarelli, R.E.: An Approximate Analytical Model of the Vapor Deposition and Aggregation Growth of Snowflakes., *J. Atmos. Sci.*, 35, 118–124, 1978
- Phillips, V.T.J., Yano, J.-I., and Khain, A.: Ice multiplication by breakup in ice-ice collisions. Part I: Theoretical formulation, *J. Atmos. Sci.*, 74, 1705–1719, <https://doi.org/10.1175/JAS-D->

865 16-0224.1, 2017a.

Phillips, V.T.J., Yano, J.-I., Formenton, M., Ilotoviz, E., Kanawade, V., Kudzotsa, I., Sun, J., Bansemer, A., Detwiler, A.G., Khain, A., and Tessendorf, S.A.: Ice Multiplication by Breakup in Ice–Ice Collisions. Part II: Numerical Simulations. *J. Atmos. Sci.*, 74, 2789–
870 2811, <https://doi.org/10.1175/JAS-D-16-0223.1>, 2017b.

Prein, A. F., Langhans, W., Fosser, G., Ferrone, A., Ban, N., Goergen, K., Keller, M., Tlle, M., Gutjahr, O., Feser, F., Brisson, E., Kollet, S., Schmidli, J., Lipzig, N. P. M., and Leung, R.: A review on regional convection-permitting climate modeling: Demonstrations, prospects,
875 and challenges, *Reviews of Geophysics*, 53(2), 323–361, doi:10.1002/2014RG000475, 2015.

Pruppacher, H.R., and Klett, J.D. : *Microphysics of Clouds and Precipitation*. 2nd Edition, Kluwer Academic, Dordrecht, 954 p., 1997

880 Qu, Y., Khain, A., Phillips, V., Ilotoviz, E., Shpund, J., Patade, S., and Chen, B. : The role of ice splintering on microphysics of deep convective clouds forming under different aerosol conditions: Simulations using the model with spectral bin microphysics. *J. Geophys. Res. Atmos.*, 125, e2019JD031312. <https://doi.org/10.1029/2019JD031312>, 2020

885 Rangno, A.L., and Hobbs, P.V., Ice particles in stratiform clouds in the Arctic and possible mechanisms for the production of high ice concentrations, *J. Geophys. Res.*, 106, 15, 065–15,075, 2001.

Reisner, J., Rasmussen, R. M., and Bruintjes, R. T.: Explicit forecasting of supercooled liquid
890 water in winter storms using the MM5 mesoscale model, *Quart. J. Roy. Meteor. Soc.*, 124(548), 1071–1107, doi: 10.1002/qj.49712454804, 1998.

Skamarock, W. C., and Klemp, J. B.: A time-split nonhydrostatic atmospheric model for weather research and forecasting applications. *J. Comp. Phys.*, 227(7), 3465–3485.
895 <https://doi.org/10.1016/j.jcp.2007.01.037>, 2008

Schmale, J., Baccarini, A., Thurnherr, I., Henning, S., Efraim, A., Regayre, L., Bolas, C., Hartmann, M., Welti, A., Lehtipalo, K., Aemisegger, F., Tatzelt, C., Landwehr, S., Modini, R.

- L., Tummon, F., Johnson, J., Harris, N., Schnaiter, M., Toffoli, A., Derkani, M., Bukowiecki,
900 35 N., Stratmann, F., Dommen, J., Baltensperger, U., Wernli, H., Rosenfeld, D., Gysel-Beer,
M., and Carslaw, K.: Overview of the Antarctic Circumnavigation Expedition: Study of
Preindustrial-like Aerosols and Their Climate Effects (ACE-SPACE), *Bull. Amer. Meteor.
Soc.*, 100 (11): 2260–2283, DOI:10.1175/BAMSD-18-0187.1, 2019
- 905 Schwarzenboeck, A., Shcherbakov, V., Lefevre, R., Gayet, J.-F., Duroure, C., and Pointin, Y.:
Indications for stellar-crystal fragmentation in Arctic clouds, *Atmos. Res.*, 92, 220
228, <https://doi.org/10.1016/j.atmosres.2008.10.002>, 2009.
- Sotiropoulou, G., Sullivan, S., Savre, J., Lloyd, G., Lachlan-Cope, T., Ekman, A. M. L., and
Nenes, A.: The impact of Secondary Ice Production on Arctic Stratocumulus, *Atmos. Chem.
910 Phys.*, 20, 1301–1316, <https://doi.org/10.5194/acp-2019-804>, 2020.
- Stephens, G.L.: Radiation profiles in extended water clouds. II. Parameterization schemes. *J.
Atmos. Sci.*, 35, 2123–2132, 1978
- 915 Sullivan, S. C., Kiselev, A., Leisner, T., Hoose, C., and Nenes, A.: Initiation of secondary ice
production in clouds, *Atmos. Chem. Phys.*, 18, 1593–1610, doi:10.5194/acp-18-1593-2018,
2018a.
- Sullivan, S. C., Barthlott, C., Crosier, J., Zhukov, I., Nenes, A., and Hoose, C.: The effect of
920 secondary ice production parameterization on the simulation of a cold frontal rainband,
Atmos. Chem. Phys., 18, 16461–16480, <https://doi.org/10.5194/acp-18-16461-2018>, 2018b.
- Takahashi, T., Nagao, Y., and Kushiyama, Y.: Possible high ice particle production during
graupel-graupel collisions, *J. Atmos. Sci.*, 52, 4523–4527, doi:10.1175/1520-0469, 1995.
- 925 Vardiman, L.: The generation of secondary ice particles in clouds by crystal-crystal collision,
J. Atmos. Sci., 35, 2168–2180, doi:10.1175/1520-0469, 1978.
- Verlinde, J., Flatau, P.J., and W.R. Cotton, W.R.: Analytical Solutions to the Collection
Growth Equation: Comparison with Approximate Methods and Application to Cloud

930 Microphysics Parameterization Schemes. *J. Atmos. Sci.*, 47, 2871–
2880, [https://doi.org/10.1175/1520-0469\(1990\)047<2871:ASTTCG>2.0.CO;2](https://doi.org/10.1175/1520-0469(1990)047<2871:ASTTCG>2.0.CO;2)

Verlinde, J. and W.R. Cotton, 1993: Fitting Microphysical Observations of Nonsteady
Convective Clouds to a Numerical Model: An Application of the Adjoint Technique of Data
935 Assimilation to a Kinematic Model. *Mon. Wea. Rev.*, 121, 2776–
2793, [https://doi.org/10.1175/1520-0493\(1993\)121<2776:FMOONC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1993)121<2776:FMOONC>2.0.CO;2)

Vignon, É., Besic, N., Jullien, N., Gehring, J., & Berne, A. Microphysics of snowfall over
coastal East Antarctica simulated by Polar WRF and observed by radar. *J. Geophys. Res.:*
Atmospheres, 124, 11452–11476, 2019.

940 Welti, A., Bigg, E. K., DeMott, P. J., Gong, X., Hartmann, M., Harvey, M., Henning, S.,
Herenz, P., Hill, T. C. J., Hornblow, B., Leck, C., Löffler, M., McCluskey, C. S., Rauker, A.
M., Schmale, J., Tatzelt, C., van Pinxteren, M., and Stratmann, F.: Shipbased measurements
of ice nuclei concentrations over the Arctic, Atlantic, Pacific and Southern Ocean, *Atmos.*
945 *Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-466>, in review, 2020.

Yano, J.-I. and Phillips, V. T. J.: Ice-ice collisions: an ice multiplication process in
atmospheric clouds, *J. Atmos. Sci.*, 68, 322–333, doi:10.1175/2010JAS3607.1, 2011.

950 Yano, J.-I., Phillips, V. T. J., and Kanawade, V.: Explosive ice multiplication by mechanical
break-up in-ice-ice collisions: a dynamical system-based study, *Q. J. Roy. Meteor. Soc.*, 142,
867–879, <https://doi.org/10.1002/qj.2687>, 2016.

Young, G., Lachlan-Cope, T., O'Shea, S. J., Dearden, C., Listowski, C., Bower, K. N.,
955 Choularton T.W., and Gallagher M.W.: Radiative effects of secondary ice enhancement in
coastal Antarctic clouds. *Geophys. Res. Lett.*, 46, 23122321,
<https://doi.org/10.1029/2018GL080551>, 2019.


Young, K. C., 1974: The Role of Contact Nucleation in Ice Phase Initiation in Clouds. *J.*
Atmos. Sci., 31, 768–776, [https://doi.org/10.1175/1520-
960 *0469\(1974\)031<0768:TROCNI>2.0.CO;2*.](https://doi.org/10.1175/1520-0469(1974)031<0768:TROCNI>2.0.CO;2)

Tables:

965 **Table 1:** Mean modeled downward and upward shortwave (SWD_{SFC} , SWU_{SFC}) and longwave (LWD_{SFC} , LWU_{SFC}) surface radiation, along with upward shortwave and longwave (SWU_{TOA} , LWU_{TOA}) radiation at the top of the atmosphere, during flights M218 and M219. Model results are averaged over the dashed rectangular area in Fig. 1.


Simulation	SWD_{SFC} (Wm^{-2})	SWU_{SFC} (Wm^{-2})	LWD_{SFC} (Wm^{-2})	LWU_{SFC} (Wm^{-2})	SWU_{TOA} (Wm^{-2})	LWU_{TOA} (Wm^{-2})
CNTRL	323.9	182.1	244.3	304.6	255.8	218.4
PHIL0.2	328.6	184.5	244.1	304.6	254.8	218.5
PHIL0.3	322.3	181.0	247.4	305.3	256.6	217.9
PHIL0.4	339.7	190.8	243.3	304.9	251.1	219.8
FRAG1	354.1	198.6	236.7	303.8	246.9	221.5
FRAG1siz	335.7	188.8	244.0	304.6	250.5	220.7
TAKAH	365.9	206.5	229.8	303.3	242.5	221.2
TAKAHsc	349.5	194.5	237.0	304.2	246.6	222.0

Figures:



975 **Figure 1:** Map of Antarctic domains. Colors indicate terrain heights (green to yellow) and
sea-ice concentrations (blue to white), whereas the purple contours correspond to 500 hPa
geopotential heights from the CNTRL simulation at 18:00 UTC, 27 November 2015. The
black solid line delimits the 1-km horizontal grid spacing domain, while the dashed one
980 outlines the subset of the nest used for direct comparison with the aircraft data. Orange and
blue lines indicate the flight tracks, while the red circle represents Halley station. The small
figure in the top right corner indicates the location of the 1-km horizontal grid spacing domain
relative to the Antarctic continent.

985



990

Figure 2: (a) Mean ice number concentrations (cloud ice + snow + graupel, N_{ice}) as a function of temperature for the whole MAC campaign (pink), our case study (grey) and the eight model simulations. Grey dots indicate point observations. (b) Relative frequency distribution of N_{ice} , binned in $0.5 L^{-1}$ intervals, scaled with maximum frequency. Ice properties are calculated for particles $> 80 \mu m$ and for $N_{ice} > 0.005 L^{-1}$ within the lowest 1.5 km a.s.l.

995

1000

1005





Figure 3: Mean vertical profiles of number concentrations of modeled (a) cloud ice, (b) graupel, (c) snow and (d) total ICNCS for six simulations. Grey lines represent measured concentrations with diameters (a) smaller and (c) larger than 250 μm . Graupel concentrations cannot be distinguished in the measurements (hence no grey profile in panel b). Ice properties from the model are calculated for $N_{\text{isg}} > 0.005 \text{ L}^{-1}$. For consistency with observations, only particles with sizes $> 80 \mu\text{m}$ are included in the modeled profiles in panels (a), (c) and (d).

1010

1015




1020

Figure 4: (a) Mean liquid water content (LWC) as a function of temperature for our case study (grey) and the eight model simulations. (b) Relative frequency distribution of LWC, binned in 0.01 g m^{-3} intervals, scaled with maximum frequency. Only values greater than 0.01 g m^{-3} within the lowest 1.5 km a.s.l. are included in the analysis.

1025

1030

1035



1040

Figure 5: Total ice number concentrations (N_{ice}) for particles $> 80 \mu\text{m}$ as a function of temperature for the while MAC campaign (pink), our case study (grey) and the nine sensitivity simulations with varying treatment of the H-M process (see section 4.3). Mean values and the 95th percentile are shown in panels (a) and (b), respectively.

1045

1050

1055

1060




Figure 6: Same as in Fig. 5 but for sensitivity simulations with varying INP conditions (see section 4.4)

1065

1070

1075

1080