

---

Editor Decision: Reconsider after major revisions (03 Sep 2020) by Jason Surratt

Comments to the Author:

Dear Authors,

The reviewers mostly feel you have addressed the comments initially raised on your manuscript. However, one reviewer noted that the English grammar was very poor in many sections of the manuscript. I concur with this assessment and feel further English grammar editing is needed before full publication can be considered. I kindly ask the authors to either reach out to a native-English speaking colleague to edit your manuscript or use a professional service. The number of grammar writing errors make the article difficult to read. Please let me know if you have any questions. One of the reviewers noted many places where the English writing needs to be fixed, but I should note to the authors it is not the duty of the reviewers to help with rewriting of the manuscript.

Most sincerely, Jason Surratt

**Response**

Dear Surratt,

Thanks very much for your kindness and your valuable advice. Now this paper has been edited for grammar, phrasing, and punctuation by American Journal Experts. In addition, many edits were made to further improve the flow and readability of the text. I very much hope that the manuscript can meet the requirements of your journal so that it can continue to be reviewed and published. Thanks again.

The proof is as following:

September 16, 2020

Dear Hongling Yin,

Thank you for choosing American Journal Experts. This manuscript, titled "Measurement report: Seasonality, distribution and sources of organophosphate esters in PM<sub>2.5</sub> from an inland urban city in Southwest China," is very interesting. The paper was edited for grammar, phrasing, and punctuation. In addition, many edits were made to further improve the flow and readability of the text. Below, we highlight the areas of this paper that we focused on in our edit.

In cases where the meaning of the text was not clear, revisions were made to convey the information with increased clarity and reduced ambiguity.

Articles are an important aspect of the English language, including the definite article "the" and the indefinite articles "a" and "an." Our edits focused on improving article use, which is often strongly dependent on context and field conventions.

Some edits were made to improve conciseness by trimming unnecessary words and streamlining the flow of your manuscript.

Comments were left if further clarification would be helpful or confirmation of the meaning of the text was necessary. Please review these comments and all our changes carefully for more detailed suggestions, as well as to ensure that the final version of the manuscript is fully accurate.

Thank you again for using our editing services; we wish you the best of luck with your submission.

Best regards,

Katie D.  
Senior Editor  
American Journal Experts

Best Regards.

Yours,

Sincerely,

Hongling Yin

2020-9-22

## Measurement report: Seasonality, distribution and sources of organophosphate esters in PM<sub>2.5</sub> from an inland urban city in Southwest China

Hongling Yin, Jiangfeng Liang, Di Wu, Shiping Li, Yi Luo, Xu Deng

College of Resources and Environment, Chengdu University of Information Technology, Chengdu, Sichuan, 610025, China

Correspondence: Hongling Yin ([yhl@cuit.edu.cn](mailto:yhl@cuit.edu.cn))

**Abstract.** Organophosphate esters (OPEs) are contaminants of emerging concern, and studies have concluded that urban areas are a significant source of OPEs. Samples were collected from six ground-based sites located in Chengdu, a typical rapidly developing metropolitan area in Southwest China, and

删除的内容: southwest

删除的内容: a kind of

删除的内容: were

删除的内容: fast

删除的内容: of

删除的内容: southwest

were analysed for seven OPEs in atmospheric PM<sub>2.5</sub> ( $\Sigma_7$  OPEs). The concentrations of  $\Sigma_7$  OPEs in PM<sub>2.5</sub> ranged from 5.83 to 6.91 ng m<sup>-3</sup>, with a mean of 6.6 ± 3.3 ng m<sup>-3</sup>, and the primary pollutants were tris-(2-butoxyethyl) phosphate (TBEP), tri-n-butyl phosphate (TnBP), tris-(2-chloroethyl) phosphate (TCEP) and tris-(2-chloroisopropyl) phosphate (TCPP), which together made up more than 80% of the  $\Sigma_7$  OPEs. The concentrations of  $\Sigma_7$  OPEs were higher in autumn/winter than in summer. Nonparametric tests showed that there was no significant difference in  $\Sigma_7$  OPE concentrations among the six sampling sites, but the occurrence of unexpectedly high levels of individual OPEs at different sites in autumn might indicate noteworthy emissions. A very strong correlation ( $R^2 = 0.98$ ,  $p < 0.01$ ) between the OPEs in soil and in PM<sub>2.5</sub> was observed. Backward trajectory analysis indicated that the OPEs in PM<sub>2.5</sub> were mainly affected by local sources. Principal component analysis (PCA) revealed that the OPEs in PM<sub>2.5</sub> were largely sourced from the plastic industry/interior decoration/traffic emission (34.5%) and the chemical, mechanical and electrical industries (27.8%), while the positive matrix factorization (PMF) model revealed that the main sources were the plastics industry/indoor source emissions, the food/cosmetics industry, and industrial emissions. In contrast to coastal cities, sustained and stable high local emissions in the studied inland city were identified, which is particularly noteworthy. Chlorinated phosphates, especially TCPP and TCEP, had a high content, and their usage and source emissions should be controlled.

## 1. Introduction

With the prohibition of brominated flame retardants, the production of and demand for organophosphate esters (OPEs) have rapidly increased in recent years (Wang et al., 2012). OPEs are widely distributed in the environment and have been detected in air (Guo et al., 2016; Li et al., 2017), water (Wang et al., 2013; Li et al., 2014), soil (Yin et al., 2016), sediment (Cristale J. et al., 2013; Celano R, et al., 2014) and organisms (Kim et al., 2011). However, many scholars have found that OPE residues in the environment can cause toxic effects on organisms (WHO, 1991, 1998, 2000; Kanazawa et al., 2010; Van der Veen and de Boer, 2012; Du et al., 2015). Some countries have enacted legislature to restrict the usage of OPEs (Blum et al., 2019; Exponent, 2018; State of California, 2020). Nevertheless, the production and usage of OPEs in China are still on the rise.

删除的内容: analyzed...for seven OPEs in atmospheric PM<sub>2.5</sub> ( $\Sigma_7$  OPEs). The concentrations of  $\Sigma_7$  OPEs in PM<sub>2.5</sub> ranged from 5.83 to 6.91 ng m<sup>-3</sup>, with a mean of 6.6 ± 3.3 ng m<sup>-3</sup>, and the primary pollutants were tris-(2-butoxyethyl) phosphate (TBEP), tri-n-butyl phosphate (TnBP)TnBP... tris-(2-chloroethyl) phosphate (TCEP) TCEP ...nd tris-(2-chloroisopropyl) phosphate (TCPP)TCPP

删除的内容: in...f the  $\Sigma_7$  OPEs. The concentrations of  $\Sigma_7$  OPEs were higher in autumn/winter than that ...n summer. Nonparametric test...ests showed that there was no significant difference in  $\Sigma_7$  OPEs...PE concentrations among the six sampling sites, but the occurrence of unexpected...expectedly high level

删除的内容: that there was a

删除的内容: Very

删除的内容: The b

删除的内容: displayed

删除的内容: The p...rincipal component analysis (PCA) identified

删除的内容: decoration /traffic

删除的内容: y...(27.8%), while the positive matrix factorization (PMF) model found

删除的内容: Differed from the ...n contrast to coastal cities, the ...sustained and stable high local emissions in the the...studied inland city

删除的内容: ies...were identified, which were

删除的内容: The chlorinated ...hlorinated phosphates, especially TCPP and TCEP have

删除的内容: s... a high content, whose

删除的内容:

删除的内容: the

删除的内容: of

删除的内容: To date,

删除的内容: the ...PE residues of OPEs ...n the environment could

删除的内容:

删除的内容: legislated ...nacted legislature to restrict the usage of OPEs (Blum et al., 2019; Exponent, 2018; State of

As OPEs are synthetic substances, the only source of OPEs in the environment is anthropogenic emissions. The detection of OPEs in Arctic and Antarctic snow samples and atmospheric particulate matter samples demonstrated that OPEs can be transported over long distances (Moller et al., 2012; Li et al., 2017). Many studies on OPEs in oceans have been carried out, and the concentrations of particle-bound OPEs range from tens to thousands of ng m<sup>-3</sup> (Moller et al., 2011; 2012; Cristale & Lacorte, 2013; Li et al., 2017; McDonough et al., 2018). Some researchers noted that high concentrations of OPEs (thousands of ng m<sup>-3</sup>) originated from air flow from the mainland (Moller et al., 2012; Lai et al., 2015). In addition, studies have proven that urban areas have the highest OPE pollution. However, until now, only a few papers have reported the concentration and distribution of OPEs in urban atmospheric PM<sub>2.5</sub>. Concentrations of atmospheric OPEs in most cities were lower than 10 ng m<sup>-3</sup>; a higher concentration of 19.2 ng m<sup>-3</sup> was observed at a suburban site in Shanghai, and a concentration of 49.1 ng m<sup>-3</sup> was observed in Hong Kong (Ohura et al., 2006; Salamova et al., 2014b; Marklund et al., 2005; Shoeib et al., 2014; Yin et al., 2015; Liu et al., 2016; Ren et al., 2016; Guo et al., 2016; Wong et al., 2018). To date, most studies in China have focused on OPEs in the Yangtze River Delta and Pearl River Delta, especially eastern coastal cities, while little attention has been paid to western inland cities.

Chengdu is a typical inland city located in Southwest China. This megacity is the capital of Sichuan Province, covers an area of 14,335 square kilometres and has a permanent population of 16.33 million. As an important national high-tech industrial base, commercial logistics centre and comprehensive transportation hub designated by the State Council, Chengdu is the most important central city in the western region (<https://en.wikipedia.org/wiki/Chengdu>). Liu et al. (2016) investigated three chlorinated OPEs in the atmosphere at 10 urban sites in China during 2013 – 2014 and observed the highest annual mean concentrations in Chengdu (1,300 ± 2,800 ng m<sup>-3</sup>). However, there is still a lack of information regarding the levels, sources, and fate of OPEs in Southwest China, which may obviously differ from those of coastal cities or ocean locations. Our previous study investigated OPE concentrations in PM<sub>2.5</sub> at two sites (urban and suburban sites) in Chengdu (a city experiencing fast economic growth in Southwest China) and found that the OPE concentrations and profiles were similar at the two sites (Yin et al., 2015). However, the influencing factors and potential sources of OPEs in PM<sub>2.5</sub> in Chengdu are still unclear. Therefore, in this study, PM<sub>2.5</sub> was collected over one year (October 2014 to September 2015) at six sites in Chengdu to (a) report the levels and composition profiles of OPEs in urban air in a typical inland city; (b) obtain the seasonal and spatial variations in OPEs in PM<sub>2.5</sub>; (c) investigate the

删除的内容: S

删除的内容: were...have been carried out a lot... and the concentrations of particle-bound OPEs ranged

删除的内容: R...searchers noted that the contribution of air flow originated from the mainland when...high concentrations of OPEs (thousands of ng m<sup>-3</sup>) appeared...originated from air flow from the mainland (Moller et al., 2012; Lai et al., 2015). In addition, studies proved...ave proven that the...urban areas was...ave the highest OPE pollution area of OPEs... However, until now, only a few papers have reported the concentration and distribution of OPEs in urban atmospheric PM<sub>2.5</sub>. Concentrations of atmospheric OPEs in most cities were lower than 10 ng m<sup>-3</sup>; a...higher concentrations...of 19.2 ng m<sup>-3</sup> were...as observed at a suburban site in Shanghai, and a concentration of 49.1 ng m<sup>-3</sup> were

删除的内容: Hongkong...ong Kong (Ohura et al., 2006; Salamova et al., 2014b; Marklund et al., 2005; Shoeib et al., 2014; Yin et al., 2015; Liu et al., 2016; Ren et al., 2016; Guo et al., 2016; Wong et al., 2018). To date, most of...studies in China focus...ave focused on the...OPEs in the Yangtze River Delta and Pearl River Delta, especially eastern coastal cities, while little attention was...has been paid to the...estern inland cities.

删除的内容: the

删除的内容: southwest

删除的内容: of

删除的内容: It is...his megacity is the capital and megacity...f Sichuan Province, which...overs an area of

删除的内容: differed...iffer from those of the

删除的内容: over the sea

删除的内容: has

删除的内容: the

删除的内容: OPEs

删除的内容: n economically fast -growing...city experiencing fast economic growth in Southwest southwest

删除的内容: of...hina)...and found that the OPEs...PE concentrations and profile...rofiles were similar at the two

删除的内容: of

relationships and correlations among the target compounds or with influencing factors; and (d) illustrate the potential sources of OPEs in PM<sub>2.5</sub>.

删除的内容: influence

## 2. Materials and Methods

### 2.1 Chemicals

The main reagents, such as ethyl acetate, acetone, hexane and acetonitrile, were high-performance liquid chromatography (HPLC) grade (Kelon Chemical Corp., China). The standard solution (Sigma Aldrich Corp., USA) included tri-n-butyl phosphate (TnBP), tris-(2-ethylhexyl) phosphate (TEHP), tris-(2-butoxyethyl) phosphate (TBEP), triphenyl phosphate (TPhP), tris-(2-chloroethyl) phosphate (TCEP), tris-(2-chloroisopropyl) phosphate (TCPP), and tris-(2,3-dichloropropyl) phosphate (TDCIPP). Copper, aluminium oxide, silica gel, Na<sub>2</sub>SO<sub>4</sub> and other chemicals were purchased from Kelon Chemical Corp., China. Deionized water was obtained from Milli-Q equipment.

删除的内容: High Performance Liquid Chromatography

删除的内容: -

删除的内容: -

删除的内容:

删除的内容: supplied

删除的内容: a

### 2.2. Sample collection

The atmospheric sampling sites were located in the main city area (site B: downtown; site C: south; site D: east; site E: north; site F: west) and suburban area (site A) of Chengdu, as shown in Fig. S1. The atmospheric samples were collected by a KC-6120 medium-flow atmospheric comprehensive sampler with quartz film. The speed was set at 100 L min<sup>-1</sup>, and each collection campaign lasted 23 hours. The sampling campaign was carried out between October 2014 and September 2015. In each season, continuous sampling was carried out for approximately one week, except for rainy days. In autumn, the sampling duration was from October 23 to October 29, 2014 (no samples were obtained due to rain on October 26 and 27); in winter, the sampling duration was from December 22 to December 30, 2014 (no samples were obtained due to rain on October 25 and 26); in spring, the sampling duration was from March 25 to March 30, 2015; and in summer, the sampling duration was from July 16 to July 24, 2015 (no sample was obtained due to rain on July 21). A total of 149 samples were obtained. Most of the weather conditions were cloudy days, with south/north winds at ≤5.5 m/s. The temperature ranged from 0 to 35 °C. The weather conditions represented typical seasonal weather conditions.

删除的内容:

删除的内容: about

删除的内容: was

删除的内容: the

删除的内容: was

删除的内容: the

删除的内容: the

删除的内容: wind

删除的内容: Temperature

删除的内容: °C

删除的内容: W

删除的内容: could

删除的内容: of the season

### 2.3. Sample preparation and analysis

The shredded PM<sub>2.5</sub> sample film was placed in a test tube and incubated in 20 mL ethyl acetate/acetone (v:v, 3:2) for 12 hours. After ultrasonic extraction for 30 minutes, the liquid was separated, and the residue was further extracted with 10 mL ethyl acetate/acetone (v:v, 3:2) by ultrasonic extraction for 15 minutes. The extracts were combined, concentrated with vacuum-condensing equipment (Buchi Syncore Q-101, Switzerland) to approximately 1 mL, and then loaded onto an activated aluminium oxide/silica gel (v:v, 3:1) column. The column was eluted first with 20 mL hexane to remove impurities, then with 20 mL ethyl acetate/acetone (v:v, 3:2), and the latter eluate (ethyl acetate/acetone) was collected. The eluate was concentrated to nearly dry by vacuum-condensing equipment and then fixed to 200  $\mu$ L with hexane for gas chromatography-mass spectrometry (GC-MS) (Shimadzu 2010 plus, Japan) analysis.

The GC was equipped with an Rti-5MS capillary column (30 m  $\times$  0.25  $\mu$ m  $\times$  0.25 mm, Kelong), and operated with a 280  $^{\circ}$ C inlet temperature using splitless injection. The MS source was electron impact (EI), and it was operated in selected ion monitoring (SIM) mode. Helium was used as the carrier gas with a flow rate of 1.00 mL min<sup>-1</sup>. The GC oven temperature was held at 50  $^{\circ}$ C for 1 minute, increased to 200  $^{\circ}$ C at 15  $^{\circ}$ C min<sup>-1</sup> and held for 1 minute, increased to 250  $^{\circ}$ C at 4.00  $^{\circ}$ C min<sup>-1</sup>, and then increased to 300  $^{\circ}$ C at 20  $^{\circ}$ C min<sup>-1</sup> and held for 4 minutes. The interface temperature was 280  $^{\circ}$ C and the ion source temperature was 200  $^{\circ}$ C. The respective characteristic ion and reference ions (m/z) of the 7 target compounds were, 155/99, 211, 125 (TnBP), 249/63, 143, 251 (TCEP), 125/99, 201, 277, 157 (TCCP), 75/99, 191, 209, 381 (TDCPP), 326/325, 77, 215 (TPhP), 85/100, 199, 299 (TBEP), 99/113 and 211 (TEHP).

### 2.4. QA/QC

Thorough QA/QC procedures for OPE analysis were conducted to ensure data quality. To evaluate the recovery efficiencies of the analytical procedures, all samples were spiked with an internal standard (TDCPP-d<sub>15</sub> and TPhP-d<sub>15</sub>), and the accuracy was evaluated by their recoveries. The concentrations of the 7 OPEs were determined by an external standard method. The correlation coefficients of the standard curves of the seven OPE monomers were all greater than 0.990. The recoveries of the 7 OPEs and the internal standard were between 78.9% and 122.5%. A matrix blank was analysed with each batch of samples. Only TnBP was detected in the blanks, and the level of TnBP found in the blanks was <5% of the concentrations measured in all samples, which meant it was negligible. Field blanks were prepared

删除的内容: and

删除的内容: by

删除的内容:

删除的内容:

删除的内容: first

删除的内容:

删除的内容:

删除的内容: volume

删除的内容: is

删除的内容: capillary column

删除的内容: ,

删除的内容:  $^{\circ}$ C

删除的内容: a

删除的内容:  $^{\circ}$ C

删除的内容:  $^{\circ}$ C

删除的内容:  $^{\circ}$ C

删除的内容:

删除的内容:  $^{\circ}$ C

删除的内容:  $^{\circ}$ C

删除的内容:  $^{\circ}$ C

删除的内容:  $^{\circ}$ C

删除的内容:  $^{\circ}$ C

删除的内容:  $^{\circ}$ C

删除的内容: :

删除的内容: /

删除的内容: added

删除的内容: standard

删除的内容: s

删除的内容:

删除的内容: done

at each site to evaluate the background contamination in the field. TBEP, TnBP and TEHP were detected in the field blanks. The levels found in the blanks were <15% of the concentrations measured in all samples. The instrument precision was in the range of 1.9%-8.3%.

删除的内容: in it

删除的内容: level of them

### 3. Results and Discussion

#### 3.1. Levels of OPEs in PM<sub>2.5</sub>

删除的内容:

OPEs were present in PM<sub>2.5</sub> samples collected across the study area (Fig. S1). The seven OPEs were found in 96.7% - 100% of the samples (n=149). The high detection frequencies of most OPEs indicated that OPE contamination was ubiquitous in the air of Chengdu city.

删除的内容: S

The concentrations of  $\Sigma_7$  OPEs in PM<sub>2.5</sub> across the six sites were in the range of 3.5 - 11.5 ng m<sup>-3</sup>, and the annual median concentration of  $\Sigma_7$  OPEs was 6.5 ± 3.3 ng m<sup>-3</sup> (Fig. 1). The seasonal average value of OPEs in PM<sub>2.5</sub> at each site was almost at the same level (5.8 ± 1.3 ng m<sup>-3</sup> - 6.9 ± 2.5 ng m<sup>-3</sup>). Nonparametric tests showed that there was no significant difference in  $\Sigma_7$  OPE concentrations among the six sampling sites, indicating that the atmosphere was evenly mixed, and there was no particularly heavy- or light-polluted area in Chengdu city. These data were quite consistent with our previous study, which reported the annual median concentration of OPEs in PM<sub>2.5</sub> from December 2013 to October 2014 (Yin et al., 2015). Interestingly, the concentration of  $\Sigma_7$  OPEs at the suburban site was similar to, or even higher than those at some urban sites, which indicated more local sources of these compounds in the suburban area.

删除的内容: C

删除的内容: test

删除的内容: OPEs

删除的内容: mixed

删除的内容:

删除的内容: ,

The concentrations of OPEs in the particles of Chengdu were comparable to those reported for Beijing (0.257 - 8.36 ng m<sup>-3</sup>) (Wang et al., 2018),  $\Sigma_6$  OPEs for a Shanghai urban site (6.6 ng m<sup>-3</sup>) (Ren et al., 2016), and  $\Sigma_6$  OPEs for Bursa (6.5 ng m<sup>-3</sup>), but higher than those in Houston, US ( $\Sigma_{12}$  OPEs, 0.16 - 2.4 ng m<sup>-3</sup>) (Clark et al., 2017), Dalian ( $\Sigma_9$  OPEs, 0.32-3.46 ng m<sup>-3</sup>, 1.21 ± 0.67 ng m<sup>-3</sup>) (Wang et al., 2019), the European Arctic (0.033 - 1.45 ng m<sup>-3</sup>) (Salamova et al., 2014), the Northern Pacific and Indian Oceans (0.23 - 2.9 ng m<sup>-3</sup>) (Moller et al., 2012), the Yellow Sea and Bohai Sea (0.044 - 0.52 ng m<sup>-3</sup>) (Li et al., 2017), the South China Sea (0.047 - 0.161 ng m<sup>-3</sup>) (Lai et al., 2015), and the North Atlantic and Arctic Oceans (0.035 - 0.343 ng m<sup>-3</sup>) (Li et al., 2017). The detected OPE concentrations were also lower than those in Guangzhou and Taiyuan ( $\Sigma_{11}$  OPEs, 3.10 - 544 ng m<sup>-3</sup>) (Chen et al., 2020), Bursa, Turkey ( $\Sigma_6$  OPEs, 0.53 - 19.14 ng m<sup>-3</sup>) (Kurtkarakus et al., 2018), and 20 industrial sites in an urban region in Guangzhou, China ( $\Sigma_{12}$  OPEs, 0.52 - 62.75 ng m<sup>-3</sup>) (Wang et al., 2018).

删除的内容: that

删除的内容: from

删除的内容: 6.6 ng m<sup>-3</sup> (

删除的内容: )

删除的内容:

删除的内容: 6.5 ng m<sup>-3</sup> (

删除的内容: )

删除的内容: that

删除的内容: And they

删除的内容: that

删除的内容: 544ng

删除的内容: in

删除的内容: in Guangzhou, China

### 3.2. Composition profiles of OPEs in PM<sub>2.5</sub>

Nonchlorinated OPEs were the predominant OPEs across Chengdu city. The annual median concentrations of total OPEs were fairly uniform at the six sites and were influenced mainly by alkylated OPEs. As listed in Table 1, the general trend was that TBEP was the most abundant OPE (annual median concentration: 2.3 ng m<sup>-3</sup>, 35.3% of Σ<sub>7</sub> OPEs), followed by TCEP (1.1 ng m<sup>-3</sup>, 16.3%) ≈ TnBP (1.0 ng m<sup>-3</sup>, 15.6%) ≈ TCPP (1.0 ng m<sup>-3</sup>, 15.0%) > TPhP (0.5 ng m<sup>-3</sup>, 8.4%) > TEHP (0.3 ng m<sup>-3</sup>, 5.1%) > TDCPP (0.3 ng m<sup>-3</sup>, 4.3%), with the concentrations of TBEP being approximately 7 - 10 times higher than those of TDCPP and TEHP. The composition profile of OPEs was similar at all sites except for the east site, which had a higher contribution of TnBP. However, TBEP, TCEP, TCPP and TnBP were the dominant OPEs across the city and contributed more than 80% to Σ<sub>7</sub> OPEs. This profile was similar to that in Longyearbyen, Norway, with the primary pollutants being TnBP and TBEP (Möller et al., 2012), as well as the profiles of OPEs in outdoor urban air (TBEP > TCPP > TCEP > TnBP > TPhP) in Stockholm, Sweden (Wong et al., 2018) and Turkey (TBEP > TCPP > TPhP > TEHP > TCEP) (Kurtkarakus et al., 2018). However, these results substantially differed from the report of an urban site in Shanghai that showed TCEP (0.1 - 10.1 ng m<sup>-3</sup>, 1.8 ng m<sup>-3</sup>) > TCPP (0.1 - 9.7 ng m<sup>-3</sup>, 1.0 ng m<sup>-3</sup>) > TPhP (0.06 - 14.0 ng m<sup>-3</sup>, 0.5 ng m<sup>-3</sup>) > TBP (0.06 - 2.1 ng m<sup>-3</sup>, 0.4 ng m<sup>-3</sup>) > TDCPP (Nd. - 23.9 ng m<sup>-3</sup>, 0.3 ng m<sup>-3</sup>) and only detected TBEP in 3 out of 116 samples (Nd. - 0.7 ng m<sup>-3</sup>, Nd.) (Ren et al., 2016) and reported data from the Bohai and Yellow Seas showing TCPP (43 - 530 ng m<sup>-3</sup>; 100 ng m<sup>-3</sup>, 50 ± 11%) > TCEP (27 - 150 ng m<sup>-3</sup>; 71 ng m<sup>-3</sup>, 25 ± 7%) > TiBP (19 - 210 ng m<sup>-3</sup>; 57 ng m<sup>-3</sup>, 14 ± 12%) > TnBP (3.0 - 37 ng m<sup>-3</sup>; 13 ng m<sup>-3</sup>). Li et al. (2014) determined that the primary pollutant of outdoor air in Nanjing was TCEP, and TBEP was not detected. These differences reflected significant differences in OPE production and usage in different regions, even in the same country. It should be noted that the concentrations of TCPP and TCEP were at the same level in this study, failing to indicate the industrial replacement of TCEP by TCPP, in Southwest China, this result differed from the higher concentration of TCPP than TCEP observed due to the industrial replacement of TCEP by TCPP in Europe (Quednow and Pittmann, 2009). This absence of industrial replacement was confirmed by the fact that there are manufacturers and sellers of TCEP and TCPP in Chengdu (<https://show.guidechem.com/hainuowei>, <http://www.sinostandards.net/index.php>), indicating the production of and demand for both TCPP and TCEP in this region.

删除的内容: The c

删除的内容: Non-chlorinated

删除的内容: the

删除的内容: found

删除的内容: that

删除的内容: has

删除的内容: But

删除的内容: who

删除的内容: being

删除的内容: in Turkey

删除的内容: , whereas TBEP was

删除的内容: ,

删除的内容: the

删除的内容: over t

删除的内容: showed

删除的内容: that there were

删除的内容: in

删除的内容: suggesting that

删除的内容: wasn't was not identified

删除的内容: the

删除的内容: southwest

删除的内容: ,

删除的内容: which

删除的内容: that

删除的内容: in comparison with

删除的内容: was

删除的内容: that there is

删除的内容: for



Combined with the data from 2013-2014 (Yin et al., 2015), TBEP was always the dominant OPE during the two sampling periods (2013 - 2014 and 2014 - 2015). The Kruskal-Wallis test revealed that TnBP and TCPP had no significant difference between the two sampling periods, but there were significant differences in other kinds of OPEs between the two sampling periods. This result indicated that the production and usage of individual OPEs have changed to a certain degree, suggesting that OPEs should be better investigated and governed with respect to individual compounds.

OPEs can be categorized by whether they are halogenated, alkylated or aryl OPEs. Of the OPEs measured in this study, TCEP, TCPP and TDCPP are halogenated OPEs, TBEP, TnBP and TEHP are alkylated OPEs, and TPhP is an aryl OPE. The OPEs in PM<sub>2.5</sub> at all sites were dominated by alkylated compounds (55.9 ± 10.1%), followed by halogenated OPEs (35.8 ± 9.9%) and aryl OPEs (8.3 ± 4.1%). Our results are similar to those observed in Bursa, Turkey (Kurtkarakus et al., 2018), where alkylated OPEs accounted for 68% - 95% of total OPEs, while halogenated OPEs accounted for 3.1% - 29%, and aryl OPEs accounted for 1.4% - 3.7%. Wu et al. (2020) also reported that alkyl OPEs dominated the OPE compositional profiles of urban air collected from Chicago and Cleveland. At Longyearbyen, the nonchlorinated OPE concentrations comprised 75% of the Σ<sub>8</sub>OPE concentrations (Salamova et al., 2014a). However, our results are obviously different from those of many studies, with the atmospheric samples collected in urban areas being dominated by chlorinated OPEs (50 - 80%) (Salamova et al., 2014b; Liu et al., 2016; Guo et al., 2016). In our study, nonchlorinated OPEs were dominant in urban and suburban areas across the city.

### 3.3. Seasonal and spatial variations in OPEs in PM<sub>2.5</sub>

The mean seasonal concentrations are plotted for the six sampling sites in Fig. 1. The data were quite consistent with our previous study from December 2013 to October 2014 (Yin et al., 2015). The concentrations of OPEs in PM<sub>2.5</sub> were fairly uniform throughout the three studied years. As shown in Fig. 2, the general order of decreasing average Σ<sub>7</sub>OPE concentrations in the suburban area was autumn (8.4 ± 4.3 ng m<sup>-3</sup>) ≈ winter (8.4 ± 4.5 ng m<sup>-3</sup>) > spring (7.6 ± 2.2 ng m<sup>-3</sup>) > summer (3.5 ± 1.1 ng m<sup>-3</sup>), while in the urban area, the order was autumn (9.30 ± 3.89 ng m<sup>-3</sup>) > winter (6.63 ± 3.65 ng m<sup>-3</sup>) > spring (6.36 ± 1.72 ng m<sup>-3</sup>) > summer (4.60 ± 1.91 ng m<sup>-3</sup>). The average concentration of Σ<sub>7</sub>OPEs in autumn/winter was approximately 2 times that in summer. In summer, turbulent flow accelerated the diffusion of pollutants, leading to the lowest concentration, while higher concentrations of OPEs appeared in autumn

删除的内容: of

删除的内容: year

删除的内容: OPEs

删除的内容:

删除的内容: was used and found

删除的内容: change

删除的内容: for

删除的内容: OPEs

删除的内容: the

删除的内容: whose

删除的内容: covered

删除的内容: covered

删除的内容: covered

删除的内容: of total OPEs

删除的内容: non-chlorinated

删除的内容: OPEs

删除的内容: -

删除的内容: area

删除的内容: of

删除的内容: were

批注 [Ed1]: Please ensure that the intended meaning has been maintained in this edit.

删除的内容: have been

删除的内容: in

删除的内容: past

删除的内容: the

删除的内容: OPEs

删除的内容: the

删除的内容: the

and winter because the inversion layer appeared more frequently in autumn and winter, making the diffusion and dilution of pollutants more difficult. This seasonal variation was mostly in line with that at the Shanghai urban site, with an order of autumn ( $8.4 \text{ ng m}^{-3}$ ) > winter ( $7.6 \text{ ng m}^{-3}$ ) > spring ( $5.5 \text{ ng m}^{-3}$ ) > summer ( $4.4 \text{ ng m}^{-3}$ ); in that case, the maximum value was also approximately twice the minimum (Ren et al., 2016). In addition, this finding was similar to that in Xinxiang, which showed no significant seasonal changes and only exhibited individual high values in winter. In contrast, Wang et al. (2019) found that the  $\text{PM}_{2.5}$ -bound fractions of OPEs varied significantly between seasons in Dalian, China, with their concentrations being higher in hot seasons, which may be due to temperature-driven emissions or gas-particle partitioning. Wong et al. (2018) reported that most OPEs in outdoor urban air in Stockholm, Sweden, showed seasonality, with increased concentrations during the warm period. Süring et al. (2016) reported the temperature dependence of chlorinated OPEs and 2-ethylhexyl diphenyl phosphate (EHDPP) in Arctic air. Wu et al. (2020) reported that median concentrations of  $\Sigma$  OPEs for summer samples were up to 5 times greater than those for winter samples. Similar seasonal patterns were reported by Salamova et al. (2014) for atmospheric particle-phase OPE concentrations in samples collected from the Great Lakes in 2012. A reasonable explanation is that OPEs are not chemically bound to the materials in which they are used and higher temperatures may facilitate their emission from buildings and vehicles. However, Liu et al. (2014) did not observe any temperature dependence for OPEs in urban air in Toronto, Canada. Thus, previous reports of the temperature dependence of OPEs are not consistent. In our study, correlation analysis between temperature, wind speed, wind direction and  $\Sigma_7$  OPE concentrations was performed. The results showed statistically significant negative correlations between temperature and  $\Sigma_7$  OPEs ( $R = -0.355$ ,  $p < 0.01$ ). The lowest concentrations of  $\Sigma_7$  OPEs and individual compounds were observed in summer, suggesting that the OPE level was not dominated by temperature-driven emissions. Gas-particle partitioning and local emission sources may contribute to this variation.

The most obvious difference between these results and those for coastal cities was that the concentrations of almost all OPE monomers in this study were highest in autumn/winter and lowest in summer, suggesting sustained and stable high local emissions in the studied inland city, which is particularly noteworthy. No point sources were identified in summer, and the OPE levels were diluted and diffused in summer due to the higher wind speed than that in winter in the studied inland city. This behaviour was different from that of coastal cities: Liu et al. (2016) observed the highest TCPP and TCEP concentrations,

删除的内容: resulting in the pollutants being...making more difficult to diffuse and dilute...he diffusion and dilution of pollutants more difficult. This seasonal variation was mostly in line with that at the Shanghai urban site, with an order of autumn ( $8.4 \text{ ng m}^{-3}$ ) > winter ( $7.6 \text{ ng m}^{-3}$ ) > spring ( $5.5 \text{ ng m}^{-3}$ ) > summer ( $4.4 \text{ ng m}^{-3}$ );...of which

删除的内容: that... which showed no significant seasonal changes and only exhibited individual high values in winter. On the contrary...n contrast, Wang et al. (2019) found that the  $\text{PM}_{2.5}$ -bound fractions of OPEs varied significantly between seasons in Dalian, China, with their concentrations being higher in hot seasons, which may be due to the...temperature-driven emissions or gas-particle partitioning. Wong et al. (2018) reported that most of

删除的内容: in Stockholm, Sweden

删除的内容: The similar

删除的内容: the ...atmospheric particle-phase OPE concentrations in samples collected from the Great Lakes in 2012. A reasonable explanation is that OPEs are not chemically bound to the materials in which they are used and higher temperatures may facilitate their emission from buildings and vehicles. However, Liu et al. (2014) did not

删除的内容: the

删除的内容: ...correlation analysis between the

删除的内容: OPEs...PE concentrations has been done...as performed. The results showed statistically significant

删除的内容: driven

删除的内容: the...temperature-driven emission

删除的内容: the

删除的内容: Compared to the coastal cities, t

删除的内容: OPEs

删除的内容: and concentrated

删除的内容: the

删除的内容: were ...s particularly noteworthy. No point sources was

删除的内容: OPEs

删除的内容: was ...ere diluted and diffused in summer due to the higher wind speed than that in winter in the studied

in the summer in Guangzhou, and Javier et al. (2018) found that the OPEs in spring generally exhibited the lowest concentrations in Bizerte, Tunisia, probably linked to the influence of local meteorological conditions and, to a lesser extent, air mass trajectories.

Although the Kruskal-Wallis test showed no significant variation in  $\Sigma_7$  OPE concentrations across the city, spatial differences were identified in this study. For example, TnBP and TCPP had significant differences among the six sites. In addition, higher concentrations and more dispersed patterns of most OPEs were observed in autumn and winter than in summer (Fig. 3). The concentrations of TEHP in autumn at the eastern and northern sampling sites were more dispersed than those at other sites. The same dispersion pattern was observed for TBEP in winter at the western sampling site, TPhP in autumn at the suburban sampling site, and TnBP in autumn at the eastern sampling site, suggesting that extra emission sources existed in autumn or winter. Considering the layout of Chengdu, which spreads out from the central area along the loop line (the first ring road, the second ring road and the third ring road), the uniform patterns of OPE levels and distribution across the city are understandable. Different types of industrial parks in different directions in Chengdu may be the reason for the spatial differences in OPEs. For example, in eastern Chengdu, there are automobile industrial parks and other large industrial parks, while logistics and shoemaking industrial parks are located in the suburbs. The occurrence of unexpectedly high levels of individual OPEs at different sites in autumn might indicate noteworthy emissions. The spatial and seasonal variations in individual OPEs suggest that OPE control and management measures should be taken. Interestingly, in this study, alkyl OPEs dominated at both urban and suburban sites. This finding was extremely different from the results reported by Wu et al. (2020), in which alkyl OPEs dominated at urban sites, chlorinated OPEs were prevalent at rural sites, and aryl OPEs were most abundant at remote locations.

Many studies have focused on halogenated OPEs due to their persistence, bioaccumulation, and potential human health effects, and they dominate the OPE profile in the air of many cities and other areas (Li et al., 2017). Liu et al. (2016) reported that the sum of the concentrations of three halogenated OPEs at 10 urban sites ranged from 0.05 to 12 ng m<sup>-3</sup>, suggesting that the highest production volume and widest application of OPEs have led to large emissions of OPEs in China in recent years. However, in our study, the mean concentrations of halogenated, alkylated and aryl OPEs were 2.4 ± 1.4 ng m<sup>-3</sup>, 3.7 ± 2.1 ng m<sup>-3</sup>, and 0.5 ± 0.4 ng m<sup>-3</sup>, respectively, which showed that alkylated OPEs dominated the profile of OPEs in PM<sub>2.5</sub> in Chengdu. The most notable seasonal variation was observed for alkyl phosphates, followed

删除的内容: to a lesser extent

删除的内容: .  
Though

删除的内容:

删除的内容: that there was

删除的内容: of ...n  $\Sigma_7$  OPEs ...PE concentrations across the city, the

删除的内容: the

删除的内容: difference ...ifferences among the six sites. In addition, the ...higher concentrations and more dispersed patterns of most OPEs were observed in autumn and winter than in summer (Fig. 3). The concentrations of TEHP in autumn at the eastern and northern sampling site

删除的内容: others

删除的内容: there existed the

批注 [Ed2]: Please note that the font changes here. Please consider using a consistent font type and size throughout the manuscript.

删除的内容: develops ...preads out from the central area with

删除的内容: OPEs ...PE levels and distribution across the city is ...re understandable. Different types of industrial parks in different directions in Chengdu may be the reason for the spatial differences of ...n OPEs. For example, in the east of ...astern Chengdu, there are automobile industrial parks and other large industrial parks, while logistics and shoemaking industrial parks are located in the suburbs. The

删除的内容: that there was a ...oteworthy emissions. The spatial and seasonal variations of

删除的内容: OPE suggest

删除的内容: the ...PE control and management of OPEs ...easures should be taken. Interestingly, in this study

删除的内容: the ...alogenated OPEs due to their persistence, bio-accumulation ...ioaccumulation, and potent

删除的内容: the ...hree halogenated OPEs at 10 urban sites ranged from 0.05 to 12 ng m<sup>-3</sup>, suggesting that the highest

删除的内容: the

删除的内容: In different seasons, t

by halogenated OPEs and aryl OPEs. These results were significantly different from those in other studies [that](#) reported that halogenated OPEs had the maximum seasonal variability (Guo et al., 2016; Shoeib et al., 2014).

删除的内容: which  
删除的内容: the

### 3.4. Correlation analysis of OPEs

#### 3.4.1 Linkage to environmental factors

Most OPE [monomer](#) concentrations in PM<sub>2.5</sub> have a strong linear correlation ( $R^2 = 0.79$ ) with vapor pressure (Fig. 4), suggesting that vapor pressure is an important factor controlling the levels of OPEs in PM<sub>2.5</sub>, except for TBEP. Generally, the greater the vapor pressure of [an OPE is](#), the [more easily the compound can](#) be released into the environment. Therefore, the [main](#) sources of most OPEs in Chengdu atmospheric PM<sub>2.5</sub> are production processes [including](#) OPEs and the phase transition process before they enter into the atmosphere. The boiling points of OPEs are relatively high, so they tend to be adsorbed [onto](#) PM<sub>2.5</sub> after being released to the environment (Wang et al., 2019), and their gas-particle distributions determine their concentrations in PM<sub>2.5</sub>. Interestingly, the vapor pressure of TBEP is lower than [that of](#) other OPEs, but its concentration in PM<sub>2.5</sub> was higher, which indicated [that](#) sustained and stable high emission sources, [possibly including traffic emission](#), [kept](#) its concentration at a high level (Chen et al., 2020). S [thring](#) et al. (2016) reported [that nonhalogenated](#) OPE concentrations in Canadian Arctic air appeared to have diffuse sources or local sources close to the land-based sampling stations.

删除的内容: of  
删除的内容: monomers  
删除的内容: their  
删除的内容: the  
删除的内容: speaking  
删除的内容: OPEs  
删除的内容: easier it is to  
删除的内容: mainly both from the  
删除的内容: containing  
删除的内容: in  
删除的内容: that there were  
删除的内容: to keep  
删除的内容: , which may include the traffic emission source  
删除的内容: non-halogenated

#### 3.4.2 Correlation between target analytes

Spearman's [rank](#) correlation coefficients were used to investigate the potential emission sources for OPEs [according to](#) the relationship between individual OPEs in PM<sub>2.5</sub> (Fig. 5, Table 2). Fig. 5 [shows](#) no statistically significant positive correlations between OPE monomers ( $r < 0.50$ ,  $p < 0.01$ ). However,  $\Sigma_7$  OPE concentrations were closely related to TBEP, TCEP and TnBP ( $r = 0.53-0.61$ ,  $p < 0.01$ ), which further [indicated that](#) OPE levels were influenced mainly by the [dominant OPE](#) compounds. Comparatively, weak correlations between most OPEs were observed in urban regions (Wang et al., 2018) and Turkey (KurtKarakus et al., 2018). However, strong correlations between individual OPEs were found in Guangzhou and Taiyuan (Chen et al., 2020). Further analysis results [are](#) shown in Table 2. Significant [correlations](#) between [only](#) TCP and TCEP [at](#) both downtown ( $r = 0.82$ ,  $p < 0.01$ ) and suburban sites ( $r = 0.85$ ,  $p < 0.01$ ) were observed, indicating the high

删除的内容: ranks  
删除的内容: by  
删除的内容: OPE  
删除的内容: showed  
删除的内容: OPEs  
删除的内容: identified  
删除的内容: the OPEs  
删除的内容: dominated OPEs  
删除的内容: of  
删除的内容:  
删除的内容: were  
删除的内容: Only s  
删除的内容: correlation  
删除的内容: at

homology between these two compounds. The studied inland city in China still uses a large number of products containing chlorinated flame retardants, which was confirmed by our previous study of house dust (Liu et al. 2017; Yin et al., 2019). At the downtown site, another significant correlation existed between TEHP and TCEP ( $r=0.50, p<0.01$ ), while other compounds had weak to moderate correlations ( $r<0.46, p<0.01$ ). The downtown area is mainly focused on light industry and software development, and TCPP, TCEP, TnBP, TBEP and TPhP are used in textiles, leather, electronic products and other fields. However, the correlations of each OPE monomer at site A (suburb) were stronger than those in the urban area. The correlations between TnBP and TCEP, TnBP and TDCPP, TCEP and TCPP, TCEP and TDCPP, TCEP and TBEP, TCPP and TDCPP, and TBEP were all extremely significant. This result indicated that the pollution in the suburbs was commixed and influenced by many kinds of pollution sources.

#### 3.4.3 Correlation analysis of OPEs and PM<sub>2.5</sub> concentrations

SPSS software was used to produce scatter plots to analyse the relationship between the concentrations of OPE monomers and PM<sub>2.5</sub>. As displayed in Fig. S2, only weak to moderate correlations were observed between most OPEs and PM<sub>2.5</sub>, except for a significant correlation between TDCPP and PM<sub>2.5</sub> ( $r=0.53, p<0.01$ ), which suggests that continuous and relatively constant local emissions were the main sources. This result was similar to that reported for Taiyuan (Guo et al., 2016), where no correlation was found between the concentrations of OPEs and the concentration of particulate matter. However, this result differed from that in Xinxiang (Shen et al., 2016), where the concentrations of OPEs and PM<sub>2.5</sub> had a significant correlation ( $r=0.85c$ ), and high values of OPEs/PM<sub>2.5</sub> were related to the contribution of air masses from heavily polluted areas (Henan and Jiangsu Provinces), while low OPEs/PM<sub>2.5</sub> values were, due to air masses from Shanxi-Gansu and Neimenggu Provinces. Chen et al. (2020) found that there was a significant correlation ( $p < 0.05$ ) between the concentrations of  $\Sigma_{11}$  OPEs and PM<sub>2.5</sub> at some sampling sites but not at a site located in the urban region in Guangzhou with potential additional pollution sources.

#### 3.4.4 Correlation analysis of OPEs in PM<sub>2.5</sub> and soil

Due to the low detection frequency of TCPP and TDCPP in the soil (Yin et al., 2016), the relationships of the other five OPE monomers in the soil and in atmospheric PM<sub>2.5</sub> are presented in Fig. 6. A very strong linear relationship was obtained between the OPEs in soil and in PM<sub>2.5</sub> ( $R^2 = 0.98, p<0.01$ ),

删除的内容: The ...he studied inland city in China is ...till using ...ses a large number of products containing chlorinated flame retardants, which was confirmed by our previous study of house dust (Liu et al. 2017; Yin et al., 2019). At the downtown site, another significant correlation existed between TEHP and TCEP ( $r=0.50, p<0.01$ ), while others

删除的内容: have

删除的内容: s...on the ...ight industry and software development, and TCPP, TCEP, TnBP, TBEP and TPhP are used in textiles, leather, electronic products and other fields. However, the correlations of each OPE monomer at site A (suburb) was ...ere stronger than that

批注 [Ed3]: Please note that some text appears to be missing here. Please add any missing information.

删除的内容: suburb...uburbs was commixed and influenced by many kinds of pollution sources.

删除的内容:

删除的内容: The ...PSS software was used to produce scatter diagram was ...lotsused

删除的内容: correlation...orrelations were observed between most of

删除的内容: was found

删除的内容: suggest the...uggests that continuous and relative

删除的内容: sources

删除的内容: found ...imilar with

删除的内容: from ...or Taiyuan (Guo et al., 2016), where no correlation was found between the concentrations of OPEs and the concentration of particulate matter. However, this result differed from that in Xinxiang (Shen et al., 2016), which showed that ...here the concentrations of OPEs and

删除的内容: province

删除的内容: was ...due to the

删除的内容: Province

删除的内容: in

删除的内容:

删除的内容: relationship of...relationships of the other five OPE monomers in the soil and in atmospheric PM<sub>2.5</sub>

indicating that atmospheric PM<sub>2.5</sub> settling is an important source of OPEs in the soil, just as soil is a source for OPEs in the air.

### 3.4.5 Correlation analysis of OPEs in indoor and outdoor air

The OPE profile in outdoor air in this study was TBEP > TCEP > TnBP > TCPP > TPhP > TEHP > TDCPP, which was different from that of indoor dust reported in our previous study (Liu et al., 2017): TPhP > TCPP > TnBP > TDCPP > TBEP > TCEP > TEHP. TPhP is used as an important alternative to decabrominated diphenyl ether (deca-BDE), which is typically used as a flame retardant in electrical and electronic products. In addition, plastic films and rubber may be important sources of TPhP. Thus, OPEs in indoor dust mainly come from the indoor environment and are related to human activities, not from outdoor air. In addition to the different usages of OPEs, many factors may also lead to differences between indoor and outdoor OPEs. For example, TBEP has the shortest atmospheric half-life, which may explain why its dominance in indoor samples was not observed for its outdoor counterparts. Studies in Sweden (Wong et al., 2018) reported that the concentrations of OPEs in indoor air followed the order TCPP > TCEP > TBEP > TnBP > TPhP, and those in outdoor urban air followed the order TBEP > TCPP > TCEP > TnBP > TPhP (Wong, 2018), which also indicated differences in OPE profiles in indoor and outdoor air. They found that activities in buildings, e.g., floor cleaning, polishing, construction, introduction of new electronics and changes in ventilation rate, could be key factors controlling the concentration of indoor air pollutants, while the observed seasonality for OPEs in outdoor air was due to changes in primary emissions.

## 3.5 Source apportionment of OPEs

### 3.5.1 Backward trajectory model analysis

Backward trajectory cluster analysis (HYSPLIT4), which combines the horizontal and vertical motion of the atmosphere, and can analyse the transport, migration and diffusion of atmospheric pollutants, was used in this study. The height of 500 m above ground level (AGL) can best represent the characteristics of the wind field associated with this process, and HYSPLIT4 was used to obtain the 500 m AGL 24-hour backward trajectory during the sampling period in Chengdu. During the sampling period, the air masses were mainly from the north-eastern and southern parts of Sichuan Province, including Mianyang, Deyang, Renshou and Chengdu, and a few of the trajectories came from Chongqing and other places in

- 删除的内容: the ...tmospheric PM<sub>2.5</sub> settlement
- 删除的内容: so
- 删除的内容: does the
- 删除的内容: be
- 删除的内容: OPEs profile
- 删除的内容: were
- 删除的内容: ...TBEP> TCEP> TnBP> TCPP > TPhP> TEHP> TDCPP, which was different with...rom that of indoor dust reported from...n our previous study (Liu et al., 2017); TPhP>TCPP>TnBP> TDCPP >TBEP> TCEP > TEHP. TPhP is used as one of important alternatives for
- 删除的内容: technical ...ecabrominated diphenyl ether (deca-BDE) product... which is typically used as a flame retardant in electrical and electronic products. In addition, the use of ...lastic films and rubber may be an
- 删除的内容: comes...ome from the indoor environment and are related to human activities, not from outdoor air. Except for...n addition to the different usage
- 删除的内容: lives
- 删除的内容: the
- 删除的内容: Swedish ...weden (Wong et al., 2018) reported that the concentrations of OPEs in indoor air were ...ollowed the order TCPP > TCEP > TBEP > TnBP> TPhP, and those in outdoor urban air were ...ollowed the order TBEP > TCPP > TCEP > TnBP > TPhP (Wong, 2018), which also indicated the
- 删除的内容: of OPEs profile
- 删除的内容: the ...uildings, e.g., floor cleaning, polishing, construction, introduction of new electronics and changes in
- 删除的内容: Analysis of b
- 删除的内容: The b...ackward trajectory cluster analysis (HYSPLIT4), which combines the horizontal and vertical
- 删除的内容: AGL500m
- 删除的内容: AGL...00 m above ground level (AGL) can better ...est represent the characteristics of the process
- 删除的内容: hours
- 删除的内容: of AGL500m AGL500 m ...uring the sampling period of ...n Chengdu. During the sampling

Gansu Province. Therefore, during the sampling period, Chengdu was mainly affected by air masses from eastern Sichuan.

删除的内容: the

删除的内容: of

删除的内容: the

删除的内容: the

删除的内容: mass

In different seasons, the air sources always came from the southern or northern regions of Chengdu. In spring, Chengdu was influenced by air masses from the southern region, which could be divided into three paths: (a) from Ya'an through Renshou to Chengdu; (b) from Leshan and Yibin; and (c) from Chongqing through Ziyang to Chengdu. The concentrations of OPEs at the northern and suburban sites

删除的内容: site

were relatively high in spring. During the summer period, Chengdu was mainly influenced by air masses from both the southern areas (Yibin, Zigong and others) and the northern areas (Gansu Province, Guangyuan and Mianyang), but there was no significant difference in OPE concentrations at each sampling site, nor in autumn and winter. Combined with the results of backward trajectory cluster analysis and the concentrations of OPEs at each sampling site, the concentrations of OPEs had no obvious change. This result suggested that OPEs were not affected by exogenous pollution but were mainly affected by local sources in Chengdu. These results are consistent with the meteorological and topographic conditions. Chengdu's wind has always been breezy with a much lower strength than those of coastal cities or other inland cities

删除的内容: the

删除的内容: of

删除的内容: smaller

(<https://baike.baidu.com/item/%E6%88%90%E9%83%BD/128473?fr=aladdin>). The wind direction is relatively constant, mainly from the south and the north. In addition, Chengdu is a city located in the interior of China, surrounded by the Qinghai-Tibet Plateau, the Qinling Mountains, etc. These topographic and meteorological conditions block the influence of foreign sources on Chengdu's atmosphere, which further explains why the pollution of OPEs in PM<sub>2.5</sub> was controlled by endogenous pollution, not by exogenous pollution.

删除的内容: which was

删除的内容: explained that

### 3.5.2 Principal component analysis

删除的内容: Component Analysis

Principal component analysis (PCA) of OPEs was carried out by SPSS. The normalized correlation coefficient matrix of the original data for each sampling site showed that there was a strong correlation between TCPP and TCEP, TCEP and TBEP, and TnBP and TPhP, which satisfied the condition of dimensionality reduction of PCA. Two principal component factors were obtained in this study. The cumulative contribution of the two principal component factors was 62.3%, which can basically explain the data. The results are shown in Table S1. For factor 1, there was a large load on TCEP, TCPP, and TBEP and a moderate load on TDCPP. Factor 1 can represent sources of OPEs from the plastic industry,

删除的内容: The p

删除的内容: of

删除的内容: reflect

删除的内容: information

删除的内容: were

删除的内容: the

interior decoration and traffic emissions, with a contribution ratio of 34.5% (Marklund et al., 2005; Regnery et al., 2011; CEFIC, 2002). Factor 2 has a higher load on TnBP, TEHP and TPhP. The highest load was on TnBP, which is often used as a high-carbon alcohol defoamer, mostly in industries that do not come into contact with food and cosmetics, as well as in antistatic agents and extractants of rare earth elements. TEHP can be used as an antifoaming agent, hydraulic fluid and so on. TPhP is typically used in electrical and electronic products, plastic films and rubber (Esch, 2000; Stevens et al., 2006; Wei et al., 2015). Factor 2 can be attributed to the chemical, mechanical and electrical industries, and its contribution ratio was 27.8%.

删除的内容: emission

删除的内容: the

删除的内容: or

删除的内容: considered

删除的内容: y

### 3.5.3 PMF model analysis

The basic principle of the PMF method is to decompose a sample matrix into a factor contribution matrix and factor component spectrum. The source type of the factor is judged according to the factor component spectrum, and then the contribution ratio of the source is determined. The uncertainty is estimated by three methods: bootstrapping (BS), displacement (DISP) and BS-DISP. From the 149 samples collected in Chengdu, 132 valid samples were selected to participate in the model calculation, and three factors were determined. TPhP was the only chemical with a residual (4.0) greater than 3. The concentrations of OPEs satisfied the normal distribution. The components of factor 1 were complex. Factor 1 contributed d 71.0%, 70.7% and 70.9% to TCEP, TCPP and TEHP, respectively, and 58.3% to TPhP. Factor 1 was deduced to be the plastics/electrical industry and indoor source emissions (Esch, 2000; Stevens et al., 2006). Factor 2 contributed the most to TBEP (78.0%), followed by TDCPP (44.7%), while it did not contribute to TnBP. Therefore, factor 2 was deduced to be the food/cosmetics industry and traffic emissions (Marklund et al., 2005). Factor 3 contributed d 71.7% of the total TnBP, and was deduced to be a chemical industrial source (Regnery et al., 2011).

删除的内容: the

删除的内容: disp

删除的内容: bs-disp

删除的内容: Concentrations

删除的内容: s

删除的内容: as

删除的内容: s

删除的内容: which can be

删除的内容: as

删除的内容: Implications

## 4. Conclusions and implications

Compared to the levels of OPEs in other cities, the levels of OPEs measured in this study were comparable to or even higher than those in most other studies. This result suggests that during the shift of labour-intensive manufacturing from v coastal developed areas to inland regions, OPEs were widely used in industrial and manufacturing processes in Southwest China, which should arouse concern.

删除的内容: of

删除的内容: the

删除的内容: southwest

删除的内容: needs



This intensive sampling campaign of urban and suburban areas found no significant spatial variability in  $\Sigma_7$  OPEs across Chengdu, China, but the most notable seasonal variation was observed for alkyl phosphate, followed by halogenated OPEs and aryl OPEs. Higher concentrations and more dispersed patterns of OPEs were observed in autumn/winter than in summer, with TBEP, TCEP, TCPP and TnBP being the dominant compounds. The occurrence of unexpectedly high levels of individual OPEs at different sites in autumn might indicate that there were noteworthy emissions. PCA showed that the main sources of OPEs in PM<sub>2.5</sub> include plastic industry/interior decoration/traffic emission (34.5%) and the chemical, mechanical and electrical industries (27.8%). PMF showed that the main sources were the plastics/electrical industry and indoor source emissions. OPEs have a wide range of physical and chemical properties, and combined with the differences in their behaviour, identified in this study, the management of OPEs as individual compounds instead of a single chemical class should be considered. In addition, due to the special topography and meteorological conditions of the studied inland city, the distribution and seasonal variation of OPEs in the air in this study were significantly different from those of most coastal cities and ocean locations. The sustained and stable high local emissions are particularly noteworthy. Chlorinated phosphates, especially TCPP and TCEP, which are highly toxic and persistent in the environment, had high concentrations in this study. Their usage and source emissions should be controlled.

删除的内容: area

删除的内容: of

删除的内容: pattern

删除的内容: that

删除的内容: unexpected

删除的内容: level

删除的内容: was a

删除的内容: analysis

删除的内容: decoration /traffic

删除的内容: y

删除的内容: its

删除的内容: behavior

删除的内容: that

删除的内容: over the sea

删除的内容: The chlorinated

删除的内容: have