

Pre-deliquescent water uptake in deposited nanoparticles observed with *in situ* ambient pressure X-ray photoelectron spectroscopy

Jack J. Lin¹, Kamal Raj R¹, Stella Wang², Esko Kokkonen³, Mikko-Heikki Mikkilä³, Samuli Urpelainen¹, and Nønne L. Prisle¹

¹Nano and Molecular Systems Research Unit, P. O. Box 3000, FI-90014 University of Oulu, Finland

²Division of Physics, Math, and Astronomy, California Institute of Technology, Pasadena, California, 91125, USA

³MAX IV Laboratory, Lund University, Box 118, SE-22100 Lund, Sweden

Correspondence: Nønne L. Prisle (nonne.prisle@oulu.fi); Samuli Urpelainen (samuli.urpelainen@oulu.fi)

Abstract. We study the adsorption of water onto deposited inorganic sodium chloride and organic malonic acid and sucrose nanoparticles at ambient water pressures corresponding to relative humidities (RH) from 0 to 16 %. To obtain information about water adsorption at conditions which are not accessible with typical aerosol instrumentation, we use surface-sensitive ambient pressure X-ray photoelectron spectroscopy (APXPS), which has a detection sensitivity from parts per thousand. Our results show that water is already adsorbed on sodium chloride particles at RH well below deliquescence, and that the chemical environment on the particle surface is changing with increasing humidity. While the sucrose particles exhibit only very modest changes on the surface at these relative humidities, the chemical composition and environment of malonic acid particle surfaces is clearly affected. Our observations indicate that water uptake by inorganic and organic aerosol particles could already have an impact on atmospheric chemistry at low relative humidities. We also establish the APXPS technique as a viable tool for studying chemical changes on the surfaces of atmospherically relevant aerosol particles which are not detected with typical online mass- and volume-based methods.

Copyright statement. TEXT

1 Introduction

The interaction between atmospheric particulate matter and water is one of the most important processes in Earth's atmosphere. The amount of water associated with an aerosol particle is a function of its intrinsic hygroscopicity and the ambient relative humidity (RH). Liquid water comprises a significant fraction of the global aerosol mass with field measurements documenting the global presence of the metastable aerosol phase state (Rood et al., 1989; Nguyen et al., 2016). Condensed water in the atmosphere critically influences both direct and indirect climate effects of aerosols, governed by aerosol growth and light scattering and via the activation of aerosol particles into cloud droplets (Kreidenweis and Asa-Awuku, 2014).

The surfaces of aerosol particles and droplets are distinct physical and chemical environments compared to their associated bulk phases. Reaction rates in micrometer-scale droplets have been measured (Jacobs et al., 2017; Marsh et al., 2019; Zhang

et al., 2020b) and modeled (Benjamin, 2019; Mallick and Kumar, 2020) to be higher than those in bulk water with some reactions even proceeding spontaneously (Lee et al., 2019). For cloud and fog systems where the interfacial region makes up a significant fraction of the condensed aqueous phase, the reaction rate at the surface can be the rate-limiting step in multi-phase
25 OH oxidation involving surface-active organic species such as pinonic acid (Huang et al., 2018). Interfacial water molecules can promote reactions between organic acids and SO_3 , which are distinct from those that occur in the gas phase and important for heterogeneous formation of H_2SO_4 and subsequent new particle formation (Zhong et al., 2019; Lv and Sun, 2020). The bulk/surface partitioning of saccharide molecules in aqueous solution affects their ability to be oxidized by OH (Fan et al., 2019). The presence of surface-enriched nonanoic acid also increases the surface concentration of Fe(III) in aqueous mixtures
30 which mediates enhanced photochemical release of volatile organic compounds (Huang et al., 2020).

The composition of the droplet surface can influence the mass transport and chemical reactions that occur at the surface (e.g. Cosman et al., 2008; Park et al., 2009; Roy et al., 2020). The acidity of organic acids on water surfaces has been measured to be much lower than predicted for the bulk aqueous phase (Enami et al., 2010; Prisle et al., 2012; Öhrwall et al., 2015; Werner et al., 2018). Uptake of glyoxal has been shown to be enhanced at acidic interfaces (Shi et al., 2020). Changes in
35 droplet size via evaporation or condensation can affect the chemical reactions that take place at the surface via changes in the relative dimensions and chemical environments of the interface and associated bulk phases (Prisle et al., 2012; Djikaev and Ruckenstein, 2019; Petters et al., 2020). The presence of surface-active organic molecules on droplet surfaces can also affect droplet surface tension (Shulman et al., 1996; Prisle et al., 2010b; Bzdek et al., 2020) and morphology (Kwamena et al., 2010) that affect both warm (Sareen et al., 2013; Ovadnevaite et al., 2017) and ice cloud nucleation (Knopf and Forrester, 2011;
40 Perkins et al., 2020) as well as droplet coalescence (Pak et al., 2020).

The formation of an aqueous aerosol phase can lead to the partitioning of water-soluble gases to the condensed phase (Prisle et al., 2010a), including many reactive oxidants (Donaldson and Valsaraj, 2010), that can initiate a wide range of aqueous phase chemistry (McNeill, 2015). The mobilization of ions in aqueous solution has the ability to influence atmospheric chemistry (Cwiertny et al., 2008). In particular, the mobilization of chloride ion from sea salt aerosol is a significant source of chlorine gas
45 in the troposphere, with the subsequent formation of chlorine radicals affecting the budgets of important atmospheric species such as volatile organic compounds, ozone, OH, and nitrogen oxides (Wang et al., 2019). Recent field studies have detected the depletion of chloride and bromide from marine aerosol particles under the influence of acidic species from wildfire emissions (Braun et al., 2017). A number of aqueous phase reactions occur between inorganic salt species and organic compounds. The hygroscopic properties of sodium halide particles coated with fatty acids depend on both the salt anion and the carboxylic acid,
50 with some mixtures showing barriers to water uptake while others do not (Miñambres et al., 2014). Enhanced production of sulfate aerosol via nitrate photolysis was observed to be facilitated by the presence of surface-active halide ions (Zhang et al., 2020a).

A wide range of experimental techniques are available to study the hygroscopic properties of particles in sub-saturated (< 100 % RH) conditions, including humidity-tandem differential mobility analyzers (H-TDMA), various optical extinction and
55 scattering methods, physisorption analyzers, quartz crystal microbalances and various microscopic and spectroscopic methods (Kreidenweis and Asa-Awuku, 2014; Tang et al., 2019). A drawback of many of the otherwise highly valuable traditional

aerosol techniques, such as the H-TDMA or optical methods, is that they are based on detection of growth in mass or volume. When applied alone, these methods are not sensitive enough for detecting minute amounts of water adsorbed onto the aerosols at low humidity or below the deliquescence relative humidity (DRH). Furthermore, most of these methods do not provide molecular-level information about the system and are insensitive to surface-specific chemistry and the phase state of adsorbed water. This can lead to underestimating the amount and impact of water adsorption at lower RH. The equilibrium between water and the particle phase has been shown experimentally to depend on the particle phase state and humidity history, with important implications for particle hygroscopic growth and cloud droplet activation (Bilde and Svenningsson, 2004).

A number of spectroscopic techniques have been applied to systems of environmental and atmospheric relevance (see e.g. the reviews of Ault and Axson, 2016; Tang et al., 2019). Surface-sensitive techniques such as X-ray photoelectron spectroscopy (XPS) provide molecular-level information on the chemical composition and properties of surfaces. So far, most of these studies, in particular those employing XPS, have adopted a surface science approach using single crystal surfaces as simplified model systems (Tang et al., 2019). For example, XPS has been used to study the photochemistry of TiO₂ (110) (Lampimäki et al., 2015), ionic mobility on NaCl (001) surfaces induced by water adsorption (Verdaguer et al., 2008) and reaction of water vapor with MgO (001) surfaces (Kaya et al., 2011). XPS on a liquid micro-jet (Winter, 2009) has also been used to study surface-specific chemistry of aqueous solutions comprising atmospherically relevant organic surfactants (Prisle et al., 2012; Werner et al., 2014; Öhrwall et al., 2015; Walz et al., 2015, 2016; Toribio et al., 2018; Werner et al., 2018; Ammann et al., 2018) as macroscopic model systems for atmospheric microscopic water droplets. Experiments on single aerosol particles are, on the other hand, scarce (Antonsson et al., 2015; Shakya et al., 2016; Ouf et al., 2016). As some of the few examples of atmospheric relevance, XPS has been employed to study the surface composition of size-segregated ambient aerosol collected from an urban environment (Cheng et al., 2013). We have previously studied solvation of RbBr in free flying water clusters (Hautala et al., 2017b), as well as size-dependent structural phase changes in CsBr (Hautala et al., 2017a), as model systems for salt clusters of higher environmental relevance. Surface-enhanced Raman spectroscopy (SERS) has also been employed to study the surface composition of atmospherically relevant particles generated via electrospray (Gen and Chan, 2017; Gen et al., 2019). Using environmental transmission electron microscopy, Wise et al. (2008) observed significant amounts of water uptake onto NaCl particles at humidities as low as 70 % RH, well below the deliquescence RH. With the advent of high-brilliance synchrotron radiation and recent developments in electron analyzer technology, previous barriers of too low sample density can be overcome, enabling XPS studies under realistic conditions, *in situ* and *operando*, using the so-called ambient pressure XPS (APXPS) technique (Salmeron and Schlögl, 2008; Ogletree et al., 2009; Starr et al., 2013; Kong et al., 2020).

In this work, we study the pre-deliquest water adsorption to laboratory-generated, deposited sodium chloride (NaCl), malonic acid, and sucrose particles. NaCl is a major component of sea salt, which is the most abundant aerosol species by mass in the atmosphere (Murphy et al., 2019). Salt aerosols, apart from being important light scatterers, also take part in atmospheric chemistry by interacting with atmospheric trace gases as a source of halogens (Rossi, 2003), especially in the presence of water in the aerosols. The water uptake and deliquescence of salt aerosol particles can be affected by the presence of other inorganic or organic matter. For example, the presence of malonic acid in NaCl aerosol will not only lower the DRH of the aerosol, but also facilitate the depletion of chlorine (Laskin et al., 2012). Organic material comprises a large fraction of the

ambient aerosol mass (Kanakidou et al., 2005), and malonic acid and sucrose represent hygroscopic organic compounds with different chemical functionalities. Dicarboxylic acids such as malonic acid have been identified in ambient aerosols samples where they can dominate the water-soluble organic fraction (Khwaja, 1995; Yu et al., 2005; Decesari et al., 2000, 2001) and are
95 hygroscopic in both sub- and supersaturated conditions (Prenni et al., 2001; Hori et al., 2003; Rissman et al., 2007; Pope et al., 2010). Sucrose is studied as a model carbohydrate that can form a glassy state in response to changing relative humidity with effects on its ability to act as either cloud or ice nuclei (Zobrist et al., 2011; Estillore et al., 2017). Utilizing APXPS, interaction with water vapor at the particle surface was studied at relative humidities between 0 and 16 %. To the best of our knowledge, this is the first time XPS has been used on sampled nano-scale particles of immediate atmospherically relevant composition at
100 ambient pressure conditions.

2 Experimental

We employed the APXPS technique to obtain chemically specific information about the composition of the nanoparticle surfaces. Photoelectron spectroscopy utilizes the photoelectric effect, by which the sample is ionized from inelastic collisions with photons and the emitted electrons are detected and characterized in terms of their kinetic energy (E_k). When the ionizing photon energy ($h\nu$) is known, the binding energy (E_b) of electrons within the sample can be determined simply as $E_b = h\nu - E_k$.
105 By using X-ray photons, core-level atomic-like orbitals can be ionized, and the electron binding energy gives a very sensitive fingerprint of the chemical composition and environment of the sampled region. XPS is furthermore a highly surface-sensitive technique, because the resulting kinetic energies of the photoelectrons yield very short characteristic attenuation lengths and the detected photoelectron signal therefore originates mainly from the topmost few nanometers of the sample. An XPS measurement consists of measuring the intensity of photoelectrons emitted from the sample as a function of the characterized electron
110 kinetic energy. Typically, an XPS spectrum presents the photoelectron signal intensity as function of the orbital binding energy and consists of a collection of peaks, each corresponding to a different chemical species or specific environment, which is identified by the spectral position in terms of the binding energy. Here, we quantify the spectral peaks in terms of their areas, which are directly proportional to the relative abundances of each species or environment on the surface of the sample. Spectral
115 fitting techniques are employed to obtain accurate results for both binding energies and peak areas.

Experiments were carried out at the APXPS end station (Schnadt et al., 2012; Knudsen et al., 2016) of the SPECIES beamline (Urpelainen et al., 2017) at the MAX IV Laboratory in Lund, Sweden. The end station is equipped with a hemispherical SPECS Phoibos NAP-150 electron energy analyzer and allows for measurements both at ultra-high vacuum (UHV) conditions, as well as up to 25 mbar in an ambient pressure (AP) cell. During the experiments, the SPECIES beamline was still under construction,
120 and we used a double anode (SPECS XR-50) X-ray source with Al (1486.6 eV) and Mg (1253.6 eV) anodes for exciting the samples instead of synchrotron radiation.

Samples were prepared at the Lund University Aerosol Lab and deposited on either silicon or gold substrates. Deposited samples were kept and transferred to the SPECIES end station in a desiccator in order not to expose them to ambient humidity for extended periods of time prior to the experiments. For the XPS measurements, the substrates were mounted onto stainless

125 steel sample holders using an adhesive Cu tape in a clean tent environment and loaded into the end station. The experiments consisted of recording C 1s, O 1s (sucrose and malonic acid), Na 1s and Cl 2p (NaCl) core level XPS spectra of the deposited aerosol particles at UHV and *in situ* at varying RH conditions. The Al anode was used to measure spectra from NaCl while the Mg anode was used to measure spectra from the malonic acid and sucrose samples.

A detailed account of the sample preparation process and experimental conditions is given in the following sections.

130 2.1 Sample preparation

Aerosol samples were generated by nebulizing aqueous solutions of either sodium chloride (NaCl), sucrose ($C_{12}H_{22}O_{11}$), or malonic acid ($CH_2(COOH)_2$). Solutions were prepared using ultra-pure Milli-Q water. All chemicals were obtained from Sigma-Aldrich and used without further treatment.

The air flow containing solution droplets (3 L min^{-1}) was mixed with dry, particle-free air (3 L min^{-1}) in a 3 L aerosol mixing chamber. From the mixing chamber, $1\text{--}1.5\text{ L min}^{-1}$ was sent through a diffusion dryer followed by a ^{63}Ni bipolar charger. The dried and charge neutralized aerosol flow was divided between a scanning mobility particle sizer (SMPS; TSI 3936) for size distribution characterization and a nanometer aerosol sampler (NAS; TSI 3089) for sample collection of the entire dry aerosol particle size distribution. Sodium chloride particles were collected onto silicon (with native oxide) wafers, while sucrose and malonic acid particles were deposited onto a gold film substrate.

140 The surface coverage on the substrate in the NAS is a function of the particle concentration in the gas flow, the drift velocity of the particles in the electric field, and the total sampling time (Preger et al., 2020). Particles are assumed to be spherical across the entire size distribution with a charge distribution described by Boltzmann statistics. We further assume uniform, 100 % deposition efficiency of the positively charged fraction with values above 100 % indicating the presence of more than one monolayer of deposited particles. Details of the sample generation are given in Table 1.

Table 1. Aerosol sampling data including generated size distribution information (geometric mean number μ_N , geometric number standard deviation σ_g , total number N , and geometric mean surface area μ_{SA}) and sample collection parameters (sampler flow rate Q , collection time t , substrate, and coverage).

Compound	Mean Size Distribution				Sample Collection			
	μ_N (nm)	σ_g	N (cm^{-3})	μ_{SA} (nm^2)	Q (L min^{-1})	t (min)	Substrate	Coverage (%)
Sodium chloride	72	2.0	3.94×10^5	195	1.1	42	Si	232
Sucrose	79	2.2	3.59×10^4	202	1.1	90	Au	56
Malonic acid	52	2.1	1.02×10^5	193	1.2	275	Au	245

145 2.2 XPS measurements

UHV XPS spectra were recorded using a pass energy of 50 eV and an entrance slit of 3 mm × 20 mm. This contributes to an experimental broadening of approximately 500 meV in addition to the natural broadening from the excitation source (850 meV for Al K_{α} and 680 meV for Mg K_{α} radiation).

For measurements at humid conditions, samples were transferred from the UHV manipulator to the AP cell. Milli-Q water vapor was let into the AP cell through a high precision leak valve. The pressure inside the cell was kept constant using a valve connected to a pump in a feedback loop with an absolute capacitance manometer. All measurements were made at 25 °C at which the saturation vapor pressure of water is $P_{\text{sat}} = 31.73$ mbar. The relative humidity was calculated using the (water) vapor pressure inside the AP cell so that $\text{RH} = P_{\text{cell}}/P_{\text{sat}}$. While samples were inside the cell, spectra were recorded at different water vapor pressure conditions (NaCl: 0, 2, 5, and again at 0 mbar; sucrose 0.2, 1, and 5 mbar; malonic acid 0.2 and 1 mbar). The water vapor pressures of 0, 0.2, 1, 2, and 5 mbar correspond to relative humidities of 0, 0.63, 3.2, 6.3, and 16 %, respectively. These relative humidities are well below the DRH for NaCl (75.3 % (Tang and Munkelwitz, 1993)), malonic acid (72.1 % (Parsons et al., 2004)), and sucrose (85.7 % (Yao et al., 2011)). Any remaining air was removed from the water by several freeze-pump-thaw cycles before introducing the vapor into the AP cell. The cell was also purged with dry nitrogen gas in order to remove any excess water after the experiment. In the case of NaCl particles, the sample was heated up to 125 °C using a button heater placed behind the sample holder. This was done after the sample was dosed up to 10 mbar to see if the changes incurred on exposure to water vapor were reversible. Spectra obtained at 10 mbar, however, are not included in the analysis below due to very low signal to noise ratios.

The electron count rates inside the AP cell are reduced when compared to measurements in UHV conditions due to the attenuation of X-ray intensity by the SiN_3/Al window, the limited aperture of the differential pumping of the electron analyzer and scattering of the photoelectrons from the vapor (Knudsen et al., 2016). In order to increase the count rates, spectra recorded inside the cell were acquired at a pass energy of 100 eV instead of 50 eV as used in UHV conditions in order to compensate for the reduced intensity. The analyzer broadening when using 100 eV pass energy was approximately 1.00 eV. Each spectrum was recorded 50-100 times in static conditions for increased statistics. To check the reproducibility of the measurements, we first verified that the individual spectra had not drifted in energy with time, after which all spectra recorded for a given sample and humidity condition were averaged for the final representation.

2.3 Data Analysis

Recorded XPS spectra were fitted in order to accurately determine the binding energies of the core electrons and the relative amounts of the probed elements under different conditions. Chemical identities and environments were identified from binding energies obtained from the XPS peak positions and their relative amounts by determining the corresponding peak areas. Data analysis was performed using the Igor Pro software (WaveMetrics, Inc, Lake Oswego, OR, USA). A Shirley-type background was removed from the data before fitting the peaks using the SPANCF curve fitting macro package (Kukk et al., 2001, 2005). All spectra were fitted using symmetric Voigt line shapes. A linear background was included in all the fits to remove any

residual background after the Shirley-type background removal. The energy scales of the spectra were calibrated using the well-known binding energies of Si 2p and Au 4f recorded for the silicon and gold substrates, respectively, as the low aerosol
180 particle coverage allowed simultaneous measurement of the substrate.

The aim of the spectral fitting procedure is to characterize the measured XPS spectra in terms of peak position and area. The position of a given peak gives the binding energy of the core electron for each probed element. Changes in the binding energy as well as the width of the fitted peaks—or peak broadening—can indicate changes in the chemical environment or physical state of the sampled surface. The area of the peak is directly proportional to the amount of the element being measured. For the
185 analysis here, we determine the elemental composition of particle surfaces as the relative ratios of the core level peak areas. The peak area of the XPS signal depends on a number of factors, including experimental parameters of the incident radiation and electron spectrometer as well as physical and environmental properties affecting the orbital from which the photoelectron originated. If all of these parameters are known, the XPS signal can be used to quantify the amount of species *i*. While these parameters are not always known, comparison of XPS signals is still possible to quantify relative differences in elemental
190 abundances and chemical states between experimental conditions. Before extracting relative ratios of the peaks, all spectra were normalized to the photoionization cross section (Yeh and Lindau, 1985) of the given core electron. The attenuation of photoelectron intensity due to scattering of the photoelectrons from the water vapor was estimated by using the kinetic theory formulation (Ogletree et al., 2009) and measured electron scattering cross section data (Muñoz et al., 2007). The attenuation must be taken into account, because the fixed excitation energy from the X-ray source leads to significantly different kinetic
195 energies of the emitted photoelectrons and consequently different mean free paths in the vapor environment.

3 Results and discussion

Below we present experimental XPS spectra together with details and results of curve fitting for deposited NaCl, sucrose and malonic acid aerosol nanoparticles. Binding energy shifts of varying degrees from the dry conditions are observed with increasing relative humidity, indicating changes in the chemical environment of the target element. In addition, changes in the
200 stoichiometry of molecules on the particle surface can be seen from the intensity ratios between different core level XPS peaks. This is a sign of changes in the chemical composition on the surface of the deposited particles. These observations are clear indications that water is already being adsorbed onto the particles and interacting with molecules on their surfaces at these low relative humidity conditions. The implications and possible interpretations of these observations are discussed below.

3.1 NaCl

205 Figures 1 and 2 show the recorded Na 1s and Cl 2p spectra of NaCl aerosol particles deposited on a Si substrate measured at 0, 2, 5, and again at 0 mbar (0, 6.3, 16, and 0 % RH, respectively) water vapor pressure. Spectra recorded at UHV are shown in Figs. S2 and S3 of the supplement, since they provide the same information as the measurements made in the AP chamber at 0 mbar water vapor pressure. The Na 1s spectra were fitted using a single, symmetric Voigt peak, except for the 0 mbar spectrum after water exposure, where two peaks were required in order to obtain a fit that represents the measured spectrum. The Cl 2p

210 spectra were fitted with two peaks representing the 1/2 and 3/2 spin-orbit components. The O 1s region for the NaCl sample was also monitored during the experiments, but the spectra are dominated by the signals from the native oxide of the Si wafer and the vapor phase water, and no direct signs of adsorbed or liquid water could therefore be observed from these spectra at low RH. These spectra are therefore omitted from the discussion below.

After calibrating the energy scale using the substrate Si 2p peaks, the binding energy of the Na 1s core level at UHV
215 conditions was determined to be 1073.48 eV and is indicated by the dashed vertical line in Fig. 1. The signal-to-noise ratio in the spectra recorded at 5 mbar (16 % RH) is very low and the fitted line should be regarded more as guiding the eye than an accurate fit to the peak. The binding energy of the Cl 2p 3/2 component at UHV conditions was determined to be 200.3 eV shown by the dashed vertical line in Fig. 2.

The binding energies measured here for Na 1s and Cl 2p in deposited NaCl nanoparticles at UHV conditions are higher than
220 the values typically reported in the literature (e.g. Beard, 1993). When binding energy shifts are determined for humidified conditions, they are calculated relative to the binding energy measured here at 0 mbar, and not relative to the literature value. The loss of electrons due to photoionization from the X-ray beam leads to positive charging of the sample surface since the silicon substrate is a semiconductor. The charging of the sample causes an apparent increase in the binding energies seen here. The addition of ambient gas during APXPS measurements can offset the effect of surface charging since photoelectrons from
225 the photoionization of the gas can travel to the sample surface and compensate for the positive charge.

3.1.1 Binding energy shifts: changes in chemical environment

Figure 3 shows the binding energy shifts relative to dry conditions of the Na 1s and Cl 2p core level peaks (red diamonds and dots, respectively) as a function of water vapor pressure at 25 °C. We observe a significant shift of approximately 1.1 eV in both Na 1s and Cl 2p binding energies already when water vapor is introduced into the system at very low relative humidities.
230 The observed shifts are greater than the error estimates in the peak positions shown in Fig. 3 and therefore cannot be entirely explained by measurement or fitting uncertainty. After this first rapid decrease, the Cl 2p binding energies reach a plateau, when the relative humidity is increased to 16 %. The binding energy shifts for Na 1s and Cl 2p between 0 % and 6.3 % RH are nearly identical, but the subsequent shift for Na 1s between 6.3 % and 16 % RH is towards larger binding energies by 0.17 eV, while Cl 2p shifts towards lower binding energies by -0.26 eV.

235 Some of the shift in binding energy can be attributed to the introduction of water vapor lessening the effect of charging of the sample compared to the measurements at 0 mbar, but since the amount of ambient gas changes between the measurements, we cannot deconvolute the individual effects of condensed water and gas-phase water on the binding energy shift. However, the binding energy shifts observed here are in good agreement with those reported for Na 2s and Cl 2p in NaCl (001) single crystal surfaces by Verdaguer et al. (2008), even showing a similar plateau at intermediate pressures. The measurements by
240 Verdaguer et al. (2008) were done at constant water vapor pressure and the similarities in the observed binding energy shifts with relative humidity between their work and ours therefore suggest that condensed water also plays a role in the binding energy shift observed in the present measurements. As Verdaguer et al. (2008) did not report the magnitude of the binding energy shift from 0 to 5 % RH, we have shifted their 5 % RH data point to coincide with our data in Fig. 3 for easier qualitative




Figure 1. Na 1s XPS spectra of NaCl aerosol particles recorded at different water vapor pressures (relative humidities, RH). The point markers show the recorded data and the solid lines the fit envelope curve. The spectra are fit using a single, symmetric Voigt peak shown in red with an additional peak in blue necessary to explain the spectrum at 0 mbar after water exposure. The dashed vertical line shows the binding energy of Na 1s at 0 mbar pressure (0 % RH) at the beginning of the experiments. Error bars show the estimated uncertainty in the peak position from Monte Carlo analysis. Photon energy was 1486.6 eV from the Al anode.

comparison of the two data sets. The shift of the Na 1s peak for deposited aerosol in the present work towards larger binding
 245 energies at 16 % RH is not entirely in line with the observations on NaCl (001), which show monotonic shifts towards lower
 binding energies both for Na 2s and Cl 2p peaks. However, the poorer quality of the experimental spectrum for aerosol Na
 1s at 16 % RH in the present work, compared to spectra recorded at lower RH, leads to larger uncertainty in the peak fitting
 procedure. Furthermore, the shifts observed for NaCl nanoparticles in the present work are larger for Cl 2p than for Na 1s, as
 opposed to larger shifts (by approximately 50 meV) for Na 2s than Cl 2p in the NaCl (001) single crystal.

250 In our experiments, the water vapor was removed from the AP cell after the measurements at 16 % RH and another set of
 spectra was recorded at 0 % RH. We see that when RH decreases from 16 % to 0 %, the binding energies of the Na 1s and Cl




Figure 2. Cl 2p XPS spectra of NaCl aerosol particles recorded at different water vapor pressures (relative humidities, RH). The dots show the recorded data, the solid lines the fit envelope curve and the red and blue peaks the fitted 3/2 and 1/2 spin-orbit components, respectively. The dashed vertical line shows the binding energy of Cl 2p 3/2 component at 0 mbar pressure (0 % RH) at the beginning of the experiments. Error bars show the estimated uncertainty in the peak position from Monte Carlo analysis. Photon energy was 1486.6 eV from the Al anode.

2p do not shift back to their original values, but a memory effect of approximately 0.2 eV for both Na 1s and Cl 2p is observed. Since this observed memory effect is greater than the energy step size of 0.1 eV used in the acquisition of the spectra which is in turn greater than the energy accuracy of the electron analyzer when operating at 50 or 100 eV pass energy, we consider it to
 255 be a real effect that cannot be explained solely by experimental uncertainty. This is in line with the observations by Verdaguer et al. (2008) who also report a memory effect of less than 0.5 eV for single crystal NaCl (001).

3.1.2 Changes in spectral peak width

In addition to the shifts in binding energies, we also investigated possible changes in the width of the XPS peaks due to the effects of water adsorption to the particle surfaces. Changes in the spectral peak widths can indicate changes in the chemical




Figure 3. Binding energy shifts relative to dry conditions of the core level peaks for NaCl, sucrose and malonic acid particles, as functions of water vapor pressure (relative humidity). The orange traces extending above 60 % RH show the data from Verdaguer et al. (2008) for single NaCl (001) crystal that has been aligned to the shifts obtained in this study.

260 environment or charge carrier mobility on the aerosol surfaces. While we found clear binding energy shifts (Fig. 3), we did not observe any significant changes in the peak width of either the Na 1s or the Cl 2p peaks. In contrast, the data obtained by Verdaguer et al. (2008) show decreasing total widths of Na 2s and Cl 2p peaks for NaCl (001) single crystal with increasing relative humidity, qualitatively following the behavior of their observed core level binding energy shifts.

The difference between deposited aerosols and single crystal spectra may, at least partially, be explained by the experimental 265 resolution, which is lower in the present study with peak widths dominated by experimental factors. In the NaCl (001) case, the observed binding energy shifts are attributed to the discharging of the surface, where increasing relative humidity causes the surface to be gradually discharged due to ionized water vapor and secondary electrons. As water is adsorbed onto the surface, it leads to solvation of surface ions and increased mobility of these charge carriers. This ion mobility makes discharging more efficient, leading to further shifts in observed binding energies. According to Verdaguer et al. (2008) these solvation effects 270 become more significant above approximately 35 % RH.

This interpretation is further supported by scanning force microscopy studies and infrared studies that show modifications in the surface structure and sudden increase of water coverage at around 40 % RH (Dai et al., 1997; Peters et al., 1997; Foster and Ewing, 2000). Kelvin probe microscopy (KPM) experiments by Cabrera-Sanfelix et al. (2007), Verdaguer et al. (2005), and Verdaguer et al. (2008) show how water vapor affects the surface potential of the NaCl (001) surface. They found variations on 275 the order of 0.1 V to 0.25 V across the RH range of their experiments, which clearly is not sufficient to explain the energy shifts they observed. The variations in surface potential also cause peak broadening due to slight differences to the kinetic energies of photoelectrons from different regions of the surface. The decrease in peak broadening observed for single crystals is ultimately

attributed to adsorbed water reducing inhomogeneities in the surface potential. These inhomogeneities originate from potential differences between and within the step and terrace sites of the crystal. The KPM experiments show that the inhomogeneities are removed as water adsorbs at very low surface coverage below that of one monolayer. In the present work, the deposited NaCl aerosol particle samples are most likely very far from being perfect crystals, possibly containing a large number of steps, terraces and kinks leading to larger variations in the surface potential and thus increased broadening of the peaks in any humidity condition compared to a NaCl (001) crystal. This could, together with the moderate experimental resolution, explain why no decrease in the broadening of the peaks is observed for the aerosol particles at low RH.

Results of our study show, not surprisingly, that the aerosol particle samples investigated have more complex morphology than the simple single crystal surfaces previously studied by XPS. Several previous studies have observed that the process of drying an aerosol can indeed affect its crystalline form. For example, studies of NaCl aerosol particles generated from drying of aqueous droplets have inferred a non-crystalline structure with pores or pockets that trap liquid water (Weis and Ewing, 1999; Darr et al., 2014; Braun and Krieger, 2001). This is explained by the presence of liquid water detected below the deliquescence relative humidity but still at much higher RH than in our study. Furthermore, the morphology of NaCl particles expressed via the shape factor has been shown to be controlled by the drying rate (Wang et al., 2010). A recent study (Archer et al., 2020) has explained the morphology of particles formed from drying of a colloid as a competition between diffusion of solute in solution versus loss of solvent, with higher solvent loss rate compared to solute diffusivity leading to more complex morphologies. For atmospheric samples, microscopy studies on sea salt particles have shown them to have complex morphologies (e.g. Cheng et al., 1988), similarly to what was found for the laboratory generated aerosol samples in the present study.

Atmospheric aerosols are likely to undergo multiple drying and humidification cycles under a wide range of conditions and thus to exhibit a range of morphologies related to drying. Our measurements on aerosol particle samples generated from nebulization and subsequent dessication are therefore expected to much more closely represent the morphologies of atmospheric aerosols, compared to the simple uniform morphologies of single-crystal samples.

3.1.3 Changes in peak area ratios: chemical changes in surface composition

Considering the relative amounts of Na and Cl on the particle surfaces, we observe a decrease in the Na 1s to Cl 2p peak area ratio, even after taking differences in photoionization cross section and attenuation due to scattering in the water vapor into account. This change as a function of increasing RH is presented in Fig. 4. The extracted Na to Cl ratio of 0.68 at 0 % RH and decreases slightly to 0.51 and 0.58 when the RH is increased to 6.3 % and 16 %, respectively. The ratio measured at 0 % RH after dosing was determined to be 0.62 which is close to the value before exposure to water vapor. As the spectra for all RH are recorded with a constant excitation energy from the Al $K\alpha$ anode, different transmission through the electron analyzer at different electron kinetic energies cannot explain the difference between the Na to Cl peak area ratio and the NaCl stoichiometric ratio. Due to the constant relative transmission, this could lead only to a constant ratio differing from the stoichiometric ratio. Furthermore, the reported transmission of the analyzer is very nearly the same for the Cl 2p and Na 1s regions (SPECS Surface Nano Analysis GmbH). This correlation between changes in the chemical composition of the surface layer and RH is therefore a clear indication of water being adsorbed to the NaCl particle surfaces. Assuming that

the thickness of the layer of water molecules adsorbed on the NaCl surfaces is approximately 2.4 Å (Cabrera-Sanfeliix et al., 2007), photoelectrons emitted from the NaCl particles need to pass through this distance to escape from the sample and would be attenuated by collisions with the water molecules in the process. In our experimental conditions, the photoelectrons have kinetic energies (rounded to the nearest ten) of approximately 410 eV and 1290 eV for Na 1s and Cl 2p, respectively. The two core level signals will therefore be attenuated differently due to the different inelastic mean free paths of the electrons, approximately 2 nm and 5 nm, respectively (Emfietzoglou and Moscovitch, 2002).

To quantify the attenuation of the photoelectron signal, we use an exponential decay function $I_n = I_n^0 e^{-x/\lambda_n}$, where I_n is the attenuated intensity of peak n , I_n^0 is the corresponding unattenuated intensity, x is depth into the sample from where the signal originates, and λ_n the energy-dependent inelastic mean free path of photoelectrons contributing to peak n . The depth of origin can be expressed as $x = \frac{\lambda_1 \lambda_2}{\lambda_1 - \lambda_2} \ln R$, where $R = \frac{I_1 I_2^0}{I_2 I_1^0}$ is the relative ratio of attenuated and unattenuated signals from two separate peaks $n = 1, 2$. We here use the total integrated peak areas to represent signal intensities. In our experiments, the unattenuated signal ratio (measured without water vapor) for Na 1s and Cl 2p is $I_{\text{Na}}^0 / I_{\text{Cl}}^0 = 0.9$. With this, from the corresponding signal ratios at elevated humidities, the simple attenuation model gives depths of photoelectron origin (or water layer thickness) of approximately 14 and 4 Å for 6.3 and 16 % RH, respectively. This is much more than the previous observations of 2.4 Å by Cabrera-Sanfeliix et al. (2007) and also counterintuitive as it would mean decreasing layer thickness with increasing RH. Furthermore, this simple attenuation model implicitly assumes at least a full monolayer coverage so that all signal is uniformly attenuated by the adsorbed water layer. Previous studies on NaCl (100) crystals (Peters et al., 1997; Peters and Ewing, 1997; Foster and Ewing, 2000) have shown that a full water monolayer does not develop until approximately 35 % RH. On the other hand, it has been shown that NaCl (100) does not adsorb water strongly whereas small particles (1-10 μm) have a propensity for adsorbing large amounts of water (Ghosal and Hemminger, 2004). This strong adsorption has also been connected to observations of water remaining on the particle surface, even at elevated temperatures in vacuum. Indeed, this is also observed in the present work in the form of a memory effect in the Na 1s and Cl 2p peak shifts as the humidity is decreased to 0 % RH: the peak shift does not completely disappear, even after heating the sample to 125 °C (Fig. S1 in the supplement). The signs of stronger water adsorption in the present work could therefore be attributed to the small size of our sampled NaCl particles, contributing to larger surface area (as well as more crystal imperfections) and therefore to a larger number of possible surface adsorption sites.

While attenuation observed due to the adsorbed water layer is one possible explanation for the decreasing Na to Cl ratio observed in this work, another, complementary, explanation could be the preferential dissolution of Cl⁻ ions into the water film. Spatial segregation of the Na⁺ and Cl⁻ ions could further enhance the difference in the attenuation between Na 1s and Cl 2p XPS signals and is likely to be further enhanced as a more complete water layer forms on the particle surfaces. In this case, the Cl 2p photoelectrons only need to pass through part of the water layer while the Na 1s electrons will be attenuated by both a layer of water as well as the dissolved Cl⁻ ions. This interpretation is supported by density functional theory (DFT) calculations and contact potential measurements (Cabrera-Sanfeliix et al., 2007), which showed that Cl⁻ can indeed be lifted out of a NaCl (100) crystal surface already at one monolayer water coverage. Using XPS on free-flying sub-2 nm CsBr water clusters, Hautala et al. (2017b) found that Br⁻ ions were closer to the surface than their counter cations. These experimental

results further support the interpretation of our present observations that pre-deliquescent water adsorption enhances chloride relative to sodium in the aerosol surface layer.

The presence of halide ions, especially Cl^- and Br^- , at the air-water interface has been connected to increased photochemical activity (e.g. George and Abbatt, 2010, and references therein). The mobilization of ions can lead to release of gaseous halogen compounds from sea salt aerosol due to reactions in the aqueous phase (Mozurkewich, 1995; Vogt et al., 1996; Kerminen et al., 1998; Keene et al., 1999). In the atmosphere, formation of solvated halogen ions even at very dry conditions via similar pre-deliquescent adsorption of water onto the surfaces of sea salt aerosol as seen for laboratory generated aerosol in this work could therefore have significant implications for the halogen cycle, including ozone chemistry in the polar regions (Simpson et al., 2007).

Another possible explanation for the difference in the Na to Cl ratio could be beam damage. Cl has been shown to be more prone to the effect of irradiation, and in the course of the experiments this could mean an accumulating loss of Cl due to desorption or diffusion (Verdaguer et al., 2008). Our present results, however, show exactly the opposite: the Cl is enhanced with respect to Na. Furthermore, the radiation from the X-ray anode is far less intense than that produced by a synchrotron, making the effects due to beam damage less likely. This leads us to conclude that the observed effect cannot be due to beam damage. Finally, it is also possible that the photoelectron signal is attenuated by so-called adventitious carbon, a thin layer of carbon-containing material that may form on the samples when exposed to ambient air (Barr and Seal, 1995), in addition to adsorbed water. During our experiments, we unfortunately did not measure C 1s spectra for the NaCl particles and we are therefore not able to quantify the potential influence of adventitious carbon on the Na 1s and Cl 2p peak area ratios. Effects of adventitious carbon is discussed in more detail for the sucrose and malonic acid samples where C 1s spectra were measured.




Figure 4. Relative peak areas for NaCl (Na:Cl, red), sucrose (O:C, black) and the malonic acid carboxyl groups (COOH:COOH, blue). The dashed horizontal red, black and blue lines show the expected stoichiometric ratio for NaCl, sucrose and malonic acid, respectively. Relative ratios differing from the stoichiometric ratio indicate changes in the chemical composition of the surface of the aerosol particles.

3.2 Sucrose

Figure 5 shows the C 1s XPS spectra of deposited sucrose aerosol particles at 0, 0.2, 1 and 5 mbar water vapor pressure. Measurements were repeated at 0 mbar water vapor pressure after dosing but are not shown here since no significant changes were observed. The C 1s spectra were fitted with three peaks corresponding to distinct carbon atoms in C-C/C-H, C-O and O-C-O bonds at 285.90, 287.36 and 288.76 eV, respectively, at 0 % RH. Sucrose molecules contain only C-O and O-C-O bound carbon and the largest peak corresponding to C-C/C-H bound carbon is therefore attributed to adventitious carbon on the substrate. The obtained peak energies at 0 % RH for C-O and O-C-O carbons are in good agreement with values reported by Stevens and Schroeder (2009), 286.7 and 288.1 eV, respectively. Since the gold substrate is highly conductive, we do not expect any charging effect as for the NaCl samples deposited on Si. Differences between the binding energies measured here and those of Stevens and Schroeder (2009) may instead arise from the choice of reference for calibrating the energy scale. Whereas we here calibrated the energy scale using the well-known Au 4f peak, Stevens and Schroeder (2009) used the adventitious carbon peak, which may have a wide range of binding energies (Greczynski and Hultman, 2018). Additional details on the peak fitting can be found in the supplement.

Figure 6 shows the O 1s XPS spectra of the deposited sucrose aerosol particles. The O 1s spectra were fitted with two peaks for sucrose corresponding to C-O and O-C-O bound oxygen. With the introduction of water vapor into the high pressure cell, a third peak from the water vapor (Patel et al., 2019) appears in the XPS spectra, as evidenced by the increase in signal with increasing humidity. The binding energies of the C-O and O-C-O peaks at 0 % RH were determined to be 533.02 eV and 533.72, respectively. These values match closely with those (533.0 eV and 533.7 eV, respectively) reported by Stevens and Schroeder (2009). Additional peak fitting details can be found in the supplement.

The binding energy shifts of sucrose C 1s and O 1s peaks with respect to 0 % RH are shown in the black traces in Fig. 3. Because the peak separation in each case is fixed, only a single shift is reported for spectra of each element. Unlike with NaCl, the binding energy shifts in the sucrose spectra do not suggest adsorption of water to the sucrose particles at the investigated humidities. The C 1s shows initially a small shift and then returns to 0 at 16 % RH. The O 1s peaks on the other hand seem to shift slightly towards lower binding energies (580 meV at 16 % RH). Figure 4 shows the O:C ratio of sucrose as a function of RH. The ratio is in all conditions very close to the stoichiometric value of O:C= 11/12 \approx 0.92. Looking specifically at the relative ratios of C-O and O-C-O bound carbon C 1s and oxygen O 1s peaks, we see only minor changes with RH. The C-O C 1s signal seems to increase slightly from 0 % to 3.2 % RH, but is again lower at 16 % RH. The same is seen for the O 1s O-C-O signal, which increases slightly from dry conditions towards 16 % RH. Taken together, the observations that both the O 1s and C 1s peaks only show a small binding energy shift with increasing RH, the O:C ratio remains nearly constant and the changes in relative ratios between C-O and O-C-O bound carbon C 1s and oxygen O 1s signals are small, suggest that either there is no significant amounts of water adsorbed onto the particles at RH below 16 % or that, if water is adsorbed, there is no significant chemical interaction between the water and the sucrose nanoparticle.

The high surface sensitivity of the XPS spectra enables us to confirm previous measurement results that rely on detecting changes in volume (Zobrist et al., 2011) or mass (Yao et al., 2011). Zobrist et al. (2011) used optical techniques to determine the




Figure 5. C 1s XPS spectra of sucrose aerosol particles. The dots show the recorded data, the solid lines the fit envelope curve and the red, blue and green peaks the C-C/C-H, C-O and O-C-O bound carbon components, respectively. The dashed vertical line shows the binding energy of the C-O component at 0 mbar pressure (0 % RH) at the beginning of the experiments. Error bars show the estimated uncertainty in the peak position from Monte Carlo analysis. Photon energy was 1253.6 eV from the Mg anode.

400 size of single sucrose microparticles levitated in an electrodynamic balance (EDB) under varying RH conditions. Their results show that crystalline sucrose barely adsorbs any water before the deliquescence point of 85.6 %. They also show that sucrose does not exhibit any efflorescence upon drying, but instead assumes an amorphous glassy state. On subsequent humidification cycles the amorphous sucrose particle did not show any significant water uptake before RH of 35-40 %. Our present work also supports this observation on the molecular level. In contrast to NaCl particles, the absence of changes in the sucrose C 1s and O

405 1s binding energies and O to C ratio with increasing RH, together with this high surface sensitivity of the XPS measurements, indicate insignificant water adsorption by sucrose aerosol particles at humidities up to 16 % RH.




Figure 6. O 1s XPS spectra of sucrose aerosol particles. The dots show the recorded data, the solid lines the fit envelope curve and the red, blue and green peaks the water vapor, and sucrose O-C-O and C-O components, respectively. The dashed vertical line shows the binding energy of the C-O component at 0 mbar pressure (0 % RH) at the beginning of the experiments. Error bars show the estimated uncertainty in the peak position from Monte Carlo analysis. Photon energy was 1253.6 eV from the Mg anode.

In the atmosphere, these results could mean that pre-deliquescent water adsorption and resulting chemical changes in very dry conditions could be much less likely to occur for aerosol comprising significant amounts of sugar functionalities, such as those produced in biomass burning (Scaramboni et al., 2015; Bhattarai et al., 2019), compared to sea salt aerosol.

410 3.3 Malonic Acid

Figures 7 and 8 show the recorded C 1s and O 1s XPS spectra, respectively, of deposited malonic acid aerosol particles at 0, 0.2, and 1 mbar water vapor. Due to measurement time constraints, spectra were not recorded for 5 mbar or again at 0 mbar water vapor after dosing. The C 1s spectra were fitted with three main peaks corresponding to C-C/C-H bound carbon originating from adventitious carbon, and COOH and C-C carbon from malonic acid particles at 285.75, 290.02 and 286.02

415 eV, respectively, in UHV conditions. As for sucrose, we also do not expect any charging effects for the malonic acid samples due to the high conductivity of the gold substrate. Malonic acid has been shown to be prone to beam damage when exposed to radiation from laboratory X-ray sources. This can give rise to additional peaks in the XPS spectra arising from malonic acid molecules damaged by the X-ray beam (Ferreira Jr. et al., 2017a, b). The decomposition peaks DP1 and DP2 observed by Ferreira Jr. et al. (2017b) were included in the fit for 0 % RH (UHV), where a good fit could not be produced without their
420 inclusion. In UHV conditions, the DP1 and DP2 peaks were observed at 291.52 eV and 287.22 eV. The spectra recorded with water vapor did not show clear indication of the DP2 feature, but an additional peak needs to be fitted close to the DP1 feature in both 0.6 % and 3.2 % RH spectra. The best fit was acquired with the feature at 287.05 eV and 286.50 eV for 0.6 % and 3.2 % RH, respectively. This is 1.4 eV higher than the C-C carbon peak of malonic acid in both cases, in good agreement with the position of the DP1 peak of Ferreira Jr. et al. (2017b). Additional details of the fitting procedure may be found in the
425 supplement.

The malonic acid O 1s spectra were fitted with two main peaks for the C=O and C-OH bound oxygen of the two carboxyl groups. For UHV conditions, the best fit was acquired with a splitting of approximately 1.0 eV between the C=O and C-OH peaks (at 532.33 and 533.28 eV, respectively). This splitting is smaller than the previously reported values of 1.1 and 1.3 eV by Ferreira Jr. et al. (2017b). For our measurements with water vapor, the best fits were obtained with a splitting of 1.2 eV
430 between the C=O and C-OH oxygen peaks (at 531.84 and 533.04 eV at 3.2 % RH, respectively), now in good agreement with the work of Ferreira Jr. et al. (2017b). The peak at highest binding energy in the spectra recorded at humid conditions is the O 1s peak from water vapor. While peaks due to possible beam damage were observed in the C 1s spectra of malonic acid (DP1 and DP2 in Fig. 7), fitting corresponding peaks in the O 1s spectrum did not yield a better representation of the measured XPS spectra. However, this is also in line with the previous works of Ferreira Jr. et al. (2017a), showing that the signal due to beam
435 damage is much less pronounced in the O 1s spectra than in the C 1s spectra.

The binding energy shifts of malonic acid particle C 1s and O 1s peaks with respect to 0 % RH are shown in the blue traces in Fig. 3. When RH is increased, the COOH (and thus also C-C) peaks of malonic acid are seen to shift towards lower binding energies. While the the shift for malonic acid O 1s is not as dramatic as for NaCl particles, the malonic acid C 1s binding energies shift is even more pronounced than that of Na 1s or Cl 2p, indicating that water is being adsorbed onto the surface of
440 the malonic acid particles and changing the chemical environment of the surface molecules. In addition to the peaks shifts, the O 1s to C 1s signal ratio within the malonic acid carboxyl group, COOH:COOH, changes with increasing RH. We focus on the carboxyl groups as the C-C/C-H contribution is under the strong C-C/C-H signal from adventitious carbon and therefore cannot be fitted reliably. The stoichiometric O:C ratio of a carboxyl group is 2. Interestingly, already at 0 % RH (UHV), the extracted ratio COOH:COOH= 0.94 is far from this stoichiometric value. When introducing water vapor into the system at
445 0.6 % RH, the ratio changes to 2.15. At 3 % RH, it increases even further to 2.95. While we cannot offer a clear explanation for this observation at the present, based on the observed binding energy shifts and the changes in the COOH:COOH ratio, it is safe to conclude that water is adsorbed on nanoscale malonic acid aerosols particles already at the lowest RH measured and that it has an effect on the chemical composition of the particle surface.




Figure 7. C 1s spectra of malonic acid aerosol particles. The dots show the recorded data and the solid lines the fit to the data. The blue and purple peaks correspond to COOH and C-C of malonic acid, respectively while the yellow and green peaks correspond to DP1 and DP2 components, respectively, attributed to beam damage. The red peak corresponds to C-C/C-H of adventitious carbon. The dashed vertical line shows the binding energy of the COOH component at 0 mbar pressure (0 % RH) at the beginning of the experiments. Error bars show the estimated uncertainty in the peak position from Monte Carlo analysis. Photon energy was 1253.6 eV from the Mg anode.

On closer inspection of the malonic acid spectra, it is clear that the C 1s spectrum at 0 % RH (UHV) resembles that of a
 450 beam damaged sample described by Ferreira Jr. et al. (2017a, b), but not entirely: the DP2 feature in our particle spectra is stronger than reported by Ferreira Jr. et al. (2017b) and the DP1 weaker. When increasing the RH in our measurements, the stronger DP2 peak disappears and only the extra feature attributed to DP1 is seen. Our malonic acid particle O 1s spectra do not show any signs of beam damage even at increased RH. Instead, the relative intensities between the C=O and C-OH components change toward the expected C=O:C-OH ratio of 1 from the C=O:C-OH = 1.4 observed at 0 % RH (UHV). This indicates an
 455 increase in the number of C-OH and decrease in the number of C=O groups on the malonic acid particle surface with increasing RH. Indeed, the binding energy of the C 1s (DP2) feature is close to the typical binding energy (approximately 286.5 eV) of a




Figure 8. O 1s spectra of malonic acid aerosol particles. The dots show the recorded data, the solid lines the fit envelope curve and the red, blue and green peaks the water vapor, C-OH and C=O components, respectively. The dashed vertical line shows the binding energy of the C=O component at 0 mbar pressure (0 % RH) at the beginning of the experiments. Error bars show the estimated uncertainty in the peak position from Monte Carlo analysis. Photon energy was 1253.6 eV from the Mg anode.

hydroxyl group carbon, C-OH. If this were purely a result of beam damage, the DP2 feature in the C 1s spectrum should also be visible, not to mention the beam damage component in the O 1s spectra. It could therefore be a sign of significant chemical changes in the malonic acid particles.

460 Malonic acid is known to exhibit keto-enol tautomerism and it has been theoretically predicted that the transition between the two forms can take place through an intramolecular transition in a system of just six water molecules and one malonic acid molecule (see e.g. Yamabe et al. (2004)). Ghorai et al. (2011) performed spectromicroscopic experiments combining scanning transmission X-ray microscopy with near-edge X-ray absorption fine structure spectroscopy (STXM/NEXAFS) and Fourier transform infrared spectroscopy (FTIR) on deliquesced submicron malonic acid aerosol particles at varying RH up to 90 %.

465 Their bulk-sensitive experiments show that while the keto form dominates in dilute aqueous solutions of malonic acid, the

equilibrium shifts toward the enol form in saturated solutions. This observation is also supported by the theoretical calculations of Dick-Pérez and Windus (2017) for concentrated malonic acid solution particles. The results of Ghorai et al. (2011) show that the amount of enol form increases with increasing relative humidity as water is absorbed by deliquesced malonic acid aerosols. Furthermore, they demonstrate water uptake by malonic acid aerosols already at RH as low as 2 %, which is in good
470 agreement with the conclusion of our surface-sensitive XPS experiments in the present work.

The single particle carbon K-edge NEXAFS spectra of Ghorai et al. (2011) show a clear increase in C=C and C-OH resonances, while C-COOH and C-CH are decreased with increasing RH. Their oxygen K-edge spectra also shows a clear decrease in the -COOH resonance. This is similar to the observations of the present work, where the O 1s C-OH signal from malonic acid particle surfaces increases with increasing RH. While a similar increase is also observed here in the C 1s for the C-OH
475 (DP1) component, caution is necessary when attributing this to the enol form of malonic acid. The C-OH peak at approximately 286.5 eV is a sign of carbon with a single hydroxyl group, while the malonic acid enol form would have a carbon with two hydroxyl groups bound to it (HO-C-OH or O-C-O), increasing the binding energy from 286.5 eV to approximately 288 eV (Lannon Jr. and Meng, 1999). Although no such peak has been fit to the data in the present work, it also cannot be ruled out completely. However, if the spectral changes observed in our XPS experiments are due to the formation of the enol tautomer
480 on the malonic acid particle surface, this could have profound implications in terms of atmospheric chemistry as the keto and enol forms have differing physical and chemical stabilities and reaction pathways in both the gas and aqueous phases (Yamabe et al., 2004).

4 Conclusions

The main goal of this work has been to investigate the potential for obtaining meaningful results with the APXPS technique
485 for aerosol samples comprising simple but atmospherically relevant chemical components. We studied water adsorption onto NaCl, sucrose and malonic acid aerosol particles deposited on silicon and gold substrates using the APXPS with a laboratory X-ray source (Al K α and Mg K α). To our knowledge, this is the first time surface sensitive and chemically specific XPS has been used in AP conditions on nano-scale particles with compositions of immediate atmospheric relevance.

The samples were exposed to relative humidities between 0 % and 16 % at room temperature and the XPS spectra of Na
490 1s, Cl 2p, C 1s and O 1s core levels were recorded. The results show that water can be adsorbed onto the aerosol particles already at very low relative humidities. Corresponding changes were observed in the chemical environment, and possibly even composition, of the particle surfaces with increasing relative humidity. For NaCl particles, we find signs of Cl⁻ ions dissolving into the thin surface water layer, as well as a memory effect in binding energies when drying samples back to 0 % RH. This water uptake memory persists even after heating the particles up to 125 °C. Although some water seems to remain on the
495 particles, the Cl appears to return to the crystal lattice of the aerosol as the particle is dried. Sucrose particles show very few changes when exposed to water vapor at the conditions of these experiments. Malonic acid particles on the other hand show a dramatic shift in the C 1s binding energies even when exposed to very small amounts of water (0.2–3 % RH), as well as a shift in the O 1s binding energies. Changes are also seen in the relative amounts of C=O and C-OH bound oxygen on the surface of

malonic acid particles, together with the emergence of C-OH bound carbon at the expense of COOH. These surface specific
500 changes are in line with earlier bulk-sensitive observations of water uptake by malonic acid particles at very low RH (Ghorai
et al., 2011) and could even indicate a change in the keto-enol equilibrium of malonic acid within the surface adsorbed water
layer in our experiments.

The present observations suggest that while sucrose particles do not interact appreciably with water vapor at low RH con-
ditions, the adsorption of small amounts of water onto both NaCl and malonic acid particles could already have implications
505 for their surface chemistry on a molecular level. These differences between chemically simple components emulating sea salt,
biomass burning, and carboxylic acid aerosol could represent major differences in their atmospheric chemistry in response
to water in dry conditions. The presence of water at the aerosol surface could impact water-catalyzed chemical reactions, as
well as reactions involving mobilization of charged species such as organic and inorganic ions or components with acid-base
functionalities. For example, Cl^- ions in sea salt aerosols are a source of chlorine in the troposphere through heterogeneous
510 processes (see e.g. Wang et al. (2019) and references therein). With sea salt being one of the most abundant inorganic aerosol
components in the global atmosphere, our present observations could imply significant reactive uptake and processing of at-
mospheric trace gases even at low RH. Dicarboxylic acids like malonic acid are also ubiquitous in atmospheric aerosols in
the troposphere. The formation of concentrated solutions with the enol form present even in relatively dry conditions may
have important implications for the reactivity of these aerosol surfaces. The role of particle surfaces is enhanced for smaller
515 particles with large surface-to-bulk ratios (Prisle et al., 2010b; Bzdek et al., 2020). The effects of water uptake detected at very
low relative humidities in this work could not have been detected with standard mass or volume based aerosol techniques, nor
attributed specifically to the surface with bulk sensitive spectroscopic methods.

Our results demonstrate the viability of the APXPS technique for *in situ* studies of vapor interaction with aerosol particle
surfaces. XPS can provide surface-specific, molecular-level chemical information on uptake and reactions through changes in
520 the presence, position or intensity of spectral peaks, and can detect chemical changes on aerosol surfaces which likely cannot be
resolved with even high-resolution bulk methods such as aerosol mass spectrometry. We have shown that APXPS can be used
on sampled aerosol particles comprising both organic and inorganic components of high atmospheric relevance. Our samples
were generated with standard aerosol laboratory methods and exposed to a well-defined vapor phase during experiments. The
present experiments were carried out using a conventional laboratory X-ray source which limited the ambient pressure at which
525 the measurements could be done. While the experiments were conducted at atmospherically relevant water vapor pressures, they
were still far below the deliquescent relative humidities of the compounds studied and we cannot immediately conclude that
the interaction between water vapor and the nanoparticles studied here will remain the same for all relative humidities below
deliquescence. Combining APXPS with high brilliance synchrotron radiation now available at modern synchrotron facilities
therefore shows great promise for studying aerosol surfaces at even higher ambient pressure conditions up to deliquescence, in
530 more complex and reactive environments, as well as with high time resolution.

Author contributions. Conceptualization: NLP; Funding acquisition: NLP; Investigation: JLL, KRR, SW, EK, M-HM, and SU; Analysis: SU, JLL, SW, and NLP; Visualization: SU and JLL; Writing of first version: JLL, SU, and NLP with input from all co-authors. Writing of revised version: JLL and NLP with input from all co-authors. Authors' response: JLL, NLP, and SU with input from all co-authors; Supervision: NLP; Project administration: NLP.

535 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme, Project SURFACE (Grant Agreement No. 717022). The authors also gratefully acknowledge the financial contribution from the Academy of Finland, including Grant Nos. 308238, 314175, 335649, 290145, 326291, and 331532. We acknowledge MAX IV Laboratory for time on Beamline SPECIES. Research conducted at MAX IV, a Swedish national user facility, is supported by the Swedish Research council under contract 2018-07152, the Swedish Governmental Agency for Innovation Systems under contract 2018-04969, and Formas under contract 2019-02496. We warmly thank Jenny Rissler and Birgitta Svenningsson for assistance with preparation of samples and estimation of surface coverage.

References

- 545 Ammann, M., Artiglia, L., and Bartels-Rausch, T.: X-Ray Excited Electron Spectroscopy to Study Gas–Liquid Interfaces of Atmospheric Relevance, in: *Physical Chemistry of Gas-Liquid Interfaces*, pp. 135–166, Elsevier, 2018.
- Antonsson, E., Patanen, M., Nicolas, C., Neville, J. J., Benkoula, S., Goel, A., and Miron, C.: Complete Bromide Surface Segregation in Mixed NaCl/NaBr Aerosols Grown from Droplets, *Physical Review X*, 5, 011 025–7, 2015.
- Archer, J., Walker, J., Gregson, F. K. A., Hardy, D. A., and Reid, J. P.: Drying Kinetics and Particle Formation from Dilute Colloidal Suspensions in Aerosol Droplets, *Langmuir*, pp. acs.langmuir.0c01 830–49, 2020.
- 550 Ault, A. P. and Axson, J. L.: Atmospheric Aerosol Chemistry: Spectroscopic and Microscopic Advances, *Analytical Chemistry*, 89, 430–452, 2016.
- Barr, T. L. and Seal, S.: Nature of the use of adventitious carbon as a binding energy standard, *Journal of Vacuum Science & Technology A: Vacuum, Surfaces, and Films*, 13, 1239–1246, 1995.
- Beard, B. C.: Fresh Cleaved Single Crystal NaCl, XPS spectra, Mg Source, *Surface Science Spectra*, 2, 128–132, 1993.
- 555 Benjamin, I.: Chemical Reaction Dynamics at Liquid Interfaces: A Computational Approach, *Progress in Reaction Kinetics and Mechanism*, 27, 87–126, 2019.
- Bhattacharai, H., Saikawa, E., Wan, X., Zhu, H., Ram, K., Gao, S., Kang, S., Zhang, Q., Zhang, Y., Wu, G., Wang, X., Kawamura, K., Fu, P., and Cong, Z.: Levoglucosan as a tracer of biomass burning: Recent progress and perspectives, *Atmospheric Research*, 220, 20–33, 2019.
- Bilde, M. and Svenningsson, B.: CCN activation of slightly soluble organics: the importance of small amounts of inorganic salt and particle phase, *Tellus B*, 56, 128–134, 2004.
- 560 Braun, C. and Krieger, U. K.: Two-dimensional angular light-scattering in aqueous NaCl single aerosol particles during deliquescence and efflorescence, *Optics Express*, 8, 314–8, 2001.
- Braun, R. A., Dadashazar, H., MacDonald, A. B., Aldhaif, A. M., Maudlin, L. C., Crosbie, E., Aghdam, M. A., Hossein Mardi, A., and Sorooshian, A.: Impact of Wildfire Emissions on Chloride and Bromide Depletion in Marine Aerosol Particles, *Environmental Science & Technology*, 51, 9013–9021, 2017.
- 565 Bzdek, B. R., Reid, J. P., Malila, J., and Prisle, N. L.: The surface tension of surfactant-containing, finite volume droplets, *Proceedings of the National Academy of Sciences*, 117, 8335–8343, 2020.
- Cabrera-Sanfeliix, P., Sanchez Portal, D., Verdaguer, A., Darling, G. R., Salmeron, M., and Arnau, A.: Spontaneous Emergence of Cl⁻ Anions from NaCl(100) at Low Relative Humidity, *The Journal of Physical Chemistry C*, 111, 8000–8004, 2007.
- 570 Cheng, R. J., Blanchard, D. C., and Cipriano, R. J.: The formation of hollow sea-salt particles from the evaporation of drops of seawater, *Atmospheric Research*, 22, 15–25, 1988.
- Cheng, W., Weng, L.-T., Li, Y., Lau, A., Chan, C. K., and Chan, C.-M.: Surface Chemical Composition of Size-Fractionated Urban Walkway Aerosols Determined by X-Ray Photoelectron Spectroscopy, *Aerosol Science and Technology*, 47, 1118–1124, 2013.
- Cosman, L. M., Knopf, D. A., and Bertram, A. K.: N₂O₅ Reactive Uptake on Aqueous Sulfuric Acid Solutions Coated with Branched and Straight-Chain Insoluble Organic Surfactants, *The Journal of Physical Chemistry A*, 112, 2386–2396, 2008.
- 575 Cwiertny, D. M., Young, M. A., and Grassian, V. H.: Chemistry and Photochemistry of Mineral Dust Aerosol, *Annual Review of Physical Chemistry*, 59, 27–51, 2008.
- Dai, Q., Hu, J., and Salmeron, M.: Adsorption of Water on NaCl (100) Surfaces: Role of Atomic Steps, *The Journal of Physical Chemistry B*, 101, 1994–1998, 1997.

- 580 Darr, J. P., Davis, S. Q., Kohno, Y., McKenna, K., and Morales, P.: Morphological effects on the hygroscopic properties of sodium chloride–sodium sulfate aerosols, *Journal of Aerosol Science*, 77, 158–167, 2014.
- Decesari, S., Facchini, M. C., Fuzzi, S., and Tagliavini, E.: Characterization of water-soluble organic compounds in atmospheric aerosol: A new approach, *Journal of Geophysical Research*, 105, 1481–1489, 2000.
- Decesari, S., Facchini, M. C., Matta, E., Lettini, F., Mircea, M., Fuzzi, S., Tagliavini, E., and Putaud, J. P.: Chemical features and seasonal
585 variation of fine aerosol water-soluble organic compounds in the Po Valley, Italy, *Atmospheric Environment*, 35, 3691–3699, 2001.
- Dick-Pérez, M. and Windus, T. L.: Computational Study of the Malonic Acid Tautomerization Products in Highly Concentrated Particles, *The Journal of Physical Chemistry A*, 121, 2259–2264, 2017.
- Djikaev, Y. S. and Ruckenstein, E.: Formation and evolution of aqueous organic aerosols via concurrent condensation and chemical aging, *Advances in Colloid and Interface Science*, 265, 45–67, 2019.
- 590 Donaldson, D. J. and Valsaraj, K. T.: Adsorption and Reaction of Trace Gas-Phase Organic Compounds on Atmospheric Water Film Surfaces: A Critical Review, *Environmental Science & Technology*, 44, 865–873, 2010.
- Emfietzoglou, D. and Moscovitch, M.: Inelastic collision characteristics of electrons in liquid water, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 193, 71–78, 2002.
- Enami, S., Hoffmann, M. R., and Colussi, A. J.: Proton Availability at the Air/Water Interface, *The Journal of Physical Chemistry Letters*, 1,
595 1599–1604, 2010.
- Estillore, A. D., Morris, H. S., Or, V. W., Lee, H. D., Alves, M. R., Marciano, M. A., Laskina, O., Qin, Z., Tivanski, A. V., and Grassian, V. H.: Linking hygroscopicity and the surface microstructure of model inorganic salts, simple and complex carbohydrates, and authentic sea spray aerosol particles, *Physical Chemistry Chemical Physics*, 19, 21 101–21 111, 2017.
- Fan, H., Wenyika Masaya, T., and Goulay, F.: Effect of surface–bulk partitioning on the heterogeneous oxidation of aqueous saccharide
600 aerosols, *Physical Chemistry Chemical Physics*, 21, 2992–3001, 2019.
- Ferreira Jr., J. M., Trindade, G. F., Tshulu, R., Watts, J. F., and Baker, M. A.: Introduction to a series of dicarboxylic acids analyzed by x-ray photoelectron spectroscopy, *Surface Science Spectra*, 24, 011 001–5, 2017a.
- Ferreira Jr., J. M., Trindade, G. F., Tshulu, R., Watts, J. F., and Baker, M. A.: Dicarboxylic acids analysed by x-ray photoelectron spectroscopy, Part I - propanedioic acid anhydrous, *Surface Science Spectra*, 24, 011 101–8, 2017b.
- 605 Foster, M. C. and Ewing, G. E.: Adsorption of water on the NaCl(001) surface. II. An infrared study at ambient temperatures, *The Journal of Chemical Physics*, 112, 6817–6826, 2000.
- Gen, M. and Chan, C. K.: Electro spray surface-enhanced Raman spectroscopy (ES-SERS) for probing surface chemical compositions of atmospherically relevant particles, *Atmospheric Chemistry and Physics*, 17, 14 025–14 037, 2017.
- Gen, M., Kuniyama, R., Matsuki, A., and Chan, C. K.: Electro spray surface-enhanced Raman spectroscopy (ES-SERS) for studying organic
610 coatings of atmospheric aerosol particles, *Aerosol Science and Technology*, 53, 760–770, 2019.
- George, I. J. and Abbatt, J. P. D.: Heterogeneous oxidation of atmospheric aerosol particles by gas-phase radicals, *Nature Publishing Group*, 2, 713–722, 2010.
- Ghorai, S., Laskin, A., and Tivanski, A. V.: Spectroscopic Evidence of Keto–Enol Tautomerism in Deliquesced Malonic Acid Particles, *The Journal of Physical Chemistry A*, 115, 4373–4380, 2011.
- 615 Ghosal, S. and Hemminger, J. C.: Surface Adsorbed Water on NaCl and Its Effect on Nitric Acid Reactivity with NaCl Powders, *The Journal of Physical Chemistry B*, 108, 14 102–14 108, 2004.

- Greczynski, G. and Hultman, L.: Reliable determination of chemical state in x-ray photoelectron spectroscopy based on sample-work-function referencing to adventitious carbon: Resolving the myth of apparent constant binding energy of the C 1s peak, *Applied Surface Science*, 451, 99–103, 2018.
- 620 Hautala, L., Jänkälä, K., Löytynoja, T., Mikkilä, M. H., Prisle, N., Tchapyguine, M., and Huttula, M.: Experimental observation of structural phase transition in CsBr clusters, *Physical Review B*, 95, 045 402, 2017a.
- Hautala, L., Jänkälä, K., Mikkilä, M.-H., Turunen, P., Prisle, N. L., Patanen, M., Tchapyguine, M., and Huttula, M.: Probing RbBr solvation in freestanding sub-2 nm water clusters, *Physical Chemistry Chemical Physics*, 19, 25 158–25 167, 2017b.
- Hori, M., Ohta, S., Murao, N., and Yamagata, S.: Activation capability of water soluble organic substances as CCN, *Journal of Aerosol*
625 *Science*, 34, 419–448, 2003.
- Huang, D., Wang, J., Xia, H., Zhang, Y., Bao, F., Li, M., Chen, C., and Zhao, J.: Enhanced Photochemical Volatile Organic Compounds Release from Fatty Acids by Surface-Enriched Fe(III), *Environmental Science & Technology*, p. acs.est.0c03793, 2020.
- Huang, Y., Barraza, K. M., of, C. K. T. J., and 2018: Probing the OH oxidation of pinonic acid at the air–water interface using Field-Induced Droplet Ionization Mass Spectrometry (FIDI-MS), *ACS Publications*, 122, 6445–6456, 2018.
- 630 Jacobs, M. I., Davies, J. F., Lee, L., Davis, R. D., Houle, F., and Wilson, K. R.: Exploring Chemistry in Microcompartments Using Guided Droplet Collisions in a Branched Quadrupole Trap Coupled to a Single Droplet, *Paper Spray Mass Spectrometer, Analytical Chemistry*, 89, 12 511–12 519, 2017.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis,
635 K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmospheric Chemistry and Physics*, 5, 1053–1123, 2005.
- Kaya, S., Kendelewicz, T., Porsgaard, S., Salmeron, M. B., Brown Jr., G. E., and Nilsson, A.: Autocatalytic Surface Hydroxylation of MgO(100) Terrace Sites Observed under Ambient Conditions, *The Journal of Physical Chemistry C*, 115, 12 864–12 872, 2011.
- Keene, W. C., Khalil, M. A. K., Erickson III, D. J., McCulloch, A., Graedel, T. E., Lobert, J. M., Aucott, M. L., Gong, S. L., Harper, D. B.,
640 Kleiman, G., Midgley, P., Moore, R. M., Seuzaret, C., Sturges, W. T., Benkovitz, C. M., Koropalov, V., Barrie, L. A., and Li, Y. F.: Composite global emissions of reactive chlorine from anthropogenic and natural sources: Reactive Chlorine Emissions Inventory, *Journal of Geophysical Research*, 104, 8429–8440, 1999.
- Kerminen, V.-M., Teinilä, K., Hillamo, R., and Pakkanen, T.: Substitution of chloride in sea-salt particles by inorganic and organic anions, *Journal of Aerosol Science*, 29, 929–942, 1998.
- 645 Khwaja, H. A.: Atmospheric concentrations of carboxylic acids and related compounds at a semiurban site, *Atmospheric Environment*, 29, 127–139, 1995.
- Knopf, D. A. and Forrester, S. M.: Freezing of Water and Aqueous NaCl Droplets Coated by Organic Monolayers as a Function of Surfactant Properties and Water Activity, *The Journal of Physical Chemistry A*, 115, 5579–5591, 2011.
- Knudsen, J., Andersen, J. N., and Schnadt, J.: A versatile instrument for ambient pressure x-ray photoelectron spectroscopy: The Lund cell
650 approach, *Surface Science*, 646, 160–169, 2016.
- Kong, X., Castarède, D., Boucly, A., Artiglia, L., Ammann, M., Bartels-Rausch, T., Thomson, E. S., and Pettersson, J. B. C.: Reversibly Physisorbed and Chemisorbed Water on Carboxylic Salt Surfaces Under Atmospheric Conditions, *The Journal of Physical Chemistry C*, 124, 5263–5269, 2020.

- Kreidenweis, S. M. and Asa-Awuku, A.: Aerosol Hygroscopicity: Particle Water Content and Its Role in Atmospheric Processes, in: Treatise on Geochemistry, edited by Holland, H. D. and Turekian, K. K., pp. 331–361, Elsevier, Oxford, 2014.
- 655 Kukk, E., Snell, G., Bozek, J. D., Cheng, W.-T., and Berrah, N.: Vibrational structure and partial rates of resonant Auger decay of the $N\ 1s \rightarrow 2\pi$ core excitations in nitric oxide, *Physical Review A*, 63, 062 702, 2001.
- Kukk, E., Ueda, K., Hergenhanh, U., Liu, X. J., Prümper, G., Yoshida, H., Tamenori, Y., Makochekanwa, C., Tanaka, T., Kitajima, M., and Tanaka, H.: Violation of the Franck-Condon Principle due to Recoil Effects in High Energy Molecular Core-Level Photoionization, 660 *Physical Review Letters*, 95, 133 001–4, 2005.
- Kwamena, N. O. A., Buajareng, J., and Reid, J. P.: Equilibrium Morphology of Mixed Organic/Inorganic/Aqueous Aerosol Droplets: Investigating the Effect of Relative Humidity and Surfactants, *The Journal of Physical Chemistry A*, 114, 5787–5795, 2010.
- Lampimäki, M., Schreiber, S., Zelenay, V., Křepelová, A., Birrer, M., Axnanda, S., Mao, B., Liu, Z., Bluhm, H., and Ammann, M.: Exploring the Environmental Photochemistry on the $TiO\ 2(110)$ Surface in Situ by Near Ambient Pressure X-ray Photoelectron Spectroscopy, 665 *The Journal of Physical Chemistry C*, 119, 7076–7085, 2015.
- Lannon Jr., J. M. and Meng, Q.: Analysis of a Poly(oxymethylene) Polymer by XPS, *Surface Science Spectra*, 6, 99–102, 1999.
- Laskin, A., Moffet, R. C., Gilles, M. K., Fast, J. D., Zaveri, R. A., Wang, B., Nigge, P., and Shutthanandan, J.: Tropospheric chemistry of internally mixed sea salt and organic particles: Surprising reactivity of NaCl with weak organic acids, *Journal of Geophysical Research*, 117, n/a–n/a, 2012.
- 670 Lee, J. K., Samanta, D., Nam, H. G., and Zare, R. N.: Micrometer-Sized Water Droplets Induce Spontaneous Reduction, *Journal of the American Chemical Society*, 141, 10 585–10 589, 2019.
- Ly, G. and Sun, X.: The role of air-water interface in the SO_3 hydration reaction, *Atmospheric Environment*, 230, 117 514, 2020.
- Mallick, S. and Kumar, P.: $OH\bullet + HCl$ Reaction at the Surface of a Water Droplet: An Ab Initio Molecular Dynamical Study, *The Journal of Physical Chemistry B*, 124, 2465–2472, 2020.
- 675 Marsh, B. M., Iyer, K., and Cooks, R. G.: Reaction Acceleration in Electrospray Droplets: Size, Distance, and Surfactant Effects, *Journal of The American Society for Mass Spectrometry*, 30, 1–9, 2019.
- McNeill, V. F.: Aqueous Organic Chemistry in the Atmosphere: Sources and Chemical Processing of Organic Aerosols, *Environmental Science & Technology*, 49, 1237–1244, 2015.
- Miñambres, L., Méndez, E., Sánchez, M. N., Castaño, F., and Basterretxea, F. J.: The effect of low solubility organic acids on the hygroscopicity of sodium halide aerosols, *Atmospheric Chemistry and Physics*, 14, 11 409–11 425, 2014.
- 680 Mozurkewich, M.: Mechanisms for the release of halogens from sea-salt particles by free radical reactions, *Journal of Geophysical Research*, 100, 14 199–9, 1995.
- Muñoz, A., Oller, J. C., Blanco, F., Gorfinkiel, J. D., Limão-Vieira, P., and García, G.: Electron-scattering cross sections and stopping powers in H_2O , *Physical Review A*, 76, 707–7, 2007.
- 685 Murphy, D. M., DiGangi, J. P., Diskin, G., Dollner, M., Kupc, A., Weinzierl, B., and Yu, P.: The distribution of sea-salt aerosol in the global troposphere, *Atmospheric Chemistry and Physics*, 19, 4093–4104, 2019.
- Nguyen, T. K. V., Zhang, Q., Jimenez, J. L., Pike, M., and Carlton, A. G.: Liquid Water: Ubiquitous Contributor to Aerosol Mass, *Environmental Science & Technology Letters*, p. acs.estlett.6b00167, 2016.
- Ogletree, D. F., Bluhm, H., Hebenstreit, E. L. D., and Salmeron, M.: Photoelectron spectroscopy under ambient pressure and temperature 690 conditions, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 601, 151–160, 2009.

- Öhrwall, G., Prisle, N. L., Ottosson, N., Werner, J., Ekholm, V., Walz, M.-M., and Björneholm, O.: Acid–Base Speciation of Carboxylate Ions in the Surface Region of Aqueous Solutions in the Presence of Ammonium and Aminium Ions, *The Journal of Physical Chemistry B*, 119, 4033–4040, 2015.
- 695 Ouf, F. X., Parent, P., Laffon, C., Marhaba, I., Ferry, D., Marcillaud, B., Antonsson, E., Benkoula, S., Liu, X. J., Nicolas, C., Robert, E., Patanen, M., Barreda, F. A., Sublemontier, O., Coppalle, A., Yon, J., Miserque, F., Mostefaoui, T., Regier, T. Z., Mitchell, J. B. A., and Miron, C.: First in-flight synchrotron X-ray absorption and photoemission study of carbon soot nanoparticles, *Nature Publishing Group*, 6, 1–12, 2016.
- 700 Ovadnevaite, J., Zuend, A., Laaksonen, A., Sanchez, K. J., Roberts, G., Ceburnis, D., Decesari, S., Rinaldi, M., Hodas, N., Facchini, M. C., Seinfeld, J. H., and O’ Dowd, C.: Surface tension prevails over solute effect in organic-influenced cloud droplet activation, *Nature*, 546, 637–641, 2017.
- Pak, C. Y., Li, W., and Steve Tse, Y.-L.: Free Energy and Dynamics of Organic-Coated Water Droplet Coalescence, *The Journal of Physical Chemistry C*, 124, 8749–8757, 2020.
- 705 Park, S.-C., Burden, D. K., and Nathanson, G. M.: Surfactant Control of Gas Transport and Reactions at the Surface of Sulfuric Acid, *Accounts of Chemical Research*, 42, 379–387, 2009.
- Parsons, M. T., Mak, J., Lipetz, S. R., and Bertram, A. K.: Deliquescence of malonic, succinic, glutaric, and adipic acid particles, *Journal of Geophysical Research*, 109, n/a–n/a, 2004.
- Patel, D. I., Shah, D., Bahr, S., Dietrich, P., Meyer, M., Thißen, A., and Linford, M. R.: Water vapor, by near-ambient pressure XPS, *Surface Science Spectra*, 26, 014 026–6, 2019.
- 710 Perkins, R. J., Vazquez de Vasquez, M. G., Beasley, E. E., Hill, T. C. J., Stone, E. A., Allen, H. C., and DeMott, P. J.: Relating Structure and Ice Nucleation of Mixed Surfactant Systems Relevant to Sea Spray Aerosol, *The Journal of Physical Chemistry A*, p. acs.jpca.0c05849, 2020.
- Peters, S. J. and Ewing, G. E.: Water on Salt: An Infrared Study of Adsorbed H₂O on NaCl(100) under Ambient Conditions, *The Journal of Physical Chemistry B*, 101, 10 880–10 886, 1997.
- 715 Peters, S. J., Langmuir, G. E., and 1997: Thin film water on NaCl (100) under ambient conditions: An infrared study, *ACS Publications*, 13, 6345–6348, 1997.
- Petters, S. S., Hilditch, T. G., Tomaz, S., Miles, R. E. H., Reid, J. P., and Turpin, B. J.: Volatility Change during Droplet Evaporation of Pyruvic Acid, *ACS Earth and Space Chemistry*, 4, 741–749, 2020.
- Pope, F. D., Dennis-Smith, B. J., Griffiths, P. T., Clegg, S. L., and Cox, R. A.: Studies of Single Aerosol Particles Containing Malonic Acid, Glutaric Acid, and Their Mixtures with Sodium Chloride. I. Hygroscopic Growth, *The Journal of Physical Chemistry A*, 114, 5335–5341, 720 2010.
- Preger, C., Overgaard, N. C., Messing, M. E., and Magnusson, M. H.: Predicting the deposition spot radius and the nanoparticle concentration distribution in an electrostatic precipitator, *Aerosol Science and Technology*, 1, 1–11, 2020.
- Prenni, A. J., DeMott, P. J., Kreidenweis, S. M., Sherman, D. E., Russell, L. M., and Ming, Y.: The Effects of Low Molecular Weight 725 Dicarboxylic Acids on Cloud Formation, *The Journal of Physical Chemistry A*, 105, 11 240–11 248, 2001.
- Prisle, N. L., Engelhart, G. J., Bilde, M., and Donahue, N. M.: Humidity influence on gas-particle phase partitioning of α -pinene + O₃ secondary organic aerosol, *Geophysical Research Letters*, 37, L01 802, 2010a.
- Prisle, N. L., Raatikainen, T., Laaksonen, A., and Bilde, M.: Surfactants in cloud droplet activation: mixed organic-inorganic particles, *Atmospheric Chemistry and Physics*, 10, 5663–5683, 2010b.

- 730 Prisle, N. L., Ottosson, N., Öhrwall, G., Söderström, J., Dal Maso, M., and Björneholm, O.: Surface/bulk partitioning and acid/base speciation of aqueous decanoate: direct observations and atmospheric implications, *Atmospheric Chemistry and Physics*, 12, 12227–12242, 2012.
- Rissman, T. A., Varutbangkul, V., Surratt, J. D., Topping, D. O., McFiggans, G., Flagan, R. C., and Seinfeld, J. H.: Cloud condensation nucleus (CCN) behavior of organic aerosol particles generated by atomization of water and methanol solutions, *Atmospheric Chemistry and Physics*, 7, 2949–2971, 2007.
- 735 Rood, M. J., Shaw, M. A., Larson, T. V., and Covert, D. S.: Ubiquitous nature of ambient metastable aerosol, *Nature*, 337, 537–539, 1989.
- Rossi, M. J.: Heterogeneous Reactions on Salts, *Chemical Reviews*, 103, 4823–4882, 2003.
- Roy, S., Diveky, M. E., and Signorell, R.: Mass Accommodation Coefficients of Water on Organics from Complementary Photoacoustic and Light Scattering Measurements on Laser-Trapped Droplets, *The Journal of Physical Chemistry C*, 124, 2481–2489, 2020.
- Salmeron, M. and Schlögl, R.: Ambient pressure photoelectron spectroscopy: A new tool for surface science and nanotechnology, *Surface*
- 740 *Science Reports*, 63, 169–199, 2008.
- Sareen, N., Schwier, A. N., Lathem, T. L., Nenes, A., and McNeill, V. F.: Surfactants from the gas phase may promote cloud droplet formation, *Proceedings of the National Academy of Sciences*, 110, 2723–2728, 2013.
- Scaramboni, C., Urban, R. C., Lima-Souza, M., Nogueira, R. F. P., Cardoso, A. A., Allen, A. G., and Campos, M. L. A. M.: Total sugars in atmospheric aerosols: An alternative tracer for biomass burning, *Atmospheric Environment*, 100, 185–192, 2015.
- 745 Schnadt, J., Knudsen, J., Andersen, J. N., Siegbahn, H., Pietzsch, A., Hennies, F., Johansson, N., Mårtensson, N., Öhrwall, G., Bahr, S., Mahl, S., and Schaff, O.: The new ambient-pressure X-ray photoelectron spectroscopy instrument at MAX-lab, *J. Synchrotron Rad* (2012). 19, 701–704 [doi:10.1107/S0909049512032700], 19, 1–4, 2012.
- Shakya, K. M., Liu, S., Takahama, S., Russell, L. M., Keutsch, F. N., Galloway, M. M., Shilling, J. E., Hiranuma, N., Song, C., Kim, H., Paulson, S. E., Pfaffenberger, L., Barmet, P., Slowik, J., Prévôt, A. S. H., Dommen, J., and Baltensperger, U.: Similarities in STXM-NEXAFS
- 750 *Spectra of Atmospheric Particles and Secondary Organic Aerosol Generated from Glyoxal, α -Pinene, Isoprene, 1,2,4-Trimethylbenzene, and d-Limonene*, 2016.
- Shi, Q., Zhang, W., Ji, Y., Wang, J., Qin, D., Chen, J., Gao, Y., Li, G., and An, T.: Enhanced uptake of glyoxal at the acidic nanoparticle interface: implications for secondary organic aerosol formation, *Environmental Science: Nano*, 7, 1126–1135, 2020.
- Shulman, M. L., Jacobson, M. C., Charlson, R. J., Synovec, R. E., and Young, T. E.: Dissolution behavior and surface tension effects of
- 755 *organic compounds in nucleating cloud droplets*, *Geophysical Research Letters*, 23, 277–280, 1996.
- Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., ARIYA, P., Bottenheim, J., Burrows, J., Carpenter, L. J., Friß, U., Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H. W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A., Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in polar boundary-layer ozone depletion, *Atmospheric Chemistry and Physics*, 7, 4375–4418, 2007.
- 760 SPECS Surface Nano Analysis GmbH: Calculated Transmission for PHOIBOS NAP 150 R2, https://www.specs-group.com/fileadmin/user_upload/products/technical-note/TNote-PHOIBOS_150_NAP_Calculated_Transmission_function.pdf, accessed: 2020-04-18.
- Starr, D. E., Liu, Z., Hävecker, M., Knop-Gericke, A., and Bluhm, H.: Investigation of solid/vapor interfaces using ambient pressure X-ray photoelectron spectroscopy, *Chemical Society Reviews*, 42, 5833–25, 2013.
- Stevens, J. S. and Schroeder, S. L. M.: Quantitative analysis of saccharides by X-ray photoelectron spectroscopy, *Surface and Interface*
- 765 *Analysis*, 41, 453–462, 2009.
- Tang, I. N. and Munkelwitz, H. R.: Composition and temperature dependence of the deliquescence properties of hygroscopic aerosols, *Atmospheric Environment. Part A. General Topics*, 27, 467–473, 1993.

- Tang, M., Chan, C. K., Li, Y. J., Su, H., Ma, Q., Ge, M., Hu, M., and Wang, X.: A review of experimental techniques for aerosol hygroscopicity studies, *Atmospheric Chemistry and Physics*, 19, 12 631–12 686, 2019.
- 770 Toribio, A. R., Prisle, N. L., and Wexler, A. S.: Statistical Mechanics of Multilayer Sorption: Surface Concentration Modeling and XPS Measurement, *The Journal of Physical Chemistry Letters*, 9, 1461–1464, 2018.
- Urpelainen, S., Sâthe, C., Grizzolli, W., Agâker, M., Head, A. R., Andersson, M., Huang, S.-W., Jensen, B. N., Wallén, E., Tarawneh, H., Sankari, R., Nyholm, R., Lindberg, M., Sjöblom, P., Johansson, N., Reinecke, B. N., Arman, M. A., Merte, L. R., Knudsen, J., Schnadt, J., Andersen, J. N., and Hennies, F.: The SPECIES beamline at the MAX IV Laboratory: a facility for soft X-ray RIXS and APXPS, *Journal of Synchrotron Radiation*, 24, 344–353, 2017.
- 775 Verdaguer, A., Sacha, G. M., Luna, M., Ogletree, D. F., and Salmeron, M.: Initial stages of water adsorption on NaCl (100) studied by scanning polarization force microscopy, *The Journal of Chemical Physics*, 123, 124 703, 2005.
- Verdaguer, A., Segura, J. J., Fraxedas, J., Bluhm, H., and Salmeron, M.: Correlation between Charge State of Insulating NaCl Surfaces and Ionic Mobility Induced by Water Adsorption: A Combined Ambient Pressure X-ray Photoelectron Spectroscopy and Scanning Force
- 780 Microscopy Study, *The Journal of Physical Chemistry C*, 112, 16 898–16 901, 2008.
- Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, *Nature*, 383, 327–330, 1996.
- Walz, M. M., Coleman, C., Werner, J., Ekholm, V., Lundberg, D., Prisle, N. L., Öhrwall, G., and Björneholm, O.: Surface behavior of amphiphiles in aqueous solution: a comparison between different pentanol isomers, *Physical Chemistry Chemical Physics*, 17, 14 036–
- 785 14 044, 2015.
- Walz, M. M., Werner, J., Ekholm, V., Prisle, N. L., Öhrwall, G., and Björneholm, O.: Alcohols at the aqueous surface: chain length and isomer effects, *Physical Chemistry Chemical Physics*, 18, 6648–6656, 2016.
- Wang, X., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Zhu, L., Chen, Q., Alexander, B., Sherwen, T., Evans, M. J., Lee, B. H., Haskins, J. D., Lopez-Hilfiker, F. D., Thornton, J. A., Huey, G. L., and Liao, H.: The role of chlorine in global tropospheric chemistry, *Atmospheric*
- 790 *Chemistry and Physics*, 19, 3981–4003, 2019.
- Wang, Z., King, S. M., Freney, E., Rosenoern, T., Smith, M. L., Chen, Q., Kuwata, M., Lewis, E. R., Pöschl, U., Wang, W., Buseck, P. R., and Martin, S. T.: The Dynamic Shape Factor of Sodium Chloride Nanoparticles as Regulated by Drying Rate, *Aerosol Science and Technology*, 44, 939–953, 2010.
- Weis, D. D. and Ewing, G. E.: Water content and morphology of sodium chloride aerosol particles, *Journal of Geophysical Research: Atmospheres*, 104, 21 275–21 285, 1999.
- 795 Werner, F., Ditas, F., Siebert, H., Simmel, M., Wehner, B., Pilewskie, P., Schmeissner, T., Shaw, R. A., Hartmann, S., Wex, H., Roberts, G. C., and Wendisch, M.: Twomey effect observed from collocated microphysical and remote sensing measurements over shallow cumulus, *Journal of Geophysical Research: Atmospheres*, 119, 1534–1545, 2014.
- Werner, J., Persson, I., Björneholm, O., Kaweckî, D., Saak, C.-M., Walz, M.-M., Ekholm, V., Unger, I., Valtl, C., Coleman, C., Öhrwall, G.,
- 800 and Prisle, N. L.: Shifted equilibria of organic acids and bases in the aqueous surface region, *Physical Chemistry Chemical Physics*, 20, 23 281–23 293, 2018.
- Winter, B.: Liquid microjet for photoelectron spectroscopy, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 601, 139–150, 2009.
- Wise, M. E., Martin, S. T., Russell, L. M., and Buseck, P. R.: Water Uptake by NaCl Particles Prior to Deliquescence and the Phase Rule,
- 805 *Aerosol Science and Technology*, 42, 281–294, 2008.

- Yamabe, S., Tsuchida, N., and Miyajima, K.: Reaction Paths of Keto–Enol Tautomerization of β -Diketones, *The Journal of Physical Chemistry A*, 108, 2750–2757, 2004.
- Yao, W., Yu, X., Lee, J. W., Yuan, X., and Schmidt, S. J.: Measuring the Deliquescence Point of Crystalline Sucrose as a Function of Temperature Using a New Automatic Isotherm Generator, *International Journal of Food Properties*, 14, 882–893, 2011.
- 810 Yeh, J. J. and Lindau, I.: Atomic subshell photoionization cross sections and asymmetry parameters: $1 \leq Z \leq 103$, *Atomic Data and Nuclear Data Tables*, 32, 1–155, 1985.
- Yu, L. E., Shulman, M. L., Kopperud, R., and Hildemann, L. M.: Characterization of Organic Compounds Collected during Southeastern Aerosol and Visibility Study: Water-Soluble Organic Species, *Environmental Science & Technology*, 39, 707–715, 2005.
- Zhang, R., Gen, M., Huang, D., Li, Y., and Chan, C. K.: Enhanced Sulfate Production by Nitrate Photolysis in the Presence of Halide Ions in
815 Atmospheric Particles, *Environmental Science & Technology*, 54, 3831–3839, 2020a.
- Zhang, Y., Apsokardu, M. J., Kerecman, D. E., Achtenhagen, M., and Johnston, M. V.: Reaction Kinetics of Organic Aerosol Studied by Droplet Assisted Ionization: Enhanced Reactivity in Droplets Relative to Bulk Solution, *Journal of The American Society for Mass Spectrometry*, p. jasm.0c00057, 2020b.
- Zhong, J., Li, H., Kumar, M., Liu, J., Liu, L., Zhang, X., Zeng, X. C., and Francisco, J. S.: Mechanistic Insight into the Reaction of Organic
820 Acids with SO₃ at the Air–Water Interface, *Angewandte Chemie*, 115, 3803–5, 2019.
- Zobrist, B., Soonsin, V., and Marcolli, C.: Ultra-slow water diffusion in aqueous sucrose glasses, *Physical Chemistry Chemical Physics*, 13, 3514–13, 2011.