

# African Dust Particles over the Western Caribbean Part I:

## Impact on air quality over the Yucatan Peninsula

Carolina Ramírez-Romero<sup>1</sup>, Alejandro Jaramillo<sup>1</sup>, María F. Córdoba<sup>1,2</sup>, Graciela B. Raga<sup>1</sup>,  
Javier Miranda<sup>3</sup>, Harry Alvarez<sup>4</sup>, Daniel Rosas<sup>5</sup>, Talib Amador<sup>5</sup>, Jong Sung Kim<sup>6</sup>,  
5 Jacqueline Yakobi-Hancock<sup>6</sup>, Darrel Baumgardner<sup>7</sup>, and Luis A. Ladino<sup>1,\*</sup>

<sup>1</sup>Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México, México City, México.

<sup>2</sup>Posgrado en Ciencias Químicas, Universidad Nacional Autónoma de México, México City, México.

<sup>3</sup>Instituto de Física, Universidad Nacional Autónoma de México, México City, México.

10 <sup>4</sup>Facultad de Ciencias, Universidad Nacional Autónoma de México, México City, México.

<sup>5</sup>Universidad Autónoma de Yucatán, Mérida, Yucatán, México.

<sup>6</sup>Dalhousie University, Halifax, Nova Scotia, Canada.

<sup>7</sup>Droplet Measurement Technologies, Colorado, USA

\*Correspondence to: L.A.L ([luis.ladino@atmosfera.unam.mx](mailto:luis.ladino@atmosfera.unam.mx))

15

**Abstract.** On a global scale, African dust is known as one of the major sources of mineral dust particles as they can be efficiently transported to different parts of the planet. Several studies have suggested that the Yucatan Peninsula could be influenced by such particles, especially in July, associated with the strengthening of the Caribbean low level jet. Although these particles have the potential to impact the local air quality significantly,  
20 as shown elsewhere (especially particulate matter, PM), the arrival and the impact of African dust into Mexican territory has not been quantitatively reported to date.

Two short-term field campaigns were conducted to confirm the arrival of African dust onto the Yucatan Peninsula in July 2017 and July 2018 at the city of Merida atmospheric observatory (20.98N 89.64W). Aerosol  
25 particles were monitored at the ground level by different on-line and off-line sensors. Several PM<sub>2.5</sub> and PM<sub>10</sub> peaks were observed during both sampling periods, with a relative increase in the PM levels ranging between 200% and 500% with respect to the normal background. Given that these peaks were found to highly correlate with super micron particles and chemical elements typically found in mineral dust particles, such as Al, Fe, Si, and K, they are linked with African dust. This conclusion is supported by combining back trajectories with  
30 vertical profiles from radiosondes, reanalysis, and satellite images to show that the origin of the air masses arriving at Merida was the Saharan Air Layer (SAL). The good agreement found between the measured PM<sub>10</sub> concentrations and the estimated dust mixing ratio content from MERRA-2 (Version 2 of the Modern-Era Retrospective analysis for Research and Applications) corroborates the conclusion that the degradation of the local (and likely regional) air quality in Merida is a result of the arrival of African dust.

35

### 1. Introduction

The second largest natural contribution of atmospheric particles, worldwide, after sea spray, is mineral dust (Pey et al., 2013). Although volcanoes and soil dust from agricultural activities are significant sources of mineral dust (Walker, 1981; Tegen et al., 2004), the largest sources are the deserts distributed around the world (Goudie

40 and Middleton, 2006). Africa is considered one of the most important of these sources as it emits ca. 800 Tg  
yr<sup>-1</sup>, corresponding to ca. 70% of the global dust (Prospero et al., 2014; Ryder et al., 2019). Therefore, African  
dust particles play a significant role in the climate system as they can affect the planetary radiative balance and  
the hydrological cycle. Their optical properties, i.e. scattering and absorption, modulate radiative forcing and  
45 as cloud condensation nuclei and/or ice nucleating particles they will impact cloud formation and evolution  
(Zhang et al., 2007; Hoose and Möhler, 2012; DeMott et al., 2015; Kanji et al. 2017). Additionally, several  
studies have shown that the presence of mineral dust can influence tropical cyclone formation (Dunion and  
Velden, 2004; Evan et al., 2006), and human health as these particles degrade air quality (Carlson and Prospero,  
1972; Prospero, 1999; [Prospero et al., 2014](#); Venero-Fernández, 2016).

50 African dust particles are efficiently transported far from their emission source (Perry et al, 1997; Chiapello et  
al., 1997). According to Middleton and Goudie (2001), there are different trajectories that African dust  
experiences around the world. Among the most important, African dust particles can be transported to the  
western Mediterranean and Europe (Karanasiou et al., 2012; Perez et al., 2008; Prodi and Fea, 1979; Salvador  
et al., 2014); to the eastern Mediterranean and the Middle East (Ganor and Mamane, 1982; Ganor et al., 2010;  
55 Athanasopoulou et al., 2016), and towards the South of the African continent (d’Almeida, 1986; Resch et al.,  
2008). Additionally, African dust is transported across the Atlantic Ocean to the United States, Mexico, the  
Caribbean region, and South America ([Prospero et al. 1981](#); Bravo et al., 1982; Perry et al, 1997; Chiapello et  
al., 1997; Prospero and Lamb, 2003; Venero-Fernández, 2016; [Barkley et al., 2019](#); Kramer et al., 2020). The  
long-range transport of African dust over the Atlantic represents 25% of the total emissions from the Saharan  
60 desert (Shao et al., 2011). This transport is favored in the Northern hemisphere during the summer (i.e., from  
June to September) within a dry and hot elevated layer called the Saharan Air Layer (SAL) (Carlson and  
Prospero, 1972; Prospero and Carlson, 1972; Karyampudi and Carlson, 1988; Tsamalis et al., 2013; Weinzierl  
et al., 2016).

65 During the summer, the SAL ascends to altitudes between 5-7 km through interactions with cool marine air  
masses (Adams et al., 2012; Chouza et al., 2016; Korte et al., 2018). Dunion and Velden (2004), Dunion and  
Marron (2008), and Dunion (2011) studied the characteristics of the air masses that reach the North Atlantic  
and the Caribbean region during the boreal summer months. They found that there are three distinct air masses:  
moist tropical (MT), Saharan air layer (SAL), and mid-latitude dry air intrusions (MLDAIs). Each type of air  
70 mass is associated with unique thermodynamic and kinematic characteristics, and have a wide range of possible  
origins. However, the SAL and MLDAl air masses have distinct flow patterns across the North Atlantic, which  
allows differentiating these masses by tracking their origin. In contrast, their distinctly unique moisture  
characteristics allow differentiating the MT from SAL air masses (Dunion, 2011).

75 There are different methods for the detection of the long-range transport of African dust and its presence in  
different regions around the world. For several decades the tracking of dust events has been studied by remote  
sensing (Chiapello et al., 1999; Dunion and Velden, 2004; Foltz and McPhaden, 2008; Prospero et al., 2002~~5~~;

Liu et al., 2008; Voss and Evan, 2020). Ground- and space-based tools such as the light detection and ranging (LIDAR) and satellite sensors (e.g., the moderate resolution imaging spectroradiometer (MODIS) and the visible infrared imaging radiometer suite (VIIRS)) provide the aerosol spatial distribution with altitude in terms of the aerosol optical depth (AOD), mass concentration, and particle size distribution (Zhang and Reid, 2006; Jackson et al., 2013). ~~Additionally, the cloud aerosol lidar and infrared pathfinder satellite observations (CALIPSO) can quantify (in three dimensions) the trans-Atlantic transport of African dust (Liu et al., 2008; Adams et al., 2012; Chouza et al., 2016; Korte et al., 2018).~~

Another useful tool is the reanalysis from global climate models that assimilates, in a statistically optimal way, satellite and ground observations. The reanalysis produces continuous, four-dimensional fields of different atmospheric variables of interest, contrasting with the observations that may be spatially and temporally sparse (Cohn, 1997; Kalnay, 2003; Rienecker et al., 2011; Schutgens et al., 2010). The use of reanalysis, considering its inherent uncertainties, has become an essential tool in the atmospheric research community (Gelaro et al., 2017). For example, the hybrid single-particle lagrangian integrated trajectory (HYSPLIT) model has been successfully used to track the transport of African dust particles (e.g., Ashrafi et al., 2014; Prospero et al., 2005). HYSPLIT uses meteorological data from different modeling sources, including the NCEP-NCAR National Centers for Environmental Prediction (NCEP)–National Center for Atmospheric Research (NCAR) reanalysis model (Stein et al., 2015).

The transport of African dust can also be evaluated with NASA's Global Modeling and Assimilation Office (GMAO) MERRA-2 reanalysis (Prospero et al., 2020). MERRA-2 (Version 2 of the Modern-Era Retrospective analysis for Research and Applications) is the first multidecadal reanalysis that assimilates both meteorological and aerosol data from various ground- and space-based remote sensing sources (Gelaro et al., 2017; Randles et al., 2017). Despite some deficiencies, previous studies have demonstrated that the MERRA-2's aerosol assimilation system does indeed show considerable skill in simulating numerous observable aerosol properties (e.g., Buchard et al., 2015, 2016, 2017; Randles et al., 2017). MERRA-2 has been previously used to study the effects of aerosol particles in the earth system, in several studies focused on dust-related phenomena. For example, Buchard et al. (2017) showed the benefit of the MERRA-2 assimilation for the retrieval of the seasonality, vertical distribution, and magnitude of the dust surface concentrations during an episode of dust transport from Africa to the Caribbean. Later on, Veselovskii et al. (2018) showed the consistency of the MERRA-2 aerosol products with MIE-Raman lidar observations performed in West Africa during a smoke and dust mixing event. Similarly, Grogan and Thornicrof (2019) studied the characteristics of African easterly waves and their relationship with synoptic-scale plumes of Saharan mineral dust. More recently, Bibi et al. (2020) studied atmospheric dust load and deposition fluxes along the North-African coast of the Mediterranean Sea; and Aldhaif et al. (2020) studied dust events impacting the United States East Coast.

The *in situ* monitoring of aerosol properties, such as aerosol size and mass distribution, is very useful to determine their influence on local air quality and human health (Querol et al., 2019). Hence, different studies

120 have been carried out in the Caribbean islands and Florida to quantify the impact of African dust on the local  
air quality (Prospero, 1999; Prospero and Mayol-Bracero, 2013; Prospero et al., 2014). In Barbados, the  
monitoring of the atmospheric aerosol mass began in 1965, while in Miami, Florida, it began in 1974 and  
continues to the present (Prospero and Mayol-Bracero, 2013). In Barbados, it is estimated that 50% of the PM<sub>2.5</sub>  
(i.e., with an aerodynamic diameter  $d < 2.5 \mu\text{m}$ ) and ca. 90% of the PM<sub>10</sub> (i.e.,  $d < 10 \mu\text{m}$ ) consist of African dust  
(Li-Jones and Prospero, 1998; Prospero et al., 2001; Reid et al., 2003a). In Miami, the mean daily mass  
concentration of mineral dust during the summer typically ranges between 10 and 100  $\mu\text{g m}^{-3}$ , with a large  
interannual variability (Prospero et al., 2001). During the Puerto Rico Dust Experiment (PRIDE) campaign  
125 ~~carried out between 28 June and 24 July, 2000~~, the mineral dust concentration at the ground level was found to  
exceed 70  $\mu\text{g m}^{-3}$  (Reid et al., 2003b). In the aforementioned studies, the African dust particles transported over  
the Atlantic affected the local air quality, exceeding the World Health Organization (WHO) guidelines for PM<sub>2.5</sub>  
and PM<sub>10</sub> (i.e., above 25  $\mu\text{g m}^{-3}$  for PM<sub>2.5</sub> and 50  $\mu\text{g m}^{-3}$  for PM<sub>10</sub>). According to the WHO, air pollution and its  
effects are considered a global health priority (WHO, 2002). Several studies have linked high concentrations of  
mineral dust (in terms of PM<sub>2.5</sub> and PM<sub>10</sub>) to brain, [skin](#), [lung](#), cardiovascular, [cerebrovascular](#), and respiratory  
130 diseases ([Alessandrini et al., 2013](#); [Goudie, 2014](#); [Wilker et al., 2015](#); [Brook et al., 2010](#); [Dominici et al., 2006](#);  
[Zhang et al., 2016](#)). ~~According to Goudie (2014) and Zhang et al. (2016), inhaled dust particles can cause  
damage to the lungs, in addition to other parts of the body such as the heart, skin, and brain. For example, in  
Rome, Italy, the effects of PM<sub>10</sub> on cerebrovascular diseases were found to increase by 5% due to the presence  
of African dust particles (Alessandrini et al., 2013). Similarly, Rodríguez Cotto et al. (2013) found that PM<sub>2.5</sub>  
and PM<sub>10</sub> can cause asthma and allergic reactions mainly to children during the epochs of African dust in Puerto  
Rico.~~ Additionally, African dust particles have been found to serve as carriers for biological material. Griffin  
et al. (2001) reported the presence of viable bacteria and fungi associated with the arrival of African dust over  
the U.S. Virgin Islands. Similarly, Rodríguez-Gomez et al. (2020) found a higher concentration of viable  
bacteria and fungal propagules during summer than during winter in the Yucatan Peninsula, with summer being  
135 the season when African dust intrusions are more frequent.

The chemical and mineralogical composition of particles also plays an important role in the identification of  
African dust in the receptor regions (Nenes et al., 2014). The most abundant minerals present in these particles  
are silicates (quartz), clay minerals (kaolinite, illite, chlorite, palygorskite), feldspars (albite, anorthite) and  
145 carbonates (calcite) (Goudie and Middleton, 2006; Querol et al., 2019; Broadley et al., 2012). The major oxides  
in Saharan dust are SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, Fe<sub>2</sub>O<sub>3</sub>, CaO, MgO and K<sub>2</sub>O and to a lesser extent P<sub>2</sub>O<sub>5</sub> and TiO<sub>2</sub> (Goudie and  
Middleton, 2006; Linke et al., 2006). Several studies in the Caribbean have identified high levels of Fe and Al  
in dust events (Prospero et al., 2001; Rosinski et al., 1988). Additionally, Rosinski et al. (1988) reported high  
percentages of Si and Mg in particles collected in the Gulf of Mexico (GoM) during July.

150 Although the arrival of African dust in Mexico has been suggested for decades (e.g., Bravo et al., 1982;  
Prospero, 1999; Lenes et al., 2012), to our knowledge, there has not been a comprehensive study, published in  
the open, peer-reviewed literature, that documents this atmospheric phenomenon. For the first time, in this study

we document the arrival of African dust in the Yucatan Peninsula for two consecutive years (i.e., 2017 and 2018) using *in situ* and remote sensing measurements, reanalysis, back trajectory analysis, and complementary meteorological observations.

## 2. Materials and Methods

### 2.1 Sampling site and field campaigns

160 The Yucatan Peninsula is located in the southeast of Mexico. It borders with the [Gulf of Mexico \(GoM\)](#) to the north, the Atlantic Ocean to the east, and the Caribbean Sea, Guatemala, and Belize to the south. The Yucatan has characteristics that are unique to this region (Plasencia, 1998). For example, its uniform terrain, the absence of rivers, and the type of soil, formed by Cretaceous sediments that do not present mineralization and are rich in calcium, commonly called “Laja de Yucatán” (Plasencia, 1998) sets the Yucatan aside from other regions of Mexico. The average temperature of the Yucatan Peninsula ranges from 25°C to 35°C (World Resource Institute, 2018) with an average annual relative humidity of 79% (INEGI, 2009). The Peninsula has a warm, semi-dry climate on the coast and a warm, sub-humid climate throughout the rest of the region, with a rainy season between summer and autumn (June-October) (Orellana et al., 2009; ProAire, 2018). Precipitation in this region is mainly due to convective activity and it is influenced by the moisture advection by the trade winds (Orellana et al., 2009; ProAire, 2018). *In situ* measurements were made in the city of Mérida, situated in the northeast sector of the Yucatan Peninsula (20.98 °N, 89.64 °W), that is the capital of the Yucatan state. Mérida has 892,363 inhabitants (INEGI, 2015) and is 23 km away from the coast. The most representative activities in the region are tourism, commerce, and the textile industry (INEGI, 2017).

175 Aerosol particles were continuously monitored with sensors installed at the School of Chemistry of the Universidad Autónoma de Yucatán (FC-UADY), located in the central-western part of the city (Figure 1), as part of the African Dust and Biomass Burning Over Yucatan (ADABBOY) project. Table 1 lists the instrumentation used to characterize particle physical, optical and chemical properties. The Partisol and MiniVol were installed on the rooftop of the FC-UADY while the other instruments were maintained in an environmentally controlled area where they sampled from inlets connected to a ventilated chimney that extended approximately 1.5 m above the roof. This measurement site is part of the University Network of Atmospheric Observatories (RUOA) supported by the National University of Mexico (UNAM). Two intensive sampling periods were conducted between July 11 - July 31, 2017 and June 30 - July 17, 2018.

### 185 2.2. Aerosol concentration and particle size distribution

The particulate mass concentration was monitored continuously with PM<sub>2.5</sub> and PM<sub>10</sub> analyzers providing real-time measurements (FH 62 C14 Thermo Scientific Inc) with a temporal resolution of one minute at a sampling flow rate of 16.7 L min<sup>-1</sup> (Thermo Fisher Scientific Inc, 2007).

190 The total number concentration of particles with sizes approximately larger than 50 nm was measured by a  
condensation particle counter (CPC 3010, TSI) at a sampling rate of 1 hz with a flow rate of 1.0 L min<sup>-1</sup> and  
the aerosol number concentration as a function of particle size was monitored by an optical particle counter  
(LasAir II 310A, MSP). The LasAir has six different size bins (0.3, 0.5, 1.0, 5.0, 10.0, and 25 µm), a flow rate  
of 28.3 L min<sup>-1</sup> and a time resolution of 11 s.

195

### 2.3 Aerosol collection and chemical composition analysis

PM<sub>2.5</sub> and PM<sub>10</sub> aerosol particles were collected for 24 h with a Partisol model 2525 (Thermo Fisher Scientific  
Inc.) and for 48 h with a Minivol (3380, Air metrics) on 47 mm teflon filters (Pall Science). The MiniVol and  
Partisol flow rates were 5.0 L min<sup>-1</sup> and 16.7 L min<sup>-1</sup>, respectively. After the sampling periods, the filters were  
200 placed in 60 mm Petri dishes and stored at 4°C prior to the chemical analysis.

Elemental analysis was performed on each filter using X-Ray Fluorescence (XRF) with the X-ray spectrometer  
at Laboratorio de Aerosoles, Instituto de Fisica, UNAM (Espinosa et al., 2012). The X-ray tube was made by  
Oxford Instruments (Scotts Valley, CA, USA), an Rh anode and an Amptek X-123SDD spectrometer (Bedford,  
205 MA, USA) were used. The samples were irradiated for 900 s working with a current of 500 µA and resulting in  
a spectrum that was analyzed using the WinQXAS computer code (IAEA, 1997). The product of this analysis  
derived mass concentrations of Fe, Al, Si, Ca, Na, P, Mg, Mn, Ti, Cl, P, Zn, K, S, Cu, and Ni along with their  
associated uncertainties, as described by Espinosa et al. (2010).

### 2.4 Meteorological and satellite data

The local and regional meteorological conditions were monitored using different approaches. The RUOA  
meteorological sensors were placed at the rooftop of the FC-UADY (Table 1) continuously measuring the wind  
speed and direction, air temperature, relative humidity, solar radiation, and precipitation. To derive the regional  
and vertical distribution of meteorological conditions, radiosondes and reanalysis were used. The information  
215 provided by the radiosondes launched was from three World Meteorological Organization (WMO) stations in  
the Yucatan Peninsula, as shown in Figure 1. These locations are: Merida (Mérida International Airport,  
Mexico. (WMO index: 76644), Cancún (WMO index: 76595), and Belize (Philip SW Goldson International  
Airport, Belize. WMO index: 78583). The processed radiosonde data is obtained from the University of  
Wyoming (<http://weather.uwyo.edu/upperair/sounding.html>).

220

Hourly total precipitable water vapor and three-dimensional 3-hourly aerosol mixing ratio data were obtained  
from the MERRA-2 reanalysis (GMAO, 2015a, 2015b). The aerosol properties in MERRA-2 were simulated  
with the Goddard Chemistry Aerosol Radiation and Transport model (GOCART), which takes into account the  
sources, sinks, and chemistry of 15 externally-mixed aerosol mass mixing ratio tracers: dust (five non-  
225 interacting size bins), sea salt (five non-interacting size bins), hydrophobic and hydrophilic black and organic  
carbon (BC and OC, respectively; four tracers), and sulfate (SO<sub>4</sub>) (Randles et al., 2017; Buchard et al., 2017).

The air mass back trajectories were calculated using the HYSPLIT model from the National Oceanic and Atmospheric Administration (NOAA). In conjunction with the *in situ* measurements, the back trajectories were calculated considering the maximum concentration of PM reported by the PM<sub>2.5</sub> and PM<sub>10</sub> analyzers. The trajectories were initiated at 50, 250 and 500 m above ground level going backward in time for 13 days. Although Kramer et al. (2020) reported that mineral dust particles arrive in Miami ca. 10 days after they are produced in North Africa, they may take a longer time to reach the Yucatan Peninsula. The different heights of the HYSPLIT back-trajectories runs were chosen to determine the rate of descent of dust air masses to the surface. Also considered was the aerosol optical depth (AOD) measured with MODIS instruments in the Aqua and Terra satellites.

### 3. Results and Discussion

#### 3.1. Local evidence

Several studies have shown that air quality (PM<sub>2.5</sub> and PM<sub>10</sub>) significantly deteriorates upon the arrival of African dust plumes (Prospero and Lamb, 2003; Prospero et al., 2001; Prospero and Mayol-Bracero, 2013; Prospero et al., 2014). Figure 2 shows the time series of the PM<sub>2.5</sub> and PM<sub>10</sub> concentrations for the July-August period of 2017 and 2018. Some high concentration PM peaks are clearly identified, with PM<sub>2.5</sub> and PM<sub>10</sub> values as high as 54  $\mu\text{g m}^{-3}$  and 135  $\mu\text{g m}^{-3}$ , respectively. Henceforth those peaks will be referred to in our study as African dust peaks (ADPs). Note that the background concentration ([defined as the lowest values within the sampling period where the chemical composition was available](#)) of PM<sub>2.5</sub> is  $\sim 4 \mu\text{g m}^{-3}$  and of PM<sub>10</sub> is  $\sim 10 \mu\text{g m}^{-3}$ . The ADPs found in 2017 (i.e., July 22-24, 27-28, and August 4, 6-7) resulted in an increase of 300% in PM<sub>2.5</sub> and 500% in PM<sub>10</sub> with respect to the background. In 2018, the ADPs (i.e., July 10-11, 13-15, 16-17, 23-26, and August 9-10) exceeded 200% and 300% of the background levels of PM<sub>2.5</sub> and PM<sub>10</sub>, respectively. The aforementioned ADPs not only exceeded the PM<sub>2.5</sub> and PM<sub>10</sub> thresholds suggested by the WHO (i.e., PM<sub>2.5</sub>=25  $\mu\text{g m}^{-3}$  and PM<sub>10</sub>=50  $\mu\text{g m}^{-3}$ , 24-h mean) but more than double them, as was the case for the August 9-12, 2018 event. Similar behavior has been previously observed in Puerto Rico, Miami, and Barbados during the arrival of African dust particles (Reid et al., 2003b; Prospero et al., 2005, 2014). The mass concentrations of PM<sub>2.5</sub> and PM<sub>10</sub> were found to be 49% and 54% higher in 2018 than in 2017, respectively, suggesting a higher frequency or intensity of African dust plumes arriving over Merida in 2018.

Figure 2 also shows the elemental composition obtained from the XRF analysis (16 elements) for five ADPs observed during the 2017 and 2018 field campaigns. In addition, one day from each field campaign was selected to determine the elemental background composition. The selected days are July 10-11 in 2017 and July 6 in 2018. These days were chosen because the PM<sub>2.5</sub> and PM<sub>10</sub> concentrations were within the mean background values and did not include any atypical peak nor apparent external influence.

High levels of sodium (Na, pink), chlorine (Cl, turquoise blue), sulfur (S, dark orange), and calcium (Ca, light green) were found in the background samples, corresponding to >70% of the total mass. The presence of Na

265 and Cl are expected in airborne particles at this site given the city's proximity to the GoM (i.e., 23 km away). Cerón et al. (2002) reported large concentrations of Na, Cl, and Mg that originated from sea salt, when analyzing the composition of rainwater from the Yucatan Peninsula. The high levels of S can be associated with local anthropogenic activities such as vehicular, ship, and industrial emissions (e.g., Corbett and Fischbeck, 1997; Cerón-Bretón et al., 2018). Additionally, given the short distance between Merida and the GoM, it is possible  
270 that dimethylsulfide (DMS) production from plankton in the GoM can be a natural source of S, as has been shown in other studies (e.g., Rosinski et al., 1988; Kloster et al., 2006; Vallina and Simó, 2007). Finally, the presence of Ca could be related to the limestone soil prevalent in the Yucatan Peninsula and the resuspension of road dust (Plasencia, 1998; Querol et al., 2019).

275 Interestingly, the elemental composition of the airborne particles collected during the ADPs showed higher concentrations of silica (Si, dark yellow), aluminum (Al, light purple), and iron (Fe, dark purple) than those in the background particles. While the Si concentrations are approximately three times larger than the baseline, Al and Fe increased by eight and 12 times, respectively. To corroborate the relationship between the increase in PM and the African dust, each of the 16 elements analyzed by XRF were correlated with the PM<sub>2.5</sub> and PM<sub>10</sub>  
280 concentrations. Out of the 16 elements Al, Si, K, and Fe were the only ones with correlation coefficients  $r > 0.6$  ( $p < 0.05$ ) for both years, as shown in Figure S1. The present results are in agreement with previous studies that showed high correlation coefficients between the aforementioned elements (e.g., Caquineau et al., 1998; Guieu et al., 2002; Trapp et al., 2010). Also, the concentration of aerosol particles with diameters between 0.5  $\mu\text{m}$  and 25  $\mu\text{m}$ , as measured by the LasAir, were found to highly correlate with the PM<sub>2.5</sub> and PM<sub>10</sub> concentrations,  
285  $r = 0.79$  and  $r = 0.87$ , respectively (Figure S2). Finally, the typical background particle size distribution showed significant changes during the arrival of ADPs for particles ranging between 0.5  $\mu\text{m}$  and 5.0  $\mu\text{m}$  (Figure S3). It is widely known that the typical size of African dust particles transported over long distances ranges from 0.1  $\mu\text{m}$  to 20  $\mu\text{m}$  (e.g., Bégue et al., 2012; Denjean et al. 2016). Overall, the high concentration of coarse particles and the increase of Al, Si, and Fe during the ADPs, together with the good correlations found between the PM<sub>2.5</sub>  
290 and PM<sub>10</sub> concentrations with Al, Si, K, Fe, and particles larger than 0.5  $\mu\text{m}$  strongly suggests that the ADPs are mineral particles associated with dust transported from Africa to Mexico.

Additionally, it is important to note that Al, Si, K, and Fe are common oxides found in African dust composed of minerals and clays such as: Quartz ( $\text{SiO}_2$ ), Kaolinite ( $\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4$ ), Illite  
295 ( $(\text{K},\text{H}_3\text{O})(\text{Al},\text{Mg},\text{Fe})_2(\text{Si},\text{Al})_4\text{O}_{10}[(\text{OH})_2,(\text{H}_2\text{O})]$ ), Chlorite ( $(\text{MgFe})_5\text{Al}(\text{AlSi}_3)\text{O}_{10}(\text{OH})_8$ ), Palygorskite ( $(\text{Mg},\text{Al})_2\text{Si}_4\text{O}_{10}(\text{OH})_4(\text{H}_2\text{O})$ ), and Feldspars such as albite ( $\text{NaAlSi}_3\text{O}_8$ ), Anorthite ( $\text{CaAl}_2\text{Si}_2\text{O}_8$ ) and orthoclase  $\text{KAlSi}_3\text{O}_8$ , among others (A. Goudie and Middleton, 2006; Linke et al., 2006; Broadley et al., 2012; Querol et al., 2019). Rosinski et al. (1988) reports that up to 90% of the collected airborne particles in the presence of  
300 dust events in the GoM contained Al, Fe, Si. In Puerto Rico, Reid et al., (2003b) found that during dust events that reached the island, the concentrations of Si and Al on aerosol particles ( $> 0.74 \mu\text{m}$ ) were above  $10 \mu\text{g m}^{-3}$



and  $5 \mu\text{g m}^{-3}$ , respectively. Similarly, Prospero et al., (2001) reports concentrations of Al  $> 1.0 \mu\text{g m}^{-3}$  and Fe  $> 0.5 \mu\text{g m}^{-3}$  on  $\text{PM}_{2.5}$  on days when high concentrations of African dust particles were reported in Miami.

To confirm that no aerosol sources other than the African dust were the origin of the high PM peaks observed in Merida in July 2017 and 2018, the  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  concentrations were correlated with other measured variables. As shown in Figures S2 and S4,  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  concentrations are poorly correlated ( $r < 0.09$ ) with local pollution emissions such as [particle-bound polycyclic aromatic hydrocarbons \(pPAHs\)](#), black carbon (inferred from the absorption coefficient), and [nitrogen oxides \(NOx\)](#). Note that those gases and particles can be considered as proxies of anthropogenic pollutants generated by the incomplete combustion of fossil fuels and biomass burning, as previously demonstrated for Merida (Muñoz-Salazar et al., 2020; Alvarez-Ospina et al., 2020). Also, correlation coefficients below 0.29 were found between  $\text{O}_3$  and solar radiation with the  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  concentrations [indicating that it is very unlikely that secondary organic particles could be the source of the ADPs observed in Merida, as was the case in high particle concentration events shown by Muñoz-Salazar et al. \(2020\).](#) ~~Muñoz-Salazar et al., (2020) found in Merida that ultrafine aerosol particles of secondary origin are correlated with  $\text{O}_3$ , a proxy of photochemical activity and hence, of secondary particle production. Therefore, it is very unlikely that secondary organic particles could be the source of the ADPs observed in Merida.~~

Finally, although none of the different meteorological variables monitored at the surface level were found to correlate with the  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  concentrations, as shown by the wind roses in Figure S5, easterly winds were prevalent when ADPs were observed. This is relevant since African dust can only be transported by easterly winds.

### 3.2. Larger scale observations

To evaluate the source of the ADPs observed in Merida from a large-scale perspective, we focus on the classification of tropical air masses in the North Atlantic and the Caribbean region during the boreal summer months proposed by Dunion (2011). We used HYSPLIT to estimate the trajectories of different air masses that reached Merida in the periods of July-August 2017 and 2018. HYSPLIT trajectories for the 2017 and 2018 ADPs point to an African origin and, therefore, suggest that these air masses are either MT or SAL (See Figure S6). To differentiate the MT from SAL, we focused on their distinctly unique moisture characteristics. Dunion (2011) proposes that a threshold of 45 mm of total precipitable water vapor (PWV), which corresponds to the total amount of water vapor contained in the atmospheric column from the surface to the top of the troposphere (AMS, 2000), can be used to differentiate dry from moist air masses. This value is consistent with other studies that use PWV to identify dry-air days (e.g., Hanks and Marinaro, 2016), and since deep tropical convection begins to increase above a critical PWV value of 50 mm (Holloway and Neelin, 2009). [Note that PWV is given by the vertical integral of the mixing ratio  \$x\(p\)\$  at the pressure level  \$p\$  in the layer bounded by pressures  \$p\_1\$  and  \$p\_2\$  and can be calculated using Equation 1 \(AMS, 2000\):](#)

$$PWV = \frac{1}{\rho g} \int_{p_1}^{p_2} x dp \quad \text{Equation (1)}$$

340 [where  \$\rho\$  represents the density of water and  \$g\$  is the acceleration of gravity.](#)

Figure 3 shows the time series of PWV for the July-August 2017 and 2018 periods at each WMO radiosonde site. The black solid line shows PWV from MERRA-2, [available from the Vertically Integrated Diagnostics](#) (GMAO, 2015a), together with PWV estimated [using Equation 1](#) from the available radiosonde profiles shown as the dashed blue lines. One caveat is that in the periods of interest, there is a striking lack of radiosonde data. Nevertheless, we can see a good agreement between the available observed PWV and that of PWV from MERRA-2. Therefore, the latter can be used as a good approximation for PWV in the region to differentiate moist from dry air masses. In Figure 3, the periods where PWV is less than 45 mm are highlighted in red. These periods show dry air masses that coincide with air mass trajectories with an African origin (i.e., in 2017: July 22-24, July 27-28, August 04, and August 6-7; and in 2018: July 10-12, July 13-15, July 16-17, July 23-26 and August 9-12), allowing us to conclude that these dry air masses have mainly SAL characteristics.

The arrival of African dust in Merida was also explored from the MERRA-2 dataset. Figure 4 shows the time series of the estimated vertical profiles of the 3-hour time series of dust mixing ratio from MERRA-2 at Merida for 2017 and 2018. It shows that the events corresponding to the arrival of dry air masses from Africa shown in Fig. 3 nicely correlate with high dust mixing ratios, strongly supporting the hypothesis of SAL air reaching the Yucatan Peninsula. Figure 4 also shows that the July-August period of 2018 was particularly active with frequent arrivals of dust in the region, in agreement with the higher  $PM_{2.5}$  and  $PM_{10}$  concentrations measured in 2018, as depicted in Fig. 2. Figure S7 focuses on the vertical profiles of dust mixing ratio for the periods of July 21 to 25, 2017 and July 12 to 16, 2018. These periods show the increase of dust in the atmospheric column in the studied region, supporting the hypothesis that the source of the ADPs shown in Figure 2 is likely African dust.

Finally, the arrival of African dust plumes over the Yucatan Peninsula was confirmed by investigating the AOD detected by the MODIS Aqua/Terra satellite for July 2017 and 2018, as shown in Figures S8 and S9. Although this information cannot be used to perform quantitative analysis, the AOD images allow us to confirm the arrival of African dust plumes into the Yucatan Peninsula. Additionally, as in Figures 2, 3, and 4, the AOD images also show that the African dust plumes activity was higher in 2018 than in 2017. Previous studies have also used the AOD from MODIS to identify the arrival of African dust. For example, Koren et al. (2006) tracked the long-range transport of dust from the Bodélé depression (North-Central Africa) to the Amazon basin. Similarly, Kalashnikova and Kahn (2008) demonstrated that with the MODIS it is possible to observe the evolution of African dust plumes over the Atlantic Ocean. Additionally, Kaufman et al., (2005) identified and quantified the transport and deposition of mineral dust over the Atlantic Ocean using MODIS data.

### 375 3.3. Comparison of *in situ* observations and reanalysis

The daily mean PM<sub>10</sub> from MERRA-2 was estimated using the method proposed by Provençal et al. (2017). The black line in Fig. 5 shows the estimated PM<sub>10</sub>, which is compared to the PM<sub>10</sub> measured by the RUOA station in Merida (blue line). The estimated surface dust mixing ratio from MERRA-2 is also shown in red. Figure 5 shows that MERRA-2 overestimates compared to the ground-based measurements of PM<sub>10</sub>.  
380 Nevertheless, it should be clarified that the reanalysis information corresponds to a region 0.5° x 0.625°, implying that the MERRA-2 is estimating the regional average, while the station corresponds to a local measurement.

Despite these differences, Figure 5 shows that the observations at the RUOA station have variations similar to those of MERRA-2. Figure 6 shows the dispersion diagram of the daily mean surface dust mixing ratio from MERRA-2 vs. PM<sub>10</sub> measured from RUOA station for the periods indicated in Figure 5. It shows a good high correlation between the estimated dust and the measured PM<sub>10</sub> in particular for the 2018 period, which was particularly active with constant arrivals of African dust to the region, as shown in Figure 4. A similar analysis was performed for the 3-h estimated and measured PM<sub>10</sub>, as shown in Figures S10 and S11, with identical  
390 conclusions as for 24-h averages.

### 4. Conclusions

For the first time, the arrival of African dust into Mexican territory is quantitatively verified. The arrival of African dust particles in Merida significantly degraded the local air quality as PM<sub>2.5</sub> and PM<sub>10</sub> concentrations  
395 increased up to 500% with respect to the background. Therefore, the presence of African dust in Merida and the Yucatan Peninsula could be has at the potential health threat to their inhabitants to trigger or exacerbate several diseases, as has been reported elsewhere (Wilker et al., 2015; Brook et al., 2010; Dominici et al., 2006). Although the African dust intrusions caused an increase in particulate matter in Merida (Mexico), this increase is lower than those reported in other places closer to the Sahara such as Barbados and the Mediterranean. The arrival of African dust in other regions of the world has led to even higher PM concentrations than those found in the present study. For example, Querol et al. (2009) reported in Spain peak concentrations of PM<sub>10</sub> during dust events that reached up to 250 µg m<sup>-3</sup>. Similarly, in Nicosia (Cyprus), concentrations of PM<sub>10</sub> up to 470 µg m<sup>-3</sup> were measured (Achilleos et al., 2014). Moreover, In addition to the impacts on air quality, African dust particles can also be a serious health threat as they serve as the carrier of biological material originating in  
400 Africa as reported by Rodriguez-Gomez et al. (2020). If the foreign biological particles are opportunistic pathogens, they can cause a variety of diseases in the receptor regions, such as the Yucatan Peninsula. Finally, those particles can impact the development of precipitation affecting the regional hydrological cycle when they serve as efficient ice nucleating particles (e.g., Rosinski et al., 1988; Córdoba et al., 2020) Hoese and Möhler, 2012; Murray et al., 2012; Kanji et al., 2017.

410

As shown in the present study, combining ground-based off-line and on-line sensors provides robust evidence of the arrival of African dust; however, we also show that the combination of back trajectories with radiosondes, and the estimated surface dust mixing ratio from MERRA-2 are powerful tools that can be exploited when *in situ* information is missing, especially in developing countries where the necessary instrumentation is scarce.

415

Continuous monitoring of the arrival of African dust is of high importance not only in the Caribbean islands but also at other sites in Latin America such as Mexico, Belize, Guatemala, Honduras, etc. Additionally, epidemiological and statistical studies to track down the number of hospital admissions caused by respiratory issues before and after the arrival of African dust is urgently needed in the Yucatan Peninsula. This will allow policy-makers and local authorities to understand how strong the African dust impact is on local health and the need for better forecasting of such events.

420

*Data availability.* Data are available upon request to the corresponding author.

425

*Author contributions.* CRR, GBR, and LAL designed the field campaigns and the experiments. CRR, MFC, HAO, DR, TA, and LAL carried out the aerosol measurements. CRR and AJ analyzed the remote sensing data. JM and HAO performed the chemical analyses. GBR, DB, DR, JSK, JYH, and LAL installed the equipment and provided the infrastructure for the ADABBOY project. CRR, AJ, and LAL wrote the paper, with contributions from all coauthors.

430

*Competing interest.* The authors declare that they have no conflict of interest.

*Acknowledgments.* This study was financially supported by the Consejo Nacional de Ciencia y Tecnologia (Conacyt) and the Universidad Autónoma de Yucatán through the FC-2164 and SISPROY-FQUI-2018-0003 grants, respectively. The authors thank the University Network of Atmospheric Observatories (RUOA) for providing meteorological and criteria pollution data. A. Jaramillo acknowledges the fellowship from DGAPA at UNAM. Also, the authors express their gratitude to Elizabeth Garcia, Juan Carlos Pineda, Aline Cruz, and Javier Juarez for their invaluable help and support.

435

## 440 **References**

Achilleos, S., Evans, J. S., Yiallourou, P. K., Kleanthous, S., Schwartz, J., and Koutrakis, P.: [PM<sub>10</sub> concentration levels at an urban and background site in Cyprus: The impact of urban sources and dust storms](https://doi.org/10.1080/10962247.2014.923061), *J. Air. Waste. Manage.*, 64, 1352–1360, <https://doi.org/10.1080/10962247.2014.923061>, 2014.

445

Adams, A. A., Prospero, J. M., and Zhang, C.: CALIPSO- Derived Three-Dimensional Structure of Aerosol over the Atlantic Basin and Adjacent Continents, *J. Climate*, 25, 6862–6879, <https://doi.org/10.1175/JCLI-D-11-00672.1>, 2012.

Aldhaif, A. M., Lopez, D. H., Dadashazar, H., and Sorooshian, A.: Sources, frequency, and chemical nature of dust events impacting the United States East Coast, *Atmos. Environ.*, 117456, <https://doi.org/10.1016/j.atmosenv.2020.117456>, 2020.

450

- Alessandrini, E. R., Stafoggia, M., Faustini, A., Gobbi, G. P., and Forastiere, F.: Saharan dust and the association between particulate matter and daily hospitalizations in Rome, Italy. *J. Occup. Environ. Med.*, 70, 432-434, <http://dx.doi.org/10.1136/oemed-2012-101182>, 2013.
- 455 Alvarez-Ospina, H., Giordano, S., Ladino, L.A., Raga, G.B., Muñoz-Salazar, J., Leyte-Lugo, M., Rosas D., and Carabali G. Particle-bound polycyclic aromatic hydrocarbons (pPAHs) in Merida, Mexico. Under review, 2020.
- AMS: Glossary of Meteorology (T. S. Glickman, Ed.), American Meteorological Society, <https://books.google.com.mx/books?id=kyhSAQAIAAJ>, 2000.
- Ashrafi, K., Shafiepour-Motlagh, M., Aslemand, A., and Ghader, S.: Dust storm simulation over Iran using HYSPLIT, *J. Environ. Health Sci. Eng.*, 12, 1-9, <https://doi.org/10.1186/2052-336X-12-9>, 2014.
- 460 Athanasopoulou, E., Protonotariou, A., Papangelis, G., Tombrou, M., Mihalopoulos, N., and Gerasopoulos, E.: Long-range transport of Saharan dust and chemical transformations over the Eastern Mediterranean, *Atmos. Environ.*, 140, 592–604, <https://doi.org/10.1016/j.atmosenv.2016.06.041>, 2016.
- 465 [Barkley, A. E., Prospero, J. M., Mahowald, N., Hamilton, D. S., Pependorf, K. J., Oehlert, A. M., Pourmand, A., Gatineau, A., Panechou-Pulcherie, K., Blackwelder, P., and Gaston, C. J.: African biomass burning is a substantial source of phosphorus deposition to the Amazon, Tropical Atlantic Ocean, and Southern Ocean, \*P. Natl. Acad. Sci.\*, 116, 16216–16221, <https://doi.org/10.1073/pnas.1906091116>, 2019.](#)
- Bibi, M., Saad, M., Masmoudi, M., Laurent, B., and Alfaro, S. C.: Long-term (1980–2018) spatial and temporal variability of the atmospheric dust load and deposition fluxes along the North-African coast of the Mediterranean Sea, *Atmos. Res.*, 234, 104689, <https://doi.org/10.1016/j.atmosres.2019.104689>, 2020.
- 470 Bègue, N., Tulet, P., Chaboureau, J.-P., Roberts, G., Gomes, L., and Mallet, M.: Long-range transport of Saharan dust over northwestern Europe during EUCAARI 2008 campaign: Evolution of dust optical properties by scavenging, *J. Geophys. Res.*, 117, D17201, <https://doi.org/10.1029/2012JD017611>, 2012.
- Bravo J.L., Salazar S., and Muhlia A.: Mineral and sea salt aerosol concentrations over low latitude tropical Atlantic and Pacific oceans during FGGE. *Geo. Int.*, 20, 303-317, 1982.
- 475 Broadley, S. L., Murray, B. J., Herbert, R. J., Atkinson, J. D., Dobbie, S., Malkin, T. L., Condliffe, E., and Neve, L.: Immersion mode heterogeneous ice nucleation by an illite rich powder representative of atmospheric mineral dust, *Atmos. Chem. Phys.*, 12, 287–307, <https://doi.org/10.5194/acp-12-287-2012>, 2012.
- 480 Brook, R. D., Rajagopalan, S., Pope, C. A., Brook, J. R., Bhatnagar, A., Diez-Roux, A. V., Holguin, F., Hong, Y., Luepker, R. V., Mittleman, M. A., Peters, A., Siscovick, D., Smith, S. C., Whitsel, L., Kaufman, J. D., and American Heart Association Council on Epidemiology and Prevention, Council on the Kidney in Cardiovascular Disease, and Council on Nutrition, Physical Activity and Metabolism: Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association, *Circulation*, 121, 2331–2378, <https://doi.org/10.1161/CIR.0b013e3181dbee1>, 2010.
- 485 Buchard, V., da Silva, A. M., Colarco, P. R., Darmenov, A., Randles, C. A., Govindaraju, R., Torres, O., Campbell, J., and Spurr, R.: Using the OMI aerosol index and absorption aerosol optical depth to evaluate the NASA MERRA Aerosol Reanalysis, *Atmos. Chem. Phys.*, 15, 5743–5760, <https://doi.org/10.5194/acp-15-5743-2015>, 2015.
- 490 Buchard, V., da Silva, A. M., Randles, C. A., Colarco, P., Ferrare, R., Hair, J., Hostetler, C., Tackett, J., and Winker, D.: Evaluation of the surface PM<sub>2.5</sub> in Version 1 of the NASA MERRA Aerosol Reanalysis over the United States, *Atmos. Environ.*, 125, 100–111, <https://doi.org/10.1016/j.atmosenv.2015.11.004>, 2016.
- Buchard, V., Randles, C. A., Silva, A. M. da, Darmenov, A., Colarco, P. R., Govindaraju, R., Ferrare, R., Hair, J., Beyersdorf, A. J., Ziemba, L. D., and Yu, H.: The MERRA-2 Aerosol Reanalysis, 1980 Onward. Part II: Evaluation and Case Studies, *J. Climate*, 30, 6851–6872, <https://doi.org/10.1175/jcli-d-16-0613.1>, 2017.

- 495 Caquineau, S., Gaudichet, A., Gomes, L., Magonthier, M. C., and Chatenet, B.: Saharan dust: Clay ratio as a relevant tracer to assess the origin of soil-derived aerosols, *Geophys. Res. Lett.*, 25, 983–986, <https://doi.org/10.1029/98GL00569>, 1998.
- Carlson, T. N., and Prospero, J. M.: The Large-Scale Movement of Saharan Air Outbreaks over the Northern Equatorial Atlantic, *J. Appl. Meteorol.*, 11, 283–297, [https://doi.org/10.1175/1520-0450\(1972\)011<0283:TLSMOS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1972)011<0283:TLSMOS>2.0.CO;2), 1972.
- 500 Cerón, R. M. B., Padilla, H. G., Belmont, R. D., Torres, M. C. B., Garcíá, R. M., and Báez, A. P.: Rainwater chemical composition at the end of the mid-summer drought in the Caribbean shore of the Yucatan Peninsula, *Atmos. Environ.*, 36, 2367–2374, [https://doi.org/10.1016/S1352-2310\(02\)00169-3](https://doi.org/10.1016/S1352-2310(02)00169-3), 2002.
- 505 Cerón-Bretón, R., Cerón-Bretón, J., Muriel-García, M., Lara-Severino, R., Rangel-Marrón, M., Ramírez-Lara, E., López-Jiménez, D., Rodríguez-Guzmán, A., and Uc-Chi, M.: Mapping of the atmospheric deposition of sulfur and nitrogen during the dry season 2016 in the Metropolitan zone of Merida, Yucatan, Mexico. *AIP. Conf. Proc.* 1982, 020021, <https://doi.org/10.1063/1.5045427>, 2018.
- Chiapello, I., Bergametti, G., Chatenet, B., Bousquet, P., Dulac, F., and Soares, E.S.: Origins of African dust transported over the northeastern tropical Atlantic, *J. Geophys. Res.*, 102, 13701– 13709, <https://doi.org/10.1029/97JD00259>, 1997.
- 510 Chiapello, I., Prospero, J. M., Herman, J. R., and Hsu, N. C.: Detection of mineral dust over the North Atlantic Ocean and Africa with the Nimbus 7 TOMS, *J. Geophys. Res.-Atmos.*, 104, 9277–9291. <https://doi.org/10.1029/1998JD200083>, 1999.
- 515 Chouza, F., Reitebuch, O., Benedetti, A., and Weinzierl, B.: Saharan dust long-range transport across the Atlantic studied by an airborne Doppler wind lidar and the MACC model, *Atmos. Chem. Phys.*, 16, 11581–11600, <https://doi.org/10.5194/acp-16-11581-2016>, 2016.
- Cohn, S. E.: An Introduction to Estimation Theory (gtSpecial Issue>Data Assimilation in Meteorology and Oceanography: Theory and Practice), *J. Meteorol. Soc. Jpn. Ser. II*, 75, 257–288, [https://doi.org/10.2151/jmsj1965.75.1B\\_257](https://doi.org/10.2151/jmsj1965.75.1B_257), 1997.
- 520 Corbett, J. J., and Fischbeck, P.: Emissions from Ships, *Science*, 278, 823–824. <https://doi.org/10.1126/science.278.5339.823>, 1997.
- 525 [Córdoba, F., Ramirez-Romero, C., Cabrera, D., Raga, G. B., Miranda, J., Alvarez, H., Rosas, D., Figueroa, B., Kim, J. S., Yakobi-Hancock, J., Amador, T., Gutierrez, W., García, M., Bertram, A. K., Baumgardner, D., and Ladino, L.A.: Ice nucleating abilities of biomass burning, African dust, and sea spray aerosol particles over the Yucatan Peninsula. \*Atmospheric Chemistry and Physics Discussions\*. <https://doi.org/10.5194/acp-2020-783>, 2020.](https://doi.org/10.5194/acp-2020-783)
- d’Almeida, G. A.: A Model for Saharan Dust Transport, *J. Climate Appl. Meteor.*, 25, 903–916. [https://doi.org/10.1175/1520-0450\(1986\)025<0903:AMFSDT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1986)025<0903:AMFSDT>2.0.CO;2), 1986.
- 530 DeMott, P. J., Prenni, A. J., McMeeking, G. R., Sullivan, R. C., Petters, M. D., Tobo, Y., Niemand, M., Möhler, O., Snider, J. R., Wang, Z., and Kreidenweis, S. M.: Integrating laboratory and field data to quantify the immersion freezing ice nucleation activity of mineral dust particles, *Atmos. Chem. Phys.*, 15, 393–409, <https://doi.org/10.5194/acp-15-393-2015>, 2015.
- 535 Denjean, C., Formenti, P., Desboeufs, K., Chevaillier, S., Triquet, S., Maillé, M., Cazaunau, M., Laurent, B., Mayol-Bracero, O. L., Vallejo, P., Quiñones, M., Gutierrez-Molina I. E., Cassola, F., Prati, P., Andrews, E., and Ogren, J.: Size distribution and optical properties of African mineral dust after intercontinental transport, *J. Geophys. Res. Atmos.*, 121, 7117–7138, <https://doi.org/10.1002/2016JD024783>, 2016.

- Dominici, F., Peng, R. D., Bell, M. L., Pham, L., McDermott, A., Zeger, S. L., and Samet, J. M.: Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases, *JAMA*, 295, 1127–1134, <https://doi.org/10.1001/jama.295.10.1127>, 2006.
- 540 Dunion, J. P.: Rewriting the climatology of the tropical North Atlantic and Caribbean Sea atmosphere. *J. Climate*, 24, 893–908, <https://doi.org/10.1175/2010JCLI3496.1>, 2011.
- Dunion, J. P., and Marron, C. S.: A Reexamination of the Jordan Mean Tropical Sounding Based on Awareness of the Saharan Air Layer: Results from 2002. *J. Climate*, 21, 5242–5253, <https://doi.org/10.1175/2008JCLI1868.1>, 2008.
- 545 Dunion, J. P., and Velden, C. S.: The Impact of the Saharan Air Layer on Atlantic Tropical Cyclone Activity, *Bull. Amer. Meteor. Soc.*, 85, 353–366, <https://doi.org/10.1175/BAMS-85-3-353>, 2004.
- Espinosa, A., Miranda, J., and Pineda, J.: Uncertainty evaluation in correlated quantities: application to elemental analysis of atmospheric aerosols, *Rev. Mex. Fis.*, E56, 134–140, 2010.
- Espinosa, A., Reyes-Herrera, J., Miranda, J., Mercado, F., Veytia, M., Cuaute, M., and Cruz, J.: Development of an X-ray fluorescence spectrometer for environmental science applications, *Instrum. Sci. Technol.*, 40, 603–617, <https://doi.org/10.1080/10739149.2012.693560>, 2012.
- 550 Evan, A. T., Dunion, J., Foley, J. A., Heidinger, A. K., and Velden, C. S.: New evidence for a relationship between Atlantic tropical cyclone activity and African dust outbreaks, *Geophys. Res. Lett.*, 33, L19813, <https://doi.org/10.1029/2006GL026408>, 2006.
- Foltz, G. R., and McPhaden, M. J.: Trends in Saharan dust and tropical Atlantic climate during 1980–2006, *Geophys. Res. Lett.*, 35, L20706, <https://doi.org/10.1029/2008GL035042>, 2008.
- 555 Ganor, E., and Mamane, Y.: Transport of Saharan dust across the eastern Mediterranean. *Atmos. Environ.*, 16, 581–587, [https://doi.org/10.1016/0004-6981\(82\)90167-6](https://doi.org/10.1016/0004-6981(82)90167-6), 1982.
- Ganor, E., Osetinsky, I., Stupp, A., and Alpert, P.: Increasing trend of African dust, over 49 years, in the eastern Mediterranean, *J. Geophys. Res.*, 115, D07201, <https://doi.org/10.1029/2009JD012500>, 2010.
- 560 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 565 GMAO: MERRA-2 inst1\_2d\_int\_Nx: 2d, 1-Hourly, Instantaneous, Single-Level, Assimilation, Vertically Integrated Diagnostics V5.12.4. Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC). <https://doi.org/10.5067/LTVB4GPCOTK2>, 2015a.
- 570 GMAO: MERRA-2 inst3\_3d\_aer\_Nv: 3d, 3-Hourly, Instantaneous, Model-Level, Assimilation, Aerosol Mixing Ratio V5.12.4. Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC). <https://doi.org/10.5067/LTVB4GPCOTK2>, 2015b.
- Goudie, A., and Middleton, N. J.: *Desert Dust in the Global System*, Springer-Verlag, <https://doi.org/10.1007/3-540-32355-4>, 2006.
- 575 Goudie, A. S. Desert dust and human health disorders. *Environ. Int.*, 63, 101–113. <https://doi.org/10.1016/j.envint.2013.10.011>, 2014.
- Griffin, D.V., Garrison, V.H., Herman, J.R., and Shinn, E.: African desert dust in the Caribbean atmosphere: Microbiology and public health, *Aerobiologia*, 17, 203–213, <https://doi.org/10.1023/A:1011868218901>, 2001.

- Grogan, D. F. P., and Thorncroft, C. D.: The characteristics of African easterly waves coupled to Saharan mineral dust aerosols, *Q. J. R. Meteorol. Soc.*, 145, 1130–1146, <https://doi.org/10.1002/qj.3483>, 2019.
- 580 Guieu, C., Loÿe-Pilot, M. D., Ridame, C., and Thomas, C.: Chemical characterization of the Saharan dust end-member: Some biogeochemical implications for the western Mediterranean Sea, *J. Geophys. Res.*, 107, ACH 5-1-ACH 5-11, <https://doi.org/10.1029/2001JD000582>, 2002.
- Hankes, I., and Marinaro, A.: The impacts of column water vapour variability on Atlantic basin tropical cyclone activity, *Q. J. R. Meteorol. Soc.*, 142, 3026–3035. <https://doi.org/10.1002/qj.2886>, 2016.
- 585 Holloway, C. E., and Neelin, J. D.: Moisture Vertical Structure, Column Water Vapor, and Tropical Deep Convection. *J. Atmos. Sci.*, 66, 1665–1683, <https://doi.org/10.1175/2008JAS2806.1>, 2009.
- Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 9817–9854, <https://doi.org/10.5194/acp-12-9817-2012>, 2012.
- IAEA: Manual for QXAS, International Atomic Energy Agency, Vienna, 1997.
- 590 (INEGI: Instituto Nacional de Estadística y Geografía (México). Anuario estadístico de Yucatán 2009 / Instituto Nacional de Estadística y Geografía, Gobierno del Estado de Yucatán. México, 2009.
- INEGI: Principales resultados de la Encuesta intercensal 2015: Estados Unidos Mexicanos, 2015.
- INEGI: Anuario estadístico y geográfico de Yucatán 2017, 1–711, [https://www.datatur.sectur.gob.mx/ITxEF\\_Docs/YUC\\_ANUARIO\\_PDF.pdf](https://www.datatur.sectur.gob.mx/ITxEF_Docs/YUC_ANUARIO_PDF.pdf), 2017.
- 595 Jackson, J. M., Liu, H., Laszlo, I., Kondragunta, S., Remer, L. A., Huang, J., and Huang, H. C.: Suomi-NPP VIIRS aerosol algorithms and data products, *J. Geophys. Res. Atmos.*, 118, 12673–12689, <https://doi.org/10.1002/2013JD020449>, 2013.
- Kalashnikova, O. V., and Kahn, R. A.: Mineral dust plume evolution over the Atlantic from MISR and MODIS aerosol retrievals, *J. Geophys. Res.*, 113, D24204, <https://doi.org/10.1029/2008JD010083>, 2008.
- 600 Kalnay, E.: Atmospheric modeling, data assimilation and predictability, Cambridge University Press, 2003.
- Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Krämer, M.: Overview of ice nucleating particles, *Meteorol. Monogr.*, 58, 1.1–1.33, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0006.1>, 2017.
- Karanasiou, A., Moreno, N., Moreno, T., Viana, M., de Leeuw, F., and Querol, X.: Health effects from Sahara dust episodes in Europe: Literature review and research gaps, *Environ. Int.*, 47, 107–114, <https://doi.org/10.1016/j.envint.2012.06.012>, 2012.
- Karyampudi, V. M., and Carlson, T. N. Analysis and Numerical Simulations of the Saharan Air Layer and Its Effect on Easterly Wave Disturbances, *J. Atmos. Sci.*, 45, 3102–3136. [https://doi.org/10.1175/1520-0469\(1988\)045<3102:AANSOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<3102:AANSOT>2.0.CO;2), 1988.
- 610 Kaufman, Y. J., Koren, I., Remer, L. A., Tanré, D., Ginoux, P., and Fan, S.: Dust transport and deposition observed from the Terra-Moderate Resolution Imaging Spectroradiometer (MODIS) spacecraft over the Atlantic Ocean, *J. Geophys. Res.*, 110, D10S12, <https://doi.org/10.1029/2003JD004436>, 2005.
- Kloster, S., Feichter, J., Maier-Reimer, E., Six, K. D., Stier, P., and Wetzell, P.: DMS cycle in the marine ocean-atmosphere system – a global model study, *Biogeosciences*, 3, 29–51, <https://doi.org/10.5194/bg-3-29-2006>, 2006.
- 615



- Koren, I., Kaufman, Y. J., Washington, R., Todd, M. C., Rudich, Y., Martins, J. V., and Rosenfeld, D.: The Bodélé depression: A single spot in the Sahara that provides most of the mineral dust to the Amazon forest. *Environ. Res. Lett.*, 1, 014005. <https://doi.org/10.1088/1748-9326/1/1/014005>, 2006.
- 620 Korte, L. F., Pausch, F., Trimborn, S., Brussaard, C. P. D., Brummer, G.-J. A., van der Does, M., Guerreiro, C. V., Schreuder, L. T., Munday, C. I., and Stuut, J.-B. W.: Effects of dry and wet Saharan dust deposition in the tropical North Atlantic Ocean, *Biogeosciences Discuss.*, <https://doi.org/10.5194/bg-2018-484>, 2018.
- Kramer, S. J., Kirtman, B. P., Zuidema, P., and Ngan, F.: Subseasonal variability of elevated dust concentrations over South Florida. *J. Geophys. Res. Atmos.*, 125, e2019JD031874. <https://doi.org/10.1029/2019JD031874>, 2020.
- 625 Lenes, J., Prospero, J., Landing, W., Virmani, J., and Walsh, J.: A model of Saharan dust deposition to the eastern Gulf of Mexico, *Mar. Chem.*, 134-135, 1–9, <https://doi.org/10.1016/j.marchem.2012.02.007>, 2012.
- Li- Jones, X., and Prospero, J. M.: Variations in the size distribution of non-sea-salt sulfate aerosol in the marine boundary layer at Barbados: Impact of African dust, *J. Geophys. Res. Atmos.*, 103, 16073–16084. <https://doi.org/10.1029/98JD00883>, 1998.
- 630 Linke, C., Möhler, O., Veres, A., Mohácsi, Á., Bozóki, Z., Szabó, G., and Schnaiter, M.: Optical properties and mineralogical composition of different Saharan mineral dust samples: a laboratory study, *Atmos. Chem. Phys.*, 6, 3315–3323, <https://doi.org/10.5194/acp-6-3315-2006>, 2006.
- [Liu, Z., Omar, A., Vaughan, M., Hair, J., Kittaka, C., Hu, Y., Powell, K., Trepte, C., Winker, D., Hostetler, C., Ferrare, R., and Pierce, R.: CALIPSO lidar observations of the optical properties of Saharan dust: A case study of long range transport, \*J. Geophys. Res.\*, 113, D07207, <https://doi.org/10.1029/2007JD008878>, 2008.](#)
- 635 [Middleton, N. J., and Goudie, A. S.: Saharan dust: Sources and trajectories, \*Trans. Inst. Br. Geogr.\*, 26, 165–181, <https://doi.org/10.1111/1475-5661.00013>, 2001.](#)
- Muñoz-Salazar J., Raga, G.B., Yakobi-Hancock, J., Kim, J.S., Rosas, D., Caudillo, L., Alvarez-Ospina, H., and Ladino, L.A.: Ultrafine Aerosol Particles in the Western Caribbean: A first case study in Merida, Under review, 2020.
- 640 [Murray, B., O'sullivan, D., Atkinson, J., and Webb, M.: Ice nucleation by particles immersed in supercooled cloud droplets, \*Chem. Soc. Rev.\*, 41, 6519–6554, <https://doi.org/10.1039/C2CS35200A>, 2012.](#)
- Nenes, A., Murray, B., and Bougiatioti, A.: Mineral dust and its microphysical interactions with clouds, in: *Mineral Dust*, 287–325, Springer, 2014.
- 645 Orellana, R., Espadas, C., Conde, C., and Gay, C.: Atlas escenarios de cambio climático en la Península de Yucatán. Mérida, Unidad de Recursos Naturales, Centro de Investigación Científica de Yucatán y Centro de Ciencias de la Atmósfera-UNAM, Mérida, Yucatán, México, 2009.
- Perez, L., Tobias, A., Querol, X., Künzli, N., Pey, J., Alastuey, A., Viana, M., Valero, N., González-Cabré, M., and Sunyer, J.: Coarse Particles From Saharan Dust and Daily Mortality, *Epidemiology*, 19, 800–807, <https://doi.org/10.1097/EDE.0b013e31818131cf>, 2008.
- 650 [Perry, K.D., Cahill, T.A., Eldred, R.A., Dutcher, D.D., Gill, T.E.: Long-range transport of North African dust to the eastern United States, \*J. Geophys. Res.\*, 102 \(D10\), 11225-11238, <https://doi.org/10.1029/97JD00260>, 1997.](#)
- 655 [Pey, J., Querol, X., Alastuey, A., Forastiere, F., and Stafoggia, M.: African dust outbreaks over the Mediterranean Basin during 2001–2011: PM<sub>10</sub> concentrations, phenomenology and trends, and its relation with](#)

synoptic and mesoscale meteorology, *Atmos. Chem. Phys.*, 13, 1395–1410, <https://doi.org/10.5194/acp-13-1395-2013>, 2013.

Plasencia, A. P.: La Península de Yucatán en el Archivo General de la Nación, UNAM, 1998.

ProAire.: Programa de Gestión para mejorar la calidad del aire del estado de Yucatán, 1–159, 2018.

660 Prodi, F., and Fea, G.: A case of transport and deposition of Saharan dust over the Italian Peninsula and southern Europe. *J. Geophys. Res. Oceans*, 84, 6951–6960. <https://doi.org/10.1029/JC084iC11p06951>, 1979.

Prospero, J. M., and Carlson, T. N.: Vertical and areal distribution of Saharan dust over the western equatorial north Atlantic Ocean, *J. Geophys. Res.*, 77, 5255–5265, <https://doi.org/10.1029/JC077i027p05255>, 1972.

665 [Prospero, J. M., Glaccum, R. A. and Nees, R. T.: Atmospheric transport of soil dust from Africa to South America, \*Nature\*, 289, 570-572, <https://doi.org/10.1038/289570a0>, 1981.](#)

Prospero, J. M.: Long-term measurements of the transport of African mineral dust to the southeastern United States: Implications for regional air quality, *J. Geophys. Res.*, 104, 15917–15927, <https://doi.org/10.1029/1999JD900072>, 1999.

670 Prospero, J. M., Olmez, I., and Ames, M.: Al and Fe in PM<sub>2.5</sub> and PM<sub>10</sub> Suspended Particles in South-Central Florida: The Impact of the Long Range Transport of African Mineral Dust, *Water Air Soil Poll.*, 125, 291–317, <https://doi.org/10.1023/A:1005277214288>, 2001.

675 [Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., and Gill, T. E.: Environmental characterization of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer \(TOMS\) absorbing aerosol product, \*Reviews of Geophysics\*, 40, 1-31, <https://doi.org/10.1029/2000RG000095>, 2002.](#)

Prospero, J. M., and Lamb, P. J.: African Droughts and Dust Transport to the Caribbean: Climate Change Implications, *Science*, 302, 1024–1027, <https://doi.org/10.1126/science.1089915>, 2003.

680 Prospero, J. M., Blades, E., Mathison, G., and Naidu, R.: Interhemispheric transport of viable fungi and bacteria from Africa to the Caribbean with soil dust, *Aerobiologia*, 21, 1–19, <https://doi.org/10.1007/s10453-004-5872-7>, 2005.

Prospero, J. M., and Mayol-Bracero, O. L.: Understanding the Transport and Impact of African Dust on the Caribbean Basin, *Bull. Amer. Meteor. Soc.*, 94, 1329–1337, <https://doi.org/10.1175/BAMS-D-12-00142.1>, 2013.

685 Prospero, J. M., Collard, F.-X., Molinié, J., and Jeannot, A.: Characterizing the annual cycle of African dust transport to the Caribbean Basin and South America and its impact on the environment and air quality, *Global Biogeochem. Cy.*, 29, 757–773, <https://doi.org/10.1002/2013GB004802>, 2014.

690 [Prospero, J. M., Barkley, A. N., Gaston, C. J., Gatineau, A., Campos y Sansano, A., and Panechou, K.: Characterizing and Quantifying African Dust Transport and Deposition to South America: Implications for the Phosphorus Budget in the Amazon Basin, \*Global Biogeochem. Cy.\*, 34, e2020GB006536, <https://doi.org/10.1029/2020GB006536>, 2020.](#)

Provençal, S., Buchard, V., da Silva, A. M., Leduc, R., and Barrette, N.: Evaluation of PM surface concentrations simulated by Version 1 of NASA's MERRA Aerosol Reanalysis over Europe, *Atmos. Pollut. Res.*, 8, 374–382, <https://doi.org/10.1016/j.apr.2016.10.009>, 2017.

695 [Querol, X., Pey, J., Pandolfi, M., Alastuey, A., Cusack, M., Moreno, T., Viana, M., Mihalopoulos, N., Kallos, G., and Kleanthous, S.: African dust contributions to mean ambient PM<sub>10</sub> levels across the Mediterranean Basin, \*Atmos. Environ.\*, 43, 4266–4277, <https://doi.org/10.1016/j.atmosenv.2009.06.013>, 2009.](#)

- Querol, X., Tobías, A., Pérez, N., Karanasiou, A., Amato, F., Stafoggia, M., Pérez García-Pando, C., Ginoux, P., Forastiere, F., Gumy, S., Mudu, P., and Alastuey, A.: Monitoring the impact of desert dust outbreaks for air quality for health studies, *Environ. Int.*, 130, 104867. <https://doi.org/10.1016/j.envint.2019.05.061>, 2019.
- 700 Randles, C. A., da Silva, A. M., Buchard, V., Colarco, P. R., Darmenov, A., Govindaraju, R., Smirnov, A., Holben, B., Ferrare, R., Hair, J., Shinozuka, Y., and Flynn, C. J.: The MERRA-2 Aerosol Reanalysis, 1980 Onward. Part I: System Description and Data Assimilation Evaluation, *J. Climate*, 30, 6823–6850. <https://doi.org/10.1175/jcli-d-16-0609.1>, 2017.
- 705 Reid, J. S., Jonsson, H. H., Maring, H. B., Smirnov, A., Savoie, D. L., Cliff, S. S., Reid, E. A., Livingston, J. M., Meier, M. M., Dubovik, O., and Tsay, S. C.: Comparison of size and morphological measurements of coarse mode dust particles from Africa, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2002JD002485>, 2003a.
- Reid, E. A., Reid, J. S., Meier, M. M., Dunlap, M. R., Cliff, S. S., Broumas, A., Perry, K., and Maring, H.: Characterization of African dust transported to Puerto Rico by individual particle and size segregated bulk analysis, *J. Geophys. Res.*, 108, 8591, <https://doi.org/10.1029/2002JD002935>, 2003b.
- 710 Resch, F., Sunnu, A., and Afeti, G.: Saharan dust flux and deposition rate near the Gulf of Guinea, *Tellus B*, 60, 98–105. <https://doi.org/10.1111/j.1600-0889.2007.00286.x>, 2008.
- Rienecker, M.M., Suarez, M.J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M.G., Schubert, S.D., Takacs, L., Kim, G-K, Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R.D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C.R., Reichle, R., Robertson, F.R., 715 Ruddick, A.G., Sienkiewicz, M., and Woollen, J.: NASA’s Modern-Era Retrospective Analysis for Research and Applications. *J. Climate*, 24, 3624–3648. <https://doi.org/10.1175/JCLI-D-11-00015.1>, 2011.
- ~~Rodríguez Cotto, R. I., Ortiz Martínez, M. G., Rivera Ramírez, E., Méndez, L. B., Dávila, J. C., and Jiménez-Vélez, B. D.: African Dust Storms Reaching Puerto Rican Coast Stimulate the Secretion of IL-6 and IL-8 and Cause Cytotoxicity to Human Bronchial Epithelial Cells (BEAS-2B). *Health*, 5, 14–28; <https://doi.org/10.4236/health.2013.510A2003>, 2013.~~
- 720 Rodríguez-Gomez, C., Ramirez-Romero, C., Cordoba, F., Raga, G. B., Salinas, E., Martinez, L., Rosas, I., Quintana, E. T., Maldonado, L. A., Rosas, D., Amador, T., Alvarez, H., and Ladino, L. A.: Characterization of culturable airborne microorganisms in the Yucatan Peninsula, *Atmos. Environ.*, 223, 117183, <https://doi.org/10.1016/j.atmosenv.2019.117183>, 2020.
- 725 Rosinski, J., Haagenson, P., Nagamoto, C., Quintana, B., Parungo, F., and Hoyt, S.: Ice-forming nuclei in air masses over the Gulf of Mexico, *J. Aerosol Sci.*, 19, 539–551, [https://doi.org/10.1016/0021-8502\(88\)90206-6](https://doi.org/10.1016/0021-8502(88)90206-6), 1988.
- Ryder, C. L., Highwood, E. J., Walser, A., Seibert, P., Philipp, A., and Weinzierl, B.: Coarse and giant particles are ubiquitous in Saharan dust export regions and are radiatively significant over the Sahara, *Atmos. Chem. Phys.*, 19, 15353–15376, <https://doi.org/10.5194/acp-19-15353-2019>, 2019.
- 730 Salvador, P., Alonso-Pérez, S., Pey, J., Artíñano, B., de Bustos, J. J., Alastuey, A., and Querol, X.: African dust outbreaks over the western Mediterranean Basin: 11-year characterization of atmospheric circulation patterns and dust source areas, *Atmos. Chem. Phys.*, 14, 6759–6775, <https://doi.org/10.5194/acp-14-6759-2014>, 2014.
- Schutgens, N. A. J., Miyoshi, T., Takemura, T., and Nakajima, T.: Applying an ensemble Kalman filter to the assimilation of AERONET observations in a global aerosol transport model, *Atmos. Chem. Phys.*, 10, 2561–2576, <https://doi.org/10.5194/acp-10-2561-2010>, 2010.
- 735 Shao, Y., Wyrwoll, K.-H., Chappell, A., Huang, J., Lin, Z., McTainsh, G. H., Mikami, M., Tanaka, T. Y., Wang, X., and Yoon, S.: Dust cycle: An emerging core theme in Earth system science, *Aeolian Res.*, 2, 181–204, <https://doi.org/10.1016/j.aeolia.2011.02.001>, 2011.


- 740 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bull. Am. Meteorol. Soc.*, 96, 2059–2077. <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- Tegen, I., Werner, M., Harrison, S. P., and Kohfeld, K. E.: Relative importance of climate and land use in determining present and future global soil dust emission, *Geophys. Res. Lett.*, 31, L05105, <https://doi.org/10.1029/2003GL019216>, 2004.
- Thermo Fisher Scientific Inc.: Instruction Manual for Thermo Scientific Model FH62C14 Ambient Particulate Monitor. <https://assets.thermofisher.com/TFS-Assets/null%7Cnull/Package-Inserts/EPM-manual-FH62C14.pdf>, 2007.
- 750 Trapp, J. M., Millero, F. J., and Prospero, J. M.: Temporal variability of the elemental composition of African dust measured in trade wind aerosols at Barbados and Miami. *Mar. Chem.*, 120, 71–82, <https://doi.org/10.1016/j.marchem.2008.10.004>, 2010.
- Tsamalis, C., Chédin, A., Pelon, J., and Capelle, V.: The seasonal vertical distribution of the Saharan Air Layer and its modulation by the wind, *Atmos. Chem. Phys.*, 13, 11235–11257, <https://doi.org/10.5194/acp-13-11235-2013>, 2013.
- 755 Vallina, S. M., and Simó, R.: Strong Relationship Between DMS and the Solar Radiation Dose over the Global Surface Ocean. *Science*, 315, 506–508, <https://doi.org/10.1126/science.1133680>, 2007.
- Venero-Fernández, S. J.: Saharan Dust Effects on Human Health: A Challenge for Cuba's Researchers. *MEDICC Rev.*, 18, 32–34, <https://doi.org/10.1590/MEDICC.2016.18300011>, 2016.
- 760 Veselovskii, I., Goloub, P., Podvin, T., Tanre, D., da Silva, A., Colarco, P., Castellanos, P., Korenskiy, M., Hu, Q., Whiteman, D. N., Pérez-Ramírez, D., Augustin, P., Fourmentin, M., and Kolgotin, A.: Characterization of smoke and dust episode over West Africa: comparison of MERRA-2 modeling with multiwavelength Mie–Raman lidar observations, *Atmos. Meas. Tech.*, 11, 949–969, <https://doi.org/10.5194/amt-11-949-2018>, 2018.
- Voss, K. K., and Evan, A. T.: A New Satellite-Based Global Climatology of Dust Aerosol Optical Depth. *J. Appl. Meteorol. Clim.*, 59, 83–102. <https://doi.org/10.1175/JAMC-D-19-0194.1>, 2020.
- 765 Walker, G. P. L.: Generation and dispersal of fine ash and dust by volcanic eruptions. *J. Volcanol. Geotherm. Res.*, 11, 81–92. [https://doi.org/10.1016/0377-0273\(81\)90077-9](https://doi.org/10.1016/0377-0273(81)90077-9), 1981.
- Weinzierl, B., Ansmann, A., Prospero, J. M., Althausen, D., Benker, N., Chouza, F., Dollner, M., Farrell, D., Fomba, W. K., Freudenthaler, V., Gasteiger, J., Groß, S., Haarig, M., Heinold, B., Kandler, K., Kristensen, T. B., Mayol-Bracero, O. L., Müller, T., Reitebuch, O., Sauer, D., Schäfler, A., Schepanski, K., Spanu, A., Tegen, I., Toledano, C., and Walser, A.: The Saharan Aerosol Long-Range Transport and Aerosol–Cloud-Interaction Experiment: Overview and Selected Highlights. *Bull. Amer. Meteor. Soc.*, 98, 1427–1451. <https://doi.org/10.1175/BAMS-D-15-00142.1>, 2016.
- WHO: Quantifying Selected Major Risks to Health. In *The world health report 2002: Reducing risks, promoting healthy life*. World Health Organization, 2002.
- 775 Wilker, E. H., Preis, S. R., Beiser, A. S., Wolf, P. A., Au, R., Kloog, I., Li, W., Schwartz, J., Koutrakis, P., DeCarli, C., Seshadri, S., and Mittleman, M. A.: Long-term exposure to fine particulate matter, residential proximity to major roads and measures of brain structure, *Stroke*, 46, 1161–1166, <https://doi.org/10.1161/STROKEAHA.114.008348>, 2015.
- 780 World Resource Institute.: Inventario Municipal de Gases de Efecto Invernadero Municipio de Mérida, 1–29, [http://www.merida.gob.mx/municipio/sitiosphp/sustentable/contenidos/doc/Inventario\\_Municipal\\_GEL.pdf](http://www.merida.gob.mx/municipio/sitiosphp/sustentable/contenidos/doc/Inventario_Municipal_GEL.pdf), 2018.

Zhang, H., McFarquhar, G. M., Saleeby, S. M., and Cotton, W. R.: Impacts of Saharan dust as CCN on the evolution of an idealized tropical cyclone, *Geophys. Res. Lett.*, 34, L14812, <https://doi.org/10.1029/2007GL029876>, 2007.


785 Zhang, J., and Reid, J. S.: MODIS aerosol product analysis for data assimilation: Assessment of over- ocean level 2 aerosol optical thickness retrievals, *J. Geophys. Res.*, 111, D22207, <https://doi.org/10.1029/2005JD006898>, 2006.

Zhang, X., Zhao, L., Tong, D. Q., Wu, G., Dan, M., and Teng, B.: A systematic review of global desert dust and associated human health effects, *Atmosphere*, 7, 158, <https://doi.org/10.3390/atmos7120158>, 2016.


790



**Figure 1.** Location of the sampling site at the School of Chemistry of the Autonomous University of Yucatan (FC-UADY) and the three World Meteorological Organization (WMO) radiosonde stations located in the Yucatan Peninsula: Mérida International Airport, Mexico (WMO index: 76644); Cancun, Mexico (WMO index: 76595) and Philip S. W. Goldson International Airport, Belize (WMO index: 78583).




800 **Figure 2.** a) Mass concentrations of PM<sub>2.5</sub> and PM<sub>10</sub> in 2017, and b) in 2018. The green horizontal line depicts the World Health Organization Air Quality Guideline for 24 h mean PM<sub>2.5</sub> (Y axis right) and PM<sub>10</sub> (Y axis left) concentrations of 25  $\mu\text{g m}^{-3}$  and 50  $\mu\text{g m}^{-3}$ , respectively (WHO, 2002). The pie charts represent the elemental composition from the XRF for the ADPs.




805

**Figure 3.** MERRA-2 precipitable water (black solid line) and estimated from radiosonde measurements (blue dashed line), for the three WMO radiosonde stations located in the Yucatan Peninsula. a) July-August 2017, and b) July-August 2018. The red areas represent the periods where PWV is less than 45 mm.




810

**Figure 4.** 3-hour time series of the vertical profile of the estimated dust content from MERRA-2 for the July 1 to August 14 period for a) 2017 and b) 2018 for the Merida region.



**Figure 5.** Daily mean of  $PM_{10}$  estimated from MERRA-2 (black line), measured by the RUOA station (blue line), and the estimated dust mixing ratio content from MERRA-2 (red line) for a) 2017 and b) 2018.



**Figure 6.** Dispersion diagrams of surface dust mixing ratio from MERRA-2 (y-axis) vs. the  $PM_{10}$  from the RUOA station for the periods shown in Figure 5 for a) 2017 and b) 2018.

815



**Table 1.** Summary of the used instrumentation and the measured variables.

<b>Measured variable</b>	<b>Instrumentation</b>
Particle mass concentration	PM <sub>2.5</sub> and PM <sub>10</sub> analyzer (FH 62 C14 Thermo Scientific Inc.)
Total particle concentration (d>50 nm)	Condensation Particle Counter (CPC, 3010, TSI)
Particle size distribution (d>300 nm)	Optical Particle Counter (LasAir, II 310A, MSP)
Aerosol collection (PM <sub>10</sub> and PM <sub>2.5</sub> )	Partisol (2525, Thermo Fisher Scientific Inc) and MiniVol (3380, Air metrics)
Nitrogen oxides (NO <sub>x</sub> ) Ozone (O <sub>3</sub> )	NO <sub>x</sub> analyzer (42 i Thermo Scientific Inc.) O <sub>3</sub> analyzer (49 i Thermo Scientific Inc.)
Temperature, relative humidity precipitation Wind direction and wind speed  Solar radiation	T and HR Sensor ( VAISALA HMP 115) Pluviometer (Texas Electronics, TR-525M) Wind direction and wind speed sensor (Gill, 1405-PK-100) Radiation pyrometer (Intertek, 20W)
Absorption coefficient  Particle-bound polycyclic aromatic hydrocarbons (pPAHs) concentration	Soot Absorption Photometer PSAP (Radiance Research) Photoacoustic spectroscopy (PAS 2000, Ecochem)