Manuscript prepared for Atmos. Chem. Phys. with version 2014/09/16 7.15 Copernicus papers of the LATEX class copernicus.cls. Date: 25 August 2020

A Microphysics Guide to Cirrus – Part II: Climatologies of Clouds and Humidity from Observations

Martina Krämer^{1,2}, Christian Rolf¹, Nicole Spelten¹, Armin Afchine¹, David Fahey³, Eric Jensen⁵, Sergey Khaykin⁶, Thomas Kuhn⁷, Paul Lawson⁸, Alexey Lykov⁹, Laura L. Pan⁵, Martin Riese¹, Andrew Rollins³, Fred Stroh¹, Troy Thornberry^{3,4}, Veronika Wolf^{7,‡}, Sarah Woods⁸, Peter Spichtinger², Johannes Quaas¹⁰, and Odran Sourdeval¹¹

¹Institute for Energy and Climate Research (IEK-7), Research Center Jülich, Jülich, Germany ²Institute for Atmospheric Physics (IPA), Johannes Gutenberg University, Mainz, Germany ³NOAA ESRL CSD, Boulder, USA;

⁴CIRES, University of Colorado Boulder, USA;

⁵NCAR, Atmospheric Chemistry Observations and Modeling Laboratory, Boulder, USA; ⁶LATMOS/IPSL, UVSQ, Sorbonne Université, CNRS, Guyancourt, France

⁷Luleå University of Technology, Division of Space Technology, Kiruna, Sweden ⁸SPEC Inc., Boulder, CO, USA;

⁹Central Aerological Observatory (CAO), Department of Upper Atmospheric Layers Physics, Moscow, Russia

¹⁰Leipzig Institute for Meteorology (LIM), Universität Leipzig, Leipzig, Germany

¹¹Univ. Lille, CNRS, UMR 8518 - LOA - Laboratoire d'Optique Atmosphérique, F-59000 Lille, France

[‡]now at Brandenburg Univ. of Technology (BTU), Atmospheric Processes, Cottbus, Germany

Correspondence to: M. Krämer (m.kraemer@fz-juelich.de)

Abstract. This study presents airborne in situ and satellite remote sensing climatologies of cirrus clouds and humidity. The climatologies serve as a guide to the properties of cirrus clouds, with the new in situ database providing detailed insights into boreal mid-latitudes and the tropics, while the satellite-borne data set offers a global overview.

- 5 To this end, an extensive, quality checked data archive, the Cirrus Guide II in situ database, is created from airborne in situ measurements during 150 flights in 24 campaigns. The archive contains meteorological parameters, ice water content (IWC), ice crystal number concentration (N_{ice}), ice crystal mean mass radius (Rice), relative humidity with respect to ice (RHice) and water vapor mixing ratio (H₂O) for each of the flights. Depending on the parameter, the database has extended by about a factor of 5-10 compared to earlier studies.
- 10

As one result of our investigations, we show that the medians of $N_{\rm ice}$, $R_{\rm ice}$ and $RH_{\rm ice}$ have distinct patterns in the IWC-T parameter space. Lookup tables of these variables as functions of IWC and T can be used to improve global model cirrus representation and remote sensing retrieval methods. Another outcome of our investigations is, that across all latitudes, the thicker liquid-origin cirrus

- 15 predominate at lower altitudes, while at higher altitudes the thinner in situ-origin cirrus prevail. Further, exemplary investigations of the radiative characteristics of in situ-origin and liquid-origin cirrus show that the in situ-origin cirrus only slightly warm the atmosphere, while liquid-origin cirrus have a strong cooling effect.
- An important step in completing the Cirrus Guide II is the provision of the global cirrus $N_{\rm ice}$ climatology, derived by means of the retrieval algorithm DARDAR-Nice from ten years of cirrus remote 20 sensing observations from satellite. The in situ measurement database has been used to evaluate and adjust the satellite observations. We found that the global median $N_{\rm ice}$ from satellite observations is almost two times higher than the in situ median and increases slightly with decreasing temperature. Nice medians of the most frequently occuring cirrus sorted by geographical regions are highest in the
- tropics, followed by austral/boreal mid-latitudes, Antarctica and the Arctic. Since the satellite cli-25 matologies enclose the entire spatial and temporal N_{ice} occurrence, we could deduce that half of the cirrus are located in the lowest, warmest (224-242 K) cirrus layer and contain a significant amount of liquid-origin cirrus.
- A specific highlight of the study is the in situ observations of cirrus and humidity in the Asian 30 monsoon anticyclone and the comparison to the surrounding tropics. In the convectively very active Asian monsoon, peak values of $N_{\rm ice}$ and IWC of 30 ppmv and 1000 ppmv are detected around the cold point tropopause (CPT). Above the CPT, ice particles that are convectively injected can locally add a significant amount of water available for exchange with the stratosphere. We found IWCs of up to 8 ppmv in the Asian monsoon in comparison to only 2 ppmv in the surrounding tropics. Also,
- the highest RH_{ice} (120-150%) inside of clouds and in clear sky are observed around and above the 35 CPT. We attribute this to the high H₂O mixing ratios (3 - 5 ppmv) observed in the Asian monsoon compared to 1.5 to 3 ppmv typically found in the tropics. The observed supersaturations above the

CPT suggest that the water transport into the stratosphere is 10-20% higher than expected in regions of weak convective activity and up to about 50% in the Asian monsoon.

40 1 Introduction

In Part 1 of the study (Krämer et al., 2016), a detailed guide to cirrus cloud formation and evolution is provided, compiled from extensive model simulations covering the broad range of atmospheric conditions and portrayed in the same way as field measurements in the Ice Water Content-Temperature (IWC-T) parameter space. The study was motivated by the continuing lack of understanding of the

- 45 microphysical and radiative properties of cirrus clouds, which remains one of the greatest uncertainties in predicting the Earth's climate (IPCC, 2013). An important result is the classification of two types of cirrus clouds that differ in formation mechanisms and microphysical properties: relatively thin cirrus that form on-site below -38°C (in situ-origin), and thicker cirrus originating from freezing in liquid clouds (liquid-origin cirrus) that are uplifted from warmer layers farther below.
- 50 Since then, a number of studies have been published that shed further light on the exploration of the high ice clouds. For example, some new studies, mostly based on aircraft or lidar observations, provide overviews and climatologies of cirrus cloud properties (Kienast-Sjögren et al., 2016; Petzold et al., 2017; Heymsfield et al., 2017a, b; Woods et al., 2018; Lawson et al., 2019) while others present a more specific view (Urbanek et al., 2017, 2018). Overviews of the properties of cirrus
- 55 derived from global satellite remote sensing observations were also recently enhanced to include ice crystal number concentrations (Sourdeval et al., 2018; Gryspeerdt et al., 2018; Mitchell et al., 2018). Several studies make use of the concept of in situ-origin and liquid-origin cirrus, e.g. Wernli et al. (2016), investigating the occurrence of in situ-origin and liquid-origin cirrus over the North Atlantic by analyzing ERA-interim data, Gasparini and Lohmann (2016) simulating, amongst other things,
- 60 the global distribution of liquid-origin cirrus and Gasparini et al. (2018) presenting climatologies of in situ-origin and liquid-origin cirrus as seen by the CALIPSO satellite and the ECHAM-HAM global climate model. Wolf et al. (2018) studied the microphysical properties of Arctic in situ-origin and liquid-origin cirrus from balloon-borne observations, and Wolf et al. (2019) provide a cirrus parametrization demonstrating the dependence on the origin of the clouds.
- The wealth of earlier (see e.g. references in Krämer et al., 2016) and new studies has provided insights into formation processes, life cycles and appearance of cirrus. Nevertheless, there are still gaps that need to be filled, on the one hand in the understanding of ice processes, and on the other in the representation of cirrus clouds in climate prediction. A way to accomplish this task requires large and high quality observational databases that can serve, for example, to evaluate global models
- 70 or other data sets and be used to derive parameterizations for improved representation of different types of cirrus clouds in models (see e.g. Wolf et al., 2019). In addition, such databases allow detailed studies of special types of cirrus that are still poorly understood, e.g. cirrus in fast updrafts as

orographic cirrus or cirrus at the top of strong convection.

- 75 In this study, we approach these requirements as follows: we first compile a data archive of airborne in situ observations which is extended with respect to earlier versions (Schiller et al., 2008; Krämer et al., 2009; Luebke et al., 2013; Krämer et al., 2016) in terms of the size of the data set that contains all parameters needed for the desired studies, i.e. meteorological parameters, ice water content (IWC), number concentration of ice crystals (N_{ice}), ice crystal mean mass radius (R_{ice}¹),
- 80 relative humidity with respect to ice (RH_{ice}) and water vapor mixing ratio (H₂O). However, though airborne in situ measurements best represent detailed microphysical properties of cirrus and their environment, they are always snapshots of specific situations that are also limited by the possibilities of the flight patterns and thus not suitable to derive spatial geographical or seasonal views of cirrus clouds. For this purpose, a globally complete dataset of remote sensing observations from satellite
- 85 observations are the better option. Hence, as a next step of the study we use in situ climatologies to evaluate cirrus N_{ice} from satellite observations and, based on this, derive a global climatology of cirrus N_{ice}. From the portrayal of the two Cirrus Guide II data sets emerging from this study together with some more detailed analyses, we show that the combined evaluation of airborne in situ and satellite remote sensing observations enhances the insights into cirrus properties. The in situ
- 90 observations are best suitable for the investigation of specific, smaller scale phenomena and for the evaluation of satellite observations or model simulations. Satellite-borne observations, on the other hand, allow a view of the larger scale and seasonal properties.

The article is structured as follows: The Cirrus Guide II in situ databases and the methods used are
described in Section 2 and Appendix A. As an overview, in Section 3 we portray the in situ cirrus cloud and humidity database with respect to altitude for the latitudes covered by the observations. The usefulness of the data set is shown by discussing the characteristics and occurrences of in situorigin and liquid-origin cirrus. Section 4 first presents cirrus and humidity climatologies of the extended Cirrus Guide II in situ data base in terms of temperature in comparison to the earlier
studies mentioned above. Further, characteristic properties of mid-latitude and tropical climatologies are presented.

In Section 5 we show another example of a specific analysis extracted from the Cirrus Guide II in situ data base: the data set includes recent unique measurements in the tropical tropopause layer (TTL) region of the Asian monsoon anticyclone, where cirrus clouds and humidity are of special interest and observations are rare. The topic is briefly introduced before the special observations

105

in the Asian monsoon are presented and compared with the conditions found in the surrounding tropical regions.

The last part of the study (Section 6) is the step to global climatology of cirrus N_{ice} from satellite remote sensing observations. For this purpose, the Cirrus Guide II in situ database is used to evaluate

¹ mean mass radius
$$\mathbf{R}_{ice} = \left(\frac{3 \cdot IWC}{4\pi\rho \cdot N_{ice}}\right)^{1/3}$$
 with $\rho = 0.92$ g/cm³

110 remote sensing cirrus observations. Based on this, a global N_{ice} climatology is derived and first analyses of the global and also regional N_{ice} are presented.

2 Databases

2.1 in situ data set

- The observations presented here include the ice water content IWC, the ice crystal number concen-115 tration N_{ice}, the mean mass radius R_{ice} in-cloud as well as the clear sky RH_{ice} and clear sky water vapor volume mixing ratio H₂O. The complete in situ data set comprises 24 field campaigns: the 17 experiments shown in Part I of this study (Krämer et al., 2016, ; the campaigns were performed between 1999 and 2014 over Europe, Africa, Seychelles, Brazil, Australia, USA and Costa Rica), extended by the field campaigns SPARTICUS 2010 and START 2008 over Central US, LTU 2012-
- 120 2018 over Kiruna, CONTRAST and ATTREX in 2014 and POSIDON 2016 over the tropical Pacific as well as StratoClim 2017 out of Nepal.

A map of flights during the various campaigns (extended map of Cirrus Guide: Part I) is shown in Figure 1. In the Appendix A, a summary of the field campaigns and deployed instrumentation is given in Table 4; Table 5 lists all campaigns and the measured parameters. Also, a discussion of new

- 125 data evaluation methods, data quality and data coverage is presented. An overview of each campaign is given in the Supplementary Material (SM in the following). Twenty campaigns are chosen to be included in the climatologies; four campaigns (marked in Table 5), where the data volume is very low (START 2008, LTU 2012-2018) or very massive (SPARTICUS in 2010, CONTRAST 2014), so that their contribution to frequency occurrences is either negligible or dominant, are shown in the
- 130 overview of measurements only (SM).

The climatologies are advanced in several aspects in comparison to the compilations of IWC by Schiller et al. (2008), Luebke et al. (2013) and Krämer et al. (2016) and N_{ice} , R_{ice} and RH_{ice} by Krämer et al. (2009):

- The number of flights and total time in cirrus increased from 104 flights / 94 hours (Krämer
- et al., 2016) to a total of 150 flights / 168 hours, i.e. the database disporportionally has extended by about a factor of 5-10 depending on the specific parameter.
 - For IWC, a new data product has been developed that increases the observed data volume (Appendix A2.1).
 - For N_{ice}, observations from advanced and extended instrumentation could be added to the database; further, a new correction of the occurrence frequencies is applied (Appendix A2.2).
 - The geographical spread of the observations has broadened, so that a portrayal of cirrus and humidity with respect to the geographical regions mid-latitude and tropics, and also latitude/altitude seems worthwhile.

140

- As in the earlier climatologies, all data underwent strict quality control.

145 2.2 Satellite data set

DARDAR-Nice provides observational-based estimates of N_{ice} obtained from CALIPSO and Cloud-Sat measurements (Sourdeval et al., 2018). This unique approach uses the sensitivity of lidar and radar measurements to small and large particles, respectively, to constrain two parameters of a particle size distribution (PSD) parametrization and N_{ice} is subsequently estimated by direct integration,

- 150 from a minimum threshold size set in this study to 5 μ m. DARDAR-Nice uses the parametrization by (Delanoë et al., 2005), in which two normalization parameters (a slope parameter N₀^{*} and the volume-weighted diameter $D_{\rm m}$) are used to predict the shape of a PSD. Sourdeval et al. (2018) have demonstrated that the method of Delanoë et al. (2005) is theoretically capable of predicting N_{ice} from recent in situ campaigns, by comparing its prediction based on in situ N₀^{*} and $D_{\rm m}$ measurements to
- 155 the actual in situ N_{ice} measurements. Good agreements were found, although it was noted that the inability of the modified gamma distribution to match the frequently bi-modal shape of the measured PSDs could lead to an overestimation of N_{ice} in DARDAR-Nice.

This typically occurs at temperature above -50° C and is expected to be cloud-type dependent. Nevertheless, Sourdeval et al. (2018) showed that the satellite N_{ice} remains in reasonable agreement

160

with the in situ (within a factor of 2) from $T = -30^{\circ}C$ down to $-90^{\circ}C$, which should cover the entire cirrus temperature ranges in this study.

This evaluation framework is here repeated on the basis of 5 in situ campaigns archived in the Cirrus Guide II in situ database: COALESC2011, ACRIDICON2014, ATTREX2014, MLCIR-RUS2014, STRATOCLIM2017. Based on the agreement between DARDAR-Nice and the in situ ob-

servations, a global N_{ice} climatology is derived from 10 years of satellite observations. Regional N_{ice} climatologies for the Arctic (90N - 67.7N), Northern mid-latitudes (67.7N-23.3N), Tropics (23.3N - 23.3S), Southern mid-latitudes (23.3S - 67.7S) and Antarctica (67.7S - 90S) are analysed in more detail. The results are presented in Section 6.

3 Vertical distribution of cirrus and humidity from in situ observations

- 170 As an introduction, atmospheric temperature profiles in the Arctic, at mid-latitudes and in the tropics are shown in Figure 2, (left panel, adopted from Schiller et al., 2008). Exemplarily, vertical profiles inside of cirrus clouds are shown for 28 flights, using blueish colors for Arctic, greenish for midlatitude and reddish for tropical observations. It can be seen that in the tropics, with the strongest warming of the Earth's surface by the sun, the air is much warmer at higher altitudes than at mid-
- 175 latitudes or in the Arctic. The coldest atmospheric temperatures are found at the points where the slopes of the temperature profile reverses: the cold point tropopause (CPT). This is the region where the transition from the troposphere to stratosphere occurs. Above the CPT, in the stratosphere, almost

no cirrus clouds are observed because it is too dry for ice formation (Schiller et al., 2009). The relations between temperature as well as altitude and the potential temperature Θ are shown in

180 Figure 2, middle and right panel for the different geographical regions. Θ is often used in upper troposphere–lower stratosphere (UT/LS) research, since it allows a clear assignment of air parcels to the associated atmospheric layer, in contrast to the temperature, whose course reverses above the CPT. The range of the tropical tropopause layer (TTL) and the tropical CPT are marked by magenta lines (after Fueglistaler et al., 2009).

185 3.1 Latitude - altitude distributions

As a first application of the Cirrus Guide II data set the distribution of cirrus clouds and humidity is shown with respect to latitude and altitude in Figure 3. Plotted are IWC (color coded by volume mixing ratio), $N_{\rm ice}$ (color coded by concentration), $R_{\rm ice}$ (color coded by size), in-cloud as well as the clear sky humidity (color coded by RH_{ice}) and H₂O (color coded by volume mixing ratio). The

- 190 color codes range from yellow to blue with increasing amount of the respective parameter; note that the data points are plotted in the order of the colors from yellow to blue. The data were collected in the latitude range from about 70°North to around 20°South, i.e. the northern mid-latitudes and the tropics are covered by the observations. The altitude range is between about 5 to 20km. The times of data sampling are displayed in the respective panels; a more detailed description of the database
- 195 is given in Section 4. The way the data is presented here as individual points was chosen because the entire range of measurements is visible. Although data overlap occurs in this type of display, it is possible to identify cirrus types and microphysical processes, especially based on extreme values. As additional overview information, we have created latitude-altitude intervals (500 m altitude, 0.5° latitude) and calculated the 25, 50 (median) and 75 % percentiles for all variables. These additional

200 altitude-latitude climatologies are shown in the SM (Figures 1–3).

Cirrus clouds are found at lower altitudes at mid-latitudes and reach higher levels in the tropical region (Figure 2). Though this structure in the measurements is influenced by the maximum or minimum height the engaged aircraft can reach, it corresponds well with the CALIPSO latitudinal height distribution of cirrus clouds, which is largely caused by the decrease in tropopause height with increasing latitude (shown in the review article by Heymsfield et al., 2017a, original figure by

205 with increasing latitu Sassen et al. (2008)).

3.1.1 in situ-origin and liquid-origin cirrus

Krämer et al. (2016), Luebke et al. (2016) and Wernli et al. (2016) describe two different cirrus types: (1) in situ-origin cirrus that form (by heterogeneous or homogeneous ice nucleation on ice

210 nucleating particles, INP, or soluble solution aerosol particles, respectively) from water vapor directly as ice at T < 235K, $RH_{ice} > 100 \%$ and $RH_w < 100 \%$; (2) liquid-origin cirrus that evolve (also heterogeneously or homogeneously) from freezing of liquid drops in clouds at T $\gtrsim 235K$ and $RH_w \sim 100 \%^2$. In other words, in situ-origin cirrus are observed at the altitudes where they are formed, whereas liquid-origin cirrus are glaciated liquid clouds from further below which are lifted to the cirrus temperature region where liquid water no longer exists.

Microphysical characteristics: In the new in situ data set, containing advanced measurements and extended by several field campaigns in comparison to the earlier studies, some typical characteristics of the cirrus types and hints to ice nucleation mechanisms are visible. In the following, the cirrus types are briefly introduced, and, using Figures 3 and 4, the types and freezing mechanisms are discussed and summarized in Table 1.

220

215

IN SITU-ORIGIN cirrus can be devided in two sub-classes depending on the strength of the updraft: in *slow updrafts* 3 , in situ-origin cirrus form mostly heterogeneously and are rather optically thin with lower IWCs and few but large ice crystals. We like to note here that in an atmosphere free

- of INPs, cirrus clouds forming homogeneously would have similar characteristics, since also only a small number of ice crystals nucleate homogeneously in slow updrafts (see e.g. Spreitzer et al., 2017). Hence, in this regime, the freezing process is not relevant for determining the cirrus properties. In *fast updrafts*³, homogeneous freezing mostly occurs regardless of the presence of INPs, since fast updrafts cause RH_{ice} to reach the homogeneous freezing threshold even after heterogeneous freezing.
- 230 The in situ-origin cirrus that emerge in these situations are optically thicker with higher IWCs and more but smaller ice crystals.

LIQUID-ORIGIN cirrus stem from lower altitudes, where more water is available, so they generally consist predominantly of thicker cirrus with higher IWC, together with more and larger ice crystals that are frozen heterogeneously at T > 235K in *slower updrafts*. In *fast updrafts*, liquid-origin cirrus

- with very high N_{ice} appear. The reason is that in such updrafts RH_w and RH_{ice} are both > 100 % and thus the Wegener-Bergeron-Findeisen process, where liquid drops evaporate and ice crystals remain at $RH_w < 100$ % and $RH_{ice} > 100$ %, is suppressed (Korolev, 2007). Thus, the numerous supercooled drops reach the altitude (temperature ~ 235K) where they freeze homogeneously (Costa et al., 2017).
- 240 The meteorological situations where slow updraft in situ-origin cirrus frequently occur (see Krämer et al., 2016) are low- and high-pressure systems (frontal/synoptic cirrus). The warm conveyor belt in low pressure systems can also produce liquid-origin cirrus. Fast updraft in situ-origin and liquidorigin cirrus occur in gravity waves, often orographically induced, in jet streams, mesoscale convective systems and anvils.
- As outlined in the Cirrus Guide I (Krämer et al., 2016), to a certain extent cirrus types can be identified by their typical characteristics. This applies to the initial stage, after which the clouds lose the signature of the formation process e.g. by sedimentation: as long as an updraft prevails (corre-

²For a more detailed description of the freezing mechanisms see Vali et al. (2015); Heymsfield et al. (2017a) and references therein.

 $^{^{3}}$ slow updrafts: ≤ 10 cm/s; fast updrafts: 10 cm/s - 10 m/s (based on simulations of Kärcher and Lohmann, 2002; Krämer et al., 2016)

sponding $RH_{ice} > 100\%$), smaller ice crystals in the size range $\lesssim 20\mu m$ grow to larger sizes on a time scale of tens of minutes. The larger ice crystals sediment to lower altitudes, thus removing ice

- surface from the cloud volume which consequently reduces the depletion of H_2O_{gas} ($\propto RH_{ice}$) by water vapor transport to the ice (for more detail see Spichtinger and Cziczo, 2010). The ice crystals that have fallen out of the layer deepen the cirrus extent to lower altitudes (fall streaks can extend the cirrus to several km below the nucleation level Jensen et al., 2012; Murphy, 2014), while at the same time, large ice crystals from above could sediment into the cloud volume. Altogether, the cirrus
- evolution is a dynamical process and the cirrus properties change in the course of a cirrus lifetime. At the final cirrus stage, i.e. when the temperature increases, the environment becomes subsaturated ($RH_{ice} < 100\%$) and ice crystals evaporate, disappearing the faster the smaller they are (timescales of growth and evaporation of ice crystals are shown in Kübbeler et al., 2011, their Figure 12).
- $260 \quad Cirrus types with the most striking features (high IWC / N_{ice}) are the easiest to identify. They could be liquid-origin or in situ-origin cirrus in fast updrafts or also liquid-origin cirrus in slow updrafts. This will be shown on the basis of Figure 3 and Figure 4.$

In Figure 4, the relation between $N_{\rm ice}$, $R_{\rm ice}$ and IWC is shown for ~ 87 h of cirrus cloud observations. It can be nicely seen that in the cases where $N_{\rm ice}$ is high, the ice crystals are small – because

- the numerous ice crystals consumed all of the available vapor, thereby suppressing further growth while low N_{ice} are related to large ice crystals for a given IWC. The IWC (in mg/m³) is indicated by the color code and result from the combination of both N_{ice} and R_{ice} . The thin black lines are isolines of IWC (in ppmv) that appear in the order shown the legend. The scheme at the right side illustrates the partitioning of the clouds between liquid and in situ-origin: the thickest cirrus (blue
- 270 points) are of liquid-origin, the thinnest (yellow points) of in situ-origin. As the thickness of the cirrus decreases, the fraction contributed by liquid-origin cirrus becomes smaller and smaller while an increasing fraction is due to in situ-origin cirrus.

Most of the highest IWCs (> 10 mg/m^3 , dark red and blue diamonds in Figure 3, top left panel) are of liquid-origin. They appear in the lower parts of the clouds, i.e. they are uplifted from farther

- 275 below. A good example of this is the field campaign SPARTICUS in 2010, which is separately plotted in Figure 7 of the SM (right panel). About 23 hours of sampling in mostly liquid-origin clouds were performed over central USA. The clouds were observed in the temperature range 210-240 K, corresponding to altitudes between 5-10 km, i.e. they were rather low cirrus clouds. The IWCs are mostly high (red to blue colors) and, as shown by Muhlbauer et al. (2014), ice particles of
- greater than a thousand μ m diameter were frequently encountered, indicative of liquid-origin cirrus (see Table 1). In the tropics (Figure 3), some blue points are also detected at high altitudes of about 17 km from measurements made above the Asian monsoon in strong convective updrafts (see also Section 5) and consist of many ice crystals (0.1-10 cm⁻³) with medium R_{ice} (20-70 μ m).

Generally, the IWC roughly shows a vertical structure of decreasing IWC with increasing altitude. 285 This is caused on the one hand by the amount of available water that decreases with decreasing temperature, but also because cirrus of liquid-origin predominate in lower layers, whereas cirrus with in situ-origin become more abundant at higher altitudes, i.e. colder temperatures. This is in accordance with the findings of Luebke et al. (2016); Wernli et al. (2016) and Wolf et al. (2018), where Luebke et al. (2016) and Wolf et al. (2018) experimentally investigated the two cloud types in

290 mid-latitudes and in the Arctic, respectively, while Wernli et al. (2016) analyzed 12 years of ERAinterim data in the North Atlantic region (see also Section 6.2).

As IWC is an indication for the cirrus type, N_{ice} (Figure 3, middle left panel) provide a hint to the freezing mechanism, either heterogeneous or homogeneous: high ice crystal numbers N_{ice} ($\gtrsim 0.5$ cm⁻³, dark red and blue diamonds, middle left panel of Figure 3) are an indicator for homogeneous

- 295 ice formation in fast updrafts caused by waves or convection, both for in situ-origin as well as for liquid-origin cirrus. At mid-latitudes, they are found e.g. at the tops of mountain wave clouds (the ice nucleation zone), which were observed for example behind the Norwegian mountains at around 62°North. High N_{ice} values have also been observed in tropical deep convection around 10°North, corresponding to measurements reported by Jensen et al. (2009).
- 300 Another source of mid-latitude high N_{ice} are young contrails. Note however, that these high N_{ice} exist only for a short time. This is because high ice crystal numbers are associated with small ice crystal sizes (yellow diamonds in the bottom left panel, see also Figure 4) that grow/evaporate quickly. At lower altitudes, N_{ice} tends to be lower and the crystals are larger, because large crystals having lower concentrations sediment out from the cloud tops and, as mentioned above, liquid-origin
- 305 cirrus with characteristic large ice crystals are also common in this altitude range. In the tropics, high N_{ice} at high altitudes are induced by in situ homogeneous freezing in convection or gravity waves (see also Jensen et al., 2013a). At cloud bases, such high N_{ice} are most probably of liquid-origin, initiated by homogeneous freezing of supercooled drops.

A more detailed discussion of the microphysical properties of cirrus including frequencies of oc-310 currence of specific signatures which can not be seen from Figure 3, is given in Section 4.

Radiative characteristics: A motivation to study cirrus clouds is to investigate the radiative properties on the basis of the findings on their microphysical properties. In the Cirrus Guide I, Krämer et al. (2016) speculated that the physically and optically thinner in situ slow updraft cirrus cause a

- 315 warming effect, while thicker fast updraft in situ-origin and, particularly, thick liquid-origin cirrus have the potential to cool. Here, we show a first estimate of the radiative forcing of typical in situorigin slow and fast updraft as well as liquid-origin cirrus (Figure 5). To this end, radiative transfer calculations for idealized scenarios under specific conditions are realized, at noon at the equinox, which are briefly described in the caption of Figure 5.
- In the left panel of Figure 5, the radiative forcing of the slow (light green) and fast (dark green) in situ-origin cirrus is displayed with respect to optical depth. This panel is expanded from the right panel, where the forcing of the liquid-origin cirrus is shown. Obviously, the net radiative forcing of

the in situ-origin cirrus is much smaller than that of liquid-origin and, moreover, change the sign from warming to cooling.

- In more detail, the slow in situ-origin cirrus have only small optical depth (τ) between 0.001 0.05, resulting in a slight net warming effect of not larger than about 1.5 W/m². The optical depth of fast in situ-origin cirrus is larger (τ : 0.05 1), but most of them are also warming (2-10 W/m²). The thickest fast-updraft in situ-origin cirrus at the lowest altitudes change the sign of their net forcing, they switch to a slight cooling effect. The reason is the warmer temperature at lower altitude that
- 330 reduces the warming effect of the longwave infrared radiation. The results of the radiative forcing calculations for the slow and fast updraft cirrus are in agreement with investigations from lidar observations reported by Kienast-Sjögren et al. (2016) and Campbell et al. (2016), who observed cirrus with optical depth up to 1 and 3, respectively, and found a decreasing warming effect with decreasing optical depth. Campbell et al. (2016) even reported a slight cooling effect at the warmest
- observed cirrus. The liquid-origin cirrus, however, mostly found in the warmest cirrus layers, have large optical depths (τ : 1 - 12), which is larger than the range of cirrus optical depth reported in many studies (the maximum optical depth is often found to be 1-3, e.g. Sassen et al., 2008; Kienast-Sjögren et al., 2016; Campbell et al., 2016; Mitchell et al., 2018, ; note that this is likely because the lidar technique, often use to investigate cirrus cloud optical properties, has restrictions in the τ
- 340 determination of thicker ice clouds). A consequence of the large optical thickness is a quite strong net cooling effect (- 15 to -250 W/m²) of liquid-origin cirrus. These values are of the same order of magnitude as reported from direct measurements inside of cirrus clouds (Wendisch et al., 2007; Joos, 2019).
- Thus, from this first and very idealized simulations, we can conclude that in situ formed cirrus 345 clouds are most likely to warm the atmosphere, whereas liquid-origin ice clouds have the potential for strong cooling. Note here, that we only investigate local time 12 h, where the cooling is probably most pronounced. For lower sun position (i.e. larger zenith angle) the cooling is probably reduced and during night time, cirrus clouds can only warm the atmosphere (due to the thermal greenhouse effect). Thus, the net effect of cirrus clouds averaged over the whole daily cycle is not yet clear. Such 350 investigations go beyond the scope of this study and are subject of future work.

3.1.2 Humidity

The distribution of in-cloud and clear sky RH_{ice} as well as water vapor H₂O with latitude and altitude is shown in the right column of Figure 3. It can nicely be seen how the amount of H₂O decreases with altitude (bottom panel). The clear sky RH_{ice} (middle panel) ranges from very dry conditions 355 (< 70% green and orange diamonds) up to highly supersaturated regions (> 130%, dark red and blue diamonds), which mainly are found at high altitudes in the tropics. Such high supersaturations are also found inside of the tropical cirrus clouds (top panel). The behavior of RH_{ice} will be further discussed in Sections 4 and 5.

4 In situ climatologies

360 With the term climatologies we refer here to statistical evaluations of the available variables with regard to temperature or potential temperature.

4.1 The IWC-T parameter space: median N_{ice}, R_{ice}, RH_{ice}

In the Cirrus Guide I (Krämer et al., 2016), observations and model simulations are portrayed in the Ice Water Content- Temperature (IWC-T) parameter space. One result of the simulations is that IWC and N_{ice} are correlated with each other. This relationship has been confirmed for some meteorological situations by examples from six individual field campaigns. The database of combined IWC-N_{ice} measurements has grown considerably since then, and with it the covered temperature range of the

With the extended Cirrus Guide II data set, we further investigate the appearance of N_{ice} , but also 370 the corresponding mean mass size R_{ice} and the in-cloud RH_{ice} in the IWC-T parameter space. In Figure 6, medians of the respective parameters are presented in intervals (five iwc intervals per order of magnitude, 1K temperature intervals) covering the entire IWC-T parameter space. The variability of the parameters can be seen in Figures 1–3 of the SM, where the 25, 50 (median) and 75 % percentiles for the variables are shown.

observations. In the Cirrus Guide I, only few observations below 200 K were available.

375 4.1.1 IWC-T-N_{ice} and R_{ice} relation

Observational evidence for the correlation between IWC and N_{ice} can nicely be seen in the upper panel of Figure 6. Almost symmetrical colored bands of N_{ice} can be seen across the entire IWC-T parameter space (N_{ice} concentration increases from yellow to blue). With decreasing temperature, the same N_{ice} numbers cause lower IWC values, which is caused by the likewise decreasing available water content of the air. This finding might be of importance for parametrizations used in global

- 380 able water content of the air. This finding might be of importance for parametrizations used in global models or satellite retrieval algorithms, where IWC is often the only available parameter that characterizes cirrus clouds, but functions are used to assign N_{ice} to specific IWCs. This new analysis can be used (after some smoothing of the bands, which will be subject of a follow up study) as lookup table to derive N_{ice} from the information of temperture and IWC.
- The ice crystal sizes also form colored bands in the IWC-T parameter space (Figure 6, middle panel; R_{ice} inreases from yellow to blue, running diagonally through the IWC-T room with the size of the ice crystals increasing with increasing temperature (and thus decreasing amount of available water). This is because R_{ice} is calculated from the third root of the IWC divided by N_{ice}, as described in Section 1. However, the R_{ice} bands are somewhat less clear delineated from each other than the
- 390 N_{ice} bands. They might be smoothed in the follow up study, so that R_{ice} can also be assigned to the IWC-T data points.

One could have expected to find distinct pattern for in situ-origin and liquid-origin cirrus in this type of analysis to quantify the characteristics of the cirrus types shown in Table 1. However, the differences are merged by the calculation of median values in the overlap regions of the types in

395 the IWC-T parameter space. However, we believe that the more heterogeneous structure of the R_{ice} bands is caused by the different cirrus types, because the ice crystal size is their most pronounced difference. A longer term research goal is to derive the IWC-T-N_{ice}-R_{ice} relations seperately for in situ-origin and liquid-origin cirrus.

4.1.2 IWC-T-RH_{ice} relation

- 400 In the bottom panel of Figure 6, median in-cloud RH_{ice} are shown in the IWC-T parameter space. Also for RH_{ice} patterns are visible: RH_{ice} decreases with decreasing IWC. Above the median IWC (black solid line), RH_{ice} is mostly between 90-110 % (orange data points), i.e. around saturation. This corresponds to the existence phase of the cirrus (between 110-100 / 100-90 %, the ice crystals slowly grow /evaporate). Below the median IWC, the median RH_{ice} decrease with decreasing IWC
- 405 (green and yellow data points), reflecting the evaporation phase of the cirrus clouds. Interesting is the distribution of the higher supersaturations (red and blue points): at low temperatures, the clouds tend to higher supersaturations. This has been discussed by (Krämer et al., 2009) for thin cirrus (low IWC) and explained by the corresponding low N_{ice} , whose small ice surface cannot efficiently deplete the water vapor. However, the supersaturations are also seen at high IWC (~ high N_{ice}). The
- 410 reason for that is the vertical velocity, which is -together with the ice surface and temperature- a driver of the RH_{ice} (RH_{ice}= H₂O/H₂O_{sat,ice}(T)). The air inside of clouds is supersaturated in cases the depletion of H₂O on the available ice surface (decrease of RH_{ice}) cannot compensate for the increase of RH_{ice} caused by the cooling of the air (decrease of H₂O_{sat,ice}(T) with decreasing temperature). High IWC, coinciding always with high N_{ice}, appear in high updrafts, in in situ-origin as well as
- 415 in liquid-origin cirrus. These high updrafts allow that the supersaturations remain high despite the large ice surfaces available for deposition of excess water vapour. This was also reported by Petzold et al. (2017), who observed high RH_{ice} together with high N_{ice} in tropical convective cirrus on board of passenger aircraft (IAGOS).

4.2 Entire in situ climatologies

420 The Cirrus Guide II data set shown in Figure 3 is now displayed as frequencies of occurrence as a function of temperature (binned in 1 K intervals, Figure 7) and discussed in comparison to the earlier in situ climatologies presented by Schiller et al. (2008), Krämer et al. (2009) and Luebke et al. (2013) (Figure 6, SM). Further, as mentioned above, the new data set is large enough to split it in mid-latitude and tropical cirrus. The differing cirrus cloud properties are presented in Figures 8 and the respective clear sky and in-cloud RH_{ice} in Figures 9.

4.2.1 Ice water content (IWC-T)

Figure 7 (top left panel) depicts the IWC. The black solid and dotted lines represent the median, minimum and maximum IWC of the core IWC band, that is the envelope of the most frequent IWC (>5% per IWC-T bin, Schiller et al., 2008).

- 430 The number of hours spent sampling in cirrus clouds increased from 27 hours in Schiller et al. (2008), 38 hours in Luebke et al. (2013) and 94 hours in (Krämer et al., 2016) to 168 hours in the new extended data set. Part of the additional data is due to the new IWC data product that is applied to all campaigns and combines the IWC from total water measurements as well as the IWC derived from cloud particle size distributions (see Appendix A2.1).
- However, the median IWC and the core IWC band -decreasing with temperature as described by Schiller et al. (2008)- is still valid, showing that the IWC measurement techniques are robust and that the IWC is a stable parameter describing cirrus clouds. Note here that at temperatures ≤ 200 K data points underneath the lower dotted line are not unambiguously identified as clouds, while above about 200 K this threshold is 0.05 ppmv. For more detail see also Appendix A2.1.

440 4.2.2 Ice crystal number (Nice-T)

455

About 90 hours of $N_{\rm ice}$ observations are shown in Figure 7 (middle left panel ⁴), which is an increase of about a factor of ten in comparison to the data set of Krämer et al. (2009), who compiled 8.5 hours (Figure 6, SM). For $N_{\rm ice}$, the picture has greatly changed when comparing the old and the new data sets. This is on the one hand due to an extension of the lower detection limit of $N_{\rm ice}$ from

- 445 4.10⁻³ to 10⁻⁴ cm⁻³ (see Appendix A2.2), but also because the new data set represents a better mixture of different meteorological situations. For example, the higher N_{ice} at warmer temperatures in the Krämer et al. (2009) data set (Figure 6, SM) were caused by flights where lee wave cirrus behind the Norwegian mountains were probed (see also Figure 3, blue diamonds at around 60°North). Also, at temperatures colder than about 200 K, N_{ice} was most often very low. Further, the enhanced
- 450 occurrence frequencies at the lowest concentrations seen in the earlier data set are corrected in the new data evaluation procedures (Figures 17 and Figure 6, SM, middle left panel).

90 hours of aircraft N_{ice} observations within cirrus clouds is a tremendous amount when taking into account the necessary effort. However, this is still far from being representative for the distribution of N_{ice} in the atmosphere. We nevertheless calculated 10, 25, 50, 75 and 90% percentiles, which are shown as thin, dotted and solid lines in Figure 7 (middle left panel). Note that the 10 and 90%

are shown as thin, dotted and solid lines in Figure 7 (middle left panel). Note that the 10 and 90% percentiles enclose the core region of N_{ice} , i.e. the envelope of the most frequent N_{ice} (>5% per N_{ice} -T bin). Fits through these percentiles and the median N_{ice} reveal no temperature dependence of N_{ice} (10%, median and 90% N_{ice} : 0.002, 0.03 and 0.3 cm⁻³). This is different to the slight decrease with

 $^{^{4}}$ Note that for $N_{\rm ice}$ - and thus $R_{\rm ice}$ - the number of hours spent in clouds in Figure 7 is less than in Figure 3. For details see figure caption.

temperature of the minimum, middle and maximum Nice lines shown by Krämer et al. (2009), which

- 460 was, as discussed above, caused by two flights with high N_{ice} at comparatively warm temperatures. It is a question why the N_{ice} do not increase with decreasing temperature, as would be expected because the homogeneous ice nucleation rate then increases. This has been investigated by Gryspeerdt et al. (2018) based on a 10 year global data set retrieved from satellite observations (DARDAR-Nice, Sourdeval et al., 2018, see also Section 6). Gryspeerdt et al. (2018) analyzed the N_{ice} (>5µm) only at
- cloud tops and also those throughout the cirrus clouds. From the cloud top analysis, a clear increase 465 of N_{ice} with temperature was obvious, while integrating throughout the cirrus this temperature dependence becomes much weaker, though it is still present. Gryspeerdt et al. (2018) propose that the missing temperature dependence in the in situ results could be due to a lack of in situ measurements near the cloud top, where the temperature dependence is strongest. Another reason could be that the higher $N_{\rm ice}$ are short-lived (see Section 3.1.1) and thus not easy to trace by aircraft. 470

A further consideration of Nice frequencies of occurrence on a global as well as regional scales derived from satellite remote sensing will be presented in Section 6.

4.2.3 Ice crystal mean mass radius (R_{ice}-T)

The ice crystal mean size is calculated as mean mass radius $R_{\rm ice}$ as shown in Footnote 1. $R_{\rm ice}$ is 475 close to the common effective cloud particle radius $R_{\rm eff}$ (~ ice Volume/Area), but can be calculated without knowing details of the ice particle size distribution (see also Krämer et al., 2016).

A total of 84 hours of observations are compiled in Figure 7 (bottom left panel). Overall, the mean mass Rice ranges from about 1 to 100 µm, while individual ice crystals in cirrus can reach sizes up to 1000 μm or even larger. The $R_{\rm ice}$ core band (frequencies $\gtrsim 5\%$ per $R_{\rm ice}$ -T bin) decreases slightly

as the temperature decreases, which is caused by the decrease of the IWC core band, since the N_{ice} 480 band is not dependent on temperature (see previous section).

The 10, 25, 50, 75 and 90% percentiles of the data set are plotted as thin, dotted and solid red lines. The black lines represent the minimum, middle and maximum R_{ice} shown by Krämer et al. (2009) (see Figure 6, SM). The range of Rice has shifted slightly to larger sizes in the new data set, which

is caused by the Nice range extended towards lower concentrations that mainly consist of larger ice 485 crystals.

Remarkable is the drop of the most frequent Rice from larger to smaller ice crystals seen at around 215 K. This is probably the approximate temperature up to which liquid-origin clouds are detected (see Luebke et al., 2016; Sourdeval et al., 2018, and also Section 6.2), that are characterized by

490 larger ice crystals than in situ-origin cirrus (Krämer et al., 2016). At higher temperatures where both liquid and in situ-origin cirrus prevail, higher IWCs - and thus also a larger Rice - occur more often than at lower temperature where only the thinner in situ-origin clouds exist. This is especially true at mid-latitudes, in the tropics the liquid-origin cirrus reach also lower temperatures and thus no sudden drop of R_{ice} is observed (see Figure 8, left and right bottom panel). The bifurcated structure 495 of the most frequent R_{ice} which can be seen for temperatures ≤ 190 K is discussed in Section 4.3.1 (vii).

4.2.4 Clear sky and in-cloud RH_{ice} (RH_{ice}-T)

The new RH_{ice} clear sky and in-cloud data sets are displayed in the middle and top right panel of Figure 7. The respective earlier data sets of Krämer et al. (2009) are shown in Figure 6, SM. The overall picture of the RH_{ice} distributions has not changed substantially, though the amount of in-cloud data in the database has increased from 10 hours of measurements to about 96 hours, and the clear sky observation time from 16 to 186 hours. The only small difference is found in the in-cloud RH_{ice}, potentially caused by the larger amount of data: below about 200 K, where high supersaturations (>120%) occur more often and low subsaturations (<80%) less often. At higher temperatures, the peak of RH_{ice} frequencies at 100% is more pronounced.

For the new data set, in addition to the clear sky RH_{ice} the absolute water vapor volume mixing ratio H_2O in the bottom right panel of Figure 7 is plotted. To guide the eye, water vapor saturation wrt ice, $H_2O_{sat,ice}$, is drawn as black solid line. The decrease of H_2O with temperature is nicely seen in this panel and clear sky supersaturations appear in this portrayal as data points above $H_2O_{sat,ice}$.

510

4.3 Mid-latitudes and tropics

4.3.1 Mid-latitude and tropical cirrus clouds

The data sets of mid-latitude/tropical cirrus consist of 67/101 hours of IWC, 29/61 hours of N_{ice} and 28/56 hours of R_{ice} measurements. This section describes some of the pronounced characteristics of
 mid-latitudes and tropical cirrus, which are displayed in Figure 8 (greenish/ reddish colors represent mid-latitude/tropical cirrus).

(i) *Temperature ranges:* Comparing mid-latitude and tropical cirrus, the first obvious difference -as expected when looking at the temperature profiles in Figure 2- is the temperature range. The
 520 observed mid-latitude cirrus rarely occur below 200 K, while tropical cirrus are detected down to temperatures of 182 K. The core IWC/N_{ice}/R_{ice}range of both mid-latitude and tropical cirrus corresponds to the total climatology (see Figure 7).

(ii) *Mid-latitude WCBs and MCS:* At European mid-latitudes, the most frequent cirrus can be
assigned to slow updrafts in frontal systems (WCBs: warm conveyor belts) containing both liquid-origin and in situ-origin cirrus. High IWCs stem mostly from liquid-origin WCB cirrus. Above the central USA, mesoscale convective systems (MCS) with faster updrafts are more frequent. The resulting liquid-origin cirrus are thicker than the European, i.e. the ice crystals are larger and the IWC is higher (see also Krämer et al., 2016).

(iii) *Contrails:* A striking feature in the cirrus observations are N_{ice} of up to several hundreds per cm³, that are found at mid-latitudes in the temperature range of about 210-220 K, which corresponds to about 10 km altitude (see Figure 2), the typical cruising level of passenger aircraft. They can be attributed to young contrails, which were a topic of investigation during COALESC 2011 (Jones et al., 2012) and also ML-CIRRUS 2014 (Voigt et al., 2017). Higher mid-latitude N_{ice} at higher temperatures are most probably in situ-origin cirrus caused by stronger updrafts in e.g. mountain waves.

(iv) *Drop freezing:* High N_{ice} between ten and hundred per cm³ or even more are found above 235
K. Such high concentrations together with small R_{ice} are typical for supercooled liquid cloud drops

540 that might be frozen by spontaneous homogeneous drop freezing (that occurs latest at around 235 K in the atmosphere) or not. In any case, these cloud particles are an indication for liquid-origin clouds caused in tropical or mid-latitude convective systems with fast updrafts.

(v) *Tropical convection:* In contrast, in tropical cirrus, and particularly for temperatures ≥ 220 K,
545 high IWCs and N_{ice} above the core range -corresponding to convective liquid-origin cirrus- become more frequent, while R_{ice} tends to be smaller. This is most likely caused by the fast convective vertical velocities that often let clouds in the mixed-phase temperature regime rise to the cirrus altitude range.

- (vi) *Tropical deep convection:* Also remarkable is that in the tropics, massive convective liquidorigin cirrus carrying a high IWC -often accompanied by high N_{ice} - are detected down to very cold temperatures (< 200 K), which corresponds to high altitudes up to 17 km. The thickest cirrus with very high IWC and N_{ice} can also be seen in Figure 3 (upper and middle left panel: distribution of cirrus with latitude; blue data points represent liquid-origin with a high IWC/ N_{ice}) at around
- 555 25° northern latitude and 16-18 km altitude. These exceptional thick and cold cirrus at high altitudes were observed in the Asian monsoon tropical tropopause layer (TTL). The observed many small N_{ice} are generated most likely by in situ homogeneous ice nucleation, triggered either by fast updrafts in gravity waves (see also Spichtinger and Krämer, 2013; Jensen et al., 2017) or deep convection. They often occur in the tops of massive liquid-origin cirrus with very high IWC; a theoretical description
- 560 of such clouds is given by Jensen and Ackerman (2006).

(vii) *TTL cirrus:* The discussion of cirrus clouds in the tropical tropopause layer is presented in Section 5.

4.3.2 Mid-latitude and tropical humidity

535

565 The mid-latitude/tropical humidity data sets include 24/72 hours of in-cloud RH_{ice} and 51/132 hours of clear sky RH_{ice}/H_2O measurements, which are displayed in Figure 9 with the same color code as in Figure 8.

In clear sky at temperatures higher than about 200 K, RH_{ice} is most often below saturation and randomly distributed in both mid-latitudes and tropics (Figure 9, middle row). Clear sky supersatu-

570 rations occurs less frequently, simply because they only take place in those periods when moist air parcels are cooled towards the ice nucleation thresholds (heterogeneous or homogeneous), which are rare compared to drier conditions of the atmosphere.

Below about 200 K, i.e. in the TTL (see also Section 5), the clear sky RH_{ice} distribution looks very different. In this region, H_2O is low and its variability is only small (Figure 9, bottom row). We

- 575 plotted lines of constant H₂O (1.5, 3 and 5 ppmv) in the clear sky RH_{ice} panels to illustrate that for constant H₂O, RH_{ice} increases only due to the decrease in temperature, i.e. H₂O_{sat,ice}. Thus it can be seen that at mid-latitudes RH_{ice} mostly represent H₂O values around 3 ppmv and in the tropics between 1.5 and 3 ppmv. Since in the tropics much colder temperatures are reached, the respective RH_{ice} ranges from 10 up to about 150% or even more.
- 580 Clear sky RH_{ice} above the homogeneous freezing line are under discussion, because they would indicate that no supercooled liquid aerosol particles are present to initiate freezing, or that the homogeneous freezing is prevented, for example by organic material contained in the aerosol particles. More probably, these few data points are outliers; note also that the uncertainty of RH_{ice} rises from approximately 10% at warmer temperatures to about 20% at colder temperatures (Krämer et al., 585 2009).

Inside of cirrus, the peak of the RH_{ice} frequencies is mostly around the thermodynamical equilibrium value of 100% (saturation) at mid-latitudes as well as in the tropics (Figure 9, upper row). However, in the TTL, at the coldest prevailing temperatures (≤ 190 K), supersaturation increasingly

- 590 becomes the most common condition, which is discussed in more detail in Section 5. High supersaturations at low temperatures were also reported by (Krämer et al., 2009) and Jensen et al. (2013a), and the reason given for the existence of such high supersaturation is low N_{ice} concentrations, which were mostly present at low temperatures in these observations (Figure 6, SM). But, Jensen et al. (2013a) also showed that RH_{ice} rapidly drops to saturation in the presence of many ice crystals. As
- 595 can be seen from Figure 8, middle right panel (and discussed in Section 4.2.2), in the new data set N_{ice} cover a broader concentration range in comparison to the earlier data (Figure 6, SM), while RH_{ice} is supersaturated in most cases. This is not straightforward to understand due to the complex relation between RH_{ice} and N_{ice} . We will investigate the TTL supersaturations in cirrus clouds in a follow-up study.

600 5 TTL in situ climatologies

in- and outside of the Asian monsoon anticyclone

The tropical tropopause layer is the region above the upper level of main convective outflow, where the transition from the troposphere to stratosphere occurs. It is placed at temperatures <205 K between ~150 hPa/355 K potential temperature/14 km and 70 hPa/425 K/18.5 km (Fueglistaler et al.,

605 2009, see also Figure 2). The coldest temperatures are found here at the point where the slope of

the temperature profile reverses (cold point tropopause, CPT). In the TTL, the prevailing dynamical conditions are very slow large-scale updrafts superimposed by a spectrum of high-frequency gravity waves (i.e. Spichtinger and Krämer, 2013; Dinh et al., 2015; Jensen et al., 2017; Podglajen et al., 2017). In addition, deep convection with fast updrafts that occasionally overshoot into the TTL.

- 610 Cirrus clouds and humidity in the TTL deserve a special consideration, because this region represents the main pathway by which water vapor enters the upper troposphere and lower stratosphere (UT/LS) where it is further distributed over long distances (e.g. Brewer, 1949; Rolf et al., 2018; Vogel et al., 2016; Ploeger et al., 2013). This is of importance, because water vapor is a greenhouse gas that has a significant impact on the climate, with greatest sensitivity in the tropical UT/LS (Solomon
- et al., 2010; Riese et al., 2012), but also in the LS at high latitudes where it is being transported from the tropics. Cirrus clouds are of particular relevance as regulators of the partitioning of H₂O between gas and ice phase. Furthermore, they have a climate feedback themselves by influencing the Earth's radiation balance (e.g. Boucher et al., 2013). Thus, the simultaneous observation and analysis of both cirrus clouds and humidity (H₂O) in this climatically crucial region are of special interest.

This section specifically reports cirrus clouds together with humidity (from the climatologies presented in Section 4), that was recently observed for the first time in the TTL in the Asian monsoon anticyclone (June to September) in comparison to observations in the surrounding tropical regions. As the Asian monsoon is characterized by strong convective activity, this comparison will show

- 625 the difference to less active, calmer areas. Observations of Asian monsoon TTL cirrus clouds and humidity are of particular importance because, amongst other trace species, large amounts of H₂O and also cloud particles are convectively transported upwards from far below, where the additional H₂O in turn often cause cirrus formation (e.g. Ueyama et al., 2018, and references therein). Directly injected H₂O or H₂O from sublimated ice crystals can then be mixed up into the stratosphere. Thus,
- 630 the Asian monsoon anticyclone represents a significant gateway for H₂O between UT and LS (e.g. Fueglistaler et al., 2009; Ploeger et al., 2013) and currently it is under discussion to what extent cirrus cloud particles contribute to the amount of H₂O entering the stratosphere (e.g. Ueyama et al., 2018, and references therein).

The airborne measurements in the Asian monsoon (see Figure 1 and Tables 4 and 5) were performed during July-August 2017 out of Khatmandu, Nepal, during a field campaign embedded in the StratoClim project (http://www.stratoclim.org/). An overview of the observations is given in Figures 10, 11 and 12, where the frequencies of IWC, N_{ice}, R_{ice} and in-cloud, clear sky RH_{ice} as well as the clear sky H₂O volume mixing ratio are shown in the temperature and also the potential temperature Θ parameter space. ⁵ Most of the measurements during StratoClim are performed at

temperatures ≤ 205 K, corresponding to potential temperatures ≥ 355 K and altitudes about ≥ 14 km, i.e. in the TTL (marked in the middle panels of the Figures).

⁵ The additional Θ portrayal provides a more detailed impression of the distribution of cirrus clouds and humidity around the CPT, i.e. at the transition between troposphere and stratosphere (see Figure 2).

The surrounding, in most (but not all) cases calmer tropical TTL regions are represented by observations during the campaigns shown in Figure 1 and listed in Tables 4 and 5. The majority of the data were sampled during ATTREX_2014 and POSIDON_2016. The TTL measurements in the

645

temperature parameter space are shown in Figures 8 and 9; in these plots, the Asian monsoon observations are included, but since the measurements represent only a small part of all TTL observations, excluding the StratoClim campaign only slightly change the frequency distributions. Thus, the Figures are representative for the TTL outside of the Asian monsoon anticyclone, so we refrain from showing an additional Figure. The measurements in the Θ parameter space (StratoClim excluded) are presented in the left panels of Figures 11 and 12.

650

5.1 TTL cirrus clouds

5.1.1 **Temperature parameter space**

In the tropics outside of the Asian monsoon (Figure 8, right column), cirrus IWC and Nice range from very low to quite high values. We want to draw attention to a special feature of the most frequently 655 occurring cirrus N_{ice} (middle right column). Two main branches of most frequent N_{ice} are found, one at very low ($\sim 10^{-3}~cm^{-3})$ and the other at moderate ($\sim 5\cdot 10^{-2}~cm^{-3})~N_{\rm ice}.$ These two branches are also reflected in R_{ice}, where the larger ice crystals are found together with lower concentrations and vice versa, as shown in Figure 4. More precisely, cirrus with $N_{\rm ice} \lesssim \! 1.5 \cdot 10^{-2} \ cm^{-3}$ consist of ice particles larger than about 20 μ m 6 . That means that the ice crystal spectra of the low N_{ice} cirrus

- 660 most likely represent aged clouds (in situ-origin or liquid-origin) where the smaller nucleation mode ice crystals are either grown to larger sizes in supersaturations or evaporated in case of subsaturated conditions (see also Section 3.1.1). The middle N_{ice} cirrus containing ice crystals smaller than 20 μ m are most likely young cirrus that have formed in situ, because these small crystals quickly (on a time scale of 10 - 20 minutes) grow to larger sizes. It is impossible to speculate if they have formed homo-
- or heterogeneously, since both pathways might produce such N_{ice} in the slow updrafts prevailing 665 in the TTL. The reason that the aged cirrus marked in Figure 8 (middle right column) become visible only in the TTL, though they certainly occur in all cirrus regions, is probably the calmer dynamic environment with -in comparison to lower altitudes- less frequently occurring temperature fluctuations. These fluctuations likely cause new ice nucleation events superimposed on the aged 670 cirrus.

675

The clouds observed in the Asian monsoon include in situ formed cirrus as well as cirrus clouds from overshooting deep convection. In the much more convectively unstable Asian monsoon conditions, IWC and N_{ice} are most frequently above the median lines derived from the entire climatology (Figure 10, left column) and also high in comparison to the tropical climatology (Figure 8, right column). Here, the highest observed values (at temperatures <205 K) are found with IWC mixing ratios

⁶It should be noted that concentrations $\leq 0.1 \text{ cm}^{-3}$ of cloud particles $\leq 20 \mu \text{m}$ are below the detection limit of cloud spectrometers, i.e. small ice crystals with such low concentrations could be present (see Appendix A2.2).

of up to 1000 ppmv and a maximum N_{ice} as high as 30 cm⁻³ (note that the ice crystal shattering was significantly minimized, see Section A2.2). Also, the ice crystals mean mass size, R_{ice} , is above the median, especially at very low temperatures, which means that large ice crystals are found around the cold point. These exceptional findings are recorded during flights in strong convection, where

680 liquid-origin clouds from far below are detected in the upper part of the Asian monsoon anticyclone simultaneously with freshly homogeneously nucleated ice crystals. The observations were possible due to the pilot of the Geophysica aircraft, who dared to fly into the strong updrafts. Because of the dangerous nature of measurements under such conditions, the frequency of convective – and also orographic wave cirrus – is under-represented in the entire in situ climatology.

685

690

5.1.2 Potential temperature (Θ) parameter space

The distribution of Asian monsoon cirrus clouds in the Θ parameter space is shown in the right panel of Figure 11. TTL average upper and lower boundaries and the CPT are marked in the middle panel of the Figure following the definition of Fueglistaler et al. (2009). In the left panel of Figure 11, the climatologies of the tropical observations without Asian monsoon are shown.

From Figure 11, upper row, it can be nicely seen how steeply the IWC increases during the transition from the TTL to the free troposphere. In the tropics outside of the Asian monsoon, the maximum Θ where ice is detected is about 420 K (\sim 19 km). The mixing ratios of these highest cirrus are about 0.05 ppmv. At about 380 K (CPT, \sim 16-18 km), the range of most frequent IWCs

695 broadens, ranging between about 0.005 and 1 ppmv. Some higher IWCs are also detected in the upper TTL, indicating that overshooting convection (liquid-origin cirrus) is also embedded in these measurements. Below about 380 K the most frequent IWC increases steadily up to values of around 10 ppmv at 355 K and 1000 ppmv at 340 K and farther below.

In the Asian monsoon, the maximum Θ where ice is detected is about 415 K (~0.5 ppmv) and at 400 K the IWC ranges between 0.05 - 0.1 ppmv, as outside of the Asian monsoon. But, at 380 K (CPT), the range of most frequent IWCs rises to 0.5 - 2 ppmv, then steadily increasing below about 380 K to values of around 50 - 500 ppmv at 355 K. Obviously, IWC near and below the CPT within the Asian monsoon anticyclone is enhanced by a factor of ten or more by injection of liquid-origin cirrus in overshooting events. The highest TTL IWCs (up to 1000 ppmv) are detected in the Asian

monsoon up to 390 K.

These overshooting cirrus around and below the CPT are also seen in high ice crystal numbers N_{ice} ($\gtrsim 0.5 \text{ cm}^{-3}$) when comparing the Asian monsoon with the other tropical regions (Figure 11 middle panels). Striking are again the high N_{ice} up to 30 cm⁻³ in the Asian monsoon, already discussed with respect to the temperature parameter space. Here it is visible that this burst of in situ homogeneous

710 ice nucleation in a strong convective event (theoretically described by Jensen and Ackerman, 2006) took place below the CPT. Above 390 K, at the transition to the stratosphere, N_{ice} is very low, mostly lower than 0.1 cm⁻³. Recall that these low concentrations only contain particles > 20 µm (Footnote 6 and Appendix A2.2). This means that in the Asian monsoon (middle right panel of Figure 11) low concentrations of larger

715 ice crystals (together wit higher IWCs) are more often present at such altitudes in comparison to the surrounding tropics.

In the TTL outside of the Asian monsoon, the two branches of more frequent N_{ice} (and R_{ice}) – discussed with respect to the temperature parameter space (Figure 8, right middle panel) – are also very clearly visible. The smaller/larger ice crystals with higher/lower concentrations were identified

720 as young and aged cirrus clouds, which most probably have formed in situ. Farther below, N_{ice} further increases with decreasing Θ /altitude and more and more small liquid-origin cloud particles (frozen or liquid) with higher concentrations appear.

5.2 TTL humidity

5.2.1 Temperature parameter space

- 725 Considering the in-cloud and clear sky RH_{ice} in the Asian monsoon (Figure 10, right column) in comparison to the entire tropical climatologies (Figure 9, right column), significant differences are visible. Below about 200 K, the most frequent in-cloud RH_{ice} are found in supersaturation in the Asian monsoon cirrus. In the entire climatology (Figure 9, right column) this occurs only below about 185 K. The same is seen in clear sky RH_{ice}: higher supersaturations occur frequently already
- 730 at higher temperatures in the Asian monsoon comparison to the entire tropical climatology. These higher supersaturations reflect a higher amount of water vapor in the Asian monsoon TTL, visible through the dashed lines in the middle right panel of Figure 10. They show the increase of RH_{ice} caused by the decrease of H₂O_{sat,ice} during cooling of air at a constant H₂O mixing ratio. These lines correspond to H₂O between 3 and 5 ppmv in the Asian monsoon, while in the total tropical
- 735 climatology the most frequent H₂O ranges only between 1.5 and 3 ppmv (Figure 9; compare also the bottom panels of the Figures, where H₂O is plotted). Due to the higher water vapor mixing ratios, supersaturation occurs already at higher temperatures. Implications of the observed supersaturations are further discussed in Section 5.3.

5.2.2 Potential temperature (Θ) parameter space

- The high RH_{ice} in and outside of cirrus clouds in the Asian monsoon are also visible in the Θ representation of the humidity (see Figure 12, top and middle right panels). In the tropical climatology outside of the Asian monsoon (top left panel of Figure 12), RH_{ice} most frequently center around saturation inside of the cirrus clouds. In the cloud free TTL outside of the Asian monsoon (middle left panel of Figure 12), a humidification of the layer between ~ 360 and 380 K to RH_{ice} around 90%
- ran be seen (for more detail see Schoeberl et al., 2019).

In the Asian monsoon, on the other hand, the in-cloud RH_{ice} in the TTL exceeds saturation (top right panel of Figure 12). Especially, beween 380 and 400 K the most frequent RH_{ice} is around 130%. Also outside of clouds (middle right panel of Figure 12), around the CPT supersaturation is frequently detected and in general the humidification is higher than in the surrounding tropical

- regions. This is in agreement with Schoeberl et al. (2019), who reported high RH_{ice} coincident with the Himalaya Monsoon during summer and closely associated with convection. The higher amount of water vapor in the Asian monsoon in comparison to the other tropical regions can also be seen in the H₂O volume mixing ratios (bottom panels of Figure 12). In the tropics outside the Asian monsoon, the most frequent H₂O between 365-410 K is 1.5-4 ppmv, while in the Asian monsoon
- 755 we found 3-8 ppmv, as also seen in the temperature parameter space. This finding is in accordance to other studies, but has been observed in situ from aircraft directly in the Asian monsoon for the first time. For example, analyzing the air mass histories of higher in situ H₂O observations at other locations (highest values of 8 ppmv), Schiller et al. (2009) found that those air masses has passed the Asian monsoon region. Also, Ueyama et al. (2018) (and references therein) reported 5-7 ppmv at
- 760 100 hPa from MLS observations and extensive model simulations over the Asian summer monsoon region.

5.3 H_2O and IWC for transport to the stratosphere

The H₂O transport to the stratosphere is regulated by the coldest temperature an air parcel experiences during transition through the tropopause region. The water amount passing the tropopause region is set by the freeze-drying process associated with this transition (Jensen and Pfister, 2004) and is

discussed to be as low as H₂O saturation at the minimum temperature (e.g. Schiller et al., 2009).

However, Rollins et al. (2016) showed for the ATTREX 2014 observations, that the water vapor at the stratospheric entry point is higher by $\sim 10\%$, because the water vapor depletion by ice crystals becomes increasingly inefficient at temperatures below 200 K (note that this is a rough estimate of

- 770 the excess water vapor at the stratospheric entry point based on the actual temperature, which could be somewhat different from the minimum temperature of the air parcel's back trajectory). This is of importance, since already small amounts of H₂O can influence the stratospheric radiation budget (Solomon et al., 2010; Riese et al., 2012).
- High supersaturations at the coldest points of the TTL are also discussed in Section 5.2.1. The
 ATTREX 2014 and also POSIDON 2016 observations are included in Figure 9, right panels, where
 the most frequent in-cloud and clear sky RH_{ice} at these temperatures can be seen to increase with decreasing temperature as discussed by Rollins et al. (2016). But, in the Asian Monsoon measurements
 (Figure 10, right panels), RH_{ice} at the coldest temperatures is even higher above saturation than in
 the calmer tropics outside, the saturation is exceeded by about 20-50% below 195 K.
- 780 Therefore, we extend the conclusion of Rollins et al. (2016): taking saturation at the stratospheric entry of an air mass as set point for water vapor transport to the stratosphere, the transport is underes-

timated by $\sim 10\%$ in regions of weak convective activity. In convective regions the underestimation increased to 20-50% in our observations.

- 785 As mentioned earlier, the question of how much H₂O from convectively injected ice crystals is transported from above the CPT further into the lower stratosphere is a subject of recent research. In the study of Ueyama et al. (2018), it is concluded that over the Asian monsoon at 100 hPa, convection is the dominant driver of the localized H₂O and that nearly all of the convective enhancements in H₂O is due to the effect of convective humidification, while convectively detrained ice crystals have
 700 paglicible impost
- 790 negligible impact.

From our measurements in and above the Asian monsoon anticyclone, however, it is difficult to estimate to what extent cirrus cloud particles contribute to the amount of H_2O that might enter the stratosphere. Nevertheless, from Section 5.1 we know that the most frequent IWCs between about 365-400 K are 10 to 0.5 ppmv. Above 400 K, up to 415 K some overshoots with IWCs between

- 795 0.5 to 8 ppmv are detected (Figure 11, upper right panel). These IWCs are similar to the amount of gas phase H₂O, which is between 3 to 8 ppmv in this region above the CPT (see last Section). Such amounts of H₂O and IWC indicate that the air 'masses originate' from an altitude of about 15 km (Figure 3), which corresponds to the lower bound of the TTL (see Figure 2).
- The question is if the ice particles found in the upper TTL will further grow and sediment out in supersaturated conditions (dehydration) or evaporate and add H₂O to the gas phase H₂O in subsaturation (hydration, see Jensen et al., 2007; Schoeberl et al., 2018). Inspecting the in-cloud RH_{ice} (see Figure 12, upper right panel) related to the respective IWCs, it can be seen that the air in the overshoots \geq 400 K with the fairly high IWCs (0.5 to 8 ppmv) is subsaturated, while at lower altitudes both super- and subsaturation occur. This means that convective overshoots in the Asian
- 805 monsoon can locally contribute a significant amount of water that might be further transported into the stratosphere. Unfortunately, we can not say anything about the frequency of these events in this study.

Above the CPT in the surrounding tropics outside of the Asian monsoon, the in-cloud RH_{ice} at $\gtrsim 400$ K are also subsaturated. Here, the IWCs range only between 0.02 and 2 ppmv, with some single data points up to about 2 ppmv. The related H_2O is 1.5 to 3 ppmv.

Comparing gas phase H_2O and IWC above the Asian monsoon CPT (both 3-8 ppmv) with the surrounding tropical regions (1.5-3/0.02-2 ppmv) might be an indication by in situ observations that, driven by overshooting convection, the Asian monsoon could by an important source for transport

815 of H_2O to the stratosphere.

6 Global cirrus Nice climatology from satellite remote sensing

Though the new in situ cirrus climatologies presented in Section 4 represent a considerable data set from research aircraft from which substantial insights are gained, it is still a mixture of different me-

- 820 teorological situations encountered during various field campaigns that does not necessarily display a statistically representative overall picture of the distribution of cirrus cloud properties. For this purpose, long-term global satellite remote sensing observations would be the method of choice, though retrievals of cirrus microphysical properties also have their own difficulties and limitations. This is especially true for the ice crystal number N_{ice} , which is particularly challenging to estimate from
- 825

satellite remote sensing. The new retrieval method for Nice introduced in Section 2.2, DARDAR-Nice, has however demonstrated the ability to satisfactorily reproduce in situ ice concentrations (Sourdeval et al., 2018).

In this section, we show a global climatology of N_{ice} derived from DARDAR-Nice in the same presentation as the in situ $N_{\rm ice}$ climatology. Median $N_{\rm ice},\,25/75^{\rm th}$ and 10/90^{\rm th} percentiles are provided for a global data set. In addition, median N_{ice} are discussed for the Arctic region, the northern

830

Comparison of remote sensing and in situ $N_{\rm ice}$ 6.1

mid-latitudes, the tropics, the southern mid-latitudes and Antarctica.

As a base for reliable N_{ice} climatologies from satellite remote sensing, in situ measurements of PSDs (ice particle size distributions) from five campaigns of the Cirrus Guide II are used for an evaluation

- 835 of the DARDAR-Nice retrieval algorithm (see Section 2.2). This directly follows the evaluation of DARDAR-Nice presented in Sourdeval et al. (2018), a subset of campaigns from the Cirrus Guide II data set was used. This section therefore focuses on these five campaigns, which are nevertheless representative of a wide range of mid-latitude and tropical ice clouds (see Figure 1 and caption of Figure 13). Figure 13 (top panel) shows the N_{ice} frequencies of occurrence in 1K temperature
- 840 bins of the in situ measurements, the corresponding frequencies obtained from DARDAR-Nice are shown in the bottom panel. The latter are derived on the basis of two input parameters which are extracted from the in situ PSDs of the five campaigns listed in the figure caption, rather than being constrained from lidar-radar measurements during usual retrievals. This approach allows to identify inherent incompatibilities between the satellite retrieval assumptions and the in situ measurements,
- by assuming that the in situ PSD parameters are perfectly constrained by the lidar-radar. Therefore, 845 possible differences should only be attributed to other retrieval assumptions, such as the PSD shape. Sourdeval et al. (2018) showed that this approach is efficient for identifying algorithmic limitations while still being representative of actual satellite retrievals. Note that DARDAR-Nice provides concentrations of ice crystals $>5\mu m$, as it has not been evaluated for smaller sizes, while $N_{\rm ice}$ from in
- 850 situ observations includes sizes $> 3\mu m$; however, the resulting difference in the ice concentrations is

negligible. Also, the 1-Hz in situ PSDs are here sampled into 10-s averages to simulate the 1.7-km horizontal resolution of the DARDAR-Nice retrievals (assuming a flight speed of about 170 m s^{-1}).

The black solid and dotted lines of the in situ climatology indicate the 50, 25 and 75th percentiles, which agree well with those of the entire in situ climatology (see Figure 7, middle left panel). This demonstrates that the selected subset is statistically representative of the entire database (indeed, the entire N_{ice} climatology contains only one additional campaign, see Table 5). A satisfactory agreement between DARDAR-Nice and the in situ observations is also seen, demonstrating that DARDAR-Nice

N_{ice} retrievals very well match the in situ database for known size distibutions.

- A detailed comparison of the DARDAR-Nice percentiles (black solid and dashed lines) with those of the in situ observations, however, yields a small offset by a factor of 1.73, which is visualized in Figure 13 by the average medians of DARDAR-Nice (straight black line, 0.064 cm^{-3}) and in situ (straight blue line, 0.037 cm^{-3}). One reason for this offset lies in the method of the retrieval. The two parameters, N_0^* and D_m , calculated here from the in situ PSDs (usually from lidar-radar observations) are feeded into a predefined four-parameter gamma-modified function to calculate ice particle size
- distributions (PSDs, see Section 2.2). N_{ice} is then calculated by summing up individual ice concentrations over a grid of size bins distributed over the gamma-shaped PSD for the entire range of observed ice crystal sizes (usually, a continuous integration of the PSD is performed). However, in the in situ measurements, there are often PSDs that do not contain ice particles $< 20 \,\mu$ m, i.e. these size bins are empty. Such PSDs represent aged cirrus after the ice nucleation phase where the smaller ice crys-
- 870 tals have grown to larger sizes (see Section 4.3.1). DARDAR-Nice, however, assumes a modified Gamma distribution including all size bins, which partly explains the described positive offset. This behavior was also found by Wolf et al. (2019) who parameterized in situ-origin and liquid-origin cirrus from balloon-borne measurements by Gamma functions. It is of note that when subtracting the ice concentrations of the 'empty bins' of in situ PSDs from the retrieved DARDAR-Nice in a
- 875 direct intercomparison, the agreement of N_{ice} is improving (not shown here). This is an important finding, as gamma functions are often used to represent cirrus PSDs, both in remote sensing retrieval algorithms and in global models. On the other hand it has to be noted that an offset on the order of a factor of 1.73 is tolerable given the variability of N_{ice} (six orders of magnitude) and other possible error sources in the measurements, both in situ and remote sensing.
- Other assumptions on the PSD shape by the satellite remote sensing method might also contribute to this bias. The PSD shape indeed is provided by 4 parameters, 2 of which are fixed and 2 are retrieved (see Section 2.2). Delanoë et al. (2014) showed that the two fixed PSD parameters defined by Delanoë et al. (2005) and used in DARDAR-Nice might lead to a too steep representation of the small ice mode (i.e. too high N_{ice}) and should be updated in future algorithm versions. Also,
- 885 the bi-modality of the PSD towards temperature where growth processes become important is not accounted for and usually leads to small positive N_{ice} biases (Sourdeval et al., 2018). The cause of these assumptions are difficult to account for, as they most likely depend on the cloud-type and on the

thermodynamical environment, but they should on a first order be reasonably captured by the 1.73 adjustment factor. Higher Nice in DARDAR-Nice towards low temperatures are also not surprising,

- as Sourdeval et al. (2018) reported that the PSD parametrization used by DARDAR-Nice predicts 890 higher concentrations of small particles ($D_{ice} < 25 \,\mu m$) than the in situ comparative measurements. The overestimation increases with decreasing temperature, as the small ice particles dominate the PSDs more and more. This might be caused by a sharper representation of small ice concentrations in the PSD parametrization in comparison to the in situ small ice particle measurements below ~ 210
- 895 K.

A further expected difference between the data sets arises from their detection limits. The lowest N_{ice} that can be detected by the in situ instruments is 10^{-4} cm⁻³, for the highest N_{ice} there is no limitation. Detection limits for DARDAR-Nice depend on lidar-radar sensitivity but are also influenced by the instrumental resolution, which may cause specific features to remain undetected. This effect,

- 900 however, is represented in Fig. 13 by the 10-s sampling of 1-Hz in situ PSDs. Indeed, DARDAR-Nice misses the high N_{ice} in the temperature range 210-220 K, but these are young, line shaped contrails which are too small scale to be detected from satellite. Also, part of the thinnest cirrus are not represented by DARDAR-Nice. This will be further discussed in the next section describing the global N_{ice} distribution from DARDAR-Nice.
- 905 Overall, following the above-mentioned arguments, an overestimation within a factor of 2 in DARDAR-Nice by comparison to the in situ data set is expected, although there is still some uncertainty as to how this overestimation will propagate globally and over various ice cloud regimes. The ratio of 1.73 in N_{ice} between DARDAR-Nice and the in situ data set, found from Figure 13, should therefore be thought of as a minimum expected bias. Finally, the bias is expected to be stronger at
- 910 low temperatures than at high temperatures, due to the increased likelihood of missing bins as well as the higher importance of the representation of the small ice particle mode in the PSDs. A correction that depends on temperature, and possibly IWC, might therefore be optimal but a simpler first-order correction of 1.73 for all T and IWC range should here be sufficient for the needs of this study and considering the multitude of processes that can lead to this bias. Future studies will be required to 915
- precisely understand such inherent differences between satellite and in situ dataset.

6.2 Global cirrus N_{ice}

The global frequency distribution of N_{ice} from ten years of satellite observations is shown in Figure 14. The data was collected twice a day, approximately at midday and at midnight (satellite equator-crossing time is 1.30 am/pm), from June 2006 to December 2016. It should still be noted

920 that satellite retrievals are not continuously available throughout this period due to instrumental limitations and night-time retrievals are only available until 2011. This does not impact the conclusions presented here as diurnal cycles in N_{ice} are not considered this study. Such variations are also difficult to quantify lidar/radar products, although a diurnal signature has been investigated in the DARDAR-

Nice N_{ice} data for tropical anvils (Wall et al., 2020). Overall, the global satellite data set consists of

925 nea

nearly $2\times10^{10}~N_{\rm ice}$ retrievals. The color code represent frequencies of occurrence, the black contours the 25, 50 and 75^{\rm th} percentiles.

From the median (solid black line), a slight increase of N_{ice} with decreasing temperature is visible, which is somewhat different to the in situ median N_{ice} (solid blue line, from Figure 7, middle left panel), where no temperature dependence is found. The DARDAR-Nice temperature dependence

930 was already noted and discussed by Gryspeerdt et al. (2018). Another difference is that at T \gtrsim 210 K, the maximum N_{ice} of DARDAR-Nice reaches up to 100 cm⁻³, while the maximum in situ N_{ice} is only about 10 cm⁻³ (outside of young contrails).

We attribute the slightly increasing median $N_{\rm ice}$ with decreasing temperature to homogeneous ice nucleation events, because homogeneous ice nucleation rates increase with decreasing temperature,

935 but their appearance in space and time is transient, as discussed in Section 4.2.2. Thus, such events are difficult to find by research aircraft. Hence, these events are likely underrepresented in the aircraft observations.

The thinnest cirrus with low N_{ice} are represented by DARDAR-Nice for temperatures $\gtrsim 190$ K. At $\lesssim 190$ K, however, a decreasing detectability of thin cirrus becomes apparent in the DARDAR-

- 940 Nice climatology: in contrast to the in situ climatology, lesser or no thin cirrus are measured with decreasing temperature. The reason is that the colder the cirrus clouds, the smaller are the ice crystals and the lesser their IWC (Figure 10, top and bottom left panel). In the in situ N_{ice} climatology, in the TTL (≤ 205 K), an increased occurrence of very thin cirrus with N_{ice} around 0.001 cm⁻³ is reported in Section 4.3.1 and explained as aged cirrus consisting of only larger ice crystals. Though this
- 945 type of clouds is partly missing in the DARDAR-Nice climatology, the median N_{ice} decreases again at these temperatures. This might reflect the frequent presence of aged thin cirrus. Note, however, that the statistics at these temperatures is based on a lower number of observations than at higher temperatures (see Figure 15).

The median N_{ice} across all temperatures of the 10 years DARDAR-Nice climatology is about 0.100 cm⁻³ (Figure 14, plain cyan line). Adjusting this number to the offset factor of 1.73 between DARDAR-Nice and the in situ observations reported in the previous section yields an in situ average median of 0.056 cm⁻³ (dashed cyan line, see also Table 3). It is nevertheless important to note that such median is by construction computed assuming that all temperature bins have equal weight. When considering their actual relative occurrence as function of temperature (see Figure 15

and later discussion), the in situ adjusted global DARDAR-Nice median falls down to 0.046 cm^{-3} . This is still larger than the average median of 0.03 cm^{-3} reported from the N_{ice} in situ climatology (blue line; from Figure 7). Nevertheless, in the light of the offset between the data sets, the decreasing detectability of thin cirrus by DARDAR-Nice and other sources of error in both methods, the agreement between the DARDAR-Nice and the in situ N_{ice} climatology is good. 960 Altogether, both data sets have advantages and disadvantages: DARDAR-Nice has the advantage of the long, global time series where all atmospheric situations above the detection limit showing up at the times of observations are sampled. However, the thinnest cirrus at cold temperatures are not detected. Further, an offset is found between DARDAR-Nice and the in situ observations. On the other hand, the in situ N_{ice} climatology is based on a much smaller data set which is randomly distributed between different atmospheric situations. But, nearly the complete range of possible N_{ice}

can be detected by the in situ instruments.

For comparisons with other $N_{\rm ice}$ data sets or results from global models, the percentiles of the adjusted DARDAR-Nice as well as the average in situ $N_{\rm ice}$ are shown in Figure 14 and listed in Table 2 (note that an in situ data set slightly smaller than the data set presented here, but extended

970

0 with regard to Krämer et al. (2009) is already used for this purpose by Bacer et al., 2018; Penner et al., 2018; Righi et al., 2019).

The DARDAR-Nice data set does not only provide the N_{ice} distribution with temperature as shown in Figure 14, but, as it contains the entire spatial and temporal N_{ice} information, also the actual appearance of cirrus clouds in the N_{ice}-T parameter space. This is shown in Figure 15, where the occurrences of DARDAR-Nice N_{ice} retrievals are indicated by the colored background. It can be noticed that, in regions of high frequency of occurrence (for T ≥ 210 K), the agreement between DARDAR-Nice and the in situ data set is best.

From this portrayal it can be seen that the cirrus clouds do not spread evenly across all tempera-

- 980 tures, but that about 50% of the cirrus appear at temperatures above about 225 K (see also Table 3, temperature range of global 'most frequent' median N_{ice}). This clustering of cirrus at warmer temperatures is likely because in situ-origin and liquid-origin cirrus clouds are both found in this warmest and lowest cirrus layer, while at colder temperatures and higher altitudes only in situ-origin cirrus are present (see also Section 3.1 and Figure 3). This vertical structure of the cloud types is also reported
- 985 from observations by Krämer et al. (2016); Luebke et al. (2016); Wolf et al. (2018), from global modeling by Gasparini and Lohmann (2016), from satellite and global modeling by Gasparini et al. (2018) and also from 12 years ERA-interim data analysis in the North Atlantic region by Wernli et al. (2016).
- Gasparini et al. (2018) provide global relative frequencies of liquid-origin cirrus with respect to 990 temperature: more than 50% of the cirrus clouds in the range between 235 K and 220 K are of liquidorigin, decreasing rapidly to about 20% between 200 K and 180 K. These findings from satellite observations and global modeling are reasonably consistent with those from trajectory analysis by Wernli et al. (2016), who reported the relative frequencies of liquid-origin cirrus with respect to pressure to decrease from about 55% of the cirrus at 500 hPa (roughly 240 K), to 50, 30, 5% at 400,
- 995 300, 200 hPa (roughly 230, 215, 200 K; see Figure 2).

The analysis presented here provides the information that globally, half of the cirrus are in the lowest, warmest cirrus layer between about 220 K and 240 K, where, as mentioned above, about half

of the cirrus are of liquid-origin. This is of importance because of the differing radiative properties of in situ-origin and liquid-origin cirrus shown in Section 3.1.1 (see Figure 5): if the thick liquid-origin

cirrus clouds occur with such a high frequency, their strong cooling effect might exceed the much

1000

smaller warming effect of the in situ-origin cirrus which could lead to a general cooling by cirrus. Note that an analysis of $N_{\rm ice}$ in the IWC-T space, similarly to Figure 6, is shown in Figure 5 of

the SM. This figure shows a good climatological agreement between the satellite product and the Cirrus Guide II data set, with very similar distribution of N_{ice} in the IWC-T space. Differences could
be attributed to lack of statistics in Figure 6 (noisy patterns). However, one notable difference is the slope of the IWC-T relation, which appears much flatter in the satellite product than in the in situ data, as indicated by the density isolines.

6.3 Regional cirrus N_{ice}

- Lastly, we provide an impression of the regional variations of N_{ice} (see Table 3). To this end, subsets of the DARDAR-Nice data for five regional latitude bands between 90°, 67.7°, 23.3°, -67.7°, -90°, corresponding to the Arctic, Mid-lat North, the Tropics, Mid-lat South and Antarctica, are considered, and three different N_{ice} medians are computed for each region and for the global data set. A first median (labeled "all Temp.") corresponds to the median N_{ice} value across all temperatures and considering all temperature bins to be equiprobable, i.e. the same as used in Figure 14. A second
- 1015 (labeled "T-int weighted") represents the median using the relative occurrence of each temperature interval as a weight. The last ("most frequent") shows the most frequently appearing N_{ice}, considering those temperature intervals that contain 50% of the total N_{ice} occurrence; T-range denotes the respective minimum and maximum temperatures (note that in the in situ data set all three methods would yield to an identical median due to the flat temperature dependence).
- 1020 Comparing the three median N_{ice}, it is obvious that in most cases the 'T-int weighted' and 'most frequent' medians are smaller than those for 'all temperatures'. This behavior is found because the cirrus clouds appear often at warmer temperatures (see previous subsection) where N_{ice} is moderate, which can be seen particularly in the 'most frequent' medians, where the temperature range of the cirrus occurrence is given.

The global 'most frequent' median N_{ice} is 0.031 cm⁻³ (in comparison to 0.056 and 0.046 cm⁻³ for 'all temperatures' and 'T-int weighted') in the temperature range between 224-242 K (note that, to avoid representing mixed-phase clouds in the DADAR-Nice analysis, 242 K is chosen as maximum temperature). The lowest N_{ice} are found in the Arctic ('most frequent' median 0.016 cm⁻³ between

1030 230-242 K), which is most probably because the updrafts in the Arctic regions are generally lower than in other regions. In Antarctica, the 'most frequent' median is higher (0.029 cm⁻³ between 224 and 242 K). This could be traced back to stronger orographic uplifts in austral winter. The boreal and austral mid-latitude cirrus clouds are similar to the global, with 'most frequent' medians of 0.030 and 0.031 cm⁻³ in the temperature range of 227-242 K. This points to moderate updrafts on average,

1035 though regionally orography and convection can cause higher updrafts and high N_{ice} , which is also shown by Sourdeval et al. (2018).

Tropical cirrus clouds are different from those in the other regions. The 'most frequent' cirrus have the highest median (0.074 cm⁻³) and appear at much colder temperatures (197-221 K), i.e. higher altitudes. This finding corresponds to the in situ observations presented in Section 3.1 (see Figure 3)

1040 and also the DARDAR-Nice analyses of the vertical distribution of N_{ice} shown by Sourdeval et al. (2018). The reason is the strong convection prevailing in tropical regions, lifting liquid-origin clouds up to high altitudes and cold temperatures while homogeneously nucleating a large number of in situ ice crystals in addition (see also Section 5).

For all regions, except the tropics, more than half of the cirrus clouds are found at temperatures

1045 warmer than about 225 K and contain a considerable amount of liquid-origin cirrus, as discussed in the previous subsection. The same is true in the tropics, but for colder temperatures.

7 Summary and Conclusions

The Cirrus Guide II aims to represent cirrus clouds and their environment from the perspective of in situ and satellite remote sensing observations. To this end, an in situ database is created, which is

- 1050 based on measurements with state-of the art instrumentation and extended in comparison to earlier studies (Schiller et al., 2008; Krämer et al., 2009; Luebke et al., 2013; Krämer et al., 2016). The in situ database consists now of about 168 hours of ice water content IWC, 90/84 hours of ice crystal number concentration N_{ice} and R_{ice}, 96/186 hours of in-cloud and clear sky RH_{ice} and, correspondingly, 186 hours of clear sky water vapor H₂O. The measurements span the altitude range between 5
- 1055 and 20 km and cover the latitude band from 75 °North to 20 °South. The remote sensing database includes 10 years (2006-2016) of global recording of ice crystal number concentrations $N_{\rm ice}$. The main results form the evaluation of the two data sets are summarized in the following.

7.1 Characteristics and distribution of in situ-origin and liquid-origin cirrus (Section 3)

From the extended Cirrus Guide II in situ data set, we generally *confirm the typical characteristics of in situ-origin and liquid-origin cirrus* introduced by Krämer et al. (2016) with some additions (see Table 1):

- in situ-origin as well as liquid-origin cirrus consist of two sub-classes determined by the updraft (1:slow updrafts - few large ice crystals form from heterogeneous freezing; 2:fast updrafts
 - many small ice crystals nucleate homogeneously).
 New in this concept is that also liquid-origin cirrus differ in the two updraft regimes. Often, the two subclasses occur in succession and form a bimodal size distribution.
 - liquid-origin cirrus are thicker (higher IWC) than in situ-origin cirrus and are usually characterized by larger ice crystals.
 - The differences between the cirrus types are most pronounced in the formation phase of the clouds and are blurred with increasing lifetime due to ice crystal growth and sedimentation or additional ice formation.

In addition, we present a *picture of the distribution of cirrus with respect to altitude and latitude*, 1075 including an impression of the vertical structure of liquid-origin and in situ-origin cirrus (Figure 3):

- Across all latitudes, the thicker liquid-origin cirrus predominate at lower altitudes, while at higher altitudes the thinner in situ-origin cirrus prevail. In between, the two cirrus types overlap. This finding is in agreement with Luebke et al. (2016) and Wernli et al. (2016) for midlatitude and Wolf et al. (2018) for Arctic cirrus.
- 1080 Finally, a first estimate of the radiative characteristics of typical, specific and idealized in situ-origin and liquid-origin cirrus scenarios is given (Figure 5):

1070

- slow in situ-origin cirrus have a small optical depth (τ : 0.001 0.05), resulting in a slight net warming effect of not larger than about 1.5 W/m².
- the optical depth of fast in situ-origin cirrus is larger (τ : 0.05 1), but most of them are also warming (2-10 W/m²). The thickest fast in situ-origin cirrus at the lowest altitudes can change the sign of their net forcing, they switch to a slight cooling effect.
 - liquid-origin cirrus have large optical depths (τ : 1 12), and consequently exhibit a quite strong net cooling effect (-15 to -250 W/m²).

1090 7.2 Cirrus and humidity in the tropical tropopause layer (TTL) (Section 5)

The new in situ data set is extended by observations in the tropical TTL outside, but also, for the first time, inside of the Asian monsoon anticyclone. Therefore, we put special emphasis on the analysis of the TTL environment.

- 1095 *TTL cirrus clouds* (Figure 11):
 - Two types of most likely in situ formed cirrus are identified in slow large scale updrafts at low temperatures (T $\lesssim 205$ K, $\Theta\gtrsim 355$ K). The first is interpreted as young cirrus (N_{ice} around 0.1-1 cm⁻³ $\gtrsim 3$ µm diameter), the second as aged cirrus (N_{ice} around 0.001 cm⁻³ $\gtrsim 20$ µm diameter) where the smaller ice crystals have grown to larger sizes.
- The highest N_{ice} (up to 30 cm⁻³ around the cold point tropopause, CPT) and IWC (up to 1000 ppmv around the CPT) are found in deep convective systems in the Asian monsoon. Such systems represent massive liquid-origin cirrus (very high IWC) superimposed by a fresh strong homogeneous in situ ice nucleation event (many small ice crystals ≤ 20 µm) in fast updrafts.
- 1105 *TTL humidity* (Figure 12):
 - In the Asian monsoon, in-cloud and clear sky RH_{ice} is higher (often supersaturated around and above the CPT) than in the surrounding tropics. This is caused by a higher amount of H₂O in the Asian monsoon (most frequently 3 to 5 ppmv) in comparison to the tropics outside (1.5 to 3 ppmv).
- 1110 Taking saturation at the stratospheric entry of an air mass as set point for water vapor transport to the stratosphere, the transport is underestimated by ~ 10% in regions of weak convective activity (see also Rollins et al., 2016). In convective regions, the underestimation increased to 20-50% in our observations in the Asian monsoon.
- Convectively injected ice over the Asian monsoon CPT (Θ ≥ 400 K, RH_{ice} ≤ 100%) can
 locally contribute a significant amount of water (up to an IWC of 8 ppmv, in comparison to only 2 ppmv in the surrounding tropics) that might be evaporated and further transported into the stratosphere.

7.3 in situ and satellite climatologies (Sections 4 and 6)

Cirrus Guide II in situ cirrus and humidity climatologies:

– Median N_{ice} , R_{ice} and RH_{ice} are presented in the IWC-T parameter space (Figure 6).

Distinct patterns are found: N_{ice} form bands parallel to IWC, while R_{ice} are sorted diagonally in the IWC-T space. At tropical cold temperatures, supersaturations occur in thin as well as in thick cirrus.

1125 thick cirru

- The entire extended data set (Figure 7) is compared to the earlier studies (Figure 6, SM).

IWC: the median IWC and the core IWC band is the same in both data sets, showing that the in situ IWC measurement techniques are robust and that the IWC is a stable parameter describing cirrus clouds.

RH_{ice}: the overall picture of the in-cloud and clear sky RH_{ice} distributions has also not changed, demonstrating that high altitude water vapor measurements that were under discussion earlier have improved and stabilized.

 N_{ice} : an extended view is presented for N_{ice} , which is due to the better lower N_{ice} detection limit and a better mixture of dynamical situation. The new N_{ice} -T percentiles are lower and show no distinct temperature dependence (average 10%, median and 90% N_{ice} percentiles: 0.002, 0.03 and 0.3 cm⁻³) in comparison to the earlier observations, that show a slight decrease of N_{ice} with temperature and an average median N_{ice} of about 0.1 cm⁻³.

The in situ data set is subdivided into mid-latitude and tropical climatologies (Figures 8 and 9) and typical cirrus and humidity characteristics of the respective regions are presented.

DARDAR-Nice satellite global cirrus N_{ice} climatology

(Figures 14 and 15):

- A global climatology of N_{ice} from 10 years (2006-2016) of satellite observations is provided that can be used for comparison with global models or other data sets.
- 1145 The N_{ice} from satellite observations are validated by and adjusted to in situ measurements from a subset of five campaigns of the Cirrus Guide II.
 - The global median N_{ice} from satellite observations is almost 2 times higher than the in situ median and increases slightly with decreasing temperature.
 - N_{ice} medians sorted by geographical regions are highest in the tropics, followed by austral/boreal mid-latitudes, Antarctica and the Arctic.
 - In the satellite climatologies of N_{ice} , half of the cirrus are located in the lowest, warmest cirrus layer and contain a significant amount of liquid-origin cirrus. Their global median N_{ice} is 0.031 cm⁻³.

1150

1135

- Regarding the frequent appearance of liquid-origin cirrus together with strong cooling effect

1155

is a motivation to investigate their influence on the overall cirrus radiative feedback on climate in future studies.

Acknowledgments

This paper is dedicated to our colleague Cornelius Schiller, who passed away much too early in 2012. He was the initiator of the StratoClim project, including the tropical aircraft campaign. Through

1160 this campaign, he intended to complement his work on hydration and dehydration in the tropical tropopause layer, based on observations over Brazil, Australia and Africa (Schiller et al., 2009) with measurements over Asia. The campaign took place successfully out of Kathmandu, Nepal, in Summer 2017, after several years of planning and many hurdles that had to be overcome.

1165 References

- Afchine, A., Rolf, C., Costa, A., Spelten, N., Riese, M., Buchholz, B., Ebert, V., Heller, R., Kaufmann, S., Minikin, A., Voigt, C., Zöger, M., Smith, J., Lawson, P., Lykov, A., Khaykin, S., and Krämer, M.: Ice particle sampling from aircraft – influence of the probing position on the ice water content, Atmospheric Measurement Techniques, 11, 4015–4031, doi:10.5194/amt-11-4015-2018, 2018.
- 1170 Bacer, S., Sullivan, S. C., Karydis, V. A., Barahona, D., Krämer, M., Nenes, A., Tost, H., Tsimpidi, A. P., Lelieveld, J., and Pozzer, A.: Implementation of a comprehensive ice crystal formation parameterization for cirrus and mixed-phase clouds in the EMAC model (based on MESSy 2.53), Geoscientific Model Development, 11, 4021–4041, doi:10.5194/gmd-11-4021-2018, https://www.geosci-model-dev.net/11/4021/2018/, 2018.
- 1175 Baumgardner, D., Jonsson, H., Dawson, W., O'Connor, D., and Newton, R.: The cloud, aerosol and precipitation spectrometer (CAPS): A new instrument for cloud investigations, Atmos. Res., 59-60, 251–264, 2001.
 - Baumgardner, D., Abel, S. J., Axisa, D., Cotton, R., Crosier, J., Field, P., Gurganus, C., Heymsfield, A., Korolev,
 A., Krämer, M., Lawson, P., McFarquhar, G., Ulanowski, Z., and Um, J.: Cloud Ice Properties: In Situ Measurement Challenges; Chapter 9 of 'Ice Formation and Evolution in Clouds and Precipitation: Measurement
- 1180 and Modeling Challenges', Meteorol. Monographs, doi:10.1175/AMSMONOGRAPHS-D-16-0011.1, 2017. Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S., Sherwood, S., B., S., Zhang, X.Y.; Eds. Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.: Clouds and aerosols. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the
- 1185 Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, doi:0.1017/CBO9781107415324.016, 2013.
 - Brewer, A. W.: Evidence for a world circulation provided by the measurements of helium and water vapour distribution in the stratosphere, Quarterly Journal of the Royal Meteorological Society, 75, 351–363, doi:10.1002/qj.49707532603, https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49707532603, 1949.
- 1190 Buchholz, B., B., K., Smit, H., and Ebert, V.: Validation of an extractive, airborne, compact TDL spectrometer for atmospheric humidity sensing by blind intercomparison., Applied Physics B, 110, 249–262, 2013.
 - Campbell, J. R., Lolli, S., Lewis, J. R., Gu, Y., and Welton, E. J.: Daytime Cirrus Cloud Top-of-the-Atmosphere Radiative Forcing Properties at a Midlatitude Site and Their Global Consequences, Journal of Applied Meteorology and Climatology, 55, 1667–1679, doi:10.1175/JAMC-D-15-0217.1, 2016.
- 1195 Costa, A., Meyer, J., Afchine, A., Luebke, A., Günther, G., Dorsey, J. R., Gallagher, M. W., Ehrlich, A., Wendisch, M., Baumgardner, D., Wex, H., and Krämer, M.: Classification of Arctic, midlatitude and tropical clouds in the mixed-phase temperature regime, Atmospheric Chemistry and Physics, 17, 12219–12238, doi:10.5194/acp-17-12219-2017, 2017.
 - Delanoë, J., Protat, A., Testud, J., Bouniol, D., Heymsfield, A. J., Bansemer, A., Brown, P. R. A., and Forbes,
- 1200 R. M.: Statistical properties of the normalized ice particle size distribution, Journal of Geophysical Research: Atmospheres, 110, doi:10.1029/2004JD005405, https://agupubs.onlinelibrary.wiley.com/doi/abs/10. 1029/2004JD005405, 2005.
 - Delanoë, J. M. E., Heymsfield, A. J., Protat, A., Bansemer, A., and Hogan, R. J.: Normalized particle size distribution for remote sensing application, Journal of Geophysical Research: Atmo-

- 1205 spheres, 119, 4204–4227, doi:10.1002/2013JD020700, https://agupubs.onlinelibrary.wiley.com/doi/abs/10. 1002/2013JD020700, 2014.
 - Dinh, T., Podglajen, A., Hertzog, A., Legras, B., and Plougonven, R.: Effect of gravity wave temperature fluctuations on homogeneous ice nucleation in the tropical tropopause layer, Atmos. Chem. Phys. Discuss., 15, 8771–8799, doi:0.5194/acpd-15-8771-2015, 2015.
- 1210 Fahey, D., Gao, R., Möhler, O., Saathoff, H., Schiller, C., Ebert, V., Krämer, M., Peter, T., Amarouche, N., Avallone, L. M., Bauer, R., Bozóki, Z., Christensen, L. E., Davis, S. M., Durry, G., Dyroff, C., Herman, R. L., Hunsmann, S., Khaykin, S. M., Mackrodt, P., Meyer, J., Smith, J. B., Spelten, N., Troy, R. F., Vömel, H., Wagner, S., and Wienhold, F. G.: The AquaVIT-1 intercomparison of atmospheric water vapor measurement techniques, AMT, 7, 3177–3213, doi:10.5194/amtd-7-3177-2014, 2014.
- 1215 Finger, F., Werner, F., Klingebiel, M., Ehrlich, A., Jäkel, E., Voigt, M., Borrmann, S., Spichtinger, P., and Wendisch, M.: Spectral optical layer properties of cirrus from collocated airborne measurements – a feasibility study, Atmos. Chem. Phys. Discuss., 15, 19045–19077, doi:10.5194/acpd-15-19045-2015, 2015.
 - Fu, Q. and Liou, K.: Parameterization of the Radiative Properties of Cirrus Clouds, J. Atmos. Sci., 50, 2008–2025, doi:10.1175/1520-0469(1993)050<2008:POTRPO>2.0.CO;2, 1993.
- 1220 Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, Rev. Geophys., 47, 1–31, doi:10.1029/2008RG000267, 2009.
 - Gasparini, B. and Lohmann, U.: Why cirrus cloud seeding cannot substantially cool the planet, Journal of Geophysical Research: Atmospheres, 121, 4877–4893, doi:10.1002/2015JD024666, 2016.
 - Gasparini, B., Meyer, A., Neubauer, D., Münch, S., and Lohmann, U.: Cirrus Cloud Properties as Seen by
- 1225 the CALIPSO Satellite and ECHAM-HAM Global Climate Model, Journal of Climate, 31, 1983–2003, doi:10.1175/JCLI-D-16-0608.1, 2018.
 - Gryspeerdt, E., Sourdeval, O., Quaas, J., Delanoë, J., Krämer, M., and Kühne, P.: Ice crystal number concentration estimates from lidar–radar satellite remote sensing – Part 2: Controls on the ice crystal number concentration, Atmospheric Chemistry and Physics, 18, 14351–14370, doi:10.5194/acp-18-14351-2018, 2018.
- 1230 Heymsfield, A., Krämer, M., Luebke, A., Brown, P., Cziczo, D., Franklin, C., Lawson, P., Lohmann, U. Mc-Farquhar, G., Ulanowski, Z., and Van Tricht, K.: Ice Formation and Evolution in Clouds and Precipitation: Measurement and Modeling Challenges, Chapter 2: Cirrus Clouds, Eds. Baumgardner, D., McFarquhar, G. and Heymsfield, A., Meteor. Monogr., 58, doi:10.1175/AMSMONOGRAPHS-D-16-0010.1, 2017a.
- Heymsfield, A., Krämer, M., Wood, N. B., Gettelman, A., Field, P. R., and Liu, G.: Dependence of the Ice
 Water Content and Snowfall Rate on Temperature, Globally: Comparison of in Situ Observations, Satellite Active Remote Sensing Retrievals, and Global Climate Model Simulations, JOURNAL OF APPLIED METEOROLOGY AND CLIMATOLOGY, 56, 189–215, doi:10.1175/JAMC-D-16-0230.1, 2017b.
 - Jensen, E. and Pfister, L.: Transport and freeze-drying in the tropical tropopause layer, J. Geophys. Res., 109, doi:10.1029/2003JD004022, 2004.
- 1240 Jensen, E., Lawson, P., Baker, B., Pilson, B., Mo, Q., Heymsfield, A., Bansemer, A., Bui, T., McGill, M., Hlavka, D., Heymsfield, G., Platnick, S., Arnold, G., and Tanelli, S.: On the importance of small ice crystals in tropical anvil cirrus, Atmos. Chem. Phys., 9, 5519–5537, 2009.

Jensen, E., Diskin, G., Lawson, P. Lance, S., Bui, T., Hlavka, D., McGill, M., Pfister, L., Toon, O., and Gao, R.: Ice nucleation and dehydration in the Tropical Tropopause Layer, PNAS, 110, 2041–2046,

1245 doi:10.1073/pnas.1217104110, 2013a.

- Jensen, E., Lawson, P., Bergman, J. W., Pfister, L., Bui, T. P., and Schmitt, C. G.: Physical processes controlling ice concentrations in synoptically forced, midlatitude cirrus, J. Geophys. Res., 118, 5348–5360, doi:10.1002/jgrd.50421, 2013b.
- Jensen, E. J. and Ackerman, A. S.: Homogeneous aerosol freezing in the tops of high-altitude tropical
 cumulonimbus clouds, Geophysical Research Letters, 33, doi:10.1029/2005GL024928, https://agupubs.
 onlinelibrary.wiley.com/doi/abs/10.1029/2005GL024928, 2006.
 - Jensen, E. J., Ackerman, A. S., and Smith, J. A.: Can overshooting convection dehydrate the tropical tropopause layer?, Journal of Geophysical Research: Atmospheres, 112, 1407–1413, doi:10.1029/2006JD007943, 2007.
 Jensen, E. J., Pfister, L., and Bui, T. P.: Physical processes controlling ice concentrations in cold cir-
- 1255 rus near the tropical tropopause, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 117, doi:10.1029/2011JD017319, 2012.
 - Jensen, E. J., Pfister, L., Jordan, D. E., Bui, T. V., Ueyama, R., Singh, H. B., Thornberry, T. D., Rollins, A. W., Gao, R.-S., Fahey, D. W., Rosenlof, K. H., Elkins, J. W., Diskin, G. S., DiGangi, J. P., Lawson, R. P., Woods, S., Atlas, E. L., Rodriguez, M. A. N., Wofsy, S. C., Pittman, J., Bardeen, C. G., Toon, O. B., Kindel, B. C.,
- 1260 Newman, P. A., McGill, M. J., Hlavka, D. L., Lait, L. R., Schoeberl, M. R., Bergman, J. W., Selkirk, H. B., Alexander, M. J., Kim, J.-E., Lim, B. H., Stutz, J., and Pfeilsticker, K.: THE NASA AIRBORNE TROPI-CAL TROPOPAUSE EXPERIMENT High-Altitude Aircraft Measurements in the Tropical Western Pacific, BULLETIN OF THE AMERICAN METEOROLOGICAL SOCIETY, 98, 129+, doi:10.1175/BAMS-D-14-00263.1, 2017.
- 1265 Jones, H., Haywood, J., Marenco, F., O'Sullivan, D., Meyer, J., Thorpe, R., Gallagher, M., Krämer, M., Bower, K., Rädel, G., Rap, A., Woolley, A., Forster, P., and Coe, H.: A methodology for in-situ and remote sensing of microphysical and radiative properties of contrails as they evolve into cirrus, Atmospheric Chemistry and Physics, 12, 8157–8175, doi:10.5194/acp-12-8157-2012, cited By 1, 2012.

Joos, H.: Warm Conveyor Belts and Their Role for Cloud Radiative Forcing in the Extratropical Storm Tracks, Journal of Climate, 32, 5325–5343, doi:10.1175/JCLI-D-18-0802.1, 2019.

- Joos, H., Spichtinger, P., Reutter, P., and Fusina, F.: Influence of heterogeneous freezing on the microphysical and radiative properties of orographic cirrus clouds, Atmospheric Chemistry and Physics, 14, 6835–6852, doi:10.5194/acp-14-6835-2014, https://www.atmos-chem-phys.net/14/6835/2014/, 2014.
- Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: homogeneous freezing of super cooled aerosols, J. Geophys. Res., 107, 7433 7462, 2002.
- Kaufmann, S., Voigt, C., Heller, R., Jurkat-Witschas, T., Krämer, M., Rolf, C., Zöger, M., Giez, A., Buchholz, B., Ebert, V., Thornberry, T., and Schumann, U.: Intercomparison of midlatitude tropospheric and lower-stratospheric water vapor measurements and comparison to ECMWF humidity data, Atmospheric Chemistry and Physics, 18, 16729–16745, doi:10.5194/acp-18-16729-2018, https://www.atmos-chem-phys.net/ 18/16729/2018/, 2018.

- Kienast-Sjögren, E., Rolf, C., Seifert, P., Krieger, U. K., Luo, B. P., Krämer, M., and Peter, T.: Climatological and radiative properties of midlatitude cirrus clouds derived by automatic evaluation of lidar measurements, Atmospheric Chemistry and Physics, 16, 7605–7621, doi:10.5194/acp-16-7605-2016, 2016.
- Korolev, A.: Limitations of the Wegener–Bergeron–Findeisen Mechanism in the Evolution of Mixed-Phase
 Clouds , J. Atmos. Sci., 64, 3372–3375, doi:10.1175/JAS4035.1, 2007.
 - Korolev, A. and Field, P. R.: Assessment of the performance of the inter-arrival time algorithm to identify ice shattering artifacts in cloud particle probe measurements, Atmospheric Measurement Techniques, 8, 761– 777, doi:10.5194/amt-8-761-2015, https://www.atmos-meas-tech.net/8/761/2015/, 2015.
 - Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N.,
- 1290 Borrman, S., de Reus, M., and Spichtinger, P.: Ice supersaturations and cirrus cloud crystal numbers, ACP, 9, 3505–3522, 2009.
 - Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Meyer, J., Zoeger, M., Smith, J., Herman,
 R. L., Buchholz, B., Ebert, V., Baumgardner, D., Borrmann, S., Klingebiel, M., and Avallone, L.: A micro-physics guide to cirrus clouds Part 1: Cirrus types, Atmospheric Chemistry and Physics, 16, 3463–3483,

doi:10.5194/acp-16-3463-2016, 2016.

- Kübbeler, M., Hildebrandt1, M., Meyer, J., Schiller, C., Hamburger, T., Jurkat, T., Minikin, A., Petzold, A., Rautenhaus, M., Schlager, H., chumann, U., Voigt, C., Spichtinger, P., Gayet, J.-F., Gourbeyre, C., and Krämer, M.: Thin and subvisible cirrus and contrails in a subsaturated environment, ACP, 11, 5853–5865, doi:10.5194/acp-11-5853-2011, 2011.
- 1300 Lawson, R., O'Connor, D., Zmarzly, P., Weaver, K., Baker, B., Mo, Q., and Jonsson, H.: The 2D-S (Stereo) probe: Design and preliminary tests of a new airborne, high speed, high resolution particle Imaging probe, J. Atmos. Ocean. Technol., 23, 1462 1477, 2006.
 - Lawson, R. P., Woods, S., Jensen, E., Erfani, E., Gurganus, C., Gallagher, M., Connolly, P., Whiteway, J., Baran, A. J., May, P., Heymsfield, A., Schmitt, C. G., McFarquhar, G., Um, J., Protat, A., Bailey, M.,
- 1305 Lance, S., Muehlbauer, A., Stith, J., Korolev, A., Toon, O. B., and Krämer, M.: A Review of Ice Particle Shapes in Cirrus formed In Situ and in Anvils, Journal of Geophysical Research: Atmospheres, 124, 10049–10090, doi:10.1029/2018JD030122, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/ 2018JD030122, 2019.
 - Luebke, A., Avallone, L., Schiller, C., Meyer, J., Rolf, C., and Krämer, M.: Ice water content of Arctic, midlati-
- tude, and tropical cirrus Part 2: Extension of the database and new statistical analysis, ACP, 13, 6447–6459, 2013.
 - Luebke, A. E., Afchine, A., Costa, A., Grooss, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Krämer, M.: The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, Atmospheric Chemistry and Physics, 16, 5793–5809, doi:10.5194/acp-16-5793-2016,
- 1315 2016.
 - May, R.: Open path, near infrared tunable diode laser spectrometer for atmospheric measurements of H₂O,
 J. Geophys. Res., 103, 19 161 19 172, doi:10.1029/98jd01678, 1998.
- McFarquhar, G., Junshik, U., Freer, M., Baumgardner, D., Kok, G., and Mace, G.: The Importance of Small Ice Crystals to Cirrus Properties: Observations from the Tropical Warm Pool International Cloud Experiment
- 1320 (TWP-ICE), Geophys. Res. Lett., 34, doi:10.1029/2007GL029865, 2007.

Meyer, J.: Ice Crystal Measurements with the New Particle Spectrometer NIXE-CAPS, Schriften des Forschungszentrums Jülich. Reihe Energie und Umwelt / Energy and Environment; 160; ISBN:9783893368402, 2012.

Meyer, J., Rolf, C., Schiller, C., Rohs, S., Spelten, N., Afchine, A., Zöger, M., Sitnikov, N., Thornberry, T.,

1325 Rollins, A., Gao, R., Bozoki, Z., Tatrai, D., Buchholz, B., Ebert, V., Mackrodt, P., Möhler, O., Saathoff, H., Rosenlof, K., and Krämer, M.: Two decades of water vapor measurements with the FISH fluorescence hygrometer: A review., ACP, 15, 8521–8538, doi:10.5194/acp-15-8521-2015, 2015, 2015.

Mitchell, D. L., Garnier, A., Pelon, J., and Erfani, E.: CALIPSO (IIR–CALIOP) retrievals of cirrus cloud iceparticle concentrations, Atmospheric Chemistry and Physics, 18, 17 325–17 354, doi:10.5194/acp-18-17325-2018, https://www.atmos-chem-phys.net/18/17325/2018/, 2018.

- Muhlbauer, A., Ackerman, T., Comstock, J., Diskin, G., Evans, S., Lawson, R., and Marchand, R.: Impact of large-scale dynamics on the microphysical properties of midlatitude cirrus, Journal of Geophysical Research: Atmospheres, 119, 3976–3996, doi:10.1002/2013JD020035, 2014.
 - Murphy, D. M.: Rare temperature histories and cirrus ice number density in a parcel and a one-dimensional
- 1335 model, Atmospheric Chemistry and Physics, 14, 13013–13022, doi:10.5194/acp-14-13013-2014, https://www.atmos-chem-phys.net/14/13013/2014/, 2014.
 - Pan, L. L., Bowman, K. P., Atlas, E. L., Wofsy, S. C., Zhang, F., Bresch, J. F., Ridley, B. A., Pittman, J. V., Homeyer, C. R., Romashkin, P., and Cooper, W. A.: The Stratosphere-Troposphere Analyses of Regional Transport 2008 (START08) Experiment, Bulletin of the American Meteorological Society, 91, 327–342,
- 1340 doi:10.1175/2009BAMS2865.1, 2010.

1330

- Pan, L. L., Atlas, E. L., Salawitch, R. J., Honomichl, S. B., Bresch, J. F., Randel, W. J., Apel, E. C., Hornbrook,
 R. S., Weinheimer, A. J., Anderson, D. C., Andrews, S. J., Baidar, S., Beaton, S. P., Campos, T. L., Carpenter,
 L. J., Chen, D., Dix, B., Donets, V., Hall, S. R., Hanisco, T. F., Homeyer, C. R., Huey, L. G., Jensen, J. B.,
 Kaser, L., Kinnison, D. E., Koenig, T. K., Lamarque, J.-F., Liu, C., Luo, J., Luo, Z. J., Montzka, D. D., Nicely,
- 1345 J. M., Pierce, R. B., Riemer, D. D., Robinson, T., Romashkin, P., Saiz-Lopez, A., Schauffler, S., Shieh, O., Stell, M. H., Ullmann, K., Vaughan, G., Volkamer, R., and Wolfe, G.: Convective Transport of Active Species in the Tropics (CONTRAST) experiment, Bull. Amer. Meteor. Soc., 98, 106–128, doi:10.1175/BAMS-D-14-00272.1, 2017.
 - Penner, J. E., Zhou, C., Garnier, A., and Mitchell, D. L.: Anthropogenic Aerosol Indirect Effects in Cirrus
- Clouds, Journal of Geophysical Research: Atmospheres, 123, 11,652–11,677, doi:10.1029/2018JD029204, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD029204, 2018.
 - Peter, T., Marcolli, C., Spichtinger, P., Corti, T., Baker, M., and Koop, T.: When dry air is too humid, Science, 314, 1399 1401, 2006.
 - Petzold, A., Krämer, M., Neis, P., Rolf, C., Rohs, S., Berkes, F., Smit, H. H., Gallagher, M., Beswick, K.,
- Lloyd, G., Baumgardner, D., Spichtinger, P., Nédélec, P., Ebert, V., Buchholz, B., Riese, M., and Andreas,
 W.: Upper tropospheric water vapour and its interaction with cirrus clouds as seen from IAGOS longterm routine in-situ observations, Faraday Discussions, 200, 229–249, doi:10.1039/C7FD00006E, http://dx.doi.org/10.1039/C7FD00006E, 2017.

Ploeger, F., Günther, G., Konopka, P., Fueglistaler, S., Müller, R., Hoppe, C., Kunz, A., Spang, R., Grooß,

1360 J.-U., and Riese, M.: Horizontal water vapor transport in the lower stratosphere from subtropics to

high latitudes during boreal summer, Journal of Geophysical Research: Atmospheres, 118, 8111–8127, doi:10.1002/jgrd.50636, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/jgrd.50636, 2013.

- Podglajen, A., Bui, T. P., Dean-Day, J. M., Pfister, L., Jensen, E. J., Alexander, M. J., Hertzog, A., Kärcher, B., Plougonven, R., and Randel, W. J.: Small-Scale Wind Fluctuations in the Tropical Tropopause Layer from
- 1365 Aircraft Measurements: Occurrence, Nature, and Impact on Vertical Mixing, Journal of the Atmospheric Sciences, 74, 3847–3869, doi:10.1175/JAS-D-17-0010.1, 2017.
 - Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., and Forster, P.: Impact of uncertainties in atmospheric mixing on simulated UTLS composition and related radiative effects, JOURNAL OF GEO-PHYSICAL RESEARCH-ATMOSPHERES, 117, doi:10.1029/2012JD017751, 2012.
- 1370 Righi, M., Hendricks, J., Lohmann, U., Gerhard Beer, C., Hahn, V., Heinold, B., Heller, R., Krämer, M., Rolf, C., Tegen, I., and Voigt, C.: A new approach to simulate aerosol effects on cirrus clouds in EMAC v2.54, GMD Discussion, xx, xxx, doi:xxx, 2019.
 - Rolf, C., Vogel, B., Hoor, P., Afchine, A., Günther, G., Krämer, M., Müller, R., Müller, S., Spelten, N., and Riese, M.: Water vapor increase in the lower stratosphere of the Northern Hemisphere due to the Asian
- 1375 monsoon anticyclone observed during the TACTS/ESMVal campaigns, Atmospheric Chemistry and Physics,
 18, 2973–2983, doi:10.5194/acp-18-2973-2018, https://www.atmos-chem-phys.net/18/2973/2018/, 2018.
 - Rollins, A. W., Thornberry, T. D., Gao, R. S., Smith, J. B., Sayres, D. S., Sargent, M. R., Schiller, C., Krämer, M., Spelten, N., Hurst, D. F., Jordan, A. F., Hall, E. G., Vömel, H., Diskin, G. S., Podolske, J. R., Christensen, L. E., Rosenlof, K. H., Jensen, E. J., and Fahey, D. W.: Evaluation of UT/LS hygrometer accuracy by
- 1380 intercomparison during the NASA MACPEX mission, JGR, 119, 1915–1935, doi:10.1002/2013JD020817, 2014.
 - Rollins, A. W., Thornberry, T. D., Gao, R. S., Woods, S., Lawson, R. P., Bui, T. P., Jensen, E. J., and Fahey, D. W.: Observational constraints on the efficiency of dehydration mechanisms in the tropical tropopause layer, Geophysical Research Letters, 43, 2912–2918, doi:10.1002/2016GL067972, 2016.
- 1385 Sassen, K., Wang, Z., and Liu, D.: Global distribution of cirrus clouds from CloudSat/Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) measurements, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 113, doi:10.1029/2008JD009972, 2008.
 - Schiller, C., Krämer, M., Afchine, A., Spelten, N., and Sitnikov, N.: Ice water content in Arctic, midlatitude and tropical cirrus, J. Geophys. Res., 113, D24208, doi:10.1029/2008JD010342., 2008.
- 1390 Schiller, C., Grooß, J.-U., Konopka, P., Plöger, F., Silva dos Santos, F. H., and Spelten, N.: Hydration and dehydration at the tropical tropopause, Atmospheric Chemistry and Physics, 9, 9647–9660, doi:10.5194/acp-9-9647-2009, 2009.
 - Schoeberl, M., Jensen, E., Pfister, L., Ueyama, R., Wang, T., Selkirk, H., Thornberry, T., Dessler, A., and Avery,
 M.: Water Vapor, Clouds, and Saturation in the Tropical Tropopause Layer, Journal of Geophysical Research:
 Atmospheres, accepted article, 1–30, doi:10.1029/2018JD029849, 2019.
 - Schoeberl, M. R., Jensen, E. J., Pfister, L., Ueyama, R., Avery, M., and Dessler, A. E.: Convective Hydration of the Upper Troposphere and Lower Stratosphere, Journal of Geophysical Research: Atmospheres, 123, 4583–4593, doi:10.1029/2018JD028286, 2018.

Sitnikov, N., Yushkov, V., Afchine, A., Korshunov, L., Astakhov, V., Ulanovskii, A., Krämer, M., Mangold,

- 1400 A., Schiller, C., and Ravegnani, F.: The FLASH Instrument for Water Vapor Measurements on Board the High-Altitude Airplane, Instruments and Experimental Techniques, 50, 113121, 2007.
 - Solomon, S., Rosenlof, K. H., Portmann, R. W., Daniel, J. S., Davis, S. M., Sanford, T. J., and Plattner, G.-K.: Contributions of Stratospheric Water Vapor to Decadal Changes in the Rate of Global Warming, SCIENCE, 327, 1219–1223, doi:10.1126/science.1182488, 2010.
- 1405 Sourdeval, O., Gryspeerdt, E., Krämer, M., Goren, T., Delanoë, J., Afchine, A., Hemmer, F., and Quaas, J.: Ice crystal number concentration estimates from lidar–radar satellite remote sensing – Part 1: Method and evaluation, Atmospheric Chemistry and Physics, 18, 14 327–14 350, doi:10.5194/acp-18-14327-2018, 2018. Spichtinger, P. and Cziczo, D. J.: Impact of heterogeneous ice nuclei on homogeneous freezing events, J. Geo-

phys. Res., 115, D14208, doi:10.1029/2009JD012168, 2010.

- 1410 Spichtinger, P. and Krämer, M.: Tropical tropopause ice clouds: a dynamic approach to the mystery of low crystal numbers, Atmos. Chem. Phys., 13, doi:10.5194/acp-13-9801-2013, 2013.
 - Spreitzer, E. J., Marschalik, M. P., and Spichtinger, P.: Subvisible cirrus clouds a dynamical system approach, Nonlinear Processes in Geophysics, 24, 307–328, doi:10.5194/npg-24-307-2017, https://www.nonlin-processes-geophys.net/24/307/2017/, 2017.
- 1415 Thornberry, T., Rollins, A., Avery, M., Woods, S., Lawson, R., Bui, T., and Gao, R.-S.: Ice water contentextinction relationships and effective diameter for TTL cirrus derived from in situ measurements during ATTREX 2014, J. Geophys. Res., 122, 4494–4507, doi:10.1002/2016JD025948, 2017.
 - Thornberry, T. D., Rollins, A. W., Gao, R. S., Watts, L. A., Ciciora, S. J., McLaughlin, R. J., and Fahey, D. W.: A two-channel, tunable diode laser-based hygrometer for measurement of water vapor and cirrus cloud ice
- water content in the upper troposphere and lower stratosphere, Atmospheric Measurement Techniques, 8, 211–224, doi:10.5194/amt-8-211-2015, https://www.atmos-meas-tech.net/8/211/2015/, 2015.
 - Ueyama, R., Jensen, E. J., and Pfister, L.: Convective Influence on the Humidity and Clouds in the Tropical Tropopause Layer During Boreal Summer, Journal of Geophysical Research: Atmospheres, 123, 7576–7593, doi:10.1029/2018JD028674, 2018.
- 1425 Urbanek, B., Groß, S., Schäfler, A., and Wirth, M.: Determining stages of cirrus evolution: a cloud classification scheme, Atmospheric Measurement Techniques, 10, 1653–1664, doi:10.5194/amt-10-1653-2017, https://www.atmos-meas-tech.net/10/1653/2017/, 2017.
 - Urbanek, B., Groß, S., Wirth, M., Rolf, C., Krämer, M., and Voigt, C.: High Depolarization Ratios of Naturally Occurring Cirrus Clouds Near Air Traffic Regions Over Europe, Geophysical Research Letters, 45, 13,166–
- 1430 13,172, doi:10.1029/2018GL079345, 2018.
 - Vali, G., DeMott, P. J., Möhler, O., and Whale, T. F.: Technical Note: A proposal for ice nucleation terminology, Atmospheric Chemistry and Physics, 15, 10263–10270, doi:10.5194/acp-15-10263-2015, https: //www.atmos-chem-phys.net/15/10263/2015/, 2015.
 - Vogel, B., Günther, G., Müller, R., Grooß, J.-U., Afchine, A., Bozem, H., Hoor, P., Krämer, M., Müller, S.,
- 1435 Riese, M., Rolf, C., Spelten, N., Stiller, G. P., Ungermann, J., and Zahn, A.: Long-range transport pathways of tropospheric source gases originating in Asia into the northern lower stratosphere during the Asian monsoon season 2012, Atmospheric Chemistry and Physics, 16, 15 301–15 325, doi:10.5194/acp-16-15301-2016, https://www.atmos-chem-phys.net/16/15301/2016/, 2016.

Voigt, C., Schumann, U., Minikin, A., Abdelmonem, A., Afchine, A., Borrmann, S., Boettcher, M., Bucuchholz,

- B., Bugliaro, L., Costa, A., Curtius, J., Dollner, M., Doernbrack, A., Dreiling, V., Ebert, V., Ehrlich, A., Fix, A., Forster, L., Frank, F., Fuetterer, D., Giez, A., Graf, K., Grooss, J.-U., Gross, S., Heimerl, K., Heinold, B., Hueneke, T., Jaervinen, E., Jurkat, T., Kaufmann, S., Kenntner, M., Klingebiel, M., Klimach, T., Kohl, R., Krämer, M., Krisna, T. C., Luebke, A., Mayer, B., Mertes, S., Molleker, S., Petzold, A., Pfeilsticker, K., Port, M., Rapp, M., Reutter, P., Rolf, C., Rose, D., Sauer, D., Schaefer, A., Schlage, R., Schnaiter, M., Schneider,
- 1445 J., Spelten, N., Spichtinger, P., Stock, P., Walser, A., Weigel, R., Weinzierl, B., Wendisch, M., Werner, F., Wernli, H., Wirth, M., Zahn, A., Ziereis, H., and Zöger, M.: ML-Cirrus the airborne experiment on natural cirrus and contrail cirrus with the high-altitude long-range research aircraft HALO, Bulletin of the American Meteorological Society, 98, 271–288, doi:10.1175/BAMS-D-15-00213.1, 2017.
- Wall, C. J., Norris, J. R., Gasparini, B., Smith, William L., J., Thieman, M. M., and Sourdeval, O.: Observational Evidence that Radiative Heating Modifies the Life Cycle of Tropical Anvil Clouds, J. Clim., pp. 1–68, doi:10.1175/JCLI-D-20-0204.1, 2020.
 - Wendisch, M., Yang, P., and Pilewskie, P.: Effects of ice crystal habit on thermal infrared radiative properties and forcing of cirrus, Journal of Geophysical Research, 112, doi:10.1029/2006JD007899, 2007.
- Wendisch, M., Poeschl, U., Andreae, M. O., Machado, L. A. T., Albrecht, R., Schlager, H., Rosenfeld, D.,
 Martin, S. T., Abdelmomonem, A., Afchine, A., Araujo, A. C., Artaxo, P., Aufmhoff, H., Barbosa, H. M. J.,
 Borrmann, S., Braga, R., Buchholz, B., Cecchini, M. A., Costa, A., Curtius, J., Dollner, M., Dorf, M., Dreiling, V., Ebert, V., Ehrlich, A., Ewald, F., Fisch, G., Fix, A., Frank, F., Futterer, D., Heckl, C., Heidelberg, F.,
 Hueneke, T., Jakel, E., Jarvinen, E., Jurkat, T., Kanter, S., Kaestner, U., Kenntner, M., Kesselmeier, J., Klimach, T., Knecht, M., Kohl, R., Koelling, T., Krämer, M., Krueger, M., Krisna, T. C., Lavric, J. V., Longo,
- 1460 K., Mahnke, C., Manzi, A. O., Mayer, B., Mertes, S., Minikin, A., Molleker, S., Munch, S., Nillius, B., Pfeilsticker, K., Pohlker, C., Roiger, A., Rose, D., Rosenowow, D., Sauer, D., Schnaiter, M., Schneider, J., Schulz, C., de Souza, R. A. F., Spanu, A., Stock, P., Vila, D., Voigt, C., Walser, A., Walter, D., Weigel, R., Weinzierl, B., Werner, F., Yamasoe, M. A., Ziereis, H., Zinner, T., and Zoeger, M.: ACRIDICON-CHUVA CAMPAIGN Studying Tropical Deep Convective Clouds and Precipitation over Amazonia Using the New
- German Research Aircraft HALO, BULLETIN OF THE AMERICAN METEOROLOGICAL SOCIETY, 97, 1885–1908, doi:10.1175/BAMS-D-14-00255.1, 2016.
 - Wernli, H., Boettcher, M., Joos, H., Miltenberger, A. K., and Spichtinger, P.: A trajectory-based classification of ERA-Interim ice clouds in the region of the North Atlantic storm track, Geophysical Research Letters, 43, 6657–6664, doi:10.1002/2016GL068922, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/ 2016GL 060022, 2016

```
1470 2016GL068922, 2016.
```

- Wolf, V., Kuhn, T., Milz, M., Voelger, P., Krämer, M., and Rolf, C.: Arctic ice clouds over northern Sweden: microphysical properties studied with the Balloon-borne Ice Cloud particle Imager B-ICI, Atmospheric Chemistry and Physics, 18, 17 371–17 386, doi:10.5194/acp-18-17371-2018, 2018.
- Wolf, V., Kuhn, T., and Krämer, M.: On the dependence of cirrus parametrizations on the cloud origin, Geo physical Research Letters, 46, 12565–12571, doi:10.1029/2019GL083841, 2019.
 - Woods, S., Lawson, R. P., Jensen, E., Bui, T. P., Thornberry, T., Rollins, A., Pfister, L., and Avery, M.: Microphysical Properties of Tropical Tropopause Layer Cirrus, Journal of Geophysical Research: Atmospheres, 123, 6053–6069, doi:10.1029/2017JD028068, 2018.

Zondlo, M. A., Paige, M. E., Massick, S. M., and Silver, J. A.: Vertical cavity laser hygrometer for the National

1480 Science Foundation Gulfstream-V aircraft, J. Geophys. Res., 115, D02309, doi:10.1029/2010JD014445, 2010.



Figure 1. Aircraft flight paths during the 24 campaigns listed in Table 5. 185 flights, 192 h IWC measurements, 90 h N_{ice} , 84 h R_{ice} , 116/331 h RH_{ice} in/outside cirrus (campaign names: red – GfD Learjet, green – BAe-146, blue – Geophysica, purple – WB-57, black – HALO, light-purple – Global Hawk, cyan – GV HIAPER, orange – Spec Learjet). In comparison: Schiller et al. (2008) / Krämer et al. (2009) – 52 flights, 27 h IWC measurements, 8.5 h N_{ice} , 8.5 h R_{ice} , 10/16 h RH_{ice} in/outside cirrus.



Figure 2. Left panel: Temperature vs. altitude inside of cirrus clouds (adopted from Schiller et al., 2008), middle panel: Temperature vs. potential temperature Θ , right panel: clear sky Θ vs. altitude. Plotted are 28 flights from the data base, with blueish colors for Arctic, greenish colors for mid-latitude and reddish colors for tropical observations. Magenta lines: range of the TTL (tropical tropopause layer) and tropical CPT (cold point tropopause) (after Fueglistaler et al., 2009).



Figure 3. Cirrus cloud distribution with latitude and altitude of **Left column:** Ice water content (IWC, top panel), ice crystal number (N_{ice} , size > 3 μ m diameter, middle panel) and mean mass radius (R_{ice} , calculated from IWC/ N_{ice} , bottom panel). **Right column:** In-cloud and clear sky relative humidity wrt ice (RH_{ice} , top and middle panel), and clear sky water vapor volume mixing ratio (H_2O , bottom panel). The field campaigns are listed in Table 4, data evaluation methods and detection ranges of the parameters are described in Appendix A. The color codes range from yellow to blue with increasing amount of the respective parameter; for R_{ice} , the color code is reversed to indicate that high / low N_{ice} belong to small / large R_{ice} ; note that the data points are plotted in the order of the colors from yellow to blue; frequencies of occurence of the parameters can be seen in Figure 7.

Typical characteristics of cirrus types						
in the initial stage						
ORIGIN IWC N _{ice} R _{ice} weather system						
slow updraft (heterogeneous ice nucl.)						
IN-SITU	low	few	large			
LIQUID	high	more	larger	frontals systems (WCBs)		
fast updraft (homogeneous ice nucl.)						
IN-SITU	high	many	small			
LIQUID	high	more	small & larger	gravity waves, convection		

slow updraft: $\lesssim 10$ cm/s; fast updraft: $\gtrsim 10$ cm/s (Kärcher and Lohmann 2002) Krämer et al. 2016) IWC high/low: above/below the IWC median (see Figure 7).

 $N_{\rm ice}$ few/more/many: below/in-between/above the 10 and 90% $N_{\rm ce}$ percentiles (see Figure 7).

 $R_{\rm ice}$ small/large/larger:

ice particles $\lesssim 20 \mu m$ dominate the PSD /

ice particles $\gtrsim 20 \mu m$ dominate the PSD, max. size several hundred μm diameter / ice particles $\gtrsim 20 \mu m$ dominate the PSD, max. size up to thousand μm diameter, PSD: particle size distribution.





Figure 4. Relation between ice crystal concentration N_{ice} and mean mass radius R_{ice} , color coded by the ice water content IWC in mg/m⁻³, from ~87 h of cirrus cloud observations (R_{ice} is calculated by dividing IWC/ N_{ice}). The thin black lines are isolines of IWC in ppmv (in the order of the legend). The scheme at the right side illustrates the partitioning of the clouds between 'liquid origin' and 'in situ origin': the thickest cirrus (blue points) are of 'liquid origin', the thinnest (yellow points) of 'in situ origin'. As the thickness decreases, the portion of 'liquid origin' cirrus becomes smaller and smaller while more and more 'in situ origin' cirrus appear.



Figure 5. Simulated radiative forcing versus optical depth for exemplary 'in situ origin' slow and fast updraft (light and dark green dots) as well as 'liquid origin' (blue dots) cirrus; the idealized scenarios are summarized in the table.

Description of the idealized scenarios: a temperature profile is prescribed representative for midlatitude conditions, the surface temperature is set to 288 K (at p = 1000 hPa, p exponentially decreases with height). The vertical profile of relative humidity over ice is also established, with a saturated layer of 1000 m thickness, constant subsaturation below the layer, and a strong decrease in humidity towards the stratosphere. A cirrus cloud with constant ice mass and number concentration is placed in the saturated layer. From the measurements (see Figure 4), typical values of N_{ice}, R_{ice} and IWC are chosen for the three cirrus types (see table above). The vertical profiles are adjusted with respect to tropopause height (z tropop.) and placement of the cirrus layers in accordance with the cirrus types. For the radiative transfer calculations, the well-known two-stream radiative transfer model for ice particles by Fu and Liou (1993), with 6 bands in the solar and 12 bands in the thermal infrared regime, is used. The simulations are realized for a geographic latitude of $\varphi = 50^{\circ}$, solar surface albedo of 0.3, infrared surface emissivity of 1 and solar constant S = 1340 W/m²; we assume equinox conditions (e.g. end of March) at local time t = 12 h (for these settings, we use the modified model of Joos et al., 2014). The net cloud radiative forcing (CRF) is calculated using the fluxes at the top of atmosphere in the short wave and long wave ranges in comparison with a clear sky case.



Figure 6. Median N_{ice} , R_{ice} and RH_{ice} in intervals in the IWC-T parameter space (five iwc intervals per order of magnitude, 1K temperature intervals). The color codes of N_{ice} and RH_{ice} range from yellow to blue with increasing amount of the respective parameter; for R_{ice} , the color code is reversed to indicate that high / low N_{ice} belong to small / large R_{ice} . Black solid/dotted lines: median, min/max IWC of the core IWC band of Schiller et al. (2008). The variability of the parameters (25, 50 (median) and 75 % percentiles) is shown in Figures 1–3 of the supplementary material.



Figure 7. Frequencies of occurrence in dependence on temperature, binned in 1K intervals of: Ice Water Content (**IWC**; top left panel; black solid/dotted lines: median, min/max IWC of the core IWC band of Schiller et al. (2008)). Ice crystal number (N_{ice} , size > 3 µm diameter, middle left panel; red lines: 10, 25, 50, 75, 90% N_{ice} percentiles; blue line: fit through median N_{ice} values (median N_{ice}: 0.03 cm⁻³; 10, 25, 75, 90% percentiles: 0.002, 0.007, 0.102, 0.3 cm⁻³); black lines: minimum/middle/maximum N_{ice} of Krämer et al. (2009)). Mass mean radius (R_{ice} : calculated from IWC/N_{ice}, bottom left panel; red lines: 10, 25, 50, 75, 90% R_{ice} percentiles; black lines: minimum/middle/maximum R_{ice} of Krämer et al. (2009)). In-cloud and clear sky relative humidity wrt ice (RH_{ice} , top and middle right panel), and water vapor volume mixing ratio (H_2O , bottom right panel). The field campaigns included in the data analysis are listed in Table 5. Data evaluation methods and detection ranges of the parameters are described in Appendix A. Note that for N_{ice} -and thus R_{ice}- the hours spent in clouds is less than in Figure 3. The reason is that for the calculation of data frequency distributions only measurements covering the same detection range are used (see also Appendix A2).



Figure 8. Same as Figure 7 (left column), but for mid-latitudes and tropics (WCBs: Warm Conveyor Belts; MCS: Mesoscale Convective Systems; TTL: Tropical Tropopause Layer). The field campaigns are listed in Table 5.



Figure 9. Same as Figure 7 (right column), but for mid-latitudes and tropics. The field campaigns are listed in Table 5.



Figure 10. Same as Figure 7, but for the field campaign StratoClim 2017 above the Asian monsoon.



Figure 11. Climatology of cirrus clouds in dependence on potential temperature Θ (left panels: tropics without Asian monsoon, right panels: Asian monsoon). In the middle panel, the range of the TTL (after Fueglistaler et al., 2009) and the cold point tropopause (CPT, derived from the observed temperature profiles) are marked; the corresponding altitude range is ~14 to 20 km. Note that in the tropics 355–330 K \approx 235–275 K (-38–0 C) and that the TTL spans from about 425–355 K. The field campaigns are listed in Table 5.



Figure 12. Climatology of humidity in dependence on potential temperature Θ (left panels: tropics without Asian monsoon, right panels: Asian monsoon). In the middle panel, the range of the TTL (after Fueglistaler et al., 2009) and the cold point temperature (CPT, derived from the observed temperature profiles) are marked; the corresponding altitude range is ~14 to 20 km. Note that in the tropics 355–330 K \approx 235–275 K (-38–0 C) and that the TTL spans from about 425–355 K. The field campaigns are listed in Table 5.



DARDAR $N_{\rm ice} > 5~\mu m$ diameter



Figure 13. N_{ice} – temperature climatologies from (top panel) in situ measurements during the field campaigns COALESC2011, ATTREX2014, ACRIDICON2014, ML-CIRRUS2014 and STRATOCLIM2017, (bottom panel) satellite remote sensing, applying the algorithm DARDAR-Nice $f(N_0^*, D_m)$ to N_0^* , D_m derived from the in situ observations. The black contours represent the 25, 50 and 75th percentiles. Straight black line: DARDAR-Nice average median N_{ice} (0.064 cm⁻³), straight blue line: in situ average median N_{ice} (0.037 cm⁻³). The excess of N_{ice} by a factor of 1.73 in DARDAR is caused by the retrieval method, for more information see text.

DARDAR $N_{ice} > 5 \ \mu m$ diameter 10 years climatology (2006 - 2016) Global 10³ 10² 0.0-2.0 2.0-5.0 10¹ 5.0-7.5 $N_{ice} \left(cm^{-3} \right)$ 7 5-10 0 10⁰ 10.0-12.5 10⁻¹ 12.5-15.0 15.0-20.0 10⁻² 20.0-30.0 30.0-40.0 10⁻³ 40.0-100 10 210 180 190 200 220 230 240 Temperature (K)

Figure 14. N_{ice} – temperature climatology from 10 years satellite remote sensing observations using DARDAR-Nice. The data was collected twice a day, at midday and at midnight. Thick black lines: 25, 50 and 75th percentiles; thin plain cyan line: DARDAR-Nice median N_{ice} ; thin dotted cyan line: DARDAR-Nice median N_{ice} adjusted to the offset factor of 1.73 between DARDAR-Nice and the in situ observations (see Figure 13); blue line: in situ average median N_{ice} (from Figure 7). Temperature intervals containing less than 10 retrievals were excluded (results non-sensitive to exact threshold).



Figure 15. N_{ice} – temperature climatology across the entire N_{ice} -T parameter space. Red lines: global DARDAR-Nice percentiles (adjusted to the in situ observations by the offset factor of 1.73, see Figure 13). The 10/90th, 25/75th and 50th percentiles are indicated by dotted, dashed and plain lines, respectively. Blue lines: in situ average percentiles (from Figure 7). The percentiles are recommended for intercomparison with other data sets (the corresponding numbers are listed in Table 2); regional median N_{ice} are shown in Table 3. The shaded background indicates the occurrence of retrievals, i.e. cirrus clouds. The cirrus appear most often at higher temperatures above about 225 K.

Temperature	in-situ adj. DARDAR-Nice				
(K)	$N_{ice} (cm^{-3})$				
	10% perc.	25% perc.	median	75% perc.	90 % perc.
180-190	0.034	0.051	0.087	0.149	0.243
190-200	0.033	0.055	0.100	0.173	0.274
200-210	0.028	0.051	0.091	0.163	0.265
210-220	0.017	0.034	0.068	0.133	0.239
220-230	0.011	0.022	0.049	0.102	0.195
230-240	0.007	0.015	0.033	0.068	0.130
Temperature		In-situ			
all	$N_{ice} (cm^{-3})$				
	10% perc.	10% perc. 25% perc. median 75% perc. 90 % perc			
	0.002	0.007	0.03	0.102	0.3

Table 2. DARDAR-Nice and in situ observation N_{ice} percentiles (as portrayed in Figure 15) for intercomparison with global models or other data sets.

Region	in-situ adj. DARDAR-Nice					
	$N_{ice} (cm^{-3})$					
	medians					
	all Temp.	T-int weighted	most frequent T-rang			
Arctic	0.036	0.023	0.016	230-242 К		
Mid-lat North	0.057	0.040	0.030	227-242 K		
Tropics	0.070	0.067	0.074	197-221 K		
Mid-lat South	0.057	0.043	0.031	227-242 K		
Antarctica	0.050	0.045	0.029	222-242 K		
Global	0.056	0.046	0.031	224-242 K		

Table 3. DARDAR-Nice median N_{ice} (as portrayed in Figure 15) for five regional latitude bands: Arctic: 90° – 67.7° , Mid-lat North: 67.7° – 23.3° , Tropics: 23.3° – -23.3° , Mid-lat South: -23.3° – -67.7° , Antarctica: -67.7° – -90° as well as global median N_{ice} . All Temp.: median without temperature binning; T-int weighted: median using the relative occurrence of each temperature interval as a weight; most frequent: median considering those temperature intervals that contain 50% of the total N_{ice} occurrence, T-range denotes the respective minimum and maximum temperatures.

Appendix A: Methods

A1 Field campaigns and instrumentation

The field campaigns, locations and deployed instruments are listed in Table 4, which is an extended
version of the respective Table in the Cirrus Guide: Part I. For a brief description of the instruments already introduced there, we refer to Krämer et al. (2016). Other instruments used during START 2008, CONTRAST 2014, ATTREX 2014, POSIDON 2016 and LTU 2012-2018 are briefly introduced here.

- 1490 ATTREX 2014 and POSIDON 2016: IWC and RH_{ice} are measured with the dual channel NOAA-H₂O instrument, a dual channel TDL hygrometer which detects both H₂O_{tot} (= H₂O_{gas} + IWC) and H₂O_{gas} by using a forward and a backward directed inlet. The detection limit of the IWC is 0.03 ppmv ($\approx 0.003 \text{ mg/m}^3$). Cloud particle size distributions (PSDs) and thus N_{ice} are recorded by a FCDP and a 2D-S. The size range of the FCDP is 1–50 µm diameter and the 2D-S counts cloud 1495 particles between 25–3005 µm
- 1495 particles between 25–3005 μm.

Start-08 and CONTRAST: IWC and N_{ice} are both derived from PSD measurements by a CDP and a 2D-C. The size range of the CDP is 2–50 µm diameter and the 2D-C counts cloud particles between 60–1100 µm.

1500

LTU 2012-2018: IWC and N_{ice} are both derived from PSD measurements by the Balloon-borne Ice Cloud particle Imager (B-ICI), detecting ice particle sizes between 10 and 1200 μ m diameters. Note that the measurements are not from aircraft, but from balloons launched at Kiruna, Sweden.

A2 Data evaluation methods

- 1505 The data evaluation methods, detection ranges and data quality criteria of the measurements of IWC, N_{ice}, R_{ice}, RH_{ice} and H₂O are described in the following. The instruments measuring the respective parameters are listed in Table 4, the estimated uncertainties are mentioned in the Table caption. For the calculation of frequency distributions of the data, only those campaigns where the measured parameters cover the same detection range are used, since mixing of data with differing detection
- 1510 ranges influences the calculation of percentiles. Thus, the list of analyzed campaigns varies for each parameter, since not all parameters are available for the same detection ranges in all campaigns, or the data quality is considered as suspicious. In Table 5 the available parameters are listed for each campaign. The resulting data coverage of the parameters over the entire temperature range is shown in Figure 16.

1515 A2.1 Ice water content (IWC)

In the earlier studies of Schiller et al. (2008) and Luebke et al. (2013), the IWC was detected as difference between total water (H_2O_{tot} : water vapor + evaporated ice crystals) and gas phase water (H₂O_{gas}); this product is named here IWC(H₂O). Krämer et al. (2016) also determined the IWC by integrating the ice particle size distributions (PSDs) measured by cloud spectrometers in cases where

no total water measurements were available; we call this IWC(PSD). A detailed description of both 1520 methods is given in Afchine et al. (2018). Also, Afchine et al. (2018) demonstrated good agreement between the two methods under reliable sampling conditions. Agreement between IWC(H₂O) and IWC(PSD) is also a criterion for the quality of the IWC data (see also Thornberry et al., 2017), as well as for the measurements of H_2O and the PSDs (for the data quality of H_2O and N_{ice} see also the following subsections). 1525

Unlike in the earlier studies, we here combine the two methods, IWC(H₂O) and IWC(PSD). In case both measurements are available, IWC(H2O) is used as first choice for each flight second. But if IWC(H₂O) is zero or missing, IWC(PSD) is used. This combination of IWCs can fill times of measuring failures or signals beyond detection limits of one instrument and also compensates for

some of the inhomogeneities in cirrus. If only one instrument is available, the respective IWC is

1530

1545

taken.

As discussed by Schiller et al. (2008) and Afchine et al. (2018), the lower detection limit of IWC(H₂O) depends on the temperature and can not be clearly defined. Data points below the lower dotted line drawn in the IWC-T parameter space in Figure 16 (upper left panel) represent obser-

vations where the difference between both the H_2O_{tot} and H_2O_{gas} measurements is not significant 1535 to unambiguously identify a cloud. This is where the ratio $H_2O_{enh}/H_2O_{gas} < 1.07$; note here that H₂O_{enh} is the originally measured total water quantity, which is enhanced due to the sampling characteristics of the system (for more detail see Schiller et al., 2008; Afchine et al., 2018). The lower detection limit of IWC(PSD) is 0.05 ppmv ($\approx 0.005 \text{ mg/m}^{-3}$) for NIXE-CAPS and 0.01 ppmv (\approx 0.001 mg/m^{-3}) for the combination of FCDP and 2D-C. 1540

A2.2 Ice crystal number (N_{ice})

From the campaigns where cloud particle size distributions (PSDs) are measured, we chose those where cloud particles between 3 and $\sim 1000 \ \mu m$ diameter are recorded when calculating N_{ice} (see Table 5). We calculate N_{ice} as the sum of ice crystal concentrations over all size bins larger than 3 µm on a 1 Hz time resolution, because this time span represents around 200 m flight distance - which might already not resolve cirrus inhomogeneities. Thus, when averaging Nice over longer

flight times, cloud free segments might influence the ice crystal concentrations.

Two instruments are needed to be deployed to cover the complete cirrus ice particle size range, one for smaller and the other for larger cloud particles. The PSDs from NIXE-CAPS (CAS-Depol + CIPgs) and FCDP + 2D-S are merged between 20 and 25 µm to avoid overlap of particle sizes, 1550

those of CDP + 2D-C at 55 μ m. In earlier campaigns where only FSSP measurements (3–30 μ m) are available, we calculate N_{ice} without the larger particles, since they contribute only a negligible part to N_{ice} (these campaigns are marked as y^{*} in Table 5). However, for the calculation of occurrence frequencies only the campaigns covering the whole size range are used.

1555

Concentrations and frequencies of occurrence of small ice crystals: the effect of limited sampling volumes

 N_{ice} is given by the number of cloud particle events recorded in the cloud volume that is sampled. Thus, the lowest concentration that can be detected is when only 1 particle is recorded. For the particle imaging probes (CIPgs, 2D-S, 2D-C, used size range ~ 20-1000 µm) that results to ~10⁻⁴ cm⁻³, while for the light scattering probes (CAS-Depol, FCDP, CDP, used size range ~ 3-20 µm) it reaches the higher value of ~0.015 cm⁻³. The difference in the recorded concentration range is caused by the differing sampling volumes of the two instrument types. As a consequence, N_{ice} concentrations ≤ 0.015 cm⁻³ contain only ice particles ≥ 20 µm. Lower concentrations of smaller 1565 particles can not be detected with current particle measurement techniques (see also Krämer et al.,

2016; Baumgardner et al., 2017).

Figure 17 (left panel) shows exemplarily N_{ice} in dependence on temperature, measured with NIXE-CAPS (NIXE-CAPS combines a CAS-Depol and a CIPgs) during the field campaign ML-Cirrus 2014, color coded by their frequencies of occurrence. CAS-DPOL samples 47.5cm⁻³@190m/s

- 1570 aircraft cruising speed in 1 second while CIPgs probes about 1000-16000cm⁻³ in dependence on particle size in the same time interval and aircraft speed (Costa et al., 2017). Thus, one particle event in the CAS-DPOL corresponds to a higher concentration as in the CIPgs. The red line in the left panel of Figure 17 represents the lowest detectable N_{ice} of CAS-DPOL and the blue line the same for CIPg. A notable feature in the figure is that directly above the lowest detectable N_{ice} of
- 1575 CAS-DPOL (red line) the N_{ice} frequencies jump to a higher level than below. The reason for this peak is an effect caused by the small sampling volumes of these instruments, which is often called 'bad statistics'. It concerns instruments whose lowest N_{ice} detection limit is above the naturally occurring particle concentrations. When clouds with concentrations smaller than the lowest detectable N_{ice} are probed, these concentrations appear in the sampling volume as 'single particles events' (in
- 1580 the size range $\lesssim 20 \ \mu\text{m}$) and are assigned to the minimum detectable concentration of the respective instrument, since the occurrence of one particle per time unit can not be related to the true larger air volume it belongs to. Therefore, the data sets of this study are cleared from the artificial concentration feature at the instrument's detection limit by excluding N_{ice} caused by one single particle in the respective instruments.
- The effect of this correction on the occurrence frequencies can be seen in Figure 17 (right panel), which shows the same data set as the left panel, but with 'single particle events' removed: the sharp edge at the lowest detectable N_{ice} of the CAS-DPOL has disappeared, instead more lower concen-

trations from the CIPg become visible, which were hidden by the artificial concentrations from the CAS-DPOL 'single particle events' before.

1590 'Single particle events' are less common if the measurements are averaged over larger time periods. The considered time interval can be adapted according to the required spatial resolution of the measurements. However, as outlined above, for this study we choose to show the lowest time interval (1 s) to achieve the highest possible spatial resolution.

1595 *Ice crystal shattering*

Fragmentation of large ice crystals at the cloud probes housings distorted a correct recording of the ice crystal concentrations in earlier times. Nowadays, new inlets and also postprocessing algorithms based on the interarrival times of the crystals in the sampling volume (Korolev and Field, 2015) have minimized this effect. Nevertheless, older data sets might be contaminated by small artifacts

1600 of shattered ice particles and needs to be marked in the data set and not used in the data analyses. Shattering can be recognized when looking at the N_{ice} frequencies of occurrence plotted versus the temperature, as can be seen in Figure 18, where an example of a data set including shattered ice crystals is shown. Clearly, high frequencies of occurrence appear at N_{ice} concentrations between 10-100 cm⁻³ for all temperatures. This ice particle 'mode' is not present in data sets not influenced by shattered ice crystals, as can be seen when comparing Figures 18 and 17.

A2.3 Water vapor and Relative Humidity wrt ice (H₂O and RH_{ice})

Water vapor measurements where, as ice crystal concentrations, under discussion in the last decade (Peter et al., 2006). After some efforts to improve the quality of the measurements (e. g. Fahey
1610 et al., 2014), the water vapor instruments today provide high-precision data (see Rollins et al., 2014;

Meyer et al., 2015; Thornberry et al., 2015; Buchholz et al., 2013; Kaufmann et al., 2018).

Nevertheless, we check the H_2O data quality for each campaign, including the recent ones, best by comparison of several H_2O instruments as done by e.g. Kaufmann et al. (2018). In case this is not possible, the frequencies of occurrence of in-cloud RH_{ice} are used to evaluate the quality of

1615 the measurements: inside of cirrus clouds, at temperatures $\gtrsim 200$ K, the frequencies should center around saturation, as can be seen in Figure 7 (upper right panel). Figure 19 gives an example of a data set with a bias in the in-cloud RH_{ice}. We should note that such a distortion might also be caused by a shift in the temperature measurements, which we, however, have not yet observed. In any case, such data sets are not used for the data analysis.

Aircraft	Locations	IWC	RH _{ice}	N _{ice}
Geophysica	Seychelles ¹ , Europe ² , Brazil ³ , Australia ⁴ , Africa ⁵	FISH ^a	FLASH ^c	FSSP-100 ⁱ
Geophysica	Nepal ¹⁴	FISH ^a	FLASH ^c	NIXE-CAPS ^m
Learjet	Europe ⁶	FISH ^a	OJSTER ^d , SEAL ^e	FSSP-300 ^j
BAe-146	UK ⁹	NIXE-CAPS ^m	-	NIXE-CAPS ^m
HALO	Europe ¹⁰ , Brazil ¹¹	NIXE-CAPS ^m	SHARC ^h	NIXE-CAPS ^m
Balloon	Europe ¹⁵	B-ICI ⁿ	-	B-ICI ⁿ
WB-57	USA ^{7,16} , Costa Rica ⁸	CLH ^b	HWV ^f , JLH ^g	CAPS ^k , 2D-S ¹
WB-57	Guam ¹³	NOAA-H ₂ O ^o	NOAA-H ₂ O ^o	FCDP ^p , 2D-S ¹
Global Hawk	Guam ¹²	NOAA-H ₂ O ^o	NOAA-H ₂ O ^o	FCDP ^p , 2D-S ¹
GV HIAPER	USA ¹⁷ , Guam ¹⁹	CDP ^q , 2D-C ^r	VCSEL ^{s:}	CDP ^q , 2D-C ^r
Learjet	USA ¹⁸	$2D-S^1$	_	FSSP ⁱ , 2D-S ¹
 TROCCINOX 2 SCOUTO-2, 200 AMMA 2006 CIRRUS 2003, AIRTOSS-ICE MidCix 2004, N. TC-4 2007 COALESC 201 MC-TRRUS 2014 POSIDON 2016 STRUE/COMPACTION 2016 STRUE/COMPACTION 2016 STRUE/COMPACTION 2016 STRUE/COMPACTION 2016 SPARTICUS 2012-2015 CRYSTAL-FAG SPARTICUS 2012-2015 SPARTICUS 2012-2015 SPARTICUS 2012-2015 	2005 b: Tunable diode laser hygror 15 c: Lyman-cf fluorescence hyg 15 c: Lyman-cf fluorescence hyg 16 c: Lyman-cf fluorescence hyg 2013 f: Lyman-cf fluorescence hyg 17 i: Light scattering cloud prob 1014 j: Light scattering cloud prob 2014 i: Optical imaging cloud prob 2015 o: Tunable diode laser hygror 1014 i: Light scattering cloud prob 2015 o: Tunable diode laser hygror 2014 i: Optical imaging cloud prob 2015 o: Tunable diode laser hygror 2016 o: Tunable diode laser hygror 21 o: Tunable diode laser hygror 22 p: Light scattering cloud prob 23 c: Light scattering cloud prob 24 c: Dynical imaging cloud prob 25 c: Light scattering cloud prob 26 c: Dynical imaging cloud prob 27 c: Light scattering cloud prob <td< td=""><td>neter (Luckke et al., 20 rometer (Stinikov et al., neter (Krämer et al., 20 neter (Buchholz et al., 2 neter (May, 1998) neter (May, 1998) (Baumgardner et al., 2 e (Baumgardner et al., i imaging cloud probe (Lawson et al., 2006) al imaging cloud probe (I al., 2018) neter (Thornberry et al. e (Baumgardner et al., e (Baumgardner et al., e (Baumgardner et al., neter (Thordh ot al., 2016)</td><td>13) 2007) 09) 2013) 2014) 5) 2017) Baumgardner et al., 2001) (Meyer, 2012; Luebke et al., 20 , 2017) 007) 2017) 2017) 2017) 10)</td><td>16)</td></td<>	neter (Luckke et al., 20 rometer (Stinikov et al., neter (Krämer et al., 20 neter (Buchholz et al., 2 neter (May, 1998) neter (May, 1998) (Baumgardner et al., 2 e (Baumgardner et al., i imaging cloud probe (Lawson et al., 2006) al imaging cloud probe (I al., 2018) neter (Thornberry et al. e (Baumgardner et al., e (Baumgardner et al., e (Baumgardner et al., neter (Thordh ot al., 2016)	13) 2007) 09) 2013) 2014) 5) 2017) Baumgardner et al., 2001) (Meyer, 2012; Luebke et al., 20 , 2017) 007) 2017) 2017) 2017) 10)	16)
The campaigns under	914 Schuler et al. (2008). Knimer et al. (2008). Knimer et al. (2009) and Finger of 9 Jones et al. (2012), 10 Voigt et al. (2017), 11 Wendisch et al. (2016), 12 Ja 14 Stroh et al. (2017), in prep., 15 Wolf et al. (2018), 16 https://espo.nasa.gov 19 Pan et al. (2017).	et al. (2015), 7-8 by Luebl ensen et al. (2017), 13 http /crystalface, 17 Pan et al.	xe et al. (2013); Jensen et al. (20 55://espo.nasa.gov/posidon, (2010), 18 Muhlbauer et al. (20)13), 14),

Table 4. Overview of campaigns and instruments. IWC: ice water content (uncertainties ~20%), RH_{ice} : relativehumidity over ice (uncertainties ~10-15%), N_{ice} : ice crystal number concentration (uncertainties ~10-100%);the sampling rate of all aircraft instruments is 1 Hz.

Campaign	data sets †	IWC	$N_{\rm ice}$	$RH_{\rm ice}/H_2O$				
	Ar	ctic						
EUPLEX_2003	SKL	У	у*	У				
ENVISAT_2003	_1 skl	У	y*	У				
ENVISAT_2003	_2 skl	У	y*	У				
LTU_2012-18		У	$\boldsymbol{y}^{\rm spe}$	-				
	Mid-latitude							
ENVISAT_2002	_2 sк	У	у*	У				
CIRRUS_2003	SKL	У	у*	У				
CIRRUS_2004	SKL	У	y*	У				
CIRRUS_2006	SKL	У	y*	У				
START_2008		У	$\boldsymbol{y}^{\rm spe}$	У				
MIDCIX_2004	Lk	У	y>15					
-								
SPARTICUS_201	0	У	y>15	-				
MACPEX_2011	k	У	y>15	У				
COALESC_2011	k	У	$\boldsymbol{y}^{\rm spe}$	_				
AIRTOSS_2013	k	У	y>15	У				
ML-CIRRUS_20	14 k	У	$\boldsymbol{y}^{\rm spe}$	У				
Tropics								
APE_THESEO_	1999 ski	У	y*	У				
CRYSTAL-FACE	Е_2002 l	У	_	-				
TROCCINOX_2	005 skl	У	у*	У				
SCOUT_2005	SKL	У	y*	У				
TC-4_2007	L	У	y>15	-				
CONTRAST_201	4	У	$\boldsymbol{y}^{\rm spe}$	_				
ACRIDICON_20)14 k	У	$\boldsymbol{y}^{\rm spe}$	У				
ATTREX_2014		у	$\boldsymbol{y}^{\rm spe}$	У				
POSIDON_2016		У	$\boldsymbol{y}^{\rm spe}$	У				
STRATOCLIM_	2017	у	$\boldsymbol{y}^{\rm spe}$	У				

Table 5. Cirrus Guide II data base for the IWC, N_{ice} and RH_{ice} climatologies. y: quality checked measurements; -: no/questionable measurements; an overview of each campaign is given in the Supplementary Material; y^{spe}: N_{ice} is corrected for single particle events (spe, see Section A2.2); y^{*}: N_{ice} from size limited detection range (3 - 30 μ m \rightarrow contains \geq 90% of the ice particles), not spe corrected; y>15: N_{ice} from size limited detection range (> 15 μ m \rightarrow underestimating the full ice particle concentration), not included in the data analysis presented here; the campaigns written in *italics* are not contained in the occurrence frequencies. [†] S K L k: campaigns contained in the data sets of Schiller et al. (2008); Krämer et al. (2009); Luebke et al. (2013); Krämer et al. (2016) (for S&K see Figure 5 of the Supplementary Material



Figure 16. Data coverage of the Cirrus Guide II in situ data base: IWC, N_{ice} , R_{ice} , in-cloud and clear sky RH_{ice} frequencies of occurrence in dependence on temperature.



Figure 17. N_{ice} frequencies of occurrence for ML-Cirrus 2014; left: original data set, right: corrected for 'single particle events'. A further example of an uncorrected data set can be seen in Figure 5 of the Supplementary Material, (middle left panel). For more details see text.



Figure 18. Example of N_{ice} measurements biased by shattering of large ice particles, visible in the frequencies of occurrence: high frequencies appear at N_{ice} concentrations between 10-100 cm⁻³ for all temperatures, which are not present in undisturbed measurements (see Figure 17). The black lines denote the middle and maximum N_{ice} lines from Krämer et al. (2009).



Figure 19. Example of biased H_2O measurements, visible in the in-cloud RH_{ice} frequencies: inside of cirrus clouds, at temperatures $\gtrsim 200$ K, the frequencies should center around saturation (see Figure 7 upper right panel).

1620 In Table CirrusGuide_Table_Observations.tex:

Krämer et al. (2009) Schiller et al. (2008) Meyer (2012); Meyer et al. (2015) Finger et al. (2015) Sitnikov et al. (2007) (Rollins et al., 2014) (Buchholz et al., 2013) (Baumgardner et al., 2001, 2017) Lawson et al. (2006) Luebke et al. (2013, 2016) Thornberry et al. (2017) McFarquhar et al. (2007) (Zondlo et al., 2010) (May, 1998) (Jones et al., 2012) (Voigt et al., 2017) (Wendisch et al., 2016)

(Muhlbauer et al., 2014) Wolf et al. (2018) Jensen et al. (2013b, 2017) Pan et al. (2017) Pan et al. (2010)