Reply to Reviewer 1

We thank the reviewer for their useful comments on our manuscript. Below we provide detailed response to your comments. In the following, the comments of the reviewer are presented in *italic blue*. Our responses are in normal black font. Changes to the text are in red.

Comment 1:

From my understanding of the manuscript, I do not believe that the primary conclusion "... most of the convectively moistened air is then transported to the center of the NA anticyclone and the anticyclonic structure helps maintain high water vapor content there." is necessarily supported by the evidence provided. What is shown is that there is an offset between the region that corresponds to the authors chosen metric of deep convection, and the region of maximum H2O anomaly as measured by MLS.

Even if one accepts the chosen convective proxy, there seems to be another plausible explanation for this geographic mismatch. Air simply resides in the Southern US region for a longer period before being transported downwind to other longitudes, and therefore has greater convective moisture efficiency. Thus, it is not clear to me to that transport from higher latitudes is required. The claim that can easily be made is that there is a strong correlation between air that has a long residence time over NA and air with a large H2O anomaly.

We agree that we need to show more analysis to support the conclusion, so we will add a new row to Fig. 3 in the paper; the revised figure is shown below. To produce the new row (the fourth one), we take MLS observations that our back-trajectory analysis shows were convectively enhanced and show the location where those trajectories encountered convection. We find that most of the convective encounters happened over the regions where GridRad data show deep convection is most frequently occurring. This supports our conclusion that the offset between convection and high water vapor is due to transport.

We acknowledge that we cannot rule out a contribution from unobserved convection in regions of high water vapor, but we see no evidence to support that.

In the revised text, we will add this row to Fig. 3 in the paper and insert a paragraph after line 118:

We identify the locations where parcels encounter deep convection in the back trajectories and grid the number of trajectories encountering convection into 2° x 2° boxes (Fig. 3J-L). Most of the locations of convective encounters occur over the region where NEXRAD data show deep convection frequently occurs, e.g., over the Central Plain region, and over Florida during August. The geographical distribution of convective encounters does not match with the convective influence ratio, indicating that convective moisture is transported to the region of high water vapor by the dynamics of the monsoon.

Comment 2: I am also skeptical about the relevance of the fact that in June the back trajectories

travel through very cold regions. These are back trajectories, so these low temperatures are influencing the parcels before they are moistened by the deep convection. Perhaps the authors are suggesting that these parcels are moistened over the US, travel to the tropics, and then come back to the US, but I would have thought that such a trajectory path would only be followed by only a very small fraction of parcels and the authors have provided no evidence to the contrary.

Yes, our argument is that parcels encounter deep convection over the US, travel to the (sub)tropics, are dehydrated along the way, and then travel back over the US. However, as the reviewer correctly suggested, only a minority of parcels take this path. If we regard 20°N-20°S as the tropics, then 33%, 13%, and 9% of the convectively influenced parcels follow this pathway during June, July, and August over 2005-2016, respectively. This reflects the fact that the monsoon circulation is weaker during June, and so a small but important fraction of parcels can follow this path; during July and August, when the monsoon is better established, this pathway is nearly shut off.

As we argue in the paper, year-to-year variations around these average values are responsible for observable variations in water vapor. During June 2011, 44% of the convectively influenced parcels traveled to the tropics, while during June 2010, only 20%. Per the Clausius-Claperyon equation, a 1-K change in minimum temperature will change water vapor by 20% at TTL temperatures. So, 20% of the parcels in 2011 experiencing temperatures a few degrees colder than those in 2010 (as suggested by Fig. 6 of the original paper) can change the water vapor over the monsoon region by the amount observed.

However, we acknowledge that our text could be clearer. We will make the following change to the text:

Line 137 - The second reason is also connected to the changing dynamics during June, July, and August. Parcels tend to travel to lower latitudes during June (Fig. 4a). 33% of the convectively moistened parcels travels to the tropics (20°N-20°S) during June, and 13% during July and 9% during August. Traveling to the tropics leads them to experience colder temperatures at 100 hPa (Fig. 4b). As a result, the median of the water vapor mixing ratio of the parcels that stays in the mid-latitudes is 5.98 ppmv, while it is 5.36 ppmv for those parcels that travels to the tropics. This means that convectively moistened air experiences subsequent dehydration more frequently in June than in later months (Randel et al., 2015).

Additional comment 1:

Abstract line 8– I'm not sure what the "hypothesis" is, nor is there a particular need to mention or have one.

Changes have been made in the text: We have replaced "Our hypothesis" with "This".

Additional comment 2:

Line 52 – *The convective radar data play a fundamental role in this paper. Although the reference to Cooney et al. is good, please devote a few lines to describing the reflectivity*

observations and what they mean. For instance, is a reflectivity Zn over 10 dBz a well-accepted value for tropopause overshooting convection?

Changes have been made in the text:

Line 52 - Cooney et al. (2018) used GridRad data to calculate the deep convective echo top, and found out a highest level that the reflectivity over 10 dBz is a representative threshold that balances the sensitivity and noise. In our analysis, we also use this strategy and identify deep convection as observations of reflectivity over 10 dBz.

Additional comment 3:

Figure 1 - I understand that in panel 'A' different very different quantities are being plotted, but just putting a y-axis that says "normalized" is unacceptable. There needs to be some way for the reader to connect the plotted value with a physical quantity (ppmv, fractional occurrence of convection, etc.). Also, the two reddish lines in panel 'A' (apparently one is orange) are very difficult to distinguish.

Fig. 1A has been revised. We now have multiple y-axes, with each showing the values of each physical quantity. A green line replaces the orange line, which should improve the figures clarity. The revised Fig. 1 is shown below.

Additional comment 4:

Figure 3 – This is an interesting figure, and the fact that the convective influence ratio looks not dissimilar from the MLS H2O anomaly is very interesting. The black line single value GridRad contour is useful because it makes it easier to see the offset between the maximum convection and the maximum H2O anomaly, and if this specific

convection contour represented all convection, then one would be forced into the conclusion presented by the authors, i.e. that high H2O air is being transported from these regions to the lower latitude regions. But there is nothing special about the specific convection level contour that the authors have chosen. While some of the moist air may

have been brought down from the North, some of the moistening in the Southern US is almost certainly caused by local convection in this region. This figure therefore include a full row showing a color contour of the convective occurrence by month throughout the US.

We agree that there is nothing special about the 1e-5 contour. We have therefore taken the reviewer's suggestion and added a row with the maps of convective occurrence over the NA.

Changes have been made in the text:

Line 106: From June to August every year, deep convection frequently occurs during boreal summer, especially over the central US (Fig.3A-C, see also Cooney et al. (2018)). The water vapor mixing ratio over NA also shows positive anomalies relative to the zonal mean (Fig. 3D-F). However, there is a discrepancy between the spatial distribution of the water vapor anomaly

and deep convective occurrence: The deep convection occurs mainly over the Central Plains region, centered around 40°N. Large positive water vapor anomalies are observed over a broader longitude range south of 40°N latitude.



Figure 1. (A) Time series of normalized 100-hPa (red line) MLS water vapor anomaly (zonal mean removed), (blue line) convective occurrence from GridRad, and (green line) the convective influence ratio in the back trajectory experiments. All data are 5-day averages over the NA region (25°N-50°N, 70°W-130°W) during June, July, August 2005-2016. For the convective frequency, we use linear interpolation to estimate the value at 100 hPa. The convective influence ratio is the fraction of the MLS observations that encounter deep convection, as determined by the back trajectory calculations. (B-D) Joint distribution of convection and water vapor time series during 2005-2016 divided into (B) June, (C) July, and (D) August. Solid lines show the linear fit, and dashed lines show the 95% significant level margin of error of the slope bar (accounting for auto-correlation). To account for the time for the water vapor to spread out, each data point is a 10-day average of convection, with water vapor averaged over the last five days of the averaging period for convection.



Figure 3. (Top row) Distribution of the 100-hPa GridRad deep convection occurrence averaged over 2005-2016 in (A) June, (B) July, and (C) August. The black contour in each panel (repeated in each row) is the 10⁻⁵ contour of GridRad convective occurrence, averaged over that month. (Second row) Geographical distribution of the MLS 100-hPa water vapor anomaly (after removal of the zonal mean), averaged over 2005-2016 in (D) June, (E) July, and (F) August. (Third row) Geographical distribution of the convection influence ratio over the NA region during (G) June, (H) July and (I) August 2005-2016. The black dashed contour (repeated in panels G-L) is the water vapor anomaly contours matching the shading in the corresponding upper panel. (Fourth row) Location where convectively influenced parcels encounter convection during 2005-2016 (J) June, (K) July and (L) August; (Bottom row) Geographical distribution of the parcel time spent over the NA region during (M) June, (N) July, and (O) August. (The stream lines are horizontal velocities interpolated onto 100 hPa using the cubic spline method from ERAi data averaged over the same period.

References

Cooney, J. W., Bowman, K. P., Homeyer, C. R. and Fenske, T. M.: Ten Year Analysis of Tropopause-Overshooting Convection Using GridRad Data, J. Geophys. Res. Atmos., 123(1), 329–343, doi:10.1002/2017JD027718, 2018.

Randel, W. J., Zhang, K. and Fu, R.: What controls stratospheric water vapor in the NH summer monsoon regions?, J. Geophys. Res. Atmos., 120(15), 7988–8001, doi:10.1002/2015JD023622, 2015.

Reply to Reviewer 2

We thank the reviewer for their useful comments on our manuscript. Below we provide detailed response to your comments. In the following, the comments of the reviewer are presented in *italic blue*. Our responses are in normal black font. Changes to the text are in red.

Comment 1:

The paper presents a hypothesis that deep overshooting convection over the great plains of north America moistens the lower stratosphere and is transported and trapped in the North American anticyclone. they use trajectory calculations to see if water vapor measurements from MLS encountered convection or not. They show that during July and August that the difference between convective and non convective trajectories is nearly 1 ppmv. In June it is much less. The take away message i get from this is that the establishment of the North American anticyclone is the dominant factor for having high lower stratospheric water vapor over North America.

This is not the impression we were trying to get across. Convection is they key reason that water vapor is higher in the monsoon region. Dynamics and the structure of the monsoon can mediate the impact of convection by, for example, regulating the temperature that air in the monsoon experiences. In fact, one should not really separate these phenomena. After all, it is convection that fundamentally causes the anticyclonic monsoonal circulation. Changes like this will be made in the text:

Line 4 - We use a Lagrangian trajectory model to demonstrate that the structure and the location of the NA anticyclone, as well as the tropical upper tropospheric temperature, mediate the moistening impact of convection.

It completely dominates convective activity in the Junes of 2010 and 2011 where the former year is high with low NA convection and low the following year despite having more NA convection. I have difficulty believing that in 2010 before the anticyclone is set up that the convectively moistened air moves to the colder tropics where it gets freeze dried first.

While this might be counterintuitive, we provide evidence in the paper that this is indeed happening. We examine Lagrangian trajectories (e.g., Fig. 6 in the original paper and associated text) and show that parcels are indeed reaching deep into the tropics. If we regard 20°N-20°S as the tropics, then 20% and 44% of the convectively influenced parcels travel into the tropics and back during June 2010 and 2011, respectively, and this difference can lead to significant interannual variability in the monsoon region.

We acknowledge that we should add more numerical information to the paper to make it clear. Changes have been made in the text:

Line 137 - The second reason is also connected to the changing dynamics during June, July, and August. Parcels tend to travel to lower latitudes during June (Fig. 4a). 33% of the convectively moistened parcels travels to the tropics (20°N-20°S) during June, and 13% during July and 9% during August. Traveling to the tropics leads them to experience colder temperatures at 100 hPa

(Fig. 4b). As a result, the median of the water vapor mixing ratio of the parcels that stays in the mid-latitudes is 5.98 ppmv, while it is 5.36 ppmv for those parcels that travels to the tropics. This means that convectively moistened air experiences subsequent dehydration more frequently in June than in later months (Randel et al., 2015).

Line 157 - In June 2010, 20% of the convectively influenced parcels travel to the tropics (20°N-20°S) in 5 days, while in June 2011, 44% do (Fig. 6a).

It seems from looking at the figure 3 and figure 5 that the horizontal tape recorder signature of water transport is playing a big role here too. i certainly agree that deep convection over the NA plains is adding water but it might be a small perturbation on top of the large scale transport coming up from the tropics that is also becoming more moist during the summer months. I think this could be disentangled with some modelling studies where one could artificially hold the tropical tropopause temperature constant all year thus removing the tape recorder signatures from the tropics and seeing what NA enhancements occur just due to local convection.

There is abundant previously published work that show that the high water vapor found in this region is the result of local convection in the region (Hanisco et al., 2007; Herman et al., 2017; Schwartz et al., 2013; Smith et al., 2017). This can also be seen in Fig. S1, which shows that the water vapor over the monsoon region is a local maximum, meaning that it could not be caused by horizontal transport out of the deep tropics.

To provide further support, Fig. S2 shows the horizontal distribution of parcels initially located in the NA monsoon after 10 day back trajectories. This shows that many of the parcels had been within the NA monsoon for more than 10 days, and those that were not had been transported to the monsoon by westerly winds from the Pacific — not from the tropics.

We will add some discussion to the revised paper discussing how the water vapor maximum in the monsoon is due to local convection:

Line 18 - However, over the Asian monsoon and North American monsoon, higher water vapor mixing ratios, sometimes as high as 12 ppmv, are observed. This value is much higher than the water vapor mixing ratio in the tropics, indicating that the air did not go through the TTL or is moistened further after leaving the TTL (Anderson et al., 2012; Schwartz et al., 2013; Randel et al., 2015; Smith et al., 2017).

Line 80 - We have subtracted the zonal mean value (based on the JJA mean from 70°W to 130°W at each degree of latitude) from NA the 100-hPa water vapor content before normalizing, allowing us to focus on the variability in the NA region relative to the zonal average value and minimize the impact of transport from the tropics.

Minor recommendation 1:

minor recommendations page 1 line 18 replace sometimes as high as with exceeding ... MLS has seen higher values as has Anderson.

Changes have been made in the text:

Line 18- However, over the Asian monsoon and North American monsoon, higher water vapor mixing ratios, sometimes exceeding 12 ppmv, are observed, indicating that the air did not go through the TTL or is moistened further after leaving the TTL.

Minor recommendation 2: page 3 line 81 from NA the 100 to from the NA 100...

Changes have been made in the text:

Line 81 - We have subtracted the zonal mean value (based on the JJA mean from 70°W to 130°W at each degree of latitude) from the NA 100-hPa water vapor content before normalizing, allowing us to focus on the variability in the NA region relative to the zonal average value.

Minor recommendation 3:

Throughout the manuscript lower case letters a, b, .. are used to refer to panels in the figures but the figures use upper case letters A, B, ... Please make this consistent.

Changes have been made in the text.

Minor recommendation 4:

page 5 line 135 I would write that sentence as As a result moistening from deep convection becomes less diluted by zonal mean flow later in the summer.

Changes have been made in the text:

As a result, moistening from deep convection becomes less diluted by zonal mean flow later in the summer.



Figure S1. MLS 100-hPa water vapor mixing ratio during 2005-2016 (A) June, (B) July, and (C) August.



Figure S2. Number of the parcels in each $2^{\circ}2^{\circ}$ grid box after 10 days in the back trajectory model during 2005-2016 (A) June, (B) July and (C) August. In the back trajectory experiments, we initiate the parcels $1^{\circ}x1^{\circ}$ $1_{\circ}x$ 1_{\circ} grid over NA ($25^{\circ}N - 50^{\circ}N$, $70^{\circ}W - 130^{\circ}W$) every day during each month, and track back their position in 10 days. (Black contour) Geographical distribution of the MLS 100-hPa water vapor anomaly (after removal of the zonal mean).

References

Hanisco, T. F., Moyer, E. J., Weinstock, E. M., St. Clair, J. M., Sayres, D. S., Smith, J. B., Lockwood, R., Anderson, J. G., Dessler, A. E., Keutsch, F. N., Spackman, J. R., Read, W. G. and Bui, T. P.: Observations of deep convective influence on stratospheric water vapor and its isotopic composition, Geophys. Res. Lett., 34(4), L04814, doi:10.1029/2006GL027899, 2007.

Herman, R. L., Ray, E. A., Rosenlof, K. H., Bedka, K. M., Schwartz, M. J., Read, W. G., Troy, R. F., Chin, K., Christensen, L. E., Fu, D. J., Stachnik, R. A., Bui, T. P. and Dean-Day, J. M.: Enhanced stratospheric water vapor over the summertime continental United States and the role of overshooting convection, Atmos. Chem. Phys., 17(9), 6113–6124, doi:10.5194/acp-17-6113-2017, 2017.

Randel, W. J., Zhang, K. and Fu, R.: What controls stratospheric water vapor in the NH summer monsoon regions?, J. Geophys. Res. Atmos., 120(15), 7988–8001, doi:10.1002/2015JD023622, 2015.

Schwartz, M. J., Read, W. G., Santee, M. L., Livesey, N. J., Froidevaux, L., Lambert, A. and Manney, G. L.: Convectively injected water vapor in the North American summer lowermost stratosphere, Geophys. Res. Lett., 40(10), 2316–2321, doi:10.1002/grl.50421, 2013.

Smith, J. B., Wilmouth, D. M., Bedka, K. M., Bowman, K. P., Homeyer, C. R., Dykema, J. A., Sargent, M. R., Clapp, C. E., Leroy, S. S., Sayres, D. S., Dean-Day, J. M., Paul Bui, T. and Anderson, J. G.: A case study of convectively sourced water vapor observed in the overworld stratosphere over the United States, J. Geophys. Res. Atmos., doi:10.1002/2017JD026831, 2017.

List of relevant changes:

To be consistent with the figures, all the letters in the manuscript referring to the panels in the figures are changed from lower case to upper case.

Line 4: change the text

We use a Lagrangian trajectory model to demonstrate that the structure and the location of the NA anticyclone, as well as the tropical upper tropospheric temperature, mediate the moistening impact of convection.

Line 8: change 'our hypothesis' to this

This explains both the summer seasonal cycle and inter-annual variability of the convective moistening efficiency in the NA region, and can provide valuable insight on modeling stratospheric water vapor.

Line 18, now line 17: change the text

However, over the Asian monsoon and North American monsoon, higher water vapor mixing ratios, sometimes exceeding 12 ppmv, are observed. This value is much higher than the water vapor mixing ratio in the tropics, indicating that the air did not go through the TTL or is moistened further after leaving the TTL (Anderson et al., 2012; Schwartz et al., 2013; Randel et al., 2015; Smith et al., 2017).

Line 52, now line 50: add more details

Cooney et al. (2018) used GridRad data to calculate the deep convective echo top, and found out a highest level that the reflectivity over 10 dBz is a representative threshold that balances the sensitivity and noise. In our analysis, we also use this strategy and identify deep convection as observations of reflectivity over 10 dBz.

Line 80, now line 81: add more details

We have subtracted the zonal mean value (based on the JJA mean from 70°W to 130°W at each degree of latitude) from the NA 100-hPa water vapor content before normalizing, allowing us to focus on the variability in the NA region relative to the zonal average value and minimize the impact of transport from the tropics.

Line 93, now line 94: delete irrelevant sentence

Fig. 1a shows the convective influence, the fraction of MLS observations that encountered convection and Fig. 2 shows the probability density function (PDF) of water vapor mixing ratio during June, July, and August 2005-2016 in the two groups.

Line 106: change the text

From June to August every year, deep convection frequently occurs during boreal summer, especially over the central US (Fig.3A-C, see also Cooney et al. (2018)). The water vapor mixing ratio over NA also shows positive anomalies relative to the zonal mean (Fig. 3D- F). However, there is a discrepancy between the spatial distribution of the water vapor anomaly and deep convective occurrence: The deep convection occurs mainly over the Central Plains region, centered around 40°N. Large positive water vapor anomalies are observed over a broader longitude range south of 40°N latitude.

Line 118: insert a paragraph

We identify the locations where parcels encounter deep convection in the back trajectories and grid the number of trajectories encountering convection into 2° x 2° boxes (Fig. 3J-L). Most of the locations of convective encounters occur over the region where NEXRAD data show deep convection frequently occurs, e.g., over the Central Plain region, and over Florida during August. The geographical distribution of convective encounters does not match with the convective influence ratio, indicating that convective moisture is transported to the region of high water vapor by the dynamics of the monsoon.

Line 135, now line 140: change the text

As a result, moistening from deep convection becomes less diluted by zonal mean flow later in the summer.

Line 137, now line 141: Add two sentences

Line 137 - The second reason is also connected to the changing dynamics during June, July, and August. Parcels tend to travel to lower latitudes during June (Fig. 4a). 33% of the convectively moistened parcels travels to the tropics (20°N-20°S) during June, and 13% during July and 9% during August. Traveling to the tropics leads them to experience colder temperatures at 100 hPa (Fig. 4b). As a result, the median of the water vapor mixing ratio of the parcels that stays in the mid-latitudes is 5.98 ppmv, while it is 5.36 ppmv for those parcels that travels to the tropics. This means that convectively moistened air experiences subsequent dehydration more frequently in June than in later months (Randel et al., 2015).

Line 157, now line 163: change the text

Line 157 - In June 2010, 20% of the convectively influenced parcels travel to the tropics (20°N-20°S) in 5 days, while in June 2011, 44% do (Fig. 6A).

Figure 1A: change normalized value to absolute value, change line color:

Figure 1. (A) Time series of normalized 100-hPa (red line) MLS water vapor anomaly (zonal mean removed), (blue line) convective occurrence from GridRad, and (green line) the convective influence ratio in the back trajectory experiments. All data are 5-day averages over the NA

region (25°N-50°N, 70°W-130°W) during June, July, August 2005-2016. For the convective frequency, we use linear interpolation to estimate the value at 100 hPa. The convective influence ratio is the fraction of the MLS observations that encounter deep convection, as determined by the back trajectory calculations. (B-D) Joint distribution of convection and water vapor time series during 2005-2016 divided into (B) June, (C) July, and (D) August. Solid lines show the linear fit, and dashed lines show the 95% significant level margin of error of the slope bar (accounting for auto-correlation). To account for the time for the water vapor to spread out, each data point is a 10-day average of convection, with water vapor averaged over the last five days of the averaging period for convection.

Figure 3: add two new rows

Figure 3. (Top row) Distribution of the 100-hPa GridRad deep convection occurrence averaged over 2005-2016 in (A) June, (B) July, and (C) August. The black contour in each panel (repeated in each row) is the 10⁻⁵ contour of GridRad convective occurrence, averaged over that month. (Second row) Geographical distribution of the MLS 100-hPa water vapor anomaly (after removal of the zonal mean), averaged over 2005-2016 in (D) June, (E) July, and (F) August. (Third row) Geographical distribution of the convection influence ratio over the NA region during (G) June, (H) July and (I) August 2005-2016. The black dashed contour (repeated in panels G-L) is the water vapor anomaly contours matching the shading in the corresponding upper panel. (Fourth row) Location where convectively influenced parcels encounter convection during 2005-2016 (J) June, (K) July and (L) August; (Bottom row) Geographical distribution of the parcel time spent over the NA region during (M) June, (N) July, and (O) August. (The stream lines are horizontal velocities interpolated onto 100 hPa using the cubic spline method from ERAi data averaged over the same period.

Influence of convection on stratospheric water vapor in the North American Monsoon region

Wandi Yu¹, Andrew E. Dessler¹, Mijeong Park², and Eric J. Jensen² ¹Department of Atmospheric Sciences, Texas A&M University, College Station, TX, USA ²National Center for Atmospheric Research, Boulder, Colorado, USA

Correspondence: Andrew Dessler (adessler@tamu.edu)

Abstract.

We quantify the connection between deep convective occurrence and summertime 100-hPa water vapor anomaly over the North America (NA) region and find substantial consistency of their inter-annual variations and that the water vapor mixing ratio over the NA region is up to ~1 ppmv higher when deep convection occurs. We use a Lagrangian trajectory model to

5 demonstrate that the structure and the location of the NA anticyclone, as well as the tropical upper tropospheric temperature, mediate the moistening impact of convection. The deep convection mainly occurs over the Central Plains region, and most of the convectively moistened air is then transported to the center of the NA anticyclone and the anticyclonic structure helps maintain high water vapor content there. This explains both the summer seasonal cycle and inter-annual variability of the convective moistening efficiency in the NA region, and can provide valuable insight on modeling stratospheric water vapor.

10 Copyright statement. TEXT

1 Introduction

Stratospheric water vapor influences both the climate (Forster and Shine, 1997, 2002; Smith et al., 2001; Solomon et al., 2010; Dessler et al., 2013) and chemistry of the atmosphere (Ramaswamy et al., 1996; Evans et al., 1998; Dvortsov and Solomon, 2001; Shindell, 2001; Stenke and Grewe, 2005). Most of the air reaching 100 hPa has traversed the tropical tropopause layer
(TTL), where low temperatures dehydrate the air to typically 3-5 ppmv (Mote et al., 1996; Sherwood and Dessler, 2000; Randel et al., 2004; Fueglistaler, 2005; Fueglistaler et al., 2009). However, over the Asian monsoon and North American monsoon, higher water vapor mixing ratios, sometimes exceeding 12 ppmv, are observed. This value is much higher than the water vapor mixing ratio in the tropics, indicating that the air did not go through the TTL or is moistened further after leaving the TTL (Anderson et al., 2012; Schwartz et al., 2013; Randel et al., 2015; Smith et al., 2017).

20 Convection penetrating the tropopause plays a potentially important role in moistening the stratosphere (Dessler and Sherwood, 2004; Dessler et al., 2007; Hanisco et al., 2007; Ueyama et al., 2015, 2018). Previous case studies have shown that deep convection over North America (NA) can reach the lowermost stratosphere (between the local tropopause and the 380 K isentropic surface), and can even enter the stratospheric overworld (above 380 K), thereby bringing a high water vapor content to the stratosphere (Hanisco et al., 2007; Herman et al., 2017; Smith et al., 2017). However, previous studies on the long-term

25 behavior of NA stratospheric water vapor and deep convection conclude that there is little connection (Randel et al., 2015; Sun and Huang, 2015; Kim et al., 2018). Model simulations, using prescribed global satellite-derived deep convection, also cannot reproduce a high water vapor mixing ratio over NA (Ueyama et al., 2018; Wang et al., 2019).

One crucial issue in these analyses is how high convection extends into the stratosphere. Previous studies based on the infrared satellite cloud-top height measurements are low biased (Ueyama et al., 2018; Schoeberl et al., 2019) owing to the

- 30 isothermal nature of the lowermost stratosphere as well as the fact that convective clouds rapidly sublimate in the dry stratosphere and therefore may be missed in observations from polar orbiting satellites. A better option for estimating cloud-top height over NA is ground-based radar (Liu and Liu, 2016; Cooney et al., 2018). The hourly interval GridRad data, derived from NEXRAD radar data, captures convective overshooting events over most parts of the contiguous United States (Solomon et al., 2016; Cooney et al., 2018), sometimes extending as high as 22 km.
- In this study, we focus on the NA region and use the GridRad data as a convective proxy to study the impact from deep convection on stratospheric water vapor. In the following analysis, we will first show the relationship between the inter-annual variability of the water vapor and deep convection over the NA region. Then we use a back trajectory model to illustrate the processes that influence whereby deep convection moistens the NA stratosphere, from the perspectives of spatial distribution, seasonal cycle, and inter-annual variability.

40 2 Data and methods

2.1 Data

45

The Microwave Limb Sounder (MLS) on board the NASA Aura satellite (Waters et al., 2006) has provided high-quality daily near-global observations of water vapor in the upper troposphere and stratosphere since August 2004. In our research, we use the version 4.2 level 2 product from 2005 to 2016 and focus on water vapor at 100 hPa over the NA region (25°N - 50°N, 70°W- 130°W). MLS makes roughly 2500 observations over this region every month and we bin and average these MLS data into a 4°latitude by 8°longitude grid after applying quality control (Livesey et al., 2018).

The GridRad dataset archives national NEXRAD WSR-88D radar data (version 3.1). Gridrad v3.1 has horizontal resolution 0.02°longitude and 0.02°latitude, and covers 25°N to 49°N, and 70°W to 105°W, or most parts of the contiguous United States (Homeyer, 2014; Homeyer and Kumjian, 2015; Solomon et al., 2016; Cooney et al., 2018). The vertical resolution of

50 the GridRad data is 1 km, from 1 to 24 km above sea level and it has a temporal resolution of one hour. Cooney et al. (2018) used GridRad data to calculate the deep convective echo top, and found out a highest level that the reflectivity over 10 dBz is a representative threshold that balances the sensitivity and noise. In our analysis, we also use this strategy and identify deep convection as observations of reflectivity over 10 dBz.

Finally, we bin the occurrence of convective cloud into a 1°latitude by 1°longitude (horizontal) by 1 km (vertical) grid box, dividing the number of observations that encounters convection by the total number of observations in each grid box. Temperature, pressure, and tropopause information comes from the European Centre for Medium-Range Weather Forecast (ECMWF) Re-Analysis (ERA)-Interim (ERAi) archived 6-hourly model fields (Dee et al., 2011). The ERAi data is gridded into a 1°x 1°horizontal grid with 37 vertical layers. To identify the tropopause, we linearly interpolate ERAi temperature into 2 hPa pressure intervals and define the lowest level where the lapse rate is less than 2 K/km to be the tropopause (World Meteorological Organization, 1957).

2.2 Back trajectory model

60

To identify water vapor observations that had previously encountered convection, we use back trajectory analyses. These use temperature, wind, and diabatic heating rate from the ERAi to drive the TRAJ3D Bowman trajectory code (Bowman, 1993; Bowman and Carrie, 2002). Vertical velocities in isentropic coordinates are computed from 6-hourly average diabatic heating

65 rates. Horizontal velocities come from 6-hourly instantaneous two-dimensional horizontal wind fields. Several previous studies have successfully identified convectively influenced air masses based on trajectory model driven by reanalysis data (Wright et al., 2011; Bergman et al., 2012, 2013; Smith et al., 2017).

In these experiments, we initialize air parcels at and surrounding the MLS observations made during June -August (JJA) over 2005-2016. The initiation of parcel positions follows the same strategy as Smith et al. (2017): We initiate a cluster of 27 parcels

on a 3 x 3 x 3 gird surrounding each MLS measurement ($\pm 0.25^{\circ}$ latitude, $\pm 0.25^{\circ}$ longitude, and ± 5 K potential temperature around and at 100 hPa). We advect the parcels back 5 days and record latitude, longitude, potential temperature, pressure, and temperature every hour.

After performing the back trajectory model, we then divide the trajectories into two groups, depending whether the parcels encountered deep convection along the path or not. The definition of encountering deep convection is if the trajectory is within

 $\pm 0.25^{\circ}$ latitude and $\pm 0.25^{\circ}$ longitude of a GridRad deep convection observation and when the parcel is below the convection top and above the local tropopause at the time the observation is made. If at least one parcel in the cluster of parcels encounters deep convection, the corresponding observation is categorized as encountering convection; otherwise, it is in the 'no-convection' group.

3 Relationship between area-average water vapor and convection

80 In Fig. 1.A, we show the time series of 5-day average 100-hPa water vapor anomaly and 10-day average convective occurrence, both averaged over the NA region. We have subtracted the zonal mean value (based on the JJA mean from 70°W to 130°W at each degree of latitude) from the NA 100-hPa water vapor content before normalizing, allowing us to focus on the variability in the NA region relative to the zonal average value and minimize the impact of transport from the tropics.

Overall, there is a high correlation between the water vapor anomaly and convective occurrence, which suggests that deep convection moistens the stratosphere. There are exceptions, however — *e.g.*, during June 2010 (marked by the arrow), the water vapor anomaly is high, despite deep convection being relatively infrequent. To make this correlation clearer, Figs. 1B-D show scatter plots of the 5-day water vapor anomaly and deep convective occurrence over NA in June, July, and August. We find that deep convection increases 100-hPa NA stratospheric water vapor by up to ~1 ppmv. The slope of the linear fit in Figs. 1.B-D represents the moistening efficiency, which is defined as the amount of water vapor content added per unit of deep convective occurrence in the stated month added. This moistening efficiency is

significantly lower during June than July and August, which we will be explained in the next section.

90

95

110

One must be careful not to confuse correlation with causality. We therefore use the back trajectory model to demonstrate the causal relationship implied in Fig. 1. As discussed in section 2.2, we divided the 100 hPa MLS observations into two groups depending on whether they encountered the deep convection during the 5-day back trajectory or not. Fig. 2 shows the probability density function (PDF) of water vapor mixing ratio during June, July, and August 2005-2016 in the two groups.

We see that the no-convection group has a similar PDF shape in June, July, and August, with peak values around 4 ppmv. For the MLS measurements that encountered convection, the peak of the PDF is 5-6 ppmv during July and August, and 4-5 ppmv during June, 0.37 ppmv, 0.62 ppmv, and 0.69 ppmv higher than the no-convection group during June, July, and August, respectively.

100 Our work in this section establishes that deep convection is increasing water vapor over the NA region. However, three questions remain to be answered: First, can deep convection explain the spatial distribution of the water vapor anomaly? Second, why is the convection more effective in July and August than in June? Third, why is there inter-annual variability in the effectiveness of moistening (for example, June 2010 *vs.* June 2011)? These are three key questions we answer in the following sections.

105 4 Differences between June, July, and August

From June to August every year, deep convection frequently occurs during boreal summer, especially over the central US (Fig. 3A-C, see also Cooney et al. (2018)). The water vapor mixing ratio over NA also shows positive anomalies relative to the zonal mean (Fig. 3D- F). However, there is a discrepancy between the spatial distribution of the water vapor anomaly and deep convective occurrence: The deep convection occurs mainly over the Central Plains region, centered around 40°N. Large positive water vapor anomalies are observed over a broader longitude range south of 40°N latitude.

- Our back trajectory calculations show that regions with high convective influence ratios (Fig. 3G-I) tend to be collocated with large positive water vapor anomalies (Fig. 3 A-C, also black dashed contours in Fig. 3G-I). Here, we define the convective influence ratio as the number of MLS observations in each grid box that encountered deep convection during the past 5 days, divided by the total number of MLS observations in that grid box. This collocation suggests that the pattern of enhanced water
- 115 vapor seen by MLS can be explained by frequent convection. It is worth mentioning that previous studies have also suggested that the water vapor maximum over the NA region cannot be reproduced without the inclusion of convection in the model (Ueyama et al., 2018, their Fig. 3C); (Wang et al., 2019, their Fig. 2F).

We identify the locations where parcels encounter deep convection in the back trajectories and grid the number of trajectories encountering convection into 2°x 2°boxes (Fig. 3J-L). Most of the locations of convective encounters occur over the region

120 where GridRad data show deep convection frequently occurs, *e.g.*, over the Central Plain region, and over Florida during August. The geographical distribution of convective encounters does not match with the convective influence ratio, indicating that convective moisture is transported to the region of high water vapor by the dynamics of the monsoon.

Also shown in Fig. 3 (panels M, N, and O) is the NA residence time, as calculated by back trajectory analyses. We initialize the parcel evenly on a 1°x 1°grid over NA (25°N - 50°N, 70°W- 130°W) every hour during each month, and track their

125 positions back 10 days. We then calculate the time from when it entered the NA region to the initialization point, and then grid the time of these parcels according to their location of initiation. The NA residence time indicates how long the air parcels in each grid box have been in the NA region. These figures show that parcels at the center of the monsoon have the longest history over NA a week or longer.

There is also a similarity between the distribution of the time spent over NA (Fig. 3M-O) and the convective influence ratio, indicating that the monsoon circulation tends to hold air that has flowed over convection in the NA region. This provides an explanation for the observations in Figs. 3E-G showing that convection tends to be located north of the 100-hPa water vapor maximum.

The monsoon dynamics are also an essential factor in the seasonal cycle of water vapor anomaly. Here, seasonal cycle refers to the increase in water vapor mixing ratio through the summer, from June to July to August. There are two reasons for this.

- 135 First, the North American monsoon anticyclone (NAMA) forms in June and enlarges and becomes stable during July and August (Clapp et al., 2019). This leads to increases in the average NA residence time from 3.4 to 4.5 to 4.9 days from June to August. This increases the convective influence (the fraction of MLS observations that encountered convection), with values in June, July, and August of 0.038, 0.081, and 0.091, respectively. What is happening here is that, later in the summer, the convectively moistened air is more likely to be confined within the NA region instead of being transported downwind by the
- 140 zonal mean flow. As a result, moistening from deep convection becomes less diluted by zonal mean flow later in the summer. The second reason is also connected to the changing dynamics during June, July, and August. Parcels tend to travel to lower latitudes during June (Fig. 4A). 33% of the convectively moistened parcels travels to the tropics (20°N-20°S) during June, and 13% during July and 9% during August. Traveling to the tropics leads them to experience colder temperatures at 100 hPa (Fig. 4B). As a result, the median of the water vapor mixing ratio of the parcels that stays in the mid-latitudes is 5.98 ppmv, while
- 145 it is 5.36 ppmv for those parcels that travels to the tropics. This means that convectively moistened air experiences subsequent dehydration more frequently in June than in later months (Randel et al., 2015).

The PDFs of the minimum saturation water vapor mixing ratio, which limits the amount of water in the parcel, indicates that parcels in June tend to have lower values (Fig. 4C). If we choose a minimum saturation water vapor mixing ratio of 5 ppmv as a threshold of effective moistening (stratospheric water vapor mixing ratio commonly won't exceed this value), then

150 88.0%, 97.8%, and 97.5% of the observations that encountered deep convection are effectively moistened in June, July and August, respectively. We calculate the effective convective influence ratio by dividing the number of convectively moistened observations that have a minimum saturation water vapor mixing ratio over 5 ppmv by the total number of observations. The effective convective influence ratio is 0.039, 0.082, and 0.092 during June, July, and August, respectively.

5 Interannual variability

- 155 Figure 1A shows times series for water vapor, deep convection, and the convective influence ratio for June-August of 2005-2016. The correlation coefficient between water vapor and the effective convective influence ratio time series is 0.74, and 0.60 between water vapor and deep convection. Despite the high correlation between these time series, there are also clear outliers, *e.g.*, June 2010 has lower convection, but much higher water vapor than June 2011. In this section, we compare these two months to illustrate the factors that contribute to the interannual variability of convective moistening.
- During June 2010, a stable anticyclone forms over the eastern NA region from June 11 to June 30 (Fig. 5, upper panel). In June 2011, the monsoon anticyclone is located further south, and the NA region is dominated by the westerly winds. Because of the difference in locations of the monsoon anticyclone, the parcels influenced by convection experience different pathways. In June 2010, 20% of the convectively influenced parcels travel to the tropics (20°N-20°S) in 5 days, while in June 2011, 44% do (Fig. 6A). This means that parcels influenced by convection in June 2011 on average experience colder temperatures (Fig.
- 165 6B). Finally, the tropics were slightly cooler during June 2011 compared to June 2010, which further contributed to lower water vapor in NA. The net result of these differences is that convectively influenced parcels retain more water vapor in June 2010 than in June 2011 (Fig. 6C) due to differing monsoon dynamics. Thus, monsoon dynamics variability plays a significant role in the generating inter-annual variability of stratospheric water vapor in the NA region.

6 Conclusions

170 In this study, we investigated the contribution of convection to the water vapor in the North American (NA) monsoon region, including the seasonal cycle and interannual variation of convective contributions during the Northern Hemisphere summer. We have shown that the deep convection moistens the lower stratosphere, adding on up to ~1 ppmv to the summertime NA water vapor at 100 hPa based on the observations from MLS.

We have also shown that it is not the amount of convection alone that determines the impact on water vapor — NA monsoon dynamics also play a role. We note that the location of deep convection is not collocated with the maximum water vapor in NA, and this is due to high water vapor content being transported downstream by the monsoon circulation. The maximum water vapor content appears near the center of the NA anticyclone.

We also analyzed the seasonal cycle of convective influence. During June, the NA monsoon circulation is located further south than during July and August, so air influenced by convection during June experiences colder temperatures while travel-

180 ing to the tropics. Subsequent dehydration reduces the net moistening from convection during June compared to those other months. Variations in the monsoon dynamics can also lead to interannual variations in convective moistening through a similar mechanism. We compare June 2010 and June 2011 and show that a more northerly monsoon circulation in June 2010 leads to convectively influenced air encountering warmer temperatures, leading to higher water vapor than in June 2011.

Our use of GridRad data as a source of convection is a limitation in our analysis because it only covers the continental US. 185 Much of the monsoonal deep convection also occurs over the Gulf of Mexico (Clapp et al., 2019), out of range of the NEXRAD stations. Future studies including convective data with a larger spatial extent may find that the deep convection over the Gulf of Mexico influences NA stratospheric water vapor, but we do not expect this will conflict with the main conclusions from our paper.

Data availability. Deep convection data from GridRad data: http://gridrad.org/data.html (Homeyer, 2014)

Water vapor observed by MLS: https://mls.jpl.nasa.gov/products/h2o_product.php (Waters et al., 2006)
 Wind, Temperature, and heating rate are from ERAinterm: https://www.ecmwf.int/en/forecasts/datasets/ (Dee et al., 2011)

Author contributions. WY performed the analysis and wrote the original draft. AED provided guidance on the outline of the research, and also edit the paper. MP and EJ provided guidance and discussion on section 3&4 and participate in editing.

Competing interests. The authors declare that they have no conflict of interest.

195 Acknowledgements. We thank William J. Randel, Laura Pan and Mark Schoeberl for their helpful discussion. We thank the reviewers for their comments. This work was supported by NASA grants 80NSSC18K0134 and 80NSSC19K0757. This material is also based upon work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under Cooperative Agreement No. 1852977. Any opinions, findings and conclusions or recommendations expressed in this material do not necessarily reflect the views of the National Science Foundation.

200 References

- Anderson, J. G., Wilmouth, D. M., Smith, J. B., and Sayres, D. S.: UV dosage levels in summer: Increased risk of ozone loss from convectively injected water vapor, Science, 337, 835–839, https://doi.org/10.1126/science.1222978, 2012.
- Bergman, J. W., Jensen, E. J., Pfister, L., and Yang, Q.: Seasonal differences of vertical-transport efficiency in the tropical tropopause layer: On the interplay between tropical deep convection, large-scale vertical ascent, and horizontal circulations, Journal of Geo-
- 205 physical Research: Atmospheres, 117, https://doi.org/10.1029/2011JD016992, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/ 2011JD016992, 2012.
 - Bergman, J. W., Fierli, F., Jensen, E. J., Honomichl, S., and Pan, L. L.: Boundary layer sources for the Asian anticyclone: Regional contributions to a vertical conduit, Journal of Geophysical Research: Atmospheres, 118, 2560–2575, https://doi.org/10.1002/jgrd.50142, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/jgrd.50142, 2013.
- 210 Bowman, K. P.: Large-Scale Isentropic Mixing Properties of the Antarctic Polar Vortex from Analyzed Winds, Journal of Geophysical Research-Atmospheres, 98, 23 013–23 027, https://doi.org/Doi 10.1029/93jd02599, 1993.
 - Bowman, K. P. and Carrie, G. D.: The mean-meridional transport circulation of the troposphere in an idealized GCM, Journal of the Atmospheric Sciences, 59, 1502–1514, https://doi.org/Doi 10.1175/1520-0469(2002)059<1502:Tmmtco>2.0.Co;2, 2002.
 - Clapp, C., Smith, J., Bedka, K., and Anderson, J.: Identifying Source Regions and the Distribution of Cross-Tropopause Convec-
- 215 tive Outflow Over North America During the Warm Season, Journal of Geophysical Research: Atmospheres, 124, 13750–13762, https://doi.org/10.1029/2019JD031382, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD031382, 2019.
 - Cooney, J. W., Bowman, K. P., Homeyer, C. R., and Fenske, T. M.: Ten Year Analysis of Tropopause-Overshooting Convection Using GridRad Data, Journal of Geophysical Research: Atmospheres, 123, 329–343, https://doi.org/10.1002/2017JD027718, http://doi.wiley. com/10.1002/2017JD027718, 2018.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, Quarterly Journal of the Royal Meteorological Society, 137, 553–597,
- https://doi.org/10.1002/qj.828, https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.828, 2011.
 - Dessler, A. E. and Sherwood, S. C.: Effect of convection on the summertime extratropical lower stratosphere, Journal of Geophysical Research-Atmospheres, 109, https://doi.org/Artn D23301 10.1029/2004jd005209, 2004.
 - Dessler, A. E., Hanisco, T. F., and Fueglistaler, S.: Effects of convective ice lofting on H2O and HDO in the tropical tropopause layer, Journal of Geophysical Research-Atmospheres, 112, https://doi.org/Artn D18309 10.1029/2007jd008609, 2007.
- 230 Dessler, A. E., Schoeberl, M. R., Wang, T., Davis, S. M., and Rosenlof, K. H.: Stratospheric water vapor feedback, Proceedings of the National Academy of Sciences, 110, 18087–18091, https://doi.org/10.1073/pnas.1310344110, http://www.pnas.org/cgi/doi/10.1073/pnas. 1310344110, 2013.
 - Dvortsov, V. L. and Solomon, S.: Response of the stratospheric temperatures and ozone to past and future increases in stratospheric humidity, Journal of Geophysical Research-Atmospheres, 106, 7505–7514, https://doi.org/Doi 10.1029/2000jd900637, 2001.
- 235 Evans, S. J., Toumi, R., Harries, J. E., Chipperfield, M. P., and Russell, J. M.: Trends in stratospheric humidity and the sensitivity of ozone to these trends, Journal of Geophysical Research-Atmospheres, 103, 8715–8725, https://doi.org/Doi 10.1029/98jd00265, 1998.

- Forster, P. M. and Shine, K. P.: Radiative forcing and temperature trends from stratospheric ozone changes, Journal of Geophysical Research: Atmospheres, 102, 10841–10855, https://doi.org/10.1029/96JD03510, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/ 96JD03510, 1997.
- 240 Forster, P. M. d. F. and Shine, K. P.: Assessing the climate impact of trends in stratospheric water vapor, Geophysical Research Letters, 29, 10–1–10–4, https://doi.org/10.1029/2001GL013909, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2001GL013909, 2002.
 - Fueglistaler, S.: Stratospheric water vapor predicted from the Lagrangian temperature history of air entering the stratosphere in the tropics, Journal of Geophysical Research, 110, D08107, https://doi.org/10.1029/2004JD005516, http://doi.wiley.com/10.1029/2004JD005516, 2005.
- 245 Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, Reviews of Geophysics, 47, https://doi.org/10.1029/2008RG000267, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008RG000267, 2009.
- Hanisco, T. F., Moyer, E. J., Weinstock, E. M., St. Clair, J. M., Sayres, D. S., Smith, J. B., Lockwood, R., Anderson, J. G., Dessler, A. E., Keutsch, F. N., Spackman, J. R., Read, W. G., and Bui, T. P.: Observations of deep convective influence on stratospheric water vapor and its isotopic composition, Geophysical Research Letters, 34, L04814, https://doi.org/10.1029/2006GL027899, http://doi.wiley.com/ 10.1029/2006GL027899, 2007.
 - Herman, R. L., Ray, E. A., Rosenlof, K. H., Bedka, K. M., Schwartz, M. J., Read, W. G., Troy, R. F., Chin, K., Christensen, L. E., Fu, D. J., Stachnik, R. A., Bui, T. P., and Dean-Day, J. M.: Enhanced stratospheric water vapor over the summertime continental United States and the role of overshooting convection, Atmospheric Chemistry and Physics, 17, 6113–6124, https://doi.org/10.5194/acp-17-6113-2017, 2017.
- 255 Homeyer, C. R.: Formation of the Enhanced-V Infrared Cloud-Top Feature from High-Resolution Three-Dimensional Radar Observations, Journal of the Atmospheric Sciences, 71, 332–348, https://doi.org/10.1175/JAS-D-13-079.1, http://journals.ametsoc.org/doi/abs/10.1175/ JAS-D-13-079.1, 2014.
 - Homeyer, C. R. and Kumjian, M. R.: Microphysical Characteristics of Overshooting Convection from Polarimetric Radar Observations, Journal of the Atmospheric Sciences, 72, 870–891, https://doi.org/10.1175/JAS-D-13-0388.1, https://journals.ametsoc.org/doi/pdf/10.1175/
- 260 JAS-D-13-0388.1, 2015.
 - Kim, J., Randel, W. J., and Birner, T.: Convectively Driven Tropopause-Level Cooling and Its Influences on Stratospheric Moisture, Journal of Geophysical Research: Atmospheres, 123, 590–606, https://doi.org/10.1002/2017JD027080, https://agupubs.onlinelibrary.wiley.com/ doi/abs/10.1002/2017JD027080, 2018.

Liu, N. and Liu, C.: Global distribution of deep convection reaching tropopause in 1 year GPM observations, Journal of Geophysical Re-

search, 121, 3824–3842, https://doi.org/10.1002/2015JD024430, 2016.

Livesey, N., Read, W., Wagner, P., Froidevaux, L., Lambert, A., Manney, G., Millán, L., Pumphrey, H., Santee, M., Schwartz, M., et al.: Earth Observing System (EOS) Aura Microwave Limb Sounder (MLS), Tech. rep., NASA Jet Propulsion Laboratory, https://doi.org/JPL D-33509, http://disc.gsfc.nasa.gov/, 2018.

Mote, P. W., Rosenlof, K. H., Mcintyre, E., Carr, E. S., Gille, J. C., Holton, R., Kinnersley, S., Pumphrey, H. C., Russell, M., and Wal, J. W.:

- 270 An Atmospheric Tape Recorder, Journal of Geophysical Research: Atmospheres, 101, 3989–4006, https://doi.org/10.1029/95JD03422, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/95JD03422, 1996.
 - Ramaswamy, V., Schwarzkopf, M. D., and Randel, W. J.: Fingerprint of ozone depletion in the spatial and temporal pattern of recent lowerstratospheric cooling, Nature, 382, 616, https://doi.org/10.1038/382616a0, 1996.

Randel, W. J., Wu, F., Oltmans, S. J., Rosenlof, K., and Nedoluha, G. E.: Interannual changes of stratospheric water vapor and corre-

- 275
 lations with tropical tropopause temperatures, Journal of the Atmospheric Sciences, 61, 2133–2148, https://doi.org/10.1175/1520-0469(2004)061<2133:ICOSWV>2.0.CO;2, http://journals.ametsoc.org/doi/abs/10.1175/1520-0469(2004)061{%}3C2133:ICOSWV{%}3E2.0.CO;2, 2004.
 - Randel, W. J., Zhang, K., and Fu, R.: What controls stratospheric water vapor in the NH summer monsoon regions?, Journal of Geophysical Research: Atmospheres, 120, 7988–8001, https://doi.org/10.1002/2015JD023622, http://doi.wiley.com/10.1002/2015JD023622, 2015.
- 280 Schoeberl, M. R., Jensen, E. J., Pfister, L., Ueyama, R., Wang, T., Selkirk, H., Avery, M., Thornberry, T., and Dessler, A. E.: Water Vapor, Clouds, and Saturation in the Tropical Tropopause Layer, Journal of Geophysical Research: Atmospheres, 124, 3984–4003, https://doi.org/10.1029/2018JD029849, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD029849, 2019.
 - Schwartz, M. J., Read, W. G., Santee, M. L., Livesey, N. J., Froidevaux, L., Lambert, A., and Manney, G. L.: Convectively injected water vapor in the North American summer lowermost stratosphere, Geophysical Research Letters, 40, 2316–2321, https://doi.org/10.1002/grl.50421, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/grl.50421, 2013.
 - Sherwood, S. C. and Dessler, A. E.: On the control of stratospheric humidity, Geophysical Research Letters, 27, 2513–2516, 2000.

285

290

Shindell, D. T.: Climate and ozone response to increased stratospheric water vapor, Geophysical Research Letters, 28, 1551–1554, https://doi.org/10.1029/1999GL011197, 2001.

Smith, C. A., Haigh, J. D., and Toumi, R.: Radiative forcing due to trends in stratospheric water vapour, Geophysical Research Letters, 28, 179–182, 2001.

- Smith, J. B., Wilmouth, D. M., Bedka, K. M., Bowman, K. P., Homeyer, C. R., Dykema, J. A., Sargent, M. R., Clapp, C. E., Leroy, S. S., Sayres, D. S., Dean-Day, J. M., Paul Bui, T., and Anderson, J. G.: A case study of convectively sourced water vapor observed in the overworld stratosphere over the United States, Journal of Geophysical Research: Atmospheres, 122, 9529–9554, https://doi.org/10.1002/2017JD026831, 2017.
- 295 Solomon, D. L., Bowman, K. P., Homeyer, C. R., Solomon, D. L., Bowman, K. P., and Homeyer, C. R.: Tropopause-Penetrating Convection from Three-Dimensional Gridded NEXRAD Data, Journal of Applied Meteorology and Climatology, 55, 465–478, https://doi.org/10.1175/JAMC-D-15-0190.1, http://journals.ametsoc.org/doi/10.1175/JAMC-D-15-0190.1, 2016.

Solomon, S., Rosenlof, K. H., Portmann, R. W., Daniel, J. S., Davis, S. M., Sanford, T. J., and Plattner, G. K.: Contributions of stratospheric water vapor to decadal changes in the rate of global warming, Science, 327, 1219–1223, https://doi.org/10.1126/science.1182488, 2010.

- 300 Stenke, A. and Grewe, V.: Simulation of stratospheric water vapor trends: impact on stratospheric ozone chemistry, Atmospheric Chemistry and Physics, 5, 1257–1272, https://doi.org/10.5194/acp-5-1257-2005, https://www.atmos-chem-phys.net/5/1257/2005/, 2005.
 - Sun, Y. and Huang, Y.: An examination of convective moistening of the lower stratosphere using satellite data, Earth and Space Science, 2, 320–330, https://doi.org/10.1002/2015EA000115, http://doi.wiley.com/10.1002/2015EA000115, 2015.
- Ueyama, R., Jensen, E. J., Pfister, L., and Kim, J.-E.: Dynamical, convective, and microphysical control on wintertime distributions
 of water vapor and clouds in the tropical troppause layer, Journal of Geophysical Research: Atmospheres, 120, 10,483–10,500, https://doi.org/10.1002/2015JD023318, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JD023318, 2015.
 - Ueyama, R., Jensen, E. J., and Pfister, L.: Convective Influence on the Humidity and Clouds in the Tropical Tropopause Layer During Boreal Summer, Journal of Geophysical Research: Atmospheres, 123, 7576–7593, https://doi.org/10.1029/2018JD028674, http://doi.wiley.com/ 10.1029/2018JD028674, 2018.

- 310 Wang, X., Dessler, A. E., Schoeberl, M. R., Yu, W., and Wang, T.: Impact of convectively lofted ice on the seasonal cycle of water vapor in the tropical tropopause layer, Atmospheric Chemistry and Physics, 19, 14621–14636, https://doi.org/10.5194/acp-19-14621-2019, https://www.atmos-chem-phys.net/19/14621/2019/, 2019.
 - Waters, J. W., Froidevaux, L., Harwood, R. S., Jarnot, R. F., Pickett, H. M., Read, W. G., Siegel, P. H., Cofield, R. E., Filipiak, M. J., Flower,D. A., Holden, J. R., Lau, G. K., Livesey, N. J., Manney, G. L., Pumphrey, H. C., Santee, M. L., Wu, D. L., Cuddy, D. T., Lay, R. R., Loo,
- M. S., Perun, V. S., Schwartz, M. J., Stek, P. C., Thurstans, R. P., Boyles, M. A., Chandra, K. M., Chavez, M. C., Chen, G. S., Chudasama, B. V., Dodge, R., Fuller, R. A., Girard, M. A., Jiang, J. H., Jiang, Y., Knosp, B. W., Labelle, R. C., Lam, J. C., Lee, K. A., Miller, D., Oswald, J. E., Patel, N. C., Pukala, D. M., Quintero, O., Scaff, D. M., Van Snyder, W., Tope, M. C., Wagner, P. A., and Walch, M. J.: The Earth Observing System Microwave Limb Sounder (EOS MLS) on the aura satellite, IEEE Transactions on Geoscience and Remote Sensing, 44, 1075–1092, https://doi.org/10.1109/TGRS.2006.873771, 2006.
- 320 World Meteorological Organization: A three-dimensional science: Second session of the commission for aerology, WMO bull, 4, 134–138, 1957.
 - Wright, J. S., Fu, R., Fueglistaler, S., Liu, Y. S., and Zhang, Y.: The influence of summertime convection over Southeast Asia on water vapor in the tropical stratosphere, Journal of Geophysical Research: Atmospheres, 116, https://doi.org/10.1029/2010JD015416, 2011.



Figure 1. (A) Time series of 100-hPa (red line) MLS water vapor anomaly (zonal mean removed), (blue line) convective occurrence from GridRad, and (green line) the convective influence ratio in the back trajectory experiments. All data are 5-day averages over the NA region (25°N-50°N, 70°W-130°W) during June, July, August 2005-2016. For the convective frequency, we use linear interpolation to estimate the value at 100 hPa. The convective influence ratio is the fraction of the MLS observations that encounter deep convection, as determined by the back trajectory calculations. (B-D) Joint distribution of convection and water vapor time series during 2005-2016 divided into (B) June, (C) July, and (D) August. Solid lines show the linear fit, and dashed lines show the 95% significant level margin of error of the slope bar (accounting for auto-correlation). To account for the time for the water vapor to spread out, each data point is a 10-day average of convection, with water vapor averaged over the last five days of the averaging period for convection.



Figure 2. Probability density function of MLS water vapor observations at 100 hPa averaged over NA $(25^{\circ}N-50^{\circ}N, 70^{\circ}W-130^{\circ}W)$ in (A) June, (B) July, and (C) August 2005-2016. The observations are divided into two groups: (red) those whose back trajectory encounters convection and (grey) those that do not.



Figure 3. (Top row) Distribution of the 100-hPa GridRad deep convection occurrence averaged over 2005-2016 in (A) June, (B) July, and (C) August. The black contour in each panel (repeated in each row) is the 10^{-5} contour of GridRad convective occurrence, averaged over that month. (Second row) Geographical distribution of the MLS 100-hPa water vapor anomaly (after removal of the zonal mean), averaged over 2005-2016 in (D) June, (E) July, and (F) August. (Third row) Geographical distribution of the convection influence ratio over the NA region during (G) June, (H) July and (I) August 2005-2016. The black dashed contour (repeated in panels G-L) is the water vapor anomaly contours matching the shading in the corresponding upper panel. (Fourth row) Location where convectively influenced parcels encounter convection during (M) June, (N) July, and (O) August. (The stream lines are horizontal velocities interpolated onto 100 hPa using the cubic spline method from ERAi data averaged over the same period.



Figure 4. (A) PDF of lowest latitude; (B) PDF of hourly ERAi daily temperature (K) interpolated to the location of the parcels; (C) PDF of minimum saturated water vapor (ppmv). The minimum water saturated water vapor mixing ratio and the lowest latitude are the minimum values along the path after the parcels encounter deep convection in the back trajectory model and prior to being observed by MLS.



Figure 5. Ten day average water vapor anomaly (ppmv) at 100 hPa (after removing the zonal mean) during (top) June 2010 and (bottom) June 2011. Thick black contours are the 10^{-5} contours of GridRad deep convection, and stream lines are horizontal velocity at 100 hPa obtained from ERAi averaged over the same period.



Figure 6. PDF of (A) lowest latitude along the path in the back trajectory experiments (B) mean tropical temperature (K) obtained from ERAi ($0^{\circ}-20^{\circ}$ N, 70° W-130°W), and (C) minimum saturation water vapor mixing ratio (ppmv) in the back trajectory experiment results. Minimum saturation water vapor mixing ratios and lowest latitude are the minimum values along the path after the parcels encounter deep convection in the back trajectory model and prior to being observed by MLS.