

Author's response by Wenjie Wang et al.

Corresponding to li_xin@pku.edu.cn.

We greatly appreciate the time and efforts that the Referees spent in reviewing our manuscript. The comments are really thoughtful and helpful to improve the quality of our paper. We have addressed each comment below, with the Referee comment in black text, our response in blue text, and relevant manuscript changes noted in red text.

1. Lines 541-558. This part is inserted upon comment/revision but is a bit difficult to read. In this part ozone "DECREASE" is described as sensitivities with aerosol "INCREASES", while in the previous part ozone change (mostly increase) is described with aerosol "TEMPORAL" change (i.e., aerosol decrease). Although most of the sentences here were taken directly from Abstract of Gao et al. (2020), I would suggest rewriting them in the context of this manuscript, to avoid any confusion or misunderstanding.

Response: Many thanks, I have revised it according to your suggestions.

Line 542-561:

Quantitative studies suggested that, the impact of aerosols via affecting photolysis rates on net ozone production (Cai, 2013; Wang et al., 2019; Castro et al., 2001) is more than on surface ozone concentrations (Jacobson, 1998; Li et al., 2011a; Li et al., 2011b; Wang et al., 2016a). Moreover, several different transport model studies (Xing et al., 2017; Li et al., 2018a; Li et al., 2019c) also show that the impact of PM_{2.5} on summer surface ozone concentrations is not important, which is different from the relatively large change of net ozone production in this study. There are several possible causes of this difference: (1) The net ozone production characterizes the local ozone production level but can't completely represent ozone concentrations that are also influenced by regional transport of ozone (Streets et al., 2007; Wang et al., 2020; Moghani et al., 2018). (2) The light-extinction effect of aerosols lead to ozone at the top of PBL being entrained by turbulence to the surface to partly counteract the reduction in surface ozone photochemical production induced by aerosols (Gao et al., 2020). (3) AOD decreased by 58% in Beijing in this study, which is higher than that of regional AOD decrease in eastern China (20%) (Li et al.,

2019a). (4) The photolysis rates in this study are determined using the observed AOD SSA, and confirmed by ground-level measurements of photolysis frequencies, while photolysis rates are determined using online algorithm in other transport model studies. The different calculation procedures also contribute to the different results between our study and other transport model studies.

2. Line 533. above PBL (not below PBL)?

Response: Many thanks, I have revised it.

Line: 533-535:

The crossing point between $j(\text{NO}_2)$ profile of 2006 and zero AOD profile is above PBL, while in 2016 the $j(\text{NO}_2)$ profile crosses the zero AOD profile within the PBL.

3. Line 567 (previous Line 494): Please respond to the comment from the referee #1: Please take caution when saying "ozone increase". you mean surface ozone concentrations?

Response: Many thanks, I have revised it.

Line 567-570:

Consistent with the implementation of stringent emission control measures, concentrations of $\text{PM}_{2.5}$ and ozone precursors (VOCs and NO_x) decreased rapidly, but in contrast surface O_3 and O_x concentrations increased.

4. Line 278. Remove "was"

Response: Many thanks, I have revised it.

**Exploring the drivers of the increased ozone production in Beijing in
summertime during 2005-2016**

Wenjie Wang¹, David D. Parrish², Xin Li^{1,3,4*}, Min Shao^{2,1}, Ying Liu¹, Sihua Lu¹, Min
Hu¹, Yusheng Wu^{1,#}, Limin Zeng¹, Yuanhang Zhang¹

¹ State Key Joint Laboratory of Environmental Simulation and Pollution Control,
College of Environmental Sciences and Engineering, Peking University, Beijing,
China

² Institute for Environmental and Climate Research, Jinan University,
Guangzhou 511443, China

³ International Joint Laboratory for Regional Pollution Control, Ministry of
Education, Beijing, 100816, China

⁴ Collaborative Innovation Centre of Atmospheric Environment and Equipment
Technology, Nanjing University of Information Science & Technology, Nanjing,
210044, China

[#] now at Department of Physics, University of Helsinki, Helsinki, Finland

* Corresponding author.

Address: College of Environmental Sciences and Engineering, Peking
University, Beijing 100871, China

Phone: 86-10-62757973

Email: li_xin@pku.edu.cn

Abstract

In the past decade, average $\text{PM}_{2.5}$ concentrations decreased rapidly under the strong pollution control measures in major cities in China; however, ozone (O_3) pollution emerged as a significant problem. Here we examine a unique (for China) 12-year data set of ground-level O_3 and precursor concentrations collected at an urban site in Beijing (PKUERS), where the maximum daily 8 h average (MDA8) O_3 concentration and daytime Ox ($\text{O}_3 + \text{NO}_2$) concentration in August increased by 2.3 ± 1.2 ppbv ($+3.3 \pm 1.8\%$) yr^{-1} and 1.4 ± 0.6 ($+1.9 \pm 0.8\%$) yr^{-1} respectively from 2005 to 2016. In contrast, daytime concentrations of nitrogen oxides (NO_x) and the OH reactivity of volatile organic compounds (VOCs) both decreased significantly. Over this same time, the decrease of particulate matter, and thus the aerosol optical depth, led to enhanced solar radiation and photolysis frequencies, with near-surface $j(\text{NO}_2)$ increasing at a rate of $3.6 \pm 0.8\%$ yr^{-1} . We use an observation based box model to analyze the combined effect of solar radiation and ozone precursor changes on ozone production rate, $P(\text{O}_3)$. The results indicate that the ratio of the rates of decrease of VOCs and NO_x (about 1.1) is inefficient in reducing ozone production in Beijing. $P(\text{O}_3)$ increased during the decade due to more rapid atmospheric oxidation caused to a large extent by the decrease of particulate matter. This elevated ozone production was driven primarily by increased actinic flux due to $\text{PM}_{2.5}$ decrease and to a lesser extent by reduced heterogeneous uptake of HO_2 . Therefore, the influence of $\text{PM}_{2.5}$ on actinic flux and thus on the rate of oxidation of VOCs and NO_x to ozone and to secondary aerosol (i.e., the major contributor to $\text{PM}_{2.5}$) is important for determining the atmospheric effects of controlling the emissions of the common precursors of $\text{PM}_{2.5}$ and ozone when attempting to control these two important air pollutants.

1 Introduction

Tropospheric ozone (O_3) plays a key role in the oxidizing capacity of the atmosphere and affects the global climate; high concentrations of ground-level ozone are harmful to human health and ecosystems (Monks et al., 2015; Fiore et al., 2009). Ozone is produced rapidly in sun-lit polluted air by photochemical oxidation of volatile organic compounds (VOCs) in the presence of nitrogen oxides ($NO_x \equiv NO + NO_2$) (Atkinson, 2000). In recent years, China has undergone rapid economic development, resulting in higher demand for energy, and greater usage of fossil fuels. As a result, high emissions to the atmosphere produce heavy pollution in eastern China, which now suffers from severe ozone pollution, especially in urban areas, where the daily maximum 8 h average (MDA8) ozone level often exceeds the standard of 80 ppb (Jinfeng et al., 2014; Wang et al., 2011; Zhang et al., 2014; Lu et al., 2018; Li et al., 2019a). A recent study reported that the national warm-season (April–September) fourth highest MDA8 ozone level (86.0 ppb) and the number of days with MDA8 values of > 70 ppb was much higher than regional averages in Japan, South Korea, Europe, or the United States (Lu et al., 2018). Satellite observations found that regional ozone concentrations in eastern China increased by 7% between 2005 and 2010 (Verstraeten et al., 2015). From 2013 to 2017, the O_3 concentrations in 74 cities as a whole showed an upward trend with Beijing-Tianjin-Hebei region being the most serious (Li et al., 2019a; Lu et al., 2018). Better understanding of the causes of elevated ozone in China is important for developing effective emission control strategies to reduce the ozone pollution problem.

Aerosols impact ozone production primarily in two ways: alteration of photolysis rates by aerosol radiative influence and heterogeneous reactions occurring on the aerosol surface. The reduction of photolysis frequencies by the extinction effect of aerosol and thus its influence on ozone production has been explored in the past (Dickerson et al., 1997; Castro et al., 2001; Real and Sartelet, 2011; Gerasopoulos et al., 2012; Wang et al., 2019). Absorbing aerosols reduce photolysis frequencies

throughout the boundary layer, and as a result decrease near-surface photochemical ozone production (de Miranda et al., 2005; Jacobson, 1998; Wendisch et al., 1996; Raga et al., 2001). Conversely, scattering aerosols in the boundary layer increase photolysis frequencies throughout the troposphere, and thereby increase ozone production aloft (Jacobson, 1998; Tian et al., 2019; Dickerson et al., 1997). The importance of aerosol heterogeneous reactions in ozone photochemistry in China has been previously investigated in model studies (Lou et al., 2014; Li et al., 2018b; Xu et al., 2012; Li et al., 2019a). The effects of NO₂, NO₃, and N₂O₅ heterogeneous reactions showed opposite O₃ concentration changes in VOC-limited and NO_x-limited regions. In a VOC-limited region, NO₂, NO₃, and N₂O₅ heterogeneous reactions lead to ozone concentration increases (Lou et al., 2014; Xu et al., 2012). The heterogeneous reaction of HO₂ decreases ozone production in both VOC-limited and NO_x-limited regions by decreasing the reaction rate of HO₂ with NO (Lou et al., 2014; Li et al., 2019a).

In the past decade, Eastern China has experienced severe fine particulate matter (PM_{2.5}) pollution in winter (Zhang et al., 2016), and this issue has been the main focus of the government's air pollution control strategy. These stringent emission control measures have significantly decreased the concentrations of particulate matter in many Chinese cities. During 2008-2013, ground-level PM_{2.5} estimated from satellite-retrieved aerosol optical depth (AOD) in China declined at a rate of 0.46 μg m⁻³ year⁻¹ (Ma et al., 2016b). Another study indicated that the annual average concentration of PM_{2.5} in Beijing decreased by 1.5 μg m⁻³ year⁻¹ and 27% in total from 2000 to 2015 under the implementation of 16 phases' air pollution control measures (Lang et al., 2017). Hu et al (2017) reported that PM_{2.5} in Beijing declined significantly from 2006 to 2016, and meanwhile solar radiation increased (Hu et al., 2017). However, despite the reduction in emissions of particulate matter (PM) and ozone precursors, ozone concentrations increased, even while PM concentrations decreased.

In Beijing, the second largest city in China, with rapid economic development and urbanization in recent years, ozone pollution is one of the worst among China's cities. Thus, Beijing is a representative city in which to study urban ozone pollution in China. Despite extensive study of the relationship between ozone and its precursors in Beijing

and other mega cities in China (Zhang et al., 2014;Chou et al., 2011;Lu et al., 2019;Liu et al., 2012), there remains a lack of understanding of the cause of the long-term surface ozone concentration increase that accompanied reductions in precursor emissions. In this study, we utilize measurements from a representative urban site in Beijing to explore how the variations in solar radiation and heterogeneous reactions influence the trend of ozone and the coupling effect of aerosol and ozone precursor changes on ozone production. Our overall goal is to determine the extent to which increasing actinic flux caused by the decline in PM contributed to the observed increase in ozone concentrations. This research provides a clearer understanding of how efforts to reduce PM concentrations affect ozone concentrations, and thus informs air quality improvement efforts in China's urban areas.

2 Materials and methods

2.1 Measurements of air pollutants, photolysis frequencies and aerosol surface concentration

Ambient air pollutants and photolysis frequencies were measured at an urban site in Beijing in August between 2005 and 2016. The site (39.99° N, 116.31°E) was located on the roof of a six story building (~20m above the ground level) on the campus of Peking University (PKUERS) near the 4th Ring Road with high density of traffic, but without obvious industrial or agricultural sources (Wehner et al., 2008). Temporal trends of air pollutants and composition of VOCs are thought to be representative for the whole of Beijing (Wang et al., 2010;Xu et al., 2011;Zhang et al., 2012). Measured parameters include O₃, NO_x, CO, SO₂, C₂ - C₁₀ VOCs, photolysis frequencies and aerosol surface concentration. The measurement techniques are included in the Table 1.

During 2006 and 2008, ambient levels of VOCs were measured using an online GC-FID system built by the Research Center for Environmental Changes (RCEC; Taiwan). A detailed description of this system and QA/QC procedures can be found in Wang et al. (Wang et al., 2004). During August 2007 and 2009, ambient VOCs were

measured using a commercial GC-FID/PID system (Syntech Spectra GC955 series 600/800 analyzer) (Xie et al., 2008;Zhang et al., 2014). From 2010 to 2016, VOCs were measured using a cryogen-free online GC-MS/FID system developed by Peking University. A detailed description of this system and QA/QC procedures can be found in Yuan et al. and Wang et al. (Yuan et al., 2012;Wang et al., 2014). Formaldehyde (HCHO) concentrations were measured by a Hantzsch fluorimetry.

Photolysis frequencies (including $j(\text{O}^1\text{D})$, $j(\text{NO}_2)$, $j(\text{HONO})$, $j(\text{HCHO})_{\text{M}}$, $j(\text{HCHO})_{\text{R}}$, $j(\text{H}_2\text{O}_2)$) were calculated from solar actinic flux spectra measured by a spectroradiometer as described by Bohn et al. (Bohn et al., 2008). The particle number size distributions were measured by a system consisting of a Nano-SMPS (TSI DMA3085 + CPC3776) and a SMPS (TSI DMA3081 + CPC3775). Aerosol surface concentration (Sa) during 2006-2016 was calculated from the measured particle number size distributions between 3 nm and 700 nm by assuming the particles are spherical in shape.

2.2 Estimate of photolysis frequencies

Photolysis frequencies were measured in August 2011-2014 and 2016. The Tropospheric Ultraviolet and Visible (TUV) radiation model (version 5.3) was used to calculate photolysis frequencies in August over the entire 2006-2016 period under clear-sky conditions. TUV uses the discrete-ordinate algorithm (DISORT) with four streams and calculates the actinic flux spectra with a wavelength range of 280 – 420 nm in 1 nm steps and resolution. We used observed aerosol optical properties including AOD, single scattering albedo (SSA) and Ångström exponent (AE), total ozone column to constrain the TUV model (Madronich, 1993). The calculated values agree well with measured results as shown in Figure 1 indicating that the TUV model accurately calculated the photolysis frequencies. Data of photolysis frequencies under cloudless conditions were selected according to the presence of AOD data since AOD measurements were not possible under cloudy conditions.

2.3 Measurements of aerosol optical properties

Aerosol optical properties were measured with a CIMEL Sun photometer (AERONET level 1.5 and level 2.0 data collection, <http://aeronet.gsfc.nasa.gov/>) at the Beijing-CAMS site (39.933°N, 116.317°E) and at the Beijing site (39.977N, 116.381E). The instrumentation, data acquisition, retrieval algorithms and calibration procedure, which conform to the standards of the AERONET global network, are described in detail by Fottiadi et al. (Fottiadi et al., 2006). The solar extinction measurements taken every 3 minutes within the spectral range 340 – 1020 nm were used to compute AOD at 340, 380, 440, 500, 675, 870, 970 and 1020 nm. The overall uncertainty in AOD data under cloud-free conditions was 0.02 at a wavelength of 440 nm (Dubovik and King, 2000). In this study, AOD at the wavelength of 380nm was chosen for analysis. This wavelength was selected as it is more representative of $j(\text{NO}_2)$. In addition to AOD, that network also provided single scattering albedo (SSA) and Ångström exponent (AE) data.

Cloud optical thickness (COT) was acquired from Aura satellite measurements with a time resolution of 24 hours. Total ozone column was obtained by OMI (Ozone Monitoring Instrument), using overpass data.

2.4 Trend analysis method

A simple linear regression (the least-squares method) was implemented to investigate temporal trends of ozone, precursors, aerosol optical properties, $\text{PM}_{2.5}$ and photolysis frequencies. The null hypothesis is that air pollutants and time have no linear relationship and this was tested using the standard F-statistic test (ratio of the mean-square regression to the mean-square residual). The p value associated with the F-statistic is the probability of mistakenly rejecting the null hypothesis (** $p < 0.01$; * $p < 0.05$). The p values for the trends of different parameters are summarized in Table 2.

2.5 Chemical box model

Ozone production rate, $P(O_3)$, is calculated by a chemical box model. This model is based on the compact Regional Atmospheric Chemical Mechanism version 2 (RACM) described by Goliff et al. (Goliff et al., 2013), which includes 17 stable inorganic species, 4 inorganic intermediates, 55 stable organic compounds and 43 intermediate organic compounds. Compounds that are not explicitly treated in the RACM are lumped into species with similar functional groups. The isoprene mechanism includes a more detailed mechanism based on the Leuven Isoprene Mechanism (LIM) proposed by Peeters et al. (Peeters et al., 2009). A detailed description of this model can be found in Tan et al. (Tan et al., 2017).

In this study, the model was constrained by measured hourly average CO, NO₂, O₃, SO₂, NMHCs (56 species), HCHO, photolysis frequencies, temperature, pressure, and relative humidity. HONO was not measured. HONO concentrations are generally underestimated by the gas phase reaction source of HONO ($OH + NO \rightarrow HONO$) in urban areas due to the emission of HONO and the heterogeneous reaction of NO_x at surfaces to form HONO, both of which are related to NO_x concentration. As a result, the HONO concentration was calculated according to the concentration of NO₂ and the observed ratio of HONO to NO₂ at an urban site in Beijing, which had a marked diurnal cycle (Hendrick et al., 2014). For the model calculation, the ratio of HONO to NO₂ is equal to 0.08 at 6:00 and decreases linearly from 0.08 to 0.01 during 6:00 - 10:00 reflecting increasing photolysis of HONO, and maintains the value of 0.01 during 10:00-18:00. In this study, we focused on daytime $P(O_3)$ (6:00 - 18:00), thus the nocturnal HONO concentrations were not required.

RO₂, HO₂, OH were simulated by the box model to calculate the ozone production and loss rates as shown in Equations E1 and E2 as derived by Mihelcic et al. (Mihelcic et al., 2003).

$$P(O_3) = k_{HO_2+NO} [HO_2][NO] + \sum (k_{RO_2+NO}^i [RO_2^i][NO]) - k_{OH+NO_2} [OH][NO_2] - L(O_3) \quad E1$$

$$L(O_3) = (\theta j(O^1D) + k_{OH+O_3} [OH] + k_{HO_2+O_3} [HO_2] + \sum (k^j_{alkene+O_3} [alkene^j]) [O_3] \quad E2$$

where θ is the fraction of O^1D from ozone photolysis that reacts with water vapor. i and j represent the number of species of RO_2 and alkenes, respectively.

The model runs were performed in a time-dependent mode with two days' spin-up. A 24 h lifetime was introduced for all simulated species, such as secondary species and radicals, to approximately simulate dry deposition and other losses of these species (Lu et al., 2013). This lifetime corresponds to an assumed deposition velocity of 1.2 cm s^{-1} and a well-mixed boundary layer height of about 1 km. Sensitivity tests show that this assumed deposition lifetime has a relatively small influence on the reactivity of modeled oxidation products and RO_x radicals.

Aerosols can influence O_3 production by heterogeneous reactions such as uptake of HO_2 , NO_2 , N_2O_5 and NO_3 . For these species, the heterogeneous uptake of HO_2 is expected to have the largest effect on rapid ozone production in summertime and VOC-limited conditions (Li et al., 2019a). Thus, the effect of heterogeneous reaction of HO_2 on ozone production was simulated in the chemical box model using RH corrected aerosol surface concentration (S_{aw}) and uptake coefficient of HO_2 . The rate of change in HO_2 due to irreversible uptake is expressed by E3.

$$\frac{dC}{dt} = \frac{\gamma_{HO_2} \times S_{aw} \times v \times C}{4} \quad E3$$

Where C , v , and γ_{HO_2} are the gas phase concentration, mean molecular velocity, and uptake coefficient, respectively. To derive S_{aw} we used the measured hygroscopic factor (Liu et al., 2009) and measured RH to correct the measurement-derived S_a to ambient conditions. In this study, we chose $\gamma_{HO_2} = 0.2$ provided by laboratory measurements of HO_2 uptake by aerosol particles collected at two mountain sites in eastern China (Taketani et al., 2012). The effects of HO_2 uptake on $P(O_3)$ in Beijing in 2006 were simulated assuming that the product of HO_2 uptake by aerosols is either H_2O or H_2O_2 . The results indicate that the two scenarios showed no significant difference because the recycling of HO_x radicals from H_2O_2 is inefficient (Li et al., 2019a). In the following simulations in this study, the product of HO_2 uptake by aerosols is taken to be H_2O .

Heterogeneous uptake of N_2O_5 , NO_2 and NO_3 was included in the chemical box

model. This includes $\gamma_{\text{N}_2\text{O}_5} = 0.007$ for converting N_2O_5 to HNO_3 (Wang et al., 2017), $\gamma_{\text{NO}_2} = 1 \times 10^{-5}$ for conversion of NO_2 to HONO and HNO_3 (which yields a good simulation of HONO/ NO_2 concentration ratios in China (Shah et al., 2020)) and $\gamma_{\text{NO}_3} = 1 \times 10^{-3}$ for conversion of NO_3 to HNO_3 (Jacob, 2000).

3 Results and discussion

3.1 Trend of ozone

Ozone pollution levels can be characterized by a number of metrics. Table 3 lists 10 ozone metrics and their definition summarized by Lu et al. (2018). We classify these indicators into four categories: (1) metrics that characterize general levels of ozone: median value of hourly ozone concentrations (median), daily maximum 8 h average ozone concentration (MDA8) and daytime average ozone concentration (DTAvg); (2) metrics that characterize extreme levels of ozone: daily maximum 1 h average ozone concentration (MDA1), 98th percentile of hourly ozone concentrations (Perc98) and 4th highest MDA8 (4MDA8); (3) metrics that characterize ozone exposure: cumulative hourly ozone concentrations of >40 ppb (AOT40) and sum of positive differences between MDA8 and a cutoff concentration of 35 ppb (SOMO35); (4) The metrics that characterize the days when the ozone exceeds the standard: total number of days with MDA8 values of >70 ppb (NDGT70) and number of days with the ozone concentration exceeding the Chinese grade II national air quality standard (Exceedance). Figure 2 presents variations in these four categories of ozone metrics at PKUERS site during the study periods. The results show that overall all metrics increased during the 12 year period. However, the percent increase, the p value and the correlation coefficient vary between metrics. The median, DTAvg, and MDA8 indicators, which characterize the general concentration levels of ozone, increased at rates of 2.8% - 5.7% yr^{-1} . The metrics that characterize the extreme concentration levels of ozone increased more slowly (1.2% - 2.7% yr^{-1}). Among them, Perc98 had the smallest rate of increase, only 1.2% yr^{-1} , and the correlation is not significant ($p = 0.29$, $r^2 = 0.11$). This indicates that

increases in the extreme ozone pollution was less significant. In contrast, the increase rates of the ozone exposure metrics AOT40 and SOMO35 are faster, $8.4\% \text{ yr}^{-1}$ and $8.3\% \text{ yr}^{-1}$, respectively, than the metrics that characterize ozone concentrations. The NDGT70 and Exceedance metrics, related to the number of days of ozone exceeding the standard, showed the fastest increases, $10\% \text{ yr}^{-1}$ and $9.8\% \text{ yr}^{-1}$, respectively.

As shown in Figure 3, from 2005 to 2016 MDA8 O₃ concentrations increased at a rate of $2.3 \pm 1.2 \text{ ppbv}$ ($3.3 \pm 1.8 \%$) yr^{-1} ($r^2 = 0.66$) at the PKUERS site, which corresponds to a total MDA8 ozone concentration increase of 25.3 ppbv. Meanwhile, O_x (O₃+NO₂) concentrations increased at a slower rate of $1.4 \pm 0.6 \text{ ppbv}$ ($1.9 \pm 0.8 \%$) yr^{-1} , due to the decrease in NO_x concentrations (Figure 5).

Temperature and wind speed, which can directly influence ozone production and concentrations, showed no significant trend during 2005-2016 (Figure 4). The average temperatures in summer were between 26 and 31°C. The temperature in 2005 was the lowest and in 2007 it was the highest. The average wind speeds were less than 2.5 m s^{-1} in all years. The average relative humidity may have decreased slightly ($\sim 1.5\% \text{ yr}^{-1}$). In summary, we believe that meteorological factors did not play more than a minor role in the overall Beijing O₃ trend. Therefore, our discussion focuses on photochemical processes.

The ozone concentration observed at a receptor site depends on two contributions: regional background ozone and local photochemical production. We have no direct measurements of the long-term trend of regional background ozone in Beijing, but others have reported measurements of ozone at regional background sites in China. At a baseline Global Atmospheric Watch (GAW) station in the northeastern Tibetan Plateau region (Mt Waliguan, 36.28° N, 100.9° E) the average annual daytime ozone concentration increased at a rate of 0.24 ppb yr^{-1} , over the 1994 to 2013 period, but there was no significant trend in summer (Xu et al., 2018). The measurement at a rural station (Dingling site) in Beijing (116.22° E, 40.29° N, 34 km northwest of the observation site in this study) showed a decrease of ozone at a rate of $-0.47 \text{ ppb yr}^{-1}$ over the 2004 to 2015 period (Zheng et al., 2016). The MDA8 ozone concentration at

the Shangdianzi site, a background station in Beijing, showed an increasing trend of 1.1 ppb yr⁻¹ during 2004-2014 (Ma et al., 2016a). Additionally, there were very small trends of O₃ concentrations at the background site (Dongtan) in Shanghai, located to the south of the North China Plain (Gao et al., 2017). However, these background sites in Beijing and Shanghai may be strongly affected by local emissions. MDA8 ozone concentrations at the Changdao site, a background site in the east of the North China Plain that is much less influenced by local emissions, increased slowly (+1.2 ppbv yr⁻¹, r²=0.11), but that rate is not statistically significant (p = 0.25) during 2013-2019 (Figure S1). Based on these reports of smaller and variable trends, we assume that the trend in regional background ozone in the North China Plain made only a minor contribution to the relatively larger ozone trend observed at the PKUERS site (+2.3 ± 1.2 ppbv yr⁻¹, r²=0.66, p = 0.001). We thus surmise that the increase in O₃ at the PKUERS site was mainly due to “local” photochemistry driven by emissions of ozone precursors from the central urban and surrounding suburban areas of Beijing.

3.2 Trend of gaseous precursors

This increase in ozone concentrations is opposite to the decreasing trend of its precursors, including VOCs, CO and NO_x (Figure 5). The overall change of the total OH loss rate due to VOCs (VOC reactivity) was -0.36 s⁻¹ (-6.0%) yr⁻¹. For anthropogenic VOCs, the highest reactivity was generally contributed by alkene species, with an average value over the eleven years of 2.00 ± 0.43 s⁻¹, followed by aromatics and alkanes, with average reactivities of 1.51 ± 0.74 s⁻¹ and 0.92 ± 0.60 s⁻¹, respectively. Thus, the alkenes and aromatics are more important for O₃ production than are alkanes. The trends for alkenes, aromatics, and alkanes were a decrease of 0.14 s⁻¹ (7.1%), 0.12 s⁻¹ (7.9%), and 0.065 s⁻¹ (7.0%) yr⁻¹, respectively, indicating that alkenes and aromatics also played the dominant role in the reduction of anthropogenic VOC reactivity. The rate of decrease in VOCs at PKUERS site is similar to that reported for Los Angeles by Warneke et al. and Pollack et al. (7.3-7.5% yr⁻¹ over 50 years) (Warneke et al., 2012; Pollack et al., 2013). The decrease in anthropogenic

VOCs in Los Angeles was predominantly attributed to decreasing emissions from motor vehicles due to increasingly strict emissions standards. Similarly, a previous study at the PKUERS site indicated that the decreasing anthropogenic VOC was mainly attributed to the reduction of gasoline evaporation and vehicular exhaust under the implementation of stricter emissions standards for new vehicles and specific control measures for in-use vehicles (Wang et al., 2015a). For naturally emitted VOCs, mainly isoprene, the OH reactivity had little trend with large fluctuations, as the emissions of plants vary greatly with temperature and light intensity. Therefore, the decrease in total VOCs reactivity was dominated by the decrease in anthropogenic VOCs. Similarly, CO, which is mainly contributed by anthropogenic emissions, decreased rapidly ($9.3\% \text{ yr}^{-1}$) during 2006–2016.

NO_x data in 2005 were not available. Therefore, the trend of NO_x during 2006–2016 was analyzed. Daytime concentrations of NO_x at the PKUER site also decreased significantly from 2006 to 2016 (Figure 5), with a slope (excluding 2008, which had a much lower NO_x concentration due to enhanced emission controls implemented during the Olympic Games) of $-1.48 \text{ ppbv yr}^{-1}$ ($-5.5\% \text{ yr}^{-1}$, $r^2 = 0.81$). The decrease in NO_x was mainly due to the reduction in vehicle exhaust and coal combustion (Zhao et al., 2013). The decrease in NO_x was significantly faster than that found in Los Angeles by Pollack et al. ($2.6\% \text{ yr}^{-1}$ over 50 years) (Pollack et al., 2013). In contrast to Beijing, Los Angeles O₃ concentrations have continuously decreased from 1980 to 2010 (Parrish et al., 2016). The ratio of the rates of decrease of VOCs and NO_x in Los Angeles (2.9) is significantly greater than unity and larger than that at the PKUER site (1.1), which possibly can be a contributing cause of the opposite trends of ozone in the two regions. It worth noting that the precursor concentrations in 2008, the Olympic Games year, were particularly low, but that ozone was nevertheless on the regression line. The monthly average ratio of VOC reactivity to NO_x concentration in 2008 is $0.28 \text{ s}^{-1} \text{ ppbv}^{-1}$, higher than the average ratio of VOC reactivity to NO_x concentration during 2006–2016 ($0.24 \text{ s}^{-1} \text{ ppbv}^{-1}$). The adverse reduction ratio of VOC to NO_x is the main cause of inefficient reduction in O₃ level in 2008, which is consistent with the study of Chou et al. (2011).

Since 2013, under the implementation of the Action Plan on Air Pollution Prevention and Control (http://www.gov.cn/zwggk/2013-09/12/content_2486773.htm), more stringent emission control measures were implemented to restrict industrial and vehicle emission. As a result, there are indications that both VOCs and NO_x decreased faster over the 2013 to 2016 period: $0.81 \text{ s}^{-1} \text{ yr}^{-1}$ ($16\% \text{ yr}^{-1}$, $r^2 = 0.71$) and $1.94 \text{ ppbv yr}^{-1}$ ($9.3\% \text{ yr}^{-1}$, $r^2 = 0.78$) for VOC reactivity and NO_x, respectively. This could be the cause of the decline in O₃ concentrations from 2014 to 2016.

3.3 Trend of particulate matter

From 2009 to 2016, PM_{2.5} concentrations declined rapidly, achieving the air quality standard of China ($35 \mu\text{g}/\text{m}^3$) in 2016 (Figure 6). Since 2000, Beijing had implemented 16 phases' air pollution control measures, mainly including the controlling of industry, motor vehicle, coal combustion and fugitive dust pollution, which was effective for the reduction in PM_{2.5} (Lang et al., 2017). Especially the strengthening of the reduction in coal combustion, which was gradually replaced by natural gas since 2004, favored improved visibility in Beijing (Zhao et al., 2011).

As shown in Figure 6, from 2006 to 2016 AOD decreased at a rate of $9.3\% \text{ yr}^{-1}$. The correlation between AOD and PM_{2.5} can be determined from the observations of PM_{2.5} and AOD in August during 2009-2016 at the PKUERS site (Figure 7). AOD and PM_{2.5} are linearly correlated with a correlation coefficient of +0.74. This result indicates that the decrease in PM_{2.5} was the primary cause of the reduction in AOD. In addition to PM_{2.5}, relative humidity also has an important effect on AOD. The decrease in relative humidity during 2006-2016 (Figure 4) would reduce the hygroscopic growth of aerosol, leading to a weakened extinction effect of particulate matter on solar radiation (Qu et al., 2015). It is worth noting that although PM_{2.5} in 2011 was lower than that in 2010, AOD in 2011 was higher than that in 2010 (Figure 6). For one reason, the relative humidity in 2011 was higher. Additionally, the aerosol type, atmospheric boundary layer height and the vertical structure of aerosol distribution also affects the dependence of AOD on PM_{2.5} (Zheng et al., 2017),

probably contributing to the scatter about the AOD versus PM_{2.5} relationship shown in Figure 7.

Monthly mean AE (380/550 nm) in August showed no overall trend during 2006-2016 (Figure 8). The monthly AE means were between 0.87 and 1.2, suggesting that the size-distribution of aerosol was generally stable during this period. Monthly mean SSA (440 nm) in August showed an upward trend of $+0.004 \text{ yr}^{-1}$ ($+0.45\% \text{ yr}^{-1}$, $p = 0.001$) during 2006-2016 (Figure 8), indicating the proportion of the light-absorbing component of aerosols (e.g. black carbon) has decreased, due to the stringent and effective controls on the burning of biomass/biofuel and coal (Ni et al., 2014; Cheng et al., 2013). This result is consistent with the studies of Lang et al. and Wang et al., which indicated that black carbon in China's mega cities has decreased rapidly over the past decade (Wang et al., 2016b; Lang et al., 2017).

3.4 Trend of photolysis frequencies

The influence of solar radiation on O₃ photochemistry can be described by actinic flux (or photolysis frequencies). We chose $j(\text{NO}_2)$ as a representative photolysis frequency to analyze the trend of actinic flux. Wang et al (2019) studied the quantitative relationship between $j(\text{NO}_2)$ and AOD at the PKUERS site, and found that $j(\text{NO}_2)$ and AOD showed a clear nonlinear negative correlation at a given SZA, with slopes ranging from -1.3 to $-3.2 \times 10^{-3} \text{ s}^{-1}$ at $\text{AOD} < 0.7$, indicating a significant extinction effect of AOD on actinic flux near the ground.

The $j(\text{NO}_2)$ calculated by the TUV model under clear-sky conditions shows an upward trend of $3.6\% \text{ yr}^{-1}$ from 2005 to 2016 and agrees well with the 5 years of observed values from 2011 to 2016 (Figure 6). According to sensitivity analysis of TUV, the decrease in AOD plays a dominant role in the $j(\text{NO}_2)$ increase, contributing about 80% of the total. Additionally, the increase in SSA also contributes significantly to $j(\text{NO}_2)$ increase, contributing about 17%.

In addition to aerosol optical properties, the photolysis frequency in the planetary boundary layer is affected by other factors, including cloud extinction, ground

reflection, absorption by gases such as O_3 , and Rayleigh scattering by gases. The ground reflection is relatively stable for different years in the same city with stable ground covering. The change in Rayleigh scattering of gases and absorption of NO_2 , SO_2 and $HCHO$ plays a negligible role in the variation in photolysis frequencies according to sensitivity analysis of TUV model. This is consistent with the results of Barnard et al. (Barnard et al., 2004). As shown in Figure 9, the total ozone column fluctuated between 285-307 DU without a significant overall trend. The magnitude of total ozone column variation (22 DU) can change $j(O^1D)$ by about 10%, but plays a negligible role in changing other photolysis frequencies according to sensitivity analysis using the TUV model. The cloud optical thickness (COT) for most years was relatively stable, ranging from 6 to 8, but in 2005, 2012 and 2015 COT was significantly larger (Figure 9). As there was no significant trend of COT, we surmised that the light-extinction effect of clouds did not play a key role in determining the trend of photolysis frequencies.

3.5 Combined effect of changes in ozone precursors and aerosols on ozone production

We investigated the overall effect of the changes in VOCs, NO_x , photolysis frequency, and aerosol uptake of HO_2 on ozone production rate using the chemical box model. We focus on the period during 2006-2016 due to the lack of NO_x data in 2005. By testing the response of $P(O_3)$ as calculated from Equation E1 to the changes of VOCs and NO_x concentrations (Figure 10), we concluded that photochemical environment of the PKUERS site was, on average, in the VOC-limited regime. This result is consistent with previous studies (Zhang et al., 2014; Chou et al., 2011). Under this condition, the long-term decrease in VOCs in Beijing has contributed to a decrease in $P(O_3)$, while the decrease in NO_x has tended to increase $P(O_3)$. As shown in Figure 11, when the increase in photolysis frequencies and aerosol uptake of HO_2 were not included in the calculation, the simulated daytime average $P(O_3)$ decreased slightly at a rate of $1.1\% \text{ yr}^{-1}$. This indicates that the ratio of the rates of decrease of

VOCs and NO_x (about 1.1) is nearly inefficient in reducing ozone production in Beijing. However, when the increase in photolysis frequencies was included in the model calculation, the calculated daytime average P(O₃) showed an increasing trend of 2.2% yr⁻¹. This result indicates that the increase in photolysis frequencies more than compensated for the downward trend of O₃ production driven by decreased VOCs and NO_x, leading to increasing O₃ production through the decade. The photochemical box model calculations indicate that the increase in photolysis frequencies has two major impacts on P(O₃) - an increase in primary production of OH through accelerated photolysis of O₃, HONO, HCHO and other carbonyl compounds, and an accelerated radical recycling of OH as VOCs are oxidized. As particulate matter has decreased and photolysis frequencies correspondingly have increased, a more rapidly decreasing rate of the VOC to NO_x ratio is required to achieve a significant reduction in O₃ in the future.

The simulated P(O₃) in the afternoon hour (12:00-15:00) when ozone production is active and HO_x levels are high increased at a rate of 1.3% yr⁻¹, which is lower than the increasing rate of daytime average P(O₃) (2.2% yr⁻¹) (Figure S2). Hollaway et al. (2019) show that the impacts of aerosols on the summertime photolysis of NO₂ and ozone at surface in Beijing are important before 11:00 am and after 3:00 pm but very limited in afternoon hours due to smaller SZA and lower light absorption of aerosol (i.e. higher SSA) in the afternoon. However, the diurnal variation of simulated P(O₃) in this study indicates that the influence of aerosols on P(O₃) is still significant in the afternoon, leading to average P(O₃) decreased by ~18% (Figure S3), which is slightly lower than the mean daytime decrease (26%). This is because the average AOD in the afternoon (1.4) is significantly higher than that before 11:00 am (0.94) and after 3:00 pm (1.1) despite the smaller SZA and higher SSA.

When we include heterogeneous uptake of HO₂ in the model, the calculated P(O₃) increases at a faster rate of 2.9% yr⁻¹ due to the overall reduced aerosol surface concentration (S_a), which reduces heterogeneous uptake of HO₂ (Figure 11). This result indicates that the effect of heterogeneous uptake of HO₂ contributed roughly

0.7% yr⁻¹ to the P(O₃) increase. Hence, our result indicates that the increase in photolysis rates due to PM decrease plays a more important role than the decrease in heterogeneous uptake of HO₂ by aerosols in accelerating ozone production in Beijing. Previous measurements indicate that the uptake coefficient varies widely from 0.003 to 0.5 with a strong dependence on the aerosol concentration of transition metal ions such as Cu(II) (Zou et al., 2019;Taketani et al., 2008;Lakey et al., 2015;Matthews et al., 2014;Lakey et al., 2016). This strong dependence on aerosol composition implies that a single assumed value for $\gamma_{\text{HO}_2} = 0.2$ has large uncertainty. $\gamma_{\text{HO}_2} = 0.2$ used in our simulation is likely an overestimate of the effect of heterogeneous uptake of HO₂ on ozone production rate at PKUERS site.

A few heterogeneous chemical reactions of nitrogen oxides are thought to be potential influential factors of ozone production. For example, the heterogeneous uptake of NO₂ to produce HNO₃ and HONO, and the heterogeneous uptake of NO₃ and N₂O₅ to produce HNO₃. Our simulation indicates that the reduced heterogeneous uptake of NO_x caused P(O₃) to increase by only ~2.2 % during 2006-2016. Li et al. (2019b) also reported that the effect of heterogeneous uptake of nitrogen oxides on ozone is very small under VOC-limited and summertime conditions in North China Plain. Our simulated result in summertime Beijing, where VOC-limited photochemistry dominates, is consistent with the result of Li et al. (2019b).

In summertime, PM in the Beijing urban area is mainly formed by the secondary conversion of gaseous precursors (Han et al., 2015;Guo et al., 2014), indicating that VOCs and NO_x are not only the precursors of ozone, but also the main precursors of PM in this urban area. In addition, observations in Beijing have shown that the secondary components of PM, including secondary organic matter, ammonium sulfate and ammonium nitrate, dominate the light extinction of PM (Han et al., 2014;Han et al., 2017;Wang et al., 2015b). As a result, reductions of VOCs and NO_x are expected to lead to a decrease in secondary PM formation, and thus to further enhancement in solar radiation (or actinic flux). Therefore, in order to reduce ozone effectively, the contribution of VOCs and NO_x to secondary PM formation and their effect on solar radiation must be comprehensively considered. However, the summertime formation of

PM is quite complex; the conversion efficiency of gaseous precursors to aerosols and the resulting influence on ozone production is a research area that requires further study.

3.6 Additional considerations

One limitation of this study is that the photochemical box model is constrained by surface observations, and hence may not accurately represent some aspects of the photochemistry through the full depth of the planetary boundary layer over Beijing. Here we briefly consider several of these aspects: (1) The treatment of ozone and VOC and NO_x precursor concentrations likely are accurately represented, because rapid daytime vertical mixing ensures that there is only a small vertical gradient in the concentrations of these relatively long-lived species. (2) In daytime, the HONO lifetime is so short that it may be largely confined to near the surface, where it has surface sources (heterogeneous reaction of H₂O and NO₂ and emissions on surfaces). Therefore, the estimated HONO based on near-surface NO₂ concentrations may overestimate average boundary layer HONO concentrations; however, in this study the influence of HONO on the calculation is relatively small, so this is not a large source of error. (3) The model is constrained by surface measurements of photolysis frequencies, but these surface measurements do not accurately quantify the actinic flux throughout the boundary layer. Figure 12 presents the vertical profiles of $j(\text{NO}_2)$ simulated by the TUV model for aerosol properties representative of Beijing. A thick layer of aerosol effectively reduces radiation at the bottom of the layer, but not at the top, where radiation may be enhanced due to upward scattering from the aerosol below (Dickerson et al., 1997; Jacobson, 1998). Overall, vertical average $j(\text{NO}_2)$ increased by 32% from 2006 to 2016, which is comparable to the surface increase (36%). These simulations indicate that the increased trend of $j(\text{NO}_2)$ derived from surface observations do approximate the trend through the entire boundary layer. However, there is a shift in the vertical profile of $j(\text{NO}_2)$ that is important. The crossing point between $j(\text{NO}_2)$ profile of 2006 and zero AOD profile is above PBL, while in 2016 the $j(\text{NO}_2)$ profile crosses the zero AOD profile within the PBL. This

means that as the AOD is reduced further, changes in the vertical average $j(\text{NO}_2)$ will be limited, since increases in $j(\text{NO}_2)$ near the top of the PBL will compensate for decreases near the surface. Additionally, this also denotes that the role of $\text{PM}_{2.5}$ may be more important under condition like 2006, but will be limited under condition like 2016 when there is offsetting effect for PBL ozone by vertical mixing caused by larger ozone vertical gradient (Gao et al., 2020).


Quantitative studies suggested that, the impact of aerosols via affecting photolysis rates on net ozone production (Cai, 2013; Wang et al., 2019; Castro et al., 2001) is more than on surface ozone concentrations (Jacobson, 1998; Li et al., 2011a; Li et al., 2011b; Wang et al., 2016a). Moreover, several different transport model studies (Xing et al., 2017; Li et al., 2018a; Li et al., 2019c) also show that the impact of $\text{PM}_{2.5}$ on summer surface ozone concentrations is not important, which is different from the relatively large change of net ozone production in this study. There are several possible causes of this difference: (1) The net ozone production characterizes the local ozone production level but can't completely represent ozone concentrations that are also influenced by regional transport of ozone (Streets et al., 2007; Wang et al., 2020; Moghani et al., 2018). (2) The light-extinction effect of aerosols lead to ozone at the top of PBL being entrained by turbulence to the surface to partly counteract the reduction in surface ozone photochemical production induced by aerosols (Gao et al., 2020). (3) AOD decreased by 58% in Beijing in this study, which is higher than that of regional AOD decrease in eastern China (20%) (Li et al., 2019a). (4) The photolysis rates in this study are determined using the observed AOD SSA, and confirmed by ground-level measurements of photolysis frequencies, while photolysis rates are determined using online algorithm in other transport model studies. The different calculation procedures also contribute to the different results between our study and other transport model studies.

4 Conclusion

During the past decade, China has devoted very substantial resources to improving the environment. These efforts have improved atmospheric particulate matter loading, but ambient ozone levels have continued to increase. Based on the long-term measurements at a representative site in Beijing, we explored the factors driving the increase in ozone production. Consistent with the implementation of stringent emission control measures, concentrations of PM_{2.5} and ozone precursors (VOCs and NO_x) decreased rapidly, but in contrast surface O₃ and Ox concentrations increased. This investigation finds that the primary cause of the O₃ increase is that decreasing PM concentrations led to an increase in actinic flux, which in turn increased the photochemical production of ozone. This result indicates that the influence of aerosol on ozone production is important for determining the full manifold of atmospheric effects that result from reducing the emissions of the O₃ and PM precursors.

ACKNOWLEDGEMENTS

This work was supported by the Major Program of the National Natural Science Foundation of China [Grant number 91644222, Grant number 91644108]. We thank Hongbin Chen and Philippe Goloub for data management of AOD and other aerosol optical properties on AERONET.



581

582 Figure 1. Correlation between Observed and calculated $j(\text{NO}_2)$ by TUV model in

583 Beijing in summer time during 2012 - 2015.

584





Figure 2. Variations in multiple O₃ metrics at the PKUERS site in Beijing in August between 2005 and 2016.

591
592



593
594
595
596

Figure 3. Variations in average MDA8 O₃ and daytime (7:00-19:00) average O_x in Beijing, August between 2005 and 2016.




Figure 4. Variations in daytime (7:00-19:00) averages of meteorological conditions including temperature (T), relative humidity (RH), wind direction (WD) and wind speed (WS) in Beijing, August during 2005 - 2016.




Figure 5. Variations in geometric mean of daytime NO_x, CO and VOCs reactivity in Beijing, August between 2005 and 2016. VOCs reactivity is depicted by reactivity of each species (left axis) and total VOC reactivity (right axis). On the y-axes, a log-scale is used.





Figure 6. Variations in daytime (7:00-19:00) averages of AOD (380 nm), PM_{2.5}, S_a, j(NO₂) Calculated j(NO₂) by TUV in Beijing, August between 2006 and 2016. AOD and j(NO₂) are both corresponding to cloudless weather. On the y-axes, a log-scale is used for PM_{2.5}, AOD and S_a and a linear scale is used for j(NO₂).



617

618 Figure 7. Correlation between AOD and PM_{2.5} in Beijing, summertime during 2009 -

619 2016.

620




Figure 8. Variation in monthly mean single scattering albedo (SSA) and Ångström exponent (AE) in Beijing for the month of August during 2005 - 2016.






Figure 9. Variations in mean total ozone column and cloud optical thickness (COT) in Beijing for the month of August during 2005 - 2016.



633

634 Figure 10. Sensitivity of monthly daytime mean $P(O_3)$ to VOCs and NO_x simulated
 635 by box model during 2006 - 2016. VOCs and NO_x is increased by 50% or decreased
 636 by 50% to test the fractional change of monthly daytime mean $P(O_3)$.

637



639

640 Figure 11. Trend of monthly daytime mean $P(O_3)$ simulated by the chemical box
 641 model. Red dots: Only the gas-phase reactions are considered in the box model
 642 constrained by observed photolysis frequencies from 2006 for all eleven years. Blue
 643 dots: the box model as above, but constrained by the photolysis frequencies derived
 644 for each year. Green dots: the box model constrained by the photolysis frequencies
 645 derived for each year with the changing aerosol uptake of HO_2 also considered.
 646




Figure 12. Vertical profiles of $j(\text{NO}_2)$ simulated by the TUV model in Beijing. Three scenarios are simulated: The model parameters are: (1) AOD=2.2, SSA=0.91 in August 2006; (2) AOD=0.8, SSA=0.96 in August 2016; (3) AOD=0. The daytime average SZA=53° is used for all simulations. Dotted line represent the top of boundary layer.

654 Table 1. Instruments deployed in the measurement undertaken in August during 2005 -
655 2016 and used for data analysis.

Parameters	Measurement technique	Time resolution	Detection limit	Accuracy
Photolysis frequencies	Spectroradiometer	10 s	/	± 10%
O ₃	UV photometry	60 s	0.5 ppbv	± 5%
NO	Chemiluminescence	60 s	60 pptv	± 20%
NO ₂	Chemiluminescence	60 s	300 pptv	± 20%
CO	IR photometry	60 s	4 ppb	± 5%
SO ₂	Pulsed UV fluorescence	60 s	0.1 ppbv	± 5%
HCHO	Hantzsch fluorimetry	60 s	25 pptv	± 5%
C2-C10VOCs	GC-FID/MS	1 h	20-300 pptv	± 15~20%
PM _{2.5}	TH-2000	60s	1µg m ⁻³	± 5%
S _a	SMPS	60s	/	±3%
AOD, SSA, AE	CIMEL Sun photometer	5min	0.01	±5%

656

657

658 Table 2. p value of temporal trends for different parameters.

Parameter	Period	r ²	p value	P value <0.01?	P value <0.05?
median	2005-2016	0.63	0.002	yes	yes
perc98	2005-2016	0.11	0.288	no	no
DTAvg	2005-2016	0.47	0.014	no	yes
MDA1	2005-2016	0.32	0.057	no	no
MDA8	2005-2016	0.66	0.001	yes	yes
4MDA8	2005-2016	0.42	0.023	no	yes
AOT40	2005-2016	0.67	0.001	yes	yes
NDGT70	2005-2016	0.56	0.005	yes	yes
SOMO35	2005-2016	0.57	0.004	yes	yes
exceedance	2005-2016	0.32	0.054	no	no
Ox	2005-2016	0.38	0.044	no	yes
CO	2005-2016	0.87	0.001	yes	yes
VOC reactivity	2005-2016	0.52	0.006	yes	yes
NO _x	2006-2016	0.81	0.001	yes	yes
Calculated j(NO ₂)	2006-2016	0.94	0.000	yes	yes
AOD (380 nm)	2006-2016	0.78	0.000	yes	yes
PM _{2.5}	2009-2016	0.93	0.000	yes	yes
Sa	2006-2016	0.51	0.010	yes	yes
SSA	2005-2016	0.70	0.001	yes	yes
AE	2005-2016	0.03	0.593	no	no
COT	2005-2016	0.003	0.875	no	no
Total O ₃ column	2005-2016	0.15	0.215	no	no

659

660

661

662

663

664

665

666

667

668

669

670

Table 3. Description of Ozone Metrics used in this study.

categories	metric	definition
general level	median (ppb)	50th percentile of hourly concentrations
	MDA8 (ppb)	daily maximum 8 h average; the mean MDA8 O ₃ in August of each year is used in this study.
	DTAvg (ppb)	daytime average ozone is the average of hourly ozone concentrations for the 12 h period from 07:00 to 19:00 local time
extreme level	MDA1 (ppb)	daily maximum 1 h average; the mean MDA1 O ₃ in August of each year is used in this study.
	Perc98 (ppb)	98th percentile of hourly concentrations
	4MDA8 (ppb)	4th highest MDA8
ozone exposure	AOT40 (ppb h)	cumulative hourly ozone concentrations of >40 ppb
	SOMO35 (ppb day)	sum of positive differences between MDA8 and a cutoff concentration of 35 ppb
Exceedance days	NDGT70 (day)	total number of days with MDA8 values of >70 ppb
	Exceedance (day)	number of days with the ozone concentration exceeding the Chinese grade II national air quality standard, defined as $\text{MDA8} > 160 \mu\text{g m}^{-3}$

674 Reference

- 675 Atkinson, R.: Atmospheric chemistry of VOCs and NO_x, *Atmos. Environ.*, 34, 2063-
676 2101, 2000.
- 677 Barnard, J. C., Chapman, E. G., Fast, J. D., Schmelzer, J. R., Slusser, J. R., and Shetter,
678 R. E.: An evaluation of the FAST-J photolysis algorithm for predicting nitrogen
679 dioxide photolysis rates under clear and cloudy sky conditions, *Atmos. Environ.*,
680 38, 3393-3403, <https://doi.org/10.1016/j.atmosenv.2004.03.034>, 2004.
- 681 Bohn, B., Corlett, G. K., Gillmann, M., Sanghavi, S., Stange, G., Tensing, E.,
682 Vrekoussis, M., Bloss, W. J., Clapp, L. J., Kortner, M., Dorn, H. P., Monks, P. S.,
683 Platt, U., Plass-Dulmer, C., Mihalopoulos, N., Heard, D. E., Clemmshaw, K. C.,
684 Meixner, F. X., Prevot, A. S. H., and Schmitt, R.: Photolysis frequency
685 measurement techniques: results of a comparison within the ACCENT project,
686 *Atmospheric Chemistry and Physics*, 8, 5373-5391, 10.5194/acp-8-5373-2008,
687 2008.
- 688 Cai, Y. F., Wang, T. J., and Xie, M.: Impacts of atmospheric particles on surface ozone
689 in Nanjing (In Chinese), *Climatic Environment Research*, 18, 251-260,
690 10.5194/acp-8-6155-2008, 2013.
- 691 Castro, T., Madronich, S., Rivale, S., Muhlia, A., and Mar, B.: The influence of aerosols
692 on photochemical smog in Mexico City, *Atmos. Environ.*, 35, 1765-1772,
693 [https://doi.org/10.1016/S1352-2310\(00\)00449-0](https://doi.org/10.1016/S1352-2310(00)00449-0), 2001.
- 694 Cheng, Y., Engling, G., He, K. B., Duan, F. K., Ma, Y. L., Du, Z. Y., Liu, J. M., Zheng,
695 M., and Weber, R. J.: Biomass burning contribution to Beijing aerosol, *Atmos.*
696 *Chem. Phys.*, 13, 7765-7781, 10.5194/acp-13-7765-2013, 2013.
- 697 Chou, C. C. K., Tsai, C. Y., Chang, C. C., Lin, P. H., Liu, S. C., and Zhu, T.:
698 Photochemical production of ozone in Beijing during the 2008 Olympic Games,
699 *Atmos. Chem. Phys.*, 11, 9825-9837, 10.5194/acp-11-9825-2011, 2011.
- 700 de Miranda, R. M., Andrade, M. D., and Fattori, A. P.: Preliminary studies of the effect
701 of aerosols on nitrogen dioxide photolysis rates in the city of Sao Paulo, Brazil,
702 *Atmospheric Research*, 75, 135-148, 10.1016/j.atmosres.2004.12.004, 2005.
- 703 Dickerson, R. R., Kondragunta, S., Stenchikov, G., Civerolo, K. L., Doddridge, B. G.,
704 and Holben, B. N.: The Impact of Aerosols on Solar Ultraviolet Radiation and
705 Photochemical Smog, *Science*, 278, 827-830, 10.1126/science.278.5339.827,
706 1997.
- 707 Dubovik, O., and King, M. D.: A flexible inversion algorithm for retrieval of aerosol
708 optical properties from Sun and sky radiance measurements, *Journal of*
709 *Geophysical Research-Atmospheres*, 105, 20673-20696, 10.1029/2000jd900282,
710 2000.
- 711 Fiore, A. M., Dentener, F. J., Wild, O., Cuvelier, C., Schultz, M. G., Hess, P., Textor, C.,
712 Schulz, M., Doherty, R. M., Horowitz, L. W., MacKenzie, I. A., Sanderson, M. G.,
713 Shindell, D. T., Stevenson, D. S., Szopa, S., Van Dingenen, R., Zeng, G., Atherton,
714 C., Bergmann, D., Bey, I., Carmichael, G., Collins, W. J., Duncan, B. N., Faluvegi,
715 G., Folberth, G., Gauss, M., Gong, S., Hauglustaine, D., Holloway, T., Isaksen, I.

- S. A., Jacob, D. J., Jonson, J. E., Kaminski, J. W., Keating, T. J., Lupu, A., Marmer, E., Montanaro, V., Park, R. J., Pitari, G., Pringle, K. J., Pyle, J. A., Schroeder, S., Vivanco, M. G., Wind, P., Wojcik, G., Wu, S., and Zuber, A.: Multimodel estimates of intercontinental source-receptor relationships for ozone pollution, *Journal of Geophysical Research-Atmospheres*, 114, 21, 10.1029/2008jd010816, 2009.
- Fotiadi, A., Hatzianastassiou, N., Drakakis, E., Matsoukas, C., Pavlakis, K. G., Hatzidimitriou, D., Gerasopoulos, E., Mihalopoulos, N., and Vardavas, I.: Aerosol physical and optical properties in the Eastern Mediterranean Basin, Crete, from Aerosol Robotic Network data, *Atmospheric Chemistry and Physics*, 6, 5399-5413, 10.5194/acp-6-5399-2006, 2006.
- Gao, J., Li, Y., Zhu, B., Hu, B., Wang, L., and Bao, F.: What have we missed when studying the impact of aerosols on surface ozone via changing photolysis rates?, *Atmos. Chem. Phys. Discuss.*, 2020, 1-28, 10.5194/acp-2020-140, 2020.
- Gao, W., Tie, X. X., Xu, J. M., Huang, R. J., Mao, X. Q., Zhou, G. Q., and Chang, L. Y.: Long-term trend of O₃ in a mega City (Shanghai), China: Characteristics, causes, and interactions with precursors, *Science of the Total Environment*, 603, 425-433, 10.1016/j.scitotenv.2017.06.099, 2017.
- Gerasopoulos, E., Kazadzis, S., Vrekoussis, M., Kouvarakis, G., Liakakou, E., Kouremeti, N., Giannadaki, D., Kanakidou, M., Bohn, B., and Mihalopoulos, N.: Factors affecting O₃ and NO₂ photolysis frequencies measured in the eastern Mediterranean during the five-year period 2002-2006, *J. Geophys. Res.-Atmos.*, 117, 14, 10.1029/2012jd017622, 2012.
- Goliff, W. S., Stockwell, W. R., and Lawson, C. V.: The regional atmospheric chemistry mechanism, version 2, *Atmospheric Environment*, 68, 174-185, 10.1016/j.atmosenv.2012.11.038, 2013.
- Guo, S., Hu, M., Zamora, M. L., Peng, J., Shang, D., Zheng, J., Du, Z., Wu, Z., Shao, M., Zeng, L., Molina, M. J., and Zhang, R.: Elucidating severe urban haze formation in China, *Proceedings of the National Academy of Sciences*, 111, 17373-17378, 10.1073/pnas.1419604111, 2014.
- Han, T. T., Liu, X. G., Zhang, Y. H., Qu, Y., Gu, J. W., Ma, Q., Lu, K. D., Tian, H. Z., Chen, J., Zeng, L. M., Hu, M., and Zhu, T.: Characteristics of Aerosol Optical Properties and Their Chemical Apportionments during CAREBeijing 2006, *Aerosol Air Qual. Res.*, 14, 1431-1442, 10.4209/aaqr.2013.06.0203, 2014.
- Han, T. T., Liu, X. G., Zhang, Y. H., Qu, Y., Zeng, L. M., Hu, M., and Zhu, T.: Role of secondary aerosols in haze formation in summer in the Megacity Beijing, *Journal of Environmental Sciences*, 31, 51-60, 10.1016/j.jes.2014.08.026, 2015.
- Han, T. T., Xu, W. Q., Li, J., Freedman, A., Zhao, J., Wang, Q. Q., Chen, C., Zhang, Y. J., Wang, Z. F., Fu, P. Q., Liu, X. G., and Sun, Y. L.: Aerosol optical properties measurements by a CAPS single scattering albedo monitor: Comparisons between summer and winter in Beijing, China, *Journal of Geophysical Research-Atmospheres*, 122, 2513-2526, 10.1002/2016jd025762, 2017.
- Hendrick, F., Muller, J. F., Clemer, K., Wang, P., De Maziere, M., Fayt, C., Gielen, C., Hermans, C., Ma, J. Z., Pinardi, G., Stavrou, T., Vlemmix, T., and Van Roozendaal, M.: Four years of ground-based MAX-DOAS observations of HONO

and NO₂ in the Beijing area, *Atmospheric Chemistry and Physics*, 14, 765-781, 10.5194/acp-14-765-2014, 2014.

Hollaway, M., Wild, O., Yang, T., Sun, Y., Xu, W., Xie, C., Whalley, L., Slater, E., Heard, D., and Liu, D.: Photochemical impacts of haze pollution in an urban environment, *Atmospheric Chemistry and Physics*, 19, 9699-9714, 2019.

Hu, B., Zhao, X., Liu, H., Liu, Z., Song, T., Wang, Y., Tang, L., Xia, X., Tang, G., Ji, D., Wen, T., Wang, L., Sun, Y., and Xin, J.: Quantification of the impact of aerosol on broadband solar radiation in North China, *Scientific Reports*, 7, 44851, 10.1038/srep44851, 2017.

Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, 34, 2131-2159, 2000.

Jacobson, M. Z.: Studying the effects of aerosols on vertical photolysis rate coefficient and temperature profiles over an urban airshed, *Journal of Geophysical Research: Atmospheres*, 103, 10593-10604, 10.1029/98jd00287, 1998.

Jinfeng, Keding, Liuju, Zhong, Yubo, Duohong, Chen, Huang, Yuanhang, and Zhang: Fast increasing of surface ozone concentrations in Pearl River Delta characterized by a regional air quality monitoring network during 2006-2011, *Journal of Environmental Sciences*, 26, 23-36, 2014.

Lakey, P. S. J., George, I. J., Whalley, L. K., Baeza-Romero, M. T., and Heard, D. E.: Measurements of the HO₂ Uptake Coefficients onto Single Component Organic Aerosols, *Environ. Sci. Technol.*, 49, 4878-4885, 10.1021/acs.est.5b00948, 2015.

Lakey, P. S. J., George, I. J., Baeza-Romero, M. T., Whalley, L. K., and Heard, D. E.: Organics Substantially Reduce HO₂ Uptake onto Aerosols Containing Transition Metal ions, *Journal of Physical Chemistry A*, 120, 1421-1430, 10.1021/acs.jpca.5b06316, 2016.

Lang, J. L., Zhang, Y. Y., Zhou, Y., Cheng, S. Y., Chen, D. S., Guo, X. U., Chen, S., Li, X. X., Xing, X. F., and Wang, H. Y.: Trends of PM_{2.5} and Chemical Composition in Beijing, 2000-2015, *Aerosol Air Qual. Res.*, 17, 412-425, 10.4209/aaqr.2016.07.0307, 2017.

Li, G., Bei, N., Tie, X., and Molina, L. T.: Aerosol effects on the photochemistry in Mexico City during MCMA-2006/MILAGRO campaign, *Atmos. Chem. Phys.*, 11, 5169-5182, 10.5194/acp-11-5169-2011, 2011a.

Li, J., Wang, Z., Wang, X., Yamaji, K., Takigawa, M., Kanaya, Y., Pochanart, P., Liu, Y., Irie, H., and Hu, B.: Impacts of aerosols on summertime tropospheric photolysis frequencies and photochemistry over Central Eastern China, *Atmos. Environ.*, 45, 1817-1829, 2011b.

Li, J., Chen, X., Wang, Z., Du, H., Yang, W., Sun, Y., Hu, B., Li, J., Wang, W., and Wang, T.: Radiative and heterogeneous chemical effects of aerosols on ozone and inorganic aerosols over East Asia, *Science of the Total Environment*, 622, 1327-1342, 2018a.

Li, J., Chen, X. S., Wang, Z. F., Du, H. Y., Yang, W. Y., Sun, Y. L., Hu, B., Li, J. J., Wang, W., Wang, T., Fu, P. Q., and Huang, H. L.: Radiative and heterogeneous chemical effects of aerosols on ozone and inorganic aerosols over East Asia, *Science of the Total Environment*, 622, 1327-1342,

10.1016/j.scitotenv.2017.12.041, 2018b.

Li, K., Jacob, D. J., Liao, H., Shen, L., Zhang, Q., and Bates, K. H.: Anthropogenic drivers of 2013–2017 trends in summer surface ozone in China, *Proceedings of the National Academy of Sciences*, 116, 422–427, 2019a.

Li, K., Jacob, D. J., Liao, H., Shen, L., Zhang, Q., and Bates, K. H.: Anthropogenic drivers of 2013–2017 trends in summer surface ozone in China, *Proceedings of the National Academy of Sciences of the United States of America*, 116, 422–427, 10.1073/pnas.1812168116, 2019b.

Li, K., Jacob, D. J., Liao, H., Zhu, J., Shah, V., Shen, L., Bates, K. H., Zhang, Q., and Zhai, S.: A two-pollutant strategy for improving ozone and particulate air quality in China, *Nature Geoscience*, 12, 906–910, 10.1038/s41561-019-0464-x, 2019c.

Liu, X., Zhang, Y., Jung, J., Gu, J., Li, Y., Guo, S., Chang, S.-Y., Yue, D., Lin, P., Kim, Y. J., Hu, M., Zeng, L., and Zhu, T.: Research on the hygroscopic properties of aerosols by measurement and modeling during CAREBeijing-2006, *Journal of Geophysical Research-Atmospheres*, 114, 10.1029/2008jd010805, 2009.

Liu, Z., Wang, Y., Gu, D., Zhao, C., Huey, L. G., Stickel, R., Liao, J., Shao, M., Zhu, T., Zeng, L., Amoroso, A., Costabile, F., Chang, C. C., and Liu, S. C.: Summertime photochemistry during CAREBeijing-2007: RO_x budgets and O₃ formation, *Atmospheric Chemistry and Physics*, 12, 7737–7752, 10.5194/acp-12-7737-2012, 2012.

Lou, S. J., Liao, H., and Zhu, B.: Impacts of aerosols on surface-layer ozone concentrations in China through heterogeneous reactions and changes in photolysis rates, *Atmospheric Environment*, 85, 123–138, 10.1016/j.atmosenv.2013.12.004, 2014.

Lu, K., Fuchs, H., Hofzumahaus, A., Tan, Z., Wang, H., Zhang, L., Schmitt, S. H., Rohrer, F., Bohn, B., Broch, S., Dong, H., Gkatzelis, G. I., Hohaus, T., Holland, F., Li, X., Liu, Y., Liu, Y., Ma, X., Novelli, A., Schlag, P., Shao, M., Wu, Y., Wu, Z., Zeng, L., Hu, M., Kiendler-Scharr, A., Wahner, A., and Zhang, Y.: Fast Photochemistry in Wintertime Haze: Consequences for Pollution Mitigation Strategies, *Environmental Science & Technology*, 53, 10676–10684, 10.1021/acs.est.9b02422, 2019.

Lu, K. D., Hofzumahaus, A., Holland, F., Bohn, B., Brauers, T., Fuchs, H., Hu, M., Haseler, R., Kita, K., Kondo, Y., Li, X., Lou, S. R., Oebel, A., Shao, M., Zeng, L. M., Wahner, A., Zhu, T., Zhang, Y. H., and Rohrer, F.: Missing OH source in a suburban environment near Beijing: observed and modelled OH and HO₂ concentrations in summer 2006, *Atmospheric Chemistry and Physics*, 13, 1057–1080, 10.5194/acp-13-1057-2013, 2013.

Lu, X., Hong, J., Zhang, L., Cooper, O. R., Schultz, M. G., Xu, X., Wang, T., Gao, M., Zhao, Y., and Zhang, Y.: Severe Surface Ozone Pollution in China: A Global Perspective, *Environmental Science & Technology Letters*, 5, 487–494, 10.1021/acs.estlett.8b00366, 2018.

Ma, Z. Q., Xu, J., Quan, W. J., Zhang, Z. Y., Lin, W. L., and Xu, X. B.: Significant increase of surface ozone at a rural site, north of eastern China, *Atmospheric Chemistry and Physics*, 16, 3969–3977, 10.5194/acp-16-3969-2016, 2016a.

- Ma, Z. W., Hu, X. F., Sayer, A. M., Levy, R., Zhang, Q., Xue, Y. G., Tong, S. L., Bi, J., Huang, L., and Liu, Y.: Satellite-Based Spatiotemporal Trends in PM_{2.5} Concentrations: China, 2004-2013, *Environmental Health Perspectives*, 124, 184-192, 10.1289/ehp.1409481, 2016b.
- Madronich, S.: The atmosphere and UV-B radiation at ground level, in: *Environmental UV photobiology*, Springer, 1-39, 1993.
- Matthews, P. S. J., Baeza-Romero, M. T., Whalley, L. K., and Heard, D. E.: Uptake of HO₂ radicals onto Arizona test dust particles using an aerosol flow tube, *Atmos. Chem. Phys.*, 14, 7397-7408, 10.5194/acp-14-7397-2014, 2014.
- Mihelcic, D., Holland, F., Hofzumahaus, A., Hoppe, L., Konrad, S., Musgen, P., Patz, H. W., Schafer, H. J., Schmitz, T., Volz-Thomas, A., Bachmann, K., Schlomski, S., Platt, U., Geyer, A., Alicke, B., and Moortgat, G. K.: Peroxy radicals during BERLIOZ at Pabstthum: Measurements, radical budgets and ozone production, *Journal of Geophysical Research-Atmospheres*, 108, 17, 10.1029/2001jd001014, 2003.
- Moghani, M., Archer, C. L., and Mirzakhilili, A.: The importance of transport to ozone pollution in the US Mid-Atlantic, *Atmos. Environ.*, 191, 420-431, 2018.
- Monks, P. S., Archibald, A. T., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G. E., Stevenson, D. S., Tarasova, O., Thouret, V., von Schneidemesser, E., Sommariva, R., Wild, O., and Williams, M. L.: Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, *Atmospheric Chemistry and Physics*, 15, 8889-8973, 10.5194/acp-15-8889-2015, 2015.
- Ni, M. J., Huang, J. X., Lu, S. Y., Li, X. D., Yan, J. H., and Cen, K. F.: A review on black carbon emissions, worldwide and in China, *Chemosphere*, 107, 83-93, 10.1016/j.chemosphere.2014.02.052, 2014.
- Parrish, D. D., Xu, J., Croes, B., and Shao, M.: Air quality improvement in Los Angeles-perspectives for developing cities, *Frontiers of Environmental Science & Engineering*, 10, 10.1007/s11783-016-0859-5, 2016.
- Peeters, J., Nguyen, T. L., and Vereecken, L.: HO_x radical regeneration in the oxidation of isoprene, *Physical Chemistry Chemical Physics*, 11, 5935-5939, 10.1039/b908511d, 2009.
- Pollack, I. B., Ryerson, T. B., Trainer, M., Neuman, J. A., Roberts, J. M., and Parrish, D. D.: Trends in ozone, its precursors, and related secondary oxidation products in Los Angeles, California: A synthesis of measurements from 1960 to 2010, *Journal of Geophysical Research-Atmospheres*, 118, 5893-5911, 10.1002/jgrd.50472, 2013.
- Qu, W. J., Wang, J., Zhang, X. Y., Wang, D., and Sheng, L. F.: Influence of relative humidity on aerosol composition: Impacts on light extinction and visibility impairment at two sites in coastal area of China, *Atmospheric Research*, 153, 500-511, 10.1016/j.atmosres.2014.10.009, 2015.
- Raga, G. B., Castro, T., and Baumgardner, D.: The impact of megacity pollution on local climate and implications for the regional environment: Mexico City, *Atmospheric Environment*, 35, 1805-1811, 10.1016/s1352-2310(00)00275-2,

- 2001.
- Real, E., and Sartelet, K.: Modeling of photolysis rates over Europe: impact on chemical gaseous species and aerosols, *Atmos. Chem. Phys.*, 11, 1711-1727, 10.5194/acp-11-1711-2011, 2011.
- Shah, V., Jacob, D., Li, K., Silvern, R., Zhai, S., Liu, M., Lin, J., and Zhang, Q.: Effect of changing NO_x lifetime on the seasonality and long-term trends of satellite-observed tropospheric NO₂ columns over China, 2020.
- Streets, D. G., Fu, J. S., Jang, C. J., Hao, J. M., and Yu, C.: Air quality during the 2008 Beijing Olympic Games, *Atmospheric Environment* (1967), 41, 4800-4492, 2007.
- Taketani, F., Kanaya, Y., and Akimoto, H.: Kinetics of heterogeneous reactions of HO₂ radical at ambient concentration levels with (NH₄)₂SO₄ and NaCl aerosol particles, *The Journal of Physical Chemistry A*, 112, 2370-2377, 2008.
- Taketani, F., Kanaya, Y., Pochanart, P., Liu, Y., Li, J., Okuzawa, K., Kawamura, K., Wang, Z., and Akimoto, H.: Measurement of overall uptake coefficients for HO₂ radicals by aerosol particles sampled from ambient air at Mts. Tai and Mang (China), *Atmospheric Chemistry and Physics*, 12, 11907-11916, 10.5194/acp-12-11907-2012, 2012.
- Tan, Z., Fuchs, H., Lu, K., Hofzumahaus, A., Bohn, B., Broch, S., Dong, H., Gomm, S., Häsel, R., He, L., Holland, F., Li, X., Liu, Y., Lu, S., Rohrer, F., Shao, M., Wang, B., Wang, M., Wu, Y., Zeng, L., Zhang, Y., Wahner, A., and Zhang, Y.: Radical chemistry at a rural site (Wangdu) in the North China Plain: observation and model calculations of OH, HO₂ and RO₂ radicals, *Atmos. Chem. Phys.*, 17, 663-690, 10.5194/acp-17-663-2017, 2017.
- Tian, R., Ma, X., Jia, H., Yu, F., Sha, T., and Zan, Y.: Aerosol radiative effects on tropospheric photochemistry with GEOS-Chem simulations, *Atmospheric Environment*, 208, 82-94, 2019.
- Verstraeten, W. W., Neu, J. L., Williams, J. E., Bowman, K. W., Worden, J. R., and Boersma, K. F.: Rapid increases in tropospheric ozone production and export from China, *Nature geoscience*, 8, 690, 2015.
- Wang, B., Shao, M., Lu, S. H., Yuan, B., Zhao, Y., Wang, M., Zhang, S. Q., and Wu, D.: Variation of ambient non-methane hydrocarbons in Beijing city in summer 2008, *Atmospheric Chemistry and Physics*, 10, 5911-5923, 10.5194/acp-10-5911-2010, 2010.
- Wang, H., Lu, K., Tan, Z., Sun, K., Li, X., Hu, M., Shao, M., Zeng, L., Zhu, T., and Zhang, Y.: Model simulation of NO₃, N₂O₅ and ClNO₂ at a rural site in Beijing during CAREBeijing-2006, *Atmospheric Research*, 196, 97-107, 2017.
- Wang, J., Allen, D. J., Pickering, K. E., Li, Z., and He, H.: Impact of aerosol direct effect on East Asian air quality during the EAST-AIRE campaign, *Journal of Geophysical Research: Atmospheres*, 121, 6534-6554, 2016a.
- Wang, J. L., Din, G. Z., and Chan, C. C.: Validation of a laboratory-constructed automated gas chromatograph for the measurement of ozone precursors through comparison with a commercial analogy, *Journal of Chromatography A*, 1027, 11-18, 10.1016/j.chroma.2003.08.099, 2004.
- Wang, M., Zeng, L., Lu, S., Shao, M., Liu, X., Yu, X., Chen, W., Yuan, B., Zhang, Q.,

- Hu, M., and Zhang, Z.: Development and validation of a cryogen-free automatic gas chromatograph system (GC-MS/FID) for online measurements of volatile organic compounds, *Analytical Methods*, 6, 9424-9434, 10.1039/c4ay01855a, 2014.
- Wang, M., Shao, M., Chen, W., Lu, S., Liu, Y., Yuan, B., Zhang, Q., Zhang, Q., Chang, C. C., Wang, B., Zeng, L., Hu, M., Yang, Y., and Li, Y.: Trends of non-methane hydrocarbons (NMHC) emissions in Beijing during 2002-2013, *Atmospheric Chemistry and Physics*, 15, 1489-1502, 10.5194/acp-15-1489-2015, 2015a.
- Wang, P., Wang, T., and Ying, Q.: Regional source apportionment of summertime ozone and its precursors in the megacities of Beijing and Shanghai using a source-oriented chemical transport model, *Atmos. Environ.*, 224, 10, 10.1016/j.atmosenv.2020.117337, 2020.
- Wang, Q. Q., Sun, Y. L., Jiang, Q., Du, W., Sun, C. Z., Fu, P. Q., and Wang, Z. F.: Chemical composition of aerosol particles and light extinction apportionment before and during the heating season in Beijing, China, *Journal of Geophysical Research-Atmospheres*, 120, 12708-12722, 10.1002/2015jd023871, 2015b.
- Wang, W., Li, X., Shao, M., Hu, M., Zeng, L., Wu, Y., and Tan, T.: The impact of aerosols on photolysis frequencies and ozone production in Beijing during the 4-year period 2012–2015, *Atmos. Chem. Phys.*, 19, 9413-9429, 10.5194/acp-19-9413-2019, 2019.
- Wang, X. M., Chen, W. H., Chen, D. H., Wu, Z. Y., and Fan, Q.: Long-term trends of fine particulate matter and chemical composition in the Pearl River Delta Economic Zone (PRDEZ), China, *Front. Env. Sci. Eng.*, 10, 53-62, 10.1007/s11783-014-0728-z, 2016b.
- Wang, Y., Zhang, Y., Hao, J., and Luo, M.: Seasonal and spatial variability of surface ozone over China: contributions from background and domestic pollution, *Atmospheric Chemistry and Physics*, 11, 3511-3525, 10.5194/acp-11-3511-2011, 2011.
- Warneke, C., de Gouw, J. A., Holloway, J. S., Peischl, J., Ryerson, T. B., Atlas, E., Blake, D., Trainer, M., and Parrish, D. D.: Multiyear trends in volatile organic compounds in Los Angeles, California: Five decades of decreasing emissions, *Journal of Geophysical Research-Atmospheres*, 117, 10.1029/2012jd017899, 2012.
- Wehner, B., Birmili, W., Ditas, F., Wu, Z., Hu, M., Liu, X., Mao, J., Sugimoto, N., and Wiedensohler, A.: Relationships between submicrometer particulate air pollution and air mass history in Beijing, China, 2004-2006, *Atmospheric Chemistry and Physics*, 8, 6155-6168, 10.5194/acp-8-6155-2008, 2008.
- Wendisch, M., Mertes, S., Ruggaber, A., and Nakajima, T.: Vertical profiles of aerosol and radiation and the influence of a temperature inversion: Measurements and radiative transfer calculations, *J. Appl. Meteorol.*, 35, 1703-1715, 10.1175/1520-0450(1996)035<1703:vpoaar>2.0.co;2, 1996.
- Xie, X., Shao, M., Liu, Y., Lu, S., Chang, C.-C., and Chen, Z.-M.: Estimate of initial isoprene contribution to ozone formation potential in Beijing, China, *Atmospheric Environment*, 42, 6000-6010, 10.1016/j.atmosenv.2008.03.035, 2008.
- Xing, J., Wang, J., Mathur, R., Wang, S., Sarwar, G., Pleim, J., Hogrefe, C., Zhang, Y.,

- Jiang, J., and Wong, D. C.: Impacts of aerosol direct effects on tropospheric ozone through changes in atmospheric dynamics and photolysis rates, *Atmospheric chemistry and physics*, 17, 9869, 2017.
- Xu, J., Ma, J. Z., Zhang, X. L., Xu, X. B., Xu, X. F., Lin, W. L., Wang, Y., Meng, W., and Ma, Z. Q.: Measurements of ozone and its precursors in Beijing during summertime: impact of urban plumes on ozone pollution in downwind rural areas, *Atmospheric Chemistry and Physics*, 11, 12241-12252, 10.5194/acp-11-12241-2011, 2011.
- Xu, J., Zhang, Y. H., Zheng, S. Q., and He, Y. J.: Aerosol effects on ozone concentrations in Beijing: A model sensitivity study, *J. Environ. Sci.*, 24, 645-656, 10.1016/s1001-0742(11)60811-5, 2012.
- Yuan, B., Shao, M., de Gouw, J., Parrish, D. D., Lu, S., Wang, M., Zeng, L., Zhang, Q., Song, Y., Zhang, J., and Hu, M.: Volatile organic compounds (VOCs) in urban air: How chemistry affects the interpretation of positive matrix factorization (PMF) analysis, *Journal of Geophysical Research-Atmospheres*, 117, 10.1029/2012jd018236, 2012.
- Zhang, J. P., Zhu, T., Zhang, Q. H., Li, C. C., Shu, H. L., Ying, Y., Dai, Z. P., Wang, X., Liu, X. Y., Liang, A. M., Shen, H. X., and Yi, B. Q.: The impact of circulation patterns on regional transport pathways and air quality over Beijing and its surroundings, *Atmospheric Chemistry and Physics*, 12, 5031-5053, 10.5194/acp-12-5031-2012, 2012.
- Zhang, L., Shao, J. Y., Lu, X., Zhao, Y. H., Hu, Y. Y., Henze, D. K., Liao, H., Gong, S. L., and Zhang, Q.: Sources and Processes Affecting Fine Particulate Matter Pollution over North China: An Adjoint Analysis of the Beijing APEC Period, *Environ. Sci. Technol.*, 50, 8731-8740, 10.1021/acs.est.6b03010, 2016.
- Zhang, Q., Yuan, B., Shao, M., Wang, X., Lu, S., Lu, K., Wang, M., Chen, L., Chang, C. C., and Liu, S. C.: Variations of ground-level O₃ and its precursors in Beijing in summertime between 2005 and 2011, *Atmospheric Chemistry and Physics*, 14, 6089-6101, 10.5194/acp-14-6089-2014, 2014.
- Zhao, B., Wang, S. X., Liu, H., Xu, J. Y., Fu, K., Klimont, Z., Hao, J. M., He, K. B., Cofala, J., and Amann, M.: NO_x emissions in China: historical trends and future perspectives, *Atmospheric Chemistry and Physics*, 13, 9869-9897, 10.5194/acp-13-9869-2013, 2013.
- Zhao, P. S., Zhang, X. L., and Xu, X. F.: Long-term visibility trends and characteristics in the region of Beijing, Tianjin, and Hebei, China, *Abstr. Pap. Am. Chem. Soc.*, 242, 1, 2011.
- Zheng, C., Zhao, C., Zhu, Y., Wang, Y., Shi, X., Wu, X., Chen, T., Wu, F., and Qiu, Y.: Analysis of influential factors for the relationship between PM_{2.5} and AOD in Beijing, *Atmospheric Chemistry and Physics*, 17, 13473-13489, 2017.
- Zou, Q., Song, H., Tang, M., and Lu, K.: Measurements of HO₂ uptake coefficient on aqueous (NH₄)₂SO₄ aerosol using aerosol flow tube with LIF system, *Chin. Chem. Lett.*, 30, 2236-2240, <https://doi.org/10.1016/j.cclet.2019.07.041>, 2019.

Supplementary Information




Figure S1. The trend of average MDA8 ozone in Changdao during 2013-2019. These data are acquired from “Blue book on prevention and control of atmospheric ozone pollution in China (in Chinese)” reported by Chinese Society of Environmental Sciences in 2020 (http://www.ep-serve.com/forepart/zxnr_index.do?oid=51478637&tid=26378242).





Figure S2. Trend of monthly afternoon (12:00-15:00) mean $P(O_3)$ simulated by the chemical box model. Red dots: Only the gas-phase reactions are considered in the box model constrained by observed photolysis frequencies from 2006 for all eleven years. Blue dots: the box model as above, but constrained by the photolysis frequencies derived for each year without the changing aerosol uptake of HO_2 considered.

1075
1076
1077



1078
1079
1080
1081
1082

Figure S3. Diurnal variation of simulated $P(O_3)$ in Beijing in August during 2006- 2016. $P(O_3)_{j_obs}$ represents ozone production rate under observed photolysis frequencies; $P(O_3)_{j_AOD=0}$ represents ozone production rate under calculated photolysis frequencies when AOD is equal to 0.