# Supplement of Mixing states of Amazon-basin aerosol particles transported over long distances using transmission electron microscopy

Kouji Adachi<sup>1</sup>, Naga Oshima<sup>1</sup>, Zhaoheng Gong<sup>2,a</sup>, Suzane de Sá<sup>2</sup>, Adam P. Bateman<sup>2</sup>, Scot T. Martin<sup>2</sup>, Joel F. de Brito<sup>3,b</sup>, Paulo Artaxo<sup>3</sup>, Glauber G. Cirino<sup>4,c</sup>, Arthur J. Sedlacek III<sup>5</sup>, Peter R. Buseck<sup>6</sup>

 <sup>1</sup> Department of Atmosphere, Ocean and Earth System Modelling Research, Meteorological Research Institute, Tsukuba, Japan
 <sup>2</sup> School of Engineering and Applied Sciences & Department of Earth and Planetary Sciences, Harvard University, Cambridge, Massachusetts, USA
 <sup>3</sup> Institute of Physics, University of São Paulo, São Paulo, Brazil

- <sup>4</sup> National Institute for Amazonian Research, Manaus, Amazonas, Brazil
   <sup>5</sup> Brookhaven National Laboratory, Upton, New York, USA
   <sup>6</sup> School of Earth and Space Exploration & School of Molecular Sciences, Arizona State University, Tempe, Arizona, USA
   a: Now at: Harvard Business School, Boston, Massachusetts, USA, b: now at: IMT Lille Douai,
- 15 University of Lille, SAGE, Lille 59000, France, c: now at: Geosciences Institute, Federal University of Para, Belem, Brazil.

Correspondence to: Kouji Adachi (adachik@mri-jma.go.jp)

#### Supporting figures 1 to 10.



Supporting Figure 1. Flow chart for the particle classification. All analysed particles were classified based on their compositions obtained using STEM-EDS. For Al, Fe, and P, we chose the threshold values
that are just above their detection limits because they are unique but minor components within particles. For K and S, we used larger threshold values (2 weight %) than that of Na (1 weight %), as they have heavier atomic masses than Na.







(c) Element mapping images



30 Supporting Figure 2. Shapes and compositions of mineral particles. (a) TEM and (b) STEM and (c) element mapping images of mineral particles. These mineral particles contain Al-Si-Mg-O-, Al-Si-Na-O-, or Al-Si-K-O-rich grains. There are Ca- or Fe-rich grains within the mineral particles. Ns-soot and sulfate particles also occur. The sample was collected from 12:00-12:30, 29 March, 2014.



Supporting Figure 3. Shape and composition of PBA particle with K-rich grains. (a) TEM and (b)
STEM and (c) element mapping images of the PBA particle. A mineral particle attaches to the PBA particle. The sample was collected from 3:00-3:30, 15 March, 2014.



#### (b) STEM image



(c) Element mapping images



**Supporting Figure 4.** Shape and composition of PBA particle with P-rich grains. (a) TEM and (b) STEM and (c) element mapping images of the PBA particle. The sample was collected from 0:00-0:30, 27 February, 2014.







## (c) Element mapping images



**Supporting Figure 5.** Shapes and compositions of PBA, mineral, and KCl particles. (a) TEM and (b) STEM and (c) element mapping images of these particles. The sample was collected from 0:00-0:30, 27 February, 2014.



**Supporting Figure 6.** Shapes and compositions of mineral particles that attach sea-salt (NaCl) particles. (a) TEM and (b) STEM images and (c) element mapping images of these particles. The sample was collected from 18:00-18:30, 2 February, 2014.



(c) Element mapping images



Supporting Figure 7. Shapes and compositions of K-bearing particles that attach ns-soot particles. (a) 55 TEM and (b) STEM and (c) element mapping images of the particles. These particles consist of K and S, suggesting that they are potassium sulfate. These K-bearing particles occur either with or without ns-soot particles. The sample was collected from 0:00-0:30, 25 February, 2014.





**Supporting Figure 8.** Shapes and compositions of sulfate particles. (a) TEM and (b) STEM and (c) element mapping images of these particles. These sulfate particles include N, O, S, and K, implying that they are ammonium or potassium sulfate. Sulfate particles deform over the substrates and decompose after intense electron beam irradiation. The sample was collected during 12:55-13:25, 26 March, 2014.

(a) TEM image



(b) STEM image

(c) Element mapping images

70



**Supporting Figure 9.** Shapes and compositions of mineral and carbonaceous particles, including tarball, ns-soot, and organic matter. (a) TEM and (b) STEM and (c) element mapping images of these particles. The mineral particles include those with Al- and Si-rich particles. The sample was collected from 12:00-12:30, 14 February, 2014.



## (b) STEM image







**Supporting Figure 10.** Shapes and compositions of PBA particles coated with sea salt. (a) TEM and (b) STEM and (c) element mapping images of these particles. The sea-salt coating mainly consists of NaCl with a thin sulfur layer. PBOA is characterized by C and P. The sample was collected from 12:00-12:30,

2 February, 2014.