

Responses to Referee 3:

The authors thank the reviewer for her/his excellent comments.

In this study, Hao et al. studied the photochemical reaction of α -pinene in a smog chamber by using HR-ToF-PTRMS and HR-ToF-AMS. They mainly focused on the formation of ammonium. It was found the concentration of ammonium was in good correlation with gas phase organic acids. Thus, they pointed the important role of organic acids in the formation of ammonium. Considering the ubiquitous presence and its alkalinity in the atmosphere, this work provided useful information to realize the environmental impacts of NH_3 and has important enlightenment for the future research on SOA formation. I think it is suitable for publication on ACP. However, there are still several concerns needed to be addressed before acceptance.

We thank the referee for the positive comment.

Major concerns.

1. The title. The “CCN-size” in the title is not necessary. The title of a paper should contain informative key words to represent the main content. However, no measurement about the CCN activity of particles was conducted. In fact, the size of particles reported in this study were based on the measurement limit of size-range of AMS, which has no direct link with CCN-size.

Reply:

We agree with the reviewer that the “CCN-size” in the title is not necessary. We have removed it and accordingly we modified the manuscript.

2. Since the work focused on the conversion of gas phase NH_3 to particulate NH_4^+ , the concentration of NH_3 in the reaction system should be measured. In all experiments, ammonia was introduced to the chamber as an impurity. This condition makes the research loose and unscientific. The source of NH_3 may include the evaporation from the wall of reactors due the previous deposition of ammonium. However, its sources may be not stable and fluctuate the concentration of NH_3 . Then there are errors in the analysis of ammonium as a function of time due to the uncertainty of NH_3 . At least the author needs to prove that the concentration of ammonia is constant or varies regularly during the experiment.

Reply:

The NH_3 was introduced to the chamber as an impurity. We lacked its measurement device in our lab and its concentration was estimated from AMS measurement. We agree with the referee that the missed NH_3 measurement could lead to some uncertainties of our interpretation to the NH_4^+ results. We need to point out that prior to each experiment, the chamber was continuously flushed overnight with laboratory clean air produced by a zero air generator to minimize any possible impurities from chamber air and wall. There is no physical reason for the fluctuation of the source of impurity: the temperature and other relevant conditions in the chamber were stable over the experiment. It is possible thought that the NH_3 concentration is gradually changing over the experiment, but this doesn't affect our conclusions. Fig C1A shows the time series of total organic acids measured by PTRMS (the sum of C1-C5 organic acids after excluding the background), and also the estimated mean NH_3 concentration of 19 ppb based on a statistics study on the ammonia in 14 Finnish office buildings (Salonen *et al.*, 2009). After 50 mins, the concentration of organic acids is higher than that of NH_3 , and NH_3 is a limiting factor for the base-acid neutralization. Meanwhile, we still observe the formation of NH_4^+ in a continuous and smooth way in three experiments (Fig. C1B), suggesting a constant and regular source of NH_3 .




Figure C1 (A) The time series of total C1-C5 organic acids after excluding the background measured by PTRMS in the nucleated SOA experiments, and the estimated mean concentration of NH₃ (in cyan) and the highest concentration (in orange), based on a statistics study on the ammonia in 14 Finnish office buildings in a national scale. (B) The measured ammonium concentration by AMS.

3. The interpretation about the difference between experimental results with and without seeds is lack. The authors designed two sets of experiments and found distinctly different results. However, the reasons were not provided. Why the VOC concentrations are so different between these two sets of experiments? What's the role and effects of seeds on the formation of ammonium? Moreover, it is difficult to understand the difference in the consumption of NO_x ($\Delta\text{NO}_x\text{-E314} > \Delta\text{NO}_x\text{-E327}$ in Fig. S2) and the corresponding formation of nitrate ($\Delta\text{NO}_3\text{-E314} < \Delta\text{NO}_3\text{-E327}$ in Fig. 1).

Reply:

We thank the reviewer for the nice comments. In the seeded experiments alpha-pinene concentration was about 20 times lower than in the nucleation experiment. The relatively low alpha-pinene concentration was used to mimic an atmospherically relevant monoterpene mixing ratios, for example, in Hyytiälä forest area in Finland (e.g. Kourtchev *et al.*, 2006). We aimed to study if the ammonia plays a similar role as in the high alpha-pinene concentration case. Also, the nucleation experiments can't be conducted with such low concentrations because the nucleation experiments requires higher concentration due to the Kelvin effect to produce detectable amount of SOA mass in the size range of aerosol mass spectrometer (AMS). Both sets of experiments have consistently shown that organic acids were required to neutralize the observed ammonium salt and gas-phase organic acids contribute to the SOA formation.







Fig. C2. Time series of NO_x and formed O₃ in E0314 and E0327.

Fig. C2 shows the added NO_x and formed O₃ in the chamber in E0314 and E0327. The added NO was rapidly converted to NO₂ by RO₂ radicals derived from photooxidation of alpha-pinene. The fate of NO₂ is a competition among the reactions (1)-(3):



In Fig. C2A, we have observed a large amount of O₃ formation because of reaction (1). Meanwhile, we also observed an increase in N:C ratios in both E0314 and E0327 either because of reaction (2) or NH₃ reaction as we have discussed in the manuscript. The results show the participation of NO_x in different reactions, besides producing ammonium nitrate, which qualitatively explains the different amount of nitrate observed in two experiments (table C1).




Fig. C3. N:C ratios measured by AMS in E0314 and E0327.

Table C1. Changes of NO_x , and formed O_3 and nitrated aerosol in E0314 and E0327.

Exp.	ΔNO_x , ppb	ΔO_3 , ppb	NO_3 , $\mu\text{g m}^{-3}$
E0314	16	316	0.12
E0327	7.3	52.3	2.5

Accordingly, we added in line89 in the manuscript that “The α -pinene concentration in the seeded experiments was about 20 times lower than in the nucleation experiment. The relatively low α -pinene concentration was used to mimic an atmospherically relevant monoterpene mixing ratios, for example, in Hyytiälä forest area in Finland (e.g. Kourtchev *et al.*, 2006). We aimed to study if the ammonia plays a similar role as in the high α -pinene concentration case.”

4. removal Fig. 3 to SI. The data in fig. 3 were directly derived from the subtraction of data in Fig. 2, then no new meaning was provided in Fig. 3 and it could be removed to SI.

Reply:

The data in Fig. 3 was estimated from the conversion of the difference between predicted and measured ammonium in Fig. 2. The graph shows directly the amount of organic acids required to neutralize ammonium and the time at which the organic acids started to play a role in ammonium formation. Hence, we believe that the figure is informative and would like to keep it in the main text.

5. line 200: the statement “The delay is caused by the effect of nitric acid arising from the background NO_x photooxidation.” is not correct. According to Fig. 1-B2, it seems the main reason may be the formation of sulfate. However, no obvious consumption in SO_2 in Fig. S2 make the yield of sulfate hard to understand. Why the formation of ammonium in the seed experiments is earlier than sulfate, nitrate, and light on? Again, the effect of seeds needed to be explained.

Reply:

As the referee pointed out that no obvious consumption in SO_2 in E0314-E0316, the formation of sulfate was minor in the experiments. The increase in ammonium and sulphate concentrations before UV were switched on in Fig 1-B2&A2 is because we added ammonium sulfate seeds to the chamber before the lights were turned on. We would like to note that the used ammonium sulphate seed and low organic concentrations makes the ammonium formation more difficult to distinguish from the data in the experiments E0314-E0316. Overall, the seed dominated the NH_4^+ signal in AMS. But e.g. in the case of E0316 we see the increase in NH_4^+ (after UV lights were turned on) and simultaneously a decrease in SO_4^{2-} (SO_4^{2-} signal is originated from the neutral ammonium sulphate seed) meaning that there is a NH_4^+ formation taking place in particle phase. Also at the same time the NO_3^- and Org signals are increased suggesting that the increase in NH_4^+ is related to nitric acids and organic acids. The ammonium sulfate was added as seeds for condensational growth of SOA. Since the seeds were introduced to the chamber in a neutral state, we didn’t observe other effects of seed aerosol, e.g., acid-catalysis accretion, on SOA formation.




Fig. C4. The time series of org, nitrate, sulfate and nitrate measured by AMS in the seeded SOA experiments.

After checking the authors' response to the former version, I find some critical concerns still present. A more convincing reply is needed to address these concerns.

Minor concerns:

1. line 9: ubiquitously

Reply: We have fixed it.

2. The Sequence of Fig. 3 appears in the main text is earlier than Fig. 4

Reply: We have fixed it.

3. line 168: Fig. 2?

Reply: We have fixed it.

4. line 247: AMSSurprisingly?

Reply: We have fixed it.

References:

- Kourtchev, I., Giorio, C., Manninen, A., Wilson, E., Mahon, B., Aalto, J., Kajos, M., Venables, D., Ruuskanen, T., Levula, J., Lopenon, M., Connors, S., Harris, N., Zhao, D., Kiendler-Scharr, A., Mentel, T., Rudich, Y., Hullquist, M., Doussin J., Maenhaut, W., Back, J., Petaja, T., Wenger, J., Kulmala, M. and Kalberer, M., Enhanced volatile organic compounds emissions and organic aerosol mass increase the oligomer content of atmospheric aerosols, *Sci. Reports*, 6, 35038, 2016.
- Salonen, H., Pasanen, A., Lappalainen, S.K., Riuttala, H.M., Tuomi, T.M., Pasanen, P.O., Back, B.C. and Reijula, K.E., Airborne concentrations of volatile organic compounds, formaldehyde and ammonia in Finnish office buildings with suspected indoor air problems, *Journal of Occupational and Environmental Hygiene*, 6, 200-209, 2009.

Direct contribution of ammonia to ~~CCN-size~~ α -pinene secondary organic aerosol formation

Liqing Hao¹, Eetu Kari^{1,a}, Ari Leskinen^{1,2}, Douglas R. Worsnop^{1,3}, Annele Virtanen¹

¹Department of Applied Physics, University of Eastern Finland, Kuopio, Finland

²Finnish Meteorological Institute, Kuopio, Finland

³Aerodyne Research Inc., Billerica, MA 08121-3976, USA

^anow at: Neste Oyj, Porvoo, Finland

Correspondence to: Liqing Hao (hao.liqing@uef.fi) and Annele Virtanen (annele.virtanen@uef.fi)

Abstract. Ammonia (NH₃), a gaseous compound ubiquitously present in the atmosphere, is involved in the formation of secondary organic aerosol (SOA), but the exact mechanism is still not well known. This study presents the results of SOA experiments from the photooxidation of α -pinene in the presence of NH₃ in the reaction chamber. SOA was formed in nucleation experiment and in seeded experiment with ammonium sulfate particles as seeds. The chemical composition and time-series of compounds in the gas- and particle- phase were characterized by an on-line high-resolution time-of-flight proton transfer reaction mass spectrometer (HR-ToF-PTRMS) and a high-resolution time-of-flight aerosol mass spectrometer (HR-ToF-AMS), respectively. Our results show that ~~for the aerosol particles in cloud condensation nuclei (CCN) size,~~ the mass concentration of ammonium (NH₄⁺) was still rising even after the mass concentration of organic component started to decrease due to aerosol wall deposition and evaporation, implying the continuous new formation of particle phase ammonium in the process. Stoichiometric neutralization analysis of aerosol indicates that organic acids have a central role in the formation of particle phase ammonium. Our measurements show a good correlation between the gas phase organic mono- and di-carboxylic acids formed in the photooxidation of α -pinene and the ammonium in the particle phase, thus highlighting the contribution of gas-phase organic acids to the ammonium formation ~~in the CCN-size SOA particles~~. The work shows that the gas-phase organic acids contribute to the SOA formation by forming organic ammonium salts through acid-base reaction. The changes in aerosol mass, particle size and chemical composition resulting from the NH₃-SOA interaction can potentially alter the aerosol direct and indirect forcing and therefore alter its impact on climate change.

1 Introduction

The largest uncertainty in forward projection of global warming is related to our limited knowledge of ~~negative~~ solar radiative forcing associated with aerosols (IPCC, 2013). Formation of secondary organic aerosols (SOA) is one of the main processes that affects the composition and properties of atmospheric aerosols. Formation of SOA occurs through two distinct mechanisms: by increasing the mass of the existing aerosol and through the formation of new particles. Estimate on the SOA formation shows its significance as a source of atmospheric organic aerosol: about 60% of the organic aerosol mass is SOA on the global scale and regionally even more (Hallquist *et al.*, 2009; Jimenez *et al.*, 2009; Kanakidou *et al.*, 2005). Hence, SOA plays an important role in the direct scattering of solar radiation, cloud formation and precipitation, and visibility reduction, and may also have a direct impact on human health.

Ammonia (NH₃) is ubiquitously present in the atmosphere as a dominant volatile base. The majority of its sources is accounted by emissions from agriculture, livestock, soil and traffic (Huang *et al.*, 2012; Grönroos *et al.*, 2009; Battye *et al.*, 2003). NH₃ governs the neutralization of atmospheric aerosol by reacting with the inorganic acids such as sulfuric acid and nitric acid, leading to transformation of a substantial amount of ammonium sulfate (and

derivatives) and ammonium nitrate (Seinfeld and Pandis, 2016). These inorganic salts play a vital role in contributing to the fine particle matter (PM_{2.5}) and altering the chemical and physical properties of aerosol particles in the atmosphere. The mechanism between NH₃ and the inorganic acids leading to the secondary inorganic aerosol has been well recognized.

A number of studies have shown that NH₃ is one of the key species for the new particle formation through ternary and binary nucleation with water and sulfuric acid (e.g. Lehtipalo et al., 2018; Jokinen et al., 2018; Bianchi et al., 2016; Kirkby et al., 2011; Kurten et al., 2007; Kulmala et al., 2000). The nucleated particles are a significant source of atmospheric SOA particles and subsequent growth to a larger size (>50nm) allows them to serve as cloud condensation nuclei (CCN). However, the role of NH₃ for the large size (e.g. CCN-size) SOA particles are still rarely studied. A study, conducted more than a decade ago, demonstrated that NH₃ increased the number and volume concentrations of CCN-sized SOA particles from α -pinene-ozone system by 15% and 8%, respectively (Na et al., 2007). Similar results have also been observed in the photooxidation and ozonolysis of α -pinene SOA experiments that SOA mass yield increased by 13% as a response to NH₃ addition (Babar et al., 2017). Besides, the addition of NH₃ could also promote the SOA formation from photooxidation of vehicle exhaust (Chen et al., 2019; Liu et al., 2015), anthropogenic VOCs (Wang et al., 2018; Huang et al., 2018) and acrolein (Li et al., 2019). The promotion mechanism of NH₃ to SOA formation can be both through base-acid reaction (Schlag et al., 2017; Babar et al., 2017; Rubach et al., 2013; Mensah et al., 2012; Na et al., 2007) and by the NH₃ uptake to the carbonyl group (Zhu et al., 2018; Liu et al., 2015). As a consequence, the changes in particle size and chemical composition could alter the CCN ability and hygroscopicity of SOA particles (Dinar et al., 2008). Moreover, the reaction of NH₃ with SOA decreases the volatility of SOA particle (Paciga et al., 2014), and also results in production of light-absorbing brown carbon compounds that modify the optical properties of the aerosols (Huang et al., 2018; Updyke et al., 2012; Bones et al., 2010). Additionally, the uptake of NH₃ by SOA can deplete ambient NH₃ concentrations, causing indirect reductions in the amount of inorganic ammonium salts in particulate matter (Horne et al., 2018). Therefore, the interaction of NH₃ and SOA could alter both direct and indirect aerosol radiative forcing and potentially alter its impact on climate change.

This work presents the results of SOA formation from photooxidation of α -pinene in the presence of NH₃ in the nucleation and seeded experiments. The chemical composition of gas-phase and particle-phase compounds were characterized with a high-resolution time-of-flight proton transfer reaction mass spectrometer (HR-ToF-PTRMS) and a high-resolution time-of-flight aerosol mass spectrometer (HR-ToF-AMS), respectively. Our experiments show the formation of ammonium salt (NH₄⁺) in the CCN-size SOA particles, and our gas phase measurements indicate organic acids are responsible for their formation. The results will have potential applications in studying SOA formation mechanism and the impact of SOA on climate forcing.

2 Experimental

2.1 Reaction chamber experiments

SOA experiments were conducted in the ILMARI environmental chamber infrastructure in University of Eastern Finland. The experimental system and experimental procedure have been described in details in Leskinen et al. (2015) and Kari et al. (2019a), respectively. The chamber consists of a 29 m³ TeflonTM FEP film bag. Ultraviolet (UV) lights with a spectrum centered at 340 nm around the chamber enable photochemical reactions. Two sets of experiments were designed (Table 1): (1) SOA nucleation experiments in the absence of seed aerosols; (2) SOA

formation experiments in the presence of ammonium sulfate seeds. Prior to each experiment, the chamber was continuously flushed overnight with laboratory clean air produced by a zero air generator (Model 737-15, Aadco Instruments Inc., USA) and conditioned in a humidifier (Model FC125-240-5MP-02, Perma Pure LLC., USA), aiming at 50 % relative humidity at the typical temperature of 20 °C during the course of each experiment. Independent on the type of experiments (nucleation and seeded), the reactants were added into the chamber in the sequence described below. The (NH₄)₂SO₄ (ammonium sulfate, 99 %, Sigma Aldrich) seed was injected to the chamber first. The seed was generated using an atomizer (Topas ATM226, Germany). In two experiments, nitrogen oxide (NO) was added to chamber, and ozone (O₃) was introduced into the chamber to convert NO to nitrogen dioxide (NO₂) to reach the atmospherically relevant NO₂-to-NO ratio of ~ 3 (Kari et al., 2019a). After O₃ and NO_x were fed, 3 µl 9-fold butanol (butanol-d9, 98 %, Sigma Aldrich) was injected to the chamber, from whose consumption the hydroxyl radical (OH) exposure was estimated in each experiment (Kari et al., 2019a). Next, 1 µl and 18 µl of α-pinene (≥ 99%, Sigma Aldrich) were added into the chamber for the seeded and nucleation SOA experiments, respectively, corresponding to concentrations of ~5 ppb and ~100 ppb of α-pinene in the chamber. The α-pinene concentration in the seeded experiments was about 20 times lower than in the nucleation experiment. The relatively low α-pinene concentration was used to mimic an atmospherically relevant monoterpene mixing ratios, for example, in Hyytiälä forest area in Finland (e.g. Kourtshev et al., 2006). We aimed to study if the ammonia plays a similar role as in the high α-pinene concentration case. Last, 5 ml of hydrogen peroxide (H₂O₂, 30 wt. % in H₂O, Sigma Aldrich) was conducted into the chamber by mixing it with purified air flowing at 10 lpm (liter per minute), as a precursor for OH radicals to be generated under UV radiation. The H₂O₂ concentration was roughly 30 ppm estimated from the amount of its injection and the volume of chamber. In two sets of experiments, ammonia was introduced to the chamber as an impurity. The likely sources of ammonia were the pressurized air, possible leakage of the lab air and water used for humidification. After all the compounds were introduced into the chamber, the chamber was closed, and the compounds were allowed to stabilize for 15 min. Then the UV lights were switched on to initiate photochemistry.

2.2 Analytical methods and instrumentation

The size-resolved chemical composition and mass concentration of aerosol particles were measured directly with an on-line high-resolution time-of-flight aerosol mass spectrometer (HR-ToF-AMS, abbreviated as AMS) (DeCarlo et al., 2006). A detailed description of AMS operational procedure is provided in previous publications (Canagaratna et al., 2007; Jayne et al., 2000). In brief, AMS was operated in V - mode in EI - mode. Calibration of ionization efficiency (IE) followed the standard protocol using dried and size-selected ammonium nitrate particles. The data were analyzed using standard AMS data analysis toolkits (Squirrel V1.62D and Pika V1.22D) in Igor Pro Software (version 6.37, WaveMetrics Inc.). For determining mass concentrations, the default relative ionization efficiency (RIE) values were 1.4, 1.1 and 1.2, for organics, nitrate and sulfate, respectively. The RIE for ammonium was 2.95, as determined in the IE calibration. After a comparison to the volume concentration derived from a scanning mobility particle sizer (SMPS TSI 3081 DMA + 3775 CPC) measurement (Fig. S1), a collection factor of 100 % was applied to determine the aerosol mass concentration in the reported results in this work. The positive matrix factorization (PMF) analysis was performed on the high-resolution mass spectra by using the PMF Evaluation Tool V2.08 (Paatero and Tapper, 1994; Ulbrich et al., 2009). The standard error matrices were processed following the principles of applying minimum error estimate, downweighting weak variables, removing bad variables, and downweighting m/z 44 related fragments

(Ulbrich et al., 2009). The PMF was evaluated with 1 to 6 factor. Rotation (Fpeak) varied from -1 to 1 at a step of 0.1.

The concentrations of α -pinene and oxidized organic products were quantified by an on-line high-resolution time-of-flight proton transfer reaction mass spectrometer (HR-ToF-PTRMS, abbreviated as PTRMS) (Ionicon Analytik). A detailed description of the instrument and operation procedure has been provided in Jordan et al. (2009) and Kari et al. (2019a; 2019b). In brief, the PTRMS instrumental setting were the same as in Kari et al. (2019a). During measurement, the mass calibration was conducted using protonated water isotope signals at m/z 21 and internal instrumental signals of diiodobenzene and its fragment ions (protonated integer m/z 331). The instrumental transmission calibration was conducted using a standard gas mixture containing 8 different VOCs (protonated integers ranging from m/z 79 to m/z 181). The concentration of VOCs were determined according to the principles by Hansel et al. (1995). The selected ions that we are interested in this study are consistent with α -pinene (m/z 136+1), formic acid (m/z 46+1), acetic acid (m/z 60+1), propionic acid (m/z 74+1), pinonic acid (m/z 184+1), butyric acid (m/z 88+1), pentanoic acid (m/z 102+1), malonic acid (m/z 104+1) and succinic acid (m/z 118+1).

For supporting information we measured the particle concentration and size distribution in a diameter range of 7-800 nm with an SMPS (TSI 3081 DMA + 3775 CPC) and the concentrations of NO, nitrogen oxides (NO_x , Fig. S2), ozone (O_3), and sulfur dioxide (SO_2 , Fig. S2), as well as the relative humidity and temperature inside the chamber. In this study, we lacked the measurement of NH_3 concentration in the chamber but estimated it from our AMS measurement results, by assuming that the particulate ammonium salt (NH_4^+) was converted from the gas-phase NH_3 (Fig. 1). The maximum NH_4^+ concentration was in the range of $1.17\sim 1.51 \mu\text{g m}^{-3}$, which corresponds to a minimum NH_3 concentration level of $1.6 \sim 2.1 \text{ ppbV}$ in our chamber. The NH_3 mixing ratio over continental range is typically between 0.1 and 10 ppb (Seinfeld and Pandis, 2016). A statistics study on the ammonia concentration in 14 Finnish office buildings shows a range of $1\text{-}49 \mu\text{g m}^{-3}$ and a geometric mean concentration of $14 \mu\text{g m}^{-3}$ (corresponds 19 ppbV) (Salonen et al., 2009). Hence, our method should provide a lowest boundary of ammonia mixing ratio in the chamber.

3 Results and discussion

3.1 Time series of aerosol species

After the UV lamps were switched on, the photooxidation reaction produced oxidized gas-phase compounds and SOA particles in both nucleation and seeded experiments. The time series of mass concentrations of formed SOA (in green) and ammonium (in orange) along with sulfate and nitrate components measured by AMS are presented in Fig. 1. In the nucleation experiments (the left panels in Fig. 1), we observed rapid increase in the SOA mass concentrations after the photooxidation reaction had started. After reaching the maximum concentration of $390 - 476 \mu\text{g m}^{-3}$, the SOA concentration declined by $27.9 \pm 9.2 \%$ at the end of experiment because of particle deposition on the chamber wall and/or aerosol evaporation. The O:C ratio (oxygen to carbon ratio) of the SOA particles were slightly increased from the initial 0.39 ± 0.015 to the final 0.44 ± 0.01 because of aerosol aging. In a distinct contrast, the mass concentrations of ammonium component were still rising at the stage of decreasing SOA masses. Together taking into account the fact that aerosol wall deposition loss was present resulting in decreasing organic mass (and decreasing sulphate mass in the seeded experiments), our results suggest new production of ammonium salts. The newly formed ammonium can be partly attributed to the co-generated nitrate and sulfate as the photooxidized products of NO and SO_2 in the chamber. However, the amount of the two inorganic species can't fully interpret the ammonium and we will elaborate this in more detail in Sec. 3.2. Similar phenomena were also observed in the seeded experiments (Fig.

1 and Fig. 2). Note that as the measured mass is centered on the larger sizes of AMS measurement range (35-1000nm, Jayne et al., 2000; Zhang et al., 2004), we can state that the measurements represents CCN relevant particle sizes (in general it is often assumed that atmospheric aerosol particles larger than 100 nm are able to active to cloud droplets). Additionally, the organics, nitrate and ammonium aerosols showed similar mass-based size distributions from our AMS measurement (refer to Sec. 3.2.1), and thus internally-mixed aerosol can be assumed in our study, indicating that the aerosol components were lost in a similar rate to the chamber wall. Therefore, no loss correction to the aerosol wall disposition was conducted in this work.

3.2 Participation of ammonia in SOA formation

3.2.1 Reaction of ammonia with organic acids

To investigate the monotonically increasing profile of the ammonium salt in our chamber experiment, we studied the stoichiometric neutralization of formed SOA particles. The approach proposed by Zhang et al (2007) was adopted, in which the ammonium mass concentrations measured ($\text{NH}_4^+_{\text{mea}}$) in the particles were compared to the stoichiometric ammonium concentrations required ($\text{NH}_4^+_{\text{pre}}$) to fully neutralize the measured concentrations of SO_4^{2-} , NO_3^- and Cl^- :

$$\text{NH}_4^+_{\text{pre}} = 18 \times (2 \times \text{SO}_4^{2-}/96 + \text{NO}_3^-/62 + \text{Cl}^-/35.5) \quad (1)$$

where NH_4^+ , SO_4^{2-} , NO_3^- and Cl^- represent the mass concentrations (in $\mu\text{g m}^{-3}$) of the species and the denominators correspond to their molecular weights. The factor 18 is the molecular weight of NH_4^+ .

A comparison between the predicted and measured ammonium masses is displayed in Fig. 2 and S3. In both sets of experiments, the measured ammonium mass concentration was systematically greater than the predicted value. The trend doesn't show a dependence on the presence of NO_x in the chamber. On average, $\text{NH}_4^+_{\text{mea}}$ is $400 \pm 156\%$ and $21 \pm 11\%$ greater than $\text{NH}_4^+_{\text{pre}}$ at the end of nucleation and seeded experiments, respectively. The large discrepancy between the measured and predicted ammonium concentrations suggested that the current amount of sulfate, nitrate and chloride is insufficient to neutralize the ammonium formed in the particle phase, which indicates that organic component must have played a role in this process. Considering the nature of NH_3 as a base compound, the candidate species of organic compounds are attributed to organic acids.

Since a vast variety of molecular compositions of organic acids may be present in the photooxidation products of α -pinene, it isn't possible to define the amount of individual organic acid required to neutralize the NH_3 . Therefore, we use the CO_2^+ ion measured by AMS to represent carboxylic functional group of organic acids. The CO_2^+ is not only considered as a reliable marker of oxygenated organic aerosol (e.g. Zhang et al., 2005), but is also tightly associated with the formation of organic mono- and di-acids shown in laboratory and field measurements (Yatavelli et al., 2015; Takegawa et al., 2007; Alfarra et al., 2004).

Taking into account the contribution of organic acids to ammonium salt, we reformulate Eq. (1) to:

$$\text{NH}_4^+_{\text{pre_CO}_2} = 18 \times (2 \times \text{SO}_4^{2-}/96 + \text{NO}_3^-/62 + \text{Cl}^-/35.5 + \text{CO}_2^+_{\text{NH}_4}/44) \quad (2)$$

where $\text{NH}_4^+_{\text{pre_CO}_2}$ is the predicted ammonium mass concentration after taking into account contribution of organic acids. $\text{CO}_2^+_{\text{NH}_4}$ is the mass concentration of carboxylic function group ($-\text{CO}_2$) representing organic acids which were required to neutralize the ammonium. The denominator 44 is the molecular weight of carboxylic functional group.

To interpret $\text{NH}_4^+_{\text{mea}}$, $\text{NH}_4^+_{\text{pre_CO}_2}$ should equal to $\text{NH}_4^+_{\text{mea}}$:

$$\text{NH}_4^+_{\text{pre_CO}_2} = \text{NH}_4^+_{\text{mea}} \quad (3)$$

We can estimate the amount of $\text{CO}_2^+_{\text{NH}_4}$ by combining Eqs. (2-3):

$$\text{NH}_4^+_{\text{mea}} = 18 \times (2 \times \text{SO}_4^{2-}/96 + \text{NO}_3^-/62 + \text{Cl}^-/35.5 + \text{CO}_2^+_{\text{NH}_4}/44) \quad (4)$$

$$\text{So, } \text{CO}_2^+_{\text{NH}_4} = (\text{NH}_4^+_{\text{mea}} / 18 - 2 \times \text{SO}_4^{2-} / 96 - \text{NO}_3^- / 62 - \text{Cl}^- / 35.5) \times 44 \quad (5)$$

The time series of estimated $\text{CO}_2^+_{\text{NH}_4}$ over each experiments is shown in Fig. 3. On average, the $\text{CO}_2^+_{\text{NH}_4}$ concentration required to explain the observed ammonium concentrations was 48.6 times lower for seeded experiments than for the nucleation experiments. Fig. 3 indicates that organic acids participated in reacting with NH_3 much earlier in the low- NO_x test (red and green mark) than in the high- NO_x experiment (blue mark). Based on the time series in Fig. 3, in the high- NO_x test, the time at which the organic acids started to play a role in ammonium formation was delayed by 31 and 100 minutes for the nucleated and seeded experiments, respectively. This observation can be associated to the formation of nitric acid (HNO_3) from photooxidation of NO_x compounds in the high- NO_x conditions. The reaction of HNO_3 and NH_3 takes precedence over the reaction between organic acids and NH_3 . In the low- NO_x test, a time lag for $\text{CO}_2^+_{\text{NH}_4}$ is also observed, especially in the seeded SOA experiments. The delay is caused by the effect of nitric acid arising from the background NO_x photooxidation. In addition, the slower formation of particulate organic acids makes their reaction with ammonia delayed in the seeded experiments compared to the nucleated cases. In general, the required $\text{CO}_2^+_{\text{NH}_4}$ accounted for the 27.0 ± 3.1 % of total CO_2^+ mass in the nucleation experiments and 18.7 ± 6.0 % in the seeded SOA experiments. The amount of $\text{CO}_2^+_{\text{NH}_4}$ is assumed to be related to: (1) the photooxidation and ozonolysis capacity in our chamber, which determines the amount of particulate organic acids (total CO_2^+), (2) the relative amount of low- and high- molecule weight organic acids and their physicochemical properties, (3) the relative amount of inorganic and organic acids, and (4) the relative amount of ammonia and total acids, which needs to be thoroughly characterized in the future study.

To further verify our conclusion that organic acids are the drivers of the ammonium formation, we explored the size distribution of organic acids (represented by CO_2^+ ion), ammonium and nitrate at the end of nucleation experiments (Fig. 4). The mode diameters of the three species, determined by performing log-normal fitting on the size distributions, are listed in Table S1. The mode diameter of CO_2^+ ion is about 5-13 nm greater than that of ammonium and nitrate in the three individual experiments. The slight difference in two species mode diameters might be associated with the lower evaporation rate of organic CO_2^+ than ammonium ions on the AMS vaporizer. Anyhow, the similarity in the mode diameters and size distributions of three chemical species suggests that they are internally mixed in the physical phase and are originated from the similar formation sources.

3.2.2 Connecting gas compounds by PTRMS to ammonium ion by AMS

To investigate the organic acid species which may potentially contribute to the ammonium formation in our experiments, we compared the gas phase organic acids to the particle phase ammonium attributed to the neutralization by organic acids ($\text{NH}_4^+_{\text{orgacid}}$, defined as the difference between $\text{NH}_4^+_{\text{mea}}$ and $\text{NH}_4^+_{\text{pre}}$). We first examined the organic monocarboxylic acids. Fig. 5(a1) shows the correlation between the concentration of butyric acid ($\text{C}_4\text{H}_8\text{O}_2$) (or its isomer) measured by PTRMS and the $\text{NH}_4^+_{\text{orgacid}}$ calculated from AMS measurement in the nucleated SOA experiments. The excellent linear correlation (coefficient $R^2 \approx 0.88-0.95$) of these two species implies that the formation of butyric acid is associated with the ammonium formation. In addition, we also identified a molecular ion $\text{C}_4\text{H}_8\text{O}_2^+$ in the AMS mass spectrum, which also shows an excellent correlation to the formed ammonium (Fig. 5c1) and thus indicates its simultaneous formation with ammonium ion. The simultaneous observation of the same ion molecules in both gas and particle phases gives us confidence to speculate that this ion is derived from gas-phase butyric acid. The observation suggests a reaction between butyric acid and NH_3 which enables the production of ammonium butyrate salts ($\text{NH}_4\text{C}_4\text{H}_7\text{O}_2$). The formation of the salts favors the condensation of butyric acid on the

particle phase contributing to the observed ammonium ions. Although we can't overlook the fact that ammonium butyrate is severely fragmented inside AMS, it is believed that the detected $C_4H_8O_2^+$ ion signal is a residual of parental molecule. In the same way, we also explored pentanoic acid ($C_5H_{10}O_2$) (or its isomer) (Fig. 3b1 and 3d1) and obtained similar results as with butyric acid.

A similar comparison has also been made in the seeded SOA experiments (Fig. 5a2-d2). We have observed a good correlation of butyric acid to the organic acid-driven ammonium ($NH_4^{+,orgacid}$) ($r^2 = 0.76-0.83$), and a moderate correlation of pentanoic acid to $NH_4^{+,orgacid}$ ($r^2 = 0.27-0.62$). Compared with the nucleation experiments, the correlation relationship in the seeded SOA experiments is worse, mainly because the signals in both PTRMS and AMS measurements are weak in the seeded SOA experiments. Similar to the observation in the nucleated SOA experiments, the ion $C_4H_8O_2^+$ in AMS mass spectrum was also identified to correlate to $NH_4^{+,orgacid}$. The ion has the same molecular formula as butyric acid. The concentration of $C_5H_{10}O_2^+$ of the same molecular formula as pentanoic acid was in a concentration level of $10^{-3} \text{ ug m}^{-3}$ (Fig. 5d2). Such signal is at the same level as the background noise in AMS measurement, making it challenging to correlate with $NH_4^{+,orgacid}$. In general, the results in the seeded SOA experiments are consistent with those in the nucleated SOA experiments, confirming the role of butyric and pentanoic acids in the formation of ammonium salt.

We also extend our study to other four types of gas phase organic acids measured by the PTR-MS. Fig. 6 shows the relationship of the formic acid, acetic acid, propionic acid and pinonic acid measured in the gas phase with the ammonium ion measured by AMS. Surprisingly, the gas phase formic acid, acetic acid and propionic acid were in good agreement with particle phase ammonium concentration with excellent linear correlation coefficients $r^2 \geq 0.95$ in the nucleated SOA experiments and $r^2 \geq 0.72$ in the seeded SOA experiments. The good correlation of gas phase organic acids with the particle phase ammonium salt suggests that these acids played a role in the formation of ammonium. Formic acid and acetic acid are the most abundant organic monoacids in the atmosphere (Chebbi and Carlier, 1996), whose one significant source is from photooxidation of α -pinene and other alkenes and terpenes (Friedman and Farmer, 2018). Propionic and butyric acids are also other important organic acids in this process (Nah et al., 2018a; Nah et al., 2018b; Chebbi and Carlier, 1996). The formation mechanisms of these low molecular weight organic acids are unclear in the photochemical reaction of α -pinene (e.g. Friedman and Farmer, 2018). It can be speculated that they were produced as products of stabilized Criegee intermediate associated with ozonolysis of α -pinene and double carbon bond-containing products (Jacob and Wofsy, 1988; Orzechowska and Paulson, 2005). The statement is verified by the fact that we have observed the highest O_3 formation at the highest NO_x/VOC_x initial input in E0314 (Fig. S4). Therefore, in E0314 we have also observed higher concentrations of C1-C5 monoacids corresponding to higher ozonolysis reactivity (Fig. S5). These low molecular weight monoacids have a vapor pressure of $6.8 \sim 8.1 \text{ } \mu\text{g m}^{-3} (\log_{10}(C^*))$ (Friedman and Farmer, 2018), and are generally considered to be too volatile to be distributed substantially to the particle phase. However, their presence in the aerosol particles is ubiquitous in various areas over the world, although the levels of these monoacids are one or two order of magnitude lower than those in the gas phase (Nah et al., 2018a; Fisseha et al., 2004; Chebbi and Carlier, 1996). A study conducted in a forest-agriculture area in Atlanta showed that the acetic acid and formic acid are the second and third richest water-soluble organic acids in the particle phase and their molar fractions to the total individual acid concentrations in the particulate phase were $5.8 \pm 5.0\%$ and $3.6 \pm 3.6\%$, respectively (Nah et al., 2018a). In addition, the presence of NH_3 as a strong base facilitates the shift of the equilibrium of these monocarboxylic acids and NH_3 to the particle phase (Barsanti et al., 2009). Under a base environment, a higher molar fraction of formic acid and acetic acid has also been observed in the particle phase (Nah et al., 2018a). These organic acids could exist in the particle phase in chemical forms of

ammonium formate, ammonium acetate and ammonium propionate salts (Barsanti et al., 2009; Smith et al., 2008; Becker and Davidson, 1963), and thus contribute to the observed ammonium ions.

In an obvious contrast to the observation of C1-C5 monoacids above, the pinonic acids didn't show a linear correlation with the ammonium at the later stage of the nucleated SOA experiments (pink region, Fig. 6d1), and there was no connection in the seeded SOA experiments. Pinonic acid is a typical first-generation marker product in the photooxidation reaction of α -pinene (Hallquist et al., 2009). Na et al. (2007) reported that the addition of NH_3 to the aerosolized pinonic acid in an environmental reactor led to dramatic increase in both number and volume concentrations of pinonic acid. They attributed the observation to the reaction between NH_3 and pinonic acids. However, the lack of positive correlation between the pinonic acid and ammoniums can be caused by several reasons in this study. Firstly, the vapor pressure of pinonic acid is about 4 ~ 5 orders of magnitude lower than those of C1-C5 monoacids (Jimenez et al., 2009), so pinonic acid can condense on the particle phase independent on NH_3 . Meanwhile, the concentration of pinonic acid accounts for less than 1% of the total mono-acids based on the PTR-MS results. Hence, the contribution of pinonic acid to the formed ammonium was estimated to be less than 1%, considering that the acidity strength of pinonic acid is similar to other C1-C5 monoacids (acid dissociation constant (pKa) of pinonic acid is 4.8 (Howell and Fisher, 1958), close to the pKa values of formic acid (3.75), acetic acid (4.75), propionic acid (4.86), butyric acid (4.83) and pentanoic acid (4.84) (Lide, 2007)). Therefore, after α -pinene was completely consumed and pinonic acid formation had ceased in the gas phase (pink region, Fig. 6), the rapid condensation of pinonic acid on the particle phase or chamber wall causes a non-linear correlation of pinonic acid to ammonium. Secondly, the gas-phase pinonic acids could be further reacted away by OH radicals, which also contributes to the non-linear observation. Field experiment has shown the relatively low atmospheric PM concentrations of pinonic acid measured in summer because of the consumption of pinonic acid by OH radicals (Szmigielski et al., 2007).

Analogously, the scattering plots for the gas-phase organic dicarboxylic acids and particle-phase ammonium salts are shown in Fig. 7. We chose malonic acid and succinic acid as representative organic diacids, two of the most abundant dicarboxylic acids measured in the atmospheric aerosols (Chebbi and Carlier, 1996). The nice correlations of the gas-phase malonic and succinic acids to the particle-phase ammonium study suggest that diacids contribute to the formation of ammonium in both nucleated and seeded SOA experiments. It also indicates the continuous formation of those diacids at the later stage of photooxidation in nucleated and seeded experiments. The formation mechanism of these low molecule-weight diacids remains unclear for us and might be similar to the C1-C5 monoacids.

Our results qualitatively demonstrate that in the photooxidation of α -pinene, the presence of NH_3 drives the gas-phase mono- and dicarboxylic acids to the particle phase and promotes the SOA mass concentration ~~in the CCN size~~. Previous studies have shown that the presence of NH_3 can significantly enhance SOA formation from the α -pinene/ozone/photooxidation system because of the interaction of NH_3 with gas-phase organic acids (Na et al., 2007; Babar et al., 2017), which is consistent with our results. Carboxylic acid is one of the key species in determining SOA physico-chemical property. Our results may prompt us to reconsider the pathway of gas phase organic acids involving in SOA formation in the atmosphere, whether they directly participate partitioning between the gas and particle phases, or they undergo secondary conversion via reaction with NH_3 before they condensate on the particle phase in the atmosphere. After SOA are formed, carbonyl group of chemical compounds in SOA particles can also uptake NH_3 heterogeneously to form nitrogen-containing compounds (Zhu et al., 2018; Liu et al., 2015; Updyke et al., 2012; Dinar et al., 2007) and organic ammonium salts (Schlag et al., 2017). In this study the N:C ratios measured by AMS were nearly constant or slightly increasing in two sets of experiments (Fig. S6), which could suggesting the formation

of nitrogen-containing compounds via carbonyl-NH₃ heterogeneous reaction. However, we can't rule out that NO_x also participated in the formation of SOA, resulting in an increase in the N:C ratios. The interaction of NH₃ and SOA affects the cloud condensation nuclei (CCN) and hygroscopic growth of SOA particles and may have a potential impact on climate change (Dinar et al., 2007).

3.3 Ammonium relevant to PMF-solved SOA component

PMF analysis on the high-resolution organic mass spectra resolved the organic component to two factors in the nucleation experiments: MO-OOA (more oxidized oxygenated organic aerosol) and LO-OOA (less oxidized OOA). The oxidation level of MO-OOA is represented by an O:C ratio of 0.47 in this study, which is close to the value of 0.48 for MO-OOA determined in New York City in Summer (Sun et al., 2012). Its time series shows a good correlation to the measured gas-phase monocarboxylic and di- acids such as butyric and succinic acids. The second factor LO-OOA is featured by an O:C ratio of 0.38. Its time series is related to the gas-phase oxidant products such as pinonaldehyde. The mass spectra profiles and time series of the two factors are shown in Fig.S7. In this study, the formation of ammonium salts is consistent with MO-OOA factor and also organic mono- and di- acids (Fig. 8). A higher oxidized organic factor is usually associated with the formation of organic mono- and di-acids. These results further suggest that the ammonium has a close relation to the organic acids. Our observation is in agreement with the study by Schlag et al. (2017) where they showed by field data that NH₄ is associated with a more oxygenated organic aerosol factor.

4 Conclusions

The SOA experiments were carried out from photooxidation reaction of α -pinene in the presence of NH₃ in a 29m³ indoor simulation chamber. Experiments were designed for SOA formation in the presence of ammonium sulfate seeds and at the absence of seed aerosols. The chemical composition and time-series of compounds in the gas- and particle-phase were characterized by an on-line high-resolution time-of-flight proton transfer reaction mass spectrometer (PTRMS) and a high-resolution time-of-flight aerosol mass spectrometer (AMS), respectively.

After the precursor α -pinene was consumed in the chamber, the mass concentration of organic aerosol ~~in CCN-size~~ was decreased because of aerosol wall deposition or evaporation, the ammonium concentration was still rising, suggesting the continuous formation of ammonium. AMS results showed that organic acids were required to neutralize the observed ammonium salt. The CO₂⁺ ion was selected to represent organic acids. The amount of CO₂⁺ required for neutralizing ammonium accounted for the 27.0 ± 3.1 % of total CO₂⁺ mass in the nucleated SOA experiments, and 18.7 ± 6.0 % in the seeded SOA experiments. The good correlation of organic monocarboxylic acids (such as formic acid, acetic acid, propionic acid, butyric acid and pentanoic acid (or their isomers)) in the gas phase to the ammonium salts further qualitatively confirms an effective role of organic acids for the ammonium formation. The same conclusion is also applied to the organic dicarboxylic acids such as malonic and succinic acids. In addition, the formed ammonium salts correlated well to the more-oxidized oxygenated organic aerosol (MO-OOA), which is consistent with the conclusion that organic acids contributed to the observed particulate ammonium.

Our work firmly shows the direct contribution of NH₃ to the ~~CCN-size~~ SOA formation through the organic acids-base reaction. The increase in SOA mass and the change of chemical composition due to NH₃-SOA interaction could change the hygroscopicity, CCN ability and optical property of aerosol particles, which may alter the aerosol impact on climate change and need to be studied in the future.

Code/Data availability.

The data included in this paper can be obtained by contacting the authors.

Author contribution.

LH, EK, AL and AV designed and conducted the experiments. LH and EK performed the data analysis with contributions by DW and AV. LH and AV wrote the paper with contributions from all co-authors.

Competing interests.

The authors declare that they have no conflict of interest.

Acknowledgements

The authors thank Mr. Ilkka Summanen for assisting the chamber experiments. This work was supported by European Union's Horizon 2020 research and innovation program through the EUROCHAMP-2020 Infrastructure Activity (grant no. 730997) and the Academy of Finland Center of Excellence programme (grant no 307331).

References

- Alfarra, R., Insights Into Atmospheric Organic Aerosols Using An Aerosol Mass Spectrometer, Ph.D. Thesis at the University of Manchester, 2004.
- Babar, Z.B., Park, J., and Lim, H.: Influence of NH₃ on secondary organic aerosols from the ozonolysis and photooxidation of α -pinene in a flow reactor, *Atmos. Environ.*, 164, 71-84, 2017.
- Barsanti, K. C., McMurry, P. H., and Smith, J. N.: The potential contribution of organic salts to new particle growth, *Atmos. Chem. Phys.*, 9, 2949-2957, <https://doi.org/10.5194/acp-9-2949-2009>, 2009.
- Battye, W., Aneja, V.P., and Roelle, P.A.: Evaluation and improvement of ammonia emission inventories, *Atmos. Environ.*, 37, 3873-3883, 2003.
- Becker, B., and Davidson, A.W.: the systems formic acid-ammonia and propionic acid-ammonia, *J. Am. Chem. Soc.*, 85(2), 57-159, <https://doi.org/10.1021/ja00885a010>, 1963.
- Bianchi, F., Trostl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Herrmann, E., Adamov, A., Bukowiecki, N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kurten, A., Manninen, H. E., Munch, S., Perakyla, O., Petaja, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J., Worsnop, D. R., Kulmala, M., Dommen, J., and Baltensperger, U.: New particle formation in the free troposphere: A question of chemistry and timing, *Science*, 352, 1109-1112, 2016.
- Bones, D.L., Henricksen, D.K., Mang, S.A., Sams, M., Bateman, A.P., Nguyen, T.B., Cooper, W.J., and Nizkorodov, S.A.: Appearance of strong absorbers and fluorophores in limonene-O₃ secondary organic aerosol due to NH₄⁺-mediated chemical aging over long time scales, *J. Geophys. Res.*, 115, D05203, doi: 10.1029/2009JD012864, 2010.
- Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and Microphysical Characterization of Ambient Aerosols with the Aerodyne Aerosol Mass Spectrometer, *Mass Spectrometry Reviews*, 26, 185– 222, 2007.
- Chebbi, A. and Carlier, P.: Carboxylic acids in the troposphere, occurrence, sources, and sinks: a review, *Atmos. Environ.*, 30, 4233-4249, 1996.

Chen, T., Liu, Y., Ma, Q., Chu, B., Zhang, P., Liu, C., Liu, J., and He, H.: Significant source of secondary aerosol: formation from gasoline evaporative emissions in the presence of SO₂ and NH₃, *Atmos. Chem. Phys.*, 19, 8063–8081, <https://doi.org/10.5194/acp-19-8063-2019>, 2019.

DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K., Worsnop, D. R. and Jimenez, J. L.: Field-Deployable, High-Resolution, Time-of-Flight Aerosol Mass Spectrometer, *Anal. Chem.*, 78, 8281–8289, 2006.

Dinar, E., Anttila, T., and Rudich, Y.: CCN activity and hygroscopic growth of organic aerosols following reactive uptake of ammonia, *Environ. Sci. Technol.*, 42, 793-799, 2008.

Fisseha, R., Dommen, J., Sax, M., Paulsen, D., Kalberer, M., Maurer, R., Höfler, F., Weingartner, E., and Baltensperger, U.: Identification of organic acids in secondary organic aerosol and the corresponding gas phase from chamber experiments, *Anal. Chem.*, 76, 6535-6540, 2004.

Friedman, B. and Farmer, D.K.: SOA and gas phase organic acid yields from the sequential photooxidation of seven monoterpenes, *Atmos. Environ.*, 187, 335-345, 2018.

Grönroos, J., Mattila, P., Regina, K., Nousiainen, J., Perälä, P., Saarinen, K., and Mikkola-Pusa, J.: Development of the ammonia emission inventory in Finland. Revised model for agriculture, Finnish Environment Institute, Helsinki, 2009.

Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155-5236, <https://doi.org/10.5194/acp-9-5155-2009>, 2009.

Hansel, A., Jordan, A., Holzinger, R., Prazeller, P., Vogel, W., and Lindinger, W.: Proton-Transfer Reaction Mass-Spectrometry – Online Trace Gas-Analysis at the Ppb Level, *Int. J. Mass Spectrom.*, 149, 609-619, [DOI: 10.1016/0168-1176\(95\)04294-U](https://doi.org/10.1016/0168-1176(95)04294-U), 1995.

Huang, X., Song, Y., Li, M.M., Li, J.F., Huo, Q., Cai, X.H., Zhu, T., Hu, M., and Zhang, H.S.: A high-resolution ammonia emission inventory in China, *Global Biogeochemical Cycles*, 26, GB1030, [DOI: 10.1029/2011GB004161](https://doi.org/10.1029/2011GB004161), 2012.

Howell, H. and Fisher, G.S.: The dissociation constants of some of the terpene acids, *J. Am. Chem. Soc.*, 80(23), 6316-6319, 1958.

Horne, J.R., Zhu, S., Montoya-Aguilera, J., Hinks, M.L., Wingen, L.M., Nizkorodov, S.A. and Dabdub, D.: Reactive uptake of ammonia by secondary organic aerosols: Implications for air quality, *Atmos. Environ.*, 189, 1-8, 2018.

Huang M.Q., Xu, J., Cai, S.Y., Liu, X.Q., Zhao, W.X., Hu, C.J., Gu, X.J., Fang, L., and Zhang, W.J.: Characterization of brown carbon constituents of benzene secondary organic aerosol aged by ammonia, *J. Atmos. Chem.*, 75, 205-218, 2018.

IPCC 2013. Climate change 2013: The physical science basis. Intergovernmental panel on Climate Change, Cambridge University Press, New York.

Jayne, J. T., Leard, D. C., Zhang, X. F., Davidovits, P., Smith, K. A., Kolb, C. E., and Worsnop, D. R.: Development of an Aerosol Mass Spectrometer for size and composition analysis of submicron particles, *Aerosol Sci. Technol.*, 33, 49–70, 2000.

- Jacob, D. J. and Wofsy, S. C.: Photochemistry of biogenic emissions over the Amazon forest, *J. Geophys. Res.*, 93, 1477–1486, doi:10.1029/JD093iD02p01477, 1988.
- Jimenez, J.L., Canagaratna, M.R., Donahue, N.M., Prevot, A.S.H., Zhang, Q., Kroll, J.H., DeCarlo, P.F., Allan, J.D., Coe, H., Ng, N.L., Aiken, A.C., Docherty, K.S., Ulbrich, I.M., Grieshop, A.P., Robinson, A.L., Duplissy, J., Smith, J.D., Wilson, K.R., Lanz, V.A., Hueglin, C., Sun, Y.L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J.M., Collins, D. R., Cubison, M. J., Dunlea, E.J., Huffman, J. A., Onasch, T.B., Alfarra, M.R., Williams, P.I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J.Y., Zhang, Y.M., Dzepina, K., Kimmel, J.R., Sueper, D., Jayne, J.T., Herndon, S.C., Trimborn, A.M., Williams, L.R., Wood, E.C., Middlebrook, A.M., Kolb, C.E., Baltensperger, U., and Worsnop, D.R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, 2009.
- Jordan, A., Haidacher, S., Hanel, G., Hartungen, E., Mark, L., Seehauser, H., Schottkowsky, R., Sulzer, P., and Mark, T. D.: A high resolution and high sensitivity proton-transfer-reaction time-of-flight mass spectrometer (PTR-TOF-MS), *Int. J. Mass Spectrom.*, 286, 122–128, 10.1016/j.ijms.2009.07.005, 2009.
- Jokinen, T., Sipila, M., Kontkanen, J., Vakkari, V., Tisler, P., Duplissy, E. M., Junninen, H., Kangasluoma, J., Manninen, H. E., Petaja, T., Kulmala, M., Worsnop, D. R., Kirkby, J., Virkkula, A., and Kerminen, V. M.: Ion-induced sulfuric acid-ammonia nucleation drives particle formation in coastal Antarctica, *Sci. Adv.*, 4, eaat9744, DOI: 10.1126/sciadv.aat9744, 2018.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, <https://doi.org/10.5194/acp-5-1053-2005>, 2005.
- Kari, E., Hao, L., Ylisirniö, A., Buchholz, A., Leskinen, A., Yli-Pirilä, P., Nuutinen, I., Kuusalo, K., Jokiniemi, J., Faiola, C., Schobesberger, S., and Virtanen, A.: Potential dual effect of anthropogenic emissions on the formation of biogenic secondary organic aerosol (BSOA), *Atmos. Chem. Phys.*, 19, 15651–15671, <https://doi.org/10.5194/acp-19-15651-2019>, 2019a.
- Kari, E., Faiola, C.L., Isokääntä, S., Miettinen, P., Yli-Pirilä, P., Buchholz, A., Kivimäenpää, M., Mikkonen, S., Holopainen, J.K., and Virtanen, A.: Time-resolved characterization of biotic stress emissions from Scots pines being fed upon by the pine weevil by means of PTR-ToF-MS, *Boreal Environ. Res.*, 24, 25–49, 2019b.
- Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L., Kuurten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., J. H. Seinfeld, Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429–433, 2011.
- Kourtchev, I., Giorio, C., Manninen, A., Wilson, E., Mahon, B., Aalto, J., Kajos, M., Venables, D., Ruuskanen, T., Levula, J., Loponen, M., Connors, S., Harris, N., Zhao, D., Kiendler-Scharr, A., Mentel, T., Rudich, Y., Hullquist,

- [M., Doussin J., Maenhaut, W., Back, J., Petaja, T., Wenger, J., Kulmala, M. and Kalberer, M.: Enhanced volatile organic compounds emissions and organic aerosol mass increase the oligomer content of atmospheric aerosols, *Sci. Reports*, 6, 35038, 2016.](#)
- Kulmala, M., Pirjola, L., and Mäkelä, J.M.: Stable sulphate clusters as a source of new atmospheric particles, *Nature*, 404, 66-69, 2000.
- Kurten, T., Torpo, L., Ding, C.G., Vehkamäki, H., Sundberg, M. R., Laasonen, K., and Kulmala, M.: A density functional study on water-sulfuric acid-ammonia clusters and implications for atmospheric cluster formation, *J. Geophys. Res.-Atmospheres*, 112, D4, DOI: 10.1029/2006JD007391, 2007.
- Lehtipalo, K., Yan, C., Dada, L., Bianchi, F., Xiao, M., Wagner, R., Stolzenburg, D., Ahonen, L. R., Amorim, A., Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A., Bernhammer, A.-K., Breitenlechner, M., Brilke, S., Buchholz, A., Mazon, S. B., Chen, D., Chen, X., Dias, A., Dommen, J., Draper, D. C., Duplissy, J., Ehn, M., Finkenzeller, H., Fischer, L., Frege, C., Fuchs, C., Garmash, O., Gordon, H., Hakala, J., He, X., Heikkinen, L., Heinritzi, M., Helm, J. C., Hofbauer, V., Hoyle, C. R., Jokinen, T., Kangasluoma, J., Kerminen, V.-M., Kim, C., Kirkby, J., Kontkanen, J., Kürten, A., Lawler, M. J., Mai, H., Mathot, S., Mauldin, R. L., Molteni, U., Nichman, L., Nie, W., Nieminen, T., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Piel, F., Pospisilova, V., Quéléver, L. L. J., Rissanen, M. P., Rose, C., Sarnela, N., Schallhart, S., Schuchmann, S., Sengupta, K., Simon, M., Sipilä, M., Tauber, C., Tomé, A., Tröstl, J., Väisänen, O., Vogel, A. L., Volkamer, R., Wagner, A. C., Wang, M., Weitz, L., Wimmer, D., Ye, P., Ylisirniö, A., Zha, Q., Carslaw, K. S., Curtius, J., Donahue, N. M., Flagan, R. C., Hansel, A., Riipinen, I., Virtanen, A., Winkler, P. M., Baltensperger, U., Kulmala, M., and Worsnop, D. R.: Multicomponent new particle formation from sulfuric acid, ammonia, and biogenic vapors, *Sci. Adv.*, 4, eaau5363, <https://doi.org/10.1126/sciadv.aau5363>, 2018.
- Leskinen, A., Yli-Pirilä, P., Kuusalo, K., Sippula, O., Jalava, P., Hirvonen, M.-R., Jokiniemi, J., Virtanen, A., Komppula, M., and Lehtinen, K. E. J.: Characterization and testing of a new environmental chamber, *Atmos. Meas. Tech.*, 8, 2267-2278, <https://doi.org/10.5194/amt-8-2267-2015>, 2015.
- Lide, D. R.: CRC handbook of chemistry and physics, 88th edition, CRC press, 2007.
- Liu, Y., Liggitto, J., Staebler, R., and Li, S.-M.: Reactive uptake of ammonia to secondary organic aerosols: kinetics of organonitrogen formation, *Atmos. Chem. Phys.*, 15, 13569-13584, <https://doi.org/10.5194/acp-15-13569-2015>, 2015.
- Li, Z., Nizkorodov, S. A., Chen, H., Lu, X., Yang, X., and Chen, J.: Nitrogen-containing secondary organic aerosol formation by acrolein reaction with ammonia/ammonium, *Atmos. Chem. Phys.*, 19, 1343-1356, <https://doi.org/10.5194/acp-19-1343-2019>, 2019.
- Liu, T. Y., Wang X.M., Deng, W., Zhang, Y.L., Chu, B.W., Ding, X., Hu, Q.H., He, H., and Hao, J.M.: Role of ammonia in forming secondary aerosols from gasoline vehicle exhaust, *Science China Chemistry*, 58, 1377-1384, 2015.
- Mensah, A. A., Holzinger, R., Otjes, R., Trimborn, A., Mentel, Th. F., ten Brink, H., Henzing, B., and Kiendler-Scharr, A.: Aerosol chemical composition at Cabauw, The Netherlands as observed in two intensive periods in May 2008 and March 2009, *Atmos. Chem. Phys.*, 12, 4723-4742, <https://doi.org/10.5194/acp-12-4723-2012>, 2012.
- Nah, T., Guo, H., Sullivan, A. P., Chen, Y., Tanner, D. J., Nenes, A., Russell, A., Ng, N. L., Huey, L. G., and Weber, R. J.: Characterization of aerosol composition, aerosol acidity, and organic acid partitioning at an agriculturally

- intensive rural southeastern US site, *Atmos. Chem. Phys.*, 18, 11471-11491, <https://doi.org/10.5194/acp-18-11471-2018>, 2018a.
- Nah, T., Ji, Y., Tanner, D. J., Guo, H., Sullivan, A. P., Ng, N. L., Weber, R. J., and Huey, L. G.: Real-time measurements of gas-phase organic acids using SF_6^- chemical ionization mass spectrometry, *Atmos. Meas. Tech.*, 11, 5087-5104, <https://doi.org/10.5194/amt-11-5087-2018>, 2018b.
- Na, K., Song, C., Switzer, C., and Cocker, D.R.: Effect of ammonia on secondary organic aerosol formation from α -pinene ozonolysis in dry and humid conditions, *Environ. Sci. Technol.*, 41, 6096-6102, 2007.
- Odum, J.R., Hoffmann, T., Bowman, F.M., Collins, D., Flagan, R.C., and Seinfeld, J.H.: Gas/particle partitioning and secondary organic aerosol yields, *Environ. Sci. Technol.*, 30, 2580-2585, 1996.
- Orzechowska, G.E., and Paulson, S.E.: Photochemical sources of organic acids. 1. Reaction of ozone with isoprene, propene, and 2-butenes under day and humid conditions using SPME, *J. Phys. Chem. A*, 109, 5358-5365, 2005.
- Paatero, P. and Tapper, U.: Positive Matrix Factorization: a non-negative factor model with optimal utilization of error estimates of data values. *Environmetrics*, 5, 111–126, 1994.
- Paciga, A.L., Riipinen, I., and Pandis, S.N.: Effect of ammonia on the volatility of organic diacids, *Environ. Sci. Technol.*, 48, 13769-13775, 2014.
- Reddy L. N. Yatawelli, R.L., Mohr, C., Stark, H., Day, D.A., Thompson, S.L., Lopez-Hilfiker, F.D., Campuzano-Jost, P., Palm, B.B., Vogel, A.L., Hoffmann, T., Heikkinen, L., Äijälä, M., Ng, N.L., Kimmel, J.R., Canagaratna, M.R., Ehn, M., Junninen, H., Cubison, M.J., Tuukka Petäjä, T., Kulmala, M., Jayne, J.T., Worsnop, D.R., and Jimenez, J. L.: Estimating the contribution of organic acids to northern hemispheric continental organic aerosol, *Geophys. Res. Lett.*, 42, 6084–6090, doi: 10.1002/2015GL064650, 2015.
- Rubach, F.: Aerosol processes in the planetary boundary layer: High resolution aerosol mass spectrometry on a Zeppelin NT airship, Forschungszentrum Jülich GmbH, 2013.
- Salonen, H., Pasanen, A., Lappalainen, S.K., Riuttala, H.M., Tuomi, T.M., Pasanen, P.O., Back, B.C. and Reijula, K.E.: Airborne concentrations of volatile organic compounds, formaldehyde and ammonia in Finnish office buildings with suspected indoor air problems, *Journal of Occupational and Environmental Hygiene*, 6, 200-209, 2009.
- Schlag, P., Rubach, F., Mental, T.F., Reimer, D., Canonaco, F., Henzing, J.S., Moerman, M., Otjes, R., Prevot, A.S.H., Rohrer, F., Rosati, B., Tillmann, R., Weingartner, E., and Kiendler-Scharr, A.: Ambient and laboratory observations of organic ammonium salts in PM₁, *Faraday Discussions*, 200, 331-351, 2017.
- Seinfeld, J. H. and Pandis, S. N.: Atmospheric chemistry and physics: from air pollution to climate change, John Wiley & Sons, New York, 2016.
- Smith, J.N., Dunn, M.J., VanReken, T.M., Iida, K., Stolzenburg, M.R., McMurry, P.h., and Hyey, L.G.: Chemical composition of atmospheric nanoparticles formed from nucleation in Tecamac, Mexico: Evidence for an important role for organic species in nanoparticle growth, *Geophys. Res. Lett.*, 35, 104808, doi: 10.1029/2007GL032523, 2008.
- Sun, Y. L., Zhang, Q., Schwab, J. J., Yang, T., Ng, N. L., and Demerjian, K. L.: Factor analysis of combined organic and inorganic aerosol mass spectra from high resolution aerosol mass spectrometer measurements, *Atmos. Chem. Phys.*, 12, 8537-8551, <https://doi.org/10.5194/acp-12-8537-2012>, 2012.
- Szmigielski, R., Surratt, J.D., Gomez-Gonzalez, Y., Van der Veken, P., Kourtchev, I., Vermeylen, R., Blockhuys, F., Jaoui, M., Kleindienst, T.E., Lewandowski, M., Offenberg, J.H., Edney, E.O., Seinfeld, J.H., Maenhaut, W., and

- 553 Claeys, M.: 3-methyl-1,2,3-butanetricarboxylic acid: an atmospheric tracer for terpene secondary organic aerosol,
554 Geophys. Res. Lett., 34, L24811, doi: 10.1029/2007GL031338, 2007.
- 555 Takegawa, N., Miyakawa, T., Kawamura, K., and Kondo, Y.: Contribution of Selected Dicarboxylic and ω -
556 Oxocarboxylic Acids in Ambient Aerosol to the m/z 44 Signal of an Aerodyne Aerosol Mass Spectrometer,
557 Aerosol Sci. Technol., 41, 418-437, DOI: 10.1080/02786820701203215, 2007.
- 558 Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic
559 components from Positive Matrix Factorization of aerosol mass spectrometric data, Atmos. Chem. Phys., 9, 2891-
560 2918, <https://doi.org/10.5194/acp-9-2891-2009>, 2009.
- 561 Updyke, K.M., Nguyen, T.B., and Nizkorodov, S.A.: Formation of brown carbon via reactions of ammonia with
562 secondary organic aerosols from biogenic and anthropogenic precursors, Atmos. Environ., 63, 22-31, 2012.
- 563 Wang, Y., Liu, P.F., Li, Y.J., Bateman, A.P., Martin, S.T., and Hung, H.M.: The reactivity of toluene-derived
564 secondary organic material with ammonia and influence of water vapor, J. Phys. Chem. A., 122, 7739-7747, 2018.
- 565 Zhang, Q., Alfarra, M. R., Worsnop, D. R., Allan, J. D., Coe, H., Canagaratna, M. R., and Jimenez, J. L.:
566 Deconvolution and Quantification of Hydrocarbon-Like and Oxygenated Organic Aerosols Based on Aerosol
567 Mass Spectrometry, Environ. Sci. Technol., 39, 4938–4952, doi: 10.1021/es048568l, 2005.
- 568 Zhang, Q., Stanier, C. O., Canagaratna, M. R., Jayne, J. T., Worsnop, D. R., Pandis, S. N., and Jimenez, J. L.:
569 Insights into the chemistry of new particle formation and growth events in Pittsburgh based on aerosol mass
570 spectrometry, Environ. Sci. Technol., 38, 4797-4809, 2004.
- 571 Zhu, S., Horne, J. R., Montoya-Aguilera, J., Hinks, M. L., Nizkorodov, S. A., and Dabdub, D.: Modeling reactive
572 ammonia uptake by secondary organic aerosol in CMAQ: application to the continental US, Atmos. Chem. Phys.,
573 18, 3641-3657, <https://doi.org/10.5194/acp-18-3641-2018>, 2018.

Table 1 Initial experimental parameters and results in two sets of experiments

	Exp. ID	α -Pinene (ppb)	NO _x (ppb) ⁽¹⁾	R.H. (%)	Temp (°C)	AS seed surface ⁽¹⁾ ($\mu\text{m}^2/\text{cm}^3$)	OH exposure ($\#/\text{cm}^3 \text{ s}$)	$\Delta\text{NH}_4^{(2,3)}$ mass ($\mu\text{g m}^{-3}$)	SOA ⁽³⁾ mass ($\mu\text{g m}^{-3}$)
Nucleation	E0322	91.7	bg (0.5)	67.2	19.0	bg	1.2e11	1.2	391.8
SOA Exp.	E0326	100.9	bg (0.3)	50.5	21.4	bg	1.3e11	1.5	389.7
	E0327	107.2	13.3	51.4	21.5	bg	1.3e11	1.6	476.1
Seeded SOA	E0314	4.1	61.3	56.2	22.6	3.6e7	1.2e11	0.02	5.5
Exp.	E0315	4.2	bg (2.6)	57.2	22.1	4.0e7	1.2e11	0.02	7.5
	E0316	4.8	bg (1.1)	56.5	21.8	2.4e7	1.3e11	0.03	7.7

⁽¹⁾ bg=background concentration inside the chamber.

⁽²⁾ In the seeded SOA experiment, because of the presence of ammonium sulfate seeds, the maximum mass concentration of newly formed ammonium salt was estimated from the difference between $\text{NH}_4^+_{\text{pre}}$ and $\text{NH}_4^+_{\text{mea}}$, refer to the text for details.

⁽³⁾ The formed aerosol mass at maximum.




Figure 1. The mass concentrations of organics (left axis), ammonium (top panels, right axis), sulfate and nitrate (bottom panels) as a function of irradiation time in nucleation experiments (left panels) and in seeded SOA experiments (right panels). The irradiation time of 0 min marks the start of photooxidation reactions after UV lights were switched on.




Figure 2. The time series of measured and predicted ammonium in the nucleated SOA (top panels) and in the seeded SOA experiments (bottom panels).





Figure 3. The estimated amount of organic carboxylic acid needed to fully neutralize SOA particles ($\text{CO}_2^+_{\text{NH}_4}$). CO_2^+ group is chosen to represent organic acids. (A) nucleation experiments (B) seeded experiments. The time zero marks the beginning of photooxidation reactions when UV lights were switched on.

764
765
766



767
768 **Figure 4.** The averaged 30-minute size distribution of organic acids (represented by CO_2^+ ion measured by AMS),
769 nitrate and ammonium in the last 30 minutes prior to the end of the nucleated SOA experiments.

770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804




Figure 5. The comparison of measured organic monoacids in the gas phase by PTRMS and the ammonium attributed to the neutralization by organic acids ($\text{NH}_4^+, \text{orgacid}$) in the particle phase by AMS in the nucleated SOA (left four panels) and seeded SOA experiments (right four panels). Top panels: Linear correlation between butyric ($\text{C}_4\text{H}_8\text{O}_2$) and pentanoic ($\text{C}_5\text{H}_{10}\text{O}_2$) monoacids measured by PTRMS and the $\text{NH}_4^+, \text{orgacid}$ by AMS; Bottom panels: Linear correlation of two fragmental ions $\text{C}_4\text{H}_8\text{O}_2^+$ and $\text{C}_5\text{H}_{10}\text{O}_2^+$ to the $\text{NH}_4^+, \text{orgacid}$ by AMS. Note that the two fragmental ions have identical ion molecular formula to those of butyric and pentanoic acids..





Figure 6. Correlations of the ammonium attributed to the neutralization by organic acids ($\text{NH}_4^+, \text{orgacid}$) in the particle phase by AMS to the (a) formic acid, (b) acetic acid, (c) propionic acid, (d) pinonic acid in the gas phase measured by PTRMS in the nucleated SOA experiments (left four panels) and in the seeded SOA experiments (right four panels). The light pink area in panel d1 indicates a non-positive correlation regime between two species.


891
892
893
894



895
896 **Figure 7.** Correlations of the ammonium attributed to the neutralization by organic acids ($\text{NH}_4^+, \text{orgacid}$) in the particle
897 phase by AMS to two organic dicarboxylic acids: (a) malonic acid, (b) succinic acid in the gas phase measured by
898 PTRMS. Both of the diacids were not observed in the experiment E0322.

899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934

935
936
937
938
939
940
941
942



943
944
945
946
947

Figure 8. The relevance of particulate ammonium salt to the resolved MO-OOA component by PMF, and to the gas-phase butyric monocarboxylic acid and succinic diacid.